

Constraining the budget of NO_x and VOCs at a remote Tropical island using multi-platform observations and WRF-Chem model simulations

5 Catalina Poraicu¹, Jean-François Müller¹, Trissevgeni Stavrakou¹, Crist Amelynck^{1,2}, Bert W. D. Verreyken^{1,2,3,4}, Niels Schoon¹, Corinne Vigouroux¹, Nicolas Kumps¹, Jérôme Brioude³, Pierre Tulet^{3,5}, Camille Mouchel-Vallon^{5,6}

1 Royal Belgian Institute for Space Aeronomy (BIRA-IASB), Ringlaan 3, 1180 Brussels, Belgium

2 Department of Chemistry, Ghent University, 9000 Ghent, Belgium

10 3 Laboratoire de l'Atmosphère et des Cyclones (LACy), UMR 8105, CNRS, Université de La Réunion, Météo France, 97744 Saint-Denis, France

4 University of Liège - Gembloux Agro-Biotech, Biosystems Dynamics and Exchanges (BIODYNE), 8 Avenue de la faculté, 5030 Gembloux, Belgium

5 Laboratoire d'Aérodynamique (LAERO), UMR 5560, CNRS, Université Paul Sabatier, IRD, Toulouse, 31400, France

6 Barcelona Supercomputing Center (BSC), Barcelona, Spain

15 *Correspondence to:* Catalina Poraicu (catalina.poraicu@aeronomie.be)

Abstract. Volatile organic compounds (VOCs) act as precursors to ozone and secondary organic aerosols, which have significant health and environmental impacts, and they can reduce the atmospheric oxidative capacity. However, their budget remains poorly quantified, especially over remote areas such as the Tropical oceans. Here, we present high-resolution simulations of atmospheric composition over Réunion Island, in the Indian Ocean, using the Weather Research and Forecasting model coupled with Chemistry (WRF-Chem). The coexistence and spatial heterogeneity of anthropogenic and biogenic emission sources in this region present a valuable but challenging test of the model performance. The WRF-Chem model is evaluated against several observational datasets, including Proton Transfer Reaction Mass Spectrometry (PTR-MS) measurements of VOCs and oxygenated VOCs (OVOCs) at the Maïdo Observatory (2160m above sea level) in January and July 2019, representing austral summer and winter, respectively, and capturing the seasonal extremes for the region. While the primary goal of our study is a better understanding of (O)VOC budget at remote Tropical latitudes, important model refinements are made to improve the model performance, including the implementation of high-resolution anthropogenic and biogenic isoprene emissions, updates to the chemical mechanism, and adjustments to the boundary conditions. These refinements are supported by comparisons with PTR-MS data as well as with meteorological measurements at Maïdo, in situ NO_x and O₃ measurements from the air quality Atmo-Réunion network, Fourier Transform Infrared Spectroscopy (FTIR) measurements of O₃, CO, ethane and several OVOCs, also at Maïdo, and satellite retrievals from the Tropospheric Monitoring Instrument (TROPOMI).

TROPOMI NO₂ data suggests that anthropogenic emissions, particularly from power plants near Le Port, dominate NO_x levels over the island. Both TROPOMI and in situ surface NO₂ comparisons are used to adjust the power plant emissions at Le Port.

Surface ozone concentrations are overestimated by ~6 ppbv on average, likely due to the neglect of halogen chemistry in the model, though other factors may also contribute. Whereas modelled NO₂ over oceans is too low in summer when the lightning source is turned off, the inclusion of this source results in model overestimations corroborated by comparisons with upper tropospheric NO₂ mixing ratios derived from TROPOMI using the cloud-slicing technique (Marais et al., 2021). The model generally succeeds in reproducing the PTR-MS isoprene and its oxidation products (Iox), except for a moderate underestimation (~30%) of noontime isoprene concentration and for modelled concentration peaks near dawn and dusk, that are not seen in the observations. The ratio of Iox to isoprene (0.8 at noon in January) is fairly well reproduced by the model. The methanol and monoterpenes observations both suggest overestimations of their biogenic emissions, by factors of about 2 and 5, respectively. Acetaldehyde anthropogenic emissions are likely strongly overestimated, due to the lumping of higher aldehydes into this compound. Without this lumping, the modelled acetaldehyde would be underestimated by almost one order of magnitude, suggesting the existence of a large missing source, likely photochemical. The comparisons suggest the existence of a biogenic source of MEK equivalent to about 3% of isoprene emissions, likely associated with the dry deposition and conversion to MEK of key isoprene oxidation products. A strong model underestimation of the PTR-MS signal at mass 61 is also found, by a factor of 3-5 during daytime, consistent with previously reported missing sources of acetic and peracetic acid.

1 Introduction

Volatile organic compounds (VOCs) play a key role in important chemical processes throughout the atmosphere. They are precursors of ozone and secondary organic aerosols, both responsible for negative effects on human health (Jerrett et al., 2009; Pye et al., 2021) and the environment (e.g. Sicard et al., 2017), while also having climatic consequences (Mickley et al., 2004; Kanakidou et al., 2005; Shindell et al., 2006; Shrivastava et al., 2017). VOCs are mostly emitted by biogenic, anthropogenic and pyrogenic sources. The emissions of biogenic VOCs (BVOCs) due to terrestrial vegetation are affected by meteorological conditions (Guenther et al., 2006). VOCs undergo oxidation mainly by reaction with the hydroxyl radical (OH). OH is responsible for the removal of numerous pollutants in the atmosphere, initiating oxidation processes that usually transform airborne species into more oxygenated, and therefore more water soluble compounds (Comes, 1994). The reaction between VOCs and OH can effectively deplete OH and diminish the oxidative capacity of the atmosphere, thereby increasing the lifetime of pollutants and greenhouse gases such as methane (Zhao et al., 2019). Following the initial reaction of a VOC with OH, the further degradation of oxidation products can lead to additional OH loss, especially in remote regions with low levels of NO_x (NO_x = NO + NO₂) (Di Carlo et al., 2004; Read et al., 2012; Travis et al., 2020), where the reaction with NO is not the dominant sink of peroxy radicals (Logan, 1985; Atkinson, 2000).

Field measurements of OH reactivity in the remote troposphere have revealed the existence of a “missing” OH sink, primarily attributed to unknown organic compounds, in particular over boreal (Sinha et al., 2010) and tropical forests (Nölscher et al., 2016; Pfannerstill et al., 2021), as well as over the remote marine boundary layer (MBL) (Thames et al., 2020). In addition, models underestimate the concentrations of several known oxygenated VOCs (OVOCs) contributing to OH reactivity over the

remote ocean, most importantly acetaldehyde (Travis et al., 2020). The high observed abundances of acetaldehyde were recently supported by unexpectedly high measured concentrations of peroxyacetic acid (PAA) over the remote ocean (Wang et al., 2019), since PAA is photochemically produced almost exclusively from acetaldehyde oxidation under low NO_x conditions. The underestimation of acetaldehyde in models (e.g. Millet et al., 2010; Read et al., 2012; Travis et al., 2020) indicates a missing source likely due to air/sea exchange and secondary photochemical formation from oceanic precursors (Singh et al., 2003; Millet et al., 2010; Wang et al., 2019). Acetone is a substantial source of HO_x radicals (HO_x=OH+HO₂) in the upper troposphere and lower stratosphere (UT/LS) (Müller and Brasseur, 1999; Wang et al., 2020). Like acetaldehyde, it is a compound strongly regulated by the ocean, but which appears to be better simulated by models (Wang et al., 2020), although the observations suggest uncertainties in its sea-air exchanges, continental emissions and photochemical production and photodissociation rates (Fischer et al., 2012; Wang et al., 2020). Formic and acetic acid make up more than half of rain acidity in the remote atmosphere (Keene and Galloway, 1984), but their global budget remains poorly understood, and the significant discrepancies between modelled and measured distributions indicate large missing sources of both species in both polluted and remote regions (e.g. Paulot et al., 2011; Stavrakou et al., 2012; Millet et al., 2015; Khan et al., 2018). Although methanol is the most abundant tropospheric nonmethane compound, there are still uncertainties in its source apportionment (e.g. Jacob et al., 2005). Biogenic emissions are the largest source of methanol over continental areas (Stavrakou et al., 2011), and secondary photochemical sources appear to be the main contribution to methanol abundances over remote oceanic areas (Bates et al., 2021). Finally, although methyl ethyl ketone (MEK) is much less abundant than acetone, it is also more reactive (Brewer et al., 2020). Besides diverse sources including photochemical production and anthropogenic emissions, there is evidence of significant biogenic and oceanic sources of MEK that can affect its overall abundance, especially in more remote locations (Yáñez-Serrano et al., 2016; Brewer et al., 2020). Recently, the deposition of isoprene oxidation products on vegetation and subsequent conversion to MEK and other products has been proposed to be the largest contribution to the MEK budget at the global scale (Canaval et al., 2020), although this estimate relies on relatively few measurements.

In this study, we confront multiple chemical observational datasets from the remote island of Réunion, in the southern Indian Ocean, with regional atmospheric composition simulations using the high-resolution Weather Research and Forecasting model coupled with chemistry (WRF-Chem). More specifically, we make use of Proton Transfer Reaction - Mass Spectrometry (PTR-MS) measurements of VOC and OVOC concentrations (Verreyken et al., 2021) performed at the high-altitude site of Maïdo Observatory (21.1° S, 55.4° E, 2160m a.s.l.). Despite its small size, Réunion Island has significant diversity in emission sources, and it undergoes the influence of oceanic and continental emissions (Baray et al., 2013). For these reasons, Réunion Island is an area of continuous interest, where long-term observations as well as dedicated measurement campaigns were conducted and used to validate large-scale models (e.g. Vigouroux et al., 2012; Callewaert et al., 2022) and analyze factors influencing local atmospheric composition (e.g. Rocco et al., 2020; Rocco et al., 2022; Verreyken et al., 2020; Verreyken et al., 2021; Dominutti et al., 2022; Dufлот et al., 2022). Due to these multiple influences and to the pronounced orography of the island, atmospheric chemistry modelling over Réunion is especially challenging. In particular, the important role and strong spatial heterogeneity of anthropogenic and biogenic emissions (Verreyken et al., 2022) needs to be considered. Equally, the

steep topography of the island makes high resolution indispensable in order to forecast the local circulation patterns (El Gdachi et al., 2024). In this work, the WRF-Chem model is therefore enhanced with high-resolution (1 km^2) emission datasets for anthropogenic emissions and biogenic isoprene emissions. Furthermore, the model is evaluated and further refined based on comparisons with meteorological and air quality in situ data, with Fourier Transform Infrared Spectroscopy (FTIR) column measurements at Maïdo and with spaceborne (TROPOspheric Monitoring Instrument, TROPOMI) observations. Given the importance of meteorology for the simulation of transport, biogenic emissions and photochemistry, the model is evaluated against meteorological observations at Maïdo. FTIR and PTR-MS observations of long-lived compounds are particularly useful to evaluate the background atmospheric composition and to constrain the lateral boundary conditions of the regional model. TROPOMI column observations and air quality measurements are essential to test and constrain the emissions, in particular for NO_x . The PTR-MS dataset at Maïdo is expected not only to validate the model and better constrain the local emissions of important (O)VOCs, but also to identify potential shortcomings that should be considered in future studies of atmospheric composition in environments similar to Réunion Island.

The set-up and configuration of the WRF-Chem model, the chemical mechanism, the initial and boundary conditions and the emissions used in the simulations presented in this study are described in Sect. 2.1-2.3. Sect. 2.4 presents the observational datasets used to evaluate the model, including meteorological observations, surface chemical concentrations data (Sect. 2.4.1), the PTR-MS dataset of VOC and OVOC concentrations at Maïdo (Sect. 2.4.2), Fourier-Transform infrared spectroscopy (FTIR) column measurements at Maïdo (Sect. 2.4.3) and finally, spaceborne columns from TROPOMI (Sect. 2.4.4). Sect. 3 evaluates the model performance relative to the various measurement datasets. The results are recapitulated before concluding remarks in Sect. 4. Complementary figures and statistics can be found in the Supplement.

2. Methodology

2.1 Simulation area

Réunion is a French overseas island located in the Indian Ocean, $\sim 700 \text{ km}$ off the coast of Madagascar (Fig. 1). The island covers an area of 2512 km^2 (63 km long and 45 km wide), and roughly spans from -21.39° S to -20.87° S , and 55.22° E to 55.84° E . Réunion is a volcanic island with mountainous orography, with a maximum altitude of 3070 m above sea level (Piton de la Fournaise). The region is largely covered by native vegetation ($100\,000 \text{ ha}$) and is mostly free of strong anthropogenic emission sources (Duflot et al., 2019). The main anthropogenic emission sources are fossil fuel combustion in industrial and transport sectors (responsible for 87.1% of energy generation, Praene et al., 2012). The three principal industrial emission hotspots are due to biomass power plants (Le Gol and Bois-Rouge) and a diesel powerplant (Le Port) (Fig. 2).




Figure 1. Map of Africa and surrounding Indian Ocean indicating two model domains in blue and red (with 12.5 and 2.5 km horizontal resolution, respectively).

Réunion Island has a large concentration of endemic species (Myers et al., 2000). The island is largely dominated by vegetation and the plant species distribution changes with altitude (Fig. 2 in Strasberg et al., 2005; Foucart et al., 2018; Dufлот et al., 2019). Over 80% of the human population is concentrated in the coastal regions.

2.2 WRF-Chem

The Weather and Research Forecasting model coupled with chemistry is used to calculate meteorological and chemical atmospheric processes (WRF-Chem; Grell et al., 2005). Model version 4.1.2 of WRF-Chem was used in conjunction with its preprocessing system, WPS (WRF Preprocessing System) version 4.1.

2.2.1 Model configuration

To minimize computational demand, the simulations were conducted at 12.5 and 2.5 km horizontal resolution in the parent and nested domains, denoted d01 and d02, respectively (Fig. 1). Note that a 2 km horizontal resolution was found appropriate for simulating FTIR and in situ observations in a model study of greenhouse gases above Réunion, also using WRF-Chem but with the chemistry turned off (Callewaert et al., 2022). The projection is set to Mercator, as is the recommended set-up for low-latitude simulations close to the Equator. Thirty-day simulations were conducted for January and July 2019, each starting on the first day of the month at 00:00 UT. The local time at Réunion is UT + 4h. January and July correspond to summer and winter in the southern hemisphere, although the difference in meteorology between the two seasons is relatively small under the tropical climate of the island. January and July in 2019 were relatively free of events interfering with data collection (weather, volcanic activity, maintenance, etc.; see Table 1 in Verreyken et al., 2021). Lightning NO_x emissions are ignored in the reference model simulation, but are included in a sensitivity run, using the updated Price and Rind parameterization scheme

based on cloud-top height (Price and Rind, 1992; Wong et al., 2013). Since Barten et al. (2020) noted that the standard WRF-
150 Chem settings lead to a large overestimation of lightning emissions, we also downscale the number of flashes (adopted flash
rate factor of 0.1 and 0.02 for d01 and d02, respectively, to account for the difference in resolution) and the production of NO
per lightning strike is set to 250 moles, instead of 500 in the standard setting (Barten et al., 2020).
Simulations were conducted using Silicon Graphics (SGI) high-performance computing (HPC), equipped with an Intel
processor, using 72 cores (accounting for 5 h walltime for a two-day simulation in a 2-domain set-up). The simulations were
155 conducted sequentially in 2-day intervals, whereby the initial chemical conditions of each run are obtained from the previous
run, except in the case of the first day of the month. The meteorology is re-initialized at the start of each 2 day simulation. The
physical parameterizations incorporated in the simulations are listed in Table 1.

Table 1. List of WRF-Chem physical parameterizations adopted for this study.

Model component	Name	Reference
Microphysics	Morrison 2-moment scheme	Morrison et al. (2009)
Longwave radiation	RRTMG	Iacono et al. (2008)
Shortwave radiation	RRTMG	Iacono et al. (2008)
Planetary boundary layer	Shin-Hong scale-aware scheme	Shin and Hong (2015)
Surface layer	Revised MM5 scheme	Jiménez et al. (2012)
Land surface	Unified Noah land surface model	Tewari et al. (2004)
Cumulus parameterization	Grell 3D ensemble scheme	Grell (1993) Grell and Dévényi (2002)
Urban surface	Single-layer urban canopy model	Chen et al. (2011)

160




Figure 2. Map of Réunion Island with population density at 500 m x 500 m resolution. Key locations are denoted on the plot; cities (with 50,000+ population, based on the 2020 Population Document, <https://www.insee.fr/fr/statistiques/fichier/4265439/dep974.pdf>, last access: 29 November 2023) and the principal power plants. The blue triangle, representing the Maïdo Observatory, is the location of the PTR-MS and FTIR instruments. The locations of the air quality monitoring sites are represented by the red circles.

The land use dataset utilized in WRF is provided by MODIS 21 class (Friedl et al., 2002; Hulley et al., 2016), and has a resolution of 30 seconds (roughly 1km). The distribution of the MODIS land use index at the resolution of the nested domain is displayed on Fig. 3.

The dominant surface type at Maïdo is Woody Savannas (8), which is characterized by a tree cover between 30 and 60% and a tree canopy higher than 2 meters. There is spatial variability in the pixels adjacent to Maïdo, ranging from Evergreen Broadleaf Forests (2), Savannas (9) and Grasslands (10), each with different specifications regarding major vegetation type and tree cover percentage. Note that the Model of Emissions of Gases and Aerosols from Nature (MEGAN) v2.04 module utilized to calculate biogenic emissions (Guenther et al., 2006) in WRF-Chem recognizes only four vegetation types: broadleaf trees (BT), needle leaf tree (NT), shrub and brush (SB), and herbs, crops and grasses (HB). Their spatial distribution is provided by an independent dataset developed specifically for Réunion Island (see further below).




Figure 3. Land use index used for defining surface features in WRF, following the classification of the MODIS Land Cover Type Product. The resolution shown is $2.5 \times 2.5 \text{ km}^2$ (d02).

2.2.2 Chemical mechanism

The reference gas-phase mechanism used in the model simulations is the Model for Ozone and Related chemical Tracers, version 4 (MOZART-4) mechanism (Emmons et al., 2010), with the Kinetic PreProcessor (KPP) (Damian et al., 2002). Several updates to the mechanism were tested and implemented, as described further below. Aerosol chemistry is simulated using the Global Ozone Chemistry Aerosol Radiation and Transport model (GOCART) (Chin et al., 2002). The MOZART-4 mechanism was chosen in lieu of its more updated counterpart, MOZART-T1 (Emmons et al., 2020) for computational reasons, despite the improvements of the new version, namely on isoprene, monoterpene and aromatic chemistry. Nevertheless, the isoprene chemistry of the MOZART-4 mechanism used in the model has been updated to reflect recent mechanistic updates regarding OH-recycling and formation of methyl vinyl ketone (MVK) and methacrolein (MACR) (Sect. 2.2.3).

2.2.3 Mechanistic updates



2.2.3.1 Isoprene

The MOZART-4 isoprene mechanism lacks the OH recycling mechanisms which were shown to occur through the reactions of isoprene hydroxy hydroperoxides (Paulot et al., 2009) and the unimolecular reactions of isoprene peroxy radicals (Peeters et al., 2009, 2014; Wennberg et al., 2018; Müller et al., 2019). The reactions of isoprene peroxy radicals (ISOPO₂) and isoprene hydroxy hydroperoxides (ISOPOOH) in MOZART-4 are amended based on those studies and adjusted in order to match the results of a more up-to-date mechanism (MAGRITTEv1.1, Müller et al., 2019) through box model comparisons at both high NO_x (1 ppb NO_x) and low NO_x (0.1 ppb NO_x). More specifically, as seen in Table 2, the yields of MVK and MACR in the ISOPO₂ reaction with NO are increased at the detriment of the hydroxy aldehydes (HYDRALD), and OH is formed in reactions that play a dominant role at low NO_x (ISOPO₂+HO₂ and ISOPOOH+OH), in order to mimic OH recycling processes

involving compounds that are missing from the mechanism. Note that the mechanistic changes were chosen to avoid the introduction of additional species to keep a similar overall computational cost.

2.2.3.2 MEK oxidation

The mechanism of methyl ethyl ketone (MEK) is revised with the aim to avoid overestimation of the photochemical production of acetaldehyde through the oxidation of MEK and higher alkanes, which are MEK precursors. The oxidation of MEK by OH forms an intermediate radical product (MEKO2 in MOZART-4) which leads to the formation of acetaldehyde in the presence of NO, with a unit yield in MOZART-4 (reaction MO1 in Table 2). MEKO2 actually represents a lumping of three isomers denoted MEKAO2, MEKBO2 and MEKCO2 in the comprehensive Master Chemical Mechanism (MCM) v3.3.1 (Saunders et al., 2003; <https://mcm.york.ac.uk/MCM/>). Only one of the three radicals leads to acetaldehyde in the presence of NO according to MCM, and the overall chemistry can be summed up as follows



where RO2 denotes the acetonyl peroxy radical, CH3C(O)CH2O2, and EO2 denotes HOCH2CH2O2. Note that intermediary MCM reactions leading to the final products above were skipped as they involved compounds that are not defined in the MOZART-4 mechanism. These compounds were assumed to react rapidly according to their major sink reaction in presence of NO. Combining the equations above leads to the updated MO1 reaction shown in Table 2. The oxidation of MEK at low NO is oversimplified in the MOZART-4 mechanism and leads to the same final products as at high NO. However, the reaction of the peroxy radicals MEKAO2, MEKBO2 and MEKCO2 with HO2 form ketohydroperoxides that are expected to photolyse rapidly, leading in part to very different products (enols) than in the high NO case (Liu et al., 2018).

Table 2. List of mechanistic changes adopted in the MOZART-4 mechanism.

	Isoprene oxidation	
	In MOZART-4	In this work
IO1	ISOPO2 + NO → .08 ONITR + .92 NO2 + .23 MACR + .32 MVK + .37 HYDRALD + .55 CH2O + HO2 Rate = 4.4E-12 exp(180/T)	ISOPO2 + NO → .10 ONITR + .9 NO2 + .30 MACR + .53 MVK + .07 HYDRALD + .83 CH2O + .92 HO2 Rate = 4.4E-12 exp(180/T)
IO2	ISOPO2 + NO3 → HO2 + NO2 + .6 CH2O + .25 MACR + .35 MVK + .4 HYDRALD Rate = 2.4E-12	ISOPO2 + NO3 → HO2 + NO2 + .93 CH2O + .33 MACR + .59 MVK + .08 HYDRALD Rate = 2.4E-12

IO3	ISOPO2 + HO2 → ISOPOOH Rate = 8E-3 exp(700/T)	ISOPO2 + HO2 → ISOPOOH + OH Rate = 8E-3 exp(700/T)
IO4	ISOPOOH + OH → .5 XO2 + .5 ISOPO2 Rate = 1.52E-11 exp(200/T)	ISOPOOH + OH → .8 XO2 + .1 ISOPO2 + 1.7 OH + .2 CH3COCHO Rate = 1.1E-10
IO5	ONITR + OH → HYDRALD + HO2 +.4 NO2 Rate = 4.5E-11	ONITR + OH → HYDRALD + HO2 + .4 NO2 Rate = 3E-11
	MEK oxidation	
MO1	MEKO2 + NO → CH3CO3 + CH3CHO + NO2 Rate = 4.2E-12 exp(180/T)	MEKO2 + NO → 0.11 RO2 + 0.11 HO2 + 0.428 HCHO + 0.349 EO2 + 0.462 CH3CO3 + 0.462 CH3CHO + 0.079 C2H5O2 + NO2 Rate = 4.2E-12 exp(180/T)

2.2.4 Initial and boundary conditions

The model is initialized at the start of each run using input data from the Community Atmosphere Model with Chemistry (CAM-Chem; (Lamarque et al., 2012; Emmons et al., 2020)), which is a global model utilizing the MOZART-4 mechanism and providing output at 0.9° x 1.25° resolution. This allows for a direct representation of the initial and boundary conditions in the run. For species where the global Copernicus Atmosphere Monitoring Service (CAMS) reanalysis data (Inness et al., 2019) is available (NO_x, CO, O₃, HNO₃, H₂O₂, HCHO, peroxyacetyl nitrate or PAN, C₂H₆ and C₃H₈), we replace the CAM-Chem initial conditions with these higher resolution data (0.75° x 0.75°). Both inputs were taken at 6 hour intervals. Adjustments to these initial and boundary conditions were made based on comparisons with ground-based and FTIR measurements, which are detailed in Sect. 3.4. More specifically, the concentrations of methanol and PAN were multiplied by 0.6. Toluene, MEK and C_{>3} alkanes (represented as the lumped compound BIGALK) were multiplied by 0.4 in both seasons, while the same factor was only applied to acetone in July. Ethane was increased by a factor of 1.6 in July and left unchanged in January.

2.3 Emissions

The source apportionment of the main gaseous pollutants and VOC precursors is summarized in Fig. 4, which provides the total emission for every species over the island (average of January and July). In addition, Fig. 5 displays the emission

distribution of the four most emitted VOCs over the island, namely isoprene, methanol, monoterpenes and acetaldehyde, for the month of January. The assumptions and datasets used to derive these emissions are given in the following subsections.

2.3.1 Anthropogenic Emissions

240 The horizontal resolution of global anthropogenic emission inventories such as the Emissions Database for Global Atmospheric Research (EDGAR) (0.1° x 0.1°) (Crippa et al., 2022) is too coarse for air quality simulations over Réunion Island. The low-resolution dataset cannot accurately represent the transition from highly polluted areas (e.g. cities) to remote zones such as Maïdo. We utilized a 1 x 1 km² emission inventory estimated for NO_x, SO₂, CO, and NMVOCs based on data from the local air quality agency (Atmo-Réunion). The NMVOC species used in this study are listed in Table 3 below. The
245 NMVOC speciation follows the Regional Atmospheric Chemistry Mechanism, Version 2 (RACM2) (Goliff et al., 2013). Traffic emissions were provided by Atmo-Réunion for the main roads of the island. Industrial emissions, mostly cane sugar refining, rum distillation and diesel-electric power production, were estimated as point sources. Agricultural emissions are distributed in cultivation areas.

The Atmo-Réunion emissions are complemented with the remaining sectors (namely shipping, residential burning, non-road
250 transport, solvent processing, and waste management) from the EDGAR inventory, as well with species not included in the high-resolution inventory (ammonia, black and organic carbon and particulate matter). EDGARv6.1 (Crippa et al., 2022) was used for the trace gases and particulate matter, and EDGAR Hemispheric Transport of Air Pollution (EDGAR-HTAP) v4.3.2 (Crippa et al., 2018; Huang et al., 2017) was used for the non-methane VOCs (NMVOCs). Those inventories are also used for all species and sectors in the rest of the model domain. Emissions from EDGARv6.1 reflect values from 2018, while emissions
255 from EDGAR-HTAP v4.3.2 are based on values from 2012.

Temporal variations in the emissions are applied in accordance with Poraicu et al. (2023), based on EDGAR-specific temporal profiles (Crippa et al., 2020). Réunion being a French territory, the temporal profile follows the French specifications, which is defined based on French mainland regions. The temporal variation for some activities might not be representative of Réunion Island. For example, the seasonal temperature variation in mainland France is higher than in the tropics, with cold winters and
260 hot summers, which determines a different residential heating behavior. The seasonal dependence of residential sector emissions was therefore omitted.

Table 3: MOZART-4 VOC precursors and correspondence with the species classes of the Atmo-Réunion and EDGAR-HTAP V4.3.2 inventories.

MOZART-4	Atmo-Réunion inventory (RACM2 compound classes, Goliff et al., 2013)	EDGAR-HTAP V4.3.2 VOC classes (Huang et al., 2017)
BIGALK	0.6 * HC3 + HC5 + HC8	voc4 + voc5 + voc6 + voc18 + voc19
BIGENE	OLI + 0.5 * OLT	voc12

C ₂ H ₄	ETE	voc7
C ₂ H ₅ OH	0.85 * ROH	0.85*voc1
C ₂ H ₆	ETH	voc2
C ₃ H ₆	0.5 * OLT	voc8
C ₃ H ₈	0.4 * HC3	voc3
CH ₂ O	HCHO	voc21
CH ₃ CHO	ACD	voc22
CH ₃ COCH ₃	0.253 * KET	0.2 * voc23
CH ₃ OH	0.15 * ROH	0.15 * voc1
MEK	MEK	0.8 * voc23
TOLUENE	TOL + BEN + XYM + XYO + XYP	voc14 + voc15 + voc13
ISOP	ISO	voc10




Figure 4. Percentage contributions of different emission sources (anthropogenic and biogenic) to total emission of different chemical compounds considered in this work. The total, shown on the right-hand y-axis, is calculated from the average of the two months. Results shown for the optimal run (R0), after adjustment of the anthropogenic and biogenic sources described in text (Sects. 2.2, 2.3 and 3).




Figure 5. Total emissions of isoprene, methanol, monoterpenes and acetaldehyde over the inner model domain at 2.5km resolution (d02), averaged over the month of January 2019. The white circle indicates the location of the Maïdo observatory.

2.3.1.1 Anthropogenic emission adjustments

275 Given the coarse resolution (0.1°) of the EDGAR inventory used for residential emissions, those emissions were redistributed spatially on the model grid using a distribution of population density, specific to Réunion (<https://public.opendatasoft.com/explore/dataset/population-francaise-par-departement-2018/table/?disjunctive.departement>, last access: 8 December 2023). The population distribution is on a latitude-longitude grid at a resolution of 30m, and was regridded to the model resolution in the nested domain. In addition, since traffic activity occurs not only over highways but
280 also in cities and generally where people live, the traffic sector was redistributed by assuming that 70% of the emissions follow the population map, and 30% of the emissions are distributed on the principal highways defined in the Atmos-Réunion inventory. The resulting high-resolution distribution is compared with EDGAR on Fig. 6.

The NO_x industrial emissions (primarily energy production) were redistributed based on the EDGARv6.1 emissions for this species and sector. Both the EDGARv6.1 dataset and TROPOMI NO₂ column data (see Sect. 3.5) suggest a maximum emission
285 in the northwestern region of the island, around the Port-Est power plant. Therefore, the total industrial NO_x emissions were distributed amongst the 4 point source regions using percentage contributions based on the NO_x industrial EDGARv6.1 emission dataset. EDGAR-HTAP v4.3.2 reports a total VOC emission three times higher than those from Atmo-Réunion, implying significantly lower VOC/NO_x emission ratio in the latter inventory. Although both inventories exhibit similar spatial distributions, the Atmo-Réunion emissions are higher around the largest city (Saint-Denis) and lower in the other industrialized
290 areas.




Figure 6. Anthropogenic emissions of (a-b) NO_x and (c-d) total VOC over the inner model domain at 2.5km resolution (d02), averaged over the month of January 2019. (a) and (c) display low-resolution EDGAR emissions, while (b) and (d) show the high-resolution Atmo-Réunion dataset, with adjustments described in this Section.

2.3.2 Biogenic Emissions


Emissions from biogenic sources, mainly from vegetation, are calculated using MEGAN version 2.04. The emissions are calculated on-line (at the same time step as the model), based on the simulated meteorological fields and vegetation types defined by the land use map. This model utilizes 4 general vegetation types (broadleaf trees, needleleaf trees, shrubs, herbaceous land) together with standard emission factors and other parameters needed to estimate the emission at each timestep. MEGAN calculates the emissions for 20 compounds/compound classes, based on which the emission of 138 individual species can be estimated. The emissions of WRF-Chem species classes (e.g. monoterpenes) are obtained by lumping of those individual compounds. The biogenic emission for each class (E) is calculated using

$$E = \varepsilon \gamma \rho, \quad (1)$$

i.e., E is calculated as a product of the emission factor at standard conditions (ε), the emission activity factor (γ) and a factor accounting for production and loss within the plant canopy (ρ). The latter factor is not considered here, i.e. $\rho=1$. The standard emission factor (in $\mu\text{g m}^{-2} \text{hr}^{-1}$) is obtained from independent field and laboratory studies.

The dimensionless emission activity factor γ accounts for the dependence on environmental conditions such as the leaf area index (LAI), the photosynthetic photon flux density, i.e. the amount of visible light at leaf level (γ_P), leaf temperature (γ_T), leaf age (γ_A), soil moisture (γ_{SM}) and CO_2 inhibition (γ_C). For more detail on the MEGAN algorithm, see Guenther et al. (2006; 2012).

MEGAN relies on a 1 km resolution global map of plant functional types and leaf-area index. Since the standard database provided by NCAR (<https://www.acom.ucar.edu/wrf-chem/download.shtml>, last access: 1 December 2023) has zero values for all these variables over the Réunion area, we replaced it with a detailed cartography of plant functional types and isoprene standard emission factors (100m resolution) for Réunion Island constructed from an extensive survey of natural habitats (ZNIEFF, 2013) and phytosociology studies (Strasberg et al., 2005). This is completed by measurements of the leaf area index (LAI) for representative plant species (Dufлот et al., 2019). The resulting distributions of plant functional types, standard isoprene emission factor and leaf area index are shown on Fig. 7 at the model resolution (inner domain). As seen on this figure, broadleaf trees are by far the dominant PFT over the island. On average over the island (except bare soil), the isoprene emission factor is about $2900 \mu\text{g m}^{-2} \text{h}^{-1}$ ($42.7 \text{ mol km}^{-2} \text{h}^{-1}$). Around Maïdo observatory, the emission factor ranges between 3000 and 6000 $\mu\text{g m}^{-2} \text{h}^{-1}$. Those values are much lower than the default values in the WRF-Chem model for the dominant PFTs (13000 and 11000 $\mu\text{g m}^{-2} \text{h}^{-1}$ for broadleaf trees and for shrubs, respectively). This illustrates the importance of accounting for the local plant species distribution, when available.



325 **Figure 7. (a-d) Plant functional type coverage (%) used to calculate biogenic emissions with MEGAN in WRF-Chem, (e) isoprene standard emission factor ($\text{mol km}^{-2} \text{hr}^{-1}$) and (f) leaf area index at 2.5 km horizontal resolution. The white circle represents the location of the Maïdo Observatory.**

Except for isoprene, the emission parameters for each compound (or compound class) are assumed constant within each of the 4 vegetation types, despite possibly important variations. In this work, the MEGAN parameters for several compounds (methanol, monoterpenes, methyl ethyl ketone, acetaldehyde/ethanol) were updated based on previous work and on
330 comparisons with the PTR-MS measurements at Maïdo, which will be further discussed in Sect. 3.3. More specifically, the methanol emission factor was decreased from 800 to 400 $\mu\text{g m}^{-2} \text{h}^{-1}$ for broadleaf trees and herbaceous vegetation, in agreement with Stavrakou et al. (2011). The emission factors for monoterpenes were decreased by a factor of 5, and the light-dependent fraction (LDF) was increased to 0.9 (from initial values ranging between 0.05 and 0.8 for different monoterpenes). This high LDF value implies an emission algorithm similar to isoprene, with little emission at night. It might reflect the
335 dominance of broadleaf forests and the marginal extent of coniferous trees over the island (Fig. 7), suggesting a minor role of temperature-controlled emission of VOC from storage pools (see e.g., Derstroff et al. 2017; Yáñez-Serrano et al., 2015). The MEGAN2.1 emissions of methyl ethyl ketone (MEK) were enhanced, since recent studies showed an important biogenic contribution to its budget (Yáñez-Serrano et al., 2016). Since this biogenic source is thought to result from the partial conversion of isoprene oxidation products (MVK and ISOPOOH) deposited on plant leaves (Canaval et al., 2020), the biogenic
340 source of MEK is assumed proportional to that of isoprene. Although this assumption might underestimate MEK emission at night, since non-zero concentrations of isoprene oxidation products might be sustained during nighttime (e.g. Langford et al., 2010), the error is likely small since the dry deposition of those compounds is usually very slow at night (Nguyen et al., 2015). The scaling factor used for calculating the emission has been adjusted as discussed in Sect. 3.3.

2.3.3 Biomass burning emissions

345 The fire emissions from the Fire INventory from NCAR (FINN) emission inventory (Wiedinmyer et al., 2011) were zero over both domains in January and July 2019. FINN is a global, 1 km-resolution dataset providing daily estimates of biomass burning emissions. However, as for the biogenic emissions input files, the biomass burning emissions could be artificially missing in this inventory due to the small size of the island. In any case, there was no significant biomass burning activity detected at Maïdo in January and July 2019 (Verreyken et al., 2020). The fire emissions were therefore omitted in the simulations.

350 2.4 Measurements

2.4.1 In situ surface chemical observations

Common air pollutants are measured hourly at 18 air quality stations around Réunion Island. These stations are operated by Atmo-Réunion (<https://atmo-reunion.net/le-dispositif-de-surveillance>, last access: 1 December 2023). The stations measuring NO₂, O₃ or CO are shown on Fig. 2. In the comparison of modelled and observed NO₂ concentrations, a correction factor is
355 applied to account for known interferences in the NO₂ measurement (Lamsal et al., 2008). These interferences are due to NOy

reservoir compounds (such as HNO_3 and PAN) contributing to the measured signal and leading to higher NO_2 values than are actually present. The correction factor is calculated using NO_2 , PAN and HNO_3 concentrations obtained from the model, and applied to the simulated NO_2 , as detailed in Poraicu et al. (2023).

2.4.2 PTR-MS measurements

360 The PTR-MS instrument is deployed for the measurement of VOC species (Hewitt et al., 2002). The dataset (Verreyken et al., 2021) includes 13 molecules, or more precisely, 13 mass-to-charge ratios (m/z). The instrument (hs-PTR-MS, Ionicon Analytik GmbH, Austria) was located at the Maïdo Observatory high-altitude site (2160 a.s.l) (21.079°S, 55.384°E). Further technical information on the PTR-MS measurements can be found in Verreyken et al. (2021). The major isoprene oxidation products at high NO_x , methacrolein (MACR) and methylvinyl ketone (MVK), are measured conjointly due to their identical molar mass

365 (70 g mol^{-1}) and molecular formula ($\text{C}_4\text{H}_6\text{O}$). Isoprene hydroxy hydroperoxides (ISOPOOH), which are major isoprene oxidation products at low NO_x , are known to react heterogeneously in the PTR-MS instrument and decompose to either MVK or MACR (+HCHO) (Rivera-Rios et al., 2014). Therefore the PTR-MS signal corresponding to MVK+MACR includes a contribution of ISOPOOH, which is significant at low NO_x conditions. Although the exact conversion efficiency of ISOPOOH into MVK+MACR is uncertain, a complete conversion is assumed here, and the corresponding signal will be referred to as

370 Iox (isoprene oxidation product, MVK+MACR+ISOPOOH).

The PTR-MS signal for acetic acid (CH_3COOH) has interferences from other chemical species, namely glycolaldehyde, peroxyacetic acid, propanols and ethyl acetate (Baasandorj et al., 2015). The contribution of propanol is very low (<1%) and is not considered. Protonated glycolaldehyde has the same mass-to-charge ratio as protonated acetic acid (61 m/z), while peroxyacetic acid and ethyl acetate were shown to fragment upon protonation with a resultant fragment ion (61 m/z) (Španel

375 et al., 2003, Fortner et al., 2009). We therefore opt to compare the observations with the sum of the modelled concentrations of acetic acid, glycolaldehyde, and peroxyacetic acid, assuming a similar sensitivity for all three compounds. Ethyl acetate is not considered in the MOZART-4 mechanism although this compound has both direct emissions and formation from the chemical oxidation of ethers (Orlando and Tyndall, 2010 and references within). Formic acid (46 g mol^{-1}), also measured by the PTR-MS, is not defined in the MOZART-4 mechanism, and is thus not considered further in this study. The signal at m/z

380 73 has a major contribution from methyl ethyl ketone (MEK) (Verreyken et al., 2021) but other compounds, namely butanal isomers and methylglyoxal (MGLY), also contribute (Yáñez-Serrano et al., 2016). The contribution of butanal is neglected here, being not calculated by the model. The estimated sensitivity of the instrument to MGLY being only a fraction (~0.7) of the sensitivity to MEK (Koss et al., 2018), the PTR-MS concentrations will be compared to the sum $\text{MEK} + 0.7 \times \text{MGLY}$ calculated using the model.

385 Benzene, toluene and xylenes are all measured at the site. However, in the MOZART-4 mechanism they are lumped and treated as toluene. Benzene, toluene and xylenes having different chemical lifetimes (Atkinson, 2000) and different emission distributions, the PTR-MS observations of aromatics cannot be evaluated with the model. In situ meteorological data (2-m

temperature, relative humidity, wind speed and direction and solar radiation) is measured at and provided for the PTR-MS site (Maïdo Observatory) at the same temporal resolution as the chemical concentrations (2.7 minutes).

390 2.4.3 FTIR measurements

Ground-based FTIR measurements have been performed at Réunion Island (Maïdo and Saint-Denis) since 2002, first on a campaign basis (Senten et al. 2008). Long-term measurements of VOCs and other compounds started in 2009 at Saint-Denis (Vigouroux et al., 2012). Since 2011, FTIR measurements in the prescribed NDACC (Network for the Detection of Atmospheric Change) spectral range ($600\text{--}4500\text{ cm}^{-1}$) are performed at the Maïdo Observatory (Baray et al. 2013), using a
395 Bruker 125 HR (since 2013).

The FTIR being a remote sensing technique which measures the absorption of solar light by atmospheric species along the line-of-sight (instrument-sun), the primary product of the FTIR retrievals is the total column of the absorbing gases. In addition, low resolution vertical profile information can also be derived by using the pressure and temperature dependence of the line shape. The choice of spectral windows and spectroscopic parameters is optimized for each target species, and preferably within
400 the whole NDACC FTIR community to ensure a consistency within the network. Table S1 summarizes the main retrieval settings for the species used in this paper.

The HCHO retrieval has been optimized recently at more than twenty FTIR stations (Vigouroux et al., 2018) and is now an official NDACC target species. Methanol is not an official NDACC species, and is measured at only a few sites (see e.g. Wells et al., 2024) but not in a harmonized way. Ethane is an official NDACC species and has been harmonized within the network
405 as described in Franco et al. (2015). At Maïdo, the humidity being important, the third window suggested for harmonization in Franco et al. (2015) is not used to avoid strong interference with water vapor lines. Carbon monoxide is also an NDACC target species, and as such the windows and spectroscopy are harmonized and found in the NDACC InfraRed Working Group (IRWG) documentation (<https://www2.acom.ucar.edu/irwg>). The current IRWG retrieval strategies for official species are currently being reprocessed, with the aim of using improved settings. Ozone is the FTIR species with the most degrees of
410 freedom for signal ($\text{DOFS} = 4$ to 5), allowing the retrieval of several independent partial columns: one in the troposphere and three in the stratosphere (Vigouroux et al., 2008).

Peroxyacetyl nitrate (PAN) is not an official target species. It has been measured at only a few stations (Mahieu et al., 2021) because its weak spectral signature makes it difficult to detect and retrieve. This is the first time that PAN time-series at the Maïdo station are presented. From the two stronger PAN absorptions used in Mahieu et al. (2021), only the band at 1163 cm^{-1}
415 1 is useful at Maïdo, the other one having too low signal-to noise-ratio. As shown in Fig. S1, the spectral signature of PAN is much weaker than the other absorbing gases. Therefore, the interfering species must first be carefully taken into account. In addition to the gases shown in Fig. S1, HCFC-22 also interferes. We pre-retrieve HCFC-22 in a dedicated window ($828.62\text{--}829.35\text{ cm}^{-1}$), and then, for each individual spectrum, the retrieved profile of HCFC-22 is used as fixed values in the PAN retrieval. For each spectrum, profile retrievals are also made for H_2O , HDO , O_3 , N_2O , and CFC-12 and used as a priori values

420 in the PAN retrievals, in which PAN and H₂O profiles are retrieved, while the other species are scaled from their a priori pre-retrieved profiles (with the exception of HCHC-22 which is not scaled, but fixed).

2.4.4 TROPOMI observations

The TROPospheric Monitoring Instrument (TROPOMI) is a spaceborne instrument aboard the European Space Agency (ESA) Sentinel-5P (S5P) satellite (Veefkind et al., 2012), which monitors the global distribution of multiple air pollutants, among
425 which nitrogen dioxide, formaldehyde, ozone and carbon monoxide. The S5P satellite is sun-synchronous and the retrieved data has a daily global coverage with (typically) one TROPOMI measurement at ~13:30 local time, with a spatial resolution of 7 x 3.5 km² (updated to 5.5 x 3.5 km² in August 2019). The main species of interest for this study are NO₂ and HCHO. The measurement of multiple species is possible due to the large spectral range of the TROPOMI spectrometer, specifically ultraviolet (UV), visible (VIS), near-infrared (NIR) and shortwave infrared (SWIR) ranges (ranging from 267 to 2389 nm).
430 The instrument is a push-broom spectrometer that scans the Earth while the satellite moves and measures the composition of the atmosphere using differential optical absorption spectroscopy (DOAS). The light travels from the Sun through the atmosphere and is reflected back to space, where it is measured by the spectrometer. Molecules absorb photons at well-defined windows, depending on their molecular structure (primarily 405 - 465 nm for NO₂ and 328-359 nm for HCHO). Comparison of the observed spectrum to a reference spectrum enables the retrieval of the slant columns through a fitting procedure
435 involving multiple compounds that are active in the absorption bands of the species of interest. For both compounds, TROPOMI retrieves a slant column density (SCD) from the Level-1b radiance and irradiance spectra, that represents the total amount of compound present along the effective solar light path (van Geffen et al., 2020).

In the case of NO₂, the total SCD combines both tropospheric and stratospheric distributions. The tropospheric slant column is obtained by subtracting the stratospheric contribution from the total SCD. This contribution is obtained from a global
440 chemical transport model (TM5-MP, Williams et al., 2017). The tropospheric vertical column density (VCD) is obtained by dividing the slant column by an air mass factor (AMF) (Palmer et al., 2001) which is dependent on the vertical profile of the considered compound. AMFs are obtained from radiative transfer calculations and can introduce a large source of uncertainty in the VCD calculation (30-40%, Lorente et al., 2017), especially in the presence of clouds. The vertical distributions of NO₂ and HCHO are taken from the global chemistry transport model TM5-MP (van Geffen et al., 2022a; Williams et al., 2017; De
445 Smedt et al., 2018) at a resolution of 1° x 1°. In the HCHO VCD algorithm, a background correction is applied on a daily basis. Since the reference spectrum is obtained from Earth radiances in the equatorial Pacific, the slant column retrieved from the fit corresponds to an excess over the remote background, where the principal HCHO source is methane oxidation. The TM5-MP columns over the same region are therefore added to the vertical columns to account for this background (De Smedt et al., 2021).

450 Due to its role for the calculation of the AMF, the choice of vertical profile has an impact on the VCD retrieval, and it is necessary to consider the vertical sensitivity of the TROPOMI instrument when comparing with other datasets. NO₂ VCD biases against independent measurement datasets decreased when the TM5-MP a priori was replaced with model profiles at

higher resolution (e.g Tack et al., 2021; Judd et al., 2020; Douros et al., 2023) or with measured profiles (e.g. Dimitropoulou et al., 2020). In our study, the averaging kernels of the satellite data are applied to the model profiles to calculate a “smoothed” vertical column for comparison with the satellite retrieved columns (e.g. Boersma et al., 2016). The averaging kernel represents the sensitivity of the measurement to the tracer concentrations at different altitudes, weighted by the assumed vertical profile of the tracer (Eskes and Boerma, 2003). It is provided alongside the measurement in the TROPOMI products.

We present comparisons between the reprocessed version (RPRO) of the NO₂ and HCHO retrievals from TROPOMI (v3.2) and simulated tropospheric columns. For both compounds, the averaging kernels are applied to the model profiles vertically interpolated to the TM5 vertical pressure grids. Both WRF-Chem and TM5 vertical pressure levels are calculated using hybrid sigma-pressure coordinates and the surface altitude using values from WRF-Chem and TM5, respectively. Quality filtering (QF) follows Algorithm Theoretical Basis Document (ATBD) recommendations (QF > 0.75 for NO₂ and QF > 0.5 for HCHO) (van Geffen et al., 2022b; De Smedt et al., 2022).

In addition, we applied the oversampling technique to the NO₂ and HCHO data in order to gain insight into the fine-resolution distribution of those compounds (e.g., de Foy et al., 2009). This technique consists in the long-term averaging of TROPOMI measurements on a very fine grid, taken here to be $0.01^\circ \times 0.01^\circ$ ($\sim 1 \times 1 \text{ km}^2$). The measurement from a given TROPOMI pixel is taken to apply to a circle defined by the center of the pixel and a radius of 3.5 km. In this way, each $0.01^\circ \times 0.01^\circ$ pixel accumulates >500 measurements over the considered time period (May 2018 - July 2022). This technique takes advantage of the variable offset of TROPOMI observations from day to day and achieves a high signal-to-noise ratio at high resolution but is not intended for direct comparison with the model, given the long-term averaging. It aims to present the average pollutant distribution at finer scales to inform about emission hotspots that are potentially lost when considering data over short periods of time.

3. Results and discussion

The optimal model set-up for this region was obtained through multiple sensitivity runs testing various changes to the emissions, lateral boundary conditions and chemical mechanism used in the model. In this section, the reference run (R0) represents the simulation set-up incorporating all model updates. The sections below highlight the impact of the updates through model comparisons with the observations. The list of sensitivity runs portrayed in this study is provided in Table 4.

Table 4: Sensitivity runs conducted in this study, with the shorthand notation.

Shorthand	Description
R0	Best run with following adjustments: <ul style="list-style-type: none">• downscaling of principal NOx power plant emission by a factor of 5 (Sects. 3.2 and 3.5)• adjustment of lateral boundary conditions (Sect. 2.2.4)

	<ul style="list-style-type: none"> • adjustment of biogenic VOC emissions (Sect. 2.3.2) • updates to the MEK and isoprene chemistry within the MOZART-4 mechanism (Sect. 2.2.3)
S1	As R0, without the downscaling of NO _x power plant emission by a factor of 5
S2	As R0, without the adjustment of lateral boundary conditions
S3	As R0, without the biogenic VOC emission adjustments
S4	As R0, without updates to the MEK and isoprene chemistry within the MOZART-4 mechanism
S5	As R0, with direct anthropogenic and biogenic emissions of acetaldehyde, acetone, MEK, formaldehyde and acetic acid set to zero
S6	As R0, including lightning emissions (Sect. 2.2.1)

480 3.1 Evaluation against meteorological measurements

The model is evaluated against meteorological measurements at the Maïdo site on Fig. 8. Surface temperature and solar radiation are both represented by the model in both January and July. The observed diurnal profile of temperature is well reproduced, except for a low nighttime bias on many instances. On most days in both months, the midday modelled value matches the observation well. Exceptions occur, e.g. during the first days of January, when the model underestimates

485 temperature and overestimates relative humidity and cloudiness. Errors in the WRF-Chem cloud parameterization were shown to impact the model performance for many physical and chemical processes (Zhao et al., 2012; Berg et al., 2015; Ryu et al., 2018). Furthermore, cloudy periods (when observed solar radiation is lower, e.g. on Jan. 8-9) are typically not well represented as such. The higher solar radiation fluxes in January (maximum of about 1000 W m⁻² in January compared to 750 W m⁻² in

490 month, as indicated by both model and measurement data in Fig. 8. Nevertheless, the seasonal variation of meteorological parameters is well reproduced by the model. January is warmer (by 5.5 and 6.4 K according to the measurements and the model, respectively), more humid (by 14 and 11%) and has higher radiation fluxes (by 51 and 65 W m⁻²) than the month of July. On average, the wind speed is slightly overestimated in both seasons. It is generally close to the observations, except during several predicted high-wind episodes (e.g. on 7-9 January and 24 July) that are not present in the observations. The

495 predominant wind direction is correctly predicted (mostly westerly winds during January, and an alternance of westerly, northerly and easterly winds during July), but the model often fails to reproduce the short-term variability of that direction. For example, the sporadic occurrence of easterly winds (270°) in January is missed by the model, whereas sporadic southerly

winds are simulated in January and July but are seldom observed. The observed circulation likely results from the competition between overflowing trade winds and meso-scale dynamics induced by anabatic thermal flow coupled with upslope transport to the Maïdo Observatory, a direct consequence of orography (Duflot et al., 2019; El Gdachi et al., 2024). This will directly affect transport patterns and model comparisons with chemical measurements over the site. Seasonal averages are given in Table S2.




Figure 8. Evolution of observed and modelled surface meteorology at Maïdo observatory for (a-e) January and (f-j) July. The observations are shown in black, while the cyan line represents the WRF-Chem results. The wind direction is represented in degrees, relative to north.

3.2 Evaluation against air quality station measurements




Figure 9. Averaged modelled surface concentrations of NO_x, OH, O₃ and HCHO (ppb) in January (top row) and July (bottom row), obtained from simulation R0.

Figure 9 presents the monthly averaged distributions of key chemical compounds calculated by the model over Réunion Island, for January and July 2019. These distributions will be useful for the discussion of model comparisons with in situ and satellite observations (see next subsections). The strong heterogeneity of anthropogenic emissions and their high intensity over urban and industrial centers lead to strong contrasts between chemical regimes over different regions of the island. For example, the NO_x mean surface concentration ranges between 0.12 ppb at the most remote part of the island to typically 5-15 ppb over most urban/industrial areas, and over 30 ppb near the Le Port thermal plant. As seen on Fig. 9, other important compounds like OH and O₃ are strongly impacted by anthropogenic emissions. Over a large portion of the island (southwestern part as well as a small portion of the northern coast) but also over a large oceanic region surrounding almost the entire island (especially to the northwest due to the influence of Le Port emissions), OH levels are enhanced, primarily because NO_x promotes the conversion of HO₂ to OH (Spivakovsky et al., 2000) through the reaction



As a result, OH increases as NO_x increases when NO ranges between ca. 10 ppt and ca. 500 ppt (Logan et al., 1981). At very high NO_x levels (> ~5 ppb NO_x), OH decreases as NO_x increases, mainly due to the sink of HO_x (=OH+HO₂) due to the radical termination reaction



which explains the depletion of OH levels over the main NO_x hotspots. NO_x has a relatively weak effect on the patterns of the ozone distribution, except for the clear titration effect near Le Port and the main cities, such as Saint Denis and Saint Pierre.

530 Formaldehyde is clearly strongly affected by anthropogenic activities, as seen from the high surface concentration levels around the NMVOC emission areas (Fig. 6). The distribution also partly reflects the complex impacts of NO_x on OH, since the NMVOC oxidation processes leading to HCHO formation are mainly driven by their reaction with OH. Note that the NO_x concentrations rapidly decrease with altitude. Therefore the OH depletion effect due to NO_x seen on Fig. 9 becomes rapidly negligible at higher altitudes. Therefore, the main effect of anthropogenic NO_x emissions on the oxidative conditions above
 535 the island is a strong enhancement of OH concentrations, except over localized hotspots, very close to the surface. NO_x plays a major role in the OH budget, and inaccurate predictions of its abundances can affect the model comparisons for VOCs and their oxidation products, in particular at Maïdo. Besides the insights provided by TROPOMI NO₂ column data (see Sect. 3.5), network measurements of in situ concentrations of NO₂, NO_x (=NO+NO₂) and O₃ are available mostly in polluted areas, in cities and in the vicinity of industrial sources (Fig. 2). Due to representativeness issues, caution is required when comparing
 540 model results with measurements often obtained very close to strong pollution sources. For this reason, model underestimation of NO_x levels is to be expected at many stations. A useful indicator is the [NO_x]/[NO₂] ratio. At photochemical steady state (PSS), it is given by

$$\left(\frac{[\text{NO}_x]}{[\text{NO}_2]}\right)_{\text{PSS}} = 1 + \frac{J_{\text{NO}_2}}{k_1 \cdot [\text{O}_3] + \dots}, \quad (6)$$

where J_{NO_2} is the photolysis rate of NO₂, and k_1 ($=1.95 \times 10^{-14} \text{ molec.}^{-1} \text{ cm}^3 \text{ s}^{-1}$) is the rate of the reaction



During the night, the expected value of the ratio is unity. However, close to a pollution source, photochemical steady state is not achieved, since directly emitted NO can travel over a few hundred meters within its chemical lifetime, of the order of several minutes for typical nighttime ozone levels (a few ppb, see Fig. 10). Furthermore, ozone is titrated to even lower levels in direct vicinity of strong sources. Given the model resolution and the distribution of NO_x sources (Fig. 6), strong and
 550 systematic ozone titration occurs only in the region of Le Port (Figs. 2 and 9), with its strongly emitting power plants. Elsewhere, the nighttime [NO₂]/[NO_x] ratio is below 1.1 in the model. In the measurements, however, the observed ratio is usually ~1.3, based on monthly averaged nighttime concentrations, and it even exceeds a factor 2 at several stations (Table S3), namely the stations 10-12 on the southern coast of the island (see Fig. S2 for station locations). These high values suggest the presence of very close NO_x sources that are unresolved by the model.

555 Figure 10 presents a comparison of observed and modelled surface concentrations of NO_x and O₃ for a 30-day simulation in January 2019. The stations were divided into two classes – the two stations near the Le Port power plants (LP) and the other stations, excluding the least representative stations, i.e. those for which the observed nighttime [NO_x]/[NO₂] ratio exceeds a factor of 2 (Table S3). Figures 10 and 11 display the averaged performance for the two classes of stations. Table S4 in the Supplement provides a statistical analysis for each station with available observations, including the correlation coefficient,
 560 root mean square error (RMSE), and index of agreement.




Figure 10. Evaluation of the average modelled surface NO₂, NO_x and O₃ in January, from simulations S1 (cyan) and R0 (red), against in situ network observations (dashed line) (a-b) in the region of Le Port (LP) and (c-e) at the other stations, for which the observed nighttime ratio $[\text{NO}_x]/[\text{NO}_2] < 2$ (stations 3-9 of Table S3, see Fig. S2). The model NO₂ concentrations have been corrected to account for interferences in the measurements using the corresponding modelled concentrations of PAN and HNO₃ (Poraicu et al., 2023).




Figure 11. Average diurnal profile of NO₂, NO_x and O₃ in January, from simulations S1 (cyan) and R0 (red), shown for the Le Port (LP) stations, and at the other stations for which the observed nighttime ratio [NO_x]/[NO₂] < 2. Note that ozone measurements were not available at LP stations. The black dashed line represents the observations at the same stations.

The LP stations (Fig. 2, Terrain de Sel and Centre Pénitentiaire) are strongly impacted by the NO_x emissions due to the power plants located near Le Port, on the northeastern coast of the island (Fig. 9). However, the observed nighttime NO_x concentrations are largely overestimated by the model at those stations. During daytime, a large overestimation is also found for simulation S1, which adopted the original estimation of the NO_x emissions due to the Le Port power plants. On average over the 30 day period, the model overestimation for NO_x (NO₂) against the LP measurements, a factor of 3 (4) during daytime in the S1 simulation, is strongly reduced (to a factor of 1-1.4) when the power plant emission is reduced by a factor of 5 (run R0) (Fig. 11). This finding suggests that this reduction of the power plant emission is justified, in agreement with the comparison of NO₂ columns with TROPOMI data (Sect. 3.5). The large model overestimation of nighttime NO₂ and NO_x at the LP stations stands in contrast with the fairly good agreement at the other stations (see further below). A possible reason might be that all emissions are injected at surface level in the model, whereas the actual injection heights of power plant emissions might be above the (usually stable) nighttime boundary layer. The impact of the 5-fold reduction of NO_x emissions is small (-25%) on the nighttime NO₂ levels at the two LP stations, despite their close proximity to the power plants. This is explained by the fact that NO (not NO₂) is emitted by the power plants, and that a very large emission leads to O₃ titration

585 through the reactions of NO_x with O₃. The much stronger ozone titration in the S1 run, compared to R0, explains that NO is much less rapidly converted to NO₂ in S1, compensating the higher emission. The titrating effect of the strong NO_x emissions on the ozone distribution is clearly seen on Fig. 9, especially at and near Le Port and to a lesser extent at other industrialized centers.

By comparison, the effect of this emission reduction is small at the other stations, except during the morning traffic peak (4
590 UT i.e. 8 LT, Fig. 11). The model correlates very well with the observed NO₂ and NO_x time series, although the model performance is lower on the first and last days of the month, coinciding with unusually large errors in meteorological parameters at Maïdo (e.g. solar radiation and wind direction, see Fig. 8). The average diurnal cycle of concentrations shows a good model agreement for NO₂ and NO_x during night and day, except for a moderate overestimation of NO₂ (Figs. 10 and 11). This discrepancy between the model agreement for NO_x and for NO₂ is likely due to the model overestimation of surface
595 O₃, by more than 10 µg m⁻³ on average, leading to an overestimated [NO₂]/[NO] ratio (reaction r7). Although the S1 run achieves a better agreement with O₃ data at the morning peak hour, the performance of the two simulations is similar at other times of the day, and the shape of the diurnal profile is better represented by the R0 run. The reasons for the ozone overestimation (by ca. 12 µg m⁻³ on average) are unclear, but could be due to the neglect of halogen chemistry in the WRF-Chem version used in this study. For example, recent studies indicated that the inclusion of halogen (Cl, Br, I) chemistry in a
600 regional or global model might deplete near-surface ozone levels by as much as 7 ppb (~14 µg m⁻³) in the marine tropical troposphere (Badia et al., 2019; Caram et al., 2023). This reduction of ozone is partly due to a shortening of its lifetime, primarily attributed to iodine chemistry, and partly due to a reduction of ozone production, which is a consequence of the depleting effect of halogens on NO_x levels (Caram et al., 2023; Iglesias-Suarez et al., 2020; Sherwen et al., 2016). Since this influence of halogens is ignored in our model, the good agreement of the model with in situ NO_x measurements (Figs. 10 and
605 11) might mean that our NO_x emissions are actually too low, although a definitive assessment is not possible at this stage. Although the absence of halogen chemistry likely contributes to the ozone overestimation, other factors like uncertainties in emissions, model resolution, and vertical mixing may also influence the bias.

3.3 Comparisons with PTR-MS measurements

Figs. 12 and 13 show modelled VOC concentrations (from simulations R0, R2, S3, S4 and S5) against observations at the
610 Maïdo observatory for the months of January and July 2019. The averaged diurnal cycles of observed and simulated concentrations are shown on Fig. 14. For most species, the general evolution is well represented by the model. Both the observed and modelled diurnal cycles display a pronounced daytime maximum for almost all species. This general behavior reflects the dominance of daytime sources for most compounds but also the weaker influence of surface emissions during the night, when the station of Maïdo is frequently located in or near the free troposphere (Verreyken et al., 2021).

615 3.3.1 Formaldehyde

Both the model and the measurements display a pronounced diurnal cycle of formaldehyde. The first and last days of each month show the best agreement with measurements, matching both day- and nighttime values closely. On the other days, there is a consistent underestimation, especially at night, when it can reach a factor of two. This underestimation is unexpected, given the model overestimation of FTIR columns (Sect. 3.4) and the fact that nighttime observations at Maïdo reflect mostly
620 free tropospheric air. Previous model calculations suggest very little contrast between daytime and nighttime HCHO columns at Maïdo, due to the long photochemical lifetime of this compound during the night (Stavrakou et al., 2015). Formaldehyde remains mostly stable between the different sensitivity runs, with similar statistics (Table S5). Being mainly produced from the oxidation of other VOCs, including methane, both the simulated and observed formaldehyde concentrations are highest in January (austral summer), due to higher solar radiation fluxes and biogenic VOC emissions at that time.

625 3.3.2 Methanol

The simulated methanol from the standard run (R0) generally matches the observations very well, except for a slight average positive bias, very similar in January and July (0.2 ppb, see Table S5). This good agreement and the much higher bias (~0.6 ppb) of the S3 run which adopted unadjusted biogenic emission factors ($800 \mu\text{g m}^{-2} \text{h}^{-1}$ for all PFTs) appear to validate the halving of the emission factor for the dominant PFTs in run R0 (see Sect. 2.3.2). The lower biogenic emissions are in line with
630 the MEGAN recommendation (Stavrakou et al., 2011). The boundary condition adjustment is also validated, since the simulation adopting unadjusted values (S2) also displayed higher bias (0.3 ppb) and root mean square error (RMSE) than the R0 simulation. The budget of methanol over the island is largely dominated by the biogenic source, since anthropogenic emissions represent only ca. 1% of the surface emissions of methanol in the model (Fig. 4), and the photochemical production of methanol from the reaction of methylperoxy radicals (CH_3O_2) with itself and with other peroxy radicals (Jacob et al., 2005)
635 is only a minor source (Müller et al., 2016; Bates et al., 2021). Note that the formation of methanol from the reaction of CH_3O_2 with OH (Archibald et al., 2009), which was only recently shown to be a significant source (Bossolasco et al., 2014; Müller et al., 2016; Bates et al., 2021; Caravan et al., 2018), is not included in the model.

Although the R0 simulation performs very well on most days, large positive biases are found during specific periods (e.g. 29-30 January and 2-9 July). Those discrepancies are likely not related to the emissions, since other compounds (e.g. Iox and
640 acetaldehyde) are similarly overestimated during the same period, which suggests an issue with the representation of meteorology and possibly tracer transport. Cloudiness (in January) and relative humidity (in both months) are overestimated at Maïdo during those periods of low model performance (Fig. 8).

3.3.3 Isoprene and monoterpenes

Isoprene is generally well reproduced in WRF-Chem during both months, giving credence to the standard emission factor
645 distribution used in the model. The seasonal variation is correctly predicted, with a factor of 3 higher concentrations in January

compared to July in both the model and the observations. This difference is primarily due to the higher temperatures and radiation fluxes during January (Table S2). Model underpredictions of isoprene are found on days with very low simulated solar radiation due to excessive cloudiness, in particular on Jan. 1-3, Jan. 28-30 and July 27, 29 and 30 (Figs. 12 and 13). Conversely, on days with high observed cloudiness and overestimated modelled radiation fluxes, e.g. on July 7-8, the simulated isoprene is too high. Those patterns reflect the strong control of solar radiation on isoprene emissions in MEGAN. Dufлот et al. (2019) and Verreyken et al. (2021) reported that westerlies are generally associated with higher isoprene abundances at Maïdo, compared to easterlies, presumably because of the closer proximity of vegetation west of Maïdo. However, easterlies were frequently associated with low solar radiation fluxes due to cloudiness, especially in January (1-3 and 27-30 Jan., and to a lesser extent on 9 and 12 July), such that the influence of wind direction on isoprene cannot be established for the period considered here. There is essentially no variability in isoprene concentration between the different sensitivity runs, except for simulation S4, which predicts slightly higher concentrations around midday (+5% at noon), due to the much weaker OH-recycling in isoprene oxidation in the original MOZART mechanism, compared to the modified mechanism used in R0 (see Sect. 2.2.3). Whereas the average midday isoprene concentration is about 30% too low in the model, possibly due to a small underestimation of the isoprene emission factor, the concentrations in early morning (5-6 UT i.e. 9-10 LT) and late afternoon (17-18 LT) are too high (Fig. 14). This feature, also found for other biogenic compounds such as methanol, Iox (MVK+MACR+ISOPOOH) and monoterpenes, might be due to insufficient boundary layer mixing.

The model performance for the isoprene oxidation products Iox (MVK+MACR+ISOPOOH) is similar as for isoprene. On many days for which isoprene is too low or too high, so are its oxidation products. There are exceptions to that pattern, such as the first and last days of January, when isoprene is strongly underestimated due to excessive cloud cover, whereas Iox is not. This difference might be related to wind transport errors in the model, as suggested by the poor model performance for wind direction on those days (see above), or it might be due to the longer lifetime of Iox compared to isoprene (Verreyken et al., 2021) implying that Iox is influenced by more distant sources.

Examination of the average diurnal cycle of Iox concentrations (Fig. 14) shows that, while the PTR-MS data show an almost complete disappearance of Iox during the night, WRF-Chem calculates low, but non-negligible concentrations, especially in winter. This discrepancy is even larger in the sensitivity simulation S3. While isoprene remains relatively unchanged between the sensitivity runs, Iox shows important changes in the average diurnal shape between R0 and the S3 run (unadjusted biogenic emissions) and S4 run (unadjusted chemistry). As seen in Table 2, the high-NO_x yield of the sum MVK+MACR is much higher (+50%) in the updated chemical mechanism of R0 than in the MOZART-4 mechanism used in S4, which explains the higher Iox concentrations of R0 (by about 10%). Conversely, the simulation S3 predicts significantly higher Iox levels than R0, by ~30% near midday in July and up to a factor of 5 during the night. The improved diurnal cycle of R0 is due to the strong decrease of the monoterpenes emission factor and their increased light-dependence factor in the R0 simulation, leading to reduced emissions, especially during the night. Indeed, the MOZART-4 mechanism includes a source of MVK and MACR originating from the reaction of lumped monoterpenes with ozone, OH and NO₃:





Since monoterpenes are not a source of MVK or MACR (see e.g., <https://mcm.york.ac.uk/MCM/>), this artificial source of Iox causes a model overestimation of Iox in both simulations, but especially in simulation S4, and even more during the night and
685 in July, when the lower radiation fluxes cause a stronger decrease of isoprene emissions, compared to monoterpenes. Both the RMSE and mean bias of R0 are strongly reduced relative to run S4, and an even better agreement would likely be achieved without the artificial source of MVK and MACR from monoterpenes in the model.

The observed ratio of Iox to isoprene concentrations (0.74, based on concentrations averaged between 12 and 16 LT) is reproduced fairly well by the R0 run in January (0.82), whereas a higher ratio (1.05) is derived from run S3, due to its higher
690 monoterpenes emissions and therefore higher Iox production from their oxidation. In July, the Iox to isoprene ratios are more dissimilar (0.62 based on observations, 1.0 and 1.54 for the R0 and S3 runs), due to the strong reduction of isoprene emissions and higher share of monoterpenes to the the Iox budget in winter, compared to summer.

The much improved agreement for monoterpenes after the biogenic emission adjustments (Table S5 and Fig. 14) appears to validate the lower emissions, especially during the night. A precise assessment is difficult given the large difference between
695 the PTR-MS concentrations derived from the two signals (m/z 137 and 81, corresponding with protonated monoterpenes and for their main fragments, respectively). These differences might result from e.g. temporal variations in monoterpene distribution or from contributions of other compounds to the m/z signal. The LDF correction improved the temporal correlation between model and observation through the strong reduction of nighttime concentrations, although, as for isoprene, the model simulates early morning and late afternoon concentration peaks that are not seen in the observations. On average, the modelled
700 values in R0 fall between the two measured signals, i.e. they overestimate (by 8-17 pptv) the monoterpenes detected at m/z 137 and underestimate (by 2-11 pptv) the determination based on the m/z 81 signal.

3.3.4 Methyl ethyl ketone and methylglyoxal (m/z 73)

Although both methyl ethyl ketone (MEK) and methylglyoxal (MGLY) contribute to the 73 m/z signal, MEK is dominant according to the model results. As seen on Fig. S3, the modelled MGLY concentrations are highest near local noon, and
705 account for at most 25% (15%) of the total signal in January (July) in simulation R0, taking into account that the PTR-MS is less sensitive to MGLY (by a factor of 0.7) compared to MEK (Sect. 2.4.2). The observed 73 m/z signal displays a pronounced diurnal cycle (Fig. S34), with daytime concentrations about 3-5 times higher than during nighttime. Although the model simulation without any direct MEK emission (run S5) reproduces this large night/day difference in January, this run strongly underestimates the observations, and it underestimates the amplitude of the diurnal cycle in July (Fig. 14). The photochemical
710 production of MEK in the model originates exclusively from the oxidation of the surrogate compound BIGALK by OH. The emissions of BIGALK includes the sources of all $\text{C}_{\geq 4}$ alkanes, but its oxidation mechanism is that of $n\text{-C}_4\text{H}_{10}$, a well-known major precursor of MEK (Yáñez-Serrano et al., 2016; Jenkin et al., 1997; Sommariva et al., 2011). Among the higher alkanes,

only n-butane and 3-methyl propane are significant MEK precursors (Sommariva et al., 2011; <https://mcm.york.ac.uk/MCM/>). Therefore, since n-butane and 3-methyl propane emissions make up only a fraction of total BIGALK emissions, of the order of 34% (Stavrakou et al., 2015), the MEK photochemical production in the model is likely overestimated. MGLY has no direct source in the model, but it is photochemically produced, mainly from the oxidation of isoprene and other BVOCs (Mitsunishi et al., 2018). Given its short lifetime (~1 hour), primarily due to photolysis (Fu et al., 2007), MGLY displays a pronounced diurnal cycle with a noon maximum. The molar yield of MGLY in isoprene oxidation is of the order of 0.25 (Fu et al., 2007), and it is unlikely to be much underestimated. Therefore, the high MEK/MGLY mixing ratios observed during the day, ca. 0.07 and 0.05 ppb in January and July, respectively, are best explained by the presence of a substantial biogenic source of MEK. This source has been taken equal to 3% of the biogenic isoprene emissions in run R0 (see Sect. 2.3.2). The diurnal shape of modelled MEK (run R0) in this run is similar to the observations, except that it underestimates the observed diurnal amplitude in July (Fig. S3). Furthermore, in January, the modelled concentrations rise too early in the morning and decrease too rapidly in the afternoon, suggesting that the biogenic emissions of MEK are delayed compared to those of isoprene. This delay is of the order of 2 hours and is qualitatively in line with the proposal that MEK is released through the uptake of isoprene oxidation products (MVK and the 1,2-ISOPROOH isomer) by vegetation and their subsequent conversion and re-emission as MEK and other compounds (Cappellin et al., 2019; Tani et al., 2020; Canaval et al., 2020). The PTR-MS signal for Iox (the sum MVK+MACR+ISOPROOH) is slightly delayed (by 1-2 h) compared to isoprene, as shown on Fig. S4, and the delay for MVK is expected to be longer than for Iox, due to the lower rate of the reaction of MVK with OH, compared to MACR and ISOPROOH (e.g. Müller et al., 2019). The adopted ratio between MEK and isoprene biogenic emissions (3% on a mass basis) is larger than the value of 1.5% derived by Canaval et al. (2020) based on enclosure measurements on gray poplar trees and field eddy-covariance measurements at two forested sites. This discrepancy could be partly due to uncertainties and natural variability of MVK/ISOPROOH deposition velocities and conversion rates to MEK in plant leaves. In addition, the good model agreement of the R0 run could possibly be achieved with a lower MEK/isoprene emission ratio, e.g. if the contribution of MGLY to the PTR-MS signal would be higher. In addition, the transport of chemical compounds from source regions (e.g. cities) to Maïdo is imperfectly represented in the model due to its relatively coarse resolution (2.5km), which may cause errors in the diurnal cycle of advected species, in particular for MEK which has a significant anthropogenic component (Fig. 4) (Bon et al., 2011; Brito et al., 2015). Finally, note that the model ignores the potentially significant oceanic source of MEK, which is however very uncertain (Brewer et al., 2020). The S4 run leads to higher MEK/MGLY mixing ratios than R0 around midday (Fig. 14), due to differences in the isoprene chemical mechanism (Sect. 2.2.3); the original MOZART-4 mechanism (used in S4) has a higher yield of C₅-hydroxyaldehydes (HYDRALD), which generate slightly more MGLY than other isoprene oxidation products such as MVK and MACR (Emmons et al., 2010).

In July, the model overestimates both nighttime and daytime observations in run S2, with unadjusted lateral boundary conditions. This overestimation motivated the adjustment (multiplication by 0.4) of initial and boundary conditions for MEK (Sect. 2.2.4). This adjustment has much less effect in January, due to its shorter photochemical lifetime in summer, compared to winter.

3.3.5 Acetaldehyde

Regarding acetaldehyde, WRF-Chem largely underestimates the PTR-MS concentrations in January, whereas the model is essentially unbiased on average in July (-0.009 ppb), but correlates very poorly with the observations ($r=0.39$, Table S5). The primary sources of acetaldehyde over the island, including anthropogenic and biogenic direct emissions, and photochemical production, are detailed in Table S6 of the Supplement. The temporal correlation improves markedly (to ca. 0.75 in January and July) when the direct emissions of acetaldehyde and other VOCs are turned off (run S5), even though the mean bias increases. This finding suggests that direct emissions of acetaldehyde are overestimated, while photochemical production might be too low. Acetaldehyde emissions in the model are largely due to the residential sector (Fig. 4). Aldehydes are the main NMVOC class released by the residential sector, and these emissions are mainly due to biomass burning combustion (Huang et al., 2017). The model uses the combined emissions of all higher ($C_{\geq 2}$) aldehydes as anthropogenic emissions of CH_3CHO . However, acetaldehyde accounts for only a small fraction of total aldehyde emission from boilers (7-17%) (Macor and Pavanello, 2009) and from the heating of cooking oils (at most ~1%) (Katragadda et al., 2010; Takhar et al., 2023). The anthropogenic emissions of CH_3CHO used in the model are therefore strongly overestimated, possibly by one order of magnitude or more. Without this lumping, we expect that the model would underestimate the PTR-MS data by almost one order of magnitude, suggesting that other sources of acetaldehyde are likely strongly underestimated. This would be in line with previous assessments based on in situ data at remote marine locations (e.g., Wang et al., 2019; Travis et al., 2020). For example, at Cape Verde in the Atlantic Ocean, the CAM-Chem model was found to underestimate in situ concentration measurements by a factor 30 (Read et al., 2012), and the discrepancy was attributed to both oceanic acetaldehyde emissions and photochemical production from unknown, relatively long-lived precursors. Direct oceanic emissions might contribute significantly to the signal at Maïdo (Millet et al., 2010; Read et al., 2012), given the relative proximity of the sea (~20km) and the relatively long lifetime of CH_3CHO (2-9 hours, with $[OH] = (2-10) \times 10^6 \text{ molec. cm}^{-2}$, see Fig. 9). Photochemical production of acetaldehyde in the model proceeds mainly from the oxidation of alkanes, alkenes and ethanol (Millet et al., 2010). The total production of CH_3CHO calculated using the model (R0 simulation) over Réunion Island averaged over January and July amounts to 19, 0.92, 14, 13, 4.2, 3.86 and 18 tons per month for ethane, propane, propene, higher alkenes (BIGENE), higher alkanes (BIGALK), MEK and ethanol, respectively. Unfortunately, only MEK and ethane were measured at Maïdo, and large model underestimations cannot be excluded for any of the other precursors. For example, the average mixing ratios of propane and BIGALK calculated by the model at Maïdo are only ca. 20 and 30 ppt, respectively, whereas long-term in situ measurements at other southern tropical marine sites (Seychelles and Ascension) are considerably higher, of the order of 100 ppt in January and 50 ppt in July for propane, and about 150-250 ppt in January and 100-200 ppt in July for the total concentration of measured butanes and pentanes ($n\text{-C}_4\text{H}_{10}$, $i\text{-C}_4\text{H}_{10}$, $n\text{-C}_5\text{H}_{12}$, $i\text{-C}_5\text{H}_{12}$) (Helmig et al., 2021). Biogenic emissions of acetaldehyde and ethanol are included in the model but are very uncertain (Millet et al., 2010) and might be underestimated. As suggested by Travis et al. (2020), the oceanic emissions of various acetaldehyde precursors (primarily ethanol, alkanes,

alkenes) might partially explain the negative model biases against aircraft in situ data across the globe and particularly at remote locations and in the free troposphere. These emissions are currently ignored in the model simulations.

3.3.6 Acetone


Due to its relatively long lifetime (several weeks) (Jacob et al., 2002; Fisher et al., 2012), acetone is sensitive to the lateral boundary conditions, especially in July when solar radiation and OH levels (Fig. 9) are at their minimum. The large positive model bias of the S2 run in July is reduced by a factor of 10 after adjustment of boundary conditions (R0 run). Acetone is relatively insensitive to this change in January, likely due to its shorter lifetime and larger biogenic emissions during summer. Acetone is sensitive to a removal of its direct emissions (S5), which have a small anthropogenic component (~3 tons/month) and a dominant biogenic contribution (55 tons/month, Fig. 4). The secondary source of acetone from photochemical production is comparatively weak over the island (17 tons/month on average for the two months). The principal precursors are BIGENE, BIGALK, propane and monoterpenes, with productions amounting to 8.5, 3.4, 3.5 and 1.7 tons, respectively. As in the case of acetaldehyde, the contribution of propane and BIGALK oxidation is likely strongly underestimated. In agreement with previous studies, (e.g. Wang et al., 2020), the direct biogenic source of acetone is dominant over the secondary production due to monoterpenes and anthropogenic VOCs. This source is poorly constrained and therefore very uncertain, the acetone emission factor being constant for all PFTs in MEGAN, except for grasses and crops (Guenther et al., 2012). The ocean/atmosphere exchange of acetone plays a substantial role over the remote troposphere (Wang et al., 2020) but is neglected in our study. The diurnal cycle of acetone mixing ratios is underestimated by the model during both months (Fig. 14). Since photochemical production plays only a minor role for this compound, this underestimation might be due to either a misrepresentation of transport patterns (Fig. 8) or an underestimation of biogenic emissions during daytime. The light-dependent fraction (LDF) for biogenic acetone is only 0.25 in the model, implying little diurnal variability of the emissions. Comparison of the model performance for acetone (Fig. 12) with the meteorological evaluation (Fig. 8) does not suggest a clear link between the modelled diurnal cycle and meteorological parameters, in particular wind direction.

3.3.7 Acetic acid, peracetic acid and glycolaldehyde (m/z 61)

The m/z 61 signal exhibits a pronounced diurnal shape in January and July, with a clear daytime maximum, and nighttime values approximately 5 times lower than the midday peak (Fig. 14). Although this signal is mainly attributed to acetic acid, there are interferences, mainly from glycolaldehyde (GLYALD) and peracetic acid (PAA). Here, this measurement is compared against the sum of acetic acid, GLYALD and PAA from WRF-Chem. The contributions of the three species to the combined modelled concentration are displayed on Fig. 15. The model strongly underestimates the magnitude of the signal, by a factor of 3-4, but it replicates the observed diurnal and seasonal variation. All three species have their highest concentrations during daytime in January, when photochemical activity and biogenic emissions are at their highest. Acetic acid has only a weak biogenic source (5 tons/month on average over the island, Fig. 4), in agreement with previous studies (e.g. Paulot et al., 2011), while all three compounds have significant photochemical production terms. GLYALD is produced

at a high yield (~0.3 at high NO_x) by the oxidation isoprene by OH and has a short chemical lifetime (a few hours during the day), which explains that its nighttime concentrations are very low (Fig. 15). Given the large isoprene emissions, GLYALD is the main contribution to m/z 61 during the day according to the model. The oxidation of monoterpenes also leads to GLYALD formation in the MOZART-4 mechanism, which explains the higher GLYALD concentrations of the S3 run, which uses
815 unadjusted (e.g. 5-times higher) emissions of monoterpenes.

The contribution of the two acids shows less diurnal variation than GLYALD, particularly in July. Photochemical production of these two compounds over the island amounts to 19.5 and 62.4 tons/month for acetic acid and PAA, respectively, on average for the two months. It is mainly due to the reaction of the acetylperoxy radical (CH₃CO₃) with HO₂, represented in the model as




This reaction alone accounts for 79% and 94% of the total production of acetic and peracetic acid over the island, respectively. The model ignores the radical-forming channel,




(followed by CH₃CO₂ (+O₂) → CH₃O₂ + CO₂), which was shown to account for about half of the total reaction (Jenkin et al.,
825 2007). The total rate constant of the reaction, 1.4×10⁻¹¹ molec.⁻¹ cm³ s⁻¹ at 298 K in MOZART-4, is also likely underestimated by a factor of ca. 1.5 compared to more recent estimates (Gross et al., 2014). These approximations might contribute to a moderate overestimation of acetic and peracetic acid production from the reaction above, although the probable underestimation of alkanes and other VOCs noted above likely implies an underestimation of the production rate of CH₃CO₃ radicals. In addition, many reactions of CH₃CO₃ radicals with other organic peroxy radicals are neglected in the mechanism,
830 although they might contribute significantly to acetic acid production (Khan et al., 2018). Furthermore, the rate constant of the reaction of peracetic acid with OH was recently shown in a combined experimental and theoretical study (Berasategui et al., 2020) to be much lower (factor of ~30) than previously determined, implying a much longer lifetime in the troposphere.

The model underestimation of the m/z 61 signal is consistent with (1) the suggestion of underestimated or unknown sources of acetic acid based on model comparison with in situ measurements from aircraft and at surface stations (Paulot et al., 2011;
835 Khan et al., 2018), and (2) the observations of high PAA concentrations over remote oceans during the Atmospheric Tomography Mission (ATom) (Wang et al., 2019). Given that acetaldehyde is a known major precursor of PAA, box model calculations by Wang et al. (2019) showed that these high levels of PAA are fully consistent with the (simultaneously observed) acetaldehyde measurements. As noted above, there is abundant evidence of a large source of acetaldehyde over the oceans.



840 **Figure 12.** Time series of observed and modelled concentrations of PTR-MS measured species in January. All concentrations are shown in ppb. The model results are shown for simulations R0, S2, S3, S4 and S5 (see Table 4). Both monoterpene measurements are depicted, differentiated by solid (m/z 81) and dashed black lines (m/z 137).



845 **Figure 13.** Same as Fig. 12, for July 2019.




Figure 14. Average diurnal cycle of organic compounds measured at Maïdo observatory, and corresponding modeled concentrations from the 30-day runs defined in Table 4 (except S1). The concentrations are shown in ppb. Both monoterpene measurements are depicted on the same figure, differentiated by solid (m/z 81) and dashed black lines (m/z 137).




Figure 15. Averaged diurnal cycle of the m/z 61 signal in January (left) and July (right), and comparison with the modelled concentrations of acetic acid, GLYALD and PAA, which are all assumed to contribute to the observed signal. The PTR-MS sensitivity is assumed identical for the three compounds.

3.4 Evaluation against FTIR columns

The model is evaluated against FTIR data in Fig. 16. The model values are sampled at the same times as the measurements and smoothed using the averaging kernels and a priori profile of the FTIR retrieval. The results of two simulations are shown: R0, the standard run with all model updates, and S2, identical to R0 but without the adjustments to the initial and boundary conditions (ICBC). The results of the other simulations differ only marginally from those of the R0 run, and are therefore not shown. ICBC adjustments were made only for CH_3OH , PAN and C_2H_6 ; for the other compounds, the discrepancies are generally small.

For HCHO, a model overestimation is noted during both months, amounting to 8 and 2×10^{14} molec. cm^{-2} in January and July, respectively. In January, this bias exceeds the reported systematic uncertainty of the FTIR column ($\sim 2.5 \times 10^{14}$ molec. cm^{-2}) (Vigouroux et al., 2018) and is not likely due to ICBC errors for HCHO, given the short lifetime of this compound (a few hours). The monthly averaged retrieved columns (3.2 and 1.61×10^{15} molec. cm^{-2} in January and July, respectively) are similar to the reported FTIR columns at Maïdo for the same months in previous years (Vigouroux et al., 2018; 2020). The weak sensitivity of the modelled columns to changes in local VOC emissions (e.g. from run S5) suggests that the cause of the model overestimation is due either to HCHO precursors that are not well represented in the model, or to background HCHO, i.e. primarily the contribution of methane oxidation, above ~ 2 km altitude (the altitude of Maïdo). The addition of lightning NO emissions (run S6) aggravates the bias, as it increases the HCHO column (+24% in January, see Fig. S5). The HCHO enhancement due to lightning NO is mostly located in the mid-troposphere, between 4 and 11 km (Fig. S6) and is primarily caused by the increase in OH consecutive to the added NO_x emissions in this altitude range, as the OH radical promotes

methane oxidation and its associated HCHO production. At lower altitudes, OH levels are less impacted, whereas at higher levels, HCHO is more controlled by deep convection of low-altitude pollutants to higher altitudes (Bozem et al., 2017; Fried et al., 2008).




Figure 16. Time series of FTIR-measured and modelled HCHO, CH₃OH, C₂H₆, CO, PAN and tropospheric O₃ columns in January (top rows) and July (bottom rows) 2019. The symbols denote the individual FTIR measurements and corresponding WRF-Chem results, while the solid lines (red for R0, cyan for S2) show the 24-hr averages.

Given the longer atmospheric lifetimes of the other compounds shown on Fig. 16, the model performance for these species is very dependent on their boundary and initial conditions, obtained from either CAM-Chem (for CH₃OH) or CAMS (for the other compounds). Without any adjustment, the model agreement with observed CO and tropospheric O₃ columns is reasonable, with biases generally well below 10% for CO and 20% for O₃. The simple ICBC adjustments, presented in Sect.

2.2.4, succeed generally very well in reducing the gap between model and observations for the other compounds. Since the ICBC adjustments consist of a single scaling factor for each species, the short-term variability of the observations remains
885 poorly reproduced by the model after adjustment, e.g. for CH₃OH. Nevertheless, the model correlates well with the daily-averaged data for O₃ (Pearson's correlation coefficient $r=0.6$ and 0.7 in January and July), and to a lesser extent also for C₂H₆ ($r=0.7$ and 0.3).

The monthly averaged FTIR PAN columns, ca. 1×10^{15} molec. cm⁻², are lower, but of the same order of magnitude as those typically observed above the Jungfraujoch station in the Swiss Alps during summer (up to 4×10^{15} molec. cm⁻²) (Mahieu et al.,
890 2021). They are also significantly lower than those observed by the Infrared Atmospheric Sounder Interferometer (IASI), typically $>2 \times 10^{15}$ molec. cm⁻² above oceanic areas around Réunion Island (Franco et al., 2018), although this difference is likely partly due to the high altitude of the station. The spatial distribution displayed by IASI suggests a substantial influence of long-range transport of African emissions on PAN levels in the region. Although PAN is short-lived in the lower troposphere due to fast thermal decomposition, it is much longer-lived in the cold upper troposphere, where it can be transported over long
895 distances. Due to this long lifetime, a large fraction of the column lies at those high altitudes, which explains why the modelled PAN column at Maïdo responds strongly to the ICBC change between simulations S1 and R0. Note that this does not imply a strong influence of ICBC on PAN and NO_x close to the surface, where the PAN lifetime is short. The modelled columns of the S1 run overestimate the FTIR data by about a factor 2, providing the justification for the adjusted ICBC for PAN in simulation R0. However, both FTIR data and the R0 run underestimate IASI columns by a factor of 2 or more. The
900 acknowledged uncertainties of the PAN retrievals by FTIR and IASI suggest that the ICBC adjustment for PAN should be considered as very uncertain.

3.5 TROPOMI

The distribution of NO₂ and HCHO columns obtained from the oversampling of TROPOMI retrievals over a period of more than four years is displayed on Fig. 17. Despite the much lower NO₂ columns retrieved over and around the island ($<1.5 \times 10^{15}$
905 molec. cm⁻²) in comparison with e.g. Europe (typically $(2-10) \times 10^{15}$ molec. cm⁻²) (Poraicu et al., 2023; Lange et al., 2023), sharp gradients are observed, reflecting the strong heterogeneity of NO_x emissions. A very well-localized maximum (1.5×10^{15} molec. cm⁻²) is found at the precise location of the Le Port power plants (Fig. 17), whereas the other thermal power plants, located near Sainte-Suzanne (Bois Rouge) on the northern coast, and near Saint-Pierre (Le Gol) on the southern coast, are not clearly detected. The emissions from the Le Port power plants (and to a lesser extent from the cities of Le Port, Saint-Denis
910 and Saint-Paul) generate a broad hot spot of NO₂ columns exceeding 1×10^{15} molec. cm⁻², which extends over land as well as over sea over distances of the order of 5-10 km. This distribution confirms the dominance of the Le Port power plants emissions over those from the other point sources, and justifies the redistribution of industrial emissions based on the EDGAR 6.1 dataset described in Sect. 2.3.1. The NO₂ column values around Maïdo are significantly lower, of the order of 0.5×10^{15} molec. cm⁻². Even lower columns are observed in the southeastern part of the island, around the volcano (Piton de la Fournaise).




Figure 17. Oversampled NO₂ and HCHO columns from TROPOMI, covering 2018-2022. The resolution is 0.01°×0.01°.

The oversampled distribution of TROPOMI HCHO is noisier compared to NO₂, but also shows well-defined features and gradients. The largest columns ($\sim 5 \times 10^{15}$ molec. cm⁻²) are found in a large area along the northwestern coast and in particular near the Le Port power plants. Since the power plants are not a large VOC source, and since the HCHO distribution shows little correlation with the anthropogenic VOC emission distribution (Fig. 6), these high columns clearly do not reflect the distribution of VOC emissions. Instead, they are likely due to the influence of anthropogenic NO_x leading to higher OH levels and therefore to enhanced HCHO production rates from the oxidation of long-lived hydrocarbons, especially methane. Methane oxidation is by far the largest source of HCHO at the global scale (Stavrakou et al., 2009), and to a large extent also above Réunion Island (see further below), even though non-methane hydrocarbons are the dominant source over most continental areas. This dominance of the background is the main reason for the many common features shared by the HCHO and NO₂ distributions from Fig. 17, such as the high values to the west and northwest of the island, and the lower columns to the east and southeast of the island and over several specific areas in the island interior. Besides this likely dominant contribution of the methane background to HCHO, the TROPOMI distribution also shows evidence of elevated columns due to isoprene emissions, in particular along the eastern coast (see Fig. 5).

The model is evaluated against NO₂ columns from TROPOMI in January and July 2019 in Fig. 18. The domain shown on Fig. 18 encompasses both Réunion and Mauritius Island, located to the Northeast of Réunion, at a distance of ca. 200 km. The results of two simulations are shown: the reference run without lightning (R0) and the simulation with lightning emissions (S6). The modelled columns of run S1 (with higher NO_x emissions from the Le Port power plants) are given in the Supplement (Fig. S7). The TROPOMI averaging kernels were applied for both compounds, and the model averages were calculated from values sampled at the same times as the TROPOMI monthly averages. Both modelled and retrieved columns were regridded to 0.1° resolution.

Both the model and TROPOMI data display higher NO₂ columns on Mauritius Island, compared to Réunion. Based on EDGARv6.1 emission data, the main hotspot on Mauritius is due to the energy sector, with several fossil fuel power plants located in Port-Louis, the capital city. The model comparison with TROPOMI strongly suggests that the EDGAR NO_x

940 emissions are overestimated. Those emissions generate a large plume extending towards the Southwest (in January) or the West (in July), but its impact on Réunion appears to be limited, based on the distributions shown on Fig. 18.

Over Réunion, both R0 and S6 model simulations reproduce the main spatial patterns observed in the measurements, namely, the maximum at Le Port and over adjacent regions, primarily the eastern and (to a lesser extent) the southeastern coast in January, and the northern coast, in July. A secondary NO₂ maximum is also seen along the southern coast near Saint-Pierre

945 and especially the thermal power plant of Le Gol. The model also reproduces the minimum seen in the eastern part of the island, in January, and near the southeastern extremity of the island, in January. Overall, both R0 and S6 overestimate the observations in the winter, but there is an underestimation for R0 during summer (-11% and +64% for the island-averaged column of R0 and S6 in January, respectively and +25% for both in July). The discrepancy is however locally much higher, e.g. near Maïdo in July (factor of ~1.8 at the two pixels nearest to Maïdo), likely due to the export of pollution from the Le

950 Port area. Caution is warranted when comparing model and satellite over remote locations, as the satellite columns are very noisy, despite their spatial (0.1°) and monthly averaging, especially over the mountainous island interior. The TROPOMI NO₂ uncertainties are of the order of 5×10^{14} molec. cm⁻² (see Fig. S8), i.e. they are of the same order as the retrieved columns over most of the island except the strongest anthropogenic hotspots. The model correlates better with the data in July ($r = 0.72$ for R0 over the inner domain, see Fig. 18) than in January ($r = 0.43$), likely due to the higher values in July. The model reproduces

955 the observed seasonal variation of NO₂ columns, with higher values found in July (winter) compared to January (summer), in response to changes in OH radical concentrations (Fig. 9). The R0 run provides a significantly better match with TROPOMI than the S1 run, which strongly overestimates the columns along the western and southwestern coast (Fig. S7), by a factor of 2, as well as over the sea, west and southwest of the island. Those overestimations essentially disappear in the R0 run, which assumes a 5-fold reduction of the NO_x emissions due to the Le Port power plants.

960 The causes for the slight NO₂ model overestimation of the R0 run in July are unclear. The NO_x emissions could be overestimated, the NO_x lifetime could be too long, or the TROPOMI NO₂ columns might be too high. Unfortunately, most TROPOMI NO₂ validation studies were conducted at mid-latitudes, in regions with strong pollution sources (Verhoelst et al., 2021; Poraicu et al., 2023; Lange et al., 2023; van Geffen et al., 2022a). Those studies have generally reported the slope (s) and intercept (i) of linear regressions of the type $C = i + s C'$, where C denotes the TROPOMI column and C' the co-located

965 independent measurement. Most validation studies reported positive values of the intercept i and slope values s lower than unity, suggesting that TROPOMI underestimates high values and overestimates very low values. More studies are needed to better characterize the potential biases of TROPOMI NO₂ in remote areas.




Figure 18. Monthly-averaged NO₂ columns from TROPOMI and WRF-Chem (simulations R0 and S6) for January (top row) and July (bottom row), regridded to 0.1° resolution. The WRF-Chem averages account for the TROPOMI sampling times and for the averaging kernel corresponding to each overpass. The white line denotes the boundary of the inner model domain (d02).

Implementation of lightning emissions (run S6) leads to a strong column enhancement in January, while its impact is negligible in July. The S6 run worsens the model overestimation over land, and leads to overestimated columns over ocean. For example, the average column from run S6 over an oceanic area north of Réunion is 6.3×10^{15} molec. cm⁻², a factor of 1.75 above the TROPOMI average of 3.61×10^{15} molec. cm⁻², whereas the R0 run is a factor of 2 too low (1.83×10^{15} molec. cm⁻²). The lightning emissions flip the seasonality of NO₂ columns over ocean, with higher values predicted in January, while both TROPOMI and the R0 run suggests a wintertime maximum. In January, lightning NO_x is responsible for increasing the average mixing ratio of NO₂ in the upper troposphere by a factor of 4, compared to the R0 simulation (Fig. S6). Note that the default WRF-Chem settings would lead to much larger lightning emissions since, following Barten et al. (2020), we decreased the number of flashes by a factor of 10, and we also reduced the NO production per flash to 250 moles. Nevertheless, the S6 run substantially overestimates the upper tropospheric (UT) NO₂ mixing ratios obtained from TROPOMI using the cloud slicing technique (Marais et al., 2021; Horner et al., 2024). Indeed, the TROPOMI-based UT mixing ratios (between 180 and 450 hPa) are typically 40-50 pptv above the Indian ocean around Réunion Island during Dec. 2019 - Jan. 2020 (Marais et al., 2021), whereas the model averages over this region are 28 and 120 pptv in the R0 and S6 runs, respectively. Both the model comparison with the tropospheric columns (Fig. 18) and with the UT mixing ratios based on TROPOMI indicate that the lightning source of the

S6 run is too high, despite the reduction of flash count and number of moles per strike. In July, lightning emissions are largely insignificant, with less than 2.5% increase in NO_x in the upper troposphere.

Much like for TROPOMI NO₂, the model evaluation against TROPOMI HCHO data shows a relatively good agreement with
990 regards to spatial representation (Fig. 19), even though the spatial correlation coefficients are very low (e.g. $r = 0.2$ for the R0
run in January), largely due to the high noise in the data. In January, relatively high columns (of the order of 8×10^{15} molec.
 cm^{-2} in the R0 run, slightly less in the TROPOMI columns) are found to the southwest of the two islands, as well as along the
western coast of Réunion. These patterns mirror the NO₂ distribution, which also shows enhancements in those areas (Fig. 18).
Nevertheless, HCHO columns from both R0 and S6 show overestimations when compared to the observations. For example,
995 TROPOMI shows values close to 6×10^{15} molec. cm^{-2} along the northwestern coast of the island in January, slightly lower than
the modelled values close to 8×10^{15} molec. cm^{-2} . The discrepancy is close to the uncertainty provided with the retrieval product,
 2×10^{15} molec. cm^{-2} for the systematic component (trueness) and ca. 1×10^{15} molec. cm^{-2} for the random part (precision), when
accounting for the number of measurements used in the shown averages (see Fig. S9). This model overestimation is consistent
with the model overestimation against FTIR HCHO columns at Maïdo (see Sect. 3.4). As pointed out above, the main source
1000 of HCHO in this region is methane oxidation. Based on the model output, this production is estimated at about 3.0 and 1.2 Gg
per month over the land area of Réunion for January and July, respectively. The overestimation of HCHO levels in the model,
in comparison with TROPOMI and FTIR data, can probably be explained by an overestimation of OH levels in the model
simulations. Note that the addition of lightning in S6 increases the OH levels, and therefore worsens the overestimation of
HCHO columns in January. There might be several causes of the HCHO overestimation. NO_x might be too high, as suggested
1005 by the moderate overestimation of NO₂ columns against TROPOMI data, and this could impact OH. In addition, the neglect
of halogen chemistry in the model likely also leads to OH overestimation (Sherwen et al., 2016). Finally, it was recently found
out that the absorption of UV radiation by water vapor is more significant than previously assumed, and slows down the rate
of ozone photolysis and therefore the production and concentrations of OH radicals in the lower troposphere (Prather and Zhu,
2024). This process is particularly efficient at tropical latitudes where water vapor is most abundant. Note however, that this
1010 effect might be partially counterbalanced by the reduction of HCHO photolysis rates, following the decrease of UV radiation
levels.

The HCHO production due to methane oxidation might be compared with the contribution of isoprene, a major source of
HCHO at the global scale. Taking an average isoprene flux of $0.25 \text{ Gg month}^{-1}$ over the island (Fig. 4) and assuming 2.5
HCHO molecules produced for one molecule of isoprene (Stavrakou et al., 2009), the HCHO production from this source is
1015 estimated to be $0.28 \text{ Gg month}^{-1}$, i.e. almost an order of magnitude lower than the contribution of methane oxidation (see
previous paragraph). The secondary source of HCHO from anthropogenic VOC oxidation might be significant, but the
lifetimes of these compounds being much longer compared to isoprene, the HCHO production over the island is diluted due
to mixing, and partially exported away from the island.

Whereas the model overestimation of HCHO columns with respect to TROPOMI might be partially due to model errors, the TROPOMI columns are very low and noisy, and therefore close to the detection limit of the instrument. Taking the detection limit as three times the precision, most frequently in the range $(0.5-1.5) \times 10^{15}$ molec. cm^{-2} for 0.1° pixels over the island (Fig. S9), and taking into account that the TROPOMI precision was found to be underestimated by a factor of about 2 (1.6-2.3) in comparisons against FTIR data from a wide network of stations (Vigouroux et al., 2020), almost all TROPOMI monthly averages over the island (except around the Le Port hotspot) fall below the detection limit, which rationalize the high noise seen on the TROPOMI HCHO maps (Fig. 19).




Figure 19. Monthly-averaged formaldehyde columns from TROPOMI and WRF-Chem (simulations R0 and S6) for January (top row) and July (bottom row), regridded to 0.1° resolution. The WRF-Chem averages account for the TROPOMI sampling times and for the averaging kernel corresponding to each overpass. The white dotted line denotes the borders of the inner model domain (d02).

6. Conclusions

The WRF-Chem model has been used to compute the atmospheric composition in a region surrounding Réunion Island, in January and July 2019. The oversampled TROPOMI NO_2 distribution demonstrates the predominant influence of anthropogenic emissions on NO_x abundances over the island. In particular, the power plants near Le Port cause by far the strongest NO_x hotspot over the island, implying that their emissions largely exceed the emissions of any other point source or major city over Réunion. The model (R0 simulation) is only moderately biased against TROPOMI NO_2 columns (-11% in January, +25% in July) and reproduces the main observed patterns. To reach this agreement, the high-resolution (1 km^2)

anthropogenic emission inventory used as input to the model has been adjusted with regard to the repartition of energy sector emissions among the different power plants (in accordance with EDGAR v6.1), and furthermore, the NO_x emissions due to the Le Port power plants were downscaled by a factor of 5. This reduces the industry/energy sector emission estimates from
1040 Atmo-Réunion and EDGAR v6.1 by factors of 3.8 and 1.9, respectively. The emissions from the other power plants could not be constrained based on TROPOMI data. The 5-fold reduction above is a crude adjustment, but it is comforted by the model comparisons with both TROPOMI NO₂ and in situ NO_x measurements in the direct vicinity of the Le Port power plants. The fairly good model agreement against in situ NO_x measurements at other air quality stations lends confidence in the anthropogenic NO_x emission estimates for the other sectors (mostly traffic). In situ O₃ concentrations are overestimated by
1045 the model at the air quality stations, by ca. 6 ppb on average. This is likely due to the neglect of halogen (Cl, Br, I) emissions and chemistry in the model. These compounds were shown to deplete O₃ substantially in the tropical marine troposphere, by about 7 ppb, in line with previous work.

Over the ocean, the model comparisons with TROPOMI (for January) show the importance of the lightning source of NO. Accounting for this source (S6 run), or neglecting it (R0) leads to either overestimated or underestimated modelled NO₂
1050 columns in comparison to TROPOMI over sea; the comparison against upper tropospheric NO₂ mixing ratios based on cloud-sliced TROPOMI NO₂ data follows the same trend, and confirms the significant impact of lightning in this region in January. However, the lightning emissions are strongly overestimated by WRF-Chem with its current parametrization, even when the flash count is reduced by an order of magnitude in the model.

The model performance against the PTR-MS VOC measurements is species-dependent, and the model comparisons prompted
1055 several adjustments to the MEGAN model-calculated emissions. Most importantly, the biogenic emissions of methanol and monoterpenes were downscaled by factors of about 2 and 5, respectively, and the light-dependent fraction for monoterpenes was increased to 0.9 (see Sect. 2.3.2), in order to provide a fair match with the measurements. In addition, the chemical mechanism of the model was updated to better account for OH recycling in isoprene oxidation and for MEK degradation mechanisms. The model performs quite well for isoprene and for its oxidation products (Iox), but their diurnal shape displays
1060 unrealistic peaks in the early morning and late afternoon (a feature also seen for other compounds), likely due to an underestimation of vertical mixing in the model. The ratio of Iox to isoprene (ca. 0.8 around noon in January) is fairly well reproduced. This is reassuring regarding the model performance, since the ratio is strongly dependent on oxidative conditions, i.e. on OH levels.

Despite a good reproduction of the seasonal variation and diurnal shape of the formaldehyde concentrations observed by PTR-
1065 MS, the model underestimates the measurements, by up to 25% during the day and a factor of 2 during the night. This stands in contrast with the model overestimation of HCHO column measurements by FTIR (also at Maïdo) and TROPOMI, for reasons unclear. The discrepancy against the column measurements could be due to an overestimation of the background HCHO production, primarily due to methane oxidation, which would result from overestimated OH levels in the free troposphere. The underestimation against PTR-MS could possibly indicate direct HCHO emissions; however, the nighttime

1070 measurements at Maïdo are primarily influenced by the free troposphere. More work will be needed to elucidate the factors influencing formaldehyde concentrations around Maïdo.

The model fails to reproduce both the seasonal cycle of acetaldehyde (higher values observed in summer compared to winter) and its diurnal variation (strong midday peak). The anthropogenic source of CH_3CHO in the model is almost certainly too high, due to the lumping of higher aldehydes into this compound. Lower anthropogenic emissions and a much higher
1075 photochemical production would be needed to match the observations. The modelled concentration of $\text{C}_{>3}$ alkanes is very likely underestimated, as shown by the model evaluation against measurements in similar environments. Therefore, as proposed in previous studies, a large part of the missing production of acetaldehyde may arise from the oxidation of alkanes, alkenes and alcohols, presumably released by the oceans.

Methyl ethyl ketone (MEK) being the largest contribution to the 73 m/z signal from the PTR-MS, the large observed daytime
1080 mixing ratio suggests the presence of a significant biogenic source of MEK, well in line with the proposal that the deposition of isoprene oxidation products (MVK and 1,2-ISOPROOH) on vegetation is followed by their partial conversion and re-emission as MEK. The ratio between MEK and isoprene emission which achieves the best agreement with the data is 3% (mass basis), higher than the value of this ratio derived by Canaval et al. (2020) based on more direct emission measurements, for different environments. The difference with the latter study might be due to natural variability and/or to model uncertainties related e.g.
1085 to other contributions to the observed signal.

Clearly, more work will be needed to understand and quantify the budget of key compounds (O_3 , OH, NO_x , VOCs and OVOCs) around Réunion Island and similar environments. From the modelling perspective, efforts should be made to improve the representation of NMVOC speciation, with more explicit chemical mechanisms (e.g. for monoterpenes and aromatics) and more detailed emission inventories (e.g. for higher aldehydes). The advection of chemical compounds should be improved,
1090 e.g. with finer resolution modelling to better represent transport in a mountainous environment, and the vertical mixing of the WRF-Chem model (especially during nighttime) should be addressed in future model studies. The ocean/atmosphere exchange of important VOC should be implemented in the model, as it is a well-recognized source of several OVOCs and their precursors (alkanes, etc.). The ocean is also a large source of halogens, which have far-reaching impacts on ozone, OH, NO_x and VOC levels, and their emissions and chemistry should be implemented in WRF-Chem. Efforts should also be made to improve the
1095 representation of boundary conditions of the regional atmospheric models. FTIR column data were used in this work to adjust the boundary conditions for several compounds; other species should be adjusted, based on current knowledge relying on campaign data and network in situ measurements.

Code availability.

The WPS and WRF-Chem model code is provided by the National Center for Atmospheric Research (NCAR)
1100 (<https://doi.org/10.5065/D6MK6B4K>, NCAR, 2020). The WRF-Chem preprocessing tools can be found at

<https://www2.acom.ucar.edu/wrf-chem/wrf-chem-tools-community> (last access: 1 October 2024, NCAR, 2023). Python scripts used in this work can be provided upon request.

Data availability.

Geographical static data used in the preprocessing step of the WRF-Chem simulations was downloaded from the WRF Users Page, hosted by the University Corporation for Atmospheric Research (UCAR), which can be found at https://www2.mmm.ucar.edu/wrf/users/download/get_sources_wps_geog.html (last access: 22 October 2024, UCAR, 2020). CAM-Chem files are distributed by NCAR and available at <https://www.acom.ucar.edu/cam-chem/cam-chem.shtml> (last access: 22 October 2024; <https://doi.org/10.5065/NMP7-EP60>; Buchholz et al., 2019). CAMS global reanalyses, provided by the Copernicus Atmosphere Monitoring Service, were taken from <https://ads.atmosphere.copernicus.eu/cdsapp#!/dataset/cams-global-reanalysis-eac4?tab=form> (last access: 22 October 2024; Inness et al., 2019). Global emissions from the Emission Database for Global Atmospheric Research (EDGAR) v6.1, published by the Joint Research Center (JRC) at the European Commission, are available at <https://data.jrc.ec.europa.eu/dataset/df521e05-6a3b-461c-965a-b703fb62313e> (last access: 22 October 2024; Monforti Ferrario et al., 2022). Speciated NMVOC emissions were obtained at https://data.jrc.ec.europa.eu/dataset/jrc-edgar-edgar_v432_voc_spec_timeseries (last access: 22 October 2024; Janssens-Maenhout et al., 2017). Temporal profiles for the EDGAR inventories have been detailed by Crippa et al. (2020), provided by the JRC and accessed at https://edgar.jrc.ec.europa.eu/dataset_temp_profile (last access: 22 October 2024; Crippa et al., 2020). The population density map for the region of Réunion Island is publicly available at <https://public.opendatasoft.com/explore/dataset/population-francaise-par-departement-2018/table/?disjunctive.departement> (last access: 22 October 2024; French National Institute of Statistics and Economic Studies (INSEE), 2018). Air quality measurements of surface NO₂, NO and O₃ were obtained from the Atmo-Réunion website at <https://atmo-reunion.net/> (last access: 22 October 2024). The PTR-MS measurements are available at <https://data.aeronomie.be/dataset/long-term-in-situ-o-voc-measurements-at-the-maïdo-observatory-reunion-island-v2> (last access: 22 October 2024; Amelynck et al., 2024). FTIR observations at the Maïdo Observatory can be accessed at <https://www-air.larc.nasa.gov/missions/ndacc/data.html?station=la.reunion.maïdo/hdf/ftir/> (last access: 22 October 2024), except for PAN (which can be provided on request). TROPOMI S5P V3.2 data, as well as TM5 profiles, can be accessed at the ESA's public dataspace <https://browser.dataspace.copernicus.eu/> (last access: 22 October 2024; European Space Agency, 2023).

Author contributions.

CP was responsible for running the WRF-Chem simulations, compiling the necessary data, conducting the model validation, and writing the initial version of the manuscript. JFM and TS developed the project concept, oversaw the work, and contributed

to interpreting the findings on a regular basis. CA, BWDV and NS provided the PTR-MS measurements, refined the relevant methodology text and helped with clarifications regarding comparisons with this data. CV and NK provided the FTIR data. CV wrote the text describing the FTIR methodology. JB supplied the 100m resolution dataset of plant functional types and isoprene emission factors. CMV and PT produced the 1 km resolution anthropogenic emission inventory. JFM handled the revision and editing of the manuscript. CP implemented co-author comments and edits and formatted the final version of the manuscript.

Competing interests.

The authors declare that they have no conflict of interest.

Acknowledgements.

We thank Guillaume Peris and Dr. Marion Haramboure from Atmo-Réunion for helping us access ancillary air quality data from the Atmo-Réunion website.

Financial support.

This research received funding from the Belgian Federal Science Policy Office (Belspo) through the European-Space-Agency-funded ProDex TROVA-E2 (2020-2023) and TROVA-3 (2024-2025) projects, as well as via the OCTAVE project (grant no. BR/175/A2/OCTAVE) also funded by Belspo. This work was also supported by the EU Horizon 2020 programme (grant no. ACTRIS-2, 654109).

References

- Amelynck, C., Schoon, N., and Verreyken, B.: Long-term in situ (O)VOC measurements at the Maïdo Observatory (Reunion Island) v2, Version 2 [Data set], Royal Belgian Institute for Space Aeronomy, <https://doi.org/10.18758/7B97H9G4>, 2024.
- Archibald, A. T., Petit, A. S., Percival, C. J., Harvey, J. N. and Shallcross, D. E.: On the importance of the reaction between OH and RO₂ radicals, *Atmospheric Science Letters*, 10(2), 102–108, doi:10.1002/asl.216, 2009.
- Atkinson, R.: Atmospheric Chemistry of VOCs and NO_x, *Atmospheric Environment*, 34(12–14), 2063–2101, doi:10.1016/s1352-2310(99)00460-4, 2000.

Baasandorj, M., Millet, D. B., Hu, L., Mitroo, D., and Williams, B. J.: Measuring acetic and formic acid by proton-transfer-reaction mass spectrometry: sensitivity, humidity dependence, and quantifying interferences, *Atmos. Meas. Tech.*, 8, 1303–1321, <https://doi.org/10.5194/amt-8-1303-2015>, 2015.

1160

Badia, A., Reeves, C. E., Baker, A. R., Saiz-Lopez, A., Volkamer, R., Koenig, T. K., Apel, E. C., Hronbrook, R. S., Carpenter, L. J., Andrews, S. J., Sherwen, R., and Glasow, R.: Importance of reactive halogens in the tropical marine atmosphere: a regional modelling study using WRF-Chem, *Atmos. Chem. Phys.*, 19, 3161–3189, <https://doi.org/10.5194/acp-19-3161-2019>, 2019.

1165

Baray, J.-L., Courcoux, Y., Keckhut, P., Portafaix, T., Tulet, P., Cammas, J.-P., Hauchecorne, A., Godin Beekmann, S., De Mazière, M., Hermans, C., Desmet, F., Sellegri, K., Colomb, A., Ramonet, M., Sciare, J., Vuillemin, C., Hoareau, C., Dionisi, D., Dufлот, V., Vèrèmes, H., Porteneuve, J., Gabarrot, F., Gaudo, T., Metzger, J.-M., Payen, G., Leclair de Bellevue, J., Barthe, C., Posny, F., Ricaud, P., Abchiche, A., and Delmas, R.: Maïdo observatory: a new high-altitude station facility at Reunion Island (21° S, 55° E) for long-term atmospheric remote sensing and in situ measurements, *Atmos. Meas. Tech.*, 6, 2865–2877, <https://doi.org/10.5194/amt-6-2865-2013>, 2013.

1170

Barten, J. G. M., Ganzeveld, L. N., Visser, A. J., Jiménez, R., and Krol, M. C.: Evaluation of nitrogen oxides (NO_x) sources and sinks and ozone production in Colombia and surrounding areas, *Atmos. Chem. Phys.*, 20, 9441–9458, <https://doi.org/10.5194/acp-20-9441-2020>, 2020.

1175

Bates, K. H., Jacob, D. J., Wang, S., Hornbrook, R. S., Apel, E. C., Kim, M. J., Millet, D. B., Wells, K. C., Chen, X., Brewer, J. F., Ray, E. A., Commane, R., Diskin, G. S. and Wofsy, S. C.: The Global Budget of atmospheric methanol: New constraints on secondary, Oceanic, and terrestrial sources, *Journal of Geophysical Research: Atmospheres*, 126(4), doi:10.1029/2020jd033439, 2021.

1180

Berasategui, M., Amedro, D., Vereecken, L., Lelieveld, J., and Crowley, J. N.: Reaction between CH₃C(O)OOH (peracetic acid) and OH in the gas phase: a combined experimental and theoretical study of the kinetics and mechanism, *Atmos. Chem. Phys.*, 20, 13541–13555, <https://doi.org/10.5194/acp-20-13541-2020>, 2020.

1185

Berg, L. K., Shrivastava, M., Easter, R. C., Fast, J. D., Chapman, E. G., Liu, Y., and Ferrare, R. A.: A new WRF-Chem treatment for studying regional-scale impacts of cloud processes on aerosol and trace gases in parameterized cumuli, *Geosci. Model Dev.*, 8, 409–429, <https://doi.org/10.5194/gmd-8-409-2015>, 2015.

- 1190 Brewer, J.F., Fischer, E. V., Commane, R., Wofsy, S. C., Daube, B. C., Apel, E. C., Hills, A. J., Hornbrook, R. S., Barletta, B., Meinardi, S., Blake, D. R., Ray, E. A., and Ravishankara, A. R.: Evidence for an oceanic source of methyl ethyl ketone to the atmosphere, *Geophys. Res. Lett.*, 47, e2019GL086045, <https://doi.org/10.1029/2019GL086045>, 2020.
- Boersma, K. F., Vinken, G. C. M., and Eskes, H. J.: Representativeness errors in comparing chemistry transport and chemistry
1195 climate models with satellite UV–Vis tropospheric column retrievals, *Geosci. Model Dev.*, 9, 875–898, <https://doi.org/10.5194/gmd-9-875-2016>, 2016.
- Bon, D. M., Ulbrich, I. M., de Gouw, J. A., Warneke, C., Kuster, W. C., Alexander, M. L., Baker, A., Beyersdorf, A. J., Blake, D., Fall, R., Jimenez, J. L., Herndon, S. C., Huey, L. G., Knighton, W. B., Ortega, J., Springston, S., and Vargas, O.:
1200 Measurements of volatile organic compounds at a suburban ground site (T1) in Mexico City during the MILAGRO 2006 campaign: measurement comparison, emission ratios, and source attribution, *Atmos. Chem. Phys.*, 11, 2399–2421, <https://doi.org/10.5194/acp-11-2399-2011>, 2011.
- Bossolasco, A., Faragó, E. P., Schoemaeker, C. and Fittschen, C.: Rate constant of the reaction between CH_3O_2 and OH
1205 radicals, *Chemical Physics Letters*, 593, 7–13, doi:10.1016/j.cplett.2013.12.052, 2014.
- Bozem, H., Pozzer, A., Harder, H., Martinez, M., Williams, J., Lelieveld, J., and Fischer, H.: The influence of deep convection on HCHO and H_2O_2 in the upper troposphere over Europe, *Atmos. Chem. Phys.*, 17, 11835–11848, <https://doi.org/10.5194/acp-17-11835-2017>, 2017.
- 1210 Brito, J., Wurm, F., Yáñez-Serrano, A. M., de Assunção, J. V., Godoy, J. M. and Artaxo, P.: Vehicular emission ratios of VOCs in a megacity impacted by extensive ethanol use: Results of ambient measurements in São Paulo, Brazil, *Environmental Science and Technology*, 49(19), 11381–11387, doi:10.1021/acs.est.5b03281, 2015.
- 1215 Buchholz, R. R., Emmons, L. K., Tilmes, S., and The CESM2 Development Team: CESM2.1/CAM-chem Instantaneous Output for Boundary Conditions, UCAR/NCAR – Atmospheric Chemistry Observations and Modeling Laboratory. (e.g. Lat: -30 to -10, Lon: 355 to 70, 10 June 2019–01 July 2019), UCAR [dataset], <https://doi.org/10.5065/NMP7-EP60> (last access: 22 October 2024), 2019.
- 1220 Callewaert, S., Brioude, J., Langerock, B., Dufлот, V., Fonteyn, D., Müller, J.-F., Metzger, J.-M., Hermans, C., Kumps, N., Ramonet, M., Lopez, M., Mahieu, E., and De Mazière, M.: Analysis of CO_2 , CH_4 , and CO surface and column concentrations observed at Réunion Island by assessing WRF-Chem simulations, *Atmos. Chem. Phys.*, 22, 7763–7792, <https://doi.org/10.5194/acp-22-7763-2022>, 2022.

- 1225 Canaval, E., Millet, D. B., Zimmer, I., Nosenko, T., Georgii, E., Partoll, E. M., Fischer, L., Alwe, H. D., Kulmala, M., Karl,
T., Schnitzler, J.-P. and Hansel, A.: Rapid conversion of isoprene photooxidation products in terrestrial plants,
Communications Earth & Environment, 1(1), doi:10.1038/s43247-020-00041-2, 2020.
- Cappellin, L., Loreto, F., Biasioli, F., Pastore, P., and McKinney, K.: A mechanism for biogenic production and emission of
1230 MEK from MVK decoupled from isoprene biosynthesis, Atmos. Chem. Phys., 19, 3125–3135, <https://doi.org/10.5194/acp-19-3125-2019>, 2019.
- Caram, C., Szopa, S., Cozic, A., Bekki, S., Cuevas, C. A., and Saiz-Lopez, A.: Sensitivity of tropospheric ozone to halogen
chemistry in the chemistry–climate model LMDZ-INCA vNMHC, Geosci. Model Dev., 16, 4041–4062,
1235 <https://doi.org/10.5194/gmd-16-4041-2023>, 2023.
- Caravan, R. L., Khan, M. A., Zádor, J., Sheps, L., Antonov, I. O., Rotavera, B., Ramasesha, K., Au, K., Chen, M.-W., Rösch,
D., Osborn, D. L., Fittschen, C., Schoemaeker, C., Duncianu, M., Grira, A., Dusanter, S., Tomas, A., Percival, C. J.,
Shallcross, D. E. and Taatjes, C. A.: The reaction of hydroxyl and methylperoxy radicals is not a major source of atmospheric
1240 methanol, Nature Communications, 9(1), doi:10.1038/s41467-018-06716-x, 2018.
- Chen, F., Kusaka, H., Bornstein, R., Ching, J., Grimmond, C. S., Grossman-Clarke, S., Loridan, T., Manning, K. W., Martilli,
A., Miao, S., Sailor, D., Salamanca, F. P., Taha, H., Tewari, M., Wang, X., Wyszogrodzki, A. A. and Zhang, C.: The integrated
WRF/Urban Modelling System: Development, evaluation, and applications to urban environmental problems, International
1245 Journal of Climatology, 31(2), 273–288, doi:10.1002/joc.2158, 2011.
- Chin, M., Ginoux, P., Kinne, S., Torres, O., Holben, B. N., Duncan, B. N., Martin, R. V., Logan, J. A., Higurashi, A. and
Nakajima, T.: Tropospheric aerosol optical thickness from the GOCART model and comparisons with satellite and Sun
Photometer Measurements, J. Atmos. Sci., 59(3), 461–483, [https://doi.org/10.1175/1520-0469\(2002\)059<0461:taotft>2.0.co;2](https://doi.org/10.1175/1520-0469(2002)059<0461:taotft>2.0.co;2), 2002.
- 1250 Comes, F. J.: Recycling in the Earth’s atmosphere: The OH radical—its importance for the chemistry of the atmosphere and
the determination of its concentration, Angewandte Chemie International Edition in English, 33(18), 1816–1826,
doi:10.1002/anie.199418161, 1994.

- Crippa, M., Guizzardi, D., Muntean, M., Schaaf, E., Dentener, F., van Aardenne, J. A., Monni, S., Doering, U., Olivier, J. G. J., Pagliari, V., and Janssens-Maenhout, G.: Gridded emissions of air pollutants for the period 1970–2012 within EDGAR v4.3.2, *Earth Syst. Sci. Data*, 10, 1987–2013, <https://doi.org/10.5194/essd-10-1987-2018>, 2018.
- 1260 Crippa, M., Solazzo, E., Huang, G., Guizzardi, D., Koffi, E., Muntean, M., Schieberle, C., Friedrich, R., and Janssens-Maenhout, G.: High resolution temporal profiles in the Emissions Database for Global Atmospheric Research, *Sci. Data*, 7, 121, <https://doi.org/10.1038/s41597-020-0462-2>, 2020 (data available at: https://edgar.jrc.ec.europa.eu/dataset_temp_profile, last access: 1 January 2023).
- 1265 Crippa, M., Guizzardi, D., Muntean, M., Schaaf, E., Monforti-Ferrario, F., Banja, M., Pagani, F. and Solazzo, E., EDGAR v6.1 global air pollutant emissions, European Commission, JRC129555, 2022.
- Damian, V., Sandu, A., Damian, M., Potra, F. and Carmichael, G. R.: The kinetic preprocessor KPP-a software environment for solving chemical kinetics, *Computers & Chemical Engineering*, 26(11), 1567–1579, doi:10.1016/s0098-1270 1354(02)00128-x, 2002.
- de Foy, B., Krotkov, N. A., Bei, N., Herndon, S. C., Huey, L. G., Martínez, A.-P., Ruiz-Suárez, L. G., Wood, E. C., Zavala, M., and Molina, L. T.: Hit from both sides: tracking industrial and volcanic plumes in Mexico City with surface measurements and OMI SO₂ retrievals during the MILAGRO field campaign, *Atmos. Chem. Phys.*, 9, 9599–9617, 1275 <https://doi.org/10.5194/acp-9-9599-2009>, 2009.
- De Smedt, I., Theys, N., Yu, H., Danckaert, T., Lerot, C., Compennolle, S., Van Roozendaal, M., Richter, A., Hilboll, A., Peters, E., Pedernana, M., Loyola, D., Beirle, S., Wagner, T., Eskes, H., van Geffen, J., Boersma, K. F., and Veefkind, P.: Algorithm theoretical baseline for formaldehyde retrievals from S5P TROPOMI and from the QA4ECV project, *Atmos. Meas. Tech.*, 11, 2395–2426, <https://doi.org/10.5194/amt-11-2395-2018>, 2018. 1280
- De Smedt, I., Pinardi, G., Vigouroux, C., Compennolle, S., Bais, A., Benavent, N., Boersma, F., Chan, K.-L., Donner, S., Eichmann, K.-U., Hedelt, P., Hendrick, F., Irie, H., Kumar, V., Lambert, J.-C., Langerock, B., Lerot, C., Liu, C., Loyola, D., Piders, A., Richter, A., Rivera Cárdenas, C., Romahn, F., Ryan, R. G., Sinha, V., Theys, N., Vlietinck, J., Wagner, T., Wang, 1285 T., Yu, H., and Van Roozendaal, M.: Comparative assessment of TROPOMI and OMI formaldehyde observations and validation against MAX-DOAS network column measurements, *Atmos. Chem. Phys.*, 21, 12561–12593, <https://doi.org/10.5194/acp-21-12561-2021>, 2021.

- De Smedt, I., Theys, N., Yu, H., Vlietinck, J., Lerot, C., and Van Roozendael, M.: S5P/TROPOMI HCHO ATBD, S5P-BIRA-L2-400F-ATBD, 2.2.0, BIRA-IASB Royal Belgian Institute for Space Aeronomy, Brussels, Belgium, <https://sentinels.copernicus.eu/documents/247904/2476257/Sentinel-5P-ATBD-HCHO-TROPOMI> (last access: 23 October 2024), 2022.
- Derstroff, B., Hüser, I., Bourtsoukidis, E., Crowley, J. N., Fischer, H., Gromov, S., Harder, H., Janssen, R. H. H., Kesselmeier, J., Lelieveld, J., Mallik, C., Martinez, M., Novelli, A., Parchatka, U., Phillips, G. J., Sander, R., Sauvage, C., Schuladen, J., Stönnner, C., Tomsche, L., and Williams, J.: Volatile organic compounds (VOCs) in photochemically aged air from the eastern and western Mediterranean, *Atmos. Chem. Phys.*, 17, 9547–9566, <https://doi.org/10.5194/acp-17-9547-2017>, 2017.
- Di Carlo, P., Brune, W. H., Martinez, M., Harder, H., Leshner, R., Ren, X., Thornberry, T., Carroll, M. A., Young, V., Shepson, P. B., Riemer, D., Apel, E. and Campbell, C.: Missing OH reactivity in a forest: Evidence for unknown reactive biogenic VOCs, *Science*, 304(5671), 722–725, doi:10.1126/science.1094392, 2004.
- Dimitropoulou, E., Hendrick, F., Pinardi, G., Friedrich, M. M., Merlaud, A., Tack, F., De Longueville, H., Fayt, C., Hermans, C., Laffineur, Q., Fierens, F., and Van Roozendael, M.: Validation of TROPOMI tropospheric NO₂ columns using dual-scan multi-axis differential optical absorption spectroscopy (MAX-DOAS) measurements in Uccle, Brussels, *Atmos. Meas. Tech.*, 13, 5165–5191, <https://doi.org/10.5194/amt-13-5165-2020>, 2020.
- Dominutti, P. A., Renard, P., Vaïtilingom, M., Bianco, A., Baray, J.-L., Borbon, A., Bourrianne, T., Burnet, F., Colomb, A., Delort, A.-M., Duflot, V., Houdier, S., Jaffrezo, J.-L., Joly, M., Lereboure, M., Metzger, J.-M., Pichon, J.-M., Ribeiro, M., Rocco, M., Tulet, P., Vella, A., Leriche, M., and Deguillaume, L.: Insights into tropical cloud chemistry in Réunion (Indian Ocean): results from the BIO-MAÏDO campaign, *Atmos. Chem. Phys.*, 22, 505–533, <https://doi.org/10.5194/acp-22-505-2022>, 2022.
- Douros, J., Eskes, H., van Geffen, J., Boersma, K. F., Compennolle, S., Pinardi, G., Blechschmidt, A.-M., Peuch, V.-H., Colette, A., and Veeffkind, P.: Comparing Sentinel-5P TROPOMI NO₂ column observations with the CAMS regional air quality ensemble, *Geosci. Model Dev.*, 16, 509–534, <https://doi.org/10.5194/gmd-16-509-2023>, 2023.
- Duflot, V., Tulet, P., Flores, O., Barthe, C., Colomb, A., Deguillaume, L., Vaïtilingom, M., Perring, A., Huffman, A., Hernandez, M. T., Sellegri, K., Robinson, E., O'Connor, D. J., Gomez, O. M., Burnet, F., Bourrianne, T., Strasberg, D., Rocco, M., Bertram, A. K., Chazette, P., Totems, J., Fournel, J., Stamenoff, P., Metzger, J.-M., Chabasset, M., Rousseau, C., Bourrianne, E., Sancelme, M., Delort, A.-M., Wegener, R. E., Chou, C., and Elizondo, P.: Preliminary results from the FARCE

- 2015 campaign: multidisciplinary study of the forest–gas–aerosol–cloud system on the tropical island of La Réunion, *Atmos. Chem. Phys.*, 19, 10591–10618, <https://doi.org/10.5194/acp-19-10591-2019>, 2019.
- 1325 Dufлот, V., Bègue, N., Pouliquen, M.-L., Goloub, P. and Metzger, J.-M.: Aerosols on the tropical island of La Réunion (21°S, 55°E): Assessment of climatology, origin of variability and trend, *Remote Sensing*, 14(19), 4945, doi:10.3390/rs14194945, 2022.
- 1330 El Gdachi, S., Tulet, P., Réchou, A., Burnet, F., Mouchel-Vallon, C., Jambert, C. and Leriche, M.: Thermodynamic processes driving thermal circulations on slopes: Modeling anabatic and katabatic flows on Reunion Island, *Journal of Geophysical Research: Atmospheres*, 129(17), doi:10.1029/2023jd040431, 2024.
- 1335 Emmons, L. K., Walters, S., Hess, P. G., Lamarque, J.-F., Pfister, G. G., Fillmore, D., Granier, C., Guenther, A., Kinnison, D., Laepple, T., Orlando, J., Tie, X., Tyndall, G., Wiedinmyer, C., Baughcum, S. L., and Kloster, S.: Description and evaluation of the Model for Ozone and Related chemical Tracers, version 4 (MOZART-4), *Geosci. Model Dev.*, 3, 43–67, <https://doi.org/10.5194/gmd-3-43-2010>, 2010.
- 1340 Emmons, L. K., Schwantes, R. H., Orlando, J. J., Tyndall, G., Kinnison, D., Lamarque, J., Marsh, D., Mills, M. J., Tilmes, S., Bardeen, C., Buchholz, R. R., Conley, A., Gettelman, A., Garcia, R., Simpson, I., Blake, D. R., Meinardi, S. and Pétron, G.: The chemistry mechanism in the Community Earth System Model Version 2 (CESM2), *Journal of Advances in Modeling Earth Systems*, 12(4), doi:10.1029/2019ms001882, 2020.
- 1345 Eskes, H. J. and Boersma, K. F.: Averaging kernels for DOAS total-column satellite retrievals, *Atmos. Chem. Phys.*, 3, 1285–1291, <https://doi.org/10.5194/acp-3-1285-2003>, 2003.
- 1350 Fisher, E. V., Jacob, D. J., Millet, D. B., Yantosca, R. M., and Mao, J.: The role of the ocean in the global atmospheric budget of acetone, *Geophys. Res. Lett.*, 39, L01807, <https://doi.org/10.1029/2011GL050086>, 2012.
- 1350 Fortner, E. C., Zheng, J., Zhang, R., Berk Knighton, W., Volkamer, R. M., Sheehy, P., Molina, L., and André, M.: Measurements of Volatile Organic Compounds Using Proton Transfer Reaction – Mass Spectrometry during the MILAGRO 2006 Campaign, *Atmos. Chem. Phys.*, 9, 467–481, <https://doi.org/10.5194/acp-9-467-2009>, 2009.

- 1355 Foucart, B., Sellegri, K., Tulet, P., Rose, C., Metzger, J.-M., and Picard, D.: High occurrence of new particle formation events at the Maïdo high-altitude observatory (2150 m), Réunion (Indian Ocean), *Atmos. Chem. Phys.*, 18, 9243–9261, <https://doi.org/10.5194/acp-18-9243-2018>, 2018.
- 1360 Franco, B., Bader, W., Toon, G. C., Bray, C., Perrin, A., Fischer, E. V., Sudo, K., Boone, C. D., Bovy, B., Lejeune, B., Servais, C. and Mahieu, E.: Retrieval of ethane from ground-based FTIR solar spectra using improved spectroscopy: Recent burden increase above jungfrauoch, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 160, 36–49, doi:10.1016/j.jqsrt.2015.03.017, 2015.
- 1365 Franco, B., Clarisse, L., Stavrakou, T., Müller, J.-F., Van Damme, M., Whitburn, S., Hadji-Lazaro, J., Hurtmans, D., Taraborrelli, D., Clerbaux, C. and Coheur, P. -F: A general framework for global retrievals of trace gases fromiasi: Application to methanol, formic acid, and Pan, *Journal of Geophysical Research: Atmospheres*, 123(24), doi:10.1029/2018jd029633, 2018.
- 1370 Fried, A., Olson, J. R., Walega, J. G., Crawford, J. H., Chen, G., Weibring, P., Richter, D., Roller, C., Tittel, F., Porter, M., Fuelberg, H., Halland, J., Bertram, T. H., Cohen, R. C., Pickering, K., Heikes, B. G., Snow, J. A., Shen, H., O’Sullivan, D. W., Brune, W. H., Ren, X., Blake, D. R., Blake, N., Sachse, G., Diskin, G. S., Podolske, J., Vay, S. A., Shetter, R. E., Hall, S. R., Anderson, B. E., Thornhill, L., Clarke, A. D., McNaughton, C. S., Singh, H. B., Avery, M. A., Huey, G., Kim, S. and Millet, D. B.: Role of convection in redistributing formaldehyde to the upper troposphere over North America and the North Atlantic during the Summer 2004 INTEx campaign, *Journal of Geophysical Research: Atmospheres*, 113(D17), doi:10.1029/2007jd009760, 2008.
- 1375 Friedl, M. A., McIver, D. K., Hodges, J. C. F., Zhang, X. Y., Muchoney, D., Strahler, A. H., Woodcock, C. E., Gopal, S., Schneider, A., Cooper, A., Baccini, A., Gao, F. and Schaaf, C.: Global land cover mapping from MODIS: Algorithms and early results, *Remote Sensing of Environment*, 83(1–2), 287–302, doi:10.1016/s0034-4257(02)00078-0, 2002.
- 1380 Goliff, W. S., Stockwell, W. R. and Lawson, C. V.: The regional atmospheric chemistry mechanism, version 2, *Atmospheric Environment*, 68, 174–185, doi:10.1016/j.atmosenv.2012.11.038, 2013.
- Grell, G. A.: Prognostic evaluation of assumptions used by Cumulus Parameterizations, *Monthly Weather Review*, 121(3), 764–787, [https://doi.org/10.1175/1520-0493\(1993\)121<0764:PEOAUB>2.0.CO;2](https://doi.org/10.1175/1520-0493(1993)121<0764:PEOAUB>2.0.CO;2), 1993.
- 1385 Grell, G. A. and Dévényi, D.: A generalized approach to parameterizing convection combining ensemble and data assimilation techniques, *Geophysical Research Letters*, 29(14), doi:10.1029/2002gl015311, 2002.

- Grell, G. A., Peckham, S. E., Schmitz, R., McKeen, S. A., Frost, G., Skamarock, W. C. and Eder, B.: Fully coupled “online” chemistry within the WRF model, *Atmospheric Environment*, 39(37), 6957–6975, doi:10.1016/j.atmosenv.2005.04.027, 2005.
- 1390 Gross, C. B. M., Dillon, T. J., Schuster, G., Lelieveld, J., and Crowley, J. N.: Direct kinetic study of OH and O₃ formation in the reaction of CH₃C(O)O₂ with HO₂, *J. Phys. Chem. A*, 118, 974-985, <https://doi.org/10.1021/jp412380z>, 2014.
- Guenther, A., Karl, T., Harley, P., Wiedinmyer, C., Palmer, P. I., and Geron, C.: Estimates of global terrestrial isoprene emissions using MEGAN (Model of Emissions of Gases and Aerosols from Nature), *Atmos. Chem. Phys.*, 6, 3181–3210, 1395 <https://doi.org/10.5194/acp-6-3181-2006>, 2006.
- Guenther, A. B., Jiang, X., Heald, C. L., Sakulyanontvittaya, T., Duhl, T., Emmons, L. K., and Wang, X.: The Model of Emissions of Gases and Aerosols from Nature version 2.1 (MEGAN2.1): an extended and updated framework for modeling biogenic emissions, *Geosci. Model Dev.*, 5, 1471–1492, <https://doi.org/10.5194/gmd-5-1471-2012>, 2012.
- 1400 Helmig, D., Hueber, J., and Tans, P.: University of Colorado Institute of Arctic and Alpine Research (INSTARR), NOAA GML CCGG Group, University of Colorado Institute of Arctic and Alpine Research (INSTARR) Flask-Air Sample Measurements of Atmospheric Non Methane Hydrocarbons Mole Fractions from the NOAA GML Carbon Cycle Surface Network at Global and Regional Background Sites, 2004-2016 (Version 2021.05.04), NOAA Global Monitoring Laboratory 1405 [data set], <https://doi.org/10.15138/6AV8-GS57>, 2021.
- Hewitt, C. N., Hayward, S. and Tani, A.: The application of proton transfer reaction-mass spectrometry (PTR-MS) to the monitoring and analysis of volatile organic compounds in the atmosphere, *Journal of Environmental Monitoring*, 5(1), 1–7, doi:10.1039/b204712h, 2002.
- 1410 Horner, R. P., Marais, E. A., Wei, N., Ryan, R. G., and Shah, V.: Vertical profiles of global tropospheric nitrogen dioxide (NO₂) obtained by cloud-slicing TROPOMI, *EGUsphere* [preprint], <https://doi.org/10.5194/egusphere-2024-1541>, 2024.
- Huang, G., Brook, R., Crippa, M., Janssens-Maenhout, G., Schieberle, C., Dore, C., Guizzardi, D., Muntean, M., Schaaf, E., 1415 and Friedrich, R.: Speciation of anthropogenic emissions of non-methane volatile organic compounds: a global gridded data set for 1970–2012, *Atmos. Chem. Phys.*, 17, 7683–7701, <https://doi.org/10.5194/acp-17-7683-2017>, 2017.
- Hulley, G., Malakar, N., and Freepartner, R.: Moderate Resolution Imaging Spectroradiometer (MODIS) Land Surface Temperature and Emissivity Product (MxD21) Algorithm Theoretical Basis Document Collection-6, 1420 https://modis.gsfc.nasa.gov/data/atbd/atbd_mod21.pdf (last access: 6 February 2024), 2016.

- Iacono, M. J., Delamere, J. S., Mlawer, E. J., Shephard, M. W., Clough, S. A. and Collins, W. D.: Radiative forcing by long-lived greenhouse gases: Calculations with the AER radiative transfer models, *Journal of Geophysical Research: Atmospheres*, 113(D13), doi:10.1029/2008jd009944, 2008.
- 1425 Iglesias-Suarez, F., Badia, A., Fernandez, R. P., Cuevas, C. A., Kinnison, D. E., Tilmes, S., Lamarque, J.-F., Long, M. C., Hossaini, R. and Saiz-Lopez, A.: Natural halogens buffer tropospheric ozone in a changing climate, *Nature Climate Change*, 10(2), 147–154, doi:10.1038/s41558-019-0675-6, 2020.
- 1430 Inness, A., Ades, M., Agustí-Panareda, A., Barré, J., Benedictow, A., Blechschmidt, A.-M., Dominguez, J. J., Engelen, R., Eskes, H., Flemming, J., Huijnen, V., Jones, L., Kipling, Z., Massart, S., Parrington, M., Peuch, V.-H., Razinger, M., Remy, S., Schulz, M., and Suttie, M.: The CAMS reanalysis of atmospheric composition, *Atmos. Chem. Phys.*, 19, 3515–3556, <https://doi.org/10.5194/acp-19-3515-2019>, 2019.
- 1435 Jacob, D. J., Field, B. D., Jin, E. M., Bey, I., Li, Q., Logan, J. A., Yantosca, R. M. and Singh, H. B.: Atmospheric budget of acetone, *Journal of Geophysical Research: Atmospheres*, 107(D10), doi:10.1029/2001jd000694, 2002.
- Jacob, D. J., Field, B. D., Li, Q., Blake, D. R., de Gouw, J., Warneke, C., Hansel, A., Wisthaler, A., Singh, H. B. and Guenther, A.: Global budget of methanol: Constraints from atmospheric observations, *Journal of Geophysical Research: Atmospheres*, 110(D8), doi:10.1029/2004jd005172, 2005.
- 1440 Janssens-Maenhout, G., Crippa, M., Guizzardi, D., Muntean, M., and Schaaf, E.: JRC-EDGARv432_VOC_spec_timeseries, European Commission, Joint Research Centre (JRC), Dataset, https://data.europa.eu/89h/jrc-edgar-edgar_v432_voc_spec_timeseries, 2017.
- 1445 Jenkin, M. E., Saunders, S. M. and Pilling, M. J.: The tropospheric degradation of volatile organic compounds: A protocol for mechanism development, *Atmospheric Environment*, 31(1), 81–104, doi:10.1016/s1352-2310(96)00105-7, 1997.
- Jenkin, M. E., Hurley, M. D., and Wallington, T. J.: Investigation of the radical product channel of the $\text{CH}_3\text{C}(\text{O})\text{O}_2 + \text{HO}_2$ reaction in the gas phase, *Phys. Chem. Chem. Phys.*, 9, 3149–3162, <https://doi.org/10.1039/870275E>, 2007.
- 1450 Jerrett, M., Burnett, R. T., Pope, C. A., Ito, K., Thurston, G., Krewski, D., Shi, Y., Calle, E. and Thun, M.: Long-term ozone exposure and mortality, *New England Journal of Medicine*, 360(11), 1085–1095, doi:10.1056/nejmoa0803894, 2009.

- 1455 Jiménez, P. A., Dudhia, J., González-Rouco, J. F., Navarro, J., Montávez, J. P. and García-Bustamante, E.: A revised scheme for the WRF surface layer formulation, *Monthly Weather Review*, 140(3), 898–918, doi:10.1175/mwr-d-11-00056.1, 2012.
- Kanakidou, M., Seinfeld, J. H., Pandis, S. N., Barnes, I., Dentener, F. J., Facchini, M. C., Van Dingenen, R., Ervens, B., Nenes, A., Nielsen, C. J., Swietlicki, E., Putaud, J. P., Balkanski, Y., Fuzzi, S., Horth, J., Moortgat, G. K., Winterhalter, R., Myhre, C. E. L., Tsigaridis, K., Vignati, E., Stephanou, E. G., and Wilson, J.: Organic aerosol and global climate modelling: a review, *Atmos. Chem. Phys.*, 5, 1053–1123, <https://doi.org/10.5194/acp-5-1053-2005>, 2005.
- 1460 Katragadda, H. R., Fullana, A., Sidhu, S., and Carbonell-Barrachina, A. A.: Emissions of volatile aldehydes from heated cooking oils, *Food Chem.*, 120, 59-65, <https://doi.org/10.1016/j.foodchem.2009/09/070>, 2010.
- 1465 Keene, W. C. and Galloway, J. N.: A note on acid rain in an amazon rainforest, *Tellus B: Chemical and Physical Meteorology*, 36(2), 137, doi:10.3402/tellusb.v36i2.14883, 1984.
- Khan, M. A., Lyons, K., Chhantyal-Pun, R., McGillen, M. R., Caravan, R. L., Taatjes, C. A., Orr-Erwing, A. J., Percival, C. J., and Shallcross, D. E.: Investigating the tropospheric chemistry of acetic acid using the global 3-D chemistry transport model, *STOCHEM-CRI, J. Geophys. Res.*, 123, 6267-6281, <https://doi.org/10.1029/2018JD028529>, 2018.
- 1470 Koss, A. R., Sekimoto, K., Gilman, J. B., Selimovic, V., Coggon, M. M., Zarzana, K. J., Yuan, B., Lerner, B. M., Brown, S. S., Jimenez, J. L., Krechmer, J., Roberts, J. M., Warneke, C., Yokelson, R. J., and de Gouw, J.: Non-methane organic gas emissions from biomass burning: identification, quantification, and emission factors from PTR-ToF during the FIREX 2016 laboratory experiment, *Atmos. Chem. Phys.*, 21, 3299-3319, <https://doi.org/10.5194/acp-18-3299-2018>, 2018.
- 1475 Lamarque, J.-F., Emmons, L. K., Hess, P. G., Kinnison, D. E., Tilmes, S., Vitt, F., Heald, C. L., Holland, E. A., Lauritzen, P. H., Neu, J., Orlando, J. J., Rasch, P. J., and Tyndall, G. K.: CAM-chem: description and evaluation of interactive atmospheric chemistry in the Community Earth System Model, *Geosci. Model Dev.*, 5, 369–411, <https://doi.org/10.5194/gmd-5-369-2012>, 2012.
- 1480 Lamsal, L. N., Martin, R. V., van Donkelaar, A., Steinbacher, M., Celarier, E. A., Bucsela, E., Dunlea, E. J. and Pinto, J. P.: Ground-level nitrogen dioxide concentrations inferred from the satellite-borne Ozone Monitoring Instrument, *Journal of Geophysical Research: Atmospheres*, 113(D16), doi:10.1029/2007jd009235, 2008.
- 1485 Lange, K., Richter, A., Schönhardt, A., Meier, A. C., Bösch, T., Seyler, A., Krause, K., Behrens, L. K., Wittrock, F., Merlaud, A., Tack, F., Fayt, C., Friedrich, M. M., Dimitropoulou, E., Van Roozendaal, M., Kumar, V., Donner, S., Dörner, S., Lauster,

- B., Razi, M., Borger, C., Uhlmannsiek, K., Wagner, T., Ruhtz, T., Eskes, H., Bohn, B., Santana Diaz, D., Abuhassan, N.,
 1490 Schüttemeyer, D., and Burrows, J. P.: Validation of Sentinel-5P TROPOMI tropospheric NO₂ products by comparison with
 NO₂ measurements from airborne imaging DOAS, ground-based stationary DOAS, and mobile car DOAS measurements
 during the S5P-VAL-DE-Ruhr campaign, *Atmos. Meas. Tech.*, 16, 1357–1389, <https://doi.org/10.5194/amt-16-1357-2023>,
 2023.
- 1495 Langford, B., Misztal, P. K., Nemitz, E., Davison, B., Helfter, C., Pugh, T. A. M., MacKenzie, A. R., Lim, S. F., and Hewitt,
 C. N.: Fluxes and concentrations of volatile organic compounds from a South-East Asian tropical rainforest, *Atmos. Chem.
 Phys.*, 10, 8391–8412, <https://doi.org/10.5194/acp-10-8391-2010>, 2010.
- Liu, Z., Nguyen, V. S., Harvey, J., Müller, J.-F. and Peeters, J.: The photolysis of α -hydroperoxycarbonyls, *Physical Chemistry
 Chemical Physics*, 20(10), 6970–6979, doi:10.1039/c7cp08421h, 2018.
- 1500
- Logan, J. A.: Tropospheric ozone: Seasonal behavior, trends, and anthropogenic influence, *Journal of Geophysical Research:
 Atmospheres*, 90(D6), 10463–10482, doi:10.1029/jd090id06p10463, 1985.
- Lorente, A., Folkert Boersma, K., Yu, H., Dörner, S., Hilboll, A., Richter, A., Liu, M., Lamsal, L. N., Barkley, M., De Smedt,
 1505 I., Van Roozendaal, M., Wang, Y., Wagner, T., Beirle, S., Lin, J.-T., Krotkov, N., Stammes, P., Wang, P., Eskes, H. J., and
 Krol, M.: Structural uncertainty in air mass factor calculation for NO₂ and HCHO satellite retrievals, *Atmos. Meas. Tech.*, 10,
 759–782, <https://doi.org/10.5194/amt-10-759-2017>, 2017.
- Macor, A. and Pavanello, P.: Performance and emissions of biodiesel in a boiler for residential heating, *Energy*, 34(12), 2025–
 1510 2032, doi:10.1016/j.energy.2008.08.021, 2009.
- Mahieu, E., Fischer, E. V., Franco, B., Palm, M., Wizenberg, T., Smale, D., Clarisse, L., Clerbaux, C., Coheur, P.-F., Hannigan,
 J. W., Lutsch, E., Notholt, J., Cantos, I. P., Prignon, M., Servais, C. and Strong, K.: First retrievals of peroxyacetyl nitrate
 (PAN) from ground-based FTIR solar spectra recorded at remote sites, comparison with model and satellite data, *Elementa:
 1515 Science of the Anthropocene*, 9(1), doi:10.1525/elementa.2021.00027, 2021.
- Marais, E. A., Roberts, J. F., Ryan, R. G., Eskes, H., Boersma, K. F., Choi, S., Joiner, J., Abuhassan, N., Redondas, A., Grutter,
 M., Cede, A., Gomez, L., and Navarro-Comas, M.: New observations of NO₂ in the upper troposphere from TROPOMI,
Atmos. Meas. Tech., 14, 2389–2408, <https://doi.org/10.5194/amt-14-2389-2021>, 2021.
- 1520

- Mickley, L. J., Jacob, D. J., Field, B. D. and Rind, D.: Climate response to the increase in tropospheric ozone since preindustrial times: A comparison between ozone and equivalent CO₂ forcings, *Journal of Geophysical Research: Atmospheres*, 109(D5), doi:10.1029/2003jd003653, 2004.
- 1525 Millet, D. B., Guenther, A., Siegel, D. A., Nelson, N. B., Singh, H. B., de Gouw, J. A., Warneke, C., Williams, J., Eerdekens, G., Sinha, V., Karl, T., Flocke, F., Apel, E., Riemer, D. D., Palmer, P. I., and Barkley, M.: Global atmospheric budget of acetaldehyde: 3-D model analysis and constraints from in-situ and satellite observations, *Atmos. Chem. Phys.*, 10, 3405–3425, <https://doi.org/10.5194/acp-10-3405-2010>, 2010.
- 1530 Millet, D. B., Baasandorj, M., Farmer, D. K., Thornton, J. A., Baumann, K., Brophy, P., Chaliyakunnel, S., de Gouw, J. A., Graus, M., Hu, L., Koss, A., Lee, B. H., Lopez-Hilfiker, F. D., Neuman, J. A., Paulot, F., Peischl, J., Pollack, I. B., Ryerson, T. B., Warneke, C., Williams, B. J., and Xu, J.: A large and ubiquitous source of atmospheric formic acid, *Atmos. Chem. Phys.*, 15, 6283–6304, <https://doi.org/10.5194/acp-15-6283-2015>, 2015.
- 1535 Mitsuishi, K., Iwasaki, M., Takeuchi, M., Okochi, H., Kato, S., Ohira, S.-I. and Toda, K.: Diurnal variations in partitioning of atmospheric glyoxal and methylglyoxal between gas and particles at the ground level and in the free troposphere, *ACS Earth and Space Chemistry*, 2(9), 915–924, doi:10.1021/acsearthspacechem.8b00037, 2018.
- Monforti-Ferrario, F., Crippa, M., Guizzardi, D., Muntean, M., Schaaf, E., Banja, M., Pagani, F., and Solazzo, E.: EDGAR
 1540 v6.1 Global Air Pollutant Emissions, European Commission, Joint Research Centre (JRC), Dataset, <https://data.europa.eu/89h/df521e05-6a3b-461c-965a-b703fb62313e>, 2022.
- Morrison, H., Thompson, G. and Tatarskii, V.: Impact of cloud microphysics on the development of trailing stratiform precipitation in a simulated squall line: Comparison of one- and two-moment schemes, *Monthly Weather Review*, 137(3),
 1545 991–1007, doi:10.1175/2008mwr2556.1, 2009.
- Müller, J.-F., and Brasseur, G. P.: Sources of upper tropospheric HO_x: A three-dimensional study, *J. Geophys. Res.*, 104, 1705, <https://doi.org/10.1029/1998JD10005>, 1999.
- 1550 Müller, J.-F., Liu, Z., Nguyen, V. S., Stavrakou, T., Harvey, J. N. and Peeters, J.: The reaction of methyl peroxy and hydroxyl radicals as a major source of atmospheric methanol, *Nature Communications*, 7(1), doi:10.1038/ncomms13213, 2016.

- Müller, J.-F., Stavrou, T., and Peeters, J.: Chemistry and deposition in the Model of Atmospheric composition at Global and Regional scales using Inversion Techniques for Trace gas Emissions (MAGRITTE v1.1) – Part 1: Chemical mechanism, *Geosci. Model Dev.*, 12, 2307–2356, <https://doi.org/10.5194/gmd-12-2307-2019>, 2019.
- Myers, N., Mittermeier, R. A., Mittermeier, C. G., da Fonseca, G. A. and Kent, J.: Biodiversity Hotspots for Conservation Priorities, *Nature*, 403(6772), 853–858, doi:10.1038/35002501, 2000.
- Nguyen, T. B., Crounse, J. D., Teng, A. P., St. Clair, J. M., Paulot, F., Wolfe, G. M. and Wennberg, P. O.: Rapid deposition of oxidized biogenic compounds to a temperate forest, *Proceedings of the National Academy of Sciences*, 112(5), doi:10.1073/pnas.1418702112, 2015.
- Nölscher, A. C., Yañez-Serrano, A. M., Wolff, S., de Araujo, A. C., Lavrič, J. V., Kesselmeier, J. and Williams, J.: Unexpected seasonality in quantity and composition of Amazon Rainforest air reactivity, *Nature Communications*, 7(1), doi:10.1038/ncomms10383, 2016.
- Orlando, J. J. and Tyndall, G. S.: The atmospheric oxidation of ethyl formate and ethyl acetate over a range of temperatures and oxygen partial pressures, *International Journal of Chemical Kinetics*, 42(7), 397–413, doi:10.1002/kin.20493, 2010.
- Palmer, P. I., Jacob, D. J., Chance, K., Martin, R. V., Spurr, R. J., Kurosu, T. P., Bey, I., Yantosca, R., Fiore, A. and Li, Q.: Air mass factor formulation for spectroscopic measurements from satellites: Application to formaldehyde retrievals from the Global Ozone Monitoring Experiment, *Journal of Geophysical Research: Atmospheres*, 106(D13), 14539–14550, doi:10.1029/2000jd900772, 2001.
- Paulot, F., Crounse, J. D., Kjaergaard, H. G., Kürten, A., St. Clair, J. M., Seinfeld, J. H. and Wennberg, P. O.: Unexpected epoxide formation in the gas-phase photooxidation of isoprene, *Science*, 325(5941), 730–733, doi:10.1126/science.1172910, 2009.
- Paulot, F., Wunch, D., Crounse, J. D., Toon, G. C., Millet, D. B., DeCarlo, P. F., Vigouroux, C., Deutscher, N. M., González Abad, G., Notholt, J., Warneke, T., Hannigan, J. W., Warneke, C., de Gouw, J. A., Dunlea, E. J., De Mazière, M., Griffith, D. W. T., Bernath, P., Jimenez, J. L., and Wennberg, P. O.: Importance of secondary sources in the atmospheric budgets of formic and acetic acids, *Atmos. Chem. Phys.*, 11, 1989–2013, <https://doi.org/10.5194/acp-11-1989-2011>, 2011.
- Peeters, J., Nguyen, T. L. and Vereecken, L.: HOx radical regeneration in the oxidation of isoprene, *Physical Chemistry Chemical Physics*, 11(28), 5935, doi:10.1039/b908511d, 2009.

- Peeters, J., Müller, J.-F., Stavrou, T. and Nguyen, V. S.: Hydroxyl radical recycling in isoprene oxidation driven by hydrogen bonding and hydrogen tunneling: The upgraded LIM1 mechanism, *The Journal of Physical Chemistry A*, 118(38), 8625–8643, doi:10.1021/jp5033146, 2014.
- Pfannerstill, E. Y., Reijrink, N. G., Edtbauer, A., Ringsdorf, A., Zannoni, N., Araújo, A., Ditas, F., Holanda, B. A., Sá, M. O., Tsokankunku, A., Walter, D., Wolff, S., Lavrič, J. V., Pöhlker, C., Sörgel, M., and Williams, J.: Total OH reactivity over the Amazon rainforest: variability with temperature, wind, rain, altitude, time of day, season, and an overall budget closure, *Atmos. Chem. Phys.*, 21, 6231–6256, <https://doi.org/10.5194/acp-21-6231-2021>, 2021.
- Poraicu, C., Müller, J.-F., Stavrou, T., Fonteyn, D., Tack, F., Deutsch, F., Laffineur, Q., Van Malderen, R., and Veldeman, N.: Cross-evaluating WRF-Chem v4.1.2, TROPOMI, APEX, and in situ NO₂ measurements over Antwerp, Belgium, *Geosci. Model Dev.*, 16, 479–508, <https://doi.org/10.5194/gmd-16-479-2023>, 2023.
- Praene, J. P., David, M., Sinama, F., Morau, D. and Marc, O.: Renewable energy: Progressing towards a Net Zero Energy Island, the case of Reunion Island, *Renewable and Sustainable Energy Reviews*, 16(1), 426–442, doi:10.1016/j.rser.2011.08.007, 2012.
- Prather, M. J. and Zhu, L.: Resetting tropospheric OH and CH₄ lifetime with ultraviolet H₂O absorption, *Science*, 385(6705), 201–204, doi:10.1126/science.adn0415, 2024.
- Price, C. and Rind, D.: A simple lightning parameterization for calculating Global Lightning distributions, *Journal of Geophysical Research: Atmospheres*, 97(D9), 9919–9933, doi:10.1029/92jd00719, 1992.
- Pye, H. O., Ward-Caviness, C. K., Murphy, B. N., Appel, K. W. and Seltzer, K. M.: Secondary Organic Aerosol Association with cardiorespiratory disease mortality in the United States, *Nature Communications*, 12(1), doi:10.1038/s41467-021-27484-1, 2021.
- Read, K. A., Carpenter, L. J., Arnold, S. R., Beale, R., Nightingale, P. D., Hopkins, J. R., Lewis, A. C., Lee, J. D., Mendes, L. and Pickering, S. J.: Multiannual observations of acetone, methanol, and acetaldehyde in remote tropical Atlantic Air: Implications for atmospheric OVOC budgets and oxidative capacity, *Environmental Science & Technology*, 46(20), 11028–11039, doi:10.1021/es302082p, 2012.

- 1620 Rivera-Rios, J. C., Nguyen, T. B., Crounse, J. D., Jud, W., St. Clair, J. M., Mikoviny, T., Gilman, J. B., Lerner, B. M., Kaiser, J. B., de Gouw, J., Wisthaler, A., Hansel, A., Wennberg, P. O., Seinfeld, J. H. and Keutsch, F. N.: Conversion of hydroperoxides to carbonyls in field and laboratory instrumentation: Observational bias in diagnosing pristine versus anthropogenically controlled atmospheric chemistry, *Geophysical Research Letters*, 41(23), 8645–8651, doi:10.1002/2014gl061919, 2014.
- 1625 Rocco, M., Colomb, A., Baray, J.-L., Amelynck, C., Verreyken, B., Borbon, A., Pichon, J.-M., Bouvier, L., Schoon, N., Gros, V., Sarda-Esteve, R., Tulet, P., Metzger, J.-M., DufLOT, V., Guadagno, C., Peris, G. and Brioude, J.: Analysis of volatile organic compounds during the OCTAVE campaign: Sources and distributions of formaldehyde on Reunion Island, *Atmosphere*, 11(2), 140, doi:10.3390/atmos11020140, 2020.
- 1630 Rocco, M., Baray, J. -L., Colomb, A., Borbon, A., Dominutti, P., Tulet, P., Amelynck, C., Schoon, N., Verreyken, B., DufLOT, V., Gros, V., Sarda-Estève, R., Péris, G., Guadagno, C. and Leriche, M.: High resolution dynamical analysis of Volatile Organic Compounds (VOC) measurements during the Bio-Maïdo Field Campaign (Réunion Island, Indian Ocean), *Journal of Geophysical Research: Atmospheres*, 127(4), doi:10.1029/2021jd035570, 2022.
- 1635 Ryu, Y.-H., Hodzic, A., Barre, J., Descombes, G., and Minnis, P.: Quantifying errors in surface ozone predictions associated with clouds over the CONUS: a WRF-Chem modeling study using satellite cloud retrievals, *Atmos. Chem. Phys.*, 18, 7509–7525, <https://doi.org/10.5194/acp-18-7509-2018>, 2018.
- 1640 Saunders, S. M., Jenkin, M. E., Derwent, R. G., and Pilling, M. J.: Protocol for the development of the Master Chemical Mechanism, MCM v3 (Part A): tropospheric degradation of non-aromatic volatile organic compounds, *Atmos. Chem. Phys.*, 3, 161–180, <https://doi.org/10.5194/acp-3-161-2003>, 2003.
- 1645 Senten, C., De Mazière, M., Dils, B., Hermans, C., Kruglanski, M., Neefs, E., Scolas, F., Vandaele, A. C., Vanhaelewyn, G., Vigouroux, C., Carleer, M., Coheur, P. F., Fally, S., Barret, B., Baray, J. L., Delmas, R., Leveau, J., Metzger, J. M., Mahieu, E., Boone, C., Walker, K. A., Bernath, P. F., and Strong, K.: Technical Note: New ground-based FTIR measurements at Ile de La Réunion: observations, error analysis, and comparisons with independent data, *Atmos. Chem. Phys.*, 8, 3483–3508, <https://doi.org/10.5194/acp-8-3483-2008>, 2008.
- 1650 Sherwen, T., Evans, M. J., Carpenter, L. J., Andrews, S. J., Lidster, R. T., Dix, B., Koenig, T. K., Sinreich, R., Ortega, I., Volkamer, R., Saiz-Lopez, A., Prados-Roman, C., Mahajan, A. S., and Ordóñez, C.: Iodine's impact on tropospheric oxidants: a global model study in GEOS-Chem, *Atmos. Chem. Phys.*, 16, 1161–1186, <https://doi.org/10.5194/acp-16-1161-2016>, 2016.

- Shin, H. H. and Hong, S.-Y.: Representation of the subgrid-scale turbulent transport in convective boundary layers at Gray-Zone Resolutions, *Monthly Weather Review*, 143(1), 250–271, doi:10.1175/mwr-d-14-00116.1, 2015.
- Shindell, D., Faluvegi, G., Lacis, A., Hansen, J., Ruedy, R. and Aguilar, E.: Role of tropospheric ozone increases in 20th-Century climate change, *Journal of Geophysical Research: Atmospheres*, 111(D8), doi:10.1029/2005jd006348, 2006.
- Shrivastava, M., Cappa, C. D., Fan, J., Goldstein, A. H., Guenther, A. B., Jimenez, J. L., Kuang, C., Laskin, A., Martin, S. T., Ng, N. L., Petaja, T., Pierce, J. R., Rasch, P. J., Roldin, P., Seinfeld, J. H., Shilling, J., Smith, J. N., Thornton, J. A., Volkamer, R., Wang, J., Worsnop, D. R., Zaveri, R. A., Zelenyuk, A. and Zhang, Q.: Recent advances in understanding secondary organic aerosol: Implications for global climate forcing, *Reviews of Geophysics*, 55(2), 509–559, doi:10.1002/2016rg000540, 2017.
- Sicard, P., Anav, A., De Marco, A., and Paoletti, E.: Projected global ground-level ozone impacts on vegetation under different emission and climate scenarios, *Atmos. Chem. Phys.*, 17, 12177–12196, <https://doi.org/10.5194/acp-17-12177-2017>, 2017.
- Singh, H. B., Tabazadeh, A., Evans, M. J., Field, B. D., Jacob, D. J., Sachse, G., Crawford, J. H., Shetter, R. and Brune, W. H.: Oxygenated volatile organic chemicals in the oceans: Inferences and implications based on atmospheric observations and air-sea exchange models, *Geophysical Research Letters*, 30(16), doi:10.1029/2003gl017933, 2003.
- Sinha, V., Williams, J., Lelieveld, J., Ruuskanen, T. M., Kajos, M. K., Patokoski, J., Hellen, H., Hakola, H., Mogensen, D., Boy, M., Rinne, J. and Kulmala, M.: Oh reactivity measurements within a Boreal Forest: Evidence for unknown reactive emissions, *Environmental Science & Technology*, 44(17), 6614–6620, doi:10.1021/es101780b, 2010.
- Sommariva, R., de Gouw, J. A., Trainer, M., Atlas, E., Goldan, P. D., Kuster, W. C., Warneke, C., and Fehsenfeld, F. C.: Emissions and photochemistry of oxygenated VOCs in urban plumes in the Northeastern United States, *Atmos. Chem. Phys.*, 11, 7081–7096, <https://doi.org/10.5194/acp-11-7081-2011>, 2011.
- Španěl, P., Diskin, A. M., Wang, T. and Smith, D.: A sift study of the reactions of H_3O^+ , no^+ and O_2^+ with hydrogen peroxide and peroxyacetic acid, *International Journal of Mass Spectrometry*, 228(2–3), 269–283, doi:10.1016/s1387-3806(03)00214-8, 2003.
- Spivakovsky, C. M., Logan, J. A., Montzka, S. A., Balkanski, Y. J., Foreman-Fowler, M., Jones, D. B., Horowitz, L. W., Fusco, A. C., Brenninkmeijer, C. A., Prather, M. J., Wofsy, S. C. and McElroy, M. B.: Three-dimensional climatological distribution of tropospheric oh: Update and evaluation, *Journal of Geophysical Research: Atmospheres*, 105(D7), 8931–8980, doi:10.1029/1999jd901006, 2000.

- Stavrakou, T., J.-F. Müller, I. De Smedt, M. Van Roozendaal, G. R. van der Werf, L. Giglio, and A. Guenther: Evaluating the performance of pyrogenic and biogenic emission inventories against one decade of space-based formaldehyde columns, *Atmos. Chem. Phys.*, 9, 1037–1060, <https://doi.org/10.5194/acp-9-1037-2009>, 2009.
- Stavrakou, T., Guenther, A., Razavi, A., Clarisse, L., Clerbaux, C., Coheur, P.-F., Hurtmans, D., Karagulian, F., De Mazière, M., Vigouroux, C., Amelynck, C., Schoon, N., Laffineur, Q., Heinesch, B., Aubinet, M., Rinsland, C., and Müller, J.-F.: First space-based derivation of the global atmospheric methanol emission fluxes, *Atmos. Chem. Phys.*, 11, 4873–4898, <https://doi.org/10.5194/acp-11-4873-2011>, 2011.
- Stavrakou, T., Müller, J.-F., Peeters, J., Razavi, A., Clarisse, L., Clerbaux, C., Coheur, P.-F., Hurtmans, D., De Mazière, M., Vigouroux, C., Deutscher, N. M., Griffith, D. W., Jones, N. and Paton-Walsh, C.: Satellite evidence for a large source of formic acid from boreal and tropical forests, *Nature Geoscience*, 5(1), 26–30, doi:10.1038/ngeo1354, 2012.
- Stavrakou, T., Müller, J.-F., Bauwens, M., De Smedt, I., Van Roozendaal, M., De Mazière, M., Vigouroux, C., Hendrick, F., George, M., Clerbaux, C., Coheur, P.-F., and Guenther, A.: How consistent are top-down hydrocarbon emissions based on formaldehyde observations from GOME-2 and OMI?, *Atmos. Chem. Phys.*, 15, 11861–11884, <https://doi.org/10.5194/acp-15-11861-2015>, 2015.
- Strasberg, D., Rouget, M., Richardson, D. M., Baret, S., Dupont, J. and Cowling, R. M.: An assessment of habitat diversity and transformation on La Réunion Island (Mascarene Islands, Indian Ocean) as a basis for identifying broad-scale conservation priorities, *Biodiversity and Conservation*, 14(12), 3015–3032, doi:10.1007/s10531-004-0258-2, 2005.
- Takhar, M., Li, Y., Ditto, J. C., and Chan, A. W. H.: Formation pathways of aldehydes from heated cooking oils, *Environ. Sci.: Process. Impacts*, 25, 165-175, <https://doi.org/10.1039/D1EM00532D>, 2023.
- Tani, A., Muramatsu, K. and Mochizuki, T.: Emission of methyl ethyl ketone and 2-butanol converted from methyl vinyl ketone in plant leaves, *Atmosphere*, 11(8), 793, doi:10.3390/atmos11080793, 2020.
- Tewari, M., Chen, F., Wang, W., Dudhia, J., LeMone, M. A., Mitchell, K., Ek, M., Gayno, G., Wegiel, J., and Cuenca, R. H.: Implementation and verification of the unified NOAA land surface model in the WRF model, in: 20th conference on weather analysis and forecasting/16th conference on numerical weather prediction, Seattle, WA, USA, 10–16 January 2004, Abstract 14.2A, https://ams.confex.com/ams/84Annualtechprogram/paper_69061.htm (last access: 01 February 2024), 2004.

- Thames, A. B., Brune, W. H., Miller, D. O., Allen, H. M., Apel, E. C., Blake, D. R., Bui, T. P., Commane, R., Crounse, J. D., Daube, B. C., Diskin, G. S., DiGangi, J. P., Elkins, J. W., Hall, S. R., Hanisco, T. F., Hannun, R. A., Hints, E., Hornbrook, R. S., Kim, M. J., McKain, K., Moore, F. L., Nicely, J. M., Peischl, J., Ryerson, T. B., St. Clair, J. M., Sweeney, C., Teng, A., Thompson, C. R., Ullmann, K., Wennberg, P. O. and Wolfe, G. M.: Missing OH reactivity in the global marine boundary layer, *Atmospheric Chemistry and Physics*, 20(6), 4013–4029, doi:10.5194/acp-20-4013-2020, 2020.
- Travis, K. R., Heald, C. L., Allen, H. M., Apel, E. C., Arnold, S. R., Blake, D. R., Brune, W. H., Chen, X., Commane, R., Crounse, J. D., Daube, B. C., Diskin, G. S., Elkins, J. W., Evans, M. J., Hall, S. R., Hints, E. J., Hornbrook, R. S., Kasibhatla, P. S., Kim, M. J., Luo, G., McKain, K., Millet, D. B., Moore, F. L., Peischl, J., Ryerson, T. B., Sherwen, T., Thames, A. B., Ullmann, K., Wang, X., Wennberg, P. O., Wolfe, G. M., and Yu, F.: Constraining remote oxidation capacity with ATom observations, *Atmos. Chem. Phys.*, 20, 7753–7781, <https://doi.org/10.5194/acp-20-7753-2020>, 2020.
- van Geffen, J., Boersma, K. F., Eskes, H., Sneep, M., ter Linden, M., Zara, M., and Veefkind, J. P.: S5P TROPOMI NO₂ slant column retrieval: method, stability, uncertainties and comparisons with OMI, *Atmos. Meas. Tech.*, 13, 1315–1335, <https://doi.org/10.5194/amt-13-1315-2020>, 2020.
- van Geffen, J., Eskes, H., Compernelle, S., Pinardi, G., Verhoelst, T., Lambert, J.-C., Sneep, M., ter Linden, M., Ludewig, A., Boersma, K. F., and Veefkind, J. P.: Sentinel-5P TROPOMI NO₂ retrieval: impact of version v2.2 improvements and comparisons with OMI and ground-based data, *Atmos. Meas. Tech.*, 15, 2037–2060, <https://doi.org/10.5194/amt-15-2037-2022>, 2022a.
- van Geffen, J., Eskes, H. J., Boersma, K., and Veefkind, J.: TROPOMI ATBD of the total and tropospheric NO₂ data products, Royal Netherlands Meteorological Institute, https://sentinels.copernicus.eu/web/sentinel/data-products/-/asset_publisher/fp37fc19FN8F/content/sentinel-5-precursor-level-2-nitrogen-dioxide (last access: 3 February 2023), 2022b.
- Veefkind, J. P., Aben, I., McMullan, K., Förster, H., de Vries, J., Otter, G., Claas, J., Eskes, H. J., de Haan, J. F., Kleipool, Q., van Weele, M., Hasekamp, O., Hoogeveen, R., Landgraf, J., Snel, R., Tol, P., Ingmann, P., Voors, R., Kruizinga, B., Vink, R., Visser, H. and Levelt, P. F.: Tropomi on the ESA sentinel-5 precursor: A GMES mission for global observations of the atmospheric composition for climate, air quality and ozone layer applications, *Remote Sensing of Environment*, 120, 70–83, doi:10.1016/j.rse.2011.09.027, 2012.
- Verreyken, B., Amelynck, C., Brioude, J., Müller, J.-F., Schoon, N., Kumps, N., Colomb, A., Metzger, J.-M., Lee, C. F., Koenig, T. K., Volkamer, R., and Stavrou, T.: Characterisation of African biomass burning plumes and impacts on the

- 1755 atmospheric composition over the south-west Indian Ocean, *Atmos. Chem. Phys.*, 20, 14821–14845,
<https://doi.org/10.5194/acp-20-14821-2020>, 2020.
- Verreyken, B., Amelynck, C., Schoon, N., Müller, J.-F., Brioude, J., Kumps, N., Hermans, C., Metzger, J.-M., Colomb, A.,
 and Stavrakou, T.: Measurement report: Source apportionment of volatile organic compounds at the remote high-altitude
 1760 Maïdo observatory, *Atmos. Chem. Phys.*, 21, 12965–12988, <https://doi.org/10.5194/acp-21-12965-2021>, 2021.
- Vigouroux, C., De Mazière, M., Demoulin, P., Servais, C., Hase, F., Blumenstock, T., Kramer, I., Schneider, M., Mellqvist,
 J., Strandberg, A., Velasco, V., Notholt, J., Sussmann, R., Stremme, W., Rockmann, A., Gardiner, T., Coleman, M., and
 Woods, P.: Evaluation of tropospheric and stratospheric ozone trends over Western Europe from ground-based FTIR network
 1765 observations, *Atmos. Chem. Phys.*, 8, 6865–6886, <https://doi.org/10.5194/acp-8-6865-2008>, 2008.
- Vigouroux, C., Stavrakou, T., Whaley, C., Dils, B., Duflot, V., Hermans, C., Kumps, N., Metzger, J.-M., Scolas, F.,
 Vanhaelewyn, G., Müller, J.-F., Jones, D. B. A., Li, Q., and De Mazière, M.: FTIR time-series of biomass burning products
 (HCN, C₂H₆, C₂H₂, CH₃OH, and HCOOH) at Reunion Island (21° S, 55° E) and comparisons with model data, *Atmos. Chem.*
 1770 *Phys.*, 12, 10367–10385, <https://doi.org/10.5194/acp-12-10367-2012>, 2012.
- Vigouroux, C., Bauer Aquino, C. A., Bauwens, M., Becker, C., Blumenstock, T., De Mazière, M., García, O., Grutter, M.,
 Guarin, C., Hannigan, J., Hase, F., Jones, N., Kivi, R., Koshelev, D., Langerock, B., Lutsch, E., Makarova, M., Metzger, J.-
 M., Müller, J.-F., Notholt, J., Ortega, I., Palm, M., Paton-Walsh, C., Poberovskii, A., Rettinger, M., Robinson, J., Smale, D.,
 1775 Stavrakou, T., Stremme, W., Strong, K., Sussmann, R., Té, Y., and Toon, G.: NDACC harmonized formaldehyde time series
 from 21 FTIR stations covering a wide range of column abundances, *Atmos. Meas. Tech.*, 11, 5049–5073,
<https://doi.org/10.5194/amt-11-5049-2018>, 2018.
- Vigouroux, C., Langerock, B., Bauer Aquino, C. A., Blumenstock, T., Cheng, Z., De Mazière, M., De Smedt, I., Grutter, M.,
 1780 Hannigan, J. W., Jones, N., Kivi, R., Loyola, D., Lutsch, E., Mahieu, E., Makarova, M., Metzger, J.-M., Morino, I., Murata,
 I., Nagahama, T., Notholt, J., Ortega, I., Palm, M., Pinardi, G., Röhling, A., Smale, D., Stremme, W., Strong, K., Sussmann,
 R., Té, Y., van Roozendaal, M., Wang, P., and Winkler, H.: TROPOMI–Sentinel-5 Precursor formaldehyde validation using
 an extensive network of ground-based Fourier-transform infrared stations, *Atmos. Meas. Tech.*, 13, 3751–3767,
<https://doi.org/10.5194/amt-13-3751-2020>, 2020.
- 1785 Wang, S., Hornbrook, R. S., Hills, A., Emmons, L. K., Tilmes, S., Lamarque, J.-F., Jimenez, J. L., Capuzano-Jost, P., Nault,
 B. A., Crounse, J. D., Wennberg, P. O., Thompson, C. R., Peischl, J., Moore, F., Nance, D., Hall, B., Elkins, J., Tanner, D.,
 Huey, L. G., Hall, S. R., Ullmann, K., Orlando, J. L., Tyndall, G. S., Flocke, F., Ray, E., Hanisco, T. F., Wolfe, G. M., St Clair,

- J., Commane, R., Daube, B., Barletta, B., Weinzierl, B., Dollner, M., Conly, A., Vitt, F., Wofsy, S. C., Riemer, D. D., and
 1790 Apel, E. C.: Atmospheric acetaldehyde: Importance of air-sea exchange and a missing source in the remote troposphere,
Geophys. Res. Lett., 48, 5601–5613, <https://doi.org/10.1029GL082034>, 2019.
- Wang, S., Apel, E. C., Schwantes, R. H., Bates, K. H., Jacob, D. J., Fischer, E. V., Hornbrook, R. S., Hills, A. J., Emmons, L.
 K., Pan, L. L., Honomichl, S., Tilmes, S., Lamarque, J.-F., Yang, M., Marandino, C. A., Saltzmann, E. S., de Bruyn, W.,
 1795 Kameyama, S., Tanimoto, H., Omori, Y., Hall, S. R., Ullmann, K., Ryerson, T. B., Thompson, C. R., Peischl, J., Daube, B.
 C., Commane, R., McKain, K., Sweeney, C., Thames, A. B., Miller, D. O., Brune, W. H., Diskin, G. S., DiGangi, J., and
 Wofsy, S. C.: Global atmospheric budget of acetone: Air-sea exchange and the contribution to hydroxyl radicals, *J. Geophys.*
Res., 125, e2020JD032553, <https://doi.org/10.1029/2020JD032553>, 2020.
- 1800 Wells, K., Millet, D., Brewer, J., Payne, V., Cady-Pereira, K., Pernak, R., Kulawik, S., Vigouroux, C., Jones, N., Mahieu, E.,
 Makarova, M., Nagahama, T., Ortega, I., Palm, M., Strong, K., Schneider, M., Smale, D., Sussmann, R., and Zhou, M.: Long-
 term global measurements of methanol, ethene, ethyne, and HCN from the Cross-track Infrared Sounder, *EGUsphere*
 [preprint], <https://doi.org/10.5194/egusphere-2024-1551>, 2024.
- 1805 Wennberg, P. O., Bates, K. H., Crounse, J. D., Dodson, L. G., McVay, R. C., Mertens, L. A., Nguyen, T. B., Praske, E.,
 Schwantes, R. H., Smarte, M. D., St Clair, J. M., Teng, A. P., Zhang, X. and Seinfeld, J. H.: Gas-phase reactions of isoprene
 and its major oxidation products, *Chemical Reviews*, 118(7), 3337–3390, doi:10.1021/acs.chemrev.7b00439, 2018.
- Williams, J. E., Boersma, K. F., Le Sager, P., and Verstraeten, W. W.: The high-resolution version of TM5-MP for optimized
 1810 satellite retrievals: description and validation, *Geosci. Model Dev.*, 10, 721–750, <https://doi.org/10.5194/gmd-10-721-2017>,
 2017.
- Wong, J., Barth, M. C., and Noone, D.: Evaluating a lightning parameterization based on cloud-top height for mesoscale
 numerical model simulations, *Geosci. Model Dev.*, 6, 429–443, <https://doi.org/10.5194/gmd-6-429-2013>, 2013.
 1815
- Yáñez-Serrano, A. M., Nölscher, A. C., Williams, J., Wolff, S., Alves, E., Martins, G. A., Bourtsoukidis, E., Brito, J., Jardine,
 K., Artaxo, P., and Kesselmeier, J.: Diel and seasonal changes of biogenic volatile organic compounds within and above an
 Amazonian rainforest, *Atmos. Chem. Phys.*, 15, 3359–3378, <https://doi.org/10.5194/acp-15-3359-2015>, 2015.
- 1820 Yáñez-Serrano, A. M., Nölscher, A. C., Bourtsoukidis, E., Derstroff, B., Zannoni, N., Gros, V., Lanza, M., Brito, J., Noe, S.
 M., House, E., Hewitt, C. N., Langford, B., Nemitz, E., Behrendt, T., Williams, J., Artaxo, P., Andreae, M. O., and Kesselmeier,

J.: Atmospheric mixing ratios of methyl ethyl ketone (2-butanone) in tropical, boreal, temperate and marine environments, *Atmos. Chem. Phys.*, 16, 10965–10984, <https://doi.org/10.5194/acp-16-10965-2016>, 2016.

1825 Zhao, Z., Pritchard, M. S. and Russell, L. M.: Effects on precipitation, clouds, and temperature from long-range transport of idealized aerosol plumes in WRF-Chem simulations, *Journal of Geophysical Research: Atmospheres*, 117(D5), doi:10.1029/2011jd016744, 2012.

1830 Zhao, Y., Saunio, M., Bousquet, P., Lin, X., Berchet, A., Hegglin, M. I., Canadell, J. G., Jackson, R. B., Hauglustaine, D. A., Szopa, S., Stavert, A. R., Abraham, N. L., Archibald, A. T., Bekki, S., Deushi, M., Jöckel, P., Josse, B., Kinnison, D., Kirner, O., Marécal, V., O'Connor, F. M., Plummer, D. A., Revell, L. E., Rozanov, E., Stenke, A., Strode, S., Tilmes, S., Dlugokencky, E. J., and Zheng, B.: Inter-model comparison of global hydroxyl radical (OH) distributions and their impact on atmospheric methane over the 2000–2016 period, *Atmos. Chem. Phys.*, 19, 13701–13723, <https://doi.org/10.5194/acp-19-13701-2019>, 2019.

1835