



Machine Learning Assisted Inference of the Particle Charge Fraction and the Ion-induced Nucleation Rates during New Particle Formation Events

Pan Wang¹, Yue Zhao¹, Jiandong Wang^{2,3}, Veli-Matti Kerminen⁴, Jingkun Jiang⁵, Chenxi Li^{1*}

5

¹School of Environmental Science and Engineering, Shanghai Jiao Tong University, 200240, Shanghai, China

²Collaborative Innovation Center on Forecast and Evaluation of Meteorological Disasters, Nanjing University of Information Science and Technology, Nanjing, China

10 ³China Meteorological Administration Aerosol-Cloud-Precipitation Key Laboratory, School of Atmospheric Physics, Nanjing University of Information Science and Technology, Nanjing, China

⁴Institute for Atmospheric and Earth System Research / Physics, Faculty of Science, University of Helsinki, 00014 Helsinki, Finland

⁵State Key Joint Laboratory of Environment Simulation and Pollution Control, School of Environment, Tsinghua University, 100084 Beijing, China

15 *Correspondence to: Chenxi Li (chenxi20@sjtu.edu.cn)



Abstract: The charge state of atmospheric new particles is controlled by both their initial charge state upon formation and subsequent interaction with atmospheric ions. By measuring the charge state of growing particles, the fraction of ion-induced nucleation (F_{IIN}) within total new particle formation (NPF) can be inferred, which is critical for understanding NPF mechanisms. However, existing theoretical approaches for predicting particle charge states suffer from inaccuracies due to simplifying assumptions, hence their ability to infer F_{IIN} is sometimes limited. Here we develop a numerical model to explicitly simulate the charging dynamics of new particles. Our simulations demonstrate that both particle growth rate and ion concentration substantially influence the particle charge state, while ion-ion recombination becomes important when the charged particle concentrations are high. Leveraging a large set of simulations, we constructed two regression models using residual neural networks. The first model (ResFWD) predicts the charge state of growing particles with known F_{IIN} values, while the second model (ResBWD) operates in reverse to estimate F_{IIN} based on the charge fraction of particles at prescribed sizes. Good agreement between the regression models and benchmark simulations demonstrates the potential of our approach for analysing ion-induced nucleation events. Sensitivity analysis further reveals that ResFWD and the benchmark simulations exhibit similar sensitivity to input noises, but the robustness of ResBWD requires that the information of initial particle charge state is retained at the prescribed sizes. Our study provides insights on charging dynamics of atmospheric new particles and introduces a new method for assessing ion-induced nucleation rates.



1 Introduction

In the low atmosphere, ions are continuously produced by galactic cosmic rays and radioisotope decay at the earth surface (Stozhkov, 2003; Eisenbud and Gesell, 1997). Due the high abundance of N_2 and O_2 in the atmosphere, the initially formed ions (primary ions) typically include N_2^+ , O_2^+ , NO^+ , O^- , and O_2^- . These primary ions subsequently go through ion-molecule reactions to form a large set of organic and inorganic secondary ions, e.g., NH_4^+ , NO_3^- , HSO_4^- and $C_3H_3O_4^-$ (Yin et al., 2023). Once formed, ions can be lost by condensing on the aerosol particles, deposition to surfaces, and ion-ion recombination, leading to an typical ion concentration of $100\text{-}5000\text{ cm}^{-3}$ globally (Hirsikko et al., 2011). The ions contribute to atmospheric electricity and play an important role in the formation of aerosol particle (Golubenko et al., 2020; Kerminen et al., 2018; Yu et al., 2012).

New particle formation (NPF) is the conversion of gas molecules to nascent nanoparticles and is estimated to contribute about half of the cloud condensation nuclei on a global scale (Gordon et al., 2017; Zhao et al., 2024). Atmospheric ions can participate in NPF events during both the nucleation stage (i.e., the process in which the stable clusters are formed from gas phase precursors) and the growth stage (in which the clusters further grow due to vapor condensation and coagulation). During nucleation, ions can induce NPF at lower vapor concentrations than neutrals by stabilizing the embryonic clusters by the presence of the charge (Kirkby et al., 2016; Yu et al., 2020). Measurements even suggest that ion-induced nucleation (IIN) might be the main mechanism for NPF in the higher troposphere and the stratosphere (Yu et al., 2008; Lee et al., 2003; Zhao et al., 2024). Atmospheric ions also play a role in particle growth by altering the charge state of the particles and affect their growth in several ways. Firstly, charged particles tend to have higher condensational growth rates (GR) due to enhanced ion and neutral vapor condensation, which are caused by Coulombic, charge-dipole and charge-induced dipole interactions between the particles and the condensing species (Svensmark et al., 2017; Nadykto and Yu, 2003). Second, particle charging promotes coagulation between charged particle of opposite polarities and oppress coagulation between particles of the same polarity. Third, the coagulation sink (CoagS) for charged particles can be different from that of the neutral particles, which makes charged particles less likely to grow larger.

An accurate estimation of IIN rates is a prerequisite to assess the role of ions in NPF. However, IIN rates are often challenging to measure directly because the INN pathway must be distinguished from particle formation through neutral pathways that proceed simultaneously. Additionally, the constant interaction with atmospheric ions alter the particle charge state and makes it difficult to determine whether a given particle is charged upon formation or during growth. Therefore, the IIN rates is often deduced by comparing the charge fraction of nucleated particles to the so-called steady state particle charge distribution, using a model that relates these two quantities given other measurables (e.g., the particle growth rate, the ion concentration) (Iida et al., 2006). Towards this end, Kerminen et al. (2007) developed an analytical equation to calculate the charge fraction of particles at a given size. By fitting the theoretical values with measured particle charge fraction at several sizes (Laakso et al., 2007), the IIN fraction can be obtained. This equation was further extended to deal with situations with different positive and negative ions concentrations (Gagné et al., 2012). However, as shown by comparison with numerical simulation (Leppä et al., 2011; Leppä et al., 2009), the accuracy of the theoretical approach is sometimes limited by its



underlying assumptions, e.g., the particle population is monodisperse and the charged fraction of the particles is substantially below unity.

Machine learning (ML) is increasingly applied in atmospheric sciences due to its capability to deal with complex and non-linear processes. In the study of atmospheric NPF, ML has been applied to identify NPF and non-NPF days (Su et al., 2022; Joutsensaari et al., 2018), to speed up configurational sampling of embryonic clusters (Kubečka et al., 2023), and to train force fields used in molecular dynamics simulations of NPF (Jiang et al., 2022). Conceivably, ML can also be applied to calculate the charge fraction of atmospheric new particle in lieu of the theoretical equations, with potentially higher accuracy and less restrictive assumptions. An even more ambitious goal is to directly calculate F_{IIN} with measurable particle charge fractions using a trained ML model, hence circumventing the data fitting procedure.

In this work, we present an initial exploration of machine learning (ML) models to infer particle charge fractions and ion-induced nucleation (IIN) rates during NPF events. To achieve this goal, we couple dynamic charging simulations with a sectional model (Li et al., 2023) to simulate NPF under typical atmospheric conditions. The data generated from these benchmark simulations are then utilized to train and validate ML models. Both the accuracy and sensitivity of the ML models to input noises are discussed and compared with benchmark simulations.

2 Methods

2.1 The sectional model

We applied a two-dimensional sectional model (Fig. 1) to simulate the evolution of the particle size distribution and particle charge fraction during the NPF events. We refer to this model as CDMS-ion (cluster dynamics multicomponent section model-with ions) for brevity. CDMS-ion divides the particles into mass sections, and the particles within a mass section are further divided into subsections according to their charge states. All particles in the same mass section are assumed to have the same chemical composition (internally mixed). The simulated processes include particle charging, coagulation, growth or shrinkage due to vapor condensation-evaporation and losses to pre-existing particles (i.e., coagulation sink (CoagS)). Particle nucleation is not explicitly simulated; rather, prescribed nucleation rates at 1 nm are specified in the model as an input.

The processes under consideration were simulated using an operator splitting approach, where the differential equations for distinct processes were solved sequentially within each time step, h_{step} . In our previous work, which did not include particle charging (Li et al., 2023), these equations were solved simultaneously. However, incorporating an additional dimension—the particle charge state—significantly increases computational costs. Therefore, in this study, we employed the operator splitting method to enhance simulation efficiency. To determine an optimal h_{step} , we conducted a convergence study, gradually reducing h_{step} and observing its effect on simulation outcomes. We found that values lower than 20 seconds had a negligible impact on our results. For all simulations, we utilized 126 mass sections with a geometric factor of 1.1, covering a particle size range from 1.17 to 100.50 nm. A uniform particle density of 1.4 g cm^{-3} was assumed.




Figure 1. (a) Illustration of the CDMS-ion model. (b) The evolution of the particle size distribution in a simulated NPF event. The structures of (c) ResFWD and (d) ResBWD.

100 2.2 Simulation of particle charging

The interaction between particles and atmospheric ions was simulated with a dynamic particle charging module. Neutral particles collide with ions to generate charged particle, and charged particles increase/decrease its charge by colliding with ions of the same/opposite polarity. This dynamic process is described by the following equation,

$$n_{d_p,k} = \beta_{d_p,k-1,+} n_{d_p,k-1} n_+ - \beta_{d_p,k,+} n_{d_p,k} n_+ + \beta_{d_p,k+1,-} n_{d_p,k+1} n_- - \beta_{d_p,k,-} n_{d_p,k} n_- \quad (1)$$

105 where $n_{d_p,k}$ is the concentration of particles with a diameter of d_p and k charges, $\beta_{d_p,k,\pm}$ is the collision rate constant between these particles and positive/negative ions, and n_{\pm} is the concentrations of positive or negative ions. Since we are interested in particles formed during NPF events with sizes smaller than 100 nm, we set the maximum particle charge to be ± 5 . The concentrations of particle with more charges are negligible under atmospherically relevant conditions (Wiedensohler, 1988).



To solve Eq. (1), the collision rate constant $\beta_{d_p,k,\pm}$ needs to be calculated accurately. In this work, we used the rate
 110 coefficients developed by López-Yglesias and Flagan (2013), who considered both three body trapping and image potential in
 their calculations. The collision rate constant was calculated with the following expression:

$$\beta_{d_p,k,\pm} = 10^{\sum_{q=0}^Q B_{q,\pm}(k)(\log_{10}(\frac{d_p}{2}))^q} \quad (2)$$

where $B_{q,\pm}(k)$ are fit coefficients, q is number of charges on the particle, and $Q = 23$ is the maximum of q .

2.3 Vapor condensation-evaporation

115 Particle growth due condensation-evaporation of sulfuric acid and oxygenated organic molecules (OOMs) was simulated
 according to the following equation:

$$\frac{dm_p}{dt} = \sum_i m_i (\beta_i n_i - E_i) \quad (3)$$

where m_p is the particle mass, m_i is the molecular mass of the species i , β_i is the collision constant of species i with the particle,
 n_i is the gas phase concentration of species i , E_i is the evaporation rate of species i from the particle.

120 To calculate β_i in Eq. (3), we first calculated the collision rate coefficients with the regression equation developed by
 Gopalakrishnan and Hogan (2011) and subsequently multiply these coefficients with an enhancement factor to account for
 charge-dipole interactions between the particles and the vapors. The expression for the enhancement factor is given by Nadykto
 and Yu (2003) :

$$EF = 1 + \frac{lE(d_p+d_v)L(\frac{lE(d_p+d_v)}{2bT})+0.5\alpha\epsilon_0E^2(d_p+d_v)}{3bT} \quad (4)$$

125 where l is dipole moment of the vapor, $E(r) = (\frac{1}{\epsilon_g} - \frac{1}{\epsilon_p}) \frac{ke_0}{4\pi\epsilon_0r^2}$ is the electrical field of the charged particle, ϵ_g and ϵ_p are
 relative permittivity of air and the particle, respectively, ϵ_0 is the vacuum permittivity, e_0 is the elementary charge, k is the
 number of charges of the particle, d_p and d_v are the diameters of the particle and the vapor molecule, respectively, $L(z) =$
 $\frac{e^z+e^{-z}}{e^z-e^{-z}} - \frac{1}{z}$ is the Langevin function, and α is the polarizability of the molecules. In the calculation of enhancement factors
 involving sulfuric acid molecules, we set the dipole moment and polarizability to 2.84 Debye and 6.2 \AA^3 , respectively (Nadykto
 130 and Yu, 2003). For collisions involving OOMs molecules, due to the lack of information on the average dipole moment and
 polarizability, we calculated the enhancement factor of EF_{OOMs} with an empirical relation developed by Kirkby et al. (2016):

$$EF_{OOMs} = \frac{EF_{SA}-1}{f_{OOMs,SA}} + 1 \quad (5)$$

where $f_{OOMs,SA} = 4$ is a fitting parameter.

Within each simulation interval (20 s), we calculated the growth of particles in each subsection with a Lagrangian
 135 approach using Eq. (3) (Zaveri et al., 2008). At the end of an interval, the grown particles are distributed into different mass
 bins using the Linear Discrete Method (Simmel and Wurzler, 2006). The particle charge state was preserved during particle
 growth.



2.4 Particle coagulation

We considered the effect of Coulombic interactions on particle coagulation. The coagulation rate coefficients were calculated with the equations developed by Gopalakrishnan and co-workers (Ouyang et al., 2012; Gopalakrishnan and Hogan, 2012, 2011; Chahl and Gopalakrishnan, 2019), who derived the rate coefficients using Langevin dynamics simulations. The expressions for the collision rate coefficients are given by

$$\beta(d_{p1}, k_1, d_{p2}, k_2) = \frac{H f_r (d_1 + d_2)^3 \eta_{FM}^2}{m_r \eta_C} \quad (6)$$

$$H = \exp(\mu) \frac{4\pi K n_D^2 + 25.836 K n_D^3 + (8\pi)^{1/2} 11.211 K n_D^4}{1 + 3.502 K n_D + 7.211 K n_D^2 + 11.211 K n_D^3} \quad (7)$$

$$K n_D = \frac{2(k_b T m_r)^{1/2} \eta_C}{f_r (d_1 + d_2) \eta_{FM}} \quad (8)$$

where H is the dimensionless collision rate constant, $K n_D$ is the diffusive Knudsen number, k_b is Boltzmann's constant, T is the ambient temperature, d_{p1} and d_{p2} are the diameters of two colliding particles, respectively, m_r is the reduced mass of the colliding particles (defined as $m_r = m_1 m_2 / (m_1 + m_2)$), f_r is the reduced friction factor (defined as $f_r = f_1 f_2 / (f_1 + f_2)$), η_C is the continuum limit enhancement factor due to the presence of charge, and η_{FM} is the free molecular limit enhancement factor (Gopalakrishnan and Hogan, 2012), μ is a function of the electrostatic energy to thermal energy ratio and the diffusive Knudsen number. Expressions for η_C , η_{FM} and μ are found in Gopalakrishnan and Hogan (2012) and Chahl and Gopalakrishnan (2019).

The explicit simulation of particle charge state significantly increases the computational cost of coagulation simulation as the number of coagulation pairs is proportional to the number of subsections squared. To speed up the simulation, we used the coagulation algorithm developed by Matsui, which is a simplified version of Jacobson's semi-implicit approach (Matsui, 2013; Matsui, 2017; Jacobson, 1994). In this algorithm, after a coagulation time step Δt , the mass concentration $M_{i,k}^{t+\Delta t}$ of particles with k charges in the i^{th} mass section is given by:

$$M_{i,k}^{t+\Delta t} = \frac{M_{i,k}^t + f_{corr} P_{i,k}^t}{1 + L_{i,k}^t} \quad (9)$$

where $P_{i,k}$ and $L_{i,k}$ are the mass production and loss rates of particles due to coagulation at time t , respectively, f_{corr} is a correction factor to ensure mass conservation. The expressions for $P_{i,k}$, $L_{i,k}$ and f_{corr} are given in Sect. S1 in the supplementary information (SI). Overall, this coagulation algorithm is non-iterative for any time step, conserves total particle mass but leads to slight inaccuracies in particle number distribution (Matsui and Mahowald, 2017; Matsui, 2017).

2.5 Coagulation sink

In addition to newly formed particles, the atmospheric ions also condition the charge distribution of the pre-existing atmospheric particles and affect the magnitude of the coagulation sink (CoagS). To account for this influence of charge on CoagS, we calculated CoagS with the assumption that the pre-existing particles are at steady state charge distribution due to interaction with atmospheric ions. This assumption is supported by field observations conducted by Li et al. (2022), which



show good agreement between the particle size distributions measured by the SMPS with and without a neutralizer. Additionally, background particles (larger in size) have a shorter characteristic charging time (see Fig. 2 below) and longer residence time in the atmosphere compared with newly formed particles, which further justifies the steady state assumption.

The coagulation sink $CoagS_{d_p,k}$ for particles with a diameter of d_p and k charges was calculated with the following equation,

$$CoagS_{d_p,k} = \sum_{k'=-6}^6 \int \beta_{d_p,k,d'_p,k'} f(d'_p,k') \frac{dN_{pre}}{dd'_p} dd'_p \quad (10)$$

where $\beta_{d_p,k,d'_p,k'}$ is the collision rate coefficient of particles with a diameter of d_p and k charges with background particles with a diameter of d'_p and k' charges, $f(d'_p,k')$ is the steady state fraction of particles with k' charges among all particles with size d'_p , $\frac{dN_{pre}}{dd'_p}$ is the number-based particle size distribution of pre-existing particles. $f(d'_p,k')$ can vary with time since the properties of atmospheric ions constantly change (Chen and Jiang, 2018). In this work, however, we assume that $f(d'_p,k')$ is independent of time for the pre-existing particles since the variation of ion properties is relatively small. To be consistent with the particle charging simulations (Sect. 2.2), the values of $f(d'_p,k')$ were calculated with the rate coefficients given by López-Yglesias and Flagan (2013) by solving Eq. (1). As to $\frac{dN_{pre}}{dd'_p}$, we assume that the background particles are lognormally distributed with a geometric mean diameter of 100 nm and a geometric standard deviation of 1.4.

2.6 Simulation setup and key metrics

We set up our simulations to mimic typical NPF events. Specifically, we assume that new particle formation lasts for 3 hours with a constant nucleation rate J . The newly formed particles enter the smallest section (particle size is about 1 nm) and start to grow due the condensation of sulfuric acid (SA) and oxygenated organic molecules (OOMs). The SA and OOMs concentrations are assumed to be constant in the simulation. SA is assumed to be non-evaporative and the OOMs are classified into 6 bins by saturation vapor concentration C^* ($\log_{10}C^* = -9, -7, -5, -3, -1, 0$ C^* in unit of $\mu\text{g m}^{-3}$). Simultaneous to condensational growth, the particles coagulate with other particles or lose to pre-existing particles. All simulations were conducted at 298.15 K and 1 atm. A typical PSD obtained from such simulations is shown in Fig. 1b.

Several factors influence the charge state of the new particles, including the atmospheric ion properties (e.g., mobility), the ion concentrations (N_{ion}), the particle growth rate (GR), and the coagulation sink (CoagS), the total nucleation rate (J) and the fraction of ion induced nucleation (F_{IIN} , which is equal to $\frac{IIN \text{ rates}}{IIN \text{ rates} + \text{neutral NPF rates}}$ and ranges from 0 to 1). The atmospheric ion properties used in this work is listed in Table 1. We set positive and negative ions to have the same mass and mobility. The properties of positive and negative ions can be different (e.g., in a neutralizer), but in the atmosphere the positive and negative ions often exhibit similar mobilities (Li et al., 2022; Gautam et al., 2017). The value of the other factors (N_{ion} , GR, CoagS and J) spanned ranges of typical NPF events in the atmospheric boundary layer (also shown in Table 1) (Chu et al., 2019; Kerminen et al., 2018). The ion concentration N_{ion} and the nucleation rate J were directly specified as simulation parameters, while GR



and CoagS were controlled indirectly in the simulation by scaling the SA and OOMs concentrations while maintaining their relative concentration. The reported GR values in the following was obtained by first simulating particle growth (Sect. S2 in the SI) and subsequently fitting the particle size as a function of time with a linear function. Therefore, the GR values reported in this study are a measure of particle growth rates due to neutral vapor condensation. We note that although GR defined in this way neglects the effect of coagulation on particle growth, it can be retrieved from the evolution of particle size distribution (Li and McMurry, 2018; Stolzenburg et al., 2005). To control CoagS, we scaled the concentration of the pre-existing aerosols while maintaining their distribution (lognormal distribution with a geometric mean diameter of 100 nm and a geometric standard deviation of 1.4).

To analyse the simulation results, we mainly focus on the charge ratio r_c , which is defined as the ratio of the simulated fraction of singly charged particles to the steady state value. This metric indicates to what extent the particle charge distribution deviates from the steady state: $r_c < 1$ indicates that the particles are undercharged and $r_c > 1$ indicates that the particles are overcharged. The second and third metrics are the maximum number of particles during a simulation (N_{\max}) and the particle mode diameter (d_m). Comparisons of these two metrics between simulations with and without particle charging shows the effect of charging on particle survival and growth.

Table 1. Simulation parameters

Symbol	Meaning	Value or range
μ_{\pm}	Ion mobility	$1.8 \times 10^{-4} \text{ m}^2 \text{ V}^{-1} \text{ s}^{-1}$
M_{\pm}	Ion mass	150 Da
GR	Growth rate	1-15 nm/h
J	The nucleation rate at 1 nm	$0.1\text{-}1000 \text{ \# cm}^{-3}$
CoagS	Coagulation sink (defined with respect to sulfuric acid molecules)	$0.001\text{-}0.02 \text{ s}^{-1}$
N_{ion}	Ion concentration	$50\text{-}5000 \text{ cm}^{-3}$

2.7 Analytical equation for particle charge state

Kerminen et al. (2007) derived a theoretical equation to calculate the charge state of a monodisperse nucleation mode:

$$r_{c,d_p} = 1 - \frac{1}{Kd_p} + \frac{1+(r_{c,0}-1)Kd_0}{Kd_p} \exp[-K(d_p - d_0)], \quad (11)$$

where r_{c,d_p} and $r_{c,0}$ are the charging state at size d_p and d_0 (i.e., the initial particle size), respectively. K is expressed as:

$$K = \frac{\alpha N_{ion}}{GR} \quad (12)$$

where α is the association rate between ions and particles of opposite polarity. In this work, we use a constant value of $1.8 \times 10^{-6} \text{ cm}^3 \text{ s}^{-1}$ for α , which is the collision rate constant between ions and particles 1 nm in diameter (calculated with



220 Eq. (2)). According to Eq. (11), the particle charge state is governed by both the initial charge state $r_{c,0}$ and the parameter K , which is directly proportional to ion concentration and inversely proportional to particle growth rate. We note that $r_{c,0}$ is a different concept from F_{IIN} : the former is the ratio of the particle charge fraction to the equilibrium charge fraction at the initial particle size, while the latter refers to the ratio of particle concentration fluxes past a threshold size. These two ratios can be significantly different (Leppä et al., 2013).

225 2.8 Regression Models with Neutral Network

In this study, we used a residual neural network (ResNet)-based architecture to construct regression models. ResNet addresses the problem of vanishing gradients through residual connections and can accelerate network convergence (He et al., 2016). Initially introduced to enhance image recognition performance, ResNet has demonstrated broad applicability across various fields, including emulation of atmospheric chemistry solvers (Kelp et al., 2018; Liu et al., 2021).

230 Our first application of ResNet was to determine the charge state of new particles, assuming that J , GR , N_{ion} , $CoagS$ and F_{IIN} are already known (Fig. 1c). The network consists of six fully connected layers with 64, 128, 256, 128, 64, and 1 node, respectively, with residual connections introduced between each layer. The input layer has 5 nodes corresponding to the $\log_{10}(J)$, GR , $\log_{10}(N_{\text{ion}})$, $CoagS$ and F_{IIN} , and the output layer has 1 node corresponding to $\log_{10}(r_c)$ at a specific size. \log_{10} values of N_{ion} , J , and r_c were used because their significant variation across approximately two orders of magnitude. Each fully
235 connected layer is followed by a ReLU activation function, with shortcut connections mapping the input of each layer directly to its output. The trained model is referred to as ResFWD (FWD denotes ‘forward’) and serves as an alternative of the Eq. (11).

In our second application of ResNet, we aimed to predict F_{IIN} based on r_c values at multiple sizes (2.2 nm, 3 nm, 4 nm, 5 nm, 6 nm, 7 nm, 8 nm), alongside $\log_{10}(J)$, GR , $\log_{10}(N_{\text{ion}})$, and $CoagS$ (Fig. 1d). This model's input layer consists of 11
240 nodes, which is more complex compared with ResFWD. Consequently, we expanded the number of fully connected layers to eight, with node counts of 64, 128, 256, 512, 256, 128, 64, and 1, respectively. Batch normalization layers were incorporated to accelerate training and enhance the model's generalization ability, while other configurations remained consistent with the first model. The resulting trained model is termed ResBWD (BWD denotes ‘backward’).

The dataset used to train ResFWD consists of ~ 4 million CDMS-ion simulations, but this dataset was reduced in the
245 training of ResBWD by removing sets of simulations (each set corresponds to a specific combination of J , GR , N_{ion} and $CoagS$) in which the information of F_{IIN} is almost lost before the particles reach 2.2 nm due to interaction with atmospheric ions (discussed in Sect. 3.3). In training all ResNet models, 80% of the data were used for the model training and 20% were used for model validation. The max-min normalization method was used for data pre-processing of all input and output features. The models were trained with PyTorch with mean squared error (MSE) as the loss function. The optimizer was Adam, with a
250 learning rate set to 0.001. The batch size was set to 2048, and the training was conducted over at least 50 epochs.

3 Results and discussion

3.1 Evolution of particle charge state

In this section, we discuss some general characteristics of particle interaction with atmospheric ions, including the time scale for particles to reach steady state charge distribution (Sect. 3.1.1), how particle charge state evolves after formation (Sect. 3.1.2 and 3.1.3) and the influence of charging on particle number concentration and growth, which is included in the SI (Sect. S5).

3.1.1 Characteristic time to reach steady state charge distribution

To estimate the time scale for particles of a given size to achieve a steady-state charge distribution under ion influence, we calculated the characteristic time (τ_{ss}) for particle (dis)charging by solving Eq. (1). τ_{ss} is defined as the times it takes for the singly charged fraction of initially neutral particles reaches $(1 - 1/e)$ of the steady-state value, or for the singly charged fraction of initially charged particles reaches $(1 + 1/e)$ of the steady-state value. We neglect multiply charged particles in this calculation, as their occurrence is typically low for ultrafine particles (Wiedensohler, 1988). Additionally, an analytical analysis of τ_{ss} is presented in Sect. S3 in the SI to elucidate certain features of τ_{ss} .




Figure 2. Contour plots of the characteristic time (in seconds) for particles to reach the steady state distribution as a function of particle diameter and ion concentration. (a) The particles are initially neutral. (b) The particles are initially singly charged (50% positive, 50% negative).

Figures 2a and 2b show contour plots of τ_{ss} at NPF-relevant particle sizes (1-100 nm) at atmospherically relevant ion concentrations ($50-10^4 \text{ cm}^{-3}$, note that throughout this work the ion concentration N_{ion} refers to the sum of positive and negative ion concentration) for initially neutral and initially charged particles, respectively. Apparently, τ_{ss} is dependent on both the



particle size and the ion concentration. On the one hand, for particles of all sizes, τ_{ss} decreases as ion concentration increases since the characteristic time is inversely proportional to the ion concentration (Eq. (S12) in the SI). On the other hand, at a fixed ion concentration, τ_{ss} stays relatively constant or decreases with increasing particle size. This trend is caused by the increased collision rates between ions and larger particles, to which τ_{ss} is inverse proportional (Eq. (S12) in the SI). Further
275 comparison between Fig. 2a and Fig. 2b reveals that τ_{ss} is smaller for initially neutral particles than initially charged particles. Such a difference is caused by the asymmetry of the collision rate coefficients of ion-neutral collisions and ion-charged particle collisions (see Eq. (S12) and relevant text in the SI).

An uncertainty regarding τ_{ss} stems from the collision rate coefficients in its calculation. Pfeifer et al. (2023) showed that experimental and theoretical collision rate coefficients between ions and singly charged particles can differ by one to two
280 orders of magnitude (Pfeifer et al., 2023; López-Yglesias and Flagan, 2013; Gopalakrishnan and Hogan, 2012; Gatti and Kortshagen, 2008). The rate coefficients used in this study (i.e., López-Yglesias and Flagan (2013)) is on the higher end of these rates. If the rate expressions developed by Gatti and Kortshagen (2008) or Gopalakrishnan and Hogan (2012) had been utilized, we would anticipate a longer characteristic charging time. Additionally, we neglected the van der Waals potential between colliding entities, as its interplay with the Coulomb potential in influencing collision rates remains unclear.

285 The time scale τ_{ss} , as illustrated in Fig. 2, ranges from tens of seconds to several hours, overlapping with the time scale for new particles to grow to a few or tens of nanometers in NPF events. Thus, during NPF and subsequent growth events the newly formed particles cannot be assumed to be at the steady-state charge distribution without verification.

3.1.2 Neutral NPF

We next examine how the charge distribution of particles evolves when newly formed particles are electrically neutral.
290 To understand the effect of ion concentration (N_{ion}), the coagulation sink (CoagS), particle growth rate (GR), and nucleation rate (J) on particle charge distribution during NPF events, we calculated the ratio (r_c) of the simulated fraction of singly charged particles to the steady-state value under different NPF conditions. Results at representative conditions are shown in Fig. 3. Some data points are omitted due to the exclusion of exceedingly low particle number concentrations ($<1 \text{ cm}^{-3}$), which occur when both J and particle survival probability (primarily determined by GR/Coag (Kulmala et al., 2017)) are both low, resulting
295 in very few particles surviving to sizes of interest. Additionally, in Fig. 3 we do not distinguish between positive and negative particles since they have the same charge fraction (we have assumed that positive and negative ions have the same concentration and properties and hence the simulation is ‘symmetric’ with respect to particle polarity).




Figure 3. Ratio of simulated singly charged fraction to the steady-state value (r_c) as a function of d_p at different simulation conditions. Different color corresponds to different size ranges. (a) $J=100 \text{ cm}^{-3} \text{ s}^{-1}$, $\text{CoagS} = 0.01 \text{ s}^{-1}$, $\text{GR} = 4 \text{ nm h}^{-1}$ (b) $J=1 \text{ cm}^{-3} \text{ s}^{-1}$, $\text{CoagS} = 0.001 \text{ s}^{-1}$, $\text{GR} = 4 \text{ nm h}^{-1}$ (c) $J=100 \text{ cm}^{-3} \text{ s}^{-1}$, $\text{CoagS} = 0.005 \text{ s}^{-1}$, $N_{\text{ion}} = 200 \text{ cm}^{-3}$ (d) $J=100 \text{ cm}^{-3} \text{ s}^{-1}$, $\text{CoagS} = 0.005 \text{ s}^{-1}$, $N_{\text{ion}} = 1000 \text{ cm}^{-3}$; (e) $J=100 \text{ cm}^{-3} \text{ s}^{-1}$, $\text{GR} = 10 \text{ nm h}^{-1}$, $N_{\text{ion}} = 250 \text{ cm}^{-3}$, (f) $J=100 \text{ cm}^{-3} \text{ s}^{-1}$, $\text{GR} = 4 \text{ nm h}^{-1}$, $N_{\text{ion}} = 250 \text{ cm}^{-3}$, (g) $\text{CoagS} = 0.005 \text{ s}^{-1}$, $\text{GR} = 10 \text{ nm h}^{-1}$, $N_{\text{ion}} = 250 \text{ cm}^{-3}$, (h) $\text{CoagS} = 0.005 \text{ s}^{-1}$, $\text{GR} = 4 \text{ nm/h}$, $N_{\text{ion}} = 250 \text{ cm}^{-3}$.



305 Figures 3a and 3b illustrate the variation of r_c as a function of N_{ion} for selected particle sizes at two conditions typical of polluted ($J=100 \text{ cm}^{-3} \text{ s}^{-1}$, $\text{CoagS} = 0.01 \text{ s}^{-1}$, $\text{GR} = 4 \text{ nm/h}$) and clean environments ($J = 1 \text{ cm}^{-3} \text{ s}^{-1}$, $\text{CoagS} = 0.001 \text{ s}^{-1}$, $\text{GR} = 4 \text{ nm/h}$). Both figures demonstrate that during NPF events, r_c depends on both the particle size and the ion concentration. At a fixed particle size, r_c increases with N_{ion} , which is expected as higher N_{ion} reduces the characteristic charging time (Fig. 2) and promotes the particle charge distribution to reach the steady state. Moreover, larger particles have r_c closer to 1, indicating that as particle grow, their charge fraction gradually approaches the steady state value.

310 Figures 3c and 3d show how GR affects r_c at low ($N_{ion} = 200 \text{ cm}^{-3}$) and high ($N_{ion} = 1000 \text{ cm}^{-3}$) ion concentrations, respectively. As GR increases, r_c for a given particle size decreases, which is due to the decreased charge conditioning time by atmospheric ions (the time for particles to reach size d_p is approximately $\frac{d_p}{GR}$). Similar to Figs. 3a and 3b, smaller particles have lower r_c due to their shorter interaction time with ions and longer characteristic charging time (Fig. 2a). Furthermore, r_c is larger at higher ion concentrations, corroborating the trend shown in Figs. 3a and 3b.

315 The effect of CoagS on r_c is shown in Figs. 3e and 3f for two particle growth rates. At a higher growth rate (10 nm/h, Fig. 3e), r_c remains largely unchanged as CoagS varies from 0.001 to 0.02 s^{-1} . At a lower growth rate (4 nm/h, Fig. 3f), CoagS has a more pronounced effect on r_c , although changes at a given particle size is still smaller than 0.1. Compared with the impact of N_{ion} and GR on r_c , the influence of CoagS on r_c is minor or even negligible.

320 Finally, Figs. 3g and 3h show the influence of nucleation rate J on r_c at two particle growth rates. Similar CoagS, J has an almost negligible effect on the r_c at both fast ($\text{GR} = 10 \text{ nm h}^{-1}$, Fig. 3g) and slow ($\text{GR} = 4 \text{ nm h}^{-1}$, Fig. 3h) particle growth conditions. However, as J increases, there is a slightly decreasing trend of r_c in Fig. 3h. This small but noticeable trend is caused by the increased coagulation between new particles, which elevates the particle growth rate and decreases the time for the particles to reach a certain size.

325 Overall, Figure 3 indicates that the charge distribution of new particles deviates from the steady-state distribution during new particle formation (NPF) events. Among the four factors considered—ion concentration, particle growth rate, coagulation sink, and nucleation rate—the first two exert a strong influence on r_c , while the latter two have a minor impact.

The interaction between aerosol particles and atmospheric ions can be leveraged to measure the particle size distribution (PSD). In this approach, atmospheric ions serve as aerosol neutralizers in the scanning mobility particle sizer (Li et al., 2022; Chen and Jiang, 2018), reducing both the cost and safety risks associated with the instrument. However, a prerequisite for this method is that aerosol particles must reach a steady-state charge distribution at the time of measurement. Our analysis demonstrates that during NPF events, freshly formed neutral particles require tens of minutes to hours to achieve this steady-state distribution. To establish a characteristic size d_b above which the PSD can be measured without a neutralizer, we formulated a regression equation for d_b as a function of GR, N_{ion} , J , and CoagS, defining d_b as the size at which the singly charged fraction of new particles reaches 63% (i.e., $(1 - 1/e)$) of the steady-state value. The functional form of this regression, along with comparisons to simulations, is detailed in Sect. S4 of the SI.

330

335



3.1.3 Initially charged particles




Figure 4. r_c as a function of particle diameter at different IIN conditions: (a) $GR = 4 \text{ nm h}^{-1}$, $CoagS = 0.005 \text{ s}^{-1}$, $J = 100 \text{ cm}^{-3} \text{ s}^{-1}$, (b) $N_{ion} = 250 \text{ cm}^{-3}$, $CoagS = 0.005 \text{ s}^{-1}$, $J = 100 \text{ cm}^{-3} \text{ s}^{-1}$, (c) $N_{ion} = 250 \text{ cm}^{-3}$, $GR = 4 \text{ nm h}^{-1}$, $J = 100 \text{ cm}^{-3} \text{ s}^{-1}$, (d) $N_{ion} = 250 \text{ cm}^{-3}$, $CoagS = 0.005 \text{ s}^{-1}$, $GR = 4 \text{ nm h}^{-1}$. The nucleation rate J is the sum of the formation rates of the positive and negative particles.

To understand the evolution of initially charged particles, we examine a limiting case where all particles are formed via ion-induced nucleation (i.e., $F_{IIN}=1$). We simplify our discussion by assuming equal IIN rates for both polarities. Figure 4 illustrates the behavior of the charge fraction ratio r_c as a function of particle size under selected new particle formation (NPF) conditions. The evolution of r_c can be categorized into three stages, as depicted by the red curve in Figure 4a. In stage 1, r_c rapidly decreases until it reaches unity. In stage 2, r_c drops further to a minimum value $r_{c,min}$. In stage 3, r_c rebounds towards 1. This behavior can be attributed to two main effects: the collisions between particles and atmospheric ions (termed the 'ion effect'), which drives the charge distribution towards $r_c = 1$, and the coagulation of oppositely charged particles, also known as ion-ion recombination, (termed the 'coagulation effect'), which reduces the r_c value. During stage 1, newly formed particles experience both the ion and coagulation effects, leading to a rapid decrease in charge fraction towards the steady-state value of $r_c = 1$. In stage 2, as particles grow, the coagulation effect becomes dominant due to the increased concentration of charged



particles from nucleation, resulting in a further reduction of r_c to below 1. In comparison, r_c remain above 1 if coagulation between particles are turned off in the simulation (Fig. S3). In stage 3, the coagulation effect diminishes for two reasons: (1) the IIN terminates and the generation of charged particles stops; (2) the charged particles already formed are overall more
355 neutralized as they grow. Because of the diminished coagulation effect, the ion effect drives r_c towards 1.

Variation of different simulation parameters alter the r_c curves to different extents. Figure 4a indicates that higher N_{ion} increases $r_{c,min}$ and restores r_c to the steady state values faster than lower N_{ion} . This phenomenon occurs because as N_{ion} increases, the ion effect becomes greater and the coagulation between oppositely charged particles become comparatively less important. Figure 4b shows that the increase of GR causes $r_{c,min}$ to move to the right, but does not significantly change the value of $r_{c,min}$.
360 This relationship between GR and $r_{c,min}$ means that GR does not strongly impact the coagulation effect. Figure 4c shows that CoagS increases $r_{c,min}$ to a smaller extent. Higher CoagS corresponds to larger consumption of the particles and lower particle concentration, hence depressing the coagulation effect. Lastly, Fig. 4d indicates that the larger the J values, the smaller $r_{c,min}$ becomes. As J increases, the coagulation effect becomes appreciably stronger because it is proportional to the particle number concentration squared. The minima $r_{c,min}$ also appear at smaller sizes as the higher particle concentration cause the coagulation
365 between the charged particle to proceed at a faster rate.

In addition to charge state, the interactions between particles and atmospheric ions can also influence particle number concentration and size during NPF events. To quantify such effects, we compared the particle number concentration and mode diameters in simulations with and without considering particle charging (Sect. S5 in the SI). This comparison suggests that particle charging has almost a negligible influence on the mode diameter. However, although the particle number
370 concentrations also remain largely unaffected during neutral NPF, it can experience a considerable decrease during IIN due the strong coagulation between oppositely charged particles.



3.2 Prediction of particle charge fraction with ResNet




Figure 5. (a)-(d) Comparison between the simulated r_c ($r_{c,sim}$), the ResFWD-predicted r_c ($r_{c,ML}$) and the r_c calculated with Eq. (11) ($r_{c,Anal}$) at particle diameters of 2.2 nm, 3nm, 5 nm and 8 nm. The numbers in the subscript of r_c denote the particle size. The R^2 and MSE obtained from testing the ResFWD model against $r_{c,sim}$ are shown in the panels. (e)-(i) Sensitivity of $r_{c,2.2}$ to 10%-10% variations of model input. The color bar indicates the degree of variation quantified by $S_{2.2} = \frac{r_{c,2.2}^{noise} - r_{c,2.2}}{r_{c,2.2}}$. The colored dots are calculated with ResFWD, while grey dots are obtained by changing the CDMS-ion input by either 10% or -10%.

Using simulated results as training data, we developed several ResNet-based regression models, collectively referred to as ResFWD. Each of these models can predict r_c value for a specific particle size. Figures 5a–5d compare the r_c values calculated with ResFWD, the analytical expression (i.e., Eq. (11)) and CDMS-ion for $F_{IIN} < 0.2$ and $J < 10 \text{ cm}^{-3}\text{s}^{-1}$. The scope of F_{IIN} and J are limited to small values because Eq. (11) was developed to cope with situations with a low fraction of charged particles (Kerminen et al., 2007). The r_c values calculated with ResFWD ($r_{c,ML}$) align closely with those simulated by CDMS-ion ($r_{c,sim}$), demonstrating the neural network's ability to capture the nonlinear relationship between r_c and the key parameters including F_{IIN} , GR, N_{ion} , J, and CoagS. In contrast, the values calculated with Eq. (11) ($r_{c,Anal}$) deviate significantly from $r_{c,sim}$, and this discrepancy grows larger with particle size. This suggests that as the particles grow, the simulation conditions deviate farther away from the underlying assumption of Eq. (11). As shown in Figure S7 in the SI, $r_{c,Anal}$ tends to be larger than $r_{c,sim}$ in the entire range of F_{IIN} and J. Such overestimation of r_c by the analytical equation may arise from its inability to account for the strong coagulation between charged particles, especially when a large fraction of the particle population are charged. Another cause for the overestimation could be that Eq. (11) was developed based on the charge state of the smallest particles



rather than F_{IIN} (the former is the ratio of charged particle concentration to the total particle concentration in the smallest size bin, while the latter is a ratio of fluxes). A comparison of these two values is shown in Fig. S8 in the SI.

Figure 5e-i present sensitivity analysis of $r_{c,2.2}$ in response to variations of different model inputs. This analysis is crucial for (1) assessing whether ResFWD overfits the training data and (2) evaluating its susceptibility to input noise—an inevitable factor in field data—compared to the benchmark model CDMS-ion. In these figures, colored dots represent the fractional change in $r_{c,ML,2.2}$ (denoted as $S_{2.2}^{ML}$) when ResFWD inputs are randomly varied between -10% and +10%, while grey dots reflect the fractional change in $r_{c,sim,2.2}$ (denoted as $S_{2.2}^{Sim}$) resulting from variations in CDMS-ion inputs at two extreme values, i.e., +10% and -10%. Figures 5e–5i demonstrate that $S_{2.2}^{Sim}$ envelops $S_{2.2}^{ML}$ (the grey dots put a limit to the colored dots), indicating that ResFWD exhibits a response to input noise similar to that of CDMS-ion. Moreover, both $S_{2.2}^{Sim}$ and $S_{2.2}^{ML}$ display comparable variations as functions of $r_{c,2.2}$.

Fig. 5e shows that $S_{2.2}$ initially increases with $r_{c,2.2}$ and subsequently stabilizes. This behavior suggests that when initial particle charge information is obscured by interactions with atmospheric ions during growth (leading to low $r_{c,2.2}$ values), F_{IIN} has a minimal effect on $r_{c,2.2}$. However, when charge information is preserved during growth (higher $r_{c,2.2}$ values), $r_{c,2.2}$ scales near linearly with F_{IIN} and also vary between -10% to 10%. Conversely, when varying N_{ion} , GR, CoagS and J, $S_{2.2}$ initially increases with $r_{c,2.2}$ and then decreases (Figs. 5f–5i). This indicates that $r_{c,2.2}$ is relatively insensitive to variations of these parameters when particle interactions with atmospheric ions are either highly effective (resulting in low $r_{c,2.2}$ values) or ineffective (leading to high $r_{c,2.2}$ values). Comparisons between Figures 5e–5i further reveal that $r_{c,2.2}$ is sensitive to variations in GR, N_{ion} , and J, but not to CoagS: a 10% variation in CoagS results in less than an 8% change in $r_{c,2.2}$. This finding aligns with Figure 4, which indicates that simulations with differing GR, N_{ion} , and J values yield well-separated curves, while varying CoagS values only change the curves slightly.

Despite good agreement with the benchmark model, the applicability of the ResFWD is limited by the data used for its training. In this study, we did not consider scenarios where the mobilities and concentrations of atmospheric positive and negative ions differ, restricting the direct application of ResFWD in these cases. The applicability of ResFWD can be further expanded by training the neural network with a larger dataset that includes these variations.


3.3 ResNet assisted inference of F_{IIN}

During field measurements of atmospheric NPF, the charge fraction and its ratio to the steady-state charge fraction (r_c) can be measured across different particle sizes (Leppä et al., 2013; Iida et al., 2006). To infer F_{IIN} from these measurements, the traditional approach involves identifying the optimal F_{IIN} value that best fits Eq. (11) to the measured r_c . In this study, we utilize simulated r_c values at 2.2 nm, 3 nm, 4 nm, 5 nm, 6nm, 7nm and 8 nm as inputs to directly infer F_{IIN} using ResBWD. Alongside particle charge fractions, additional inputs to the ResNet model include GR, N_{ion} , J, and CoagS (Fig. 1d).

In some of the CDMS-ion simulations, interactions with atmospheric ions may obscure initial charge information as new particles grow to the smallest measurable size (assumed to be 2.2 nm in this work). To address this, we applied a screening criterion for the data used for training ResBWD: for all sets of GR, N_{ion} , J, and CoagS combinations, we only keep the



425 combinations where the change of F_{IIN} by 1% leads to particle charge fraction at 2.2 nm to vary by at least χ . Below we illustrate the impact of this screening criterion for $\chi = 0.11\%$ and $\chi = 0.55\%$, corresponding to 10% and 50% of the *state steady* charge fraction of 2.2 nm particles (which is 1.12%).



430 **Figure 6.** (a-b) Comparison between the predicted F_{IIN} by ResBWD and the true F_{IIN} used in CDMS-ion simulation. The upper and lower panels correspond to χ values of 0.11% and 0.55%, respectively. (c-h) Sensitivity of F_{IIN} to noises of inputs including $r_{c,2.2}$, $r_{c,5}$, and $r_{c,8}$. The sensitivity is defined as $S_x = \frac{F_{\text{IIN}}^{\text{noise}} - F_{\text{IIN}}}{F_{\text{IIN}}}$, with the subscript x denoting the size at which r_c is varied. The green reference lines indicate S_x values of ± 0.2 .

435 Figures 6a and 6b compare ResBWD predicted and the true F_{IIN} for the two χ values, demonstrating good agreement regardless of the χ employed. This indicates that ResBWD effectively captures the nonlinear relationship between the charge state of grown particles and F_{IIN} , even when initial charge information is largely lost (Fig. 6a, $\chi = 0.11\%$). However, further sensitivity tests (Figs. 6c-h) reveal that noise in input parameters to ResBWD (i.e., random noises of $r_{c,2.2}$, $r_{c,5}$, $r_{c,8}$ within -10% to +10%) results in F_{IIN} variations primarily ranging from -10% to +10% for $\chi = 0.55\%$ (lower panels), whereas this variation increases to -20% to +20% for $\chi = 0.11\%$ (upper panels). This suggests that as initial particle charge information is more obscured due to stronger particle interactions with atmospheric ions, the deduction of F_{IIN} from measured charge fractions
440 becomes increasingly uncertain. At low F_{IIN} values (~ 0.01), high sensitivity for both $\chi = 0.11\%$ and $\chi = 0.55\%$ is observed in Figs. 6c-h. This is as expected since the screening criterion ensures the training data has an F_{IIN} resolution on the order of 1%, hence at low F_{IIN} values (close to 1%) ResBWD is more sensitive to noises. Further comparisons of panels d, f and h (or panels



c, e and g) indicate that $r_{c,2.2}$ is a more critical parameter for F_{IIN} inference than $r_{c,5}$ and $r_{c,8}$, as it retains the most information about F_{IIN} .

445 Overall, predicting F_{IIN} from known r_c values necessitates more stringent conditions than the reverse process. This challenge stems from the loss of initial charge information as particles increase in size. To find parameter sets of GR, N_{ion} , J, and CoagS which meet the screening criteria, the ResFWD model can be employed to calculate χ .

4. Conclusions

In this study, we developed a two-dimensional sectional model, CDMS-ion, to simulate particle growth and their interactions with atmospheric ions. Using this model, we first explored the general characteristics of particle charge state evolution. Our findings reveal that particle growth rate and ion concentration have the most significant effects on particle charge. Notably, when the number concentration of charged particles is high, the ratio of the particle charge fraction to the steady-state value can drop substantially below 1 due to coagulation between oppositely charged particles. Furthermore, atmospheric new particles cannot be treated as if they are at steady state charge distribution until they grow to a certain size
455 (Eq. (S13) in the SI).

Using the extensive dataset generated by CDMS-ion, we trained two types of neural network models. The first model, ResFWD, predicts the particle charge state as they grow, under the assumption that the fraction of ion-induced nucleation is known. This model effectively captures the non-linearity of the particle charging process and shows good agreement with model simulations. Compared to existing analytical equations, ResFWD demonstrates improved accuracy, particularly at high
460 F_{IIN} and J conditions. Therefore, this approach can serve as a reliable alternative to the analytical equation when the assumptions inherent in the training data are met.

The second model, ResBWD, predicts the fraction of ion-induced nucleation using particle charge fractions measured at several sizes. This prediction is more challenging compared to ResFWD because the initial charge information of new particles may be lost as they grow. However, by restricting the application of the model to cases where initial charge information is relatively well preserved (this screening can be completed with ResFWD), we can achieve accurate predictions of F_{IIN} with
465 reasonable sensitivity to input noise.

This work represents a pioneering effort to describe the dynamic charging process of atmospheric new particles with machine learning tools. With these tools, one can calculate the charge state of the new particles as they growth, or use observed particle charge state to deduce the rates of ion induced nucleation, which is a major particle formation mechanism on the global
470 scale. Note that our simplifications of the NPF processes include constant nucleation rates, constant atmospheric ion concentrations, and equal ion concentrations and mobilities. Future endeavors to develop more comprehensive ML models should take these complexities into account.

Code and data availability: The simulation data related to the figures are available from the corresponding author upon
475 request.



Author Contribution: C.L. proposed this study. P.W. and C.L. wrote the simulation program and performed the simulation. All authors participated in the discussion of the results and contributed to writing the paper.

Competing Interests: I declare that I or my co-authors have competing interests as follows: Some authors are members of the editorial board of Atmospheric Chemistry and Physics.

480 **Acknowledgments:** This research has been supported by the National Key R&D Program of China (grant no. 2022YFC3704100), the National Natural Science Foundation of China (grant no. 22206120), the State Key Joint Laboratory of Environmental Simulation and Pollution Control, Samsung PM2.5 SRP, and the ACCC Flagship funded by the Academy of Finland (grant no. 337549).

References

- 485 Chahl, H. S. and Gopalakrishnan, R.: High potential, near free molecular regime Coulombic collisions in aerosols and dusty plasmas, *Aerosol Sci. Technol.*, 53, 933-957, 10.1080/02786826.2019.1614522, 2019.
- Chen, X. and Jiang, J.: Retrieving the ion mobility ratio and aerosol charge fractions for a neutralizer in real-world applications, *Aerosol Sci. Technol.*, 52, 1145-1155, 10.1080/02786826.2018.1498587, 2018.
- 490 Chu, B., Kerminen, V. M., Bianchi, F., Yan, C., Petäjä, T., and Kulmala, M.: Atmospheric new particle formation in China, *Atmos. Chem. Phys.*, 19, 115-138, 10.5194/acp-19-115-2019, 2019.
- Eisenbud, M. and Gesell, T. F.: *Environmental radioactivity from natural, industrial and military sources: from natural, industrial and military sources*, Elsevier 1997.
- Gagné, S., Leppä, J., Petäjä, T., McGrath, M. J., Vana, M., Kerminen, V. M., Laakso, L., and Kulmala, M.: Aerosol charging state at an urban site: new analytical approach and implications for ion-induced nucleation, *Atmos. Chem. Phys.*, 12, 4647-4666, 10.5194/acp-12-4647-2012, 2012.
- 495 Gatti, M. and Kortshagen, U.: Analytical model of particle charging in plasmas over a wide range of collisionality, *Physical Review E*, 78, 046402, 10.1103/PhysRevE.78.046402, 2008.
- Gautam, A. S., Siingh, D., and Kamra, A. K.: Statistical analysis of the atmospheric ion concentrations and mobility distributions at a tropical station, Pune, *Quarterly Journal of the Royal Meteorological Society*, 143, 2116-2128, 500 <https://doi.org/10.1002/qj.3071>, 2017.
- Golubenko, K., Rozanov, E., Mironova, I., Karagodin, A., and Usoskin, I.: Natural Sources of Ionization and Their Impact on Atmospheric Electricity, *Geophys. Res. Lett.*, 47, e2020GL088619, <https://doi.org/10.1029/2020GL088619>, 2020.
- Gopalakrishnan, R. and Hogan, C. J.: Determination of the Transition Regime Collision Kernel from Mean First Passage Times, *Aerosol Sci. Technol.*, 45, 1499-1509, 10.1080/02786826.2011.601775, 2011.
- 505 Gopalakrishnan, R. and Hogan, C. J.: Coulomb-influenced collisions in aerosols and dusty plasmas, *Physical Review E*, 85, 026410, 10.1103/PhysRevE.85.026410, 2012.
- Gordon, H., Kirkby, J., Baltensperger, U., Bianchi, F., Breitenlechner, M., Curtius, J., Dias, A., Dommen, J., Donahue, N. M., Dunne, E. M., Duplissy, J., Ehrhart, S., Flagan, R. C., Frege, C., Fuchs, C., Hansel, A., Hoyle, C. R., Kulmala, M., Kürten, A., Lehtipalo, K., Makhmutov, V., Molteni, U., Rissanen, M. P., Stozkhov, Y., Tröstl, J., Tsagkogeorgas, G., Wagner, R., 510 Williamson, C., Wimmer, D., Winkler, P. M., Yan, C., and Carslaw, K. S.: Causes and importance of new particle formation in the present-day and preindustrial atmospheres, *J. Geophys. Res. Atmos.*, 122, 8739-8760, 10.1002/2017JD026844, 2017.
- He, K., Zhang, X., Ren, S., and Sun, J.: Deep residual learning for image recognition, *Proceedings of the IEEE conference on computer vision and pattern recognition*, 770-778,
- Hirsikko, A., Nieminen, T., Gagné, S., Lehtipalo, K., Manninen, H. E., Ehn, M., Hörrak, U., Kerminen, V. M., Laakso, L., 515 McMurry, P. H., Mirme, A., Mirme, S., Petäjä, T., Tammet, H., Vakkari, V., Vana, M., and Kulmala, M.: Atmospheric ions and nucleation: a review of observations, *Atmos. Chem. Phys.*, 11, 767-798, 10.5194/acp-11-767-2011, 2011.
- Iida, K., Stolzenburg, M., McMurry, P., Dunn, M. J., Smith, J. N., Eisele, F., and Keady, P.: Contribution of ion-induced nucleation to new particle formation: Methodology and its application to atmospheric observations in Boulder, Colorado, *J. Geophys. Res. Atmos.*, 111, <https://doi.org/10.1029/2006JD007167>, 2006.



- 520 Jiang, S., Liu, Y.-R., Huang, T., Feng, Y.-J., Wang, C.-Y., Wang, Z.-Q., Ge, B.-J., Liu, Q.-S., Guang, W.-R., and Huang, W.: Towards fully ab initio simulation of atmospheric aerosol nucleation, *Nat. Commun.*, 13, 6067, 10.1038/s41467-022-33783-y, 2022.
- Joutsensaari, J., Ozon, M., Nieminen, T., Mikkonen, S., Lähivaara, T., Decesari, S., Facchini, M. C., Laaksonen, A., and Lehtinen, K. E.: Identification of new particle formation events with deep learning, *Atmos. Chem. Phys.*, 18, 9597-9615, 2018.
- 525 Kelp, M. M., Tessum, C. W., and Marshall, J. D.: Orders-of-magnitude speedup in atmospheric chemistry modeling through neural network-based emulation, arXiv preprint arXiv:1808.03874, 2018.
- Kerminen, V.-M., Chen, X., Vakkari, V., Petäjä, T., Kulmala, M., and Bianchi, F.: Atmospheric new particle formation and growth: review of field observations, *Environ. Res. Lett.*, 13, 103003, 10.1088/1748-9326/aadf3c, 2018.
- 530 Kerminen, V.-M., Anttila, T., Petäjä, T., Laakso, L., Gagné, S., Lehtinen, K. E. J., and Kulmala, M.: Charging state of the atmospheric nucleation mode: Implications for separating neutral and ion-induced nucleation, *J. Geophys. Res. Atmos.*, 112, <https://doi.org/10.1029/2007JD008649>, 2007.
- Kirkby, J., Duplissy, J., Sengupta, K., Frege, C., Gordon, H., Williamson, C., Heinritzi, M., Simon, M., Yan, C., Almeida, J., Tröstl, J., Nieminen, T., Ortega, I. K., Wagner, R., Adamov, A., Amorim, A., Bernhammer, A.-K., Bianchi, F., Breitenlechner, M., Brilke, S., Chen, X., Craven, J., Dias, A., Ehrhart, S., Flagan, R. C., Franchin, A., Fuchs, C., Guida, R., Hakala, J., Hoyle, C. R., Jokinen, T., Junninen, H., Kangasluoma, J., Kim, J., Krapf, M., Kürten, A., Laaksonen, A., Lehtipalo, K., Makhmutov, V., Mathot, S., Molteni, U., Onnela, A., Peräkylä, O., Piel, F., Petäjä, T., Praplan, A. P., Pringle, K., Rap, A., Richards, N. A. D., Riipinen, I., Rissanen, M. P., Rondo, L., Sarnela, N., Schobesberger, S., Scott, C. E., Seinfeld, J. H., Sipilä, M., Steiner, G., Stozhkov, Y., Stratmann, F., Tomé, A., Virtanen, A., Vogel, A. L., Wagner, A. C., Wagner, P. E., Weingartner, E., Wimmer, D., Winkler, P. M., Ye, P., Zhang, X., Hansel, A., Dommen, J., Donahue, N. M., Worsnop, D. R., Baltensperger, U., Kulmala, M., Carslaw, K. S., and Curtius, J.: Ion-induced nucleation of pure biogenic particles, *Nature*, 533, 521-526, 10.1038/nature17953, 2016.
- 540 Kubečka, J., Neeffjes, I., Besel, V., Qiao, F., Xie, H.-B., and Elm, J.: Atmospheric Sulfuric Acid–Multi-Base New Particle Formation Revealed through Quantum Chemistry Enhanced by Machine Learning, *J. Phys. Chem. A*, 127, 2091-2103, 10.1021/acs.jpca.3c00068, 2023.
- 545 Kulmala, M., Kerminen, V. M., Petäjä, T., Ding, A. J., and Wang, L.: Atmospheric gas-to-particle conversion: why NPF events are observed in megacities?, *Faraday Discuss.*, 200, 271-288, 10.1039/C6FD00257A, 2017.
- Laakso, L., Gagné, S., Petäjä, T., Hirsikko, A., Aalto, P. P., Kulmala, M., and Kerminen, V. M.: Detecting charging state of ultra-fine particles: instrumental development and ambient measurements, *Atmos. Chem. Phys.*, 7, 1333-1345, 10.5194/acp-7-1333-2007, 2007.
- 550 Lee, S. H., Reeves, J. M., Wilson, J. C., Hunton, D. E., Viggiano, A. A., Miller, T. M., Ballenthin, J. O., and Lait, L. R.: Particle Formation by Ion Nucleation in the Upper Troposphere and Lower Stratosphere, *Science*, 301, 1886-1889, 10.1126/science.1087236, 2003.
- Leppä, J., Anttila, T., Kerminen, V. M., Kulmala, M., and Lehtinen, K. E. J.: Atmospheric new particle formation: real and apparent growth of neutral and charged particles, *Atmos. Chem. Phys.*, 11, 4939-4955, 10.5194/acp-11-4939-2011, 2011.
- 555 Leppä, J., Gagné, S., Laakso, L., Manninen, H. E., Lehtinen, K. E. J., Kulmala, M., and Kerminen, V. M.: Using measurements of the aerosol charging state in determination of the particle growth rate and the proportion of ion-induced nucleation, *Atmos. Chem. Phys.*, 13, 463-486, 10.5194/acp-13-463-2013, 2013.
- Leppä, J., Kerminen, V.-M., Laakso, L., Korhonen, H., Lehtinen, K. E., Gagné, S., Manninen, H. E., Nieminen, T., and Kulmala, M.: Ion-UHMA: a model for simulating the dynamics of neutral and charged aerosol particles, *Boreal Environment Research*, 14, 559, 2009.
- 560 Li, C. and McMurry, P. H.: Errors in nanoparticle growth rates inferred from measurements in chemically reacting aerosol systems, *Atmos. Chem. Phys.*, 18, 8979-8993, 10.5194/acp-18-8979-2018, 2018.
- Li, C., Li, Y., Li, X., Cai, R., Fan, Y., Qiao, X., Yin, R., Yan, C., Guo, Y., Liu, Y., Zheng, J., Kerminen, V. M., Kulmala, M., Xiao, H., and Jiang, J.: Comprehensive simulations of new particle formation events in Beijing with a cluster dynamics–multicomponent sectional model, *Atmos. Chem. Phys.*, 23, 6879-6896, 10.5194/acp-23-6879-2023, 2023.
- 565 Li, Y., Chen, X., and Jiang, J.: Measuring size distributions of atmospheric aerosols using natural air ions, *Aerosol Sci. Technol.*, 56, 655-664, 10.1080/02786826.2022.2060795, 2022.



- 570 Liu, C., Zhang, H., Cheng, Z., Shen, J., Zhao, J., Wang, Y., Wang, S., and Cheng, Y.: Emulation of an atmospheric gas-phase chemistry solver through deep learning: Case study of Chinese Mainland, *Atmospheric Pollution Research*, 12, 101079, <https://doi.org/10.1016/j.apr.2021.101079>, 2021.
- López-Yglesias, X. and Flagan, R. C.: Ion–Aerosol Flux Coefficients and the Steady-State Charge Distribution of Aerosols in a Bipolar Ion Environment, *Aerosol Sci. Technol.*, 47, 688–704, [10.1080/02786826.2013.783684](https://doi.org/10.1080/02786826.2013.783684), 2013.
- Matsui, H.: Development of a global aerosol model using a two-dimensional sectional method: 1. Model design, *Journal of Advances in Modeling Earth Systems*, 9, 1921–1947, <https://doi.org/10.1002/2017MS000936>, 2017.
- 575 Matsui, H. and Mahowald, N.: Development of a global aerosol model using a two-dimensional sectional method: 2. Evaluation and sensitivity simulations, *Journal of Advances in Modeling Earth Systems*, 9, 1887–1920, [10.1002/2017MS000937](https://doi.org/10.1002/2017MS000937), 2017.
- Nadykto, A. B. and Yu, F.: Uptake of neutral polar vapor molecules by charged clusters/particles: Enhancement due to dipole-charge interaction, *J. Geophys. Res. Atmos.*, 108, <https://doi.org/10.1029/2003JD003664>, 2003.
- 580 Ouyang, H., Gopalakrishnan, R., and Hogan, C. J.: Nanoparticle collisions in the gas phase in the presence of singular contact potentials, *J. Chem. Phys.*, 137, 064316, [10.1063/1.4742064](https://doi.org/10.1063/1.4742064), 2012.
- Pfeifer, J., Mahfouz, N. G. A., Schulze, B. C., Mathot, S., Stolzenburg, D., Baalbaki, R., Brasseur, Z., Caudillo, L., Dada, L., Granzin, M., He, X. C., Lamkaddam, H., Lopez, B., Makhmutov, V., Marten, R., Mentler, B., Müller, T., Onnela, A., Philippov, M., Piedehierro, A. A., Rörup, B., Schervish, M., Tian, P., Umo, N. S., Wang, D. S., Wang, M., Weber, S. K., Welti, A., Wu, Y., Zauner-Wieczorek, M., Amorim, A., El Haddad, I., Kulmala, M., Lehtipalo, K., Petäjä, T., Tomé, A., Mirme, S., Manninen, H. E., Donahue, N. M., Flagan, R. C., Kürten, A., Curtius, J., and Kirkby, J.: Measurement of the collision rate coefficients between atmospheric ions and multiply charged aerosol particles in the CERN CLOUD chamber, *Atmos. Chem. Phys.*, 23, 6703–6718, [10.5194/acp-23-6703-2023](https://doi.org/10.5194/acp-23-6703-2023), 2023.
- 585 Simmel, M. and Wurzler, S.: Condensation and activation in sectional cloud microphysical models, *Atmospheric Research*, 80, 218–236, <https://doi.org/10.1016/j.atmosres.2005.08.002>, 2006.
- Stolzenburg, M. R., McMurry, P. H., Sakurai, H., Smith, J. N., Mauldin, R. L., Eisele, F. L., and Clement, C. F.: Growth rates of freshly nucleated atmospheric particles in Atlanta, *J. Geophys. Res. Atmos.*, 110, n/a–n/a, [10.1029/2005JD005935](https://doi.org/10.1029/2005JD005935), 2005.
- Stozhkov, Y. I.: The role of cosmic rays in the atmospheric processes, *J. Phys. G: Nucl. Part. Phys.*, 29, 913, [10.1088/0954-3899/29/5/312](https://doi.org/10.1088/0954-3899/29/5/312), 2003.
- 595 Su, P., Joutsensaari, J., Dada, L., Zaidan, M. A., Nieminen, T., Li, X., Wu, Y., Decesari, S., Tarkoma, S., and Petäjä, T.: New particle formation event detection with Mask R-CNN, *Atmos. Chem. Phys.*, 22, 1293–1309, 2022.
- Svensmark, H., Enghoff, M. B., Shaviv, N. J., and Svensmark, J.: Increased ionization supports growth of aerosols into cloud condensation nuclei, *Nat. Commun.*, 8, 2199, [10.1038/s41467-017-02082-2](https://doi.org/10.1038/s41467-017-02082-2), 2017.
- Wiedensohler, A.: An approximation of the bipolar charge distribution for particles in the submicron size range, *J. Aerosol Sci.*, 19, 387–389, [https://doi.org/10.1016/0021-8502\(88\)90278-9](https://doi.org/10.1016/0021-8502(88)90278-9), 1988.
- 600 Yin, R., Li, X., Yan, C., Cai, R., Zhou, Y., Kangasluoma, J., Sarnela, N., Lampilahti, J., Petäjä, T., Kerminen, V. M., Bianchi, F., Kulmala, M., and Jiang, J.: Revealing the sources and sinks of negative cluster ions in an urban environment through quantitative analysis, *Atmos. Chem. Phys.*, 23, 5279–5296, [10.5194/acp-23-5279-2023](https://doi.org/10.5194/acp-23-5279-2023), 2023.
- Yu, F., Nadykto, A. B., Luo, G., and Herb, J.: H₂SO₄–H₂O binary and H₂SO₄–H₂O–NH₃ ternary homogeneous and ion-mediated nucleation: lookup tables version 1.0 for 3-D modeling application, *Geosci. Model Dev.*, 13, 2663–2670, [10.5194/gmd-13-2663-2020](https://doi.org/10.5194/gmd-13-2663-2020), 2020.
- 605 Yu, F., Wang, Z., Luo, G., and Turco, R.: Ion-mediated nucleation as an important global source of tropospheric aerosols, *Atmos. Chem. Phys.*, 8, 2537–2554, [10.5194/acp-8-2537-2008](https://doi.org/10.5194/acp-8-2537-2008), 2008.
- Yu, F., Luo, G., Liu, X., Easter, R. C., Ma, X., and Ghan, S. J.: Indirect radiative forcing by ion-mediated nucleation of aerosol, *Atmos. Chem. Phys.*, 12, 11451–11463, [10.5194/acp-12-11451-2012](https://doi.org/10.5194/acp-12-11451-2012), 2012.
- Zaveri, R. A., Easter, R. C., Fast, J. D., and Peters, L. K.: Model for Simulating Aerosol Interactions and Chemistry (MOSAIC), *J. Geophys. Res. Atmos.*, 113, <https://doi.org/10.1029/2007JD008782>, 2008.
- Zhao, B., Donahue, N. M., Zhang, K., Mao, L., Shrivastava, M., Ma, P.-L., Shen, J., Wang, S., Sun, J., Gordon, H., Tang, S., Fast, J., Wang, M., Gao, Y., Yan, C., Singh, B., Li, Z., Huang, L., Lou, S., Lin, G., Wang, H., Jiang, J., Ding, A., Nie, W., Qi, X., Chi, X., and Wang, L.: Global variability in atmospheric new particle formation mechanisms, *Nature*, [10.1038/s41586-024-07547-1](https://doi.org/10.1038/s41586-024-07547-1), 2024.