

Transformation Processes in the Oder Lagoon as seen from a Model Perspective

Thomas Neumann¹, Gerald Schernewski¹, and René Friedland¹

¹Leibniz Institute for Baltic Sea Research Warnemünde, 18119 Rostock, Germany

Correspondence: Thomas Neumann (thomas.neumann@io-warnemuende.de)

Abstract. The Oder/Szczecin Lagoon is one of the largest Baltic Sea lagoons and is subject to very high nutrient loads from the Oder/Odra River. For our study, we employ a modified, high-resolution 3D ecosystem model specifically adapted for this shallow lagoon. The model demonstrates stable and reliable performance over 25 years of simulation, enabling a detailed assessment of lagoon processes under various scenarios. Our model simulations indicate that changes in riverine nutrient inputs have an immediate impact on the lagoon's water quality, affecting parameters such as phytoplankton biomass and water transparency.

Hypoxia is a prevalent phenomenon, affecting most parts of the lagoon which promotes internal eutrophication. On average, the lagoon retains 12% (253 t/year) of the phosphorus and 40% (17,278 t/year) of the nitrogen riverine inputs. The primary sink process for phosphorus is sediment burial, and for nitrogen, it is denitrification. Nitrogen retention decreases with increasing riverine loads, dropping to around 30% during years with exceptionally high inputs. The nutrient retention capacity of the lagoon has significant implications for Baltic Sea eutrophication but is not currently accounted for in major policies and Baltic Sea models.

Although recent nutrient loads from the Oder River comply with policy targets, such as the Baltic Sea Action Plan's maximum allowable inputs and Germany's river targets, these levels are insufficient to improve the lagoon's ecological state sufficiently. The Oder Lagoon remains in a highly eutrophic condition, making the achievement of a good ecological status unlikely.

1 Introduction

Marine ecosystems around the world suffer from increasing oxygen deficiencies (Diaz and Rosenberg, 2008). Especially coastal zones are affected by permanent or seasonal low oxic conditions (Fennel and Testa, 2019; Conley et al., 2011; Carstensen et al., 2014). The drivers of deoxygenation are land and airborne nutrient depositions and increased temperature (Kabel et al., 2012; Börgel et al., 2023). Coastal features such as lagoons and bays can significantly reduce terrestrial nutrient loads. This is due to their long residence time and shallow waters. Shallow waters facilitate a close interaction between sedimentary processes and those in the euphotic zone, thereby speeding up biogeochemical cycles (e.g. Asmala et al., 2017).

The Baltic Sea includes a diverse array of coastal waters, bays, lagoons, and estuaries, each with unique features and behaviors. These waters host specialized flora and fauna and act as transformers and retention units for external nutrient loads.

Asmala et al. (2017) classified Baltic Sea coastal ecosystems into different categories: lagoons, estuaries, embayments, open coast, and archipelago. For nitrogen removal (denitrification), lagoons have the highest rate, followed by estuaries. Within the lagoons, the Oder Lagoon is the region with the highest denitrification rate. In the case of phosphorus retention (burial in their study), archipelagos show the highest potential, which is driven by the import of phosphorus from the open sea.

30 Systems with prolonged water residence times and high nutrient loads are particularly important for Baltic Sea pollution. Only a few coastal waters can significantly alter nutrient loads transported from rivers to the Baltic Sea. Notable examples include the Curonian and Vistula Lagoons, the Gulf of Riga, coastal waters near St. Petersburg, and some Scandinavian estuaries. Among these, the Oder Lagoon is likely the most critical system in terms of quantitative nutrient retention and transformation (Vybernaite-Lubiene et al., 2022; Müller-Karulis and Aigars, 2011; Almroth-Rosell et al., 2016; Edman et al., 2018; Kuliński
35 et al., 2022).

Asmala et al. (2017) studied the coastal filter for the entire Baltic Sea by observations of denitrification and sediment cores. Their results suggest a removal of 16% nitrogen and 53% of phosphorus from the land based loads. Swedish coastal waters exhibit a relatively high nutrient retention capability. Edman et al. (2018) reported a mean phosphorus retention of 69% and a mean nitrogen retention of 53% estimated from model simulations. These high values, particularly for phosphorus, are due to
40 the oxic water conditions, which favor the trapping of phosphorus in the sediment.

The shallow Oder Lagoon, located at the German/Polish border in the southern Baltic Sea, is one of the largest lagoons in Europe. With an average water discharge of about 500 m³/s and a drainage area of about 120,000 km², the Oder River is one of the most important rivers in the Baltic Sea catchment. Due to similarities with other lagoons, it can be expected that insights into retention and transformation processes can be related to driving factors, and simplified relationships can be transferred
45 to other systems. By studying the most important coastal water systems in detail, load calculations to the Baltic Sea can be systematically improved, enhancing regional ecosystem state and water quality assessments.

The European Water Framework Directive (WFD) is the primary legislation aimed at achieving good ecological status in European coastal and transitional waters. Under the WFD, these systems are classified as distinct water bodies, which undergo regular monitoring. Consequently, long-term data on Baltic coastal waters are available, with some records dating
50 back to the 1970s for most countries. However, these data are typically collected fortnightly from a single station intended to represent the entire system. While this is adequate for assessing averaged overall states and long-term changes, the limited spatial and temporal resolution hinders the analysis and understanding of major processes within these systems and their annual dynamics. Increasing the frequency of sampling and the number of locations would raise costs beyond feasible levels. Therefore, combining 3D ecosystem models with field data is necessary.

55 Additionally, the Baltic Sea Action Plan (BSAP) (HELCOM, 2021a) establishes the maximum allowable inputs (MAI) for nitrogen and phosphorus required to achieve a good environmental status (Schernewski et al., 2015) in the Baltic Sea. These inputs describe loads to the open Baltic Sea and do not consider possible modifications within the coastal filter, particularly lagoons where large rivers enter.

Recent high-resolution 3D ecosystem models of the Baltic Sea, such as presented by Piehl et al. (2023), effectively and
60 sufficiently describe processes for practical purposes like policy implementation and state assessments. However, these models

fall short in semi-enclosed, enclosed, and several transitional waters due to inadequate spatial resolution. Currently, neither data nor models adequately account for retention and transformation processes in these systems, leading to overestimated nutrient loads to the Baltic Sea.

These deficiencies have been compensated for, for example, by making assumptions about the bioavailability of riverine nutrients. Eilola et al. (2011) state that the three considered models differ in their assumptions of bioavailability and discuss uncertainties in these assumptions. Ruvalcaba Baroni et al. (2024) use reduction factors for nutrient inputs in their model study because nutrient removal for different coastal types is poorly quantified. They also argue that river-specific organic matter retention factors in coastal waters would improve the input of organic matter from rivers.

The quantitatively unknown nutrient retention in coastal waters could potentially result in inaccurate policy settings, such as maximum allowable inputs or water quality thresholds and targets. Our objectives are to:

- (a) set up a spatially high resolved ecosystem model for the Oder Lagoon;
- (b) apply and validate the 3D ecosystem model in the Oder Lagoon using long-term data;
- (c) quantify the retention of nitrogen and phosphorus and their inter-annual variability in the lagoon;
- (d) analyze the driving parameters; and
- (e) assess the consequences for policy implementation, namely water quality thresholds, acceptable riverine loads, as well as lagoon and Baltic Sea management.

2 Model Setup

We employ a numerical modeling technique to evaluate the Oder Lagoon ecosystem. This model integrates both biogeochemical and circulation components. The circulation aspect utilizes the Modular Ocean Model MOM5.1 (Griffies, 2004) coupled with a sea ice model (Winton, 2000), while the biogeochemical part is based on ERGOM version 1.2 (Leibniz Institute for Baltic Sea Research, 2015). A detailed explanation and validation of the model can be found in (Neumann et al., 2022). For this study, the model was specifically set up for the Oder Lagoon area within the Baltic Sea. Figure 1 illustrates the bathymetry of the model. A notable aspect is the navigation channel, which has a depth of 10 m. Modifications to the Świna River began in 1721, and in 1880, a shortened and deepened artificial channel was completed. Subsequent dredging projects increased the depth to 9.6 meters in 1939 and to 10.5 meters in 1984 across the entire lagoon. Most recently, between 2018 and 2023, the entire waterway through the Oder Lagoon was deepened to 12.5 meters (Schernewski et al., 2025). Beyond the channel, the lagoon remains relatively shallow with depths predominantly under 5 m. The horizontal grid resolution is 150 m. Vertically, the model is divided into 28 layers, starting with a layer thickness of 25 cm at the top and 50 cm at the bottom. Three open boundary conditions (OBC), Peene, Swina, and Dziwna in Fig. 1, link the model to the Baltic Sea. Data from a coarser Baltic Sea model with a 2 km resolution (Piehl et al., 2023) are applied at the OBC locations. The Oder River enters into the lagoon from the southern edge.

The ERGOM model describes cycles of nitrogen, phosphorus, carbon, oxygen, and partially sulfur. Primary production is driven by photosynthetically active radiation, facilitated by four functional groups of phytoplankton (large cells, small cells,




Figure 1. Bathymetric map of the Oder Lagoon model (a). The green dots indicate two stations used for validation. Peene, Świna, and Dziwna mark the locations of the open boundary conditions, and Oder, Uecker, and Zarow Rivers represent the river mouths. The red rectangle in (b) shows the location of the Oder Lagoon within the Baltic Sea. The map was created using the software package GrADS 2.1.1.b0 (<http://cola.gmu.edu/grads/>, last access: 28 November 2024), using published bathymetry data (Seifert et al., 2008).

limnic phytoplankton, and cyanobacteria). The optical sub-model estimates the light climate based on chlorophyll and CDOM (Colored Dissolved Organic Matter) concentrations (Neumann et al., 2021). Dead organic matter accumulates in the detritus state variable. Bulk zooplankton grazes on phytoplankton and represents the highest trophic level considered in the model. Particulate organic matter (POC: phytoplankton, detritus, and other POC species) have the capability to sink into the water column and accumulate within a sediment layer. Detritus undergoes mineralization into dissolved inorganic nitrogen and phosphorus both in the water column and in the sediment. The mineralization process is influenced by water temperature and oxygen concentration. In oxygen-rich conditions, phosphate becomes bound to iron oxide and is subsequently retained as particles within the sediments. Erosion resuspends these particles, which are then transported by currents to deposition areas. In anoxic conditions, iron oxide is reduced, releasing phosphate into the water as dissolved phosphate (Neumann and Schernewski, 2008). Oxygen is produced through primary production and consumed by processes such as metabolism and mineralization. Furthermore, the extracellular excretion of dissolved organic matter by phytoplankton results in non-Redfield carbon uptake.

In addition to the model proposed by Neumann et al. (2022), we have introduced a fourth phytoplankton group (limnic phytoplankton). This new group is specifically designed for low salinity and turbid coastal waters, realizing growth limitations due to high salinity levels as well as increased light sensitivity.

All organic particles (phytoplankton, detritus, etc.) are counted in nitrogen units. To compare the model phytoplankton with chlorophyll observations, we sum up all phytoplankton groups and multiply them by a constant chlorophyll-to-carbon ratio.

At the open boundaries, the model is forced by data from a coarse grained model as noted above. Meteorological forcing data are from the coastDat-3 dataset (Geyer and Rockel, 2013). It is the same data set forcing the coarse grained model. Nutrient loads and runoff into the lagoon from rivers Oder, Uecker, and Zarow were provided by Polish and German national agencies




Figure 2. Riverine forcing of the model. Upper panel: Annual mean runoff. Lower panel: Annual mean loads for nitrogen (black) and phosphorus (red).

(see *code and data availability*). The riverine forcing is shown in Fig. 2. Loads are correlated with runoff, that is, the interannual runoff variability predominantly controls the load’s variability.

115 We start the model simulations in 1995 initialized with data from Piehl et al. (2023) and run it until 2019. For analysis of the simulations, we diagnosed two-day means of all relevant state variables, processes, and transports. For the analysis of our results, we derived diagnostic variables from the model state variables:

- (a) Secchi depth as a function of water constituents: Neumann et al. (2015).
- (b) Chlorophyll: The sum of phytoplankton model variables multiplied with a constant chlorophyll to carbon mass ratio of 40
- 120 (e.g., Neumann et al., 2015).

3 Results

First, we will assess the model’s performance, followed by presenting the results from its runs.

3.1 Model Skill Evaluation

The model reasonable reproduces the climatology as well as the interannual variability at stations KHM and C (Fig. 1). For

125 bottom oxygen, the model predicts lower values than in the observations. Reasons are that (i) measurements are not as close to the bottom as the model data, (ii) the measurement platform (vessel) itself disturbs the stratification, and (iii) at station C commercial ship traffic induces strong vertical mixing which is not part of the model.




Figure 3. The mean number of days of anoxia per year in the bottom layer (a) and the mean duration of anoxia (b). Numbers are calculated as mean over the simulation period 1995–2019. The map was created using the software package GrADS 2.1.1.b0 (<http://cola.gmu.edu/grads/>, last access: 28 November 2024), using published bathymetry data (Seifert et al., 2008).

The mass transport through the Dziwna channel (Fig. 1) is elevated compared to known values. The reason is the truncated Dziwna channel in the model reducing the hydraulic resistance and facilitating an enhanced discharge at the expense of a lower discharge through the Swina channel.

The stratification of the water column at station KHM (Fig.1) compares well with observed stratification. Stratification establishes as events, such as those in summer, while during winter the water is well mixed. Stratification results in oxygen deficiencies, which yield phosphate liberation from the sediment. This process is not directly observed but is indirectly indicated by elevated phosphorus concentrations in summer (Fig.A6).

An extended analysis of the model performance is provided in Appendix A.

3.2 Oxygen dynamics

Due to the shallow bathymetry, the water column is well-mixed during winter. However, in summer, stratification may occur, and elevated temperatures accelerate metabolic processes, potentially leading to anoxic conditions in bottom waters. Figure 3a illustrates the total number of oxygen-depleted days, while Figure. 3b shows the average duration of these anoxic events. A notable region of anoxia is found in the navigation channel, which is 10 meters deep – approximately 5 meters deeper than the surrounding area. This deeper channel acts as a sediment trap, increasing oxygen demand for sediment respiration. Additionally, ventilation from the surface occurs less frequently here compared to other regions due to the greater water depth. Aside from the navigation channel, the eastern part of the lagoon is more affected by anoxia than the western part. This is likely due to the high nutrient loads from the Oder River entering the eastern lagoon.




Figure 4. The relative retention capacity of the Oder lagoon for nitrogen (a) and phosphors (b). The blue line is the sensitivity simulation with halved loads.

145 3.3 Filter function

Lagoons play an important ecological role, particularly through their filtering function, which is crucial for nutrient management. Biogeochemical processes within the lagoon help retain or remove nutrients from the system. Consequently, the amount of nutrients entering the open sea is reduced compared to the amount entering the lagoon.

In this section, we relate nutrient sources to their sinks to determine the proportion of nutrient loads removed within the lagoon. Additionally, the retention capacity of the Oder Lagoon may be sensitive to variations in nutrient loads. To evaluate this sensitivity, we performed an additional simulation with a 50% reduction in nutrient input, allowing us to assess how such changes influence the lagoon's filtering capacity. The results are presented in Fig. 4.

Sources of nitrogen and phosphorus primarily include riverine loads and atmospheric deposition, with riverine loads being the dominant contributor. An additional source of nitrogen is nitrogen fixation by cyanobacteria. Phosphorus has a single sink in the system – burial in the sediment. In contrast, nitrogen has an additional sink through denitrification, occurring both in the water column and the sediment. These factors help explain the contrasting responses of phosphorus and nitrogen retention capacities to changes in nutrient loads. For phosphorus, reduced loads lead to a decrease in primary production, which results in less organic matter reaching the sediment for burial. For nitrogen, burial also decreases; however, denitrification in the sediment increases because of the greater availability of oxygen.

In the next step, we are testing whether there exists a robust relationship between the retention capacity and the loads for nitrogen and phosphorus. For this purpose, we combined data from the reference run and the run with halved loads to increase the range of loads and consider the annual means. Figure 5a,c shows the absolute numbers of loads (source) into the system and the retention (sink). The retention increases with the loads, but is this retention at the same rate for all load realizations? Figure 4 suggests that the rate changes with the loads. A detailed dependence of the retention rate on loads is given in Figure 5b,d. In the case of nitrogen, the retention rate significantly decreases with increasing loads. In contrast, a significant relationship between




Figure 5. Annual means of nitrogen and phosphorus retention (a, c) and relative retention (b, d) in dependence on the loads. Red squares are from the 50% reduced loads experiment, black dots from the control run.

the retention rate and phosphorus loads does not exist, as Figure 4b suggests. A t-test for the regression coefficient confirms the visual indication. Consequently, the difference seen in Fig. 4b is statistically not significant.

In summary, the nitrogen retention capacity is stronger than that of phosphorus, primarily due to denitrification. Approximately 40% of nitrogen and 12% of phosphorus are retained in the lagoon. Reduced nutrient loads to the lagoon increase the nitrogen retention capacity, while the phosphorus retention capacity remains largely independent of load variations. It is important to note that this statement is valid only for the load range applied in the simulations. Mean numbers of sources and sinks are summarized in Tab. 1.

Table 1. Sinks and sources (loads, N-fixation) of nitrogen and phosphorus in the Oder Lagoon in kt/a. Shown are mean values for the simulation period.

	N [kt/a]	P [kt/a]
Loads	46.5	2.2
N-Fixation	1.3	
Denitrification Sed.	13.7	
Denitrification Pelag.	1.8	
Burial	1.8	0.25

4 Discussion

4.1 Quality of the model results

175 The Oder Lagoon model demonstrates strong performance in simulating physical and biogeochemical processes. The assessment was conducted at two central stations located in the lagoon’s subregions Kleines Haff (western part) and Großes Haff (eastern part), where the best coverage of observations was available.

The model successfully reproduces temperature and salinity patterns at both sites, particularly capturing the interannual salinity variability. However, it fails to replicate the highest observed salinity values. We attribute this limitation to the truncated
180 Dziwna channel in the model (Figure A5), which results in overestimated transport through the Dziwna channel at the expense of transport through the Swina channel, thereby introducing less saline water into the lagoon. Additional uncertainty stems from the meteorological forcing and consequently the open boundary conditions.

Transport estimates through the connecting channels (Mohrholz and Lass, 1999) suggest that mass transport in the Peene and Dziwna channels is equal, or possibly even lower in the Dziwna channel. However, our simulation yields a mass transport
185 in the Dziwna channel that is twice that of the Peene channel. One possible explanation for this discrepancy could be the incomplete representation of the Dziwna channel in our lagoon bathymetry (see Fig. 1). The absence of the Dziwna channel’s full hydraulic resistance in the model could account for the enhanced mass transport observed in the simulation.

A notable characteristic of the simulated nutrient concentrations is the absence of extremely high phosphate values (Fig.A6). It is hypothesized that these high values are caused by the release of sedimentary phosphate under low-oxygen conditions.
190 Although this process is included in the model (Fig.A7), the amount of phosphate released is evidently insufficient to elevate the surface concentration to observed levels. This discrepancy could be attributed to the two-dimensional sediment module used in this study. This kind of model does not have a long memory in terms of, for example, the storage of iron-phosphate complexes. Therefore, our model does not ‘remember’ the peak eutrophication period in the 1980s and early 1990s. A hint

for this explanation is the reduced phosphate peak frequency in recent years. Implementing a more sophisticated, vertically
195 resolved sediment module (Radtke et al., 2019) could potentially improve the model's performance.

Oxygen deficiency in the near-bottom water is widespread in the Oder Lagoon. The most affected area is the artificial
navigation channel. Being the deepest part of the lagoon, it acts as an accumulation area for sediment and especially for
organic matter. Degradation processes consume near-bottom oxygen, and the depth of about ten meters hinders ventilation
from the top water layers. Besides the navigation channel, the eastern part of the lagoon experiences anoxia more frequently
200 and for longer periods. This is caused by the high nutrient loads from the Oder River entering this part of the lagoon (Fig. 1).
Nevertheless, the model does not account for ship traffic, which causes regular vertical mixing down to the bottom. Thus, in
contrast to the model, observations do rarely show hypoxia in the navigation channel.

The newly introduced phytoplankton functional group (limnic phytoplankton) is by far the most abundant model phyto-
plankton group. This new group was necessary to achieve realistic biomass concentrations. In combination with the other three
205 groups, we ensure that the biogeochemical model ERGOM can be applied in coastal waters as well as in the open Baltic Sea
without parameter tuning. This is especially important when the model is set up for the entire Baltic Sea at a high spatial reso-
lution, for example, one nautical mile. In this case, the ecosystem model provides reasonable results in coastal waters, lagoons,
and the open Baltic Sea. Thus, simulations with a coarse-grained model deliver nearly seamless data for the open boundaries
of the local model setups.

210 Coarse-grained models, usually used for long-term simulations of the entire Baltic Sea, do not sufficiently resolve coastal
features and therefore the coastal filter function. Consequently, it is advisable to use corrected riverine loads, especially for large
rivers entering a lagoon before reaching the open Baltic Sea. For such a procedure, the retention capacity for the contributing
lagoons should be estimated with the aid of a local model. Preferably, the dependence of the retention capacity on the loads
should also be estimated.

215 Further discussion of the model performance is provided in Appendix A.

4.2 Filter function for nutrients

An important ecological function of the Oder Lagoon is, *inter alia*, the filtration and retention of nutrients. Acting as a system
between the Oder River mouth and the Baltic Sea, it reduces the amount of nutrients entering the open Baltic Sea. Budget
calculations based on observations in Lampe (1999) yield a low retention capacity of 2%—5%, mainly caused by dredging
220 of the navigation channel. Grelowski et al. (2000) found that 12%—29% of total nitrogen and 11%—27% of total phosphorus
is retained in the Oder Lagoon. Asmala et al. (2017) showed that denitrification in lagoons of the Baltic Sea is highest, while
phosphorus burial in lagoons is small compared to other coastal systems. Our model based approach yields a retention capacity
of 40% for nitrogen and 12% for phosphorus. The lower retention for phosphorus is in line with Asmala et al. (2017). In contrast
to Asmala et al. (2017), who only considered denitrification, we considered additional sources and sinks for nitrogen: nitrogen
225 fixation and nitrogen burial. Figure 6 illustrates the nitrogen sources and sinks in our model. The dominant source is riverine
nutrient loads, while the primary sink is denitrification in the sediment. Dredging is not included in the model. However, its




Figure 6. Nitrogen sources (blue) and sinks (red) in the Oder Lagoon. N-Sources are riverine loads and atmospheric deposition while N-Fixation is shown as a separate source (blue, dashed line). Tot N-Sink is the total sink of nitrogen; Red dashed and dotted lines are the contributing processes.

absence is nearly compensated for by the model's sediment burial process, which occurs when the sediment thickness exceeds a specified threshold. The only sink for phosphorus is sediment burial as shown in Tab. 1.

Models offer the advantage of enabling experiments to be performed on the system. We conducted an experiment with halved
 230 riverine nutrient loads. The reduced loads resulted in decreased retention for phosphorus, while the retention for nitrogen increased (see Fig. 4). The cause of the lesser phosphorus retention is the reduced primary production due to lower nutrient concentrations. This, in turn, leads to less sedimentation and hence, less burial. However, this dependence is not statistically significant based on our model simulations. Simulations with a broader range of loads could potentially establish a statistically significant relationship if it exists. Nevertheless, the loads we applied are within a reasonable range for realistic scenarios. In
 235 the case of nitrogen, the contributing sinks behaved differently (see Fig.7). While burial is reduced, similar to the phosphorus case, the relative sedimentary denitrification increases. Denitrification in the sediment is more effective with higher oxygen concentrations at the sediment-water interface. The higher oxygen concentration results from less primary production due to reduced loads and hence less organic matter sinks to the sediment. Overall, nitrogen retention is more effective in the case of lower nutrient loads. A significant correlation could be established for this relationship (Fig. 5).

240 On average over the 25 simulation years, 40% of the riverine nitrogen (N) loads were retained in the lagoon. Instead of an average total load of 46,266 t/a N entering the lagoon, only 28,988 t/a N entered the Baltic Sea, resulting in a difference of 17,278 t/a N that was retained in the lagoon (Figure 8a,b). With respect to phosphorus, on average over the 25 years, 12% of the riverine phosphorus was retained in the lagoon and buried in sediments. Instead of an average of 2,198 t/a P entering with the rivers, only 1,945 t/a entered the Baltic Sea, resulting in a load reduction of 253 t/a P in the lagoon. Taking the nutrient




Figure 7. Change in relative nitrogen retention capacity between control run and 50% load reduction run.




Figure 8. Relationship between annual riverine nitrogen and phosphorus loads and the nutrient retention in the Oder lagoon. c,d) Relationship between average annual water residence time and the nitrogen and phosphorus retention in the Oder lagoon. Residence time has been estimated following Vollenweider (1976); Monsen et al. (2002) by volume over runoff.

245 retention in the lagoon into account, the loads to the Baltic Sea between 1995 and 2019 (28,988 t/a N; 1,296 t/a P) were already far below the BSAP requirements.

The lagoon is an efficient nutrient trap, particularly for nitrogen, and serves as an important purification unit for the Baltic Sea. Especially with respect to nitrogen, a close relationship exists between the relative nitrogen retention in the lagoon and the

lagoon's water exchange time. This relationship also exists for phosphorus, but it is less pronounced (Figure 8c,d). In general,
250 we can state that the longer water remains in the lagoon, the higher the relative retention of nutrients in the lagoon.

In summary, the filter function appears as follows: The fraction of nitrate removal is larger than that for phosphorus. A dependence of the retention rate on loads exists for nitrogen. An additional dependence exists on the water residence time. For phosphorus, we did not find a similar dependence. We think that these findings can be transferred to similar coastal regions like the Curonian Lagoon, especially the dependence of the nitrogen removal rate on the water residence time. However,
255 transferring our findings to other regions requires additional sound scientific studies. This is particularly true for semi-enclosed and open coastal systems.

4.3 Annual discharge, nutrient loads and water quality targets

The Oder River contributes about 98% of the direct water discharge to the lagoon, while the remaining 2% is contributed by the Zarow and Uecker rivers. The total water discharge into the lagoon, along with the loads of nitrogen (N) and phosphorus (P), exhibits similar temporal behavior (Figure 9a). Between 1995 and 2019, the average water discharge was $518 \text{ m}^3/\text{s}$, with
260 average annual total loads of 46,266 t N and 2,198 t P to the lagoon. Between 1995 and 1999, the average water discharge was $643 \text{ m}^3/\text{s}$, which is higher than the average over the 25-year model simulation period. Consequently, the annual N loads (62,534 t N) and P loads (3,600 t P) were also higher. In contrast, during the recent years between 2015 and 2019, the average annual discharge was only $413 \text{ m}^3/\text{s}$, resulting in N loads of 37,077 t/a and P loads of only 1,449 t/a. The close relationship
265 between discharge and nutrient loads is illustrated in Figures 9b and c. This relationship emphasizes the dependency of annual riverine nutrient loads to the lagoon on the water discharge of the Oder River. The concentration of both nutrients, nitrogen and phosphorus, is largely independent from river discharge. This has been observed already earlier and is the reason why the long-term assessment of critical loads used a discharge correction/normalization. Load reduction due to the changes in riverine nutrient concentrations was more dominant in the 1990's (Friedland et al., 2019). The conclusion is that hydrological processes
270 in the approximately 120,000 km^2 Oder River catchment basin exert a stronger control over the nutrient loads to the Oder Lagoon than annual changes in nutrient inputs. Consequently, future climate change effects on the catchment water budget will significantly impact the riverine nutrient loads and the ecological state of the lagoon. However, it is important to note that this conclusion may not apply to long-term perspectives, particularly when significant changes in the catchment begin to take effect.

275 In Germany, target or threshold values for a good ecological status in rivers exist, with concentrations below these values indicating a good status. The riverine target values are 2.6 mg/L for nitrogen (N) and 0.1 mg/L for phosphorus (P). The nutrient concentrations in recent years (2015-2019) were already close to these target concentrations. Using the average riverine water discharge to the lagoon of $518 \text{ m}^3/\text{s}$ (1995–2019) and the German nutrient target concentrations, the resulting target loads would be 42,506 t/a for N and 1,635 t/a for P. The HELCOM Baltic Sea Action Plan (BSAP) defines the maximum allowable
280 nutrient loads that enable the Baltic Sea to reach a good status. According to the BSAP, the maximum allowable loads are approximately 45,000 t/a for N and 1,500 t/a for P. Despite following different approaches, the German target values in rivers and the HELCOM maximum allowable loads result in comparable targets for the Oder River. In contrast, the Polish targets




Figure 9. Average annual water discharge and annual loads of total nitrogen and total phosphorus to the Oder (Szczecin) Lagoon during a 25 years period (1995–2019), always including the rivers Oder/Odra, Uecker and Zarow. Relationship between riverine water discharge annual riverine nitrogen loads (b), annual riverine phosphorus loads (c), annual average phytoplankton biomass (chlorophyll-concentrations) (d) and annual average water transparency (Secchi depth) (e) in the Oder Lagoon (spatially integrated). (f) Relationship between riverine phosphorus and nitrogen concentrations.

would allow much higher Oder River nutrient loads to the Oder Lagoon (63,661 t/a for N and 4,615 t/a for P), as shown by Friedland et al. (2019). However, during recent years (2015–2019), the N loads to the lagoon were 37,077 t/a and the P loads were 1,449 t/a. For both nutrients, the loads were below the maximum allowable inputs (HELCOM, 2013).

According to our model simulations, riverine nutrient loads have an immediate effect on major water quality indicators, namely phytoplankton biomass (chlorophyll-a) and water transparency (Secchi depth), and show a close correlation (Figure 9d,e). Increased discharge, resulting in increased nutrient loads, causes an increase in chlorophyll-a concentrations in the lagoon and a decrease in Secchi depth. Calculated chlorophyll-a and Secchi depth values are average annual values over the entire lagoon and cannot be directly compared with existing ecological target values for central stations in the lagoon (Schernewski et al., 2015). However, the changes in chlorophyll-a concentrations and Secchi depth that would result from fully meeting the nutrient load targets (42,506 t/a N and 1,635 t/a P) are only minor (Figure 9d,e). Compared to the average values over the period 1995–2019, chlorophyll-a concentrations would decrease by 5%, and Secchi depth would increase by 8%. Although, the nutrient loads are below the BSAP values, the lagoon remains in a highly eutrophic state with a bad ecological status, according to the official HELCOM HEAT HOLAS 3 (HELCOM, 2021b) eutrophication status assessment (BMU, 2024). Meeting water quality targets in rivers and the Baltic Sea does not guarantee that a good status in transitional ecosystems, such as lagoons, will be achieved. A good ecological status in the lagoon would require significant additional nutrient load reductions and the implementation of measures in the river catchment at high and likely unrealistic costs. It is likely that the lagoon must be regarded as a naturally eutrophic ecosystem with limited management possibilities. A re-evaluation of water quality targets in the lagoon requires a more detailed study that includes neighboring coastal waters to address interrelationships, relates model data to field data with a focus on the assessment stations, and carefully considers evaluation aspects such as water depth and evaluation period.

4.4 Interannual variability of discharge and loads and its consequences

The interannual variability of water discharge and nutrient loads is high (Figure 9a). Annual discharges vary between 291m³/s in 2015, a dry and hot year, and 781m³/s in 2010, the year with one of the largest Oder River floods ever recorded, which occurred in May. Consequently, nutrient loads also show high variability, ranging from 20,309 t/a N in 2015 to 72,333 t/a N in 2010. For P, the range is from 4,683 t/a in 1997, another major river flood year, to 1,094 t/a in 2018, another hot and dry year. Over the 25-year period, riverine concentrations of N and P do not show a close relationship; the concentrations of both elements behave differently in different years.

The differing interannual variability between N and P concentrations indicates that both nutrients enter via different pathways and that these pathways play different roles in different years. This suggests that seasonal discharge and nutrient emission patterns need to be analyzed separately and in depth. It is likely that the interannual variability is controlled by extreme events lasting weeks to a few months, which can strongly affect the annual values and the lagoon ecosystem.

The Baltic Sea environment is projected to experience extreme conditions more frequently in the future (Rutgersson et al., 2022). These include, but are not limited to, increased occurrences of flooding and drought events. Given that these events substantially influence nutrient loads, they are likely to amplify the interannual variability of ecological conditions. Consequently, the increased amplitude of environmental fluctuations will intensify stress on lagoon fauna and flora.

Table 2. Loads for different periods and targets into the Oder Lagoon and Baltic Sea. Loads and targets are in kt/a, runoff in m³s⁻¹. N and P retention is estimated from the period 1995 to 2019. Loads to the Baltic Sea for other periods are estimated from the mean retention.

	N-load Lagoon	N-load Baltic	P-load Lagoon	P-load Baltic	Runoff	N-retention	P-retention
1995–2019	46,266	28,988	2,198	1945	518	0.37	0.12
1995–1999	62,534	39,181	3,600	3,186	643		
2015–2019	31,077	23,231	1,449	1,282	413		
German target Oder River	42,506		1,635				
BSAP MAI to Baltic		45,000		1,500			

5 Conclusions

We developed a local model for the Oder Lagoon that realistically reproduces its physical and biogeochemical properties. This model serves as a tool for conducting studies in this area and, specifically, for quantifying the lagoon’s nutrient retention capacity. This is a crucial step in adjusting riverine loads for coarse-grained models, which often do not adequately resolve lagoons. Our approach can be readily applied to other lagoons in the Baltic Sea, such as the Curonian Lagoon, and can also be adapted for regions beyond the Baltic Sea. A general finding is the relationship between water residence time and retention of nitrogen.

The analysis shows that the nutrient retention in the lagoon already reduces nutrient loads to the open Baltic Sea in agreement with the BSAP (Table 2). Furthermore, riverine loads into the Oder Lagoon also meet the German targets for the Oder River. However, nutrient concentrations in the Oder Lagoon do not achieve the intended target values for good ecological status. This points to the problem that quality standards for inner and outer waters are not harmonized, making it unrealistic to achieve a good ecological status for the Oder Lagoon.

The simulation data generated in this study enable several additional analyses that could be the focus of future studies. For example, these could include short- and long-term responses to extreme events such as floods and droughts. Further improvements should include a more realistic representation of the Dziwna channel length and coupling the model with a more advanced sediment model than the one used in this study. Another weakness is the lack of ventilation of the deep water in the navigation channel due to ship traffic, which should be properly parameterized in an upcoming model version.

Code and data availability. Observations from the Oder Lagoon monitoring program were provided upon request by the German State Agency for Environment, Nature Conservation and Geology Mecklenburg-Vorpommern (Mario von Weber, LUNG-MV) and Główny Inspektorat Ochrony Środowiska (Adam Czugała). Nutrient loads and runoff data for the Oder River were obtained from from the Polish Główny Inspektorat Ochrony Środowiska. Corresponding data for the rivers Zarow and Ücker were available from LUNG-MV.

The meteorological forcing is archived at https://doi.org/10.1594/WDCC/coastDat-2_COSMO-CLM (last access: 26 November 2024,
340 Geyer and Rockel (2013)).
All model output that was analyzed in this paper, the model code, and data to run the model has been published at <https://doi.org/10.5281/zenodo.14236528> (Neumann et al., 2024).

Appendix A: Detailed analysis of the model performance

In this section, we compare model data with observations, demonstrate some model properties not present in observational data,
345 and discuss the results of this analysis. Observations are available on request from national agencies (see *data availability*).

A1 Station data

A1.1 Climatology

We start with the evaluation of data from stations KHM and C shown in Fig. 1. Figure A1 shows the climatologies (1996-2019)
for several surface variables and bottom oxygen. The model is able to reproduce the observations, and the spread of the model
350 and observations mostly matches. However, a few details need to be mentioned and discussed.

The Oder Lagoon was covered by sea ice in several winters during the simulation period. No observations exist for these
time periods. Thus, cold temperatures are missing in the observational dataset, and the number of winter values for nutrients is
reduced.

The model overestimates the summer temperature and underestimates the salinity. Several reasons could be responsible for
355 these biases. However, within the scope of this study, we cannot verify which factors drive the observed biases. Thus, we can
only suggest some possibilities that could be the subject of further investigations. (i) The meteorological forcing impacts both
temperature and salinity, and in addition, the mixing depth. A mixing depth that is too shallow could also be responsible for
the positive temperature bias in summer. Beyond the meteorological forcing, the choice and tuning of the vertical sub-scale
parameterization determines the mixing depth. Observations are only available for the surface and bottom layers. Thus, there is
360 hardly any constraint for the mixing depth. For salinity, the freshwater budget in the lagoon is also an important factor, which is
controlled by the Oder River discharge (Sec. 4.3). The overestimated volume transport through the Dziwna channel (Sec. A1.2)
may also contribute to the lower salinity values in the model.

The observed chlorophyll has two distinct peaks, denoting spring and late summer blooms. In the model, only one peak
appears in early summer. Chlorophyll is not a model's state variable and is diagnostically estimated with a constant carbon-to-
365 chlorophyll ratio. It is well known (Jakobsen and Markager, 2016) that this ratio has a seasonal cycle, which we do not consider
(see our remarks in Section A1.2).

The near-bottom oxygen levels exhibit clear differences between the model and observations. We assume the model is closer
to reality due to common shortcomings in observational techniques. The model data are collected approximately 20 cm above
the seafloor, while observations are typically taken from 1 m above the seafloor. Fredriksson et al. (2024) demonstrate that




Figure A1. Climatology of surface data for temperature (a), salinity (b), DIN (c), phosphate (d), and chlorophyll (e). Bottom oxygen is shown in (f). Blue color are data from the model green color are data from observations. The shaded area is the range between 10th and 90th percentile. All data are from station KHM (see Fig. 1).

370 strong oxygen gradients exist above the seafloor in coastal waters, which cannot be resolved with traditional CTD instruments. Furthermore, the measurement platform - typically a vessel - may disrupt the vertical structure of the water column.

The assessment for station C (Fig. A2) mirrors that of station KHM. Unfortunately, temperature data are unavailable for this station. Simulated bottom oxygen levels are even lower than those at station KHM, which can be attributed to the greater water depth in the navigation channel. As previously discussed for station KHM, discrepancies between model and observations stem from limitations in the observations themselves.

375




Figure A2. Climatology of surface data salinity (a), DIN (b), phosphate (c), and chlorophyll (d). Bottom oxygen is shown in (e). Blue color are data from the model green color are data from observations. The shaded area is the range between 10th and 90th percentile. All data are from station C (see Fig. 1).

Moreover, the model does not incorporate the substantial ship traffic to and from Szczecin harbor. According to Schernewski et al. (2025), approximately 3,300 cargo ships arrive at Szczecin harbor annually. The deep draught of these vessels, which nearly reaches the seafloor in the navigation channel, induces regular mixing of the water column.

Another difference between the model and reality involves the dredging of the navigation channel. Functioning as a sediment trap, the channel requires regular dredging to maintain its depth (Schernewski et al., 2024). While the model's burial process partially simulates this dredging effect, it represents only an approximation of the actual process.




Figure A3. Climatology of Secchi depth at stations KHM (a) and C (b) (see Fig. 1).

Table A1. Root mean square error (RMSE), standard deviation (STD) of the observations, and normalized RMSE (RMSD).

	RMSE	STD	RMSD
DIN KHM	36	44	0.8
DIP KHM	1.6	1.6	1.0
Salt KHM	0.56	0.69	0.82
Temp KHM	3.3	6.3	0.5
DIN C	47	67	0.7
DIP C	1.4	1.7	0.8
Salt C	0.59	0.65	0.9

Figure A3 presents the climatology of Secchi depth at stations KHM and C. The model data demonstrate more pronounced seasonality compared to the observations. This discrepancy may stem from the model’s exclusion of mineral sediment particles. The resuspension of these particles, particularly during winter when wind mixing is stronger, serves to limit Secchi depth. Additionally, it is important to note the absence of data during sea ice winters.

Finally, Table A1 presents the root mean square error (RMSE) and the RMSE normalized by the standard deviation of observations (RMSD). An RMSD value below unity indicates that the RMSE falls within the range of natural variability (Kärnä et al., 2021).

A1.2 Time series

Figure A4 presents surface salinity measurements at stations C (panel b) and KHM (panel c). The simulations at both stations successfully capture the observed interannual variability, though the model exhibits a negative salinity bias for some




Figure A4. Surface temperature and salinity at station KHM (a,c) and salinity at station C (b) (see Fig. 1). Red diamonds: Observations; Blue line: Model simulation.

periods. Corresponding sea surface temperature (SST) data are shown in Figures A4a for station KHM. Winter observations are frequently unavailable due to sea ice coverage, while during summer months, the model tends to overestimate SST values compared to observations.

395 Water exchange between the lagoon and the Baltic Sea occurs through three primary channels: the Peene Stream, the Świna, and the Dziwna (Fig. 1). The relative contribution of each channel to the overall water exchange with the Baltic Sea has been estimated in multiple studies using both observational data and modeling approaches. Mohrholz and Lass (1999) provides a comprehensive overview of these estimates, reporting the following contribution ranges: Peene Stream: 14%–20%, Świna: 60%–75%, and Dziwna: 9%–20%. Figure A5 illustrates the contributions of the three channels to water exchange in the model
400 simulation. The transport through the Dziwna channel appears higher than in other estimates, while transport through the Świna channel is reduced. This discrepancy occurs because the truncated Dziwna channel reduces hydraulic resistance. This issue should be addressed in future, improved model setups.

Figure A6 presents time series of nutrient concentrations, including phosphate, dissolved inorganic nitrogen (DIN), and chlorophyll, at stations C and KHM. The model successfully captures the decreasing trend in winter phosphate concentra-
405 tions at both stations. However, it fails to reproduce the exceptionally high observed concentrations during summer months. These peak values are assumed to result from very low oxygen conditions at the sediment surface, which liberate iron-bound




Figure A5. The relative contribution of the three outlet channels of the Oder Lagoon to the net water exchange with the Baltic Sea.

phosphate from the sediment. Although this process is included in the model, the amount of phosphate released is insufficient to significantly elevate surface concentrations. In certain periods and regions, the model indicates oxygen depletion (see Chapter 3.2), suggesting that the preconditions for phosphate release are indeed met. We hypothesize that the two-dimensional sediment module of the ERGOM model cannot sufficiently parameterize vertical processes, which might be important for the long-term storage of phosphorus in the sediment.

The model underestimates chlorophyll concentrations at station KHM. This discrepancy may be attributed to our simple chlorophyll estimation method, which uses a constant Chl:C mass ratio (see Chapter 2). This approach neglects the annual dynamics of the Chl:C ratio, which represents a phytoplankton response to changing ambient light conditions and Chl:C may vary between 23 and 60 (Jakobsen and Markager, 2016). Furthermore, the simulated data exhibits a slight decreasing trend, whereas the observations at station KHM do not display a significant trend.

A2 Stratification and phosphorus liberation from sediments

Particulate iron-phosphate complexes, primarily accumulated in sediments, dissolve under anoxic conditions, consequently liberating phosphate from the sediment. This process, also known as internal eutrophication (Vahtera et al., 2007), is included in our ecosystem model ERGOM, as demonstrated in Figure A7.

The mixed layer depth reaches its minimum in summer, coinciding with strong stratification that results in low bottom oxygen concentrations (Figure A1f). Simultaneously, phosphate is liberated from sediments (Figure A7a), yielding maximum phosphate concentrations in summer (Figures A1d and A2c). However, this summer phosphate peak is less pronounced in the model compared to observations.

Mixed layer depth cannot be derived from observations because only surface and near-bottom data are available. Instead, we present the density differences between bottom and surface water as a measure of stratification in Figure A7b,c. Observations




Figure A6. Surface concentration of phosphate, DIN, and chlorophyll at station C (a,c,e) and station KHM (b,d,f) (see Fig. 1). Red diamonds: Observations; Blue line: Model simulation. Straight lines show the trend if significant.

are compiled as monthly data, which can be compared to the red line in Figure A7b. The higher values of the two-day means (black line) indicate that strong stratification, and consequently anoxia, occurs in the form of discrete events.

A3 Horizontal patterns

430 Simulated data are evaluated against monitoring data, enabling the calculation of the root mean square deviation (RMSD) for each station and parameter. Following Kärnä et al. (2021), we normalize the root mean square error by the standard deviation of the observations to facilitate comparisons across different stations and parameters. A dimensionless normalized RMSD value




Figure A7. Climatology of model mixed layer depth and PO₄ liberation (a), stratification in model simulation (b), and in observations (c). Blue color in (a) are the mixed layer depth and green color the PO₄ liberation. The shaded area is the range between 10th and 90th percentile. The red line in (b) is from monthly mean and the black line from two-day mean model data. For (c), monthly mean observations are used. All data are from station KHM (see Fig. 1).

below one is typically considered indicative of good model performance, as it suggests that the RMSD falls within the range of natural variability observed in the data.

435 Figure A8 presents horizontal surface patterns of model variables and their corresponding RMSD. For the winter season, observations were unavailable for all stations. Dissolved inorganic nitrogen (DIN) exhibits a clear gradient in both seasons, with the highest values near the mouth of the Oder River and concentrations decreasing sharply toward the western lagoon and the open Baltic Sea. The RMSD for the monitoring stations is predominantly below one, indicating good model performance. Only at station E (near the mouth of the Oder River) does it slightly exceed this threshold.

440 Dissolved inorganic phosphorus (DIP) displays an opposite gradient during winter, with the highest concentrations in the western lagoon. During the growing season, DIP concentrations remain elevated. While the RMSD during the growing season is mostly below one (except at station E near the Oder River), the model overestimates DIP concentrations in winter compared to observed values. For the remainder of the year, DIP concentrations fall within the range of observations (see Figures A1 and A2).




Figure A8. Near surface concentrations of dissolved inorganic nitrogen A: Winter average, C: May to September average; dissolved inorganic phosphorus B: Winter average, D: May to September average; Chlorophyll-a E: May to September average, and Secchi depth F: May to September average. All parameters are averaged from 2010 to 2019. The model results are color-coded and circles indicate the normalized RMSD for each monitoring station and parameter. The map was created using the software ESRI ArcGIS Pro (Version 3.3.2; <https://pro.arcgis.com>).

445 Summer chlorophyll-a concentrations and Secchi depth (Figure A8) exhibit comparable gradients due to their close relationship. The lowest chlorophyll-a concentrations occur in the western part of the lagoon, where Secchi depth is greatest. Although chlorophyll-a is underestimated in the western part, the RMSD remains within the range observed at eastern stations. For Secchi depth, the RMSD is even better for western stations (values between 1.1 and 1.2), while it reaches up to two for eastern stations. Despite the good agreement between modeled and observed chlorophyll-a in the eastern Oder Lagoon, the model
 450 simulates too rapid a decline in chlorophyll-a at station C during September, resulting in an overly rapid increase in Secchi depth during that month (see Figure A2). Additionally, the model does not account for resuspended mineral sediments, which impact Secchi depth as outlined in Section A1.1.

Author contributions. TN set up the model and conducted the simulations. All authors contributed to the study design, data analysis, and the writing of the manuscript.

455 *Competing interests.* The corresponding author has declared that none of the authors have any competing interests.

Acknowledgements. The authors gratefully acknowledge the computing time made available to them on the high-performance computer "Emmy" at the NHR Center NHR-NORD@GÖTTINGEN. This center is jointly supported by the Federal Ministry of Education and Research and the state governments participating in the NHR (www.nhr-verein.de/unsere-partner). We thank LUNG-MV (Mario von Weber) and Główny Inspektorat Ochrony Środowiska (Adam Czugala) for providing German and Polish monitoring data. This work was financially supported by the German Federal Ministry of Education and Research, projects "Coastal Futures II" (grant number 03F0980B), "Prime Prevention" (grant number 03F0953D) and UBA-MoSea (FKZ 3723252040). We acknowledge the contributions of Sarah Piehl, who collected and prepared data from German and Polish authorities.

References

- Zustand der deutschen Ostseegewässer 2024, Bundesministerium für Umwelt, Naturschutz, nukleare Sicherheit und Verbraucherschutz (BMUV) (Hrsg.), https://mitglieder.meeresschutz.info/de/berichte/zustandsbewertungen-art8-10.html?file=files/meeresschutz/berichte/art8910/zyklus2024/Zustandsbericht_Ostsee_2024.pdf, last accessed on 2024-11-20, 2024.
- Almroth-Rosell, E., Edman, M., Eilola, K., Meier, H. E. M., and Sahlberg, J.: Modelling nutrient retention in the coastal zone of an eutrophic sea, *Biogeosciences*, 13, 5753–5769, <https://doi.org/10.5194/bg-13-5753-2016>, 2016.
- Asmala, E., Carstensen, J., Conley, D. J., Slomp, C. P., Stadmark, J., and Voss, M.: Efficiency of the coastal filter: Nitrogen and phosphorus removal in the Baltic Sea, *Limnology and Oceanography*, 62, S222–S238, <https://doi.org/https://doi.org/10.1002/lno.10644>, 2017.
- Börgel, F., Neumann, T., Rooze, J., Radtke, H., Barghorn, L., and Meier, H. E. M.: Deoxygenation of the Baltic Sea during the last millennium, *Frontiers in Marine Science*, 10, <https://doi.org/10.3389/fmars.2023.1174039>, 2023.
- Carstensen, J., Andersen, J. H., Gustafsson, B. G., and Conley, D. J.: Deoxygenation of the Baltic Sea during the last century, *Proceedings of the National Academy of Sciences*, 111, 5628–5633, <https://doi.org/10.1073/pnas.1323156111>, 2014.
- Conley, D. J., Carstensen, J., Aigars, J., Axe, P., Bonsdorff, E., Eremina, T., Haahti, B.-M., Humborg, C., Jonsson, P., Kotta, J., Lännegren, C., Larsson, U., Maximov, A., Medina, M. R., Lysiak-Pastuszek, E., Remeikaitė-Nikienė, N., Walve, J., Wilhelms, S., and Zillén, L.: Hypoxia Is Increasing in the Coastal Zone of the Baltic Sea, *Environ. Sci. Technol.*, 45, 6777–6783, <https://doi.org/10.1021/es201212r>, 2011.
- Diaz, R. J. and Rosenberg, R.: Spreading Dead Zones and Consequences for Marine Ecosystems, *Science*, 321, 926–929, <https://doi.org/10.1126/science.1156401>, 2008.
- Edman, M., Eilola, K., Almroth-Rosell, E., Meier, H. E. M., Wahlström, I., and Arneborg, L.: Nutrient Retention in the Swedish Coastal Zone, *Frontiers in Marine Science*, 5, <https://doi.org/10.3389/fmars.2018.00415>, 2018.
- Eilola, K., Gustafson, B. G., Kuznetsov, I., Meier, H. E. M., Neumann, T., and Savchuk, O. P.: Comparison of observed and simulated dynamics of biogeochemical cycles in the Baltic Sea during 1970–2005 using three state-of-the-art numerical models, *Journal of Marine Systems*, 88, 267–284, <https://doi.org/10.1016/j.jmarsys.2011.05.004>, 2011.
- Fennel, K. and Testa, J. M.: Biogeochemical Controls on Coastal Hypoxia, *Annual Review of Marine Science*, 11, 105–130, <https://doi.org/10.1146/annurev-marine-010318-095138>, 2019.
- Fredriksson, J. P., Attard, K., Stranne, C., Koszalka, I. M., Glud, R. N., Andersen, T. J., Humborg, C., and Brüchert, V.: Hidden seafloor hypoxia in coastal waters, *Limnology and Oceanography*, 69, 2489–2502, <https://doi.org/https://doi.org/10.1002/lno.12607>, 2024.
- Friedland, R., Schernewski, G., Gräwe, U., Greipsland, I., Palazzo, D., and Pastuszek, M.: Managing Eutrophication in the Szczecin (Oder) Lagoon-Development, Present State and Future Perspectives, *Frontiers in Marine Science*, 5, <https://doi.org/10.3389/fmars.2018.00521>, 2019.
- Geyer, B. and Rockel, B.: coastDat-2 COSMO-CLM Atmospheric Reconstruction [data set], https://doi.org/10.1594/WDCC/coastDat-2_COSMO-CLM, 2013.
- Grelowski, A., Pastuszek, M., Sitek, S., and Witek, Z.: Budget calculations of nitrogen, phosphorus and BOD5 passing through the Oder estuary, *Journal of Marine Systems*, 25, 221–237, [https://doi.org/10.1016/S0924-7963\(00\)00017-8](https://doi.org/10.1016/S0924-7963(00)00017-8), 2000.
- Griffies, S. M.: *Fundamentals of Ocean Climate Models*, Princeton University Press, Princeton, NJ, ISBN 9780691118925, 2004.
- HELCOM: HELCOM Baltic Sea Action Plan - 2021 update, <https://helcom.fi/wp-content/uploads/2021/10/Baltic-Sea-Action-Plan-2021-update.pdf>, last accessed on 2024-11-20, 2021a.

HELCOM: Eutrophication - Thematic assessment 2016–2021, Baltic Sea Environment Proceedings 192, <https://helcom.fi/wp-content/uploads/2023/06/HELCOM-Thematic-assessment-of-eutrophication-2016-2021.pdf>, last accessed on 2024-11-20, 2021b.

Jakobsen, H. H. and Markager, S.: Carbon-to-chlorophyll ratio for phytoplankton in temperate coastal waters: Seasonal patterns and relationship to nutrients, *Limnology and Oceanography*, 61, 1853–1868, <https://doi.org/10.1002/lno.10338>, 2016.

Kabel, K., Moros, M., Porsche, C., Neumann, T., Adolphi, F., Andersen, T. J., Siegel, H., Gerth, M., Leipe, T., Jansen, E., and Sinninghe Damsté, J. S.: Impact of climate change on the Baltic Sea ecosystem over the past 1,000 years, *Nature Climate Change*, 2, 871–874, <https://doi.org/10.1038/nclimate1595>, 2012.

Kärnä, T., Ljungemyr, P., Falahat, S., Ringgaard, I., Axell, L., Korabel, V., Murawski, J., Maljutenko, I., Lindenthal, A., Jandt-Scheelke, S., Verjovkina, S., Lorkowski, I., Lagema, P., She, J., Tuomi, L., Nord, A., and Huess, V.: Nemo-Nordic 2.0: operational marine forecast model for the Baltic Sea, *Geoscientific Model Development*, 14, 5731–5749, <https://doi.org/10.5194/gmd-14-5731-2021>, 2021.

Kuliński, K., Rehder, G., Asmala, E., Bartosova, A., Carstensen, J., Gustafsson, B., Hall, P. O. J., Humborg, C., Jilbert, T., Jürgens, K., Meier, H. E. M., Müller-Karulis, B., Naumann, M., Olesen, J. E., Savchuk, O., Schramm, A., Slomp, C. P., Sofiev, M., Sobek, A., Szymczycha, B., and Undeman, E.: Biogeochemical functioning of the Baltic Sea, *Earth System Dynamics*, 13, 633–685, <https://doi.org/10.5194/esd-13-633-2022>, 2022.

Lampe, R.: The Odra Estuary as a Filter and Transformation Area, *Acta hydrochimica et hydrobiologica*, 27, 292–297, [https://doi.org/10.1002/\(SICI\)1521-401X\(199911\)27:5<292::AID-AHEH292>3.0.CO;2-Z](https://doi.org/10.1002/(SICI)1521-401X(199911)27:5<292::AID-AHEH292>3.0.CO;2-Z), 1999.

Leibniz Institute for Baltic Sea Research: ERGOM: Ecological Regional Ocean Model, <https://ergom.net/>, last access: 10 March 2022, 2015.

Mohrholz, V. and Lass, H. U.: Transports between Oderhaff and Pomeranian bight – a simple barotropic box model, *Deutsche Hydrographische Zeitschrift*, 50, 371–383, <https://doi.org/10.1007/BF02764231>, 1999.

Monsen, N. E., Cloern, J. E., Lucas, L. V., and Monismith, S. G.: A comment on the use of flushing time, residence time, and age as transport time scales, *Limnology and Oceanography*, 47, 1545–1553, <https://doi.org/10.4319/lo.2002.47.5.1545>, 2002.

Müller-Karulis, B. and Aigars, J.: Modeling the long-term dynamics of nutrients and phytoplankton in the Gulf of Riga, *Journal of Marine Systems*, 87, 161–176, <https://doi.org/10.1016/j.jmarsys.2011.03.006>, 2011.

Neumann, T. and Schernewski, G.: Eutrophication in the Baltic Sea and shifts in nitrogen fixation analyzed with a 3D ecosystem model, *Journal of Marine Systems*, 74, 592–602, <https://doi.org/10.1016/j.jmarsys.2008.05.003>, 2008.

Neumann, T., Siegel, H., and Gerth, M.: A new radiation model for Baltic Sea ecosystem modelling, *Journal of Marine Systems*, 152, 83–91, <https://doi.org/10.1016/j.jmarsys.2015.08.001>, 2015.

Neumann, T., Koponen, S., Attila, J., Brockmann, C., Kallio, K., Kervinen, M., Mazeran, C., Müller, D., Philipson, P., Thulin, S., Väkevä, S., and Ylöstalo, P.: Optical model for the Baltic Sea with an explicit CDOM state variable: a case study with Model ERGOM (version 1.2), *Geoscientific Model Development*, 14, 5049–5062, <https://doi.org/10.5194/gmd-14-5049-2021>, 2021.

Neumann, T., Radtke, H., Cahill, B., Schmidt, M., and Rehder, G.: Non-Redfieldian carbon model for the Baltic Sea (ERGOM version 1.2) – implementation and budget estimates, *Geoscientific Model Development*, 15, 8473–8540, <https://doi.org/10.5194/gmd-15-8473-2022>, 2022.

Neumann, T., Schernewski, G., and Friedland, R.: Model code and output for "Transformation Processes in the Oder Lagoon as seen from a Model Perspective" paper [code], <https://doi.org/10.5281/zenodo.14236528>, last access: 28 November 2024, 2024.

Piehl, S., Friedland, R., Heyden, B., Leujak, W., Neumann, T., and Schernewski, G.: Modeling of Water Quality Indicators in the Western Baltic Sea: Seasonal Oxygen Deficiency, *Environmental Modeling & Assessment*, 28, 429–446, <https://doi.org/10.1007/s10666-022-09866-x>, 2023.

- Radtke, H., Lipka, M., Bunke, D., Morys, C., Woelfel, J., Cahill, B., Böttcher, M. E., Forster, S., Leipe, T., Rehder, G., and Neumann, T.: Ecological ReGional Ocean Model with vertically resolved sediments (ERGOM SED 1.0): coupling benthic and pelagic biogeochemistry of the south-western Baltic Sea, *Geoscientific Model Development*, 12, 275–320, <https://doi.org/10.5194/gmd-12-275-2019>, 2019.
- 540 Rutgersson, A., Kjellström, E., Haapala, J., Stendel, M., Danilovich, I., Drews, M., Jylhä, K., Kujala, P., Larsén, X. G., Halsnæs, K., Lehtonen, I., Luomaranta, A., Nilsson, E., Olsson, T., Särkkä, J., Tuomi, L., and Wasmund, N.: Natural hazards and extreme events in the Baltic Sea region, *Earth System Dynamics*, 13, 251–301, <https://doi.org/10.5194/esd-13-251-2022>, 2022.
- Ruvalcaba Baroni, I., Almroth-Rosell, E., Axell, L., Fredriksson, S. T., Hieronymus, J., Hieronymus, M., Brunnabend, S.-E., Gröger, M., Kuznetsov, I., Fransner, F., Hordoir, R., Falahat, S., and Arneborg, L.: Validation of the coupled physical–biogeochemical ocean model NEMO–SCOBI for the North Sea–Baltic Sea system, *Biogeosciences*, 21, 2087–2132, <https://doi.org/10.5194/bg-21-2087-2024>, 2024.
- 545 Schernewski, G., Friedland, R., Carstens, M., Hirt, U., Leujak, W., Nausch, G., Neumann, T., Petenati, T., Sagert, S., Wasmund, N., and von Weber, M.: Implementation of European marine policy: New water quality targets for German Baltic waters, *Marine Policy*, 51, 305–321, <https://doi.org/10.1016/j.marpol.2014.09.002>, 2015.
- Schernewski, G., Jekat, M., Kösters, F., Neumann, T., Steffen, S., and von Thenen, M.: Ecosystem Services Supporting Environmental Impact Assessments (EIAs): Assessments of Navigation Waterways Deepening Based on Data, Experts, and a 3D Ecosystem Model, *Land*, 13, <https://doi.org/10.3390/land13101653>, 2024.
- 550 Schernewski, G., Neumann, T., Piehl, S., and Swer, N. M.: Ecosystem-Model-Based Valuation of Ecosystem Services in a Baltic Lagoon: Long-Term Human Technical Interventions and Short-Term Variability, *Environments*, 12, <https://doi.org/10.3390/environments12020035>, 2025.
- 555 Seifert, T., Tauber, F., and Kayser, B.: Digital topography of the Baltic Sea [data set], <https://www.io-warnemuende.de/topography-of-the-baltic-sea.html>, last access: 27 July 2020, 2008.
- Vahtera, E., Conley, D. J., Gustafsson, B. G., Kuosa, H., Pitkänen, H., Savchuk, O. P., Tamminen, T., Viitasalo, M., Voss, M., Wasmund, N., and Wulff, F.: Internal ecosystem feedbacks enhance nitrogen-fixing cyanobacteria blooms and complicate management in the Baltic Sea, *Ambio*, 36, 186–194, 2007.
- 560 Vollenweider, R. A.: Advances in defining critical loading levels for phosphorus in lake eutrophication., <https://api.semanticscholar.org/CorpusID:131391058>, 1976.
- Vybernaite-Lubiene, I., Zilius, M., Bartoli, M., Petkuvienė, J., Zemlys, P., Magri, M., and Giordani, G.: Biogeochemical Budgets of Nutrients and Metabolism in the Curonian Lagoon (South East Baltic Sea): Spatial and Temporal Variations, *Water*, 14, <https://doi.org/10.3390/w14020164>, 2022.
- 565 Winton, M.: A Reformulated Three-Layer Sea Ice Model, *Journal of Atmospheric and Oceanic Technology*, 17, 525–531, [https://doi.org/10.1175/1520-0426\(2000\)017<0525:ARTLSI>2.0.CO;2](https://doi.org/10.1175/1520-0426(2000)017<0525:ARTLSI>2.0.CO;2), 2000.