

The Scenario Model Intercomparison Project for CMIP7 (ScenarioMIP-CMIP7)

Detlef P. Van Vuuren^{1,2}, Brian C. O'Neill³, Claudia Tebaldi³, Benjamin M. Sanderson⁴, Louise P. Chini⁵, Pierre
5 Friedlingstein^{6,7}, Tomoko Hasegawa^{8,9}, Keywan Riahi¹⁰, Govindasamy Bala¹¹, Nico Bauer¹², Veronika Eyring^{13,14},
Cheikh M.N. Fall¹⁵, Katja Frieler^{12,16}, Matthew J. Gidden^{3,10}, Laila K. Gohar¹⁷, Annika Högner¹⁰, Andrew D. Jones^{19,20},
Jarmo Kikstra¹⁰, Andrew King²¹, Reto Knutti²², Elmar Kriegler^{12,22}, Peter Lawrence²⁴, Chris Lennard²⁵, Jason Lowe¹⁷,
Camilla Mathison^{17,26}, Shahbaz Mehmood²⁷, Zebedee Nicholls^{28,10,29}, Luciana F. Prado³⁰, Qiang Zhang³¹, Steven K.
10 Rose³², Alex C. Ruane³³, Marit Sandstad⁴, Carl-Friedrich Schleussner^{10,34}, Roland Seferian³⁵, Jana Sillmann^{4,36}, Chris
Smith^{37,10}, Anna A. Sörensson^{38,39,40}, Swapna Panickal⁴¹, Kaoru Tachiiri⁴², Naomi Vaughan⁴³, Saritha
S.Vishwanathan^{9,44}, Tokuta Yokohata⁴⁵, Marco Zecchetto¹⁰, Tilo Ziehn⁴⁶

- 1 PBL Netherlands Environmental Assessment Agency, Bezuidenhoutseweg 30, The Hague, the Netherlands
- 2 Copernicus Institute of Sustainable Development, Utrecht University, Utrecht, The Netherlands
- 15 3 Joint Global Change Research Institute, Pacific Northwest National Laboratory and University of Maryland, College
Park, MD, USA
- 4 CICERO, Oslo, Norway
- 5 Department of Geographical Sciences, University of Maryland, College Park, MD 20742, USA
- 6 Faculty of Environment, Science and Economy, University of Exeter, Exeter EX4 4QF, UK
- 20 7 Laboratoire de Météorologie Dynamique, Institut Pierre-Simon Laplace, CNRS, Ecole Normale Supérieure, Université
PSL, Sorbonne Université, Ecole Polytechnique, Paris, France
- 8 Research Organization of Science and Technology, Ritsumeikan University, Kusatsu, Shiga, 525-8577, Japan
- 9 Department of Environmental Engineering, Kyoto University (Katsura campus), Kyoto, Japan
- 10 International Institute for Applied Systems Analysis (IIASA), Energy, Climate and Environment Program (ECE), A-
25 2361 Laxenburg, Austria
- 11 Center for Atmospheric and Oceanic Sciences, Indian Institute of Science, Bengaluru – 560 012, Karnataka, India
- 12 Potsdam Institute for Climate Impact Research, Member of the Leibniz Association, Potsdam, Germany
- 13 Deutsches Zentrum für Luft- und Raumfahrt (DLR), Institut für Physik der Atmosphäre, Oberpfaffenhofen, Germany
- 14 University of Bremen, Institute of Environmental Physics (IUP), Bremen, Germany
- 30 15 Laboratoire de Physique de l'Atmosphère et de l'Océan Siméon Fongang (LPAOSF), École Supérieure Polytechnique
(ESP), Univ. Cheikh Anta Diop, Dakar, Senegal
- 16 University of Potsdam, Institute for Environmental Science and Geography, Potsdam, Germany
- 17 Met Office · Hadley Centre, Exeter, UK.
- 18 Integrative Research Institute on Transformations of Human-Environment Systems (IRI THESys) and the Geography
35 Department, Humboldt-Universität zu Berlin, Berlin, Germany
- 18 Climate and Ecosystem Sciences Division, Lawrence Berkeley National Laboratory; Berkeley, CA, 94720, USA
- 19 Energy and Resources Group, University of CA, Berkeley; Berkeley, CA, 94720, USA
- 20 ARC Centre of Excellence for 21st Century Weather and School of Geography, Earth and Atmospheric Sciences,
University of Melbourne, Australia
- 40 21 Institute for Atmospheric and Climate Science, ETH Zurich, Switzerland
- 22 University of Potsdam, Faculty of Economics and Social Sciences, Potsdam, Germany
- 23 National Center for Atmospheric Research, Boulder, Colorado, USA

- 24 Climate System Analysis Group, University of Cape Town, South Africa.
25 School of Geography, University of Leeds
- 45 26 Global Climate-Change Impact Studies Centre (GCISC), Ministry of Climate Change & Environmental Coordination
(MoCC&EC), Govt. of Pakistan, Islamabad - 44000, Pakistan.
- 27 Climate Resource S, Berlin, Germany
28 School of Geography, Earth and Atmospheric Sciences, The University of Melbourne, Melbourne, Victoria, Australia
29 Faculty of Oceanography, Department of Physical Oceanography and Meteorology, Rio de Janeiro State University,
50 Campus Maracanã, Rio de Janeiro, Brazil
- 30 Department of Earth System Science, Tsinghua University, China
31 Energy Systems and Climate Analysis Research, EPRI, USA
32 National Aeronautics and Space Administration Goddard Institute for Space Studies, New York, USA 10709
33 Integrative Research Institute on Transformations of Human-Environment Systems (IRI THESys) and the Geography
55 Department, Humboldt-Universität zu Berlin, Berlin, Germany.
- 34 CNRM, Université de Toulouse, Météo-France/CNRS, Toulouse, France
35 University of Hamburg, Research Unit Sustainability and Climate Risks, Faculty of Mathematics, Informatics and
Natural Sciences, Germany
36 Department of Water and Climate, Vrije Universiteit Brussel, Belgium
- 60 37 Universidad de Buenos Aires. Facultad de Ciencias Exactas y Naturales. Buenos Aires, Argentina.
38 CONICET – Universidad de Buenos Aires. Centro de Investigaciones del Mar y la Atmósfera (CIMA). Buenos Aires,
Argentina.
39 Instituto Franco-Argentino de Estudios sobre el Clima y sus Impactos (IFAECI) – IRL 3351 – CNRSCONICET-IRD-
UBA. Buenos Aires, Argentina.
- 65 40 Indian Institute of Tropical Meteorology, Pune, India
41 Research Institute for Global Change, Japan Agency for Marine-Earth Science and Technology (JAMSTEC),
Yokohama, Japan
42 Tyndall Centre for Climate Change Research, School of Environmental Sciences, University of East Anglia, UK
43 Public Systems Group, Indian Institute of Management Ahmedabad, Ahmedabad, Gujarat, India
- 70 44 National Institute for Environmental Studies, Japan
45 CSIRO Environment, Aspendale, Australia

Correspondence to: detlef.vanvuuren@pbl.nl

75 **Abstract.** Scenarios serve as a critical tool in climate change analysis, enabling the exploration of future evolution of the
climate system, climate impacts, and the human system (including mitigation and adaptation actions). This paper describes the
scenario framework for ScenarioMIP as part of CMIP7. The design process has involved various rounds of interaction with
the research community and user groups at large. The proposal covers a set of scenarios exploring high levels of climate change
(to explore high-end climate risks), medium levels of climate change (anchored to current policy), and low levels of climate

change (aligned with current international agreements). These scenarios follow very different trajectories in terms of emissions, with some likely to experience peaks and subsequent declines in greenhouse gas concentrations in this century. An important innovation is that most scenarios are intended to be run, if possible, in emission-driven mode, providing a better representation of the Earth system uncertainty space. The proposal also includes plans for long-term extensions (up to 2500 AD) to study long-term impacts, climate change-related processes on long timescales, and (ir)reversibility. This proposal forms the basis for further implementation of the framework in terms of the derivation of emissions and land use pathways for use by Earth system models and additional variants for adaptation and mitigation studies.

1 Introduction

Scenarios serve as a critical tool in climate change analysis. Defined as plausible alternative descriptions of how the future may develop based on a coherent set of assumptions, they are used by different research communities to explore potential future avenues of socio-economic conditions, assess the effects of different drivers of climate change, characterize future climatic conditions, and assess impacts of climate change as well as adaptation and mitigation responses. As such, scenarios are also useful to bridge across different research communities. The World Climate Research Programme's (WCRP) Coupled Model Intercomparison Project (CMIP) has organized several rounds of the development and use of such scenarios. The CMIP scenarios play not only a pivotal role in climate research but also act as integrating tools for scientific assessment processes and policy analysis. In the most recent phase 6 (CMIP6), CMIP delegated the organization of experiments targeting specific research areas and questions to the scientific community, and CMIP6-Endorsed Model Intercomparison Projects were established (1). The Scenario Model Intercomparison Project (ScenarioMIP) formed a primary activity within CMIP6 that facilitated multi-model climate projections based on alternative plausible forcing scenarios that are directly relevant to societal concerns regarding climate change mitigation, adaptation, and future climate impacts and risks¹ (2, 3).

In this continuing role, ScenarioMIP's goal for CMIP7 is the design of a limited set of scenario-based experiments to be run by climate models, that 1) serve direct science questions, 2) function as input to other (science) communities and 3) support policy. With regard to the direct science contribution, the scenario information is used to study and understand climate processes and how their response to anthropogenic forcings emerges from internal variability and model structural uncertainties. To accomplish this aim, it is important that the set of forcing variables considered is internally consistent and varies over a plausible range of forcing levels, and that the set explores futures that may result in very different climate process dynamics (such as increasing and declining concentrations). With regard to the contribution to other (science) communities,

¹ In this paper, we are specifically concerned with those scenarios that are used as external forcings to climate models, including Earth System Models (ESMs), General Circulation Models (GCMs), Climate Models of Intermediate or Reduced Complexity (CMICs) and Simple Climate Models (SCMs). These external forcings encompass elements such as emissions and atmospheric concentrations of greenhouse gases, chemically reactive gases, aerosols, and land use.

ScenarioMIP ensures that data becomes available about future changes in climate variables (such as temperature, precipitation, humidity, etc.) together with information on human forcings/drivers (such as population, economic activity, land use, etc.) to a diverse set of user communities beyond the physical climate sciences. This aim is to facilitate further understanding of climate change, its impacts, risks and response options, including adaptation and mitigation choices. Targeted communities include, for instance, researchers on impacts and mitigation, but also practitioners, who might use this information for national or sub-national or local risk assessment, climate finance, mitigation policy or adaptation planning. Finally, regarding policy, ScenarioMIP aims to provide relevant outputs to support climate policy development, including through their use in IPCC assessments. Despite these important roles, climate modeling groups participating in ScenarioMIP can only run a limited set of scenarios. Computational expenses associated with setting up, running and archiving output from climate model experiments pose strict constraints on the number of scenarios that ScenarioMIP's protocol can include. Therefore, a small set of scenarios needs to be selected such that the set satisfies the three critical goals but does not require resources that are beyond reach for current scientific and computational capacity.

In preparation of the ScenarioMIP experimental design, the first meeting of the ScenarioMIP project under CMIP7 was held on June 20-22, 2023, in Reading, UK. Based on the meeting report, the Scientific Steering Committee (SSC) of ScenarioMIP formed several task groups, including external experts, and continued to work on an experimental design for the next round of ScenarioMIP. The results are captured in this document. The meeting also led to the expansion of the SSC for ScenarioMIP as well as the creation of a large advisory group(4). Both these changes responded to the desire to make ScenarioMIP more inclusive and broaden awareness of diverse viewpoints and concerns. Since then, the ScenarioMIP proposal was elaborated through various rounds of review and participation. In September 2023, the first ideas were shared with the research community in two webinars, asking for direct feedback. Subsequently, draft versions of the proposal were sent out for review in late 2023 (to a group of around 80 people worldwide that expressed interest in being involved as an advisory group) and in May 2024 (through an open review process to scientists and other user groups). Both rounds resulted in a large set of review comments (with traceable responses). The paper was also subject to public review as part of the EGU sphere review process. Development of IAM emissions and land use scenarios based on the experimental design described in this proposal began in September 2024 and is currently nearing completion. For each CMIP7 scenario, a marker was selected that accurately represented the description of the scenario. The results of this selection were discussed with representatives of the CMIP panel as well as potential scenario users. Earth system model simulations are planned to start in January 2026 for some scenarios and be completed about three months later. Additional scenario simulations are planned to begin in June 2026.

2 Overall experimental design

140 2.1 Role of ScenarioMIP in CMIP6

Before discussing the design of the current scenario round, it is useful to briefly evaluate the role of ScenarioMIP in CMIP6 (2, 3). In CMIP6, ScenarioMIP specified four Tier 1 and four Tier 2 scenarios to be run by ESMs, coordinating the production of these scenarios by the IAM community and the process to deliver them in the form that ESMs need to make use of them as external forcings. These experiments (especially those in Tier 1) were run by most modeling teams participating in CMIP6 and are by far the most used scenario-based simulations of CMIP6. The results of the ScenarioMIP experiments led to many physical science² papers describing changes in climate characteristics, but also to many papers characterizing the impacts of those changes on society and ecosystems. Further, ScenarioMIP results contributed to the assessment reports of all Working Groups of IPCC, supplying a dimension of integration that is reflected in the Synthesis Report of AR6 (5). The most direct use was in WGI, where ScenarioMIP climate model simulations formed the backbone of the assessment (6). The use in WGII (7) was more limited because of a lag in production of literature on impacts and adaptation based on ScenarioMIP results. In WGIII, ScenarioMIP results made an indirect but fundamental contribution via the calibration of simple climate models (SCMs) that allowed characterization of probabilistic global temperature projections and the resulting classification of a large set of baseline and mitigation scenarios produced by Integrated Assessment Models that were not run by ESMs (8, 9). There were some difficulties related to the process. This includes the relatively long period from data production (IAM model runs and harmonization) until final climate model runs (from 2015 to 2020). This means that studies on climate impacts (relying on ESM runs) could even start later given the need for additional downscaling and bias adjustment. This even meant that the underlying projections from IAMs were somewhat outdated by the time climate model output became available (several projected years (2015-2020) had already become history by that time). Also, over time, critiques emerged about the plausibility of the most extreme scenarios (SSP5-8.5 and its precursor, RCP8.5; SSP1-1.9) (10-12).

160

2.2 General design principles

In view of the multiple aims of the ScenarioMIP scenarios, the experimental design was created with a set of general principles in mind. The principles will be further discussed below:

- The scenarios form illustrative, internally consistent descriptions of future emissions and land use.
- The scenario set covers a wide and plausible range of emissions and climate forcing.
- Most scenarios will be run in emission-driven mode (for CO₂) in ESMs.
- The scenarios cover the period up to 2100 AD, and long-term extensions are stylized continuations of the narrative without large discontinuities.

² Various model types exist to explore the physical climate system, including Earth System Models (ESMs), General Circulation Models (GCMs), EMICs (Earth System Models of Intermediate Complexity) and SCMs (Simple Climate Models). Here, we use the term ESM to refer to all types of climate models.

170 *The scenarios form illustrative, internally consistent descriptions of future emissions and land use.*

The primary purpose of the ScenarioMIP scenarios is to provide emissions and land use pathways to drive ESMs. ScenarioMIP will produce these pathways based on plausible, internally consistent socio-economic and technological scenarios³. It should be noted, however, that there are many different scenarios that could produce similar emissions and land use outcomes. The particular storylines and quantitative drivers constituting the ScenarioMIP scenarios are therefore illustrative. In fact, the
175 resulting ESM simulations are intended to be used in future studies in combination with many different pathways describing future socioeconomic development (13). In this context, it is useful to note that even the regional patterns of emissions of short-lived forcers and land use, found to have effects within single model studies (14, 15) do not necessarily translate to a unique climate change pathway, since they, for now, have been shown to lack a robust regional climate signal in a multi-model context (16, 17).

180

In CMIP6, the Shared Socioeconomic Pathways (SSPs) provided the socio-economic and technological storylines and drivers underlying the emissions and land use pathways in ScenarioMIP (18). The SSPs continue to be in wide use and recently the demographic and economic drivers for these scenarios have been updated (19). However, other storylines and drivers could be adopted or created as a basis for the ScenarioMIP emissions and land use pathways. In practice, the IAM teams have based
185 their current scenarios on various SSPs, as it was generally deemed pragmatic as these come with already available, suitably rich quantifications and were implemented by the participating modelling teams within the given timeline.

The scenario set covers a wide and plausible range of emissions and climate forcing

The scenarios should encompass a wide range of policy-relevant emission trajectories considered to be plausible (i.e. that have
190 a non-negligible likelihood of occurring; see Box 1 for a definition of a plausible scenario and other related terms). As a set, the ScenarioMIP scenarios should thus cover plausible outcomes ranging from a high level of climate change (in the case of policy failure) to low levels of climate change resulting from stringent policies. For the 21st century, this range will be smaller than assessed before: on the high-end of the range, the CMIP6 high emission levels (quantified by SSP5-8.5) have become implausible, based on trends in the costs of renewables, the emergence of climate policy and recent emission trends (10). At
195 the low end, many CMIP6 emission trajectories have become inconsistent with observed trends during the 2020-2030 period.

Plausibility is a subjective judgment. Moreover, in several cases our plausibility judgments are conditional on assumptions that are themselves hypothetical. For example, in all scenarios we assume that there are no climate change impacts (see further

³ We use the term scenarios here for internally consistent, comprehensive descriptions of the future. The term pathways is sometimes used to refer to specific scenario elements (such as socio-economic development). We adopt this use of the term pathways here. In other literature, the term pathways is sometimes used to refer to goal-oriented scenarios or the terms scenarios and pathways are sometimes used interchangeably.

in this document). This can be justified by the overall scenarios framework in which impact analyses will be carried out by impact models using ScenarioMIP ESM simulations (and human drivers) as inputs. In other words, the ScenarioMIP scenarios are judged to be plausible conditional on the assumption of no climate change impacts. This conditionality is most consequential for the high climate change scenarios, in which impacts would be largest. The medium scenarios make subjective assumptions on current policies (see further). For low scenarios, we take into account geophysical and techno-economic limits, particularly regarding ramp-up rates of emission reduction and CDR, and technology and policy trends/constraints in the short-run. Views on plausibility evolve over time, as evidenced by the changing scenarios assumptions in successive IPCC WGIII assessment reports. It should therefore be acknowledged that there might be potential futures outside the ScenarioMIP scenario range.

Box 1: Terms characterizing scenario likelihood

There are several terms commonly used to describe the likelihood of a scenario. Likelihood is often expressed as the probability of occurrence of a selected event, e.g. a projected change lying in a given range (20). Applied to scenarios, a likely (or probable) scenario has a relatively high probability of occurring based on knowledge of current trends and common expectations about future developments and actions. In the ScenarioMIP experimental design, we focus on plausibility. A plausible scenario encompasses a range of outcomes that have a non-negligible likelihood of occurring (21). We do not quantitatively define the distinction between a plausible and implausible scenario in terms of a threshold likelihood (i.e., a “non-negligible” likelihood). Rather, we define it qualitatively by considering a scenario to be implausible if its likelihood is too low to be considered relevant to the purposes of the ScenarioMIP exercise. In the end, plausibility is a subjective judgment that can be based on several criteria. One way of making this distinction is that plausible scenarios should have a causal pathway to an outcome that aligns with current scientific understanding of natural and social processes (22). In some cases, further criteria can be adopted from the related concept of feasibility. Feasibility is defined as the potential for an action to be implemented (8) or as “doable under realistic assumptions” (22) and is most relevant for scenarios featuring strong climate policy or action. Therefore, when making judgments about the plausibility of mitigation scenarios we can borrow from approaches used in feasibility assessments; plausible scenarios should have a significant degree of feasibility. For example, the IPCC used multiple dimensions of feasibility – geophysical, technological, economic, sociocultural and institutional – to inform its judgments (22-26). Other approaches include historical analogies and mapping scenarios to a feasibility space (22).

Recent attempts have been made in the scenario literature to produce probabilistic emissions projections (27-29). While such studies can provide valuable insights, they require several methodological choices and subjective judgments that are not widely shared across the community. We therefore do not provide probabilities of occurrence for the scenarios of this experimental design. We only prescribe that all scenarios should be plausible (see Box 1). That said, the design criteria specify that the *High*

215 scenario should be "as high as plausible" while the *Very Low* scenario should be "as low as plausible," particularly in its degree of exceedance of 1.5°C warming. To meet these criteria, assumptions about one or more of the elements driving emissions outcomes in these scenarios will need to push the boundaries of plausibility. By construction, then, we anticipate that these scenarios will be considered less likely than those in the interior of the emissions range. However, we leave likelihood judgments as a research question to be addressed by the community on the basis of the emissions scenarios ultimately produced.

220 One aspect of the future also concerns elements of equity and justice. Given the focus of the ScenarioMIP scenarios to explore the broad, global relationships between the main drivers of climate change and the resulting global climate outcomes, the scenarios described here do not explicitly address a range of justice assumptions. However, it is critically important that the wider scenario literature, related to the ScenarioMIP scenarios, pays more attention to equity and justice issues. This is discussed further in Box 2.

225

Most scenarios will be run in emission-driven mode for CO₂

Up to now, ScenarioMIP experiments were driven by CO₂ concentrations. As this does not account for uncertainty in the carbon cycle response to climate, the decision was made for CMIP7 to run most simulations preferably in emission-driven mode (30). This means that for Earth System Models (ESMs), their carbon-cycle representation determines the concentration of CO₂ in the atmosphere that ensues from the prescribed emissions. This may lead to a wider concentration range as processes related to carbon cycle-climate feedbacks are newly involved in the scenario simulations, with their uncertainties. While the wider range may lead to some challenges in interpretation, it will better represent the uncertainty range resulting from both the carbon cycle and the climate system. The climate model simulations would also take better advantage of current ESM capabilities, especially regarding the outcomes of land-based mitigation solutions, which are heavily dependent on feedbacks that are not represented in concentration-driven experiments.

Concentration data will also be provided for all scenarios for ESMs that can only run in concentration-driven mode (without an active carbon cycle; see for a discussion of the current capabilities of state-of-the-art ESMs (31, 32). For the concentration-driven simulations, the median values of the concentrations as estimated by the carbon cycle emulators calibrated to CMIP6 included in Simple Climate Models (SCMs) can be used (33). This means that the concentration-driven models will likely have a narrower outcome space compared to the emission-driven set, which will have consequences for interpretation and use of certain variables. Regarding CDR options, only afforestation and reforestation will be based on endogenous representation of land use in ESMs. For all other CDR options, we will include their emission impact within the IAM emission output (see Section 5). To better assess the impact of running in emission-driven mode over the range of climate system outcomes produced by the multi-model ensemble, we propose that modeling groups adopting the emission-driven mode also run at least one scenario in concentration-driven mode, for comparison.

It is proposed that under the ScenarioMIP protocol models be run in emission-driven mode for CO₂ only. Other well-mixed greenhouse gases (CH₄, N₂O, halocarbons) remain with prescribed concentrations in the core experiments. Aerosol precursor emissions (e.g., SO₂, NO_x, VOCs, black carbon, organic carbon) will be provided for models with interactive aerosol schemes, though tropospheric and stratospheric 3d optical aerosol properties will be provided for those models which require it. Similarly, ozone concentrations will be provided for models which require this as an input (while models with full interactive chemistry can calculate it based on emissions).

Running historical simulations in an emission-driven configuration presents a challenge, as a model's simulated CO₂ concentration can drift from observations. However, this approach is a deliberate and core feature of the CMIP7 experimental design (30). Rather than taking measures to artificially correct this bias, we encourage ESM groups to quantify CO₂ concentration evolution as a key diagnostic of model performance and carbon cycle sensitivity. To facilitate this analysis, the ScenarioMIP protocol includes at least one parallel concentration-driven simulation specifically to allow for a clean diagnosis of the impact of these emergent carbon cycle feedbacks on climate outcomes.

The scenarios cover the period 2025-2100 (AD) with long-term extensions up to 2500 (AD)

The expectation is that the CMIP results will inform the process of the upcoming IPCC Seventh Assessment Report and the 2028 global stocktake as well as future research and assessments in subsequent years. As IAM narratives do not start until after 2025, ideally emissions would be harmonized until that point. However, historical emissions data for all relevant emissions species are not immediately available. Moreover, the historical emissions dataset used for harmonization must align with what is used for historical emissions in CMIP7. Emissions were therefore harmonized for the year 2023. For the 2024-2025 period, IAMs were asked to stay close to current trends – but it is based on individual model output. This is a major update compared to CMIP6 that had historical data up to 2015, and means that several critical periods (Covid pandemic, trends in natural gas use) will be included. IAM modelled emissions changes following the scenario narratives cover the period 2026-2100. It is also expected that differences among emission scenarios will remain within a relatively narrow plausibility range until 2030 (see Box 1 for a definition of plausibility).

There are important reasons to model long-term climate system dynamics beyond the end of the century, which serves the needs of communities studying long-term impacts and (ir)reversibilities over multi-century time scales (e.g., ice sheet, sea level rise, and species extinction researchers). The period after 2100 AD is covered by long-term extensions of emissions and land use to 2500 AD based on simple extension narratives and harmonization logic (See Section 4). We recognize that the period up to the end of the century, traditionally covered by future scenarios, becomes significantly shorter with every new phase of CMIP. Future phases will see IAM-based scenarios reach beyond 2100, but that technical development was not possible within the current timeline. Rather, for this iteration of ScenarioMIP, we request that the ESM models run all scenarios

at least to 2150 AD, using the 2100-2150 AD forcing from the extension protocol (see Section 4). In addition, teams are also requested to run at least two long-term extensions up to 2500 (preferably H-ext and VL-ext).

Box 2: The Role of Equity and Justice Considerations

285 Considerations of equity and justice are a cornerstone of the global climate discourse. These considerations are increasingly
also raised regarding how justice issues have been addressed in the scenarios used by the climate research community (e.g.
34, 35, 36). These authors highlight that in many scenarios used in the IPCC assessments, global inequalities in aspects of
the scenarios such as income and energy use persist through time and that equitable sharing of emissions reductions efforts,
290 carbon dioxide removal and adaptation implementation across countries is often insufficiently considered. While these
concerns are very important, the type of scenarios discussed in this paper do not address equity and justice explicitly. The
overall goal here is to explore the broad, global relationships between the main drivers of climate change and the resulting
climate outcomes, covering a wide range of plausible futures and connecting various research communities. ESMs using the
scenarios are sensitive to outcomes such as global emission levels and land use, but are not, when considered in a multi-
model ensemble, consistently responsive to underlying considerations related to equity and justice, which will strongly
295 affect the regional or sectoral distribution of these outcomes (see 37). So far, the multi-model differences in climate due to
different regional patterns of land use or emissions of short-lived forcers are still beset with uncertainty (see Section 2.2,
main text).

300 While the ScenarioMIP experiments described in this paper refrain from exploring equity and justice assumptions explicitly,
they include a regional differentiation of mitigation effort in the first few decades, motivated by plausibility considerations
based on the role various parties are currently playing. In addition, there are other features of IAMs that can have implications
for equity between regions, societal groups and generations (e.g. 38). For example, the timing of emissions reductions in IAMs
is influenced by the assumed discount rate (39), while also assumption on mitigation potential can have consequences for the
305 timing and location of mitigation efforts – with possible implications for equity.

The ScenarioMIP climate projections are expected to be used in combination with a range of socio-economic pathways. It is
important that questions related to equity and justice are addressed prominently in future research on mitigation and impact
analysis (as has been done to some degree in the past). Research on just transitions will complement the work of ScenarioMIP
310 by combining the climate outcomes (directly or based on climate emulators) with a range of socio-economic development and
policy information. We strongly recommend that both the mitigation and impact research communities consider a wide range
of socio-economic scenarios and policy assumptions that are consistent with equity and justice considerations in their research.
This can be seen as a subsequent phase of scenario development - both to inform policy making and assessment.

315 Another aspect of justice concerns the development process of the ScenarioMIP protocol (40). To enhance transparency and
inclusiveness, we have expanded the ScenarioMIP scientific steering committee to better represent relevant regions and
disciplines. We also established a broader advisory group consisting of over 80 people from different parts of the world.
Finally, the proposal, before this paper was submitted, underwent two rounds of review, one of which was public, to gather
extensive feedback. A last open review took place as part of the submission to this journal.

320 **2.3 Scenarios**


Based on the design principles and interactions with relevant communities, the following scenarios are proposed. Two
scenarios populate the high-end of the emission range to explore risks in case the world does very little to combat climate
change throughout the century and beyond (in one case) or until the second half of the century (in the second case). The

325 storylines for these scenarios include substantially slowing current observed trends towards rapid expansion of renewable
energy. A pair of medium scenarios explore the impact of current policies either remaining at current levels or, after a delay,
leading to a pathway towards net zero emissions. On the low side of the spectrum, three scenarios explore the temperature
range that has been associated with the Paris climate goals. In the full set, several scenarios explore overshoot dynamics at
high and low concentrations levels. This leads to the following proposal (see further details in Section 3):

- 330 • *High emission scenario*: A scenario with emissions as high as judged to be plausible, based on assuming developments
that include a rollback of current mitigation policies. This scenario is expected to result in forcings below SSP5-8.5.
- *High-to-Low emission scenario*: A scenario that follows approximately the same emissions pathway as the High, but
changes course in the second half of the century, applying strong mitigation measures to reach net zero CO₂ emissions
by 2100.
- 335 • *Medium emission scenario*: A middle scenario exploring consequences of extending current policies and trends into
the future.
- *Medium-to-Low emission scenario*: A scenario exploring a delayed increase in mitigation efforts, short of the Paris
temperature goal but achieving net-zero CO₂ emissions by the end of the century, with a period of net negative CO₂
emissions thereafter to achieve 1.5°C on a multi-century timescale.
- 340 • *Low emission scenario*: The Low emission scenario is designed to be consistent with the pursuit of holding warming
to a level likely below 2°C, without returning to 1.5°C before the end of the century. The scenario is extended beyond
2100 by an emissions trajectory leading to a slow decrease of warming afterwards.
- *Very Low emission scenario*: The *Very Low* emission scenario is designed to keep the temperature level as low as
plausible given feasibility constraints. This scenario is thus relevant for the low end of the Paris range (staying as
close as plausible to 1.5°C at the time of peak warming and limiting warming to 1.5°C by the end of the century).
- 345 • *Low-to-Negative emission scenario*: The last trajectory is a scenario with a higher overshoot of the 1.5°C goal,
followed by stringent climate policies resulting in net-negative greenhouse gas emissions to return to lower warming
levels, thus supporting research into the reversibility of climate outcomes and their impacts.

Note that all scenarios are named after emission trends. The global mean temperature associated with these scenarios will be
350 determined by ESM experiments and will form a range of outcomes. The temperature levels discussed above only guide the
scenario design and are being assessed during the design phase using SCMs. The overall logic of the current design shares key
characteristics with the proposal of Meinshausen et al. (2024). A summary of the scenario definitions is given in Table 1 while
Fig 1 illustrates the potential outcomes, based on preliminary results from IAMs and an SCM. In Sections 3, 4 and 5 we further
explore ideas and considerations relevant to the various scenarios.

355



360 **Figure 1: Proposed scenarios for CMIP7 ScenarioMIP, showing (a) GHG emissions pathways as a function of time for each of the proposed scenarios (based on GWP-100) and (b) the associated global average temperature outcomes using the probabilistic FaIR ensemble used in IPCC AR6 (41, 42). The shaded area for emissions shows ± 8 GtCO₂ around the expected marker scenario value while for temperature outcomes, it shows the 33-67 percentile range of the distribution for the same expected marker scenario. Data for the historical period can be found here (43). Scenarios are (H) *High*, (HL) *High-to-Low*, (M) *Medium*, (ML) *Medium-to-Low*, (L) *Low*, (LN) *Low-to-Negative* and (VL) *Very Low*. The final emission trajectories will depend on the finalized IAM runs but are expected to be roughly consistent with the illustrations provided here. The final temperature outcomes will be known only at completion of Earth system models' experiments, and will, for this phase of ScenarioMIP, include also the effects of carbon cycle feedbacks. Models are requested to run the scenarios for the period 2025-2150; the textured regions for the 2100-2150 period (AD) indicate that these emissions and forcings are defined in the extension protocol.**

365

The global average temperature trajectories shown in Fig 1 were produced using an ensemble of the FaIR simple climate model calibrated during the IPCC Sixth Assessment Report (AR6) (Smith et al., 2018; IPCC, 2021). FaIR is a reduced-complexity climate model useful for scenario assessment and idealized climate runs (for a full description (44)). Data in the figures is available at zenodo(45). It should be noted, however, that the final emission and forcing data will be produced by IAM model runs that will replace the stylized data shown here. The figures include the first 50 years of the extensions discussed in Section 4, since ESMS are requested to run scenarios for the period 2025-2150. The overall set of scenarios covers a plausible range of global emissions in 2100, from around -10 to -20 GtCO₂ per year up to levels of about a third greater than today's emissions. It is expected that the temperature results from the scenarios range from around 1.5 °C to almost 3.5 °C increase in 2100 over the 1850-1900 level.

370

375

380 **Table 1: Scenarios, naming, key design criteria, and their priority (expressed as Tier 1 and 2). These priorities are meant to be for models participating in emission-driven mode. We also request that these groups run, as part of Tier 1, the *Medium* scenario in concentration driven mode in order to distinguish the effects of the active carbon-cycle on the outcomes. Two more scenarios in concentration driven mode for these models are part of Tier 2. All the scenarios listed here will be provided also as concentrations for models that want to run ScenarioMIP in concentration mode, and for which Tier 1 will include all 7 scenarios.**

Scenario group	Scenario name	Primary emissions or temperature design criteria	Additional emissions or temperature characteristics	Tier	Scenario-ID
High scenarios, emissions-driven	High (H)	Emissions as high as plausible consistent with climate policy rollback	-	1	esm-scen7-h
	High-to-Low (HL)	High emissions until second half of century, followed by rapid decline to net zero CO2 in 2100	-	1	esm-scen7-hl
Medium scenarios, emissions-driven	Medium (M)	Emissions consistent with current policies frozen as of 2025	-	1	esm-scen7-m
	Medium-to-Low (ML)	Medium emissions until 2040 followed by gradual decline to net zero CO2 in 2100	-	1	esm-scen7-ml
Low scenarios, emissions-driven	Low (L)	Emissions consistent with staying likely below 2 °C and not returning to 1.5 C before the end of the century	Emissions decline toward net zero GHG emissions by 2100	1	esm-scen7-l
	Very Low (VL)	Emissions consistent with limiting warming to 1.5°C at the end of the century with overshoot as low as plausible	Rapid emissions decline toward net zero GHG emissions	1	esm-scen7-vl
	Low-to-Negative (LN)	Emissions consistent with limiting warming to 1.5 C at the end of the century with higher overshoot compared to the VL scenario	Rapid emissions decline begins later than in VL scenario and leads to net negative GHG emissions by 2100	1	esm-scen7-ln
Concentration-driven	<i>High, Concentration driven (HC)</i>	<i>Variant of H, concentration-driven for models that also run the emission-driven variant</i>		2	scen7-hc
	<i>Medium, Concentration driven (MC)</i>	<i>Variant of M, concentration-driven for models that also run the emission-driven variant</i>		1	scen7-mc
	<i>Low, concentration driven (LC)</i>	<i>Variant of L, concentration-driven for models that also run the emission-driven variant</i>		2	scen7-lc

385

2.4 Other design issues

Natural forcings (Solar and Volcanic)

The solar forcing recommendations for CMIP7 are provided by the SOLARIS-HEPPA group. The historical forcing reconstruction covers the period up to December 31, 2023, and is extended through December 31, 2024. For future simulations from January 1, 2025, through 2299, the data uses an intermediate scenario from a stochastic ensemble of future solar activity, which is based on a surrogate analysis of cosmogenic isotope datasets. The complete recommendations and dataset construction are described in Funke et al. (46) . The CMIP7 forcing task team has developed a new historical reconstruction

for stratospheric aerosol optical properties and volcanic sulfur emissions covering the period from January 1750 to December 2023 (47). The future volcanic forcing will involve a 9-year linear ramp-up from the historical values at the end of 2023 to a new background level. This background is defined as the mean stratospheric aerosol optical depth (SAOD) calculated over the 1850–2021 period (a value that results to be higher than the average between 2000 and 2020, a relatively quiescent volcanic eruption period). The SAOD will ramp up until 2033 and then hold steady at this piControl climatological value for the remainder of the future simulations. Notably, the major eruption of Hunga Tonga-Hunga Ha'apai in 2022 is deliberately excluded from the future forcing scenarios to avoid its large and anomalous cooling signal from complicating the analysis of the anthropogenic warming signal in the scenario runs. Full details of the methodology and dataset are forthcoming (47).

Air pollution control

All scenarios include air pollution representation (Short Lived Climate Forcers (SLCFs), among which aerosols including sulfur, black carbon (BC) and organic carbon (OC) emissions) and assumptions related to air quality measures. The effects of air quality measures on global radiative forcing can be significant (48). Despite lack of agreement among different ESMs over the response of regional climate to aerosol forcings, and a low signal to noise ratio of these responses, aerosol emissions have been observed to shape regional climate and will be one of the major drivers to influence climate change in coming decades (Persad et al., 2022). In the low scenarios, sulfur and aerosol emissions will be low as a result of the reduction in fossil fuel use. For the higher GHG emission scenarios, the emissions of air pollutants are also strongly influenced by specific air pollution control policies (as observed in many regions historically). In the High and High-to-low scenarios, the levels of air pollution control vary consistent with underlying assumptions on socio-economic development. Alternative scenarios could be run in AerChemMIP and RAMIP (49)

415 *Ensembles*

Recommendations on the use and size of initial condition ensembles are particularly relevant at the low end of the scenario range where the emergence of a climate signal relative to other low scenarios is expected to require relatively larger ensembles but are also important to enable sampling of longer return period events (rarer events) at all levels of forcing. For CMIP7, we encourage running ensembles of at least 5 members or more for each scenario ideally, to quantify the role of initial condition/internal variability uncertainty in scenario outcomes. If necessary, the three low scenarios could be prioritized, according to modeling centers' capacities. Sufficient representation of natural variability is decisive to inform climate risk assessments and adaptation decisions (50, 51).

IAM model runs

425 Based on the mostly qualitative formulation of the seven scenarios in this document, ScenarioMIP asked the IAM community to provide a database of alternative (i.e., ideally more than one) quantitative interpretations of each of the scenarios (see also Box 3). The Scenario Working Group of IAMC organized the planning, development, and vetting of the emissions and land

use scenarios, including the selection of single “marker” implementations of each scenario for provision to ESMs. This process and its results were nearly finished at the time of the final publication of this protocol and will be documented in further publications. The final set of IAM scenarios is fully consistent with the scenarios described in this paper. The full database of alternative scenarios will provide a basis for the broader IAM community to explore variants that alter key parameters, such as underlying socio-economic assumptions, climate policy and equity considerations and CDR use. These variants could also explore the implications of climate change impacts for the scenarios (see below under “Impacts and adaptation” for a discussion of this particular issue). Additional quantifications from models beyond those doing the initial quantification is strongly encouraged.

Box 3: Different emission reduction and carbon dioxide removal strategies

Emission reduction strategies can differ in timing, geographic location, the choice of technologies and other mitigation actions (including the use of CDR). Such differences in mitigation strategies form the main focus of mitigation research, including relationships with justice. Some of these differences also lead to different climate outcomes, including timing and overshoot. Several land-related mitigation options such as bioenergy with carbon capture and sequestration (BECCS) and afforestation may not only impact greenhouse gas concentration but also, for instance, albedo. These differences are also important for ScenarioMIP. This does also inform us about one element of the scenario matrix, i.e. the degree to which different socio-economic scenarios can be combined with all climate scenarios. Carbon dioxide removal strategies play a central role – as they can influence the timing of mitigation action, the ambition, but also land use. In this paper, for this reason we have separate section on the representation of CDR. It should be noted that solar radiation management is not included in these experiments, but is covered in a separate MIP (GeoMIP) (52). Novel CDR methods (such as ocean alkalinity enhancement) will be a focus in CDRMIP.

Impacts and adaptation

ScenarioMIP requests that the IAM teams produce simulations that do not include climate change impacts on society or ecosystems (e.g. on agriculture, energy use, economic growth, or biodiversity). There are two main reasons for this. First, one of the main uses of the scenarios and their climate outcomes is to drive impacts estimation by the impact modeling community, which uses both the climate projections and the direct human drivers (such as land use and agricultural systems changes) as input to their analyses. If the IAM scenarios (and therefore the climate projections based on them) already include impacts, further impact modeling based on these scenarios would lead to double counting. Second, IAMs currently do not represent a full range of potential impacts and generally lack the required detail needed to represent many regional impacted systems and adaptation strategies. Including impacts in the IAM scenarios would therefore only provide a partial and somewhat arbitrary accounting of potential climate effects. The IAM and ESM scenarios are therefore not intended to provide complete pictures of potential future worlds. Rather, they must be augmented by impact and adaptation studies that complete that picture so that it includes climate shifts, mitigation, impacts, adaptation, and development. At the same time, demand for fully consistent scenarios is growing. It is, therefore, encouraged that IAM developers undertake research projects to produce additional scenarios in which impacts are accounted for. This work may also lead to different scenario protocols for future ScenarioMIP exercises.

Modeling assumptions

The modeling paradigms and assumptions underlying IAM implementations relate to questions of socio-economic development, technological progress, mitigation preferences and climate justice. Exploring the implications of these assumptions, and alternative implementations, is of critical importance to provide policy relevant science to inform the deliberations on mitigation efforts and their regional distribution. These important questions, however, are outside the scope of ScenarioMIP, which is an exercise of CMIP that is focused on providing scenario forcing data for ESMs (see Box 2). The use of IAMs within ScenarioMIP is limited to providing emissions and land use forcing time series that allow for the exploration of different global climate futures. An exploration of alternative implementations of the ScenarioMIP scenario narratives using different modeling paradigms and normative assumptions is explicitly encouraged. Meanwhile, our working assumptions, from a climate modeling perspective, are that the regional distribution of carbon emissions do not matter to ESM outcomes, having long been characterized by their “well-mixed” nature, and therefore do not produce regional responses linked to the location of the sources of emissions. Regional distributions of land use, aerosols and other localized forcings have been shown to produce local responses that are, for now, model dependent and result in non-robust patterns of climatic changes in a multi-model context, such as CMIP-type ensemble experiments (16, 17).

480

The role of complex climate models vs emulators

Some further exploration is needed of the role of different tools at different levels of the modeling hierarchy, especially ESMs vs emulators of climate model output. By emulators we refer to computationally efficient tools that, when driven by a scenario, are able to provide impact-relevant variables akin to climate model output in spatial resolution and time frequency, bypassing the need to run ESMs. Emulators need to critically rely on available ESM ensembles over a wide range of scenarios for training. Their use can be attractive both to fill gaps in the design and to accelerate the uptake of some of the outcomes of new scenarios by the wider research community. Thus, it is useful to consider how emulators can further reduce the computational load on complex climate models for scenario exercises and the expectation is that, given the rapid developments in the emulation space of the last few years, especially with the deployment of machine learning and more broadly AI climate models, the use of emulators to augment or even substitute for ESM output may become more feasible in the not-so-distant future (53). As of now, however, no emulator can address the provision of all outputs from an ESM and for all types of scenarios. High frequency (e.g. daily) output, jointly simulated variables (respecting correlations between them), and more generally variables other than average temperature and precipitation still present a challenge to emulators. In the scenario space, overshoot/peak-and decline scenarios (intended here as temperature trajectories that exceed a global warming level of interest for a limited time and later experience a decline, due to declining GHG concentrations including from negative emissions) constitute particularly open questions, given the scarcity of this type of scenario simulation by ESMs on which emulators could be trained. The ScenarioMIP protocol therefore requests all scenarios to be run using ESMs. However, we expect emulator to be useful in filling the scenario space. Emulators may also be useful to boost ensemble sizes of existing ESM simulations.

ScenarioMIP will provide data as external forcing for Earth system model simulations and additional information on socio-economic development and related parameters for the Impacts, Adaptation and Vulnerability research community. Table 2 illustrates what type of data could be made available, although further parameters may also be provided. We note here that some variables, such as gridded population and urban land cover, are the result of dedicated efforts by the larger research community, rather than being a byproduct of IAM simulations.

Table 2: Input data for Earth System Model (ESMs) and Vulnerability, Impact and Adaptation (VIA) research communities (illustrative list)

	ESMs	VIA
Data provided during CMIP6	<ul style="list-style-type: none"> – CO₂ emissions (fossil + land use) + concentrations (harmonized with historical data) – Land cover change (harmonized with historical data) – CH₄, N₂O, CO, NO_x, H₂, VOC, SO₂, halogenated gases emissions and concentrations (harmonized with historical data) – Aerosol optical properties and ozone concentrations (based on run via atmospheric chemistry model driven by scenario emissions data) 	<ul style="list-style-type: none"> – Gridded population – Energy system parameters – Gridded land use/crop data (in addition to land cover) – Gridded water consumption and irrigation
Additional data	<ul style="list-style-type: none"> – Gridded urban land cover – Data on CDR activity (afforestation and reforestation areas; net-negative emissions) – Gridded water consumption – Fertilizer use – Crop yields – Gridded energy consumption 	<ul style="list-style-type: none"> – Gridded urban land cover – Income distribution and poverty/inequality – Fertilizer use – Crop yields – Gridded energy consumption – Air pollution

510 Further, the CMIP7 Forcings Task Team is in place to address some of these issues (required forcing input files, harmonization) and coordinate the provision of ESM forcings through the input4mips effort. This includes, for instance, harmonization of historical emission data and provision of consistent gridded land use data.

Consistency with earlier scenario sets

515 In CMIP6, one of the scenario design's stated goals was to facilitate comparison with CMIP5 and some studies were published
that quantified the relative contribution of different scenario composition versus different models to the changes in temperature
range under comparable global radiative forcing pathways. However, we believe that for the study of consistencies and
differences due to model development, the experiments prescribed as part of CMIP's Diagnostics, Evaluation and
Characterization of Klima (DECK) are more suitable. Therefore, no design choice was added to produce global forcings
520 comparable to CMIP6 scenarios.

2.5 Timeline

One of the goals of ScenarioMIP is to produce scenarios that can be useful to the 7th Assessment Report of IPCC. This means
that studies forming the basis of the assessment and relying on the new scenarios outcomes need to start appearing in the peer-
525 reviewed literature in the 2026-2027 time-frame. We recognize that although the IPCC and CMIP7 timelines will facilitate a
rigorous assessment of CMIP7 simulations by Working Group I, the research time available for the vulnerability, impacts and
adaptation (VIA) research using CMIP7 data is quite constrained. Therefore, several of the IAM scenarios based on this
proposal and the subsequent data steps to create input data for ESMs are intended to be completed around the end of 2025/early
2026, so that the climate model simulations can start early in 2026. This should allow for ESM experiments based on the Tier1
530 scenarios to start becoming available in mid-2026.

3 Further elaboration of the design of the emission scenarios for CMIP7

The ScenarioMIP scenarios are elaborated using IAMs. As indicated in Table 2, the key output data for ESM model runs
include emissions of CO₂, non-CO₂ greenhouse gases and air pollutants and land use. Subsequently, the emission data is
535 harmonized with historical data (43) (following similar methods as used in CMIP6 (54)). The output is run through small
climate models and an atmospheric chemistry model to provide concentration data for ESM that need such input. For land use,
the data is harmonised to a historical data set, again following a procedure similar to CMIP6, but now also including additional
information for CDR (see Section 5) (55). Below, further guidance is given to the elaboration of the individual scenarios and
their role in the overall set.

540

3.1 Design of the High emission scenario (H)

The *High* emission scenario explores a future world that weakens or even abandons mitigation actions and policies – in
combination with other development trends that could lead to high emissions. A scenario exploring the high-end of the
plausibility range is important for addressing questions such as: what are the impacts associated with a scenario in which

545 climate mitigation policy largely fails? What is the risk of reaching potential tipping points in the Earth system at a relatively
high level of future warming? How do extreme events look like at such warming levels? And how far beyond current conditions
are known adaptations viable?

The scenario is based on trends that may not be the most likely (based on a current assessment), but that are still plausible (see
550 Box 1). These trends can be characterized as policy roll-back, lack of cooperation in addressing global environmental concerns,
increased interest in fossil fuel resources, adoption of resource- and energy-intensive technologies and lifestyles and lack of
development in low-emission technology. Clearly, this scenario is not a “business-as-usual” scenario nor the no-policy
reference scenario for the other scenarios. The scenario is intended to explore the upper end of GHG emissions resulting from
deep political, technological, and structural deviation from current trends. There are various reasons why such a scenario could
555 emerge. For instance, a rollback of climate policies could result from a lack of public support for the energy transition. This
could be related to, for instance, local opposition to building new wind farms or concerns about impacts on fossil industries
related to jobs and national energy security. Also, the rapid cost decrease in renewable energy of the past decade could be
discontinued, possibly as a result of regional scarcity and limited tradability in materials for solar and wind technologies and
EV batteries (56, 57).

560 In terms of socio-economic development, a *High*-emission could be consistent with a range of pathways, including high
economic growth scenarios (e.g. SSP5) and regional competition scenarios (such as SSP3). In the development process for
CMIP7, IAM teams have explored several possibilities, showing that the emissions associated with SSP3 and SSP5 based
variants could be reasonably similar, with SSP5 often being slightly above SSP3 scenarios. The high challenges to adaptation
565 in SSP3 would make these variants more interesting from an impact perspective. In SSP3-based *High* scenario, resurgent
nationalism, concerns about competitiveness and security, and regional conflicts push countries to increasingly focus on
domestic or regional issues. Policies shift to becoming increasingly oriented toward national and regional security issues,
including barriers to trade. A low international priority for addressing climate concerns leads to a collapse of international and
national climate policies. This means that the SSP3 variant was preferred to the *High* emission scenario. For the *High-to-Low*
570 marker scenario (discussed below), instead the SSP5 scenario was used as the SSP5 storyline is more amendable for a shift in
policy trends. However, the exercise demonstrates that multiple development pathways can lead to high emissions. It is
important to realize that the climate outcomes resulting from the *High* scenarios can still be combined with different socio-
economic development pathways.

575 The *High* emission scenario, by design, does not include climate impacts (Section 2). This implies that the judgment of
plausibility of this scenario is conditional on this assumption. Another plausibility question relates to the volume of fossil fuel
reserves and resources. Clearly, the cumulative amount of fossil fuel use in the *High* emission scenario is considerably larger
than the estimated total reserves (known deposits that are extractable at current prices and technologies) (58, 59). However, it

is also considerably lower than total resources estimates (estimates of undiscovered deposits and/or those not recoverable at current prices) meaning that future technologies and price trends could make the resource trend possible.

3.2 Design of the High-to-Low emission scenario (HL)

The *High-to-Low* scenario represents a global emission trend that is similar to the *High* scenario up to the middle of the century, but that is followed by a deep reduction towards net zero CO₂ by 2100. Such a trend could result from climate policy based on observed impacts. The scenario can address questions like: What would the climate consequences be of substantially delayed mitigation after following a high emissions pathway? And, assuming that emissions of CO₂ become net negative beyond 2100 (which is part of the scenario extensions design, see Section 4), how large of a temperature overshoot might this lead to, and what would its consequences be for natural, managed, and human systems?

This scenario has similar considerations as the *High* scenario regarding the socio-economic pathway that could lead to high emissions and, in addition, must consider what type of pathway would also produce the capacity for rapid mitigation late in the century. The SSP5 pathway could provide a consistent story as it assumes rapid technological development and high economic growth. Based on this, the scenario would assume little effort to avoid global environmental concerns up to the second half of this century; after that increased concerns and technology development may lead to deep reductions. The high technological capacity for mitigation in this scenario, combined with strong international institutions that can support coordinated action, fosters rapid reductions to net zero CO₂ emissions by the end of the century.

3.3 Design of the Medium emission scenario (M)

The *Medium* emission scenario is a benchmark that shows the consequences of the current policy situation (as of 2025) and trends continuing over the century. In the scenario, it is assumed that policy effort is continued at the current level. As for other scenarios, also this scenario should not be considered as a “most likely” scenario. It can be used to address questions such as: what future physical, socio-economic, and ecological risks are implied by current levels of climate change policy (60, 61)? When comparing lower scenarios, what are the relative benefits and costs of taking further mitigation actions? What are the needs for adaptation implied by current policy levels? What are limits to adaptation if mitigation actions are not strengthened? A key assumption in the *Medium* scenario is that only policies are considered that are actually officially being implemented, similar to current policies scenarios in the literature (60). The scenario does not include future policy goals if not backed up (yet) by actual policy. This means that also the pledges of the Nationally Determined Contributions (NDC) and the net zero announcements are not included, if not backed up by explicit policies. As a result, the *Medium* scenario does not meet the aggregated NDCs expressed for 2030, as current policies are less ambitious.

In determining the impact of current policies, we assume that they stay in place during the announced period. For the period beyond this, the IAM community has developed several methods to extend current policies (typically beyond 2030) (62). Here,

it is assumed that current policies stay in place, while no new policies are accounted for. This does provide a clearer baseline against which the effect of new long-term policies can be evaluated in future studies. The scenario can be seen as an updated version of the reasoning from the “CurPol” scenario assumptions used in Working Group III of IPCC AR6 (8).

615 The marker implementation of this scenario is based on SSP2, with its middle of the road assumptions about socio-economic development. However, it would be interesting to explore these scenarios under a wider set of socio-economic assumptions. While we assume that climate policies remain at current levels, other policy areas could still develop. In addition, underlying technology assumptions are allowed to evolve, and the sensitivity of results to these assumptions should be assessed.

3.4 Design of the Medium to Low emission scenario (ML)

620 The *Medium to Low emission* scenario (see Fig.1) describes a future in which strengthened mitigation efforts are delayed, falling short of the levels required to meet the objectives of the Paris Agreement. As such, the scenario can be used to explore the impacts of temperature levels between the *Medium* and *Low* scenarios. Moreover, after 2100 AD, when the scenario’s extension transitions to net negative CO₂ emissions, the scenario explores an overshoot trajectory to assess the possibly related impacts. The underlying logic of the scenario is that political constraints limit rapid near-term action; however, these lessen over time as the magnitude of observed climate impacts increases and mitigation costs decline with technological progress. As a result, the scenario is expected to reach peak warming and net-zero CO₂ emissions at approximately the end of the century, with modest levels of net-negative CO₂ emissions thereafter aiming at the 1.5°C level on a multi-century time scale. It is based on the same socio-economic development pathway assumed for the *Medium* scenario. In terms of underlying socio-economic development, multiple SSPs could be possible. In the elaboration of the IAMs, SSP2 was mostly used.

630

3.5 Design of the Low emission scenario (L)

The *Low*, *Very Low* and *Low-to-Negative* emission scenarios are all aimed to explore relevant trajectories in the context of the Paris Agreement. In all three scenarios, emissions are reduced rapidly, but with different levels of ambition and with different timing. Specifically, in the *Low* scenario, the aim is to stay likely (>66% probability) below 2°C at all times, comparable to the C3 category of IPCC AR6 WGIII (9, 25). Questions related to this scenario are the associated climate impacts (vis-à-vis scenario with higher emissions, but also those with even lower emissions), as well as the required emission reduction measures.

640 In 2030, the emissions of the *Low* emission scenario will be similar to the current emission pledges as captured by the NDCs for 2030 (thus going beyond current policies). After that, emissions are projected to be reduced further and reach net-zero CO₂ emissions around 2070. Before 2070, some carbon dioxide removal (CDR) use might offset hard-to-abate emissions. After 2070, emissions reductions continue, reaching net-zero GHG emissions before 2100 and staying at approximately that level in the long term. This design helps to better understand the long-term climate implications of sustained net-zero GHG emissions.

To conform with the design specifications and plausibility criteria for short-term developments in 2030, carbon prices need to be regionally differentiated in 2030 (leading to higher taxes in high-income countries). After 2030, regional carbon prices should converge until they reach a globally uniform level in 2070. The exact elaboration is described further in the IAM papers related to ScenarioMIP.

3.5. Design of the Very Low emission scenario (VL)

At this point of time, some overshoot of the 1.5°C seems unavoidable (63.) The *Very Low (VL)* emission scenario is designed to keep climate change at the time of peak warming as low as can still be plausibly achieved (see Box 1 on plausibility) and to return warming below 1.5°C by the end of the century. Key questions therefore are to what level can an overshoot be constrained and how fast can warming be reduced after it peaks? As for other scenarios, the actual temperature trajectory will be determined by the ESM model runs.

Critical design elements of the VL scenario are reducing CO₂ emissions rapidly and deeply and reaching net-zero CO₂ emissions as quickly as possible. Also, non-CO₂ emissions are reduced deeply, including rapid CH₄ emissions reductions in the near-term to limit peak warming levels as much as plausible. After the point of net-zero CO₂ emissions, the pathway is designed to transition to sustained net-negative CO₂ emissions to increase the likelihood of limiting warming to 1.5°C in the second half of the century as computed by SCMs. The scenario also considers other Sustainable Development Goals (SDGs), such as protecting biodiversity. The scenario also has ambitious assumptions about air pollution controls. This is not to say that consideration of SDGs is unimportant in other scenarios, but that they may be of particular concern in the VL scenario, also given the possible concerns related to rapid emissions reduction in relation to development goals. IAM teams are strongly encouraged to explore this further in subsequent research on VL-type scenarios.

The VL scenario includes a range of measures and underlying trends leading to rapid emissions reductions, based on plausible assumptions about the underlying pace of the system transformations (see e.g. (23), general characteristics of low-carbon technology innovation (64, 65) and the dynamics of sociotechnical innovation (22). An important contribution to such very low emission pathway can also be formed by a shift towards low greenhouse gas emitting diets (e.g. the Lancet Planetary diet) (66-68). Rapid emission reductions also limit the deployment of CDR in the VL scenario (compared to the *Low-to-Negative, LN* scenario discussed in the next Section).

Plausibility considerations play an important role in the VL scenario. In the short-term (up to 2030), it should include ambitious but realistic reductions, also to make sure that the scenario can still represent a viable policy future in 2028-30 when the scenarios are assessed and used. Concretely, IAM teams were asked to make an assessment of “as low as plausible”, based on plausibility considerations with regard to how fast current technology and policy trends / constraints could change until 2030 taking into account stated policy objectives up to 2030 (including commitments beyond NDCs such as the Renewable Energy

and Energy Efficiency pledge, the deforestation pledge, the Global Methane Pledge, etc.). After 2030, the assumption is that policies can be sped up to limit peak warming and reach the long-term climate target of limiting warming to below 1.5°C. This ambition is bounded by considerations of techno-economic feasibility of low carbon technology deployment and, where relevant, sustainable development goals.

680

A range of emissions abatement measures (not exhaustive) are included in the VL scenario to reduce emissions: 1) reduction in final energy demand, 2) rapid decarbonization of electricity supply (as measured by carbon intensity of electricity based on gross CO₂ emissions), 3) deep electrification of industry, transport and buildings, 4) deep decarbonization of residual non-electric fuel mix in industry, transport and buildings, 5) widespread behavioral changes in diet, transportation and consumption, 6) deep reduction of industrial process emissions, including also reducing fluorinated greenhouse gases in line with the Kigali amendment, 7) deep reduction of non-CO₂ gases, in particular methane, 8) elimination of net CO₂ emissions from land use and rapid deployment of land-based CDR measures (within sustainability limits) to move to net-negative CO₂ emissions from land use in the medium to long term, and 9) deployment of CDR measures assuming plausible deployment rates, as well as keeping geological storage (or storage in materials) within technological and sustainability limits. Similar to the Low emission Scenario, also this scenario is run with differentiated carbon prices. The suggested convergence year, however, is 2050.

690

A number of particularly relevant scenario dimensions for ESMs were identified that could be explored in variants of the VL scenario in the future including: 1) land use and afforestation/reforestation policy, 2) land- and ocean-based CDR strategies, and 3) regionally defined renewable energy production. Future coupling on these dimensions between IAMs and ESMs (beyond this ScenarioMIP round) could permit improved linkages in research.

695

3.6 Design of the Low-to-Negative emission scenario (LN)

While the VL scenario aims to limit the overshoot of the 1.5°C warming level, an alternative pathway would include more overshoot. Looking into scenarios with more substantial overshoot and compare them with less overshoot is certainly relevant, also considering that as of today, global fossil-fuel related greenhouse gas emissions continue to rise (69). There are several key questions related to a scenario that aims to achieve the 1.5°C target, but with more overshoot. The first is related to how reversible the climate system is (or whether a lot of hysteresis occurs). This provides information on the viability of such an approach. Second, the scenario can facilitate the assessment of the impacts of a larger temperature overshoot, compared to the more limited overshoot. This includes understanding the benefits, costs, and trade-offs of achieving declining temperatures in the long term. This means that the scenario can be used to gain a better understanding of the near- and long-term consequences of delaying emission reductions. This will help inform ongoing policy discussions around plausibility and implications of overshoot resulting from delayed actions.


700

705

There are several considerations regarding the design of the *Low-to-Negative* (LN) scenario. First, the scenario should be based on plausible levels of CDR. To compensate for the overshoot, this pathway has much higher CDR levels than the *Very Low* (VL) pathway, but this needs to be within the assessed plausible range in the literature. Second, the emission trajectories and scenario temperature outcome of the *Low-to-Negative* scenario is still targeted to be consistent with the Paris climate objectives (thus limiting warming overshoot to well-below 2°C). Third, the scenario needs to be sufficiently different from other scenarios in ScenarioMIP, in terms of resolving differences between ESM runs. Such differences include the timing of the CO₂ emission profile and differences in emissions of short-lived climate forcers. Earlier, for CMIP6, a separation of 0.25-0.3 °C was proposed (70); in the current set, this difference might be somewhat smaller. As we suggest in Section 6, it would be useful to explore whether a smaller separation would still produce distinguishable outcomes in ESMs (71) also given the use of emission-driven scenarios (Mckenna et al., 2021). The *Very Low* and *Low-to-Negative* (LN) scenarios differ along dimensions in addition to peak warming, including policies targeting methane and those related to land use. The VL pathway also includes a sustainable land future in line with the SDG narrative, including reduced pressure from agricultural land and considering environmental constraints. In contrast, the LN scenario has very large scale and rapid upscaling of CDR in order to reduce warming after its higher peak. Finally, also this scenario is run with differentiated carbon prices (like the other mitigation scenarios). The suggested convergence year is 2070.

725 **4. Scenario extensions beyond 2100 AD**

Several climate science research communities expressed the desire to consider a set of scenario extensions going beyond the 21st century. The purpose of these extensions is twofold: to explore the long-term Earth System dynamics in response to warming at different stabilized levels, including the risk of breaching tipping points and triggering large scale irreversible changes, and also to explore reversibility of the system under different long term net-negative emissions pathways, exploring overshoots from different peak warming levels and different target long term climate states.



735 **Figure 2: Extensions for ScenarioMIP in CMIP7. Top row shows CO₂ emissions and cumulative emissions. Second row shows methane and sulfur dioxide emissions. Third row shows GHG emissions (calculated using GWP100) and total radiative forcing. Fourth row shows atmospheric CO₂ concentrations and projected temperature outcomes. Simulations are conducted with a calibrated ensemble of the FaIR simple climate model (41, 42), where solid lines show the 50th percentile outcome and shaded regions show the 5-95 percentile range. Data for the historical period is based on (43). Scenarios are (H) High, (HL) High-to-Low, (M) Medium, (ML) Medium-to-Low, (L) Low, (LN) Low-to-Negative and (VL) Very Low. The final emission trajectories will depend on**
 740 **the finalized IAM runs but are expected to be consistent with the illustrations provided here.**

Table 3: Main characteristics of the scenario extensions

	H-ext	HL-ext	M-ext	ML-ext	L-ext	VL-ext	LN-ext

	esm-scen7-h-ext	esm-scen7-hl-ext	esm-scen7-m-ext	esm-scen7-ml-ext	esm-scen7-l-ext	esm-scen7-vl-ext	esm-scen7-vl-ext
Tier	1	2	2	2	2	1	2
Purpose	Assessment of risk of large irreversible changes in slow components of the Earth system	Assessment of reversibility from large warming in the late 21st century to achieve 1.5°C on a multi-century timescale before 2500.	Assessment of long-term implications of current policy, including large overshoot and reversibility	Assessment of potential to stabilize temperatures at 1.5°C Paris target on a multicentury timescale given insufficient action for 2°C in 21st century	Assessment of potential to stabilize temperatures at 1.5°C if 21st century peak warming held is below 2C	Assessment of long-term response to net-zero GHG emissions, following highly ambitious 21st century mitigation	Assessment of reversibility following ambitious 21st century mitigation, with a target of climate restoration back to preindustrial temperatures
CO ₂ Storyline (for total combined fossil, industrial and AFOLU fluxes)	Constant total CO ₂ emissions plateau slightly above 2100 levels by 2120, remaining constant until 2180, after which there is linear reduction reaching net-zero CO ₂ by 2280 and remaining at this level thereafter.	Radical emissions reductions after 2070 in HL are continued into the 22nd century, reaching -22GtCO ₂ /yr net total CO ₂ flux by 2150. This level is maintained until 2200 and reduced to net-zero CO ₂ by 2300	CO ₂ emission reduction begins after 2100 with reductions to net zero CO ₂ by 2250.	Net zero CO ₂ is achieved in ML in 2100. In ML-ext, emissions reduction trends are maintained, reaching net-negative CO ₂ fluxes of -13GtCO ₂ by 2150. This level is maintained until 2250, and phased out to net-zero CO ₂ levels by 2300	Net CO ₂ emissions are maintained at 2100 levels of approximately -10GtCO ₂ /yr from 2100 until 2170, phased out by 2250.	Net CO ₂ emissions are maintained at approximately -3.5GtCO ₂ /yr in 2100 until 2450. This results in approximately net-zero GHG emissions using GWP100 for the period 2100-2450.	Net CO ₂ emissions reach a floor of approximately -25GtCO ₂ by 2100, and maintained at this level until 2200, with a phasing out of removals thereafter to achieve net-zero CO ₂ by 2300.

Note: All dates mentioned in the Table refer to AD.

745

As has been the case under the CMIP6 ScenarioMIP design, the scenario extensions will consist of emission and concentration trajectories to 2500 AD that are decoupled from the outcomes of IAM simulations. While IAMs are useful in generating the plausible evolution of greenhouse gas emissions in the shorter-term, beyond the end of this century the uncertainties that increasingly affect the socio-economic drivers of these trajectories end up limiting the usefulness of IAMs for scenario design.

750 The longer time-period of the extensions (relative to the 2300 AD time horizon in CMIP6) is proposed to allow for a simulation of climate stabilization at different warming levels, and to provide sufficient time to allow for a range of diverse overshoot trajectories. The period 2100-2150 covered by the extensions is meant to be run as a continuation of each 21st century scenario simulation in Tier 1.

755 We extend the seven marker scenarios (H, HL, M, ML, L, LN, VL) from 2101 to 2500 using differentiated methodologies tailored to the characteristics of each emission category. For fossil and industrial CO₂, a storyline-based approach has been applied combined with a functional representation of post-2100 emissions. This has been harmonized to produce continuous emissions trends during the transition and, after that, fit the storylines outlined in Table 3. The CO₂ emission trends described in the table are disaggregated by defining post-2100 AFOLU CO₂ emissions as a linear ramp-down approach from 2100 levels,

760 such that land use fluxes are reduced to have zero net effect on emissions post 2150. Non-CO₂ species (including air pollution)
are extended regionally, preserving spatial emission patterns from the IAM scenarios towards a global relaxation level
consistent with each species lowest emission value through the harmonized history and scenario trajectory. The extensions of
methane and sulfur emissions follow a similar logic for regional patterns, but tend toward scenario narrative specific global
relaxation targets. Several scenarios include CDR options (described in more detail in Section 5). The long-term extensions
765 are all designed to achieve temperature stabilization post-2300 AD. The rationale and proposed GHG emissions trajectories
for the extensions of the main scenarios (Fig. 2) are described below and summarized in Table 3.

The *High* (H) and *Very Low* (VL) scenario extensions are put in Tier 1. The *High* scenario extension (H-ext) simulates the
multi-century implications of the highest plausible emissions levels, while the *Very Low* extension (VL-ext) simulates a long-
770 term future relevant to the Paris agreement goal. The other scenarios are requested with lower priority and are part of Tier 2;
Each extension pathway is designed to maintain narrative consistency with each scenario's underlying socioeconomic and
technological assumptions. All scenarios (except VL-ext) ultimately reach net-zero CO₂ by 2300, which results in temperature
stabilization on multi-centennial timescales (26).

775 The *Very Low* (VL) scenario represents ambitious action to meet 1.5°C-consistent targets leading to net-zero greenhouse gas
emissions. Modest net negative CO₂ fluxes are maintained in the extension to compensate for emissions of other greenhouse
gases for the duration of the extension in the GWP100 calculation, and this results in a gradual, multi-century cooling. The
Low (L-ext), *Medium-to-Low* (ML-ext) and *High-to-Low* (HL-ext) extensions explore increasingly extreme overshoot
scenarios, requiring protracted periods of net CO₂ removal to achieve long term 1.5°C-consistent targets. Each scenario
780 continues reducing emissions trends in 2100, reaching significant net-negative CO₂ fluxes in the mid-22nd century (-9, -11
and -22 GtCO₂ respectively), which are phased out post-2200 to achieve net-zero CO₂ in the mid-23rd century. In FaIR
ensemble experiments, these three extensions each reach likely stable warming levels of 1.5°C by 2300 (Fig 2). A more extreme
overshoot is explored in LN-ext, which maintains very large net carbon removal rates to achieve net-zero *cumulative* CO₂ by
2300. This 'climate reversal' experiment explores the potential to return temperatures to near pre-industrial levels from 2300
785 onwards. Stabilization at higher warming levels is explored in the *Medium* (M-ext) and *High* (H-ext) scenarios. In each case,
the extension narrative describes an eventual reduction to net-zero CO₂ which allows the assessment of climate impacts at
higher, stable warming levels. As such, the *Medium* scenario describes little or no advancement from current policy until 2100,
but M-ext assumes that CO₂ emissions will be reduced to net-zero emissions throughout the 22nd century. H-ext maintains
emissions at slightly above 2100 levels until 2180, reducing emissions to net zero by 2250. These narratives result in estimates
790 of long-term stable warming levels of around 4 and 6°C above pre-industrial levels, respectively, for M-ext and H-ext.

The extension parameters are calibrated to maintain these narrative distinctions across the full 2101-2500 period. For AFOLU
emissions, all scenarios converge toward declining land-use emissions as agricultural intensification and ecosystem restoration


795 reduce deforestation and increase carbon sequestration, though at rates proportional to each scenario's overall ambition. Efforts
will be made to implement this assumption in the workflow for producing spatial land use emissions (LUH2) through a linear
phase-down of 2100 afforestation/deforestation rates to maintenance levels in 2150.

800 Non-CO₂ extensions similarly reflect scenario-specific assumptions: methane targets range from 95 Mt/yr in ambitious
scenarios to 520 Mt/yr in the *High* emission pathway, representing varying degrees of agricultural transformation and fossil
fuel phase-out; sulfur dioxide follows air quality policy stringency, declining to 10 Mt/yr in scenarios with strong
environmental governance while remaining higher in the high-emission cases. As is the case for the 21st century scenarios in
ScenarioMIP, emission-driven simulations are favored for the extensions to allow carbon-climate dynamics to be simulated in
the Earth System Models, with prescribed CO₂ emissions, prescribed land cover change, and prescribed non-CO₂
concentrations.

805

4.1 Geophysical plausibility of extensions

The extension methodology incorporates extended pathways for carbon dioxide removal (CDR) technologies beyond 2100.
Fig. 3 presents a comprehensive analysis of CO₂ fluxes including historical data (1850-2100) and future extensions (2101-
2500) for each scenario, displaying both annual gross emission and removal rates and cumulative totals.



810

Figure 3: Annual and cumulative CO₂ fluxes for extended climate scenarios (1850-2500). Left panels show annual emission rates, right panels show cumulative totals. Brown areas represent gross positive emissions from fossil fuel combustion and industrial

815 processes. Green areas show emissions and removals from agriculture, forestry and other land use (AFOLU). Yellow areas represent aggregated carbon dioxide removal (CDR) technologies including bioenergy with carbon capture and storage, direct air capture, ocean-based CDR, and enhanced weathering. Black lines indicate net emissions (positive values) or net removals (negative values). The red dashed vertical line marks 2100, separating historical data and original scenario projections (left) from extensions (right). Horizontal reference lines on cumulative plots (right panels) show: green line = prudent CDR storage limit (72), red solid line = proven fossil fuel reserves, red dashed line = proved plus probable fossil fuel reserves (73). Scenario codes: VL = *Very Low*, LN = *Low-to-Negative*, L = *Low*, ML = *Medium-to-Low*, M = *Medium*, H = *High*, HL = *High-to-Low* (see Table 1 for full scenario definitions).

820 Carbon dioxide removal encompasses multiple technologies including bioenergy with carbon capture and storage (BECCS), direct air capture with carbon storage (DACCS), ocean-based CDR, and enhanced weathering (see Section 5). In the extended scenarios, CDR deployment is scaled to maintain consistent 2100 baseline ratios. Gross unabated fossil fuel emissions are defined using functional forms which are continuous with 2100 values and continued according to the scenario narrative
825 (though all scenarios are defined to have zero unabated fossil fuel emissions by 2300). Required gross CDR fluxes are calculated as the difference between the net fossil fuel and industrial flux trajectory, and the gross positive fluxes.

This approach ensures technological deployment patterns remain consistent with the original scenario logic while extending to the end-of-millennium timeframe. The methodology preserves regional heterogeneity in CDR deployment, maintaining the
830 spatial distribution patterns established in the 2100 baseline year. This regional consistency is crucial for Earth system model implementations, as some CDR technologies have spatially dependent climate impacts and implementation constraints.

Fig. 3 shows both cumulative emissions and removals for each scenario in the context of geophysical constraints as assessed in the literature. The prudent cumulative CDR limit of -1460 Gt CO₂ estimated by Gidden et al. (72) represents an assessment
835 of potential limits of geologic carbon storage, reflecting both geological storage capacity and risk management considerations. This indicates that both the ML and HL scenario assumptions risk exceeding limits of prudent use of sequestration capacity.

Similarly, the cumulative emission analysis provides critical context by comparing net emissions trajectories with estimates of fossil fuel reserves. Proven fossil fuel reserves, 2866 Gt CO₂ (73), represent currently economically extractable resources,
840 while proved plus probable reserves (8032 Gt CO₂) include additionally discovered resources that may become economically viable. Cumulative emissions to 2300 in M-ext, HL-ext and H-ext all exceed proven reserves, while H-ext may also exceed proven plus probable reserves.

This consideration of geophysical limits to both fossil fuel availability and sequestration capacity underlines that the judgment
845 of the plausibility of the extended scenarios is limited to their geophysical dimension and is not as comprehensive as the plausibility judgments for the pre-2100 scenarios. In addition, some outlier scenarios challenge current estimates of geophysical feasibility. However, they are included for three reasons: first, assessments of geophysical feasibility are highly uncertain and are expected to evolve on the century timescale. Second, one of the goals of the extensions is to assess

850 nonlinearities in Earth System responses; as such, it is advantageous to include scenarios such as H-ext, which explore the upper limits of feasible fossil carbon emissions, and large overshoots such as LN-ext, which explore the upper feasible limits of removals. Third, extensions are designed to ultimately serve as narratives for interactive CDR deployment experiments in Earth System Models (for example, in CDRMIP), and scenarios in which the limits of CDR are challenged are useful for identifying where Earth System Models are unable to deliver the scenario negative emissions rates (for example if bioenergy yields in the ESM are lower than the IAM estimates).

855

5 Representation of carbon dioxide removal

Carbon dioxide removal (CDR) methods are an important component of climate action and mitigation plans and have a unique role in reducing greenhouse gas concentrations via their potential to enable net-negative emissions. How these methods are deployed will affect both land use and land management, as well as energy system compositions, impacting broader sustainable development and biodiversity considerations (Mace et al., 2021). Currently, a broad range of CDR methods is being discussed within the policy communities and considered as part of climate action plans. IAMs only represent a subset of these approaches. The main CDR methods represented in IAMs are Bioenergy with Carbon Capture and Storage (BECCS), Direct Air Capture with Carbon Storage (DACCS), and afforestation and reforestation. In addition, IAMs are exploring new CDR methods such as biochar, soil carbon sequestration, enhanced weathering, storage in long-lived materials, agroforestry, improved forest management, and ocean-based CDR, although only a subset of those are included in most ScenarioMIP scenarios. CDR methods will be investigated in ScenarioMIP scenarios, as well as within other related MIPs such as CDR-MIP and LUMIP.

The IAM land-use and emissions data that is provided to ESM teams, including data related to CDR, will be harmonized to ensure continuity with historical datasets and to ensure a consistent data format for all scenarios. Both IAMs and ESMs are expected to report gross as well as net emissions for each sector to help with analysis of each scenario. It is important to ensure a high level of consistency between the IAM and ESM models regarding land-use change and CDR activities. This requires increasing our understanding of the way relevant processes are implemented in various models. This also means that underlying information on drivers of land-use change (especially food production vs bioenergy crop production) will be reported. Below, we briefly discuss how the various CDR methods are implemented.

875

5.1 Reforestation and Afforestation

For reforestation and afforestation, the IAMs provide gridded representations of where they intend trees to be planted (and protected) to store carbon as vegetation biomass. Since ESMs typically have capabilities for modeling tree/forest growth and carbon storage, in ScenarioMIP the IAM information on land use and land-cover change and hence afforestation/reforestation

880 is forwarded (after harmonization) to the ESM frameworks (with land use representation) to create a process-based
 representation of the uptake and storage of carbon by these biomes in the ESMs. This will allow the calculation of the net
 emissions associated with reforestation and afforestation in ESMs that can be compared with the IAM information. However,
 since forest growth rates, forest management, potential biomass density as well as the representation of wildfires are likely to
 differ between individual ESMs and IAMs, there will still inevitably be differences between the carbon stored in trees/forests
 885 between these models (e.g. 74). In-depth comparison is possible if IAMs also report the carbon storage in a geographically
 explicit way, similar to the ‘secondary mean biomass density’ variable used in historical data for LUH3 (including information
 about the intended species and management practices used).

In the exchange between IAMs and ESMs, information on afforestation and reforestation will be provided as gridded areas of
 890 land-use for new forested areas in previously non-forested locations (afforestation) and/or previously forested areas
 (reforestation). This will be reported as two separate areas, along with existing forest areas. This information (from both IAMs
 and ESMs) will allow comparison of the differences in carbon storage from afforestation/reforestation between IAMs and
 ESMs and also will be used in biodiversity and impacts analysis. For ESMs, it should be noted that afforested and reforested
 area need to be represented as managed forests.

895

5.2 Bioenergy with Carbon Capture and Storage (BECCS)

IAMs typically include a detailed description of bioenergy and BECCS. This includes the representation of a variety of
 feedstocks, including bioenergy crops, forest biomass, and agricultural/forest residues. Moreover, in IAMs bio-energy yields
 are typically improving over time resulting from research, development, and innovation. This means that the carbon content
 900 for a given plot of bioenergy crop can increase over time and that a smaller area is needed to produce the same amount of bio-
 energy (or CDR).

Table 4: Results of a survey among ESMs participating in CMIP7 on their capabilities of BECCS representation

BECCS processes needed within ESMs	Percentage of ESMs ready to compute/model these processes (out of 19 ESM teams responding)
BECCS-specific crop types	26%
Technological yield improvements	42%
Anthropogenic carbon storage pools	47%
Bioenergy generation	26%

Efficiency of carbon capture at bioenergy plants	11%
Geological carbon storage	33%

905 Table 4 (based on a recent survey of ESMs in preparation for CMIP7) summarizes the current readiness of ESMs to model CDR processes, including BECCS. Based on this survey, it was concluded that most ESMs are not yet ready to simulate the specific crop types used for BECCS and to include the bioenergy crop management and harvest schemes needed to compute the associated emissions and storage. Therefore, it has been decided that for BECCS, the carbon storage and/or net emissions from BECCS will be provided to ESMs as part of the emissions data from IAMs (rather than computing these emissions in the
910 ESMs). Still, it is useful to try and harmonize as much as possible the representation of bio-energy and BECCS in the two model types.

ESMs will still need to model the emissions and biogeophysical climate impacts associated with land-use change and management for bioenergy (and other) crops. To facilitate this, they will be provided with harmonized gridded areas of
915 bioenergy crops from IAMs. However, since many ESMs are actually preparing to fully calculate the net BECCS emissions as well, as much information as possible will be transferred from IAMs to ESMs so that those analyses and experiments can be undertaken in separate MIPs or research projects that will take place over longer time periods than ScenarioMIP. In principle, the results of the ESM BECCS calculations can feed into future versions of IAMs to highlight areas of highest BECCS carbon uptake potential.

920 To relay key information around BECCS to ESMs, IAMs will report the land-use change areas associated with first and second-generation bioenergy crop deployment at the gridded level. Also irrigation and fertilizer usage associated with bioenergy crops will be provided. These data will enable ESMs to model the climate impacts of land-use change associated with increasing or decreasing areas of bioenergy crops, along with the climate impacts of managing these agricultural systems. Regional BECCS-
925 related carbon removals will also be gridded and reported to enable ESMs to capture the effects of the emissions from bioenergy that replace other emissions in the energy system and the emissions. Essentially, the ESMs will use the harmonized land-use data to calculate emissions associated with changes in land carbon pools and they will be provided with IAM data on emissions (including the reductions associated with BECCS carbon storage). Table 5 further summarizes the AFOLU processes associated with BECCS and whether the emissions/storage for each process will be provided by IAMs or computed by ESMs
930 in ScenarioMIP.

In comparing the IAM and ESM output, it is important to realize that IAMs models report the impact of BECCS in the region in which the bioenergy is used, not where the biomass is grown. At the same time, IAMs report the biomass growth in the original location. Therefore, biomass growth as a driver of carbon dioxide removal will be consistent with the underlying IAM,

935 but emissions flux field could be inconsistent because of biomass trade. However, since CO₂ is a well-mixed gas, for emissions fluxes, the importance of the spatial pattern is not very important..

Table 5: the role of IAMs and ESMs in representing individual processes associated with BECCS.

BECCS sub-processes	Emissions/storage computed by ESM or provided by IAM for ScenarioMIP
Land-use change associated with BECCS	ESM
Management of land used for BECCS	ESM
Emissions from bioenergy production	IAM
Storage of carbon from CCS	IAM

940 **5.3 Direct Air Capture with Carbon Storage (DACCS) and other forms of CDR**

Outputs from IAMs for other CDR methods (such as DACCS, enhanced rock weathering, biochar, soil carbon management, and ocean-based CDR) will be provided by IAMs as part of the total emissions data as well. Work is ongoing to provide gridded DACCS and “Other CDR” flows as an output from IAMs for future simulations and studies outside of ScenarioMIP. In some cases, IAMs may also model carbon removals related to agroforestry and improved forest management. As these

945 flows relate closely to the woody carbon stocks included in the harmonized land-use forcing product, they are not included in the emissions data – similar to other land-use CO₂ emissions. However, since the harmonized land-use forcing product, and ESMs generally do not model the processes related to CDR methods such as enhanced rock weathering, biochar, and soil carbon management, they are provided as part of the net emissions flux.

950 **5.4 Overall information flows on CDR**

Table 6 summarizes the data passed between IAMs and ESMs for CDR. In addition to the required datasets that are passed from IAMs to ESMs, as much additional information as possible on the IAM intentions and assumptions associated with CDR should be provided for additional analysis by ESMs and impact models. Moving forward, our goal is for IAMs and ESMs to work towards enabling full calculations of land-based CDR emissions/storage within ESMs for future MIPs and research

955 projects.

Table 6. CDR data passed from IAMs to ESMs annually as part of ScenarioMIP in CMIP7, along with intended ESM usage of these data.

CDR Method	IAM data passed to ESMs (via harmonization)	ESM use of this data
BECCS	Gridded areas of first- and second-generation bioenergy crops, regional net emissions associated with BECCS, regional carbon storage associated with BECCS.	Compute emissions associated with land-use change and management of bioenergy crops but do not remove and store carbon. Use BECCS net emissions or BECCS storage (depending on ESM needs), together with other emissions data inputs.
Afforestation/Reforestation	Gridded areas of afforestation and reforestation.	Plant, grow, and protect trees. Compute carbon emissions and storage associated with tree growth.
Other CDR, i.e. DACCS, enhanced weathering, ocean-based CDR, biochar, soil carbon sequestration, and other non-land CDR (where available)	Gridded net emissions associated with CDR methods included as part of the total emissions provided to ESMs.	Use along with other emissions data.

960 **6. Discussion and conclusions**

We have proposed a limited set of scenarios that, modeled by IAMs, would provide pathways of emissions, concentrations, and land use for use as external forcings for ESMs participating in CMIP7. These scenarios with their climate system outcomes are intended to facilitate studies of climate processes; provide inputs to a wide range of studies of impacts, adaptation, and mitigation; and support climate policy development and communication. They cover a wide range of plausible outcomes, from a *High* scenario representing policy failure and strong drivers of greenhouse gas emissions, to a *Very Low* scenario with dramatic emissions reductions.

The implementation of these scenarios in IAMs and ESMs poses a number of challenges, not all of which can be addressed or fully explored within the ScenarioMIP process. As a result, in a number of cases it became necessary to make design choices on the basis of current literature. Throughout the article, we have indicated where further research could advance the field and summarize important directions here.

Produce wider set of IAM scenarios:

- Produce idealized or counterfactual emissions and land use scenarios to complement the plausible scenarios proposed here. The ScenarioMIP design is limited to scenarios judged to be plausible, but other scenarios could also provide useful insight. For example, scenarios that are higher or lower than those proposed here, or that imagine counterfactual historical outcomes (Meinshausen et al., 2024), can be of scientific interest or policy relevant.
- Develop alternative implementations of the scenarios requested here from the IAM community using different modeling paradigms and normative assumptions. There are a number of questions of wide interest to the scientific

and policy communities that would be unlikely to imply forcing pathways different enough to substantially affect climate system outcomes, either regionally or globally. They therefore fall outside the scope of ScenarioMIP but are worth pursuing through other means. For example, different normative bases for approaches to mitigation may lead to different regional or sectoral emissions pathways, within approximately the same global forcing pathway. Similarly, alternative socio-economic pathways such as sustainable development scenarios with rapid convergence of regional incomes and living standards are useful to pursue but, if coupled with a target of limiting warming to (for example) likely below 2°C, would lead to similar global forcing pathways to other 2°C scenarios.

- Produce versions of the scenarios described here that account for climate change impacts. The SSP-RCP scenario framework by design does not include impacts in community scenarios. This enables an independent assessment of impact-climate relationships conditional on a given SSP in the impact research community. However, incorporating impacts in these scenarios would also be of interest, in particular to test whether this would substantially alter global emissions and land use pathways.
- Consider options to better integrate adaptation and mitigation. There are several interactions between impacts, adaptation and mitigation. At the moment, these are not considered given the way information is forwarded from one community to another. It will become increasingly important to see how this issue can be overcome.
- Explore methods for further characterizing the relative likelihood of different scenarios. In this proposal we have limited likelihood judgments to whether scenarios are plausible or not, with some suggestion that the H and VL scenarios may be considered less likely than the others, by construction. It will be worth undertaking a variety of approaches to judge the relative likelihood of the various scenarios more broadly to help inform risk assessments.

Explore climate (model) processes

- Explore the effect of carbon cycle uncertainty, including climate feedbacks, on climate system outcomes. ESMs are encouraged to run all scenarios in CO₂ emissions-driven mode, so that the carbon cycle is modeled within the ESMs. The multi-model ensemble will then capture uncertainty in the carbon cycle across models. A complementary analysis could test uncertainties in the carbon cycle parameters in the emulators (derived from the behavior of the CMIP6 ESMs) by deriving high and low CO₂ concentration pathways for a given emissions scenario using emulators. These concentration variants could then be run with ESMs (in concentration-driven mode) in addition to the median concentration pathway recommended as part of the ScenarioMIP protocol, to investigate the sensitivity of the climate system to the emulators' implementation.
- Explore the effect of atmospheric chemistry uncertainty on climate system outcomes. The ScenarioMIP design calls for non-CO₂ greenhouse gases and air pollutants to be incorporated primarily in the form of atmospheric concentrations, with a single atmospheric chemistry model providing concentration fields based on the emissions scenarios. A fuller representation of uncertainty would run emissions through additional atmospheric chemistry

1015 models and then run through ESMs to assess the sensitivity of climate outcomes to the representation of atmospheric chemistry.

Explore climate model sensitivities:

- 1020 • Further explore whether different mitigation strategies achieving approximately the same global forcing pathways would produce detectable differences in climate outcomes. The overall SSP-RCP scenario framework is based on the premise that in a multi-model setting ESM projections resulting from a given pathway of global average forcing can be considered consistent with alternative emissions and land use scenarios producing a similar global forcing pathway (van Vuuren et al., 2017). In that way, ESM simulations based on one of the ScenarioMIP scenarios can be used in studies with alternative assumptions about socio-economic pathways or mitigation strategies, as long as global forcing is similar. Limited existing research supports this assumption, but further study of the degree to which variations in regional forcing due to short-lived climate forcings or land use could produce significantly different climate and impact outcomes would be useful.
- 1025 • Further explore what minimum difference in global forcing is required to produce significantly different climate system outcomes. Global forcing pathways that are not sufficiently separated from each other may not produce multi-model ESM ensembles with statistically significant differences in important climate variables. It may be that a forcing difference sufficient to produce a difference of 0.25-0.3°C global average temperature is required (Tebaldi et al., 2015), but further analysis with the latest generation of ESMs is desirable.

1030

Foster development of ESM emulators:

- 1035 • Develop spatially explicit, multi-variable emulators that can emulate ESM outcomes, especially for temperature overshoot scenarios. Progress is being made in the development of ESM emulators, but currently there are not enough overshoot scenarios run by ESMs to train these emulators and test their accuracy in overshoot conditions. Emulators can subsequently be used to explore the outcomes of a much wider range of scenarios and can help better quantify uncertainties, even facilitating a probabilistic representation of climate outcomes.

1040 ***Improve ESM representation of mitigation strategies:***

- 1045 • Develop and explore methods for ESMs to be able to compute CDR-related emissions and storage (including BECCS). Because the implementation of CDR directly in ESMs is not yet a widespread practice, the ScenarioMIP design calls for IAMs to estimate the emissions associated with these measures and pass these results to ESMs. In future CMIP activities it would be desirable to implement CDR measures directly in ESMs. This requires research on what CDR related output is important for IAMs to provide, and how ESMs can implement it in a way that is consistent with the original IAM intentions.

2 Appendices

1050 Appendix A: Acronyms

	AerChemMIP	Aerosol Chemistry Model Intercomparison Project
	AFOLU	Agriculture, Forestry and Other Land Use
	BECCS	Bioenergy with Carbon Capture and Storage
	CCS	Carbon Capture and Storage
1055	CMIC	Climate Model of Intermediate Complexity
	CMIP	Coupled Model Intercomparison Project
	CDR	Carbon Dioxide Removal
	DAC	Direct Air Capture
	DACCS	Direct Air Capture with Carbon Storage
1060	DECK	Diagnostics, Evaluation and Characterization of Klima
	ESM	Earth System Model
	EV	Electric Vehicle
	GCM	Global Circulation Model/Global Climate Model
	GHG	Green-house gas
1065	GMST	Global Mean Surface Temperature
	GSAT	Global-mean Surface Air Temperature
	GWP100	Global Warming Potential over 100 years
	H	High scenario
	H-ext	High extension
1070	H-ext-OS	High overshoot extension
	IAM	Integrated Assessment Model
	IAMC	Integrated Assessment Modeling Consortium
	IEA	International Energy Agency
	input4mip	CMIP activity tasked with processing and availability of input data for ESM experiments under CMIP
1075	IPCC	Intergovernmental Panel on Climate Change
	L	Low scenario
	L-ext	Low extension

	LN	Low to negative emission scenario
	LN-ext	Low to negative emission scenario extension
1080	LUMIP	Land Use Model Intercomparison Project
	M	Medium scenario
	M-ext	Medium extension
	ML	Medium-to-Low scenario
	ML-ext	Medium-to-Low extension
1085	MIP	Model Intercomparison Project
	NDC	Nationally Determined Contributions
	RAMIP	Regional Aerosol Model intercomparison Project
	RCP	Representative Concentration Pathway
	SCM	Simple Climate Model
1090	SDG	Sustainable Development Goal
	SLCF	Short-Lived Climate Forcer
	SSC	Scientific Steering Committee
	SSP	Shared Socio-economic Pathways
	TCRE	Transient Climate Response to cumulative Emissions
1095	VIA	Vulnerability, Impacts and Adaptation
	VIACCs	Vulnerability, Impacts, Adaptation and Climate Services
	VL	Very Low emission scenario
	VL-ext	Very Low emission scenario extension
	WGI/II/III	Working Group I/II/III.

1100 **Code, data, or code and data availability**

Data on the figures is available at zenodo: DOI: 10.5281/zenodo.14382495

Author contributions

1105 DvV, BCO and CT coordinated the development process of the manuscript and took a lead in the writing. LC, TH, KR, PF and BMS coordinated subgroups in the writing process that worked on particular sections. BS and CS created the figures. All authors contributed to the design and the writing of the proposal. Competing interests
At least one of the authors is a member of the editorial board of GMD.

Competing interests

At least one of the authors is a member of the editorial board of GMD.

1110 Acknowledgements

This proposal and the scenario development process was endorsed by WCRP JSC. We would like to thank Helene Hewitt, John Dunne and Eleanor O'Rourke for their support in developing this proposal, and Aleh Cherp for valuable input.

Financial support

DVV is supported by the EERIE project (Grant Agreement No 101081383) and the ELEVATE project (under grant agreement
1115 No 101056873) funded by the European Union's Horizon Europe programme. NB is supported by the RESCUE and
OptimESM projects funded by the European Union's Horizon Europe programme under grant number 101056939 and
101081193. JS is funded by the German Research Foundation (DFG) under Germany's Excellence Strategy— EXC
2037: "CLICCS—Climate, Climatic Change, and Society"—Project Number: 390683824, contribution to the Center
for Earth System Research and Sustainability (CEN) at the University of Hamburg. LFP was supported by National Council
1120 for Scientific and Technological Development (CNPq grant number 405607/2023-7) and Carlos Chagas Foundation for
Research Support of the State of Rio de Janeiro (FAPERJ grant number E-26/210.291/2024). CM, LKG, CS and JL were
supported by the Met Office Hadley Centre Climate Programme funded by DSIT. LPC was supported by the NASA Carbon
Monitoring System (grant number 80NSSC21K1059) and the University of Maryland Grand Challenges program. CS and JS
were supported by the European Union's Horizon Europe programme project WorldTrans under grant number 101081661.
1125 V.E.'s work is supported by the European Research Council (ERC) Synergy Grant 'Understanding and Modeling the Earth
System with Machine Learning' (USMILE) under the Horizon 2020 Research and Innovation programme (grant agreement
no. 855187) and the Deutsche 1079 Forschungsgemeinschaft (DFG, German Research Foundation) through the Gottfried
Wilhelm Leibniz Prize awarded to V.E. (reference no. EY 22/2-1). CM was supported by both OptimESM and the TipESM
projects funded by the European Union's Horizon Europe research and innovation program under grant agreement no.
1130 101081193 and 101137673 and is therefore funded by UK Research and Innovation (UKRI) under the UK government's
Horizon Europe funding Guarantee. BMS and MS were supported by the Norwegian Research Council through the project
TRIFECTA (grant number 334811) and the European Union Horizon Europe Research and Innovation programme projects
DIAMOND (grant-no. 101081179) and ESM2025 (grant-no. 101003536).

1135 References

1. Eyring V, Bony S, Meehl GA, Senior CA, Stevens B, Stouffer RJ, et al. Overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6) experimental design and organization. *Geoscientific Model Development*. 2016;9(5):1937-58.
2. O'Neill BC, Tebaldi C, Van Vuuren DP, Eyring V, Friedlingstein P, Hurtt G, et al. The Scenario Model Intercomparison Project (ScenarioMIP) for CMIP6. *Geoscientific Model Development*. 2016;9(9):3461-82.
- 1140 3. Tebaldi C, Debeire K, Eyring V, Fischer E, Fyfe J, Friedlingstein P, et al. Climate model projections from the Scenario Model Intercomparison Project (ScenarioMIP) of CMIP6. *Earth System Dynamics*. 2021;12(1):253-93.
4. CMIP. ScenarioMIP 2026 [Available from: <https://wcrp-cmip.org/model-intercomparison-projects-mips/scenariomip/>].
- 1145 5. IPCC. AR6 Synthesis Report: Climate Change 2023. Cambridge; 2023.
6. IPCC. Climate Change 2021 - The Physical Science Basis. Summary for Policymakers. Working Group I contribution to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change; 2021.
7. IPCC. Climate Change 2022: Impacts, Adaptation and Vulnerability. Contribution of Working Group II to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change Cambridge, UK 2022.
- 1150 8. IPCC. Climate change - Mitigation of Climate Change. Contribution of Working Group III to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge: Cambridge University Press; 2022.
9. Kikstra JS, Nicholls ZRJ, Smith CJ, Lewis J, Lamboll RD, Byers E, et al. The IPCC Sixth Assessment Report WGIII climate assessment of mitigation pathways: from emissions to global temperatures. *Geoscientific Model Development*. 2022;15(24):9075-109.
- 1155 10. Hausfather Z, Peters GP. Emissions – the ‘business as usual’ story is misleading. *Nature*. 2020;29 January 2020.
11. Engels A, Marotzke J, Ratter BMW, Gonçalves Gresse E, López-Rivera A, Pagnone A, et al. Hamburg Climate Futures Outlook. Conditions for Sustainable Climate Change Adaptation. Bielefeld: Cluster of Excellence Climate, Climatic Change, and Society (CLICCS); 2024.
- 1160 12. Ritchie J, Dowlatabadi H. Why do climate change scenarios return to coal? *Energy*. 2017;140:1276-91.
13. van Vuuren DP, Kriegler E, O'Neill BC, Ebi KL, Riahi K, Carter TR, et al. A new scenario framework for Climate Change Research: Scenario matrix architecture. *Climatic Change*. 2014;122(3):373-86.
- 1165 14. Lau WK-M, Kim K-M. Competing influences of greenhouse warming and aerosols on Asian summer monsoon circulation and rainfall. *Asia-Pacific Journal of Atmospheric Sciences*. 2017;53(2):181-94.
15. Lin L, Wang Z, Xu Y, Fu Q, Dong W. Larger Sensitivity of Precipitation Extremes to Aerosol Than Greenhouse Gas Forcing in CMIP5 Models. *Journal of Geophysical Research: Atmospheres*. 2018;123(15):8062-73.
- 1170 16. Westervelt DM, Mascioli NR, Fiore AM, Conley AJ, Lamarque JF, Shindell DT, et al. Local and remote mean and extreme temperature response to regional aerosol emissions reductions. *Atmos Chem Phys*. 2020;20(5):3009-27.
17. Tebaldi C, Wehner M, Leung R, Lawrence D. Is land use producing robust signals in future projections from Earth system models, all else being equal? *Environmental Research Letters*. 2023;18:084009.
- 1175 18. Riahi K, van Vuuren DP, Kriegler E, Edmonds J, O'Neill BC, Fujimori S, et al. The Shared Socioeconomic Pathways and their energy, land use, and greenhouse gas emissions implications: An overview. *Global Environmental Change*. 2017;42:153-68.
19. IIASA. SSP Scenario Explorer 2024 [Available from: <https://data.ece.iiasa.ac.at/ssp/#/login?redirect=%2Fworkspaces>].

- 1180 20. Mastrandrea MD, Mach KJ, Plattner GK, Edenhofer O, Stocker TF, Field CB, et al. The IPCC AR5 guidance note on consistent treatment of uncertainties: A common approach across the working groups. *Climatic Change*. 2011;108(4):675-91.
21. Carter TR, Jones RN, Lu X, Bhadwal S, Conde C, Mearns LO, et al. New Assessment Methods and the Characterisation of Future Conditions. In: Parry ML, Canziani OF, Palutikof JP, van der Linden PJ, Hanson CE, editors. *Climate Change 2007 Impacts, Adaptation and Vulnerability Contribution of Working Group II to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge, UK, : Cambridge University Press, ; 2007. p. 133-71.
- 1185 22. Jewell J, Cherp A. The feasibility of climate action: Bridging the inside and the outside view through feasibility spaces. *WIREs Climate Change*. 2023;14 (5).
- 1190 23. Brutschin E, Pianta S, Tavoni M, Riahi K, Bosetti V, Marangoni G, et al. A multidimensional feasibility evaluation of low-carbon scenarios. *Environmental Research Letters*. 2021;16(6).
24. Ju Y, Sugiyama M, Shiraki H. Perceived feasibility and potential barriers of a net-zero system transition among Japanese experts. *Communications Earth & Environment*. 2023;4(1):431.
- 1195 25. Riahi K, Schaeffer R, Arango J, Calvin K, Guivarch C, Hasegawa T, et al. Chapter 3: Mitigation Pathways Compatible with Long-Term Goals. In: Skea J, et al, editors. *IPCC Working Group III contribution to the Sixth Assessment report IPCC2022*.
26. Allen MR, Dube OP, Solecki W, Aragón-Durand F, Cramer W, Humphreys S, et al. Framing and Context. In: Masson-Delmotte V, Zhai P, Pörtner H-O, Roberts D, Skea J, Shukla PR, et al., editors. *Global Warming of 15°C An IPCC Special Report on the impacts of global warming of 15°C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the global response to the threat of climate change, sustainable development, and efforts to eradicate poverty*. Cambridge, UK and New York, NY, USA: Cambridge University Press; 2018. p. 49-92.
- 1200 27. Rennert K, Errickson F, Prest BC, Rennels L, Newell RG, Pizer W, et al. Comprehensive evidence implies a higher social cost of CO₂. *Nature*. 2022;610(7933):687-92.
- 1205 28. Moore FC, Lacasse K, Mach KJ, Shin YA, Gross LJ, Beckage B. Determinants of emissions pathways in the coupled climate–social system. *Nature*. 2022;603(7899):103-11.
29. Liu PR, Raftery AE. Country-based rate of emissions reductions should increase by 80% beyond nationally determined contributions to meet the 2 °C target. *Communications Earth & Environment*. 2021;2(1):29.
- 1210 30. Sanderson BM, Booth BBB, Dunne J, Eyring V, Fisher RA, Friedlingstein P, et al. The need for carbon emissions-driven climate projections in CMIP7. *EGUsphere*. 2023;2023:1-51.
31. Hajima T, Kawamiya M, Ito A, Tachiiri K, Jones C, Arora V, et al. Consistency of global carbon budget between concentration- and emission-driven historical experiments simulated by CMIP6 Earth system models and suggestion for improved simulation of CO₂ concentration. *EGUsphere*. 2024;2024:1-49.
- 1215 32. Séférian R, Berthet S, Yool A, Palmiéri J, Bopp L, Tagliabue A, et al. Tracking Improvement in Simulated Marine Biogeochemistry Between CMIP5 and CMIP6. *Current Climate Change Reports*. 2020;6(3):95-119.
33. Nicholls Z, Meinshausen M, Pflüger M, Lewis J. CMIP GHG Forcing 2025 [Available from: <https://zenodo.org/records/14892947>].
- 1220 34. Zimm C, Mintz-Woo K, Brutschin E, Hanger-Kopp S, Hoffmann R, Kikstra JS, et al. Justice considerations in climate research. *Nature Climate Change*. 2024;14(1):22-30.
35. Kanitkar T, Mythri A, Jayaraman T. Equity assessment of global mitigation pathways in the IPCC Sixth Assessment Report. *Climate Policy*. 2024;24(8):1129-48.
- 1225 36. Hickel J, Slamersak A. Existing climate mitigation scenarios perpetuate colonial inequalities. *The Lancet Planetary Health*. 2022;6(7):e628-e31.

37. Bauer N, Bertram C, Schultes A, Klein D, Luderer G, Kriegler E, et al. Quantification of an efficiency-sovereignty trade-off in climate policy. *Nature*. 2020;588(7837):261-6.
38. Rubiano Rivadeneira N, Carton W. (In)justice in modelled climate futures: A review of integrated assessment modelling critiques through a justice lens. *Energy Research & Social Science*. 2022;92:102781.
- 1230 39. Emmerling J, Drouet L, van der Wijst K-I, Vuuren D, Bosetti V, Tavoni M. The role of the discount rate for emission pathways and negative emissions. *Environmental Research Letters*. 2019.
40. IPCC. IPCC Workshop on the Use of Scenarios in the Sixth Assessment Report and Subsequent Assessments. Bangkok, Thailand Intergovernmental Panel on Climate Change; 2023.
41. Smith C. fair calibration data (1.5.0) [Data set]. Zenodo. . 2025.
- 1235 42. Smith C, Cummins DP, Fredriksen HB, Nicholls Z, Meinshausen M, Allen M, et al. fair-calibrate v1.4.1: calibration, constraining, and validation of the FaIR simple climate model for reliable future climate projections. *Geosci Model Dev*. 2024;17(23):8569-92.
43. Nicholls Z, Kikstra J, Zecchetto M, Hoegner A. CMIP7 ScenarioMIP historical timeseries for harmonisation and simple climate model workflow 2025 [Available from: <https://zenodo.org/records/17527153>].
- 1240 44. FaIR team. FaIR: Finite Amplitude Impulse Response simple climate model 2025 [Available from: <https://docs.fairmodel.net/en/latest/>].
45. Sanderson B, Smith C. Indicative ScenarioMIP emissions and climate projections 2025 [Available from: <https://zenodo.org/records/14382495>].
46. Funke B, Dudok de Wit T, Ermolli I, Haberreiter M, Kinnison D, Marsh D, et al. Towards the definition of a solar forcing dataset for CMIP7. *Geosci Model Dev*. 2024;17(3):1217-27.
- 1245 47. Aubry TJ, Toohey M, Khanal S, Chim MM, Verkerk M, Johnson B, et al. Stratospheric aerosol forcing for CMIP7 (part 1): Optical properties for pre-industrial, historical, and scenario simulations (version 2.2.1). *EGUsphere*. 2025;2025:1-53.
48. Skeie RB, Byrom R, Hodnebrog Ø, Jouan C, Myhre G. Multi-model effective radiative forcing of the 2020 sulfur cap for shipping. *Atmos Chem Phys*. 2024;24(23):13361-70.
- 1250 49. Wilcox LJ, Allen RJ, Samset BH, Bollasina MA, Griffiths PT, Keeble J, et al. The Regional Aerosol Model Intercomparison Project (RAMIP). *Geosci Model Dev*. 2023;16(15):4451-79.
50. Lehner F, Deser C. Origin, importance, and predictive limits of internal climate variability. *Environ Res: Climate*. 2023;2:023001.
- 1255 51. Pfleiderer P, Frölicher TL, Kropf CM, Lamboll RD, Lejeune Q, Capela Lourenço T, et al. Reversal of the impact chain for actionable climate information. *Nature Geoscience*. 2025;18(1):10-9.
52. Vioni D, Robock A, Haywood J, Henry M, Tilmes S, MacMartin DG, et al. G6-1.5K-SAI: a new Geoengineering Model Intercomparison Project (GeoMIP) experiment integrating recent advances in solar radiation modification studies. *Geosci Model Dev*. 2024;17(7):2583-96.
- 1260 53. Eyring V, Collins WD, Gentine P, Barnes EA, Barreiro M, Beucler T, et al. Pushing the frontiers in climate modelling and analysis with machine learning. *Nature Climate Change*. 2024;14(9):916-28.
54. Gidden MJ, Riahi K, Smith SJ, Fujimori S, Luderer G, Kriegler E, et al. Global emissions pathways under different socioeconomic scenarios for use in CMIP6: A dataset of harmonized emissions trajectories through the end of the century. *Geoscientific Model Development*. 2019;12(4):1443-75.
- 1265 55. Hurtt GC, Chini L, Sahajpal R, Frohking S, Bodirsky BL, Calvin K, et al. Harmonization of global land use change and management for the period 850-2100 (LUH2) for CMIP6. *Geoscientific Model Development*. 2020;13(11):5425-64.
56. IEA. The Role of Critical Minerals in Clean Energy Transitions. Paris: IEA; 2021.
- 1270 57. Schlichenmaier S, Naegler T. May material bottlenecks hamper the global energy transition towards the 1.5 °C target? *Energy Reports*. 2022;8:14875-87.

58. Bauer N, Hilaire J, Brecha RJ, Edmonds J, Jiang K, Kriegler E, et al. Assessing global fossil fuel availability in a scenario framework. *Energy*. 2016;111:580-92.
59. Rogner HH. An assessment of world hydrocarbon resources. *Annual Review of Environment and Resources*. 1997;22.
- 1275 60. Roelfsema M, van Soest HL, Harmsen M, van Vuuren DP, Bertram C, den Elzen M, et al. Taking stock of national climate policies to evaluate implementation of the Paris Agreement. *Nature Communications*. 2020;11(1).
61. Rogelj J, Fransen T, den Elzen MGJ, Lamboll RD, Schumer C, Kuramochi T, et al. Credibility gap in net-zero climate targets leaves world at high risk. *Science*. 2023;380(6649):1014-6.
- 1280 62. van Soest HL, Aleluia Reis L, Baptista LB, Bertram C, Després J, Drouet L, et al. Global roll-out of comprehensive policy measures may aid in bridging emissions gap. *Nature Communications*. 2021;12(1).
63. Reisinger A, Fuglestvedt JS, Pirani A, Geden O, Jones CD, Maharaj S, et al. Overshoot: A Conceptual Review of Exceeding and Returning to Global Warming of 1.5°C. *Annual Review of Environment and Resources*. 2025; 50:185-217.
- 1285 64. Odenweller A, Ueckerdt F, Nemet GF, Jensterle M, Luderer G. Probabilistic feasibility space of scaling up green hydrogen supply. *Nature Energy*. 2022;7(9):854-65.
65. Wilson C, Grubler A, Bento N, Healey S, De Stercke S, Zimm C. Granular technologies to accelerate decarbonization. *Science*. 2020;368(6486):36-9.
66. Willett W, Rockström J, Loken B, Springmann M, Lang T, Vermeulen S, et al. Food in the
- 1290 Anthropocene: the EAT–Lancet Commission on healthy diets from sustainable food systems. *The Lancet*. 2019;393(10170):447-92.
67. Stehfest E, Bouwman L, van Vuuren DP, den Elzen MGJ, Eickhout B, Kabat P. Climate benefits of changing diet. *Climatic Change*. 2009:1-20.
68. Beier FD, Dietrich JP, Heinke J, Abrahao G, von Jeetze P, Bodirsky BL, et al. Planetary boundaries under a land-based climate change mitigation scenario with a food demand transformation: a modelling study. *The Lancet Planetary Health*. 2025;9(10).
- 1295 69. Friedlingstein P, O'Sullivan M, Jones MW, Andrew RM, Bakker DCE, Hauck J, et al. Global Carbon Budget 2025. *Earth Syst Sci Data Discuss*. 2025;2025:1-139.
70. Tebaldi C, O'Neill B, Lamarque JF. Sensitivity of regional climate to global temperature and forcing. *Environ Res Lett* 2015;10 074001.
- 1300 71. McKenna CM, Maycock AC, Forster PM, Smith CJ, Tokarska KB. Stringent mitigation substantially reduces risk of unprecedented near-term warming rates. *Nature Climate Change*. 2021;11(2):126-31.
72. Gidden MJ, Joshi S, Armitage JJ, Christ A-B, Boettcher M, Brutschin E, et al. A prudent planetary limit for geologic carbon storage. *Nature*. 2025;645(8079):124-32.
- 1305 73. McGlade C, Ekins P. The geographical distribution of fossil fuels unused when limiting global warming to 2°C. *Nature*. 2015;517(7533):187-90.
74. Melnikova I, Boucher O, Cadule P, Tanaka K, Gasser T, Hajima T, et al. Impact of bioenergy crop expansion on climate–carbon cycle feedbacks in overshoot scenarios. *Earth Syst Dynam*. 2022;13(2):779-94.

1310

Appendix A: Acronyms

AerChemMIP Aerosol Chemistry Model Intercomparison Project

AFOLU Agriculture, Forestry and Other Land Use

- BC Black Carbon
- 1315 BECCS Bioenergy with Carbon Capture and Storage
 - CCS Carbon Capture and Storage
 - CMIC Climate Model of Intermediate Complexity
 - CMIP Coupled Model Intercomparison Project
 - CDR Carbon Dioxide Removal
- 1320 DAC Direct Air Capture
 - DACCS Direct Air Capture with Carbon Storage
 - DECK Diagnostics, Evaluation and Characterization of Klima
 - ESM Earth System Model
 - EV Electric Vehicle
- 1325 GCM Global Circulation Model/Global Climate Model
 - GHG Green-house gas
 - GMST Global Mean Surface Temperature
 - GSAT Global-mean Surface Air Temperature
 - GWP100 Global Warming Potential over 100 years
- 1330 H High scenario
 - H-ext High extension
 - H-ext-OS High overshoot extension
 - IAM Integrated Assessment Model
 - IAMC Integrated Assessment Modeling Consortium
- 1335 IEA International Energy Agency
 - input4mip CMIP activity tasked with the processing and availability of input data for ESM experiments under CMIP
 - IPCC Intergovernmental Panel on Climate Change
 - L Low scenario
 - L-ext Low extension
- 1340 LUMIP Land Use Model Intercomparison Project
 - M Medium scenario
 - M-ext Medium extension
 - ML Medium-to-Low scenario
 - ML-ext Medium-to-Low extension
- 1345 MIP Model Intercomparison Project

MOS Medium scenario with Overshoot
NDC Nationally Determined Contributions
OC Organic carbon
OS Overshoot
1350 RAMIP Regional Aerosol Model intercomparison Project
RCP Representative Concentration Pathway
SCM Simple Climate Model
SDG Sustainable Development Goal
SLCF Short-Lived Climate Forcer
1355 SSC Scientific Steering Committee
SSP Shared Socio-economic Pathways
TCRE Transient Climate Response to cumulative Emissions
VIA Vulnerability, Impacts and Adaptation
VIACCS Vulnerability, Impacts, Adaptation and Climate Services
1360 VL Very Low emission scenario
VL-ext Very Low emission scenario extension
LN Low to negative emission scenario
LN-ext Low to negative emission scenario extension
WGI/II/III Working Group I/II/III