Supplementary file for "Climate-driven Biogenic Emissions alleviate the impact of man-made emission reduction on O₃ control in Pearl River Delta region, southern China"

Nan WANG^{1,2}*, Song LIU¹, Jiawei XU³, Yanyu WANG², Chun LI¹, Hua LU⁴*, Fumo YANG¹

¹College of carbon Neutrality Future Technology, Sichuan University, Chengdu 610065, P. R. China

²State Environmental Protection Key Laboratory of Formation and Prevention of Urban Air Pollution Complex, Shanghai Academy of Environment Sciences, Shanghai 200233, P. R. China

³Centre for Geography and Environmental Science, University of Exeter, Penryn, United Kingdom

⁴Chongqing Institute of Meteorological Sciences, Chongqing 401147, P. R. China

*Correspondence: WANG Nan (<u>nan.wang@scu.edu.cn</u>) and LU Hua (vibgyor0113@163.com)

Supporting Information

Supporting Information includes 5 pages, 1 figures and 3 tables

SI Figures S1-S2, p3

SI Tables S1-S3, p4-p5




Fig S1 Annual variations of meteorological parameters in the PRD region from 2000-2020




Fig S2 Responses of O₃ concentration to NOx emission reduction under different scenarios.

Table S1: Mapping of MODIS vegetation types to MEGAN-PFT based on climatic standards and climate data

Original	Mapped vegetation type	Climate standard	
vegetation type			
Evergreen	Needleleaf evergreen temperate tree	Tc > -19°C and GDD > 1200	
Needle leaf Forests			
Deciduous	Needleleaf deciduous boreal tree	-	
Needle leaf Forests			
Evergreen	Needleleaf evergreen boreal tree	$Tc \le -19$ °C or $GDD \le 1200$	
Needle leaf Forests			
Evergreen	Broadleaf evergreen tropical tree	Tc > 15.5°C	
Broadleaf Forests			
Evergreen	Broadleaf evergreen temperate tree	Te ≤ 15.5°C	
Broadleaf Forests			
Deciduous	Broadleaf deciduous tropical tree	Te > 15.5°C	
Broadleaf Forests			
Deciduous	Broadleaf deciduous temperate tree	$-15^{\circ}\text{C} < \text{Tc} \leq 15.5^{\circ}\text{C}$, and	
Broadleaf Forests		GDD>1200	
Deciduous	Broadleaf deciduous boreal tree	$Tc \le -15$ °C or $GDD \le 1200$	
Broadleaf Forests			
Shrublands	Broadleaf evergreen temperate shrub	Tc > -19°C , $GDD > 1200$,	
		$P_{ann} > 520 \ mmand \ P_{win} > 2/3 \ P_{ann}$	
Shrublands	Broadleaf deciduous temperate shrub	Tc > -19°C , $GDD > 1200$,	
		and meeting either one of the	
		following standard	
		(1) $P_{ann} \leq 520 \text{ mm}$	
		$(2) P_{win} \le 2/3 P_{ann}$	
Shrublands	Broadleaf deciduous boreal shrub	$Tc \le -19$ °C or $GDD \le 1200$	
Grasslands	Cold C3 grass	GDD < 1000	
Grasslands	Cool C3 grass	GDD > 1000, and meeting	
		either one of the following standard	
		(1) $T_w \le 22$ °C	
		(2) For months with temperatures	
		exceeding 22°C, Pmon ≤ 25	
		mm.	
Grasslands	Warm C3 grass	GDD > 1000 , $T_c > 22$ °C and	
	-	$P_{mon} > 25$ mm in the driest month	
Croplands	Other crops	-	

Noting: T_c is the average temperature of the coldest month of the year, T_w is the average temperature of the warmest month, GDD represents the growing degree days (temperature above 5°C), P_{ann} refers to annual precipitation, P_{win} is winter precipitation, and P_{min} denotes monthly precipitation.

Table S2 Statistical validation of WRF-CMAQ performance in PRD region

	MB	RMSE	IOA
T2 (°C)	-0.6	2.1	0.91
RH(%)	-4.5	8.6	0.99
Pressure (hPa)	-15.1	28.5	0.89
WS10 (m/s)	1.6	1.8	0.97
O ₃ (ppb)	3.5	27.05	0.78

Noting: T2 indicates 2-meter temperature, WS10 indicates 10m wind speed. MB is mean bias, RMSE is root mean square and IOA is index of agreement.

Table S3 Parallel numerical simulation experiments

				1	
Impact of each	Parallel	Land	Meteorology for	Meteorology for	Anthropogenic
process	numerical	cover and	BVOC emission	Chemistry	emission
	experiment	LAI			
Man-made	EXP1	2020	2020	2020	2012
emission control	EXP2	2020	2020	2020	2020
Vegetation-	EXP1	2001	2020	2020	2020
Change BVOC	EXP2	2020	2020	2020	2020
Climate-driven	EXP1	2020	2001	2020	2020
BVOC	EXP2	2020	2020	2020	2020
Climate-driven	EXP1	2020	2020	2001	2020
meteorology	EXP2	2020	2020	2020	2020