

# Assessing the impact of riverine water on the Northwest Pacific using normalized Total Alkalinity

Tatsuki Tokoro<sup>1, 62</sup>, Shin-Ichiro Nakaoka<sup>1</sup>, Shintaro Takao<sup>1</sup>, Shu ~~Saite~~<sup>2</sup>Saito<sup>3, 34</sup>, Daisuke ~~Sasano~~<sup>3</sup>Sasano<sup>4</sup>, Kazutaka ~~Enyo~~<sup>3</sup>Enyo<sup>4</sup>, Masao ~~Ishii~~<sup>4</sup>Ishii<sup>5</sup>, Naohiro ~~Kosugi~~<sup>4</sup>Kosugi<sup>5</sup>, Tsuneo ~~Ono~~<sup>5</sup>Ono<sup>6</sup>, Kazuaki ~~Tadokoro~~<sup>5</sup>Tadokoro<sup>6</sup>, and Yukihiro Nojiri<sup>1</sup>

<sup>1</sup>~~formally~~<sup>1</sup>~~currently~~ at: Earth System Division, National Institute for Environmental Studies, Tsukuba, Japan.

<sup>2</sup>~~formally~~ at: Seto Inland Carbon-Neutral Research Center, Hiroshima University, Hiroshima, Japan.

<sup>32</sup>Administration Department, Japan Meteorological Agency, Tokyo, Japan.

<sup>42</sup>Atmosphere and Ocean Department, Japan Meteorological Agency, Tokyo, Japan.

<sup>52</sup>Department of Climate and Geochemistry Research, Meteorological Research Institute, Tsukuba, Japan.

<sup>65</sup>Fisheries Resources Institute, Japan Fisheries Research and Education Agency, Yokohama/Shiogama, Japan.

<sup>6</sup>~~currently~~ at: Seto Inland Carbon-Neutral Research Center, Hiroshima University, Hiroshima, Japan.

Corresponding author: Tatsuki Tokoro (~~tokorota@hiroshima-u.ac.jp~~, tokoro.tatsuki@nies.go.jp)

**Abstract.** The impact of riverine water was assessed using salinity-normalized Total Alkalinity observations of the Northwest Pacific, including the coastal areas of Japan (20–50°N, 120–160°E). The observational data included surface carbonate parameters obtained from decades of surveys conducted by volunteer cargo ships and research vessels in the Northwest Pacifics area. This study uses data and statistical methods (e.g., re-gridding and Fourier regression) like those in a previous study that analysed air-sea CO<sub>2</sub> flux but focuses instead on the diffusion of nTA from land. ~~Statistical processes, such as re-gridding and Fourier regression, used in a previous study were also applied in this study to improve the spatiotemporal resolution.~~ First, the seawater area affected by riverine water was identified using an Empirical Orthogonal Function analysis of normalized Total Alkalinity. The differences in normalized Total Alkalinity and Dissolved Inorganic Carbon from the surrounding area were then analysed to evaluate the potential drivers, causes, such as riverine water supply, advection effects, and biological activities. In addition, the impact of riverine water on oceanic CO<sub>2</sub> uptake and acidification in the study area was assessed. The analysis showed that riverine water was the main source-cause of the higher total coastal Alkalinity compared to the surrounding area, whereas-but its contribution to was not the dominant the increase-cause of-in Dissolved Inorganic Carbon was relatively minor. The supply of riverine water had little effect on oceanic CO<sub>2</sub> uptake throughout the year. The supply of ~~by~~ riverine water had a minor effect on pH but was a factor in-of coastal acidification in a calcification index, ~~however,~~ ~~the~~ supplied Total Alkalinity reduced the overall acidification trend this index change by 6571% overall. The results of this study are expected to be further improved by enhancing observations, such as the vertical profiles of carbonate parameters, and are expected to expand to other sea areas and be applied to global budgets.

Formatted: Font: (Default) +Body (Times New Roman)

Formatted: Subscript

Formatted: Font: (Asian) MS 明朝, Not Bold, (Asian) Japanese

## 1. Introduction

The ~~flow-input~~ of riverine water to the ocean is one of the most important flows in the Earth's system. In the carbon cycle, riverine water is ~~the a~~ major carbon source for the oceans (Aufdenkampe et al., 2011; Bauer et al., 2013; Borges et al., 2005; Cai, 2011; Chen and Borges, 2009; Chen et al., 2013). ~~Recent assessments indicate that the global carbon flux into the ocean via rivers is  $1.02 \pm 0.11$  (Ave.  $\pm$  SD) Pg-C yr<sup>-1</sup> (Liu et al., 2024). This is equivalent to about one-third of the oceanic absorption of atmospheric CO<sub>2</sub>, which is the largest air-sea carbon flux to the surface ocean ( $2.9 \pm 0.4$  Pg-C yr<sup>-1</sup>, Friedllingstein et al., 2025).~~ ~~This riverine carbon flux~~ is responsible for atmospheric CO<sub>2</sub> emissions and acidification in coastal areas (Carstensen and Durate, 2019; Duarte et al., 2013; Tranvik et al., 2009). However, strong carbon flows, such as biological pumps, impact carbonate distribution in coastal areas (Passow and Carlson, 2012; Regnier et al., 2013, 2022). ~~For example, an increased biological pump in coastal areas will enhance CO<sub>2</sub> absorption from the atmosphere, but it will also promote anoxia in the sediments.~~ ~~For example, increased biological pump in coastal areas will increase CO<sub>2</sub> absorption from atmosphere, but it will also progress the anoxia on the sediment.~~ In coastal areas, the ~~decomposition-mineralization~~ of organic matter supplied by riverine water is expected to cause more severe acidification, which may ~~eause-stimulate~~ harmful algal growth and adversely affect marine products such as bivalves (Fitzer et al., 2018; Kessouri et al., 2021; Wallace et al., 2014). Quantifying these complex flows is important for ~~predicting-anticipating~~ future climate change and ocean acidification as well as for ~~predicting-projecting~~ biogeochemical changes in coastal areas.

The Northwest Pacific Ocean, including the coastal areas of Japan, plays a crucial role in global carbon cycle due to ~~the-its~~ strong ~~sinks~~ of atmospheric CO<sub>2</sub> (Takahashi et al., 2002, 2009). While several studies on carbonate systems have been conducted in the Northwest Pacific Ocean (e.g., Ishii et al., 2001; Murata et al., 1998; Takamura et al., 2010; Tokoro et al., 2023; Yoshikawa-Inoue et al., 1995, 2014), spatiotemporal variations in riverine water contribution in this region have not yet been quantified. Riverine water tracers (e.g., salinity, stable isotope, geogenic solutes like silica) are one of the most effective methods for evaluating the influence of riverine water. Although salinity is the most frequently assessed factor in riverine water transportation, other factors such as precipitation and evaporation also impact salinity. This can be a source of major error in the assessment of water transportation in the ~~northwestern~~Northwest Pacific region, where high precipitation and evaporation by low-pressure systems and heat waves, respectively, have been observed (Kitamura et al., 2016; Miyama et al., 2021; Sugimoto et al., 2013).

The total Alkalinity (TA) normalized by salinity is a potential indicator to assess the influence of riverine water on ocean. TA ~~is a carbonate parameter in seawater and~~ is defined broadly ~~by-as~~ the charge difference between ~~the~~-proton donors and acceptors (see Zeebe and Wolf-Gladrow, 2001 for ~~a~~ more detailed definition). TA ~~ehanges-is influenced~~ by several factors, such as advection from different water masses and biological ~~metabolismactivity~~, including ~~the~~-calcification and dissolution

Formatted: English (United States)

of calcium carbonate. Because TA is also highly correlated with salinity, TA normalized to ~~the a~~ reference salinity (nTA) has been used to quantify the ~~above-mentioned~~ factors (e.g., Broecker and Peng, 1982; Lee et al., 2006; Millero et al., 1998). nTA is calculated as follows:

$$70 \quad nTA = TA \cdot \frac{S_{ref}}{S} \quad (1)$$

where,  $S$  and  $S_{ref}$  are the measured and reference salinities (traditionally 35), respectively. Similarly, Dissolved Inorganic Carbon concentration (DIC), ~~which is the sum of inorganic carbon species~~, was also normalized (nDIC). Equation (1) is formulated based on the assumption that a water mass with zero salinity has zero TA, otherwise the right-hand side would go to infinity. However, this assumption is not true for riverine water because its TA is greater than zero, even when salinity is  
75 zero, owing to the weathering of carbonate and silicate rocks (e.g., Friis et al., 2003; Lehmann et al., 2023; Rassmann et al., 2016; Taguchi et al., 2009). Therefore, when Equation (1) is applied to areas affected by riverine water, the nTA value is higher than that of the surrounding seawater. Conversely, the influence of riverine water, along with factors such as water mass advection and ~~biological-metabolism~~biological activity, can be quantified by assessing the distribution of nTA. Unlike other methods that use salinity as a tracer for riverine water, nTA defined in Equation (1) excludes the effect of precipitation  
80 and evaporation, which is advantageous because contamination by water masses with almost zero salinity and TA can be excluded. ~~Furthermore, nTA can be used to~~is also easier to quantify the influence of seawater ~~inflow~~ from different local areas because ~~distinct regional differences in nTA have been reported-the TA has different values in different local areas, even if the salinity is the same~~ (Kakei et al., 2017; Lee et al., 2006; Takatani et al., 2014).

In this study, we aimed to analyse the influence of riverine water on ~~the carbonate chemistry of the Northwest Pacific carbonate systems in the ocean~~ using nTA and other carbonate parameters measured by voluntary cargo ships and research vessels ~~in the Northwest Pacific~~. First, spatiotemporal variations in the area in which riverine water significantly affected surface seawater ~~carbonate chemistry~~ were identified using ~~an~~ Empirical Orthogonal Function (EOF) analysis of the nTA distribution. Second, we focused on the differences in ~~n~~TA and ~~n~~DIC between the riverine water-affected area and surrounding ~~unaffected~~ areas. We quantified the riverine water supply and other contributing factors that affect ~~n~~TA and  
90 ~~n~~DIC. The final step involved the evaluation of the effects of riverine water on the environment in riverine water-affected area. Seawater  $CO_2$  fugacity ( $fCO_2$ ), ~~pH~~ and the ~~ealeite-aragonite~~ saturation state of seawater ( $\Omega_{arg}$ ) were the ~~two~~-carbonate parameters that were used as the index of environmental changes caused by riverine water input. ~~The former parameter affects the air-sea  $CO_2$  flux. The TA and DIC supplied by riverine water should change~~influence seawater  $fCO_2$  and oceanic  $CO_2$  uptake. ~~Meanwhile, pH is an indicator of ocean acidification. However,  $\Omega_{arg}$ , which~~ Meanwhile,  $\Omega_{cal}$  ~~is~~ is the ratio of the concentration product of  $[Ca^{2+}]$  and  $[CO_3^{2-}]$  to the solubility product of ~~ealeite-aragonite~~, and is another index of ocean  
95 acidification ~~and has been reported to exhibit different spatiotemporal variations from pH~~ (Kwiatkowski and Orr., 2018; Xue et al., 2021). ~~Calcite is a mineral composed of  $CaCO_3$  and constitutes foraminifera, coccolithophorids, and the outer layer of bivalves.~~ Therefore, the analysis of changing ~~pH and  $\Omega_{cal-arg}$~~  is expected to lead to a more detailed assessment of coastal

acidification in the study area. This study also aimed to evaluate the effects of riverine water on future climate change and coastal acidification and to predict the effects of future environmental changes.

## 2. Methods

### 2.1 Data for analysis

The observational data in this study were collected by the National Institute for Environmental Studies (NIES), Meteorological Research Institute (MRI) of the Japan Meteorological Agency (JMA), and Japan Fisheries Research and Education Agency (FREA). The NIES data were produced as part of the Voluntary Observing Ship (VOS) programs for cargo ships (namely, M/S Alligator Hope, M/S Pyxis, M/S New Century 2, and M/S Trans Future 5). MRI, JMA, and FREA collected data from research vessels (R/V Mirai and R/V Hakuho-maru [affiliated with Japan Agency for Marine-Earth Science and Technology](#) for MRI [data](#); R/V Keifu-maru and R/V Ryofu-maru for JMA [data](#); and R/V Wakataka-maru and R/V Soyo-maru for FREA [data](#)). These data were uploaded to the Surface Ocean CO<sub>2</sub> Atlas (SOCAT; Pfeil et al., 2013; Bakker et al., 2016, <https://socat.info/index.php/data-access/>) and Global Ocean Data Analysis Project (GLODAP, Key et al., 2015; Olsen et al., 2016; Olsen et al., 2020, <https://glodap.info/>). These observations were statistically processed by re-gridding, a second-order approximation of TA, carbonate equilibrium calculations, and Fourier regression into datasets with a spatial resolution of 1° × 1° and a temporal resolution of 0.1 year, as suggested in a previous study (Tokoro et al., 2023). Similar to the previous study, the data was excluded from the analysis if there was insufficient temporal data (n >= 60, corresponding to 6 years of data) in the 1°×1° spatial grid. Although the time interval (from January 1, 2000, to December 31, 2019), covered area (latitudes of 20–50°N and longitudes of 120–160°E), and original observations in the SOCAT by NIES were the same as those in the previous study, the original observations by MRI and JMA increased slightly (287750 to 291138) with the upgrade of the product (SOCAT ver. 2019 to 2023). The FREA dataset is a new addition to those used in previous studies. As the FREA data (n = 566175) were collected every 1 minute while the other data were collected every 10 min, the FREA data were weighted 1/10 with respect to the other data and weighted averaged as the gridded data. The GLODAP data for surface (<10 m depth) TA and DIC from the MRI and JMA were updated using ver. 2.2020 to 2.2023. The number of data points for JMA data increased from 2080 to 2163. However, the effect of increase in the data was small, which did not statistically affect the result (Table S1). Additionally, observations up to a depth of 150 m within the GLODAP were used to evaluate the effects of vertical advection (Text S1).

## 2.2 EOF analysis, Cause Analysis, and Environmental Impact Assessment

In this study, areas significantly affected by riverine water were identified by applying EOF analysis to the spatiotemporal variability of nTA. The EOF analysis ~~breaks-down~~breaks spatiotemporal variations into multiple orthogonal modes with multiple spatial patterns (principal EOF patterns) and time series (principal component time series) (e.g., Denbo and Allen, 1984). In addition to the effect of riverine water, the target area in this study is assumed to be subject to a mixture of multiple fluctuations, such as seasonal fluctuations of the Kuroshio and Oyashio flows. Compared to the direct use of carbonate parameters, the EOF analysis is expected to identify the influence of riverine water because the analysis is appropriate to quantify and separate multiple fluctuations. Using the principal EOF pattern and principal component time series with respect to the spatiotemporal variation of nTA, we identified the area significantly influenced by riverine water and labelled it as “Area A”. We then labelled the surrounding ~~area-region~~ with a rather similar surface area, almost the same area as “Area B.” The influence of riverine water was quantified as the value of Area A minus that of Area B ( $dAB$ ) for all related parameters such as SST (Sea Surface Temperature), SSS (Sea Surface Salinity), nTA, and nDIC. Causal analysis of  $dAB$  of TA and DIC was performed using the following equation:

$$\frac{\partial dAB}{\partial t} = Sup_{rw} + C_{flux} + C_{res} \quad (2)$$

The left-hand side of the equation represents the time derivative of  $dAB$  of nTA and nDIC.  $Sup_{rw}$  is the TA or DIC supply by riverine water;  $C_{flux}$  is the term of difference in air-sea CO<sub>2</sub> flux between Areas A and B divided by the mixed layer depth (MLD). The MLD with the same spatiotemporal resolutions as the processed data (1° × 1° and 0.1 year) was calculated from the reanalysis data of seawater temperature profiles by Japan Agency for Marine-Earth Science and Technology (JCOPE2M; Miyazawa et al., 2017, 2019, <https://www.jamstec.go.jp/jcope/htdocs/distribution/>). We determined an isothermal depth at  $\Delta T = 0.2$  °C with linear interpolation (de Boyer Montégut et al., 2004; Holte and Talley, 2009).  $C_{flux}$  was applied only to the DIC input.  $C_{res}$  is the TA or DIC input due to other residual factors related to  $dAB$  (e.g., horizontal and vertical advection and biological-metabolismbiological activity).

$Sup_{rw}$  was estimated from river discharge and TA or DIC concentrations in river water as follows:

$$Sup_{rw} = \frac{Flow_r \cdot C_r}{MLD \cdot A} \quad (3)$$

where,  $Flow_r$  is the monthly total flow rate of the river,  $C_r$  represents the TA or DIC concentrations in the associated rivers,  $A$  is the area of Area A. The flow rate was estimated using the Water Information System database of the Ministry of Land, Infrastructure, Transport, and Tourism of Japan (<http://www1.river.go.jp/>). As TA and DIC data were not available for all rivers, the referential values for zero-salinity endmembers in the three most river-influenced inner bays (Tokyo Bay, Ise Bay, and Osaka Bay) in the relevant coastal areas were used (518–1006 and 475–1371  $\mu\text{mol kg}^{-1}$  for TA and DIC, respectively; Taguchi et al., 2009; Tokoro et al., 2021). The DIC range includes the was determined based on the maximum (1171  $\mu\text{mol kg}^{-1}$  in Tokyo Bay) and minimum (675  $\mu\text{mol kg}^{-1}$  in Ise Bay) freshwater endmember values among the three bays, with an additional  $\pm 200$   $\mu\text{mol kg}^{-1}$  to account for seasonal variation estimated in the previous study, value predicted as the

Formatted: Not Superscript/ Subscript

Formatted: Font: (Default) +Body (Times New Roman)

range-of-seasonal-variation ( $\pm 200 \mu\text{mol kg}^{-1}$ ; Tokoro et al., 2021). Although the TA range does not consider-account for the seasonal variations, the range of TA variation in global rivers due to water temperature fluctuations is reported to be less than  $10 \mu\text{mol kg}^{-1}$  (Romero-Mujalli et al., 2019). This range is one order of magnitude smaller than that among the three bays described above, thus it was decided that the seasonal variation could be disregarded in the analysis. the-range-is-estimated-to-be-an-order-of-magnitude-smaller-than-the-range-among-the-three-bays-from-the-literature-value (Romero-Mujalli et al., 2018). All data were regridded to the resolution of the processed SOCAT and GLODAP data (spatial and temporal resolution of  $1^\circ \times 1^\circ$  and 0.1 year, respectively).

Air-sea  $\text{CO}_2$  flux ( $F$ ) was determined as follows:

$$F = k \cdot K_0 \cdot (f\text{CO}_{2\text{water}} - f\text{CO}_{2\text{air}}) \quad (4)$$

where,  $k$  is transfer velocity defined by the wind speed at 10 m above the sea surface (Wanninkhof, 2014).  $K_0$  is the solubility of  $\text{CO}_2$  in seawater and was calculated using an empirical equation based on SST and SSS (Weiss, 1974).  $f\text{CO}_{2\text{water}}$  and  $f\text{CO}_{2\text{air}}$  are the  $f\text{CO}_2$  in surface seawater and air, respectively.  $C_{\text{flux}}$  was determined by the difference in air-sea  $\text{CO}_2$  flux between Areas A and B, which was calculated from the processed  $f\text{CO}_2$  data and wind velocity from the database of the Cross-Calibrated Multi-Platform (CCMP, Atlas et al., 2011; Mears et al., 2019, <https://data.remss.com/ccmp/v02.0/>, version 2.0). The details of the calculations are the same as those used in a previous study (Tokoro et al., 2023).

The effect of riverine water on air-sea  $\text{CO}_2$  flux and acidification were quantified by multivariate analysis using  $dAB$  of seawater  $f\text{CO}_2$ -and pH and  $\Omega_{\text{arg}}$  as the objective variables and  $dAB$  of SST, SSS, nTA, and nDIC as the explanatory variables. The pH  $\Omega_{\text{sat}}$  was determined as total scale from  $f\text{CO}_2$  and TA, and  $\Omega_{\text{arg}}$  was calculated using the equation of an aragonite solubility product (Mucci, 1983) and the concentration of  $\text{Ca}^{2+}$  and  $\text{CO}_3^{2-}$ . Both pH and  $\Omega_{\text{arg}}$  were estimated using the CO2SYS program (Lewis and Wallace, 1998) and the “recommend” coefficients in a literature (Zeebe and Wolf-Gladrow, 2001). These Both-sSeawater  $f\text{CO}_2$ -and  $\Omega_{\text{sat}}$ pH, and  $\Omega_{\text{arg}}$  can be unambiguously determined from equilibrium calculation using SST, SSS, nTA, and nDIC (Zeebe and Wolf-Gladrow, 2001). However, it is difficult to intuitively understand the contribution of the explanatory variables because these have a non-linear relationship. Therefore, we considered that a multivariate linear model should be useful in this study. Partial least squares regression (PLS regression; Wold et al., 2001) was used in the multivariate analysis to prevent multicollinearity, particularly especially-considering the strong correlation between SSS and nTA and nDIC, due to riverine water having high nTA and nDIC.


Formatted: Subscript

Formatted: Not Superscript/ Subscript

### 3. Results

Figure 1 represents the processed spatial distributions of SSS, TA, and nTA. The SSS showed a north-south gradient, which was attributed to the high-salinity Kuroshio current and the relatively low salinity Oyashio current. There was also an area of reduced SSS along the Pacific coast of mainland Japan ( $32\text{--}34^\circ\text{N}$ ,  $132\text{--}140^\circ\text{E}$ ; Figure 1a). These trends were similar for TA, which exhibited high correlation with SSS ( $R^2 = 0.94$ ) (Figure 1b, d). The nTA was high in the northern part of the study area, and slightly high values were also observed along the Pacific coast of Japan (Figure 1c). The intercept of the regression

line between SSS and TA was  $528.04 \pm 2.00461.09 \mu\text{mol kg}^{-1}$  (Ave  $\pm$  ~~SE~~SED), which was consistent with the TA value of large river on the continental side like the Amur River ( $589 \mu\text{mol kg}^{-1}$ ; Andreev and Pavlova, 2009) and the range of Japanese river riverine water used in this study ( $518\text{--}1006 \mu\text{mol kg}^{-1}$ ) (Figure 1d). Because the intercept value was above zero, nTA seemed to be inversely proportional to SSS ( $R^2 = 0.57$ ), with a lower SSS tending to have a higher nTA (Figure 1e). These results indicate that the study area was affected by freshwater with TA above zero, especially in the northern area and on the Pacific coast of Japan, and that nTA could be used as a tracer for ~~the~~ freshwater inputs.








Figure 1 (a–c): Spatial distribution of (a) mean Sea surface Salinity (SSS) (b) Total Alkalinity (TA) and (c) normalized TA (nTA).

The rectangular in (a) show the reduced SSS area along the Pacific coast of mainland Japan (32–34°N, 132–140°E) (d and e): Scatterplot of (d) SSS–Total Alkalinity (TA) and (e) SSS–nTA. The black line is the approximate linear regression line calculated using the least squares method between SSS and TA ( $R^2 = 0.94$ ,  $p < 0.05$ ,  $TA = (50.46 \pm 0.06) \times SSS + (528.04 \pm 2.00461.09)$  (Ave.  $\pm$  SE(SD))).

205 The spatial distribution of nTA in the three most dominant modes of the EOF analysis is shown in Figure 2(a–c). The principal EOF patterns were indicated by a parameter defined as “EOF anomaly,” denoted by red or blue in Figure 2a. The spatiotemporal variation of the nTA anomaly in each mode was expressed as the product of the EOF anomaly and the principal component time series. In the most dominant mode (Mode 1), the principal patterns exhibited a clear north-south difference at approximately 37°N (Figure 2a). The principal component of the time series in Mode 1 (Figure 2d) indicated that the annual cycle was predominant. ~~Annual~~Annual maximum of nTA was observed in summer in the area south of 37°N, while it was in winter in the area north of 37°N (data not shown). The seasonal variation south of 37°N can be explained by the fact that riverine water supply is proportional to the precipitation over land, thereby reaching its peak in summer on the Pacific side of Japan (Database of Japan Meteorological Agency, <https://www.data.jma.go.jp/stats/etrn/index.php>). For the area north of 37°N, this may be due to the southward transportation of high-nTA surface seawater in the Pacific subarctic region by north winds in winter or winter vertical mixing that supplied subsurface high-nTA seawater to the surface. In addition, the EOF anomaly south of 37°N was significantly correlated with distance from the Japanese mainland (Figure 2e). Because EOF anomaly is a parameter indicating the strength of annual cycle fluctuation in nTA, as shown in Figure 2d, Figure 2e indicates that the degree of temporal variation in nTA increased significantly closer to mainland Japan in this mode. The other major modes (Modes 2 and 3; Figure 2b, c) had an east-west distribution north of 37°N. A distribution such as this could represent seasonal or multi-year variations in the flow path of the Kuroshio Extension. After mode 4, the variation north of 37°N was also explained, and thus mode 1 was dominant for the variation south of 37°N.

Based on the above considerations, we determined that it was appropriate to use the Mode 1 EOF anomaly south of 37°N as an indicator of the influence of riverine water from mainland Japan. Although the area north of 37°N contains Japanese rivers with large flow rates (e.g., the Kitakami River), the ~~effect of riverine water~~ir influence was excluded from the analysis in this study because the influence of high nTA ~~fluctuation~~ in the subarctic gyre was too dominant to extract the influence of riverine water from Japan. The influence of subarctic Pacific seawater can be expected even around 37°N; however, this is not expected to have a significant effect on the statistics in Area A, such as the spatial mean values.

Area A ~~was located~~~~defined to be~~ south of 37°N and within 250 km ~~of from~~ land (Figure 2f). This distance ~~is was~~ ~~dedetermined~~~~ined~~ using change-point detection (Killick et al., 2012), ~~meaning indicating~~ that the mean EOF anomaly south of 37°N changes abruptly before and after this distance. The outer edge of Area A roughly aligned with the Kuroshio axis, as indicated by the JCOPE2 SSH data (SSH = 0.2 m). Although the EOF anomaly outside Area A also shows a relatively weak correlation with distance, this correlation is thought to be due to the influence of riverine water from areas other than Japan, such as East Asia, which was difficult to quantify using the measurement data in this study.

To minimize the influence of riverine water outside Japan and differences in latitude and longitude, Area B was determined as adjacent to Area A, similar in size to the area for comparison (Figure 2f).

Formatted: Font: (Asian) MS 明朝, (Asian) Japanese






Figure 2. (a–c): EOF anomaly of (a) Mode 1, (b) Mode 2, and (c) Mode 3. The percentage in the bracket (18.8%) indicates the contribution of this mode to the original variation. (d): Principal component of time series of Mode 1. (e): Scatterplot of EOF anomaly of Mode 1 versus distance from Japan mainland. The plot colours indicate the latitude. The dotted line represents the boundary of Areas A and B estimated using the changepoint analysis. (f): Distribution of Area A and B. The black curve indicates mean Kuroshio axis during the whole period in this study (2000–2019).


The monthly and annual average values of the differences in the relevant parameters (SST, SSS, nTA, and nDIC) between Areas A and B ( $dAB$ ) are shown in Figure 3. The average values for each parameter in Areas A and B are summarized in Table S2. All  $dAB$  parameters showed seasonal variation, and the absolute values of  $dAB$  were largest in winter for SST and nDIC and in summer for SSS and nTA. On a decadal scale, the absolute values of all  $dAB$  tended to increase significantly. This indicates an increase in the supply of riverine water, which is consistent with the Japan Meteorological Agency’s report (<https://www.data.jma.go.jp/cpd/cgi-bin/view/index.php>) that precipitation in relevant areas of Japan increased during the period analysed in this study.




**Figure 3. The differences in Sea Surface Temperature (SST), Sea Surface Salinity (SSS), normalized Total Alkalinity (nTA), and normalized Dissolved Inorganic Carbon (nDIC) between Areas A and B (*dAB*). (a)–(d): Seasonal variations. (e)–(h): Annual variations. Although the time step in this study was 0.1 year, the monthly values were calculated by spline interpolation of values at 1/12-year intervals. The same applies to the other figures.**

Figure 4 represents the time-series variations of each term in Equation (2) used for the causal analysis of the *dAB* of nTA and nDIC. The river discharge for  $Sup_{rw}$  was calculated as the sum of the 37 rivers bordering Area A (see Text S2). The maximum nTA and nDIC- $Sup_{rw}$  for both nTA and nDIC were observed during the summer months, when precipitation on the Pacific coast of Japan was the highest (Figure 4b and 4d). Though the air-sea  $CO_2$  flux term ( $C_{flux}$ ) also showed a clear seasonal variation, the annual average did not. The residual term ( $C_{res}$ ) tended to be negative for both nTA and nDIC; however, the seasonal patterns and ranges of variation differed considerably.







**Figure 4.** (a): Temporal variations of normalized Total Alkalinity (nTA, a-c) and normalized Dissolved Inorganic Carbon (nDIC, d-f) time derivatives and related terms in Equation (2). Panels show (a,d) temporal variations, (b, e) monthly averages, and (c, f) yearly averages. S(b): The monthly averages. (c): The yearly averages. (d)–(f): Similarly, but for nDIC. The shaded areas indicate error ranges are error ranges calculated from the upper and lower limits of the riverine Total Alkalinity ( $TA_r$  (518–1006  $\mu\text{mol kg}^{-1}$ ) for and Dissolved Inorganic Carbon ( $DIC_r$  (475–1371  $\mu\text{mol kg}^{-1}$ ) in riverine water; these ranges are not random errors, thus so the same ranges apply to have remained the same error range for monthly and/or yearly averaging. The dotted lines within shaded areas represent the average values.

280

The PLS regression showed that the explanatory variables SST, SSS, nTA, and nDIC explained well the objective variables seawater fCO<sub>2</sub> ( $r^2 = 0.996$ ), ~~and~~ pH ( $r^2 = 0.9937$ ), and  $\Omega_{arg}$  ( $r^2 = 0.996$ ) (Figure 5). The average contributions due to each explanatory variable in the PLS analysis were consistent with those in the equilibrium calculation for seawater fCO<sub>2</sub>, pH and  $\Omega_{ent-arg}$  using the averages in Area B and the respective  $dAB$  values (Table S2S3). ~~Area A had lower s~~The seawater fCO<sub>2</sub> and  $\Omega_{arg}$  in Area A were lower than those in Area B by  $-3.61 \pm 0.70 \mu atm$  and  $\Omega_{ent}-0.09 \pm 0.01$  (Ave.  $\pm$  SE), respectively. Furthermore, the pH was comparable to that in Area B ( $+3.7 \pm 0.7$ )  $\times 10^{-3}$  than Area B, and both seawater fCO<sub>2</sub> and  $\Omega_{ent}$ -All parameters showed clear seasonal variations.

Formatted: Font: (Default) +Body (Times New Roman)

Formatted: Font: (Default) +Body (Times New Roman), Subscript

Formatted: Font: (Default) +Body (Times New Roman)

Formatted: Font: (Default) +Body (Times New Roman), Not Superscript/ Subscript

Formatted: Font: (Default) +Body (Times New Roman)

Formatted: Font: (Default) +Body (Times New Roman)


Formatted: Font: (Default) +Body (Times New Roman)


Formatted: Font: (Default) +Body (Times New Roman)

Formatted: Font: +Body (Times New Roman)

Formatted: Superscript







**Figures 5. (a): Temporal contributions in the PLS analysis by of Sea Surface Temperature (SST), Sea Surface Salinity (SSS), normalized Total Alkalinity (nTA), and normalized Dissolved Inorganic Carbon (nDIC) in the PLS analysis-SST, SSS, nTA, and nDIC to the difference between Area A and B in (a-c) in-seawater  $\text{CO}_2$  fugacity ( $\text{fCO}_2$ ), (d-f) pH-between-Area-A-and-B, and (g-i) aragonite saturation state ( $\Omega_{\text{arg}}$ ).** Panels show (a, d, g) the temporal contributions, (b, e, h) monthly averages, and (c, f, i) yearly averages. Red lines indicate the sum of contributions, which closely matches the temporal variation of the differences ( $d_{AB}$ ) in each variables. The corresponding monthly averages. (e): The corresponding yearly averages. (d): Temporal contributions in the PLS analysis by SST, SSS, nTA, and nDIC to the difference in  $\Omega_{\text{calc}}$  between Area A and B. (e): The monthly averages. (f): The yearly averages. The red lines represent the sum of each contribution and is almost equal to temporal variation of  $d_{AB}$  of seawater  $\text{fCO}_2$  and  $\Omega_{\text{calc}}$ .

Formatted: Subscript

Formatted: Not Superscript/ Subscript

Formatted: Not Superscript/ Subscript

Formatted: Font: Italic

#### 4. Discussion

The extent of Area A defined using nTA (Figure 2f) was consistent with the low-salinity area of the Pacific coast of Japan (Figure 1a). The SSS south of 37°N exhibited an abrupt change in value 205.7410 km from mainland Japan (Figure S4), which is consistent with the distance of the boundary between Areas A and B (250 km). Because SSS is affected by precipitation and evaporation, the distance defined by nTA is considered more accurate as the boundary for Area A; therefore, nTA was a more accurate indicator of riverine water than SSS. However, the consistency between the two distances results provides strong evidence that the nTA in Area A was derived from riverine water is an effective indicator for riverine water. Because these distances were calculated statistically using 20 years of observational data on SSS and nTA, it would be challenging to obtain a similarly significant result from single observations. Therefore, the findings of this study, which quantify the extent of the influence of riverine water, are novel and offer new insights.

To evaluate the effect of the Kuroshio Current on riverine water distribution, another EOF analysis was performed using data from the Kuroshio Large Meander period (2017–2020 in this study) in which the Kuroshio path followed an alternative meandering path south of its usual course (Kawabe, 1985). Area A was within the range of 350 km from land, utilizing the Mode 1 EOF anomaly during the meander period, whereas it was 250 km during the entire period. This result was consistent with the observed Kuroshio path meandering path southwards, that is, away from the Pacific coast of Japan, and supports the assumption that the Kuroshio Current effects Area A. However, Area A extended further out of the open ocean than the Kuroshio axis during the entire study period (Figure 2f). Therefore, the Kuroshio Current limited the spread of riverine water to some extent but did not completely inhibit it.

The supply of TA by riverine water was higher than the time deviation of  $dAB$  for almost the entire period, suggesting that the supply was one of the main factors contributing to the increase in  $dAB$  (Figure 4). On the other hand, the annual rate of increase in the  $dAB$  of nTA was almost zero ( $-0.02 \pm 0.05 \mu\text{mol kg}^{-1} \text{ year}^{-1}$ ). Therefore, the residual term would be the negative value of the same scale with  $Sup_{rw}$  term for TA. Horizontal advection is assumed to be the primary cause of the negative residual term is assumed to be horizontal advection. The advection This effect can be estimated from using the product of the current velocity and the  $dAB$  of nTA. The fact that the period of increasing  $dAB$  of nTA (Figure 3c) is consistent coincides with the period of increasing absolute value of the residual term absolute value in (Figure 4b) supports this assumption. In addition On the other hand, the effect of vertical advection estimated from the vertical gradient of nTA and mixed layer depth was not significant could not be the main cause of the negative residual term, because the vertical gradient for calculating vertical advection was small and insignificant (see Text S1). Other causes of the residual term are differences in biological metabolism biological activity, such as calcification and, nitrate consumption, and organic Alkalinity. Calcification by coccolithophores is one of the most promising important process for changing oceanic nTA driving changes in oceanic nTA. However, no regional difference in calcification rate of up to  $10 \mu\text{mol kg}^{-1} \text{ year}^{-1}$  (Figure 4b) has not been reported between the Japanese coastal Area A area (Area A) and the surrounding sea area (Area B) (Hopkins and Balch, 2018; Krumhardt et al., 2019). Another component of TA include nitrate, which is consumed during

Formatted: Font: (Default) +Body (Times New Roman)

photosynthesis to raise nTA (Brewer and Goldman, 1976). The data on total nitrate (TN) concentration is available in  
330 GLODAP, and the mean values were  $1.19 \pm 0.11 \mu\text{mol kg}^{-1}$  and  $0.34 \pm 0.04 \mu\text{mol kg}^{-1}$  in Area A and B, respectively. As the  
monthly difference were largest in the winter ( $2.11 \mu\text{mol kg}^{-1}$ ) and almost reach zero in the summer, the main source of  
nitrate would be vertical advection rather than input from land. ~~Nonehteleess~~Nonetheless, the effect of nitrate on TA was  
probably one order of magnitude smaller than that of the riverine water supply and is negligible as the main source of nTA.  
The inflow of organic Alkalinity is assumed to be another component influencing nTA fluctuations (Cai et al., 1998; Song et  
335 al., 2020; Yang et al., 2015). Organic Alkalinity is the TA component that originates from organic acids such as humic acid,  
found in terrestrial sources like wetlands and coastal sediments. Although no data on organic Alkalinity from mainland Japan  
could be found, a reference on organic Alkalinity from rivers in southern coast of China reports that the maximum value for  
organic Alkalinity was  $18 \mu\text{mol kg}^{-1}$  under condition of salinity exceeding 30 like the analysis in this study (Song et al.,  
340 2023). Assuming the outflow of organic Alkalinity from mainland Japan is of a similar order of magnitude, this represents  
less than 1% of the oceanic TA. Furthermore, compared to the upper and lower limits of the TA concentrations in the  
associated rivers ( $518\text{--}1006 \mu\text{mol kg}^{-1}$ ), this is a sufficiently small. Therefore, in this analysis, the influence of organic  
Alkalinity was considered negligible. However, it will have a significant impact on future TA budget analysis conducted  
under lower salinity conditions or in environments with abundant wetlands, such as those found in northern Japan, including  
Hokkaido.

345 ~~▲~~  
—The DIC variation is more complicated than the TA variation because the spatiotemporal variation in ~~biological~~  
~~metabolism~~biological activity and vertical advection is larger and more significant than ~~thatthat~~ of TA, in addition to the  
effect of CO<sub>2</sub> exchange with the atmosphere. Unlike the TA, the seasonal variations in the riverine water supply were not  
consistent with those in the time derivative (Figure 4e). This trend indicated that riverine water supply was not the primary  
350 driver of DIC variation in Area A. The effect of oceanic CO<sub>2</sub> uptake was highly variable and unclear on a decadal scale.  
However, on a seasonal scale, it was more distinct, resulting in negative shifts in the summer residual terms and positive  
shifts in the winter residual terms. This seasonal variation indicated that the oceanic CO<sub>2</sub> uptake in Area A was larger than  
that in Area B in summer, and vice versa in winter. ~~This was caused by the lower seawater fCO<sub>2</sub> in Area A during the~~  
~~summer season and higher wind velocity in Area B during the winter season.~~ Consequently, the residual term of nDIC  
355 indicated a strong and complex seasonal variation, which included a maximum in the winter and two minimums in spring  
and summer with a difference of  $100 \mu\text{mol kg}^{-1} \text{ year}^{-1}$ . Contrary to TA, DIC was affected by nutrient and organic matter  
loading from terrestrial sources, and was strongly affected by differences in ~~biological metabolism~~biological activity  
between Areas A and B. ~~Furthermore, For DIC, in contrast to TA,~~ a significant ~~vertical profile trend~~ in the vertical profile of  
DIC was observed (Text S1). ~~Consequently, some of the residual terms mayight have been influenced-affected by vertical~~  
360 advection.

To quantify the effect of ~~biological-metabolism~~biological activity, the effects of horizontal and vertical advection were  
estimated based on the following assumptions: 1) the residual term for nTA ( $C_{\text{res TA}}$ ) was assumed to be equal to the

Formatted: Font: (Default) +Body (Times New Roman)

Formatted: Superscript

Formatted: Font: (Asian) MS 明朝, (Asian) Japanese, (Other) English (United States)

horizontal advection term, based on the considerations that the effects of vertical advection and ~~biological-metabolism~~ ~~mother effects~~ on nTA are negligible. 2) The horizontal advection terms ~~for nTA and nDIC~~ ~~were~~as proportional to  $dAB$  of nTA ( $dAB_{TA}$ ) and nDIC ( $dAB_{DIC}$ ), ~~respectively~~. Several previous studies have used normalized values for the calculation of TA and DIC advection (Broecker and Peng, 1992; Keeling and Peng, 1995) and have also proven their validity as an approximation (Robbins, 2001). Because the advection term is calculated as the product of concentration gradient and flow field, and the associated distance and flow velocity are the same at  $dAB_{TA}$  and  $dAB_{DIC}$ , it can be assumed that the horizontal advection effect on DIC ( $C_{hadv\_DIC}$ ) can be estimated using the following equation:

$$C_{hadv\_DIC} = K \frac{dAB_{DIC}}{D_{AB}}$$

$$C_{res\_TA} = K \frac{dAB_{TA}}{D_{AB}}$$

$$C_{hadv\_DIC} = C_{res\_TA} \cdot \frac{dAB_{DIC}}{dAB_{TA}} \quad (5)$$


where,  $K$  and  $D_{AB}$  are the index values associated with the current field and horizontal distance between Areas A and B, respectively. These parameters were assumed to be the same for TA and DIC calculations. 3) Vertical advection for nDIC ( $C_{vhadv\_DIC}$ ) was estimated from mixing with subsurface water when the mixed layer was deepened. Specifically, we used a simplified equation of the method of a previous study (Ishii et al., 2001).

$$\int C_{vhadv\_DIC}(t)dt = \frac{\{\Delta nDIC(t+1) + \Delta nDIC(t)\}}{2} \times \frac{\{MLD(t+1) - MLD(t)\}}{\{MLD(t+1) \cdot \rho_{MLD(t+1)\}}$$

$$\Delta nDIC(t) = \{nDIC(July, MLD(t)) - nDIC(t, MLD(t))\} \cdot \rho_{MLD(t)} \quad (6)$$

where,  $\Delta nDIC(t)$  is the difference between the average nDIC in July ( $nDIC(July)$ ) and the nDIC at time  $t$  at mixed layer depth ( $MLD(t)$ ) multiplied by the seawater density at  $MLD(t)$  ( $\rho_{MLD(t)}$ ). The nDIC profile at a depth below the mixed layer was assumed to be maintained at its average value in July; as the mixed layer deepened, the difference from the July profile was added to the nDIC in the mixed layer. The effect of vertical advection on nDIC was negligible when the mixed-layer depth did not change or became shallower. The details on the calculation of the nDIC profiles are provided in Text S1.

The effects of horizontal and vertical advection on nDIC are shown in Figure 6. However, due to the large error range in the vertical advection term for each time step (the median was 137%), only the monthly and annual averages are shown in the figure. The horizontal and vertical advection effects were identified mainly from autumn to winter with an annual mean value of  $-15.55 \pm 1.35 \mu\text{mol kg}^{-1} \text{ year}^{-1}$  and  $15.55 \pm 2.25 \mu\text{mol kg}^{-1} \text{ year}^{-1}$ , respectively ( $Ave \pm SE$ ). Seasonally, the residual term had a maximum in the winter and two minimums in spring and summer, with an overall mean of  $-7.30 \pm 3.24 \mu\text{mol kg}^{-1} \text{ year}^{-1}$ .



**Figures 6. Temporal variations of normalized Dissolved Inorganic Carbon (nDIC) time derivative and related terms in Equation (2) (a): The monthly averages. (b): The yearly averages. The shaded areas of horizontal advection (h-advection) and vertical advection (v-advection) are the random error calculated using the error propagation (see Text S1). Therefore, unlike the error ranges of riverine supply ( $Sup_{riv}$ ), these were reduced by  $10^{-0.5}$  or  $20^{-0.5}$  by monthly or yearly averaging, respectively.**

Although the above results were not definitive because of many assumptions and large error ranges, the fact that the final residual term on the decadal scale was almost negative supports the idea that Area A is likely under heterotrophic conditions within the mixed layer. The positive shift in the residual term before and after 2007 was due to the low vertical advection term caused by the small MLD during winter. The 2007 minimum SST in Area A ( $19.41^{\circ}\text{C}$ ) was the highest of the overall annual minimum SST ( $17.71 \pm 0.66^{\circ}\text{C}$ , Ave  $\pm$  SD), and thus the surface seawater mixing would be weaker than that in other years. Therefore, the residual term for this period does not necessarily indicate changes in biological-metabolismbiological activity. However, the high SST during winter suggests that the decomposition of biofixed carbon may be accelerated. Seasonally, the March-April minimum in the residual term was assumed to be the result of phytoplankton blooms in the open ocean area in Area A. Another minimum in July-August was assumed to be due to the primary production of phytoplankton

Formatted: Font: (Default) +Body (Times New Roman)

Formatted: Font: (Default) +Body (Times New Roman), Not Italic, Not Superscript/ Subscript

Formatted: Font: (Default) +Body (Times New Roman)

~~and submerged aquatic vegetation~~ under eutrophic conditions in the inner bays and near-shore areas. The winter maximum can be explained by the upwelling of organic matter and the decomposition of bio-fixed carbons within the mixed layer.

It should be noted that the ~~DIC~~ residual term ~~above~~ includes the effects of the water CO<sub>2</sub> flux in near-shore areas. However, unlike the air-sea CO<sub>2</sub> flux in the ocean ( $C_{flux}$  in Equation (2)), the air-water CO<sub>2</sub> flux in near-shore areas is very difficult to quantify because observations of ~~water-coastal~~ fCO<sub>2</sub> and the physical regulating factor defined as “transfer velocity” (e.g., Wanninkhof, 2014) are limited. The estimation of this CO<sub>2</sub> flux varies widely among the existing studies. For example, global average models (Aufdenkamp et al., 2011; Tranvik et al., 2009) have estimated that approximately half of the carbon supply from land is released into the atmosphere via the air-water CO<sub>2</sub> flux in near-shore areas. However, the inner bays affected by riverine water in this study showed a trend in atmospheric CO<sub>2</sub> absorption (Tokoro et al., 2021), that is unlikely to follow the global average trend. Therefore, enhanced observation and analysis of atmospheric CO<sub>2</sub> exchange in nearshore areas is essential for a more accurate assessment of the residual term as an indicator of ~~biological-metabolism~~~~biological activity~~.

Despite Area A having a higher nDIC than Area B (Figure 3), seawater fCO<sub>2</sub> in Area A tended to be lower than in Area B ( $-3.61 \pm 0.70 \mu\text{atm}$ ). This was mainly because of the lower SST and higher nTA in Area A than those in Area B (Figure 5). The low SST can likely be attributed to riverine water, which is mainly supplied by snowmelt and rainfall from the mountains and highlands and therefore tends to have lower original water temperatures. In particular, the short flow paths of Japanese rivers can likely limit the effects of heating due to solar radiation and other factors. The three inner bays with strong river influence had lower average water temperature ( $18.66\text{--}19.23^\circ\text{C}$ , Tokoro et al., 2021) than areas A ( $22.02^\circ\text{C}$ ) and B ( $22.86^\circ\text{C}$ ), which support our assumption. The decrease in ~~dAB~~ of seawater fCO<sub>2</sub> peaked in the summer ( $-15.90 \pm 4.72 \mu\text{atm}$  from July to September), when the decrease in nDIC due to ~~biological-metabolism~~~~biological activity~~ and the increase in nTA due to river supply coincided. However, this decreasing in seawater fCO<sub>2</sub> had little effect on annual oceanic CO<sub>2</sub> uptake in Area A. Compared to the hypothetical case where riverine water did not affect seawater fCO<sub>2</sub> (seawater fCO<sub>2</sub> in Area A were the same as that in Area B), the change in air-sea CO<sub>2</sub> flux would be similar ( $0.00 \pm 0.33 \text{ mol m}^{-2} \text{ year}^{-1}$ ). This was because the decrease in air-sea CO<sub>2</sub> flux in summer almost ~~offsets~~~~offsets~~ the increase in winter. Although the increase in seawater fCO<sub>2</sub> in winter ( $+7.25 \pm 1.12 \mu\text{atm}$  from December to February) was smaller than the decrease in summer ( $-15.90 \pm 4.72 \mu\text{atm}$  from July to September), the air-sea CO<sub>2</sub> fluxes in summer and winter were coincidentally balanced, owing to higher wind speeds in winter.

In the evaluation of acidification, different trends were observed for pH and  $\Omega_{arg}$ . While pH showed little difference between Areas A and B ( $+3.7 \pm 1.4 \times 10^{-3}$ ),  $\Omega_{arg}$  revealed a clear acidification trend in Area A. Regarding pH, the acidification trend in Area A caused by nDIC increase appears to be offset by the above-mentioned low SST effect and the nTA inflow (Figure 5). Meanwhile, for  $\Omega_{arg}$ , seawater in Area A was notably more acidified than in the surrounding sea area, based on a 20-year average ( $-0.43\text{--}09 \pm 0.01$ ). This result is an example where pH and  $\Omega_{arg}$  exhibit different behaviours, demonstrating that acidification by riverine water has little effect on hydrogen ion concentration, but exerts a significant influence on biological activities such as calcification. A similar trend is expected to occur in coastal areas north of the temperate zone where

Formatted: Font: Italic

Formatted: Font: (Default) +Body (Times New Roman)

riverine water cooling takes place. Although the inflow of nTA has a small effect on pH (+0.01), ~~The impact on coastal acidification was particularly significant in winter, with an average decrease of -0.25 (5.33 to 5.08).~~ nTA was only mitigation factor for  ~~$\Omega_{arg}$  coastal acidification~~ among the explanatory variables, and reduced acidification due to other factors to ~~6571%~~ (-0.20-13 to -0.1309). ~~Although no significant decadal change in pH of  $dAB$  was observed regarding acidification, This~~ coastal acidification indicated by  $\Omega_{arg}$  was found to be progressing, with  $dAB$  of  $\Omega_{enl-arg}$  decreasing by  $-0.52-04 \pm 0.0144$  per decade. This can be attributed to an increasing trend in  $dAB$  of nDIC ( $0.32 \pm 0.08 \mu\text{mol kg}^{-1} \text{year}^{-1}$ ) due to the trend of increasing precipitation in Japan. ▲

Formatted: Font: Italic

Formatted: Font: (Asian) MS 明朝, (Asian) Japanese, (Other) English (United States)

## 5. Conclusions

In this study, we identified areas affected by riverine water in the Northwest Pacific Ocean by using statistically processed observational data. We also evaluated the contribution of riverine water to oceanic CO<sub>2</sub> uptake and coastal acidification by comparing the TA and DIC values in the sea area affected by riverine water and surrounding sea areas.

The Area A affected by riverine water (Area A) was within 250 km of mainland Japan and, to some extent, along the Kuroshio axis. This area was consistent with the low-SSS area on the Pacific coast of Japan. In addition, the range increased to 350 km during the period when the Kuroshio Current meandered south, indicating that the Kuroshio Current path influences the spread of riverine water.

Both nTA and nDIC were higher ~~in surrounding sea of in~~ Area A compared with the surrounding sea area. The main source of TA was riverine water in summer, which was balanced by a decrease due to horizontal advection in autumn and winter. The DIC flows were more complex than the TA flows ~~because they were influenced not only by horizontal advection but also by vertical advection and biological activities such as photosynthesis and remineralization. Consequently, the residual term, which integrates these unquantified processes, had a greater influence on the DIC budget than the direct riverine water supply, and were more strongly affected by a residual term than by the riverine water supply.~~ The contribution of ~~biological metabolism~~ biological activity was suggested is expected to be ~~more influential~~ the primary factor on in the residual term ~~after by quantification excluding~~ the horizontal and vertical advection effects. ~~Biological metabolism~~ Biological activity showed a maximum in winter and two minima in spring and summer. The annual mean suggests heterotrophic conditions within the mixed layer in Area A. In any case, because the air-water CO<sub>2</sub> flux in near-shore areas is still difficult to estimate, a more thorough quantification of DIC flow in relation to riverine water needs to be considered. ▲

Formatted: Font: (Asian) MS 明朝, (Asian) Japanese

Seawater fCO<sub>2</sub> in Area A decreased mainly in summer because of the supplied riverine water, which has low-temperature water with high nTA. However, riverine water supply was found to have virtually no effect on oceanic CO<sub>2</sub> uptake at an annual scale. This is because CO<sub>2</sub> emission was enhanced by strong wind speeds in winter, in addition to the relatively small increase in seawater fCO<sub>2</sub> in winter. The change in the air-sea CO<sub>2</sub> flux in winter was coincidentally balanced by the change in summer, and the annual average was almost zero. In Area A, the trend of progressive acidification relative to the



475 surrounding sea area was not confirmed by pH analysis but was confirmed by  $\Omega_{arg}$ . TAlthough acidification progressed more  
in Area A than in the surrounding sea area, the supply of TA fromby riverine water mitigated the  $\Omega_{arg}$  acidification to  
6571%. However, an increase in precipitation may have led to increases-higher level of in nDIC and acidification in Area A.  
This study makes a significant contribution to the analysis of carbon flows in the boundary seawater between terrestrial and  
oceanic areas because the quantification of carbon flows in these areas is often uncertain in space and time. Enhancing the  
observational data allows the results to be spatially extended to larger regional or global scales. The analysis results are  
expected to have a smaller error range owing to the high spatiotemporal resolution of the vertical profiles of the carbonate  
480 data and air-water CO<sub>2</sub> flux data in near-shore areas.

Formatted: Not Superscript/ Subscript

Formatted: Font: (Asian) MS 明朝, (Asian) Japanese

490 **Data Availability**

The SSS, SST, and fCO<sub>2</sub> in air and seawater datasets are available in the SOCAT database (Pfeil et al., 2013; Bakker et al.,  
2016, <https://socat.info/index.php/data-access/>). The TA and DIC data were available from the GLODAP database version  
485 2.2020 (Key et al., 2015; Olsen et al., 2016; Olsen, 2020, <https://glodap.info/>). Wind speed data were obtained from the  
CCMP database (version 2.0) (Atlas et al., 2011; Mears et al., 2019; <https://data.remss.com/ccmp/v02.0/>). The river flow  
datasets were obtained from the Water Information System of the Ministry of Land, Infrastructure, Transport, and Tourism of  
Japan (<http://www1.river.go.jp/>). Vertical water temperature datasets for calculating MLD were obtained from the JCOPE2M  
dataset (Miyazawa et al., 2017, 2019, . <https://www.jamstec.go.jp/jcope/htdocs/distribution/>).

**Author Contribution**

TT designed the study and wrote the first draft on the manuscript. SN, ST, SS, SD, KE, MI, NK, TO, KT, and YN supplied  
and managed data for SOCAT and GLODAP. SN supported data processing. All authors contributed to manuscript writing  
and proofing.

495 **Competing interests**

The contact author has declared that none of the authors have any competing interests.

**Acknowledgments**

500 We appreciate the cooperation of Toyofuji Shipping Co. and Kagoshima Senpaku Co. in the NIES VOS program. We also  
thank the captains and crews of M/S Pyxis, M/S New Century 2, M/S Trans Future 5, R/V Mirai, R/V Hakuho-maru, R/V  
Keifu-maru, R/V Ryofu-maru, R/V Wakataka-maru, and R/V Soyo-maru. This research was financially supported by the  
Global Environmental Research Coordination System, Ministry of the Environment, Japan (grant numbers MOE1751 and  
MOE 2252) and CREST, Japan Science and Technology Agency (grant number JPMJCR23J4).

505

## References

Andreev, A. G. and Pavlova, G. Y.: Marginal Seas, in: Carbon and nutrient fluxes in continental margins, edited by Liu, K. K., Atkinson, L., Quiñones, R., and Talaue-McManus, L., Springer, New York, 395–406, 2009.

- 510 Atlas, R., Hoffman, R. N., Ardizzone, J., Leidner, S. M., Jusem, J. C., Smith, D. K., and Gombos, D.: A Cross-calibrated Multiplatform Ocean Surface Wind Velocity Product for Meteorological and Oceanographic Applications, *Bull. Amer. Meteor. Soc.*, 92, 157–174. <https://doi.org/10.1175/2010BAMS2946.1>, 2011.
- Aufdenkampe, A. K., Mayorga, E., Raymond, P. A., Melack, J. M., Doney, S. C., Alin, S. R., Aalto, R. E., and  
525 Yoo, K.: Riverine coupling of biogeochemical cycles between land, oceans, and atmosphere, *Front. Ecol. Environ.*, 9, 53–60. <https://doi.org/10.1890/100014>, 2011.
- Bakker, D. C. E., Pfeil, B., Landa, C. S., Metzl, N., O'Brien, K. M., Olsen, A., Smith, K., Cosca, C., Harasawa, S., Jones, S. D., Nakaoka, S., Nojiri, Y., Schuster, U., Steinhoff, T., Sweeney, C., Takahashi, T., Tilbrook, B.,  
520 Wada, C., Wanninkhof, R., Alin, S. R., Balestrini, C. F., Barbero, L., Bates, N. R., Bianchi, A. A., Bonou, F., Boutin, J., Bozec, Y., Burger, E. F., Cai, W. J., Castle, R. D., Chen, L., Chierici, M., Currie, K., Evans, W., Featherstone, C., Feely, R. A., Fransson, A., Goyet, C., Greenwood, N., Gregor, L., Hankin, S., Hardman-Mountford, N. J., Harlay, J., Hauck, J., Hoppema, M., Humphreys, M. P., Hunt, C. W., Huss, B., Ibáñez, J. S. P., Johannessen, T., Keeling, R., Kitidis, V., Körtzinger, A., Kozyr, A., Krasakopoulou, E., Kuwata, A.,  
525 Landschützer, P., Lauvset, S. K., Lefèvre, N., Lo Monaco, C., Manke, A., Mathis, J. T., Merlivat, L., Millero, F. J., Monteiro, P. M. S., Munro, D. R., Murata, A., Newberger, T., Omar, A. M., Ono, T., Paterson, K., Pearce, D., Pierrot, D., Robbins, L. L., Saito, S., Salisbury, J., Schlitzer, R., Schneider, B., Schweitzer, R., Sieger, R., Skjelvan, I., Sullivan, K. F., Sutherland, S. C., Sutton, A. J., Tadokoro, K., Telszewski, M., Tuma, M., van Heuven, S. M. A. C., Vandemark, D., Ward, B., Watson, A. J., and Xu, S.: A multi-decade record of high-quality  
530 fCO<sub>2</sub> data in version 3 of the Surface Ocean CO<sub>2</sub> Atlas (SOCAT), *Earth Syst. Sci. Data*, 8, 383–413. <https://doi.org/10.5194/essd-8-383-2016>, 2016.
- Bauer, J. E., Cai, W. J., Raymond, P. A., Bianchi, T. S., Hopkinson, C. S., and Regnier, P. A. G.: The changing carbon cycle of the coastal ocean, *Nature*, 504, 61–70. <https://doi.org/10.1038/nature12857>, 2013.

Borges, A. V., Delille, B., and Frankignoulle, M.: Budgeting sinks and sources of CO<sub>2</sub> in the coastal ocean: Diversity of ecosystems counts, *Geophys. Res. Lett.*, 32, L14601. <https://doi.org/10.1029/2005GL023053>, 2005.

Brewer, P. G. and Goldman, J. C.: Alkalinity changes generated by phytoplankton growth, *Limnol. Oceanogr.*, 21, 108–117. <https://doi.org/10.4319/lo.1976.21.1.0108>, 1976.

Broecker, W. S. and Peng, T. H.: Tracers in the sea, Eldigio Press, New York, 1982.

Broecker, W. S. and Peng, T. H.: Interhemispheric transport of carbon dioxide by ocean circulation, *Nature*, 356, 587–589. <https://doi.org/10.1038/356587a0>, 1992.

Cai, W. J. and Wang, Y.: The chemistry, fluxes, and sources of carbon dioxide in the estuarine waters of the Satilla and Altamaha Rivers, Georgia, *Limnol. and Oceanogr.*, 43(4), 657-668. <https://doi.org/10.4319/lo.1998.43.4.0657>, 1998.

Cai, W. J., Wang, Y., and Hodson, R. E.: Acid-base properties of dissolved organic matter in the estuarine waters of Georgia, USA, *Geochim. Cosmochim. Acta*, 62(3), 473-483. [https://doi.org/10.1016/S0016-7037\(97\)00363-3](https://doi.org/10.1016/S0016-7037(97)00363-3), 1998.

▲

Cai, W. J.: Estuarine and coastal ocean carbon paradox: CO<sub>2</sub> sinks or sites of terrestrial carbon incineration? *Ann. Rev. Mar. Sci.*, 3, 123–145. <https://doi.org/10.1146/annurev-marine-120709-142723>, 2011.

Carstensen, J. and Duarte, C. M.: Drivers of pH variability in coastal ecosystems, *Environ. Sci. Technol.*, 53, 4020–4029. <https://doi.org/10.1021/acs.est.8b03655>, 2019.

Chen, C. T. A. and Borges, A. V.: Reconciling opposing views on carbon cycling in the coastal ocean: Continental shelves as sinks and near-shore ecosystems as sources of atmospheric CO<sub>2</sub>, *Deep Sea Research Part II: Topical Studies in Oceanography*, 56, 578–590. <https://doi.org/10.1016/j.dsr2.2009.01.001>, 2009.

Chen, C. T. A., Huang, T. H., Chen, Y. C., Bai, Y., He, X., and Kang, Y.: Air-sea exchanges of CO<sub>2</sub> in the world’s coastal seas, *Biogeosciences*, 10, 6509–6544. <https://doi.org/10.5194/bg-10-6509-2013>, 2013.

Formatted: Default Paragraph Font, Font: 10 pt

Formatted: Default Paragraph Font, Font: 10 pt

Formatted: Font: (Asian) MS 明朝, (Asian) Japanese

de Boyer Montégut, C., Madec, G., Fischer, A. S., Lazar, A., and Iudicone, D.: Mixed layer depth over the global ocean: An examination of profile data and a profile-based climatology, *J. Geophys. Res.*, 109.

570 <https://doi.org/10.1029/2004JC002378>, 2004.

Denbo, D. W. and Allen, J. S.: Rotary empirical orthogonal function analysis of currents near the Oregon Coast, *J. Phys. Oceanogr.*, 14, 35–46. [https://doi.org/10.1175/1520-0485\(1984\)014<0035:REOFAO>2.0.CO;2](https://doi.org/10.1175/1520-0485(1984)014<0035:REOFAO>2.0.CO;2), 1984.

585 Duarte, C. M., Hendriks, I. E., Moore, T. S., Olsen, Y. S., Steckbauer, A., Ramajo, L., Carstensen, J., Trotter, J. A., and McCulloch, M.: Is ocean acidification an open-ocean syndrome? Understanding anthropogenic impacts on seawater pH, *Estuaries Coasts*, 36, 221–236. <https://doi.org/10.1007/s12237-013-9594-3>, 2013.

Fitzer, S. C., Torres Gabarda, S. T., Daly, L., Hughes, B., Dove, M., O'Connor, W., Potts, J., Scanes, P., and  
580 Byrne, M.: Coastal acidification impacts on shell mineral structure of bivalve mollusks, *Ecol. Evol.*, 8, 8973–8984. <https://doi.org/10.1002/ece3.4416>, 2018.

[Friedlingstein, P., O'Sullivan, M., Jones, M. W., Andrew, R. M., Hauck, J., Landschützer, P., Le Quéré, C., Li, H., Luijkx, I. T., Olsen, A., Peters, G. P., Peters, W., Pongratz, J., Schwingshackl, C., Sitch, S., Canadell, J. G., Ciais, P., Jackson, R. B., Alin, S. R., Arneth, A., Arora, V., Bates, N. R., Becker, M., Bellouin, N., Berghoff, C. F., Bittig, H. C., Bopp, L., Cadule, P., Campbell, K., Chamberlain, M. A., Chandra, N., Chevallier, F., Chini, L. P., Colligan, T., Decayeux, J., Djeuthouang, L. M., Dou, X., Duran Rojas, C., Enyo, K., Evans, W., Fay, A. R., Feely, R. A., Ford, D. J., Foster, A., Gasser, T., Gehlen, M., Gkritzalis, T., Grassi, G., Gregor, L., Gruber, N., Gürses, Ö., Harris, I., Hefner, M., Heinke, J., Hurtt, G. C., Iida, Y., Ilyina, T., Jacobson, A. R., Jain, A. K., Jarníková, T., Jersild, A., Jiang, F., Jin, Z., Kato, E., Keeling, R. F., Klein Goldewijk, K., Knauer, J., Korsbakken, J. I., Lan, X., Lauvset, S. K., Lefèvre, N., Liu, Z., Liu, J., Ma, L., Maksyutov, S., Marland, G., Mayot, N., McGuire, P. C., Metzl, N., Monacchi, N. M., Morgan, E. J., Nakaoka, S., Neill, C., Niwa, Y., Nützel, T., Olivier, L., Ono, T., Palmer, P. I., Pierrot, D., Qin, Z., Resplandy, L., Roobaert, A., Rosan, T. M., Rödenbeck, C., Schwinger, J., Smallman, T. L., Smith, S. M., Sospedra-Alfonso, R., Steinhoff, T., Sun, Q., Sutton, A. J., Sférian, R., Takao, S., Tatebe, H., Tian, H., Tilbrook, B., Torres, O., Tourigny, E., Tsujino, H., Tubiello, F., van der Werf, G., Wanninkhof, R., Wang, X., Yang, D., Yang, X., Yu, Z., Yuan, W., Yue, X., Zaehle, S., Zeng, N., and Zeng, J.:](#) Global Carbon Budget 2024, *Earth Syst. Sci. Data*, 17, 965–1039. <https://doi.org/10.5194/essd-17-965-2025>, 2025.

595

Formatted: Default Paragraph Font, Font: 10 pt

▲  
Friis, K., Körtzinger, A., and Wallace, D. W. R.: The salinity normalization of marine inorganic carbon chemistry  
600 data, *Geophys. Res. Lett.*, 30. <https://doi.org/10.1029/2002GL015898>, 2003.

Holte, J. and Talley, L.: A new algorithm for finding mixed layer depths with applications to argo data and  
Subantarctic mode water formation, *J. Atmos. Ocean. Technol.*, 26, 1920–1939.  
<https://doi.org/10.1175/2009JTECHO543.1>, 2009.

605 Hopkins, J. and Balch, W. M.: A new approach to estimating coccolithophore calcification rates from space, *JGR  
Biogeosciences.*, 123, 1447–1459. <https://doi.org/10.1002/2017JG004235>, 2018.

Inoue, H. Y., Matsueda, H., Ishii, M., Fushimi, K., Hirota, M., Asanuma, I., and Takasugi, Y.: Long-term trend of  
610 the partial-pressure of carbon-dioxide (pCO<sub>2</sub>) in surface waters of the western North Pacific, 1984–1993, *Tellus  
B.*, 47, 391–413. <https://doi.org/10.1034/j.1600-0889.47.issue4.2.x>, 1995.

Ishii, M., Inoue, H. Y., Matsueda, H., Saito, S., Fushimi, K., Nemoto, K., Yano, T., Nagai, H., and Midorikawa,  
T.: Seasonal variation in total inorganic carbon and its controlling processes in surface waters of the western  
615 North Pacific subtropical gyre, *Mar. Chem.*, 75, 17–32. [https://doi.org/10.1016/S0304-4203\(01\)00023-8](https://doi.org/10.1016/S0304-4203(01)00023-8), 2001.

[Kakei, S., Ito, S., Wagawa, T.: Estimating surface water mixing ratios using salinity and potential alkalinity in the  
Kuroshio-Oyashio mixed water regions, \*JGR oceans\*, 122, 1927-1942. <https://doi.org/10.1002/2016JC012268>, 2017.](https://doi.org/10.1002/2016JC012268)

620 Kawabe, M.: Sea level variations at the Izu Islands and typical stable paths of the Kuroshio, *J. Ocean. Soc. Jpn.*,  
41, 307–326. <https://doi.org/10.1007/BF02109238>, 1985.

Keeling, R. and Peng, T. H.: Transport of heat, CO<sub>2</sub>, and O<sub>2</sub> by the Atlantic’s thermohaline circulation, *Phil.  
Trans. R. Soc. Lond. B*, 348, 133–142. <https://doi.org/10.1098/rstb.1995.0055>, 1995.

625 Kessouri, F., McWilliams, J. C., Bianchi, D., Sutula, M., Renault, L., Deutsch, C., Feely, R. A., McLaughlin, K.,  
Ho, M., Howard, E. M., Bednaršek, N., Damien, P., Molemaker, J., and Weisberg, S. B.: Coastal eutrophication

Formatted: Font: (Asian) MS 明朝, (Asian) Japanese

Formatted: Default Paragraph Font, Font: 10 pt

Formatted: Font: (Asian) MS 明朝, (Asian) Japanese

drives acidification, oxygen loss, and ecosystem change in a major oceanic upwelling system, *Proc. Natl Acad. Sci. U. S. A.*, 118. <https://doi.org/10.1073/pnas.2018856118>, 2021.

630 Key, R. M., Olsen, A., van Heuven, S., Lauvset, S. K., Velo, A., Lin, X., Schirnick, C., Kozyr, A., Tanhua, T., Hoppema, M., Jutterström, S., Steinfeldt, R., Jeansson, E., Ishii, M., Perez, F. F., and Suzuki, T.: Global ocean data analysis project, version 2, (GLODAPv2), Oak Ridge National Laboratory/CDIAC-162, NDP-093, Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, United States Department of Energy, Oak Ridge, TN. [https://doi.org/10.3334/CDIAC/OTG\\_NDP093\\_GLODAPv2](https://doi.org/10.3334/CDIAC/OTG_NDP093_GLODAPv2). [dataset] 2015.

635 Killick, R., Fearnhead, P., and Eckley, I. A.: Optimal detection of changepoints with a linear computational cost, *J. Am. Stat. Assoc.*, 107, 1590–1598. <https://doi.org/10.1080/01621459.2012.737745>, 2012.

640 Kitamura, T., Nakano, T., and Sugimoto, S.: Decadal variations in mixed layer salinity in the Kuroshio Extension recirculation gyre region: Influence of precipitation during the warm season, *J. Oceanogr.*, 72, 167–175. <https://doi.org/10.1007/s10872-015-0317-1>, 2016.

Krumhardt, K. M., Lovenduski, N. S., Long, M. C., Levy, M., Lindsay, K., Moore, J. K., and Nissen, C.: Coccolithophore growth and calcification in an acidified ocean: Insights from community earth system model simulations, *J. Adv. Model. Earth Syst.*, 11, 1418–1437. <https://doi.org/10.1029/2018MS001483>, 2019.

645 Kwiatkowski, L., and Orr, J. C.: Diverging seasonal extremes for ocean acidification during the twenty-first century, *Nature Climate Change*, 8, 141–145. <https://doi.org/10.1038/s41558-017-0054-0>, 2018.

650 ▲ Lee, K., Tong, L. T., Millero, F. J., Sabine, C. L., Dickson, A. G., Goyet, C., Park, G. H., Wanninkhof, R., Feely, R. A., and Key, R. M.: Global relationships of total alkalinity with salinity and temperature in surface waters of the world’s oceans, *Geophys. Res. Lett.*, 33. <https://doi.org/10.1029/2006GL027207>, 2006.

655 Lehmann, N., Lantuit, H., Böttcher, M. E., Hartmann, J., Eulenburg, A., and Thomas, H.: Alkalinity generation from carbonate weathering in a silicate-dominated headwater catchment at Iskorasfjellet, northern Norway, *Biogeosciences*, 20, 3459–3479. <https://doi.org/10.5194/bg-20-3459-2023>, 2023.

Formatted: Default Paragraph Font, Font: 10 pt

Formatted: Font: (Asian) MS 明朝, (Asian) Japanese

Lewes, E., and Wallace, D. W. R.: Program Developed for CO<sub>2</sub> System Calculations, ORNL/CDIAC-105,  
660 Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, U.S. Department of Energy, Oak  
Ridge, TN, USA, <https://doi.org/10.15485/1464255>, 1998.

[Liu, M., Raymond, P. A., Lauerwald, R., Zhang, Q., Trapp-Müller, G., Davis, K. L., Moosdorf, N., Xiao, C.,  
Middelburg, J. J., Bouwman, A. F., Beusen, A. H. W., Peng, C., Lacroix, F., Tian, H., Wang, J., Li, M., Zhu, Q.,  
665 Cohen, S., van Hoek, W. J., Li, Y., Li, Y., Yao, Y., and Regnier, P.: Global riverine land-to-ocean carbon export  
constrained by observations and multi-model assessment, \*Nature Geoscience\*, 17, 896-904.  
<https://doi.org/10.1038/s41561-024-01524-z>, 2024.](#)

Mears, C. A., Scott, J., Wentz, F. J., Ricciardulli, L., Leidner, S. M., Hoffman, R., and Atlas, R.: A near-real-time  
670 version of the Cross-Calibrated Multiplatform (CCMP) ocean surface wind velocity data set, *JGR Oceans*, 124,  
6997–7010. <https://doi.org/10.1029/2019JC015367> [dataset], 2019.

Millero, F. J., Lee, K., and Roche, M.: Distribution of alkalinity in the surface waters of the major oceans, *Mar.  
Chem.*, 60, 111–130. [https://doi.org/10.1016/S0304-4203\(97\)00084-4](https://doi.org/10.1016/S0304-4203(97)00084-4), 1998.

675 Miyama, T., Minobe, S., and Goto, H.: Marine heatwave of sea surface temperature of the Oyashio region in  
summer in 2010–2016, *Front. Mar. Sci.*, 7. <https://doi.org/10.3389/fmars.2020.576240>, 2021.

Miyazawa, Y., Kuwano-Yoshida, A., Doi, T., Nishikawa, H., Narazaki, T., Fukuoka, T., and Sato, K.:  
680 Temperature profiling measurements by sea turtles improve ocean state estimation in the Kuroshio-Oyashio  
Confluence region, *Ocean Dyn.*, 69, 267–282. <https://doi.org/10.1007/s10236-018-1238-5> [dataset], 2019.

Miyazawa, Y., Varlamov, S. M., Miyama, T., Guo, X. Y., Hihara, T., Kiyomatsu, K., Kachi, M., Kurihara, Y.,  
and Murakami, H.: Assimilation of high-resolution sea surface temperature data into an operational  
685 nowcast/forecast system around Japan using a multi-scale three-dimensional variational scheme, *Ocean Dyn.*, 67,  
713–728. <https://doi.org/10.1007/s10236-017-1056-1> [dataset], 2017.

Mucci, A.: The solubility of calcite and aragonite in seawater at various salinities, temperatures, and one  
atmospheric pressure, *Am. J. Sci.*, 283, 780-799, <https://doi.org/10.2475/ajs.283.7.780>, 1983.

Formatted: Default Paragraph Font, Font: 10 pt

Formatted: Font: 11 pt

690

Murata, A. M., Kaneko, I., Nemoto, K., Fushimi, K., and Hirota, M.: Spatial and temporal variations of surface seawater f(CO<sub>2</sub>) in the Kuroshio off Japan, *Mar. Chem.*, 59, 189–200. [https://doi.org/10.1016/S0304-4203\(97\)00096-0](https://doi.org/10.1016/S0304-4203(97)00096-0), 1998.

695

Olsen, A., Key, R. M., van Heuven, S., Lauvset, S. K., Velo, A., Lin, X. H., Schirnack, C., Kozyr, A., Tanhua, T., Hoppema, M., Jutterström, S., Steinfeldt, R., Jeansson, E., Ishii, M., Pérez, F. F., and Suzuki, T.: The Global Ocean Data Analysis Project version 2 (GLODAPv2) - An internally consistent data product for the world ocean, *Earth Syst. Sci. Data*, 8, 297–323. <https://doi.org/10.5194/essd-8-297-2016><https://doi.org/10.5194/essd-8-297-2016> [dataset], 2016.

700

Olsen, A., Lange, N., Key, R. M., Tanhua, T., Bittig, H. C., Kozyr, A., Álvarez, M., Azetsu-Scott, K., Becker, S., Brown, P. J., Carter, B. R., Cotrim da Cunha, L., Feely, R. A., van Heuven, S., Hoppema, M., Ishii, M., Jeansson, E., Jutterström, S., Landa, C. S., Lauvset, S. K., Michaelis, P., Murata, A., Pérez, F. F., Pfeil, B., Schirnack, C., Steinfeldt, R., Suzuki, T., Tilbrook, B., Velo, A., Wanninkhof, R., and Woosley, R. J.: An updated version of the global interior ocean biogeochemical data product, GLODAPv2.2020., *Earth Syst. Sci. Data*, 12, 3653–3678. <https://doi.org/10.5194/essd-12-3653-2020>, 2020.

705

Passow, U. and Carlson, C. A.: The biological pump in a high CO<sub>2</sub> world, *Mar. Ecol. Prog. Ser.*, 470, 249–271. <https://doi.org/10.3354/meps09985>, 2012.

710

Pfeil, B., Olsen, A., Bakker, D. C. E., Hankin, S., Koyuk, H., Kozyr, A., Malczyk, J., Manke, A., Metzl, N., Sabine, C. L., Akl, J., Alin, S. R., Bates, N., Bellerby, R. G. J., Borges, A., Boutin, J., Brown, P. J., Cai, W. J., Chavez, F. P., Chen, A., Cosca, C., Fassbender, A. J., Feely, R. A., González-Dávila, M., Goyet, C., Hales, B., Hardman-Mountford, N., Heinze, C., Hood, M., Hoppema, M., Hunt, C. W., Hydes, D., Ishii, M., Johannessen, T., Jones, S. D., Key, R. M., Körtzinger, A., Landschützer, P., Lauvset, S. K., Lefèvre, N., Lenton, A., Lourantou, A., Merlivat, L., Midorikawa, T., Mintrop, L., Miyazaki, C., Murata, A., Nakadate, A., Nakano, Y., Nakaoka, S., Nojiri, Y., Omar, A. M., Padin, X. A., Park, G. H., Paterson, K., Perez, F. F., Pierrot, D., Poisson, A., Rios, A. F., Santana-Casiano, J. M., Salisbury, J., Sarma, V. V. S. S., Schlitzer, R., Schneider, B., Schuster, U., Sieger, R., Skjelvan, I., Steinhoff, T., Suzuki, T., Takahashi, T., Tedesco, K., Telszewski, M., Thomas, H., Tilbrook, B., Tjiputra, J., Vandemark, D., Veness, T., Wanninkhof, R., Watson, A. J., Weiss, R., Wong, C. S., and Yoshikawa-

720

Formatted: Default Paragraph Font, Font: 10 pt



Inoue, H.: A uniform, quality controlled Surface Ocean CO<sub>2</sub> Atlas (SOCAT), Earth Syst. Sci. Data, 5, 125–143. <https://doi.org/10.5194/essd-5-125-2013>, 2013.

Rassmann, J., Lansard, B., Pozzato, L., and Rabouille, C.: Carbonate chemistry in sediment porewaters of the Rhône River delta driven by early diagenesis (~~northwestern~~Northwest Mediterranean), Biogeosciences, 13, 5379–5394. <https://doi.org/10.5194/bg-13-5379-2016>, 2016.

Regnier, P., Friedlingstein, P., Ciais, P., Mackenzie, F. T., Gruber, N., Janssens, I. A., Laruelle, G. G., Lauerwald, R., Luyssaert, S., Andersson, A. J., Arndt, S., Arnosti, C., Borges, A. V., Dale, A. W., Gallego-Sala, A., Goddérès, Y., Goossens, N., Hartmann, J., Heinze, C., Ilyina, T., Joos, F., LaRowe, D. E., Leifeld, J., Meysman, F. J. R., Munhoven, G., Raymond, P. A., Spahni, R., Suntharalingam, P., and Thullner, M.: Anthropogenic perturbation of the carbon fluxes from land to ocean, Nature Geosci., 6, 597–607. <https://doi.org/10.1038/Ngeo1830>, 2013.

Regnier, P., Resplandy, L., Najjar, R. G., and Ciais, P.: The land-to-ocean loops of the global carbon cycle, Nature, 603, 401–410. <https://doi.org/10.1038/s41586-021-04339-9>, 2022.

Robbins, P. E.: Oceanic carbon transport carried by freshwater divergence: Are salinity normalizations useful? J. Geophys. Res., 106, 30939–30946. <https://doi.org/10.1029/2000JC000451>, 2001.

Romero-Mujalli, G., Hartmann, J., and Börker, J.: Temperature and CO<sub>2</sub> dependency of global carbonate weathering fluxes – Implications for future carbonate weathering research, Chem. Geol., <https://doi.org/10.1016/j.chemgeo.2018.08.010>, ~~2018~~2019.

Song, S., Wang, Z. A., Gonneea, M. E., Kroeger, K. D., Chu, S. N., Li, D., and Liang, H.: An important biogeochemical link between organic and inorganic carbon cycling: effects of organic alkalinity on carbonate chemistry in coastal waters influenced by intertidal salt marshes, Geochim. Cosmochim. Acta, 275, 123–139. <https://doi.org/10.1016/j.gca.2020.02.013>, 2020.

Formatted: Font: 11 pt

Formatted: Default Paragraph Font, Font: 10 pt

750 Song, S., Bellerby, R. G. J., Wang, Z. A., Wurgaft, E., and Li, D.: Organic Alkalinity as an Important Constituent  
755 of Total Alkalinity and the Buffering System in River-To-Coast Transition Zones, JGR oceans, 128.  
<https://doi.org/10.1029/2022JC019270>, 2023.

Formatted: Default Paragraph Font, Font: 10 pt

Sugimoto, S., Takahashi, N., and Hanawa, K.: Marked freshening of North Pacific subtropical mode water in  
755 2009 and 2010: Influence of freshwater supply in the 2008 warm season, *Geophys. Res. Lett.*, 40, 3102–3105.  
<https://doi.org/10.1002/grl.50600>, 2013.

Taguchi, F., Fujiwara, T., Yamada, Y., Fujita, K., and Sugiyama, M.: Alkalinity in coastal seas around Japan,  
*Bull. Coast. Oceanogr.*, 47, 71–75, 2009.

760 Takahashi, T., Sutherland, S. C., Sweeney, C., Poisson, A., Metzl, N., Tilbrook, B., Bates, N., Wanninkhof, R.,  
Feely, R. A., Sabine, C., Olafsson, J., and Nojiri, Y.: Global sea-air CO<sub>2</sub> flux based on climatological surface  
ocean pCO<sub>2</sub>, and seasonal biological and temperature effects, *Deep Sea Research Part II: Topical Studies in*  
*Oceanography*, 49, 1601–1622. [https://doi.org/10.1016/S0967-0645\(02\)00003-6](https://doi.org/10.1016/S0967-0645(02)00003-6), 2002.

765 Takahashi, T., Sutherland, S. C., Wanninkhof, R., Sweeney, C., Feely, R. A., Chipman, D. W., Hales, B.,  
Friederich, G., Chavez, F., Sabine, C., Watson, A., Bakker, D. C. E., Schuster, U., Metzl, N., Yoshikawa-Inoue,  
H., Ishii, M., Midorikawa, T., Nojiri, Y., Körtzinger, A., Steinhoff, T., Hoppema, M., Olafsson, J., Arnarson, T.  
S., Tilbrook, B., Johannessen, T., Olsen, A., Bellerby, R., Wong, C. S., Delille, B., Bates, N. R., and de Baar, H.  
770 J. W.: Climatological mean and decadal change in surface ocean pCO<sub>2</sub>, and net sea-air CO<sub>2</sub> flux over the global  
oceans, *Deep Sea Research Part II: Topical Studies in Oceanography*, 56, 554–577.  
<https://doi.org/10.1016/j.dsr2.2008.12.009>, 2009.

Takamura, T. R., Inoue, H. Y., Midorikawa, T., Ishii, M., and Nojiri, Y.: Seasonal and interannual variations in  
775 pCO<sub>2</sub>(sea) and air-sea CO<sub>2</sub> fluxes in midlatitudes of the western and eastern North Pacific during 1999–2006:  
Recent results utilizing voluntary observation ships, *J. Meteorol. Soc. Jpn*, 88, 883–898.  
<https://doi.org/10.2151/jmsj.2010-602>, 2010.

780 Takatani, Y., Enyo, K., Iida, Y., Kojima, A., Nakano, T., Sasano, D., Kosugi, N., Midorikawa, T., Suzuki, T., and  
Ishii, M.: Relationships between total alkalinity in surface water and sea surface dynamic height in the Pacific  
Ocean, *J. Geophys. Res. Oceans*, 119, 2806–2814. <https://doi.org/10.1002/2013JC009739>, 2014.

Tokoro, T., Nakaoka, S., Takao, S., Kuwae, T., Kubo, A., Endo, T., and Nojiri, Y.: Contribution of biological  
effects to carbonate-system variations and the air-water CO<sub>2</sub> flux in urbanized bays in Japan, *JGR Oceans*, 126.  
785 <https://doi.org/10.1029/2020JC016974>, 2021.

Tokoro, T., Nakaoka, S., Takao, S., Saito, S., Sasano, D., Enyo, K., Ishii, M., Kosugi, N., and Nojiri, Y.:  
Statistical analysis of spatiotemporal variations of air-sea CO<sub>2</sub> fluxes in the Kuroshio region, *JGR Oceans*, 128.  
<https://doi.org/10.1029/2023JC019762>, 2023.

790 Tranvik, L. J., Downing, J. A., Cotner, J. B., Loiselle, S. A., Striegl, R. G., Ballatore, T. J., Dillon, P., Finlay, K.,  
Fortino, K., Knoll, L. B., Kortelainen, P. L., Kutser, T., Larsen, S., Laurion, I., Leech, D. M., McCallister, S. L.,  
McKnight, D. M., Melack, J. M., Overholt, E., Porter, J. A., Prairie, Y., Renwick, W. H., Roland, F., Sherman, B.  
S., Schindler, D. W., Sobek, S., Tremblay, A., Vanni, M. J., Verschoor, A. M., von Wachenfeldt, E., and  
795 Weyhenmeyer, G. A.: Lakes and reservoirs as regulators of carbon cycling and climate, *Limnol. Oceanogr.*, 54,  
2298–2314. [https://doi.org/10.4319/lo.2009.54.6\\_part\\_2.2298](https://doi.org/10.4319/lo.2009.54.6_part_2.2298), 2009.

Wallace, R. B., Baumann, H., Grear, J. S., Aller, R. C., and Gobler, C. J.: Coastal Ocean acidification: The other  
eutrophication problem, *Estuarine Coast. Shelf Sci.*, 148, 1–13. <https://doi.org/10.1016/j.ecss.2014.05.027>, 2014.

800 Wanninkhof, R.: Relationship between wind speed and gas exchange over the ocean revisited, *Limnol. Oceanogr.*  
*Methods*, 12, 351–362. <https://doi.org/10.4319/lom.2014.12.351>, 2014.

Weiss, R. F.: Carbon dioxide in water and seawater: The solubility of a non-ideal gas, *Mar. Chem.*, 2(3), 203–  
805 215. [https://doi.org/10.1016/0304-4203\(74\)90015-2](https://doi.org/10.1016/0304-4203(74)90015-2), 1974.

Wold, S., Sjöström, M., and Eriksson, L.: PLS-regression: A basic tool of chemometrics, *Chemom. Intell. Lab.*  
*Syst.*, 58, 109–130. [https://doi.org/10.1016/S0169-7439\(01\)00155-1](https://doi.org/10.1016/S0169-7439(01)00155-1), 2001.

Formatted: Font: (Asian) MS 明朝, (Asian) Japanese

Formatted: Font: 11 pt

810 Xue, L., Cai, W. J., Jiang, L. Q., and Wei, Q.: Why are surface ocean pH and CaCO<sub>3</sub> saturation state often out of  
phase in spatial patterns and seasonal cycles? Glob. Biogeochem. Cycles, 35, <https://doi.org/10.1029/2021GB00694>,  
2021.

Formatted: Subscript

Formatted: Default Paragraph Font, Font: 10 pt

815 Yang, B., Byrne, R. H., and Lindemuth, M.: Contributions of organic alkalinity to total alkalinity in coastal  
waters: a spectrophotometric approach, Mar. Chem., 176, 199-207. <https://doi.org/10.1016/j.marchem.2015.09.008>,  
2015.

Formatted: Default Paragraph Font, Font: 10 pt

▲  
Yoshikawa-Inoue, H., Midorikawa, T., and Takamura, T. R.: Temporal and spatial variations in carbonate system  
and air-sea CO<sub>2</sub> flux in the Kuroshio and Kuroshio Extension, in: Western Pacific air-sea interaction study, edited  
820 by Uematsu, M., Yokouchi, Y., Watanabe, Y. W., Takebe, S., and Yamanaka, Y., Terra Publishing, Tokyo, 2014.  
<https://doi.org/10.5047/w-pass.a02.004>.

Formatted: Font: (Asian) MS 明朝, (Asian) Japanese

Zeebe, R. E. and Wolf-Gladrow, D.: CO<sub>2</sub> in seawater: Equilibrium, kinetics and isotopes, Elsevier Science,  
Amsterdam, 2001.

825