

Photosynthetic electron, carbon and oxygen fluxes within a mosaic of Fe limitation in the California Current Upwelling System

5 Yayla Sezginer¹, Kate Schuler¹, Emily Speciale², Adrian Marchetti², Claire Till³, Ralph Till³, Philippe Tortell^{1,4}

¹Department of Earth, Oceans and Atmospheric Sciences, University of British Columbia, Vancouver, BC, Canada

10 ²Earth Marine and Environmental Sciences, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA

³Department of Chemistry, California State Polytechnic University, Humboldt, Arcata, CA, USA

⁴Department of Botany, University of British Columbia, Vancouver, BC, Canada

Correspondence to: Yayla Sezginer (ysezginer@eoas.ubc.ca)

Abstract. We compare primary productivity estimates based on different photosynthetic ‘currencies’
15 (electrons, O₂ and carbon) measured in the dynamic coastal upwelling waters of the California Current. Fast Repetition Rate Fluorometry and O₂/N₂’ measurements were used to collect high-resolution underway estimates of photosynthetic electron transport rates and net community productivity, respectively, alongside on-station ¹⁴C uptake experiments to measure gross carbon fixation rates. Our survey captured two upwelling filaments at Cape Blanco and Cape Mendocino with distinct
20 biogeochemical signatures and iron availabilities, enabling us to examine photosynthetic processes along a natural iron gradient. Significant differences in photo-physiology, cell sizes, Si:NO₃⁻ draw-down ratios, and molecular markers of Fe-stress indicated that phytoplankton assemblages near Cape Mendocino were Fe-stressed, while those near Cape Blanco were Fe-replete. Upwelling of O₂-poor deep water to the surface complicated O₂-based net community productivity estimates, but we were able
25 to correct for these vertical mixing effects using continuous [N₂O] surface measurements and depth-profiles of $\frac{\partial [O_2]}{\partial [N_2O]}$. Vertical mixing corrections were strongly correlated to sea surface temperature, which serves as an N₂O-independent proxy for upwelling. All three productivity estimates reflected trends in Fe-stress physiology, indicating greater productivity near Cape Blanco compared to Cape Mendocino. For all phytoplankton assemblages, carbon fixation varied as a hyperbolic function of
30 photosynthetic electron transport rates, but the derived parameters of this relationship were variable and significantly correlated with physiological indicators of Fe-stress (σ_{PSII} , F_v/F_m, Si:NO₃⁻ and diatom-specific PSI gene expression), suggesting that iron availability influenced the coupling between photosynthetic electron transport and carbon fixation. Net community productivity showed strong coherence with daily-integrated photosynthetic electron transport rates across the entire cruise track,
35 with no apparent relationship with Fe-stress. This result suggests that fluorescence-based estimates of

gross photochemistry are still a good indicator for bulk primary productivity, even if Fe-limitation influences the stoichiometric relationship between different productivity currencies.

1 Introduction

40 Along the eastern boundaries of ocean basins, coastal upwelling delivers nutrient-rich deep water to the euphotic zone, sustaining high phytoplankton growth rates and primary productivity (Bograd et al., 2023). Despite representing less than 1% of the surface ocean, these productive upwelling ecosystems, support ~20% of global fishery catches (Pauly and Christensen, 1995), and play a disproportionate role in ocean carbon uptake through the ‘biological carbon pump’ (Mathis et al., 2024). Quantifying rates of
45 primary productivity (PP) within eastern boundary currents is thus vital for accurate carbon budgeting and fishery yield predictions (Marshak and Link, 2021), yet this remains challenging due to the highly dynamic nature of these systems.

The California Current system (CCS) is one of the best studied eastern boundary currents, extending
50 from British Columbia, Canada, to Baja California, Mexico. Upwelling in the CCS occurs during spring and summer when northerly winds drive Ekman transport of surface water offshore. Within the upwelling season, short-term changes in windspeed and direction can dampen or reverse upwelling signals on the scale of hours to days, while complex coastline geometry directs wind flow, creating upwelling hotspots in the lee of capes (Castelao and Luo, 2018). Underlying bathymetric features and
55 deep-water composition further influence the nature of upwelling filaments and the availability of macro and micro nutrients. In regions with shallow and wide continental shelves, sediment deposition provides a primary source Fe and other micronutrients (Deutsch et al., 2021). In contrast, waters overlying steep narrow shelves retain less Fe, and PP in these regions can be limited by Fe availability despite the presence of upwelling conditions (Biller et al., 2013). The resulting ‘mosaic of Fe limitation’
60 influences the distribution of phytoplankton biomass and productivity across the CCS (Till et al., 2019; Hutchins et al., 1998). Resolving ecosystem responses to such a heterogenous environment requires high resolution measurements.

Traditionally, primary productivity has been measured using discrete bottle incubations where the net
65 change in dissolved O₂ or particulate organic carbon is measured over time. Shorter incubations approximate gross primary productivity (GPP), whereas longer incubations allow time for respiration to act upon tracers, yielding estimates somewhere between GPP and net PP ($NPP = GPP - \text{respiration}$). Although bottle incubations are still widely used to directly observe carbon fixation rates, the resulting measurements can be ambiguous in terms of GPP vs. NPP, while also providing low sampling
70 resolution and posing potential containment artefacts (Banse, 2002). To avoid these challenges, a number of high through-put PP proxies have been developed based on advances in dissolved gas measurements, bio-optical techniques, and satellite-based ocean color observations (IOCCG, 2022). These diverse PP methodologies target different photosynthetic processes, from subcellular light absorption to ecosystem scale carbon export.

At the smallest spatial and temporal scales, Photosystem II (PSII) electron transport rates (ETR_{PSII}) quantify light absorption and conversion to chemical energy for a variety of metabolic activities including carbon fixation. Measurements of ETR_{PSII} can be obtained from active chlorophyll fluorescence techniques, which exploit the inverse relationship between PSII fluorescence and photochemical yields to enable non-invasive and high-frequency measurement from underway seawater lines (Kranz et al., 2020; Sezginer et al., 2023) or autonomous platforms (e.g. Carvalho *et al.*, 2020). Downstream of ETR_{PSII} , carbon fixation can be directly measured using bottle incubations, or approximated from empirical algorithms relating NPP to remotely sensed Chlorophyll (Chl) concentration, sea surface temperature (SST), and photosynthetically available radiance (PAR) (Behrenfeld et al., 2005; Saba et al., 2011; Behrenfeld and Falkowski, 1997). Finally, net community productivity (NCP), represents the difference between GPP and community-wide respiration and can be equated to carbon export out of the mixed layer. This term can be derived from measurements of biological oxygen saturation, $\Delta O_2/Ar$, using Ar-normalization of O_2 to correct for physical influences on gas disequilibria (e.g. temperature or salinity changes or bubble injection) to isolate the biological signal (Cassar et al., 2009; Craig and Hayward, 1987). Assuming steady-state conditions, O_2 fluxes from the mixed layer represent a balance between net biological production and sea-air exchange, allowing the calculation of NCP from sea-air flux estimates. Steady state assumptions are violated in upwelling systems, such as the CCS, where O_2 fluxes in the mixed layer are also affected by vertical mixing. However, such vertical mixing effects can be corrected using N_2O as a tracer of O_2 -depleted deep water, given the strong stoichiometric relationship between N_2O and O_2 ratios in sub-surface waters (Cassar et al., 2014).

The various measurement techniques described above yield PP estimates in a number of different ‘currencies’, i.e. carbon, oxygen and electrons, each with different integration time scales (seconds to weeks). In theory, combining PP quantification approaches can fill data gaps (e.g. cloud or ice interference with satellite data, or missing years in time series operations), and provide deeper understanding of ocean metabolism. In practice, incorporating alternative measurement approaches requires understanding of conversion rates between the various PP currencies. Predicting these conversion factors is challenging, as they vary in response to environmental conditions and phytoplankton taxonomy and physiology (Hughes et al., 2021; Hughes et al., 2018a; Schuback et al., 2017; Halsey and Jones, 2015). For example, the $ETR_{PSII} : GPP$ ratio often exceeds the theoretical stoichiometry of 4 (Halsey and Jones, 2015), implying that redox potential generated at PSII is used for functions other than carbon fixation, such as nitrogen uptake or cyclic electron transport. Similarly, differences between ETR_{PSII} and NCP represent combined O_2 consumption pathways, including community-wide respiration, chloro- and photorespiration and pseudo cyclic electron transport. Investigating the drivers of decoupling between currencies can thus improve conversion rate estimates, and also provide insights into energy transfer efficiencies between different components of the photosynthetic process.


Here we present simultaneous PP measurements in the CCS collected using high resolution, underway sampling techniques along a cruise track between Newport, OR and San Francisco, CA. Underway

measurements of ETR_{PSII} and NCP conducted during the May-June upwelling season of 2023 were complemented with parallel measurements of ETR_{PSII} and ^{14}C -GPP collected at oceanographic stations. Our results demonstrate fine-scale spatial patterns in GPP and NCP associated with variability in localized upwelling filaments, with particularly notable differences observed across gradients of dissolved iron concentrations resulting from variable coastal bathymetry. These results enable us to examine variability in productivity currency conversion factors across natural Fe availability gradients within the California Current System, with potential application to other complex coastal waters.

2 Methods and Materials

2.1 Sampling Sites

Measurements were collected along the Oregon and Northern California coast during the Phytoplankton UPwelling Cycle (PUPCYCLE II) expedition, onboard the *R/V Sally Ride* from May 27 – June 11, 2023. A key objective of the program was to examine the evolution of phytoplankton blooms in recently upwelled waters, and we specifically targeted two upwelling plumes off Cape Blanco and Cape Mendocino, which were identified by low sea surface temperature ($SST < 12^{\circ}C$; Figure 1). Along the cruise track, temperature, salinity, and chlorophyll fluorescence were monitored by the ship's underway system, supplied by a seawater supply line with a nominal intake depth of approximately 5m. Nitrate concentrations were measured continuously with a Seabird SUNA sensor calibrated against $NaNO_3$ standards immediately prior to the cruise. In addition to standard oceanographic variables, the seawater supply line was also used for continuous underway measurements of phytoplankton photo-physiology and ETR_{PSII} using a Fast Repetition Rate fluorometer (FRRF; Soliense Inc.), and NCP using a custom-built Pressure In-Situ Gas Instrument (PIGI), as described below.



140 **Figure 1. Study site map.** Location of the cruise track (white line) and discrete sampling stations
 (white circles) during the PUPCYCLE expedition. The black bathymetric contour line represents the
 200 m isobath. The study region and surrounding waters are colored by **a.** daily NASA Aqua Modis
 Level 3 satellite retrievals of sea surface temperature (SST) averaged over the cruise period (May 27 –
 145 June 11, 2024) **b.** NASA Aqua Modis Level 3 satellite retrievals of Chlorophyll *a* (Chl) averaged over
 the cruise period.

Daily CTD and rosette casts were conducted one hour before local sunrise, and samples were collected at
 four depths targeting 1%, 10%, 22%, and 46% of surface solar irradiance levels. Photosynthesis-
 Irradiance (PI) curves (measured with Fast Repetition Rate fluorometry– see below), chlorophyll *a* (Chl)
 150 and nutrient concentrations (NO_3^- , Si, PO_4 , Fe) were measured at all four depths. Samples collected from
 1% and 46% irradiance levels were also incubated for ^{14}C -based PI curves (see sect. 2.5), with sub-
 samples collected for High Performance Liquid Chromatography (HPLC)-based analysis of
 phytoplankton pigments.

2.2 Fast Repetition Rate Fluorometry

155 A bench-top Soliense Inc. Fast Repetition Rate Fluorometer (FRRF) was configured for underway
 collection of chlorophyll *a* fluorescence transients following Sezginer *et al.*, (2021). Samples were
 introduced to the measurement cuvette by an integrated peristaltic pump. The pump was used to flush
 the measurement cuvette for 2.5 minutes before isolating a sample for analysis. Following a one-minute

160 dark period to relax short lived non-photochemical quenching, five single-turnover fluorescence
transients were collected from each sample in the dark. Each single-turnover transient included a
sequence of 100 sub-saturating excitation light pulses of 1.5 μs duration, and a 1 μs interval between
pulses. The excitation phase is designed to induce photochemistry and gradually reduce the pool of
primary electron acceptor molecules, Q_a (Kolber et al., 1998). As the Q_a pool is reduced, further
electron exchange between PSII and Q_a is prevented, closing the photochemistry pathway and causing a
165 concurrent increase in PSII fluorescence yields. The excitation phase is followed by a relaxation
sequence consisting of 127 light pulses with an initial 20 μs interval. During the relaxation phase, the
interval between light pulses increases exponentially, enabling Q_a reoxidation between pulses to
gradually reopen the photochemistry pathway, such that fluorescence yields return to their basal levels.
The biophysical model described by Kolber et al. (1998) was fit to the resulting fluorescence transient
170 to derive the maximum quantum yield of PSII (F_v/F_m), functional absorption area of PSII (σ_{PSII}), and
turnover rate of the primary electron acceptor Q_a (τ_{Qa}).

2.3 Electron Transport Rates, ETR_{PSII}

Continuous FRRF sampling was interrupted every 12 samples (~3hrs) to conduct a Photosynthesis-
Irradiance (PI) curve characterizing light-dependent changes in photo-physiology and photochemistry.
175 Across the entire cruise, we collected 91 PI curves. Each PI curve was initiated with a fresh sample, and
consisted of 11 light levels, increasing from 0 – 850 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$. Light was supplied evenly by
the five actinic LEDs within the FRRF (445, 470, 505, 530, and 590 nm) calibrated against a handheld
WALZ ULM-500 PAR meter prior to deployment. At each light level, ETR_{PSII} was calculated following
Suggett and Moore (2010):

$$180 \quad ETR_{PSII} = PAR * \sigma'_{PSII} * \frac{F'_q}{F'_v} * 6.033 * 10^{-3} \quad 1.$$

In this formulation, the rate of photon delivery to the pool of PSII reaction centers (RCII) is derived as
the product of photosynthetically active radiation (PAR; units $\mu\text{mol photons m}^{-2} \text{s}^{-1}$) supplied by the
LED lamps, and the wavelength-specific functional absorption area of RCII (σ'_{PSII} ; units of $\text{\AA}^2 \text{PSII}^{-1}$).
185 The conversion efficiency from light to photochemical energy depends on the fraction of open RCII,
measured as the dimensionless ratio between the fluorescence amplitude measured under actinic light
when photochemistry is active ($F'_q = F'_m - F'$) and that measured under actinic light if all RCII were
oxidized, ($F'_v = F'_m - F'_o$). In practice, measuring the minimum fluorescence when the RCII pool is
completely open (F'_o) is challenging, as actinic light always drives some degree of photochemistry and
190 reduction of RCII. Alternatively, F'_o can be derived as $F_o/(F_v/F_m + F_o/F'_m)$, following Oxborough *et al.*,
(2012).

The prime notation (') refers to FRRF parameters derived under actinic light. The constant $6.033 * 10^{-3}$
converts PAR to units of quanta $\text{m}^{-2} \text{s}^{-1}$, and σ'_{PSII} to units of $\text{m}^2 \text{RCII}^{-1}$, yielding ETR_{PSII} units of quanta

195 $\text{s}^{-1} \text{RCII}^{-1}$. For a complete description of FRRF derived parameters, see Schuback *et al.* (2021) and Tortell *et al.* (2023).

For each light curve measured, ETR_{PSII} was plotted against PAR and fit with the photosynthesis-irradiance model of Platt *et al.* (1980):

200

$$\text{ETR}_{\text{PSII}} = P_s \left(1 - e^{-\frac{\alpha \text{PAR}}{P_s}} \right) e^{-\frac{\beta \text{PAR}}{P_s}} \quad 2.$$

During the initial light-limiting part of the curve, ETR_{PSII} increases linearly with PAR with a slope of α . As PAR increases to saturating levels, ETR_{PSII} stabilizes at maximum levels, P_{max} . The light saturation index, E_k is derived as P_{max}/α . When phytoplankton are affected by photoinhibition, β describes the decrease in ETR_{PSII} at high light levels (i.e. $\gg E_k$). In the absence of photoinhibition, $P_s = P_{\text{max}}$. When photo-inhibition is present ($\beta > 0$), P_s represents the theoretical maximum potential ETR_{PSII} . When $\beta > 0$, P_{max} is derived as:

205

$$P_{\text{max}} = P_s \left(\frac{\alpha}{\alpha + \beta} \right) \left(\frac{\beta}{\alpha + \beta} \right)^{\left(\frac{\beta}{\alpha} \right)} \quad 3.$$

To evaluate ETR_{PSII} over the cruise track, we linearly interpolated derived values of α , β , and P_s to the sampling resolution of continuous PAR measurements from the ship's meteorological tower (Biospherical Inst. QSR-240P). The mean light intensity in the mixed layer ($\text{PAR}_{\text{in situ}}$) was estimated by accounting for light attenuation with depth, quantified by the diffuse attenuation coefficient ($K_d = \ln(\text{PAR}_0/\text{PAR}_{\text{mld}})/(\text{mld} - 0)$), where PAR_0 is PAR measured at the surface and PAR_{mld} is PAR measured at the mixed layer depth (mld) (Domingues and Barbosa, 2023). For each CTD cast ($n = 28$), mld was determined using a density difference criterion of 0.125 kg m^{-3} , and PAR_{mld} was measured with a Biospherical QSP-200 PAR sensor mounted to the CTD rosette. Both K_d and mld were linearly interpolated to the resolution of continuous PAR measurements. Finally, in-situ PAR was estimated following Domingues and Barbosa (2023) as,

215

$$\text{PAR}_{\text{in situ}} = \text{PAR}_0 (1 - e^{-K_d * \text{MLD}}) (K_d * \text{mld})^{-1} \quad 4.$$

220

To compare ETR_{PSII} with NCP (Sect. 2.6), we converted ETR_{PSII} from $\text{e}^{-} \text{RCII}^{-1} \text{ s}^{-1}$ to volumetric units of $\text{mmol O}_2 \text{ m}^{-2} \text{ d}^{-1}$. This conversion requires an estimate of the chlorophyll content of RCII, which is known to vary significantly across phytoplankton in response to taxonomic and environmental influences (Greene *et al.*, 1992; Murphy *et al.*, 2017; Aardema *et al.*, 2024). Following previous authors (Schuback *et al.*, 2015; Kolber and Falkowski, 1993), we assumed a possible range of Chl to RCII ratios of 400 to 700, yielding upper and lower bounds of Chl-normalized ETR_{PSII} .

225

$$GP = ETR_{PSII} * 86400 * (Chl:RCII)^{-1} * Chl * mld * \frac{1}{4} \quad 5.$$

To obtain volumetric units, Chl-normalized ETR was multiplied by mixed layer Chl concentrations (mmol Chl m⁻³ * mld), assuming homogenous [Chl] throughout the mixed layer. Multiplying by 86400 converts from s⁻¹ to d⁻¹. Given that four charge separation events are required per O₂ evolved, ETR was divided by 4 for final gross photochemistry (GP) estimates in terms of mmol O₂ m⁻² d⁻¹.

2.4 Non-Photochemical Quenching, NPQ

Under excess irradiance, light supply to PSII outpaces maximum downstream electron transport rates, creating the potential for dangerous reactive oxygen species (ROS) to accumulate (Müller et al., 2001). To mitigate excess excitation, photoautotrophs have evolved a number of photoprotective mechanisms, including non-photochemical quenching (NPQ), which dissipates excitation absorbed by PSII as heat, thereby reducing PSII photochemical and fluorescence yields. Previously, NPQ has been quantified from FRRF data as Stern-Volmer quenching, defined as the relative decrease in PSII fluorescence in response to light exposure: $NPQ_{SV} = (F_m - F'_m)/F'_m$. However, this formulation does not account for longer-lived NPQ mechanisms that may still be active during dark measurements following recent high light exposure. To overcome this limitation, we used the normalized Stern-Volmer parameter (NPQ_{NSV}), calculated as F'_o/F'_v (McKew et al., 2013). For each PI curve measured, NPQ_{NSV} was plotted against PAR and fit with a single component exponential curve. Out of 91 curve fits, 95% had $R^2 > 0.90$ and 87% had $R^2 > 0.95$. In-situ NPQ_{NSV} was then estimated by mapping the resulting curve fit onto in-situ PAR values.

Similarly, for each PI curve, the fraction of closed RCII, F'_q/F'_v , was plotted as a function of PAR and fit with a single component exponential curve, with 92% of curves having an $R^2 > 0.90$. Following the same procedure used to evaluate in-situ NPQ_{NSV} , in-situ F'_q/F'_v was approximated by mapping the resulting curve fit onto in-situ PAR values.

2.5 ¹⁴C-uptake experiments

During daily station sampling, 200 mL were collected from 1% and 46% light level depths (approximately 50 and 10 meters, respectively) into acid washed 250 mL bottles for ¹⁴C incubations. Samples were immediately spiked with 150 μCi of H¹⁴CO₃ (Perkin Elmer), and inverted to homogenize the contents of the bottles. The homogenized media was then aliquoted into 20 mL borosilicate scintillation vials, which were incubated over 3 hours in a custom-built photosynthetron at 7 light levels from 0 – 650 μmol photons m⁻² s⁻¹. At the end of the incubation, the entire content of the vials was filtered onto 25mm GF/F filters with a nominal pore size of 0.7 μm. Filters were fumed with 10% HCl for 24 hours to remove any inorganic carbon prior measuring activity on filters with an on-board scintillation counter (Beckman LS 6500). Immediately after spiking samples, three vials were filtered

for triplicate time zero measurements. Three 100 μL aliquots were also taken from the initial 200 mL sample and treated with 100 μL of 3M NaOH to measure total ^{14}C counts. Disintegrations per minute were converted into hourly C fixation rates according to Knap *et al.*, (1996).

2.6 Net Community Productivity (NCP)

265 We measured NCP based on mixed layer concentrations of O_2 and N_2 obtained from the Pressure of In-Situ Gas Instrument (PIGI), following Izett and Tortell (2021) and Izett *et al.* (2021). This method estimates NCP from the biological oxygen saturation anomaly, $\Delta\text{O}_2/\text{N}_2'$, using N_2' as an analog for Argon (Ar) to correct for physical effects on O_2 saturation. In this method, net community productivity is equated to the sea-air flux of O_2 as determined by the biological saturation anomaly ($\Delta\text{O}_2/\text{N}_2'$)
270 scaled by the $[\text{O}_2]$ in equilibrium with the atmosphere ($[\text{O}_2]_{\text{sat}}$), and the O_2 piston velocity (k_{O_2}).

$$\text{NCP} = k_{\text{O}_2} * \Delta \frac{\text{O}_2}{\text{N}_2'} * [\text{O}_2]_{\text{sat}} \quad 6.$$

The PIGI enables cost effective measurements of $\Delta\text{O}_2/\text{N}_2'$ using an oxygen optode and a gas tension device rather than a mass spectrometer, which is more commonly used to measure $\Delta\text{O}_2/\text{Ar}$ (Izett and
275 Tortell, 2021). In this method, N_2' is derived as an approximation of Ar, using model calculations that quantify differences between Ar and N_2 concentrations due primarily to solubility changes and bubble processes. A full description of the 1D model applied to estimate N_2' is available in Izett and Tortell (2021). The model uses ancillary data, including windspeed, mixed layer depth, temperature, salinity, and sea level pressure, to estimate changes in mixed layer Ar and N_2 concentrations over one residence
280 time period prior to sampling. We applied a residence time of 14 days for this region where mixed layer gas residence times are strongly influenced by the timescales of upwelling events (Austin and Barth, 2002). Ancillary datasets required for N_2' calculations were obtained from a combination of satellite observations and model products, and are compiled in the Supplement S1, alongside descriptions of each data source. The 1D model calculations and code are available at
285 https://github.com/rizett/O2N2_NCP_toolbox with example calculations.

Additional corrections to NCP estimates were made to account for vertical mixing fluxes, which transport low O_2 water to the surface (Izett *et al.*, 2018). Previous studies have omitted $\Delta\text{O}_2/\text{Ar}$ data collected in known upwelling areas, where the assumption of limited vertical mixing fluxes on $\Delta\text{O}_2/\text{Ar}$
290 variability is violated (e.g. Stanley *et al.*, 2010). To address this limitation, Cassar *et al.* (2014) developed an approach to use surface measurements of N_2O to quantify vertical transport of low O_2 waters. In marine environments, there is a strong stoichiometric relationship between apparent oxygen utilization and N_2O , which is produced as a by-product of subsurface oxygen-consuming N remineralization pathways (Elkins *et al.*, 1978). These N_2O production pathways are thought to be
295 photo-inhibited within the euphotic zone (Olson, 1981; Horrigan *et al.*, 1981), so that excess N_2O concentrations in the mixed layer serve as a tracer for vertical influxes of O_2 -depleted subsurface water.

We thus used the approach of Cassar et al. (2014) and Izett *et al.* (2018), to correct for vertical mixing following Eq. 7.

$$\text{NCP} = k_{O_2} * \left(\frac{\Delta O_2}{N_2'} * [O_2]_{\text{sat}} - \frac{k_{N_2O}}{k_{O_2}} * \frac{\partial [O_2]^B}{\partial [N_2O]^B} * [N_2O]^B \right) \quad 7.$$

300

This mixing correction uses surface measurements of the N₂O biological concentration ([N₂O]^B), the ‘supply ratio’ of oxygen saturation, given by the vertical gradient of biological O₂ to N₂O ($\frac{\partial [O_2]^B}{\partial [N_2O]^B}$), and the ratio of N₂O and O₂ gas transfer velocities ($\frac{k_{N_2O}}{k_{O_2}}$). Biological concentrations, indicated by the superscript ‘B’, are derived by isolating and removing physical solubility effects from measured gas concentrations. The surface water biological concentration of N₂O, [N₂O]^B, was derived based on the difference between the N₂O saturation anomaly ([N₂O]_{sat}) and changes in N₂O solubility due to recent heat fluxes ([N₂O]_{meas} - [N₂O]_{sat} - [N₂O]_{thermal}). Heat flux effects on solubility, [N₂O]_{thermal}, were derived following Keeling and Shertz (1992), with corrections from Jin et al. (2017). Surface [N₂O]_{meas} was continuously measured from the surface seawater supply with an integrated cavity output spectroscopy (OA-ICOS) gas analyzer (Los Gatos Research, N₂O/CH₄ Analyzer, Model Number: 913–0055) coupled to a gas extraction module (Schuler and Tortell, 2023). The region-specific supply ratio, $\frac{\partial [O_2]^B}{\partial [N_2O]^B}$, was calculated by taking the slope of subsurface [O₂]^B plotted against subsurface [N₂O]^B. The compiled data across the cruise track resulted in a supply ratio of $-1.6 * 10^4 \pm 0.3 * 10^4 \text{ mmol O}_2 (\text{mmol N}_2\text{O})^{-1}$, which is similar to previous measurements for the Northeast Pacific ($-1.8 * 10^4$; Izett et al., 2018) and global basins ($-1.5 * 10^4$; (Cassar et al., 2014). Following Cassar et al. (2014), we assumed a constant $\frac{k_{N_2O}}{k_{O_2}}$ ratio of 0.92.

305

310

315

320

325

We note that several recent studies have observed nitrification within the euphotic zone, challenging the assumption that N₂O production is limited to subsurface waters (Grundle et al., 2013; Smith et al., 2014), and potentially leading to overestimates in our vertical mixing-corrected NCP estimates. Previous observations in the CCS reported a range of depth-integrated mixed layer nitrification rates between 0.3 – 2 mmol NH₄⁺ m⁻² d⁻¹, resulting in consumption of 0.6 – 4 mmol O₂ m⁻² d⁻¹ (Stephens et al., 2020). Following the approach of Izett et al. (2018) we used a range of realistic N₂O:O₂ stoichiometries to estimate potential upper and lower bounds of mixed layer N₂O production. We determined mixed layer N₂O production likely ranged between 0.09 – 0.23 μmol N₂O m⁻² d⁻¹, which would yield offsets in our final NCP estimates between 1.2 and 3.3 mmol O₂ m⁻² d⁻¹. Total uncertainty due to sources of error in other derived parameters was determined by following Izett (2021).

330 2.7 Nutrient concentrations

Samples collected during daily productivity casts were analysed for dissolved $\text{NO}_3^- + \text{NO}_2^-$, PO_4^{3-} , and silicic acid concentrations. Samples of 30 mL were collected from niskin bottles and filtered through GF/F filters, using acid-washed syringes into 20mL HDPE scintillation vials. Samples were kept frozen before analysis on a OI Analytical Flow Solutions IV auto analyzer by Wetland Biogeochemistry
335 Analytical Services at Louisiana State University. Detection limits were $0.09 \mu\text{mol L}^{-1}$ for nitrate + nitrate, $0.02 \mu\text{mol L}^{-1}$ for phosphate and $0.02 \mu\text{mol L}^{-1}$ for silicic acid. Reference standards for dissolved nutrients in seawater were used to ensure quality control.

Samples for iron analysis were collected with a rosette of Teflon-coated OTE-bottles during a separate
340 cast directly after the daily sampling cast. After recovery, OTE-bottles were taken directly to a trace metal-clean sampling van where they were pressurized with filtered compressed air. Surface samples (~3 m depth) were also collected with a tow-fish system plumbed into the trace metal van, as in Bruland *et al.* (2005). All samples were passed through pre-cleaned 0.2 micrometer Supor membrane Acropak capsule filters into trace metal cleaned bottles (Cutter *et al.*, 2014). Samples were acidified to pH 1.8
345 with optima HCl, and analyzed post-cruise with a flow injection analysis method (Lohan *et al.*, 2006), with modifications as in Biller *et al.* (2013). Briefly, this method involved pre-concentrating the Fe at pH 2 with Toyopearl Chelate-650 resin and eluting into a reaction stream containing the colorimetric agent N,N-dimethyl-p-phenylenediaminedihydrochloride (DPD). The absorbance of the reaction stream was measured with a flow-through spectrophotometer. Calibration was performed with a standard
350 addition curve, and blanks were assessed using acidified MilliQ. Reference samples analyzed to assess accuracy compared well to consensus values: SAFe D1 $0.70 \pm 0.04 \text{ nmol Fe/kg}$, $n=12$ compared with consensus value $0.67 \pm 0.04 \text{ nmol/kg}$, and GEOTRACES GSC $1.51 \pm 0.07 \text{ nmol/kg}$ ($n=11$) compared with consensus value $1.53 \pm 0.11 \text{ nmol/kg}$.

2.8 Transcriptomic analysis

355 Water collected from near-surface niskin bottle casts (46% surface irradiance) during daily productivity measurements was subsampled for RNA extraction. Approximately 2.5L to 4L of seawater were filtered onto $0.8 \mu\text{m}$ Pall Supor filters (142 mm) using a peristaltic pump, and then flash frozen in liquid nitrogen and stored at -80°C . RNA was extracted using the RNAqueous-4PCR kit, following manufacturer instructions with the incorporation of a bead beating step during RNA lysis. All RNA
360 samples were sent to GENEWIZ for library preparation and sequencing with PolyA tail selection. Sequencing was performed on an Illumina HiSeq 4000 with a $2 \times 150 \text{ bp}$ configuration. GENEWIZ provided raw paired-end read sequences for each sample.

Raw reads were trimmed using Trim Galore 0.6.10 (Martin, 2011) and quality control was determined with FastQC (Andrews, 2010). A *de novo* metatranscriptome assembly was conducted using rnaSPAdes
365 3.15.5 (Bushmanova *et al.*, 2019) and CD-HIT-EST (Li and Godzik, 2006). Contigs were annotated using the Marine Functional Eukaryotic Reference Taxa (MarFERReT) database (Groussman *et al.*, 2023), which provides NCBI taxonomic annotations (Federhen, 2012) and Pfam 34.0 functional

annotations (Mistry et al., 2021). Samples were mapped against the MarFERReT DIAMOND sequencing aligner and its compatible BLASTX command (e-value < 1e-06) (Buchfink et al., 2015).
370 Trimmed samples were aligned using Salmon (Patro et al., 2017). The package tximport (Soneson et al., 2016) was used to generate a comprehensive table of read count data for each sample and each contig. Only counts taxonomically mapping to Bacillariophyta (i.e., diatoms) were included. The normalized counts for all genes were then calculated using DESeq2's median of ratios method (Love et al., 2014).
375 Normalized counts of the low iron-inducible periplasmic protein (Fea1) (Allen et al., 2007), which shows high similarities to Iron Starvation-Induced Protein 2A (ISIP2A) (Behnke and LaRoche, 2020), was used as an indicator for iron stress (Marchetti et al., 2017).

2.7 Pigment Concentrations and Taxonomic Compositions

During daily productivity casts, duplicate 1L dark Nalgene bottles were filled with water from niskin bottles collected at 1% and 46% PAR level depths. Under low ambient light, samples were filtered onto
380 47mm GFF filters (Whatman, nominal pore size 0.7 μ m). Filters were immediately flash frozen and stored in an onboard -80°C freezer. Samples were shipped on dry ice to the Pinckney Estuarine Ecology Photopigment Analysis Facility at the University of Southern California. There, photopigment concentrations were determined with high performance liquid chromatography following (Pinckney et al., 2001). Pigment compositions are reported in Appendix B.

385 In addition to pigment sampling, Light microscopy was used to identify and enumerate dominant phytoplankton taxa. For microscopic cell counts, 25-50 mL subsamples preserved in Lugol's solution were concentrated by sedimentation using Utermöhl chambers for >24h (Lund et al., 1958). Cell counts of recognizable dinoflagellate and diatom genera were carried out using an Olympus CKX-31 inverted
390 microscope in at least ten fields of view per sample at 200x and 400x magnification.

3 Results

3.1 Oceanographic conditions

Across the cruise track, sea surface temperature (SST) ranged from 8.5 to 14 °C (Fig. 2), with strong coastal to offshore gradients (Fig. 1 and 2). The lowest SST was observed within near-shore upwelling
395 plumes, which were associated with high salinity (>33). Along the entire cruise transect, salinity was negatively correlated with SST ($\rho = -0.73$, $p < 0.01$), as expected for upwelling regions. Sharp hydrographic fronts were apparent along coastal to offshore transects. Moving offshore, SST rapidly increased, while salinity dropped, changing by as much as 2 degrees and 0.5, respectively, within a span of 5 km. These results indicate the presence localized on-shore upwelling plumes, as compared to more
400 homogenous off-shore waters. Within the upwelling plumes, NO_3^- concentrations were elevated, reaching maximum concentrations of 20.5 μM and displaying a positive relationship with salinity ($\rho = 0.89$, $p < 0.01$) and a negative relationship with SST ($\rho = -0.76$, $p < 0.01$). Off-shore, NO_3^- decreased

to concentrations below the SUNA detection limit ($\sim 0.3 \mu\text{M}$), highlighting the difference in nutrient availability between the oligotrophic offshore waters and productive coastal upwelling environments. Chlorophyll concentrations varied between 0.04 to 5.6 mg m^{-3} and exhibited a statistically significant (though weak) positive relationship with NO_3^- ($\rho = 0.30$, $p < 0.01$).





Figure 2. Surface water (a) SST, (b) salinity, (c) Chl, and (d) NO_3^- along the cruise track. Cape Mendocino is abbreviated as Cape Mendo.

In addition to the coastal-offshore gradient, surface water hydrography also differed between the two distinct upwelling plumes we sampled. These plumes were identified as low SST in the lees of Cape

415 Blanco and Cape Mendocino (Fig 1). Both plumes exhibited an upwelling signature, but the apparent
 intensity of upwelling (as reflected in SST, salinity, NO_3^- , and Chl) was significantly stronger within the
 northern Cape Blanco plume (Fig. 3). Most apparently, SST was several degrees cooler at Cape Blanco
 (median of 9.6 ± 0.4 °C) than at Cape Mendocino (median of 11.5 ± 0.01 °C). Nitrate concentrations
 were highly variable across both plumes, but the mean NO_3^- at Cape Blanco (9.4 ± 0.8 μM) was nearly
 420 twice as high as that of Cape Mendocino (5.2 ± 0.4 μM). Chlorophyll concentrations were elevated at
 both plumes relative to offshore waters, with a median of 1.9 ± 0.01 mg m^{-3} around Cape Blanco and
 1.4 ± 0.01 mg m^{-3} at Cape Mendocino. Both these Chl concentrations are well below that which can be
 supported by the available NO_3^- concentrations (1 μM NO_3^- can typically yield 1 $\mu\text{g chl L}^{-1}$), indicating
 that the phytoplankton blooms were likely in the early phases of development following upwelling.



425

Figure 3. Variability in SST (a), salinity (b), NO_3^- (c), and Chl (d) within the observed water masses offshore, at Cape Blanco, and Cape Mendocino. The line inside each boxplot represents the median, while whiskers display the 75th percentile. Points outside the whiskers represent outliers. Panel (e) displays the spatial distribution of offshore (off), Cape Blanco, and Cape Mendocino (Mendo), data. Samples with SST < 12°C and south of 41.5°N were considered part of the Cape Mendocino plume, while samples with <12°C and north of 41.5°N were considered part of the Cape Blanco plume.

430

Samples falling out of these criteria were designated as offshore samples. Criteria were chosen based on a visual inspection of water hydrographic distributions (Fig 1.

435

Underway surface measurements were accompanied by on-station discrete sampling for NO_3^- , PO_4 , Si, and Fe concentrations. As expected, NO_3^- was highly correlated with PO_4 ($\rho = 0.99$, $p < 0.01$) and SiO_2 ($\rho = 0.92$, $p < 0.01$). The ratio of $\text{NO}_3^-:\text{PO}_4$ was 7.4 ± 2.0 , and less than half the expected Redfield ratio of 16. This result is consistent with observations of low $\text{NO}_3^-:\text{PO}_4$ (~2-3) in the North

440

Pacific attributed to high subsurface denitrification rates (Tyrrell and Law, 1997). In contrast with the strong covariance observed among macronutrients, surface Fe distributions were not correlated with surface NO_3^- concentrations ($\rho = 0.35$, $p = 0.6$). At Cape Blanco, Fe concentrations varied from 1.2 – 2.0 nM, whereas Cape Mendocino concentrations were significantly lower, ranging from 0.21 – 0.65 nM. Offshore Fe concentrations were relatively high, with a surface concentration of 0.29-0.42 nM.

445

These waters exhibited low macronutrient concentrations. In section 4.1 we explore the potential causes for different nutrient signatures in the various water masses we sampled.

3.2 Photo-physiology

Along the cruise track, phytoplankton photophysiological properties displayed spatial variability associated with hydrographic gradients, superimposed on significant diel cycles. In particular, we

450 observed strong diel signatures in the expression of various photo-protective mechanisms. Generally, we observed decreases in the PSII functional absorption area, σ_{PSII} , throughout the day, followed by recovery overnight (Fig. 4a). The maximum photochemical efficiency of PSII (F_v/F_m) similarly decreased during the day and peaked overnight. Measurements of NPQ_{NSV} displayed an inverse diel pattern to those of σ_{PSII} and F_v/F_m , reflecting adjustments to the allocation of absorbed energy between

455 competitive photochemistry and thermal dissipation pathways. However, the magnitude of diel variability in F_v/F_m , σ_{PSII} , and NPQ_{NSV} signals displayed significant variability between subregions, as discussed below. These observations agree with previous diel cycle studies of the region (Schuback and Tortell, 2019).




Figure 4. Diel patterns in photo-physiological properties along the cruise track. Grey shading indicates night-time, when surface PAR < 5 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$. **a.** F_v/F_M (black) and σ_{PSII} (red), **b.** P_{max} (black) and α (red), **c.** E_k (black) and NPQ_{NSV} (red) are displayed with respect to sampling time. High sea states between June 4 - 6 impacted the seawater intake, resulting in the data gap.

Overall, photosynthetic parameters derived from semi-continuous PI curves also exhibited diel patterns that mirrored those of F_v/F_M and σ_{PSII} . The maximum, light-saturated, ETR_{PSII} (P_{max}) and the light utilization efficiency under light limiting conditions (α) both peaked during the midday when *in-situ* irradiance was highest. The light saturation level, E_k , tracked surface light availability, while the photoinhibition parameter (β), data not shown, peaked during midday and decreased overnight (data not shown). As with the continuous underway data, results from these discrete PI curves match the previous diel observations of Schuback and Tortell (2019). We note, however, that there is potential for some convolution of temporal and spatial variability, as the ship spent more time offshore in the night, and

on-shore during the daytime. It is thus possible, that some of the diel cycling partially reflects variable photo-physiological signals between coastal and offshore waters.

Beyond diel signals, we also observed significant gradients in photophysiological parameters in relation to oceanographic conditions. In general, F_V/F_M , P_{max} , E_k and α displayed positive relationships with upwelling indicators, i.e. high salinity and macronutrients, and low sea surface temperature (Table 1), suggesting that vertical transport of nutrient-rich water to the surface supported high photochemical yields. In contrast, upwelling signals were associated with decreased σ_{PSII} and NPQ_{NSV} . However, despite general trends between photo-physiological parameters and upwelling, there were significant differences in photo-physiological properties between the Cape Blanco and Cape Mendocino upwelling plumes. At Cape Blanco, mean values of NPQ_{NSV} and σ_{PSII} were significantly lower than at Cape Mendocino, while F_V/F_M , P_{max} , α , and β were all higher at this site, compared to Cape Mendocino (Table 1). Photophysiological properties at Cape Mendocino were much closer to those observed in offshore non-upwelling waters, with mean F_V/F_M values that were lower than offshore, despite elevated macro-nutrient concentrations. This result, combined with low Fe concentrations at Cape Mendocino, suggests that phytoplankton at Cape Mendocino were Fe-stressed despite the presence of upwelling conditions (see Discussion).

	CAPE BLANCO	CAPE MENDO	OFF- SHORE	SST N= 71604	SAL N= 71604	NO ₃ ⁻ N= 989	FE N=40	SI N= 40	PAR N= 71604
F_V/F_M N= 1438	0.47±0.0 1 ^A	0.39 ± 0.08 ^B	0.41± 0.08 ^C	-0.55*	0.27*	0.39*	0.32	0.52	-0.04
σ_{PSII} , N= 1438	545 ± 5 ^A	655 ± 11 ^B	647 ± 5 ^C	0.58*	-0.38*	-0.45*	-0.55	-0.60	-0.12
P_{MAX} , N= 91	977±83 ^A	600 ± 42 ^B	600 ± 91 ^B	-0.67*	0.40*	0.50*	0.45	0.78*	0.19
α , N= 91	4.8±0.3 ^A	3.3 ± 0.3 ^B	3.6 ± 0.4 ^B	-0.57*	0.19	0.24	0.37	0.45	-0.11
β , N= 91	1.55 ± 0.47	1.00 ± 0.36	0.98 ± 0.43	0.44*	0.19	0.31	0.19	0.56	0.09
E_K , N= 91	210±19 ^A	172 ± 20 ^B	176 ± 15 ^B	-0.38*	0.33*	0.41*	0.62	0.72*	0.37*
NPQ_{NSV} , N= 91	0.75±0.0 5 ^A	1.33 ± 0.2 ^B	1.2 ± 0.2 ^B	0.66*	-0.26	-0.46	-0.36	-0.62	0.58*
F'_q/F'_m N= 91	0.21 ± 0.10	0.31 ± 0.14	0.33 ± 0.13	0.34*	-0.18	-0.28	-0.55	-0.31	0.84*

Table 1. Summary of photophysiological properties at each sampling environment and their relationship to hydrographic parameters. Left side of the table displays the median ± median absolute deviation for each photophysiological parameter according to sampling environment.

Superscripts denote groups with significantly different medians. Right side of the table displays

spearman rank correlation coefficients for each photophysiological parameter against environmental parameters. * indicates $p < 0.05$. Due to differences in sampling frequencies, the number of observations (N) varied across different parameters. For correlation analyses, paired observations were matched in space and time to the lowest resolution measurement. Cape Mendocino is abbreviated as Cape Mendo.

3.3 Primary Productivity

3.3.1 Electron Transport Rates, ETR_{PSII}

In-situ ETR_{PSII} , followed a notable diel cycle due to its first order dependency on irradiance (Eq 1, Fig. 5a). However, the relationship between PAR and ETR_{PSII} was spatially variable across the cruise track, reflecting differences in α_{FRRF} , P_{max} , and E_k between Cape Blanco and Cape Mendocino (Table 1; Fig. 4). The amplitude of diel ETR_{PSII} was greatest near Cape Blanco, despite lower average mixed layer PAR (Fig 5a). At Cape Blanco, maximum ETR averaged $977 \pm 83 \text{ e}^- \text{ RCII}^{-1} \text{ s}^{-1}$, as compared to maximum values of only $600 \pm 42 \text{ e}^- \text{ RCII}^{-1} \text{ s}^{-1}$ at Cape Mendocino. The higher maximum ETR at Cape Blanco is in good agreement with the observations of higher photochemical yields in this region (Table 1).




Figure 5. Primary productivity time series along the cruise track. **a)** ETR_{PSII} derived from FRRF-measured PI curve parameter interpolated to match NCP sampling frequency (black line). The grey patches indicate the mean PAR available within the mixed layer ($PAR_{in situ}$). The bar on the bottom is coloured by the sampling latitude. The Cape Blanco filament is designated as latitude > 40.5 and is coloured red, while Cape Mendocino is designated as latitude < 40.5 coloured blue. **b)** Uncorrected NCP (grey) and mixing-corrected NCP (black). The dashed red line denotes the boundary between net autotrophy (NCP > 0) and net heterotrophy (NCP < 0). The color bar on the bottom illustrates the sample temperature. **c)** mixing-corrected NCP (black, right y-axis) and gross photochemistry (GP), shown in red (left y-axis). GP is calculated by converting ETR_{PSII} from units $e^- RCII^{-1} s^{-1}$ to $mmol O_2 m^{-2} d^{-1}$. The grey shading around the red line displays the range of GP based on an assumed range of 400 – 700 Chl:RCII.

3.3.2 Net Community Productivity, NCP

525 In addition to GPP, we estimated NCP from underway measurements. Prior to correcting for vertical mixing, more than 80% of derived NCP values were negative, suggesting net heterotrophic conditions over most of the cruise track. The most negative uncorrected NCP values were observed near Cape Blanco, despite the high gross photochemistry rates measured in this region (Fig 5a and b).

530 The apparent decoupling between NCP and GPP can be largely explained by vertical mixing of low O₂ waters, which artificially depress O₂-derived NCP estimates (see Sect. 2.4). After applying the N₂O-based mixing correction, we found that the majority of the cruise track (73% of measurements) exhibited net autotrophy, with the highest value recorded within the Cape Blanco filament. For the most part, net heterotrophy only existed at night in warmer off-shelf waters. The mean corrected NCP was 80

535 ± 218 mmol O₂ m⁻² d⁻¹, within range of previous observations of late-spring NCP within the California Current (Kranz et al., 2020). The large standard deviation reflects the large diel and spatial variability observed along the cruise track. The highest NCP estimates we obtained (> 500 mmol O₂ m⁻² d⁻¹) are on the upper end of previous measurements. Values above 100 mmol O₂ m⁻² d⁻¹ have only been observed in the most productive coastal waters (Wang et al., 2020; Niebergall et al., 2023), further emphasizing

540 the high productivity of the CCS.

The N₂O-derived mixing correction term was strongly correlated to N₂O-independent indicators of upwelling, namely temperature ($\rho = -0.77$, $p < 0.01$). Uncertainty in vertical-mixing corrected NCP due potential mixed layer nitrification (see sect. 2.6) represented between 1.5 – 4.2% of our mean

545 corrected NCP value. These results give confidence that high surface concentrations of N₂O are a valid marker of upwelling and transport of O₂-poor subsurface water into the mixed layer. The maximum correction factor, 1200 mmol O₂ m⁻² d⁻¹, was observed within the cold upwelling filament near Cape Blanco, where uncorrected NCP was below -500 mmol O₂ m⁻² d⁻¹. This result highlights the impact of vertical fluxes on O₂-based NCP estimates in upwelling regions.

550 3.3.3 Carbon fixation rates

At nine discrete sampling stations, ¹⁴C-based PI curves were measured in parallel with ETR_{PSII} at the surface and at the base of the euphotic zone. Volumetric carbon fixation rates varied significantly between stations and depths. Maximum carbon fixation rates (carbon-based P_{max}) ranged from 0.4 to 96

555 $\mu\text{g C L}^{-1} \text{ hr}^{-1}$. Over 85% of the variability in carbon fixation rates could be explained by differences in biomass, which varied from 0.11 to 9 mg Chl L⁻¹. Chlorophyll concentrations in near-surface waters were, on average, four times higher than those at the base of the euphotic zone, implying the bulk of carbon fixation took place in the mixed layer. On average, the bottom depth of the euphotic zone was 14 ± 12 meters deeper than the bottom depth of the mixed layer

560 To compare carbon-based GPP estimates against parallel ETR_{PSII} measurements, carbon fixation rates were normalized to chlorophyll and converted to units of C chl⁻¹ s⁻¹. Chlorophyll normalized carbon

fixation rates were positively correlated with Fe ($\rho = 0.40$, $p < .05$), F_v/F_m ($\rho = 0.56$, $p < .05$), and Si:NO₃⁻ ratios ($\rho = 0.60$, $p < .05$), and negatively correlated with σ_{PSII} ($\rho = -0.68$, $p < .05$). These results suggest that stations with low chlorophyll-normalized carbon fixation rates may have been affected by Fe and Si co-limitation. By comparison, carbon fixation rates were not significantly correlated with NO₃⁻ or PO₄ concentrations or salinity.

Notably, carbon fixation consistently saturated at lower light intensities than ETR_{PSII}. The average E_k for ETR_{PSII} was 5 ± 2.8 times greater than E_k for carbon fixation. As a result, carbon fixation did not scale linearly with ETR_{PSII}, but rather demonstrated a hyperbolic relationship at each station and depth sampled. At sub-saturating light levels, carbon fixation increased linearly with ETR_{PSII}, until approaching an asymptote as ETR_{PSII} continued to increase while carbon-fixation remained stationary (Fig 6). This result indicates that at light levels beyond the saturating index for carbon-fixation, ETR_{PSII} provides reducing power in excess Calvin-Benson Cycle requirements. Previous studies have noted a similar nonlinear relationship between carbon fixation and ETR_{PSII}, consistent with an upregulation of alternative electron pathways under high light levels (Schuback and Tortell, 2019; Suorsa, 2015; Zhu et al., 2017). This non-linear relationship between C fixation and ETR has been cited as a key limitation to the widespread use of FRRF for autonomous high resolution GPP estimates. Although our results demonstrate a clear hyperbolic relationship between these rates, the parameters describing this relationship were variable across stations. In sect. 4.3, we further examine the relationship between carbon fixation and ETR_{PSII}, and contextualize the apparent differences in carbon-electron decoupling with available physiological, and environmental data.




Figure 6. Parallel measurements of fluorescence- and ^{14}C -based photosynthesis-irradiance experiments conducted on-station. The top left displays the sample locations coloured by maximum chl-specific surface carbon fixation rates. The black contour line displays the 200m isobath. The other panels show carbon fixation rates plotted against ETR_{PSII} . Surface samples are represented by open circles while sub-surface samples are depicted as closed circles. Each curve is fit with the function, $C \text{ fixation} = \frac{P_{\text{max-C}} * \text{ETR}_{\text{PSII}}}{K_{\text{sat}} + \text{ETR}_{\text{PSII}}}$, where $P_{\text{max-C}}$ is the maximum carbon fixation rate, and K_{sat} is the saturation constant beyond which changes in carbon fixation with respect to ETR_{PSII} become increasingly non-linear. Carbon fixation data were not collected at Station 1. All curve fits had an $R^2 > 0.9$.

4 Discussion

As expected, our continuous underway measurements revealed strong spatial and temporal variability in biogeochemical properties across the California Upwelling system. In particular, we observed large diel cycles, and coastal-offshore gradients in biogeochemical properties, with two distinct upwelling filaments in the vicinity Cape Mendocino and Cape Blanco. Differences in nutrient availability between sample sites appeared to exert a strong influence on photo-physiology, gross photochemistry, gross primary productivity and net community productivity. In this section, we explore the potential underlying causes of biogeochemical differences across our survey region, with a focus on iron gradients across the two distinct upwelling filaments. We also discuss the direct and indirect influence

of iron availability and other environmental variables on phytoplankton photo-physiology, and energy
605 transfer efficiencies between photosynthetic processes.

4.1 Factors driving the contrasting biogeochemistry of Cape Blanco and Cape Mendocino filaments

The significant differences in surface oceanographic conditions between upwelling filaments may have
been driven by differences in 1) the strength and timing of upwelling at the two Capes, or 2) differences
in the nutrient content of the sub-surface upwelling source waters. We investigated these two
610 possibilities by examining the NOAA coastal upwelling transport index (CUTI) as a proxy for
upwelling strength during and prior to the sampling period, and by evaluating nutrient depth profiles to
examine the upwelling source waters at the two capes. Our analysis suggests that both factors likely
contributed to the apparent differences between Cape Blanco and Cape Mendocino biogeochemistry,
providing evidence that Fe and Si concentrations were particularly affected by bathymetric features that
615 influence Fe supply.

4.1.1 Strength of upwelling and relative age of filaments

To examine differences in the timing and strength of upwelling between the two filaments, we tracked
CUTI for a 10 day interval prior to sampling (Jacox et al., 2018). Throughout the sampling period, the
strength of upwelling at Cape Blanco varied from 1.3 to 2.4 m d⁻¹ (positive values indicate upwelling)
620 and peaked 6 days prior to our arrival, at 3.8 m d⁻¹. During our sampling period at Cape Mendocino,
upwelling varied from -0.2 to 3.3 m d⁻¹. Upwelling conditions persisted in this region, with vertical
transport rates > 2 m d⁻¹ between June 1 and 4, up to one day prior to our arrival. However, after June 4,
the vertical mixing index at Cape Mendocino rapidly reversed to weak downwelling (-0.1 to -0.3 m d⁻¹)
during the last few days of sampling, emphasizing the dynamic nature of this sampling environment.
625 These results support the hypothesis that colder and more nutrient rich water near Cape Blanco was
attributable to stronger and more consistent upwelling in this region in the interval prior to our
sampling. In contrast, Cape Mendocino was transitioning from upwelling to downwelling conditions
during our sampling period.

4.1.2 Nutrient Content of upwelling source waters

630 Depth profiles offer additional insight into the nutrient concentrations of upwelling filament source
waters. Unfortunately, nutrient samples were only collected down to the base of the euphotic zone,
missing the deeper source waters. Nonetheless, measurements at the base of the euphotic zone (40-50m)
enable us to compare subsurface nutrient concentrations. Mean concentrations of [NO₂⁻ + NO₃⁻]
between 40 and 50m were significantly (p < 0.05) higher at Cape Blanco (24.5 ± 4.3 μM) than at Cape
635 Mendocino (16.5 ± 5.7 μM). Similarly, phosphate concentrations between 40-50m were significantly
greater around Cape Blanco (2.6 ± 0.3 μM) compared to Cape Mendocino (2.1 ± 0.4 μM; p = 0.02).
Relative to nitrate and phosphate, even larger differences were observed in Fe and Si concentrations

between the two capes. The 40-50m silicic acid concentration at Cape Blanco ($30.0 \pm 8.6 \mu\text{M}$) was nearly double that observed at Cape Mendocino ($16.3 \pm 8.2 \mu\text{M}$; ($p \ll 0.01$), while, Fe concentrations
640 between 40-50m at Cape Blanco ($6.8 \pm 4.1 \text{ nM}$) were more than three-fold higher than those at Cape Mendocino ($1.8 \pm 2.3 \text{ nM}$; $p = 0.01$). These results support the hypothesis that the two upwelling plumes were seeded by different water masses with distinct nutrient concentrations.

Differences in underlying bathymetric features between Cape Blanco and Mendocino likely contributed
645 to the observed differences in Fe and Si availability. Cape Blanco sits over a broad section of the continental shelf ($> 30\text{km}$ wide) composed of highly erodible sedimentary rocks with mineral rich sand-silt layers originating from the Klamath Mountains (Spigai, 1971). The broad shelf continues south until the triple junction of the North American, Pacific, and Gorda plates which forms the submarine Mendocino escarpment, a narrow ridge extending west from Cape Mendocino along the transform fault
650 (Menard and Dietz, 1952). Importantly, the shelf rapidly narrows to less than 5 km at the latitude of Cape Mendocino (Appendix A). Differences in shelf width have important implications on sub-regional iron availability. Previous work by Biller *et al.* (2013) demonstrated that shelf width correlated with greater Fe bioavailability in the water layer directly overlying the seafloor. This trend was evident in our study as well, with a positive correlation ($\rho = 0.57$) between shelf width and Fe concentrations at the
655 bottom of the euphotic zone for on-shelf stations. Yet, with only nine stations on-shelf stations, this correlation was not statistically significant.

In contrast with NO_3^- and PO_4^{3-} , which are resupplied to the surface by upwelling, remineralized Fe is rapidly removed from the water column, such that Fe supply to the surface can be significantly
660 decoupled from macronutrients (King and Barbeau, 2011; Bruland *et al.*, 2014). As a result, differences in upwelling strength at Cape Blanco and Cape Mendocino likely account for the differences in PO_4^{3-} and NO_3^- between the two sites, while contrasting shelf features can explain the larger differences in Fe availability. These observations fit within the theory that the California Current contains a ‘mosaic’ of Fe limitation, where patches of Fe-poor water may persist even in the presence of upwelling conditions
665 (Till *et al.*, 2019; Hutchins *et al.*, 1998). In the following section, we present several lines of evidence that photo-physiological properties of phytoplankton assemblages were influenced by iron gradients.

4.2 Environmental and taxonomic influences on physiology and productivity

Environmental gradients exert direct effects on phytoplankton physiology and productivity by determining the supply of essential nutrients that support cellular growth and the maintenance of
670 photosynthetic proteins. As a cofactor in many biological redox reactions, Fe plays a particularly important role in the photosynthetic electron transport chain and nutrient uptake pathways. Several lines of evidence suggest that iron stress was a key factor shaping phytoplankton productivity and photo-physiology across our study site. As noted above, there was a significant difference in Fe concentrations between Cape Blanco (high Fe) and Mendocino (low Fe) associated with variability in the shelf width.
675 The difference in Fe-availability was strongly correlated with $\text{Si}:\text{NO}_3^-$ ($\rho = 0.85$, $p \ll 0.0.1$), likely reflecting excess Si uptake by iron-limited diatoms (Sarhou *et al.*, 2005; M. Franck *et al.*, 2000).

Further evidence of Fe-stress at Cape Mendocino was obtained from ancillary transcriptomic analysis, which demonstrated elevated expression of the Fe assimilation gene *Fea1* (Appendix C), which has previously been cited as a marker of Fe-stress (Allen et al., 2007). Together, these observations indicate the onset of Fe stress at sites with reduced Fe-availability.

Beyond direct effects, Fe gradients indirectly influence phytoplankton physiology and productivity by driving taxonomic shifts towards species that are adapted to low Fe conditions. Although diatoms were the most dominant group across the study area, their relative contribution was significantly lower around Cape Mendocino compared to Cape Blanco (Appendix B). At Cape Mendocino, we observed a taxonomic shift towards smaller phytoplankton, including smaller diatoms and dinoflagellates. Smaller cells sizes afford larger surface area to volume ratios, facilitating nutrient uptake at lower concentrations (Sunda and Huntsman, 1997). Moreover, stations near Cape Mendocino had high abundances of *Pseudo-nitzschia*, whereas this genus was not observed near Cape Blanco. *Pseudo-nitzschia* is a well-studied diatom with a number of physiological adaptations to Fe-limitation (Lampe et al., 2018). These shifts in phytoplankton assemblages towards smaller sizes and low Fe specialists suggest bottom-up environmental controls driving taxonomic composition.

Cell size and nutrient status influence the optical properties and photo-physiology of phytoplankton. Large cells are prone to pigment packaging effects, which decrease Chl-specific absorption as intracellular Chl concentrations increase and surface area to volume ratios decrease. This effect causes reduced σ_{PSII} , as was observed near Cape Blanco (Table 1). Nutrient limitation, particularly for Fe, can also lead to accumulation of photo-inactive or damaged RCII, which still absorb light but do not contribute to photochemistry (Roncel et al., 2016). This further drives high σ_{PSII} , which is proportional to the light harvesting complex absorption coefficient normalized by active RCII concentrations (Oxborough et al., 2012; Z. Li et al., 2021), and low F_v/F_m , due to inactive RCII contributing to the F_m but not F_v signal (Schuback et al., 2021). Both of these commonly cited indicators of Fe stress were observed around Cape Mendocino and in some offshore regions (Table 1).


The taxonomic and nutrient-dependent effects on photo-physiology described above directly impact ETR_{PSII} (Eq. 1). Previous studies have noted higher ETR_{PSII} among Fe-limited phytoplankton, presumably due to increased σ_{PSII} (Schuback et al., 2015). However, we observed greater ETR_{PSII} in the relatively Fe-rich waters near Cape Blanco (Table 1), likely due to high F'_q/F'_v (PAR), which represents the proportion of open RCII at a given light level (Suggett et al., 2011). Low NPQ observed in the Cape Blanco region (Fig. 4d) likely enabled F'_q/F'_v to remain high under high light levels. Indeed, F'_q/F'_v measured during underway Photosynthesis-Irradiance curves and interpolated to in-situ irradiances demonstrated that F'_q/F'_v was higher around Cape Blanco compared to Cape Mendocino (Table 1). It is well recognized that iron limitation exacerbates high light stress and NPQ (Schallenberg et al., 2020; Ryan-keogh et al., 2020), and the high NPQ at Cape Mendocino compared to Cape Blanco provides further evidence that Cape Mendocino assemblages were affected by Fe stress. Iron limitation also impacts photosynthetic processes downstream of ETR_{PSII} . In this study, maximum carbon fixation rates

(P_{\max} determined during ^{14}C PI experiments) displayed a strong correlation with Si ($\rho = 0.63, p \ll 0.01$) and the ratio of $\text{Si}:\text{NO}_3^-$ ($\rho = 0.60, p \ll 0.01$) in the water column.

720 Overall, our results suggest that Fe availability gradients between Cape Blanco and Mendocino
influenced local community composition and physiology with consequential effects on C and Si cycles.
Differences in taxonomic composition, photo-physiology, nutrient quotas and productivity all serve as
evidence that the community shifted towards Fe-limitation in proximity to Cape Mendocino. Due to the
differential sensitivity of ETR_{PSII} , carbon fixation, and NCP to Fe-availability, we hypothesized that Fe-
725 limitation would lead to a decoupling between these different PP currencies. We explore this hypothesis
below with direct comparisons of ETR_{PSII} , C-fixation, and NCP.

4.3 Energy transfer efficiencies between photosynthetic processes

Measurements of primary productivity in different ‘currencies’ (electrons, carbon, and oxygen) reflect
the rates of distinct photosynthetic processes. Comparison of these alternative productivity metrics can
730 yield information on energy transfer efficiencies across the photosynthetic pipeline (Fig 7). A minimum
of four charge separation events are required to produce one O_2 and fix one CO_2 , stoichiometrically
linking water splitting and carbon fixation. Yet a number of non-linear electron transport pathways can
divert reducing power from carbon fixation, decoupling ETR_{PSII} from GPP, while oxygen consumption
by respiration and non-linear electron transport pathways can further decouple ETR_{PSII} from NCP (Fig
735 7). The ratio between ETR_{PSII} and GPP thus provides information on the magnitude of non-linear
electron transport, while the ratio between ETR_{PSII} and NCP reflects the sum of non-linear electron
transport and respiration. In practice, interpreting the decoupling between ETR, GPP, and NCP is
complicated by differences in the temporal and spatial scales of the various measurement approaches, as
well as the different assumptions implicit in each method. In the following sections, we directly
740 compare parallel productivity measurements to examine energy transfer efficiencies across
photosynthetic processes, taking care to note important methodological considerations.



745 **Figure 7. Different primary productivity currencies and their decoupling mechanisms.** The three
 productivity currencies of interest are indicated with respect to the times scales they represent.
 Numbered red pathways denote decoupling mechanisms including 1) chloro-respiration 2) cyclic
 transport around PSII 3) cyclic transport around PSI 4) pseudo-cyclic/ Mehler reactions 5) reductant
 consuming nutrient uptake pathways 6) photo-respiration and 7) Community-wide respiration. Figure
 750 produced in Biorender.

4.3.1 Carbon fixation as a function of ETR_{PSII}

Non-linear electron transport pathways (Fig 7) can act to maintain redox homeostasis when ETR_{PSII}
 exceeds downstream energy requirements for growth and metabolism. The energy balance between PSII
 and PSI becomes disrupted under high irradiance, when PSII absorbs energy in excess of PSI electron
 transport rates, and/or under nutrient limitation, when biosynthesis of electron transporters is limited
 755 (Hughes et al., 2018b; Schuback et al., 2015; Roncel et al., 2016). Iron limitation, in particular, exerts
 acute constraints on the synthesis of Photosystem I (PSI) and Cytochrome b₆f (Cyt b₆f), which require
 12 and 5 Fe atoms each (Raven et al., 1999). As a result, Fe-limited phytoplankton have high levels of
 PSII relative to PSI, exacerbating energy imbalances between PII and PSI, and necessitating
 760 upregulation of non-linear electron transport pathways (Behrenfeld and Milligan, 2013). We therefore
 hypothesized that Fe-stress would increase decoupling between C-fixation and ETR_{PSII}.

One of our primary findings is that C-fixation varies as a hyperbolic function of ETR_{PSII} , with curves defined by the maximum carbon fixation rate (P_{max-C}) and the saturation constant (K_{sat}). These parameters were highly variable between samples (Fig. 6). Samples with high P_{max-C} and K_{sat} , (e.g. Station 6) showed more linear relationships between C-fixation and ETR_{PSII} , and a nearly constant electron requirement for carbon fixation ($\phi_e: c/n_{PSII}$, units = $\mu\text{e Chl C}^{-1} \text{RCII}^{-1}$). In contrast, in phytoplankton assemblages with low P_{max-C} and K_{sat} , carbon-fixation quickly saturated with respect to ETR_{PSII} , resulting in an increase in $\phi_e: c/n_{PSII}$ with increasing light levels (e.g. Station 7). Determining sources of P_{max-C} and K_{sat} variability therefore provides significant utility in predicting electron requirements for gross carbon fixation, a key parameter required for fluorescence-based GPP measurements.

Previous studies have documented the importance of different environmental, taxonomic and physiological parameters in driving variability in $\phi_e: c/n_{PSII}$, but efforts to develop empirical algorithms predicting $\phi_e: c/n_{PSII}$ remain ongoing (Lawrenz et al., 2013). Recent progress has been made on this front by Schuback *et al.*, (2015, 2016, 2017) who reported a consistent relationship between NPQ_{NSV} and $\phi_e: c/n_{PSII}$ in the surface waters of the Northeast Pacific and Canadian Arctic, suggesting that carbon fixation can be estimated using FRRF-based NPQ and ETR_{PSII} measurements alone. Subsequent studies have applied the $NPQ_{NSV} \propto \phi_e: c/n_{PSII}$ relationship observed by Schuback et al. (2015) to collect high-resolution fluorescence-based GPP estimates, e.g. Kranz *et al.* (2020). Others, however, have noted that the $NPQ_{NSV} \propto \phi_e: c/n_{PSII}$ relationship does not hold for all taxa (Hughes et al., 2021), light conditions (Schuback et al., 2017), or environments where phytoplankton grow on more reduced N forms like NH_4^+ (Fei et al., 2024).

In our study, NPQ_{NSV} was positively correlated with $\phi_e: c/n_{PSII}$ ($\rho = 0.55, p \ll 0.01$). However, the linear relationship proposed by Schuback et al. (2017; 2019) did not adequately predict $\phi_e: c/n_{PSII}$ for our samples ($R^2 = -0.41$), nor did we find a single line of best fit that could describe all of our data ($R^2 = 0.30$). Rather, we found that $\phi_e: c/n_{PSII}$ scaled directly with NPQ_{NSV} only for samples with low P_{max-C} ($\leq 0.3 \text{ C Chl}^{-1} \text{ s}^{-1}$). In contrast, for samples with high P_{max-C} ($> 0.3 \text{ C Chl}^{-1} \text{ s}^{-1}$), $\phi_e: c/n_{PSII}$ remained relatively constant across increasing light, NPQ_{NSV} , and ETR_{PSII} (Fig 8).




Figure 8. The relationship between $\phi_{e:C}/n_{PSII}$, NPQ_{NSV} and P_{max-C} . The electron requirement for carbon fixation, $\phi_{e:C}/n_{PSII}$, measured as the ratio of ETR_{PSII} to Chl-normalized carbon fixation, is plotted against NPQ_{NSV} . Each data point is colored by the sample's P_{max-C} , shown by the red/blue colorbar. Lines of best fit are drawn through points with $P_{max-C} \leq 0.3$ and > 0.3 mol C mol Chl⁻¹ s⁻¹, with the colorbar indicating P_{max-C} .

By analogy with the Michaelis-Menten enzyme kinetics model, variability in P_{max-C} can be explained as the product of the enzyme concentration and maximum reaction rate (Choi et al., 2017). In our case, P_{max-C} reflects the entire suite of proteins that facilitate the conversion of chemical energy to organic matter. Reduced concentrations of PSI and Cyt b₆f expected under Fe-stress would therefore reduce P_{max-C} . Indeed, P_{max-C} was significantly correlated to physiological markers of Fe-stress, σ_{PSII} ($\rho = -0.68$, $p < 0.01$), F_V/F_M ($\rho = 0.56$, $p = 0.02$), and $Si:NO_3^-$ ($\rho = 0.60$, $p = 0.01$). Additionally, meta transcriptomic analysis of diatom RNA revealed significant positive correlations ($p < 0.05$) between P_{max-C} and the expression level of different PSI subunits (*psaE*, *psaL*, *psaM*), with correlation coefficients of $\rho = 0.71$, 0.70 , and 0.93 , respectively. In contrast, there were no detected correlations between P_{max-C} and Cyt b₆f, however Cyt *b559a*, a subunit of PSII, also demonstrated a strong positive

810 correlation with $P_{\max-C}$ ($\rho = 0.76$, $p < 0.05$). While the precise functional roles of *Cyt b559a* are still not certain, previous studies have demonstrated its potential role in PSII assembly and photoprotective cyclic electron transport around PSII (Chiu and Chu, 2022). Finally, it is worth noting K_{sat} and $P_{\max-C}$ displayed a positive correlation ($\rho = 0.70$, $p < 0.01$).

815 In addition to non-linear electron transport, $\phi_{e:C}/n_{PSII}$ is also directly affected by the number of Chl energetically coupled to RCII ($1/n_{PSII}$). Directly measuring $1/n_{PSII}$ requires specialized O_2 flash yield instrumentation (Suggett *et al.*, 2009), which was unavailable for this study. Moreover, O_2 flash yield measurements may be challenging in natural plankton assemblages containing phytoplankton and bacteria. As an alternative, variability in $1/n_{PSII}$ between samples can be assessed based on

820 in-situ Chl concentrations, normalized to FRRF-derived proxies for [RCII] ($\propto F_0/\sigma_{PSII}$) following the approach of Oxborough *et al.* (2012). With a known instrument calibration factor, K_a , either provided by instrument manufacturers or determined independently by O_2 flash yield measurements, this approximation could be used to estimate the absolute value of $1/n_{PSII}$.

825 It is well established that Chl:RCII ($1/n_{PSII}$) ratios increase under low light, to maximize light absorption (Greenbaum and Mauzerall, 1991). In our measurements, the proxy for $1/n_{PSII}$ varied significantly between sample depths, with higher $1/n_{PSII}$ at the bottom of the euphotic zone compared to surface depths, confirmed by a t-test comparison of population means ($p < 0.01$). Iron limitation is also expected to increase Chl:RCII. Although iron limitation lowers total cellular Chl content, Chl is

830 more likely to be energetically coupled to RCII rather than PSI reaction centers (Greene *et al.*, 1992). Accordingly, $1/n_{PSII}$ displayed a negative correlation with *Feal* expression in surface samples ($\rho = 0.72$, $p < 0.05$, $n = 9$), which we used as a proxy for iron limitation. We thus conclude that Fe-stress likely contributed to variability in $\phi_{e:C}/n_{PSII}$ in addition to influencing non-linear electron transport. The hyperbolic relationship between carbon fixation and electron transport was unaffected by $1/n_{PSII}$,

835 which was assumed to be constant for individual samples throughout the course of photosynthesis-irradiance experiments. However, this assumption may be violated under high light, due to photoinactivation of RCII (Campbell and Serôdio, 2020). A robust understanding of $\phi_{e:C}/n_{PSII}$ variability requires direct [RCII] measurements collected in parallel with ETR_{PSII} and carbon fixation measurements.

840 4.3.2 Comparison of ETR_{PSII} and NCP

As opposite end members of the productivity spectrum (Fig. 7), ETR_{PSII} quantifies gross photochemical energy production, while NCP represents the net accumulation of photosynthetic carbon or oxygen remaining after accounting for all sources of mixed layer respiration. To directly compare ETR_{PSII} and NCP, we converted ETR_{PSII} from units of $e^- \text{ RCII}^{-1} \text{ s}^{-1}$ to $\text{mmol } O_2 \text{ m}^{-2} \text{ d}^{-1}$ (Eq. 5) by assuming each

845 RCII was functionally coupled to 400 – 700 Chl pigments (Schuback *et al.*, 2015; Kolber and Falkowski, 1993) and 4 charge separation events per gross O_2 evolved. The resulting O_2 -based gross photochemistry values varied between 0 – 4000 $\text{mmol } O_2 \text{ m}^{-2} \text{ d}^{-1}$ (Fig. 5c), coincident with the range of previously reported values for the CCS (Kranz *et al.*, 2020). On average, NCP accounted for $17 \pm 8\%$

of gross photochemistry, indicating ~80% of oxygen produced at PSII by water splitting reactions was
 850 consumed within the mixed layer through autotrophic and heterotrophic respiration. In contrast to
 $\phi_{e:C}/n_{PSII}$, there was no significant differences in NCP:ETR_{PSII} between Cape Blanco, Cape
 Mendocino, or offshore, suggesting minimal effects of nutrient limitation on decoupling between
 ETR_{PSII} and NCP. Although NCP is constrained by gross photochemistry, NCP was greater than
 855 ETR_{PSII} over 29% of the cruise track. This apparent contradiction can be explained by differences time-
 scales between instantaneous ETR_{PSII} measurements, and NCP, which is integrated over O₂ residence
 times in the mixed layer (~1–2 weeks). Sustained net autotrophy can lead to accumulation of O₂ in the
 mixed layer, such that measured O₂ fluxes indicate high levels of NCP despite short-term decreases in
 ETR_{PSII} (e.g. overnight).

860 Regardless of the large differences in integration time-scales and metabolic sources of decoupling, NCP
 showed strong coherence with ETR_{PSII} (Fig 9). Direct comparison between continuous underway
 measurements of ETR_{PSII} and NCP yielded a moderate positive correlation ($\rho = 0.43$, $p \ll 0.01$). To
 account for some of the decoupling introduced by the strong diel dependence of ETR_{PSII}, we also
 compared ETR_{PSII} and NCP measurements integrated over 24hr bins. This comparison indicated a much
 865 stronger relationship between ETR_{PSII} and NCP ($\rho = 0.92$, $p \ll 0.01$; Fig 9). For daily integrated time-
 scales, NCP linearly increased as a function of ETR_{PSII} with a ~15% energy transfer efficiency and a
 predicted NCP of -0.55 mmol O₂ m⁻² when ETR_{PSII} is zero. This efficiency estimate is within range of
 previous studies that have compared gross oxygen production and NCP using triple oxygen isotope and
 O₂/Ar methods (Howard et al., 2017; Haskell II et al., 2017), despite the differences in integration time
 870 scales between ETR_{PSII} and the triple oxygen isotope method. Further, a sensitivity analysis found no
 significant changes in the derived energy transfer efficiency between ETR and NCP integrated over 24,
 48, 72, and 96-hour bins. These results support the utility of FRRF to estimate gross oxygen
 productivity as an alternative to triple oxygen isotopes or other discrete methods, such as H₂¹⁸O tracer
 experiments.




Figure 9. Comparison between fluorescence-based gross photochemistry (GP) and NCP. Panel (a) compares continuous underway ETR_{PSII} and NCP measurements ($\rho = 0.43$, $p \ll 0.01$), panel (b) compares daily-integrated measurements of ETR_{PSII} and NCP measurements ($\rho = 0.92$, $p \ll 0.01$). A line of best fit ($y = 0.14 \cdot \text{GP} - 0.55$) is drawn through the positive NCP data points. The error bars represent the total uncertainty of each measurement determined by propagating uncertainty in each input variable. For GP, total uncertainty is represented as the uncertainty in the Chl:RCII. For NCP, uncertainty was quantified from uncertainty in the deep water supply ratio of $\text{O}_2:\text{N}_2\text{O}$, and modelled differences between $[\text{N}_2]$ and $[\text{Ar}]$. Assumption biases (e.g. no horizontal advection of O_2) also represents a potential large source of uncertainty but were not quantified.

Despite the inherent dependency of net oxygen production on gross oxygen production, the strength of the correlation between NCP and ETR_{PSII} and the consistency of $\text{ETR}_{\text{PSII}}:\text{NCP}$ across subregions is surprising given the multitude of methodological and physiological factors that can uncouple these rates (Fig 7). However, the derivations of NCP and GP both have similar dependencies on mixed layer Chl concentration. To obtain FRRF-derived GP estimates in comparable units of $\text{mmol O}_2 \text{ m}^{-2} \text{ d}^{-1}$, we multiplied in-situ ETR_{PSII} by mixed layer Chl concentrations (Eq 5). While mixed layer Chl concentrations are not explicitly included in NCP calculations (Sect 2.6), biomass is expected to be a primary driver of bulk productivity. If Chl-normalized NCP is instead compared against GP expressed in units of $\text{mmol O}_2 \text{ Chl}^{-1} \text{ d}^{-1}$, the correlation between 24h binned and instantaneous NCP and ETR_{PSII} estimates decrease to $\rho = 0.22$ and 0.35 , respectively. We therefore conclude that it remains challenging to derive gross and net carbon fluxes from FRRF measurements alone, but paired ETR_{PSII} and Chl measurements can provide useful constraints for NCP estimates.

4.3.3 Final methodological considerations: spectral corrections

900 As discussed above, decoupling between ETR_{PSII} , C-fixation and NCP is affected by methodological factors, including differences in the time scale of different photosynthetic processes, and the different normalizations for various measured rates (e.g. per volume, Chl or RCII). Additionally, differences in the spectral characteristics of the in-situ light environment and instrument light sources may contribute further to decoupling between ETR_{PSII} , C-fixation, and NCP.

905 Phytoplankton exhibit variable light use efficiencies across the photosynthetically available wavelength spectrum, due to non-uniform pigment compositions and absorption spectra across assemblages. Consequently, photosynthetic measurements are wavelength-dependent (Kyewalyanga et al., 1997). As a result, differences in the spectral distribution of light between the FRRF and photosynthetron incubator (used for ^{14}C uptake measurements) and the in-situ light environment could influence the stoichiometry we observed between ETR_{PSII} , C-fixation, and NCP.

In principal, spectral corrections, can be used to account for variability between instrument light sources and in-situ light environments to improve the inter-comparability of measurements (Schuback *et al.*, 915 2021; Tortell et al., 2023). These corrections require measurements of the spectral distribution of FRRF, photosynthetron incubator, and in-situ light, and a reconstruction of photosynthesis absorption spectra based on photosynthetic pigment concentrations, determined from HPLC analysis. These corrections rely on the assumption that absorption spectra of photosynthetic pigments accurately represent the action spectra of photosynthesis, which not always the case (Kyewalyanga et al., 1997). In our study, 920 we did not collect measurements of the spectral distribution of light in the euphotic zone, which would have likely varied significantly across depths and sampling sites (i.e. on-shore versus off-shore). For this reason, we are not able to spectrally correct in vitro measurements of ETR_{PSII} and ^{14}C -uptake to in-situ spectral fields for more direct comparison with NCP. As a result, spectral differences between instrument light sources and the in-situ light environment could contribute to some of the observed 925 variability between ETR_{PSII} , C-fixation and NCP in this study. As for the comparison of FRRF and ^{14}C data, spectral corrections would affect the absolute magnitude of $\phi_{e:C}/n_{PSII}$, but these corrections would yield a station-specific scalar and would not affect the hyperbolic relationship observed between ETR_{PSII} and C-fixation.

930 The best approach to minimize the influence of spectral variability is to match the spectral properties of instrument light sources to ambient light fields. However, this remains challenging for high-resolution underway applications across varying spectral environments.

5 Conclusion

935 Consistent with previous observations, our results indicate a patchwork of Fe-stress within the coastal upwelling waters of the California Current, with evidence for physiological Fe-stress within an upwelling filament near Cape Mendocino. Differences in iron availability between upwelling filaments

appear to be linked to bathymetric features that influence sediment loading, and variable micronutrient content of sub-surface upwelling source waters. Paired fluorescence- and ^{14}C -based photosynthesis-irradiance measurements indicated strong connectivity between ETR_{PSII} and carbon-fixation in Fe-replete phytoplankton, and greater decoupling in these rates for Fe-limited assemblages, with greater associated variability in $\phi_e: C/n_{\text{PSII}}$. Recently, there has been significant focus on understanding $\phi_e: C/n_{\text{PSII}}$ variability to expand FRRF-based GPP surveys (Hughes et al., 2018b). Our results suggest that nutrient replete phytoplankton are able maintain near constant $\phi_e: C/n_{\text{PSII}}$ under increasing excitation pressures due to their ability to efficiently transfer energy between PSI and PSII. Under these circumstances, NPQ_{NSV} is not a good predictor of $\phi_e: C/n_{\text{PSII}}$. However, where nutrient limitation necessitates enhanced non-linear electron transport pathways to maintain energy balance between PSII and PSI, $\phi_e: C/n_{\text{PSII}}$ does scale with NPQ_{NSV} and excess excitation pressure. With the accumulation of further data across a range of oceanographic conditions, it may be possible to derive more robust empirical relationships between NPQ_{NSV} and $\phi_e: C/n_{\text{PSII}}$, which could be used to derive GPP in C-based units from FRRF measurements. In addition, our results show a general coherence between daily integrated GP, derived from ETR_{PSII} , and NCP measurements, suggesting that ETR_{PSII} may have significant utility as an indicator of bulk primary productivity. We thus conclude that high-resolution, ship-board measurements hold significant potential to explore fine-scale variability in surface water primary productivity in complex coastal waters.

955 **Appendix A: Study region shelf width variability**

The shelf width was determined at each latitude as the distance between the 200m and 0m isobath calculated with the Haversine formula.





Figure A1. Bathymetric map of the study region. The black contour line indicates the 200m isobath.
 960 The horizontal bar graph demonstrates the shelf width at the aligned latitude.

Appendix B: Photopigment and species distribution by sub-region



965 **Figure B1. Pigment and taxonomic composition of study area sub-regions.** a-c) display non-Chl pigment distribution by mass for Cape Blanco (CB), Cape Mendocino (CM), and Offshore (Off). d-f) display the taxonomic distribution of diatom and dinoflagellate groups visible for microscope counts.

Appendix C: Normalized counts of diatom-specific *Fea1* transcripts

970




Figure C1. Diatom-specific expression of Fe assimilation gene *Fea1*. Bar height displays the mean of normalized *Fea1* counts mapping to Bacillariophyta by station. Normalized *Fea1* counts were calculated using DESeq2's median of ratios method in R (Love et al., 2014). Error bars display the standard deviation. Stations 1-4 were in close proximity of Cape Blanco, 6 and 10 were offshore while the remaining stations were in proximity of the Cape Mendocino upwelling filament.

7 Code and data availability

In-situ hydrographic (<https://doi.pangaea.de/10.1594/PANGAEA.977780>), dissolved gas concentration (<https://doi.pangaea.de/10.1594/PANGAEA.977807>), and phytoplankton photophysiology data (<https://doi.pangaea.de/10.1594/PANGAEA.977630>) collected during the PUPCYCLE II expedition are all publicly available on PANGAEA, an Earth Science data repository. Dissolved Fe data have been submitted to <https://www.bco-dmo.org/>. DOI names are expected to be issued shortly. Satellite SST (11 μ daytime) and Chl data used to produce Fig 1 were downloaded from NASA Aqua MODIS platform

985 (<https://oceancolor.gsfc.nasa.gov/13/>). Ancillary data required to derive N_2' were sourced from several
public databases. Wind speed and Ekman transport were taken from NOAA's Windspeed, Stress, Curl,
Divergence, and Ekman Upwelling, Metop-C ASCAT, 0.25 degree, Global, Near Real Time, 2020-
present, 1-Day Composite product. Sea level pressure was downloaded from the US Navy Global
Environmental Model (NAVGEM) 0.5 degree, 2013-present Pressure MSL
990 (<https://coastwatch.pfeg.noaa.gov/erddap/griddap/erdNavgem05DPres.html>). Modeled sea surface
temperature and salinity products were downloaded from
<https://psl.noaa.gov/data/gridded/data.noaa.oisst.v2.highres.html> and
https://podaac.jpl.nasa.gov/dataset/SMAP_JPL_L3_SSS_CAP_8DAY-RUNNINGMEAN_V5,
respectively. Model code used to calculate N_2' and NCP calculations are available at
https://github.com/rizett/O2N2_NCP_toolbox.

995 **8 Author Contributions**

Y.S. conceived the research plan, conducted FRRF and ^{14}C measurements, analysed data and wrote the
manuscript with significant contribution from all co-authors. K.S. operated the PIGI, conducted all N_2O
measurements, and assisted with NCP computations and analysis. E.S. collected macronutrient and
meta-transcriptomic data and assisted with meta-transcriptomic analysis. A.M. facilitated meta-
000 transcriptomic data collection, assisted with meta-transcriptomic analysis, and co-organized the
PUPCYCLE II expedition with C.T. Trace metal clean Fe samples were collected by C.T. and
processed by R.T. Funding was secured by A.M., C.T., and P.T. Primary advisory support and
manuscript editing was done by P.T.

9 Competing Interests

005 The authors declare they have no conflict of interest.

10 Acknowledgements

We would like to thank the crew and fellow scientists aboard the *R/V Sally Ride* who made this work
possible and created a delightful working environment. Special thanks to Sacchinandan Pillai who
provided HPLC data and mobilization assistance. We thank the Bundy lab at the University of
010 Washington for the loan of their trace metal sampling van. We thank D. Patel from the University of
North Carolina for performing microscope counts.

11 Financial Support

Funding for the PUPCYCLE II cruise was provided to AM from a National Science Foundation grant
(OCE1751805). Iron analyses were funded through the Research Corporation for Science

015 Advancement's Cottrell Scholar Award #26844 to C.T. Trace metal sampling was conducted with a shared-use rosette maintained at Skidaway and purchased with NSF Award OCE-2015430.

12 References

- Aardema, H. M., Slagter, H. A., de Angelis, I., Calleja, M. L., Dragoneas, A., Moretti, S., Schuback, N., Heins, L., Walter, D., Weis, U., Haug, G. H., and Schiebel, R.: On the Variability of Phytoplankton Photophysiology Along a Latitudinal Transect in the North Atlantic Surface Ocean, *J. Geophys. Res. Biogeosciences*, 129, e2023JG007962, <https://doi.org/https://doi.org/10.1029/2023JG007962>, 2024.
- 020 Allen, M., del Campo, J., Kropat, J., and Merchant, S.: FEA1, FEA2, and FRE1, encoding two homologous secreted proteins and a candidate ferriredutase, are expressed coordinately with FOX1 and FTR1 in iron-deficient *Chlamydomonas reinhardtii*, *Eukaryot Cell.*, [https://doi.org/10.1128/EC.00205-](https://doi.org/10.1128/EC.00205-07)
- 025 07, 2007.
- Andrews, S.: FastQC: a quality control tool for high throughput sequence data., 2010.
- Austin, J. A. and Barth, J. A.: Variation in the position of the upwelling front on the Oregon shelf, *J. Geophys. Res. Ocean.*, 107, 1–15, <https://doi.org/https://doi.org/10.1029/2001JC000858>, 2002.
- Banse, K.: SHOULD WE CONTINUE TO MEASURE 14C-UPTAKE BY PHYTOPLANKTON FOR
- 030 ANOTHER 50 YEARS?, *Limnol. Oceanogr. Bull.*, 11, 45–46, <https://doi.org/https://doi.org/10.1002/lob.200211345>, 2002.
- Behnke, J. and LaRoche, J.: Iron uptake proteins in algae and the role of Iron Starvation-Induced Proteins (ISIPs), *Eur. J. Phycol.*, 55, 339–360, <https://doi.org/10.1080/09670262.2020.1744039>, 2020.
- Behrenfeld, M. J. and Falkowski, P. G.: Photosynthetic rates derived from satellite-based chlorophyll concentration, *Limnol. Oceanogr.*, 41, 1–20, 1997.
- 035 Behrenfeld, M. J. and Milligan, A. J.: Photophysiological expressions of iron stress in phytoplankton, *Ann. Rev. Mar. Sci.*, 5, 217–246, <https://doi.org/10.1146/annurev-marine-121211-172356>, 2013.
- Behrenfeld, M. J., Boss, E., Siegel, D. A., and Shea, D. M.: Carbon-based ocean productivity and phytoplankton physiology from space, *Global Biogeochem. Cycles*, 19, 1–14, <https://doi.org/10.1029/2004GB002299>, 2005.
- 040 Biller, D. V., Coale, T. H., Till, R. C., Smith, G. J., and Bruland, K. W.: Coastal iron and nitrate distributions during the spring and summer upwelling season in the central California Current upwelling regime, *Cont. Shelf Res.*, 66, 58–72, <https://doi.org/https://doi.org/10.1016/j.csr.2013.07.003>, 2013.
- Bograd, S. J., Jacox, M. G., Hazen, E. L., Lovecchio, E., Montes, I., Pozo Buil, M., Shannon, L. J., Syde-
045 man, W. J., and Rykaczewski, R. R.: Climate Change Impacts on Eastern Boundary Upwelling Systems, *Ann. Rev. Mar. Sci.*, 15, 303–328, <https://doi.org/10.1146/annurev-marine-032122-021945>, 2023.
- Bruland, K. W., Rue, E. L., Smith, G. J., and DiTullio, G. R.: Iron, macronutrients and diatom blooms in the Peru upwelling regime: brown and blue waters of Peru, *Mar. Chem.*, 93, 81–103, <https://doi.org/https://doi.org/10.1016/j.marchem.2004.06.011>, 2005.
- 050 Bruland, K. W., Middag, R., and Lohan, M. C.: 8.2 - Controls of Trace Metals in Seawater, in: *Treatise on Geochemistry (Second Edition)*, edited by: Holland, H. D. and Turekian, K. K., Elsevier, Oxford, 19–51, <https://doi.org/https://doi.org/10.1016/B978-0-08-095975-7.00602-1>, 2014.

- Buchfink, B., Xie, C., and Huson, D. H.: Fast and sensitive protein alignment using DIAMOND, *Nat. Methods*, 12, 59–60, <https://doi.org/10.1038/nmeth.3176>, 2015.
- Bushmanova, E., Antipov, D., Lapidus, A., and Prjibelski, A. D.: RnaSPAdes: A de novo transcriptome assembler and its application to RNA-Seq data, *Gigascience*, 8, 1–13, <https://doi.org/10.1093/gigascience/giz100>, 2019.
- Campbell, D. A. and Serôdio, J.: Photoinhibition of Photosystem II in Phytoplankton: Processes and Patterns, in: *Photosynthesis in Algae: Biochemical and Physiological Mechanisms*, edited by: Larkum, A. W. D., Grossman, A. R., and Raven, J. A., Springer International Publishing, Cham, 329–365, https://doi.org/10.1007/978-3-030-33397-3_13, 2020.
- Carvalho, F., Gorbunov, M. Y., Oliver, M. J., Haskins, C., Aragon, D., Kohut, J. T., and Schofield, O.: FIRE glider: Mapping in situ chlorophyll variable fluorescence with autonomous underwater gliders, *Limnol. Oceanogr. Methods*, 18, 531–545, <https://doi.org/10.1002/lom3.10380>, 2020.
- Cassar, N., Barnett, B. A., Bender, M. L., Kaiser, J., Hamme, R. C., and Tilbrook, B.: Continuous high-frequency dissolved O₂/Ar Measurements by Equilibrator Inlet Mass Spectrometry, *Anal. Chem.*, 81, 1855–1864, <https://doi.org/10.1021/ac802300u>, 2009.
- Cassar, N., Nevison, C. D., and Manizza, M.: Correcting oceanic O₂/Ar-net community production estimates for vertical mixing using N₂O observations, *Geophys. Res. Lett.*, 41, 8961–8970, <https://doi.org/https://doi.org/10.1002/2014GL062040>, 2014.
- Castelao, R. M. and Luo, H.: Upwelling jet separation in the California Current System, *Sci. Rep.*, 8, 1–8, <https://doi.org/10.1038/s41598-018-34401-y>, 2018.
- Chiu, Y.-F. and Chu, H.-A.: New Structural and Mechanistic Insights Into Functional Roles of Cytochrome b559 in Photosystem II, *Front. Plant Sci.*, 13, <https://doi.org/10.3389/fpls.2022.914922>, 2022.
- Choi, B., Rempala, G., and Kim, J.: Beyond the Michaelis-Menten equation: Accurate and efficient estimation of enzyme kinetic parameters., *Sci Rep.*, <https://doi.org/10.1038/s41598-017-17072-z>, 2017.
- Craig, H. and Hayward, T.: Oxygen Supersaturation in the Ocean: Biological Versus Physical Contributions, *Science* (80-.), 235, 199–202, <https://doi.org/10.1126/science.235.4785.199>, 1987.
- Cutter, G., Andersson, P., Codispoti, L., Croot, P., Francois, R., Lohan, M., and Rutgers Van Der Loeff, M.: Sampling and sample-handling protocols for GEOTRACES cruises. GEOTRACES cookbook, 1–238, 2014.
- Deutsch, C., Frenzel, H., McWilliams, J. C., Renault, L., Kessouri, F., Howard, E., Liang, J.-H., Bianchi, D., and Yang, S.: Biogeochemical variability in the California Current System, *Prog. Oceanogr.*, 196, 102565, <https://doi.org/https://doi.org/10.1016/j.pocean.2021.102565>, 2021.
- DJ, S., HL, M., and TM, K.: Comparing electron transport with gas exchange: parameterising exchange rates between alternative photosynthetic currencies for eukaryotic phytoplankton , *Aquat. Microb. Ecol.*, 56, 147–162, 2009.
- Domingues, R. B. and Barbosa, A. B.: Evaluating Underwater Light Availability for Phytoplankton: Mean Light Intensity in the Mixed Layer versus Attenuation Coefficient, *Water*, 15, <https://doi.org/10.3390/w15162966>, 2023.
- Elkins, J. W., Wofsy, S. C., McElroy, M. B., Kolb, C. E., and Kaplan, W. A.: Aquatic sources and sinks for nitrous oxide, *Nature*, 275, 602–606, <https://doi.org/10.1038/275602a0>, 1978.
- Federhen, S.: The NCBI Taxonomy database, *Nucleic Acids Res.*, 40, D136–D143,

- <https://doi.org/10.1093/nar/gkr1178>, 2012.
- Fei, S., Gao, Y., Li, J., and Cao, L.: The linkage between phytoplankton productivity and photosynthetic electron transport in the summer from the Changjiang River to the East China Sea, *Front. Mar. Sci.*, 11, <https://doi.org/10.3389/fmars.2024.1383988>, 2024.
- 100 Greenbaum, N. L. and Mauzerall, D.: Effect of irradiance level on distribution of chlorophylls between PS II and PS I as determined from optical cross-sections, *Biochim. Biophys. Acta - Bioenerg.*, 1057, 195–207, [https://doi.org/https://doi.org/10.1016/S0005-2728\(05\)80102-1](https://doi.org/https://doi.org/10.1016/S0005-2728(05)80102-1), 1991.
- Greene, R. M., Geider, R. J., Kolber, Z., and Falkowski, P. G.: Iron-Induced Changes in Light Harvesting and Photochemical Energy Conversion Processes in Eukaryotic Marine Algae 1, *Plant*
- 105 *Physiol.*, 100, 565–575, <https://doi.org/10.1104/pp.100.2.565>, 1992.
- Groussman, R. D., Blaskowski, S., Coesel, S. N., and Armbrust, E. V: MarFERReT, an open-source, version-controlled reference library of marine microbial eukaryote functional genes, *Sci. Data*, 10, 926, <https://doi.org/10.1038/s41597-023-02842-4>, 2023.
- Grundle, D. S., Juniper, S. K., and Giesbrecht, K. E.: Euphotic zone nitrification in the NE subarctic
- 110 Pacific: Implications for measurements of new production, *Mar. Chem.*, 155, 113–123, <https://doi.org/https://doi.org/10.1016/j.marchem.2013.06.004>, 2013.
- Halsey, K. H. and Jones, B. M.: Phytoplankton Strategies for Photosynthetic Energy Allocation, *Ann. Rev. Mar. Sci.*, 7, 265–297, <https://doi.org/https://doi.org/10.1146/annurev-marine-010814-015813>, 2015.
- 115 Haskell II, W. Z., Prokopenko, M. G., Hammond, D. E., Stanley, R. H. R., and Sandwith, Z. O.: Annual cyclicity in export efficiency in the inner Southern California Bight, *Global Biogeochem. Cycles*, 31, 357–376, <https://doi.org/https://doi.org/10.1002/2016GB005561>, 2017.
- Horrigan, S. G., Carlucci, A. F., and Williams, P. M.: Light inhibition of nitrification in sea-surface films, *J. Mar. Res.*, 39, 1981.
- 120 Howard, E. M., Durkin, C. A., Hennon, G. M. M., Ribalet, F., and Stanley, R. H. R.: Biological production, export efficiency, and phytoplankton communities across 8000 km of the South Atlantic, *Global Biogeochem. Cycles*, 31, 1066–1088, <https://doi.org/https://doi.org/10.1002/2016GB005488>, 2017.
- Hughes, D. J., Varkey, D., Doblin, M. A., Ingleton, T., Mcinnes, A., Ralph, P. J., van Dongen-Vogels, V., and Suggett, D. J.: Impact of nitrogen availability upon the electron requirement for carbon fixation in Australian coastal phytoplankton communities, *Limnol. Oceanogr.*, 63, 1891–1910, <https://doi.org/10.1002/lno.10814>, 2018a.
- 125 Hughes, D. J., Campbell, D. A., Doblin, M. A., Kromkamp, J. C., Lawrenz, E., Moore, C. M., Oxborough, K., Prášil, O., Ralph, P. J., Alvarez, M. F., and Suggett, D. J.: Roadmaps and Detours: Active Chlorophyll- a Assessments of Primary Productivity Across Marine and Freshwater Systems, *Environ. Sci. Technol.*, 52, 12039–12054, <https://doi.org/10.1021/acs.est.8b03488>, 2018b.
- 130 Hughes, D. J., Giannini, F. C., Ciotti, A. M., Doblin, M. A., Ralph, P. J., Varkey, D., Verma, A., and Suggett, D. J.: Taxonomic Variability in the Electron Requirement for Carbon Fixation Across Marine Phytoplankton, *J. Phycol.*, 57, 111–127, <https://doi.org/https://doi.org/10.1111/jpy.13068>, 2021.
- 135 Hutchins, D. A., DiTullio, G. R., Zhang, Y., and Bruland, K. W.: An iron limitation mosaic in the California upwelling regime, *Limnol. Oceanogr.*, 43, 1037–1054, <https://doi.org/https://doi.org/10.4319/lo.1998.43.6.1037>, 1998.

- 140 Izett, R. W.: Improved Estimates of Net Community Production in the Subarctic Pacific and Canadian Arctic Ocean Using Ship-Based Autonomous Measurements and Computational Approaches, 2021.
- Izett, R. W. and Tortell, P. D.: The pressure of in situ gases instrument (Pigi) for autonomous shipboard measurement of dissolved O_2 and N_2 in surface ocean waters, *Oceanography*, 33, 156–162, <https://doi.org/10.5670/oceanog.2020.214>, 2020.
- 145 Izett, R. W. and Tortell, P. D.: $\Delta O/N'$ as a tracer of mixed layer net community production: Theoretical considerations and proof-of-concept, *Limnol. Oceanogr. Methods*, 19, 497–509, <https://doi.org/https://doi.org/10.1002/lom3.10440>, 2021.
- Izett, R. W., Manning, C. C., Hamme, R. C., and Tortell, P. D.: Refined Estimates of Net Community Production in the Subarctic Northeast Pacific Derived From $\Delta O_2/Ar$ Measurements With N_2O -Based
- 150 Corrections for Vertical Mixing, *Global Biogeochem. Cycles*, 32, 326–350, <https://doi.org/https://doi.org/10.1002/2017GB005792>, 2018.
- Izett, R. W., Hamme, R. C., McNeil, C., Manning, C. C. M., Bourbonnais, A., and Tortell, P. D.: $\Delta O_2/N_2'$ as a New Tracer of Marine Net Community Production: Application and Evaluation in the Subarctic Northeast Pacific and Canadian Arctic Ocean, *Front. Mar. Sci.*, 8, 1–19, <https://doi.org/10.3389/fmars.2021.718625>, 2021.
- 155 Jacox, M. G., Edwards, C. A., Hazen, E. L., and Bograd, S. J.: Coastal Upwelling Revisited: Ekman, Bakun, and Improved Upwelling Indices for the U.S. West Coast, *J. Geophys. Res. Ocean.*, 123, 7332–7350, <https://doi.org/https://doi.org/10.1029/2018JC014187>, 2018.
- King, A. L. and Barbeau, K. A.: Dissolved iron and macronutrient distributions in the southern
- 160 California Current System, *J. Geophys. Res. Ocean.*, 116, <https://doi.org/https://doi.org/10.1029/2010JC006324>, 2011.
- Knap, A., Michaels, A., Close, A., Ducklow, H., and Dickson, A.: Protocols for the Joint Global Ocean Flux Study (JGFOS) Core Measurements, JGOFS Reoprt Nr. 19, vi+170 pp, 198, 1996.
- Kolber, Z. and Falkowski, P. G.: Use of active fluorescence to estimate phytoplankton photosynthesis in
- 165 situ, *Limnol. Oceanogr.*, 38, 1646–1665, <https://doi.org/10.4319/lo.1993.38.8.1646>, 1993.
- Kolber, Z. S., Prášil, O., and Falkowski, P. G.: Measurements of variable chlorophyll fluorescence using fast repetition rate techniques: Defining methodology and experimental protocols, *Biochim. Biophys. Acta - Bioenerg.*, 1367, 88–106, [https://doi.org/10.1016/S0005-2728\(98\)00135-2](https://doi.org/10.1016/S0005-2728(98)00135-2), 1998.
- Kranz, S. A., Wang, S., Kelly, T. B., Stukel, M. R., Goericke, R., Landry, M. R., and Cassar, N.:
- 170 Lagrangian Studies of Marine Production: A Multimethod Assessment of Productivity Relationships in the California Current Ecosystem Upwelling Region, *J. Geophys. Res. Ocean.*, 125, e2019JC015984, <https://doi.org/https://doi.org/10.1029/2019JC015984>, 2020.
- Kyewalyanga, M. N., Platt, T., and Sathyendranath, S.: Estimation of the photosynthetic action spectrum: Implication for primary production models, *Mar. Ecol. Prog. Ser.*, 146, 207–223, <https://doi.org/10.3354/meps146207>, 1997.
- 175 Lampe, R. H., Mann, E. L., Cohen, N. R., Till, C. P., Thamtrakoln, K., Brzezinski, M. A., Bruland, K. W., Twining, B. S., and Marchetti, A.: Different iron storage strategies among bloom-forming diatoms, *Proc. Natl. Acad. Sci. U. S. A.*, 115, E12275–E12284, <https://doi.org/10.1073/pnas.1805243115>, 2018.
- Lawrenz, E., Silsbe, G., Capuzzo, E., Ylöstalo, P., Forster, R. M., Simis, S. G. H., Prášil, O.,

- 180 Kromkamp, J. C., Hickman, A. E., Moore, C. M., Forget, M. H., Geider, R. J., and Suggett, D. J.: Predicting the Electron Requirement for Carbon Fixation in Seas and Oceans, *PLoS One*, 8, <https://doi.org/10.1371/journal.pone.0058137>, 2013.
- Li, W. and Godzik, A.: Cd-hit: a fast program for clustering and comparing large sets of protein or nucleotide sequences, *Bioinformatics*, 22, 1658–1659, <https://doi.org/10.1093/bioinformatics/btl158>, 2006.
- 185 Li, Z., Li, W., Zhang, Y., Hu, Y., Sheward, R., Irwin, A. J., and Finkel, Z. V: Dynamic Photophysiological Stress Response of a Model Diatom to Ten Environmental Stresses, *J. Phycol.*, 57, 484–495, <https://doi.org/https://doi.org/10.1111/jpy.13072>, 2021.
- Lohan, M. C., Aguilar-Islas, A. M., and Bruland, K. W.: Direct determination of iron in acidified (pH 1.7) seawater samples by flow injection analysis with catalytic spectrophotometric detection: Application and intercomparison, *Limnol. Oceanogr. Methods*, 4, 164–171, <https://doi.org/https://doi.org/10.4319/lom.2006.4.164>, 2006.
- 190 Love, M. I., Huber, W., and Anders, S.: Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2, *Genome Biol.*, 15, 550, <https://doi.org/10.1186/s13059-014-0550-8>, 2014.
- 195 Lund, J. W. G., Kipling, C., and Le Cren, E. D.: The inverted microscope method of estimating algal numbers and the statistical basis of estimations by counting, *Hydrobiologia*, 11, 143–170, <https://doi.org/10.1007/BF00007865>, 1958.
- M. Franck, V., Brzezinski, M. A., Coale, K. H., and Nelson, D. M.: Iron and silicic acid concentrations regulate Si uptake north and south of the Polar Frontal Zone in the Pacific Sector of the Southern Ocean, *Deep Sea Res. Part II Top. Stud. Oceanogr.*, 47, 3315–3338, [https://doi.org/https://doi.org/10.1016/S0967-0645\(00\)00070-9](https://doi.org/https://doi.org/10.1016/S0967-0645(00)00070-9), 2000.
- 200 Marchetti, A., Moreno, C. M., Cohen, N. R., Oleinikov, I., deLong, K., Twining, B. S., Armbrust, E. V., and Lampe, R. H.: Development of a molecular-based index for assessing iron status in bloom-forming pennate diatoms, *J. Phycol.*, 53, 820–832, <https://doi.org/https://doi.org/10.1111/jpy.12539>, 2017.
- 205 Marshak, A. R. and Link, J. S.: Primary production ultimately limits fisheries economic performance, *Sci. Rep.*, 11, 12154, <https://doi.org/10.1038/s41598-021-91599-0>, 2021.
- Martin, M.: Cutadapt removes adapter sequences from high-throughput sequencing reads, *EMBnet.journal* [Online], 10–12, <https://doi.org/https://doi.org/10.14806/ej.17.1.200>, 2011.
- Mathis, M., Lacroix, F., Hagemann, S., Nielsen, D. M., Ilyina, T., and Schrum, C.: Enhanced CO₂ uptake of the coastal ocean is dominated by biological carbon fixation, *Nat. Clim. Chang.*, 14, 373–379, <https://doi.org/10.1038/s41558-024-01956-w>, 2024.
- 210 McKew, B. A., Davey, P., Finch, S. J., Hopkins, J., Lefebvre, S. C., Metodiev, M. V, Oxborough, K., Raines, C. A., Lawson, T., and Geider, R. J.: The trade-off between the light-harvesting and photoprotective functions of fucoxanthin-chlorophyll proteins dominates light acclimation in *Emiliania huxleyi* (clone CCMP 1516), *New Phytol.*, 200, 74–85, <https://doi.org/https://doi.org/10.1111/nph.12373>, 2013.
- Menard, H. W. and Dietz, R. S.: Mendocino Submarine Escarpment, *J. Geol.*, 60, 266–278, <https://doi.org/10.1086/625962>, 1952.
- 215 Mistry, J., Chuguransky, S., Williams, L., Qureshi, M., Salazar, G. A., Sonnhammer, E. L. L., Tosatto, S. C. E., Paladin, L., Raj, S., Richardson, L. J., Finn, R. D., and Bateman, A.: Pfam: The protein families database in 2021, *Nucleic Acids Res.*, 49, D412–D419, <https://doi.org/10.1093/nar/gkaa913>,
- 220

- 2021.
- Müller, P., Li, X. P., and Niyogi, K. K.: Non-photochemical quenching. A response to excess light energy, *Plant Physiol.*, 125, 1558–1566, <https://doi.org/10.1104/pp.125.4.1558>, 2001.
- 225 Murphy, C. D., Ni, G., Li, G., Barnett, A., Xu, K., Grant-Burt, J., Liefer, J. D., Suggett, D. J., and Campbell, D. A.: Quantitating active photosystem II reaction center content from fluorescence induction transients, *Limnol. Oceanogr. Methods*, 15, 54–69, <https://doi.org/10.1002/lom3.10142>, 2017.
- Niebergall, A. K., Traylor, S., Huang, Y., Feen, M., Meyer, M. G., McNair, H. M., Nicholson, D., Fassbender, A. J., Omand, M. M., Marchetti, A., Menden-Deuer, S., Tang, W., Gong, W., Tortell, P.,
- 230 Hamme, R., and Cassar, N.: Evaluation of new and net community production estimates by multiple ship-based and autonomous observations in the Northeast Pacific Ocean, *Elem. Sci. Anthr.*, 11, 107, <https://doi.org/10.1525/elementa.2021.00107>, 2023.
- Olson, R. J.: Differential photoinhibition of marine nitrifying bacteria: A possible mechanism for the formation of the primary nitrite maximum, *J. Mar. Res.*, 1981.
- 235 Oxborough, K., Moore, C. M., Suggett, D. J., Lawson, T., Chan, H. G., and Geider, R. J.: Direct estimation of functional PSII reaction center concentration and PSII electron flux on a volume basis: A new approach to the analysis of Fast Repetition Rate fluorometry (FRRf) data, *Limnol. Oceanogr. Methods*, 10, 142–154, <https://doi.org/10.4319/lom.2012.10.142>, 2012.
- Patro, R., Duggal, G., Love, M. I., Irizarry, R. A., and Kingsford, C.: Salmon provides fast and bias-aware quantification of transcript expression, *Nat. Methods*, 14, 417–419, <https://doi.org/10.1038/nmeth.4197>, 2017.
- 240 Pauly, D. and Christensen, V.: Primary production required to sustain global fisheries, *Nature*, 374, 255–257, <https://doi.org/10.1038/374255a0>, 1995.
- Pinckney, J. L., Richardson, T. L., Millie, D. F., and Paerl, H. W.: Application of photopigment
- 245 biomarkers for quantifying microalgal community composition and in situ growth rates, *Org. Geochem.*, 32, 585–595, [https://doi.org/https://doi.org/10.1016/S0146-6380\(00\)00196-0](https://doi.org/10.1016/S0146-6380(00)00196-0), 2001.
- Platt, T., Gallegos, C. L., and Harrison, W. G.: Photoinhibition of photosynthesis in natural assemblages of marine phytoplankton, 1980.
- Raven, J. A., Evans, M. C. W., and Korb, R. E.: The role of trace metals in photosynthetic electron
- 250 transport in O₂-evolving organisms, 111–149, 1999.
- Roncel, M., González-Rodríguez, A. A., Naranjo, B., Bernal-Bayard, P., Lindahl, M., Hervás, M., Navarro, J. A., and Ortega, J. M.: Iron deficiency induces a partial inhibition of the photosynthetic electron transport and a high sensitivity to light in the diatom *Phaeodactylum tricornutum*, *Front. Plant Sci.*, 7, 1–14, <https://doi.org/10.3389/fpls.2016.01050>, 2016.
- 255 Ryan-keogh, T. J., Thomalla, S. J., Robinson, C., and Smith, W.: Deriving a Proxy for Iron Limitation From Chlorophyll Fluorescence on Buoyancy Gliders, 7, 1–13, <https://doi.org/10.3389/fmars.2020.00275>, 2020.
- Saba, V. S., Friedrichs, M. A. M., Antoine, D., Armstrong, R. A., Asanuma, I., Behrenfeld, M. J., and Ciotti, A. M.: An evaluation of ocean color model estimates of marine primary productivity in coastal
- 260 and pelagic regions across the globe, 489–503, <https://doi.org/10.5194/bg-8-489-2011>, 2011.
- Sarthou, G., Timmermans, K. R., Blain, S., and Tréguer, P.: Growth physiology and fate of diatoms in the ocean: a review, *J. Sea Res.*, 53, 25–42, [https://doi.org/https://doi.org/10.1016/j.seares.2004.01.007](https://doi.org/10.1016/j.seares.2004.01.007), 2005.

- Schallenberg, C., Strzepek, R. F., Schuback, N., Clementson, L. A., Boyd, P. W., and Trull, T. W.: Diel quenching of Southern Ocean phytoplankton fluorescence is related to iron limitation, 793–812, 2020.
- Schuback, N. and Tortell, P. D.: Diurnal regulation of photosynthetic light absorption, electron transport and carbon fixation in two contrasting oceanic environments, *Biogeosciences Discuss.*, 1–33, <https://doi.org/10.5194/bg-2018-524>, 2019.
- Schuback, N., Schallenberg, C., Duckham, C., and Maldonado, M. T.: Interacting Effects of Light and Iron Availability on the Coupling of Photosynthetic Electron Transport and CO₂-Assimilation in Marine Phytoplankton, *PLoS One*, 1–30, <https://doi.org/10.1371/journal.pone.0133235>, 2015a.
- Schuback, N., Flecken, M., Maldonado, M. T., and Tortell, P. D.: Diurnal variation in the coupling of photosynthetic electron transport and Diurnal variation in the coupling of photosynthetic electron transport and carbon fixation in iron-limited phytoplankton in the NE subarctic Pacific, <https://doi.org/10.5194/bg-13-1019-2016>, 2016.
- Schuback, N., Hoppe, C. J. M., Tremblay, J. É., Maldonado, M. T., and Tortell, P. D.: Primary productivity and the coupling of photosynthetic electron transport and carbon fixation in the Arctic Ocean, *Limnol. Oceanogr.*, 62, 898–921, <https://doi.org/10.1002/lno.10475>, 2017.
- Schuback, N., Tortell, P. D., Berman-Frank, I., Campbell, D. A., Ciotti, A., Courtecuisse, E., Erickson, Z. K., Fujiki, T., Halsey, K., Hickman, A. E., Huot, Y., Gorbunov, M. Y., Hughes, D. J., Kolber, Z. S., Moore, C. M., Oxborough, K., Prášil, O., Robinson, C. M., Ryan-Keogh, T. J., Silsbe, G., Simis, S., Suggett, D. J., Thomalla, S., and Varkey, D. R.: Single-Turnover Variable Chlorophyll Fluorescence as a Tool for Assessing Phytoplankton Photosynthesis and Primary Productivity: Opportunities, Caveats and Recommendations, *Front. Mar. Sci.*, 8, <https://doi.org/10.3389/fmars.2021.690607>, 2021a.
- Schuler, K. H. and Tortell, P. D.: Impacts of vertical mixing and ice-melt on N₂O and CH₄ concentrations in the Canadian Arctic Ocean, *Cont. Shelf Res.*, 269, 105124, <https://doi.org/10.1016/j.csr.2023.105124>, 2023.
- Sezginer, Y., Suggett, D. J., Izett, R. W., and Tortell, P. D.: Irradiance and nutrient-dependent effects on photosynthetic electron transport in Arctic phytoplankton: A comparison of two chlorophyll fluorescence-based approaches to derive primary photochemistry, *PLoS One*, 16, 1–23, <https://doi.org/10.1371/journal.pone.0256410>, 2021.
- Sezginer, Y., Campbell, D., Pillai, S., and Tortell, P.: Fluorescence-based primary productivity estimates are influenced by non-photochemical quenching dynamics in Arctic phytoplankton, *Front. Microbiol.*, 14, <https://doi.org/10.3389/fmicb.2023.1294521>, 2023.
- Smith, J. M., Casciotti, K. L., Chavez, F. P., and Francis, C. A.: Differential contributions of archaeal ammonia oxidizer ecotypes to nitrification in coastal surface waters, *ISME J.*, 8, 1704–1714, <https://doi.org/10.1038/ismej.2014.11>, 2014.
- Soneson, C., Love, M. I., and Robinson, M. D.: Differential analyses for RNA-seq: transcript-level estimates improve gene-level inferences [version 2; peer review: 2 approved], *F1000Research*, 4, <https://doi.org/10.12688/f1000research.7563.2>, 2016.
- Spigai, J. J.: Marine geology of the continental margin off southern Oregon, Oregon State University, 1971.
- Stanley, R. H. R., Kirkpatrick, J. B., Cassar, N., Barnett, B. A., and Bender, M. L.: Net community production and gross primary production rates in the western equatorial Pacific, *Global Biogeochem. Cycles*, 24, <https://doi.org/10.1029/2009GB003651>, 2010.

- Stephens, B. M., Wankel, S. D., Beman, J. M., Rabines, A. J., Allen, A. E., and Aluwihare, L. I.: Euphotic zone nitrification in the California Current Ecosystem, *Limnol. Oceanogr.*, 65, 790–806, <https://doi.org/https://doi.org/10.1002/lno.11348>, 2020.
- 310 Suggett D, Moore M, G. R. J.: Estimating Aquatic Productivity from Active Fluorescence Measurements, 2010.
- Suggett, D. J., Moore, C. M., and Geider, R. J.: Chlorophyll a fluorescence in Aquatic Sciences: Methods and Applications, *Dev. Appl. Phycol.*, 2011.
- Sunda, W. G. and Huntsman, S. A.: Interrelated influence of iron, light and cell size on marine phytoplankton growth, *Nature*, 390, 389–392, <https://doi.org/10.1038/37093>, 1997.
- 315 Suorsa, M.: Cyclic electron flow provides acclimatory plasticity for the photosynthetic machinery under various environmental conditions and developmental stages, *Front. Plant Sci.*, 6, 1–8, <https://doi.org/10.3389/fpls.2015.00800>, 2015.
- Till, C. P., Solomon, J. R., Cohen, N. R., Lampe, R. H., Marchetti, A., Coale, T. H., and Bruland, K. W.: The iron limitation mosaic in the California Current System: Factors governing Fe availability in the shelf/near-shelf region, *Limnol. Oceanogr.*, 64, 109–123, <https://doi.org/https://doi.org/10.1002/lno.11022>, 2019.
- 320 Tortell, P. D., Schuback, N., and Suggett, D. J.: Application of Single Turnover Active Chlorophyll Fluorescence for Phytoplankton Productivity Measurements., <https://doi.org/http://dx.doi.org/10.25607/OBP-1084>, 2023.
- 325 Tyrrell, T. and Law, C. S.: Low nitrate:phosphate ratios in the global ocean, *Nature*, 387, 793–796, <https://doi.org/10.1038/42915>, 1997.
- Wang, S., Kranz, S. A., Kelly, T. B., Song, H., Stukel, M. R., and Cassar, N.: Lagrangian Studies of Net Community Production: The Effect of Diel and Multiday Nonsteady State Factors and Vertical Fluxes on O₂/Ar in a Dynamic Upwelling Region, *J. Geophys. Res. Biogeosciences*, 125, e2019JG005569, <https://doi.org/https://doi.org/10.1029/2019JG005569>, 2020.
- 330 Zhu, Y., Ishizaka, J., Tripathy, S. C., Wang, S., Sukigara, C., Goes, J., Matsuno, T., and Suggett, D. J.: Relationship between light, community composition and the electron requirement for carbon fixation in natural phytoplankton, *Mar. Ecol. Prog. Ser.*, 580, 83–100, <https://doi.org/10.3354/meps12310>, 2017.

335