

Response to CC 1

Community Comment text from Sayedain and O'Neill in blue.

Author (Ashpole and Wiacek) replies in black.

This paper was, in our opinion, inadequately researched and showed a discouraging level of ill-justified conclusions.

Drs. Sayedain and O'Neill took time to present some detailed comments on our work, which we appreciate. However, our goals, approaches and conclusions were not summarized in their Community Comment and there is evidence that they were not fully understood.

Most notably, the authors demonstrated a lack of awareness of the expected magnitude of Arctic DODs (and Arctic AODs in general) at the regional level coupled with an attendant lack of citations to the existing Arctic DOD literature (there is admittedly not a lot of existing literature but it is easily found*).

While we used the phrase 'persistently high dust loadings' once (L214), it should and will be changed to 'more frequent signatures of extreme dust loadings' because we do not claim anywhere in our paper that Arctic Dust Optical Depths (DOD's) are **on average** > 0.5 . This is only our threshold for an **extreme event** in the analysis, as supported by well known *in situ* measurements of DOD exceeding such values on occasion, including in Sayedain and O'Neill's own work, which we are familiar with (more on this below).

The Frequency of Occurrence (**FoO**) values shown in Figs. 1-3 are very low in absolute terms (< 100 events over 20 years) but indeed higher in relative terms (between 0 % and 50 % of observations over 20 years, depending on location). However, we also show the total number of observations as quite low, and the seasonal dependence such that the vast majority of our extreme event data comes from satellite observations made in JJA (least snow and ice in satellite retrievals, peak time of Arctic dust mobilization).

Despite higher-than-expected fractions of extreme events in some Arctic places, our main result is not a map of average or extreme DOD values but rather a map of the **frequency of occurrence** of such **extreme** events. The logic as to why this would be useful in dust emission modeling work is based on "where there's smoke, there's fire", except repeated for dust: "where there are recurring extreme dust events over 20 years, there is potentially an active dust source there".

Our interest in **active dust source areas** is also the reason why we produced a high-resolution (0.1°) MODIS DOD product, and why we are concerned with analyses over land (the source of dust emissions). Lastly, our approach is not new – we apply previously published work on locating dust **sources** in this way in warm deserts to the Canadian Arctic Archipelago (**CAA**).

It is also surprising that their analysis is predominantly top-down with virtually no investigations at the event level.

This is an interesting point – individual studies can proceed by either approach, but eventually both approaches must meet and compare notes. We chose a top-down analysis in our study to explore potential dust **source areas** in a large geographic region (CCA) through a 20-year analysis of infrequent extreme events. We view it as entirely complementary to the 'event-level' approach of Sayedain and O'Neill, and others who have done related work in their group. Instead of carrying out

event-level analyses that attempt to, e.g., validate satellite data or improve retrieval algorithms, we start from a place of accepting the big picture of what the satellite data are showing us, while understanding that there may be biases and limitations in such data. Our top-down approach yielded a map of likely dust emission hotspots (not average AOD or DOD, as the Comment claims) that agrees well in general with **bottom-up** estimates of dust 'Source Intensity' (SI) maps (as we show in Fig. 6). This comparison is the main 'validation' of our result.

Rather than studying 'events' to validate the MODIS DOD product (not our goal), we identified two 'new' dust source areas (unpublished in the literature, to our knowledge, and not found in the bottom-up SI map of Fig. 6) and we examined if they are consistent with being dust sources. Indeed, they have correct surface features for dust emission (see visible imagery we presented in Figures A1 and A2) with high average wind speeds for horizontal dust flux mobilization. We also discussed these areas with field Geologists who have visited the Québec/Labrador site during field studies and who confirm dust sources in this region. Nevertheless, we cautiously recommended in our work that additional observational studies are needed (including at the 'event-level', which is exceedingly difficult to do in remote areas, as is well known).

Finally we would underscore the fact that the authors did not use the DT MODIS option of AODs associated with dark pixels over water : this product is on much firmer ground than the DB AOD product and the DT product over land (much less dependent on the surface reflectance).

Again, we are looking for dust **sources**, which are on **land** and not over the ocean. It would be inappropriate for us to use Dark Target (DT) MODIS data, which is not optimized to perform well over brighter land surfaces. We are also not expecting or tracking dust plumes far from their source regions, including over dark water surfaces, as we are fully aware that there are no Arctic-equivalents of Saharan dust plumes – only that, very occasionally, near sources, DOD can reach levels comparable to some Saharan plume levels.


We agree that where in Section 3.6 of our paper we compare MODIS AOD to AEROET AOD in geographic boxes that include both land and ocean, neither dataset (DB nor DT) is ideal.

* including the recent publication by O'Neill et al., "Remote sensing detectability of airborne Arctic dust", (2025). That the authors apparently decided not to rebut its strikingly opposing message of very low Arctic DODs compared with their very high DOD results indicates a lack of due diligence on their part. Note also the Kawai et al. DOD simulations (that are cited in O'Neill et al): those simulations yielded CM AOD results which were even smaller than the CM AOD climatologies cited in the latter paper (DODs being, in general $< \sim$ CM AODs).

We are aware of O'Neill et al. (2025) and we read its Discussion paper in detail before our submission to ACP. We did not cite the work because it focuses on problems with using the Brightness Temperature Differences (BTD) at 11 and 12 μm to detect Arctic dust, which is quite different from the MODIS DOD dataset. Furthermore, that paper largely focuses on detecting average values of coarse mode dust, while we concern ourselves with extreme values, and the study states that it is "difficult if not impossible to detect using passive satellite-based techniques at any wavelength", which we do not disagree with. However, we also knew that one of the concluding statements of this study is: *"While it is difficult to account for all examples of strongly negative BTD_{11-12} , it is very unlikely that airborne dust plays a major RS role in any case **other than***

plumes of strong DOD ($> \sim 0.5$).” Our work is precisely a tally of such extreme dust events in one satellite dataset. We are happy to cite O’Neill et al. (2025) as it supports the validity of our search for occasional extreme events in RS data.

As to the simulations of Kawai et al., who include Arctic dust emission in their model, again, our FoO maps for extreme events are not comparable to any climatology of average CM AOD or CM DOD (neither based on models nor based on measurements). However, it is nice to see in the Supplementary Materials of Kawai et al. that our FoO maps (left below) match the spatial patterns in their simulated annual mean dust emission fluxes (right below, red is logarithmically more emission than blue).



We have inserted comments throughout this pdf to explain our point of view and are open to exchanges with the authors if they wish clarification on any of those comments. Our major high-level comments are the comment on lines 76 and 214 and the comment associated with the sentence that starts on line 295.

We are of course open to collaborative and collegial exchanges and modifications that will frame our results in the most objective light, while recognizing that our goals were not satellite dataset validation nor event-level studies of individual optically thick dust plumes far from their sources. We will directly address the main points made on lines 76, 214 and 295, and a few other points, leaving the specifics of all modifications for later in the peer-review process.

[end of high-level comments]

LINE 76 (MODIS DB vs DT): Given that there are significant areas of open water in the late-spring to early-fall in the CAA and that open water dark target (DT) AODs are clearly a more universally accepted product than deep blue (DB) AODs, why was the analysis limited to dust events over land?!

The analysis was limited to dust events over land because dust sources exist on land and that is the focus of our study. Given the low dust emissions in the Arctic, it only makes sense to look for thick plumes ($\text{AOD} > 0.5$) right over land sources, as per the available *in situ* observations of such events. Plumes transported farther from sources will be diluted and also de-coupled from their source areas.

(b.t.w: the quality flag (QA) of the pixels used for DOD pixels should have been stated).

The effect of the quality flag is shown in Fig. 2 and discussed in detail below that (on pg. 10).

We have tested the Level 2 (L2) DB algorithm in the Arctic and have found examples of significant positive biases of the latter over the former at the boundaries between the two products.

Good to know. Curious if this was in individual L2 granules or an L3 product. Either way, our FoO maps primarily show potential dust source regions away from coasts.

Conversely, we have also found examples of obvious dust events over land that the DB algorithm did not catch* (while the MAIAC algorithm at least produced AOD estimates over land). We have also seen that the DB AOD product can be very sporadic (low sampling numbers) relative to the DT algorithm. See, for e.g. the Aug. 18, 2011 event (URL below **) and scroll through, day by day, the whole summer of 2011 while turning the DT and DB AOD algorithms on and off.

This is in agreement with our low numbers of MODIS DOD (DB) observations over 20 years (Fig. 1, bottom row) and shows that our work on sources (over land) could perhaps be improved in a future study using the MAIAC product.

We also noted examples of suspicious similarities between dark reflectance patches and both high DT and DB AODs (to the point where the suspicious DT and DB AODs would "hover" over the same place from day to day [see ** below, for e.g.] while DK AODs over water would evolve in a fashion that was coherent with the natural appearance of plumes in the RGB images).


That is interesting. Perhaps local dust (of a coarser size) is indeed 'hovering' over the source area by actually being repeatedly injected into the atmosphere near the time of the satellite overpass and then settling out at night, while the transported dust fraction (of a finer size) is more mobile.

Finally, the authors should have been aware of Sayer et al.'s (2019) paper on the validation of the DB AOD algorithm: a paper which included OPAL and PEARL in the set of DB AOD validation sites (those site statistics were part of their "BOR" category) over a nominal validation period of 2000 to 2017. The marginal correlation coefficients (≤ 0.42 for Terra and ≤ 0.56 for Aqua) and the small number of validation points (we can provide those statistics upon request or the authors can contact Sayer directly) are not the type of scattergram results that one would employ to claim satellite-derived DOD statistics that are decidedly outside the norm.

Sayer et al.. Validation, stability, and consistency of MODIS Collection 6.1 and VIIRS Version 1

Thank you for this newer reference on the DB product from MODIS and VIIRS. We find that it reports the correlation coefficients between AERONET and MODIS Terra and Aqua in the BOR region as 0.75 and 0.76, respectively (Table 1), with respective numbers of coincident validation data points as 10,225 and 9,158. We see that the BOR region is **1)** very sparsely validated by AERONET (Fig. 2, red points, figure snippet reproduced below, with just one point in the Canadian Arctic), which is not news to anyone, **2)** that the fraction of days with L3 coverage is low (< 0.25) in the Canadian Arctic (Fig. 15), also not news, and **3)** that inter-sensor correlation is high for AOD (Fig. 16) but mixed for AE (Fig. 17), with AE correlation 'ok' between Aqua and Terra, but poor between either of these and VIIRS (we come back to this last point below).

Irrespective of the above, which does not send a strong signal not to use MODIS DB data in the Canadian Arctic Archipelago, we make no claims about ‘satellite-derived DOD statistics’ – only their frequency of occurrence as a spatial map, based on 20 years of ‘hits’ of extreme values.



To summarize all these critical DB AOD / DOD comments, the authors were, given the significant physical / optical complexities of the Arctic environment, found wanting in not performing a much more detailed analysis of the DB AOD product that was the basis of their dust analysis (and they notably did not report any event-based DOD analysis: the very large-DOD statistics that they reported should have motivated them to at least find a few concrete examples of such strong regional DOD values)

We already addressed the essence of this comment in our responses above.

*see, for e.g.: [Link](#)

** this was one of our events of interest [Link](#)

LINE 214: It is a significant oversight that the authors do not cite Sayedain et al., (2023) in addition to the Huck et al. paper. Sayedain et al. provided in-situ lidar as well as AERONET-derived estimates of coarse mode (CM) AODs and OPS (Optical Particle Sizer) measurements of CM particle-volume concentration near the source of strong sub-Arctic dust plumes (the Kluane Lake, Yukon site of Huck et al). They developed a means of classifying dust plumes using correlations between the lidar CM AODs and the CM OPS concentrations and the attendant correlation between the lidar CM AODs and AERONET CM AODs to respectively (i) define the presence of a dust plume on any given day and (ii) to define whether that dust plume could be remotely sensed (their class "D_{RS}"). Referring to their Figure 2 ensemble of (red-coloured) class "D_{RS}" points one can show that ~ 11% of those DODs were > 0.5." (the relatively broad (averaging) footprint of MODIS will have its own influence on the reduction of DODs within a near-source plume). These near-source DODs will systematically decrease (along with the % > 0.5) as the dust plumes are dispersed downwind. In a nutshell: 'persistently high dust loading' is largely limited to the close vicinity of the drainage basins whose wind-erosion dynamics incite the plumes.

Sayedain, S. A., O'Neill, N. T., King, J., Hayes, P. L., Bellamy, D., Washington, R., Engelstaedter, S., Vicente-Luis, A., Bachelder, J., & Bernhard, M. (2023). Detection and analysis of Lhù'ààn Man' (Kluane Lake) dust plumes using passive and active ground-based remote sensing supported by physical surface measurements. *Atmospheric Measurement Techniques*, 16(17), 4115–4135. <https://doi.org/10.5194/amt-16-4115-2023>.

As we mentioned in the second paragraph of this reply, the phrase ‘persistently high dust loading’ snuck into this manuscript and needs to change to ‘more frequent signatures of extreme dust

loadings' because we do not make any claims of average dust loadings, only their frequencies. Drs. Sayedain and O'Neill comment that our Fig. 7 represents AOD values for the Arctic and point out the 'astonishing' occurrence of AOD ~2. Indeed, that would be quite astonishing, but fortunately Fig. 7 shows the **spatial average** (in a very large geographic region) of the **frequency of occurrence** of **extreme** values of AOD (i.e., AOD > 0.5), not the spatial average of AOD itself. (The figure also shows DOD values in the same sense.) We will re-label the y-axes of this figure for improved clarity, although it is discussed in the text.

We are aware of the work of Sayedain et al. (2023) but chose not to cite it because it does not change our goals, methods or conclusions and we had already cited Huck et al. (2003), who show extreme dust events near the same dust source in the Yukon Territory. We will nevertheless cite it because, as the authors note, there are so few field measurements of dust and their study also documents extreme dust events using both *in situ* and ground-based remote sensing measurements (extreme to the point that they cause false positive cloud events in AERONET algorithms).

As Sayedain and O'Neill point out in their comment above, they estimate 11% of plumes that can be remotely sensed had DOD > 0.5, which further supports our attempts to identify the spatial mapping of the frequency of such events in MODIS DOD data (at high spatial resolution) to identify dust source regions. Our relative frequencies of such extreme events are 0-50%, depending on location in a large and unmonitored (from the ground) geographic region, which is not grossly different from their finding of 11% at one spatial location downstream of a dust source.

Their study is primarily concerned with the classification of dust events as those that can be detected remotely (with active and passive ground-based remote sensing data) and those that cannot; they also postulate the presence of dust particles of radius 11-12 μm based on their data, and they explore the detectability (by remote sensing) of fine mode dust particles that would better indicate long-range dust transport in the Arctic. It is interesting to consider how such extremely large particles (reasonable right at a dust source) that occupy a near-surface layer of 1-2 km thickness (their Fig. 3) would show up in passive MODIS measurements (based on surface-reflected solar radiation) as compared to passive AERONET measurements (no surface reflection).

LINE 295 VIIRS: Dust "accounting for 80% of all aerosol type retrievals in some hotspots" over a 3-year period (and notably over substantial regions according to Fig. 4a) is simply unrealistic. Fine mode Arctic haze and fine mode smoke AODs will, in general, dominate DODs that are not measured very near to strong local dust sources.


In light of the poor correlations between VIIRS and MODIS Terra or Aqua reported by Sayer et al., 2019, we will re-check our results based on VIIRS products of Aerosol Type == 'Dust'. Regarding (*) below, AOD > 0.3 is not 'our' threshold, but rather this is the VIIRS threshold for producing an Aerosol Type == 'Dust' product, which we used in our analysis.

W.r.t. to our own DOD experience at KLRS (see the comment above in line 214 concerning the <~ 11% FoO at the [near-source site of KLRS](#)). We also have significant experience at Eureka and other

sites in or near the Canadian Arctic Archipelago where (publication being prepared) the near source DODs are generally $< \sim$ KLRs DODs).

The VIIRS analysis of the authors shows FoO values ranging from ~ 10 to 50 over 3 years (2020-2022) in the Eureka region (their Figure 4a). An investigation of all AERONET L1.5 (cloud-screened) data at the OPAL AERONET (Eureka) site from 2020 to 2022 resulted in a grand total of 7 CM (coarse mode) L1.5 AODs $> 0.3^*$. The authors should have carried out this type of concrete detailed validation^{**}. In general they failed to produce satellite-based DOD validation at any kind of event level.

Regarding (^{**}), the documented presence of 7 CM events at Eureka (on the same day) is on the same order of magnitude as the lower bound of ~ 10 -50 events in various nearby places in our Figure 4a (over the same time periods), which we reproduce below, with a tiny **green square** at the location of Eureka (80 N, for those not familiar with its location who may have a hard time spotting the green square). Also, as we illustrate with our Fig. 10 for **Resolute Bay** (with much more data available for comparison to MODIS than at Eureka, which we show in Fig. 8), the location of a point station even in close proximity to a dust source may not guarantee detection of dust at that station, unless the station is positioned precisely downwind of the dust source. **Given that Eureka is one of the only observing stations in the Canadian Arctic Archipelago and that retrieval numbers dwindle so high north, one can hardly speak of any meaningful satellite-based DOD validation in this region** – not that this was our goal in section 3.2, called ‘Inferences about dust over Canada from VIIRS’. That section was written to document extra correlative data that shows dust in the Arctic, as was Section 3.3 for CALIOP products of Aerosol Type == ‘Dust’.



* the VIIRS retrieval threshold for AOD employed by the authors

^{**} for the record, those 7 CM AODs occurred on one single day (16:24 to 16:45 UTC on 23/03/2020). The MODIS image of that day (Worldview URL below) shows a completely snow-covered surface in the Eureka region. In effect there was no validation data available for the period of 2020 to 2022 that would sustain any claims of DOD retrieval validation in the high Canadian Arctic

At the "point" (single site) to regional (6 AERONET sites) scale of the CM AOD climatology of AboEl-Fetouh et al. (2020) (AeF) the largest (geometric mean x geometric standard deviation) summertime CM AOD (an upper limit for DODs) was $0.003 \times 2.13_{\pm 1}$ for the month of June at Resolute Bay and OPAL, Nunavut. This number is smaller by two orders of magnitude than the 0.5 threshold of

Figures 3 and 4: the attendant FoO computations using a lognormal distribution* driven by AeF's stats yielded, not surprisingly, FoO percentages that are all much below 1%)

* results which can be supplied upon request


[Worldview URL](#)

Again, we presented qualitative Aerosol Type == 'Dust' VIIRS products while the authors are referring to climatologies of CM AOD (> DOD), also from the handful of ground-based (point observation) stations operating in the remote Arctic environment.

Other comment: Our experience has been that CALIOP dust classification and attenuated extinction coefficient capabilities are at the margins of significance with respect to the weak optical signals of dust in the Arctic. Kawaii et al. (2023) tried to develop a CALIOP-derived local dust optical-depth climatology for the Arctic and found that CALIOP predictions of DOD were significantly greater than predicted by their simulations (see their supplementary material). There is no published literature on the successful validation of CALIOP retrievals of weak and low altitude Arctic dust plumes.

Our CALIOP data of Aerosol Type == 'Dust' is based on JJA observations (2006 – 2020), when dust-cirrus confusion will be least (below, right). We do not draw any critical conclusions from this figure except to note that signatures of the **increased count** of boundary layer-like dustiness (< 1 km altitude) are apparent during dust emission season in the large geographic region we call the “Canadian Arctic Dust Belt”, and also in the vicinity of Greenland. Nevertheless, we don't dispute that even the heavily-averaged (in space and time) CALIOP figure of dust classification counts may be pushing the limits of detection and we are happy to learn more about this.

As to Kawai's simulations (below, left), there are a number of reasons why their simulations could be lower than the potentially true (but unvalidated) CALIOP reality, i.e., model representation of dust source areas; model resolution ($1.9^\circ \times 2.5^\circ$ with 30 layers between 0-40 km) influencing both 3-D wind speeds *and* representations of snow and ice cover and soil moisture; moreover, the nonlinear dust flux parameterization itself can be updated. See, e.g., the work of Meng et al. (2021, [GMD](#)) who have implemented offline dust emissions in GEOS-Chem (with a source function spanning low latitudes) at the native resolution of the meteorological fields ($0.25^\circ \times 0.3125^\circ$), which lead to better agreement in comparisons to (low-latitude) AERONET; or see the high resolution ($0.25^\circ \times 0.25^\circ$) SI map that we compare to in Fig. 6 (Vukovic, 2021; 2022).



Response to CC 2

Blue font is the content of Community Comment 2, CC 2, as submitted.

Green font indicates changes we made to CC 2 to make it easier to follow.

Black and red font are our replies (AC 2).

The same community Commenters (Sayedain and O'Neill) submitted CC 1 and CC 2; our AC 1 already responded to CC 1.

The points below are our reaction (CC 2) to comments submitted by A&W (Ashpole & Wiacek) on Feb. 11, 2025 (AC 1, in response to CC 1). Details of our comments that follow had already been transmitted to Wiacek in a Feb. 11 email¹ (outside of the EGU sphere discussion forum). In this current document we react to their AC 1 by limiting ourselves to the most important critical points made in our Feb. 11 Email². Our first 3 critical comments below are the most important.

¹ that preceeded the deposition of AC 1 in the discussion forum and for which we received no answer

² this is our 2nd reaction to the A&W paper that we have submitted to the EGU sphere discussion forum. Our 1st submission (CC 1) consisted of comments (annotations) made directly on their manuscript pdf.

Before we engage in the charade of being ‘completely objective scientists’ and address the **substance** of these comments point by point, we will point out that CC1 and CC2 (both made by PhD student Sayedain with Professor O'Neill) are **biased** (only limitations are noted, without commenting on positives in any way, and without summarizing our paper at the outset, as is the reviewing norm), **hostile** (in CC 1 comments on our manuscript), and made under **two major conflicts of interest**.

First, Wiacek and O'Neill are currently on three major community grants together related to the HAWCSat project (one of these grants is currently under review), **directly competing for funding from the same funding envelopes**. A paper detailing HAWCSat has just come out in [BAMS](#), including O'Neill and Wiacek as co-authors. We were also both competitively funded from the same programs in the past (CSA's ESSDA1 & ESSDA2).

Second, it is problematic for these comments (CC 1 and CC 2) that Sayedain is currently preparing a paper on Arctic-wide dust remote sensing, although we doubt there is any overlap in our methodologies, and we recognize that anyone is free to use open satellite data to analyze the Arctic how they wish. We know this because it was stated in a poster presented at the HAWCSat meeting in Montreal (Oct 15-17, 2025), which we attended together, delivered by Sayedain, O'Neill and others (“Remote sensing of local dust across the Canadian Arctic Archipelago”). The last bullet point of this poster's *Introduction* states: “The Canadian Arctic Archipelago remote sensing analysis is the subject of a paper in prep

(Sayedain et al., 2025)”, which was promised to appear on EGU sphere imminently, but not yet as of the time of writing this AC 2. Moreover, the second bullet point in the poster’s *Key Findings* states: “Satellite RS detects optically thicker plumes across the CAA with CM AOD $< \sim 0.56$ ”. Our same focus on Arctic-wide dust phenomena appears to be highly unwelcome, triggering very high standards of proof for our work’s validity.

We confirm that we initially engaged in an **email exchange initiated by O’Neill**, in parallel to their EGU sphere comments. We stopped responding to these emails when it became clear to us that our intellectual independence was not recognized by the Commenters, which is an [obligation](#) of a Copernicus journal referee, and by extension any Commenter.

1. Re the reluctance of the authors to consider any type of validation / evaluation of their FoO (DOD) statistics

We strongly disagree with that point of view. DODs > 0.5 are easy to detect visually over water and over land near the source³. It is, given the questionable quality of DB AOD retrievals over the high Arctic, the responsibility of those who would employ DB AOD statistics (as evidence of the presence of optically thick airborne dust over the Arctic) to find concrete illustrations for a handful of their largest DODs. Such cases could be included in the supplementary material to avoid disrupting the general flow of the narrative in the main paper.

³ see, for e.g., Ranjbar et al. (2022), Sayedain et al. (2023) and Sayedain et al. (2025).

This point was already addressed in our AC 1 response to CC 1. It is misleading to say that we have not considered ‘any type of validation / evaluation’ of our FoO (DOD) statistics.

First, we worked hard to ‘validate / evaluate’ our Frequency of Occurrence (FoO) of Dust Optical Depth (DOD) statistics by 1) comparing their spatial patterns to potential dust source emission areas (G-SDS-SBM map, Fig. 6), 2) comparing with two different satellite dust products (VIIRS, CALIOP, Figs. 4 and 5), and 3) comparing AOD at a strong dust emission location in Resolute Bay to the AOD from the AERONET station (Fig. 8), noting also that there are not many AERONET stations to choose from, and 4) checking if two new dust source areas implied by our analysis (Figs. A1 and A2) are plausible (they are).

The issue is that Sayedain and O’Neill are asking us to validate the DB AOD dataset *itself* (from which DOD is derived, using also the Angstrom exponent), which is a very high standard of proof. It is far beyond the scope of this semi-quantitative work.

Second, at the start of our work we spent considerable time searching for visible dust plume events and we disagree that they are ‘easy’ to detect. Arctic cloudiness (80% of the time over open ocean in summer, when dust emission is most active, see, e.g., Zamora et al., 2022, ACP) and Arctic ice are a normal state, decimating the dataset, even in JJA. Dust

over water is easier to see, but it can still look like smoke, even in the high Arctic, and in any case the land source of the dust might then be under clouds (no satellite data), yielding no information about the emission source (our objective). Dust over land is not at all easy to see given the bright desert conditions of the Arctic when it is ice-free, which is why Sayedain and O'Neill have focused on dust over water in the analysis of single dust events (and also because DT AOD over water is *better* validated than DB AOD over land, with DT over land still inaccurate). We chose to work around these limitations by carrying out a 20-year study of elevated DOD frequency counts, working on the reasonable hypothesis that spatial features of the analysis will agree with known potential dust source emission areas (as they did).

2. *Re the reluctance of the A&W authors to look into DT (dark target) AOD retrievals over water*

In our experience, strong plumes emanating from the land in the CAA (Canadian Arctic Archipelago) can find their way to a body of water⁴. The more believable MODIS DT AOD algorithm over water and the spatial continuity with dust plumes over the land could then be employed to help confirm the all-too questionable quality of DB AODs over the Arctic.

⁴ Their statement that “We are also not expecting or tracking dust plumes far from their source regions” is an oversimplification in the CAA region.

We are not disputing these phenomena or approach. We argue that they represent a different type of analysis, supporting a different kind of study, and leading to a different final paper from the one that we have written. We do not wish to simplify the CAA region with our reluctance to track dust plumes over water, but **our work (see next paragraph)** was not to “confirm the all-too questionable quality of DB AODs over the Arctic.” We counter that here, Sayedain and O'Neill oversimplify the effort involved in what they are suggesting, i.e., starting from the DT dataset and connecting high AOD values over water to high DB AOD values over land, which requires some level of back-trajectory analysis and logical, e.g., threshold-based, processing.

Related to ‘our work’, O'Neill and Wiacek have been able to secure funding from the same funding envelopes as co-applicants precisely because of this different focus of our groups, with O'Neill specializing in the details of remote sensing retrievals and analysis, including the development of new techniques and data products, which then requires case-by-case validation analyses; in contrast, Wiacek currently makes use of existing remotely sensed data products to, e.g., learn more about dust source regions in a big picture sense. In this case we used a *sparsely* validated DB AOD product to calculate DODs and FoO statistics to infer land-based dust sources. There are no recommendations against using DB (or DT-land for that matter) datasets for **semi-quantitative studies, like ours**, which had as its goal

the satellite-based identification of potential dust source regions, not the creation of mean AOD or DOD maps. In the revised manuscript we have included text (Abstract, Conclusions) to highlight and acknowledge that these potential dust source regions should be considered for further exploration / validation, ideally through a multi-modal approach, i.e., *in situ* observations and additional or improved RS products.

3. Re the use of the “BOR” (Sayer, 2019) statistics rather than the PEARL & OPAL statistics that we recommended

In actual fact 24 out of the 27 “BOR” sites are in the low Arctic tundra⁵. Resolute Bay and Iqaluit (which were not employed by Sayer, 2019) and OPAL and PEARL make up the sites that are in the high Arctic tundra⁶. Those high Arctic tundra statistics are quite marginal⁷.

We remain, accordingly, unconvinced by the identification of very large DB AODs⁸ as being publishable indicators of large-DOD occurrences. Put another way, we don’t think it is appropriate to use largely unvalidated DB AODs over the high Arctic tundra as a form of DOD “ground truth”. One can complain about the lack of statistics in the high Arctic tundra ... but that only confirms the argument that there isn’t enough validation data to support claims of not insignificant numbers of (DOD > 0.5) instances.

⁵ The ecosystem-based delineation of the Arctic regions: see for e.g., the [Britannica website](#)

⁶ this was always the focus of all our comments: the high-Arctic tundra region that includes basically all of the CAA.

⁷ “The insignificant to marginal correlation coefficients (≤ 0.42 for Terra and ≤ 0.56 for Aqua) and the small number of validation points (we can provide those statistics upon request or the authors can contact Sayer directly) are not the type of scattergram results that one should employ to claim satellite-derived DOD statistics ...” (statement from the comment associated with line 76 in our original EGU sphere Discussion forum comments in CC 1 file)

⁸ which are transformed to DODs given the constraints that the A&W applied (we have no strong argument with those constraints)

To clarify, we did not “use” BOR statistics in our paper, but only commented on them in AC 1, while replying to CC 1, which brought these up in the context of the validation of the DB dataset. We did not fully understand this CC 1 comment (on line 76), but now we infer that Sayedain and O’Neill were providing us with the Terra and Aqua DB AOD vs. AERONET AOD correlation coefficients for the subset of high-Arctic AERONET stations only, which were weaker than for Sayer’s ‘BOR’ region ($r = 0.75$, $r = 0.76$, i.e., ‘strong’) that was overwhelmingly based on low-Arctic stations. Assuming that the high-Arctic values provided by Sayedain and O’Neill are also the smaller Spearman coefficients (this is not critical, but then they are directly comparable to the ‘strong’ values we cited above), their upper limit values ($r \leq 0.42$, $r \leq 0.56$) would typically be referred to as ‘moderate’ instead

of ‘insignificant to marginal’, with $r < 0.3$ typically referred to as ‘weak’ (because of the inequality used by Sayedain and O’Neill, it’s not clear how many high-Arctic AERONET stations have ‘weak’ correlations).

Regardless of the strength of the correlation between DB and AERONET AOD in the high Arctic, our semi-quantitative work on frequencies of occurrence of extreme AOD and DOD events (extreme but well within the range of observed high-latitude dust storm measurements) is a reasonable use of the DB dataset, i.e. to infer *dust source locations*. We have now analyzed the % of DOD values > 0.5 , 1.0 , 1.5 , 2.0 and found these to be 1.6%, 0.9%, 0.02% and 0.008%, respectively, i.e., the extreme values are not common, consistent with the low mean values of Arctic AOD and DOD found by Sayedain and O’Neill (and others).

We will once again state that validating the DB dataset in the high-Arctic is an exceedingly high standard of proof for our type of work, especially given that such validation is unlikely to happen in the foreseeable future given the prohibitive conditions of the Arctic.

4. Persistent dark spots in the imagery associated with large DB AODs


[pink color added by AW for clarity] The author’s comment⁹ ignores the more practical DB retrieval problems that can occur. A dust plume over the darker areas should, in the presence of roughly (spatially) continuous surface reflectance, have produced a brighter reflectance, not a darker reflectance. The more likely culprit is an **artifactual decrease in the DB-estimated surface reflectance (and/or a real decrease in the actual surface reflectance)** caused by a change in surface type or terrain slope (BRDF); see, for e.g., Liu et al., 2017 for the important influence of surface type in the CAA): **this would have produced an excessively large DB AOD** (we can be reasonably sure that it is excessively large because the DT AOD over water is a much more universal and dependable AOD product).

⁹“Perhaps local dust (of a coarser size) is indeed ‘hovering’ over the source area by actually being repeatedly injected into the atmosphere near the time of the satellite overpass and then settling out at night, while the transported dust fraction (of a finer size) is more mobile.”

The CC 1 comment that we were responding to in footnote ⁹ above, was: “*We also noted examples of suspicious similarities between dark reflectance patches and both high DT and DB AODs (to the point where the suspicious DT and DB AODs would “hover” over the same place from day to day [see ** below, for e.g.] while DT AODs over water would evolve in a fashion that was coherent with the natural appearance of plumes in the RGB images).*”

We understand that bright dust over a dark surface would lead to brighter, not darker reflectance. We do not dispute that retrieval problems can occur as described by Sayedain and O’Neill in CC 1 and CC 2. Variable surface reflectance over land is a challenge in

retrievals, and we included a statement about this in our Conclusions. The scene referred to by ** in CC 1 (Aug 18, [linked](#) again) is a large region and it is not clear which high AOD / low reflectance patches are thought to be ‘hovering’ by Sayedain and O’Neill, nor what they mean by ‘**high**’ AOD. The large yellow-orange-red (AOD < 0.5, DOD would necessarily be lower still) patches on Aug 17-23, covering large fractions of Baffin Island, would be excluded in our analysis (DOD > 0.5), whether they were real (e.g., transported smoke) or artifactual (reflectance, thin cirrus, other retrieval issues). What interests us in this large scene is a few pixels of **dark red** (AOD 1.13 to 1.56), which remain quite attached to one location on western Baffin Island on Aug. 12,15,16,17,22, shown below (**left** image; **dark red** within the **pink** box). This location is, by inspection, a glacial outwash plain (**right** image; which is *roughly* zoomed to within the **pink** box, and **bottom** large [image](#), with more zoom), which appears to be a dust emitting area, also captured by the 20-year mean of **extreme** (DOD > 0.5) events on our FoO maps. **We added this case to our Supplements.**



It does not seem that we will settle our differences case by case, or event by event. It is not clear to us why a patch of $1.1 < \text{AOD} < 1.6$ ‘hovering’ for a few days is ‘suspicious’ or ‘excessively large’ in a glacial drainage area. But we are quite sure that if there is a remote sensing paper to be written about what the systematic problems are with using DB AOD over land as we have used it, Sayedain and O’Neill will write it, moving us all forward. This is a different paper from our submitted work.

5. Re the use of the O’Neill et al. (2025)

That citation was taken out of context: it applies to MODTRAN-derived $\Delta\text{BTD} / \Delta\text{DOD}$ derivatives that we employed to show that it would require visible-wavelength DODs $> \sim 0.5$ to produce measurable BTD_{11-12} values¹⁰ (to “extrapolate” that sentence to the remote sensing detection of dust plumes in the visible is inappropriate¹¹).

¹⁰ Values that are also functions of the surface temperature, the amplitude of the temperature change across the inversion layer and the position of the plume relative to the inversion layer. Visible wavelength DODs (rather than TIR DODs) were employed because the visible DOD was a bit of a fixed standard in the literature.

¹¹ Which is actually what A&W themselves state at the beginning of the same paragraph in their AC 1.

To clarify, we did not “use” O’Neill et al. (2025) in our paper, but only commented on it in AC 1, while replying to CC 1, where it was brought up. Fair point – we will not extrapolate to the visible; we only meant that an ‘extreme’ dust event is noted in that work with a (visible) DOD ~ 0.5 , but it is in the work of Ranjbar et al. (2021), also with O’Neill, whom we already cite, so nothing new would be added by citing O’Neill et al. (2025).

We note also that this comment cuts both ways: O’Neill et al. (2025) cannot be used as justification for why **extreme** (> 0.5) DB AOD’s are invalid in the high Arctic in the vicinity of presumed / potential dust sources, because their study was written to highlight the limitations of the approach of using BTD_{11-12} while referring to ‘pan-Arctic’ events of **diluted** dust concentrations, as captured by a network of six AERONET stations that are not set up to detect extreme AOD’s near dust sources. The inability of one of these stations to capture even very nearby dust sources, Resolute Bay, is illustrated in our Figure 10. Also, O’Neill et al. (2025) focused on events from October to April, while our multi-year analysis focused on the period from May to September in the high Arctic.

References

- Liu N, Budkewitsch P, Treitz P. Examining spectral reflectance features related to Arctic percent vegetation cover: Implications for hyperspectral remote sensing of Arctic tundra. *Remote sensing of environment*. 2017 Apr 1;192:58-72.
- O'Neill, N. T., Ranjbar, K., Ivanescu, L., Blanchard, Y., Sayedain, S. A., Aboel-Fetouh, Y.: Remote-sensing detectability of airborne Arctic dust. *Atmospheric Chemistry and Physics*. 2025 Jan 6;25(1):27-44.
- Ranjbar, K., O'Neill, N. T., and Aboel-Fetouh, Y.: Comment on “Short-cut transport path for Asian dust directly to the Arctic: a case Study” by Huang et al. (2015) in *Environ. Res. Lett., Atmos. Chem. Phys.*, 22, 1757–1760, <https://doi.org/10.5194/acp-22-1757-60>, 2022.
- Sayedain, S. A., O'Neill, N. T., Ranjbar, K., Gauvin-Bourdon, P., Chang, R., Hayes, P. L., King, J.: Remote sensing of local dust events over the Canadian Archipelago, oral presentation, HLD workshop, Agricultural University of Iceland (AUI), Reykjavik, Iceland, Feb. 12-13, 2025.
- Sayedain, S.A., O'Neill, N.T., King, J., Hayes, P. L., Bellamy, D., Washington, R., Engelstaedter, S., Vicente-Luis, A., Bachelder, J., Bernhard, M.: Detection and analysis of Lhù'ààn Mân'(Kluane Lake) dust plumes using passive and active ground-based remote sensing supported by physical surface measurements. *Atmospheric Measurement Techniques*. 2023 Sep 12;16(17):4115-35, <https://doi.org/10.5194/amt-16-4115-2023>.
- Sayer, A. M., Hsu, N. C., Lee, J., Kim, W. V., Dutcher, S. T.: Validation, stability, and consistency of MODIS Collection 6.1 and VIIRS Version 1 Deep Blue aerosol data over land. *Journal of Geophysical Research: Atmospheres*. 2019 Apr 27;124(8):4658-88.

On the following pages we present detailed responses to all comments from CC 1 that we left for later in our quick AC 1. They have, for the most part, made our paper better, even though the high and generally constructive reviewing standard of the EGU sphere community was, in our opinion, not met.



Satellite-based evidence of dust emission over Northern Canada

Ian Ashpole¹, Aldona Wiacek^{1,2}

¹Department of Environmental Science, Saint Mary's University, Halifax, Canada

²Department of Astronomy and Physics, Saint Mary's University, Halifax, Canada

5 Correspondence to: Ian Ashpole (ian.ashpole@smu.ca)

Abstract. High latitude dust (HLD) is receiving growing research interest as its relative impact in the Arctic has been validated. Precise knowledge of HLD emission locations is limited in both field studies and satellite observations, leading to a general lack of representation in global models. Using the Frequency of Occurrence (FoO) of above-average Dust Optical Depth (DOD > 0.5) from twenty years (2002-2022) of high-resolution MODIS observations derived for this study (0.1° x 0.1°), we present quantitative evidence that dust sources are widespread across the Canadian Arctic Archipelago. Additionally, we present qualitative supporting evidence from aerosol type 'dust' classifications in VIIRS and CALIPSO satellite data products, as well as some challenges of comparing MODIS AOD to two co-located AERONET sites.

The HLD hotspots identified in the "Canadian Arctic Dust Belt" correspond to surfaces with high potential for dust emission in the G-SDS-SBM dataset. There are more areas where hotspots are observed but emission potential is low than the opposite case; additionally, two considerable areas of dust emission are identified at lower latitudes in mainland Canada. When spatially averaged across the broad dust producing region (65°N – 85°N, 125°W – 70°W), annual mean time series of FoO of MODIS DOD > 0.5 between 2003 and 2022 suggest an increase in the frequency of dustiness in the latter half of this period, consistent with our understanding that HLD emissions are increasing in a warming climate. These results further motivate model development to include HLD sources and provide an observational basis for evaluating them.

20 1 Introduction

Mineral dust aerosols (herein "dust") play an important role in the Earth System, for example through modifying the atmospheric and surface radiative budgets (Kok et al., 2018), influencing cloud processes (Tobo et al., 2019), providing nutrients to terrestrial (Nogueira et al., 2021) and oceanic (Dansie et al., 2022) ecosystems, and reducing the albedo of snow and ice when deposited (Sarangi et al., 2020) on these surfaces. Dust can also pose a significant health risk when particles are inhaled, with increased risk found for populations living downwind of source regions (Yang et al., 2022).

Dust emission depends upon high wind speeds occurring over surfaces where dust-sized particles (< ~100 µm) are available for entrainment. Therefore, the most dominant global dust hotspots are the hot subtropical deserts (e.g. Prospero et al., 2002; Washington et al., 2003), where a lack of surface moisture and vegetation create favourable conditions for emission to occur. However, dust sources in colder, mid-latitude regions have been shown to have a higher potential for interactions with high ice clouds, despite being much smaller dust emission sources (Wiacek et al., 2010). More recently, high latitude



regions have received growing research interest too, even though preferential dust source areas in the Arctic have been identified two decades ago (Tegen, 2003 and references therein). Although current best estimates suggest that they contribute only around 3 – 5%, by mass, to the global dust cycle (Bullard et al., 2016; Groot Zwaafink et al., 2016), there is evidence that high latitude dust sources have a disproportionately large contribution to radiative forcing in the Arctic in comparison to dust from lower latitudes, due to being the dominant contributor to near-surface loadings and deposition there (Groot Zwaafink et al., 2016; Kylling et al., 2018). High latitude dust originates largely in glacial processes, which efficiently produce and deliver fine sediment to barren proglacial floodplains that are the prime emission hotspots when winds (which may be accelerated by regional ice sheets and glaciers) are strong enough (Bullard et al 2016). It is likely that the high latitude emissive areas will grow as a result of Arctic climate change impacts such as glacial retreat, decreased snow cover duration, and permafrost thaw (Bullard, 2013; Meinander et al., 2022), further justifying their continued study.

Meinander et al. (2022) provide a comprehensive overview of high latitude dust, and use numerical modelling, data from the Global Sand and Dust Storms Source Base Map (G-SDS-SBM; Vukovic, 2021; 2022), and case study analysis to make the case for a northern high latitude dust belt. Of the 64 high latitude dust sources documented in that study, only 4 are in Canada. However, presented source intensity (SI) data from the G-SDS-SBM dataset (Fig. 2 in Meinander et al., 2022) suggests that large portions of the land surface in Northern Canada have high potential to emit dust under favourable wind conditions. The study's conclusion calls for an increase in observational studies aimed at identifying new high latitude dust sources, without which accurate modelling of known dust effects is not possible. That is the aim of this paper, with a focus on Northern Canada.

While products such as the G-SDS-SBM can highlight surfaces that have the potential to emit dust, observations of dust storms actually occurring are necessary to classify areas as sources. To this end, datasets derived from satellite instruments are crucial to understanding the global distribution of dust, with areas where dust is most frequently detected determined to be source locations (e.g. Prospero et al., 2002; Washington et al., 2003; Ginoux et al., 2012). Dust detection by these remote sensing methods is based on the exceedance of a threshold value in retrieval products that quantify the amount of aerosol present in the atmospheric column, such as aerosol optical depth (AOD; e.g. Ginoux et al., 2012; Baddock et al., 2016), or binary presence/absence data based on retrieval criteria related to dust (e.g. Ashpole and Washington, 2012). Information on the vertical location of dust in the atmospheric column, provided by instruments such as the Cloud-Aerosol Lidar with Orthogonal Polarization (CALIOP), can also be used to determine the presence of dust source regions, with low altitude dust likely to be proximal to a source (Todd and Cavazos-Guerra, 2016). Since satellite measurements are sensitive to the influence of all aerosol types, not just dust, results are typically verified using additional independent data sources, such as ground observations (when available), retrievals from other satellite instruments, or simulated dust fields from numerical models. Unfortunately, ground observations are few and far between in many known or likely source regions owing to their remoteness, with the combination of environmental variables required for dust emission to occur in the first place being unfavourable for human habitation.

To date, satellite data have only been used to study individual source areas in northern Canada on a case study basis (Ranjbar et al., 2021; Huck et al., 2023). The key objective of this paper is to use satellite datasets to identify where dust is



65 detected most frequently across the whole of northern Canada. The primary dataset used is derived from the Moderate
Resolution Imaging Spectroradiometer (MODIS) AOD, which has been widely used to identify dust source areas on a global
(Ginoux et al., 2012) and regional (Baddock et al., 2016) basis, but not at high latitudes. Covering a 20+ year period, data from
MODIS also enable an assessment of temporal changes in dustiness. “Aerosol Type” data from the Visible Infrared Imaging
Radiometer Suite (VIIRS) instrument, and information on the vertical location of “dust” aerosol type in the atmospheric
70 column from CALIOP are used as comparison datasets to evaluate results from MODIS. Findings are compared to SI from G-
SDS-SBM to assess whether the satellite-derived dust hotspots correspond to surfaces with high emissive capacity; and, finally,
our satellite datasets are verified against surface AOD observations from two Aerosol Robotic Network (AERONET) stations
in northern Canada.

2 Data and methods

75 2.1 Satellite-derived aerosol datasets

2.1.1 MODIS

This study uses Dust Optical Depth (DOD), derived from Aerosol Optical Depth (AOD) data, to map the distribution of dust
aerosols. DOD is a widely used quantity that estimates what portion of retrieved AOD is due to extinction by mineral dust
particles alone, as opposed to the total contribution to AOD from all aerosols. First employed to map the distribution of dust
80 sources on a global scale by Ginoux et al. (2012), DOD calculations use retrievals of Ångström exponent (α) and single
scattering albedo (ω) to isolate the dust contribution to AOD; α is a measure of how AOD changes with wavelength and is
highly sensitive to particle size (Eck et al., 1999). Dust aerosols are typically coarser (particle radius $> 1 \mu\text{m}$) than urban and
biomass burning aerosols (particle radius $< 1 \mu\text{m}$, Dubovik et al., 2002), although the dust size distribution tends towards
smaller radii as distance from source areas increases due to the gravitational settling of larger particles. For coarse mode
85 particles, values of α are low or even negative. The fraction of light that is scattered compared to the total extinction optical
depth, which is a function of both scattering and absorption, is given by ω . Dust absorbs solar radiation and is therefore
characterised by ω values < 1 , separating it from purely scattering coarse aerosols such as a sea salt.

There are various DOD formulations in the literature, and this study employs two of them:

- 90 1. DOD is calculated according to the method of Pu and Ginoux (2016; herein denoted as DOD_{PG16}), whereby DOD is
calculated from every AOD retrieval based on an empirically derived continuous function relating α to AOD
(Anderson et al., 2005):

$$\text{DOD}_{\text{PG16}} = \text{AOD} \times (0.98 - 0.5089\alpha + 0.0512\alpha^2), \text{ if } \omega < 1; \quad (1)$$



2. DOD is simply the AOD value for those retrievals where α is less than a threshold value. Different studies have used different α thresholds, and here the value of 0.3 is applied following Baddock et al. (2016, herein denoted DOD_{B16}), which employed DOD to precisely analyse dust sources in the Chihuahuan Desert, one of the most active dust regions in North America. For clarity:

$$\text{DOD}_{\text{B16}} = \text{AOD}, \text{ if } \alpha < 0.3 \text{ and } \omega < 1. \quad (2)$$

Both DOD formulations are derived to highlight the coarse-mode fraction of dust only, optimising them for dust source identification since the coarse-mode is estimated to account for more than 90% of particles at emission (Kok et al., 2017), and since coarse mode particles remain closer to source regions.

The DOD datasets used in this study are derived from MODIS aerosol products retrieved using the “Deep Blue” algorithm. Deep Blue employs radiance measurements in the blue channels to detect aerosols globally over land, even over bright surfaces such as deserts. The algorithm exploits the fact that because surface reflectance is low and less variable at blue channels, increases of reflectance and spectral contrast indicate the presence of aerosols (Hsu et al., 2013). Our study uses Deep Blue aerosol products from the MODIS instrument (Sayer et al., 2015) carried onboard the Aqua satellite, which passes the equator at around 13:30 local time (ascending node). Products (AOD, α , ω , and various quality flags – see below) are taken from the MODIS Level 2 Joint Atmosphere product (“ATML2”), collection 6.1 (Platnick et al., 2017), with aerosol parameters stored at 10 km spatial resolution in 5-minute time interval swath files. For the Aqua satellite, these files are denoted “MYDATML2”. Data were downloaded from the NASA Level-1 and Atmosphere Archive and Distribution System Distributed Active Archive Centre (<https://ladsweb.modaps.eosdis.nasa.gov/search/order/2/MYDATML2--61>, last access 11 January 2024). For computational efficiency, only files that contained data in the region spanning (150° W – 50° W, 40° N – 85° N) were obtained, for the time period spanning 2002-07-04 to 2022-12-31 (the whole period covered by the MYDATML2 dataset at the time this study was undertaken). The products used in this study were regridded from their Level 2 swath files to a regular 0.1° x 0.1° grid using nearest neighbour interpolation and mosaiced daily, thereby creating a high-resolution Level 3 product optimised for this study.

It is recommended that only aerosol products flagged as “very good” (QA = 3) are used for scientific analysis (Hsu et al., 2013). However, Baddock et al., (2016) clearly demonstrate the detrimental effect that this has for dust source detection, because of highly variable AOD values over 10 x 10 pixels of 1 km each, upon which the data quality flags are based, proximal to dust sources. Filtering for QA = 3 resulted in a fivefold loss of data and far fewer source detections in their work. This study therefore follows the recommendation of Baddock et al., (2016) to include data of all quality levels (QA = 1 – 3) in the analysis. However, a sensitivity analysis is performed comparing results to those obtained by applying the QA = 3 filter (Sect. 3.1).



2.1.2 VIIRS

Information on aerosol type is obtained from Visible Infrared Imaging Radiometer Suite (VIIRS) Deep Blue aerosol products, which include an “aerosol type” classification. The classification is based on Deep Blue AOD and Ångström Exponent (α) retrievals, as well as a series of smoke tests, based on the spectral variation of measured reflectance. The algorithm is outlined in detail and verified against other satellite products/imagery and AERONET data by Hsu et al., (2019). In brief, retrievals are given the aerosol type classification “dust” when:

- AOD > 0.3,
- Smoke tests determine the retrieval not to be smoke,
- $\alpha < 0.5$.

The AOD > 0.3 threshold is applied to minimise the effect of variable surface reflectance in the visible wavelengths; retrievals not meeting this threshold are given the “background” aerosol type. Where the AOD threshold is met, smoke is ruled out, and $\alpha > 1$, the aerosol type is set to “non-smoke fine mode”, while those retrievals with an Ångström exponent between 0.5 and 1 are determined to be “mixed type” aerosols. It should be noted that retrievals performed over clouds or snow/ice surfaces are excluded from the product, as are retrievals performed over coastal pixels, to avoid subpixel mixed surface-type issues (since the surface type determines whether aerosol retrievals are performed using the Deep Blue algorithm (land) or Satellite Ocean Aerosol Retrieval (SOAR) algorithm).

VIIRS aerosol products used in this study are from the VIIRS sensor carried onboard the Suomi National Polar-orbiting Partnership (S-NPP) satellite, which has an equator crossing time (ascending) of 13:30 local time. Level 2 “AERDB_L2_VIIRS_SNPP” files were downloaded from the NASA Level-1 and Atmosphere Archive and Distribution System Distributed Active Archive Centre (doi.org/10.5067/VIIRS/AERDB_L2_VIIRS_SNPP.002, https://ladsweb.modaps.eosdis.nasa.gov/search/order/1/AERDB_L2_VIIRS_SNPP--5200, last access 11 January 2024). Version 2.0 data are used. As with MODIS data, only files that contained data in the region spanning (150° W – 50° W, 40° N – 85° N) were obtained, for the period spanning 2020-01-01 to 2022-12-31 (a three-year subset of the period covered by MODIS data used in this study). Likewise, data were regridded from their Level 2 swath files (native resolution 6 x 6 km²) to a regular 0.1° x 0.1° grid using nearest neighbour interpolation and mosaiced daily.

2.1.3 CALIOP

To obtain information on the vertical distribution of dust in the atmosphere – specifically to address the question of whether it is close to the surface and therefore indicative of local emission – aerosol type classification data from the Cloud-Aerosol Lidar with Orthogonal Polarization (CALIOP) are used. CALIOP products have been used in global aerosol source attribution studies (Prijiath et al., 2013), and the vertical information in retrievals has been exploited to detect dust emission at the surface in the Sahara Desert (Todd and Cavazos-Guerra, 2016). CALIOP is carried onboard the Cloud-Aerosol Lidar and Infrared Pathfinder Satellite Observations (CALIPSO) platform, which also has an equator crossing time of 13:30 local time



(ascending). CALIOP has been making vertical profile measurements of the Earth's atmosphere on a global scale since June 2006 and provides information on layer types for both clouds and aerosols (Winker et al., 2010). Its aerosol classification algorithm uses altitude, location, surface type, estimated particulate depolarization ratio, and integrated attenuated backscatter to identify the aerosol subtype (Kim et al., 2018; Omar et al., 2009); aerosol type "dust" corresponds to total attenuated backscatter > 0.075 and estimated depolarization ratio > 0.20.

For this study, the CALIPSO lidar Level 3 (L3) tropospheric aerosol profile "all sky" product, Version 4.20 is used ("CAL_LID_L3_Tropospheric_APro_AllSky-Standard-V4-20" files). Parameters included in the L3 files are derived from the Version 4 CALIOP Level 2 aerosol profile product; prior to aggregating, L2 data are quality screened to reduce the impact of e.g. layer detection or classification errors. Level 2 data are aggregated monthly on a grid of dimensions 5° longitude x 2.5° latitude x 60 metres vertical (for altitudes between -0.5 and 12 km) (Tackett et al., 2018). This level of aggregation compensates for the ~70 m footprint of the laser on the ground, produced every 333 m, during each orbit (compared to MODIS and VIIRS orbital swaths of 2330 km and 3060 km, respectively). This study uses the "aerosol type" parameter from the L3 files. The "aerosol type" parameter stores the monthly sums of the number of Level 2 samples ("sample" equals the number of measurement opportunities within the grid cell) having the CALIPSO aerosol types "clean marine", "dust", "polluted continental/smoke", "clean continental", "polluted dust", "elevated smoke", and "dusty marine" within each L3 latitude/longitude/altitude grid cell. Results presented here are based only on analysis of the "dust_nsamples" parameter, i.e., "polluted dust" and "dusty marine" are excluded. Data were downloaded from the NASA Earthdata portal (<https://search.earthdata.nasa.gov/search>, last access 11 January 2024), for the period spanning June 2006 – July 2020 inclusive. Only daytime files were used in the analysis.

2.2 Other datasets

Data from the Global Sand and Dust Storms Source Base Map ("G-SDS-SBM"; Vukovic, 2021; 2022) were used as an independent source of information to verify the dust emission potential of the satellite-identified dust hotspot areas. Specifically, the source intensity parameter was used, "SI", with values from 0 (no capacity to emit dust) to 1 (maximum capacity in cases of strong surface winds). G-SDS-SBM was developed by the United Nations Convention to Combat Desertification (UNCCD) in collaboration with the United Nations Environment Programme (UNEP) and World Meteorological Organization (WMO). It is derived using information on soil texture, bare land fraction, land cover, topsoil moisture, topography, and temperature. Values of SI represent the maximum potential of the surface to emit dust under favourable wind conditions between the years 2014-2018, and are available for the months of January, April, July, and October. SI data were regridded from their native 30 arcsec resolution (~0.008°) to 0.1° to match the MODIS data used in this study.

For verification of satellite-based results, data from the Aerosol Robotic Network (AERONET; Holben et al., 1998) are used. AERONET is a network of ground-based sun photometers that retrieve AOD and other atmospheric properties in the column above the instrument, based on solar extinction within the column. This study uses data from two stations in the broad study region of northern Canada: Kluane Lake (Yukon; 61.027°N, 138.410°W) and Resolute Bay (Nunavut; 74.705°N,



94.969°W; station locations are marked on Fig. 11). The Kluane Lake AERONET station was intentionally located proximal to a known dust source with the purpose of monitoring emission processes there (e.g. Huck et al., 2023), and has been providing data between April and November since 2018 (with varying temporal coverage by year). Resolute Bay is an Arctic monitoring site, providing data since 2004 (also with varying temporal coverage, also restricted to the months of April – November). Arctic aerosol studies have demonstrated that the site detects transported Asian dust during the spring and smoke aerosols during the summer, and that during summer months the area is also subject to wind-induced sea-salt-spray aerosol and potentially dust derived from more localised sources, based on the analysis of retrieved particle size distributions (Tomasi et al., 2015; AboEl-Fetouh et al., 2020).

Data from both stations were downloaded from the AERONET website (<https://aeronet.gsfc.nasa.gov/>, last access 11 January 2024), and this study analyses Level 1.5 AOD retrievals at 500 nm, processed by the Version 3 algorithm (Giles et al., 2019). Only retrievals performed within 11 am and 5 pm local time were considered for analysis (which roughly corresponds to a 6-hour window centred on the local overpass time of both satellites). The maximum retrieved AOD each day within this time window was retained for comparison to satellite-derived parameters.

Two additional datasets are used in this study. Firstly, wind fields discussed in Sect. 3.6 are taken from the European Centre for Medium-Range Weather Forecasts Reanalysis Version 5 (ERA-5), with a spatial resolution of 0.28125 degrees (~ 31 x 31 km) and 137 vertical levels (Hersbach et al., 2020; downloaded from the Copernicus Climate Data Store at <https://doi.org/10.24381/cds.143582cf>, last access 11 January 2024). Surface elevation data, also discussed in Sect. 3.6, are taken from the Global Multi-resolution Terrain Elevation Data 2010 dataset (GMTED2010; Danielson and Gesch, 2011, downloaded from <https://www.temis.nl/data/gmted2010/index.php>, last access 11 January 2024), at 0.0625° resolution.

3 Results and discussion

3.1 Inferences about dust over Canada from MODIS

Since DOD values will be elevated by both locally sourced and long-range transported dust, this study applies a threshold to DOD data to identify areas of persistently high dust loadings, which we assume to be largely undiluted and locally sourced dust. A high frequency of occurrence (FoO) of DOD above a specified threshold relative to the surrounding region is commonly used to indicate a local dust source region (e.g. Ginoux et al., 2012; Baddock et al., 2016). The FoO is simply the total number of observations in the time series that meet a given DOD threshold, expressed as either an absolute value or a relative value given the total number of observations in the time series. DOD threshold selection is somewhat subjective and depends upon the amount of long-range transported dust present in the atmosphere as well as the intensity of local emission. For example, Ginoux et al., (2012) apply a DOD threshold of 0.2 globally and experiment with higher and lower thresholds regionally; Baddock et al., (2016) applies a DOD threshold of 0.75 to pinpoint sources on a landscape-scale; and Pu et al., (2020) experiment with DOD thresholds ranging from 0.02 for less dusty regions to 0.5 for global dust hotspots such as North Africa. This study considers results using DOD thresholds of 0.5 and 1.



Maps of FoO $\text{DOD}_{\text{PG16}} > 0.5$ show several distinct features (Fig. 1a). There is a broad swath of FoO $> \sim 50$ in southern
 225 Alberta and Saskatchewan, extending across the border with the United States; this feature is associated with croplands, as
 discussed more fully in Section 3.2. The remaining northern features are of more interest to the high latitude dust question:
 most notably, numerous spatially discrete dust hotspots (FoO > 80) scattered across the islands (north of 70°N) that make up
 Canada's Arctic Archipelago. There are a few additional isolated FoO hotspots in continental northern Canada, for example
 in southern Northwest Territories close to the border with the Yukon Territory (MacKenzie Mountains, pink square in Fig.
 230 11), scattered across northern continental Nunavut, and at the border between northern Québec and Labrador (blue square in
 Fig. 11). Many of these hotspots correlate with 'barren land' classifications in the Canadian Land Cover Atlas. Notably, the
 greatest FoO values are in the Arctic Archipelago, as opposed to further south where dust is perhaps more commonly expected
 to be present.

The pattern in FoO $\text{DOD}_{\text{B16}} > 0.5$ is very similar to that in FoO DOD_{PG16} (Fig. 1b), with two exceptions. Firstly, FoO
 235 values are around 50% lower in the DOD_{B16} dataset; secondly, the northern FoO hotspots tend to be focussed east of 95°W in
 DOD_{B16} , whereas they are distributed more evenly across the full longitudinal extent of Canada's north in DOD_{PG16} . Lower
 FoO values may be expected in DOD_{B16} than DOD_{PG16} , because the former is only calculated for subsets of AOD retrievals
 that meet certain filtering criteria ($\alpha < 0.3$), whereas the latter is calculated for all AOD retrievals with no filtering applied (see
 DOD equation outlined in Sect. 2.1.1.)

240 MODIS AOD retrievals – upon which DOD is based – are more numerous at southern latitudes than in the far north (Fig.
 1 g-i) for reasons such as cloud and snow cover, and the lack of reflected visible radiation during northern winter (polar night
 of varying duration occurs north of 67°N) which is required to make AOD retrievals. To account for this, FoO $\text{DOD} > 0.5$
 values are normalised across the study domain by dividing the FoO count by the number of available DOD retrievals. The
 resulting maps for both DOD datasets (Fig. 1d-e) show that the relative frequency of $\text{DOD} > 0.5$ is far greater in the north than
 245 for the broad DOD hotspot across the Canada-US border, with values exceeding 50% in DOD_{PG16} and reaching 100% in
 DOD_{B16} (values closer to 100% are more likely in DOD_{B16} than DOD_{PG16} due to its threshold-based derivation, as mentioned
 above).

The same threshold of 0.5 is applied to the parent AOD data (Fig 1c, f). Overall FoO $\text{AOD} > 0.5$ values are greater across
 Canada, as expected, but the northern hotspots remain with FoO exceeding 200 at some locations. The greater values for FoO
 250 $\text{AOD} > 0.5$ across Canada, in regions where FoO DOD is close to zero, shows the value of using DOD data to highlight the
 contribution of dust particles to the retrieved AOD value, against the contribution from other aerosols which are present in the
 atmosphere.




Figure 1. Frequency of Occurrence (FoO) of **a)** $\text{DOD}_{\text{PG16}} > 0.5$, **b)** $\text{FoO DOD}_{\text{B16}} > 0.5$, and **c)** $\text{FoO AOD} > 0.5$. FoO as a percentage of the number of retrievals available for **d)** $\text{DOD}_{\text{PG16}} > 0.5$, **e)** $\text{DOD}_{\text{B16}} > 0.5$, and **f)** $\text{AOD} > 0.5$. Number of retrievals available for **g)** DOD_{PG16} , **h)** DOD_{B16} , and **i)** AOD . Note the different colour scales used. Grey shading corresponds to no data. All panels are based on MODIS data for 2003 – 2022.

If the DOD (or AOD) threshold is increased to 1, FoO values decrease (maximum values < 20) and the spatial extent of hotspots contracts (compare Fig. 2a and Fig. 2b for a comparison of FoO DOD_{PG16} with the thresholds of 0.5 and 1 respectively; note that maps for $\text{FoO DOD}_{\text{B16}} > 1$ and $\text{FoO AOD} > 1$ are included in Supp. Mat. 1). However, many spatially discrete hotspots in the north remain in attenuated form (compare Fig. 2d, e).

There are no DOD products north of 70°N between the months of October and April inclusive (Fig. 3, left column). This is to be expected due to a combination of a) polar night during parts of the winter preventing satellite AOD retrievals based on visible wavelengths and b) the fact that much of the surface is frozen, which inhibits the atmospheric entrainment of surface-



derived dust particles. For these reasons, DOD retrievals in the north predominantly occur during the months of June, July, and August (Fig. 3, right column), with May and September representing the shoulder seasons (Fig. 3, middle column).




Figure 2. **a)** Frequency of Occurrence (FoO) of $\text{DOD}_{\text{PG16}} > 0.5$, using data at all quality levels (i.e. not screened for “best estimate” quality flag). Note that this is a reproduction of Fig. 1a. **b)** FoO $\text{DOD}_{\text{PG16}} > 1$, using data at all quality levels. **c)** FoO $\text{DOD}_{\text{PG16}} > 0.5$, using data screened for “best estimate” quality flag. **d-f)** as a)-c) but zoomed to better highlight northern DOD hotspots. Grey shading corresponds to no data. All panels are from MODIS data for 2003 – 2022.

The results discussed in Figs. 1 and 2 thus far are based on AOD retrievals that are not filtered using the data quality flag that is distributed with the data. As discussed at the end of Sect. 2.1.1, this is based on the findings of Baddock et al., (2016), wherein the DOD based on the unfiltered AOD dataset was of greater value for dust source detection – a core aim of this paper – as opposed to DOD calculated using only the AOD retrievals that remain after filtering for the “best estimate” quality flag. The present analysis was replicated after applying the “best estimate” quality flag filter to the AOD data to test the effect of this choice. This resulted in a big reduction in FoO DOD (and AOD) > 0.5 values and a consequent contraction of the hotspot areas (Figs. 2c and 2a show a comparison of FoO $\text{DOD}_{\text{PG16}} > 0.5$ with and without the “best estimate” quality flag filter, respectively; note that equivalent maps for DOD_{B16} and AOD are included in Supp. Mat. 2), however many of the isolated hotspots in the north remain in attenuated form (compare Figs. 2d and 2f). The spatial patterns in these maps are similar to those of FoO $\text{DOD}_{\text{PG16}} > 1$ created using data unfiltered by the quality flag (Figs. 2b and 2e).

The dust hotspots inferred in this section from MODIS DOD data will be evaluated against the dust emission potential map in Sect. 3.4, examined over time in Sect. 3.5, and evaluated against AERONET stations in Sect. 3.6.





Figure 3. Seasonal cycle in DOD_{PG16} over Canada (MODIS data spans 2003 – 2022). **Top row)** FoO $DOD_{PG16} > 0.5$, by season. **Middle row)** FoO $DOD_{PG16} > 0.5$ as a percentage of the number of DOD_{PG16} retrievals, by season. **Bottom row)** number of DOD_{PG16} retrievals, by season. **Left column)** October – April inclusive (winter season). **Middle column)** May and September (shoulder season). **Right column)** June – August inclusive (summer season). Grey shading corresponds to no data.

3.2 Inferences about dust over Canada from VIIRS

The frequency with which VIIRS aerosol retrievals were classified as aerosol type “dust” (FoO AT_{DUST}) for the period 2020 – 2022 is shown in Fig. 4a. Greatest FoO AT_{DUST} values are seen north of $65^{\circ}N$ over land, with values exceeding 80 in some
 290 hot regions for the 3-year period. When represented as a percentage of the number of all aerosol type retrievals, dust is
 clearly the dominant retrieval type over much of the land area of northern Canada (Fig. 4d), accounting for $> 80\%$ of all aerosol
 type retrievals in some hotspots. However, it should be noted that some of these high percentage areas have quite low FoO
 $AT_{DUST} (< 10)$, and a low number of aerosol type retrievals overall, too. We note that AT_{DUST} is also being retrieved over
 water (Fig. 4a) with increasing frequency towards lower latitudes and a stark land-water contrast (both over open salt water



300 and inland fresh water like the Great Lakes). This FoO feature is attenuated in the relative representation of FoO in Fig. 4c, even though the land-water contrast remains; we do not interpret it as dust.



305 **Figure 4.** Frequency of Occurrence (FoO) of **a)** VIIRS aerosol type (AT) = “dust” (AT_{DUST}), **b)** VIIRS AT = “mixed” (AT_{MIXED}), and **c)** MODIS $DOD_{PG16} > 0.5$. FoO expressed as a percentage of the number of available retrievals for **d)** AT_{DUST} , **e)** AT_{MIXED} , and **f)** MODIS $DOD_{PG16} > 0.5$. Note the different colour scales used. Grey shading corresponds to no data. All panels are from data only for the period 2020 – 2022 inclusive.

310 Aside from dust, there are large areas with an aerosol type classification of “mixed” ($FoO\ AT_{MIXED}$) north of $\sim 80^\circ N$ on Ellesmere Island and in the surrounding region (Fig. 4b), accounting for a high proportion ($> 60\%$) of total aerosol type retrievals there (Fig. 4e). Mixed aerosol type is also widespread south of $70^\circ N$, reaching peak values in south-central Canada and extending across the border with the US, where it accounts for 20 – 30% of all aerosol type retrievals (with most of the other retrievals being classified either as “smoke”, “non-smoke fine mode”, or “background” aerosol type – see Supp. Mat. 3 for an analysis of additional aerosol types not considered in this study). Mixed aerosol type ($0.5 < \alpha < 1$) is indicative of smaller particles than the large dust aerosol type ($\alpha < 0.5$), but smaller clay fraction ($< 2\ \mu m$) dust particles in fact dominate the emitted *number* size distribution (e.g., Kok et al., 2011), and they are found both close to sources and also after long-range transport (e.g., Prospero et al., 1999; Zhao et al., 2022); therefore, we interpret VIIRS mixed aerosols as likely dust indicators, where they occur as discrete source areas (as opposed to a homogeneous background) and where they coincide with areas of high emission potential in the dust emission potential map (Sect. 3.4).



320 For direct comparison, FoO $\text{DOD}_{\text{PG16}} > 0.5$ is shown for the sub-period 2020 – 2022 (Fig. 4c, f). It is apparent that in the north, peak FoO AT_{DUST} is greater than peak FoO $\text{DOD}_{\text{PG16}} > 0.5$ and that the area of FoO AT_{DUST} is larger than it is for FoO $\text{DOD}_{\text{PG16}} > 0.5$. However, such differences are to be expected since these datasets are derived differently (as outlined in Sect. 2.1). Specifically, the FoO AT_{DUST} map will include dust that did not pass the $\text{DOD}_{\text{PG16}} > 0.5$ threshold (i.e., VIIRS AT_{DUST} requires $\text{AOD} > 0.3$ with $\alpha < 0.5$, while MODIS $\text{DOD}_{\text{PG16}} > 0.5$ requires, e.g., $\text{AOD} > 0.74$ if $\alpha = 0.5$ and $\text{AOD} > 0.51$ if $\alpha =$
325 0; Eq. 1). Nonetheless, there is agreement between the two datasets on numerous hotspot areas across northern Canada. Lastly, the large southern hotspot evident in FoO $\text{DOD}_{\text{PG16}} > 0.5$ (Fig. 4c) is likely fine dust emitted and transported in the large area sources of the Great Plains of Canada and the US; this is consistent with the spatial coincidence with a peak in FoO AT_{MIXED} in the Great Plains (Fig. 4b), where croplands are found, and where finer dust is expected to be mixed with anthropogenic aerosols. The dust hotspots inferred in this section from VIIRS ‘dust’ aerosol type data will be evaluated against the dust
330 emission potential map in Sect. 3.4.

3.3 Inferences about dust over Canada from CALIOP

To analyse the vertical distribution of atmospheric dust over northern Canada, ‘aerosol type’ classification data from CALIOP were used. As outlined in Sect. 2.1.3, this product shows the number of times an aerosol of type ‘dust’ (dust aerosol types ‘polluted dust’ and ‘dusty marine’ were excluded as they are not the focus of this study) is detected in a bin of 5° longitude x
335 5° latitude x 60 m vertically, per month. The monthly totals were summed for JJA 2006 – 2020, inclusive, and these period-sum values were then averaged for the latitude band $60^\circ\text{N} - 80^\circ\text{N}$. A longitude-height plot of the resulting data (Fig. 5) shows clear evidence of enhanced dust detection frequency at low altitudes for the longitude range between $125^\circ\text{W} - 75^\circ\text{W}$, in good agreement with the longitudinal range spanned by the dust hotspot regions identified in both the MODIS and VIIRS datasets previously discussed. That dust detection frequency values are lower at higher altitudes across this region, and immediately to
340 the east and west of the region, suggests that the low-altitude enhancement in dust detection is due to increases in dustiness within the region, implying surface sources of emission there. Also evident from this analysis are near-surface enhancements in dust detection frequency between $\sim 50^\circ\text{W}$ and 25°W , as well as between 25°W and 15°W ; these indicate emission from sources in Greenland and Iceland, respectively, which have been documented elsewhere (see, e.g., Meinander et al., 2022). The dust location information from CALIOP informs our analysis but is not suitable for direct comparison against the dust
345 emission potential map (Sect. 3.4) on account of the coarse spatial aggregation necessary to compensate for the 70-m laser footprint (Sect. 2.1.3).




Figure 5. Number of times ('count') an aerosol is classified as “dust” in CALIOP L3 aerosol type data for the period JJA 2006–2020, averaged for the latitude band 60 – 80° N. The blue line shows the corresponding median surface elevation within the latitude band.

3.4 Evaluation of satellite dust hotspot results using the dust emission potential map

We turn now to comparing our satellite results with known information on dust emission potentials, the latter under favourable wind conditions. In Fig. 6 we compare MODIS FoO $DOD_{PG16} > 0.5$ and VIIRS FoO AT=Dust to source intensity (SI) for July from the G-SDS-SBM dataset. SI for July was chosen based on the results of the seasonal analysis in Sect. 3.1, where all-season FoO was dominated by June-July-August FoO in absolute terms and in spatial extent, and the fact that SI values are low for other months provided in the G-SDS-SBM dataset (SI for other months is plotted in Supp. Mat. 4). The dust emission potential is high ($SI > 0.5$) in much of Canada's Arctic Archipelago (Fig. 6a), which is broadly consistent with MODIS FoO $DOD_{PG16} > 0.5$ (Fig. 1a) and VIIRS FoO AT=Dust (Fig. 4a). We quantified areas of SI overlap with both MODIS (Fig. 6b, red colour) and VIIRS (Fig. 6c, red colour) using threshold SI and FoO values as follows. Because SI is a relative measure of dust emission potential with values between 0 (no emission potential under any wind conditions) and 1 (maximum emission potential under sufficient wind conditions), we looked for areas where SI is greater than 0.5 while, simultaneously, MODIS FoO is greater than 10%. This MODIS threshold was chosen because we are testing for the FoO of strong dust emission events ($DOD > 0.5$), and it is reasonable to assume that such conditions will be met in less than 0.5 (50%) of observational cases.

Red colours indicate agreement between dust emission potential (SI) and satellite observations, while blue and green colours both indicate disagreement. Blue disagreement areas have high dust emission potential that is not matched with observations



of dust, while green disagreement areas have low dust emission potential where satellites nevertheless see high DOD (> 0.5) in at least 10% of observations. For consistency we report here the same 10% threshold in VIIRS FoO AT=Dust data, which leads to relatively prominent green areas (low emission potential but frequent VIIRS-observed dust) because, unlike MODIS FoO data, VIIRS FoO data considers only the presence of dust (VIIRS AT=Dust) and not the intensity of events (i.e., MODIS DOD > 0.5). While the thresholds can be manipulated to an extent, influencing the level of agreement and disagreement and thus the spatial colour patterns, the key point is that all three datasets have limitations, yet all three point to non-zero dust activity north of 60°N , in contrast to many model simulations of dust emission to date that have ignored these high latitude dust sources, albeit reasonably so, e.g., on account of low model resolution or lack of input information. The key dust emission areas shared by all three datasets (shared orange and red colours) are found in a broad “Canadian Arctic Dust Belt” between the horn-like Brodeur Peninsula of westernmost Baffin Island ($\sim 71^\circ\text{N}$, $\sim 85^\circ\text{W}$) and northwest of Prince Patrick Island ($\sim 79^\circ\text{N}$, $\sim 125^\circ\text{W}$). The same area also contains few blue colours (high emission potential but few observations of dust) but many green colours, where emission potential is low (< 0.5) but satellite observations show strong (MODIS) and/or frequent (VIIRS) dust events. The latter indicates areas where dust emission potential maps require re-examination and/or verification, i.e., also with new field measurements, challenging as they are.




Figure 6. a) Source intensity (SI) from the G-SDS-SBM dataset for July. b) Combined classification map based on SI and FoO MODIS DOD_{PG16} > 0.5 as a percentage of the number of DOD_{PG16} retrievals available; c) Combined classification map based on SI and FoO VIIRS AT_{DUST} as a percentage of the number of VIIRS Aerosol Type (AT) retrievals available. Grey shading indicates no data.

The notable inconsistencies in the Arctic Archipelago include eastern Baffin Island ($\sim 70^\circ\text{N}$, $\sim 75^\circ\text{W}$), which is characterised by widespread high SI (except in areas permanently covered by snow and ice) but sparse MODIS FoO DOD_{PG16} > 0.5 hotspots (blue colours in Fig. 6b) with more frequent low-intensity VIIRS FoO AT=Dust regions (red colours in Fig. 6c). The same is true for ice-free parts of Ellesmere Island and Axel Heiberg Island ($\sim 80^\circ\text{N}$, $\sim 85^\circ\text{W}$). One explanation for such



discrepancies, apart from thresholding choices, is that high SI values only show surfaces where the *potential* for dust emission is high; other conditions still need to be met for dust emission to actually occur, i.e. surface winds that are strong enough to tain sediment. Another factor is that we have filtered MODIS data for $\text{DOD} > 0.5$, which represents strong dust emission events, on par with Saharan dust in transport across the tropical Atlantic, whereas our VIIRS data shows all observations of pure (unpolluted) dust. On the other hand, it is plausible that the satellite datasets also do not capture the full extent of dustiness across the region, especially in the far north, due to frequent cloud coverage and  misclassification as clouds (e.g., Huck et al., 2023; Ranjbar et al., 2021).

There are also a few notable areas south of the Arctic where FoO $\text{DOD}_{\text{PG16}} > 0.5$ hotspots are not matched by high SI, e.g., green domains in Fig. 6b in the Northwest Territories (MacKenzie Mountains), close to the border with the Yukon Territory ($\sim 126^\circ\text{W}$, $\sim 63^\circ\text{N}$) and at the border of northern Labrador and Québec (65°W , 58°N), as marked in Fig. 11. Reasons for such disagreements are unclear, but we stress again that the G-SDS-SBM SI mapping results are sensitive to the quality of the input data (soil texture, soil moisture, soil temperature, and bare land fraction), which all have limitations. In examining the visible imagery (Appendix), average wind speeds and geological features of these two discrepancy areas available from the Global Wind Atlas (Davis et al., 2023), there is strong indication that both, although geologically distinct, are likely unclassified dust hotspot source areas. The MacKenzie Mountains location (Fig. A1), between the Keele River and Nahanni National Park, shows intense weathering and erosion in the area of the Redstone Plateau, leading to some colluvial and landslide features, and abundant alluvial fans, terraces, and flood plains (Boydell and Rutter, 1980; Duk-Rodkin and Huntley, 2018). The latter are comprised of sand, silt and minor gravel and represent sediments from modern to post-glacial drainage. The Labrador / Québec location (Fig. A2), east-southeast of Ulittaniujalik National Park, is the exposed and weathered Torngat plateau. The area was recently deglaciated (Occhietti et al., 2004), leaving behind regionally atypical glacial sand, silty sand, and gravel (Brouard et al., 2020), with scarce vegetation and a thin soil profile that promotes dust mobilization. Clearly, further investigation is needed, including with field measurements.

Lastly, the broad swath of only occasionally high FoO $\text{DOD}_{\text{PG16}} > 0.5$ in the Great Plains of southern Canada and USA (connected to croplands and discussed in Sect. 3.2, Figs. 4b and 4c) is also largely not matched by high SI; this is reasonable because this extensive pattern of occasional dustiness disappears in a percentage-of-observations representation (Figs. 4e and 4f), i.e., the satellite data is showing infrequent events of dust transported from confined sources. SI hotspots in this region (unlike the FoO signature, found only in northwestern US and not southwestern Canada) are much more spatially restricted than the FoO signature and comprise a mixture of croplands (Washington, northern Oregon, Idaho) and arid regions (southern Oregon, eastern Montana, Wyoming), but they do not correspond to either intense MODIS $\text{DOD}_{\text{PG16}} > 0.5$ events (Fig. 6b, blue colours) or VIIRS AT=Dust events (Fig. 6c, blue colours). Instead, they are associated with VIIRS AT=Mixed (Figs. 4b and 4e), which, again, makes sense given that agricultural dust emissions are mixed with anthropogenic aerosols and transported over a broad region.



Overall, the reasonable agreement between satellite-derived dust hotspots and areas with high SI values in the G-SDS-SBM dataset supports the use of the satellite datasets in dust emission modelling development in the Canadian Arctic and in other high-latitude regions. This section also suggested regions where the SI dataset may benefit from further development.

3.5 Temporal change in intense dustiness frequency over Northern Canada in the 20-year MODIS record

The 20-year record of MODIS data enables an assessment of whether the frequency of dustiness over Northern Canada has changed over time. Time series of annual means of FoO $\text{DOD}_{\text{PG16}} > 0.5$, FoO $\text{DOD}_{\text{B16}} > 0.5$, and FoO $\text{AOD} > 0.5$, averaged for the region ($65^{\circ}\text{N} - 85^{\circ}\text{N}$, $125^{\circ}\text{W} - 70^{\circ}\text{W}$), are shown in Fig. 7. This averaging area subsumes the broad “Canadian Arctic Dust Belt” identified in Sect. 3.4. Note that this figure shows values of mean FoO rather than mean optical depth, and that left and right y axes with different scales are employed to better show the (always) larger mean FoO $\text{AOD} > 0.5$. There is clear interannual variability in each of the time series; however, the greatest annual mean FoO $\text{DOD}_{\text{PG16}} > 0.5$ values all occur in the second half of the period, with 2017, 2019, 2014, 2015 (in descending order of mean FoO) appearing to be peak dust emission years. The same is true with FoO $\text{DOD}_{\text{B16}} > 0.5$, although the pattern is much less pronounced, with no peak for 2014. By contrast, the annual mean FoO $\text{AOD} > 0.5$ time series is qualitatively different, with peaks and troughs relatively evenly distributed across the study period. This suggests that any increase in AOD across the 20-year period may be driven by coarser, dust-sized particles, which the DOD datasets emphasise. As noted in Sect. 1, an increase in high latitude dust emission is an expected consequence of climate change in the Arctic leading to glacial retreat, decreased snow cover duration, and permafrost thaw, all making exposed dust more available (Bullard, 2013; Meinander et al., 2022). Our result is consistent with such processes occurring in the Arctic (Gulev et al., 2021).




Figure 7. Time series of annual mean FoO of MODIS optical depth > 0.5 , averaged for the region ($65^{\circ}\text{N} - 85^{\circ}\text{N}$, $125^{\circ}\text{W} - 70^{\circ}\text{W}$), for the years 2003 – 2022 inclusive. MODIS optical depth datasets underlying the FoO values plotted are: $\text{DOD}_{\text{PG16}} > 0.5$, $\text{DOD}_{\text{B16}} > 0.5$, and $\text{AOD} > 0.5$.

3.6 Evaluation of MODIS AOD using AERONET



AOD data from MODIS are evaluated against AOD data from AERONET stations at Kluane Lake (AOD_{KL}) and Resolute Bay (AOD_{RB}). The station locations are marked on Fig. 11. The Kluane Lake AERONET station was intentionally located proximal to a known dust source with the purpose of monitoring emission processes there (Huck et al., 2023). Curiously, the area is not identified as a strong potential dust source area by our satellite data analysis, which is on the one hand not very surprising given that Huck et al. (2023) document large discrepancies between remotely sensed dust (both by AERONET and by MODIS-MAIAC) and *in situ* remote camera images of dust at Kluane Lake. On the other hand, their “dust event days” (DED’s) have modal AOD values of 0.5 with a high tail extending out to an AOD of 3; in the context of such thick dust plumes, one may reasonably expect that in the analysis of 20 years of MODIS data in our study (Collection 6.1 at 10-km resolution) we would detect a stronger signal in FoO – yet we do not. However, the Kluane Lake region and the nearby mountainous regions, are largely without satellite data due to the extremely variable and highest topography in North America (Mt. Logan in Canada stands at 6,000 m and Denali in Alaska to the west stands at 6,200 m), which presents difficult surface retrieval conditions (as shown in detail later in Fig. 10). Unlike Kluane Lake, the Resolute Bay AERONET station is situated adjacent to strong potential dust emission hotspots identified in our study on Cornwallis Island (as shown in detail later in Fig. 10); however, to the best of our knowledge, this is coincidental, i.e., the station location was not selected with the explicit purpose of monitoring



dust aerosol. In this section, satellite data retrieved within both a $1^\circ \times 1^\circ$ and $0.5^\circ \times 0.5^\circ$ box centred on each AERONET station were considered. From MODIS, the maximum AOD values within the search box were retained, while from VIIRS the presence or absence of each aerosol classification type (AT) was recorded. For AERONET, only retrievals performed within 11 am and 5 pm local time were considered for analysis and the maximum retrieved AOD each day within this time window was retained for the comparison.



AOD comparison results are first considered using MODIS data detected within the $1^\circ \times 1^\circ$ search box centred on the respective AERONET station. At Kluane Lake, six years of AOD_{KL} data correlates strongly with MODIS AOD ($R = 0.87$ in Fig. 8a). These AOD correlation results are at odds with the work of Huck et al. (2023) at Kluane Lake who find weak mean AOD (unlike our maximum AOD) correlations on “dust event days” in 2018 and 2019 ($R = 0.11$ and $R = 0.36$), which improve on non-DED’s in 2018 and 2019 ($R = 0.35$ and $R = 0.96$). They classify DED’s with threshold AOD and α parameters similar to our work (but from their AERONET data) and they use the mean MODIS-MAIAC AOD product at 470 nm with the mean AERONET AOD product at 500 nm. Given their *in situ* remote camera imagery verifying the presence of dust, they reasonably attribute the lack of correlation on “dust event days” to specific dust-cloud detection problems in AERONET and the lack of dust testing at high latitudes in the MODIS-MAIAC dataset (rather than to the less important 470 nm vs. 500 nm wavelength mismatch between MODIS and AERONET AOD). Their sampling strategy is different and covers just 70 km² with 1-km MODIS-MAIAC pixels as opposed to our $1^\circ \times 1^\circ$ box based on 10-km resolution L2 MODIS data. While it is not possible to compare the results directly, it is nevertheless interesting to note both the range of choices and results in this apparently ‘similar’ type of comparison carried out in our respective studies. When three years of VIIRS AT data are present




Figure 8. Maximum AERONET AOD (500 nm) between 11 am and 5 pm local time at Kluane Lake (blue open circles) and Resolute Bay (red crosses) vs. maximum MODIS AOD **a)** from a $1^\circ \times 1^\circ$ box centred on each station; **b)** from a $0.5^\circ \times 0.5^\circ$ box centred on each station. Numbers in blue (red) above each panel correspond to the Pearson’s correlation coefficient, R ,



between Kluane Lake (Resolute Bay) AERONET AOD and the MODIS dataset in that panel. The number of matched
485 maximum AOD days for each comparison is given in parentheses in the legends.




Figure 9. Percentage of days with VIIRS Aerosol Type (AT) data that have at least one AT retrieval of the type indicated on the x-axis within the $1^\circ \times 1^\circ$ and $0.5^\circ \times 0.5^\circ$ boxes centred on AERONET stations at Kluane Lake (KL; dark and light blue, respectively) and Resolute Bay (RB; dark and light red, respectively). “bg” = background, “nsfm” = non-smoke fine mode, “has” = high-altitude smoke. The number of days with VIIRS AT data in the $1^\circ \times 1^\circ$ ($0.5^\circ \times 0.5^\circ$) box is 1278 (852) at
490 Kluane Lake and 429 (228) at Resolute Bay, both for the period 2020-2022.




Figure 10. **a,c,e)** Winds at 925 hPa, 850 hPa, and 700 hPa above the region surrounding the Kluane Lake AERONET station (blue circle). Winds are the mean of ERA-5 reanalysis data for Jun. – Aug. 2004 – 2023. Underlying shading corresponds to FoO DOD_{PG16} > 0.5 for the period 2003 – 2022 (grey shading corresponds to no data). **g)** Surface elevation surrounding Kluane Lake station (red circle) at 0.0625 degrees resolution from the Global Multi-resolution Terrain Elevation Data 2010 (GMTED2010) dataset. **b,d,f,h)** Like a,c,e,g) except for Resolute Bay AERONET station.



over the $1^\circ \times 1^\circ$ region surrounding the Kluane Lake AERONET station, AT_{DUST} is detected on 18% of days while AT_{MIXED} and $AT_{BACKGROUND}$ are the two modal classifications, both detected on 27% of days (Fig. 9). Unlike at Kluane Lake, 18 years of Resolute Bay AOD_{RB} has only a weak correlation ($R = 0.32$) with MODIS AOD for the $1^\circ \times 1^\circ$ search area (Fig. 8a). As Fig. 9 shows, VIIRS AT_{DUST} is detected on 32% of days here, with $AT_{BACKGROUND}$ being detected more frequently (44% of days) and AT_{MIXED} less frequently (12% of days).




Figure 11. The location of Kluane Lake AERONET station (pink circle) and Resolute Bay AERONET station (blue circle), together with new dust source areas in the McKenzie Mountains (pink square) and Quebec/Labrador (blue square). Underlying shading corresponds to $FoO\ DOD_{PG16} > 0.5$ for the period 2003 – 2022 (grey shading corresponds to no data).

A question arises about the poor correlation at Resolute Bay between AERONET AOD_{RB} and MODIS AOD datasets given the proximity to our potential hotspot regions and the more frequent VIIRS AT_{DUST} presence there. One reason could be the deliberate positioning of an AERONET station to monitor dust at a known emission location (like Kluane Lake), which was not the case for Resolute Bay (notwithstanding the AOD disagreement at Kluane Lake found by Huck et al, 2023 using the MODIS-MAIAC product). Is it possible that coarse dust in the $1^\circ \times 1^\circ$ search area around Resolute Bay is simply not passing over the AERONET station, given that coarse dust does not generally travel far? The map of average regional winds above both stations (Fig. 10) shows that Kluane Lake is downwind of dust emission regions in the foothills of the mountains (where satellite data is increasingly lacking on account of the variable topography and surface properties) while Resolute Bay is upwind of the dust emission hotspot (identified with abundant satellite data). The elevated topography to the east of Resolute Bay may itself obscure any dust suspended in that direction from the AERONET instrument's field of view for low solar zenith



angles in the morning. As may be expected in this situation, the correlation between MODIS AOD and AOD_{RB} increases from
520 0.32 to 0.58 as the MODIS search area is decreased from 1° x 1° to 0.5° x 0.5°, while the MODIS AOD correlation with AOD_{KL}
hardly changes (0.87 vs. 0.90) under the same decrease (Fig. 8b). Since the AERONET station at Kluane Lake is already next
to and downwind of the dust, decreasing the search area is not helpful to the correlation. Interestingly, this same decrease in
search area size causes the AT_{DUST} classification frequency at Resolute Bay to decrease from 32% for 1° x 1° to 16% for 0.5°
x 0.5°, while the AT_{BACKGROUND} classification frequency increases from 44% to 64% (Fig. 9); there is no appreciable change
525 at Kluane Lake. This reduction supports the argument that dust in the broader region surrounding Resolute Bay is not within
detection range of the AERONET station and highlights the challenge of detecting even nearby dust sources given the station's
position and prevailing winds.

4 Summary and conclusions

High latitude mineral dust is receiving growing research interest, with evidence that its impacts on numerous components of
530 the Arctic environment are greater than for dust transported into the region from lower latitudes. Understanding these processes
– and how they change in the future – depends upon precise knowledge of where dust emission occurs at high latitudes, yet
field measurements are exceedingly difficult (and therefore scarce) and satellite-based retrievals of suspended dust suffer from
known high-latitude biases (polar night, increased cloudiness, bright surfaces). Using the Frequency of Occurrence (FoO) of
above-average Dust Optical Depth (DOD > 0.5) from twenty years (2002-2022) of high-resolution MODIS observations
535 derived for this study (0.1° x 0.1°), we present quantitative evidence that dust sources are widespread across the islands of the
Canadian Arctic Archipelago. Additionally, qualitative supporting evidence is presented from aerosol type ‘dust’ classification
in the VIIRS ($\alpha < 0.5$ with AOD > 0.3) and CALIPSO (total attenuated backscatter > 0.075 with depolarization ratio > 0.20)
satellite data products.

The locations identified as dust hotspots (DOD > 0.5 with FoO > 10%) in the “Canadian Arctic Dust Belt”
540 (overwhelmingly based on retrievals from June to August) correspond to surfaces with high potential for dust emission during
the summer season in the G-SDS-SBM dataset (Source Intensity, SI > 0.5). However, there are significant areas of
disagreement in this region, where MODIS data shows dust (FoO > 10%) but the G-SDS-SBM dataset shows lower emission
potential (SI < 0.5); there are fewer areas where MODIS data does not show dust (FoO < 10%), but emission potential is higher
(SI > 0.5), notably on the southern portion of the Brodeur Peninsula of Baffin Island. Additionally, two considerable areas of
545 disagreement emerge at lower latitudes in mainland Canada (MacKenzie Mountains and the border between northern Québec
and Labrador), which we show to be potential dust emitting regions based on their geological features. The main limitation of
this comparison is the use of threshold FoO and SI values, which can be manipulated to an extent, influencing the level of
agreement and disagreement. However, the key point is that both datasets point to non-zero dust activity north of 60°N, in
contrast to many model simulations of dust emission to date that have ignored these high latitude dust sources, albeit reasonably
550 so, e.g., on account of low model resolution or lack of input information.



When spatially averaged across the larger dust producing region ($65^{\circ}\text{N} - 85^{\circ}\text{N}$, $125^{\circ}\text{W} - 70^{\circ}\text{W}$), annual mean time series of FoO of MODIS DOD > 0.5 between 2003 and 2022 suggest an increase in the frequency of dustiness in the latter half of the period, which is consistent with our understanding that high latitude dust emissions are likely increasing in a warming climate.

555 The MODIS AOD product used in our study compares well ($R = 0.87$) to AOD from the AERONET station at Kluane Lake in Yukon Territory. While Kluane Lake is a known dust source area receiving focussed research attention, it is not a prominent dust emission hotspot in our study, likely because satellite retrievals are less frequent in the mountainous terrain. Nevertheless, the AERONET station was positioned downstream of known dust emission sources leading to good agreement with fewer data points. Conversely, the comparison is poor ($R = 0.32$) between our MODIS AOD product and AOD from the
560 AERONET station at Resolute Bay, Nunavut, with approximately double the data points. This AERONET station is situated close to satellite-derived dust hotspots in our study, however, its location is not well suited to detecting dust plumes emitted in the region because the emission hotspots are downstream of the station (on average). Somewhat unsurprisingly, station location is critical for verifying satellite-derived dust aerosols; this becomes even more important when the dust product specifically targets coarse-mode particles that do not travel far from their sources.

565 The results presented here confirm the dust emission potential in Northern Canada with satellite observations. The regions identified as dust hotspots are candidates for further research into processes involved in driving dust emission at high latitudes, and also provide an observational basis for evaluating and refining numerical models that are used to simulate the dust cycle and the Earth System impacts of dust.



570 Appendix




Figure A1. Visible imagery of the potential dust source area in the MacKenzie Mountains (marked by a pink square in Fig. 11). Centre of 9.0 km² box is at 63°N, 126°W with a mean wind speed of 8.13 m/s (predominantly westerly). Topography varies from 900 m to 2300 m in full region shown. Obtained from the Global Wind Atlas version 3.3, a free, web-based application developed, owned and operated by the Technical University of Denmark (DTU). The Global Wind Atlas version 3.3 is released in partnership with the World Bank Group, utilizing data provided by Vortex, using funding provided by the Energy Sector Management Assistance Program (ESMAP). For additional information: <https://globalwindatlas.info>.




Figure A2. Visible imagery of the potential dust source area at the Québec and Labrador border (marked by a blue square in Fig. 11). Centre of 9.0 km² box is at 58.4°N, 65.1°W with a mean wind speed of 11.4 m/s (predominantly westerly). Topography varies from 200 m to 900 m in full region shown. Obtained from the Global Wind Atlas version 3.3, a free, web-based application developed, owned and operated by the Technical University of Denmark (DTU). The Global Wind Atlas version 3.3 is released in partnership with the World Bank Group, utilizing data provided by Vortex, using funding provided by the Energy Sector Management Assistance Program (ESMAP). For additional information: <https://globalwindatlas.info>.



Data availability. The high-resolution MODIS AOD and DOD data products underlying this study are available upon request.

590 *Author contributions.* AW and IA both conceived of and designed the study. IA performed data processing and analysis. Both authors examined and interpreted the results. IA prepared the manuscript while AW edited it.

Competing interests. The authors declare that they have no conflict of interest.

595 *Acknowledgements.* The authors thank Dr. Deanne Van Rooyen and Dr. Jacob Hanley for helpful discussions about the geological features and dust-emission potential of the MacKenzie Mountains and Québec / Labrador. Any errors of discussion presented here are our own.

Financial support. This research has been supported by the Canadian Space Agency (grant no. 21SUASMDAC), the Canadian
600 National Sciences and Engineering Research Council through the Discovery Grants Programme (grant no. RGPIN-2014-03888, RGPIN-2022-05225), and Saint Mary's University.

References

AboEl-Fetouh, Y., O'Neill, N. T., Ranjbar, K., Hesarak, S., Abboud, I., and Sobolewski, P. S.: Climatological-scale analysis of intensive and semi-intensive aerosol parameters derived from AERONET retrievals over the Arctic, *Journal of Geophysical Research: Atmospheres*, 125, e2019JD031569, <https://doi.org/10.1029/2019JD031569>, 2020.
605

Anderson, T. L., Wu, Y. H., Chu, D. A., Schmid, B., Redemann, J., and Dubovik, O.: Testing the MODIS satellite retrieval of aerosol fine-mode fraction, *J. Geophys. Res.-Atmos.*, 110, D18204, <https://doi.org/10.1029/2005jd005978>, 2005.

610 Baddock, M. C., Ginoux, P., Bullard, J. E., and Gill, T. E.: Do MODIS-defined dust sources have a geomorphological signature?, *Geophys. Res. Lett.*, 43, 2606–2613, <https://doi.org/10.1002/2015GL067327>, 2016.

Boydell, A. N. and Rutter, N. W.: Surficial Geology and Geomorphology, Root River, District of Mackenzie, *Geological Survey of Canada, Preliminary Map*, 12-1979, 1980.

615

Brouard, E., Roy, M., Dubé-Loubert, H., Lamarche, O., and Hébert, S.: Carte des dépôts de surface de la province de Québec: rapport sur les méthodes et les données: MB 2020-10, 1:2 500 000, *Ministère de l'Énergie et Ressources naturelles, Québec*, 2020.



- 620 Bullard, J. E.: Contemporary glacigenic inputs to the dust cycle, *Earth Surf. Proc. Land*, 38,71–89,
<https://doi.org/10.1002/esp.3315>, 2013.
- Bullard, J. E., Baddock, M., Bradwell, T., Crusius, J., Darlington, E., Gaiero, D., Gasso, S., Gisladdottir, G., Hodgkins, R.,
McCulloch, R.: High-latitude dust in the Earth system, *Reviews of Geophysics*, 54, 447–485, 2016.
- 625 Danielson, J.J., and Gesch, D.B.: Global multi-resolution terrain elevation data 2010 (GMTED2010), U.S. Geological Survey
Open-File Report 2011-1073, 26 p., <http://pubs.usgs.gov/of/2011/1073/pdf/of2011-1073.pdf>, 2011.
- Dansie, A. P., Thomas, D.S.G., Wiggs, G.F.S., Baddock, M.C., and Ashpole, I.: Plumes and blooms – Locally-sourced Fe-rich
630 aeolian mineral dust drives phytoplankton growth off southwest Africa, *Science of The Total Environment*, 829,
<https://doi.org/10.1016/j.scitotenv.2022.154562>, 2022.
- Davis, N. N., Badger, J., Hahmann, A. N., Hansen, B. O., Mortensen, N. G., Kelly, M., Larsén, X. G., Olsen, B. T., Floors, R.
Lizcano, G., Casso, P., Lacave, O., Bosch, A., Bauwens, I., Knight, O. J., van Loon, A. P., Fox, R., Parvanyan, T., Krohn
635 Hansen, S. B., Heathfield, D., Onninen, M., Drummond, R.: The Global Wind Atlas: A high-resolution dataset of
climatologies and associated web-based application, *Bulletin of the American Meteorological Society*, 104 (8), E1507-E1525,
<https://doi.org/10.1175/BAMS-D-21-0075.1>, 2023.
- Dubovik, O., Holben, B., Eck, T. F., Smirnov, A., Kaufman, Y. J., King, M. D., Tanré, D. and Slutsker, I.: Variability of
640 absorption and optical properties of key aerosol types observed in worldwide locations, *J. Atmos. Sci.*, 59,590–608, 2002.
- Duk-Rodkin, A. and Huntley, D. H.: Surficial geology, Dahadinni River, Northwest Territories, NTS 95-N southwest.
prelim., *Geological Survey of Canada, Canadian Geoscience Map*, 297, 2018.
- 645 Eck, T. F., Holben, B. N., Reid, J. S., Dubovik, O., Smirnov, A., O'Neill, N. T., Slutsker, I., and Kinne, S.: Wavelength
dependence of the optical depth of biomass burning, urban, and desert dust aerosols, *J. Geophys. Res.*, 104(D24), 31,333–
31,349, 1999.
- Giles, D. M., Sinyuk, A., Sorokin, M. G., Schafer, J. S., Smirnov, A., Slutsker, I., Eck, T. F., Holben, B. N., Lewis, J. R.,
650 Campbell, J. R., Welton, E. J., Korkin, S. V., and Lyapustin, A. I.: Advancements in the Aerosol Robotic Network
(AERONET) Version 3 database – automated near-real-time quality control algorithm with improved cloud screening for Sun
photometer aerosol optical depth (AOD) measurements, *Atmos. Meas. Tech.*, 12, 169–209, <https://doi.org/10.5194/amt-12-169-2019>, 2019.



655 Ginoux, P., Prospero, J. M., Gill, T. E., Hsu, N. C., and Zhao, M.: Global-scale attribution of anthropogenic and natural dust sources and their emission rates based on MODIS Deep Blue aerosol products, *Rev. Geophys.*, *50*, RG3005, <https://doi.org/10.1029/2012RG000388>, 2012.

660 Groot Zwaafink, C. D., Grythe, H., Skov, H., and Stohl, A.: Substantial contribution of northern high-latitude sources to mineral dust in the Arctic, *Journal of Geophysical Research: Atmospheres*, *121*(22), 13,678–13,697. <https://doi.org/10.1002/2016JD025482>, 2016.

Gulev, S.K., Thorne, P.W., Ahn, J., Dentener, F.J., Domingues, C.M., Gerland, S., Gong, D., Kaufman, D.S., Nnamchi, H.C., Quaas, J., Rivera, J.A., Sathyendranath, S., Smith, S.L., Trewin, B., von Schuckmann, K., and Vose, R. S.: Changing State of the Climate System. In *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* [Masson-Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 287–422, 2021.

670 Hersbach H., Bell B., Berrisford P., et al.: The ERA5 global reanalysis, *Q. J. R. Meteorol. Soc.*, *146*, 1999–2049, <https://doi.org/10.1002/qj.3803>, 2020.

Holben, B. N., Eck, T. F., Slutsker, I., Tanré, D., Buis, J. P., Setzer, A., Vermote, E., Reagan, J. A., Kaufman, Y. J., Nakajima, T., Lavenue, F., Jankowiak, I., and Smirnov, A.: AERONET – A federated instrument network and data archive for aerosol characterization, *Remote Sens. Environ.*, *66*, 1–16, [https://doi.org/10.1016/S0034-4257\(98\)00031-5](https://doi.org/10.1016/S0034-4257(98)00031-5), 1998.

Hsu, N. C., Jeon, M.-J., Bettenhausen, C., Sayer, A. M., Hansell, R., Seftor, C. S., Huang, J., and Tsay, S.-C.: Enhanced Deep Blue aerosol retrieval algorithm: The second generation, *J. Geophys. Res. Atmos.*, *118*, 9296–9315, <https://doi.org/10.1002/jgrd.50712>, 2013.

680 Hsu, N. C., Lee, J., Sayer, A. M., Kim, W., Bettenhausen, C., and Tsay, S.-C.: VIIRS Deep Blue aerosol products over land: Extending the EOS long-term aerosol data records, *Journal of Geophysical Research: Atmospheres*, *124*, 4026–4053, <https://doi.org/10.1029/2018JD029688>, 2019.

685



- Huck, R., Bryant, R.G., and King, J.: The (mis)identification of high-latitude dust events using remote sensing methods in Yukon, Canada: a sub-daily variability analysis, *Atmos. Chem. Phys.*, (23), 6299-6318, <https://doi.org/10.5194/acp-23-6299-2023>, 2023.
- 690 Kim, M.-H., Omar, A. H., Tackett, J. L., Vaughan, M. A., Winker, D. M., Trepte, C. R., Hu, Y., Liu, Z., Poole, L. R., Pitts, M. C., Kar, J., and Magill, B. E.: The CALIPSO version 4 automated aerosol classification and lidar ratio selection algorithm, *Atmos. Meas. Tech.*, 11, 6107–6135, <https://doi.org/10.5194/amt-11-6107-2018>, 2018.
- Kok, J. F., Ward, D. S., Mahowald, N. M., & Evan, A. T.: Global and regional importance of the direct dust-climate feedback, 695 *Nature Communications*, 9(1), <https://doi.org/10.1038/s41467-017-02620-y>, 2018.
- Kok, J. F., Ridley, D. A., Zhou, Q., Miller, R. L., Zhao, C., Heald, C. L., Ward, D. S., Albani, S., and Haustein, K.: Smaller desert dust cooling effect estimated from analysis of dust size and abundance, *Nat. Geosci.*, 10, <https://doi.org/10.1038/Ngeo2912>, 2017.
- 700 Kok, J. F.: A scaling theory for the size distribution of emitted dust aerosols suggests climate models underestimate the size of the global dust cycle, *Proceedings of the National Academy of Sciences*, 108(3), 1016-1021, <https://doi.org/10.1073/pnas.1014798108>, 2011.
- 705 Kylling A., Groot Zwaftink, C. D., & Stohl, A.: Mineral dust instantaneous radiative forcing in the Arctic, *Geophysical Research Letters*, 45, 4290–4298, <https://doi.org/10.1029/2018GL077346>, 2018.
- Meinander, O., Dagsson-Waldhauserova, P., Amosov, P., Aseyeva, E., Atkins, C., Baklanov, A., et al.: Newly identified climatically and environmentally significant high-latitude dust sources, *Atmos. Chem. Phys.*, 22, 11889–11930, 710 <https://doi.org/10.5194/acp-22-11889-2022>, 2022.
- Nogueira, J., Evangelista, H., Valeriano, C.d.M. et al.: Dust arriving in the Amazon basin over the past 7,500 years came from diverse sources, *Commun Earth Environ* 2, 5, <https://doi.org/10.1038/s43247-020-00071-w>, 2021.
- 715 Occhietti, S., Govare, É., Klassen, R., Parent, M. and Vincent, J.S.: Late Wisconsinan-Early holocene deglaciation of Québec-Labrador, in *Developments in Quaternary Sciences* (Vol. 2, pp. 243-273), 2004.



- Omar, A. H., Winker, D. M., Vaughan, M. A., Hu, Y., Trepte, C. R., Ferrare, R. A., Lee, K. P., Hostetler, C. A., Kittaka, C., Rogers, R. R., and Kuehn, R. E.: The CALIPSO Automated Aerosol Classification and Lidar Ratio Selection Algorithm, *J. Atmos. Ocean. Tech.*, 26, 1994–2014, <https://doi.org/10.1175/2009JTECHA1231.1>, 2009.
- Platnick, S., King, M., Wind, B., and Ridgway, W.: MODIS Atmosphere L2 Joint Atmosphere Product. NASA MODIS Adaptive Processing System, Goddard Space Flight Center, doi:10.5067/MODIS/MODATML2.061; <https://doi.org/10.5067/MODIS/MYDATML2.061>, 2017.
- Prijith, S.S., Aloysius, M., and Mohan, M.: Global aerosol source/sink map, *Atmospheric Environment*, 80, 533–539, <https://doi.org/10.1016/j.atmosenv.2013.08.038>, 2013.
- Prospero, J.M., Ginoux, P., Torres, O., Nicholson, S.E., Gill, T. E.: Environmental characterization of global sources of atmospheric soil dust identified with the NIMBUS 7 Total Ozone Mapping Spectrometer (TOMS) absorbing aerosol product, *Reviews of Geophysics*, 40(1), 1002, <https://doi.org/10.1029/2000RG000095>, 2002.
- Prospero, J. M.: Long-range transport of mineral dust in the global atmosphere: Impact of African dust on the environment of the southeastern United States, *Proceedings of the National Academy of Sciences*, 96(7), 3396–3403, <https://doi.org/10.1073/pnas.96.7.3396>, 1999.
- Pu, B. and Ginoux, P.: The impact of the Pacific Decadal Oscillation on springtime dust activity in Syria, *Atmos. Chem. Phys.*, 16, 13431–13448, <https://doi.org/10.5194/acp-16-13431-2016>, 2016.
- Pu, B., Ginoux, P., Guo, H., Hsu, C. N., Kimball, J., Marticorena, B., Malyshev, S., Naik, V., O'Neill, N. T., García-Pando, C. P., Paireau, J., Prospero, J. M., Shevliakova, E., and Zhao, M.: Retrieving the global distribution of the threshold of wind erosion from satellite data and implementing it into the Geophysical Fluid Dynamics Laboratory land-atmosphere model (GFDL AM4.0/LM4.0), *Atmos. Chem. Phys.*, 20(1), 55–81, <https://doi.org/10.5194/acp-20-55-2020>, 2020.
- Ranjbar, K., O'Neill, N. T., Ivanescu, L., King, J., and Hayes, P. L.: Remote sensing of a high- Arctic, local dust event over Lake Hazen (Ellesmere Island, Nunavut, Canada), *Atmos. Environ.*, 246, 118102, <https://doi.org/10.1016/j.atmosenv.2020.118102>, 2021.
- Sarangi, C., Qian, Y., Rittger, K., Leung, L. R., Chand, D., Bormann, K. J., & Painter, T. H.: Dust dominates high-altitude snow darkening and melt over high-mountain Asia, *Nature Climate Change*, <https://doi.org/10.1038/s41558-020-00909-3>, 2020.



- 755 Sayer, A. M., Hsu, N. C., Bettenhausen, C., Jeong, M.-J., and Meister, G.: Effect of MODIS Terra radiometric calibration improvements on Collection 6 Deep Blue aerosol products: Validation and Terra/Aqua consistency, *J. Geophys. Res. Atmos.*, *120*, 12, 157–12, 174, <https://doi.org/10.1002/2015JD023878>, 2015.
- Tegen, I.: Modeling the mineral dust aerosol cycle in the climate system, *Quat. Sci. Rev.*, *22*, 1821–1834, 2003.
- 760 Tackett, J. L., Winker, D. M., Getzewich, B. J., Vaughan, M. A., Young, S. A., and Kar, J.: CALIPSO lidar level 3 aerosol profile product: version 3 algorithm design, *Atmos. Meas. Tech.*, *11*, 4129–4152, <https://doi.org/10.5194/amt-11-4129-2018>, 2018.
- 765 Tobo, Y., Adachi, K., DeMott, P. J., Hill, T. C. J., Hamilton, D. S., Mahowald, N. M., ... Koike, M.: Glacially sourced dust as a potentially significant source of ice nucleating particles, *Nature Geoscience*, *12*(4), 253–258. <https://doi.org/10.1038/s41561-019-0314-x>, 2019.
- Todd, M. C., and Cavazos-Guerra, C.: Dust aerosol emission over the Sahara during summertime from Cloud-Aerosol Lidar with Orthogonal Polarization (CALIOP) observations, *Atmospheric Environment*, *128*, 147–157, <https://doi.org/10.1016/j.atmosenv.2015.12.037>, 2016.
- 770 Tomasi, C., Kokhanovsky, A. A., Lupi, A., Ritter, C., Smirnov, A., O'Neill, N. T., et al.: Aerosol remote sensing in polar regions, *Earth-Science Reviews*, *140*, 108–157. <https://doi.org/10.1016/j.earscirev.2014.11.001>, 2015.
- 775 Vukovic, A.: Global high-resolution dust source map, InDust webinar, https://cost-indust.eu/sites/default/files/2021-04/vukovic_indust_webinar_april2021.pdf (last access: 11 January 2024), 21 April 2021.
- Vukovic, A.: Sand and dust storm source mapping, in United Nations Convention to Combat Desertification (UNCCD), (2022), *Sand and Dust Storms Compendium: Information and Guidance on Assessing and Addressing the Risks*, Bonn, Germany, 2022.
- 780 Washington, R. M., Todd, M., Middleton, N. J., and Goudie, A. S.: Dust-storm areas determined by the Total Ozone Monitoring Spectrometer and surface observations, *Ann. Assoc. Am. Geogr.*, *93*, 297–313, <https://doi.org/10.1111/1467-8306.9302003>, 2003.



- 785 Winker, D. M., Pelon, J., Coakley Jr, J. A., Ackerman, S. A., Charlson, R. J., Colarco, P. R., Flamant, P., Fu, Q., Hoff, R. M., Kittaka, C., Kubar, T. L., Le Treut, H., McCormick, M. P., Mégie, G., Poole, L., Powell, K., Trepte, C., Vaughan, M. A., and Wielicki, B. A.: The CALIPSO mission: A global 3D view of aerosols and clouds, *Bull. Amer. Meteor. Soc.*, *91*, 1211–1229, <https://doi.org/10.1175/2010BAMS3009.1>, 2010.
- 790 Wiacek, A., Peter, T. and Lohmann, U.: The potential influence of Asian and African mineral dust on ice, mixed-phase and liquid water clouds, *Atm. Chem. Phys.*, *10*, 8649–8667, <https://doi.org/10.5194/acp-10-8649-2010>, 2010.
- Yang, A., Tan, Q., Rajapakshe, C., and Chin, M.: Global premature mortality by dust and pollution PM_{2.5} estimated from aerosol reanalysis of the modern-era retrospective analysis for research and applications, version 2, *Front. Environ. Sci.*, *10*, <https://doi.org/10.3389/fenvs.2022.975755>, 2022.
- 795 Zhao, X., Huang, K., Fu, J. S., and Abdullaev, S. F.: Long-range transport of Asian dust to the Arctic: identification of transport pathways, evolution of aerosol optical properties, and impact assessment on surface albedo changes, *Atmos. Chem. Phys.*, *22*, 10389–10407, <https://doi.org/10.5194/acp-22-10389-2022>, 2022.

Response to CC 3

Just a quick comment. The authors might want to be aware that the CALIPSO algorithm can sometimes misclassify the diamond dust (small ice crystals in clear sky conditions) as mineral dust. This leads to an overestimation of CALIPSO "dust" in the Arctic when $RH_i > 100\%$ (e.g., see <https://acp.copernicus.org/articles/22/12269/2022/> and <https://agupubs.onlinelibrary.wiley.com/doi/10.1002/2017JD027530>).

Best regards,

-Lauren Zamora

We thank Dr. Zamora for her helpful comment on our manuscript and the two references, which we have added to our discussion of the CALIPSO data in Section 3.3. While we were in general aware of dust/ice misclassification issues for 'diamond dust' (and blowing snow), we were reluctant to enter this discussion because it is complex, and our CALIPSO data spans JJA only, when we presumed that issues of dust/ice misclassification were at a minimum. With the information in these references, we have more confidence that some of our CALIPSO aerosol *type* 'dust' data is indeed likely to be dust, and we have added a brief discussion in Section 3.3 to explain when/where this is likely, and where it is not, based on an analysis of mean atmospheric temperature fields and ice nucleation mechanisms.

This point was also raised by RC 2 and was addressed there, too, with additional details.

Response to RC 1

In this study, the authors attempted to identify HLD sources across the Canadian Arctic Archipelago and adjacent areas by using FoO of DOD from 20-yr high-resolution MODIS retrievals as well as VIIRS aerosol type product, CALIOP products, and observations from two AERONET sites. The authors also compared the spatial distribution of FoO with G-SDS-SBM dataset and showed an overall agreement. There are also significant areas of disagreement between FoO from MODIS and VIIRS and between satellite retrievals and G-SDS-SBM dataset. Overall, I think this study provides insights for identifying HLD sources and can benefit future modeling and observational studies. I have a few comments for the authors to consider.

We thank the Anonymous Reviewer for their work on our paper.

General Comments:

1. I wonder if MODIS products using the Dark Target algorithm (DT alone or combined DT/DB) can change the FoO of DOD and contribute to the uncertainties in identifying potential HLD sources, especially for coastal regions having both land and ocean.

The FoO of DOD will likely change a bit when using two satellite retrievals based on the same data, but optimized for land vs. water. How much change can be expected in FoO of DOD (or the always larger AOD) is a complex question, although we have some idea from the work of Sayer et al. (2014, JGR), which generally shows high correlations in JJA between DB and DT AOD (Figure 5), as well as small AOD differences (Figure 4), with the major caveat that Arctic ($> 60^\circ \text{N}$) satellite coverage is sparse, even in JJA. The abstract of this paper states: *“Neither algorithm consistently outperforms the other, although in many cases the retrieved AOD and the level of its agreement with AERONET are very similar. In many regions the DB, DT, and merged data sets are all suitable for quantitative applications, bearing in mind that they cannot be considered independent, while in other cases one algorithm does consistently outperform the other. Usage recommendations and caveats are thus somewhat complicated and regionally dependent.”* No recommendations are made for the Arctic because none of the 111 AERONET stations used in the validation of the DB/DT/merged products were in the Arctic (Figure 12). As far as using the merged DB/DT product over land, Figure 1 of Sayer et al. (2014) shows that some regions of the Arctic (over land) use only DB, some use only DT, and some use a merging of the two. Coastal regions having both land and ocean in data pixels are indeed tricky and would require very special treatment at the level of L2 granules. Our MODIS data has a 10-km resolution, making products less certain roughly 5-10 km inland. We have added this limitation to our discussion. We also added statements in our Conclusions about the need for future work related to the use of different satellite retrieval products for dust identification in the high Arctic.

2. I would suggest providing more details or high-level formulas to better explain how SI from G-SDS-SBM is calculated which may be helpful to explain the differences between the datasets and satellite retrievals. It seems to be mostly dependent on soil properties. It is not clear to me if SI depends on surface wind.

We agree that further explanation of how SI from G-SDS-SBM is calculated would be helpful. We added clarifications to the Introduction (1) and Other Datasets (2.2) of the manuscript. Briefly:

- friction velocity of 1 m/s assumed globally
- 5-year climatology (2014-2018) of month-specific assimilated meteorological fields at 0.25 deg resolution (ECMWF) were used to estimate minimum soil moisture values and maximum soil temperature values, gridded globally. This represents a region's maximum capacity to emit under favorable meteorological conditions (i.e., dry warm soil and high wind).

SI being 1 indicates maximum capacity in cases of strong surface winds. However, the authors state that the derivation of SI does not need surface wind. What about the surface wind/friction velocity in the large green domain in Figure 6c compared to that in the red domain in Figure 6b and 6c.

- An SI of 1 indicates a soil's potential to emit at or above the 99th percentile level of erodible soils, globally (Vukovic (2021)). The derivation of SI did not include surface wind input fields, rather, a globally constant high friction velocity of 1 m/s was assumed at all times in the emission parameterization. As such, SI represents the potential of a soil to emit under high wind conditions.
- Geographical wind speed differences do not appear to explain discrepancies in Fig6b or 6c; we included a wind climatology in the Supplement.

I also wonder how native (~0.008°) SI data were regridded to 0.1°.

- Regridding was performed using bilinear interpolation in the CDO package.

Specific comments:

Line 9, please check if multiplication sign is used instead of letter x. Please check other places as well.

Fixed throughout.

Lines 32-33, there are some more recent studies addressing this issue. I would suggest adding them. For example, Shi et al. (2022). Relative importance of high-latitude local and long-range-transported dust for Arctic ice-nucleating particles and impacts on Arctic mixed-phase clouds. *Atmospheric Chemistry and Physics*, 22, 2909-2935.

Yes, this is a great paper that we have also found since our submission to ACPD. We have referenced it and expanded the discussion around lines 32-33 to include the contributions from Shi et al., including the primarily lower tropospheric contribution of HLD. This is also relevant to the discussion of CALIOP-derived vertical dust distributions and is now mentioned in Section 3.3.

Line 235, why $\text{FoO DOD}_{\text{B16}} > 0.5$ is much lower than $\text{FoO DOD}_{\text{PG16}} > 0.5$ in the east of 95°W high-latitudes?

In the paper we stated that $\text{FoO DOD}_{\text{B16}} > 0.5$ is in general 50% lower than $\text{FoO DOD}_{\text{PG16}} > 0.5$ and we explained that this was due to the threshold nature of DOD_{B16} , which requires $\alpha < 0.3$ (or no DOD is produced). Second, we stated that DOD_{B16} hot spots are focused east of the 95°W longitude, and did not state reasons why. We clarified these points and uncertainty in the manuscript.

Line 238, for DOD_{PG16} , is it calculated when $w < 1$? Does that mean it is filtered as well?

Yes, but in practice this filtering eliminates small numbers of points because ω is > 0.99 only 0.5% of the time.

Figure 3, I would suggest changing the color bar for the three bottom figures. It is just showing black over most places and gives less info.

The color bars were adjusted in the final manuscript version.

Figure 5. I feel it gives limited information compared to lat-lon plot. How about showing similar lat-lon spatial distribution of FoO of CALIOP DOD using the algorithms from literature (e.g., Yu et al., 2015)?

It is true that Figure 5 gives more limited horizontal information compared to the lat-lon plots, however, it gives some vertical information about the location of the dust in active satellite observations. We chose this representation also because a lat-lon plot of CALIOP FoO of aerosol type “dust” was very sparsely populated, even in a 20-year mean, at the horizontal resolution of the lat-lon figures (DOD would also be sparse, if it was calculated). As discussed in Section 2.1.3, this is because of the extremely narrow footprint of the CALIOP laser.

Thank you for the reference to Yu et al. (2015). These authors carried out a quantitative analysis of transported dust flux in the Saharan outflow region, which has the most abundant and continuous plumes of dust in the world, making their quantitative work more justified than in our case of using occasional DOD extremes to point to potential dust source regions.

Lines 362-363, this sentence is not clear to me. Could you explain a bit more and rephrase it?

“This 10% MODIS threshold, which does not match the SI threshold of 0.5 (50%), was chosen because we are testing for the FoO of strong dust emission events ($\text{DOD} > 0.5$), and it is reasonable to assume that such conditions will be met in far less than ~~0.5 (50%)~~ of observational cases, i.e., 10%.”

Figure 6, I agree with the key point the authors tried to make. However, I would suggest at least trying to use different thresholds for FoO MODIS DOD and FoO VIIRS AT_{DUST} . There are significant differences between FoO MODIS DOD and FoO VIIRS.

We experimented with different thresholds before submission and there was not one that made all three datasets agree, so we decided to use the same thresholds for all datasets, for consistency, as stated in the discussion. We revised the wording to make it clear that we had experimented with thresholds.

Figure 7, I'm a bit confused about this figure. What's the unit for FoO shown here. Is it the number of occurrence or the percentage as to total observations? It seems that blue and green lines correspond to the left y-axis while the brown line corresponds to the right y-axis. For right y-axis, is it for number of occurrence or the percentage?

Indeed, this figure was confusing (also to CC 1 and CC 2) and we have revised it in several ways. It used to show the spatial mean number of occurrence, but the legend labels did not match the axis labels (there was a descriptive note in the text, but it was hard to spot). We have matched the axis and legend labels and normalized the FoO values, eliminating the need for two y axes. The normalizing factors are included in the caption.

Line 448, I would suggest showing the locations, as well as other selections of region such as the region in Figure 7, in Figure 8 or in the section 2.

Line 448 refers to Figure 11, which was created as we struggled with making the fewest number of figures while not showing Results (new dust source locations) in Methods (AERONET station locations). We have added the spatial averaging domain of Figure 7 to Figure 11.

For the conclusion part, I would suggest the authors give a high-level summary of some potential HLD sources in the order of confidence (i.e., agreements between MODIS, VIIRS, and G-SDS-SBM).

Agreed. We have expanded the summarizing statements about potential HLD sources in our Conclusions, referring to various dataset agreements.

References

Yu, H., Chin, M., Bian, H., Yuan, T., Prospero, J. M., Omar, A. H., Remer, L. A., Winker, D. M., Yang, Y., Zhang, Y., and Zhang, Z.: Quantification of trans-Atlantic dust transport from seven-year (2007–2013) record of CALIPSO lidar measurements, *Remote Sens. Environ.*, 159, 232–249, <https://doi.org/10.1016/j.rse.2014.12.010>, 2015.

Response to RC 2

The manuscript titled “Satellite-based evidence of dust emission over Northern Canada” by Ashpole and Wiacek attempted to use multiple satellite products including MODIS, VIIRS, and CALIOP to detect the frequency of occurrence of dust events for mapping possible dust sources. I think the methods to detect dust sources are generally adequate and useful for identifying dust sources. Different from another reviewer, I personally think using the Deep Blue over the land make sense since the authors are looking for dust sources from the land and not from the water bodies. I think this paper is helpful for constraining Canadian dust sources using a suite of satellite retrievals, and the study is helpful for the dust community. I have some general comments below, and I would suggest accepting this manuscript after a major revision.

We thank the Anonymous Reviewer for their work on our paper.

Major comments:

Line **224**: I would imagine that using 0.2 or an even smaller DOD threshold for FoO is needed for HLD sources like Canada. Using 0.5 as a threshold may miss out a lot of small local dust sources.

We experimented with various thresholds briefly, like other investigators that we referenced for this method, and added more to this discussion to Section 3.1. We settled on a threshold of 0.5 because it corresponds to strong dust emission events and it has DOD greater than typical values of thin cirrus cloud optical depth (COD), which ranges between 0.03 and 0.3 (Pierce et al., 2010). In future work this balance between finding weaker dust sources while avoiding thin cirrus contamination can be explored more fully; in this work, we have added a series of figures to the supplementary material that depict FoO DOD sensitivity to the threshold, i.e., 0.1, 0.2, 0.3, 0.4, 0.5, as well as the FoO of extreme DODs ($\text{DOD} > 2$ and $\text{DOD} > 3$).

The difference between DOD_{PG16} and DOD_{B16} are not clear to the reader. The authors need to give more details on the differences between them and why the occurrence of $\text{DOD}_{\text{B16}} > 0.5$ are generally lower than that of $\text{DOD}_{\text{PG16}} > 0.5$.

We revised and improved text relating to the above, also in response to RC 1.

Line **317**: I am not convinced that VIIRS mixed aerosols are mostly dust in Canada. They mentioned Kok (2011) but Kok showed that most emitted dust are coarse dust, not fine dust. They should stick with VIIRS dust (instead of VIIRS mixed aerosols) when interpreting possible dust sources.

We revised the text to note this limitation of VIIRS AT_MIXED more clearly. In Figure 6, where we compare SI, MODIS and VIIRS, only VIIRS_dust is used to help interpret possible dust sources.

Line 335: The authors used aggregated CALIOP aerosol vertical profiles with a coarse resolution of $5^\circ \times 2^\circ$. A finer resolution might be needed for detecting smaller and local dust events.

While this is a valid point, the sampling of the narrow laser (70-m footprint) makes the CALIOP dataset spatially sparse, even as it provides unprecedented ‘curtain’ views of aerosols along the orbital track. We revised the discussion of CALIOP data products’ horizontal resolution in Section 2.1.3 and also the discussion of the limitations of the CALIOP-based results in Section 3.3. This point was also raised by RC 1 and addressed in our response there.

Line 336: How did the authors exclude the diamond dust (ice crystal that are mistaken as dust) mistaken by CALIOP? There could be quite some precipitation in the summertime Arctic.

This comment was brought up also by CC 3, who pointed us to literature on the likelihood of occurrence of diamond dust as “60% of the time in winter, but hardly ever occurs in summer” (Zamora et al., 2022, ACP). We added this to our discussion of CALIOP results and noted where diamond dust may be occurring still (over Greenland), based on a consideration of mean atmospheric temperature fields and ice nucleation mechanisms.

Figure 5: It looks like to me that CALIOP dust aloft be transported from elsewhere (e.g., the lower latitudes). How do you separate local sources from regional transport?

We cannot differentiate local from transported dust based on the data; we can only make some inferences that low altitude dust, within 1-2 km of the surface is more likely to be locally sourced, especially in the dust emitting season that we sample (JJA). Dust emission and transport to North America from Asia often involves warm conveyor belt uplift mechanisms and is unlikely to be found in these low regions (e.g., Wiacek et al., 2010). We revised and expanded wording relating to the discussion of dust aloft (> 2.5 km) in the text concerning Figure 5.

Line 351: The authors used two satellite instruments (MODIS and VIIRS) to generate two FoO maps. It is not clear to the reader which one is better for use. The authors should explain more on the pros and cons of each of them and how people can combine them to better constrain the Canadian dust sources.

We revised the manuscript to stress that we originally added VIIRS and CALIPSO products to ‘verify / evaluate’ our main (semi-quantitative) results in a *qualitative* sense. We converted the quantitative MODIS AOD product to DOD, then ran the threshold analysis on DOD values to tally extreme dust events over 20 years. We only use the qualitative VIIRS and CALIOP aerosol ‘type’ products to corroborate the spatial locations of potential dust sources determined using the MODIS data.

To speak to the main point above, we agree that there are underlying VIIRS and CALIOP quantitative AOD and DOD data products that one could combine. The *selection* of the optimal sensor and retrieval algorithm for HLD detection is one complex matter, but not as complex as *combining* these products consistently, e.g., does one use MODIS AOD from the retrieval algorithms Deep Blue, Dark Target, or MAIAC? Or does one use VIIRS AOD? Often this question is settled pragmatically, depending on which time period one is interested in. CALIPSO is yet a different sensor (active lidar), providing vertical aerosol information and AOD.

This is the first study to use the Deep Blue MODIS data in the Arctic to detect potential dust sources in a systematic, long-term and spatially extensive way. Future work is needed to repeat and expand this work with other sensors and retrieval products. We have added this to our Concluding discussion.

Line 433: I would say that the climate change impacts on FoO is not clear from the time series in Fig. 7. I wonder what drives the interannual variability. Are there correlations between the wind / temperature / snow cover time series and the FoO time series?

We agree that this temporal signal is not very clear in Figure 7. We investigated the correlations between the FoO of DOD and AOD greater than 0.5 and wind, soil temperature, soil moisture and snow cover (from 20 years of ERA 5 fields) and included these results in the supplementary material. There was a significant ($p < 0.05$) and negative ($R = -0.5$) correlation only with FoO AOD (not FoO DOD) and snow cover.