

Influence of Various Criteria on Identifying the Springtime Tropospheric Ozone Depletion Events (ODEs) at Utqiagvik, Arctic

Xiaochun Zhu¹, Le Cao¹, Xin Yang², Simeng Li³, Jiandong Wang¹, and Tianliang Zhao¹

¹State Key Laboratory of Climate System Prediction and Risk Management, Nanjing University of Information Science and Technology, Nanjing, 210044, China

²British Antarctic Survey, Natural Environment Research Council, Cambridge, UK

³Institute of Environmental Sciences, Universiteit Leiden, Leiden, 2333 CA, the Netherlands

Correspondence: L. Cao
(le.cao@nuist.edu.cn)

Abstract. Tropospheric ozone depletion events (ODEs) occurring in the Arctic spring are a unique photochemical phenomenon in which the boundary layer ozone drops rapidly to near-zero levels. However, the criterion for identifying ODEs remains inconsistent among different studies, which may influence conclusions regarding the characteristics of ODEs. To address this issue, in this study, we applied various criteria used in previous studies to identify springtime ODEs at Utqiagvik, Arctic (the BRW Station), based on observational data spanning 23 years (2000-2022), and investigated the influences of implementing different criteria. We compared three types of criteria: traditional methods (fixed thresholds), variability-based methods (considering mean value and standard deviation), and machine learning methods (Isolation Forest), and found that criteria using fixed thresholds (e.g., 10 ppbv) and relative thresholds based on monthly averaged ozone levels (0.42 times the monthly average) are more suitable for capturing ODEs at BRW compared to other criteria. Results applying these appropriate criteria all reveal a significant decline in ODE occurrence frequency over the investigated 23 years, particularly in April, suggesting potential links to climate change and Arctic sea ice melting. However, implementing relative thresholds or more stringent fixed thresholds (5 ppbv and 4 ppbv) instead of the 10 ppbv threshold would display a more significant decline in the number of ODE hours across these 23 years. Further investigation of meteorological conditions indicates that ODEs at BRW are more prevalent under northerly and northeasterly winds with moderate wind speeds ($3\text{--}6\text{ m s}^{-1}$), at lower temperatures, and higher pressures, while severe ODEs are more associated with lower wind speeds and temperatures below 256 K. This research highlights the importance of selecting appropriate criteria to accurately identify ODEs and contributes to a better understanding of the complex processes driving the Arctic ODEs.

1 Introduction

The Arctic has been described as an important “window” on the global environment, as changes in the Arctic serve as a precursor for the global changes anticipated as the Earth’s temperature rises (Thoman et al., 2023). Among all the changes in the Arctic environment, the variation in the ozone concentration during spring seasons has attracted considerable attention from the scientific community. In the stratosphere, the depletion of ozone in the Arctic spring, which is caused due to the existence of

polar stratospheric clouds and halogen compounds (Crutzen and Arnold, 1986; Cairo and Colavitto, 2020; Seinfeld and Pandis, 2006; Akimoto, 2016), allows for increased UV radiation that can lead to higher rates of skin cancer, cataracts, and weakened immune systems in humans (Umar and Tasduq, 2022), while also affecting marine and terrestrial ecosystems, potentially altering weather patterns and contributing to climate change (Liu et al., 2022).


In contrast to the stratospheric ozone depletion, in the lower troposphere, a unique phenomenon, namely ozone depletion events (ODEs), has also been frequently observed in the near-surface layer of the Arctic since the 1980s (Oltmans, 1981; Barrie et al., 1988; Bottenheim et al., 1986; Anlauf et al., 1995; Bottenheim et al., 2002; Shupe et al., 2022). This phenomenon is associated with a photochemical process that can activate halogen ions (i.e., Br^- , Cl^- , I^-) from substrates such as ice/snow packs (Lehrer et al., 2004; Simpson et al., 2007b; Abbatt et al., 2012; Pratt et al., 2013; Custard et al., 2017) and sea-salt aerosols (Michalowski et al., 2000; Yang et al., 2010, 2019, 2020; Thomas et al., 2011, 2012; Huang et al., 2020) into reactive halogens that can deplete ozone. Consequently, the ozone in the polar boundary layer frequently falls from background levels (30-40 ppbv) to less than 10 ppbv or even near-zero values within a few days or even hours in the springtime of the Arctic.

The occurrence of ODEs is related to a complex photochemical process as follows (Simpson et al., 2007b),



In reaction cycle (R1), X denotes halogen species (i.e., Br, Cl and I). When the sun rises in the Arctic spring, halogen containing compounds (i.e., X_2) in the atmosphere are photodissociated into halogen atoms (i.e., X). These halogen atoms subsequently react rapidly with ozone, producing halogen monoxide XO, as depicted in reaction (R1). XO then participates in self-reactions that regenerate X atoms, thereby depleting ozone without consuming halogens.

However, in reaction cycle (R1), the total amount of halogens remains constant, which is inconsistent with measurements at Arctic coastal stations in which a substantial increase in halogen concentrations is observed during ODEs (Hausmann and Platt, 1994; Zhao et al., 2016). Furthermore, previous researches have demonstrated that ozone cannot be depleted in such a short time if only gas-phase reactions occur (Lehrer et al., 2004). Thus, another reaction cycle including heterogeneous reactions was proposed to explain the rapid ozone decline and the large amount of reactive halogens in the atmosphere during ODEs (Fan and Jacob, 1992; McConnell et al., 1992; Platt and Lehrer, 1997; Tang and McConnell, 1996; Wennberg, 1999). Using bromine as a representation of halogen species, this proposed reaction cycle can be outlined as follows:



In reaction cycle (R2), hypobromous acid (HOBr) can activate bromides (Br^-) from substrates such as snowpacks on the sea ice, and sea-salt aerosols, leading to the conversion of bromides into reactive halogen species such as Br_2 , which in turn undergo photodissociation and consume ozone. Consequently, ozone is continuously depleted and the total bromine amount in the atmosphere is explosively elevated during ODEs. Thus, this process is referred to as the “bromine explosion mechanism” (Platt and Lehrer, 1997; Wennberg, 1999).

The occurrence of ODEs is predominantly confined to the spring season due to the unique combination of meteorological and chemical conditions (Lehrer et al., 2004). First, the sunlight during spring is essential for photochemical reactions to take place, which is crucial for converting inert halogen ions into reactive halogens. Second, the strong temperature inversion that forms in the spring effectively isolates the boundary layer air from the free troposphere, preventing the downward mixing of ozone-rich air from aloft, allowing the reactive halogens to efficiently deplete the local ozone. Additionally, the snowpack above the sea ice in springtime acts as a significant source of halogen ions, as the brine layer on the sea ice is enriched with halogens that can be released into the atmosphere through photochemical processes. These factors collectively create the ideal conditions for ODEs to occur in the spring, while such conditions are not simultaneously met in other seasons.

ODEs can exert profound influence on both the Arctic environment and the global ecosystems. First, they can alter the radiation balance in the Arctic by reducing the infrared radiations absorbed by the atmosphere (Lacis et al., 1990). Moreover, during ODEs, the oxidative capacity of the Arctic atmosphere is dominated by the enhanced bromine compounds. This shift in the oxidative capacity facilitates the oxidation of elemental mercury ($\text{Hg}(0)$), followed by deposition of active mercury ($\text{Hg}(\text{II})$), a pollutant that is highly toxic to humans. The increased deposition of active mercury would ultimately jeopardize the human health in mid-latitude regions through snow melting and oceanic circulations (Ariya et al., 2004; Pratt et al., 2013; Dastoor et al., 2022; Basu et al., 2022).

Since the discovery of ODEs, our understanding of this phenomenon has been improved, such as the role of sea ice formation (Jones et al., 2011; Lehrer et al., 2004; Simpson et al., 2007a; Abbatt et al., 2012; Peterson et al., 2019), the linkage to climate variability (Koo et al., 2014), the influence of snowpack photochemical emissions (Pratt et al., 2013; Custard et al., 2017; Herrmann et al., 2021, 2022) and the Arctic blowing snow (Chen et al., 2022; Yang et al., 2010, 2019, 2020; Huang et al., 2020), as well as the connection to the total column ozone (Cao et al., 2022). However, to date, the criterion for identifying ODEs remains unclear. In previous studies, the occurrence of ODEs was mostly recognized by the volume mixing ratio of the surface ozone, when the surface ozone decreases to a level below a fixed threshold. However, the threshold is inconsistent across different studies, ranging from 20 ppbv down to 4 ppbv. For example, Tarasick and Bottenheim (2002) and Koo et al. (2012) used a threshold of 10 ppbv, while Bottenheim et al. (2009), Frieß et al. (2011) and Jacobi et al. (2010) utilized a 5 ppbv. Meanwhile, Piot and Von Glasow (2008) and Ridley et al. (2003) identified ODEs using a more stringent 4 ppbv threshold. In contrast to these studies employing a fixed threshold, Cao et al. (2022) used a criterion depending on the mean value and the standard deviation of the surface ozone across various months and years to identify ODEs. This criterion was based on the method of Bian et al. (2018) which was originally used to indicate uncommon variations in the surface ozone in polar regions. In that study, Cao et al. (2022) applied this criterion to pick out ODE hours from surface ozone measurements at the Halley

Station in Antarctica for the spring months from 2007 to 2013, and the results demonstrated that the criterion is effective in identifying ODEs from the springtime surface ozone measurements at the Halley Station.

85 Because the identification criterion for ODEs may influence conclusions regarding the characteristics of ODEs, such as the interannual variability of ODEs and the relationship between meteorological conditions and the occurrence frequency of ODEs, in this study, we employed various criteria to identify ODEs from 23 years of spring observational data detected at Utqiagvik in Alaska, and explored the impacts of using different screening criteria on the results. We then selected appropriate criteria to investigate the correlation between meteorological parameters (e.g., wind speed, wind direction, temperature, and pressure) and the occurrence of ODEs, and compared the results to assess the impact of different criteria on the conclusions.

2 Observational Data and Screening Criteria

We first used different criteria to screen out ODE hours from observational data for the springtime of 23 years (2000-2022). Then, based on a comparison of the screening results, we investigated the properties of these ODE screening criteria.

2.1 Observational Data

95 Surface ozone mixing ratio and meteorological parameters (pressure, 2-m temperature, 10-m wind speed and direction) at Utqiagvik (the BRW Station, see Fig. 1 for the geographic location) were taken from the National Oceanic and Atmospheric Administration/Oceanic and Atmospheric Research/Global Monitoring Laboratory (NOAA/OAR/GML) baseline observatory (<https://gml.noaa.gov/aftp/data/>) (McClure-Begley et al., 2024; Crocker, 2024), which are freely provided to the public and scientific community. We adopted the data for the years 2000-2022 with a 1-hr time resolution, and only focused on the three months of spring (March, April, and May) in the present study.

2.2 Criteria for Identifying ODEs

Three types of criteria (traditional methods, variability-based methods, machine learning methods) were used to screen out ODE hours from the measurements, which are listed in Table 1 and described in detail below.

2.2.1 Traditional Methods

105 This kind of methods define ODEs according to the mixing ratios of ozone. The first criterion to be tested is similar to that used by Halfacre et al. (2014), in which ODEs are defined as time periods when ozone mixing ratio falls below 10 ppbv. Moreover, the situation with an ozone value lower than 10 ppbv should last for longer than one hour (i.e., $[O_3] < 10$ ppbv at two consecutive time points). Thus, this criterion can be described as follows:

$$([O_3]_i < 10) \cap ([O_3]_{i+1} < 10), \quad (1)$$

110 in which $[O_3]_i$ is the ozone mixing ratio at the i -th time point. This criterion is referred to as TM1 in the following context (see Table 1). In previous studies (Bottenheim et al., 2009; Frieß et al., 2011; Jacobi et al., 2010; Piot and Von Glasow, 2008; Ridley



Figure 1. Location of the BRW Station in the Arctic.

Table 1. Criteria used to identify ODEs in this study and their expressions.

Type	Name	Formula
Traditional Methods	TM1	$([O_3]_i < 10 \text{ ppbv}) \cap ([O_3]_{i+1} < 10 \text{ ppbv})$
	TM1-5 ppbv	$([O_3]_i < 5 \text{ ppbv}) \cap ([O_3]_{i+1} < 5 \text{ ppbv})$
	TM1-4 ppbv	$([O_3]_i < 4 \text{ ppbv}) \cap ([O_3]_{i+1} < 4 \text{ ppbv})$
	TM2	$([O_3]_i < 10 \text{ ppbv}) \cap ([O_3]_{i+j} < 10 \text{ ppbv}), j \in \{1, 2, 3, 4, 5, 6\}$
	TM3	$([O_3]_i < \beta \overline{[O_3]}) \cap ([O_3]_{i+1} < \beta \overline{[O_3]}), \beta = 0.42$
	TM4	$([O_3]_i < \beta \overline{[O_3]}) \cap ([O_3]_{i+j} < \beta \overline{[O_3]}), \beta = 0.42, j \in \{1, 2, 3, 4, 5, 6\}$
	TM5	$TM4 \cup (-\frac{d[O_3]}{dt} > 1 \text{ ppbv h}^{-1})$
Variability-Based Method	VM	$[O_3]_i - \overline{[O_3]} < \alpha \cdot \sigma, \alpha = -1.5$
Machine Learning Method	Isolation Forest (IF)	-




Figure 2. Monthly averaged ozone mixing ratios at the BRW Station for spring months (March, April, and May) from the year 2000 to 2022.

et al., 2003), different constant thresholds (e.g., 5 ppbv and 4 ppbv) have also been utilized for identifying ODEs. Therefore, in addition to the commonly used 10 ppbv threshold, we also tested 5 ppbv and 4 ppbv thresholds in this study, and named them as TM1-5 ppbv and TM1-4 ppbv (see Table 1).

115 We then modified TM1 to obtain different screening criteria. At first, we relaxed the duration time of ODEs in the TM1 criterion. Instead of using a consecutive two-hour duration, we defined ODEs as that within the 6 hrs after the first time when ozone is lower than 10 ppbv, there should be at least one hour that ozone is below 10 ppbv (see TM2 in Table 1):

$$([O_3]_i < 10) \cap ([O_3]_{i+j} < 10), j \in \{1, 2, 3, 4, 5, 6\}. \quad (2)$$

This criterion is able to include time periods when an abrupt increase of ozone occurs during ODEs due to processes such as the stratospheric intrusion and local anthropogenic emissions.

120 We also replaced the fixed value (10 ppbv) used in TM1 with a percentage of the monthly averaged ozone value (see Fig. 2 showing the monthly averaged ozone mixing ratios in different months and years), and named this criterion as TM3:

$$([O_3]_i < \beta \overline{[O_3]}) \cap ([O_3]_{i+1} < \beta \overline{[O_3]}). \quad (3)$$

In Eq. (3), $\overline{[O_3]}$ is the averaged ozone value of the corresponding month. β is a constant that is artificially given. For a better comparison, we determined the coefficient β to be 0.42 using the least squares method, so that $\beta \overline{[O_3]}$ in Eq. (3) is closest to the constant 10 ppbv used in TM1 (see Fig. S1 in the Supplementary Information for the comparison between $\beta \overline{[O_3]}$ and the 10 ppbv constant).

After that, by integrating TM2 and TM3, we obtained TM4 (see Table 1):

$$([O_3]_i < \beta \overline{[O_3]}) \cap ([O_3]_{i+j} < \beta \overline{[O_3]}), j \in \{1, 2, 3, 4, 5, 6\}. \quad (4)$$

130 Based on TM4, we further took the depleting stage of ozone into consideration. Typically, a complete ODE can be divided into three stages. The first stage is the start of the ODE when the ozone depletes very fast. The second stage is that the

ozone remains at a low level (e.g., <10 ppbv), which is the maintenance of ODE. And the third stage is that the ozone returns to background levels, which is the termination of the ODE. The TM4 criterion described above basically accounts for time periods corresponding to the second stage. In the next criterion, other than time points that are identified by TM4, time points when the depletion rate of ozone is larger than 1 ppbv h⁻¹ were also added,

$$\text{TM4} \cup \left(-\frac{d[\text{O}_3]}{dt} > 1 \text{ ppbv h}^{-1} \right). \quad (5)$$

This criterion further considers time periods representing the first stage of the ODE, and is referred to as TM5, which is listed in Table 1.

2.2.2 Variability-Based Method

In our previous work (Cao et al., 2022), we picked out time points representing the tropospheric ODEs at the Halley Station in Antarctica according to the variability in the surface ozone mixing ratio, which included considerations of both the mean value and standard deviation. This method for identifying ODEs was proposed based on the study of Bian et al. (2018), when the ozone mixing ratio fulfills the following criterion:

$$[\text{O}_3]_i - \overline{[\text{O}_3]} < \alpha \cdot \sigma, \quad (6)$$

in which $[\text{O}_3]_i$ is the ozone mixing ratio at the i -th time point, and $\overline{[\text{O}_3]}$ is the monthly averaged ozone value. σ in Eq. (6) denotes the standard deviation. α is a constant which is set to -1.5 in that study (Cao et al., 2022) so that many partial ODEs (i.e., 10 ppbv < $[\text{O}_3]$ < 20 ppbv) can also be identified. According to the criterion described by Eq. (6), ODEs were defined as the time periods when the surface ozone drops to an uncommon low level, rather than the time periods when the surface ozone falls below a specific threshold, and we named this criterion as VM in the following context (see Table 1).

2.2.3 Machine Learning Method

Recently, machine learning methods have been used to investigate ODEs and the associated variability in the tropospheric BrO in the Arctic (Bougoudis et al., 2022). In this study, a machine learning method, Isolation Forest (Liu et al., 2008; Al Farizi et al., 2021), which can detect anomalies in data, was used to screen out ODE hours from the measurements (i.e., IF in Table 1). The Isolation Forest method is a commonly used machine learning approach for detecting anomalies in data, based on decision tree algorithms. In this method, anomalies are detected by constructing an ensemble of isolation trees, where each tree recursively isolates data points by randomly selecting split values until each point is isolated. After that, the average path length for each point across all trees is calculated to derive an average anomaly score, with higher scores indicating a greater likelihood of being an anomaly. Values of parameters utilized in this method for the current study are outlined as follows: n_estimators (number of trees), 100; max_samples (number of samples), auto (i.e., using all samples); contamination (proportion of outliers), auto (i.e., auto-select based on data arrangement); max_features (maximum number of features), 1.0 (i.e., using all features).

2.3 Regression of ODE Hours over Time

We employed linear regression analysis (Draper and Smith, 1998; Weisberg, 2005; Montgomery et al., 2021) to assess the trend of ODE occurrence over time. Linear regression is a classical statistical method which is widely used to quantify the trend of a dependent variable with respect to one or more independent variables through the method of least squares. The form of the linear regression used in this study is:

$$Y = a(X - 2000) + b, \quad (7)$$

where Y is the dependent variable (yearly averaged ODE hours in this study). X is the independent variable (year in this study). The slope a is the regression coefficient, representing the change rate of Y with $X - 2000$. The intercept b represents the value of Y when $X - 2000 = 0$.

To evaluate the significance of the regression, we also calculated the p -value. The p -value is usually used to judge whether the regression is significantly or not. Specifically, a p -value less than 0.05 ($p < 0.05$) suggests that there is less than a 5% probability that the observed trend could have occurred by random chance, indicating a significant trend. A p -value less than 0.01 ($p < 0.01$) suggests that there is less than a 1% probability, indicating a highly significant trend. In contrast, $0.05 < p \leq 0.1$ indicates a close to significant trend and $p > 0.1$ denotes an insignificant trend.

3 Results and Discussions

In this section, we first show the identification results of ODE hours after applying various criteria. These results were also compared and discussed to examine the characteristics and performance of each criterion. Subsequently, we selected several appropriate criteria for a detailed comparative analysis. This comparative examination allowed us to elucidate the effects of using different criteria on the interannual variability of ODEs and their associations with meteorological parameters.

3.1 Interannual Variability of ODE Hours Screened using Various Criteria

Figure 3 depicts the annual changes in ODE hours identified by various criteria across the 2000-2022 period, reflecting the characteristics of each screening criterion. We first focus on the results obtained by the traditional methods (i.e., TM1-TM5, see Fig. 3a). Generally, the results derived from all the traditional methods display a similar pattern, and a general decline in ODE occurrence over this 23-year period can be observed. The years with the most ODE hours were found to be 2000 and 2012. Between the year 2000 and 2012, the ODE hours exhibit a pattern of fluctuation throughout this period. There seems to have a weak upward trend between 2000 and 2012 if the year 2000 is excluded, suggesting a slight increase in ODE hours during this time period. This result is consistent with that of Oltmans et al. (2012) reporting an increase in the ODE occurrence frequency in the time period of 1973-2010. However, after the year 2012, the curves of all TM methods show a sharp decline, indicating a strong decrease in ODE hours in recent years. This findings is in accordance with the results of Law et al. (2023) in which they found a pronounced increase trend of the tropospheric ozone at BRW in the springtime of the period 1999-2019.




Figure 3. Number of ODE hours identified by (a) traditional methods (i.e., TM1-TM5), (b) the variability-based method (i.e., VM) and the Isolation Forest method (i.e., IF) from 2000 to 2022.

With respect to each TM criterion, it can be seen in Fig. 3(a) that the results using the thresholds of 5 ppbv (TM1-5 ppbv) and 4 ppbv (TM1-4 ppbv) are largely consistent with those using the 10 ppbv threshold (TM1), but with a significant reduction in the number of ODE hours, because more stringent thresholds were applied. The ODE hours identified using the 5 ppbv and 4 ppbv thresholds are 46% and 55% fewer, respectively, than those identified using the 10 ppbv threshold on average.

Figure 3(a) also shows that results obtained by TM1 and TM2 are almost identical, indicating that influence brought about by the modification of the ODE duration (from a continuous two-hour occurrence to a following one-hour occurrence within a six-hour period) is negligible. This also means that during occurrences of ODEs, ozone concentrations rarely make a sudden transition from below 10 ppbv to above 10 ppbv. ODEs typically persist for at least 2 hrs in the majority of cases. Similar behavior was also found between the results obtained by TM3 (ozone below $0.42[\overline{O_3}]$ for continuous two hours) and TM4 (ozone below $0.42[\overline{O_3}]$ for a start hour and a following hour within a six-hour period). We then focus on the differences between the results obtained by using TM1 and TM3. It is not surprising to see that when the difference between $0.42[\overline{O_3}]$ and 10 ppbv is significant, there is a considerable discrepancy in the screening results. For instance, the discrepancy in the results for the year 2012 is remarkable (see Fig. 3a), because the mean values of ozone in March and April of this year are quite low ($0.42[\overline{O_3}] = 4.95$ and 5.34 , respectively); conversely, when the difference is smaller, the discrepancy is less pronounced such as those for the years 2008 and 2014. As is well known, the background levels of Arctic ozone vary across different years and

seasons, driven by factors such as climate change and atmospheric circulation. For instance, the average ozone level in the Arctic in the springtime of 2023 was reported to be abnormally higher than historical levels (Ding et al., 2024). Considering these dramatic changes in the background levels of ozone in the Arctic, using “tailored” thresholds based on average ozone levels across different time periods (i.e., TM3, TM4) might be another appropriate choice to identify ODEs. With respect to
 210 TM5 ($\text{TM4} \cup (-\frac{d[\text{O}_3]}{dt} > 1 \text{ ppbv h}^{-1})$), it is seen in Fig. 3(a) that the overall trend of the TM5 curve is generally similar to that of TM4, while the TM5 curve is consistently above the TM4 curve by a margin (152 hrs on average), indicating the additional hours representing ozone depleting stages considered in the TM5 criterion.


Regarding to the variability-based method (i.e., VM in Fig. 3b), its behavior is remarkably different from those TM methods. Generally, it screens out the least number of ODE hours, indicating it as the most rigorous selection criterion compared with
 215 others. Moreover, it is shown in Fig. 3(b) that between the year 2000 and 2005, the trend of the VM curve is consistent with those of TM curves. However, during some other time periods such as from 2010 to 2015, the VM curve displays a trend that contrasts with the patterns observed in the TM curves. It is because that except the mean value, the VM criterion also takes the standard deviation into account. When the ozone concentration oscillates greatly, the standard deviation will be high, thus dominating the criterion given by Eq. (6). As a result, the variability of the criterion would be more closely aligned with the
 220 patterns of the standard deviation, rather than following the mean ozone value. More discussions are given in the following context.

With respect to the Isolation Forest method (see the IF curve in Fig. 3b), generally, the number of ODE hours screened by this method is comparable to those picked out using TM methods. Interestingly, it is seen that after the year 2014, the IF curve behaves similarly to those of the TM methods, while before 2008, the IF curve’s trend is more like the VM method’s trend,
 225 although the values are significantly higher, indicating a possible consideration of the standard deviation in the IF method. Because of the black box characteristics of machine learning models (Hassija et al., 2024), it is difficult for us to further explore the reasons and principles behind the screening results of this method. Further interpretability of this machine learning method is also one of areas we aim to investigate in the future.


In the subsequent analysis, we will focus on two specific years (2012 and 2021) to investigate more deeply into the characteristics of these criteria for identifying ODEs. These two years were selected because 2012 is one of years with the most ODE
 230 hours (530-950 hrs indicated by the TM methods). In contrast, 2021 is one of years with the least ODE hours (60-350 hrs), but having a frequent oscillation in ozone levels.

3.2 ODE Hours on Specific Years Identified by Different Criteria


Because the results of TM1 are almost identical to those of TM2, and the results of TM3 are largely in line with TM4. Additionally, results of TM1-5 ppbv and TM1-4 ppbv show a decrement from TM1’s results, and results of TM5 show an
 235 increment from TM4’s, while maintaining essentially a similar trend. Therefore, in this section, we only compare the results for ODE hours screened by TM1, TM4, VM, and IF methods on specific years (i.e., 2012 and 2021). Results for ODE hours identified using other criteria can be found in Figs. S2 and S3 in the Supplementary Information.




(a) TM1




(b) TM1




(c) TM4




(d) TM4




(e) VM



(f) VM



(g) IF



(h) IF

Figure 4. Screened results for the years 2012 and 2021 using various criteria. The blue curve represents the hourly time series of the ozone mixing ratio, and the red dots denote the ODE hours identified by various criteria.

3.2.1 Traditional Methods

240 Figure 4(a) and (b) illustrate the results of screened ODE hours using the TM1 criterion for the years 2012 and 2021. We found that in the year 2012, because the average ozone level is lower than that in 2021, the identified ODE hours in 2012 (i.e., 925 hrs) are significantly more than those in 2021 (i.e., 165 hrs). This implies that when TM1 is applied, the number of identified ODE hours is positively correlated with the annual average ozone concentration for a specific year, which can be easily expected since a year with a lower ozone level is more likely to meet the fixed threshold criteria.

245 The TM1 criterion is straightforward and easy to apply. However, Fig. 4(b) shows that in 2021, many time points when the ozone mixing ratio decreased sharply were not identified as ODE hours because the depletion is not strong enough to reduce the ozone level to below 10 ppbv. This is also the possible reason why another criterion for identifying “partial” ODEs (i.e., $10 \text{ ppbv} < [\text{O}_3] < 20 \text{ ppbv}$) is often suggested in many previous studies (Ridley et al., 2003; Koo et al., 2012; Halfacre et al., 2014).

250 In contrast to the TM1 method, TM4 gives different thresholds according to the monthly averaged ozone value across different months and years. Generally, the results obtained by TM4 are consistent with those obtained by TM1 (see Fig. 4c and d). In months and years that possess low ozone concentrations ($< 15 \text{ ppbv}$, e.g., year 2012, see Fig. 4c), the implementation of TM4 would exert a more stringent threshold so that less ODE hours are identified. In contrast, in month and years that possess high ozone concentrations (e.g., year 2021, see Fig. 4d), comparable or more ODE hours can be identified using this criterion.

255 3.2.2 Variability-Based Method

As discussed above, the variability-based method screened out significantly fewer ODE hours than the other methods. For instance, in 2012, the VM method identified less than 200 ODE hours, whereas the other methods each screened more than 500 hrs. Therefore, we focus on specific years to clarify the reasons. Figure 4(e) shows that for the year 2012, the VM method identified only a few ODE hours from the time series of ozone. Moreover, these identified ODE hours all reside in the month of
260 May, whereas no ODE hours were identified in March and April of 2012. Similar results were also found in some other years such as 2013 and 2022 (see Figs. S4a and S4b in the Supplementary Information). The reason for this ODE underestimation is that when the monthly averaged ozone level is low, being below 1.5 times the standard deviation, a negative threshold would be calculated using Eq. (6), which cannot be fulfilled. As a result, the VM method would give a null ODE identification for that month. In contrast, the ODE hours in 2021 screened by the VM method are more reasonable, shown in Fig. 4(f). This is
265 because the monthly averaged ozone levels for this year fall within a typical normal range ($\sim 20\text{--}30 \text{ ppbv}$), which is moderately greater than 1.5 times the standard deviation ($\sim 10\text{--}18 \text{ ppbv}$). Consequently, a more reasonable criterion is derived from Eq. (6). Thus, when using the VM method to determine ODE hours, extra caution is required for months characterized by notably low average ozone levels and pronounced oscillations in the mixing ratio.

To solve the problem of this ODE underestimation, we relaxed the constant α in Eq. (6) from the original value of -1.5
270 designed for the Halley Station to a value of -0.8. Consequently, more reasonable results were obtained for the year 2012 (see Fig.S4c in the Supplementary Information). The interannual variability of ODE hours determined by this modified criterion

is also more consistent with those determined by other criteria (see Fig. S5 in the Supplementary Information). However, in years and months with high ozone levels, this modified criterion with a smaller α seems to give an excessively high number of ODE hours. For instance, in the year 2021 (see Fig.S4d in the Supplementary Information), this method (VM with $\alpha = -0.8$) recognized many data points with ozone mixing ratios between 15 and 20 ppbv. However, the springtime average ozone mixing ratio at BRW for the year 2021 was calculated to be 23.93 ppbv. It means that many time points picked up by this criterion possess an ozone value close to the average ozone level in springtime at BRW. In that case, we feel that many of these time points cannot be viewed as ODE hours, which also indicates that this criterion may overestimate the number of ODE hours. This overestimation in ODE hours may also lead to a misunderstanding of the trend of ODE occurrence. We also tested other values of α such as 1.0, but still did not obtain satisfactory screening results for the ODEs in 2012 at BRW (not shown here). Thus, we concluded that for the variability-based method, the original value of α (i.e., -1.5) designed for the Halley Station in Antarctica may not be suitable for identifying ODEs in certain years and months at other stations, and a more appropriate value for α should be carefully determined for different years, months, and stations.

3.2.3 Machine Learning Method

Results of the IF method are very interesting. For years with normal ozone levels such as 2021 (see Fig. 4h), the IF method performs well, identifying not only hours with low ozone but also hours when ozone suddenly drops. However, for years with relatively low ozone levels such as 2012 (see Fig. 4g), the IF method gives problematic results. It was found in Fig. 4(g) that the hours with a moderately low ozone value (3-7 ppbv) were not recognized as ODE hours. The reason is that in the machine learning model, dense data were regarded as normal whereas sparse data were considered as outliers. Thus, for years with only a few ODE occurrences such as 2021, the IF method is capable of identifying ODE hours by recognizing them as outliers. However, in years with a frequent occurrence of ODEs such as 2012, hours with moderately low ozone values are regarded as normal so that they are not viewed as ODE hours by this model. Thus, the IF method exhibits a limitation in accurately identifying ODE hours in years characterized by a high frequency of ODE occurrences.

From the results discussed above, we found that TM1 and TM4 are more suitable for identifying ODEs from the time series of ozone at the BRW Station than the other criteria. We then investigated the monthly and yearly variability of ODE hours at the BRW Station based on the results applying these two criteria. Another two criteria with different constant thresholds (i.e., TM1-5 ppbv and TM1-4 ppbv) were also examined to assess the impact of varying the constant threshold on the conclusions.

3.3 Variability of ODE Hours at BRW

We first analyzed the monthly variation in ODE hours across various spring months over these 23 years (see Fig. 5). It is shown in Fig. 5(a) that when the constant 10 ppbv criterion was applied, April has the most ODE hours, with a median of approximately 210 hrs, suggesting that April is the predominant month when ODEs occur. In contrast, May possesses the least ODE hours, with a median of 57 hrs. Moreover, Fig. 5(a) shows that in March, the number of ODE hours ranges from zero to more than 400 hrs, with the majority concentrated around 170 hrs. In contrast, the range of ODE hours for April is narrower, spanning from 50 to 400 hrs, but with a more uniform distribution of hours throughout this range. It means that the ozone




Figure 5. Monthly hours of ODEs for March, April, and May across the 23-year period from 2000 to 2022. The ODE hours were screened by (a) TM1, (b) TM4, (c) TM1-5 ppbv and (d) TM1-4 ppbv.

concentration at the BRW Station fluctuates more widely in March. For May, this month possesses a relatively small range of ODE hours (0-350 hrs), with the bulk of hours centered at a low value (~ 50 hrs).

When the TM4 criterion (ozone below $0.42[\overline{O_3}]$ for a start hour and a following hour within a six-hour period) was applied (Fig. 5b), we found that the order of the medians for these three months remains unchanged (April>March>May), compared with the results of TM1. However, we found that the ODE hours in April lowered while the hours in May elevated, when TM4 was applied. The data are also more concentrated. It is because that TM4 uses a threshold that depends on the monthly averaged value. As the average ozone level in April is lower compared with those in March and May, using this relative criterion would exert a more stringent threshold than the constant 10 ppbv so that it screens out less ODE hours than TM1. For May with a high ozone level, the relative criterion is easier to achieve compared to the constant 10 ppbv criterion. Therefore, TM4 identifies

more ODE hours in May compared to TM1. In summary, the TM4 method can screen out more ODE hours in months with
315 high ozone values than TM1, and vice versa.

When 5 ppbv was used to replace the 10 ppbv threshold (i.e., TM1-5 ppbv, see Fig. 5c), we found that the order of ODE
hours across the three spring months remained unchanged (April>March>May). However, the ODE hours for each month
were significantly lower compared to those identified by TM1, due to the stricter threshold applied. Furthermore, the discrep-
ancy in ODE hours between March and April was found to be less pronounced than that in the TM1 results, rendering the
320 ODE hours in March (median: 87 hrs) and April (median: 102 hrs) very close. This indicates that the primary cause for the
significantly greater number of ODE hours in April compared to March, as identified by TM1, is the more frequent occurrence
of ozone concentrations falling within the 5-10 ppbv range in April than in March. With respect to the situation in May, the
implementation of the 5 ppbv threshold resulted in a significant reduction to almost zero ODE hours. This finding suggests that
ODEs characterized by very low ozone levels (<5 ppbv) have nearly vanished in May in recent years, which could be linked
325 to global warming, resulting in higher temperatures during May, thereby causing the ODE season to cease earlier than before
(Burd et al., 2017). Results obtained using the 4 ppbv threshold (Fig. 5d) are similar to those using the 5 ppbv threshold, except
that the discrepancy in ODE hours between March (median: 73 hrs) and April (median: 85 hrs) is even less pronounced, thereby
confirming the conclusions drawn above.

Figure 6 presents the yearly variability of the ODE hours for each spring month and the entire spring season, spanning
330 from 2000 to 2022. It is seen from Fig. 6(a) that based on the results of TM1, ODE hours in spring decrease significantly
across these 23 years ($p<0.05$). This decreasing trend is mostly caused by the decline in ODE hours in April (see Fig. 6e),
which is highly significant ($p<0.01$). In contrast to April, the drops in ODE hours in March and May are not significant ($p>0.1$,
Fig. 6c) and close to significant ($0.05<p<0.1$, Fig. 6g), respectively. Our findings are in good agreement with those of Law
et al. (2023), who reported a notable increase in observed surface ozone levels during spring from 1993 to 2019 at BRW, with
335 the most significant elevation observed in April. Hung et al. (2025) also observed an increasing trend in Arctic spring ozone
concentrations at Eureka, Nunavut, Canada (80°N , 86°W) from 2008 to 2022, further supporting the notion of declining ODE
frequency in the Arctic. Burd et al. (2017), in their study on the Arctic BrO season, found a decrease in BrO concentrations
and an early end of the BrO season at BRW from 2012 to 2016, which may also imply a reduction in the occurrence of ODEs,
aligning with our findings.

340 The results of TM4 are similar (Fig. 6b), but showing a more significant decline trend. The decrease in ODE hours in the
entire spring season transits to be highly significant ($p<0.01$) in the results of TM4, and the p-value in April is also smaller,
confirming the highly significant decline in ODE hours in April at BRW over the 23-year period. Aside from that, the drop
in ODE hours in May was found to be insignificant ($p>0.1$) in TM4's results, whereas in TM1's results, the drop in May
approached statistical significance ($0.05<p<0.1$).

345 When more rigorous thresholds (5 ppbv and 4 ppbv) were applied instead of the 10 ppbv threshold (refer to Fig.S6 in the Sup-
plementary Information), the reduction in ODE hours in the entire spring season was found to be highly significant ($p<0.01$),
denoting a more remarkable decline in the occurrence of severe ODEs with very low ozone levels during these years. This re-
markable decline is still mainly attributable to the highly significant reduction in ODE hours in April ($p<0.01$). The decrease in




Figure 6. Yearly variability of ODE hours at the BRW Station, identified by two different criteria. Subplots (a), (c), (e) and (g) show the ODE hours screened by the TM1 method for the whole spring, March, April and May, respectively, and subplots (b), (d), (f) and (h) show the hours screened by the TM4 method. Red dashed lines represent linear regressions of the ODE hours. The regression equations and p-values are also given.

March remains statistically insignificant, while the decline in May is identified as insignificant ($p > 0.1$) for the 5 ppbv threshold and close to significant ($0.05 < p < 0.1$) for the 4 ppbv threshold, respectively.




Figure 7. Wind rose diagrams during (a) the investigated spring seasons from 2000 to 2022, and ODE time periods identified by (b) TM1 and (c) TM4.

In summary, results utilizing different criteria all indicate a decline in ODE hours during the spring season over the 23-year period, primarily driven by the highly significant reduction in April. However, when employing the threshold that varies with the monthly average ($0.42[\overline{O_3}]$) or more stringent fixed thresholds (5 ppbv and 4 ppb), the results would show a more significant decline in ODE hours in spring, compared to that using the 10 ppb threshold.

355 3.4 Relationship between ODE Hours and Meteorological Parameters

We then investigated the relationship between ODE hours and surface meteorological parameters measured at the BRW Station. Figure 7 shows the wind information during the investigated spring seasons (Fig. 7a) and the time periods of ODEs identified by TM1 (Fig. 7b) and TM4 (Fig. 7c). From Fig. 7(a), we can see that the prevailing wind at BRW is northeastern in the springtime. The wind speeds are relatively higher when the winds are easterly ($9\text{--}12\text{ m s}^{-1}$ at most) and northeasterly ($>12\text{ m s}^{-1}$ at most). In contrast, wind speeds along other directions are mostly lower than 9 m s^{-1} . Compared to the wind speeds throughout the whole spring, Fig. 7(b) shows that during ODEs, the wind speeds are significantly lower (below 9 m s^{-1} , mostly $3\text{--}6\text{ m s}^{-1}$), denoting a favored moderate wind speed condition of ODEs at BRW. Moreover, the wind direction tends

to favor the northeasterly and northerly winds during ODEs, which is associated with the fresh sea ice covering the ocean to the north of BRW (Bottenheim and Chan, 2006; Gilman et al., 2010; Oltmans et al., 2012; Peterson et al., 2016). Additionally, we also found an increased proportion of westerly winds and a decreased proportion of easterly winds in Fig. 7(b), denoting a favored westerly wind condition for ODEs at BRW. In an earlier study of Oltmans et al. (2012), they suggested that ODEs at BRW are predominantly associated with easterly winds, a finding that contrasts with the conclusions achieved in this study. The discrepancy between our findings and those of Oltmans et al. (2012) may source from the different time periods investigated in these two studies. Moreover, a recent flight campaign originating from BRW (Brockway et al., 2024) also detected elevated levels of BrO being advected from the west of BRW, suggesting a potential transport of ozone-depleting air from the western direction. Additionally, in our recent modeling study on ODEs at BRW (Cao et al., 2023), we also found an ozone-depleting air mass transported to the BRW Station from the southwest under the influence of a cyclone moving eastward. The results obtained by applying the TM4 criterion (Fig. 7c) are also similar, confirming our findings.

Results utilizing the 5 ppbv and 4 ppbv thresholds (refer to Fig.S7 in the Supplementary Information) also suggested that northerly and northeasterly wind conditions would facilitate the occurrence of ODEs at BRW. However, relative fractions of the moderate wind speed ($3\text{--}6\text{ m s}^{-1}$) and the low wind speed ($0\text{--}3\text{ m s}^{-1}$) during the identified ODE periods are higher than those found in the TM1's results, indicating that lower wind speeds are conducive to the occurrence of more severe ODEs, characterized by very low ozone levels.

The connections between ODE hours and the 2-m temperature measured at BRW are shown in Fig. 8, and the results obtained by applying TM1 (Fig. 8a) and TM4 (Fig. 8b) are similar. It was found that in the range of 250–272 K, the temperature at BRW generally exhibits a uniform distribution. However, occurrences of ODEs are more frequent at a lower temperature, with the highest occurrence frequency at approximately 250 K. Moreover, when the temperature is lower than approximately 256 K, the decrease in the 2-m temperature would substantially enhance the occurrence of ODEs (see the blue dot-dash lines in Fig. 8a and b). Thus, a lower temperature condition can facilitate the occurrence of ODEs, which is possibly associated with the stability of the boundary layer (Lehrer et al., 2004; Koo et al., 2012). It could also be linked to the effective absorption of HOBr on frozen NaCl/NaBr surfaces at temperatures below the eutectic point of NaCl·2H₂O (i.e., 252 K), as reported by Adams et al. (2002). Below this temperature, a quasi-brine layer, characterized by its high acidity (Cho et al., 2002), is likely to develop on the ice/snow surface, which would promote bromine activation and subsequent ozone depletion. This lower temperature condition favored by the bromine explosion and the ozone depletion was also reported by Zilker et al. (2023), by applying a composite analysis on long-term ozonesonde data (2010–2021) and surface measurements over the Svalbard area in the Arctic.

Results obtained by applying TM1-5 ppbv and TM1-4 ppbv (Fig. 8c and d) also indicate that ODEs are favored by low temperature conditions. More interestingly, Fig. 8(c) and (d) reveal a significant reduction in the occurrence frequency of severe ODEs within the temperature range of 256–262 K, when compared to the TM1's results (Fig. 8a). This suggests that severe ODEs are more likely to occur only at temperatures below 256 K.

Figure 9 show the relationship between the surface pressure of BRW and the occurrence of ODEs. Results obtained by using various criteria are largely consistent, exhibiting only slight differences. We found the occurrence of ODEs mostly residing in the pressure range of 1012–1027 hPa. Furthermore, an increase in surface pressure at BRW is often associated with a more




Figure 8. Occurrence frequency of ODEs versus the 2-m temperature. The ratio was calculated by the number of hours within each interval divided by the total number of hours. ODE hours used to calculate the ratio in (a), (b), (c) and (d) were screened by TM1, TM4, TM1-5 ppbv and TM1-4 ppbv, respectively. The blue dot-dash lines denote the ratio of ODEs to normal conditions, and a value greater than 1.0 indicates a favorable condition for ODEs.

frequent occurrence of ODEs, possibly due to the stable and calm conditions that prevail under the control of high-pressure




Figure 9. Occurrence frequency of ODEs versus the surface pressure. The other settings for this figure are similar to those for Fig. 8.

systems. Additionally, it can be seen in Fig. 9 that when the surface pressure resides in the range of 990-995 hPa, the occurrence of ODEs is also favored (with the ratio ODE/Normal >1.0). It might be connected to the blowing snow events (Yang et al.,

2010, 2019, 2020; Huang et al., 2018, 2020) when low-pressure systems (e.g., cyclones) pass by, which are beneficial for the BrO release and the subsequent ozone depletion (Begoin et al., 2010; Zhao et al., 2016).

4 Conclusions and Future Work

In this study, we investigated the influence of applying various criteria to identify springtime tropospheric ozone depletion events (ODEs) at Utqiagvik (BRW), Arctic, using observational data from 2000 to 2022. We tested three types of criteria: traditional methods, variability-based methods, and machine learning methods, and analyzed the characteristics of these criteria in depth.

We found the criteria using a constant threshold (e.g., 10 ppbv) and using thresholds based on the monthly averaged ozone values more suitable for identifying ODEs at BRW than the other criteria. In contrast, the criterion considering both the mean value and standard deviation of ozone (i.e., the VM criterion) is able to identify time points when the surface ozone drops to an uncommon low level instead of a fixed threshold, which is more adaptive and sensitive. However, extra caution is required when determining the parameter value of this criterion. Apart from these criteria, the machine learning method adopted in this study (i.e., the IF method) can automatically detect ODE hours. But this method has a poor interpretability in screening results and sometimes is unable to correctly identify ODE hours when ODEs occur very frequently. Furthermore, criteria that are proposed as suitable for identifying ODEs all indicate an overall decreasing trend in the occurrence frequency of ODEs at BRW over this 23-year period, with the most significant decline observed in April. This suggests a potential impact of climate change such as the global warming and Arctic sea ice melting on ODE occurrences. This declining trend of ODEs can lead to a weakening of the deposition of active mercury (Hg(II)), which is highly toxic and can pose serious health risks to humans. Therefore, the decline in ODE frequency could lead to a reduction in the health hazards associated with mercury deposition in mid-latitude regions. However, results obtained by implementing a threshold that varies with the monthly average, or by applying more stringent thresholds (5 ppbv and 4 ppbv), demonstrate a more significant reduction in the occurrence of ODEs compared to those using the 10 ppbv threshold. Thus, using a different criterion to identify ODE hours may lead to different conclusions. This is why we tested various ODE criteria, compared the resulting conclusions, and attempted to propose suitable criteria for identifying ODEs in this study.

Applying suitable criteria also enables us to study the connection between meteorological conditions at BRW and the occurrence of ODEs more precisely. ODEs at BRW were found to be more likely to occur under northerly and northeasterly winds, with moderate wind speeds (mostly $3\text{--}6\text{ m s}^{-1}$) being more favorable, although ODEs were also observed at higher wind speeds ($6\text{--}12\text{ m s}^{-1}$). Lower wind speed conditions were also found to facilitate the occurrence of more severe ODEs, characterized by very low ozone concentrations. ODEs were also found to be closely linked to temperature and pressure. ODEs especially the severe ones tend to occur at temperatures lower than 256 K, which is possibly associated with the stability of the boundary layer and the effective absorption of HOBr on frozen surfaces, promoting bromine activation and subsequent ozone depletion. Additionally, ODEs at BRW tend to occur under high pressure conditions ($>1010\text{ hPa}$), indicating that the high-pressure associated weather conditions may facilitate the occurrence of ODEs.

In the future, we would like to improve the machine learning methods (e.g., Isolation Forest) so that a more reliable method
435 can be applied in ODE identification. Moreover, present analysis can also be conducted to a longer time series of data to provide
a more comprehensive understanding of the long-term trends and variability of ODEs in the Arctic. Conclusions obtained in
this study should also be verified in other Arctic stations. To achieve this goal, we gathered more observational data from 7 other
Arctic sites (Alert, Esrange, Tustervatn, Villum, Pallas, Summit, and Zeppelin) aside from the one (BRW) we initially focused
on in this study. After applying the criteria using constant values (10 ppbv and 5 ppbv) that are proposed in this study, only four
440 of these sites (Alert, BRW, Villum, and Zeppelin) were found to have ODEs occurrences (see Fig. S8 in the Supplementary
Information). We found that the ODE hours screened out by the criteria proposed in this study seem appropriate. For instance,
BRW, which has a low altitude, is featured with high frequency of ODE occurrence. In contrast, Zeppelin, which is located
at a higher altitude, has fewer ODE hours, because the air mass arriving at Zeppelin usually represents the air in the free
troposphere so that more ozone can be transported from the stratosphere and less halogens released from the surface can reach
445 the free troposphere due to the barrier at the top of the boundary layer. Additionally, the ODE curves for Villum also shows
a declining trend, which is consistent with the conclusion achieved in this study. However, more observational data for other
species such as halogen species (i.e., BrO) from satellite detection and ground-based multi-axis differential optical absorption
spectroscopy (MAX-DOAS) are still needed to validate these results and help to identify ODEs more accurately. Unfortunately,
we currently do not have access to such data, which is also a limitation of the present study.

450 *Code and data availability.* All data needed to evaluate the conclusions in the paper are present in the paper. The source code of the model
and the data of the computational results shown in this article can be acquired upon request from the authors.

Author contributions. L.C. conceptualized the study and supervised the entire research process. X.Z. conducted the simulations and processed the data. X.Y. and S.L. contributed to the interpretation of the results. J.W. provided valuable insights into the model results. T.Z. assisted with the data analysis and manuscript preparation. All authors discussed the results and contributed to the final manuscript.

455 *Competing interests.* The authors declare no conflict of interest.

Acknowledgements. This study is funded by the National Key Research and Development Program of China (Grant No. 2022YFC3701204),
the National Natural Science Foundation of China (Grant No. 41705103), and the 2023 Outstanding Young Backbone Teacher of Jiangsu
"Qinglan" Project (Grant No. R2023Q02). The authors would like to thank the National Supercomputer Center in Tianjin and High Per-
formance Computing Center at the Nanjing University of Information Science and Technology to provide the high-performance computing
460 system for calculations.

References

- Abbatt, J. P. D., Thomas, J. L., Abrahamsson, K., Boxe, C., Granfors, A., Jones, A. E., King, M. D., Saiz-Lopez, A., Shepson, P. B., Sodeau, J., Toohey, D. W., Toubin, C., von Glasow, R., Wren, S. N., and Yang, X.: Halogen activation via interactions with environmental ice and snow in the polar lower troposphere and other regions, *Atmospheric Chemistry and Physics*, 12, 6237–6271, <https://doi.org/10.5194/acp-12-6237-2012>, 2012.
- Adams, J. W., Holmes, N. S., and Crowley, J. N.: Uptake and reaction of HOBr on frozen and dry NaCl/NaBr surfaces between 253 and 233 K, *Atmospheric Chemistry and Physics*, 2, 79–91, <https://doi.org/10.5194/acp-2-79-2002>, 2002.
- Akimoto, H.: *Atmospheric Reaction Chemistry*, Springer Tokyo, 1 edn., <https://doi.org/10.1007/978-4-431-55870-5>, 2016.
- Al Farizi, W. S., Hidayah, I., and Rizal, M. N.: Isolation Forest Based Anomaly Detection: A Systematic Literature Review, in: 2021 8th International Conference on Information Technology, Computer and Electrical Engineering (ICITACEE), pp. 118–122, <https://doi.org/10.1109/ICITACEE53184.2021.9617498>, 2021.
- Anlauf, K., Mickley, R., and Trivett, N.: Measurement of ozone during Polar Sunrise Experiment 1992, *Oceanographic Literature Review*, 7, 531, 1995.
- Ariya, P. A., Dastoor, A. P., Amyot, M., Schroeder, W. H., Barrie, L., Anlauf, K., Raofie, F., Ryzhkov, A., Davignon, D., Lalonde, J., et al.: The Arctic: a sink for mercury, *Tellus B: Chemical and Physical Meteorology*, 56, 397–403, 2004.
- Barrie, L., Bottenheim, J., Schnell, R. C., Crutzen, P., and Rasmussen, R.: Ozone destruction and photochemical reactions at polar sunrise in the lower Arctic atmosphere, *Nature*, 334, 138–141, 1988.
- Basu, N., Abass, K., Dietz, R., Krümmel, E., Rautio, A., and Weihe, P.: The impact of mercury contamination on human health in the Arctic: A state of the science review, *Science of the Total Environment*, 831, 154 793, 2022.
- Begoin, M., Richter, A., Weber, M., Kaleschke, L., Tian-Kunze, X., Stohl, A., Theys, N., and Burrows, J. P.: Satellite observations of long range transport of a large BrO plume in the Arctic, *Atmospheric Chemistry and Physics*, 10, 6515–6526, <https://doi.org/10.5194/acp-10-6515-2010>, 2010.
- Bian, L., Ye, L., Ding, M., Gao, Z., Zheng, X., and Schnell, R.: Surface Ozone Monitoring and Background Concentration at Zhongshan Station, Antarctica, *Atmospheric and Climate Sciences*, 8, 1–14, <https://doi.org/10.4236/acs.2018.81001>, 2018.
- Bottenheim, J., Natcheva, S., Morin, S., and Nghiem, S.: Ozone in the boundary layer air over the Arctic Ocean: measurements during the TARA transpolar drift 2006–2008, *Atmospheric Chemistry and Physics*, 9, 4545–4557, 2009.
- Bottenheim, J. W. and Chan, E.: A trajectory study into the origin of spring time Arctic boundary layer ozone depletion, *Journal of Geophysical Research: Atmospheres*, 111, <https://doi.org/10.1029/2006JD007055>, 2006.
- Bottenheim, J. W., Gallant, A. G., and Brice, K. A.: Measurements of NO_y species and O₃ at 82 N latitude, *Geophysical Research Letters*, 13, 113–116, 1986.
- Bottenheim, J. W., Fuentes, J. D., Tarasick, D. W., and Anlauf, K. G.: Ozone in the Arctic lower troposphere during winter and spring 2000 (ALERT2000), *Atmospheric Environment*, 36, 2535–2544, 2002.
- Bougoudis, I., Blechschmidt, A.-M., Richter, A., Seo, S., and Burrows, J. P.: Simulating tropospheric BrO in the Arctic using an artificial neural network, *Atmospheric Environment*, 276, 119 032, <https://doi.org/10.1016/j.atmosenv.2022.119032>, 2022.
- Brockway, N., Peterson, P. K., Bigge, K., Hajny, K. D., Shepson, P. B., Pratt, K. A., Fuentes, J. D., Starn, T., Kaeser, R., Stirm, B. H., and Simpson, W. R.: Tropospheric bromine monoxide vertical profiles retrieved across the Alaskan Arctic in springtime, *Atmospheric Chemistry and Physics*, 24, 23–40, <https://doi.org/10.5194/acp-24-23-2024>, 2024.

- Burd, J. A., Peterson, P. K., Nghiem, S. V., Perovich, D. K., and Simpson, W. R.: Snowmelt onset hinders bromine monoxide heterogeneous recycling in the Arctic, *Journal of Geophysical Research: Atmospheres*, 122, 8297–8309, <https://doi.org/https://doi.org/10.1002/2017JD026906>, 2017.
- Cairo, F. and Colavitto, T.: Polar Stratospheric Clouds in the Arctic, *Physics and Chemistry of the Arctic Atmosphere*, pp. 415–467, 2020.
- Cao, L., Fan, L., Li, S., and Yang, S.: Influence of total ozone column (TOC) on the occurrence of tropospheric ozone depletion events (ODEs) in the Antarctic, *Atmospheric Chemistry and Physics*, 22, 3875–3890, 2022.
- Cao, L., Li, S., Gu, Y., and Luo, Y.: A three-dimensional simulation and process analysis of tropospheric ozone depletion events (ODEs) during the springtime in the Arctic using CMAQ (Community Multiscale Air Quality Modeling System), *Atmospheric Chemistry and Physics*, 23, 3363–3382, <https://doi.org/10.5194/acp-23-3363-2023>, 2023.
- Chen, D., Luo, Y., Yang, X., Si, F., Dou, K., Zhou, H., Qian, Y., Hu, C., Liu, J., and Liu, W.: Study of an Arctic blowing snow-induced bromine explosion event in Ny-Ålesund, Svalbard, *Science of the total environment*, 839, 156 335, 2022.
- Cho, H., Shepson, P., Barrie, L., Cowin, J., and Zaveri, R.: NMR Investigation of the Quasi-Brine Layer in Ice/Brine Mixtures, *Journal of Physical Chemistry B - J PHYS CHEM B*, 106, <https://doi.org/10.1021/jp020449+>, 2002.
- Crocker, I.: NOAA GML Meteorology Data, Barrow, 2000-03-01 to 2022-05-31, Data fields: 1-14, <https://gml.noaa.gov/aftp/data/barrow/meteorology/>, date accessed: 2024-11-01, 2024.
- Crutzen, P. J. and Arnold, F.: Nitric acid cloud formation in the cold Antarctic stratosphere: A major cause for the springtime ‘ozone hole’, *Nature*, 324, 651–655, 1986.
- Custard, K., Raso, A., Shepson, P., Staebler, R., and Pratt, K.: Production and Release of Molecular Bromine and Chlorine from the Arctic Coastal Snowpack, *ACS Earth and Space Chemistry*, 1, <https://doi.org/10.1021/acsearthspacechem.7b00014>, 2017.
- Dastoor, A., Wilson, S. J., Travníkov, O., Ryjkov, A., Angot, H., Christensen, J. H., Steenhuisen, F., and Muntean, M.: Arctic atmospheric mercury: Sources and changes, *Science of The Total Environment*, 839, 156 213, 2022.
- Ding, M.-H., Wang, X., Bian, L.-G., Jiang, Z.-N., Lin, X., Qu, Z.-F., Su, J., Wang, S., Wei, T., Zhai, X.-C., Zhang, D.-Q., Zhang, L., Zhang, W.-Q., Zhao, S.-D., and Zhu, K.-J.: State of polar climate in 2023, *Advances in Climate Change Research*, <https://doi.org/https://doi.org/10.1016/j.accre.2024.08.004>, 2024.
- Draper, N. R. and Smith, H.: *Applied regression analysis*, vol. 326, John Wiley & Sons, 1998.
- Fan, S.-M. and Jacob, D. J.: Surface ozone depletion in Arctic spring sustained by bromine reactions on aerosols, *Nature*, 359, 522–524, 1992.
- Frieß, U., Sihler, H., Sander, R., Pöhler, D., Yilmaz, S., and Platt, U.: The vertical distribution of BrO and aerosols in the Arctic: Measurements by active and passive differential optical absorption spectroscopy, *Journal of Geophysical Research: Atmospheres*, 116, 2011.
- Gilman, J. B., Burkhart, J. F., Lerner, B. M., Williams, E. J., Kuster, W. C., Goldan, P. D., Murphy, P. C., Warneke, C., Fowler, C., Montzka, S. A., Miller, B. R., Miller, L., Oltmans, S. J., Ryerson, T. B., Cooper, O. R., Stohl, A., and de Gouw, J.: Ozone variability and halogen oxidation within the Arctic and sub-Arctic springtime boundary layer, *Atmospheric Chemistry and Physics*, 10, 10 223–10 236, 2010.
- Halfacre, J., Knepp, T., Shepson, P., Thompson, C., Pratt, K., Li, B., Peterson, P., Walsh, S., Simpson, W., Matrai, P., et al.: Temporal and spatial characteristics of ozone depletion events from measurements in the Arctic, *Atmospheric Chemistry and Physics*, 14, 4875–4894, 2014.
- Hassija, V., Chamola, V., Mahapatra, A., Singal, A., Goel, D., Huang, K., Scardapane, S., Spinelli, I., Mahmud, M., and Husain, A.: Interpreting Black-Box Models: A Review on Explainable Artificial Intelligence, *Cognitive Computation*, 16, 45–74, <https://doi.org/10.1007/s12559-023-10179-8>, 2024.

- Hausmann, M. and Platt, U.: Spectroscopic measurement of bromine oxide and ozone in the high Arctic during Polar Sunrise Experiment 1992, *Journal of Geophysical Research: Atmospheres*, 99, 25 399–25 413, 1994.
- Herrmann, M., Sihler, H., Frieß, U., Wagner, T., Platt, U., and Gutheil, E.: Time-dependent 3D simulations of tropospheric ozone depletion events in the Arctic spring using the Weather Research and Forecasting model coupled with Chemistry (WRF-Chem), *Atmospheric Chemistry and Physics*, 21, 7611–7638, 2021.
- Herrmann, M., Schöne, M., Borger, C., Warnach, S., Wagner, T., Platt, U., and Gutheil, E.: Ozone depletion events in the Arctic spring of 2019: a new modeling approach to bromine emissions, *Atmospheric Chemistry and Physics*, 22, 13 495–13 526, 2022.
- Huang, J., Jaeglé, L., and Shah, V.: Using CALIOP to constrain blowing snow emissions of sea salt aerosols over Arctic and Antarctic sea ice, *Atmospheric Chemistry and Physics*, 18, 16 253–16 269, <https://doi.org/10.5194/acp-18-16253-2018>, 2018.
- Huang, J., Jaeglé, L., Chen, Q., Alexander, B., Sherwen, T., Evans, M. J., Theys, N., and Choi, S.: Evaluating the impact of blowing-snow sea salt aerosol on springtime BrO and O₃ in the Arctic, *Atmospheric Chemistry and Physics*, 20, 7335–7358, <https://doi.org/10.5194/acp-20-7335-2020>, 2020.
- Hung, J., Liu, L., Palm, M., Mariani, Z., Manney, G. L., Millán, L. F., and Strong, K.: Autonomous Year-Round Measurements of O₃, CO, CH₄, and N₂O in the High Arctic With the Atmospheric Emitted Radiance Interferometer, *Journal of Geophysical Research: Atmospheres*, 130, e2024JD042 847, 2025.
- Jacobi, H.-W., Morin, S., and Bottenheim, J. W.: Observation of widespread depletion of ozone in the springtime boundary layer of the central Arctic linked to mesoscale synoptic conditions, *Journal of Geophysical Research: Atmospheres*, 115, 2010.
- Jones, A. E., Wolff, E. W., Ames, D., Bauguutte, S.-B., Clemmshaw, K., Fleming, Z., Mills, G., Saiz-Lopez, A., Salmon, R. A., Sturges, W., et al.: The multi-seasonal NO_y budget in coastal Antarctica and its link with surface snow and ice core nitrate: results from the CHABLIS campaign, *Atmospheric Chemistry and Physics*, 11, 9271–9285, 2011.
- Koo, J.-H., Wang, Y., Kurosu, T. P., Chance, K., Rozanov, A., Richter, A., Oltmans, S. J., Thompson, A. M., Hair, J. W., Fenn, M. A., Weinheimer, A. J., Ryerson, T. B., Solberg, S., Huey, L. G., Liao, J., Dibb, J. E., Neuman, J. A., Nowak, J. B., Pierce, R. B., Natarajan, M., and Al-Saadi, J.: Characteristics of tropospheric ozone depletion events in the Arctic spring: analysis of the ARCTAS, ARCPAC, and AR-CIONS measurements and satellite BrO observations, *Atmospheric Chemistry and Physics*, 12, 9909–9922, <https://doi.org/10.5194/acp-12-9909-2012>, 2012.
- Koo, J.-H., Wang, Y., Jiang, T., Deng, Y., Oltmans, S. J., and Solberg, S.: Influence of climate variability on near-surface ozone depletion events in the Arctic spring, *Geophysical Research Letters*, 41, 2582–2589, 2014.
- Lacis, A. A., Wuebbles, D. J., and Logan, J. A.: Radiative forcing of climate by changes in the vertical distribution of ozone, *Journal of Geophysical Research: Atmospheres*, 95, 9971–9981, 1990.
- Law, K. S., Hjorth, J. L., Pernov, J. B., Whaley, C., Skov, H., Collaud Coen, M., Langner, J., Arnold, S. R., Tarasick, D. W., Christensen, J., Deushi, M., Effertz, P., Faluvegi, G., Gauss, M., Im, U., Oshima, N., Petropavlovskikh, I., Plummer, D., Tsigaridis, K., Tsyro, S., Solberg, S., and Turnock, S. T.: Arctic tropospheric ozone trends, *Geophysical Research Letters*, 50, e2023GL103 096, <https://doi.org/10.1029/2023GL103096>, 2023.
- Lehrer, E., Hönninger, G., and Platt, U.: A one dimensional model study of the mechanism of halogen liberation and vertical transport in the polar troposphere, *Atmospheric Chemistry and Physics*, 4, 2427–2440, <https://doi.org/10.5194/acp-4-2427-2004>, 2004.
- Liu, F. T., Ting, K. M., and Zhou, Z.-H.: Isolation forest, in: 2008 Eighth IEEE International Conference on Data Mining, pp. 413–422, IEEE, 2008.

- Liu, W., Hegglin, M. I., Checa-Garcia, R., Li, S., Gillett, N. P., Lyu, K., Zhang, X., and Swart, N. C.: Stratospheric ozone depletion and tropospheric ozone increases drive Southern Ocean interior warming, *Nature Climate Change*, 12, 365–372, 2022.
- 575 McClure-Begley, A., Petropavlovskikh, I., and Oltmans, S.: NOAA GLObal Monitoring Surface Ozone Network, Barrow, 2000-03-01 to 2022-05-31, <http://dx.doi.org/10.7289/V57P8WBF>, date accessed: 2024-11-01, 2024.
- McConnell, J., Henderson, G., Barrie, L., Bottenheim, J., Niki, H., Langford, C., and Templeton, E.: Photochemical bromine production implicated in Arctic boundary-layer ozone depletion, *Nature*, 355, 150–152, 1992.
- Michalowski, B. A., Francisco, J. S., Li, S.-M., Barrie, L. A., Bottenheim, J. W., and Shepson, P. B.: A computer model study of multi-
580 phase chemistry in the Arctic boundary layer during polar sunrise, *Journal of Geophysical Research: Atmospheres*, 105, 15 131–15 145, <https://doi.org/https://doi.org/10.1029/2000JD900004>, 2000.
- Montgomery, D. C., Peck, E. A., and Vining, G. G.: *Introduction to linear regression analysis*, John Wiley & Sons, 2021.
- Oltmans, S. J.: Surface ozone measurements in clean air, *Journal of Geophysical Research: Oceans*, 86, 1174–1180, 1981.
- Oltmans, S. J., Johnson, B. J., and Harris, J. M.: Springtime boundary layer ozone depletion at Barrow, Alaska: Meteorological influence, year-to-year variation, and long-term change, *Journal of Geophysical Research: Atmospheres*, 117,
585 <https://doi.org/https://doi.org/10.1029/2011JD016889>, 2012.
- Peterson, P. K., Simpson, W. R., and Nghiem, S. V.: Variability of bromine monoxide at Barrow, Alaska, over four halogen activation (March–May) seasons and at two on-ice locations, *Journal of Geophysical Research: Atmospheres*, 121, 1381 – 1396, 2016.
- Peterson, P. K., Hartwig, M., May, N. W., Schwartz, E., Rigor, I., Ermold, W., Steele, M., Morison, J. H., Nghiem, S. V., and Pratt, K. A.:
590 Snowpack measurements suggest role for multi-year sea ice regions in Arctic atmospheric bromine and chlorine chemistry, *Elem Sci Anth*, 7, 14, 2019.
- Piot, M. and Von Glasow, R.: The potential importance of frost flowers, recycling on snow, and open leads for ozone depletion events, *Atmospheric Chemistry and Physics*, 8, 2437–2467, 2008.
- Platt, U. and Lehrer, E.: Arctic tropospheric ozone chemistry - ARCTOC : results from field, laboratory and modelling studies : final report
595 of the EU project Contract No EV5V-V-CT93-0318(DTEF), Luxembourg, 1997.
- Pratt, K. A., Custard, K. D., Shepson, P. B., Douglas, T. A., Pöhler, D., General, S., Zielcke, J., Simpson, W. R., Platt, U., Tanner, D. J., Huey, L. G., Carlsen, M. S., and Stirm, B. H.: Photochemical production of molecular bromine in Arctic surface snowpacks, *Nature Geoscience*, 6, 351–356, 2013.
- Ridley, B. A., Atlas, E. L., Montzka, D. D., Browell, E. V., Cantrell, C. A., Blake, D. R., Blake, N. J., Cinquini, L., Coffey, M. T., Emons, L. K., Cohen, R. C., DeYoung, R. J., Dibb, J. E., Eisele, F. L., Flocke, F. M., Fried, A., Grahek, F. E., Grant, W. B., Hair, J. W., Hannigan, J. W., Heikes, B. J., Lefer, B. L., Mauldin, R. L., Moody, J. L., Shetter, R. E., Snow, J. A., Talbot, R. W., Thornton, J. A., Walega, J. G., Weinheimer, A. J., Wert, B. P., and Wimmers, A. J.: Ozone depletion events observed in the high latitude surface layer during the TOPSE aircraft program, *Journal of Geophysical Research: Atmospheres*, 108, TOP 4–1–TOP 4–22,
600 <https://doi.org/https://doi.org/10.1029/2001JD001507>, 2003.
- 605 Seinfeld, J. and Pandis, S.: *Atmospheric Chemistry and Physics: from air pollution to climate change*, A Wiley-Interscience publications, Wiley, Hoboken, NJ, USA, 2006.
- Shupe, M. D., Rex, M., Blomquist, B., Persson, P. O. G., Schmale, J., Uttal, T., Althausen, D., Angot, H., Archer, S., Bariteau, L., et al.: Overview of the MOSAiC expedition: *Atmosphere, Elem Sci Anth*, 10, 00 060, 2022.

- Simpson, W., Carlson, D., Hönninger, G., Douglas, T., Sturm, M., Perovich, D., and Platt, U.: First-year sea-ice contact predicts bromine
610 monoxide (BrO) levels at Barrow, Alaska better than potential frost flower contact, *Atmospheric Chemistry and Physics*, 7, 621–627, 2007a.
- Simpson, W. R., Von Glasow, R., Riedel, K., Anderson, P., Ariya, P., Bottenheim, J., Burrows, J., Carpenter, L., Frieß, U., Goodsite, M. E.,
et al.: Halogens and their role in polar boundary-layer ozone depletion, *Atmospheric Chemistry and Physics*, 7, 4375–4418, 2007b.
- Tang, T. and McConnell, J.: Autocatalytic release of bromine from Arctic snow pack during polar sunrise, *Geophysical Research Letters*, 23,
615 2633–2636, 1996.
- Tarasick, D. and Bottenheim, J.: Surface ozone depletion episodes in the Arctic and Antarctic from historical ozonesonde records, *Atmo-
spheric Chemistry and Physics*, 2, 197–205, 2002.
- Thoman, R. L., Moon, T. A., and Druckenmiller, M. L.: NOAA Arctic Report Card 2023: Executive Summary, Technical Report OAR ARC
; 23-01, National Oceanic and Atmospheric Administration, U.S., <https://doi.org/10.25923/5vfa-k694>, noaa:56621, 2023.
- 620 Thomas, J. L., Stutz, J., Lefer, B., Huey, L. G., Toyota, K., Dibb, J. E., and von Glasow, R.: Modeling chemistry in and above snow at Summit,
Greenland – Part 1: Model description and results, *Atmospheric Chemistry and Physics*, 11, 4899–4914, <https://doi.org/10.5194/acp-11-4899-2011>, 2011.
- Thomas, J. L., Dibb, J. E., Huey, L. G., Liao, J., Tanner, D., Lefer, B., von Glasow, R., and Stutz, J.: Modeling chemistry in and above snow
at Summit, Greenland – Part 2: Impact of snowpack chemistry on the oxidation capacity of the boundary layer, *Atmospheric Chemistry
and Physics*, 12, 6537–6554, <https://doi.org/10.5194/acp-12-6537-2012>, 2012.
- 625 Umar, S. A. and Tasduq, S. A.: Ozone layer depletion and emerging public health concerns-an update on epidemiological perspective of the
ambivalent effects of ultraviolet radiation exposure, *Frontiers in Oncology*, 12, 866 733, 2022.
- Weisberg, S.: *Applied linear regression*, vol. 528, John Wiley & Sons, 2005.
- Wennberg, P.: Bromine explosion, *Nature*, 397, 299–301, 1999.
- 630 Yang, X., Pyle, J. A., Cox, R. A., Theys, N., and Van Roozendaal, M.: Snow-sourced bromine and its implications for polar tropospheric
ozone, *Atmospheric Chemistry and Physics*, 10, 7763–7773, <https://doi.org/10.5194/acp-10-7763-2010>, 2010.
- Yang, X., Frey, M. M., Rhodes, R. H., Norris, S. J., Brooks, I. M., Anderson, P. S., Nishimura, K., Jones, A. E., and Wolff, E. W.: Sea
salt aerosol production via sublimating wind-blown saline snow particles over sea ice: parameterizations and relevant microphysical
mechanisms, *Atmospheric Chemistry and Physics*, 19, 8407–8424, <https://doi.org/10.5194/acp-19-8407-2019>, 2019.
- 635 Yang, X., Blechschmidt, A.-M., Bognar, K., McClure-Begley, A., Morris, S., Petropavlovskikh, I., Richter, A., Skov, H., Strong, K., Tarasick,
D. W., Uttal, T., Vestenius, M., and Zhao, X.: Pan-Arctic surface ozone: modelling vs. measurements, *Atmospheric Chemistry and Physics*,
20, 15 937–15 967, <https://doi.org/10.5194/acp-20-15937-2020>, 2020.
- Zhao, X., Strong, K., Adams, C., Schofield, R., Yang, X., Richter, A., Friess, U., Blechschmidt, A.-M., and Koo, J.-H.: A case study of a
transported bromine explosion event in the Canadian high arctic, *Journal of Geophysical Research: Atmospheres*, 121, 457–477, 2016.
- 640 Zilker, B., Richter, A., Blechschmidt, A.-M., von der Gathen, P., Bougoudis, I., Seo, S., Bösch, T., and Burrows, J. P.: Investigation of
meteorological conditions and BrO during ozone depletion events in Ny-Ålesund between 2010 and 2021, *Atmospheric Chemistry and
Physics*, 23, 9787–9814, <https://doi.org/10.5194/acp-23-9787-2023>, 2023.