

Visualising historical changes in air pollution with the Air Quality Stripes

Kirsty J. Pringle^{1,2}, Richard Rigby³, Steven T. Turnock^{4,5}, Carly L. Reddington³, Meruyert Shayakhmetova¹, Malcolm Illingworth¹, Denis Barclay^{1,2}, Neil Chue Hong^{1,2}, Ed Hawkins⁶, Douglas S. Hamilton⁷, Ethan Brain⁷, and James B. McQuaid³

¹EPCC, University of Edinburgh, UK

²Software Sustainability Institute, University of Edinburgh, UK

³School of Earth and Environment, University of Leeds, UK

⁴Met Office Hadley Centre, Exeter, UK

⁵University of Leeds Met Office Strategic (LUMOS) Research Group, University of Leeds, UK

⁶National Centre for Atmospheric Science, Department of Meteorology, University of Reading, Reading, UK

⁷Marine, Earth, and Atmospheric Sciences, North Carolina State University, Raleigh, USA

Correspondence: Kirsty J. Pringle (k.pringle@epcc.ed.ac.uk), James B. McQuaid (J.B.McQuaid@leeds.ac.uk)

Abstract. This paper introduces the Air Quality Stripes, a data visualisation project which presents historical changes in outdoor particulate matter air pollution (PM_{2.5}) concentrations across major cities worldwide. Inspired by the popular Warming Stripes image showing trends in surface temperature, the Air Quality Stripes aim to make complex information about air quality trends understandable and engaging for a broad audience. A historical PM_{2.5} dataset (1850–2022) was created by integrating satellite observations with model simulations (with a bias correction step to ensure a smooth time series and address known model biases). Images were produced in collaboration with a visual design specialist and revised after informal feedback from potential audiences. The images show that trends in PM_{2.5} are varied across the globe; recently there have been significant improvements in air quality in much of Europe and North America but worsening air quality in parts of Asia, Africa and South America. By showcasing historical data in easy to interpret images, the project aims to inspire dialogue among individuals, communities, and policymakers about proactive strategies to combat air pollution.

1 Introduction

Air pollution poses a major public health risk, contributing to 4.7 million premature deaths in 2021, 89% of which occurred in low- and middle-income countries (Institute for Health Metrics and Evaluation, 2021; Brauer et al., 2024; World Health Organization, 2021a). Despite strong evidence of its health impacts, improving public understanding is vital to drive policy changes, individual actions, and address the social and economic inequalities linked to poor air quality. The Air Quality Stripes is a science communication project aimed at raising public awareness and understanding of outdoor air pollution. The images show the change in air quality (PM_{2.5}) through time, from the Industrial Revolution to near present day, in cities around the world. We focus on the air pollutant PM_{2.5} because its health impact surpasses that of other pollutants due to its ability to penetrate vital organs and disrupt physiological processes (Schraufnagel et al., 2019b)

20 The project was inspired by the hugely popular Warming Stripes project (<https://showyourstripes.info/>), led by Prof Ed
Hawkins, which shows the net change in annual mean global mean temperatures as a distinctive series of blue, white and
red stripes (Hawkins et al., 2024, submitted). The Warming Stripes have very successfully translated complex scientific data
into an easily digestible format that resonates with a wide audience (e.g. The Guardian 2021; BBC Future 2023). The success
of the images has inspired other projects including the Biodiversity Stripes (<https://biodiversitystripes.info/global>, Richardson
25 (2023)) and the Ocean Acidification Stripes (<https://oceanacidificationstripes.info/>) .

1.1 Particulate matter air pollution

Air pollution encompasses various harmful substances (e.g. NO₂, O₃, metals, SO₂), but this study focuses on fine particulate
matter (PM_{2.5}), composed of airborne particles smaller than 2.5 micrometers. Exposure to PM_{2.5} is strongly linked to adverse
health effects, including asthma, lung cancer, and heart disease (Schraufnagel et al., 2019a; World Health Organization, 2021c;
30 Wan Mahiyuddin et al., 2023). ~~Its health impact surpasses that of other pollutants due to its ability to penetrate vital organs
and disrupt physiological processes.~~ Over 99% of the world's population resides in areas where PM_{2.5} levels exceed the World
Health Organization's (WHO) air quality guideline value of 5 $\mu\text{g m}^{-3}$ (World Health Organization, 2021a), but concentrations
vary significantly around the globe.

~~PM_{2.5} concentrations are influenced by factors such as industrial and agricultural activity, urbanization, air quality regulation,
35 geographic location and meteorological conditions. In some places natural (or semi-natural) sources of PM_{2.5} such as mineral
dust or black carbon from wildfires strongly influence concentrations (Reddington et al., 2014; Reddington et al., 2016; Reddington et al., 2020).
while other urban locations are affected by strong onshore winds which can influence concentrations (Pillai et al., 2002; Pillai et al., 2010).
PM_{2.5} has an atmospheric lifetime of approximately a few weeks, allowing pollution to travel between nearby cities and
countries, causing transboundary air pollution issues (Zhang et al., 2017; Zhang et al., 2022). However, this relatively short
40 lifetime means that concentrations respond quickly to effective clean air legislation (e.g. Silver et al. 2020).~~

The concentration of PM_{2.5} at any given location is governed by the balance between the magnitude of source terms
(including direct emissions, secondary formation, and transport from other regions) and removal processes (such as deposition
and atmospheric dispersion). Sources of PM_{2.5} are diverse and include combustion from vehicles, residential cooking and
heating, industry, power generation, and agriculture, as well as secondary aerosol formation from gaseous precursors like SO,
45 NO, and VOCs. Natural and semi-natural sources -such as mineral dust, sea salt, volcanic emissions, and black carbon from
wildfires - can also contribute substantially to PM_{2.5} levels (Reddington et al., 2014; Zhang et al., 2016; Graham et al., 2021;
Pai et al., 2022). Meteorological conditions strongly modulate both source and removal terms. Temperature inversions, for
instance, can suppress vertical mixing and trap pollutants near the surface, resulting in acute pollution episodes. Cities situated
in valleys or at the foot of mountain ranges can experience higher concentrations due to restricted air movement (limiting the
50 dispersion of pollutants), while some coastal cities are influenced by strong onshore winds that transport pollutants away (Pillai
et al., 2002; Dall'Osto et al., 2010). Furthermore, PM_{2.5} also displays quite significant seasonal trends due to seasonal changes
in meteorology and emission patterns. PM_{2.5} has an atmospheric lifetime of approximately a few weeks, allowing pollution
to be transported to nearby cities and countries, causing trans-boundary air pollution issues (Zhang et al., 2017; Chen et al.,

2022). At the same time, it means that PM_{2.5} concentrations can respond Relatively quickly - on the order of days to weeks -
55 to effective emissions controls and clean air legislation (e.g., Silver et al. 2020), making policy interventions both impactful
and measurable over short timescales.

2 About the data

2.1 Combined model and satellite data

Satellite observations and computer model simulations were combined to generate a historical PM_{2.5} time series from 1850 to
60 2022. Post-2000 data come from a publicly available product integrating satellite and ground-based observations into a gridded
global dataset at 0.1° resolution (Van Donkelaar et al., 2021). For pre-2000 years, historical trends rely on Earth System Model
simulations (Turnock et al., 2020; Sellar et al., 2020). These simulations are part of the Coupled Model Intercomparison Project
(CMIP6), where over 30 models provided historical climate simulations for 1850–2014 (Eyring et al., 2016), supporting the
Intergovernmental Panel on Climate Change (Arias et al., 2021). All CMIP6 models use the Community Emissions Data
65 System (CEDS) for anthropogenic and wildfire emissions (Hoesly et al., 2018; van Marle et al., 2017), while natural emissions
(dust, sea spray) are model-calculated and vary between models. Only some CMIP6 models include interactive particulate
matter and chemical processes; we used PM_{2.5} data from one of these models UKESM1 (UK Earth System Model, version
1, Sellar et al. 2020). This model uses a two-moment aerosol microphysics scheme for the main types of particulate matter
(sulfate, black carbon, organic carbon, sea salt) and a sectional (bin) scheme for mineral dust (Mulcahy et al., 2020). At present,
70 UKESM1 does not treat nitrate aerosol. Model simulations were at N96 spatial resolution (1.875° longitude × 1.25° latitude).
This model has been widely used for air quality studies (Reddington et al., 2023; Turnock et al., 2020, 2023; Allen et al., 2021;
Butt et al., 2017). CMIP6 data are freely available (Earth System Grid Federation, 2024).

Modelling global air pollutant concentrations is challenging, and models are continually refined through comparisons with
observational data to improve the accuracy of physical and chemical process representations. Previous studies have shown that
75 CMIP6 multimodel simulations tend to underestimate PM_{2.5} concentrations relative to observations (Turnock et al., 2020).
To address this issue and ensure continuity between model output and satellite data, a bias correction approach is applied.
Specifically, for each city, a three-year mean (2000–2002) of satellite-derived PM_{2.5} is compared to a three-year mean of
modelled concentrations for the same period. The ratio between these means is used to adjust the model values throughout the
historical simulation period (1850–2000), correcting for bias. This method is in line with approaches used in previous research
80 (Turnock et al., 2023; Reddington et al., 2023). More information on bias correction techniques is available in Staehle et al.
(2024). The average bias correction factor for all cities is 2.65 (St Dev = 1.6), indicating a general underestimate by the model
compared to satellite-derived observations. There was found to be low sensitivity to the averaging period chosen (e.g. 10 year
average gave a overall bias correction factor of 2.7). Plots showing the model values, satellite-derived values, and bias for
each city are publicly available (Pringle, 2024). Due to the scarcity of historical PM_{2.5} observations, evaluating the accuracy
85 of this approximation is challenging. Observations of PM_{2.5} did not become routine until after the 1990s, prior to that some
measurements of “soot” exist, which can give some indirect indication of PM_{2.5} concentrations, but even these observations




Figure 1. The four styles of image created for each city using London (UK) as an example: a) annual mean with absolute values b) annual mean PM_{2.5} concentrations with indicative descriptions, c) annual mean bar plot and d) summary statistics. Images for the other cities are available on the project website: <https://airqualitystrips.info/>.

are sparse and do not have good temporal or geographical coverage. The concentration data pre 1998 is therefore unconstrained by measurements. However, this method ensures that the historical trend in PM_{2.5} is guided by model simulations (informed by the emissions inventory), while absolute values are aligned with more recent satellite-derived observations with global coverage and higher resolution.

3 Images and Interpretation

Urban concentrations of PM_{2.5} are most relevant to public health as a large fraction of the world's population live in cities, so for the visualisations, PM_{2.5} concentrations were extracted at the location of major cities around the world. An initial set of images for around 170 cities in over 100 countries was created. To allow for different preferences, four different styles of image were created for each city. Figure 1 shows the four images for a single example city (London, UK).

3.1 The Colour of Pollution

The Warming Stripes benefit from an intuitive colour scale (blue is cool, red is warm), which reduces the cognitive load for viewers and makes the plots easy to understand. For air pollution, creating such an intuitive colour palette is more challenging. To bridge the gap between colour and its cognitive associations, it was therefore essential to take into account some of the principles of colour theory. A key aspect of colour theory is understanding the psychological associations between a specific colour, its wider representation in a sociocultural environment, and its representation within the spread of digital media (Christiansen, 2022). For air pollution, creating a simple to understand colour palette is a challenging task due to the often abstract connections between the term “air pollution” and any specific colour. To address this, multimedia designer Ethan Brain (co-author) analysed colour themes from two hundred images collected from a Google image search for “air pollution”. By taking web visuals with the tag “air pollution” and using these images as datasets, visual patterns in popular representation could be formally analysed. As one might expect, polluted images were dominated by reds, browns and greys and clean images by blues. The images were then analysed using a custom sorting and colour grouping tool to identify the dominant colour theme. Then, to hone in on images that best represented air pollution, the initial set of images was filtered only to include results that matched the dominant colours, eliminating outliers and ensuring consistency across the dataset. Dominant colour themes were then extracted from this filtered set and the resulting colours were hand-selected and optimized for web and print colour spaces. Finally, the colours were organized to create a palette that represents increasing values. In the resulting palette the lightest blue represents the cleanest value of concentrations of less than $5 \mu\text{g m}^{-3}$. Cities with years in this colour meet the World Health Organisation Air Quality Guideline (World Health Organization, 2021b), all other colours show an exceedance of the guideline value.

3.2 Refining the images

The Warming Stripes image notably presents the data without the axes and labels typically associated with scientific data representations, instead opting for a clear, compelling visual that bridges art and science. The Air Quality Stripes images differ in one key way: while the Warming Stripes (and the Biodiversity Stripes) primarily aim to highlight a single, consistent global trend - warming temperatures or biodiversity loss - the changes in $\text{PM}_{2.5}$ concentration are more complex and spatially variable. Some regions experience increased concentrations, while others see reductions. It would be challenging to create a single overall image without obscuring these important trends.

However, the project aimed to create accessible and engaging images for a wide audience. ~~To achieve this, draft images were shown to colleagues, friends, and family during an informal testing phase.~~ To achieve this, different versions of draft images were shown to 15 to 20 individuals, including colleagues (air quality researchers, public engagement specialists and researchers in different fields), friends, and family, during an informal testing phase. Feedback was gathered on aspects such as clarity, visual appeal, colour interpretation, and perceived message. Images with and without 1) a trend line, 2) axes labels, and 3) colour bar labels were tested. The trend line was found to be useful in guiding the eye and reducing the effort required to understand the data. Given the varied regional trends, axes and colour bar labels were deemed essential for clarity. Some




Figure 2. Example images for different cities annotated to highlight causes of the change in PM_{2.5} which are used on the landing page of the project website.

individuals expressed uncertainty about interpreting the data, asking questions like, "If concentrations are $10 \mu\text{g m}^{-3}$, what does that mean? Is it good or bad?". To address this, plots were updated with indicative labels ("Very Good", "Fair", "Moderate", "Poor", "Very Poor" and "Extremely Poor"). While similar labels are commonly used (e.g., European Environment Agency 2023), they typically reference daily, not annual, mean values. Instead, the WHO guideline of $5 \mu\text{g m}^{-3}$ was used for the "Very Good" category, and other categories were estimated by extrapolating from daily indicative levels. To enable direct visual comparison the mapping from PM_{2.5} concentration to the indicative label is the same for every city, for more details see Pringle (2025). White breakpoints were added to the colour bar to highlight the separation between these categories. Finally, a landing page was added to the website with images from four cities, annotated to highlight events that affected PM_{2.5} concentrations. Finally it was suggested that we add a "landing page" to the website with images from a small number of cities, to help visitors quickly grasp the format and purpose of the images. For this, images from four cities with large populations but very different trends in PM_{2.5} concentrations were selected and annotated to highlight key factors and events which have affected PM_{2.5} concentrations over different time periods (Figure 2). These images were produced manually with historical emission knowledge required for the annotations, so only a handful of annotated images were produced.

4 Discussion and Conclusion

4.1 Air Pollution Trends

~~The Air Quality Stripes project aimed to highlight the spatial and temporal variability of PM_{2.5} trends across cities worldwide. Historical PM_{2.5} data is scarce, and existing datasets are typically regional rather than city-specific.~~ The images show that PM_{2.5} levels have reduced in many cities. Concentrations in much of Europe and North America have been declining since the 1960s, and very rapid reductions have been achieved in China in the past decade. This offers real hope for public health; PM_{2.5} concentrations can respond rapidly to effective air quality legislation, and many cities have made great strides in reducing these levels. London has lower PM_{2.5} concentrations today than it has had at any point in the past 150 years! Unfortunately, many cities in Central and southern Asia face worsening air pollution. There is an urgent need to address these increasing PM_{2.5} concentrations as huge numbers of people experience high concentrations on a daily basis. The images also show the dominance of natural of PM_{2.5}. Regions with high dust and wildfire emissions (e.g. India, Pakistan and much of northern Africa) may struggle to achieve the WHO recommendations, even with very stringent emission controls (Pai et al., 2022).

It is important to note that these images do not show what is driving the changes in air quality. While some of the progress seen in the Global North has been achieved through local measures such as clean air zones, reductions in coal burning, and stricter air quality legislation, other improvements have occurred through the outsourcing of heavily polluting industries to the Global South. This dynamic has shifted some of the pollution burden from wealthier to poorer regions, contributing to persistent or worsening air quality and health issues in parts of the Global South (Nansai et al., 2020). This disparity underscores environmental and social justice concerns, as communities in lower-income regions disproportionately bear the health and environmental impacts of industrial pollution generated for the benefit of wealthier economies.

4.2 Learnings from Development of the Images

Through the process of creating and refining these images, several valuable insights were gained:

1. **Collaboration with Visual Experts:** Partnering with visual design specialists helped in the development of compelling and accessible graphics. Their expertise in colour theory helped create a final product that was visually appealing and intuitive.
2. **Informal Feedback and Review Stage:** Feedback from colleagues, friends, and family was invaluable. While this project used informal connections, many visualisation efforts could benefit from a structured feedback process to ensure accessibility and clarity for diverse audiences. ~~The visualisations have been heavily used e.g. by the EPIC Air Quality Fund community for engagement and outreach and additional cities have been added on request~~ Any further development of these images or associated works, will involve a formal structured review process which will take feedback from a diverse range of stakeholders
3. **City-Specific Focus:** Creating data for individual cities made the images relatable and grounded. This approach allows the viewer to connect with the information on a more personal level.

175 4. **Selected Annotations:** Narrative annotations on a subset of images made the data more relatable, providing context and highlighting significant points. They also helped viewers better understand the overall structure of the images.

The Air Quality Stripes project received significant media attention (Fuller, 2024; Hunter, 2024; Turns, 2024; McQuaid et al., 2024), highlighting the power of visual data to engage and inform the public. Future efforts will aim to deepen this engagement by using these images to facilitate discussions about lived experiences with air pollution worldwide, through methods such as workshops, storytelling, or other participatory approaches.

180 *Code and data availability.* The data used to create the Air Quality Stripes is available (Pringle and McQuaid, 2024) as is the visualisation code used (Pringle and Rigby, 2024).

Author contributions. JM and KP led the project with scientific guidance from CR and ST, software support from RR and MI and visualisation guidance from DH, EB, EH, MS and DB.

Competing interests. At least one of the authors is a member of the editorial board of Geoscience Communication.

185 *Ethical Statement* This article did not contain any studies involving human or animal subjects and did not need to undergo ethical review.

Acknowledgements. KP, DB and NCH were supported by UKRI grants EP/S021779/1 and AH/Z000114/1 for the Software Sustainability Institute. This work (JM) was supported by the Natural Environment Research Council [NE/T010401/1]. KP and MI are grateful for two internal grants from EPCC, University of Edinburgh. EH is supported by the National Centre for Atmospheric Science and the Ireland-UK
190 Co-Centre for Climate + Biodiversity + Water. MS was supported through Access to Industry's Access to Data program. The contributions of ST were funded by the Met Office Climate Science for Service Partnership (CSSP) China project under the International Science Partnerships Fund (ISPF). This work used JASMIN, the UK's collaborative data analysis environment (<https://www.jasmin.ac.uk>, Lawrence et al. (2013)).

References

- Allen, R. J., Horowitz, L. W., Naik, V., Oshima, N., O'Connor, F. M., Turnock, S., Shim, S., Sager, P. L., van Noije, T., Tsigaridis, K.,
195 Bauer, S. E., Sentman, L. T., John, J. G., Broderick, C., Deushi, M., Folberth, G. A., Fujimori, S., and Collins, W. J.: Significant
climate benefits from near-term climate forcer mitigation in spite of aerosol reductions, *Environmental Research Letters*, 16, 034010,
<https://doi.org/10.1088/1748-9326/abe06b>, 2021.
- Arias, P., Bellouin, N., Coppola, E., Jones, R., Krinner, G., Marotzke, J., Naik, V., Palmer, M., Plattner, G.-K., Rogelj, J., Rojas, M., Sillmann,
J., Storelvmo, T., Thorne, P., Trewin, B., Achuta Rao, K., Adhikary, B., Allan, R., Armour, K., Bala, G., Barimalala, R., Berger, S.,
200 Canadell, J., Cassou, C., Cherchi, A., Collins, W., Collins, W., Connors, S., Corti, S., Cruz, F., Dentener, F., Dereczynski, C., Di Luca,
A., Diongue Niang, A., Doblas-Reyes, F., Dosio, A., Douville, H., Engelbrecht, F., Eyring, V., Fischer, E., Forster, P., Fox-Kemper, B.,
Fuglestad, J., Fyfe, J., Gillett, N., Goldfarb, L., Gorodetskaya, I., Gutierrez, J., Hamdi, R., Hawkins, E., Hewitt, H., Hope, P., Islam,
A., Jones, C., Kaufman, D., Kopp, R., Kosaka, Y., Kossin, J., Krakovska, S., Lee, J.-Y., Li, J., Mauritsen, T., Maycock, T., Meinshausen,
M., Min, S.-K., Monteiro, P., Ngo-Duc, T., Otto, F., Pinto, I., Pirani, A., Raghavan, K., Ranasinghe, R., Ruane, A., Ruiz, L., Sallée, J.-B.,
205 Samset, B., Sathyendranath, S., Seneviratne, S., Sörensson, A., Szopa, S., Takayabu, I., Tréguier, A.-M., van den Hurk, B., Vautard, R.,
von Schuckmann, K., Zaehle, S., Zhang, X., and Zickfeld, K.: Technical Summary, p. 33-144, Cambridge University Press, Cambridge,
United Kingdom and New York, NY, USA, <https://doi.org/10.1017/9781009157896.002>, 2021.
- BBC Future: The coloured stripes that explain climate change, [https://www.bbc.com/future/article/](https://www.bbc.com/future/article/20231206-the-coloured-stripes-that-explain-climate-change)
20231206-the-coloured-stripes-that-explain-climate-change, accessed: 2024-11-21, 2023.
- 210 Brauer, M., Roth, G. A., Aravkin, A. Y., Zheng, P., Abate, K. H., Abate, Y. H., Abbafati, C., Abbasgholizadeh, R., Abbasi, M. A., Abbasian,
M., et al.: Global burden and strength of evidence for 88 risk factors in 204 countries and 811 subnational locations, 1990–2021: a
systematic analysis for the Global Burden of Disease Study 2021, *The Lancet*, 403, 2162–2203, 2024.
- Butt, E. W., Turnock, S. T., Rigby, R., Reddington, C. L., Yoshioka, M., Johnson, J. S., Regayre, L. A., Pringle, K. J., Mann, G. W., and
Spracklen, D. V.: Global and regional trends in particulate air pollution and attributable health burden over the past 50 years, *Environmental*
215 *Research Letters*, 12, 104017, <https://doi.org/10.1088/1748-9326/aa87be>, 2017.
- Chen, L., Lin, J., Martin, R., Du, M., Weng, H., Kong, H., Ni, R., Meng, J., Zhang, Y., Zhang, L., and van Donke-
laar, A.: Inequality in historical transboundary anthropogenic PM_{2.5} health impacts, *Science Bulletin*, 67, 437–444,
<https://doi.org/https://doi.org/10.1016/j.scib.2021.11.007>, 2022.
- Christiansen, J.: Building Science Graphics: An Illustrated Guide to Communicating Science through Diagrams and Visualizations, CRC
220 Press, ISBN 9781032106748, 2022.
- Dall'Osto, M., Ceburnis, D., Martucci, G., Bialek, J., Dupuy, R., Jennings, S. G., Berresheim, H., Wenger, J., Healy, R., Facchini, M. C.,
Rinaldi, M., Giulianelli, L., Finessi, E., Worsnop, D., Ehn, M., Mikkilä, J., Kulmala, M., and O'Dowd, C. D.: Aerosol properties associated
with air masses arriving into the North East Atlantic during the 2008 Mace Head EUCAARI intensive observing period: an overview,
Atmospheric Chemistry and Physics, 10, 8413–8435, <https://doi.org/10.5194/acp-10-8413-2010>, 2010.
- 225 Earth System Grid Federation: Earth System Grid Federation (ESGF), <https://esgf.github.io/index.html>, accessed: 2024-11-05, 2024.
- European Environment Agency: European Air Quality Index, <https://airindex.eea.europa.eu/AQI/index.html>, accessed: 2023-10-25, 2023.
- Eyring, V., Bony, S., Meehl, G. A., Senior, C. A., Stevens, B., Stouffer, R. J., and Taylor, K. E.: Overview of the Coupled Model
Intercomparison Project Phase 6 (CMIP6) experimental design and organization, *Geoscientific Model Development*, 9, 1937–1958,
<https://doi.org/10.5194/gmd-9-1937-2016>, 2016.

- 230 Fuller, G.: Can climate stripes change the way we think about air pollution?, *The Guardian*, <https://www.theguardian.com/environment/article/2024/aug/23/can-climate-stripes-change-the-way-we-think-about-air-pollution>, accessed: 2024-11-05, 2024.
- Graham, A. M., Pringle, K. J., Pope, R. J., Arnold, S. R., Conibear, L. A., Burns, H., and et al.: Impact of the 2019/2020 Australian megafires on air quality and health, *GeoHealth*, 5, e2021GH000454, <https://doi.org/10.1029/2021GH000454>, 2021.
- Hoesly, R. M., Smith, S. J., Feng, L., Klimont, Z., Janssens-Maenhout, G., Pitkanen, T., Seibert, J. J., Vu, L., Andres, R. J., Bolt, R. M., Bond,
235 T. C., Dawidowski, L., Kholod, N., Kurokawa, J.-i., Li, M., Liu, L., Lu, Z., Moura, M. C. P., O'Rourke, P. R., and Zhang, Q.: Historical (1750–2014) anthropogenic emissions of reactive gases and aerosols from the Community Emissions Data System (CEDS), *Geoscientific Model Development*, 11, 369–408, <https://doi.org/10.5194/gmd-11-369-2018>, 2018.
- Hunter, W.: Charts reveal deadly pollution in cities, *Daily Mail*, <https://www.dailymail.co.uk/sciencetech/article-13772213/charts-reveal-deadly-pollution-cities.html>, accessed: 2024-11-05, 2024.
- 240 Institute for Health Metrics and Evaluation: GBD Results Tool, Institute for Health Metrics and Evaluation, <https://vizhub.healthdata.org/gbd-results/>, search terms: Year: 2021, Risk: Ambient Particulate Matter Pollution, Location: Global, Cause: All Causes, Sex: Both. Accessed: 2024-11-18, 2021.
- Lawrence, B. N., Bennett, V. L., Churchill, J., Juckes, M., Kershaw, P., Pascoe, S., Pepler, S., Pritchard, M., and Stephens, A.: Storing and manipulating environmental big data with JASMIN, in: *Proceedings of the IEEE Big Data Conference*, San Francisco, CA, USA, 2013.
- 245 McQuaid, J., Pringle, K., and Illingworth, S.: These colourful diagrams show how air quality has changed in over 100 countries around the world since 1850, *The Conversation*, <https://theconversation.com/these-colourful-diagrams-show-how-air-quality-has-changed-in-over-100-countries-around-the-world-since-1850-237079>, accessed: 2024-11-05, 2024.
- Mulcahy, J. P., Johnson, C., Jones, C. G., Povey, A. C., Scott, C. E., Sellar, A., Turnock, S. T., Woodhouse, M. T., Abraham, N. L., Andrews,
250 M. B., Bellouin, N., Browse, J., Carslaw, K. S., Dalvi, M., Folberth, G. A., Glover, M., Grosvenor, D. P., Hardacre, C., Hill, R., Johnson, B., Jones, A., Kipling, Z., Mann, G., Mollard, J., O'Connor, F. M., Palmiéri, J., Reddington, C., Rumbold, S. T., Richardson, M., Schutgens, N. A. J., Stier, P., Stringer, M., Tang, Y., Walton, J., Woodward, S., and Yool, A.: Description and evaluation of aerosol in UKESM1 and HadGEM3-GC3.1 CMIP6 historical simulations, *Geoscientific Model Development*, 13, 6383–6423, <https://doi.org/10.5194/gmd-13-6383-2020>, 2020.
- 255 Nansai, K., Tohno, S., Chatani, S., Kanemoto, K., Kurogi, M., Fujii, Y., Kagawa, S., Kondo, Y., Nagashima, F., Takayanagi, W., and Lenzen, M.: Affluent countries inflict inequitable mortality and economic loss on Asia via PM_{2.5} emissions, *Environment International*, 134, 105238, <https://doi.org/https://doi.org/10.1016/j.envint.2019.105238>, 2020.
- Pai, S. J., Carter, T. S., Heald, C. L., and Kroll, J. H.: Updated World Health Organization Air Quality Guidelines Highlight the Importance of Non-anthropogenic PM_{2.5}, *Environmental Science & Technology Letters*, 9, 501–506, <https://doi.org/10.1021/acs.estlett.2c00203>, 2022.
- 260 Pillai, P. S., Suresh Babu, S., and Krishna Moorthy, K.: A study of PM, PM₁₀ and PM_{2.5} concentration at a tropical coastal station, *Atmospheric Research*, 61, 149–167, [https://doi.org/https://doi.org/10.1016/S0169-8095\(01\)00136-3](https://doi.org/https://doi.org/10.1016/S0169-8095(01)00136-3), 2002.
- Pringle, K.: Air Quality Stripes: Comparison of Model and Satellite Values, <https://doi.org/10.5281/zenodo.14392693>, 2024.
- Pringle, K.: Air Quality Stripes: Mapping Annual Mean PM_{2.5} Values to Indicative Labels, <https://doi.org/10.5281/zenodo.15363039>, 2025.
- Pringle, K. and McQuaid, J.: Air Quality Stripes (V1pt6), <https://doi.org/10.5281/zenodo.14360345>, 2024.
- 265 Pringle, K. and Rigby, R.: Air Quality Stripes: Code used to create images (V1pt6_b), <https://doi.org/10.5281/zenodo.14393148>, 2024.

- Reddington, C. L., Yoshioka, M., Balasubramanian, R., Ridley, D., Toh, Y. Y., Arnold, S. R., and Spracklen, D. V.: Contribution of vegetation and peat fires to particulate air pollution in South-east Asia, *Environmental Research Letters*, 9, 094006, <https://doi.org/10.1088/1748-9326/9/9/094006>, 2014.
- Reddington, C. L., Turnock, S. T., Conibear, L., Forster, P. M., Lowe, J. A., Ford, L. B., et al.: Inequalities in air pollution exposure and attributable mortality in a low carbon future, *Earth's Future*, 11, e2023EF003697, <https://doi.org/10.1029/2023EF003697>, 2023.
- Richardson, M.: GC Insights: Nature stripes for raising engagement with biodiversity loss, *Geoscience Communication*, 6, 11–14, <https://doi.org/10.5194/gc-6-11-2023>, 2023.
- Schraufnagel, D. E., Balmes, J. R., Cowl, C. T., De Matteis, S., Jung, S. H., Mortimer, K., Perez-Padilla, R., Rice, M. B., Riojas-Rodriguez, H., Sood, A., Thurston, G. D., To, T., Vanker, A., and Wuebbles, D. J.: Air Pollution and Noncommunicable Diseases: A Review by the Forum of International Respiratory Societies' Environmental Committee, Part 1: The Damaging Effects of Air Pollution, *Chest*, 155, 409–416, <https://doi.org/10.1016/j.chest.2018.10.042>, epub 2018 Nov 9, 2019a.
- Schraufnagel, D. E., Balmes, J. R., Cowl, C. T., De Matteis, S., Jung, S. H., Mortimer, K., Perez-Padilla, R., Rice, M. B., Riojas-Rodriguez, H., Sood, A., Thurston, G. D., To, T., Vanker, A., and Wuebbles, D. J.: Air Pollution and Noncommunicable Diseases: A Review by the Forum of International Respiratory Societies' Environmental Committee, Part 2: Air Pollution and Organ Systems, *Chest*, 155, 417–426, <https://doi.org/10.1016/j.chest.2018.10.041>, epub 2018 Nov 9, 2019b.
- Sellar, A. A., Walton, J., Jones, C. G., Wood, R., Abraham, N. L., Andrejczuk, M., Andrews, M. B., Andrews, T., Archibald, A. T., de Mora, L., Dyson, H., Elkington, M., Ellis, R., Florek, P., Good, P., Gohar, L., Haddad, S., Hardiman, S. C., Hogan, E., Iwi, A., Jones, C. D., Johnson, B., Kelley, D. I., Kettleborough, J., Knight, J. R., Köhler, M. O., Kuhlbrodt, T., Liddicoat, S., Linova-Pavlova, I., Mizieliński, M. S., Morgenstern, O., Mulcahy, J., Neining, E., O'Connor, F. M., Petrie, R., Ridley, J., Rioual, J.-C., Roberts, M., Robertson, E., Rumbold, S., Seddon, J., Shepherd, H., Shim, S., Stephens, A., Teixeira, J. C., Tang, Y., Williams, J., Wiltshire, A., and Griffiths, P. T.: Implementation of U.K. Earth System Models for CMIP6, *Journal of Advances in Modeling Earth Systems*, <https://doi.org/10.1029/2019MS001946>, 2020.
- Silver, B., Conibear, L., Reddington, C. L., Knute, C., Arnold, S. R., and Spracklen, D. V.: Pollutant emission reductions deliver decreased PM_{2.5}-caused mortality across China during 2015–2017, *Atmospheric Chemistry and Physics*, 20, 11683–11695, <https://doi.org/10.5194/acp-20-11683-2020>, 2020.
- Staehle, C., Rieder, H. E., Fiore, A. M., and Schnell, J. L.: Technical note: An assessment of the performance of statistical bias correction techniques for global chemistry–climate model surface ozone fields, *Atmospheric Chemistry and Physics*, 24, 5953–5969, <https://doi.org/10.5194/acp-24-5953-2024>, 2024.
- The Guardian: The statement shirts making climate data fashionable at COP26, <https://www.theguardian.com/environment/2021/nov/05/the-statement-shirts-making-climate-data-fashionable-at-cop26-aoe>, accessed: 2024-11-21, 2021.
- Turnock, S. T., Allen, R. J., Andrews, M., Bauer, S. E., Deushi, M., Emmons, L., Good, P., Horowitz, L., John, J. G., Michou, M., Nabat, P., Naik, V., Neubauer, D., O'Connor, F. M., Olivié, D., Oshima, N., Schulz, M., Sellar, A., Shim, S., Takemura, T., Tilmes, S., Tsigaridis, K., Wu, T., and Zhang, J.: Historical and future changes in air pollutants from CMIP6 models, *Atmospheric Chemistry and Physics*, 20, 14547–14579, <https://doi.org/10.5194/acp-20-14547-2020>, 2020.
- Turnock, S. T., Reddington, C. L., West, J. J., and O'Connor, F. M.: The air pollution human health burden in different future scenarios that involve the mitigation of near-term climate forcers, climate and land-use, *GeoHealth*, 7, e2023GH000812, <https://doi.org/10.1029/2023GH000812>, 2023.

- Turns, A.: The new climate stripes: how creativity can inspire environmental action, *The Conversation*, <https://theconversation.com/the-new-climate-stripes-how-creativity-can-inspire-environmental-action-237350>, accessed: 2024-11-05, 2024.
- 305 Van Donkelaar, A., Hammer, M. S., Bindle, L., Brauer, M., Brook, J. R., Garay, M. J., Hsu, N. C., Kalashnikova, O. V., Kahn, R. A., Lee, C., Levy, R. C., Lyapustin, A., Sayer, A. M., and Martin, R. V.: Monthly Global Estimates of Fine Particulate Matter and Their Uncertainty, *Environmental Science Technology*, 55, 15 287–15 300, <https://doi.org/10.1021/acs.est.1c05309>, 2021.
- van Marle, M. J. E., Kloster, S., Magi, B. I., Marlon, J. R., Daniau, A.-L., Field, R. D., Arneth, A., Forrest, M., Hantson, S., Kehrwald, N. M., Knorr, W., Lasslop, G., Li, F., Mangeon, S., Yue, C., Kaiser, J. W., and van der Werf, G. R.: Historic global biomass burning emissions for CMIP6 (BB4CMIP) based on merging satellite observations with proxies and fire models (1750–2015), *Geoscientific Model Development*, 10, 3329–3357, <https://doi.org/10.5194/gmd-10-3329-2017>, 2017.
- 310 Wan Mahiyuddin, W. R., Ismail, R., Mohammad Sham, N., Ahmad, N. I., and Nik Hassan, N. M. N.: Cardiovascular and Respiratory Health Effects of Fine Particulate Matters (PM_{2.5}): A Review on Time Series Studies, *Atmosphere*, 14, <https://doi.org/10.3390/atmos14050856>, 2023.
- 315 World Health Organization: Ambient (outdoor) air quality and health, [https://www.who.int/news-room/fact-sheets/detail/ambient-\(outdoor\)-air-quality-and-health](https://www.who.int/news-room/fact-sheets/detail/ambient-(outdoor)-air-quality-and-health), accessed: 2024-10-25, 2021a.
- World Health Organization: WHO Global Air Quality Guidelines. Particulate Matter (PM_{2.5} and PM₁₀), Ozone, Nitrogen Dioxide, Sulfur Dioxide and Carbon Monoxide, <https://www.who.int/publications/i/item/9789240034228>, licence: CC BY-NC-SA 3.0 IGO. Accessed: 2024-11-21, 2021b.
- 320 World Health Organization: Review of evidence on health aspects of air pollution: REVIHAAP project: technical report, Tech. rep., World Health Organization. Regional Office for Europe, <https://iris.who.int/handle/10665/341712>, 2021c.
- Zhang, Q., Jiang, X., Tong, D., and et al.: Transboundary health impacts of transported global air pollution and international trade, *Nature*, 543, 705–709, <https://doi.org/10.1038/nature21712>, 2017.
- Zhang, X., Zhao, L., Tong, D. Q., Wu, G., Dan, M., and Teng, B.: A Systematic Review of Global Desert Dust and Associated Human Health Effects, *Atmosphere*, 7, 158, <https://doi.org/10.3390/atmos7120158>, 2016.
- 325