

1 **Drivers of the spatiotemporal distribution of dissolved**
2 **nitrous oxide and air-sea exchange in a coastal**
3 **Mediterranean area**

4 Susana Flecha¹, Mercedes de la Paz², Fiz Fernández Pérez², Núria Marbà³, Carlos
5 Morell³, Eva Alou-Font⁴, Joaquín Tintoré^{3,5} and Iris E. Hendriks³

6 ¹Instituto de Ciencias Marinas de Andalucía (ICMAN-CSIC), Puerto Real, Cádiz, Spain.

7 ²Instituto de Investigaciones Marinas (CSIC), Vigo, Spain

8 ³Mediterranean Institute for Advanced Studies (CSIC-UIB), Esporles, Spain

9 ⁴King Abdullah University of Science and Technology (KAUST), Thuwal, Kingdom of Saudi Arabia.

10 ⁵Balearic Islands Coastal Observing and Forecasting System (SOCIB), Palma, Spain

11 *Correspondence to:* Susana Flecha (susana.flecha@csic.es)

12 **Abstract.** Among the well-known greenhouse gases (GHG), nitrous oxide (N₂O) is the third most
13 impactful, possessing a global warming potential approximately 300 times greater than ~~that of~~ carbon
14 dioxide (CO₂) over a century. The distribution of N₂O ~~in~~ aquatic environments exhibits notable spatial and
15 temporal variations, and emissions ~~remain~~ inadequately constrained and underrepresented in global N₂O
16 emission inventories, particularly from coastal zones. This study focuses on ~~N₂O~~ levels and air-sea fluxes
17 in ~~the coastal waters of~~ the Balearic Islands Archipelago ~~in~~ the Western Mediterranean Basin. Data were
18 gathered between 2018 and 2023 at three coastal monitoring stations: two in the ~~densely populated~~ island
19 of Mallorca and the third in the well-preserved National Park of the Cabrera Archipelago. Seawater N₂O
20 concentrations varied from 6.5 to 9.9 ~~nmol L⁻¹~~, with no significant differences detected across the sites.
21 ~~When these sink/source strengths are integrated on an annual basis, the Balearic Sea is close to equilibrium~~
22 ~~with atmospheric N₂O, resulting in a neutral atmosphere-ocean exchange (0.1±0.2 μmol m⁻² d⁻¹).~~ A
23 consistent seasonal pattern was noted ~~during~~ the study period. Machine learning analysis indicated that
24 seawater temperature was the primary factor influencing N₂O concentrations, with lesser contributions from
25 chlorophyll levels and salinity.

26
27 **1 Introduction**

28 Nitrous oxide (N₂O) is a potent greenhouse gas (GHG) with 300 times higher warming potential per mole
29 than carbon dioxide (CO₂) on a 100-year time scale (Solomon et al., 2007). Atmospheric N₂O levels have
30 ~~risen~~ by more than 18% since preindustrial times and increased by 332 ppb between 2011 and 2019
31 (Masson-Delmotte et al., 2021). The ocean N₂O budget is highly sensitive to climate changes and
32 significantly influences the climate system (~~Ravishankara et al., 2009~~). Variations in temperature, ocean
33 circulation, and biological activity can alter the production and release of N₂O from the oceans.
34 ~~Consequently~~, the concentration of ~~N₂O~~ in the atmosphere ~~influences~~ global warming and climate patterns,
35 creating a feedback loop between ~~oceanic~~ processes and the climate system. Marine N₂O sources ~~account~~
36 ~~for one-third~~ of the natural emissions to the atmosphere, yielding a net source of 3.5 (2.5-4.7) Tg N yr⁻¹.
37 ~~excluding~~ coastal contributions (Tian et al., 2024). N₂O is ~~primarily~~ produced ~~through~~ nitrification and
38 denitrification pathways (Freing et al., 2012). ~~Nitrification is an aerobic process in which ammonium~~
39 ~~(NH₄⁺) is oxidized to nitrate (NO₃⁻), producing N₂O as a byproduct, favored by low-oxygen conditions. In~~
40 ~~contrast, denitrification occurs under suboxic to anoxic conditions and involves the stepwise reduction of~~

Con formato: Color de fuente: Texto 1

Eliminado: ¶

Eliminado: from

Eliminado: still

Eliminado: the

Eliminado: coastal waters

Eliminado: high

Eliminado: inhabi

Eliminado: nM

Con formato: Color de fuente: Texto 1

Con formato: Color de fuente: Texto 1

Eliminado: The average air-sea fluxes were estimated to range from -0.3 to 0.6 μmol m⁻² d⁻¹, indicating that the study areas generally functioned as weak N₂O sources.

Eliminado: over

Con formato: Espacio Antes: 0 pto, Después: 0 pto, No agregar espacio entre párrafos del mismo estilo

Eliminado: increased

Eliminado: In turn

Eliminado: nitrous oxide

Eliminado: affect

Eliminado: the

Eliminado: represent

Eliminado: a

Eliminado:

Eliminado: part

Eliminado: 5-4.

Eliminado: ⁻¹

Eliminado: without

Eliminado: considering the

Eliminado: mainly by

Con formato: Derecha: 0.63 cm

67 nitrate to dinitrogen gas (N₂), also releasing N₂O as an intermediate. These microbial processes are
 68 especially sensitive to environmental gradients in oxygen, nutrients, and organic matter availability. The
 69 nitrogen cycle is one of the most complex regulating factors of primary production, highly dependent on
 70 dissolved oxygen concentration and the prevailing redox conditions (Codispoti, 2010). In coastal
 71 environments, significant variability exists in the nitrogen cycle, where land-derived nutrient inputs, coastal
 72 upwelling events, and complex biogeochemical processes play crucial roles (Doney, 2010), contributing
 73 significantly to the spatiotemporal variability of the N₂O concentrations (Nevison et al., 1995). In estuarine
 74 and coastal waters, the effects of climate change may be more pronounced, such as ocean acidification (OA,
 75 Carstensen and Duarte, 2019), which can enhance the generation of N₂O (Wan et al., 2023; Zhou et al.,
 76 2023). Despite the necessity for a better understanding of atmospheric and oceanic inventories of non-CO₂
 77 GHGs to provide realistic and accurate models for future scenarios, there are limited open ocean and coastal
 78 monitoring time series networks compared to CO₂ (Bakker et al., 2014; de la Paz et al., 2015; Fariás et al.,
 79 2007; Ma et al., 2019; Wilson et al., 2017).

80 The Mediterranean Sea is a semi-enclosed basin surrounded by highly sensitive coastal zones particularly
 81 vulnerable to human activities. Factors such as high population density, widespread urbanization, and
 82 intensive agriculture have escalated risks of pollution and habitat degradation in the region (UNEP/MAP,
 83 2012). Due to its distinct biogeochemical and hydrodynamic features, this basin has been identified as a
 84 “hotspot” for climate change research (Giorgi, 2006). The impacts of global warming and extreme weather
 85 events are expected to be more severe in the Mediterranean compared to other oceanic regions (Giorgi,
 86 2006; Giorgi and Lionello, 2008; Masson-Delmotte et al., 2021). Despite representing just 0.82% of the
 87 global ocean surface, the Mediterranean hosts 4-18% of the world's marine biodiversity, including
 88 numerous endemic species (Bianchi and Morri, 2000; Mouillot et al., 2011). Rising temperatures and OA
 89 threaten the biodiversity of the region (Micheli et al., 2013). Additionally, anthropogenic pressures along
 90 the Mediterranean coast have intensified due to rapid population growth and economic activities. In the
 91 Western Mediterranean, high tourism and coastal development levels have left only a small fraction of the
 92 coastline in a natural state, with even fewer areas under protection (EEA, 1999).

93 The Balearic Islands Archipelago, located in the Western Mediterranean, comprises the islands of Mallorca,
 94 Menorca, Ibiza, and Formentera, with a combined coastline of 1,723 km. Renowned as a major European
 95 tourist destination, tourism accounts for approximately 45% of the total Gross Domestic Product of the
 96 archipelago. Visitor numbers have surged dramatically over the past century, reaching nearly 18 million in
 97 2023 (Institut d'Estadística de les Illes Balears, Spain), compared to a resident population of around 1.2
 98 million. Coastal ecosystems in the Balearic Islands are vital for the local economy. Meadows of the endemic
 99 seagrass *Posidonia oceanica* extend across depths of up to 45 m in the Balearic Sea, providing critical
 100 ecosystem services such as carbon sequestration (Duarte et al., 2005), oxygen production (Hendriks et al.,
 101 2022), biodiversity support, coastal erosion prevention, sediment stabilization, and water transparency
 102 (Barbier et al., 2011). However, these ecosystems face increasing pressure from recreational activities and
 103 other anthropogenic impacts.

104 Given the increasing threats to these ecosystems, understanding the relationship between anthropogenic
 105 pressures and GHG emissions, particularly N₂O, has become urgent. The absence of long-term N₂O datasets
 106 in the Mediterranean Sea, along with the uncertainties surrounding current emissions estimates, underscores

Eliminado: regime

Eliminado: there is considerable

Eliminado: solid

Eliminado: important

Eliminado: severe

Eliminado: which also stimulates the generation of N₂O

Eliminado: ¶

Con formato: No agregar espacio entre párrafos del mismo estilo

Eliminado: '

Eliminado: ocean acidification

Con formato: Color de fuente: Texto 1

Eliminado: nsist

Con formato: Color de fuente: Texto 1

Eliminado: contributes

Con formato: Color de fuente: Texto 1

Eliminado: risen

Con formato: Color de fuente: Texto 1

Eliminado: essenti

Con formato: Color de fuente: Texto 1

Eliminado: mount

Con formato: Color de fuente: Texto 1

Eliminado: greenhouse gas

Eliminado: nitrous oxide (

Eliminado:)

Eliminado: ,

Con formato: Color de fuente: Texto 1

Eliminado: combined

Con formato: Color de fuente: Texto 1

Con formato: Derecha: 0.63 cm

the importance of assessing coastal areas with varying human impacts. These evaluations are essential for improving coastal N₂O emissions estimates and refining global ocean N₂O budgets. In this study, we evaluate the spatial and temporal N₂O concentrations in surface waters and the air-sea exchange in the coastal area of the Balearic Islands Archipelago and estimate the potential drivers of the observed variability in N₂O concentrations. We focused on three different sites in the coastal zone: a highly impacted site, a medium-impacted site, both located near the island of Mallorca, and a pristine station in the Cabrera National Park Archipelago.

2 Methods

2.1 Study area

We collected physicochemical and biogeochemical parameters from three stations in the Balearic Sea within the Western Mediterranean Basin (Fig. 1A), which are part of the Balearic Ocean Acidification Time Series (Flecha et al., 2022). The data presented here was collected from 2018 to 2023; however, data availability varies among the different stations.

Two sampling sites are fixed monitoring stations equipped with autonomous sensors. The first station, established in December 2018, is located in the Bay of Palma (PB: 39.492848°N, 2.700405°E, at a depth of approximately 30 m, Figure 1B) and is part of the fixed monitoring network of the Balearic Islands Coastal Observing and Forecasting System (Tintoré et al., 2019; 2013- SOCIB; <https://www.socib.es/>). Temperature (°C) and salinity (PSU) data were obtained from the SOCIB buoy sensors; see Tintoré and Casas (2022) for details about the sensors. Additionally, a MiniDot sensor (PME, Inc[®]) recorded DO hourly, with a manufacturer accuracy of ± 5%. The sensor location and water sampling depth were set at 1 m.

The second fixed monitoring station is located in the Bay of Santa Maria (Fig. 1B) within the Cabrera Archipelago National Park (CA: 39.151395° N, 2.950823° E, ~8 m depth) and was installed in November 2019. It is situated in an area under governmental protection and is regarded as a pristine site with no apparent human influence. Temperature (°C), salinity (PSU), and DO data were collected hourly, using SBE37-SMP-ODO (Sea-Bird Scientific Electronics[®]) and a MiniDot. Both sensors were secured to a mooring line at approximately 4 m depth, and water samples were taken at the same depth. The manufacturer accuracy of measurements was ± 0.002°C, 0.003 mS/cm, and ± 5% for temperature, conductivity, and oxygen sensors, respectively.

At PB and CA stations, samples of dissolved N₂O, dissolved oxygen (DO), dissolved organic carbon (DOC), Chlorophyll *a* (Chl *a*), inorganic nutrients, including nitrate (NO₃⁻), nitrite (NO₂⁻), phosphate (PO₄³⁻), silicate (Si (OH)₄), ammonia (NH₄), and Total Nitrogen (TN) were collected monthly.

The third sampling point is located in the coastal area near the Cape Ses Salines lighthouse (CS: 39.2649° N, 3.0535° E, Figure 1B). At this site, with a total bottom depth of 2 m, data were collected biweekly from surface water directly off the coast at approximately 0.5 m depth. Sampling began in August 2018. Temperature (°C), salinity (PSU), N₂O, DOC, Chl *a*, and inorganic nutrients were collected from the same

Eliminado: variability of ...2 ... [1]

Con formato: Color de fuente: Texto 1

Eliminado: N₂O

Con formato: Color de fuente: Texto 1

Con formato: Color de fuente: Texto 1

Eliminado: variability

Con formato: Color de fuente: Texto 1

Eliminado: and

Con formato: Color de fuente: Texto 1

Con formato: Color de fuente: Texto 1

Eliminado: the ...three stations located ... [2]

Eliminado: integrated...hich into ... [3]

Eliminado: with...quipped deployed ... [4]

Eliminado: bay ...ay of Palma (BP...B: 39.492848°N, 2.700405°E, over...t a bottom ...epth of ~ ... [5]

Eliminado: i...s part of the fixed monitoring station...etwork belonging to... ... [6]

Eliminado: ...019, ... [7]

Eliminado: PSU

Con formato: Color de fuente: Texto 1

Eliminado: sensor...etails detail ... [8]

Eliminado: ... The...ith a manufacturer accuracy of the DO measurements was ... [9]

Eliminado: loc

Eliminado: ...n ...n area under governmental protection and is considered... ... [10]

Eliminado: PSU

Eliminado: obtain...ollected hourly by ... [11]

Con formato: Color de fuente: Texto 1

Eliminado: attach

Eliminado: round

Eliminado: S...amples for...f dissolved nitrous oxide (...2 ... [12]

Con formato ... [13]

Eliminado: and Total Phosphorous (TP) and ...otal Nitrogen (TN) were collected monthly...onthly from the same depth as the sensors of the fixed stations ... [14]

Eliminado: loc

Eliminado: site

Eliminado: PSU

Con formato

Eliminado: obtain

Con formato: Derecha: 0.63 cm

242 volume of surface water. DO data were gathered using a MiniDot sensor starting in August 2018. Validation
243 of DO sensor data was performed with DO water samples from PB and CA stations, as detailed in Agueda-
244 Aramburu et al. (2024).

245 The three sites differ in their physical setting and benthic characteristics. PB is located in an open bay with
246 moderate anthropogenic influence and limited seagrass coverage within the sampling area. CA, on the other
247 hand, is located in a semi-enclosed bay within a protected marine park and is surrounded by extensive
248 *Posidonia oceanica* meadows, which are known to influence nutrient and gas dynamics. CS is a shallow
249 nearshore site with intermittent seagrass patches and high exposure to coastal land-sea interactions. These
250 contrasts are relevant to the interpretation of local biogeochemical processes, including N₂O production
251 and air-sea fluxes.

252 2.2 Data collection and analysis

253 2.2.1 Biogeochemical variables

254 To determine N₂O levels, samples were collected in duplicate using 120 mL serum vials sealed with grey-
255 butyl rubber stoppers and aluminium crimps. After being sealed, the samples were preserved with HgCl₂
256 and stored upside-down until analysis. N₂O concentrations were analyzed at the AQUANITROMET
257 laboratory ([https://www.iim.csic.es/en/about-iim/organization/aquanitromet-analysis-greenhouse-effect-](https://www.iim.csic.es/en/about-iim/organization/aquanitromet-analysis-greenhouse-effect-gases-natural-waters)
258 [gases-natural-waters](https://www.iim.csic.es/en/about-iim/organization/aquanitromet-analysis-greenhouse-effect-gases-natural-waters)) of the Instituto de Investigaciones Marinas (IIM-CSIC, Vigo, Spain), employing a
259 static headspace equilibration technique combined with gas chromatography (GC) equipped with electron
260 capture detection, following the methodology detailed by De la Paz et al. (2015). To create the headspace,
261 20 mL of nitrogen gas from a Tedlar bag at atmospheric pressure was introduced into the vials, while
262 simultaneously extracting the same volume of water sample using a double-needle setup. The vials were
263 shaken and left to equilibrate for at least 12 hours in a temperature-controlled environment. For injection
264 into the GC, a brine solution was added through one needle to displace the headspace gas into the GC via
265 the second needle.

266 The GC, an Agilent 7890 GC, was calibrated using three standard gas mixtures: a NOAA-certified primary
267 standard resembling atmospheric air composition and two additional N₂O-in-N₂ mixtures supplied by Air
268 Liquide (De la Paz et al., 2015). While participating in the first large-scale international Inter-Laboratory
269 Comparison experiment for seawater N₂O measurements (Wilson et al., 2018), an additional certified
270 standard from the Scientific Committee for Oceanographic Research (SCOR) was utilized. The precision
271 of the analysis was determined to be 0.5%, calculated from the average coefficient of variation across 400
272 replicate measurements.

273 To determine the DO concentration, samples were collected in five borosilicate bottles and then analyzed
274 using the Winkler method modified by Benson and Krause (1984), employing potentiometric titration with
275 a Metrohm 808 Titrand. The precision of the DO analysis was estimated to be less than $\pm 2 \mu\text{mol kg}^{-1}$.

276 Chl *a* samples were collected in glass bottles and three replicates were filtered in the laboratory using a
277 Whatman GF/F glass fiber filter. Chl *a* extraction was performed with 90% acetone for 24 hours in dark
278 conditions, and the samples were measured with a fluorometer (Turner Designs Instrument, Model 7200-

Eliminado: Oxygen

Con formato: Color de fuente: Texto 1

Eliminado: obtain

Eliminado: with

Eliminado: from

Eliminado: 2022

Eliminado: validation

Eliminado: perform

Eliminado: BP

Eliminado: DO water samples

Eliminado: scrib

Con formato: Normal, Interlineado: sencillo

Con formato: Fuente: Cursiva, Color de fuente: Texto 1

Con formato: Color de fuente: Texto 1

Eliminado: Nitrous oxide (

Eliminado:)

Eliminado: <https://www.iim.csic.es/en/about-iim/organization/aquanitromet-analysis-greenhouse-effect-gases-natural-waters>
<https://www.iim.csic.es/en/about-iim/organization/aquanitromet-analysis-greenhouse-effect-gases-natural-waters>

Con formato: Fuente de párrafo predeter., Color de fuente: Automático, Inglés (británico)

Eliminado: us

Con formato: Color de fuente: Texto 1

Eliminado: gas chromatograph

Con formato: Color de fuente: Texto 1

Eliminado: with a composition

Eliminado: ,

Eliminado: During

Eliminado: on

Eliminado: s

Eliminado: dissolved oxygen (

Eliminado:)

Eliminado: follow

Eliminado: by

Eliminado: ower

Eliminado: with

Eliminado: done

Con formato: Derecha: 0.63 cm

00). The fluorometer was calibrated with a pure Chl *a* standard derived from *Anacystis nidulans* algae, Sigma Chemical Company (Knap et al., 1996). DOC samples were collected in two pre-combustioned borosilicate bottles and then filtered through pre-combusted Whatman GF/F glass fiber filters and stored in two pre-combusted borosilicate vials containing 25 µL of orthophosphoric acid (H₃PO₄). The samples were analyzed with a Shimadzu TOC-L analyzer following the method described by Álvarez-Salgado and Miller (1998), which is based on catalytic oxidation at a high temperature of 680 °C. Inorganic nutrients samples were collected in four 10ml Falcom vials at the same time with the DOC sample processing. Concentrations of NO₃⁻, NO₂⁻, PO₄³⁻, Si (OH)₄, NH₄⁺ and Total Nitrogen were obtained from the analysis laboratory at the Mediterranean Center for Marine and Environmental Research (CMIMA, Barcelona, Spain) using the Autoanalyser AA3 HR (Seal Analytical, United Kingdom) through continuous flow analysis. The precision, estimated from the coefficient of variation based on replicate analyses of the same water samples (n = 10), ranged from 0.13 to 0.5 %.

2.2.2 Meteorological and atmospheric data

Wind speed at a height of 10 m was provided by the Agencia Estatal de Meteorología (AEMET) from the Sant Joan Airport station in Palma, Spain (Fig. 1B). Monthly averaged data of atmospheric N₂O molar fraction was obtained from the monitoring station at Lampedusa (LMP), Italy, as part of the NOAA (National Oceanic and Atmospheric Administration) monitoring network (<http://www.esrl.noaa.gov/gmd/dv/site/>; Lan et al. 2024).

2.3 Flux estimation and other calculations

To calculate the water-atmosphere N₂O fluxes (µmol m⁻² d⁻¹) the following equation was used:

$$F = k * (C_w - C_a^*) \quad (1)$$

Where *k* (cm h⁻¹) is the gas transfer velocity, *C_w* is the concentration of N₂O dissolved in water samples (mol⁻¹), *C_a^{*}* is the gas saturation concentration calculated as the product of the atmospheric fraction of N₂O and the solubility coefficient proposed by Weiss et al. (1980). To compute the fluxes of N₂O, we used the monthly mean atmospheric molar N₂O obtained from NOAA.

The most suitable parameterization for gas transfer (*k*) in coastal areas with seagrass ecosystems, which are characterized by limited wind fetch, representing the study area in the Balearic Sea, was utilized. This equation was described by Dobashi and Ho (2023) as follows:

$$k = 0.143 U_{10}^2 \quad (2)$$

Emissions to the atmosphere are indicated by positive values. Additionally, due to the bottom depth differences between stations and to compare with other studies, the gas transfer parameterizations described by Wanninkhof (2014) and Cole & Caraco (1998) were also used to determine water-atmosphere N₂O fluxes and differences are presented in the obtained annual N₂O fluxes.

Annual N₂O fluxes were calculated by integrating daily flux estimates over time for each station and year using the trapezoidal rule. For each group of samples, the area under the flux-time curve was computed based on the sampling dates, providing an estimate of the total annual air-sea flux. This method accounts

- Eliminado: with
- Eliminado: fixed with
- Eliminado: S
- Eliminado: by using
- Eliminado: a
- Con formato ... [17]
- Eliminado:
- Eliminado: analysis methodology
- Eliminado: (680
- Eliminado: °C)
- Eliminado: and Total Phosphorous
- Eliminado: located
- Eliminado: with
- Eliminado: by
- Eliminado: analysis
- Eliminado: height
- Eliminado: airport,
- Eliminado: Palma
- Eliminado: (Spain)
- Eliminado: statio
- Eliminado: of
- Eliminado: data
- Eliminado: <http://www.esrl.noaa.gov/gmd/dv/site/>
- Con formato ... [18]
- Con formato ... [19]
- Eliminado:
- Con formato ... [20]
- Con formato ... [21]
- Con formato ... [22]
- Con formato ... [23]
- Eliminado: that is
- Eliminado: between
- Con formato ... [24]
- Con formato ... [25]
- Con formato ... [26]
- Con formato ... [27]
- Eliminado: (
- Eliminado: ,
- Eliminado: of
- Eliminado: appropriat
- Eliminado: the
- Eliminado: a
- Eliminado: s
- Con formato ... [28]
- Con formato ... [29]
- Con formato ... [30]
- Con formato ... [31]
- Con formato ... [16]

for the uneven temporal distribution of data and offers a reliable approach to derive annual fluxes from discrete observations.

Furthermore, saturation levels for N_2O ($Sat\%N_2O$) are expressed as a percentage, determined by the ratio of the observed N_2O concentration to the calculated equilibrium concentration. ΔN_2O represents the difference between the observed N_2O concentrations. The Apparent Oxygen Utilization (AOU) is calculated using the DO values along with the measured solubility of oxygen in seawater (Benson and Krause, 1989).

2.4 Data Analysis

To test for differences in N_2O concentrations and fluxes across regions and over temporal scales, simple multifactorial general linear model analyses were conducted using MATLAB version 9.10.0 (R20211, MathWorks Inc.).

Prior to model fitting, we performed a preliminary screening of all available environmental variables to identify those with the highest potential predictive value for N_2O dynamics. We tested the influence of environmental drivers on N_2O levels using the supervised Machine Learning (ML) method known as Gradient Boosting Machine (GBM), which is based on decision tree models using a Cross-Validated Boosting (CVB). GBM utilizes a specific implementation of an ensemble method that combines sequenced base weak models to create a stronger one by applying gradient descent to minimize the model loss function. This process included correlation analysis, variance inflation factors, and variable importance metrics derived from preliminary GBM models. In each iteration, a new model adapts to the residuals of the previously combined model. The combination of all the base models is achieved by summing their predictions, and to prevent overestimation, a weighted learning factor is applied. The CVB technique allows for more precise model evaluation by repeatedly using different datasheets to train and test the model. Both GBM and CVB were implemented in Python version 3.12.3 using the XGBoost library (Chen and Guestrin, 2016). Although NH_4^+ and PO_4^{3-} were included in the initial evaluation, they were excluded from the final model due to either lower relative importance or missing data across sites, which would have compromised model consistency.

The three most influential variables identified (temperature, Chl a , and salinity) were further analyzed to explore their individual and combined effects using Ordinary Least Squares regression (OLS) and Generalized Additive Models (GAM) models. The correlation between the response of N_2O and the resulting dominant environmental variables (relative importance >10%) obtained from the CVB was assessed through OLS using the Statsmodels library (Seabold and Perktold, 2010) in Python version 3.12.3, and by employing GAM with the pyGAM library (Servén and Brummitt, 2020) in Python version 3.12.3 to address the nonlinear dependency. This multi-step approach enabled robust variable selection and the identification of mechanistic relationships driving N_2O spatiotemporal patterns.

To determine the simple linear correlation between environmental variables and FN_2O , Pearson correlation coefficients and p -values were calculated for each station using the Scipy library (Virtanen et al., 2020) in Python version 3.12.3. Furthermore, a multiple linear regression analysis was performed to assess the impact of various predictor variables on FN_2O by applying an OLS.

Eliminado: Additionally...urthermore, saturation value...evels for N_2O ($Sat\%N_2O$)...are express...xpressed as a percentage, were...etermined computed as...y the ratio between...f the observed N_2O concentration observed and...o the calculated equilibrium concentration and... ΔN_2O a...represents the difference between the observed N_2O observations and N_2O solubility concentration value...oncentrations. The Apparent Oxygen Utilization (AOU) is was ...calculated ...using ...he measured DO...O values and...ong with the ...easured solubility of oxygen gas (... [32])

Eliminado: between...cross regions and on...er temporal scales, simple multifactorial general linear model analyses were implemen...onducted in (... [33])

Eliminado: T...e tested the influence of environmental drivers on N_2O levels was tested by...sing the supervised Machine Learning (ML) method known as Gradient Boosting Machine (GBM), which is based on decision tree models using a Cross-Validated Boosting (CVB). GBM is based on...tilizes the ... specific implementation of ensemble method that combines sequenced base weak models to create a stronger one by applying gradient descent to minimize the model loss function. This process included correlation analysis, variance inflation factors, and variable importance metrics derived from preliminary GBM models. In each iteration, a new model adjus...pts to the previous combined...esiduals model...f residual...he in...reviously each...ombined iteration...odel. The combination of all the base models is perform...chieved by summing their predictions, and to avoid...revent overestimation, a weighted learning factor is applied. The CVB technique allows a...or more precise model evaluation of...y the model b...epeatedly using different datasheets to train and test the model. repeatedly. (... [34])

Con formato (... [35])

Eliminado: most ...ominant environmental variables (relative importance >10%) obtained from the CVB was analyz...essed through Ordinary Least Squares regression (...LS)...of (... [36])

Eliminado: ... and by us...mploying Generalized Additive Models (...AM)...with the pyGAM library (Servén and Brummitt, 2020) in Python version 3.12.3 to resolve (... [37])

Con formato: Color de fuente: Texto 1

Eliminado: on...nd FN_2O , Pearson correlation coefficients and p -values were calculated for each station using the Scipy library (Virtanen et al., 2020) in Python version 3.12.3. Furthermore, a multiple linear regression analysis was performed to analyze...sess the impact of numer...arious predictor variables on the (... [38])

Con formato: Derecha: 0.63 cm

3 Results and discussion

3.1 Environmental Variables Description

The observed seawater temperature patterns during the study period reflect the climatic characteristics of the Mediterranean region, with peak temperatures recorded in summer (June–August), notably reaching their maximum in August 2022, at 30.3 °C, 29.9 °C, and 29 °C in the PB, CA, and CS sites, respectively (Fig. 2A, 3C, and Table 1). Conversely, minimum temperatures recorded during winter (December–February) were 13.7 °C in 2023 for PB, 13.04 °C in 2023 for CA, and 13 °C in 2019 for CS. Over the entire study period, average ($\pm$ standard deviation) temperature values stood at 20.9 ± 5.1 , 20.7 ± 4.9 , and 20.4 ± 5.0 °C for PB, CA, and CS, respectively, with no statistical differences between sites ($p > 0.05$, Table 1), but denoted variability between months and years ($p < 0.005$, Table 1).

Surface salinity levels showed significant differences between sites ($p < 0.005$ with CS values), months ($p < 0.005$), and years ($p < 0.05$), maintaining average values of 37.5 ± 0.2 practical salinity units (PSU) for PB, 37.4 ± 0.2 PSU for CA, and 37.7 ± 0.3 PSU for CS. The highest salinity levels were recorded during the summer of 2018 at PB (38.2 PSU), the winter of 2023 at CA (38.0 PSU), and during both the summer of 2018 and winter of 2023 at CS (38.2 PSU) (Fig. 2B, Table 1). The differences in the factory precision of the sensors that measure conductivity, and thus salinity, could explain the variations observed between the CS, PB, and CA stations in terms of salinity.

Wind speed measurements taken at a height of 10 meters displayed a noticeable seasonal trend (Fig. 2C). However, no statistical differences were found between years and months ($p > 0.05$). The highest average values occurred during spring (March–May), averaging around 3.7 ± 0.4 ms⁻¹ and decreased during winter to 2.9 ± 0.5 ms⁻¹ over the study period. Maximum peak values were recorded in December and January, reaching levels of up to 19.9 ms⁻¹.

DO levels showed distinct seasonal patterns across all stations, with the highest values recorded in spring (up to 300 $\mu\text{mol kg}^{-1}$) and the lowest during late summer, particularly at CA, reaching around 195 $\mu\text{mol kg}^{-1}$. Average DO concentrations were similar at PB and CA, at 233.5 ± 22.0 and 231.8 ± 25.3 $\mu\text{mol kg}^{-1}$, respectively, while limited data were available for CS. AOU values were primarily negative, indicating net photosynthetic production. Nutrient concentrations displayed spatial and temporal variability. NO₃⁻ concentrations ranged from 0.1 to 6.3 μM , with higher values observed at CS (average 1.5 ± 1.5 μM) compared to PB and CA. NO₂⁻ levels were generally low (0.1–0.3 μM) but exhibited significant interannual differences. DOC showed seasonal variation, with values ranging from 52 to 116 μM , with the highest values in summer and lower concentrations in winter (Table 1).

3.2 Nitrous oxide and related biogeochemical variables.

Over the monitored period from 2018 to 2023, dissolved N₂O concentrations exhibited a seasonal pattern ($p < 0.005$, Figure 3A), ranging from 6.5 to 9.9 nmol L⁻¹, with no significant differences between stations and years ($p > 0.05$). The highest concentrations were recorded during winter, averaging 9.0 ± 0.2 nmol L⁻¹ (with a peak in February), followed closely by autumn (September–November) at 8.2 ± 0.6 nmol L⁻¹ and spring at 7.9 ± 0.7 nmol L⁻¹ (Table 1). In contrast, the lowest values were observed in summer, averaging 6.9 ± 0.2 nmol L⁻¹, with the minimum recorded in August 2021 at PB. Opposite trends were noted in

Eliminado: ...registering...t 30.3...29.9...C, 29.9 °C, and 29 °C in the BP...B, CA, and CS sites, respectively (Fig. 2A, 3C, and...Table 1). Conversely, minimum temperatures recorded during winter (December–February) were 13.7 °C in 2023 for BP...B, 13.04 °C in 2023 for CA, and 13 °C in 2019 for CS. Over the entire study period, average ($\pm$ standard deviation) temperature values stood at 20.9 ± 5.1 , 20.7 ± 4.9 , and 20.4 ± 5.0 °C for BP (... [39]

Eliminado: exhibit...howed significant differences between sites ($p < 0.005$ with CS values), months ($p < 0.005$), and years ($p < 0.05$), maintaining average values of 37.5 ± 0.2 practical salinity uni (... [40]

Con formato: Color de fuente: Texto 1

Eliminado: BP...B, 37.4 ± 0.2 psu (... [41]

Eliminado: ...0.3 psu (... [42]

Con formato: Color de fuente: Texto 1

Con formato: Color de fuente: Texto 1

Eliminado: BP...B (38.2 psu (... [43]

Eliminado: psu

Con formato: Color de fuente: Texto 1

Con formato: Color de fuente: Texto 1

Eliminado: psu

Con formato: Color de fuente: Texto 1

Eliminado: determin...easure conductivity, and...therefore...s salinity, could cause...xplain the differences...ariations observed between the CS and (... [44]

Eliminado: exhibit...isplayed an...observ...oticeable seasonal trend (Fig. 2C). However, no statistical differences were found between years and months ($p > 0.05$). Maximum...he averaged...ighest average values appear...occurred during spring (March–May), balanced...veraging around 3.7 ± 0.4 ms⁻¹...and decline...eased during winter to 2.9 ± 0.5 ms⁻¹...over the study period. Maximum peak values were not...ecorded in December and January, reaching levels of up to 19.9 ms⁻¹ (... [45]

Con formato (... [46]

Eliminado: display...xhibited a seasonal pattern ($p < 0.005$, Table 1, ... (... [47]

Eliminado: sparti...ferences between stations and years ($p > 0.05$). The highest levels ...oncentrations were recorded during the winter season...inter, averaging 9.0 ± 0.2 nmol L⁻¹ (with a peak in February), followed closely by autumn (September–November) with ...t 8.2 ± 0.6 nmol L⁻¹ and spring with ...t 7.9 ± 0.7 nmol L⁻¹ (Table 1). Conversely...n contrast, the lowest values were observed summer, averaging 6.9 ± 0.2 nmol L⁻¹, with the minimum recorded in August 2021 in...t PB. ...Opposite trends were observ (... [48]

Con formato: Derecha: 0.63 cm

Sat%N₂O percentages, with maximum values in summer and autumn and minimum values in winter, ranging from 93 to 116.7% (Figure 3B).

Chl *a* followed a distinct seasonal pattern (Fig. 3D, Table 1), with significant differences observed between sites, years, and months ($p < 0.005$). In January, we recorded the highest productivity with a mean Chl *a* value of $0.4 \pm 0.2 \mu\text{g L}^{-1}$, while May noted the lowest at $0.2 \pm 0.05 \mu\text{g L}^{-1}$. PB exhibited the highest levels at $0.3 \pm 0.2 \mu\text{g L}^{-1}$, whereas CA ($0.2 \pm 0.2 \mu\text{g L}^{-1}$) and CS ($0.2 \pm 0.1 \mu\text{g L}^{-1}$) displayed comparable concentrations. In 2019, all stations showed the strongest Chl *a* signal (Fig. 3D).

DO presents significant monthly differences observed across all stations ($p < 0.005$, Figure 3E and Table 1). During the spring, DO levels peaked at up to $300 \mu\text{mol kg}^{-1}$, while at the end of the summer, minimum values around $160 \mu\text{mol kg}^{-1}$ were recorded, particularly at the CA site. The average levels from 2018 to 2023 were consistent between PB and CA, at 233.5 ± 22.0 and $231.8 \pm 25.3 \mu\text{mol kg}^{-1}$, respectively. In contrast, only limited sensor data were available for CS (Fig. 3E, green dots). However, it is well-known that relevant oxygen production in the area is closely related to the presence of seagrass meadows, dominated by the endemic species *Posidonia oceanica*, as previously noted by Agueda-Aramburu et al. (2024) at the precise CA station location of this study.

AOU values were predominantly negative (Table 1), with average values of -14.1 ± 12.0 , -11.9 ± 15.3 , and $-15.1 \pm 4.9 \mu\text{mol kg}^{-1}$ for PB, CA, and CS sites, respectively. The negative AOU levels indicate the DO produced by the excess of photosynthesis compared to respiration, which aligns well with the organic matter content observed from the DOC data (Fig. 3F). In the coastal Balearic Sea, DOC showed similarities between sites and years ($p > 0.05$), with significant differences among months ($p < 0.05$). During the summer season, the highest average levels of $83.9 \pm 0.9 \mu\text{M}$ were recorded, while winter exhibited the lowest at $73.8 \pm 1.5 \mu\text{M}$ (Fig. 3F, Table 1). Variability in DOC data was notable in spring and autumn, with average values of 76.0 ± 5.9 and $76.6 \pm 3.1 \mu\text{M}$, respectively.

NO₃⁻ levels ranged from 0.1 to 6.3 μM during the study period, showing no significant differences between months but significant differences between years ($p < 0.005$) and sites ($p < 0.005$). CS exhibited the highest NO₃⁻ levels with an average of $1.5 \pm 1.5 \mu\text{M}$, followed by PB with $0.3 \pm 0.4 \mu\text{M}$, while CA had the lowest at $0.1 \pm 0.2 \mu\text{M}$ (Fig. 3G, Table 1). NO₂⁻ concentrations were significantly different between years ($p < 0.005$; Fig. 3H, Table 1) and between stations ($p < 0.05$ for CA, Table 1), with total average values of 0.03 ± 0.02 , 0.05 ± 0.05 , and $0.07 \pm 0.06 \mu\text{M}$ for CA, PB, and CS, respectively.

The analysis of the effects of environmental and biogeochemical variables on N₂O concentrations through the CVB revealed only seven main parameters associated with the variability of N₂O concentrations (Fig. 4). These parameters, in order of importance, were temperature (65 %), Chl *a* (17 %), salinity (7 %), DOC (5 %), NO₃⁻ (4 %), and NO₂⁻ (2 %).

Temperature values, the first variable modeling N₂O concentrations, exhibited a strong decreasing linear correlation with N₂O levels, with $p < 0.005$ and a R^2 of 0.94 for all evaluated stations (not shown). The solubility of N₂O declines in warmer waters, which explains the observed maximum and minimum seasonal patterns of N₂O, primarily governed by the thermodynamic effects of temperature on N₂O concentrations. This strong correlation reflects the thermodynamic influence of temperature on N₂O solubility and its broader role in regulating stratification and microbial activity. The Sat%N₂O used here as a proxy for the biological consumption/production of N₂O shows a positive linear correlation with temperature ($p < 0.005$

Eliminado: levels...ercentages, with maximum values in summer and autumn and minimum values in winter, ranging from 93 to 116.7% (Figure 3BTable 1) (... [49])

Eliminado: cernible...inct seasonal pattern (Fig. 3B...D, Table 1), with significant disparities...erences observed between sites, years, and months ($p < 0.005$). In January, we foun...ecorded the highest productivity with a mean Chl *a* value of $0.4 \pm 0.2 \mu\text{g L}^{-1}$, while May record...oted the lowest at $0.2 \pm 0.05 \mu\text{g L}^{-1}$. PB exhibited the highest levels of...t $0.3 \pm 0.2 \mu\text{g L}^{-1}$, whereas CA ($0.2 \pm 0.2 \mu\text{g L}^{-1}$) and CS ($0.2 \pm 0.1 \mu\text{g L}^{-1}$) displayed similar...omparable concentrations. In 2019, all the...stations exhibit...howed the strongest Chl *a* signal 3B... (... [50])

Eliminado: significant ...ifferences foun...bserved for...cross all the...stations ($p < 0.005$, Figure 3E and Table 1). During the spring season... DO levels reach...eaked their peak...t up to $300 \mu\text{mol kg}^{-1}$, while towards...t the end of the summer season... minimum values around $160 \mu\text{mol kg}^{-1}$ were recorded, particularly at the CA site. The average levels from 2018 to 2023 were coher...sistent between BP PB and CA, with...t 233.5 ± 22.0 and $231.8 \pm 25.3 \mu\text{mol kg}^{-1}$, respectively. In contrast, only limited sensor data were available for CS (Fig. 3D...E, green dots). However, it is well-known that the relevant oxygen production in the area is strong...losely related to the the denoted...presence of seagrass meadows, dominated by the endemic species *Posidonia oceanica*, as previously signal...oted in...y Agueda-Aramburu et al. (2024) in...t the exact (... [51])

Eliminado: d...values of -14.1 ± 12.0 , -11.9 ± 15.3 , and $-15.1 \pm 4.9 \mu\text{mol kg}^{-1}$ for PB, CA, and CS sites, respectively. AOU...he negative AOU levels indicate the DO...O produced by the excess of photosynthesis versus...ompared to respiration, which i...igns well-fitted...with the organic matter content observed from the DOC...OC data (Fig. 3C...F). In the coastal Balearic Sea, DOC reveal...howed similarities between sites and years ($p > 0.05$), with significant differences among months ($p < 0.05$). During the summer season, the highest average levels of $83.9 \pm 0.9 \mu\text{M}$ were recorded, while winter exhibited the lowest of...t $73.8 \pm 1.5 \mu\text{M}$ (Fig. 3F, Table 1). Variability in DOC data variability (... [52])

Eliminado: vari...anged from 0.1 to 6.3 μM during the study period, with...howing no significant differences between months but significant differences between years ($p < 0.005$) and sites ($p < 0.005$), with... CS showing...xhibited the highest NO₃⁻ levels with average ...n values...verage of $1.5 \pm 1.5 \mu\text{M}$, followed by PB with $\pm 0.4 \mu\text{M}$, and...hile CA had the lowest with...t $0.1 \pm 0.2 \mu\text{M}$ (Fig. 3E...G, Table 1). NO₂⁻ concentrations were significantly different between years ($p < 0.005$; Fig. 3F...H, Table 1) and between stations ($p < 0.05$ for CA, Table 1), with total averaged (... [53])

Eliminado: the...N₂O concentrations through the CVB resulted in...ealed only seven main parameters associated with the N₂O variability... (... [54])

Con formato (... [55])

Eliminado: were...arameters, in order of importance, were...temperature (65 %), Chl *a* (17 %), salinity (7 %), dissolved organic carbon (... [56])

Con formato: Color de fuente: Texto 1

Eliminado: nitrate ...4 %), and NO₂⁻nitrite (... [57])

Eliminado: the...O concentrations, presen...xhibited a strong decreasing linear correlation with N₂O levels, with $p < 0.005$ and an...R² ... of 0.94 for all the...valuated stations evaluated ...not shown). The solubility of N₂O solubility decreas...eclines in warmer waters, explaining...hich explains the observed maximum and minimum seasonal patterns of N₂O, which is ...rimarily governed by the thermodynamic influence (... [58])

Con formato (... [59])

Eliminado: of

Con formato: Derecha: 0.63 cm

and a R^2 of 0.59; Fig. 5). This positive correlation is anticipated since the microbial activities of nitrification and denitrification are expected to increase with rising temperatures, leading to greater N_2O production (Wu et al., 2018). However, our results indicate that the seasonality control of N_2O by thermodynamic solubility surpassed the enhancement of microbial activity. The dominant role of temperature observed here aligns with previous large-scale coastal assessments (e.g., Yang et al., 2020), which identified temperature as a key variable influencing both gas solubility and microbial turnover, particularly in surface waters affected by seasonal thermal stratification.

The DO availability and the predominance of negative AOU levels, with no correlation to $Sat\%N_2O$ ($p>0.05$, $R^2=0.002$), suggest that *in situ* N_2O production makes a minor contribution to the observed temporal variability of N_2O concentration. Overall, the low significance of predictors in the CVB analysis, such as Chl a , DO, DOC, NO_3^- , and NO_2^- , is consistent with limited N_2O production from nitrification, which is associated with the seasonal remineralization cycle of organic matter, particularly enhanced under high-productivity conditions. This interpretation aligns with previous studies in coastal and shelf environments that also reported low N_2O accumulation under oxygenated and nutrient-limited conditions (e.g., Babbin et al., 2015; Lin et al., 2016). These conditions can suppress canonical N_2O production through nitrification and denitrification, emphasizing the sensitivity of microbial N_2O cycling to ambient redox and substrate gradients.

DOC and Chl a also contribute to the availability of organic substrates and oxygen fluctuations, shaping microbial N_2O dynamics, especially under stratified or high-productivity conditions. In fact, DOC enrichment has been linked to elevated heterotrophic activity and subsequent shifts in nitrogen cycling pathways in productive coastal areas (Froehlich et al., 2021), potentially influencing the balance between N_2O production and consumption. Microbial production by nitrification and denitrification are generally considered the dominant pathways of N_2O biological production in the shelf sea and open ocean (Burgos et al., 2017; Chen et al., 2021; de la Paz et al., 2024; Sierra et al., 2020). However, in the oxygenated coastal Balearic Sea waters, the Balearic Sea, near-equilibrium values for $Sat\%N_2O$ and low NO_3^- and NO_2^- concentrations (Figs. 3B, G-H, 5, Table 1) suggest a negligible contribution from pelagic nitrification. Nevertheless, NO_3^- variability may still indicate diffuse anthropogenic inputs, potentially enhancing local nitrification under favorable conditions. While benthic N_2O fluxes were not measured during this study due to logistical constraints, their contribution cannot be dismissed. A follow-up study is being conducted to evaluate potential sediment-derived N_2O inputs at the same study sites.

In addition, the CBV analysis points to Chl a as the second driver explaining the observed variability of N_2O concentration since the observed seasonal cycle of Chl a is coupled to the N_2O , with maximum values in winter and minimum in summer. A hypothetical mechanism that could be contributing to the N_2O concentration variability in our study site is the poorly described production of N_2O by epipelagic photosynthetic organisms in the light through NO reduction. Given the low NO_3^- and NO_2^- concentrations, the lack of correlation between AOU and N_2O , and the seasonal coupling between Chl a and N_2O peaks, we propose that an additional, non-canonical pathway may contribute to N_2O production in our system. Recent studies have identified a light-dependent mechanism in which marine photosynthetic organisms, including green algae and cyanobacteria, reduce intracellular nitric oxide (NO) to N_2O , potentially as a detoxification strategy under fluctuating nitrogen conditions or during photosynthetic stress. This NO

- Eliminado: n
- Eliminado: R^2
- Eliminado: expect
- Eliminado: y
- Eliminado: is
- Eliminado: assum
- Eliminado: ri
- Eliminado: increa
- Eliminado: followed
- Eliminado: by
- Eliminado: control of
- Eliminado: exceed
- Eliminado: promotion
- Eliminado: with
- Eliminado: ha
- Eliminado: N_2O variability
- Eliminado: import
- Eliminado: of
- Con formato: Color de fuente: Texto 1
- Con formato: Fuente: Cursiva, Color de fuente: Texto 1
- Con formato: Color de fuente: Texto 1
- Eliminado: DOC, nitrate
- Eliminado: nitrite
- Eliminado: ,
- Eliminado: ow
- Con formato: Color de fuente: Texto 1
- Eliminado: remineralization
- Eliminado: seasonal
- Eliminado: which is
- Eliminado: in the
- Con formato: Fuente: Cursiva, Color de fuente: Texto 1
- Con formato: Color de fuente: Texto 1
- Con formato: Color de fuente: Texto 1
- Eliminado: N_2O
- Eliminado: value
- Eliminado: E-F
- Eliminado: point to
- Eliminado: minor
- Eliminado: of the
- Con formato: Color de fuente: Texto 1
- Eliminado: ¶
- Con formato: Color de fuente: Texto 1
- Eliminado: N_2O variability
- Eliminado: (Burlacot et al., 2020).
- Con formato: Fuente: Cursiva, Color de fuente: Texto 1
- Con formato: Color de fuente: Texto 1
- Con formato: Derecha: 0.63 cm

reduction pathway involves the activity of flavodiiron proteins (FLVs) or other reductases and may become relevant in oxygenated, nutrient-depleted surface waters with active phytoplankton populations (Burlacot et al., 2020). While our dataset does not allow for direct confirmation of this process, the observed seasonal patterns and weak traditional N₂O signals warrant further consideration of this alternative mechanism, especially in coastal systems dominated by epipelagic productivity.

Finally, significant unexpected negative linear correlations ($p < 0.05$) were found between N₂O concentrations and silicate only at the PB and CA sites (Table 1), with correlation coefficients of -0.41 and -0.34, respectively. This relationship and the differences in salinity values between PB and CA compared to CS may indicate the presence or absence of groundwater input pulses in these areas (Basterretxea et al., 2010; Sospedra et al., 2018), which can affect N₂O concentrations in specific coastal regions (Calvo-Martin et al., 2024). Since this relationship was only observed at two of the three stations considered, this parameter was excluded from the CVB analysis. Together, these factors illustrate a multifactorial control of N₂O production and accumulation in the coastal Balearic Sea, where solubility, productivity, and nutrient inputs act in combination.

An OLS regression was conducted to evaluate the relationship among the three variables with the highest relative importance for N₂O concentrations (Importance >5%), accounting for 89% of the variability (Temperature, Chl *a*, and salinity, Fig. 4) identified in the CVB analysis. The resulting equation is: N₂O = 7.42 - 0.18 * Temperature + 0.28 * Chl *a* + 0.11 * Salinity ($R^2 = 0.94$), with a p -value < 0.005 for temperature and p -values of 0.07 for Chl *a* and 0.152 for salinity.

To resolve the nonlinear dependency of Chl *a* and salinity on N₂O concentrations, a GAM was applied, explaining 96 % of the response variable with a p -value < 0.005 for temperature, < 0.05 for Chl *a*, and 0.1390 for salinity (Supplementary figures). The non-significance for salinity may be because GAM smooth terms are designed to capture broader nonlinear effects. In this case, salinity might not have a robust and smooth effect, or its impact is masked by other variables like temperature. Salinity also influences stratification dynamics and can indirectly affect oxygen availability and microbial processes linked to N₂O. The GAM residuals ranged from -0.6 to 0.6 nmol L⁻¹ (Supplementary figures).

When compared to other temperate and subtropical coastal regions, our results support the emerging view that N₂O dynamics in seagrass-dominated systems may diverge substantially from classic estuarine and shelf models, with solubility and light-driven processes playing an increasingly relevant role (Rosentreter et al., 2023; Al-Haj et al., 2021). Placing the obtained data in a global ocean context, specifically among studied coastal areas, the Balearic Sea presents N₂O concentrations similar to those found in 2008 in the Bohai Sea, ranging from 7.14 to 8.32 nmol L⁻¹ (Gu et al., 2022), but are low in comparison to upwelling and estuarine coastal areas. The maximum N₂O concentrations found in this study represent approximately baseline levels for the other coastal and open ocean areas evaluated previously (Bange, 2006; Bange et al., 1996; de la Paz et al., 2024; Yang et al., 2009; Zhang et al., 2008).

3.3 Air-sea N₂O exchange

FN₂O values ranged from -0.3 to 0.6 μmol m⁻² d⁻¹ during the study period, indicating that the coastal Balearic Sea functions as a minor sink or is nearly in equilibrium in winter and spring and acts as a light source in summer and autumn (Fig. 6; $p < 0.005$), closely following the seasonal temperature variations

Eliminado: unexpected ...significant unexpected negative (... [60])

Con formato (... [61])

Eliminado: levels ... concentrations and silicate only at the PB and CA sites (Table 1), with -0.41 ... correlation and ... coefficients of -0.34 ... correlation ... coefficients ... 0.34, respectively. This relationship and the obtained ... differences in salinity values between PB and CA compared to CS can ... ay deno ... indicate the presence or absence of groundwater input pulses in these areas (Basterretxea et al., 2010; Sospedra et al., 2018), that ... hich can affect N₂O concentrations in specific coastal regions (Calvo-Martin et al., 2024). As ... ince this relationship was only observed in ... t two of the three stations considered, this parameter was not ... xcluded include (... [62])

Eliminado: perform ... onducted to evaluate the relationship of ... among the three variables with the higher ... t relative importance for N₂O concentrations (Importance >5%), representing ... counting for 89% of the variability (Temperature, Chl *a*, and salinity, Fig. 4) obtained ... identified in the CVB analysis. The equation ... esulting obtained ... quation was ... s: N₂O = 7.42 - 0.18 * Temperature + 0.28 * Chl *a* + 0.11 * Salinity ($R^2 = 0.94$), with a p -value < 0.005 for temperature and a p -value (... [63])

Con formato: Color de fuente: Texto 1

Eliminado: the ... chl *a* and salinity with ... n N₂O concentrations, a GAM was applied, explaining 96 % of the response variable with a p -value < 0.005 for temperature and ... < 0.05 for Chl *a*, and 0.1390 for salinity (Supplemented ... ry figures). The non-significance for salinity could ... ay be because GAM smooth terms are designed to capture broader nonlinear effects. In this case, Salinity ... alinity might either ... ot have a robust and smooth effect, or its impact is masked by other variables like temperature. Salinity also influences stratification dynamics and can indirectly affect oxygen availability and microbial processes linked to N₂O. The GAM residuals ranged from -0.6 to 0.6 nmol L⁻¹ (Supplemented (... [64])

Eliminado: concrete ... pecifically among studied coastal areas, the Balearic Sea presents similar ... O concentrations levels to ... imilar the ... o rang ... hose found in 2008 in the Bohai Sea, 7.14- ... rom 7.14 to 8.32 nmol L⁻¹ (Gu et al., 2022), but small ... re low when ... n compared ... son to upwelling and estuarine coastal areas. M ... he maximum N₂O ... O levels ... oncentrations found in this study represent approximately baseline levels for the rest of (... [65])

Eliminado: over ... uring the study period, acting ... ndicating that the coastal Balearic Sea area ... unctions as a minor sink or is nearly in equilibrium in winter and spring and acts as a light source in summer and autumn (Fig. 6; $p < 0.005$), closely following the season (... [66])

Con formato: Derecha: 0.63 cm

(Fig. 6, grey line). No differences between years were observed. However, significant differences were noted between sites, with PB exhibiting the highest FN₂O levels and CA showing the lowest ($p < 0.05$). In the PB site, positive linear correlations were found between FN₂O and temperature ($R^2 = 0.67$; $p < 0.005$) and between FN₂O and Δ N₂O ($R^2 = 0.96$; $p < 0.005$). In the OLS analysis, wind speed also appears to exert a positive influence on FN₂O values, detectable only in PB data (OLS, $R^2 = 0.94$, $p < 0.05$). This effect is likely attributable to PB site geographical characteristics, as it represents an open bay, in contrast to the enclosed CA station in the Santa Maria Bay sampling site and the shallow coastal waters of CS. FN₂O in the most pristine station CA showed positive correlations also with temperature ($R^2 = 0.79$; $p < 0.005$), Δ N₂O ($R^2 = 0.97$; $p < 0.005$) and the Sat%N₂O ($R^2 = 0.97$; $p < 0.005$). The CS shallow site showed positive linear correlations with temperature ($r = 0.73$; $p < 0.005$), Δ N₂O ($R^2 = 0.98$; $p < 0.005$), and Sat%N₂O ($R^2 = 0.97$; $p < 0.005$). Salinity was also a predictive variable included in the OLS analysis (OLS, $R^2 = 0.96$, $p < 0.05$) and was negatively related. This characteristic may be associated with the differences in CS salinity values compared to the other stations. Annual average fluxes for the three sites were calculated, when monthly data were unavailable, linear interpolation was applied (Table 2). The different parameterizations for the gas transfer velocity indicate higher values using Cole & Caraco (1998), ranging from 8.7 to 204.2 $\mu\text{mol m}^{-2} \text{y}^{-1}$, followed by results obtained with the Wanninkhof (2014) equation, which range from 9.9 to 190.1 $\mu\text{mol m}^{-2} \text{y}^{-1}$ and the lowest values defined by Dobashi & Ho (2023), ranging from 5.4 to 107.4 $\mu\text{mol m}^{-2} \text{y}^{-1}$ (Table 2). These differences are clearly influenced by the environment since the Dobashi & Ho (2023) equation was derived from studies conducted in shallow areas at approximately 3.5 m depth, while Wanninkhof (2014) and Cole & Caraco (1998) focused on deeper ocean areas and a lake, respectively. However, since the benthic zone of this study is heavily covered with *P. oceanica* seagrass meadows and the fetch area is limited, we opted to retain Dobashi & Ho (2023) for the primary results. Similar low air-sea N₂O fluxes have also been reported in shallow vegetated environments, such as seagrass-dominated coastal lagoons in Australia (Rosentreter et al., 2023), where negative or near-zero fluxes reflect the combined effects of high primary production, oxygenation, and limited nitrogen availability. In contrast, higher fluxes, often exceeding 1 $\mu\text{mol m}^{-2} \text{d}^{-1}$, are typical of eutrophic estuaries or upwelling regions (Bange, 2006; Zhang et al., 2008), highlighting the relatively pristine nature of the Balearic coastal zone. The results showed that all the stations are significantly neutral regarding N₂O emissions to the atmosphere. However, notably higher values were detected at the PB station, as seen in the daily FN₂O values (Fig. 6). PB exhibited significant interannual variability during the study period (Table 2). The CS and CA sites followed the same trend as the PB station but had considerably lower values, particularly for CA. Observed annual N₂O fluxes in the Balearic Sea are much lower than the expected ranges for this ocean region, as described by Resplandy et al. (2024) based on global reconstruction products from 1985 to 2018 (Yang et al., 2020). These differences may relate to the substantial variability in local N₂O fluxes in coastal areas with vegetation. Coastal regions with seagrass meadows are expected to exhibit the lowest N₂O flux ranges (Rosentreter et al., 2023). These areas are considered negligible sources (Al-Haj et al., 2021) or sinks in near-pristine seagrasses, leading to significantly biased coastal global estimations (Chen et al., 2022). These low fluxes can be partly attributed to the biogeochemical effects of seagrass meadows. Furthermore,

- Eliminado: closely
- Eliminado: with
- Eliminado: r
- Eliminado: with
- Con formato ... [68]
- Con formato ... [69]
- Eliminado: r
- Eliminado: also
- Eliminado: only
- Eliminado: detectable
- Eliminado: R²
- Eliminado: ,
- Eliminado: it
- Eliminado: s
- Eliminado: PB
- Eliminado: r
- Eliminado: r
- Eliminado: r
- Eliminado: present
- Eliminado: lineal
- Eliminado: along
- Eliminado: with
- Eliminado: r
- Eliminado: the
- Eliminado: r
- Eliminado: ng
- Eliminado: present
- Eliminado: R²
- Eliminado: can
- Eliminado: rel
- Eliminado: to
- Eliminado: with
- Eliminado: , and
- Eliminado: as
- Eliminado: a
- Con formato ... [70]
- Con formato ... [71]
- Con formato ... [72]
- Con formato ... [73]
- Con formato ... [74]
- Con formato ... [75]
- Con formato ... [76]
- Con formato ... [77]
- Con formato ... [78]
- Con formato ... [79]
- Con formato ... [80]
- Con formato ... [81]
- Con formato ... [67]

elevated rates of primary production reduce the availability of inorganic nitrogen in the water column, thereby limiting N₂O generation. Additionally, the dense root systems and organic exudates of seagrass meadows support microbial communities that further facilitate N₂O consumption under oxic conditions (Rosentreter et al., 2023; Al-Haj et al., 2021). Collectively, these mechanisms contribute to the role of vegetated coastal systems as potential N₂O sinks.

These observations place the Balearic Sea at the lower end of the global N₂O flux spectrum for coastal systems. For example, in the East China Sea and Pearl River Estuary coastal areas, fluxes ranged from 5 to 17 $\mu\text{mol m}^{-2} \text{d}^{-1}$ (Zhang et al., 2008; Lin et al., 2016), often driven by hypoxia and strong anthropogenic nitrogen inputs—conditions absent in our study area.

Considering that European seagrasses represent approximately 6% of global seagrass cover, and that most of this is found in the Mediterranean Sea (Jordà et al., 2012), it is crucial to understand more about the drivers of N₂O formation, consumption seasonality, and transport pathways. Additionally, increasing the number of observations is vital since N₂O data in the Mediterranean Sea is limited. Furthermore, we must consider the ongoing coastal eutrophication linked to anthropogenic inputs from estuaries, sewage discharge from densely populated coastal areas, and industrial effluents that may significantly alter seagrass habitats, along with current emissions of N₂O from coastal zones (Bakker et al., 2014).

The low N₂O fluxes observed in our study are in line with recent assessments that emphasize the underestimated climate mitigation potential of seagrass meadows and vegetated coastal habitats (Al-Haj et al., 2021; Chen et al., 2022). Thus, understanding the ecological conditions that suppress N₂O emissions in these systems is critical for refining regional and global GHG budgets.

As coastal N₂O emissions are assumed to offset 30–58% of the net CO₂ coastal uptake radiative effect (Resplandy et al., 2024), estimating the water N₂O emission-based Global Warming Potential (GWP) can provide a clear picture by comparing it with existing data on CO₂ and methane fluxes in the area. By following the Intergovernmental Panel on Climate Change (IPCC) Assessment Report 6 (Arias et al., 2021) updated 100-year GWP, we obtained the GWP_{N₂O} (i.e., the 100-year time-integrated radiative forcing from the instant release of 1 kg of N₂O is 273 times larger than the forcing of 1 kg of CO₂).

In the coastal Balearic Sea, considering an area 1 km offshore and 1,428 km of coastal longitude, the average GWP_{N₂O} obtained in this study is $8.1 \times 10^{-7} \pm 5.7 \times 10^{-7} \text{ Pg CO}_2\text{-eq y}^{-1}$ for all sites from 2020 to 2022, which was the period with the most available data. This area estimate was used as a first-order approximation of the active N₂O exchange zone in the Balearic region, following a similar approach to other regional-scale studies (e.g., Resplandy et al., 2024; Rosentreter et al., 2023). It encompasses both vegetated and non-vegetated shallow coastal habitats and is not intended to represent specific benthic types. Therefore, it reflects a conservative estimate of the total coastal footprint, acknowledging that further habitat-specific refinement would improve spatial resolution. The associated uncertainty in the GWP estimate incorporates the observed variability in fluxes across sites and seasons.

4 Conclusions

N₂O concentrations in seawater during the study period ranged from 6.5 to 9.9 nmol L^{-1} without significant differences between the three sampling sites. Several drivers dominated the variability of N₂O concentration, with temperature as the most essential factor and less critical Chlorophyll *a* and salinity.

Eliminado: The results showed that all the stations are minor sources of N₂O to the atmosphere, with notably higher values in the PB station, as observed in the daily FN₂O values (Fig. 6). PB presented an apparent interannual variability during the study period (Table 2). The CS and CA sites followed the same trend as the PB station but with significantly lower values, especially for CA. Observed annual N₂O fluxes in the Balearic Sea are considerably lower than the expected ranges for this ocean region, as described by Resplandy et al. (2024) based on global reconstruction products from 1985 to 2018 (Yang et al., 2020). These differences could relate to the large variability in local N₂O fluxes in coastal areas with vegetation. Coastal regions with seagrass meadows are expected to have the lowest N₂O flux ranges (Rosentreter et al., 2023). These areas are considered negligible sources (Al-Haj et al., 2021) or sinks in near-pristine seagrasses, resulting in strongly biased coastal global estimations (Chen et al., 2022).

Eliminado: Considering that European seagrasses constitute 6% of the global seagrass meadows and that most of this coverage is allocated in the Mediterranean Sea, it is imperative to know more about the drivers of the formation and consumption seasonality and the transport pathways.

Eliminado: Iso, the increase

Eliminado: critic

Eliminado: becaus

Eliminado: nitrous oxide

Eliminado: scarce

Eliminado: In addition

Eliminado: have

Eliminado: to take into account

Eliminado: relat

Eliminado: sites

Eliminado: high

Eliminado: well

Eliminado: modify

Eliminado: as

Eliminado: well

Eliminado: as

Eliminado: give us

Eliminado: In the coastal Balearic Sea, considering an area of 1 km distance offshore and 1428 km of coastal longitude, the average GWP_{N₂O} obtained in this study is $8.1 \times 10^{-7} \pm 5.7 \times 10^{-7} \text{ Pg CO}_2\text{-eq y}^{-1}$ for all the sites from 2020 to 2022, which was the period with more available data. Comparing with the annual GWP_{CO₂} of $2.3 \pm 10^{-6} \pm 4.6 \times 10^{-5} \text{ g CO}_2 \text{ y}^{-1}$ obtained in the coastal Balearic Sea from the available data from Flecha et al. (2023), we can confirm that N₂O emissions in this area are not substantial yet in terms of GWP regarding other GHGs. However, the offset of 35% obtained in this study for N₂O closely follows the assumption by Resplandy et al. (2024)....

Eliminado: M

Con formato: Color de fuente: Texto 1, Superíndice

Con formato: Color de fuente: Texto 1

Eliminado: N₂O variability

Con formato: Derecha: 0.63 cm

Even with the possible biological implications in N₂O formation, atmospheric forcing may control the surface concentrations in this area. Averaged estimated N₂O fluxes oscillated between -0.3 and 0.6 μmol m⁻² d⁻¹, with the most **influenced** stations showing the highest N₂O fluxes. **Some sites appear to be weak sources of N₂O, although many fluxes are close to equilibrium and not significantly different from zero,** following a robust seasonal pattern throughout the sampling period. Reducing anthropogenic pressures in coastal regions is crucial to preserving marine ecosystems, as increasing human impacts, such as pollution, overfishing, and habitat destruction, can lead to irreversible damage, loss of biodiversity, and the degradation of vital ecosystem services. If these pressures continue to escalate, future projections indicate more frequent and severe environmental crises, including increased coastal erosion, declining fish stocks, and heightened vulnerability to climate change, threatening marine life and human coastal communities. There is a strong need to **enhance** observations of the **GHG, N₂O** in coastal areas to **better understand** dominant drivers **and make** more **accurate** predictions of future consequences under additional anthropogenic impacts. **Moreover, the values presented** for the coastal Balearic Sea will **aid in improving** global **N₂O** emissions budgets in coastal vegetated areas **overall**.

Data availability

Data was obtained from the Metocean Data Repository of the Balearic Islands Coastal Observing and Forecasting System (SOCIB). **In 2024, data** from the instruments on the Palma Bay Station platform, <https://apps.socib.es/data-catalog>, **was** consulted on 01-30-2024. Data is also **available** through Tintoré & Casas (2022) and Hendriks et al. (2023).

Supplement link

Author contribution

IE, JT, and SF conceptualized the research, data acquisition approach, and methodology. IE, SF, MdP, CM, and AEF collected the samples and **conducted** the measurements. SF, MdP, and IE analyzed the data, **while** SF wrote the manuscript draft. All authors reviewed and edited the manuscript.

Competing interests

The authors declare no competing financial interest.

Acknowledgments

We **express our gratitude** to the Cabrera National Park staff for facilitating the work **done** during this study, and to the Balearic Islands Coastal Observing and Forecasting System (SOCIB) for their invaluable **assistance** and **the** use of their fixed station in the Bay of Palma. We **extend our thanks to** Lidia Cucala (IMEDEA), Andrea Carbonero, and Juan Martínez Ayala (SOCIB) for their **support** with sample collection and analysis. This work **contributes** to CSIC Thematic Interdisciplinary Platform PTI OCEANS+. We **appreciate** the AQUANITROMET service of Instituto Investigaciones Marinas for **conducting** the methane analyses, and **we thank** the Agencia Estatal de Meteorología (AEMET) for providing the meteorological data.

Eliminado: impacted

Con formato: Fuente: Sin Cursiva, Color de fuente: Texto 1

Con formato: Fuente: Sin Cursiva, Color de fuente: Texto 1

Eliminado: All areas generally behaved as weak N₂O sources

Eliminado: increase the number of

Eliminado: Greenhouse

Eliminado: Gas

Eliminado: nitrous oxide

Eliminado: understand

Eliminado: the

Eliminado: ting

Eliminado: to

Eliminado: elaborat

Eliminado: exact

Eliminado: In addition

Eliminado: presented

Eliminado: values

Eliminado: contribute to an improve

Eliminado: estimation

Eliminado: of

Eliminado: N₂O

Eliminado: in genera

Eliminado: Data

Eliminado: ,

Eliminado: ,

Eliminado: data

Eliminado: ccessi

Eliminado: ;

Eliminado: perform

Eliminado: ;

Eliminado: ;

Eliminado: ;

Eliminado: are

Eliminado: grateful

Eliminado: performed

Eliminado: help

Eliminado: thank

Eliminado: assistance

Eliminado: is a contribution

Eliminado: thank

Con formato: Derecha: 0.63 cm

266
267 **Financial support**
268 Funding for this work was provided by the Spanish Ministry of Science (SumaEco, RTI2018–095441-B-
269 C21, CYCLE, PID2021-123723OB-C21), the Government of the Balearic Islands through la Conselleria
270 d'Innovació, Recerca i Turisme (Projecte de recerca científica i tecnològica SEPP0, PRD2018/18), and the
271 2018 call of the BBVA Foundation “Ayudas a equipos de investigación científica” for the Posi-COIN
272 Project. SF acknowledges the financial support of the “Margalida Comas-2017” and “Vicenç Munt
273 Estabilitat-2022” postdoctoral contracts, as well as project AAEE111/2017 from the Balearic Islands
274 Government. SF is staff hired under the Generation D initiative, promoted by Red.es, an organisation
275 attached to the Ministry for Digital Transformation and the Civil Service, for the attraction and retention of
276 talent through grants and training contracts financed by the Recovery, Transformation, and Resilience Plan
277 through the European Union's Next Generation funds. MP acknowledges the financial support during the
278 study period to the contracts financed by the Spanish Ministry of Science CTM2015–74510-JIN and
279 PTA2019–017983-I. F.F Pérez was supported by the FICARAM+ project (PID2023 - 148924OB - I00).

280
281 **References**

282 Agueda-Aramburu, P., Flecha, S., Lujan-Williams, C. A. M., and Hendriks, I. E.: Water column
283 oxygenation by Posidonia oceanica seagrass meadows in coastal areas: A modelling approach, Science of
284 the Total Environment, 942, 173805, <https://doi.org/10.1016/j.scitotenv.2024.173805>, 2024.
285 Al-Haj, A. N., Chidsey, T., Fulweiler, R. W., Rosentreter, J. A., Al-Haj, A. N., Fulweiler, R. W.,
286 Williamson, P., Laruelle, G. G., Bange, H. W., Bianchi, T. S., Busecke, J. J. M., Cai, W. J., Eyre, B. D.,
287 Forbrich, I., Kwon, E. Y., Maavara, T., Moosdorf, N., Najjar, R. G., Sarma, V. V. S. S., Van Dam, B., and
288 Regnier, P.: Two temperate seagrass meadows are negligible sources of methane and nitrous oxide, Nat
289 Clim Chang, 35, 579–587, <https://doi.org/10.1002/ln.12250>, 2021.
290 Álvarez-Salgado, X. A. and Miller, A. E. J.: Simultaneous determination of dissolved organic carbon and
291 total dissolved nitrogen in seawater by high temperature catalytic oxidation: Conditions for precise
292 shipboard measurements, Mar Chem, 62, 325–333, [https://doi.org/10.1016/S0304-4203\(98\)00037-1](https://doi.org/10.1016/S0304-4203(98)00037-1), 1998.
293 Arias, P., Bellouin, N., Coppola, E., Jones, R., Krinner, G., Marotzke, J., Naik, V., Palmer, M., Plattner,
294 G.-K., and Rogelj, J.: Climate Change 2021: The Physical Science Basis. Contribution of Working Group I
295 I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change; Technical Summary,
296 2021.
297 Babbin, A. R., Bianchi, D., Jayakumar, A., & Ward, B. B. Rapid nitrous oxide cycling in the suboxic ocean.
298 Science, 348(6239), 1127–1129, <https://doi.org/10.1126/science.aaa8380>, 2015.
299 Bakker, D. C. E., Bange, H. W., Gruber, N., Johannessen, T., Upstill-Goddard, R. C., Borges, A. V., Delille,
300 B., Löscher, C. R., Naqvi, S. W. A., Omar, A. M., and Magdalena Santana-Casiano, J.: Air-sea interactions
301 of natural long-lived greenhouse gases (CO₂, N₂O, CH₄) in a changing climate, in: Ocean-Atmosphere
302 Interactions of Gases and Particles, Springer Berlin Heidelberg, 113–169, [https://doi.org/10.1007/978-3-](https://doi.org/10.1007/978-3-642-25643-1_3)
303 [642-25643-1_3](https://doi.org/10.1007/978-3-642-25643-1_3), 2014.
304 Bange, H. W.: Nitrous oxide and methane in European coastal waters, Estuar Coast Shelf Sci, 70, 361–374,
305 <https://doi.org/10.1016/j.ecss.2006.05.042>, 2006.

Eliminado: as

Eliminado: nd

Eliminado: and

Con formato: Fuente: Sin Cursiva, Color de fuente: Texto 1

Eliminado: the MMT24-PTI+OCEANS-01 postdoctoral contract funded by the European Union NextGenerationEU.

Eliminado:

Con formato: Color de fuente: Texto 1

Con formato: Color de fuente: Texto 1

Con formato: Derecha: 0.63 cm

312 Bange, H. W., Rapsomanikis, S., and Andreae, M. O.: Nitrous oxide in coastal waters, Global
 313 Biogeochemical Cycles, German, 132 pp., 1996.

314 Barbier, E. B., Hacker, S. D., Kennedy, C., Koch, E. W., Stier, A. C., and Silliman, B. R.: The value of
 315 estuarine and coastal ecosystem services, Ecol Monogr, 81, 169–193,
 316 <https://doi.org/https://doi.org/10.1890/10-1510.1>, 2011.

317 Basterretxea, G., Tovar-Sanchez, A., Beck, A. J., Masqué, P., Bokuniewicz, H. J., Coffey, R., Duarte, C.
 318 M., Garcia-Orellana, J., Garcia-Solsona, E., Martinez-Ribes, L., and Vaquer-Sunyer, R.: Submarine
 319 groundwater discharge to the coastal environment of a Mediterranean island (Majorca, Spain): Ecosystem
 320 and biogeochemical significance, Ecosystems, 13, 629–643, <https://doi.org/10.1007/s10021-010-9334-5>,
 321 2010.

322 Benson, B. B. and Krause, D.: A thermodynamic treatment of dilute solutions of gases in liquids, J Solution
 323 Chem, 18, 803–821, <https://doi.org/10.1007/BF00685061>, 1989.

324 Benson, B. B. and Krause Jr., D.: The concentration and isotopic fractionation of oxygen dissolved in
 325 freshwater and seawater in equilibrium with the atmosphere, Limnol Oceanogr, 29, 620–632,
 326 <https://doi.org/https://doi.org/10.4319/lo.1984.29.3.0620>, 1984.

327 Bianchi, C. N. and Morri, C.: Marine biodiversity of the Mediterranean Sea: Situation, problems and
 328 prospects for future research, Mar Pollut Bull, 40, 367–376, [https://doi.org/10.1016/S0025-](https://doi.org/10.1016/S0025-326X(00)00027-8)
 329 326X(00)00027-8, 2000.

330 Burgos, M., Ortega, T., and Forja, J. M.: Temporal and spatial variation of N₂O production from estuarine
 331 and marine shallow systems of Cadiz Bay (SW, Spain), Science of the Total Environment, 607–608,
 332 <https://doi.org/10.1016/j.scitotenv.2017.07.021>, 2017.

333 Burlacot, A., Richaud, P., Gosset, A., Li-Beisson, Y., and Peltier, G.: Algal photosynthesis converts nitric
 334 oxide into nitrous oxide, PNAS, 117, 2704–2709, [https://doi.org/10.1073/pnas.1915276117/-](https://doi.org/10.1073/pnas.1915276117/-/DCSupplemental)
 335 /DCSupplemental, 2020.

336 Calvo-Martin, E., de la Paz, M., Álvarez-Salgado, X. A., Pazó Fernández, M. J., Vieitez Dos Santos, V.,
 337 and Ibáñez, J. S. P.: Transport and reactivity of nitrous oxide and methane in two contrasting subterranean
 338 estuaries, Limnol Oceanogr, 1–16, <https://doi.org/10.1002/lno.12622>, 2024.

339 Carstensen, J. and Duarte, C. M.: Drivers of pH Variability in Coastal Ecosystems, Environ Sci Technol,
 340 53, 4020–4029, <https://doi.org/10.1021/acs.est.8b03655>, 2019.

341 [Chen, J.-J., Wells, N. S., Erler, D. V., & Eyre, B. D. Land-use intensity increases benthic N₂O emissions](https://doi.org/10.1029/2022JG006899)
 342 [across three sub-tropical estuaries. Journal of Geophysical Research: Biogeosciences, 127,](https://doi.org/10.1029/2022JG006899)
 343 [e2022JG006899, <https://doi.org/10.1029/2022JG006899>, 2022.](https://doi.org/10.1029/2022JG006899)

344 Chen, T. and Guestrin, C.: XGBoost: A Scalable Tree Boosting System, in: Proceedings of the 22nd ACM
 345 SIGKDD International Conference on Knowledge Discovery and Data Mining, 785–794,
 346 <https://doi.org/10.1145/2939672.2939785>, 2016.

347 Chen, X., Ma, X., Gu, X., Liu, S., Song, G., Jin, H., and Zhang, G.: Seasonal and spatial variations of N₂O
 348 distribution and emission in the East China Sea and South Yellow Sea, Science of the Total Environment,
 349 775, 145715, <https://doi.org/10.1016/j.scitotenv.2021.145715>, 2021.

350 Codispoti, L. A.: Interesting Times for Marine N₂O, Science (1979), 327, 1339–1340,
 351 <https://doi.org/10.1126/science.1184945>, 2010.

Con formato: Color de fuente: Texto 1

Con formato: Color de fuente: Texto 1

Con formato: Derecha: 0.63 cm

352 Cole, J. J. and Caraco, N. F.: Atmospheric exchange of carbon dioxide in a low-wind oligotrophic lake
 353 measured by the addition of SF₆, *Limnol Oceanogr*, 43, 647–656,
 354 <https://doi.org/10.4319/lo.1998.43.4.0647>, 1998.

355 Dobashi, R. and Ho, D. T.: Air–sea gas exchange in a seagrass ecosystem—results from a ³He/SF₆ tracer
 356 release experiment, *Biogeosciences*, 20, 1075–1087, 2023.

357 Doney, S. C.: The growing human footprint on coastal and open-ocean biogeochemistry, *Science* (1979),
 358 328, 1512, 2010.

359 Duarte, C. M., Middelburg, J. J., and Caraco, N.: Major role of marine vegetation on the oceanic carbon
 360 cycle, *Biogeosciences*, 2, 1–8, <https://doi.org/10.5194/bg-2-1-2005>, 2005.

361 EEA: State and pressures of the marine and coastal Mediterranean environment, European Environment
 362 Agency, 1–44, 1999.

363 Farias, L., Paulmier, A., and Gallegos, M.: Nitrous oxide and N-nutrient cycling in the oxygen minimum
 364 zone off northern Chile, *Deep Sea Research Part I: Oceanographic Research Papers*, 54, 164–180,
 365 <https://doi.org/https://doi.org/10.1016/j.dsr.2006.11.003>, 2007.

366 Flecha, S., Giménez-Romero, À., Tintoré, J., Pérez, F. F., Alou-Font, E., Matías, M. A., and Hendriks, I.
 367 E.: pH trends and seasonal cycle in the coastal Balearic Sea reconstructed through machine learning, *Sci*
 368 *Rep*, 12, 12956, <https://doi.org/10.1038/s41598-022-17253-5>, 2022.

369 Flecha, S., Rueda, D., de la Paz, M., Pérez, F. F., Alou-Font, E., Tintoré, J., and Hendriks, I. E.: Spatial and
 370 temporal variation of methane emissions in the coastal Balearic Sea, Western Mediterranean, *Science of*
 371 *the Total Environment*, 865, <https://doi.org/10.1016/j.scitotenv.2022.161249>, 2023.

372 Giorgi, F.: Climate change hot-spots, *Geophys Res Lett*, 33,
 373 <https://doi.org/https://doi.org/10.1029/2006GL025734>, 2006.

374 Giorgi, F. and Lionello, P.: Climate change projections for the Mediterranean region, *Glob Planet Change*,
 375 63, 90–104, <https://doi.org/10.1016/j.gloplacha.2007.09.005>, 2008.

376 Gu, T., Jia, D., Wang, Z., Guo, Y., Xin, Y., Guo, C., Zhang, G., and Sun, J.: Regional distribution and
 377 environmental regulation mechanism of nitrous oxide in the Bohai Sea and North Yellow Sea: A
 378 preliminary study, *Science of the Total Environment*, 818, <https://doi.org/10.1016/j.scitotenv.2021.151718>,
 379 2022.

380 Hendriks, I. E., Escolano-Moltó, A., Flecha, S., Vaquer-Sunyer, R., Wesselmann, M., and Marbà, N.:
 381 Mediterranean seagrasses as carbon sinks: methodological and regional differences, *Biogeosciences*, 19,
 382 4619–4637, <https://doi.org/10.5194/bg-19-4619-2022>, 2022.

383 Hendriks, I. E., Flecha, S., de La Paz, M., Pérez, F. F., Morell, C., Tintoré, J., and Marbà, N.: Methane
 384 emissions in the coastal Balearic Sea between October 2019 - October 2021, *DIGITAL CISC*,
 385 <https://doi.org/10.20350/digitalCSIC/15491>, 2023.

386 Jordà, G., Marbà, N. & Duarte, C. Mediterranean seagrass vulnerable to regional climate warming. *Nature*
 387 *Clim Change* 2, 821–824, 2012. <https://doi.org/10.1038/nclimate1533>

388 Knap, A., Michaels, A., Close, A., Ducklow, H., and Dickson, A.: Protocols for the Joint Global Ocean
 389 Flux Study (JGFS) Core Measurements, *JGOFS Reoprt* Nr. 19, vi+170 pp, 198, 1996.

Con formato: Color de fuente: Texto 1, Superíndice

Con formato: Color de fuente: Texto 1

Eliminado: ¶

Eliminado: ¶

Con formato: Derecha: 0.63 cm

de la Paz, M., Huertas, I. E., Flecha, S., Ríos, A. F., and Pérez, F. F.: Nitrous oxide and methane in Atlantic and Mediterranean waters in the Strait of Gibraltar: Air-sea fluxes and inter-basin exchange, *Prog Oceanogr*, 138, 18–31, <https://doi.org/10.1016/j.pocean.2015.09.009>, 2015a.

de la Paz, M., Huertas, I. E., Flecha, S., Ríos, A. F., and Pérez, F. F.: Nitrous oxide and methane in Atlantic and Mediterranean waters in the Strait of Gibraltar: Air-sea fluxes and inter-basin exchange, *Prog Oceanogr*, 138, 18–31, <https://doi.org/10.1016/j.pocean.2015.09.009>, 2015b.

de la Paz, M., Pérez, F. F., Álvarez, M., and Bode, A.: Seasonal ventilation controls nitrous oxide emission in the NW Iberian upwelling, *Prog Oceanogr*, 224, <https://doi.org/10.1016/j.pocean.2024.103261>, 2024.

Lan, X., Mund, J. W., Crotwell, A. M., Thoning, K. W., Moglia, E., Madronich, M., Baugh, K., Petron, G., Crotwell, M. J., Neff, D., Wolter, S., Mefford, T., and DeVogel, S.: Atmospheric Nitrous Oxide Dry Air Mole Fractions from the NOAA GML Carbon Cycle Cooperative Global Air Sampling Network, 1997–2023, <https://doi.org/10.15138/53g1-x417>, 2024.

[Lin, H., Dai, M., Kao, S.-J., Wang, L., Roberts, E., Yang, J.-Y. T., Huang, T., and He, B.: Spatiotemporal variability of nitrous oxide in a large eutrophic estuarine system: The Pearl River Estuary, China, *Mar Chem*, 182, 14–24, 2016.](#)

Ma, X., Lennartz, S. T., and Bange, H. W.: A multi-year observation of nitrous oxide at the Boknis Eck Time Series Station in the Eckernförde Bay (southwestern Baltic Sea), *Biogeosciences*, 16, 4097–4111, <https://doi.org/10.5194/bg-16-4097-2019>, 2019.

Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S. L., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M. I., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J. B. R., Maycock, T. K., Waterfield, T., Yelekçi, O., Yu, R., and Zhou, B.: Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change, in: Sixth Assessment Report of the Intergovernmental Panel on Climate Change, Cambridge University Press, 2021.

Micheli, F., Halpern, B. S., Walbridge, S., Ciriaco, S., Ferretti, F., Frascchetti, S., Lewison, R., Nykjaer, L., and Rosenberg, A. A.: Cumulative human impacts on Mediterranean and Black Sea marine ecosystems: assessing current pressures and opportunities, *PLoS One*, 8, e79889, 2013.

Mouillot, D., Albouy, C., Guilhaumon, F., Ben Rais Lasram, F., Coll, M., Devictor, V., Meynard, C. N., Pauly, D., Tomasini, J. A., Troussellier, M., Velez, L., Watson, R., Douzery, E. J. P., and Mouquet, N.: Protected and threatened components of fish biodiversity in the mediterranean sea, *Current Biology*, 21, 1044–1050, <https://doi.org/10.1016/j.cub.2011.05.005>, 2011.

Nevison, C. D., Weiss, R. F., and Erickson III, D. J.: Global oceanic emissions of nitrous oxide, *J Geophys Res Oceans*, 100, 15809–15820, <https://doi.org/https://doi.org/10.1029/95JC00684>, 1995.

[Ravishankara, A. R., Daniel, J. S., & Portmann, R. W.: Nitrous oxide \(N₂O\): The dominant ozone-depleting substance emitted in the 21st century, *Science*, 326\(5949\), 123–125. <https://doi.org/10.1126/science.1176985>, 2009.](#)

Resplandy, L., Hogikyan, A., Müller, J. D., Najjar, R. G., Bange, H. W., Bianchi, D., Weber, T., Cai, W.-J., Doney, S. C., Fennel, K., Gehlen, M., Hauck, J., Lacroix, F., Landschützer, P., Le Quéré, C., Roobaert, A., Schwinger, J., Berthet, S., Bopp, L., Chau, T. T. T., Dai, M., Gruber, N., Ilyina, T., Kock, A., Manizza, M., Lachkar, Z., Laruelle, G. G., Liao, E., Lima, I. D., Nissen, C., Rödenbeck, C., Séférian, R., Toyama,

Con formato: Fuente: Sin Negrita, Color de fuente: Texto 1

Con formato: Color de fuente: Texto 1

Con formato: Color de fuente: Texto 1

Con formato: Color de fuente: Texto 1

Eliminado: ¶

Con formato: Derecha: 0.63 cm

433 K., Tsujino, H., and Regnier, P.: A Synthesis of Global Coastal Ocean Greenhouse Gas Fluxes, *Global*
434 *Biogeochem Cycles*, 38, e2023GB007803, <https://doi.org/https://doi.org/10.1029/2023GB007803>, 2024.

435 Rosentreter, J. A., Laruelle, G. G., Bange, H. W., Bianchi, T. S., Busecke, J. J. M., Cai, W. J., Eyre, B. D.,
436 Forbrich, I., Kwon, E. Y., Maavara, T., Moosdorf, N., Najjar, R. G., Sarma, V. V. S. S., Van Dam, B., and
437 Regnier, P.: Coastal vegetation and estuaries are collectively a greenhouse gas sink, *Nat Clim Chang*, 13,
438 579–587, <https://doi.org/10.1038/s41558-023-01682-9>, 2023.

439 Seabold, S. and Perktold, J.: statsmodels: Econometric and statistical modeling with python, in: 9th Python
440 in Science Conference, 2010.

441 Servén, D. and Brummitt, C.: pyGAM Documentation, 2020.

442 Sierra, A., Jiménez-López, D., Ortega, T., Ponce, R., Bellanco, M. J., Sánchez-Leal, R., Gómez-Parra, A.,
443 and Forja, J.: Distribution of N₂O in the eastern shelf of the Gulf of Cadiz (SW Iberian Peninsula), *Science*
444 *of the Total Environment*, 593–594, <https://doi.org/10.1016/j.scitotenv.2017.03.189>, 2017.

445 Sierra, A., Jiménez-López, D., Ortega, T., Gómez-Parra, A., and Forja, J.: Factors controlling the variability
446 and emissions of greenhouse gases (CO₂, CH₄ and N₂O) in three estuaries of the Southern Iberian Atlantic
447 Basin during July 2017, *Mar Chem*, 226, <https://doi.org/10.1016/j.marchem.2020.103867>, 2020.

448 Solomon, S., Qin, D., Manning, M., Averyt, K., and Marquis, M.: Climate change 2007-the physical science
449 basis: Working group I contribution to the fourth assessment report of the IPCC, Cambridge University
450 Press, Cambridge, United Kingdom and New York, NY, USA., 2007.

451 Sospedra, J., Niencheski, L. F. H., Falco, S., Andrade, C. F. F., Attisano, K. K., and Rodilla, M.: Identifying
452 the main sources of silicate in coastal waters of the Southern Gulf of Valencia (Western Mediterranean
453 Sea), *Oceanologia*, 60, 52–64, <https://doi.org/10.1016/j.oceano.2017.07.004>, 2018.

454 Tian, H., Pan, N., Thompson, R. L., Canadell, J. G., Suntharalingam, P., Regnier, P., Davidson, E. A.,
455 Prather, M., Ciais, P., Muntean, M., Pan, S., Winiwarter, W., Zechle, S., Zhou, F., Jackson, R. B., Bange,
456 H. W., Berthet, S., Bian, Z., Bianchi, D., Bouwman, A. F., Buitenhuis, E. T., Dutton, G., Hu, M., Ito, A.,
457 Jain, A. K., Jeltsch-Thömmes, A., Joos, F., Kou-Giesbrecht, S., Krummel, P. B., Lan, X., Landolfi, A.,
458 Lauerwald, R., Li, Y., Lu, C., Maavara, T., Manizza, M., Millet, D. B., Mühle, J., Patra, P. K., Peters, G.
459 P., Qin, X., Raymond, P., Resplandy, L., Rosentreter, J. A., Shi, H., Sun, Q., Tonina, D., Tubiello, F. N.,
460 van der Werf, G. R., Vuichard, N., Wang, J., Wells, K. C., Western, L. M., Wilson, C., Yang, J., Yao, Y.,
461 You, Y., and Zhu, Q.: Global nitrous oxide budget (1980–2020), *Earth Syst Sci Data*, 16, 2543–2604,
462 <https://doi.org/10.5194/essd-16-2543-2024>, 2024.

463 Tintoré, J. and Casas-Pérez, B.: Buoy Bahía de Palma Physicochemical parameters of sea water data
464 (Version x.y.z), Balearic Islands Coastal Observing and Forecasting System, SOCIB,
465 <https://doi.org/10.25704/S6JB-CK61>, 2022.

466 Tintoré, J., Vizoso, G., Casas, B., Heslop, E., Pascual, A., Orfila, A., Ruiz, S., Martínez-Ledesma, M. M.,
467 Torner, M., Cusí, S., Diedrich, A., Balaguer, P., Gómez-Pujol, L., Álvarez-Ellacuría, A., Gómara, S.,
468 Sebastian, K., Lora, S., Beltrán, J. P., Renault, L., Juzà, M., Álvarez, D., March, D., Garau, B. B., Castilla,
469 C., Cañellas, T., Roque, D., Lizarán, I., Pitarch, S., Carrasco, M. A., Lana, A. A., Mason, E., Escudier, R.,
470 Conti, D., Sayol, J. M., Barceló, B., Alemany, F. F., Reglero, P., Massuti, E., Vélez-Belchí, P., Ruiz, J.,
471 Oguz, T., Gómez, M., Álvarez, E., Ansorena, L. L., Manriquez, M., Barceló-Llull, B., Alemany, F. F.,
472 Reglero, P., Massuti, E., Vélez-Belchí, P., Ruiz Segura, J., Oguz, T., Gómez, M., Álvarez-Fanjul, E.,

Con formato: Derecha: 0.63 cm

473 Ansorena, L. L., and Manríquez, M.: SOCIB: The Balearic Islands Coastal Ocean Observing and
 474 Forecasting System Responding to Science, Technology and Society Needs, *Mar Technol Soc J*, 47, 101–
 475 117, <https://doi.org/10.4031/MTSJ.47.1.10>, 2013.

476 Tintoré, J., Pinardi, N., Álvarez-Fanjul, E., Aguiar, E., Álvarez-Berastegui, D., Bajo, M., Balbin, R.,
 477 Bozzano, R., Nardelli, B. B., Cardin, V., Casas, B., Charcos-Llorens, M., Chiggiato, J., Clementi, E.,
 478 Coppini, G., Coppola, L., Cossarini, G., Deidun, A., Deudero, S., D’Ortenzio, F., Drago, A., Drudi, M., El
 479 Serafy, G., Escudier, R., Farcy, P., Federico, I., Fernández, J. G., Ferrarin, C., Fossi, C., Frangoulis, C.,
 480 Galgani, F., Gana, S., García Lafuente, J., Sotillo, M. G., Garreau, P., Gertman, I., Gómez-Pujol, L., Grandi,
 481 A., Hayes, D., Hernández-Lasheras, J., Herut, B., Heslop, E., Hilmi, K., Juza, M., Kallos, G., Korres, G.,
 482 Lecci, R., Lazzari, P., Lorente, P., Liubartseva, S., Louanchi, F., Malacic, V., Mannarini, G., March, D.,
 483 Marullo, S., Mauri, E., Meszaros, L., Mourre, B., Mortier, L., Muñoz-Mas, C., Novellino, A., Obaton, D.,
 484 Orfila, A., Pascual, A., Pensieri, S., Pérez Gómez, B., Pérez Rubio, S., Perivoliotis, L., Petihakis, G., de la
 485 Villéon, L. P., Pistoia, J., Poulain, P. M., Pouliquen, S., Prieto, L., Raimbault, P., Reglero, P., Reyes, E.,
 486 Rotllan, P., Ruiz, S., Ruiz, J., Ruiz, I., Ruiz-Orejón, L. F., Salihoglu, B., Salon, S., Sammartino, S., Sánchez
 487 Arcilla, A., Sannino, G., Sannino, G., Santoleri, R., Sardá, R., Schroeder, K., Simoncelli, S., Sofianos, S.,
 488 Sylaios, G., Tanhua, T., Teruzzi, A., Testor, P., Tezcan, D., Torner, M., et al.: Challenges for Sustained
 489 Observing and Forecasting Systems in the Mediterranean Sea, *Front Mar Sci*, 6,
 490 <https://doi.org/10.3389/fmars.2019.00568>, 2019.

491 [United Nations Environment Programme / Mediterranean Action Plan – Barcelona Convention](#)
 492 [\(UNEP/MAP\). State of the Mediterranean Marine and Coastal Environment. UNEP/MAP – Barcelona](#)
 493 [Convention. Retrieved from https://wedocs.unep.org/20.500.11822/375, 2012.](#)

494 del Valle Villalonga, L., Pons, G. X., and Bardolet, M.: Posidonia oceanica Cartography and Evolution of
 495 the Balearic Sea (Western Mediterranean), *Remote Sens (Basel)*, 15, <https://doi.org/10.3390/rs15245748>,
 496 2023.

497 Virtanen, P., Gommers, R., Oliphant, T. E., Haberland, M., Reddy, T., Cournapeau, D., Burovski, E.,
 498 Peterson, P., Weckesser, W., Bright, J., van der Walt, S. J., Brett, M., Wilson, J., Millman, K. J., Mayorov,
 499 N., Nelson, A. R. J., Jones, E., Kern, R., Larson, E., Carey, C. J., Polat, İ., Feng, Y., Moore, E. W.,
 500 VanderPlas, J., Laxalde, D., Perktold, J., Cimrman, R., Henriksen, I., Quintero, E. A., Harris, C. R.,
 501 Archibald, A. M., Ribeiro, A. H., Pedregosa, F., van Mulbregt, P., Vijaykumar, A., Bardelli, A. Pietro,
 502 Rothberg, A., Hilboll, A., Kloeckner, A., Scopatz, A., Lee, A., Rokem, A., Woods, C. N., Fulton, C.,
 503 Masson, C., Häggström, C., Fitzgerald, C., Nicholson, D. A., Hagen, D. R., Pasechnik, D. V., Olivetti, E.,
 504 Martin, E., Wieser, E., Silva, F., Lenders, F., Wilhelm, F., Young, G., Price, G. A., Ingold, G. L., Allen, G.
 505 E., Lee, G. R., Audren, H., Probst, I., Dietrich, J. P., Silterra, J., Webber, J. T., Slavič, J., Nothman, J.,
 506 Buchner, J., Kulick, J., Schönberger, J. L., de Miranda Cardoso, J. V., Reimer, J., Harrington, J., Rodríguez,
 507 J. L. C., Nunez-Iglesias, J., Kuczynski, J., Tritz, K., Thoma, M., Newville, M., Kümmerer, M., Bolingbroke,
 508 M., Tarte, M., Pak, M., Smith, N. J., Nowaczyk, N., Shebanov, N., Pavlyk, O., Brodtkorb, P. A., Lee, P.,
 509 McGibbon, R. T., Feldbauer, R., Lewis, S., Tygier, S., Sievert, S., Vigna, S., Peterson, S., More, S., Pudlik,
 510 T., et al.: SciPy 1.0: fundamental algorithms for scientific computing in Python, *Nat Methods*, 17, 261–
 511 272, <https://doi.org/10.1038/s41592-019-0686-2>, 2020.

Eliminado: ¶

Con formato: Fuente de párrafo predeter., Color de fuente: Texto 1

Con formato: Derecha: 0.63 cm

Wan, X. S., Sheng, H. X., Dai, M., Casciotti, K. L., Church, M. J., Zou, W., Liu, L., Shen, H., Zhou, K., Ward, B. B., and Kao, S. J.: Epipelagic nitrous oxide production offsets carbon sequestration by the biological pump, *Nat Geosci*, 16, 29–36, <https://doi.org/10.1038/s41561-022-01090-2>, 2023.

Wanninkhof, R.: Relationship between wind speed and gas exchange over the ocean revisited, *Limnol Oceanogr Methods*, 12, 351–362, <https://doi.org/10.4319/lom.2014.12.351>, 2014.

Weiss, R. F., Price, B. A., Canada, E., Rochette, P., Hutchinson, G. L., Jayasundara, S., and Wagner-Riddle, C.: Nitrous oxide solubility in water and seawater, *Agric Syst*, 8, 247–286, 1980.

Wilson, S. T., Ferrón, S., and Karl, D. M.: Interannual Variability of Methane and Nitrous Oxide in the North Pacific Subtropical Gyre, *Geophys Res Lett*, 44, 9885–9892, <https://doi.org/https://doi.org/10.1002/2017GL074458>, 2017.

Wu, B., Liu, F., Weiser, M. D., Ning, D., Okie, J. G., Shen, L., Li, J., Chai, B., Deng, Y., and Feng, K.: Temperature determines the diversity and structure of N₂O-reducing microbial assemblages, *Funct Ecol*, 32, 1867–1878, 2018.

Yang, J., Zhang, G. L., Zheng, L. X., and Zhang, F.: Seasonal variations of fluxes and distributions of dissolved N₂O in the North Yellow Sea, *Huanjing Kexue/Environmental Science*, 30, 2009.

Yang, S., Chang, B. X., Warner, M. J., Weber, T. S., Bourbonnais, A. M., Santoro, A. E., Kock, A., Sonnerup, R. E., Bullister, J. L., Wilson, S. T., and Bianchi, D.: Global reconstruction reduces the uncertainty of oceanic nitrous oxide emissions and reveals a vigorous seasonal cycle, *Proc Natl Acad Sci U S A*, 117, <https://doi.org/10.1073/pnas.1921914117>, 2020.

Zhang, G., Zhang, J., Ren, J., Li, J., and Liu, S.: Distributions and sea-to-air fluxes of methane and nitrous oxide in the North East China Sea in summer, *Mar Chem*, 110, <https://doi.org/10.1016/j.marchem.2008.02.005>, 2008.

Zhou, J., Zheng, Y., Hou, L., An, Z., Chen, F., Liu, B., Wu, L., Qi, L., Dong, H., and Han, P.: Effects of acidification on nitrification and associated nitrous oxide emission in estuarine and coastal waters, *Nat Commun*, 14, 1380, 2023.


Eliminado: ¶

Eliminado: ¶

¶

Con formato: Derecha: 0.63 cm

Figure 1: (A) Map of the stations location in the Western Mediterranean Sea Basin and (B) detailed location of the Bay of Palma (PB; blue dot), the Cabrera National Park (CA, red dot), and Cape Ses Salines (CS; green dot) study sites and Palma Airport wind station (black star). Dashed lines represent bathymetry levels. Maps were developed with the Python software version 3.12.3.



Eliminado: 

Con formato: Color de fuente: Texto 1


Eliminado: Figure 1: (A) Map of the stations location in the Western Mediterranean Sea Basin and (B) detailed location of the Bay of Palma (BP; blue dot) and the Cabrera National Park (CA, red dot) and Cape Ses Salines (CS; green dot) study sites. Dashed lines represent bathymetry levels. Maps were developed with the Python software version 3.12.3. ... [82]

Con formato: Color de fuente: Texto 1


Con formato: Color de fuente: Texto 1

Con formato: Derecha: 0.63 cm

Figure 2: Time-series of A) daily averaged temperature (°C) for PB (blue dots) and CA (red dots) stations, as well as instantaneous values for the CS site (green dots); B) daily averaged salinity (PSU) for the PB (blue dots) and CA (red dots) stations, along with instantaneous values for the CS site (green dots); and C) daily averaged wind speed (m s⁻¹) at 10 m height from the Palma station (Spain). Data for PB, CA, and CS were collected at depths of 1, 4, and 0.5 m, respectively. Figures were created using Python software version 3.12.3.




Eliminado:
 Eliminado: Daily
 Eliminado: and
 Eliminado: instant
 Eliminado:),
 Eliminado: Daily
 Eliminado: psu
 Con formato: Color de fuente: Texto 1
 Eliminado: nd
 Eliminado: instant
 Eliminado: Daily
 Eliminado: -
 Eliminado: developed with
 Eliminado: the



Eliminado:
 Con formato: Color de fuente: Texto 1
 Con formato: Color de fuente: Texto 1

Con formato: Derecha: 0.63 cm

Figure 3: Time-series of A) Nitrous Oxide (N_2O) concentrations in nmol L^{-1} , B) N_2O saturation percentage ($\text{Sat\%N}_2\text{O}$, %), C) Temperature ($^{\circ}\text{C}$), D) Chlorophyll *a* (*Chl a*) in $\mu\text{g L}^{-1}$, E) Dissolved Oxygen (DO) in $\mu\text{mol Kg}^{-1}$, F) Dissolved Organic Carbon (DOC) in μM , G) Nitrate (NO_3^-) in μM and H) Nitrite (NO_2^-) in μM measured in the samples collected in PB (blue dots), CA (red dots) and CS (green dots). Data for PB, CA, and CS were collected at depths of 1, 4, and 0.5 m, respectively. Figures were developed with the Python software version 3.12.3.




Con formato: Color de fuente: Texto 1

Con formato: Color de fuente: Texto 1

Con formato: Derecha: 0.63 cm

Figure 4: The relative importance of the leading environmental and biogeochemical variables driving the variability of nitrous oxide concentrations analyzed through cross-valued boosting (CVB). These are, in decreasing order, Temperature, Chlorophyll *a* (Chla), Salinity, Dissolved Organic Carbon (DOC), Nitrate (NO₃) and Nitrite (NO₂). The figure was developed using Python software version 3.12.3.




Eliminado: Figure 3: Time-series of A) Nitrous Oxide (N₂O) levels in nmol L⁻¹, B) Chlorophyll *a* (Chl *a*) in µg L⁻¹, C) Dissolved Organic Carbon (DOC) in µM, D) Dissolved Oxygen (DO) in µmol Kg⁻¹, E) Nitrate (NO₃⁻) in µM and F) Nitrite (NO₂⁻) in µM measured in the samples collected in BP (blue dots), CA (red dots) and CS (green dots). Figures were developed with the Python software version 3.12.3. ... [83]

Eliminado: :

Con formato: Color de fuente: Texto 1

Con formato: Color de fuente: Texto 1




Eliminado:

Con formato: Color de fuente: Texto 1

Con formato: Color de fuente: Texto 1

Con formato: Derecha: 0.63 cm

Figure 5: Linear relation Saturation Percentage of N₂O (Sat%N₂O) versus Temperature (°C), from the data obtained in PB (blue dots), CA (red dots) and CS (green dots). Linear equations are represented in blue for PB, red for CA and green for CS stations. Figures were developed with the Python software version 3.12.3.



Eliminado: ¶
¶
¶

Eliminado: of Temperature (°C) with


Eliminado: levels in percentage

Eliminado: BP

Con formato: Color de fuente: Texto 1

Con formato: Derecha: 0.63 cm

Figure 6: Bar chart of monthly averaged values of air-sea Nitrous Oxide (N_2O) transfer (FN_2O) in $\mu\text{mol m}^{-2} \text{d}^{-1}$ for all study years (left axis) and averaged temperature ($^{\circ}\text{C}$) values (right axis; grey line) for the PB (upper plot), CA (middle plot), and CS (lower plot) sites. Error bars represent $\pm$ standard deviation. Analysis of individual FN_2O flux measurements revealed that none significantly differed from zero ($p > 0.5$). Figures were developed using Python software version 3.12.3.



Eliminado: ¶

Con formato: Color de fuente: Texto 1

Con formato: Color de fuente: Texto 1

Con formato: Fuente: Cursiva

Con formato: Color de fuente: Texto 1

Con formato: Color de fuente: Texto 1

Eliminado: Figure 6: Box plot of monthly averaged values of air-sea Nitrous Oxide (N_2O) transfer (FN_2O) in $\mu\text{mol m}^{-2} \text{d}^{-1}$ for all the study years (left axis) and averaged temperature ($^{\circ}\text{C}$) values (right axis; grey line) for the PB (upper plot), CA (middle plot) and CS (lower plot) sites. Error bars represent $\pm$ standard deviation. Figures were developed with the Python software version 3.12.3.¶

Con formato: Color de fuente: Texto 1

Con formato: Derecha: 0.63 cm

701 *Table 1. Seasonally averaged values of temperature (°C), salinity (PSU), chlorophyll a (Chl a; $\mu\text{g L}^{-1}$),*
702 *dissolved oxygen (DO; $\mu\text{mol kg}^{-1}$), apparent oxygen utilization (AOU; $\mu\text{mol kg}^{-1}$), nitrate (μM), nitrite*
703 *(μM), silicate (μM), dissolved organic carbon (DOC; μM), N_2O (nanomol L^{-1}), and % saturation of N_2O*
704 *measured in samples collected in PB (Palma Bay), CA (Cabrera National Park), and CS (Cape Ses*
705 *Salines). Variability is represented as $\pm$ standard deviation. * DO data is only available from August 2022*
706 *at the CS station. N_2O saturation percentages significantly differ from 100%, as evaluated by a t-test*
707 *($p>0.005$).*

Con formato: Color de fuente: Texto 1

Con formato: Derecha: 0.63 cm

708
709

	PB (n = 48)				CA (n = 42)				CS (n =91)			
	Winter	Spring	Summer	Fall	Winter	Spring	Summer	Fall	Winter	Spring	Summer	Fall
Temperature (°C)	15.5± 0.8	16.6± 2.3	26.0± 2.0	24.5± 2.6	15.3± 0.9	16.5± 1.9	25.9± 2.0	23.1± 3.3	14.9± 1.0	16.9± 1.9	25.7± 2.5	23.0± 2.9
Salinity (PSU)	37.6± 0.2	37.7± 0.3	37.4± 0.2	37.5± 0.2	37.6± 0.2	37.4± 0.2	37.3± 0.3	37.4± 0.2	37.8± 0.2	37.7± 0.3	37.6± 0.3	37.6± 0.3
Chl a (µg L ⁻¹)	0.6± 0.2	0.3± 0.1	0.2± 0.1	0.2± 0.1	0.3± 0.1	0.2± 0.1	0.1± 0.0	0.3± 0.2	0.3± 0.2	0.3± 0.2	0.2± 0.1	0.2± 0.1
DO (µmol Kg ⁻¹)	246.7± 10.1	258.5± 15.7	218.1± 7.4	211.5± 9.0	248.7± 26.8	262.4± 15.0	213.4± 9.0	218.4± 11.8	265.6± 10.4*	244.0± 6.8*	212.6± 9.6*	220.4± 5.8*
AOU (µmol Kg ⁻¹)	-5.5± 7.9	-23.0± 16.1	-17.8± 5.3	-5.4± 5.5	-6.6± 24.8	-26.5± 13.5	-12.4± 6.9	-6.5± 7.4	-17.3± 12.1*	-13.9± 4.0*	-16.7± 4.9*	-13.9± 2.3*
Nitrate (µM)	0.2± 0.4	0.2± 0.3	0.1± 0.0	0.4± 0.6	0.2± 0.2	0.1± 0.2	0.1± 0.0	0.2± 0.3	1.4± 1.6	1.9± 1.6	1.4± 1.5	1.4± 1.5
Nitrite (µM)	0.06± 0.03	0.03± 0.02	0.05± 0.04	0.07± 0.1	0.03± 0.02	0.04± 0.04	0.03± 0.02	0.03± 0.02	0.05± 0.05	0.05± 0.06	0.05± 0.03	0.05± 0.04
Silicate (µM)	2.9± 2.8	4.4± 3.6	6.7± 5.5	8.1± 5.3	4.0± 3.1	3.1± 2.3	7.6± 6.8	5.1± 3.9	0.9± 0.9	2.0± 2.6	1.4± 1.7	0.8± 0.9
DOC (µM)	76.5± 14.2	65.9± 10.7	80.8± 13.2	81.1± 16.3	75.4± 17.3	71.0± 9.5	76.6± 13.8	78.3± 11.7	74.0± 12.6	71.8± 10.5	88.0± 14.3	84.1± 14.4
N ₂ O (nmol L ⁻¹)	9.2± 0.3	9.0± 0.5	7.20± 0.5	7.4± 0.4	9.0± 0.4	8.7± 0.4	7.1± 0.3	7.4± 0.6	9.0± 0.3	8.7± 0.5	7.1± 0.5	7.4± 0.5
N ₂ O saturation (%)	101.8± 3.3	102.6± 4.4	109.6± 2.8	107.3± 3.2	98.6± 3.4	97.9± 2.7	106.6± 3.2	103.8± 3.0	99.0± 2.0	100.1± 4.9	106.9± 2.9	104.6± 2.5

Eliminado: Table 1: Seasonally averaged values of Temperature(°C), Salinity (PSU), Chlorophyll *a* (Chl *a*; µg L⁻¹), Dissolved Oxygen (DO; µmol Kg⁻¹), Apparent Oxygen Utilization (AOU; µmol Kg⁻¹), Nitrate (µM), Nitrite (µM), Silicate (µM), Dissolved Organic Carbon (DOC; µM), N₂O (nmol), % saturation N₂O measured in the samples collected in BP (Buoy of Palma), CA (Cabrera National Park) and CS (Cape Ses Salines). Variability is represented as ± standard deviation. * DO data is only available from August 2022 at the CS station.

28

Eliminado:

Con formato

Con formato: Derecha: 0.63 cm

... [84]

Table 2. Annual N₂O flux ($\mu\text{mol m}^{-2} \text{ y}^{-1}$) per station since 2019 for PB (Buoy of Palma) and CS (Cape Ses Salines) stations, and from 2020 for the CA (Cabrera National Park) site, obtained using the Dobashi & Ho, 2023; Wanninkhof, 2014; and Cole & Caraco, 1998 gas transfer parameterizations. Variability is represented as $\pm$ propagated error.

		Annual N ₂ O flux ($\mu\text{mol m}^{-2} \text{ y}^{-1}$)		
Station	Year	Dobashi & Ho, 2023	Wanninkhof, 2014	Cole & Caraco, 1998
PB	2019	105.4 $\pm$ 42.1	186.7 $\pm$ 74.4	195.7 $\pm$ 32.1
	2020	107.4 $\pm$ 25.5	190.1 $\pm$ 45.2	204.2 $\pm$ 20.2
	2021	32.4 $\pm$ 7.0	57.4 $\pm$ 12.4	68.2 $\pm$ 5.9
	2022	60.2 $\pm$ 10.0	106.8 $\pm$ 17.7	124.0 $\pm$ 8.2
	2023	44.6 $\pm$ 12.3	79.1 $\pm$ 21.7	91.3 $\pm$ 9.6
CA	2020	58.4 $\pm$ 17.7	103.4 $\pm$ 31.3	128.6 $\pm$ 14.3
	2021	15.0 $\pm$ 7.7	26.6 $\pm$ 13.7	31.7 $\pm$ 6.3
	2022	23.6 $\pm$ 7.5	42.0 $\pm$ 13.2	58.0 $\pm$ 6.0
	2023	7.3 $\pm$ 8.8	13.1 $\pm$ 15.6	12.0 $\pm$ 7.0
CS	2019	45.8 $\pm$ 13.1	81.2 $\pm$ 23.3	86.5 $\pm$ 10.4
	2020	48.4 $\pm$ 6.6	85.8 $\pm$ 11.7	98.2 $\pm$ 5.3
	2021	6.3 $\pm$ 6.2	11.1 $\pm$ 11.0	16.5 $\pm$ 5.2
	2022	49.2 $\pm$ 15.1	87.3 $\pm$ 26.7	113.3 $\pm$ 12.1
	2023	5.4 $\pm$ 6.3	9.7 $\pm$ 11.1	8.7 $\pm$ 4.9

- Eliminado: ¶
- Con formato ... [86]
- Eliminado: Table 2: Annual N₂O flux ($\mu\text{mol m}^{-2} \text{ y}^{-1}$) per ... [87]
- Con formato ... [88]
- Con formato ... [89]
- Con formato ... [91]
- Con formato ... [92]
- Con formato ... [93]
- Con formato ... [94]
- Con formato ... [95]
- Con formato ... [96]
- Con formato ... [97]
- Con formato ... [98]
- Con formato ... [99]
- Con formato ... [90]
- Con formato ... [100]
- Con formato ... [101]
- Con formato ... [102]
- Con formato ... [103]
- Con formato ... [104]
- Con formato ... [105]
- Con formato ... [107]
- Con formato ... [108]
- Con formato ... [109]
- Con formato ... [110]
- Con formato ... [111]
- Con formato ... [112]
- Con formato ... [106]
- Con formato ... [113]
- Con formato ... [114]
- Con formato ... [115]
- Con formato ... [116]
- Con formato ... [117]
- Con formato ... [118]
- Con formato ... [120]
- Con formato ... [121]
- Con formato ... [122]
- Con formato ... [123]
- Con formato ... [124]
- Con formato ... [125]
- Con formato ... [126]
- Con formato ... [127]
- Con formato ... [128]
- Con formato ... [119]
- Con formato ... [129]
- Con formato ... [130]
- Con formato ... [131]
- Con formato ... [132]
- Con formato ... [133]
- Con formato ... [134]
- Eliminado: ¶ ... [135]
- Con formato ... [136]
- Con formato ... [85]

Página 3: [1] Eliminado **Susana Flecha** **7/4/25 10:12:00**

✖
▲
Página 3: [1] Eliminado **Susana Flecha** **7/4/25 10:12:00**

✖
▲
Página 3: [2] Eliminado **Susana Flecha** **28/3/25 9:47:00**

✖
▲
Página 3: [2] Eliminado **Susana Flecha** **28/3/25 9:47:00**

✖
▲
Página 3: [3] Eliminado **Susana Flecha** **27/3/25 16:11:00**

✖
▲
Página 3: [3] Eliminado **Susana Flecha** **27/3/25 16:11:00**

✖
▲
Página 3: [4] Eliminado **Susana Flecha** **27/3/25 16:07:00**

✖
▲
Página 3: [4] Eliminado **Susana Flecha** **27/3/25 16:07:00**

✖
▲
Página 3: [4] Eliminado **Susana Flecha** **27/3/25 16:07:00**

✖
▲
Página 3: [5] Eliminado **Susana Flecha** **27/3/25 16:11:00**

✖
▲
Página 3: [5] Eliminado **Susana Flecha** **27/3/25 16:11:00**

✖
▲
Página 3: [5] Eliminado **Susana Flecha** **27/3/25 16:11:00**

✖
▲
Página 3: [5] Eliminado **Susana Flecha** **27/3/25 16:11:00**

✖
▲
Página 3: [5] Eliminado **Susana Flecha** **27/3/25 16:11:00**

✖
▲
Página 3: [5] Eliminado **Susana Flecha** **27/3/25 16:11:00**

✖
▲
Página 3: [5] Eliminado **Susana Flecha** **27/3/25 16:11:00**

✖

▲

Página 3: [6] Eliminado	Susana Flecha	27/3/25 16:11:00
--------------------------------	----------------------	-------------------------

✖

▲

Página 3: [6] Eliminado	Susana Flecha	27/3/25 16:11:00
--------------------------------	----------------------	-------------------------

✖

▲

Página 3: [6] Eliminado	Susana Flecha	27/3/25 16:11:00
--------------------------------	----------------------	-------------------------

✖

▲

Página 3: [7] Eliminado	Susana Flecha	8/4/25 14:46:00
--------------------------------	----------------------	------------------------

✖

▲

Página 3: [7] Eliminado	Susana Flecha	8/4/25 14:46:00
--------------------------------	----------------------	------------------------

✖

▲

Página 3: [8] Eliminado	Susana Flecha	27/3/25 16:07:00
--------------------------------	----------------------	-------------------------

✖

▲

Página 3: [8] Eliminado	Susana Flecha	27/3/25 16:07:00
--------------------------------	----------------------	-------------------------

✖

▲

Página 3: [9] Eliminado	Susana Flecha	27/3/25 16:07:00
--------------------------------	----------------------	-------------------------

✖

▲

Página 3: [9] Eliminado	Susana Flecha	27/3/25 16:07:00
--------------------------------	----------------------	-------------------------

✖

▲

Página 3: [9] Eliminado	Susana Flecha	27/3/25 16:07:00
--------------------------------	----------------------	-------------------------

✖

▲

Página 3: [10] Eliminado	Susana Flecha	27/3/25 16:07:00
---------------------------------	----------------------	-------------------------

✖

▲

Página 3: [10] Eliminado	Susana Flecha	27/3/25 16:07:00
---------------------------------	----------------------	-------------------------

✖

▲

Página 3: [10] Eliminado	Susana Flecha	27/3/25 16:07:00
---------------------------------	----------------------	-------------------------

✖

▲

Página 3: [11] Eliminado	Susana Flecha	27/3/25 16:08:00
---------------------------------	----------------------	-------------------------

✖

▲

Página 3: [11] Eliminado	Susana Flecha	27/3/25 16:08:00
---------------------------------	----------------------	-------------------------

✖

▲

Página 3: [12] Eliminado	Susana Flecha	25/3/25 9:21:00
---------------------------------	----------------------	------------------------

✖

▲

Página 3: [12] Eliminado	Susana Flecha	25/3/25 9:21:00
---------------------------------	----------------------	------------------------

✖

▲

Página 3: [12] Eliminado	Susana Flecha	25/3/25 9:21:00
---------------------------------	----------------------	------------------------

✖

▲

Página 3: [12] Eliminado	Susana Flecha	25/3/25 9:21:00
---------------------------------	----------------------	------------------------

✖

▲

Página 3: [13] Con formato	Susana Flecha	8/4/25 14:53:00
-----------------------------------	----------------------	------------------------

Fuente: Cursiva

▲ **Página 3: [13] Con formato** **Susana Flecha** **8/4/25 14:53:00**

Fuente: Cursiva

▲ **Página 3: [13] Con formato** **Susana Flecha** **8/4/25 14:53:00**

Fuente: Cursiva

▲ **Página 3: [13] Con formato** **Susana Flecha** **8/4/25 14:53:00**

Fuente: Cursiva

▲ **Página 3: [14] Eliminado** **Susana Flecha** **25/3/25 9:21:00**

✖
▲

Página 3: [14] Eliminado **Susana Flecha** **25/3/25 9:21:00**

✖
▲

Página 3: [14] Eliminado **Susana Flecha** **25/3/25 9:21:00**

✖
▲

Página 3: [15] Con formato **Susana Flecha** **7/4/25 10:22:00**

Color de fuente: Texto 1

▲ **Página 3: [15] Con formato** **Susana Flecha** **7/4/25 10:22:00**

Color de fuente: Texto 1

▲ **Página 1: [16] Con formato** **Susana Flecha** **6/5/25 16:58:00**

Derecha: 0.63 cm

▲ **Página 5: [17] Con formato** **Susana Flecha** **7/4/25 10:22:00**

Color de fuente: Texto 1

▲ **Página 5: [18] Con formato** **Susana Flecha** **7/4/25 10:22:00**

Color de fuente: Texto 1

▲ **Página 5: [19] Con formato** **Susana Flecha** **7/4/25 10:22:00**

Color de fuente: Texto 1

▲ **Página 5: [20] Con formato** **Susana Flecha** **7/4/25 10:22:00**

Color de fuente: Texto 1

▲ **Página 5: [21] Con formato** **Susana Flecha** **7/4/25 10:22:00**

Color de fuente: Texto 1

▲
4

Página 5: [22] Con formato	Susana Flecha	7/4/25 10:22:00
-----------------------------------	----------------------	------------------------

Color de fuente: Texto 1

Página 5: [23] Con formato	Susana Flecha	7/4/25 10:22:00
-----------------------------------	----------------------	------------------------

Color de fuente: Texto 1

Página 5: [24] Con formato	Susana Flecha	7/4/25 10:22:00
-----------------------------------	----------------------	------------------------

Color de fuente: Texto 1

Página 5: [25] Con formato	Susana Flecha	7/4/25 10:22:00
-----------------------------------	----------------------	------------------------

Color de fuente: Texto 1

Página 5: [26] Con formato	Susana Flecha	7/4/25 10:22:00
-----------------------------------	----------------------	------------------------

Color de fuente: Texto 1

Página 5: [27] Con formato	Susana Flecha	7/4/25 10:22:00
-----------------------------------	----------------------	------------------------

Color de fuente: Texto 1

Página 5: [28] Con formato	Susana Flecha	7/4/25 10:22:00
-----------------------------------	----------------------	------------------------

Color de fuente: Texto 1

Página 5: [29] Con formato	Susana Flecha	7/4/25 10:22:00
-----------------------------------	----------------------	------------------------

Color de fuente: Texto 1

Página 5: [30] Con formato	Susana Flecha	7/4/25 10:22:00
-----------------------------------	----------------------	------------------------

Color de fuente: Texto 1

Página 5: [31] Con formato	Susana Flecha	7/4/25 10:22:00
-----------------------------------	----------------------	------------------------

Color de fuente: Texto 1

Página 6: [32] Eliminado	Susana Flecha	26/3/25 13:11:00
---------------------------------	----------------------	-------------------------

Página 6: [32] Eliminado	Susana Flecha	26/3/25 13:11:00
---------------------------------	----------------------	-------------------------

Página 6: [32] Eliminado	Susana Flecha	26/3/25 13:11:00
---------------------------------	----------------------	-------------------------

Página 6: [32] Eliminado	Susana Flecha	26/3/25 13:11:00
---------------------------------	----------------------	-------------------------

Página 6: [32] Eliminado	Susana Flecha	26/3/25 13:11:00
---------------------------------	----------------------	-------------------------

Página 6: [32] Eliminado	Susana Flecha	26/3/25 13:11:00
---------------------------------	----------------------	-------------------------

▲
Página 6: [34] Eliminado Susana Flecha 28/3/25 9:57:00

▲
Página 6: [34] Eliminado Susana Flecha 28/3/25 9:57:00

▲
Página 6: [34] Eliminado Susana Flecha 28/3/25 9:57:00

▲
Página 6: [34] Eliminado Susana Flecha 28/3/25 9:57:00

▲
Página 6: [34] Eliminado Susana Flecha 28/3/25 9:57:00

▲
Página 6: [34] Eliminado Susana Flecha 28/3/25 9:57:00

▲
Página 6: [34] Eliminado Susana Flecha 28/3/25 9:57:00

▲
Página 6: [34] Eliminado Susana Flecha 28/3/25 9:57:00

▲
Página 6: [34] Eliminado Susana Flecha 28/3/25 9:57:00

▲
Página 6: [34] Eliminado Susana Flecha 28/3/25 9:57:00

▲
Página 6: [34] Eliminado Susana Flecha 28/3/25 9:57:00

▲
Página 6: [34] Eliminado Susana Flecha 28/3/25 9:57:00

▲
Página 6: [34] Eliminado Susana Flecha 28/3/25 9:57:00

▲
Página 6: [34] Eliminado Susana Flecha 28/3/25 9:57:00

▲
Página 6: [34] Eliminado Susana Flecha 28/3/25 9:57:00

▲
x

Página 6: [34] Eliminado	Susana Flecha	28/3/25 9:57:00
--------------------------	---------------	-----------------

▲
x

Página 6: [34] Eliminado	Susana Flecha	28/3/25 9:57:00
--------------------------	---------------	-----------------

▲
x

Página 6: [35] Con formato	Susana Flecha	7/4/25 10:22:00
----------------------------	---------------	-----------------

Fuente: Cursiva, Color de fuente: Texto 1

▲
x

Página 6: [35] Con formato	Susana Flecha	7/4/25 10:22:00
----------------------------	---------------	-----------------

Fuente: Cursiva, Color de fuente: Texto 1

▲
x

Página 6: [36] Eliminado	Susana Flecha	28/3/25 9:57:00
--------------------------	---------------	-----------------

▲
x

Página 6: [36] Eliminado	Susana Flecha	28/3/25 9:57:00
--------------------------	---------------	-----------------

▲
x

Página 6: [36] Eliminado	Susana Flecha	28/3/25 9:57:00
--------------------------	---------------	-----------------

▲
x

Página 6: [36] Eliminado	Susana Flecha	28/3/25 9:57:00
--------------------------	---------------	-----------------

▲
x

Página 6: [36] Eliminado	Susana Flecha	28/3/25 9:57:00
--------------------------	---------------	-----------------

▲
x

Página 6: [37] Eliminado	Susana Flecha	28/3/25 9:57:00
--------------------------	---------------	-----------------

▲
x

Página 6: [37] Eliminado	Susana Flecha	28/3/25 9:57:00
--------------------------	---------------	-----------------

▲
x

Página 6: [37] Eliminado	Susana Flecha	28/3/25 9:57:00
--------------------------	---------------	-----------------

▲
x

Página 6: [37] Eliminado	Susana Flecha	28/3/25 9:57:00
--------------------------	---------------	-----------------

▲
x

Página 6: [37] Eliminado	Susana Flecha	28/3/25 9:57:00
--------------------------	---------------	-----------------

▲
x

Página 6: [38] Eliminado	Susana Flecha	28/3/25 9:58:00
--------------------------	---------------	-----------------

▲
x

Página 6: [38] Eliminado	Susana Flecha	28/3/25 9:58:00
--------------------------	---------------	-----------------

▲	Página 6: [38] Eliminado	Susana Flecha	28/3/25 9:58:00
✖			
▲	Página 6: [38] Eliminado	Susana Flecha	28/3/25 9:58:00
✖			
▲	Página 7: [39] Eliminado	Susana Flecha	28/3/25 9:58:00
✖			◀
▲	Página 7: [39] Eliminado	Susana Flecha	28/3/25 9:58:00
✖			◀
▲	Página 7: [39] Eliminado	Susana Flecha	28/3/25 9:58:00
✖			◀
▲	Página 7: [39] Eliminado	Susana Flecha	28/3/25 9:58:00
✖			◀
▲	Página 7: [39] Eliminado	Susana Flecha	28/3/25 9:58:00
✖			◀
▲	Página 7: [39] Eliminado	Susana Flecha	28/3/25 9:58:00
✖			◀
▲	Página 7: [39] Eliminado	Susana Flecha	28/3/25 9:58:00
✖			◀
▲	Página 7: [39] Eliminado	Susana Flecha	28/3/25 9:58:00
✖			◀
▲	Página 7: [40] Eliminado	Susana Flecha	28/3/25 9:58:00
✖			
▲	Página 7: [40] Eliminado	Susana Flecha	28/3/25 9:58:00
✖			
▲	Página 7: [41] Eliminado	Susana Flecha	27/3/25 16:35:00
✖			
▲	Página 7: [41] Eliminado	Susana Flecha	27/3/25 16:35:00
✖			
▲	Página 7: [42] Eliminado	Susana Flecha	28/3/25 9:58:00
✖			
▲	Página 7: [42] Eliminado	Susana Flecha	28/3/25 9:58:00

▲	Página 7: [43] Eliminado	Susana Flecha 27/3/25 16:35:00
■		
▲	Página 7: [43] Eliminado	Susana Flecha 27/3/25 16:35:00
■		
▲	Página 7: [44] Eliminado	Susana Flecha 28/3/25 9:58:00
■		
▲	Página 7: [44] Eliminado	Susana Flecha 28/3/25 9:58:00
■		
▲	Página 7: [44] Eliminado	Susana Flecha 28/3/25 9:58:00
■		
▲	Página 7: [44] Eliminado	Susana Flecha 28/3/25 9:58:00
■		
▲	Página 7: [44] Eliminado	Susana Flecha 28/3/25 9:58:00
■		
▲	Página 7: [44] Eliminado	Susana Flecha 28/3/25 9:58:00
■		
▲	Página 7: [45] Eliminado	Susana Flecha 28/3/25 9:59:00
■		
▲	Página 7: [45] Eliminado	Susana Flecha 28/3/25 9:59:00
■		
▲	Página 7: [45] Eliminado	Susana Flecha 28/3/25 9:59:00
■		
▲	Página 7: [45] Eliminado	Susana Flecha 28/3/25 9:59:00
■		
▲	Página 7: [45] Eliminado	Susana Flecha 28/3/25 9:59:00
■		
▲	Página 7: [45] Eliminado	Susana Flecha 28/3/25 9:59:00
■		
▲	Página 7: [45] Eliminado	Susana Flecha 28/3/25 9:59:00
■		
▲	Página 7: [45] Eliminado	Susana Flecha 28/3/25 9:59:00
■		
▲	Página 7: [45] Eliminado	Susana Flecha 28/3/25 9:59:00
■		

▲
x
▲
Página 7: [45] Eliminado **Susana Flecha** **28/3/25 9:59:00**

▲
x
▲
Página 7: [45] Eliminado **Susana Flecha** **28/3/25 9:59:00**

▲
x
▲
Página 7: [45] Eliminado **Susana Flecha** **28/3/25 9:59:00**

▲
x
▲
Página 7: [45] Eliminado **Susana Flecha** **28/3/25 9:59:00**

▲
x
▲
Página 7: [45] Eliminado **Susana Flecha** **28/3/25 9:59:00**

▲
x
▲
Página 7: [46] Con formato **Susana Flecha** **7/4/25 10:22:00**

Color de fuente: Texto 1

▲
x
▲
Página 7: [46] Con formato **Susana Flecha** **7/4/25 10:22:00**

Color de fuente: Texto 1

▲
x
▲
Página 7: [47] Eliminado **Susana Flecha** **28/3/25 10:00:00**

▲
x
▲
Página 7: [47] Eliminado **Susana Flecha** **28/3/25 10:00:00**

▲
x
▲
Página 7: [48] Eliminado **Susana Flecha** **28/3/25 10:00:00**

▲
x
▲
Página 7: [48] Eliminado **Susana Flecha** **28/3/25 10:00:00**

▲
x
▲
Página 7: [48] Eliminado **Susana Flecha** **28/3/25 10:00:00**

▲
x
▲
Página 7: [48] Eliminado **Susana Flecha** **28/3/25 10:00:00**

▲
x
▲
Página 7: [48] Eliminado **Susana Flecha** **28/3/25 10:00:00**

▲
x
▲
Página 7: [48] Eliminado **Susana Flecha** **28/3/25 10:00:00**

▲
x
▲
Página 7: [48] Eliminado **Susana Flecha** **28/3/25 10:00:00**

▲	Página 7: [48] Eliminado	Susana Flecha	28/3/25 10:00:00
✖			
▲	Página 7: [48] Eliminado	Susana Flecha	28/3/25 10:00:00
✖			
▲	Página 8: [49] Eliminado	Susana Flecha	26/3/25 18:32:00
✖			
▲	Página 8: [49] Eliminado	Susana Flecha	26/3/25 18:32:00
✖			
▲	Página 8: [50] Eliminado	Susana Flecha	28/3/25 10:58:00
✖			
▲	Página 8: [50] Eliminado	Susana Flecha	28/3/25 10:58:00
✖			
▲	Página 8: [50] Eliminado	Susana Flecha	28/3/25 10:58:00
✖			
▲	Página 8: [50] Eliminado	Susana Flecha	28/3/25 10:58:00
✖			
▲	Página 8: [50] Eliminado	Susana Flecha	28/3/25 10:58:00
✖			
▲	Página 8: [50] Eliminado	Susana Flecha	28/3/25 10:58:00
✖			
▲	Página 8: [50] Eliminado	Susana Flecha	28/3/25 10:58:00
✖			
▲	Página 8: [50] Eliminado	Susana Flecha	28/3/25 10:58:00
✖			
▲	Página 8: [51] Eliminado	Susana Flecha	28/3/25 10:59:00
✖			
▲	Página 8: [51] Eliminado	Susana Flecha	28/3/25 10:59:00

▲
Página 8: [51] Eliminado Susana Flecha 28/3/25 10:59:00

▲
Página 8: [51] Eliminado Susana Flecha 28/3/25 10:59:00

▲
Página 8: [51] Eliminado Susana Flecha 28/3/25 10:59:00

▲
Página 8: [51] Eliminado Susana Flecha 28/3/25 10:59:00

▲
Página 8: [51] Eliminado Susana Flecha 28/3/25 10:59:00

▲
Página 8: [51] Eliminado Susana Flecha 28/3/25 10:59:00

▲
Página 8: [51] Eliminado Susana Flecha 28/3/25 10:59:00

▲
Página 8: [51] Eliminado Susana Flecha 28/3/25 10:59:00

▲
Página 8: [51] Eliminado Susana Flecha 28/3/25 10:59:00

▲
Página 8: [51] Eliminado Susana Flecha 28/3/25 10:59:00

▲
Página 8: [51] Eliminado Susana Flecha 28/3/25 10:59:00

▲
Página 8: [51] Eliminado Susana Flecha 28/3/25 10:59:00

▲
Página 8: [51] Eliminado Susana Flecha 28/3/25 10:59:00

▲
Página 8: [51] Eliminado Susana Flecha 28/3/25 10:59:00

▲
Página 8: [51] Eliminado Susana Flecha 28/3/25 10:59:00

▲	Página 8: [53] Eliminado	Susana Flecha	28/3/25 11:00:00
✖			
▲	Página 8: [53] Eliminado	Susana Flecha	28/3/25 11:00:00
✖			
▲	Página 8: [53] Eliminado	Susana Flecha	28/3/25 11:00:00
✖			
▲	Página 8: [53] Eliminado	Susana Flecha	28/3/25 11:00:00
✖			
▲	Página 8: [53] Eliminado	Susana Flecha	28/3/25 11:00:00
✖			
▲	Página 8: [53] Eliminado	Susana Flecha	28/3/25 11:00:00
✖			
▲	Página 8: [53] Eliminado	Susana Flecha	28/3/25 11:00:00
✖			
▲	Página 8: [53] Eliminado	Susana Flecha	28/3/25 11:00:00
✖			
▲	Página 8: [53] Eliminado	Susana Flecha	28/3/25 11:00:00
✖			
▲	Página 8: [54] Eliminado	Susana Flecha	26/3/25 18:47:00
✖			
▲	Página 8: [54] Eliminado	Susana Flecha	26/3/25 18:47:00
✖			
▲	Página 8: [54] Eliminado	Susana Flecha	26/3/25 18:47:00
✖			
▲	Página 8: [55] Con formato	Susana Flecha	7/4/25 10:22:00
	Color de fuente: Texto 1, Subíndice		
▲	Página 8: [55] Con formato	Susana Flecha	7/4/25 10:22:00
	Color de fuente: Texto 1, Subíndice		
▲	Página 8: [56] Eliminado	Susana Flecha	26/3/25 18:47:00

▲
Página 8: [56] Eliminado Susana Flecha 26/3/25 18:47:00

▲
Página 8: [56] Eliminado Susana Flecha 26/3/25 18:47:00

▲
Página 8: [57] Eliminado Susana Flecha 8/4/25 15:49:00

▲
Página 8: [57] Eliminado Susana Flecha 8/4/25 15:49:00

▲
Página 8: [58] Eliminado Susana Flecha 28/3/25 11:01:00

▲
Página 8: [58] Eliminado Susana Flecha 28/3/25 11:01:00

▲
Página 8: [58] Eliminado Susana Flecha 28/3/25 11:01:00

▲
Página 8: [58] Eliminado Susana Flecha 28/3/25 11:01:00

▲
Página 8: [58] Eliminado Susana Flecha 28/3/25 11:01:00

▲
Página 8: [58] Eliminado Susana Flecha 28/3/25 11:01:00

▲
Página 8: [58] Eliminado Susana Flecha 28/3/25 11:01:00

▲
Página 8: [58] Eliminado Susana Flecha 28/3/25 11:01:00

▲
Página 8: [58] Eliminado Susana Flecha 28/3/25 11:01:00

▲
Página 8: [58] Eliminado Susana Flecha 28/3/25 11:01:00

▲
Página 8: [59] Con formato Susana Flecha 7/4/25 10:22:00

Color de fuente: Texto 1, Subíndice

Página 8: [59] Con formato	Susana Flecha	7/4/25 10:22:00
-----------------------------------	----------------------	------------------------

Color de fuente: Texto 1, Subíndice

Página 10: [60] Eliminado	Susana Flecha	28/3/25 11:03:00
----------------------------------	----------------------	-------------------------

Página 10: [60] Eliminado	Susana Flecha	28/3/25 11:03:00
----------------------------------	----------------------	-------------------------

Página 10: [61] Con formato	Susana Flecha	7/4/25 10:22:00
------------------------------------	----------------------	------------------------

Fuente: Cursiva, Color de fuente: Texto 1

Página 10: [61] Con formato	Susana Flecha	7/4/25 10:22:00
------------------------------------	----------------------	------------------------

Fuente: Cursiva, Color de fuente: Texto 1

Página 10: [62] Eliminado	Susana Flecha	7/4/25 9:06:00
----------------------------------	----------------------	-----------------------

Página 10: [62] Eliminado	Susana Flecha	7/4/25 9:06:00
----------------------------------	----------------------	-----------------------

Página 10: [62] Eliminado	Susana Flecha	7/4/25 9:06:00
----------------------------------	----------------------	-----------------------

Página 10: [62] Eliminado	Susana Flecha	7/4/25 9:06:00
----------------------------------	----------------------	-----------------------

Página 10: [62] Eliminado	Susana Flecha	7/4/25 9:06:00
----------------------------------	----------------------	-----------------------

Página 10: [62] Eliminado	Susana Flecha	7/4/25 9:06:00
----------------------------------	----------------------	-----------------------

Página 10: [62] Eliminado	Susana Flecha	7/4/25 9:06:00
----------------------------------	----------------------	-----------------------

Página 10: [62] Eliminado	Susana Flecha	7/4/25 9:06:00
----------------------------------	----------------------	-----------------------

Página 10: [62] Eliminado	Susana Flecha	7/4/25 9:06:00
----------------------------------	----------------------	-----------------------

Página 10: [62] Eliminado	Susana Flecha	7/4/25 9:06:00
----------------------------------	----------------------	-----------------------

Página 10: [62] Eliminado	Susana Flecha	7/4/25 9:06:00
----------------------------------	----------------------	-----------------------

▲	Página 10: [62] Eliminado	Susana Flecha	7/4/25 9:06:00
✖			
▲	Página 10: [62] Eliminado	Susana Flecha	7/4/25 9:06:00
✖			
▲	Página 10: [62] Eliminado	Susana Flecha	7/4/25 9:06:00
✖			
▲	Página 10: [63] Eliminado	Susana Flecha	28/3/25 11:04:00
✖			
▲	Página 10: [63] Eliminado	Susana Flecha	28/3/25 11:04:00
✖			
▲	Página 10: [63] Eliminado	Susana Flecha	28/3/25 11:04:00
✖			
▲	Página 10: [63] Eliminado	Susana Flecha	28/3/25 11:04:00
✖			
▲	Página 10: [63] Eliminado	Susana Flecha	28/3/25 11:04:00
✖			
▲	Página 10: [63] Eliminado	Susana Flecha	28/3/25 11:04:00
✖			
▲	Página 10: [63] Eliminado	Susana Flecha	28/3/25 11:04:00
✖			
▲	Página 10: [63] Eliminado	Susana Flecha	28/3/25 11:04:00
✖			
▲	Página 10: [63] Eliminado	Susana Flecha	28/3/25 11:04:00
✖			
▲	Página 10: [64] Eliminado	Susana Flecha	28/3/25 11:04:00
✖			
▲	Página 10: [64] Eliminado	Susana Flecha	28/3/25 11:04:00
✖			
▲	Página 10: [64] Eliminado	Susana Flecha	28/3/25 11:04:00

▲
Página 10: [64] Eliminado Susana Flecha 28/3/25 11:04:00

▲
Página 10: [64] Eliminado Susana Flecha 28/3/25 11:04:00

▲
Página 10: [64] Eliminado Susana Flecha 28/3/25 11:04:00

▲
Página 10: [64] Eliminado Susana Flecha 28/3/25 11:04:00

▲
Página 10: [64] Eliminado Susana Flecha 28/3/25 11:04:00

▲
Página 10: [65] Eliminado Susana Flecha 28/3/25 11:05:00

▲
Página 10: [65] Eliminado Susana Flecha 28/3/25 11:05:00

▲
Página 10: [65] Eliminado Susana Flecha 28/3/25 11:05:00

▲
Página 10: [65] Eliminado Susana Flecha 28/3/25 11:05:00

▲
Página 10: [65] Eliminado Susana Flecha 28/3/25 11:05:00

▲
Página 10: [65] Eliminado Susana Flecha 28/3/25 11:05:00

▲
Página 10: [65] Eliminado Susana Flecha 28/3/25 11:05:00

▲
Página 10: [65] Eliminado Susana Flecha 28/3/25 11:05:00

▲
Página 10: [65] Eliminado Susana Flecha 28/3/25 11:05:00

▲
Página 10: [65] Eliminado Susana Flecha 28/3/25 11:05:00

▲ **Página 10: [65] Eliminado** Susana Flecha 28/3/25 11:05:00

▼

▲ **Página 10: [65] Eliminado** Susana Flecha 28/3/25 11:05:00

▼

▲ **Página 10: [65] Eliminado** Susana Flecha 28/3/25 11:05:00

▼

▲ **Página 10: [65] Eliminado** Susana Flecha 28/3/25 11:05:00

▼

▲ **Página 10: [66] Eliminado** Susana Flecha 28/3/25 11:06:00

▼

▲ **Página 10: [66] Eliminado** Susana Flecha 28/3/25 11:06:00

▼

▲ **Página 10: [66] Eliminado** Susana Flecha 28/3/25 11:06:00

▼

▲ **Página 10: [66] Eliminado** Susana Flecha 28/3/25 11:06:00

▼

▲ **Página 1: [67] Con formato** Susana Flecha 6/5/25 16:58:00

Derecha: 0.63 cm

▲ **Página 11: [68] Con formato** Susana Flecha 7/4/25 10:22:00

Color de fuente: Texto 1, Subíndice

▲ **Página 11: [69] Con formato** Susana Flecha 7/4/25 10:22:00

Color de fuente: Texto 1

▲ **Página 11: [70] Con formato** Susana Flecha 7/4/25 10:22:00

Fuente: Sin Negrita, Color de fuente: Texto 1

▲ **Página 11: [71] Con formato** Susana Flecha 7/4/25 10:22:00

Color de fuente: Texto 1

▲ **Página 11: [72] Con formato** Susana Flecha 7/4/25 10:22:00

Fuente: Sin Negrita, Color de fuente: Texto 1

▲ **Página 11: [73] Con formato** Susana Flecha 7/4/25 10:22:00

Color de fuente: Texto 1

▲ **Página 11: [74] Con formato** Susana Flecha 7/4/25 10:22:00

Fuente: Sin Negrita, Color de fuente: Texto 1

▲ **Página 11: [75] Con formato** Susana Flecha 7/4/25 10:22:00

Color de fuente: Texto 1

▲ **Página 11: [76] Con formato** Susana Flecha 7/4/25 10:22:00

Fuente: Sin Negrita, Color de fuente: Texto 1

▲ **Página 11: [77] Con formato** Susana Flecha 7/4/25 10:22:00

Color de fuente: Texto 1

▲ **Página 11: [78] Con formato** Susana Flecha 7/4/25 10:22:00

Fuente: Sin Negrita, Color de fuente: Texto 1

▲ **Página 11: [79] Con formato** Susana Flecha 7/4/25 10:22:00

Color de fuente: Texto 1

▲ **Página 11: [80] Con formato** Susana Flecha 7/4/25 10:22:00

Fuente: Cursiva, Color de fuente: Texto 1

▲ **Página 11: [81] Con formato** Susana Flecha 7/4/25 10:22:00

Color de fuente: Texto 1

▲ **Página 21: [82] Eliminado** Susana Flecha 27/3/25 11:37:00

▼

▲ **Página 24: [83] Eliminado** Susana Flecha 26/3/25 18:32:00

▲ **Página 28: [84] Con formato** Susana Flecha 7/4/25 10:22:00

Color de fuente: Texto 1

▲ **Página 28: [84] Con formato** Susana Flecha 7/4/25 10:22:00

Color de fuente: Texto 1

▲ **Página 1: [85] Con formato** Susana Flecha 6/5/25 16:58:00

Derecha: 0.63 cm

▲ **Página 29: [86] Con formato** Susana Flecha 7/4/25 10:22:00

Fuente: Sin Negrita, Color de fuente: Texto 1

▲ **Página 29: [87] Eliminado** Susana Flecha 27/3/25 12:48:00

▲ **Página 29: [88] Con formato** Susana Flecha 7/4/25 10:22:00

Color de fuente: Texto 1

▲ **Página 29: [89] Con formato** Susana Flecha 7/4/25 10:22:00

Color de fuente: Texto 1

Página 29: [90] Con formato	Susana Flecha	7/4/25 10:22:00
------------------------------------	----------------------	------------------------

Color de fuente: Texto 1

Página 29: [91] Con formato	Susana Flecha	7/4/25 10:22:00
------------------------------------	----------------------	------------------------

Color de fuente: Texto 1

Página 29: [91] Con formato	Susana Flecha	7/4/25 10:22:00
------------------------------------	----------------------	------------------------

Color de fuente: Texto 1

Página 29: [92] Con formato	Susana Flecha	7/4/25 10:22:00
------------------------------------	----------------------	------------------------

Color de fuente: Texto 1

Página 29: [92] Con formato	Susana Flecha	7/4/25 10:22:00
------------------------------------	----------------------	------------------------

Color de fuente: Texto 1

Página 29: [93] Con formato	Susana Flecha	7/4/25 10:22:00
------------------------------------	----------------------	------------------------

Color de fuente: Texto 1

Página 29: [93] Con formato	Susana Flecha	7/4/25 10:22:00
------------------------------------	----------------------	------------------------

Color de fuente: Texto 1

Página 29: [94] Con formato	Susana Flecha	7/4/25 10:22:00
------------------------------------	----------------------	------------------------

Color de fuente: Texto 1

Página 29: [94] Con formato	Susana Flecha	7/4/25 10:22:00
------------------------------------	----------------------	------------------------

Color de fuente: Texto 1

Página 29: [95] Con formato	Susana Flecha	7/4/25 10:22:00
------------------------------------	----------------------	------------------------

Color de fuente: Texto 1

Página 29: [95] Con formato	Susana Flecha	7/4/25 10:22:00
------------------------------------	----------------------	------------------------

Color de fuente: Texto 1

Página 29: [96] Con formato	Susana Flecha	7/4/25 10:22:00
------------------------------------	----------------------	------------------------

Color de fuente: Texto 1

Página 29: [96] Con formato	Susana Flecha	7/4/25 10:22:00
------------------------------------	----------------------	------------------------

Color de fuente: Texto 1

Página 29: [97] Con formato	Susana Flecha	7/4/25 10:22:00
------------------------------------	----------------------	------------------------

Color de fuente: Texto 1

Página 29: [97] Con formato	Susana Flecha	7/4/25 10:22:00
------------------------------------	----------------------	------------------------

Color de fuente: Texto 1

Página 29: [98] Con formato	Susana Flecha	7/4/25 10:22:00
------------------------------------	----------------------	------------------------

Color de fuente: Texto 1

▲ **Página 29: [98] Con formato** **Susana Flecha** **7/4/25 10:22:00**

Color de fuente: Texto 1

▲ **Página 29: [99] Con formato** **Susana Flecha** **7/4/25 10:22:00**

Color de fuente: Texto 1

▲ **Página 29: [99] Con formato** **Susana Flecha** **7/4/25 10:22:00**

Color de fuente: Texto 1

▲ **Página 29: [100] Con formato** **Susana Flecha** **7/4/25 10:22:00**

Color de fuente: Texto 1

▲ **Página 29: [100] Con formato** **Susana Flecha** **7/4/25 10:22:00**

Color de fuente: Texto 1

▲ **Página 29: [101] Con formato** **Susana Flecha** **7/4/25 10:22:00**

Color de fuente: Texto 1

▲ **Página 29: [101] Con formato** **Susana Flecha** **7/4/25 10:22:00**

Color de fuente: Texto 1

▲ **Página 29: [102] Con formato** **Susana Flecha** **7/4/25 10:22:00**

Color de fuente: Texto 1

▲ **Página 29: [102] Con formato** **Susana Flecha** **7/4/25 10:22:00**

Color de fuente: Texto 1

▲ **Página 29: [103] Con formato** **Susana Flecha** **7/4/25 10:22:00**

Color de fuente: Texto 1

▲ **Página 29: [103] Con formato** **Susana Flecha** **7/4/25 10:22:00**

Color de fuente: Texto 1

▲ **Página 29: [104] Con formato** **Susana Flecha** **7/4/25 10:22:00**

Color de fuente: Texto 1

▲ **Página 29: [104] Con formato** **Susana Flecha** **7/4/25 10:22:00**

Color de fuente: Texto 1

▲ **Página 29: [105] Con formato** **Susana Flecha** **7/4/25 10:22:00**

Color de fuente: Texto 1

▲ **Página 29: [105] Con formato** **Susana Flecha** **7/4/25 10:22:00**

Color de fuente: Texto 1

▲ **Página 29: [106] Con formato** **Susana Flecha** **7/4/25 10:22:00**

Color de fuente: Texto 1

▲
Página 29: [107] Con formato **Susana Flecha** **7/4/25 10:22:00**

Color de fuente: Texto 1

▲
Página 29: [107] Con formato **Susana Flecha** **7/4/25 10:22:00**

Color de fuente: Texto 1

▲
Página 29: [108] Con formato **Susana Flecha** **7/4/25 10:22:00**

Color de fuente: Texto 1

▲
Página 29: [108] Con formato **Susana Flecha** **7/4/25 10:22:00**

Color de fuente: Texto 1

▲
Página 29: [109] Con formato **Susana Flecha** **7/4/25 10:22:00**

Color de fuente: Texto 1

▲
Página 29: [109] Con formato **Susana Flecha** **7/4/25 10:22:00**

Color de fuente: Texto 1

▲
Página 29: [110] Con formato **Susana Flecha** **7/4/25 10:22:00**

Color de fuente: Texto 1

▲
Página 29: [110] Con formato **Susana Flecha** **7/4/25 10:22:00**

Color de fuente: Texto 1

▲
Página 29: [111] Con formato **Susana Flecha** **7/4/25 10:22:00**

Color de fuente: Texto 1

▲
Página 29: [111] Con formato **Susana Flecha** **7/4/25 10:22:00**

Color de fuente: Texto 1

▲
Página 29: [112] Con formato **Susana Flecha** **7/4/25 10:22:00**

Color de fuente: Texto 1

▲
Página 29: [112] Con formato **Susana Flecha** **7/4/25 10:22:00**

Color de fuente: Texto 1

▲
Página 29: [113] Con formato **Susana Flecha** **7/4/25 10:22:00**

Color de fuente: Texto 1

▲
Página 29: [113] Con formato **Susana Flecha** **7/4/25 10:22:00**

Color de fuente: Texto 1

▲
Página 29: [114] Con formato **Susana Flecha** **7/4/25 10:22:00**

Color de fuente: Texto 1

Página 29: [114] Con formato	Susana Flecha	7/4/25 10:22:00
-------------------------------------	----------------------	------------------------

Color de fuente: Texto 1

Página 29: [115] Con formato	Susana Flecha	7/4/25 10:22:00
-------------------------------------	----------------------	------------------------

Color de fuente: Texto 1

Página 29: [115] Con formato	Susana Flecha	7/4/25 10:22:00
-------------------------------------	----------------------	------------------------

Color de fuente: Texto 1

Página 29: [116] Con formato	Susana Flecha	7/4/25 10:22:00
-------------------------------------	----------------------	------------------------

Color de fuente: Texto 1

Página 29: [116] Con formato	Susana Flecha	7/4/25 10:22:00
-------------------------------------	----------------------	------------------------

Color de fuente: Texto 1

Página 29: [117] Con formato	Susana Flecha	7/4/25 10:22:00
-------------------------------------	----------------------	------------------------

Color de fuente: Texto 1

Página 29: [117] Con formato	Susana Flecha	7/4/25 10:22:00
-------------------------------------	----------------------	------------------------

Color de fuente: Texto 1

Página 29: [118] Con formato	Susana Flecha	7/4/25 10:22:00
-------------------------------------	----------------------	------------------------

Color de fuente: Texto 1

Página 29: [118] Con formato	Susana Flecha	7/4/25 10:22:00
-------------------------------------	----------------------	------------------------

Color de fuente: Texto 1

Página 29: [119] Con formato	Susana Flecha	7/4/25 10:22:00
-------------------------------------	----------------------	------------------------

Color de fuente: Texto 1

Página 29: [120] Con formato	Susana Flecha	7/4/25 10:22:00
-------------------------------------	----------------------	------------------------

Color de fuente: Texto 1

Página 29: [120] Con formato	Susana Flecha	7/4/25 10:22:00
-------------------------------------	----------------------	------------------------

Color de fuente: Texto 1

Página 29: [121] Con formato	Susana Flecha	7/4/25 10:22:00
-------------------------------------	----------------------	------------------------

Color de fuente: Texto 1

Página 29: [121] Con formato	Susana Flecha	7/4/25 10:22:00
-------------------------------------	----------------------	------------------------

Color de fuente: Texto 1

Página 29: [122] Con formato	Susana Flecha	7/4/25 10:22:00
-------------------------------------	----------------------	------------------------

Color de fuente: Texto 1

Página 29: [122] Con formato	Susana Flecha	7/4/25 10:22:00
-------------------------------------	----------------------	------------------------

Color de fuente: Texto 1

▲
Página 29: [123] Con formato **Susana Flecha** **7/4/25 10:22:00**

Color de fuente: Texto 1

▲
Página 29: [123] Con formato **Susana Flecha** **7/4/25 10:22:00**

Color de fuente: Texto 1

▲
Página 29: [124] Con formato **Susana Flecha** **7/4/25 10:22:00**

Color de fuente: Texto 1

▲
Página 29: [124] Con formato **Susana Flecha** **7/4/25 10:22:00**

Color de fuente: Texto 1

▲
Página 29: [125] Con formato **Susana Flecha** **7/4/25 10:22:00**

Color de fuente: Texto 1

▲
Página 29: [125] Con formato **Susana Flecha** **7/4/25 10:22:00**

Color de fuente: Texto 1

▲
Página 29: [126] Con formato **Susana Flecha** **7/4/25 10:22:00**

Color de fuente: Texto 1

▲
Página 29: [126] Con formato **Susana Flecha** **7/4/25 10:22:00**

Color de fuente: Texto 1

▲
Página 29: [127] Con formato **Susana Flecha** **7/4/25 10:22:00**

Color de fuente: Texto 1

▲
Página 29: [127] Con formato **Susana Flecha** **7/4/25 10:22:00**

Color de fuente: Texto 1

▲
Página 29: [128] Con formato **Susana Flecha** **7/4/25 10:22:00**

Color de fuente: Texto 1

▲
Página 29: [128] Con formato **Susana Flecha** **7/4/25 10:22:00**

Color de fuente: Texto 1

▲
Página 29: [129] Con formato **Susana Flecha** **7/4/25 10:22:00**

Color de fuente: Texto 1

▲
Página 29: [129] Con formato **Susana Flecha** **7/4/25 10:22:00**

Color de fuente: Texto 1

▲
Página 29: [130] Con formato **Susana Flecha** **7/4/25 10:22:00**

Color de fuente: Texto 1

▲
Página 29: [130] Con formato **Susana Flecha** **7/4/25 10:22:00**

Color de fuente: Texto 1

▲
Página 29: [131] Con formato **Susana Flecha** **7/4/25 10:22:00**

Color de fuente: Texto 1

▲
Página 29: [131] Con formato **Susana Flecha** **7/4/25 10:22:00**

Color de fuente: Texto 1

▲
Página 29: [132] Con formato **Susana Flecha** **7/4/25 10:22:00**

Color de fuente: Texto 1

▲
Página 29: [132] Con formato **Susana Flecha** **7/4/25 10:22:00**

Color de fuente: Texto 1

▲
Página 29: [133] Con formato **Susana Flecha** **7/4/25 10:22:00**

Color de fuente: Texto 1

▲
Página 29: [133] Con formato **Susana Flecha** **7/4/25 10:22:00**

Color de fuente: Texto 1

▲
Página 29: [134] Con formato **Susana Flecha** **7/4/25 10:22:00**

Color de fuente: Texto 1

▲
Página 29: [134] Con formato **Susana Flecha** **7/4/25 10:22:00**

Color de fuente: Texto 1

▲
Página 29: [135] Eliminado **Susana Flecha** **6/5/25 10:40:00**

▲
Página 29: [136] Con formato **Susana Flecha** **7/4/25 10:22:00**

Color de fuente: Texto 1