

Measurement report: Formation and brownness of aqueous secondary organic aerosol from the aged biomass-burning emissions in the Sichuan Basin, China

1 Chao Peng^{1,2,3,4}, Yan Ding⁴, Zhenliang Li^{1,2,3}, Tianyu Zhai⁴, Xinpeng
2 Yang⁴, Mi Tian⁶, Yang Chen⁵, Xin Long⁵, Haohui Tang⁶, Guangming
3 Shi⁷, Liuyi Zhang⁸, Kangyin Zhang⁸, Fumo Yang⁷, and Chongzhi Zhai^{1,2,3}

4 ¹Chongqing Academy of Ecology and Environmental Sciences, Chongqing, 401336,
5 China

6 ²Chongqing Branch Academy of Chinese Research Academy of Environmental
7 Sciences, Chongqing, 401336, China

8 ³Chongqing Key Laboratory of Urban Atmospheric Environment Observation and
9 Pollution Prevention, Chongqing, 401336, China

10 ⁴Chinese Research Academy of Environmental Sciences, Beijing 100012, China

11 ⁵Chongqing Institute of Green and Intelligent Technology, Chinese Academy of
12 Sciences, Chongqing, 400714, China

13 ⁶College of Environment and Ecology, Chongqing University, Chongqing, 400045,
14 China

15 ⁷College of Carbon Neutrality Future Technology, Sichuan University, Chengdu,
16 610065, China

17 ⁸Chongqing Three Gorges University, Wanzhou, 404000, China

18 **Correspondence:** Chao Peng (pengchao0623@sina.com) and Chongzhi Zhai
19 (czz66818@sina.com)

Abstract. Secondary organic aerosol (SOA) formed via complex chemical mechanisms was the major contributor to atmospheric aerosol pollution and climate forcing worldwide. The aqueous-phase oxidation was an important pathway for SOA formation and the aqueous SOA (aqSOA) exhibited absorption properties across ultraviolet to visible range. Here, we reported the formation and absorption properties of aqSOA in the Sichuan Basin, China. aqSOA was originated from the aged biomass-burning emissions via aqueous-phase reactions instead of photo-chemical reactions under high aerosol liquid water content (ALWC) conditions, especially during the polluted period. The substantial impact on brown carbon (BrC) absorption from SOA was observed from 370 nm to 660 nm (27.5%–43.2%). This study highlighted the significant contribution of aqSOA formation from aged biomass-burning emissions to the BrC budget and absorption, especially at night. The mean aerosol absorption Ångström exponents from 370 nm to 880 nm ($AAE_{370-880}$) was 1.95, higher than that observed in fresh and photo-chemically aged biomass-burning emissions. This study revealed the aqSOA formation and brownness from aged biomass-burning emissions and highlighted the importance of aqueous-phase reactions on aerosol pollution and absorption.

Keywords: Particulate matter; Secondary organic aerosol; Aqueous-phase oxidation; Aged biomass-burning emissions; Brown carbon.

1 Introduction

Organic aerosol (OA) is the dominant component (20 to 90%) of atmospheric aerosol with significantly implications for air quality and climate forcing (Jimenez et al., 2009). Numerous field observations indicated that secondary OA (SOA), formed by atmospheric oxidation of volatile organic compounds (VOCs) and primary OA (POA), accounted for most of OA worldwide (Ervens et al., 2011; Huang et al., 2014; Kourtchev et al., 2016). Recent results showed that aqueous-phase oxidation is an important pathway for SOA formation and these SOA production (aqSOA) exhibit absorption properties across ultraviolet (UV) to visible (Vis) range (Gilardoni et al., 2016; Lim et al., 2010; McNeill 2015; Powelson et al., 2014; Sun et al., 2010). However, the formation mechanisms and absorption properties of aqSOA are poorly understood, hindering to improvement of air quality and reducing the uncertainties in global climate estimations.

An increasing number of studies pointed toward aqSOA as a major SOA could form in fogs, clouds, and aerosol water (Ervens et al., 2011; Ortiz-Montalvo et al., 2012; Tan et al., 2012; Xu et al., 2022). And the oxygenated VOCs (OVOCs) with large water-soluble and low Henry's constant (i.e., methylglyoxal and glycolaldehyde) are the important aqSOA precursors (Ortiz-Montalvo et al., 2012; Tan et al., 2012). A few laboratory studies investigated the levoglucosan and phenolic species produced from biomass burning could also act as aqSOA precursors (Yu et al., 2016; Zhao et al., 2014). Gilardoni et al. (2016) reported direct ambient observations of aqSOA formation from biomass-burning emissions in fog water and wet aerosol. Additionally, recent studies indicated that aqSOA with high molecular weight (i.e., 4-ethylphenol) formed by aqueous-phase photochemical oxidation showed strong light absorptivity within UV range (Herrmann et al., 2015; Ye et al., 2019). Previous laboratory studies also demonstrated that aqSOA, such as π -conjugated compounds and imidazole with C=N bonds produced by aldol condensation and aqueous-phase carbonyl compound reactions respectively, would strongly absorb light at near-UV (Drozd and McNeill, 2014; Kampf et al., 2012; Nozière and Esteve, 2007; Powelson et al., 2014). Despite

numerous studies reported on the formation and optical properties of aqSOA, limited research on its ambient observations hinder to better understand the role of aqSOA in atmospheric chemistry and climate.

China experienced severe PM_{2.5} pollution under the stagnant high-humidity conditions, when SOA as the major component was originated from fossil fuel combustion and biomass burning (Huang et al., 2014; Wang et al., 2016; Wang et al., 2021; Xu et al., 2022). Field observations indicated that highly oxidized SOA could form through aqueous-phase processing driven by acid-catalyzed oxidation (Meng et al., 2020; Xu et al., 2017), and aqSOA is formed from biomass-burning OA (BBOA) and fossil-fuel OA via aqueous-phase reactions (Wang et al., 2021; Zhao et al., 2019). A few laboratory studies found aqueous-phase reactions were an important oxidation pathway for nitrophenol products (i.e., 5-nitrovanillin and 4-nitroguaiacol) with strong UV absorption and higher formation and transformation rates were observed in more acidic solutions (Kroflíc et al., 2015; Li et al., 2023; Pang et al., 2019; Yang et al., 2021). However, observations on aqSOA formation and optical properties in China are limited and most research concentrate on the North China Plain (NCP). Similar to NCP, the Sichuan Basin (SCB) characterized by high humidity and frequent biomass burning is also the main region with severe aerosol pollution in China (Tian et al., 2019; Wang et al., 2018; Yang et al., 2011). Previous research indicated that aqSOA from different regions exhibited distinct formation mechanisms and optical properties, due to the diverse sources and ambient conditions (Bao et al., 2023; Bao et al., 2024; Wang et al., 2021; Xu et al., 2017). For instance, Wang et al. (2021) revealed fast aqueous-phase conversion of fossil-fuel primary organic aerosol (FF-POA) to aqSOA under high-humidity conditions during a Beijing winter haze event, and found that aqSOA exhibited much lower light absorption than its primary precursor due to decreased aromaticity. Similarly, Huang et al. (2023) illustrated that the aqueous-phase oxidation of fossil fuel combustion emissions played a critical role in SOA formation under high RH conditions. Unlike these studies in NCP, the effect of aqueous-phase reactions on oxygenated OA (OOA) formation was significant when aerosol liquid water content (ALWC) was below 200 $\mu\text{g m}^{-3}$, but was insignificant

when ALWC > 200 $\mu\text{g m}^{-3}$ in SCB. Additionally, the aqueous-phase oxidation process probably did not play a role in the decay of BrC during summer in Chengdu (Bao et al., 2024). Currently, few studies explored the dynamic evolution and optical properties of aqSOA, and the knowledge of ambient aqSOA processing is still limited in SCB. Therefore, a more detailed characterization of aqSOA formation and optical properties is of great importance to reveal the key factors contributing to haze formation.

Here a time-of-flight aerosol chemical speciation monitor (ToF-ACSM) and a series of collocated instruments were used to characterize aqSOA dynamic evolution from biomass burning under real ambient conditions in a typical city with relatively serious air pollution in SCB from October 21 to November 23, 2022. We observed that the haze formation was largely driven by BBOA and aqSOA. We demonstrated aqSOA was originated from the aged BBOA via aqueous-phase reactions. Finally, we further showed that aqSOA produced from aged BBOA were strong UV absorption with positive radiative forcing. These results revealed the aqSOA formation and brownness from aged biomass-burning emissions and helped simulate the associated influences on atmospheric chemistry and climate.

2 Methods

2.1 Sampling site

An intensive field campaign on the chemical and physical properties of aerosol was conducted at a measurement site in a city affected by severe aerosol pollution (Yongchuan, 29°21'25" N, 105°54'6" E) from October 21 to November 23, 2022. This is a typical urban site surrounded by restaurants, shopping malls, and residential buildings, and the site is located in a parallel ridge-and-valley area between two megacities in SCB (Chongqing center and Chengdu) (Fig. S1). It was primarily influenced by multiple local emissions from traffic (arterial roads to the east 600 m and west 300 m) and a variety of residential sources (i.e., biomass burning and fossil fuel combustion). There was no interference of dynamics from neighboring buildings,

and measurements at the site helped understand the characteristics of haze pollution dynamic evolution.

2.2 Instrumentation

During the campaign, the non-refractory aerosol (NR-PM_{2.5}) species, including OA, ammonium (NH₄), nitrates (NO₃), sulfates (SO₄), and chlorides (Chl), were measured on-line by ToF-ACSM (Aerodyne Research Inc.). Ambient aerosols were pumped into ToF-ACSM at a flow rate of 3 L min⁻¹ through a PM_{2.5} cyclone (URG-2000-30ED) and a Nafion dryer (MD-110-48S, Perma Pure, Inc.) reducing the relative humidity to below 30%. The measurement principle was described in detail in the previous studies (Fröhlich et al., 2013; Ng et al., 2011c).

A seven-wavelength Aethalometer (AE33, Magee Scientific) was used to measure the aerosol light absorption (Abs_λ) and equivalent black carbon (BC_λ) mass concentrations in real time at 370, 470, 520, 590, 660, 880, and 950 nm. The sampled particles were dried by a Nafion dryer (MD-70024S-3, Perma Pure, Inc.) before entering into AE33. The light attenuation coefficients were converted to Abs_λ based on the real-time compensation parameter, and the nonlinear loading effects of quartz filters were dealt with on-line by the parallel measurements of attenuation values (ATN1 and ATN2) (Coen et al., 2010; Drinovec et al., 2015). The scattering effects of quartz filters were modified automatically by a fixed multiple scattering parameter (2.14). Detailed measurement methods and principles of AE33 can be found in Drinovec et al. (2015).

During the campaign, the gaseous species (including O₃, NO₂, and CO) were continuously measured by gas analyzers (49i, 42i, and 48i, Thermo Scientific), that were maintained and calibrated weekly. Hourly meteorological parameters data including temperature (T), relative humidity (RH) and PM_{2.5} mass concentrations were obtained on-line from the measurements at the National Environmental Monitoring Station, which was close to our sampling site (<http://www.cnemc.cn/>).

2.3 Data analysis

2.3.1 ToF-ACSM data analysis

The raw mass spectra data measured by ToF-ACSM were analyzed using Tofware v2.5.13 (Tofwerk AG) in Igor Pro 6.37 (WaveMetrics, Inc.). The ionization efficiency (IE) and relative ionization efficiency (RIEs) were regularly calibrated using a scanning mobility particle sizer with a differential mobility analyzer (SMPS 3081A, TSI) and a condensation particle counter (CPC 3775, TSI). The comprehensive overview of the operation and calibration procedures of ToF-ACSM can be found in Bao et al. (2023). In accordance with previous studies, the default RIEs values for OA, NO₃, and Chl were set to 1.4, 1.1, and 1.3, respectively (Canagaratna et al., 2007; Elser et al., 2016). The IE value (236 ions pg⁻¹) and RIEs of SO₄ (1.2) and NH₄ (4.3) were estimated from the calibrations of pure ammonium nitrate and ammonium sulfate, respectively. Meanwhile, a particle collection efficiency (CE) was introduced to compensate for the particle loss, as the acidity, the contribution of ammonium nitrate (ANMF) and phase state changed the particle bounce effects at the vaporiser (Matthew et al., 2008). Middlebrook et al. (2012) developed a CE algorithm for ToF-ACSM to quantify the aerosol species. Their results indicated that a constant CE value of 0.45 should be used when: (1) the ANMF is below 40%, or (2) particles are partially or fully neutralized. In this study, aerosol particles were dried by Nafion dryer (RH < 30%) before sampling by ToF-ACSM, and the ANMF was always below 40%. As shown in Fig. S2, the average ratio of the measured NH₄ to the predicted NH₄ needed to fully neutralize the SO₄, NO₃ and Chl was approximately 1. All of these conditions did not affect the CE value that had usually been used at this site. The typical default CE value (0.5) was applied during the whole sampling period, which was consistent with previous research (Bao et al., 2025; Peng et al., 2025; Sun et al., 2016a; Sun et al., 2016b; Zhao et al., 2019). While the typical default CE is 10% higher than 0.45, the difference is small considering the 30% uncertainty determined for CE (Bahreini et al., 2009). Additionally, the strong

correlation between NR-PM_{2.5} and PM_{2.5} mass concentrations supported that the CE value was reasonable (Fig. S3).

The mass spectral matrix of OA for m/z 10–120 was analyzed by positive matrix factorization (PMF) and multilinear engine (ME2) implemented with the SoFi (Source Finder) (version 6.3, Canonaco et al., 2013; Paatero 1999; Paatero and Tapper 1994). Briefly, unconstrained PMF was used to determine the numbers and types of source factors, then the restriction method ME2 was used to minimize PMF rotational ambiguity by the a -values from 0 to 1 with a step of 0.1 (Elser et al., 2016; Wang et al., 2019b; Zhong et al., 2021). The ions data with signal-to-noise (S/N) lower than 0.2 were discarded, and those S/N from 0.2–2 were downweighted by a factor of 2 (Bao et al., 2023; Paatero and Hopke 2003). Finally, five OA factors with function of the rotational parameter ($f_{\text{peak}} = 0$) were identified, including three POA factors (i.e., BBOA, coal-combustion OA (CCOA), and hydrocarbon-like OA (HOA)) and two SOA factors (i.e., OOA and aqSOA) (Fig. S9 and S10). We present detailed diagnostic plots of the PMF results in the supporting information (Fig. S4–S10). The details of OA source apportionment procedures are described in SI Text S1.

2.3.2 Aerosol liquid water content

The ALWC is controlled by meteorological conditions (T and RH) and also by inorganic and organic components. During the campaign, the ALWC with inorganic species was estimated by the ISORROPIA-II model based on the ammonium, nitrates, sulfates, and chlorides mass concentrations from ToF-ACSM and the meteorological parameters (T and RH) from National Environmental Monitoring Station (Fountoukis and Nenes, 2007). Here, the forward type and metastable mode were used in the ISORROPIA-II model (Hennigan et al., 2015). The thermodynamic equilibrium of the NH₄⁺–SO₄²⁻–NO₃⁻–Cl⁻–H₂O system was modeled and ALWC was then calculated. Consistent with previous research, the organic contribution for ALWC was calculated by Zdanovskii–Stokes–Robinson (ZSR) mixing rule as discussed in SI Text S2 (Guo et al., 2015; Huang et al., 2020; Nguyen et al., 2016; Xu et al., 2022). In this study, the

ALWC with organic species ranged from 0.1 to 35.2 $\mu\text{g m}^{-3}$, with an average of $1.9 \pm 3.0 \mu\text{g m}^{-3}$, taking up $3.7 \pm 2.2\%$ of total ALWC. As organic species had minor effects on total ALWC ($< 5\%$), the ALWC was determined only considering inorganic species (Chen et al., 2021; Guo et al., 2015; Liu et al., 2017).

2.3.3 Light absorption measurements

The Abs_λ was divided into BC and brown carbon (BrC, a group of colored OA compounds) absorption ($\text{Abs}_{\lambda,\text{BC}}$ and $\text{Abs}_{\lambda,\text{BrC}}$) ($\text{Abs}_\lambda = \text{Abs}_{\lambda,\text{BC}} + \text{Abs}_{\lambda,\text{BrC}}$) and characterized by the absorption Ångström exponents (AAE) (Laskin et al., 2015). Here, Abs_λ was determined dependent BC_λ mass concentrations ($\text{Abs}_\lambda = \text{BC}_\lambda \times \text{MAC}_\lambda$). We assumed the mass absorption cross-section of aerosols (MAC_λ) were 18.47, 14.54, 13.14, 11.58, 10.35, 7.77, and 7.19 $\text{m}^2 \text{g}^{-1}$ at 370, 470, 520, 590, 660, 880, and 950 nm, respectively (Drinovec et al., 2015; Zhu et al., 2017). Here, we assumed that Abs_{880} was sole from BC, then the following formula was used to determine $\text{Abs}_{\lambda,\text{BC}}$ values: $\text{Abs}_{\lambda,\text{BC}} = \text{Abs}_{880} \times (880/\lambda)^{-\text{AAE}_{\text{BC}}}$ (Drinovec et al., 2015; Kirchstetter and Novakov, 2004; Moosmüller et al., 2009; Qin et al., 2018; Zhu et al., 2017). The AAE of BC (AAE_{BC}) value was obtained from the equality: $\text{AAE}_{\text{BC}} = -\log_{10}(\text{Abs}_{880}/\text{Abs}_{950}) \div \log_{10}(880/950)$ (Wang et al., 2021). A detailed description of $\text{Abs}_{\lambda,\text{BC}}$ and $\text{Abs}_{\lambda,\text{BrC}}$ calculations is provided in SI Text S3. Previous research have demonstrated that AAE_{BC} is sensitive to the refractive index, size distribution, and coating of carbonaceous aerosols (Gyawali et al., 2009; Lack and Langridge, 2013; Li et al., 2019). In this study, the uncertainty associated with the estimation of Abs_{BC} and Abs_{BrC} was analyzed (Text S3). The relative uncertainty ranges of Abs_{BC} and Abs_{BrC} were $[-46\%, +21\%]$ and $[-112\%, +42\%]$ at 370 nm, respectively. $\text{Abs}_{\lambda,\text{BrC}}$ was caused by primary and secondary BrC light absorption ($\text{Abs}_{\lambda,\text{BrC,pri}}$ and $\text{Abs}_{\lambda,\text{BrC,sec}}$). The $\text{Abs}_{\lambda,\text{BrC,sec}}$ value was calculated by a minimum R-squared (MRS) method at each wavelength (Wang et al., 2019b; Wu and Yu, 2016; Wu et al., 2024). The detailed information of MRS method and $\text{Abs}_{\lambda,\text{BrC,sec}}$ estimation is provided in SI Text S3.

The multiple linear regression (MLR) method was used to analyze the light absorption of different OA components at each wavelength: $Abs_{BrC} = a \times [OOA] + b \times [BBOA] + c \times [CCOA] + d \times [aqSOA] + e \times [HOA]$ (Qin et al., 2018; Xie et al., 2019). The [OOA], [BBOA], [CCOA], [aqSOA], and [HOA] indicated the mass concentrations of OA species; the a–e were constants, used to optimize the Abs_{λ} of each OA component, and equivalent to MAC values at each wavelength (i.e., a–e at 370 nm represented $MAC_{370,OOA}$, $MAC_{370,BBOA}$, $MAC_{370,CCOA}$, $MAC_{370,aqSOA}$, and $MAC_{370,HOA}$, respectively). Here, the normalized mean bias (NMB), root mean square error (RMSE), and index of agreement (IOA) were used to evaluate the performance of the MLR method (SI Text S4) (Li et al., 2011). The IOA values of $Abs_{370,BrC}$ and $Abs_{470,BrC}$ (0.99 and 1.00) exceeded 0.95. The slopes of the relationship between $Abs_{370,BrC}$ and $Abs_{470,BrC}$ measured by AE33 and estimated by MLR method were 0.81 and 0.96 (close to unity), respectively. These results indicated a good agreement of $Abs_{370,BrC}$ between AE33 measurement and the MLR reconstruction.

3 Results and discussion

3.1 General descriptions

The temporal variations of $PM_{2.5}$ species concentrations, meteorological parameters, $Abs_{370,BrC}$ and $MAC_{370,BrC}$ during the campaign are shown in Fig. 1. The winds were weak with $0.3 \pm 0.2 \text{ m s}^{-1}$ over the whole campaign, indicating the atmosphere was in stagnant conditions. The total $PM_{2.5}$ (BC+NR- $PM_{2.5}$) mass concentrations ranged from 7.0 to 175.5 $\mu\text{g m}^{-3}$, with an average of $48.4 \pm 27.8 \mu\text{g m}^{-3}$ during the campaign. The average concentrations of OA, NO_3 , SO_4 , NH_4 , Chl, and BC were 24.1 ± 18.1 , 8.3 ± 6.2 , 6.2 ± 3.4 , 5.2 ± 2.7 , 0.2 ± 0.1 , and $4.7 \pm 2.9 \mu\text{g m}^{-3}$, taking up $46.6 \pm 10.7\%$, $17.7 \pm 8.0\%$, $13.2 \pm 4.4\%$, $11.2 \pm 2.7\%$, $0.3 \pm 0.2\%$, and $10.1 \pm 5.5\%$ of total $PM_{2.5}$, respectively. OA constituted the largest fraction of total $PM_{2.5}$, highlighting the importance of OA in $PM_{2.5}$ pollution in SCB (Bao et al., 2023; Wang et al., 2018). Meanwhile, the high values of $Abs_{370,BrC}$ and $MAC_{370,BrC}$, ranging from 5.8 to 210.2 Mm^{-1} ($42.4 \pm 28.5 Mm^{-1}$) and from 0.6 to 7.0 $m^2 g^{-1}$ ($2.1 \pm 0.9 m^2$

g⁻¹) respectively, were observed during the campaign.

According to the Chinese National Ambient Air Quality Standard (NAAQS) (GB 3095-2012) (MEP, 2012), the Grade I and Grade II levels for daily PM_{2.5} mass concentration are 35 µg m⁻³ and 75 µg m⁻³, respectively. The Chinese NAAQS Grade II level, based on WHO's Phase-1 interim target (IT-1), is higher than the WHO Air Quality Guideline (AQG) value (15 µg m⁻³), the EU daily limit (25 µg m⁻³), and U.S. 24-hour standard (35 µg m⁻³). During the campaign, the average of PM_{2.5} mass concentration was 1.4 times NAAQS Grade I level (35 µg m⁻³). Therefore, the pollution periods (PP) were defined by the daily PM_{2.5} mass concentration exceeding NAAQS Grade II level of 75 µg m⁻³. Similarly, the days with PM_{2.5} mass concentration below 75 µg m⁻³ were characterized as clean periods (CP). During PP, the mass concentrations of BC+NR-PM_{2.5} and OA were 102.3 ± 26.9 and 57.4 ± 22.5 µg m⁻³, 2.5 and 3.1 times that during CP, respectively. As shown in Fig. 2, the PM_{2.5} species were substantially different in the PP and CP. Compared with other species, a significantly higher contribution of OA was observed during PP (56.6%) than CP (46.6%) (Student's t-test, *p* < 0.001) (Fig. 2). Here, five OA factors were identified by the PMF model with detailed information in SI Text S1, and the mass spectrum of these factors is shown in Fig. S9. HOA mass spectrum was characterized by alkyl fragment ion series at C_nH⁺_{2n-1} and C_nH⁺_{2n+1} (i.e., *m/z* 41, 43, 55, and 57), they are common characteristics of primary combustion emissions (Elser et al., 2016; Lanz et al., 2007). BBOA was identified by the high signal of *m/z* 60 (mainly C₂H₄O₂⁺) and *m/z* 73 (mainly C₃H₅O₂⁺), they are the fragments of levoglucosan and mannosan emitted from incomplete biomass burning (Alfarra et al., 2007). CCOA had high correlation with the unsaturated hydrocarbon ion fragments such as PAH-related ion fragments (i.e., *m/z* 77, 91, 115), emitted from traditional coal combustion (Sun et al., 2016a). OOA was distinguished by the prominent signal of *m/z* 44 (mainly CO₂⁺) and highly correlated with the oxygenated ions (Ng et al., 2011b). aqSOA also had high correlation with the oxygenated ions (i.e., *m/z* 43 (mainly C₂H₃O⁺) and *m/z* 44). The mass spectrum of aqSOA showed a significantly higher *m/z* 29 (mainly CHO⁺) signal than other OA factors, consistent with those reported in the previous studies

(Sun et al., 2016a; Xu et al., 2019; Zhao et al., 2019; Zhong et al., 2021). Moreover, BBOA showed significant correlations with m/z 60 (mainly C₂H₄O₂⁺) and m/z 73 (Pearson's r^2 (r^2) = 0.85, 0.80, $p < 0.001$); CCOA was strongly correlated with Chl and m/z 115 (r^2 = 0.56, 0.48, $p < 0.001$); HOA was correlated with NO₂ and m/z 41 (r^2 = 0.47, 0.59, $p < 0.001$); OOA and aqSOA were significantly correlated with NO₃, NH₄ (r^2 = 0.77, 0.75, $p < 0.001$) and SO₄, ALWC (r^2 = 0.67, 0.85, $p < 0.001$), respectively (Fig. S10). These results highlighted the result of five OA factors was reasonable.

It should be noted that the contributions of BBOA and aqSOA to OA increased from CP (31.7% and 12.6%) to PP (38.6% and 14.1%), while CCOA, HOA, and OOA contributions decreased. Additionally, significantly higher RH and ALWC were observed during PP ($58.5 \pm 12.4\%$ and $69.4 \pm 30.3 \mu\text{g m}^{-3}$) than CP ($49.8 \pm 8.9\%$ and $37.1 \pm 20.8 \mu\text{g m}^{-3}$) ($p < 0.001$), but not temperature ($p > 0.1$). The wind was $0.32 \pm 0.18 \text{ m s}^{-1}$ during CP, 1.3 times that during PP. These results indicated that the atmosphere was in a stagnant state with relatively high RH and ALWC during PP, which might lead to the largely different sources and chemical processing of OA during CP and PP. Compared with CP, the obvious diurnal variation of OA concentration was exhibited during PP. As shown in Fig. 2, the OA concentration peak ($82.7 \mu\text{g m}^{-3}$) was observed at 12:00 local time (LT) in the daytime during PP, while observed at 21:00 LT at night during CP. Moreover, OA concentration rapidly increased at a rate of $7.8 \mu\text{g m}^{-3} \text{ hr}^{-1}$ from 09:00 to 12:00 LT with a significant decrease of NO₃ during PP. Meanwhile, BBOA and aqSOA concentrations showed similar diurnal patterns to OA concentration with high values in the daytime and rapidly increased from 09:00 to 12:00 LT during PP. Previous research indicated that aqSOA spectrum showed higher m/z 29 (CHO⁺) than other OA factors (Gilardoni et al., 2016; Meng et al., 2020; Wang et al., 2021). During PP, the peaks of m/z 60 and m/z 29 concentrations, tracer ion fragments of BBOA and aqSOA, were observed at 12:00 LT ($1.2 \mu\text{g m}^{-3}$) and 13:00 LT ($4.3 \mu\text{g m}^{-3}$), respectively. Additionally, the correlation between ALWC and aqSOA concentrations (r^2 = 0.86, $p < 0.001$) was stronger than BBOA concentrations (r^2 = 0.58, $p < 0.001$), and both ALWC and

aqSOA concentration peaks were observed at 13:00 LT, earlier than BBOA concentration peak (12:00 LT), supporting that ALWC might play a significant role in the chemical processing of aqSOA formation from BBOA during PP. In contrast, the odd oxygen ($O_x = O_3 + NO_2$) showed weak correlations with both OOA and aqSOA concentrations during the campaign ($p > 0.1$) (not shown). Although the average O_x concentration was higher during PP (51.1 ± 19.6 ppb) than CP (36.9 ± 14.0 ppb), no significant correlations were observed in either period (not shown). These results suggested that photochemical reactions might played a limited role in SOA formation in this study.

In summary, these results suggested that OA was the dominant component of $PM_{2.5}$, especially during PP in SCB. During PP, BBOA and aqSOA played important roles in increasing OA concentration in the daytime. Previous research indicated that the autumn harvest period – specifically October and November – is a period of intensified biomass burning in SCB, primarily due to post-harvest crop residue burning (Chen et al., 2017; Tao et al., 2014). Thus, aqSOA in the daytime during PP might be related to the high aerosol water and BBOA emissions (Bao et al., 2023; Chen et al., 2017; Chen et al., 2019; Tao et al., 2014). Additionally, Gilardoni et al. (2016) also found that aqSOA such as guaiacol dimer ($C_{14}H_{14}O_4^+$) could be formed from aged biomass-burning emissions at both in fog water and in wet aerosol, especially under high ALWC conditions. To further explore aqSOA formed from biomass-burning emissions via the aqueous-phase reactions, the next section would discuss the dynamic evolution of aqSOA in relation to BBOA.




Figure 1. Time series of (a) RH and T, (b) NR-PM_{2.5} species measured by ToF-ACSM and BC, (c) Abs_{370,BrC} and MAC_{370,BrC}, and (d) mass fraction of OA factors during the campaign. The pollution period (BC+NR-PM_{2.5} > 75 µg m⁻³) is highlighted by the shaded areas.




Figure 2. Diurnal variations of PM_{2.5} species, BC, OA factors, m/z 29, m/z 60, and m/z 115 mass

concentrations during **(a)** clean period (CP) ($\text{BC+NR-PM}_{2.5} < 75 \mu\text{g m}^{-3}$) and **(b)** polluted period (PP) ($\text{BC+NR-PM}_{2.5} > 75 \mu\text{g m}^{-3}$). The pie charts in the left side of **(a)** and **(b)** show the average mass contributions of different chemical compositions to $\text{BC+NR-PM}_{2.5}$ during CP and PP, respectively. Meanwhile, the average mass contributions of OOA, BBOA, CCOA, aqSOA, and HOA in OA are shown in the pie charts in the middle of **(a)** and **(b)**, respectively.

3.2 Biomass-burning emissions as precursors for aqSOA

Fig. 3 shows the strong correlation between the mass fraction (%) of aqSOA in total $\text{PM}_{2.5}$ and ALWC during the campaign ($r^2 = 0.64$, $p < 0.001$). The contribution of aqSOA increased with the increase of f_{29} values (normalized mass spectrum signal at m/z 29). It was important to note that the aqSOA factor showed significantly higher f_{29} and f_{60} values (normalized mass spectrum signal at m/z 60) (0.167 and 0.011) than the OOA factor (0.017 and 0.002), respectively (Fig. S9). Moreover, both aqSOA concentrations and f_{29} were well correlated with ALWC ($r^2 = 0.85$, 0.73 , $p < 0.001$) (Fig. 3). During the campaign, the average value of the oxygen-to-carbon ratio (O:C) of aqSOA factor (0.85) was 2.7 times that (0.31) of BBOA factor. However, the similar hydrogen-to-carbon ratio (H:C) values of aqSOA factor and BBOA factor were observed (1.74 and 1.81, respectively), indicating that a hydrogen atom might be replaced by a OH moiety (Lim et al., 2010; Ng et al., 2011a). These results were similar to aqSOA observed in Italy and Beijing (Gilardoni et al., 2016; Zhao et al., 2019). Additionally, previous studies indicated that aqueous-phase processes could play a role in the formation of SOA from fossil fuel emissions (Ervens et al., 2011; Huang et al., 2023; Wang et al., 2021; Xu et al., 2022; Yan et al., 2017). For example, Wang et al. (2021) and Xu et al. (2022) have highlighted the potential for aqueous-phase reactions to contribute to SOA formation, particularly in regions with high levels of anthropogenic emissions. In this study, a strong anticorrelation between the mass fraction of fossil-fuel related OA components (sum of CCOA, HOA and OOA) and ALWC at the high f_{29} values was also observed ($r^2 = 0.48$, $p < 0.001$) (not shown), consistent with recent research (Wang et al., 2021). This indicated that

aqSOA might also be produced by aqueous-phase reactions of fossil-fuel related OA components.

Fig. 4 shows the relationships between ALWC and OA factors or f_{29} during the campaign. Five OA factors mass concentrations increased with the increase of ALWC. However, compared with other OA factors, aqSOA and BBOA significantly increased from 1.1 and 4.9 $\mu\text{g m}^{-3}$ to 5.2 and 10.8 $\mu\text{g m}^{-3}$ when $20 \mu\text{g m}^{-3} < \text{ALWC} < 60 \mu\text{g m}^{-3}$, respectively. It should be noted that only aqSOA concentrations were enhancement under high ALWC conditions ($> 100 \mu\text{g m}^{-3}$). It is likely because more water-soluble organic species (i.e., glyoxal and methylglyoxal) were formed, that were further oxidized to form aqSOA via aqueous-phase reactions in aerosol liquid water (Carlton et al., 2007; Ervens et al., 2011; Tan et al., 2012). As shown in Fig. 4b, the mass fraction of aqSOA showed significant enhancement from less than 5% at $\text{ALWC} < 20 \mu\text{g m}^{-3}$ to 17–22% at $\text{ALWC} > 60 \mu\text{g m}^{-3}$ with a corresponding decrease in OOA, although POA (BBOA+CCOA+HOA) and SOA (OOA+aqSOA) contributions were fairly constant across different ALWC levels (58–68% and 32–42%). This result suggested a more intensive formation of aqSOA than OOA via aqueous-phase reactions, although aqSOA might be also formed from OOA, consistent with the recent research in northwest China (Zhao et al., 2019; Zhong et al., 2021). Additionally, the increasing f_{29} (CHO^+) from 0.010 to 0.227 as a function of ALWC was observed during the campaign (Fig. 4b). The values of f_{29} significantly increased from 0.055 to 0.210 when ALWC increased from $60 \mu\text{g m}^{-3}$ to $100 \mu\text{g m}^{-3}$ ($p < 0.001$), consistent with OA mass concentrations (13.2–109.1 $\mu\text{g m}^{-3}$) during the campaign (Fig. 4b). According to the laboratory analysis of organic standards, previous research found that the spectra of standard organic species without alcohol group showed low f_{29} (< 0.05), while high f_{29} values (0.05–0.15) were found for polyols and species with non-acid OH groups produced from biomass-burning emissions (Canagaratna et al., 2015; Gilardoni et al., 2016; Zhao et al., 2014). This further highlighted the potential formation of organic compounds with hydroxyl groups (i.e., glyoxal and methylglyoxal) under high ALWC conditions. Overall, these results pointed to the fact that the observed aqSOA could be formed from

biomass-burning emissions via aqueous-phase reactions, reinforcing the BBOA role in increasing PM_{2.5} mass concentration.




Figure 3. Scatter plot of the mass fraction of aqSOA in BC+NR-PM_{2.5} versus ALWC colored by the f_{29} (normalized mass spectrum signal at m/z 29) during the campaign. f_{29} (mainly CHO⁺) is a tracer for alcohol compounds and used to monitor the aqueous-phase oxidation of organic compounds (i.e., glyoxal).




Figure 4. Variations of (a) OA factors mass concentrations, and (b) OA mass concentrations, f_{29} (a tracer for alcohol compounds), and mass fraction of OA factors as a function of ALWC. The data

were grouped into different bins according to a $10 \mu\text{g m}^{-3}$ increment of ALWC.

To identify the formation of aqSOA and its precursors under different $\text{PM}_{2.5}$ pollution levels, the relationship between aqSOA and BBOA or OOA mass concentrations with ion fragments tracers during CP and PP was performed, respectively. The correlation r^2 between aqSOA and BBOA concentrations was higher during PP (0.64) than that during CP (0.54) ($p < 0.001$) (Fig. 5a and c). Though aqSOA and BBOA concentrations increased with the increase of ALWC during CP and PP, the correlations between ALWC and aqSOA or BBOA concentrations were relatively stronger during PP than that during CP ($p < 0.001$). Fig. 5b and d show that f_{29} was highly correlated with aqSOA formation during CP and PP. A few data points with high aqSOA and OOA concentrations had low f_{29} values (0.071–0.102) in Fig. 5d, while the average value of f_{44} (normalized mass spectrum signal at m/z 44) of these data points (0.103 ± 0.024) was 1.3 times that of all data points (0.080 ± 0.035) during PP. It was likely due to the formation of more-oxidized OOA under high ALWC values ($> 80 \mu\text{g m}^{-3}$) in this study (Xu et al., 2017). Previous research found that the f_{29} values of polyols and species with non-acid OH groups from biomass-burning emissions were lower than 0.15 (Canagaratna et al., 2015; Gilardoni et al., 2016; Zhao et al., 2014). Moreover, the mass fraction of aqSOA showed a stable increasing trend and remained high levels (from 18% to 22%) at $\text{ALWC} > 80 \mu\text{g m}^{-3}$, which was associated with a corresponding decrease in OOA (from 15% to 10%) (Fig. 4b). Compared with OOA ($p > 0.1$), the aqSOA concentrations showed strong positive correlation with ALWC ($r^2 = 0.73$, $p < 0.001$) when $\text{ALWC} > 80 \mu\text{g m}^{-3}$ during PP. In contrast, ALWC showed the weak correlations with aqSOA and OOA concentrations during CP ($p > 0.1$). It should be noted that a strong anticorrelation between aqSOA and OOA concentrations was observed during PP at $\text{ALWC} > 80 \mu\text{g m}^{-3}$ when $f_{29} > 0.15$ ($r^2 = 0.76$, $p < 0.001$), but not during CP ($p > 0.1$) (Fig. 5b and d). These results indicated that the aqSOA formation was more intensive than OOA at high ALWC levels during PP.

Previous research demonstrated that f_{44} (representation of aged OA) could be used as a tracer of aged SOA, f_{43} (normalized mass spectrum signal at m/z 43) as a

451 tracer of POA and fresh SOA, and f_{60} (presence of anhydrosugars) as a tracer of
 452 BBOA (Cubison et al., 2011; Ng et al., 2010). Additionally, m/z's 44 and 43 are
 453 usually from different functional groups and the ratio changes as a function of
 454 atmospheric aging. The triangle plot of f_{44} versus f_{43} has been widely used to
 455 characterize OA evolution, and f_{44} versus f_{60} is commonly used to investigate the
 456 aging trend of BBOA (Ortega et al., 2013; Paglione et al., 2020; Xu et al., 2017; Xu et
 457 al., 2019). As shown in Fig. 6, the bottom region of the triangle was dominated by
 458 BBOA, CCOA, and HOA with low f_{44} (0.040, 0.017, and 0.016, respectively) in this
 459 study, indicating that they were freshly emitted and less oxidized. However, the f_{44} of
 460 SOA factors (i.e., OOA and aqSOA) (0.118 and 0.117) were observably higher than
 461 POA factors, showing the freshly oxidized properties of SOA. Meanwhile, f_{44} of
 462 aqSOA was close to that observed in fogs (Gilardoni et al., 2016; Kim et al., 2019),
 463 highlighting the presence of aqueous-phase reactions in this study. The relative
 464 abundance of m/z 45 (mainly HCO_2^+), a tracer ion for carboxylic acids, was higher in
 465 the aqSOA spectra than in the OOA spectrum (Fig. S9). It was consistent with previous
 466 research which found that aqueous-phase reactions were important sources of
 467 oxygenated organic compounds, including organic acids (Ervens et al., 2011; Kim et
 468 al., 2019; McNeill, 2015; Sun et al., 2010; Yu et al., 2014). Fig. 6b shows BBOA and
 469 aqSOA with higher f_{60} values (0.019 and 0.011) than CCOA (0.009) and HOA (0.008).
 470 The f_{60} value of OOA was 0.002, lower than the typical background value (0.003) in
 471 the atmosphere without biomass burning influence (Cubison et al., 2011). The mass
 472 spectrometry feature of aqSOA showed large f_{44} and f_{60} values, laying in a schematic
 473 space of aged BBOA based on mass spectrometry features in previous research
 474 (Cubison et al., 2011; Ortega et al., 2013). Additionally, BBOA contains abundant
 475 water-soluble organic compounds (WSOC) with (i.e., sugars, phenols, and organic
 476 acids), that can form aqSOA via efficient aqueous-phase reactions (i.e., oxidation and
 477 oligomerization reactions) (Ervens et al., 2011; Gilardoni et al., 2016; Lee et al., 2013;
 478 Lei et al., 2024; Li et al., 2020; Powelson et al., 2014). In contrast, OOA formation
 479 primarily relies on gas-phase oxidation of VOCs with high-reactivity (i.e., aromatics
 480 and long-chain alkanes) (i.e., OH radicals), which has low concentrations in BBOA

(Akagi et al., 2011; Jimenez et al., 2009; Shrivastava et al., 2017; Yokelson et al., 2007). This suggested that BBOA could be the important precursors for aqSOA instead of OOA via aqueous-phase reactions. These results were consistent with previous research and most of the observation data were within the triangle space (Bao et al., 2023; Kim et al., 2019; Paglione et al., 2020). It should be noted that these processes of aqSOA formation could be more intense and important during autumn due to elevated precursor concentrations (i.e., BBOA), ALWC, and RH values, though they are chemically robust and likely occur year-round.

During PP, the f_{44} values ranging from 0.022 to 0.140 (0.080 ± 0.035) were significantly higher than that during CP (0.021–0.150, 0.064 ± 0.019) ($p < 0.001$), while the f_{43} value was slightly lower with an average of 0.062 ± 0.027 . Compared with CP ($r^2 = 0.17$, slope = -0.53), f_{44} showed a more significant increase as the decreasing of f_{43} with higher r^2 value (0.70) and the regression slope of f_{44} versus f_{43} (-1.09) was closer to -1 during PP. This indicated that more aged SOA existed in the atmosphere during PP (Fig. 6a and c). It should be noted that the points of f_{44} versus f_{43} were inside the upper boundary of the triangle region, and most points were outside the bottom boundary of the triangle region during PP. These results suggested that less oxidized SOA were formed via aqueous-phase reactions instead of photo-chemical reactions during PP (Kim et al., 2019; Zhao et al., 2019). Moreover, these points outside the bottom boundary of the triangle region with higher f_{44} (> 0.05) and lower f_{43} (< 0.06) showed relatively higher ALWC during PP, but not during CP.

Here, the triangle plots of f_{44} versus f_{60} colored by ALWC under different PM_{2.5} pollution levels were analyzed (Fig. 6b and d), when the link between aqSOA and BBOA was further stressed by a schematic representation of aged BBOA. Except for several points, the f_{60} values were ubiquitously higher than 0.003, and most points fell in the triangular region, suggesting the contribution of biomass burning to OA. During PP, the f_{60} values ranging from 0.005 to 0.019 (0.010 ± 0.004) were similar with CP (from 0.004 to 0.019, 0.010 ± 0.003). The correlation r^2 between f_{44} and f_{60} was higher during PP (0.72) than that during CP (0.31) ($p < 0.001$). Moreover, compared with all data points during PP, those in the schematic space of aged BBOA showed relatively

higher ALWC, a pattern that differed from observations during CP.

Regional transport significantly influences the aging of BBOA in the SCB. Previous studies have demonstrated that the northeast winds prevail during autumn in the SCB and facilitate the transport of pollutants along the Dazhou → Guang'an → Hechuan pathway, and this northeast-southwest transport pathway had a significant impact on Chongqing (Peng et al., 2019; Wang et al., 2018). As shown in Fig. S11, air masses predominantly originated from northeastern Chongqing, which is an area with widespread agricultural burning activities (He et al., 2015; Luo et al., 2020), and were transported over short ranges during the campaign. Compared to other air mass clusters, the highest contribution and concentration of BBOA in total PM_{2.5} were observed when air masses passed through northeastern Chongqing. The percentage of air mass trajectories that passed through biomass-burning-influenced regions was higher during PP (~57%) than CP (~35%). Unlike cluster 3 during CP, air masses originating from northeastern Chongqing (cluster 2) during PP showed significantly higher contributions and concentrations of BBOA (24.7%, 27.8 µg m⁻³) and aqSOA (9.4%, 10.6 µg m⁻³) than other clusters ($p < 0.001$). In addition, cluster 2 exhibited notably higher values of ALWC (93.7 µg m⁻³), Abs_{370,BrC,sec} (105.1 Mm⁻¹), f_{44} (0.113), and the NO₃/SO₄ ratio (2.1, a tracer for BBOA aging (Liu et al., 2024; Zhang et al., 2025)) than other clusters during PP ($p < 0.001$). During PP, approximately 68% of the trajectories in cluster 2 passed through biomass-burning-influenced regions, with transport times to the sampling site ranging between 12 and 48 hours. This time frame is consistent with previous research on the aqueous-phase aging process of biomass burning emissions (Cubison et al., 2011; Hennigan et al., 2010; Ortega et al., 2013; Zhu et al., 2023). These results suggested that regional transport of BBOA primarily originated from northeastern Chongqing, and aged to aqSOA within approximately 12 to 48 hours.




Figure 5. Scatter plots of aqSOA versus (a, b) BBOA and (c, d) OOA mass concentrations colored by ALWC during clean period and polluted period. The size of the symbols in (b) and (d) increases with the increase of the f_{29} value, which is a tracer for alcohol compounds.




Figure 6. Triangle plots of (a, c) f_{44} (normalized mass spectrum signal at m/z 44) versus f_{43} (normalized mass spectrum signal at m/z 43), and (b, d) f_{44} versus f_{60} (normalized mass spectrum signal at m/z 60) colored by ALWC (circles) during clean period and polluted period. The dashed lines in (a) and (c) were derived from Ng et al. (2010) and used to follow the aging of OA components in the atmosphere. The background space ($f_{60} < 0.003$) without biomass burning influence was also shown by the grey shaded area. The background value of secondary aged OA (brown dashed line) and the black dashed lines characterising the aging of BBOA in (b) and (d) were derived from Cubison et al. (2011). The data points (squares) included the measurements in this study (bordered in red) and previous research (Bao et al., 2023; Gilardoni et al., 2016; Kim et al., 2019; Ng et al., 2011a; Paglione et al., 2020; Xu et al., 2015; Xu et al., 2017; Xu et al., 2019; Zhao et al., 2017; Zhao et al., 2019). f_{43} (mainly $\text{C}_2\text{H}_3\text{O}^+$) is a tracer for POA and fresh SOA. f_{44} is a proxy of the OA oxygenation degree and used as a tracer for aged SOA. f_{60} is a proxy of anhydrosugars emitted from biomass burning.

3.3 Evolution of BrC Absorption

Previous research indicated that OA from fresh and aged biomass-burning emissions, which exhibited absorption properties across UV to Vis range with significantly higher AAE value than BC, might contribute to a net positive radiative forcing (Laskin et al., 2015). Therefore, the BrC absorption properties and their relationship with five OA factors were analyzed in this study. The values of $Abs_{\lambda, BrC, pri}$ and $Abs_{\lambda, BrC, sec}$ were obtained by MRS method, and the MLR method was used to estimate Abs of five OA factors at each wavelength (SI Text S3 and S4). The average value of $Abs_{370, BrC}$ was $42.4 \pm 28.5 \text{ Mm}^{-1}$ (accounting for 49.2% of Abs_{370}), much higher than $Abs_{660, BrC}$ ($2.6 \pm 1.3 \text{ Mm}^{-1}$, 10.5%), suggesting a high absorption efficiency for BrC in the near-UV wavelength. The $Abs_{\lambda, BrC, pri}$ and $Abs_{\lambda, BrC, sec}$ accounted for 56.8%–72.5% and 27.5%–43.2% of $Abs_{\lambda, BrC}$ from 370 nm to 660 nm respectively, indicating primary emissions were the main contributors to BrC absorption (Fig. S13). However, the contribution of $Abs_{\lambda, BrC, sec}$ to $Abs_{\lambda, BrC}$ increased with wavelength, suggesting the impact on Abs_{BrC} from SOA should not be ignored. Hereafter, we show that aqSOA formation from aged BBOA contributed to the BrC budget and was strong absorption across UV to Vis range.

The data at 370 nm with higher signal-to-noise ratios and Abs_{BrC} contribution was chosen to further analyze the correlations of BrC absorption with various OA components. As described in section 2.3.3, the Abs of five OA factors at each wavelength were obtained by the MLR method (Table S1). Compared with CCOA ($Abs_{370, CCOA}$), HOA ($Abs_{370, HOA}$), and OOA ($Abs_{370, OOA}$) (11.5%, 9.1%, and 11.1%), the Abs at 370 nm calculated for BBOA ($Abs_{370, BBOA}$) and aqSOA ($Abs_{370, aqSOA}$) showed higher contributions (51.9% and 16.4%) to $Abs_{370, BrC}$, consistent with the higher MAC values (Fig. S15). Fig. S16 presents the correlations between $Abs_{370, BrC}$ and the mass concentrations of OOA, BBOA, CCOA, aqSOA, HOA, and m/z 60. $Abs_{370, BrC}$ showed the strongest positive correlations with BBOA and m/z 60 (ion fragments tracers of BBOA) concentrations ($r^2 = 0.77$, $p < 0.001$), followed by aqSOA concentrations ($r^2 = 0.69$, $p < 0.001$). In contrast, the correlations with HOA

concentrations ($r^2 = 0.36$), CCOA ($r^2 = 0.25$), and OOA ($r^2 = 0.09$, $p > 0.1$) were much weaker. Compared with other OA factors (except BBOA), the contribution of Abs_{370,aqSOA} to Abs_{370,BrC} was relatively higher (Table S1), when the correlation between Abs_{370,BrC} and aqSOA concentrations was also stronger. These results might be related to the aqSOA formed from the aged BBOA via aqueous-phase reactions. Gilardoni et al. (2016) demonstrated that aqSOA formation from aged BBOA via aqueous-phase reactions in the ambient atmosphere contributed to the BrC budget and exhibited slightly higher AAE_{467–660} (AAE of aerosols from 467 nm to 660 nm) values than the fresh and processed biomass-burning emissions in laboratory experiments. The MAC values of the five resolved OA components equivalent to the a–e values in the MLR method at different wavelengths were shown in Fig. S15. Among these, BBOA showed the highest MAC value ($2.37 \text{ m}^2 \text{ g}^{-1}$), followed by aqSOA ($1.23 \text{ m}^2 \text{ g}^{-1}$) at 370 nm, indicating that the oxidation of BBOA to aqSOA decreased light absorption at short wavelengths. Previous research found that the MAC of BBOA at 365 nm was twice that of SOA, which was associated with the water-soluble BrC (Lorenzo et al., 2017; Washenfelder et al., 2015). The AAE values of OA factors, calculated by a power-law fitting of Abs for these OA factors from 370 nm to 660 nm (Qin et al., 2018; Wang et al., 2019b), were shown in Fig. S15. It should be noted that aqSOA had the lowest AAE_{370–660,aqSOA} value (3.54), while BBOA has the highest AAE_{370–660,BBOA} value (4.93). Moreover, the contribution of aqSOA to Abs_{BrC} increased from 16.4% to 26.7% from 370 to 660 nm, while the contribution from BBOA decreased from 51.9% to 39.1% from 370 to 660 nm. These suggested aqSOA formation from aged BBOA might play an important role in the light absorption of BrC across UV to Vis range.

Fig. 7 shows the ternary contour map to quantify the contribution of BBOA, CCOA, and HOA factors to Abs_{370,BrC,pri}, when the strong positive correlation ($p < 0.001$) and high slope of the linear regression (1.80) between BBOA mass concentrations and Abs_{370,BrC,pri} were observed. Among these POA factors, the high mass fractions of BBOA to POA were consistent with the high Abs_{370,BrC,pri} values (Fig. 7a). For example, the most data of Abs_{370,BrC,pri} higher than 49.1 Mm^{-1} (90th

percentile of $\text{Abs}_{370,\text{BrC}}$ fell in the region of high BBOA/POA values (> 0.5). Moreover, $\text{Abs}_{370,\text{BrC,pri}}$ significantly increased with the increases of BBOA and m/z 60 mass concentrations with higher r^2 values (0.63 and 0.55) than HOA and CCOA (0.19 and 0.14) (Fig. 7b). These results indicated the major contribution of BBOA to primary BrC light absorption.

During the campaign, the relationship between $\text{Abs}_{370,\text{BrC,sec}}$ and SOA factors mass concentrations was analyzed to understand the correlation between secondary BrC absorption and its chromophores. As shown in Fig. S17, $\text{Abs}_{370,\text{BrC,sec}}$ significantly increased with the increase of aqSOA concentrations ($r^2 = 0.44$, $p < 0.001$) and high $\text{Abs}_{370,\text{BrC,sec}}$ values were consistent with the high ALWC values, this was not the case for OOA ($p > 0.1$). The slope of the linear regression (3.50) between aqSOA mass concentrations and $\text{Abs}_{370,\text{BrC,sec}}$ was higher than OOA (Fig. S17), so was the MAC values of aqSOA across UV to Vis range (Fig. S15). To further characterize the evolution of secondary BrC absorption, $\text{Abs}_{370,\text{BrC,sec}}$ was normalized by ΔCO (the background-corrected CO mixing ratios) to minimize the effect of boundary layer height (Fig. 8) (DeCarlo et al., 2010). Here, the background CO value (400 ppb) was defined as the lowest 1.25th percentile of the CO values during the campaign (Kondo et al., 2006). Fig. 8 shows that the values of $\text{Abs}_{370,\text{BrC,sec}}/\Delta\text{CO}$ increased with the increases of aqSOA and ALWC concentrations from 17:00 to 03:00 LT ($r^2 = 0.63$, 0.57 , $p < 0.001$), while $\text{Abs}_{370,\text{BrC,pri}}/\Delta\text{CO}$ slightly decreased with the increases of BBOA and m/z 60 concentrations ($r^2 = 0.35$, 0.33 , $p < 0.001$). Additionally, the mass concentrations of NO_3 , NH_4 , and NO_2 from 17:00 to 03:00 LT were 1.2, 1.2, and 1.3 times that from 04:00 to 16:00 LT during the campaign. These results were similar to those observed in SCB during winter (Peng et al., 2025; Wu et al., 2024). As described in section 3.2, the SOA with hydroxyl groups (i.e., glyoxal and methylglyoxal) could be formed from the aged BBOA via aqueous-phase reactions under high ALWC during the campaign. Previous research have shown that oligomers (involving two glyoxal molecules) formed via aqueous reactions of glyoxal with NH_3 contain C=C or C=N bonds, exhibiting strong absorption at near-UV (Laskin et al., 2015; Lee et al., 2013; Nozière et al., 2009; Powelson et al., 2014). This suggested

secondary BrC chromophores with strong absorption at 370 nm were formed under the high ALWC from 17:00 to 03:00 LT, which might be related to the aqSOA from the aged BBOA via aqueous-phase reactions. The low values of $Abs_{370, BrC, sec}/\Delta CO$ at 12:00–14:00 LT could be related to the photolysis and/or photooxidation causing BrC photobleaching (Sareen et al., 2013; Zhao et al., 2015). Overall, we suggested that aqSOA formed from biomass-burning emissions might be important for BrC absorption, especially at night during the campaign.

Previous studies also provided evidence that spring may experience similar but less intense biomass-burning activity (Chen et al., 2014; Chen et al., 2017; Tao et al., 2014), with comparable aqueous chemistry though potentially increased photochemical bleaching of BrC (Wang et al., 2019a). Though winter features lower biomass-burning emissions, secondary BrC likely formed from BBOA through aqueous-phase reactions under high NO_x and NH_4 concentrations and stagnant conditions after sunset during winter in SCB (Peng et al., 2025; Wu et al., 2024). Elevated temperature and O_x ($O_x = NO_2 + O_3$) levels could enhance photochemical oxidation for secondary BrC formation during summer, while also promoting BrC photobleaching (Wu et al., 2024). It should be noted that the seasonal biomass-burning emissions and associated chemical processing of carbonaceous aerosols must be accounted for in climate and air quality models. This is critical to avoid underestimating aerosol impacts in autumn and overestimating them in other seasons. In general, the research of BrC chromophores is still in the early stage, and more studies of the quantitative link between the chemical composition and light absorption properties of BrC chromophores and biomass-burning emissions during different season are necessary to better understand.

$AAE_{370-880}$ was another key parameter to characterize the absorption properties of aerosols, its correlations with the mass fraction of aqSOA (f_{aqSOA}) and BBOA (f_{BBOA}) to OA, and BC-to-OA ratios were shown in Fig. 9. During the campaign, the strong positive correlation ($r^2 = 0.49$, $p < 0.001$) between $AAE_{370-880}$ and f_{aqSOA} was observed with $AAE_{370-880}$ values up to 2.65, while $AAE_{370-880}$ values increased with the slight increase of f_{BBOA} in general ($r^2 = 0.21$, $p < 0.001$) (Fig. 9a and c). AAE was

674 calculated using a power-law fitting of aerosol absorption values (Qin et al., 2018;
 675 Wang et al., 2019b). While BC concentration is linearly dependent on Abs_{BC} , OA
 676 concentration does not follow the same pattern with Abs_{BC} . The mixing state of BC
 677 and OA, influenced by combustion conditions, can also affect AAE. Previous studies
 678 have shown that biomass-burning emissions can impact absorption properties, which
 679 is reflected in the relationship between AAE and the BC-to-OA ratio (a measure of the
 680 combustion conditions) (Lu et al., 2015; Saleh et al., 2014). Thus, the relationship
 681 observed in Fig. 9b reflected the influence of biomass-burning emissions during the
 682 campaign, and the parameterized curve in this study (black) was consistent with prior
 683 research (red) using wavelengths from 370 nm to 880 nm (Lu et al., 2015). Here, 950
 684 nm and 880 nm were used as the highest wavelength respectively, and similar values
 685 were found between $AAE_{370-950}$ and $AAE_{370-880}$ (within 10.0%). It should be noted
 686 that the data points of high $AAE_{370-880}$ were consistent with the low BC-to-OA ratios
 687 and large f_{aqSOA} values in general. Moreover, the average value of $AAE_{370-880}$ observed
 688 in this study (1.95) was higher than $AAE_{370-950}$ observed in the laboratory experiments
 689 of fresh and photo-chemically aged biomass-burning emissions (i.e., 1.38 and 1.48 for
 690 fresh oak and pocosin pine, 1.42 and 1.73 for aged oak and pocosin pine) (Saleh et al.,
 691 2013).




Figure 7. (a) Ternary diagram for the mass fractions of BBOA, CCOA, and HOA in POA colored by $Abs_{370,BrC,pri}$, and **(b)** scatter plot of BBOA mass concentrations versus $Abs_{370,BrC,pri}$ colored by m/z 60 mass concentrations.




Figure 8. Scatter plots of $\text{Abs}_{370,\text{BrC,pri}}/\Delta\text{CO}$ versus **(a, b)** BBOA and $\text{m/z } 60$ mass concentrations and $\text{Abs}_{370,\text{BrC,sec}}/\Delta\text{CO}$ versus **(c, d)** aqSOA and ALWC colored by the local time.




Figure 9. Relationship between **(a)** $\text{AAE}_{370-880}$ and the mass fraction of aqSOA ($f_{\text{aqSOA}} = \text{aqSOA}/\text{OA}$) colored by the mass fraction of BBOA ($f_{\text{BBOA}} = \text{BBOA}/\text{OA}$), and **(b)** BC-to-OA ratios colored by f_{aqSOA} . **(c)** Variations of f_{BBOA} as a function of $\text{AAE}_{370-880}$. The red curve in **(b)** was the best fit curve to data taken from Lu et al. (2015) and described the Abs of fresh and aged BBOA.

4 Conclusions

This study conducted comprehensive real-time measurements of the light absorption properties and chemical composition of carbonaceous aerosols during autumn in the Sichuan Basin, China. The findings provided direct ambient evidence demonstrating that aqueous secondary organic aerosol (aqSOA) formed from the aged biomass-burning emissions under high aerosol liquid water content ($\text{ALWC} > 60 \mu\text{g m}^{-3}$) conditions significantly contributes to both aerosol pollution and light absorption. Organic aerosol (OA) was identified as the dominant component of $\text{PM}_{2.5}$ ($46.6 \pm$

10.7%) and exhibited strong absorption properties at UV wavelengths ($\text{Abs}_{370,\text{BrC}} = 42.4 \pm 28.5 \text{ Mm}^{-1}$). During pollution periods (PP), the average contribution and concentration of aqSOA to OA were 14.1% and $7.6 \mu\text{g m}^{-3}$, exhibiting enhanced oxidation ($f_{29} = 0.141 \pm 0.062$, $f_{44} = 0.080 \pm 0.035$) and substantial light absorption characteristics of OA ($\text{Abs}_{370,\text{BrC}} = 91.6 \text{ Mm}^{-1}$, $\text{AAE}_{370-880} = 2.1$). Additionally, the less oxidized aqSOA formation via aqueous-phase reactions instead of photo-chemical reactions played a key role in the haze pollution dynamic evolution during PP. Furthermore, backward trajectory analysis revealed regional transport of BBOA primarily originated from northeastern Chongqing, and aged to aqSOA within approximately 12 to 48 hours. These results underscore that aqueous-phase reactions of BBOA—particularly during the transport of biomass-burning emissions—converts primary emissions into strong light-absorbing aqSOA, substantially influencing regional haze formation and radiative effects.

Our findings align with previous laboratory studies on biomass-burning BrC formation (Lu et al., 2015; Powelson et al., 2014), while providing novel ambient quantification of these processes under realistic atmospheric conditions. The parameterized curve of $\text{AAE}_{370-880}$ versus BC-to-OA ratios in this study was consistent with the previous laboratory research on biomass-burning emissions. The $\text{AAE}_{370-880}$ values observed in this study (average = 1.95) were higher than those in the laboratory experiments of fresh and photo-chemically aged biomass-burning emissions (Saleh et al., 2013), and significantly increased with the increase of f_{aqSOA} ($r^2 = 0.49$, $p < 0.001$). Additionally, the elevated $\text{Abs}_{370,\text{BrC,sec}}$ values were consistent with the high ALWC, NO_3 , and NH_4 values, and significantly increased with the increase of aqSOA concentrations ($r^2 = 0.44$, $p < 0.001$). These results suggest that aqueous-phase reactions of BBOA in high- NO_x and high- NH_4 conditions produce secondary BrC with particularly strong light absorption. The campaign was conducted during autumn, which features intensive biomass-burning activity, and thus may not fully represent aerosol processes in other seasons. The relative uncertainty range of Abs_{BrC} was $[-112\%, +42\%]$ at 370 nm due to the choice of AAE_{BC} . Our results underscore the importance of aqueous-phase processing in transforming

biomass-burning emissions, with important implications for climate and air quality modeling. The significant contribution of aqSOA to both aerosol mass and light absorption highlights the need for improved representation of aqueous processes in models. The linkages between aging timescales, transport pathways, and aqSOA formation in this study provide a transferable framework for understanding aqSOA processing in other humid regions influenced by biomass burning. Future research should prioritize molecular-level characterization of aqSOA precursors and products, quantification of aqueous reaction rates under ambient conditions, and multi-scale modeling to assess regional climate impacts. This study highlights that aqueous processes play an important role in the evolution of biomass-burning emissions and should be adequately considered in both air quality budgets and climate forcing balance on a global scale.

Data availability. The data generated and analysed in this study are available from <https://doi.org/10.5281/zenodo.14626304> (Peng et al., 2025).

Author contributions. CZ, CP, YD, and ZL designed the experiments. Data analysis and interpretation were performed by CP, ZT, HT, KZ, ZL, and GS. CP, XY, and MT wrote the paper. ZT, YC, XL, LZ, YC, and YF contributed to the paper with useful scientific discussions or comments.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. This study was supported by the National Natural Science Foundation of China (No. 42305126), National Key Research and Development Program of China (No. 2023YFC3709301), Natural Science Foundation of Chongqing Municipality (No. CSTB2022NSCQ-MSX0504).

References

- Akagi, S. K., Yokelson, R. J., Wiedinmyer, C., Alvarado, M. J., Reid, J. S., Karl, T., Crounse, J. D., and Wennberg, P. O.: Emission factors for open and domestic biomass burning for use in atmospheric models, *Atmos. Chem. Phys.*, 11, 4039–4072, <https://doi.org/10.5194/acp-11-4039-2011>, 2011.
- Alfarra, M. R., Prévôt, A. S. H., Szidat, S., Sandradewi, J., Sandradewi, S., Lanz, V. A., Schreiber, D., Mohr, M., and Baltensperger, U.: Identification of the Mass Spectral Signature of Organic Aerosols from Wood Burning Emissions, *Environ. Sci. Technol.*, 41, 5770–5777, <https://doi.org/10.1021/es062289b>, 2007.
- Bahreini, R., Ervens, B., Middlebrook, A. M., Warneke, C., de Gouw, J. A., DeCarlo, P. F., Jimenez, J. L., Brock, C. A., Neuman, J. A., Ryerson, T. B., Stark, H., Atlas, E., Brioude, J., Fried, A., Holloway, J. S., Peischl, J., Richter, D., Walega, J., Weibring, P., Wollny, A. G., and Fehsenfeld, F. C.: Organic Aerosol Formation in Urban and Industrial Plumes Near Houston and Dallas, Texas, *J. Geophys. Res.: Atmos.*, 114, D00F16, [doi:10.1029/2008JD011493](https://doi.org/10.1029/2008JD011493), 2009.
- Bao, Z. E., Liu, Y. L., Meng, L. S., Han, Y., Tian, M., Shi, G. M., Wang, Q. Y., Huang, Y., Peng, C., Luo, B., Zhang, W., Wang, H. B., Cao, J. J., Yang, F. M., and Chen, Y.: Evaluating the effects of biomass burning on severe haze formation in a megacity of Sichuan Basin, Southwestern China, *J. Geophys. Res.: Atmos.*, 130, e2024JD042516, <https://doi.org/10.1029/2024JD042516>, 2025.
- Bao, Z. E., Zeng, X. L., Zhou, J. W., Yang, F. M., Lu, K. D., Zhai, C. Z., Li, X., Feng, M., Tan, Q. W., and Chen, Y.: Evolution of black carbon and brown carbon during summertime in Southwestern China: An assessment of control measures during the 2023 Chengdu Summer World University Games, *Environ. Pollut.*, 357, 124467, <https://doi.org/10.1016/j.envpol.2024.124467>, 2024.
- Bao, Z. E., Zhang, X. Y., Li, Q., Zhou, J. W., Shi, G. M., Zhou, L., Yang, F. M., Xie, S. D., Zhang, D., Zhai, C. Z., Li, Z. L., Peng, C., and Chen, Y.: Measurement report: Intensive biomass burning emissions and rapid nitrate formation drive severe haze formation in the Sichuan Basin, China – insights from aerosol mass spectrometry, *Atmos. Chem. Phys.*, 23, 1147–1167, <https://doi.org/10.5194/acp-23-1147-2023>,

2023.

Canagaratna, M. R., Jayne, J. T., Jimenez, J. L., Allan, J. D., Alfarra, M. R., Zhang, Q., Onasch, T. B., Drewnick, F., Coe, H., Middlebrook, A., Delia, A., Williams, L. R., Trimborn, A. M., Northway, M. J., DeCarlo, P. F., Kolb, C. E., Davidovits, P., and Worsnop, D. R.: Chemical and microphysical characterization of ambient aerosols with the aerodyne aerosol mass spectrometer, *Mass Spectrom. Rev.*, 26, 185–222, <https://doi.org/10.1002/mas.20115>, 2007.

Canagaratna, M. R., Jimenez, J. L., Kroll, J. H., Chen, Q., Kessler, S. H., Massoli, P., Hildebrandt Ruiz, L., Fortner, E., Williams, L. R., Wilson, K. R., Surratt, J. D., Donahue, N. M., Jayne, J. T., and Worsnop, D. R.: Elemental ratio measurements of organic compounds using aerosol mass spectrometry: characterization, improved calibration, and implications, *Atmos. Chem. Phys.*, 15, 253–272, <https://doi.org/10.5194/acp-15-253-2015>, 2015.

Canonaco, F., Crippa, M., Slowik, J. G., Baltensperger, U., and Prévôt, A. S. H.: SoFi, an IGOR-based interface for the efficient use of the generalized multilinear engine (ME-2) for the source apportionment: ME-2 application to aerosol mass spectrometer data, *Atmos. Meas. Tech.*, 6, 3649–3661, <https://doi.org/10.5194/amt-6-3649-2013>, 2013.

Carlton, A. G., Turpin, B. J., Altieri, K. E., Seitzinger, S., Reff, A., Lim, H. J., and Ervens, B.: Atmospheric oxalic acid and SOA production from glyoxal: results of aqueous photooxidation experiments, *Atmos. Environ.*, 41, 7588–7602, <https://doi.org/10.1016/j.atmosenv.2007.05.035>, 2007.

Chen, C. R., Zhang, H. X., Yan, W. J., Wu, N. N., Zhang, Q., and He, K. B.: Aerosol water content enhancement leads to changes in the major formation mechanisms of nitrate and secondary organic aerosols in winter over the North China Plain, *Environ. Pollut.*, 287, 117625, <https://doi.org/10.1016/j.envpol.2021.117625>, 2021.

Chen, Y., Tian, M., Huang, R. J., Shi, G. M., Wang, H. B., Peng, C., Cao, J. J., Wang, Q. Y., Zhang, S. M., Guo, D. M., Zhang, L. Y., and Yang, F. M.: Characterization of urban amine-containing particles in southwestern China: seasonal variation, source, and processing, *Atmos. Chem. Phys.*, 19, 3245–3255,

- <https://doi.org/10.5194/acp-19-3245-2019>, 2019.
- Chen, Y., Xie, S. D., Luo, B., and Zhai, C. Z.: Characteristics and origins of carbonaceous aerosol in the Sichuan Basin, China, *Atmos. Environ.*, 94, 215–223, <https://doi.org/10.1016/j.atmosenv.2014.05.037>, 2014.
- Chen, Y., Xie, S. D., Luo, B., and Zhai, C. Z.: Particulate pollution in urban Chongqing of southwest China: Historical trends of variation, chemical characteristics and source apportionment, *Sci. Total Environ.*, 584–585, 523–534, <https://doi.org/10.1016/j.scitotenv.2017.01.060>, 2017.
- Coen, M. C., Weingartner, E., Apituley, A., Ceburnis, D., Fierz-Schmidhauser, R., Flentje, H., Henzing, J. S., Jennings, S. G., Moerman, M., Petzold, A., Schmid, O., and Baltensperger, U.: Minimizing light absorption measurement artifacts of the Aethalometer: Evaluation of five correction algorithms, *Atmos. Meas. Tech.*, 3, 457–474, <https://doi.org/10.5194/amt-3-457-2010>, 2010.
- Cubison, M. J., Ortega, A. M., Hayes, P. L., Farmer, D. K., Day, D., Lechner, M. J., Brune, W. H., Apel, E., Diskin, G. S., Fisher, J. A., Fuelberg, H. E., Hecobian, A., Knapp, D. J., Mikoviny, T., Rierner, D., Sachse, G. W., Sessions, W., Weber, R. J., Weinheimer, A. J., Wisthaler, A., and Jimenez, J. L.: Effects of aging on organic aerosol from open biomass burning smoke in aircraft and laboratory studies, *Atmos. Chem. Phys.*, 11, 12049–12064, <https://doi.org/10.5194/acp-11-12049-2011>, 2011.
- DeCarlo, P. F., Ulbrich, I. M., Crounse, J., de Foy, B., Dunlea, E. J., Aiken, A. C., Knapp, D., Weinheimer, A. J., Campos, T., Wennberg, P. O., and Jimenez, J. L.: Investigation of the Sources and Processing of Organic Aerosol over the Central Mexican Plateau from Aircraft Measurements during MILAGRO, *Atmos. Chem. Phys.*, 10, 5257–5280, <https://doi.org/10.5194/acp-10-5257-2010>, 2010.
- Drinovec, L., Močnik, G., Zotter, P., Prévôt, A. S. H., Ruckstuhl, C., Coz, E., Rupakheti, M., Sciare, J., Müller, T., Wiedensohler, A., and Hansen, A. D. A.: The "dual-spot" Aethalometer: an improved measurement of aerosol black carbon with real-time loading compensation, *Atmos. Meas. Tech.*, 8, 1965–1979, <https://doi.org/10.5194/amt-8-1965-2015>, 2015.
- Drozd, G. T. and McNeill, V. F.: Organic matrix effects on the formation of

light-absorbing compounds from α -dicarbonyls in aqueous salt solution, *Environ. Sci.: Processes Impacts*, 16, 741–747, <https://doi.org/10.1039/c3em00579h>, 2014.

Elser, M., Huang, R. J., Wolf, R., Slowik, J. G., Wang, Q. Y., Canonaco, F., Li, G. H., Bozzetti, C., Daellenbach, K. R., Huang, Y., Zhang, R. J., Li, Z. Q., Cao, J. J., Baltensperger, U., El-Haddad, I., and Prévôt, A. S. H.: New insights into PM_{2.5} chemical composition and sources in two major cities in China during extreme haze events using aerosol mass spectrometry, *Atmos. Chem. Phys.*, 16, 3207–3225, <https://doi.org/10.5194/acp-16-3207-2016>, 2016.

Ervens, B., Turpin, B. J., and Weber, R. J.: Secondary organic aerosol formation in cloud droplets and aqueous particles (aqSOA): a review of laboratory, field and model studies, *Atmos. Chem. Phys.*, 11, 11069–11102, <https://doi.org/10.5194/acp-11-11069-2011>, 2011.

Fountoukis, C. and Nenes, A.: ISORROPIA II: a computationally efficient thermodynamic equilibrium model for $K^+Ca^{2+}Mg^{2+}NH_4^+Na^+SO_4^{2-}NO_3^-Cl^-H_2O$ aerosols, *Atmos. Chem. Phys.*, 7, 4639–4659, <https://doi.org/10.5194/acp-7-4639-2007>, 2007.

Fröhlich, R., Cubison, M. J., Slowik, J. G., Bukowiecki, N., Prévôt, A. S. H., Baltensperger, U., Schneider, J., Kimmel, J. R., Gonin, M., Rohner, U., Worsnop, D. R., and Jayne, J. T.: The ToF-ACSM: a portable aerosol chemical speciation monitor with TOFMS detection, *Atmos. Meas. Tech.*, 6, 3225–3241, <https://doi.org/10.5194/amt-6-3225-2013>, 2013.

Gilardoni, S., Massoli, P., Paglione, M., Giulianelli, L., Carbone, C., Rinaldi, M., Decesari, S., Sandrini, S., Costabile, F., Gobbi, G. P., Pietrogrande, M. C., Visentin, M., Scotto, F., Fuzzi, S., and Facchini, M. C.: Direct observation of aqueous secondary organic aerosol from biomass-burning emissions, *Proc. Natl. Acad. Sci. USA*, 113, 10013–10018, <https://doi.org/10.1073/pnas.1602212113>, 2016.

Guo, H. Y., Xu, L., Bougiatioti, A., Cerully, K. M., Capps, S. L., Hite Jr., J.R., Carlton, A. G., Lee, S. H., Bergin, M. H., Ng, N. L., Nenes, A., and Weber, R. J.: Fine-particle water and pH in the southeastern United States, *Atmos. Chem. Phys.*, 15, 5211–5228, <https://doi.org/10.5194/acp-15-5211-2015>, 2015.

Gyawali, M., Arnott, W. P., Lewis, K., and Moosmüller, H.: In situ aerosol optics in Reno, NV, USA during and after the summer 2008 California wildfires and the influence of absorbing and non-absorbing organic coatings on spectral light absorption, *Atmos. Chem. Phys.*, 9, 8007–8015, <https://doi.org/10.5194/acp-9-8007-2009>, 2009.

He, M., Wang, X. R., Han, L., Feng, X. D., and M, X.: Emission Inventory of Crop Residues Field Burning and Its Temporal and Spatial Distribution in Sichuan Province, *Environ. Sci.*, 36, 1208–1216, <https://doi.org/10.13227/j.hjhx.2015.04.010>, 2015.

Hennigan, C. J., Izumi, J., Sullivan, A. P., Weber, R. J., and Nenes, A.: A critical evaluation of proxy methods used to estimate the acidity of atmospheric particles, *Atmos. Chem. Phys.*, 15, 2775–2790, <https://doi.org/10.5194/acp-15-2775-2015>, 2015.

Hennigan, C. J., Sullivan, A. P., Collett, J. L., and Robinson, A. L.: Levoglucosan stability in biomass burning particles exposed to hydroxyl radicals, *Geophys. Res. Lett.*, 37, L09806, <https://doi.org/10.1029/2010GL043088>, 2010.

Herrmann, H., Schaefer, T., Tilgner, A., Styler, S. A., Weller, C., Teich, M., and Otto, T.: Tropospheric Aqueous-Phase Chemistry: Kinetics, Mechanisms, and Its Coupling to a Changing Gas Phase, *Chem. Rev.*, 115, 4259–4334, <https://doi.org/10.1021/cr500447k>, 2015.

Huang, R. J., Zhang, Y. L., Bozzetti, C., Ho, K. F., Cao, J. J., Han, Y. M., Daellenbach, K. R., Slowik, J. G., Platt, S. M., Canonaco, F., Zotter, P., Wolf, R., Pieber, S. M., Bruns, E. A., Crippa, M., Ciarelli, G., Piazzalunga, A., Schwikowski, M., Abbaszade, G., Schnelle-Kreis, J., Zimmermann, R., An, Z. S., Szidat, S., Baltensperger, U., El Haddad, I., and Prevot, A. S.: High secondary aerosol contribution to particulate pollution during haze events in China, *Nature*, 514, 218–222, <https://doi.org/10.1038/nature13774>, 2014.

Huang, R. J., He, Y., Duan, J., Li, Y. J., Chen, Q., Zheng, Y., Chen, Y., Hu, W. W., Lin, C. S., Ni, H. Y., Dai, W. T., Cao, J. J., Wu, Y. F., Zhang, R. J., Xu, W., Ovadnevaite, J., Ceburnis, D., Hoffmann, T., and O'Dowd, C. D.: Contrasting sources and

processes of particulate species in haze days with low and high relative humidity in wintertime Beijing, *Atmos. Chem. Phys.*, 20, 9101–9114, <https://doi.org/10.5194/acp-20-9101-2020>, 2020.

Huang, X. J., Liu, Z., Ge, Y. Z., Li, Q., Wang, X. F., Fu, H. B., Zhu, J., Zhou, B., Wang, L., George, C., Wang, Y., Wang, X. F., Su, J. X., Xue, L. K., Yu, S. C., Mellouki, A., and Chen, J. M.: Aerosol high water contents favor sulfate and secondary organic aerosol formation from fossil fuel combustion emissions, *npj Clim. Atmos. Sci.*, 6, 173, <https://doi.org/10.1038/s41612-023-00504-1>, 2023.

Jimenez, J. L., Canagaratna, M. R., Donahue, N. M., Prevot, A. S. H., Zhang, Q., Kroll, J. H., DeCarlo, P. F., Allan, J. D., Coe, H., Ng, N. L., Aiken, A. C., Docherty, K. S., Ulbrich, I. M., Grieshop, A. P., Robinson, A. L., Duplissy, J., Smith, J. D., Wilson, K. R., Lanz, V. A., Hueglin, C., Sun, Y. L., Tian, J., Laaksonen, A., Raatikainen, T., Rautiainen, J., Vaattovaara, P., Ehn, M., Kulmala, M., Tomlinson, J. M., Collins, D. R., Cubison, M. J., Dunlea, J., Huffman, J. A., Onasch, T. B., Alfarra, M. R., Williams, P. I., Bower, K., Kondo, Y., Schneider, J., Drewnick, F., Borrmann, S., Weimer, S., Demerjian, K., Salcedo, D., Cottrell, L., Griffin, R., Takami, A., Miyoshi, T., Hatakeyama, S., Shimo, A., Sun, J. Y., Zhang, Y. M., Dzepina, K., Kimmel, J. R., Sueper, D., Jayne, J. T., Herndon, S. C., Trimborn, A. M., Williams, L. R., Wood, E. C., Middlebrook, A. M., Kolb, C. E., Baltensperger, U., and Worsnop, D. R.: Evolution of Organic Aerosols in the Atmosphere, *Science*, 326, 1525–1529, <https://doi.org/10.1126/science.1180353>, 2009.

Kampf, C. J., Jakob, R., and Hoffmann, T.: Identification and characterization of aging products in the glyoxal/ammonium sulfate system – implications for light-absorbing material in atmospheric aerosols, *Atmos. Chem. Phys.*, 12, 6323–6333, <https://doi.org/10.5194/acp-12-6323-2012>, 2012.

Kim, H., Collier, S., Ge, X. L., Xu, J. Z., Sun, Y. L., Jiang, W. Q., Wang, Y. L., Herckes, P., and Zhang, Q.: Chemical processing of water-soluble species and formation of secondary organic aerosol in fogs, *Atmos. Environ.*, 200, 158–166, <https://doi.org/10.1016/j.atmosenv.2018.11.062>, 2019.

Kirchstetter, T. W. and Novakov T.: Evidence that the spectral dependence of light

absorption by aerosols is affected by organic carbon, *J. Geophys. Res.*, 109, D21208, <https://doi.org/10.1029/2004JD004999>, 2004.

Kondo, Y., Komazaki, Y., Miyazaki, Y., Moteki, N., Takegawa, N., Kodama, D., Deguchi, S., Nogami, M., Fukuda, M., Miyakawa, T., Morino, Y., Koike, M., Sakurai, H., and Ehara, K.: Temporal Variations of Elemental Carbon in Tokyo, *J. Geophys. Res.*, 111, D12205, <https://doi.org/10.1029/2005JD006257>, 2006.

Kourtchev, I., Giorio, C., Manninen, A., Wilson, E., Mahon, B., Aalto, J., Kajos, M., Venables, D., Ruuskanen, T., Levula, J., Lopenen, M., Connors, S., Harris, N., Zhao, D. F., Kiendler-Scharr, A., Mentel, T., Rudich, Y., Hallquist, M., Doussin, J. F., Maenhaut, W., Bäck, J., Petäjä, T., Wenger, J., Kulmala, Ma., and Kalberer, M.: Enhanced Volatile Organic Compounds emissions and organic aerosol mass increase the oligomer content of atmospheric aerosols, *Sci. Rep.*, 6, 35038, <https://doi.org/10.1038/srep35038>, 2016.

Kroftic, A., Grilc, M., and Grgic, I.: Unraveling pathways of guaiacol nitration in atmospheric waters: nitrite, a source of reactive nitronium ion in the atmosphere, *Environ. Sci. Technol.*, 49, 9150–9158, <https://doi.org/10.1021/acs.est.5b01811>, 2015.

Lack, D. A., and Langridge, J. M.: On the attribution of black and brown carbon light absorption using the Ångström exponent, *Atmos. Chem. Phys.*, 13, 10535–10543, <https://doi.org/10.5194/acp-13-10535-2013>, 2013.

Lanz, V. A., Alfarra, M. R., Baltensperger, U., Buchmann, B., Hueglin, C., and Prévôt, A. S. H.: Source apportionment of submicron organic aerosols at an urban site by factor analytical modelling of aerosol mass spectra, *Atmos. Chem. Phys.*, 7, 1503–1522, doi:10.5194/acp-7-1503-2007, 2007.

Laskin, A., Laskin, J., and Nizkorodov, S. A.: Chemistry of atmospheric brown carbon, *Chem. Rev.*, 115, 4335–4382, <https://doi.org/10.1021/cr5006167>, 2015.

Lee, A. K., Zhao, R., Li, R., Liggio, J., Li, S. M., and Abbatt, J. P.: Formation of Light Absorbing Organo-Nitrogen Species from Evaporation of Droplets Containing Glyoxal and Ammonium Sulfate, *Environ. Sci. Technol.*, 47, 12819–12826, <https://doi.org/10.1021/es402687w>, 2013.

- Lei, Y., Lei, X., Tian, G., Yang, J., Huang, D., Yang, X., Chen, C. C., and Zhao, J. C.:
Optical Variation and Molecular Transformation of Brown Carbon During
Oxidation by $\text{NO}_3\cdot$ in the Aqueous Phase, *Environ. Sci. Technol.*, 58, 3353–3362,
<https://doi.org/10.1021/acs.est.3c08726>, 2024.
- Li, C. L., He, Q. F., Fang, Z., Brown, S. S., Laskin, A., Cohen, S. R., and Rudich, Y.:
Laboratory Insights into the Diel Cycle of Optical and Chemical Transformations
of Biomass Burning Brown Carbon Aerosols, *Environ. Sci. Technol.*, 54,
11827–11837, <https://dx.doi.org/10.1021/acs.est.0c04310>, 2020.
- Li, F. H., Zhou, S. Z., Du, L., Zhao, J., Hang, J., and Wang, X. M.: Aqueous-phase
chemistry of atmospheric phenolic compounds: A critical review of laboratory
studies, *Sci. Total Environ.*, 856, 158895,
<https://doi.org/10.1016/j.scitotenv.2022.158895>, 2023.
- Li, G., Bei, N., Tie, X., and Molina, L. T.: Aerosol effects on the photochemistry in
Mexico City during MCMA-2006/MILAGRO campaign, *Atmos. Chem. Phys.*, 11,
5169–5182, <https://doi.org/10.5194/acp-11-5169-2011>, 2011.
- Li, Z. J., Tan, H. B., Zheng, J., Liu, L., Qin, Y. M., Wang, N., Li, F., Li, Y. J., Cai, M.
F., Ma, Y., and Chan, C. K.: Light absorption properties and potential sources of
particulate brown carbon in the Pearl River Delta region of China, *Atmos. Chem.
Phys.*, 19, 11669–11685, <https://doi.org/10.5194/acp-19-11669-2019>, 2019.
- Lim, Y. B., Tan, Y., Perri, M. J., Seitzinger, S. P., and Turpin, B. J.: Aqueous chemistry
and its role in secondary organic aerosol (SOA) formation, *Atmos. Chem. Phys.*,
10, 10521–10539, <https://doi.org/10.5194/acp-10-10521-2010>, 2010.
- Liu, M. X., Song, Y., Zhou, T., Xu, Z. Y., Yan, C. Q., Zheng, M., Wu, Z. J., Hu, M.,
Wu, Y. S., and Zhu, T.: Fine particle pH during severe haze episodes in northern
China, *Geophys. Res. Lett.*, 44, 5213–5221, <https://doi.org/10.1002/2017gl073210>,
2017.
- Liu, Y., Huang, R. J., Lin, C. S., Yuan, W., Li, Y. J., Zhong, H. B., Yang, L., Wang, T.,
Huang, W., Xu, W., Huang, D. D., and Huang, C.: Nitrate-Photolysis Shortens the
Lifetimes of Brown Carbon Tracers from Biomass Burning, *Environ. Sci. Technol.*,
59, 640–649, <https://doi.org/10.1021/acs.est.4c06123>, 2024.

- Lorenzo, R. A. D., Washenfelter, R. A., Attwood, A. R., Guo, H., Xu, L., Ng, N. L., Weber, R. J., Baumann, K., Edgerton, E., and Young, C. J.: Molecular-Size-Separated Brown Carbon Absorption for Biomass-Burning Aerosol at Multiple Field Sites, *Environ. Sci. Technol.*, 51, 3128–3137, <https://doi.org/10.1021/acs.est.6b06160>, 2017.
- Lu, Z., Streets, D. G., Winijkul, E., Yan, F., Chen, Y., Bond, T. C., Feng, Y., Dubey, M. K., Liu, S., Pinto, J. P., and Carmichael, G. R.: Light absorption properties and radiative effects of primary organic aerosol emissions, *Environ. Sci. Technol.*, 49, 4868–4877, <https://doi.org/10.1021/acs.est.5b00211>, 2015.
- Luo, J. Q., Zhang, J. K., Huang, X. J., Liu, Q., Luo, B., Zhang, W., Rao, Z. H., and Yu, Y. C.: Characteristics, evolution, and regional differences of biomass burning particles in the Sichuan Basin, China, *J. Environ. Sci.*, 89, 35–46, <https://doi.org/10.1016/j.jes.2019.09.015>, 2020.
- Matthew, B. M., Middlebrook, A. M., and Onasch, T. B.: Collection Efficiencies in an Aerodyne Aerosol Mass Spectrometer as a Function of Particle Phase for Laboratory Generated Aerosols, *Aerosol Sci. Technol.*, 42, 884–898, <https://doi.org/10.1080/02786820802356797>, 2008.
- McNeill, V. F.: Aqueous Organic Chemistry in the Atmosphere: Sources and Chemical Processing of Organic Aerosols, *Environ. Sci. Technol.*, 49, 1237–1244, <https://doi.org/10.1021/es5043707>, 2015.
- Meng, J. J., Liu, X. D., Hou, Z. F., Yi, Y. A., Yan, L., Li, Z., Cao, J. J., Li, J. J., and Wang, G. H.: Molecular characteristics and stable carbon isotope compositions of dicarboxylic acids and related compounds in the urban atmosphere of the North China Plain: Implications for aqueous phase formation of SOA during the haze periods, *Sci. Total Environ.*, 705, 135256, <https://doi.org/10.1016/j.scitotenv.2019.135256>, 2020.
- Middlebrook, A. M., Bahreini, R., Jimenez, J. L., and Canagaratna, M. R.: Evaluation of Composition-Dependent Collection Efficiencies for the Aerodyne Aerosol Mass Spectrometer using Field Data, *Aerosol Sci. Technol.*, 46, 258–271, <https://doi.org/10.1080/02786826.2011.620041>, 2012.

- Ministry of Environmental Protection (MEP): P.R. Of China, China National Ambient Air Quality Standard (GB 3095-2012), http://kjs.mep.gov.cn/hjbhzbz/bzwb/dqhjbh/dqhjzlbz/201203/t20120302_224165.htm, 2012.
- Moosmüller, H., Chakrabarty, R. K., and Arnott, W. P.: Aerosol light absorption and its measurement: A review, *J. Quant. Spectrosc. Ra.*, 110, 844–878, <https://doi.org/10.1016/j.jqsrt.2009.02.035>, 2009.
- Ng, N. L., Canagaratna, M. R., Jimenez, J. L., Chhabra, P. S., Seinfeld, J. H., and Worsnop, D. R.: Changes in organic aerosol composition with aging inferred from aerosol mass spectra, *Atmos. Chem. Phys.*, 11, 6465–6474, <https://doi.org/10.5194/acp-11-6465-2011>, 2011a.
- Ng, N. L., Canagaratna, M. R., Jimenez, J. L., Zhang, Q., Ulbrich, I. M., and Worsnop, D. R.: Real-Time Methods for Estimating Organic Component Mass Concentrations from Aerosol Mass Spectrometer Data, *Environ. Sci. Technol.*, 45, 910–916, <https://doi.org/10.1021/es102951k>, 2011b.
- Ng, N. L., Canagaratna, M. R., Zhang, Q., Jimenez, J. L., Tian, J., Ulbrich, I. M., Kroll, J. H., Docherty, K. S., Chhabra, P. S., Bahreini, R., Murphy, S. M., Seinfeld, J. H., Hildebrandt, L., Donahue, N. M., DeCarlo, P. F., Lanz, V. A., Prévôt, A. S. H., Dinar, E., Rudich, Y., and Worsnop, D. R.: Organic aerosol components observed in Northern Hemispheric datasets from Aerosol Mass Spectrometry, *Atmos. Chem. Phys.*, 10, 4625–4641, <https://doi.org/10.5194/acp-10-4625-2010>, 2010.
- Ng, N. L., Herndon, S. C., Trimborn, A., Canagaratna, M. R., Croteau, P. L., Onasch, T. B., Sueper, D., Worsnop, D. R., Zhang, Q., Sun, Y. L., and Jayne J. T.: An Aerosol Chemical Speciation Monitor (ACSM) for Routine Monitoring of the Composition and Mass Concentrations of Ambient Aerosol, *Aerosol Sci. Technol.*, 45, 780–794, <https://doi.org/10.1080/02786826.2011.560211>, 2011c.
- Nguyen, T. K. V., Zhang, Q., Jimenez, J. L., Pike, M., and Carlton, A. G.: Liquid Water: Ubiquitous Contributor to Aerosol Mass, *Environ. Sci. Technol. Lett.*, 3, 257–263, <https://doi.org/10.1021/acs.estlett.6b00167>, 2016.
- Nozière, B., and Esteve, W.: Light-absorbing aldol condensation products in acidic aerosols: Spectra, kinetics, and contribution to the absorption index, *Atmos.*

- Environ., 41, 1150–1163, <https://doi.org/10.1016/j.atmosenv.2006.10.001>, 2007.
- Nozière, B., Dziedzic, P., and Córdova, A.: Products and Kinetics of the Liquid-Phase Reaction of Glyoxal Catalyzed by Ammonium Ions (NH_4^+), *J. Phys. Chem. A*, 113, 231–237, <https://doi.org/10.1021/jp8078293>, 2009.
- Ortega, A. M., Ortega, D. A., Cubison, M. J., Brune, W. H., Bon, D., Gouw de, J. A., and Jimenez, J. L.: Secondary organic aerosol formation and primary organic aerosol oxidation from biomass-burning smoke in a flow reactor during FLAME-3, *Atmos. Chem. Phys.*, 13, 11551–11571, <https://doi.org/10.5194/acp-13-11551-2013>, 2013.
- Ortiz-Montalvo, D. L., Lim, Y. B., Perri, M. J., Seitzinger, S. P., and Turpin, B. J.: Volatility and Yield of Glycolaldehyde SOA Formed through Aqueous Photochemistry and Droplet Evaporation, *Aerosol Sci. Technol.*, 46, 1002–1014, <https://doi.org/10.1080/02786826.2012.686676>, 2012.
- Paatero, P.: The Multilinear Engine: A Table-Driven, Least Squares Program for Solving Multilinear Problems, Including the n-Way Parallel Factor Analysis Model, *J. Comput. Graph. Stat.*, 8, 854–888, <https://doi.org/10.1080/10618600.1999.10474853>, 1999.
- Paatero, P. and Hopke, P. K.: Discarding or downweighting high-noise variables in factor analytic models, *Anal. Chim. Acta.*, 490, 277–289, [https://doi.org/10.1016/S0003-2670\(02\)01643-4](https://doi.org/10.1016/S0003-2670(02)01643-4), 2003.
- Paatero, P. and Tapper, U.: Positive matrix factorization: A non-negative factor model with optimal utilization of error estimates of data values, *Environmetrics*, 5, 111–126, <https://doi.org/10.1002/env.3170050203>, 1994.
- Paglione, M., Gilardoni, S., Rinaldi, M., Decesari, S., Zanca, N., Sandrini, S., Giulianelli, L., Bacco, D., Ferrari, S., Poluzzi, V., Scotto, F., Trentini, A., Poulain, L., Herrmann, H., Wiedensohler, A., Canonaco, F., Prévôt, A. S. H., Massoli, P., Carbone, C., Facchini, M. C., and Fuzzi, S. C.: The impact of biomass burning and aqueous-phase processing on air quality: a multi-year source apportionment study in the Po Valley, Italy, *Atmos. Chem. Phys.*, 20, 1233–1254, <https://doi.org/10.5194/acp-20-1233-2020>, 2020.

- Pang, H. W., Zhang, Q., Lu, X. H., Li, K. N., Chen, H., Chen, J. M., Yang, X., Ma, Y. G., Ma, J. L., and Huang, C.: Nitrite-Mediated Photooxidation of Vanillin in the Atmospheric Aqueous Phase, *Environ. Sci. Technol.*, 53, 14253–14263, <https://doi.org/10.1021/acs.est.9b03649>, 2019.
- Peng, C., Tian, M., Chen, Y., Wang, H. B., Zhang, L. M., Shi, G. M., Liu, Y., Yang, F. M., and Zhai, C. Z.: Characteristics, Formation Mechanisms and Potential Transport Pathways of PM_{2.5} at a Rural Background Site in Chongqing, Southwest China, *Aerosol Air Qual. Res.*, 19, 1980–1992, <https://doi.org/10.4209/aaqr.2019.01.0010>, 2019.
- Peng, C., Tian, M., Shi, G. M., Zhang, S. M., Long, X., Che, H. X., Zhong, J., You, X. Y., Bao, Z. E., Yang, F. M., Qi, X., Zhai, C. Z., and Chen, Y.: Sources and light absorption of brown carbon in urban areas of the Sichuan Basin, China: Contribution from biomass burning and secondary formation, *Atmos. Res.*, 318, 107992, <https://doi.org/10.1016/j.atmosres.2025.107992>, 2025.
- Powelson, M. H., Espelien, B. M., Hawkins, L. N., Galloway, M. M., and Haan, D. O. D.: Brown Carbon Formation by Aqueous-Phase Carbonyl Compound Reactions with Amines and Ammonium Sulfate, *Environ. Sci. Technol.*, 48, 985–993, <https://doi.org/10.1021/es4038325>, 2014.
- Qin, Y. M., Tan, H. B., Li, Y. J., Li, Z. J., Schurman, M. I., Liu, L., Wu, C., and Chan, C. K.: Chemical characteristics of brown carbon in atmospheric particles at a suburban site near Guangzhou, China, *Atmos. Chem. Phys.*, 18, 16409–16418, <https://doi.org/10.5194/acp-18-16409-2018>, 2018.
- Saleh, R., Hennigan, C. J., McMeeking, G. R., Chuang, W. K., Robinson, E. S., Coe, H., Donahue, N. M., and Robinson, A. L.: Absorptivity of brown carbon in fresh and photo-chemically aged biomass-burning emissions, *Atmos. Chem. Phys.*, 13, 7683–7693, <https://doi.org/10.5194/acp-13-7683-2013>, 2013.
- Saleh, R., Robinson, E. S., Tkacik, D. S., Ahern, A. T., Liu, S., Aiken, A. C., Sullivan, R. C., Presto, A. A., Dubey, M. K., Yokelson, R. J., Donahue, N. M., and Robinson, A. L.: Brownness of organics in aerosols from biomass burning linked to their black carbon content, *Nat. Geosci.*, 7, 647–650, <https://doi.org/10.1038/ngeo2220>,

- 2014.
- Sareen, N., Moussa, S. G., and McNeill, V. F.: Photochemical Aging of
Light-Absorbing Secondary Organic Aerosol Material, *J. Phys. Chem. A*, 117,
2987–2996, <https://doi.org/10.1021/jp309413j>, 2013.
- Shrivastava, M., Cappa, C. D., Fan, J. W., Goldstein, A. H., Guenther, A. B., Jimenez,
J. L., Kuang, C. A., Laskin, A., Martin, S. T., Ng, N. L., Petaja, T., Pierce, J. R.,
Rasch, P. J., Roldin, P., Seinfeld, J. H., Shilling, J., Smith, J. N., Thornton, J. A.,
Volkamer, R., Wang, J., Worsnop, D. R., Zaveri, R. A., Zelenyuk, A., and Zhang,
Q.: Recent advances in understanding secondary organic aerosol: Implications for
global climate forcing, *Rev. Geophys.*, 55, 509–559, doi:10.1002/2016RG000540,
2017.
- Sun, Y. L., Du, W., Fu, P. Q., Wang, Q. Q., Li, J., Ge, X. L., Zhang, Q., Zhu, C. M.,
Ren, L. J., Xu, W. Q., Zhao, J., Han, T. T., Worsnop, D. R., and Wang, Z. F.:
Primary and secondary aerosols in Beijing in winter: sources, variations and
processes, *Atmos. Chem. Phys.*, 16, 8309–8329,
<https://doi.org/10.5194/acp-16-8309-2016>, 2016a.
- Sun, Y. L., Jiang, Q., Xu, Y. S., Ma, Y., Zhang, Y. J., Liu, X. G., Li, W. J., Wang, F., Li,
J., Wang, P. C., and Li, Z. Q.: Aerosol characterization over the North China Plain:
Haze life cycle and biomass burning impacts in summer, *J. Geophys. Res.: Atmos.*,
121, 2508–2521, <https://doi.org/10.1002/2015jd024261>, 2016b.
- Sun, Y. L., Zhang, Q., Anastasio, C., and Sun, J.: Insights into secondary organic
aerosol formed via aqueous-phase reactions of phenolic compounds based on high
resolution mass spectrometry, *Atmos. Chem. Phys.*, 10, 4809–4822,
<https://doi.org/10.5194/acp-10-4809-2010>, 2010.
- Tan, Y., Lim, Y. B., Altieri, K. E., Seitzinger, S. P., and Turpin, B. J.: Mechanisms
leading to oligomers and SOA through aqueous photooxidation: insights from OH
radical oxidation of acetic acid and methylglyoxal, *Atmos. Chem. Phys.*, 12,
801–813, <https://doi.org/10.5194/acp-12-801-2012>, 2012.
- Tao, J., Gao, J., Zhang, L., Zhang, R., Che, H., Zhang, Z., Lin, Z., Jing, J., Cao, J., and
Hsu, S. C.: PM_{2.5} pollution in a megacity of southwest China: source

apportionment and implication, *Atmos. Chem. Phys.*, c14, 8679–8699, <https://doi.org/10.5194/acp-14-8679-2014>, 2014.

Tian M., Liu Y., Yang F. M., Zhang L. M., Peng C., Chen Y., Shi G. M., Wang H. B., Luo B., Jiang C. T., Li B., Takeda N., and Koizumi K.: Increasing importance of nitrate formation for heavy aerosol pollution in two megacities in Sichuan Basin, southwest China, *Environ. Pollut.*, 250, 898–905, <https://doi.org/10.1016/j.envpol.2019.04.098>, 2019.

Wang, G. H., Zhang, R. Y., Gomez, M. E., Yang, L. X., Zamora, M. L., Hu, M., Lin, Y., Peng, J. F., Guo, S., Meng, J. J., Li, J. J., Cheng, C. L., Hu T. F., Ren, Y. Q., Wang, Y. S., Gao, J., Cao, J. J., An, Z. S., Zhou, W. J., Li, G. H., Wang, J. Y., Tian, P. F., Marrero-Ortiz, W., Secrest, J., Du, Z. F., Zheng, J., Shang, D. J., Zeng, L. M., Shao, M., Wang, W. G., Huang, Y., Wang, Y., Zhu, Y. J., Li, Y. X., Hu, J. X., Pan, B., Cai, L., Cheng, Y. T., Ji, Y. M., Zhang, F., Rosenfeld, D., Liss, P. S., Duce, R. A., Kolb, C. E., and Molina, M. J.: Persistent sulfate formation from London Fog to Chinese haze, *Proc. Natl. Acad. Sci. USA*, 113, 13630–13635, <https://doi.org/10.1073/pnas.1616540113>, 2016.

Wang, H. B., Tian, M., Chen, Y., Shi, G. M., Liu, Y., Yang, F. M., Zhang, L. M., Deng, L. Q., Yu, J. Y., Peng, C., and Cao, X. Y.: Seasonal characteristics, formation mechanisms and source origins of PM_{2.5} in two megacities in Sichuan Basin, China, *Atmos. Chem. Phys.*, 18, 865–881, <https://doi.org/10.5194/acp-18-865-2018>, 2018.

Wang, H. B., Zhang, L. M., Huo, T. T., Wang, B., Yang, F. M., Chen, Y., Tian, M., Qiao, B. Q., and Peng, C.: Application of parallel factor analysis model to decompose excitation-emission matrix fluorescence spectra for characterizing sources of water-soluble brown carbon in PM_{2.5}, *Atmos. Environ.*, 223, 117192, <https://doi.org/10.1016/j.atmosenv.2019.117192>, 2019a.

Wang, J. F., Ye, J. H., Zhang, Q., Zhao, J., Wu, Y. Z., Li, J. Y., Liu, D. T., Li, W. J., Zhang, Y. G., Wu, C., Xie, C. H., Qin, Y. M., Lei, Y. L., Huang, X. P., Guo, J. P., Liu, P. F., Fu, P. Q., Li, Y. J., Lee, H. C., Choi, H., Zhang, J., Liao, H., Chen, M. D., Sun, Y. L., Ge, X. L., Martin, S. T., and Jacob, D. J.: Aqueous production of secondary organic aerosol from fossil-fuel emissions in winter Beijing haze, *Proc.*

- Natl. Acad. Sci. USA, 118, e2022179118, <https://doi.org/10.1073/pnas.2022179118>, 2021.
- Wang, Q. Y., Ye, J. H., Wang, Y. C., Zhang, T., Ran, W. K., Wu, Y. F., Tian, J., Li, L., Zhou, Y. Q., Ho, H. S. S., Dang, B., Zhang, Q., Zhang, R. J., Chen, Y., Zhu, C. S., and Cao, J. J.: Wintertime Optical Properties of Primary and Secondary Brown Carbon at a Regional Site in the North China Plain, *Environ. Sci. Technol.*, 53, 12389–12397, <https://doi.org/10.1021/acs.est.9b03406>, 2019b.
- Washenfelder, R. A., Attwood, A. R., Brock, C. A., Guo, H., Xu, L., Weber, R. J., Ng, N. L., Allen, H. M., Ayres, B. R., Baumann, K., Cohen, R. C., Draper, D. C., Duffey, K. C., Edgerton, E., Fry, J. L., Hu, W. W., Jimenez, J. L., Palm, B. B., Romer, P., Stone, E. A., Wooldridge, P. J., and Brown, S. S.: Biomass burning dominates brown carbon absorption in the rural southeastern United States, *Geophys. Res. Lett.*, 42, 653–664, <https://doi.org/10.1002/2014gl062444>, 2015.
- Wu, C. and Yu, J. Z.: Determination of primary combustion source organic carbon-to-elemental carbon (OC/EC) ratio using ambient OC and EC measurements: Secondary OC-EC correlation minimization method, *Atmos. Chem. Phys.*, 16, 5453–5465, <https://doi.org/10.5194/acp-16-5453-2016>, 2016.
- Wu, H., Peng, C., Zhai, T. Y., Deng, J. C., Lu, P. L., Li, Z. L., Chen, Y., Tian, M., Bao, Z. E., Long, X., Yang, F. M., and Zhai, C. Z.: Characteristics of light absorption and environmental effects of Brown carbon aerosol in Chongqing during summer and winter based on online measurement: Implications of secondary formation, *Atmos. Environ.*, 338, 120843, <https://doi.org/10.1016/j.atmosenv.2024.120843>, 2024.
- Xie, C. H., Xu, W. Q., Wang, J. F., Wang, Q. Q., Liu, D. T., Tang, G. Q., Chen, P., Du, W., Zhao, J., Zhang, Y. J., Zhou, W., Han, T. T., Bian, Q. Y., Li, J., Fu, P. Q., Wang, Z. F., Ge, X. L., Allan, J., Coe, H., and Sun, Y. L.: Vertical characterization of aerosol optical properties and brown carbon in winter in urban Beijing, China, *Atmos. Chem. Phys.*, 19, 165–179, <https://doi.org/10.5194/acp-19-165-2019>, 2019.
- Xu, B. Q., Zhang, G., Gustafsson, Ö., Kawamura, K., Li, J., Andersson, A., Bikkina, S., Kunwar, B., Pokhrel, A., Zhong, G. C., Zhao, S. Z., Li, J., Huang, C., Cheng, Z.

- N., Zhu, S. Y., Peng, P. A., and Sheng, G. Y.: Large contribution of fossil-derived components to aqueous secondary organic aerosols in China, *Nat. Commun.*, 13, 5115, <https://doi.org/10.1038/s41467-022-32863-3>, 2022.
- Xu, W. Q., Han, T. T., Du, W., Wang, Q. Q., Chen, C., Zhao, J., Zhang, Y. J., Li, J., Fu, P. Q., Wang, Z. F., Worsnop, D. R., and Sun, Y. L.: Effects of Aqueous-Phase and Photochemical Processing on Secondary Organic Aerosol Formation and Evolution in Beijing, China, *Environ. Sci. Technol.*, 51, 762–770, <https://doi.org/10.1021/acs.est.6b04498>, 2017.
- Xu, W. Q., Sun, Y. L., Chen, C., Du, W., Han, T. T., Wang, Q. Q., Fu, P. Q., Wang, Z. F., Zhao, X. J., Zhou, L. B., Ji, D. S., Wang, P. C., and Worsnop, D. R.: Aerosol composition, oxidation properties, and sources in Beijing: results from the 2014 Asia-Pacific Economic Cooperation summit study, *Atmos. Chem. Phys.*, 15, 13681–13698, <https://doi.org/10.5194/acp-15-13681-2015>, 2015.
- Xu, W. Q., Sun, Y. L., Wang, Q. Q., Zhao, J., Wang, J. F., Ge, X. L., Xie, C. J., Zhou, W., Du, W., Li, J., Fu, P. Q., Wang, Z. F., Worsnop, D. R., and Coe, H.: Changes in Aerosol Chemistry From 2014 to 2016 in Winter in Beijing: Insights From High-Resolution Aerosol Mass Spectrometry, *J. Geophys. Res.: Atmos.*, 124, 1132–1147, <https://doi.org/10.1029/2018JD029245>, 2019.
- Yan, C. Q., Zheng, M., Bosch, C., Andersson, A., Desyaterik, Y., Sullivan, A. P., Collett, J. L., Zhao, B., Wang, S. X., He, K. B., and Gustafsson, Ö.: Important fossil source contribution to brown carbon in Beijing during winter, *Sci. Rep.*, 7, 43182, <https://doi.org/10.1038/srep43182>, 2017.
- Yang, J. W., Au, W. C., Law, H., Lam, C. H., and Nah, T.: Formation and evolution of brown carbon during aqueous-phase nitrate-mediated photooxidation of guaiacol and 5-nitroguaiacol, *Atmos. Environ.*, 254, 118401, <https://doi.org/10.1016/j.atmosenv.2021.118401>, 2021.
- Yang, F. M., Tan, J., Zhao, Q., Du, Z., He, K. B., Ma, Y. L., Duan, F. K., Chen, G., and Zhao, Q.: Characteristics of PM_{2.5} speciation in representative megacities and across China, *Atmos. Chem. Phys.*, 11, 5207–5219, <https://doi.org/10.5194/acp-11-5207-2011>, 2011.

- Ye, Z. L., Qu, Z. X., Ma, S. S., Luo, S. P., Chen, Y. T., Chen, H., Chen, Y. F., Zhao, Z. Z., Chen, M. D., and Ge, X. L.: A comprehensive investigation of aqueous-phase photochemical oxidation of 4-ethylphenol, *Sci. Total Environ.*, 685, 976–985, <https://doi.org/10.1016/j.scitotenv.2019.06.276>, 2019.
- Yokelson, R. J., Urbanski, S. P., Atlas, E. L., Toohey, D. W., Alvarado, E. C., Crounse, J. D., Wennberg, P. O., Fisher, M. E., Wold, C. E., Campos, T. L., Adachi, K., Buseck, P. R., and Hao, W. M.: Emissions from forest fires near Mexico City, *Atmos. Chem. Phys.*, 7, 5569–5584, <https://doi.org/10.5194/acp-7-5569-2007>, 2007.
- Yu, L., Smith, J., Laskin, A., Anastasio, C., Laskin, J., Zhang, Q.: Chemical characterization of SOA formed from aqueous-phase reactions of phenols with the triplet excited state of carbonyl and hydroxyl radical, *Atmos. Chem. Phys.*, 14, 13801–13816, <https://doi.org/10.5194/acp-14-13801-2014>, 2014.
- Yu, L., Smith, J., Laskin, A., George, K. M., Anastasio, C., Laskin, J., Dillner, A. M., and Zhang, Q.: Molecular transformations of phenolic SOA during photochemical aging in the aqueous phase: competition among oligomerization, functionalization, and fragmentation, *Atmos. Chem. Phys.*, 16, 4511–4527, <https://doi.org/10.5194/acp-16-4511-2016>, 2016.
- Zhang, X. Y., Liu, H. X., Cheng, J. L., Song, W., Wang, H. C., Zhang, Y. L., and Wang, X. M.: Gas-Phase Oxidation of Guaiacol by NO₃ Radicals: Kinetic Measurements and Implications, *ACS ES&T Air*, 2, 903–910, <https://doi.org/10.1021/acsestair.4c00353>, 2025.
- Zhao, J., Du, W., Zhang, Y. J., Wang, Q. Q., Chen, C., Xu, W. Q., Han, T. T., Wang, Y. Y., Fu, P. Q., Wang, Z. F., Li, Z. Q., and Sun, Y. L.: Insights into aerosol chemistry during the 2015 China Victory Day parade: results from simultaneous measurements at ground level and 260 m in Beijing, *Atmos. Chem. Phys.*, 17, 3215–3232, <https://doi.org/10.5194/acp-17-3215-2017>, 2017.
- Zhao, J., Qiu, Y. M., Zhou, W., Xu, W. Q., Wang, J. F., Zhang, Y. J., Li, L. J., Xie, C. H., Wang, Q. Q., Du, W., Worsnop, D. R., Canagaratna, M. R., Zhou, L. B., Ge, X. L., Fu, P. Q., Li, J., Wang, Z. F., Donahue, N. M., and Sun, Y. L.: Organic Aerosol

1279 Processing During Winter Severe Haze Episodes in Beijing, *J. Geophys. Res.:*
 1280 *Atmos.*, 124, 10248–10263, <https://doi.org/10.1029/2019JD030832>, 2019.

1281 Zhao, R., Lee, A. K. Y., Huang, L., Li, X., Yang, F., and Abbatt, J. P. D.:
 1282 Photochemical processing of aqueous atmospheric brown carbon, *Atmos. Chem.*
 1283 *Phys.*, 15, 6087–6100, <https://doi.org/10.5194/acp-15-6087-2015>, 2015.

1284 Zhao, R., Mungall, E. L., Lee, A. K. Y., Aljawhary, D., and Abbatt, J. P. D.:
 1285 Aqueous-phase photooxidation of levoglucosan – a mechanistic study using
 1286 aerosol time-of-flight chemical ionization mass spectrometry (Aerosol ToF-CIMS),
 1287 *Atmos. Chem. Phys.*, 14, 9695–9706, <https://doi.org/10.5194/acp-14-9695-2014>,
 1288 2014.

1289 Zhong, H. B., Huang, R. J., Chang, Y. H., Duan, J., Lin, C. S., and Chen, Y.:
 1290 Enhanced formation of secondary organic aerosol from photochemical oxidation
 1291 during the COVID-19 lockdown in a background site in Northwest China, *Sci.*
 1292 *Total Environ.*, 778, 144947, <https://doi.org/10.1016/j.scitotenv.2021.144947>,
 1293 2021.

1294 Zhu, C. S., Cao, J. J., Hu, T. F., Shen, Z. X., Tie, X. X., Huang, H., Wang, Q. Y.,
 1295 Huang, R. J., Zhao, Z. Z., Mocnik, G., and Hansen, A. D. A.: Spectral dependence
 1296 of aerosol light absorption at an urban and a remote site over the Tibetan Plateau,
 1297 *Sci. Total Environ.*, 590–591, 14–21,
 1298 <https://doi.org/10.1016/j.scitotenv.2017.03.057>, 2017.

1299 Zhu, L. W., Cui, Y. J., Ge, X. L., Ye, Z. L., and Zhao, Z. Z.: Aqueous-phase
 1300 photochemical oxidation of extracted WSOC in PM_{2.5} from biomass burning,
 1301 *China Environ. Sci.*, 43, 1014–1025,
 1302 <https://doi.org/10.19674/j.cnki.issn1000-6923.20220928.004>, 2023.