



Species-specific relationships between net primary productivity and forest age for subtropical China

Peng Li¹, Rong Shang^{1,2*}, Jing M. Chen^{1,3*}, Huiguang Zhang⁴, Xiaoping Zhang⁴, Guoshuai Zhao⁴, Hong Yan⁴, Jun Xiao⁴, Xudong Lin¹, Lingyun Fan¹, Rong Wang¹, Jianjie Cao¹, and Hongda Zeng¹

¹ Key Laboratory for Humid Subtropical Eco-Geographical Processes of the Ministry of Education, School of Geographical Sciences, Fujian Normal University, Fuzhou, 350007, China

² Academy of Carbon Neutrality, Fujian Normal University, Fuzhou 350007, China

³ Department of Geography and Planning, University of Toronto, Ontario, ON M5S 3G3, Canada

⁴ Fujian Forestry Survey and Planning Institute, Fuzhou 350003, China

Correspondence to: Rong Shang (shangrong@fjnu.edu.cn) and Jing M. Chen (jing.chen@utoronto.ca)

Abstract. The relationship between net primary productivity (NPP) and forest age varies among forest species, yet there were no available NPP–age relationships established for various forest species in subtropical China for use in forest carbon modeling. This study explored the NPP–age relationships for seven forest species in subtropical China using data from the Strategic Priority Project of Carbon Budget (SPPCB), National Forest Inventory (NFI) Type I (NFI-I), and Type II (NFI-II) data using the Semi-Empirical Model (SEM). Forest species included *Pinus massoniana* (*P. massoniana*), *Cunninghamia lanceolata* (*C. lanceolata*), *Eucalyptus robusta* (*Eucalyptus*), Other Coniferous Forests (OCF), Softwood Broadleaf (SWB), Hardwood Broadleaf excluding *Eucalyptus* (HWB), and Mixed Forests (MF). Based on these three datasets, we were able to derive subtropical forest species-specific NPP–age relationships. Compared with China-wide NPP–age relationships previously derived from the SPPCB dataset, these species-specific relationships derived in this study resulted in improved simulations of aboveground biomass for subtropical forests using a process-based InTEC model, suggesting that these species-specific NPP–age relationships are valuable for forest carbon modeling and management in subtropical China.

1. Introduction

Forests, recognized as one of Earth's largest carbon sinks, play a crucial role in mitigating climate change and regulating the global carbon cycle (Friedlingstein et al., 2020; Liu et al., 2012; Pan et al., 2011). Through the process of photosynthesis, forests absorb atmospheric carbon dioxide and convert it into organic carbon, thereby reducing greenhouse gas concentrations in the atmosphere (Fang et al., 2001; Chapin et al., 2006). Specifically, net primary productivity (NPP) serves as a key indicator of forest's carbon sequestration capacity, directly reflecting the biomass accumulation and carbon storage ability of forest ecosystems (Zha et al., 2013; Zhao and Zhou, 2005). NPP exhibits notable variations with forest age progression (Ben et al., 2004; Wang et al., 2007, 2011). Typically, forest NPP follows a pattern of rapid growth during the early stages, peaking in the middle ages, and gradual decline at old ages (Yu et al., 2017; He et al., 2012). However, the NPP variation pattern with age



varies with forest species and climate conditions (Yu et al., 2017; Wang et al., 2018), highlighting the importance of understanding NPP–age relationships across various forest species and climate zones for accurate forest carbon modeling (Yu et al., 2017; Wang et al., 2018; Li et al., 2024b) and effective forest management and ecological restoration strategies (Luyssaert et al., 2008; Li et al., 2024a).

35 China's subtropical region, characterized by a warm and humid climate, fertile soil, and rich biodiversity, is a vital component of global forest ecosystems (Zhou et al., 2014). This region contributes approximately 67.13% of China's terrestrial ecosystem carbon sequestration (Chen et al., 2019). The subtropical region boasts a diversity of forest types, including evergreen broad-leaved forests, coniferous forests, and mixed forests, with species such as *Pinus massoniana* (*P. massoniana*),
40 *Cunninghamia lanceolata* (*C. lanceolata*), and *Eucalyptus robusta* (*Eucalyptus*) exhibiting unique ecological characteristics that influence the forest NPP–age relationships (Huang et al., 2010). In particular, evergreen broad-leaved and coniferous forests in this region play crucial roles in the forest carbon cycle. Therefore, exploring the NPP–age relationships for diverse forest species in subtropical China is essential for formulating regional forest management strategies.

Currently, two sets of national-scale forest NPP–age curves have been established for China (Li et al., 2024a; Wang et al., 2018). However, these NPP–age curves are constructed based on broad forest cover types such as coniferous forests, broad-
45 leaved forests, or mixed forests, without considering species-specific differences within the same forest type. This limits their application in simulating forest carbon sequestration at the stand-scale and species level. Preliminary research on the NPP–age relationship in subtropical China's forests has been conducted only in Zhejiang Province, distinguishing only between coniferous and broad-leaved forest types (Zheng et al., 2019). Although this study provides a preliminary understanding of the forest carbon cycle in subtropical regions, it fails to account for the diversity of forest types in subtropical areas, especially at
50 the species level, limiting its applicability to national-scale subtropical forest carbon sequestration simulations.

Moreover, constructing species-specific NPP–age curves faces inherent limitations due to insufficient field measurements or survey sample data. For example, field survey samples from the Strategic Priority Project of Carbon Budget (SPPCB) (Fang et al., 2018) and China's National Forest Inventory (NFI) Type I (NFI-I) sample data (Lin et al., 2023) may not ensure an even
55 distribution across different forest age classes. The lack of samples from old-age forest classes hinders the accurate depiction of NPP changes with forest age, leading to biases in carbon sequestration simulations for these classes. The NFI Type II (NFI-II) stand data provides comprehensive coverage of all stands, complementing the insufficient representativeness of sample data, especially for old-age forest samples (Lin et al., 2023). However, NFI-II data typically represent the average conditions of the entire stand or dominant species, which may introduce biases in heterogeneous stands (Lin et al., 2023). Therefore, it is
60 crucial to consider the representativeness of samples and the average values of NFI-II stand data, comprehensively assessing the impact of integrating NFI-II stand data on the construction of NPP–age curves to determine the optimal approach for constructing final curves.

This study aims to explore forest NPP–age relationships for different forest species in subtropical China, with three objectives: (1) to comprehensively assess the impact of integrating NFI-II stand data on the construction of NPP–age curves; (2) to explore the NPP–age relationships of diverse forest species in subtropical China; and (3) to evaluate whether the forest



65 species-specific NPP–age relationships can improve forest aboveground biomass modeling. The resulting NPP–age curves will provide scientific support for estimating forest carbon sequestration and formulating forest management strategies in subtropical China, contributing to enhanced understanding and management of forest carbon dynamics in this region.

2. Study Area, Data and Methods

2.1. Study Area

70 Fujian Province was selected as the study area (Fig. 1a) because of its highest forest coverage in China and data availability (Shang et al., 2025). It is located on the southeastern coast, ranging from 23°33'N to 28°20'N in latitude and from 115°50'E to 120°40'E in longitude. The province is predominantly mountainous, with over 80% of its terrain comprising hills and mountains, ranging in elevation from approximately 1500 meters in the northwest to around 500 meters in the southeast. Fujian experiences a subtropical monsoon climate, characterized by mean annual temperatures ranging from 17°C to 21°C and annual precipitation between 1400 mm and 2000 mm. Fig. 1a shows the spatial distribution of the merged tree species. Three key tree species were selected for analyzing forest NPP in relation to age: *Pinus massoniana* (*P. massoniana*, 27.54% of the total studied forest species), *Cunninghamia lanceolata* (*C. lanceolata*, 23.35%), and *Eucalyptus robusta* Smith (shorten to *Eucalyptus*, 4.14%). *P. massoniana* and *C. lanceolata* were chosen for their extensive distribution within Fujian Province (Lin et al., 2023). *Eucalyptus*, although representing a smaller proportion of the forest, was included due to its artificial continuity, rapid growth, high yield, and economic value (Zhou and Wingfield, 2011). The remaining tree species were merged into four groups: Hardwood Broadleaf excluding *Eucalyptus* (HWB, 26.45%), Softwood Broadleaf (SWB, 2.01%), Other Coniferous Forests except for *P. massoniana* and *C. lanceolata* (OCF, 2.44%), and Mixed Forests (MF, 14.07%). Bamboo species were not discussed in this study.




Figure 1: The distribution of forest species in Fujian Province (a) and the distribution of NFI-I, NFI-II, and SPPCB field survey samples (b). Different colours indicate different forest species, and the grey colour is for bamboo. *P. massoniana*: *Pinus massoniana*, *C. lanceolata*: *Cunninghamia lanceolata*, *Eucalyptus*: *Eucalyptus robusta* smith, HWB: Hardwood Broadleaf excluding *Eucalyptus*, SWB: Softwood Broadleaf, OCF: Other Coniferous Forests excluding *P. massoniana* and *C. lanceolata*, MF: Mixed Forests.


2.2. Data

Forest field data from China's National Forest Inventory (NFI), comprising Type I point data (NFI-I) and Type II polygon data (NFI-II), along with SPPCB field survey samples (Fang et al., 2018), were used for building the forest NPP–age relationships. Fig. 1b shows the spatial distribution of the NFI-I, NFI-II, and SPPCB forest field samples, represented by different colors. The SPPCB field survey samples have previously been effectively used for constructing ten forest NPP–age relationships across China (Li et al., 2024a; Shang et al., 2023) and we only selected the 128 samples located in Fujian for the analysis. NFI-I samples were obtained from China's 8th (2009-2013) and 9th (2014-2018) National Forest Inventories. Each NFI-I sample records various attributes, including survey time and location, dominant tree species, forest height, diameter at breast height (DBH), forest stock volume, average forest age, and so on. After screening for different forest species, a total of 2,746 samples were retained for each period.

Given the limited availability of NFI-I and SPPCB samples, these data might be insufficient to effectively constrain the NPP–age curve in older forest age ranges. Consequently, we incorporated NFI-II polygons into our analysis. These polygons were rasterized into 30 m spatial resolution pixels using the nearest neighbor resampling method, and all pixels within a forest



105 polygon shared the same attributes (Lin et al., 2023). The NFI-II samples were then created based on the dominance of tree species, requiring a proportion of 100% (with an adjustment to “higher than 80%” for *C. equisetifolia* and SWB due to their sample sizes) and the availability of relevant attribute records necessary for establishing forest NPP–age relationships (Lin et al., 2023). To ensure sample homogeneity and confirm that each sample is positioned at the center of the forest polygon, all adjacent 11×11 pixels were required to meet both criteria (Lin et al., 2023). Finally, NFI-II samples with ages greater than those in the NFI-I and SPPCB datasets (Fig. 1b and Fig. 2) were used to construct the forest NPP–age relationships.



110 **Figure 2: Age distributions of tree species and species groups based on original NFI-I, NFI-II data, and SPPCB field survey samples in Fujian Province.** *P. massoniana*: *Pinus massoniana*, *C. lanceolata*: *Cunninghamia lanceolata*, *Eucalyptus*: *Eucalyptus robusta* smith, *HWB*: Hardwood Broadleaf excluding *Eucalyptus*, *SWB*: Softwood Broadleaf, *OCF*: Other Coniferous Forests except for *P. massoniana* and *C. lanceolata*, *MF*: Mixed Forest.



115 2.3. Methods

2.3.1. Building NPP–age relationships for different forest species

The forest NPP was calculated from the three types of forest field samples, and it consisted of four components: total biomass increment, mortality, foliage turnovers, and fine root turnovers in the soil (Chen et al., 2002; He et al., 2012; Xia et al., 2019; Li et al., 2024a):

$$120 \quad NPP = dB_c + M + L_f + L_{fr} \quad (1)$$

where dB_c is the annual increment of total living biomass (including stems, branches, and coarse roots); M is mortality ignored in this study due to a lack of observations at the ground plots and its small proportion to NPP (Li et al., 2024a); L_l is the turnover of leaves per year; and L_{fr} is the turnover of fine roots per year in the soil.

125 The annual increment of total living biomass was calculated from the annual biomass change (dB) and the ratio of carbon content (Li et al., 2011; White et al., 2000; Wu et al., 2016; Xia et al., 2019):

$$dB_c = dB \times c \quad (2)$$

where dB is the annual biomass change and c is the carbon content in biomass. Biomass was not directly provided in the NFI-I and NFI-II samples, but it could be calculated from the forest volume (V) using species-specific biomass regression equations. The coefficients for these regression equations are presented in Table 1 (Li et al., 2011; Wu et al., 2016).

130 The turnovers of leaves and fine roots per year in the soil could be calculated as follows (Chen et al., 2002; He et al., 2012; Li et al., 2024a):

$$L_l = \frac{LAI}{SLA} \times t_l \times c \quad (3)$$

$$L_{fr} = R_{fr,l} \times L_l \quad (4)$$

135 where LAI is the leaf area index (LAI), SLA is the specific leaf area, t_l is the foliage turnover ratio, and $R_{fr,l}$ represents the ratio of carbon allocated to new fine roots to carbon in new leaves. The detailed values for the coefficients of SLA , t_l , and $R_{fr,l}$ for different forest species were provided in Table 2 (Li et al., 2024a; Li et al., 2007; White et al., 2000; Xie et al., 2022; Zhou et al., 2008). The LAI values were derived from the GLOBMAP Version 3 LAI product (Liu et al., 2012).

140 The semi-empirical mathematical (SEM) function (Chen et al., 2003; He et al., 2012; Li et al., 2024a) was used to build the forest NPP–age relationships for different forest species based on the calculated forest field NPP, as it was demonstrated as the optimal method for building NPP–age curves in China (Li et al., 2024a):

$$NPP(x) = a[1 + (b(x/c)^d - 1)/e^{(x/c)}] \quad (5)$$

where $NPP(x)$ is NPP at the age of x , and a , b , c , and d are the coefficients of the SEM function. The uncertainty analysis of the built NPP–age relationships was conducted using the same method in the research of Li et al., (2024a).

Table 1: The coefficients of the species-specific biomass regression equations (Li et al., 2011; Wu et al., 2016). B: Biomass; V: Volume. P. massoniana: Pinus massoniana, C. lanceolata: Cunninghamia lanceolata, Eucalyptus: Eucalyptus robusta smith, HWB: Hardwood Broadleaf excluding Eucalyptus, SWB: Softwood Broadleaf, OCF: Other Coniferous Forests except for P. massoniana and C. lanceolata, MF: Mixed Forests.

Forest species	Biomass regression equation ($t \cdot \text{hm}^{-2}$)	Carbon content (%)
C. lanceolata	$B = 0.3999V + 22.5410$	51.27
P. massoniana	$B = 0.52V$	52.71
OCF	$B = 0.4631V + 24.2777$	51.68
Eucalyptus	$B = 0.7893V + 6.9306$	47.48
HWB	$B = 0.6255V + 91.0013$	49.01
SWB	$B = 0.4754V + 30.6034$	45.02
MF	$B = 0.8019V + 12.2799$	48.93

Table 2: The input coefficients in the calculation of forest field NPP for different forest species. SLA is the specific leaf area; t_l is the foliage turnover ratio; $R_{fr,l}$ is the ratio of NPP to fine roots and leaves. P. massoniana: Pinus massoniana, C. lanceolata: Cunninghamia lanceolata, Eucalyptus: Eucalyptus robusta smith, HWB: Other hardwood broadleaf excluding Eucalyptus, SWB: Softwood broadleaf, OCF: Other coniferous mixed forests excluding P. massoniana and C. lanceolata, MF: Mixed forest.

Forest Type	SLA ($\text{m}^2 \text{ kg C}^{-1}$)	t_l (year^{-1})	$R_{fr,l}$ (kg C kg C^{-1})
C. lanceolata	7.9	0.22	1.4
P. massoniana	6.7	0.26	1.4
OCF	8.2	0.26	1.4
Eucalyptus	26.3	0.86	1.2
HWB/SWB	32	0.86	1.2
MF	21.1	0.56	1.3

2.3.2. Forest carbon modeling using the newly built NPP–age relationships

The NPP–age relationships constructed for different forest species were integrated into the Integrated Terrestrial Ecosystem Carbon Cycle (InTEC) model for forest carbon modeling. To evaluate whether the forest species-specific NPP–age relationships can improve forest carbon modeling, the forest carbon modeling using the newly built NPP–age relationships was compared with that of using the China-wide NPP–age relationships (Shang et al., 2023; Li et al., 2024a). The InTEC model integrates multiple processes, including leaf photosynthesis (using the Farquhar biochemical model), soil carbon and nitrogen cycling, net nitrogen mineralization, and NPP–age relationships (Chen et al., 2000a, b). This model estimates forest carbon balance by accounting for atmospheric, climatic, and biological changes since the pre-industrial era. The impact of climate change on photosynthesis is modeled through changes in the growing season length and photosynthetic rate, while elevated CO_2 concentrations and leaf nitrogen content positively affect photosynthesis. Model inputs include spatially distributed data on climate, soil texture, nitrogen deposition, and vegetation parameters derived from remote sensing (Table



165

3). Since carbon flux measurements were not available in Fujian province, the performance of forest carbon modeling was indirectly validated by comparing the modeled aboveground biomass with the calculated aboveground biomass from forest field surveys or inventory data. The aboveground biomass was modeled at a spatial resolution of 30 meters from 1901 to 2023, and the period from 1901 to 1985 was used for the spin-up of the initial model parameters.

Table 3: Main Input Data of the InTEC Model. LAI: Leaf area index; BEPS: Boreal Ecosystem Productivity Simulator model; DEM: Digital elevation model.

Input data		Unit	Spatial resolution	Temporal resolution	Data source
Climate data	Precipitation	mm	0.5°	1901-2023	CRU TS 4.08
	Temperature	°C			
	Vapor pressure	hpa			
	Cloud amount	%			
Atmospheric composition data	CO ₂ concentration	mol mol ⁻¹	Site scale	1960-2021	Mauna Loa
	Nitrogen deposition	10*gN m ⁻² yr ⁻¹	1.27°×2.5°	1997-2013	(Gao et al., 2020)
Vegetation data	Forest cover types	/	30m	/	NFI-II
	LAI	m ² /m ²	500m	2015	GLOBMAP LAI V3
	Forest age	year	30m	2015	NFI-II
	Reference NPP	10 gC m ⁻² yr ⁻¹	30m	2015	BEPS
	NPP-age relationship curves	/	/	/	This study
Soil data	Sand content	%	0.0083°	/	HDSW World Soil Database
	Clay content	%	0.0083°	/	
	Soil depth	100 m	0.0083°	/	
Topographic data	Latitude/longitude	degree	30m	/	/
	DEM	m	30m	/	http://www.gsc loud.cn
	Slope and aspect	/	30m	/	Calculated from DEM
	Topographic wetness index	/	30m	/	
	Water table depth	m	30m	/	

3. Results

170

3.1. Comparisons of forest NPP–age relationships constructed with and without NFI-II samples

The NPP–age relationships constructed with and without NFI-II samples using the SEM function were compared in Fig. 3. This comparison was motivated by two considerations: (i) NFI-I and SPPCB samples alone may not provide sufficient data to



reliably constrain NPP–age curves for old forests and (ii) NFI-II samples, derived from NFI-II polygons, may introduce inherent uncertainties. In Fig. 3, the green and red lines depict the forest NPP–age curves with and without using NFI-II samples, respectively. Solid lines indicate the forest age ranges where field data are available, while dashed lines represent extrapolated curves beyond the field sample age range. Their curve-fitting performances were also quantitatively assessed using R^2 and RMSE, as shown in Fig. 4.

The peak NPP age, a critical indicator of the NPP–age relationship, remained consistent for *Eucalyptus*, *P. massoniana*, *C. lanceolata*, OCF, and MF, regardless of whether NFI-II samples were included, while only a one-year difference was observed for HWB and SWB. Notably, without using NFI-II samples, NPP values for *C. lanceolata* and SWB dropped close to zero in ages older than 150 years, suggesting a transition from forest carbon sinks to carbon sources in old ages. This finding contrasts with previous studies, which suggest that older forests continue to act as carbon sinks (Gundersen et al., 2021; Luyssaert et al., 2008). For *Eucalyptus*, the NPP reduction exceeded 70% without NFI-II samples, diverging significantly from previous studies that generally report reductions by about one-third (Luyssaert et al., 2008; Wang et al., 2011) or half (Ryan et al., 2004; Mund et al., 2002) of peak NPP. These discrepancies highlight the importance of including NFI-II samples for accurately modeling the NPP–age relationships for *Eucalyptus*, *C. lanceolata*, and SWB.

For *P. massoniana*, HWB, OCF, and MF, the inclusion of NFI-II samples had minimal effects on the overall pattern of the NPP–age curves. When NFI-II samples were included, R^2 slightly decreased by less than 0.025 for *P. massoniana*, HWB, and OCF, but slightly increased by 0.003 for MF. Similarly, RMSE values showed a minor increase (under $2 \text{ gC m}^{-2} \text{ year}^{-1}$) for *P. massoniana* and OCF, while a slight decrease (under $7 \text{ gC m}^{-2} \text{ year}^{-1}$) for HWB and MF. Therefore, we ultimately opted to consistently use NFI-II samples in constructing the NPP–age curves, as incorporating NFI-II samples can extend the age range over which the curves are constrained, thus enhancing the data coverage and consistency.




Figure 3: NPP–age curves fitted by the SEM function for different forest species with and without using NFI-II samples. The red and blue circles, with associated grey error bars, represent the average NPP values and their one standard deviation. The green and red lines depict the built forest NPP–age curves with and without using NFI-II samples. Solid lines indicate the forest age ranges where field data are available, while dashed lines represent extrapolated curves beyond the maximum age of the field samples. *P. massoniana*: *Pinus massoniana*, *C. lanceolata*: *Cunninghamia lanceolata*, *Eucalyptus*: *Eucalyptus robusta smith*, *HWB*: Hardwood Broadleaf excluding *Eucalyptus*, *SWB*: Softwood Broadleaf, *OCF*: Other Coniferous Forests except for *P. massoniana* and *C. lanceolata*, *MF*: Mixed Forest.




Figure 4: R² and RMSE of the built NPP–age curves for different forest species with and without NFI-II samples. P. massoniana: Pinus massoniana, C. lanceolata: Cunninghamia lanceolata, Eucalyptus: Eucalyptus robusta smith, HWB: Hardwood Broadleaf excluding Eucalyptus, SWB: Softwood Broadleaf, OCF: Other Coniferous Forests except for P. massoniana and C. lanceolata, MF: Mixed Forest.

3.2. Characterization of forest NPP–age curves among different forest species

The final forest NPP–age curves were built with the SEM function using the SPPCB, NFI-I, and NFI-II samples for different forest species, with model coefficients provided in Table 4. Fig. 5 shows the standardized NPP–age curves of the seven forest species and species groups. Solid lines indicate the age range supported by field data (the triangle in each line indicates the maximum age), while dashed lines indicate predicted values beyond this range using the SEM function. The NPP–age patterns were generally consistent across all species, with NPP increasing during young stages, peaking in a middle age, and then declining and stabilizing in old ages (Li et al., 2024a; He et al., 2012; Yu et al., 2017; Zheng et al., 2019; Wang et al., 2018, 2011). But there were also variations in the timing of peak NPP, as well as differences in the rate of decline in older age stages. Specifically, the peak NPP ages for Eucalyptus, P. massoniana, C. lanceolata, SWB, HWB, OCF, and MF were identified as 9, 32, 25, 22, 37, 24, and 30 years, respectively. The ratios of stabilized NPP in old ages to the maximum NPP (stabilized-to-peak NPP ratios) were 59.1%, 57.3%, 65.5%, 56.8%, 59.5%, 56.0%, and 57.9%, respectively. These values align with previous studies, which typically report reductions by approximately one-third (Luyssaert et al., 2008; Wang et al., 2011) or half (Ryan et al., 2004; Mund et al., 2002) from the peak NPP.

Broadleaf species such as HWB and SWB demonstrated later peak NPP ages and lower stabilized-to-peak NPP ratios compared to conifer species like C. lanceolata, P. massoniana, and OCF. The higher wood density and longer lifespans of broadleaf species allow them to sustain productivity and carbon absorption over an extended period (Xu et al., 2024), while conifer species, despite their rapid early growth and carbon fixation, show earlier and steeper declines, reflecting differences in their ecological and physiological strategies (Bigler and Veblen, 2009). HWB exhibited a later peak NPP age and lower



225 stabilized-to-peak NPP ratio compared to SWB. This can be explained by the fact that hardwood species maintain stronger carbon absorption during later growth stages due to their higher wood density and longer lifespans (Citations). In contrast, softwood species excel in rapid carbon sequestration during early stages but experience earlier and more significant productivity declines (Citations). Compared to *C. lanceolata*, *P. massoniana* shows a later peak NPP age and higher stabilized-to-peak NPP ratio, as *P. massoniana* supports prolonged carbon sequestration (Justine et al., 2017; Bai and Ding, 2024), while *C. lanceolata* prioritizes rapid early growth (Zhou et al., 2016).

230 The NPP–age curve for Eucalyptus forests exhibits a relatively low peak NPP age of 9 years compared to other forest species. This early peak age underscores the species' ability to achieve significant productivity at a young age, making it well-suited for fast-growing timber plantations (Zhang et al., 2023; Qin and Shangguan, 2019). While this early productivity surge is advantageous for short-rotation forestry, it often leads to a shortened early life cycle, causing a noticeable reduction in productivity soon after reaching peak levels (Zhang et al., 2023; Zhou and Wingfield, 2011).

235 **Table 4: Coefficients of Fujian Province's plantation NPP–age curves were computed using the SEM function.** a-d: the model coefficients. *P. massoniana*: *Pinus massoniana*, *C. lanceolata*: *Cunninghamia lanceolata*, *Eucalyptus*: *Eucalyptus robusta* smith, HWB: Hardwood Broadleaf excluding *Eucalyptus*, SWB: Softwood Broadleaf, OCF: Other Coniferous Forests except for *P. massoniana* and *C. lanceolata*, MF: Mixed Forest.

Tree species	Parameters			
	<i>a</i>	<i>b</i>	<i>c</i>	<i>d</i>
<i>C. lanceolata</i>	470.8	1.894	13.78	1.353
<i>P. massoniana</i>	415.3	0.132	7.61	4.136
OCF	422	1.325	9.697	2.267
<i>Eucalyptus</i>	484	1.225	3.664	2.226
HWB	507.2	1.132	14.38	2.316
SWB	468	0.532	6.847	3.104
MF	444.2	1.246	12.11	2.26




Figure 5: The normalized NPP–age curves built from the SEM function. The solid lines are for the age period with field data (the triangle in each line indicates the largest age with the field data), and the dashed lines are for the age period without field data. *P. massoniana*: *Pinus massoniana*, *C. lanceolata*: *Cunninghamia lanceolata*, *Eucalyptus*: *Eucalyptus robusta* smith, HWB: Hardwood Broadleaf excluding *Eucalyptus*, SWB: Softwood Broadleaf, OCF: Other Coniferous Forests except for *P. massoniana* and *C. lanceolata*, MF: Mixed Forest.

3.3. Comparison to the forest NPP–age curves built previously

The normalized NPP–age curves built for seven forest species and species groups (referred to as species-specific curves) in Fujian province were compared with three previously built NPP–age curves for entire China (shortened to as China-wide curves) (Fig. 6). In general, the species-specific NPP–age curves constructed exhibit earlier peak ages and faster decline in old ages, particularly for *Eucalyptus*, *C. lanceolata*, OCF, and SWB.

The peak NPP age for *Eucalyptus* is 9 years, much smaller than that from the China-wide NPP–age curve. But it aligns with the reported rapid growth and significant productivity of *Eucalyptus* at young ages (Zhang et al., 2023; Qin and Shangguan, 2019). The peak NPP age for *P. massoniana* in Fujian Province is 32 years, which agrees well with the 34 years of ENF in southern China (Li et al., 2024a). While *C. lanceolata* peaks at 25 years, earlier than ENF in southern China, but close to the 23 years found for coniferous forests in Zhejiang Province (Zheng et al., 2019). Other coniferous forests except for *P. massoniana* and *C. lanceolata* have peak NPP ages of 24 years, similar to coniferous forests in Zhejiang (Zheng et al., 2019). These relatively early peak ages indicate their efficient photosynthesis and resource utilization during the early growth stages (Lu et al., 2015; Huang et al., 2007).

The peak NPP age of HWB is 37 years, similar to that of EBF in southern and eastern China, which peaks at 30 to 40 years (Li et al., 2024a; Wang et al., 2011). In contrast, the NPP of SWB peaks at 22 years, which is relatively early compared to HWB. Soft broadleaf species generally prioritize rapid early growth in response to favorable environmental conditions (Fujita

et al., 2012). Among broadleaf species, Eucalyptus exhibits the earliest peak NPP of 9 years, making it highly suitable for short-rotation forestry, but sharp post-peak declines limit its long-term carbon storage potential. The peak NPP age of MF in Fujian is 30 years, which is similar to that of mixed forests in the south and southwest of China (Li et al., 2024a) and close to the 32 years of peak NPP age for MBF in central China (Wang et al., 2011). Mixed forests combine the fast growth of broadleaf species with the longevity of conifers, achieving a balance in productivity across growth stages. Their diverse composition enhances resource utilization efficiency and reduces competition, allowing for sustained and stable carbon absorption (Xu et al., 2024).




Figure 6: Comparison between the species-specific and China-wide normalized forest NPP-age curves. P. massoniana: Pinus massoniana, C. lanceolata: Cunninghamia lanceolata, Eucalyptus: Eucalyptus robusta smith, HWB: Hardwood Broadleaf excluding Eucalyptus, SWB: Softwood Broadleaf, OCF: Other Coniferous Forests except for P. massoniana and C. lanceolata, MF: Mixed Forest.



3.4. Forest biomass modeling using the species-specific NPP-age curves

The built species-specific NPP-age curves were incorporated into the InTEC model for forest biomass modeling. But due to the lack of field soil carbon data for validation, we primarily focused on validating the modeled forest above-ground wood biomass (AGB). We compared the simulated AGB obtained by using the newly constructed species-specific NPP-age curves with that obtained by using the previously built nationwide NPP-age curves (Fig.7). Overall, the species-specific NPP-age curves significantly outperformed the nationwide curves in simulating AGB accuracy.

For coniferous forests, the nationwide NPP-age curve tended to overestimate AGB for ages ranging between 40 and 120 years. In contrast, the species-specific curves declined more rapidly after the peak NPP year. This might be closely related to the mechanism through which the subtropical warm and humid environment accelerates plant physiological aging (Chen et al., 2024). When using species-specific curves, the accuracy of simulating AGB for *C. lanceolata* was slightly higher, while for both *C. lanceolata* and OCF, the accuracy was significantly improved, with an average reduction in RMSE ranging from 9.4 to 14.4 Mg/ha. For Eucalyptus and SWB in broadleaf forests, the nationwide curve overestimated AGB for trees older than their peak NPP age but underestimated it for Eucalyptus younger than 20 years. The accuracy of simulating AGB for HWB based on species-specific curves was slightly enhanced, but for Eucalyptus and SWB, it was significantly improved, with an average increase in R^2 greater than 0.3 and a decrease in RMSE exceeding 18.5 Mg/ha. Similarly, the accuracy for MF was also enhanced, with an average RMSE reduction of 7.56 Mg/ha. Overall, the larger the differences between species-specific and nationwide NPP-age relationships (Fig. 6), the larger improvements are found in simulated AGB values (Fig. 7).

These results demonstrate that the newly developed species-specific NPP-age curves significantly enhance the accuracy of AGB simulations in the InTEC model for subtropical forests, particularly for early-maturing species such as Eucalyptus, by capturing regional-specific growth strategies. Notably, the improvement in simulation accuracy varied across different age classes, highlighting the importance of considering age dynamics in forest carbon sink modeling and predictions.




Figure 7: Validation and comparison of the simulated aboveground wood biomass by using the species-specific and China-wide forest NPP–age curves. P. massoniana: Pinus massoniana, C. lanceolata: Cunninghamia lanceolata, Eucalyptus: Eucalyptus robusta smith, HWB: Hardwood Broadleaf excluding Eucalyptus, SWB: Softwood Broadleaf, OCF: Other Coniferous Forests except for P. massoniana and C. lanceolata, MF: Mixed Forest.

4. Discussions

This study established NPP–age relationships for seven forest species and species groups in Fujian Province based on field survey data from NFI-I, NFI-II, and SPPCB using the SEM model. It also evaluated whether the species-specific NPP–age



relationships can improve forest biomass modelling using the InTEC model. Since NFI-I and SPPCB samples alone do not provide adequate data for old forests, and NFI-II samples introduces inherent uncertainties, we compared NPP–age relationships constructed with and without NFI-II samples. Results showed that incorporating NFI-II samples was crucial for accurately modeling NPP–age relationships for *Eucalyptus*, *C. lanceolata*, and SWB, but had minimal impacts on *P. massoniana*, HWB, OCF, and MF. Nevertheless, NFI-II helped extend the data range to older ages. The constructed species-specific NPP–age relationships all three available datasets were shown to improve modelled biomass in subtropical China, highlighting the importance of species-specific parameterization in forest biomass modeling. The resulting NPP–age curves will provide scientific support for accurate estimation of forest carbon sequestration and the formulation of forest management strategies in subtropical China (Li et al., 2024c), contributing to enhanced understanding and management of forest carbon dynamics in this region with the largest sinks in China.

There were several limitations. Firstly, inherent inconsistencies may arise among the three field data sets, particularly notable discrepancies between the NFI-II stand data and the SPPCB and NFI-I sample data. The NFI-I and SPPCB field samples may lack sufficient representation within the old age classes of forests (Fig. 2), potentially leading to unconstrained NPP–age curves for certain tree species in old ages, which may exhibit an unreasonably declining trend and a transition from forest carbon sinks to carbon sources, i.e. NPP declines to values close to zero (Fig. 3). To strengthen the constraint on the curves for the old age classes, this study incorporated NFI-II stand data by converting stand attributes into point samples. However, for stands characterized by high heterogeneity, deviations may still occur despite efforts to mitigate this effect through screening based on the dominance of tree species (Lin et al., 2023). To visually indicate the data constraint on the constructed NPP–age curves, solid lines were used to denote the curve portions supported by field data, while dashed lines were employed for the curve portions lacking field data. In future studies, collecting more field data on old forests will facilitate determining the shape of the forest NPP–age curves at older ages.

Second, this study did not account for the difference of planted forests and natural forests on the NPP–age relationships, nor the impact of forest managements such as selective logging and shelterwood cutting. *Eucalyptus* in plantations often grows rapidly due to intensive management, but this can lead to ecosystem degradation, such as soil erosion and reduced biodiversity. In contrast, *Eucalyptus* in natural forests grows more slowly but supports a more stable ecosystem (Ying et al., 2010). *P. massoniana* and *C. lanceolata* in natural forests exhibit higher ecosystem complexity and biodiversity, which results in slower growth rates but longer growth cycles with higher NPP (Liu et al., 2014). The timing of selective cutting and shelterwood cutting also significantly affects forest growth. Properly timed logging practices can promote tree health, growth, and resource renewal, while mistimed logging can negatively impact growth rates and wood quality (Wu et al., 2018). Due to the lack of data, this study did not distinguish between planted and natural forests and did not consider the impact of forest management. Future research may be directed towards acquiring comprehensive data to better understand the growth differences between planted forests and natural forests and the influence of forest managements on forest NPP–age relationships.

Third, the input coefficients for specific leaf area, foliage turnover ratio, and the ratios of the turnovers of fine roots and leaves to NPP used in calculating forest field NPP for diverse forest species may introduce uncertainties into the forest NPP–



age relationships. Currently, these coefficients are primarily sourced from literature (Li et al., 2024a; Li et al., 2007; White et al., 2000; Xie et al., 2022; Zhou et al., 2008), with data originating from subtropical provinces in China such as Guangxi (Xie et al., 2022), Jiangxi (Li et al., 2007), and Guiyang (Zhou et al., 2011), as well as from other regions (White et al., 2000). Data from these regions may differ from those in subtropical China, potentially leading to biases in the calculation of forest field NPP and final built NPP–age curves. Therefore, future studies should also emphasize local field measurements of these input coefficients to improve the accuracy of the NPP–age relationship and more precisely ascertain the carbon sequestration capacity of different tree species at various ages.

Last, there were also other factors that could influence the forest NPP–age relationships, such as the site conditions and soil fertility (Li et al., 2024a). Under favorable site conditions, forests typically exhibit faster NPP growth during their early stages, attain higher peak NPP values, and undergo steeper declines in NPP as they age (Wang et al., 2018; Yu et al., 2017). Conversely, forests with poor soil fertility tend to exhibit slower NPP growth in their early stages, achieve lower peak NPP values, and undergo less dramatic declines in NPP as they mature. Notably, the rapid replacement of natural broadleaf forests with plantations dominated by species such as *P. massoniana* and *C. lanceolata* in subtropical regions has significantly reduced soil fertility (Ming et al., 2019; Ni et al., 2021; Li et al., 2023). Therefore, in future research endeavors, it is imperative to consider site conditions and soil fertility to improve the construction of forest NPP–age curves.

5. Conclusions

This study investigated the NPP–age relationships for seven forest species and species groups in subtropical China, leveraging the extensive datasets from the SPPCB, NFI-I, and NFI-II forest field surveys along with the SEM function. Forest species examined encompassed *P. massoniana*, *C. lanceolata*, *Eucalyptus*, OCF, SWB, HWB, and MF. Given that the NFI-I and SPPCB samples alone might not adequately represent old forests, while the NFI-II samples offer comprehensive coverage across all stands but could potentially introduce inherent uncertainties, we conducted a comparative analysis of the NPP–age curves with and without the inclusion of NFI-II samples. Results showed that incorporating NFI-II samples was crucial for accurately modeling NPP–age relationships for *Eucalyptus*, *C. lanceolata*, and SWB, but had minimal impacts on *P. massoniana*, HWB, OCF, and MF. Therefore, we incorporated NFI-II samples in constructing the species-specific NPP–age curves to enhance the data coverage and consistency. Significant differences are found between the species-specific and nationwide NPP–age relationships in both NPP peak age and the ratio of stabilized NPP at old ages to the peak NPP, suggesting dependence of the relationships on forest species and climate.

The built species-specific NPP–age curves were subsequently incorporated into the InTEC model for forest biomass modeling, and results demonstrate that the newly established species-specific curves significantly improved the accuracy of Above-Ground Biomass (AGB) simulations in the InTEC model for subtropical forests, particularly for early-maturing species such as *Eucalyptus*. Notably, the enhancement in simulation accuracy varied across different age classes, underscoring the significance of considering age dynamics in forest carbon sink modeling and predictions. These species-specific NPP–age



curves will serve as a fundamental basis for reliable forest carbon modeling and effective forest management in subtropical China.

Code availability

370 The codes for building the forest NPP–age relationships are available upon request from the corresponding authors.

Data availability

The coefficients of the built forest NPP–age relationships are available in Table 4.

Acknowledgments

375 Thanks to the two reviewers for their valuable comments and suggestions, as well as to the editors for their assistance with the manuscript.

Author contributions

Conceptualization, R.S.; Methodology, P.L., R.S. and J.M.C.; Validation, P.L.; Formal analysis, P.L., X.L and L.F.; Writing—original draft, R.S. and P.L.; Writing—review & editing, R.S. and J.M.C.; Funding acquisition, R.S.; Data curation, H. Z. (Huiguang Zhang), X. Z., G. Z., H. Y., J. X., W. R., J. C., and H. Z. (Hongda Zeng).

Funding sources

380 This research was supported by the National Natural Science Foundation of China (42471356, U23A2002, and 42101367), the Fujian Forestry Science and Technology Key Project (2022FKJ03), and the Natural Science Foundation of Fujian Province (2024J01469).

Competing interests

385 The authors declare that they have no conflict of interest.



References

- Alexandrov, G. A., Oikawa, T., and Esser, G.: Estimating terrestrial NPP: what the data say and how they may be interpreted?, *Ecol. Modell.*, 117, 361–369, [https://doi.org/10.1016/S0304-3800\(99\)00019-8](https://doi.org/10.1016/S0304-3800(99)00019-8), 1999.
- Bai, Y. and Ding, G.: Estimation of changes in carbon sequestration and its economic value with various stand density and rotation age of *Pinus massoniana* plantations in China, *Sci. Rep.*, 14, 1–10, <https://doi.org/10.1038/s41598-024-67307-z>, 2024.
- Ben, B.-L., Chuankuan, W., and Stith T., G.: Net primary production and net ecosystem production of a boreal black spruce wildfire chronosequence, *Glob. Chang. Biol.*, 10, 473–487, <https://doi.org/10.1111/j.1529-8817.2003.0742.x>, 2004.
- Bigler, C. and Veblen, T. T.: Increased early growth rates decrease longevity of conifers in subalpine forests, *Oikos*, 118, 1130–1138, <https://doi.org/10.1111/j.1600-0706.2009.17592.x>, 2009.
- Chapin, F. S., III, Woodwell, G. M., Randerson, J. T., Rastetter, E. B., Lovett, G. M., Baldocchi, D. D., Clark, D. A., Harmon, M. E., and Schimel, D. S.: Reconciling Carbon-Cycle Concepts, Terminology, and Methods, *Ecosystems*, 9, 1041–1050, <https://doi.org/10.1007/s10021-005-0105-7>, 2006.
- Chen, J. M., Ju, W., Cihlar, J., Price, D., Liu, J., Chen, W., Pan, J., Black, A., and Barr, A.: Spatial distribution of carbon sources and sinks in Canada's forests, *Tellus B Chem. Phys. Meteorol.*, 55, 622, <https://doi.org/10.3402/tellusb.v55i2.16711>, 2003.
- Chen, J. M., Ju, W., Ciais, P., Viovy, N., Liu, R., Liu, Y., and Lu, X.: Vegetation structural change since 1981 significantly enhanced the terrestrial carbon sink, *Nat. Commun.*, 10, 4–10, <https://doi.org/10.1038/s41467-019-12257-8>, 2019.
- Chen, W., Chen, J., and Cihlar, J.: An integrated terrestrial ecosystem carbon-budget model based on changes in disturbance, climate, and atmospheric chemistry, *Ecol. Modell.*, 135, 55–79, [https://doi.org/10.1016/S0304-3800\(00\)00371-9](https://doi.org/10.1016/S0304-3800(00)00371-9), 2000a.
- Chen, W., Chen, J., Liu, J., and Cihlar, J.: Approaches for reducing uncertainties in regional forest carbon balance, *Global Biogeochem. Cycles*, 14, 827–838, <https://doi.org/10.1029/1999GB001206>, 2000b.
- Chen, W., Chen, J. M., Price, D. T., and Cihlar, J.: Effects of stand age on net primary productivity of boreal black spruce forests in Ontario, Canada, *Can. J. For. Res.*, 32, 833–842, <https://doi.org/10.1139/x01-165>, 2002.
- Chen, X., Li, J., Peñuelas, J., Li, X., Hu, D., Wang, M., Zhong, Q., and Cheng, D.: Temperature dependence of carbon metabolism in the leaves in sun and shade in a subtropical forest, *Oecologia*, 204, 59–69, <https://doi.org/10.1007/s00442-023-05487-7>, 2024.
- Eggleston, S., Buendia, L., Miwa, K., Ngara, T., and Tanabe, K.: 2006 IPCC Guidelines for National Greenhouse Gas Inventories. V.4. Agriculture, Forestry and Other Land Use, 2006.
- Fang, J., Yu, G., Liu, L., Hu, S., and Stuart Chapin, F.: Climate change, human impacts, and carbon sequestration in China, *Proc. Natl. Acad. Sci. U. S. A.*, 115, 4015–4020, <https://doi.org/10.1073/pnas.1700304115>, 2018.
- Fang Jingyun , Ke Jinhu , Tang Zhiyao, C. E. A.: Implications and estimations of four terrestrial productivity parameters, *Acta Phytocol. Sin.*, 25, 414–419, 2001.



- Friedlingstein, P. and O'Sullivan, M. and Jones, M. W. and Andrew, R. M. and Hauck, J. and Olsen, A. and Peters, G. P. and Peters, W. and Pongratz, J. and Sitch, S. and Le Qu'ér'e, C. and Canadell, J. G. and Ciais, P. and Jackson, R. B. and Alin, S. and, S.: Global Carbon Budget 2020, *Earth Syst. Sci. Data*, 12, 3269–3340, <https://doi.org/10.5194/essd-12-3269-2020>, 2020.
- 420 Fujita, N., Noma, N., Shirakawa, H., and Kikuzawa, K.: Annual photosynthetic activities of temperate evergreen and deciduous broadleaf tree species with simultaneous and successive leaf emergence in response to altitudinal air temperature, *Ecol. Res.*, 27, 1027–1039, <https://doi.org/10.1007/s11284-012-0983-z>, 2012.
- Gao, Y., Zhou, F., Ciais, P., Miao, C., Yang, T., Jia, Y., Zhou, X., Klaus, B. B., Yang, T., and Yu, G.: Human activities
425 aggravate nitrogen-deposition pollution to inland water over China, *Natl. Sci. Rev.*, 7, 430–440, <https://doi.org/10.1093/nsr/nwz073>, 2020.
- Gundersen, P., Thybring, E. E., Nord-Larsen, T., Vesterdal, L., Nadelhoffer, K. J., and Johannsen, V. K.: Old-growth forest carbon sinks overestimated, *Nature*, 591, E21–E23, <https://doi.org/10.1038/s41586-021-03266-z>, 2021.
- Hasenauer, S., Mag, and Sulzer, W.: the significance of remote sensing in the good practice guidance for land use, land-use
430 change and forestry as specified by the kyoto protocol diploma thesis, 2004.
- He, L., Chen, J. M., Pan, Y., Birdsey, R., and Kattge, J.: Relationships between net primary productivity and forest stand age in U.S. forests, *Global Biogeochem. Cycles*, 26, 1–19, <https://doi.org/10.1029/2010GB003942>, 2012.
- Hicke, J. A., Jenkins, J. C., and Ducey, O. M.: Spatial Patterns of Forest Characteristics in the Western United States Derived from Inventories, *Ecol. Appl.*, 17, 2387–2402, <https://doi.org/https://doi.org/10.1890/06-1951.1>, 2007.
- 435 Huang, M., Ji, J., Li, K., Liu, Y., Yang, F., and Tao, B.: The ecosystem carbon accumulation after conversion of grasslands to pine plantations in subtropical red soil of South China, *Tellus, Ser. B Chem. Phys. Meteorol.*, 59, 439–448, <https://doi.org/10.1111/j.1600-0889.2007.00280.x>, 2007.
- Huang, N., Niu, Z., Wu, C., and Tappert, M. C.: Modeling net primary production of a fast-growing forest using a light use efficiency model, *Ecol. Modell.*, 221, 2938–2948, <https://doi.org/10.1016/j.ecolmodel.2010.08.041>, 2010.
- 440 Justine, M. F., Yang, W., Wu, F., and Khan, M. N.: Dynamics of biomass and carbon sequestration across a chronosequence of masson pine plantations, *J. Geophys. Res. Biogeosciences*, 122, 578–591, <https://doi.org/10.1002/2016JG003619>, 2017.
- Li, H., Lei, Y., and Zeng, W.: Forest Carbon Storage in China Estimated Using Forestry Inventory Data (In Chinese), *Sci. silvae Sin.*, 47, 7–12, 2011.
- Li, P., Shang, R., Chen, J. M., Xu, M., Lin, X., Yu, G., He, N., and Xu, L.: Evaluation of five models for constructing forest
445 NPP-age relationships in China based on 3121 field survey samples, *Biogeosciences*, 21, 625–639, <https://doi.org/10.5194/bg-21-625-2024>, 2024a.
- Li, W., Guo, W.-Y., Pasgaard, M., Niu, Z., Wang, L., Chen, F., Qin, Y., and Svenning, J.-C.: Human fingerprint on structural density of forests globally, *Nat. Sustain.*, 6, 368–379, <https://doi.org/10.1038/s41893-022-01020-5>, 2023.
- Li, W., Guo, W.-Y., Pasgaard, M., Niu, Z., Wang, L., Chen, F., Qin, Y., Qiao, H., and Svenning, J.-C.: Unmanaged naturally
450 regenerating forests approach intact forest canopy structure but are susceptible to climate and human stress, *One Earth*, 7, 1068–1081, <https://doi.org/10.1016/j.oneear.2024.05.002>, 2024b.



- Li, X., Liu, Q., Cai, Z., and Ma, Z.: Specific Leaf Area and Leaf Area Index of Conifer Plantations in Qianyanzhou Station of Subtropical China, *J. Plant Ecol. (Chinese Version)*, 31, 93–101, <https://doi.org/10.17521/cjpe.2007.0012>, 2007.
- Li, X., Zhang, H., Shang, R., Chen, J., Wang, D., Zhu, J., Huang, H., Lin, S., Pan, B., Yuan, W., and Piao, S.: It is time to
455 optimize forest management policy for both carbon sinks and wood harvest in China, *Natl. Sci. Rev.*, 12, nwae464, <https://doi.org/10.1093/nsr/nwae464>, 2024c.
- Lin, X., Shang, R., Chen, J. M., Zhao, G., Zhang, X., Huang, Y., Yu, G., He, N., Xu, L., and Jiao, W.: High-resolution forest age mapping based on forest height maps derived from GEDI and ICESat-2 space-borne lidar data, *Agric. For. Meteorol.*, 339, <https://doi.org/10.1016/j.agrformet.2023.109592>, 2023.
- 460 Liu, Y., Liu, R., and Chen, J. M.: Retrospective retrieval of long-term consistent global leaf area index (1981–2011) from combined AVHRR and MODIS data, *J. Geophys. Res. Biogeosciences*, 117, 1–14, <https://doi.org/10.1029/2012JG002084>, 2012.
- Liu, Y., Yu, D., Xun, B., Sun, Y., and Hao, R.: The potential effects of climate change on the distribution and productivity of *Cunninghamia lanceolata* in China, *Environ. Monit. Assess.*, 186, 135–149, <https://doi.org/10.1007/s10661-013-3361-6>, 2014.
- 465 Lu, Y., Coops, N. C., Wang, T., and Wang, G.: A process-based approach to estimate Chinese fir (*Cunninghamia lanceolata*) distribution and productivity in Southern China under climate change, *Forests*, 6, 360–379, <https://doi.org/10.3390/f6020360>, 2015.
- Luyssaert, S., Schulze, E. D., Börner, A., Knohl, A., Hessenmöller, D., Law, B. E., Ciais, P., and Grace, J.: Old-growth forests as global carbon sinks, *Nature*, 455, 213–215, <https://doi.org/10.1038/nature07276>, 2008.
- 470 Ming, A., Yang, Y., Liu, S., Nong, Y., Li, H., Tao, Y., Sun, D., Lei, L., Zeng, J., and An, N.: The impact of near natural forest management on the carbon stock and sequestration potential of *Pinus massoniana* (Lamb.) and *Cunninghamia lanceolata* (Lamb.) Hook. plantations, *Forests*, 10, 1–13, <https://doi.org/10.3390/f10080626>, 2019.
- Mund, M., Kummert, E., Hein, M., Bauer, G. A., and Schulze, E.-D.: Growth and carbon stocks of a spruce forest chronosequence in central Europe, *For. Ecol. Manage.*, 171, 275–296, [https://doi.org/10.1016/S0378-1127\(01\)00788-5](https://doi.org/10.1016/S0378-1127(01)00788-5), 2002.
- 475 Ni, X., Lin, C., Chen, G., Xie, J., Yang, Z., Liu, X., Xiong, D., Xu, C., Yue, K., Wu, F., and Yang, Y.: Decline in nutrient inputs from litterfall following forest plantation in subtropical China, *For. Ecol. Manage.*, 496, 119–445, <https://doi.org/10.1016/j.foreco.2021.119445>, 2021.
- Pan, Y., Birdsey, R. A., Fang, J., Houghton, R., Kauppi, P. E., Kurz, W. A., Phillips, O. L., Shvidenko, A., Lewis, S. L.,
480 Canadell, J. G., Ciais, P., Jackson, R. B., Pacala, S. W., McGuire, A. D., Piao, S., Rautiainen, A., Sitch, S., and Hayes, D.: A Large and Persistent Carbon Sink in the World’s Forests, *Sci.*, 333, 988–993, <https://doi.org/10.1126/science.1201609>, 2011.
- Qin, J. and Shangguan, Z.: Effects of forest types on leaf functional traits and their interrelationships of *Pinus massoniana* coniferous and broad-leaved mixed forests in the subtropical mountain, Southeastern China, *Ecol. Evol.*, 9, 6922–6932, <https://doi.org/10.1002/ece3.5259>, 2019.



- 485 Ryan, M. G., Binkley, D., Fownes, J. H., Giardina, C. P., and Senock, R. S.: An experimental test of the causes of forest growth decline with stand age, *Ecol. Monogr.*, 74, 393–414, <https://doi.org/10.1890/03-4037>, 2004.
- Shang, R., Chen, J. M., Xu, M., Lin, X., Li, P., Yu, G., He, N., Xu, L., Gong, P., Liu, L., Liu, H., and Jiao, W.: China's current forest age structure will lead to weakened carbon sinks in the near future, *Innovation*, 4, <https://doi.org/10.1016/j.xinn.2023.100515>, 2023.
- 490 Shang, R., Lin, X., Chen, J. M., Liang, Y., Fang, K., Xu, M., Yan, Y., Ju, W., Yu, G., He, N., Xu, L., Liu, L., Li, J., Li, W., Zhai, J., and Hu, Z.: China's annual forest age dataset at 30 m spatial resolution from 1986 to 2022, *Earth Syst. Sci. Data Discuss.*, 2025, 1–31, <https://doi.org/10.5194/essd-2024-574>, 2025.
- Wang, B., Li, M., Fan, W., Yu, Y., and Chen, J. M.: Relationship between net primary productivity and forest stand age under different site conditions and its implications for regional carbon cycle study, *Forests*, 9, 5, <https://doi.org/10.3390/f9010005>, 2018.
- 495 Wang, S., Chen, J. M., Ju, W. M., Feng, X., Chen, M., Chen, P., and Yu, G.: Carbon sinks and sources in China's forests during 1901–2001, *J. Environ. Manage.*, 85, 524–537, <https://doi.org/10.1016/j.jenvman.2006.09.019>, 2007.
- Wang, S., Zhou, L., Chen, J., Ju, W., Feng, X., and Wu, W.: Relationships between net primary productivity and stand age for several forest types and their influence on China's carbon balance, *J. Environ. Manage.*, 92, 1651–1662, <https://doi.org/10.1016/j.jenvman.2011.01.024>, 2011.
- 500 White, M. A., Thornton, P. E., Running, S. W., and Nemani, R. R.: Parameterization and Sensitivity Analysis of the BIOME–BGC Terrestrial Ecosystem Model: Net Primary Production Controls, *Earth Interact.*, 4, 1–85, [https://doi.org/10.1175/1087-3562\(2000\)004<0003:pasaot>2.0.co;2](https://doi.org/10.1175/1087-3562(2000)004<0003:pasaot>2.0.co;2), 2000.
- Wu, W., Huang, C., Zheng, D., Chen, X., Zheng, S., Kang, H., and Miao, S.: Analysis of Characteristic of Forest Carbon Storage and Destiny in Fujian Province Abstract :, *J. Sichuan Agric. Univ.*, 34, 298–303, <https://doi.org/10.16036/j.issn.1000-2650.2016.03.007>, 2016.
- 505 Wu, Z., Zhou, C., Zhou, X., Hu, X., and Gan, J.: Variability after 15 years of vegetation recovery in natural secondary forest with timber harvesting at different intensities in southeastern China: Community diversity and stability, *Forests*, 9, <https://doi.org/10.3390/f9010040>, 2018.
- 510 Xia, J., Yuan, W., Lienert, S., Joos, F., Ciais, P., Viovy, N., Wang, Y. ping, Wang, X., Zhang, H., Chen, Y., and Tian, X.: Global Patterns in Net Primary Production Allocation Regulated by Environmental Conditions and Forest Stand Age: A Model-Data Comparison, *J. Geophys. Res. Biogeosciences*, 124, 2039–2059, <https://doi.org/10.1029/2018JG004777>, 2019.
- Xie, X., Lu, F., Luo, W., Xu, Z., and Su, H.: Leaf area index estimation and correction of Eucalyptus plantation in Guangxi Autonomous Region, *Acta Ecol. Sin.*, 42, 4451–4462, <https://doi.org/10.5846/stxb202106111562>, 2022.
- 515 Xu, W., Zhang, B., Xu, Q., Gao, D., Zuo, H., Ren, R., Diao, K., and Chen, Z.: Enhanced Carbon Storage in Mixed Coniferous and Broadleaf Forest Compared to Pure Forest in the North Subtropical–Warm Temperate Transition Zone of China, *Forests*, 15, 1520, <https://doi.org/10.3390/f15091520>, 2024.



- Ying, Z., Irland, Li., Zhou, X., Song, Y., Wen, Y., Liu, J., Song, W., and Qiu, Y.: Plantation development: Economic analysis of forest management in Fujian Province, China, *For. Policy Econ.*, 12, 223–230, <https://doi.org/10.1016/j.forpol.2009.11.001>, 2010.
- Yingchun, L., Guirui, Y., Qiufeng, W., and Yangjian, Z.: Huge Carbon Sequestration Potential in Global Forests, *J. Resour. Ecol.*, 3, 193–201, <https://doi.org/10.5814/j.issn.1674-764x.2012.03.001>, 2012.
- Yu, Y., Chen, J. M., Yang, X., Fan, W., Li, M., and He, L.: Influence of site index on the relationship between forest net primary productivity and stand age, *PLoS One*, 12, 1–20, <https://doi.org/10.1371/JOURNAL.PONE.0177084>, 2017.
- Zha, T. S., Barr, A. G., Bernier, P. Y., Lavigne, M. B., Trofymow, J. A., Amiro, B. D., Arain, M. A., Bhatti, J. S., Black, T. A., and Margolis, H. A.: Gross and aboveground net primary production at Canadian forest carbon flux sites, *Agric. For. Meteorol.*, 174–175, 54–64, <https://doi.org/10.1016/j.agrformet.2013.02.004>, 2013.
- Zhang, Y., Lu, D., Jiang, X., Li, Y., and Li, D.: Forest Structure Simulation of Eucalyptus Plantation Using Remote-Sensing-Based Forest Age Data and 3-PG Model, *Remote Sens.*, 15, <https://doi.org/10.3390/rs15010183>, 2023.
- Zhao, M. and Zhou, G. S.: Estimation of biomass and net primary productivity of major planted forests in China based on forest inventory data, *For. Ecol. Manage.*, 207, 295–313, <https://doi.org/10.1016/j.foreco.2004.10.049>, 2005.
- Zheng, J., Mao, F., Du, H., Li, X., Zhou, G., Dong, L., Zhang, M., Han, N., Liu, T., and Xing, L.: Spatiotemporal simulation of net ecosystem productivity and its response to climate change in subtropical forests, *Forests*, 10, 708, <https://doi.org/10.3390/f10080708>, 2019.
- Zhou, G., Meng, C., Jiang, P., and Xu, Q.: Review of Carbon Fixation in Bamboo Forests in China, *Bot. Rev.*, 77, 262–270, <https://doi.org/10.1007/s12229-011-9082-z>, 2011.
- Zhou, G., Houlton, B. Z., Wang, W., Huang, W., Xiao, Y., Zhang, Q., Liu, S., Cao, M., Wang, X., Wang, S., Zhang, Y., Yan, J., Liu, J., Tang, X., and Zhang, D.: Substantial reorganization of China’s tropical and subtropical forests: Based on the permanent plots, *Glob. Chang. Biol.*, 20, 240–250, <https://doi.org/10.1111/gcb.12385>, 2014.
- Zhou, L., Shalom, A. D. D., Wu, P., He, Z., Liu, C., and Ma, X.: Biomass production, nutrient cycling and distribution in age-sequence Chinese fir (*Cunninghamia lanceolata*) plantations in subtropical China, *J. For. Res.*, 27, 357–368, <https://doi.org/10.1007/s11676-015-0167-0>, 2016.
- Zhou, X. and Wingfield, M. J.: Eucalypt diseases and their management in China, *Australas. Plant Pathol.*, 40, 339–345, <https://doi.org/10.1007/s13313-011-0053-y>, 2011.
- Zhou, Z., Yuan, L., and Wang, Z.: Leaf longevity of main evergreen tree species of Guiyang City, *Ecol. Sci.*, 27, 148–153, 2008.