



QBOi El Niño Southern Oscillation experiments : Teleconnections of the QBO

Naoe, Hiroaki¹, Jorge L. García-Franco², Chang-Hyun Park³, Mario Rodrigo⁴, Froila M. Palmeiro^{4,5}, Federico Serva⁶, Masakazu Taguchi⁷, Kohei Yoshida¹, James A. Anstey⁸, Javier García-Serrano^{4,9}, Seok-
5 Woo Son³, Yoshio Kawatani¹⁰, Neal Butchart¹¹, Kevin Hamilton¹², Chih-Chieh Chen¹³, Anne Glanville¹³, Tobias Kerzenmacher¹⁴, François Lott¹⁵, Clara Orbe¹⁶, Scott Osprey¹⁷, Mijeong Park¹³, Jadwiga H. Richter¹³, Stefan Versick¹⁴, Shingo Watanabe¹⁸

¹Meteorological Research Institute (MRI), Tsukuba, 305-0052, Japan

²National School of Earth Sciences (Escuela Nacional de Ciencias de la Tierra), UNAM, CDMX, 04510, Mexico

10 ³School of Earth and Environmental Sciences, Seoul National University, Seoul, 08826, Korea

⁴Group of Meteorology, Universitat de Barcelona, Barcelona, 08028, Spain

⁵CMCC Foundation - Euro-Mediterranean Center on Climate Change, Bologna, 40127, Italy

⁶Institute of Marine Sciences, National Research Council (CNR-ISMAR), 00133, Italy

⁷Department of Earth Science, Aichi University of Education, Kariya, 448-0001, Japan

15 ⁸Canadian Centre for Climate Modelling and Analysis, Environment and Climate Change Canada, V8N 1V8, Canada

⁹Barcelona Supercomputing Center (BSC), Barcelona, 08034, Spain

¹⁰Faculty of Environmental Earth Science, Hokkaido University, Sapporo, 060-0810, Japan

¹¹Met Office, Exeter, EX1 3PB, UK

¹²International Pacific Research Center (IPRC) University of Hawaii, Honolulu, 96822, USA

20 ¹³U. S. National Science Foundation National Center for Atmospheric Research (NSF NCAR), Boulder, 80305, USA

¹⁴Karlsruher Institut für Technologie (KIT), Karlsruhe, 76131, Germany

¹⁵Laboratoire de Météorologie Dynamique (LMD), Ecole Normale Supérieure, Paris, 75231, France

¹⁶National Aeronautics and Space Administration (NASA) Goddard Institute for Space Studies (GISS), New York, 10025, USA

25 ¹⁷Atmospheric, Oceanic and Planetary Physics, University of Oxford, Oxford, OX1 3PU, UK

¹⁸Japan Agency for Marine-Earth Science and Technology (JAMSTEC), Yokohama, 236-0001, Japan

Correspondence to: Hiroaki Naoe (hnaoe@mri-jma.go.jp)

30

Abstract. This study examines Quasi-Biennial Oscillation (QBO) teleconnections and their modulation by the El Niño–Southern Oscillation (ENSO), using a multi-model ensemble of the Atmospheric Processes And their Role in Climate (APARC) QBO initiative (QBOi) models. Some difficulties arise in examining observed QBO-ENSO teleconnections from distinguishing the QBO and ENSO influences outside of the QBO region, due to aliasing between the QBO and ENSO over



35 the historical record. To separate the QBO and ENSO signals, simulations are conducted with annually-repeating prescribed
sea-surface temperatures corresponding to idealized El Niño or La Niña conditions (QBOi EN and LN experiments,
respectively). In the Arctic winter climate, higher frequencies of sudden stratospheric warmings (SSWs) are found in EN than
LN. The frequency differences in SSW between QBO westerly (QBO-W) and QBO easterly (QBO-E) are indistinguishable,
suggesting that the polar vortex responses to the QBO are much weaker than those to the ENSO in these models. The Asia-
40 Pacific subtropical jet (APJ) shifts significantly equatorward during QBO-W compared to QBO-E in observations, while the
APJ-shift is not robust across models, regardless of the ENSO phases. In the tropics, these experiments do not show a robust
or coherent QBO influence on precipitation. The sign and spatial pattern of the precipitation response vary widely across
models and experiments, indicating that any potential QBO signal is strongly modulated by the prevailing phases of the ENSO.
The QBO teleconnection to the Walker circulation around boreal summer/autumn is investigated to identify the strongest
45 signal in each model. It is found that the upper-level westerly and lower-level easterly anomalies in the equatorial troposphere
over the Indian Ocean and Western Pacific are detected in the observations and most models in the La Niña year. Overall, the
QBO can modulate the zonal circulation over the tropical Indian-Pacific oceans, with its impact varying depending on the
ENSO phase.

50 **Short summary (500 characters, incl. spaces)**

This study examines links between the stratospheric Quasi-Biennial Oscillation (QBO) and large-scale atmospheric
circulations in the tropics, subtropics, and polar regions. The QBO teleconnections and their modulation by the El Niño–
Southern Oscillation (ENSO) are investigated through a series of climate model experiments. While QBO teleconnections are
qualitatively reproduced by the multi-model ensemble, they are not consistent due to modelled QBO bias and other systematic
55 model biases.

Key words: stratosphere-troposphere coupling, teleconnection, QBO, ENSO



1 Introduction

60 The Quasi-Biennial Oscillation (QBO) and the El Niño–Southern Oscillation (ENSO) are the leading modes of climate variability in the tropical stratosphere and tropical troposphere, respectively. The QBO is a semi-periodic wind variation characterized by downward propagating easterly and westerly wind regimes in the equatorial stratosphere with an average period of about 28 months (Baldwin et al., 2001; Anstey et al., 2022b). The QBO is an important source of predictability due to its long timescale and its teleconnections outside the tropical stratosphere. The QBO is primarily driven by vertical
65 momentum fluxes due to upward-propagating equatorial wave activity generated by tropospheric convective systems (Lindzen and Holton, 1968; Holton and Lindzen, 1972; Plumb and McEwan, 1978).

Over the past a couple of decades, atmospheric general circulation models (GCMs) and Earth system models (ESMs) are being increasingly developed to include an internally generated QBO to represent more realistic modes of internal variability (e.g. Butchart et al., 2018). Most of these models require parameterization of unresolved gravity waves to simulate an internally
70 generated QBO, including specific conditions of parameterized and/or resolved convection, high horizontal and vertical resolution, and weak implicit and explicit grid-scale dissipation (Anstey et al., 2022b). Although the QBO is primarily an equatorial stratospheric phenomenon, it impacts the climate system outside this region via teleconnections. We can obtain a more in-depth understanding of QBO teleconnections (extratropical impacts, tropical and subtropical impacts, and their interaction with other phenomena) by intercomparing many state-of-the-art, stratosphere-resolving models that simulate a
75 QBO-like oscillation in the tropical stratosphere.

The QBO can influence the Northern Hemisphere (NH) winter stratosphere by modulating planetary-scale waves that distort the stratospheric polar vortex. The observed statistical relationship between the QBO phase and polar vortex strength is commonly referred to as the Holton-Tan effect (Holton and Tan, 1980, 1982). When the QBO in the lower stratosphere (~50 hPa) is in its westerly phase (QBO-W), the polar vortex is observed to be stronger and colder, and the likelihood of sudden
80 stratospheric warming (SSW) events is reduced. Conversely, when the QBO is in its easterly phase (QBO-E), the stratospheric polar vortex is weaker, warmer, and more disturbed. The underlying mechanisms for this effect have been extensively examined by many observational and modeling studies. The mechanism proposed by Holton and Tan (1980) to explain this relationship involves a latitudinal shift of the zero-wind line, which acts as an effective waveguide for upward-propagating planetary waves in the NH winter stratosphere (Holton and Tan, 1980; Baldwin et al., 2001; Anstey and Shepherd 2014; Watson and Gray, 2014; Gray et al., 2018; Lu et al., 2020; Anstey et al., 2022b). A similar but distinct mechanism involves
85 planetary waves interacting with the zonal wind anomalies associated with the QBO secondary circulation, not requiring zero-wind-line-induced wave breaking (Ruzmaikin et al., 2005; Naoe and Shibata, 2010; Garfinkel et al., 2012b; White et al., 2015; Naoe and Yoshida, 2019; Rao et al., 2020; Anstey et al., 2022b). A tropospheric pathway of the Holton-Tan relationship has also been proposed. This mechanism involves Rossby waves propagating from regions of tropical convection to higher



90 latitudes, including the Aleutian low-pressure region, and the stratospheric polar vortex is disturbed by the subsequent upward wave activity flux into the stratosphere, which is modulated through tropospheric processes (Yamazaki et al., 2020). Although the relative importance of these different mechanisms remains somewhat unclear, due to the QBO's long timescale, the Holton-Tan relationship leads to increased predictability of the extratropical stratosphere on subseasonal time scales (Boer and Hamilton, 2008; Scaife et al., 2014; Garfinkel et al., 2018).

95 The QBO can affect the troposphere by modifying deep convective activity and vertical wind shear along the tropopause (Gray et al., 1992; Collimore et al., 2003). The QBO-induced zonal-mean meridional circulation modulates the temperature vertical profile in the equatorial upper troposphere and lower stratosphere (UTLS), leading to a QBO signature in tropical tropopause temperature and wind. Although the idea of a “direct effect” of the QBO on the tropical and subtropical UTLS had been discussed in the literature since the 1960s, it was not yet widely accepted until the early 2000s (Hitchman et al., 2021).
100 Recently a possible downward influence of the QBO on the tropical troposphere has been found in the Madden-Julian Oscillation (MJO) (Yoo and Son, 2016; Marshall et al., 2016; Son et al., 2017; Martin et al. 2021; Elsbury et al., submitted). For more recent reviews of stratosphere-troposphere coupling in the tropics, see Haynes et al. (2021) and Hitchman et al. (2021).

Observational and modeling studies suggest that the interannual variability of tropical precipitation is, at least partially,
105 modulated by the phase of the QBO (Collimore et al., 2003; Liess and Geller, 2012; Gray et al., 2018). In observations, the QBO signal in tropical precipitation shows zonally asymmetric patterns, e.g. wetter conditions in the eastern Pacific Intertropical Convergence Zone (ITCZ) during QBO-W compared to QBO-E (Gray et al, 2018, Serva et al., 2022). The similarity between the QBO and ENSO signals in observations could potentially be caused by the higher number of El Niño events coinciding with QBO-W than with QBO-E (García-Franco et al., 2022). Serva et al. (2022) analyzed the simulated
110 precipitation in Atmospheric Model Intercomparison Project (AMIP)-type simulations from the first phase of QBOi experiments (Butchart et al., 2018) and found that those simulations have limited ability to reproduce the observed modulation of the tropical tropopause level processes, even after subtracting the variability associated with the ENSO index. In these sea-surface temperature (SST)-forced, free-running simulations, the east Pacific ITCZ precipitation response to the QBO, which resembles the observed pattern, is simulated by many, though not all models (Fig. 11 of Serva et al. [2022]). However, the
115 simulated QBO signal on the tropopause is generally underestimated or not realistic. Using similar SST-forced simulations with a single model, García-Franco et al. (2023) suggested that the simulated precipitation response to the QBO is heavily dependent on the state of ENSO and the Walker circulation, the strength of the QBO and the ocean-atmosphere coupling.

In the subtropics, a direct influence of the QBO modulates the subtropical jet by the QBO secondary circulation. Observational studies indicate that the QBO can affect the subtropical jet variability especially in the Pacific sector (e.g.
120 Garfinkel and Hartmann, 2011a; 2011b). During QBO-W, a horseshoe-shaped zonal wind anomaly forms in the upper troposphere and lower stratosphere associated with the equatorward shift of the Asian-Pacific jet (APJ) (Crooks and Gray, 2005; Simpson et al., 2009), and the resultant response is found even in the East Asian near the surface (Park et al., 2022; Park



and Son, 2022). A study using QBO-resolving multi-model ensemble found no clear evidence of a QBO teleconnection to the subtropical Pacific-sector jet (Anstey et al., 2022c).

125 ENSO teleconnections to the NH winter stratosphere have been widely reported in a large number of observational studies (van Loon and Labitzke, 1987; Camp and Tung, 2007; Garfinkel and Hartmann, 2007; Song and Son, 2018) and in modeling studies (Taguchi and Hartmann, 2006; García-Serrano et al., 2017; Palmeiro et al., 2017, 2023; Weinberger et al., 2019). During El Niño winters, the polar vortex is weaker and the polar region is warmer than ENSO neutral years. In contrast, during strong La Niña winters, a weakening of the Aleutian low and destructive linear interference with the climatological wave pattern was identified (Iza et al., 2016). Observations show that SSW events occur preferentially during both El Niño and La Niña winters than during ENSO-neutral winters (Butler and Polvani, 2011; Garfinkel et al., 2012a). However, there may be sampling errors due to the relatively short observational record (Domeisen et al., 2019), and increased SSWs during La Niña winters are sensitive to the SSW definition (Song and Son, 2018). Observed relationships between ENSO and SSWs are often not replicated by models. Models often simulate ENSO events and teleconnections that are considerably more linear compared to the available observational data (Domeisen et al., 2019), and systematic model biases in atmospheric winds and temperatures may affect the ENSO-SSW connection (Tyrrell et al., 2022).

ENSO has significant impacts on the global atmospheric circulations, and QBO teleconnections may also be influenced by El Niño and La Niña. The QBO itself has been shown to be affected by ENSO, with weaker QBO amplitude and faster QBO phase propagation under El Niño than La Niña conditions (Taguchi, 2010a). Previous studies investigating the joint effects of QBO and ENSO on polar vortex variability in winter suggested that their interactions are nonlinear insofar as the Holton-Tan relationship is found to be significant in the La Niña phase but much weaker in the El Niño phase (Wei et al., 2007; Garfinkel and Hartmann, 2008; Calvo et al., 2009; Richter et al., 2011; Hansen et al., 2016). A recent observational study (Kumar et al., 2023) investigated the combined effects of the QBO and ENSO in modulating the extratropical winter troposphere during the 1979–2018 period. They found that during La Niña, QBO signals in the polar vortex were amplified and the polar vortex and subtropical jet were enhanced under QBO-W. During El Niño, a stronger subtropical jet and the warmer polar vortex were present under QBO-W, while QBO anomalies in the tropical stratosphere were diminished and the poleward extent and amplitude of the QBO-induced mean meridional circulation was reduced. In contrast, the polar vortex weakens more when El Niño and QBO easterly occur together than would be expected by the sum of their individual effects (Walsh et al., 2022). However, there remains lack of consensus on the nature of nonlinearity in QBO–ENSO teleconnections in the extratropical circulation of the NH winter stratosphere.

In the tropical troposphere, the QBO and ENSO teleconnections remain less understood than those in the extratropics. A relatively small number of studies have analyzed tropical tropospheric QBO teleconnections using models that simulate the QBO (Rao et al., 2020; García-Franco et al., 2022, 2023; Serva et al., 2022). As noted by García-Franco et al. (2022, 2023), the observational record is likely too short to separate QBO teleconnections in the tropical troposphere from the strong influence of ENSO, because El Niño winters often coincide with the westerly phase of the QBO.



The goal of this study is to reexamine QBO teleconnections to the extratropics and tropics but now address combined QBO-ENSO influences using a new dataset of idealized ENSO experiments. Model experiments, which are capable of separating QBO and ENSO influences on the extratropical and tropical troposphere outside of the QBO region, are a valuable tool to study the modulation of QBO teleconnections by ENSO. To isolate the QBO teleconnections from the influence of ENSO, we

160 conduct model integrations with annually-repeating prescribed SSTs characteristic of typical El Niño and La Niña conditions, removing ENSO diversity from consideration. The Quasi-Biennial Oscillation initiative (QBOi), an international project supported by the World Climate Research Programme (WCRP) core project Atmospheric Processes And their Role in Climate (APARC), has focused on assessing internally generated QBOs in climate models and improving understanding of how to simulate a realistic QBO (Butchart et al., 2018; Anstey et al., 2022a,c; Bushell et al., 2022; Richter et al., 2022). In order to

165 study QBO and ENSO teleconnections and their mutual interactions, QBOi has coordinated additional experiments building on the QBOi phase-1 experiments, referred here as the “QBOiENSO” experiments, using participating QBOi atmospheric general circulation models (AGCMs) and Earth System Models (ESMs) forced by prescribed "perpetual El Niño" and "perpetual La Niña" SSTs (Kawatani et al., in revision). In this paper, we have examined QBO teleconnections modulated by ENSO and their robustness using this multi-model ensemble of QBO-resolving models that have run the QBOiENSO

170 experiments, and evaluated them by comparison against the QBOi phase-1 “Experiment 2”, which represents the control case of ENSO-neutral conditions. Further details of how the QBOiENSO experiments are constructed can be found in Kawatani et al. (in revision). The structure of the paper is as follows. Section 2 describes datasets of the QBOiENSO experiments and observations, and the analysis methods employed. Section 3 characterizes the combined effects of QBO–ENSO teleconnections on the polar winter stratosphere (Holton-Tan relationship). Sections 4 and 5 present the subtropical and tropical impacts of the QBO modified by ENSO, respectively. Finally, Section 6 provides a summary of our findings and discussion.

175

2 Data and Methods

We use nine AGCMs and ESMs participating in the QBOi project, conducting three experiments. The first one is the QBOi Experiment 2 using climatological SST and sea ice conditions (Butchart et al., 2018). We hereafter refer to it as the control (CTL) experiment. The other two experiments are the QBOiENSO experiments, QBOiElNino and QBOiLaNina (Kawatani et al., in revision). They are also time-slice experiments consistent with the QBOi Experiment 2 design, but prescribed "perpetual El Niño" and "perpetual La Niña" SSTs are used here. They are referred to hereafter as the EN and LN experiments, respectively. The models that performed the CTL, EN, and LN experiments are EC-EARTH3.3 (hereafter EC-EARTH for short), ECHAM5sh, EMAC, GISS-E2-2G (GISS for short), LMDz6 (LMDz for short), MIROC-AGCM-LL (MIROC-AGCM for short), MIROC-ESM, MRI-ESM2.0, and CESM1(WACCM5-110L) (WACCM for short). Their characteristics have been

180

185 described in Butchart et al. (2018) and Kawatani et al. (in revision). MRI-ESM2.0 (Yukimoto et al., 2019) is an updated version of the model documented in Butchart et al. (2018), and it includes changes aimed at improving the modelled QBO (Naoe and



Yoshida, 2019). Model integration years for three experiments are presented in Table 1. Due to data availability issues, EMAC is not included in Section 4 and 5.1.

190 Table 1. Model integration years

Model name	Years	
	¹ QBOi Exp2	² QBOi ENSO
³ EC-EARTH	101-yr	101-yr
⁴ ECHAM5sh	50-yr	40-yr
EMAC	106-yr	106-yr
GISS-E2-2G	3 × 30-yr	3 × 30-yr
LMDz	70-yr	82-yr
MIROC-AGCM	3 × 30-yr	100-yr
MIROC-ESM	3 × 100-yr	100-yr
MRI-ESM2.0	30-yr	50-yr
⁵ WACCM	3 x 30-yr	100-yr

¹QBOi Experiment 2 (or CTL experiment)

²QBOi ENSO experiments (QBOiElNino and QBOiLaNina experiments)

³EC-EARTH3.1 for QBOi Exp2 and EC-EARTH3.3 for QBOi ENSO

⁴Only r2i1pi is used in ECHAM5sh.

195 ⁵CESM1 (WACCM5-110L) is abbreviated to WACCM.

Observed teleconnections are quantified using a modern reanalysis dataset, the European Centre for Medium-Range Weather Forecasts (ECMWF) fifth generation atmospheric reanalysis (ERA5; Hersbach et al., 2020) in 1959–2021. The representation of the QBO in ERA5 as compared to other reanalyses is evaluated by Naoe et al. (2025). Observed precipitation is evaluated using the dataset of the Global Precipitation Climatology Project (GPCP; Adler et al., 2003, 2016) in 1979–2022. Because the design of QBOiENSO experiments uses the Japan Meteorological Agency’s (JMA) defined NINO.3 index (https://ds.data.jma.go.jp/tcc/tcc/products/el_nino/index/index.html), the classification of ENSO phases is based on this index.

To determine if observed teleconnections are manifested in the model runs, we do model-observation comparison by applying the same QBO phase definitions to the models that are optimal for observed teleconnections. Here, we use 'standard' indices (e.g., 50-hPa equatorial wind for the QBO), without adjusting them on a model-by-model basis, for all analyses presented in this article. This will facilitate comparisons with other works. As noted by Anstey et al. (2022c), different QBO indices can maximize the response of different teleconnections (e.g. Gray et al., 2018). Thus, making these choices can account for diversity of QBO signals (tropical convection, subtropical jet response, extratropical basic-state zonal-mean flow for the



Holton-Tan effect etc.), which may lead to variations in the QBO teleconnections across the planet. Zonal wind biases need to be carefully considered when defining the QBO phases in model outputs, as noted by Serva et al. (2022). Here QBO phases are identified when the deseasonalized westerly and easterly zonal-mean zonal wind (QBO-W and QBO-E) averaged over 5° S–5° N (weighted by cosine of latitude) exceeds a given threshold value at selected pressure levels. Specifically, QBO-W and QBO-E are classified from December-January-February (DJF) zonal wind at 50 hPa using $> 0.5 \sigma$ (standard deviation) and $< -0.5 \sigma$ in Section 3.1 (Figs. 2 and 3), using $\geq 0 \text{ m s}^{-1}$ and $< 0 \text{ m s}^{-1}$ in Section 3.2 (Fig. 5), using $\geq 2 \text{ m s}^{-1}$ and $\leq -2 \text{ m s}^{-1}$ in Section 5.1 (Figs. 8, 9, and 10), and from February-March zonal wind at 70 hPa using $> 0.5 \sigma$ and $< -0.5 \sigma$ in Section 4 (Figs. 6 and 7). In the Walker circulation analysis in subsection 5.2, the strongest signal is identified from a kind of Hovmöller diagram that are assessed at three levels of 50, 70, and 85 hPa in seasons from May to October with QBO definitions provided in the legend of Figs 11, 12, and 13. This approach is used to highlight the model dependency and seasonality of the QBO signal. It accounts for model diversity and biases in the simulated QBOs and tropical convection, which may lead to variations in the simulated QBO teleconnections across models.

ENSO composites in observations are done in the extratropics and subtropics for individual seasons (Sections 3, 4, and 5.2) and in the tropics for individual months (Section 5.1). The Bonferroni correction, as described by Holm (1979), is used for the two-sided t -test when the QBO phase is not defined by the preferred 70 hPa level during June-July-August (JJA). In this method, the p -value of the statistical test is adjusted by dividing it by m , the number of tests. For instance, if the QBO definition is modified by season only, $m = 2$; if it is modified by both season and vertical level, $m = 3$. Accordingly, $\alpha' = \alpha/m$, where $\alpha = 0.025$ (the 5% significance level for a two-sided test), and α' denotes the adjusted p -value.

3 QBO teleconnections: the extratropical route

A previous study about teleconnections of the QBO in a multi-model ensemble of QBO-resolving models (Anstey et al. 2022c) suggested that QBOi models underestimated the polar vortex response to the equatorial zonal wind at 50 hPa in comparison to reanalyses. They suggested that such weak responses were likely due to model errors, such as systematically weak QBO amplitudes near 50 hPa, affecting the teleconnection. Because most of the models that have run the QBOiElNino (EN) and QBOiLaNina (LN) experiments considered here are the same models whose QBOiExp2 (CTL) runs were analyzed by Anstey et al. (2022c), EN and LN runs may similarly underestimate the polar vortex response to the QBO. This section investigates the extratropical route of the QBO teleconnection modulated by ENSO. First, we examine the Holton-Tan effect, and then show the SSW statistics.

3.1 Holton-Tan relationship

Figure 1 shows the correlation coefficient in DJF between the 50 hPa equatorial zonal wind at 5° S–5° N and the polar vortex strength at different altitudes in the CTL, EN, and LN experiments, together with ENSO-neutral, El Niño and La Niña winters for the ERA5 reanalysis. In the reanalysis, correlations are strongest for the polar vortex at 15 hPa, being roughly 0.63 during




Figure 1: Vertical profiles of correlation coefficient between the QBO zonal wind at 50 hPa averaged over 5° S–5° N and the polar-vortex zonal wind at 55°–65° N in December-January-February (DJF) for QBOi models and ERA5. Circles represent statistical significance at the 90 % level. Red and blue bars represent 5–95 % confidence ranges using a bootstrap method repeating 1000 times in EN and LN experiments for the models as well as El Niño and La Niña winters for ERA5. Number of winters available for each model run for each experiment (ENSO phase) are displayed at the upper panel. For example, 'NUE32EN15LN15' in the ERA5 panel means there are 32 ENSO-neutral, 15 El Niño, and 15 La Niña winters, respectively.

La Niña and 0.40 during El Niño, and the correlation during the ENSO-neutral winter is slightly stronger than that of El Niño. The uncertainty range (horizontal bars) shows the 5–95% range of correlation coefficients derived from bootstrap resampling. Although the confidence interval for La Niña clearly excludes zero in the stratosphere, the confidence for El Niño is close to zero at many altitudes, demonstrating large uncertainty in the strength of the correlation especially for El Niño and ENSO-neutral winters.

Most of the model correlations show smaller uncertainty than ERA5 due to having larger sample sizes. They have significant correlations over a range of altitudes only in a particular experiment. For example, at some altitudes GISS has a significant correlation in EN, MIROC-AGCM in LN, MRI-ESM2.0 in LN, WACCM in CTL, and MIROC-ESM has a significant correlation only in the lower stratosphere in CTL. LMDz has no or little correlations between the equatorial QBO winds and the polar vortex wind for any of the experiments. These results indicate that most models show weak positive correlations with the same sign as the reanalysis, but in most cases these correlations are not statistically significant. This suggests that inter-




Figure 2: Composite differences of DJF-mean zonal-mean zonal wind between QBO-W and QBO-E in the CTL, EN, and LN experiments including the ENSO neutral, El Niño, and La Niña winters for ERA5. QBO phases are classified using deseasonalized DJF zonal-mean zonal wind at 50 hPa averaged over 5° S–5° N using $> 0.5 \sigma$ for QBO-W and $< -0.5 \sigma$ for QBO-E. Contour interval is 3 m s^{-1} . Dots represent statistical significance at the 90 % level. Number of winters available for each model run, and numbers of QBO-E and QBO-W winter classification are displayed at the upper right corner of each panel. For example, 'N100E30W41' in EC-EARTH and QBOiExp2 means there are 100 winters in which 30 QBO-E winters and 41 QBO-W winters are classified.

model differences in the QBO-polar vortex relationship, or differences between experiments for the same model, may not be distinguishable.

Previous QBOi multi-model ensemble studies on the Holton-Tan relationship indicated that weak QBO amplitudes exist in the equatorial lower stratosphere. They argued that the low likelihood of obtaining the observed correlation from most models could result from model errors that impact the representation of these teleconnections (Richter et al., 2022; Anstey et al., 2022c). Figure 2 shows composite differences of zonal-mean zonal wind between QBO-W and QBO-E in the CTL, EN, and



La Niña). The vortex strength difference between QBO-W and QBO-E peaks at roughly 10 m s^{-1} in the middle stratosphere (near 10 hPa) during DJF for the Neutral and El Niño groups, and the response is strongest in La Niña with a peak value of 15 m s^{-1} . No model reproduces the observed Holton-Tan relationship in all three experiments (CTL, EN and LN). In LN, ECHAM5sh, MIROC-AGCM, and MRI-ESM2.0 tend to reproduce the observed response, peaking at $3\sim 6 \text{ m s}^{-1}$ in the polar vortex region. Some models have significant composite differences of zonal wind in a particular experiment at 60° N and 10 hPa. For example, GISS and ECHAM5sh show a significant difference in EN peaking at 7 m s^{-1} , and WACCM in CTL. In short, for any of the three experiments the models more often than not show a stronger polar vortex during NH winter when the 50-hPa QBO wind is westerly, and a weaker vortex when it is easterly, consistent with but weaker than the observed response.




Figure 3: (a) Monthly differences (QBO-W minus QBO-E) of zonal-mean zonal wind at 10 hPa averaged over $55^\circ\text{--}65^\circ \text{ N}$ as a measure of the stratospheric polar vortex strength for the CTL experiment. QBO phases are classified same as Fig. 2. Solid dots show significant differences between QBO-W and QBO-E phases at the 90 % confidence level using a Monte Carlo test. Numbers in the legend are the cases included in each QBO phase. While for the experiments, ENSO is neutral, all years in ERA5 are included in the analysis (1959–2022). (b) Same as (a) but for the EN experiment including El Niño winters for ERA5. (c) Same as (a) but for the LN experiment including La Niña winters for ERA5. Numbers of (QBO-W, QBO-E) categories for ERA5 are (12, 11) in ENSO-Neutral, (7, 4) in El Niño, and (9, 4) in La Niña winters.



Intraseasonal Holton-Tan effects are investigated in Fig. 3, which shows the composite difference (QBO-W minus QBO-E) of monthly zonal-mean zonal wind at 10 hPa, 55°–65° N in CTL, EN, and LN experiments, together with ERA5. ERA5 presents a maximum Holton-Tan effect in January with a peak of 13 m s⁻¹ for the mean (dashed black line in the top panel), but this difference is lower during ENSO-neutral years (solid black line in the top panel). Seasonal evolution of Holton-Tan effect is different between El Niño and La Niña winters; it seems stronger in early and late winters for the El Niño winters (middle panel) and in mid-winter for the La Niña winters (bottom panel), although it should be cautioned that the sample sizes (number of W/E winters) are small for both El Niño and La Niña groups. Some models show a similar seasonal cycle as ERA5 for their CTL runs (MIROC-ESM and ECHAM5sh with the statistical significance). Also, GISS in all months as well as LMDz and MIROC-AGCM in a few months exhibit an opposite sense to the observed Holton-Tan relationship for CTL. In EN, GISS, WACCM, EMAC, and ECHAM5sh capture the early-winter response in December although it is not statistically significant. The Holton-Tan relationship in El Niño years could depend on SSW occurrence because of the nonlinear joint effects of QBO and ENSO on the polar vortex as already explained in the Introduction. In LN, MRI-ESM2.0 and GISS capture the observed late-winter response relatively well, and other models do not show any response or even an opposite response.

3.2 SSW statistics

In this subsection, we examine SSW statistics modulated by ENSO and the QBO in the northern polar region. Previous observational studies indicated that the ratio of SSW frequency between La Niña and ENSO-neutral winters is dependent on details of the SSW definition (Butler and Polvani, 2011; Garfinkel et al., 2012a; Song and Son, 2018), and SSW statistics have been shown to depend on model biases (Tyrrell et al., 2022). Figure 4 shows frequencies of major and minor SSWs and final warming dates in NH for ERA5 and QBOi models. The approach to identify major, minor, and final warming dates is similar to what was proposed by previous studies (Charlton and Polvani, 2007; Butler et al., 2015). Major SSWs are identified when zonal-mean westerlies in winter are changed to easterlies at 60° N and 10 hPa. For minor SSWs, the zonal wind does not reverse but there is a change in sign of the meridional gradient of the zonal-mean temperature. Final warming date refers to the seasonal transition from westerly to easterly, with winds remaining easterly for the next months.

We consider first the influence of ENSO on SSW frequency. The frequency of major SSWs is high in ERA5 during both El Niño and La Niña years, compared to ENSO-neutral. Thus, we expect that major SSW frequencies in the QBOi models would be similar to the observations and have fewer events in CTL and more events in EN and LN experiments. LMDz and GISS reproduce the nonlinear observed ENSO response to some extent. However, most models show more SSWs during EN and they do not capture the LN response (e.g., EC-EARTH, MIROC-AGCM, MRI-ESM2.0). ECHAM5sh has similar frequencies in CTL and LN and more events in EN. GISS shows large spreads in CTL and EN, suggesting that the response is not statistically robust. In ERA5, frequencies of minor SSWs are similar in both ENSO-neutral and El Niño years and there are fewer events in La Niña years (In Fig. 4b). There is a large spread in minor SSW frequencies between the models. EC-EARTH and ECHAM5sh show high frequencies of minor SSWs in EN whereas LMDz and MRI-ESM2.0 show low frequencies of minor SSWs. MIROC-AGCM produces fewer SSWs in all the ENSO experiments, and MIROC-ESM shows relatively higher




Figure 4: SSW statistics in NH in CTL, EN, and LN experiments for QBOi models including ENSO neutral, El Niño, and La Niña years for ERA5, based on their daily data. Frequency (number of events per decade) of (a) major SSWs (reversal of zonal-mean zonal wind at 10 hPa and 60° N; U60) and of (b) minor SSWs (reversal of 90°–60° N temperature gradient at 10 hPa without U60 reversal), occurring across full seasons. Different marker signs are used to indicate ensemble members, and uncertainties are estimated at the 5–95% level based on bootstrapping of 10 years of winter months. (c) Boxplots of final SSW date (day of year), considering full seasons, i.e., from westerlies onset to their turn to easterly for ERA5 and QBOi models based on their daily data. ENSO neutral winter experiment (CTL, grey), El Niño (EN, brown), and La Niña (LN, purple).

frequencies for both major and minor SSWs in EN and LN compared to the other MIROC model. The GISS ensemble shows large spread in all three experiments, suggesting an important role for internal variability.

The final warming date is when the transition from winter westerlies to summer easterlies occurs in the polar stratosphere (Butler et al., 2015). If the stratosphere is warmer in the polar regions, the transition of zonal wind to easterlies occurs earlier, and if it is colder the transition is delayed. Hence, we assume that in El Niño (La Niña) years when the polar stratosphere would be warmer (colder) as described in the Introduction, the final warming date might happen earlier (later). Consistent with this expectation, in ERA5, the final warming date in La Niña years is more delayed than that in ENSO-neutral and El Niño years. GISS and MRI-ESM2.0 exhibit later final warming dates in LN than in EN, which is similar to the observed response. On the other hand, EC-EARTH, ECHAM5sh, LMDZ, MIROC-AGCM and MIROC-ESM do not show earlier final warming dates in EN, which is the opposite to the observed response. These results imply that the QBOi models have significant biases in reproducing observed SSWs statistics. Large inter-model variability is also diagnosed by means of the Northern Annular Mode (NAM) index (Eyring et al, 2020) compositing at 500 hPa, as shown in Fig. S1, where the geopotential heights during LN tend to be lower and there are changes in the intensity of the extratropical signature between LN and EN.




Figure 5: Scatter plots between mean vortex strength (60° N, 10 hPa) and major SSW frequency during DJF for different QBO and ENSO conditions. Major SSWs are identified as a reversal of daily zonal-mean zonal wind at 60° N and 10 hPa. QBO phases are classified using DJF-mean zonal-mean zonal wind anomalies at 50 hPa averaged over 5° S– 5° N using ≥ 0 m s $^{-1}$ for QBO-W (WLY in panel) and < 0 m s $^{-1}$ for QBO-E (ELY). The anomalies are calculated for each ensemble member of each experiment for the simulation data; those ones are calculated using all years (1959–2021 seasons) for the ERA5 data. For ERA5, El Niño and La Niña winters are when all three DJF months have the El Niño and La Niña flag, respectively. Number of (QBO-W, QBO-E) categories for ERA5 are (24, 15) in ENSO-Neutral, (5, 6) in El Niño, and (9, 4) in La Niña winters. For each condition, each model, the data are randomly resampled 100 times with replacement for the same number of the sample, and then 95% ranges are obtained and plotted.

Next, we investigate the influence of the QBO on major SSW frequencies modulated by ENSO in the NH winter. Figure 5 shows scatter plots between the climatological zonal-mean zonal wind at 60° N and 10 hPa and mean frequency of major SSWs in DJF during QBO-W and QBO-E years for three ENSO conditions. In ERA5, major SSW frequencies under QBO and ENSO conditions are likely to be distinguishable. Averaged over all QBO conditions, the NH polar vortex is stronger in La Niña than El Niño winters, while SSW frequencies are slightly higher in La Niña than El Niño winters, and both are higher than ENSO-neutral winters. Major SSW frequencies in La Niña winters are significantly higher under QBO-E and lower under QBO-W, whereas those in El Niño winters are indistinguishable between QBO-W and QBO-E. Most QBOi models are characterized by linear distributions between SSW frequencies and the polar vortex strength. The EN experiment displays



higher frequencies of SSWs than the LN experiment and SSW frequencies between QBO-W and QBO-E are indistinguishable.

380 This shows that polar vortex responses to ENSO conditions in the QBOi models are stronger than responses to the QBOs in these models. Some models, such as EMAC, MIROC-AGCM, and MIROC-ESM, have very weak responses to both ENSO and QBO.

4 The subtropical jet route of QBO teleconnection

This section examines the subtropical jet route of QBO teleconnection in the QBOi ENSO experiments. Only the late winter
385 period of February to March, when the subtropical route is strongest in the observations (Garfinkel and Hartmann, 2011a; Park et al., 2022), is considered. Since the subtropical jet change in response to the QBO is pronounced for the APJ, analyses are performed for the zonal wind averaged over the Pacific sector (150° E– 150° W). The sensitivity of the QBO-APJ connection to the ENSO phase is also examined.

The QBO-W minus QBO-E (W–E) composite differences are shown in Fig. 6 for the ENSO-neutral, El Niño, and La Niña
390 winters, for both ERA5 and QBOi ENSO experiments. During the ENSO-neutral winter, the QBO W–E anomaly exhibits a distinct horseshoe-shaped pattern extending from the tropical lower stratosphere to the subtropical lower troposphere (top-left panel in Fig. 6). It is accompanied by a quasi-barotropic easterly anomaly in the extratropics. More importantly, the zonal wind anomalies switch sign across the climatological APJ (contour). This indicates that the APJ shifts equatorward during the QBO-W winter compared to the QBO-E winter. Most models underestimate or fail to reproduce the observed QBO-APJ connection.
395 The dipolar wind anomalies are much weaker than those in observations in five models (i.e., EC-EARTH, ECHAM5sh, GISS, LMDz, and MIROC-ESM). Although one lobe of the dipolar wind anomalies is significant in ECHAM5sh and GISS, other models (i.e., EC-EARTH, LMDz, and MIROC-ESM) have statistically insignificant dipolar wind anomalies. MIROC-AGCM and MRI-ESM2.0 exhibit the opposite sign. Such large inter-model spread is consistent with a previous study (Anstey et al., 2022c). The QBO-APJ connection differs between El Niño and La Niña (top-middle and top-right panels in Fig. 6; Garfinkel
400 and Hartmann, 2010). As the APJ strengthens over the Pacific sector (150° E– 150° W) in response to El Niño (compare contours; Rasmusson and Wallace, 1983; Mo et al., 1998; Lu et al. 2008), the QBO subtropical wind anomalies become stronger near the APJ center during El Niño (top-middle panel; Ma et al., 2023). In contrast, the W–E anomalies switch sign across the climatological APJ during La Niña (top-right panel) as the APJ becomes slightly weaker (compare line contours in the top-left and top-right panels). The APJ's response to ENSO is consistently reproduced across models, whereas the ENSO
405 modulation of the QBO-APJ connection shows large inter-model spread. While all models capture a stronger APJ during EN than LN (compare line contours in the middle and right columns), they exhibit significant biases in reproducing the ENSO modulation of the QBO-APJ connection (filled contour).

The inter-model spread of the QBO subtropical route is summarized by the APJ-shift index in Fig. 7. The APJ-shift index
is derived from the QBO-W minus QBO-E wind differences shown in Fig. 6. Specifically, it is defined by the 250-hPa QBO
410 zonal wind difference from the northern flank (40° – 50° N) to the southern flank (20° – 30° N) of the climatological APJ.




Figure 6: QBO-W minus QBO-E composite differences of zonal wind averaged over the Pacific sector (150° E–150° W) for the ENSO-neutral (top-left), El Niño (top-middle), and La Niña (top-right) winters and those for the CTL, EN, and LN experiments (left to right columns). Values that are statistically significant at the 95% confidence level are cross-hatched. Contour denotes the climatological jet with zonal wind speed equal to or greater than 30 m s⁻¹. QBO-W and QBO-E phases are defined as deseasonalized February–March zonal-mean zonal wind over 5°S–5°N at 70 hPa being > 0.5 σ and < -0.5 σ , respectively. Numbers of (QBO-W, QBO-E) categories for ERA5 are (10, 9) in ENSO-Neutral, (6, 3) in El Niño, and (12, 7) in La Niña winters.

Negative values indicate that the APJ shifts equatorward during QBO-W compared to QBO-E. The observed APJ-shift index is significantly negative during the ENSO-neutral winter (black) and La Niña winter (blue), but is insignificant during El Niño (red). This is consistent with dipolar wind anomalies switching sign across the climatological APJ during ENSO-neutral and La Niña winters, while the change in APJ strength is more pronounced during El Niño winters (see Fig. 6). The APJ-shift index is not robust across models. None of the models show a statistically significant APJ shift in response to the QBO, regardless of the ENSO phase. This result suggests that QBOi models significantly underestimate or fail to reproduce the subtropical route of the QBO teleconnection and its modification by the ENSO.




Figure 7: QBO-W minus QBO-E composite difference of the Asia-Pacific Jet (APJ) shift index. The APJ-shift index is defined as the difference of the 250-hPa zonal wind anomalies averaged over the Pacific sector (150° E– 150° W) between the northern flank (40° – 50° N) and the southern flank (20° N– 30° N) of the climatological APJ core. The negative value indicates that the APJ moves toward the equator during the QBO-W phase. The composite differences are shown for the La Niña or LN experiment (blue), ENSO-neutral or CTL (black), and El Niño or EN experiment (red). The values are considered significant if the 5-95 % range of the bootstrap distribution (vertical dashed lines) does not encompass zero.

5 QBO teleconnections: the tropical route

This section investigates the tropical route of the QBO teleconnection modulated by ENSO, focusing on tropical precipitation and the Walker circulation.

5.1 Tropical precipitation

Figure 8 shows the DJF seasonal-mean precipitation differences between QBO-W and QBO-E in EN and LN, together with anomalies for El Niño and La Niña winters for GPCP. In the observations, the QBO signals are largest and statistically significant in the tropical Pacific and Indian oceans, and are in good agreement with previous analyses (Liess and Geller, 2012; García-Franco et al., 2022). The positive equatorial Pacific signal in the GPCP dataset, which resembles an El Niño anomaly (Dommenget et al., 2013; Capotondi et al., 2015), is particularly strong and statistically significant in DJF. This signal is associated with the three strongest El Niño events (1982–1983, 1997–1998, 2015–2016) coinciding with the westerly QBO phase (García-Franco et al., 2023).





Figure 8: DJF seasonal mean precipitation differences (mm day^{-1}) (QBO-W minus QBO-E) for (left) EN and (right) LN experiments for the QBOi models including El Niño and La Niña years in GPCP data. Hatching denotes statistical significance at the 95% confidence level according to a bootstrap test for the observations and a two-sided t-test for the models. The observed composite sizes in months are shown in parenthesis in the GPCP panels. QBO phases are classified using deseasonalized DJF mean zonal-mean zonal wind averaged over 5°S – 5°N at 50 hPa using $\geq 2 \text{ m s}^{-1}$ for QBO-W and $\leq -2 \text{ m s}^{-1}$ for QBO-E.

Although most models do not show such El-Niño-like precipitation anomaly patterns in either experiment, several models exhibit significant precipitation QBO-related signals. For example, GISS, ECHAM5sh and MRI-ESM2.0 show significant



545

Figure 9: (a-b) Box plots of QBO-W minus QBO-E differences in DJF precipitation (mm day^{-1}) in (a) the western equatorial Pacific (WEP) and (b) EN3.4 region ($5^{\circ}\text{ S}–5^{\circ}\text{ N}$, $170^{\circ}–120^{\circ}\text{ W}$). Error bars represent the 95 % confidence interval.

QBO responses in the EN experiment, which are comparable in magnitude to the signal diagnosed in GPCP when considering all months (Fig. S2a) but weaker than the corresponding observed signals in El Niño and La Niña conditions. However, the response in other models is generally weaker and the spatial distribution of the anomalies is not consistent between models. In the LN experiments, the models similarly do not show a clear precipitation signal in the Pacific, but EC-EARTH, ECHAM5sh, WACCM and MIROC-ESM show several precipitation signals over the Indian Ocean and Australia. A multi-model mean response is shown in Fig. S2, which illustrates the lack of model agreement characterized by a virtually zero QBO signal in a multi-model mean sense across the tropics. Thus, this result suggests that there is lack of model agreement in the spatial distribution and sign of the tropical precipitation response to the QBO phase. These results are also supported by Fig. S3, which shows DJF seasonal outgoing longwave radiation (OLR) differences between QBO-W and QBO-E in EN and LN together with ERA5. None of the models show an OLR signal comparable to observations, and models show a distinct QBO signal between EN and LN experiments. In other words, there is no consistent or robust precipitation response across models or experiments.

Previous studies have shown that the QBO signal in DJF is prominent in particular regions of the tropical Pacific: the western equatorial Pacific (WEP) region ($5^{\circ}\text{ S}–5^{\circ}\text{ N}$, $120^{\circ}–170^{\circ}\text{ E}$) and the Nino3.4 region ($5^{\circ}\text{ S}–5^{\circ}\text{ N}$, $170^{\circ}–120^{\circ}\text{ W}$) (EN3.4; Gray



et al., 2018, Serva et al., 2022, García-Franco et al., 2022). To examine the extent to which precipitation in these regions is sensitive to the QBO phase, we evaluated the area-averaged precipitation in both regions as a function of QBO and ENSO phases (Fig. S4). The QBOi models show significant spread in their climatology of precipitation amounts but all the simulations seem to reproduce the observed ENSO signal, i.e., wetter conditions in the EN3.4 region and drier in the WEP in EN and the opposite in LN, regardless of the QBO phase.

Figure 9 shows the area-averaged precipitation differences (QBO-W minus QBO-E) per region and in CTL (Neutral), El Niño and La Niña experiments/winters. In observations, the precipitation signal associated with the QBO during El Niño is opposite in sign to that of La Niña. One must consider the very small sample size (roughly 3 to 5 winters in each composite) in these observations when interpreting these results. Regardless of the sign and magnitude of the observed response, the models seem to show disagreement on the sign of the precipitation response, i.e., comparing models in the same experiment provides no consistent precipitation signal. For example, while the La Niña response is positive over the WEP in observations and most models agree, only 5 out of 7 models show a positive response.

When looking at individual models, GISS and MIROC-ESM agree that the precipitation signal (QBO-W minus QBO-E) is positive in the WEP in all their three experiments but no model agrees on the sign of the precipitation response in all three experiments for the EN3.4 region. These results emphasize that the QBO signal on tropical precipitation may strongly depend on the state of ENSO as suggested by the observations (García-Franco et al., 2023). Overall, some models show a robust and significant precipitation response to the QBO but this response is distinct from observations, and varies in sign and magnitude amongst experiments and models.

One reason for this inter-model and/or inter-experiment spread in the precipitation response could be the model spread in QBO-related temperature-associated anomalies at the equator resulting from the QBO mean meridional circulation and thermal wind balance. The QBO impact on the tropical tropopause layer (TTL) region is important for the QBO teleconnection in the tropical route (Haynes et al., 2021, Hitchman et al., 2021). Here, one common hypothesis is that if a cold QBO anomaly lies in the TTL, convective systems may grow more efficiently, penetrating to greater altitudes, locally amplifying the zonal mean QBO cold anomaly (Tegtmeier et al., 2021). Figure 10 shows a scatter plot of the QBO-W minus QBO-E difference of air temperature at 100 hPa versus the precipitation difference (QBO-W minus QBO-E) over WEP. One could reasonably question whether models or experiments with a larger TTL temperature signal or static stability may also show a larger signal in precipitation. In this panel, the W-E TTL temperature signals diagnosed from ERA5 are larger than those of the models, and are strongest for El Niño. The precipitation signal diagnosed in GPCP is also largest in El Niño, possibly due to the coincidence of the largest El Niño events with the westerly QBO phase. We confirmed an impact of removing those strongest El Niño events (1982–1983, 1997–1998, 2015–2016) on the GPCP precipitation signal (Fig. S5). It is found that the impact is more dramatic over the full composites at the top, in which the Pacific signal disappears when not considering these cases. In the El Niño winters, it is only the eastern portion of the Pacific that changes significantly. There are some models that have a strong temperature signal, such as GISS and ECHAM5sh, which also have a strong negative precipitation signal in LN. However, most models have modest positive temperature differences without a clear precipitation signal. Overall, the QBOi models




Figure 10: Scatter plot of DJF air-temperature differences at 100 hPa (QBO-W minus QBO-E in K) versus precipitation differences (QBO-W minus QBO-E in mm day⁻¹) in the western equatorial Pacific region.

show unrealistically too-weak QBO wind amplitudes in the lower stratosphere (Bushell et al., 2022), and correspondingly have temperature anomalies that are too weak in the TTL (Serva et al., 2022), which could help explain the weak precipitation signals.

5.2 Walker circulation

In this subsection, we examine whether QBO impacts on the Walker circulation can be detected across different ENSO phases. To identify the strongest signal in each model, we conduct a thorough search, defining the QBO at slightly different seasons and vertical levels. The Bonferroni correction (Holm, 1979; Section 2) is applied to the two-sided *t*-test when the QBO definition deviates from the preferred 70 hPa level during JJA. Longitude-height cross-sections of zonal wind differences (QBO-W minus QBO-E) averaged over 5° S–5° N for each model are shown in Figs. 11 and 12 for the LN and EN experiments, respectively, alongside ERA5. A recent study (Rodrigo et al., in revision) shows that in reanalyses (ERA5, NCEP-NCAR, JRA-55) the QBO teleconnection to the Walker circulation is strongest under mean state conditions in boreal summer (JJA), followed by autumn (SON), and weakest in winter. However, under different ENSO conditions this timing may slightly shift, potentially due to ENSO influence on the QBO itself (Taguchi, 2010b; Kawatani et al., in revision). Additionally, model diversity and biases in the simulated QBO (Bushell et al., 2022) could lead to inter-model variations in the simulated QBO teleconnection, which appears weak in some models and occurs at different times of the year, spanning from May to November.





Figure 11: Climatology (black contours) and QBO Westerly (W) minus Easterly (E) differences (shading and colored contours) of equatorial zonal wind profiles, averaged over 5°S – 5°N , in the LN experiment for QBOi models. Black contours are drawn at 4 m s^{-1} intervals, and the colored contour intervals match the shading scale in the color bar. The target season for each panel is indicated in the title, with the QBO definition provided in the legend. In ERA5, 15 La Niña events are identified using the NINO3 index during DJF, and they are classified into 10 QBO-W and 5 QBO-E categories by analyzing the zonal-mean zonal wind at 50 hPa in summer and autumn. Only statistically significant zonal wind differences at the 95% confidence level are shaded. For models with a QBO definition other than 70 hPa during JJA, the Bonferroni correction is applied. Note that the color bar for ERA5 differs because of the larger QBO amplitude.

Focusing on ERA5 during La Niña (Fig. 11a), the SON climatological state (black contours) features upper-level easterlies over the Eastern Hemisphere and westerlies over the Western Hemisphere, with the opposite pattern in the lower troposphere. A distinct QBO signal is observed in the equatorial troposphere over the Maritime continent and western Pacific during SON. This signal is characterized by anomalous westerlies in the upper troposphere (red contours and shading) and anomalous easterlies in the lower troposphere (blue contours and shading). Relative to the climatological state, the QBO signal appears as a westward shift in the climatological pattern over the Maritime continent–western Pacific region. Similar signals to those observed in ERA5, featuring upper-level westerly and lower-level easterly anomalies, are also found in the models (Figs. 11b–i), although their locations vary. Specifically, the strongest signals are identified in EC-EARTH during JJA, MRI-ESM2.0 during JJA, GISS during SON, LMDz during JJA, ECHAM5sh during JJA (weak and barely significant), WACCM during MJJ, MIROC-AGCM during JJA, and MIROC-ESM during SON. In all models, the QBO signal is shifted westward over the Indian Ocean relative to the ERA5 signal. This westward shift suggests a weaker climatological zonal circulation (Walker-type cell) during the westerly QBO phase over the Indian Ocean and Maritime continent. The QBO-W minus QBO-E



655

660

Figure 12: Same as Figure 11, but for EN experiments and El Niño years in ERA5. In ERA5, 14 El Niño events are identified, and they are classified into 7 QBO-W and 7 QBO-E categories.

composite in CTL shows a similar teleconnection (Fig. S6). This modulation of tropical circulation by the QBO seems robust despite differences in seasonal timing and longitudinal location, as comparable teleconnections are observed in ERA5 and most models across the CTL and LN experiments.

For El Niño in ERA5 (Fig. 12a), the QBO signal in the equatorial troposphere peaks during SON and differs notably from that of La Niña. It shows anomalous easterlies in the upper troposphere over the Pacific Ocean and anomalous westerlies in the lower troposphere. This anomalous zonal circulation can be interpreted as a weaker Walker circulation in the central-eastern tropical Pacific. Reasonably similar patterns, featuring upper-level easterly and lower-level westerly anomalies, are also found in EN experiment, such as EC-EARTH during JJA, ECHAM5sh during JA, WACCM during JJ, MIROC-AGCM during JJA, and MIROC-ESM during ASO. Signals with only significant lower-level anomalous westerlies are detected in MRI-ESM2.0 during MJ. As in the La Niña case, the QBO signal in the models is shifted westward over the Indian Ocean and Maritime continent compared to ERA5, with some model-dependent variations. In some models, such as EC-EARTH, ECHAM5sh and WACCM, these teleconnections are associated with a stronger zonal circulation over the Indian Ocean and Maritime continent (approximately 60° E to 150° E). Other models, like MRI-ESM2.0, MIROC-AGCM and MIROC-ESM show signals that can be interpreted as an eastward shift of the climatological pattern. Exceptions are GISS and LMDz, where the signals more closely resemble those of CTL and LN experiments. Consequently, the QBO W-E signal in the models is less robust for EN compared to LN and CTL.

680




Figure 13: (a) Schematic Hovmöller diagram showing statistically significant signals over the equator for (a) CTL, (b) LN, and (c) EN experiments. The QBO-W minus QBO-E zonal wind signals are assessed at three different levels and across four standard seasons. Symbols are placed between standard seasons when the strongest signal occurs in an intermediate period. In the EN experiment, the 150-hPa level is used instead of 100 hPa, as the significant signal occurs at a slightly different altitude. Filled symbols represent westerly anomalies, while empty symbols indicate easterly anomalies. The QBO definition for each model and experiment is provided in the legend.

Figure 13 shows a summary diagram of the statistically significant composite differences in zonal wind over the equator in the CTL, LN and EN experiments. This figure can be interpreted as a kind of Hovmöller diagram, where the QBO-W minus QBO-E zonal wind signals are analyzed at three representative vertical levels across four standard seasons. These statistically significant signals are identified by examining the QBO's influence on the zonal circulation, while slightly varying the vertical level for different ENSO phases and the longitudinal region for different models. An example from the EC-EARTH CTL experiment is provided in Fig. S7. The same longitudinal region is analyzed across all seasons, although it may vary slightly



between models. The QBO phase is consistently defined in the specific season indicated in the legend (i.e., it does not vary seasonally). In some models, the strongest signals occur during transitional periods between standard seasons, so the corresponding symbols are placed accordingly. In CTL and LN, almost all models, along with ERA5, display a tropospheric signal characterized by upper-level westerlies and lower-level easterlies during JJA and/or SON, suggesting a weakening of the climatological Walker circulation in the models and a longitudinal shift in ERA5 during La Niña years. By contrast, the EN experiment reveals a different pattern, with upper-level easterlies and lower-level westerlies in the troposphere during the same seasons in ERA5 and most models. Overall, this figure highlights that the QBO, defined around summer and autumn, modulates the zonal circulation over the tropical Indian–Pacific region, with its impact varying across models depending on the ENSO phase. This modulation is more consistent across models in the CTL and LN experiments compared to the EN experiment.

6 Summary and Discussion

In this paper, we have examined QBO and ENSO teleconnections in the Arctic stratosphere, the subtropical Pacific jet, and the tropical troposphere. A multi-model ensemble of QBO-resolving models that performed the QBOiENSO experiments has been used to examine the robustness of these teleconnections. Difficulties can arise in distinguishing QBO and ENSO influences on the extratropics and tropical troposphere due to the observed aliasing between the QBO and ENSO. Here we have attempted to separate these competing influences by conducting model integrations with annually-repeating prescribed SSTs that are characteristic of either strong El Niño or La Niña conditions, thereby simplifying the ENSO forcing in comparison to the diversity of observed ENSO phases. We have reexamined QBO teleconnections to the extratropics and tropics that were explored in previous QBOi studies (Anstey et al., 2022c; Serva et al., 2022) but now addressing combined QBO-ENSO influences using this new QBOi dataset of idealized ENSO experiments.

The observed strength of correlation coefficients between 50-hPa equatorial zonal wind and the polar vortex strength at stratospheric altitudes in DJF shows large uncertainty (Fig. 1a) but the confidence intervals clearly exclude zero at most altitudes during La Niña and ENSO-neutral winters, while El Niño response is statistically significant over a smaller altitude range. The models show a smaller uncertainty due to their larger sample sizes (Fig. 1). Some models have weaker correlations for a particular ENSO experiment, similar to the observations. The Holton-Tan relationship in ERA5 represents the polar vortex being significantly stronger (weaker) under QBO-W (QBO-E) for all the ENSO phases, with the strongest response occurring in the La Niña phase. Nearly half of the models exhibit stronger polar vortex during NH winter under QBO-W for each experiment, consistent with, but much weaker than the observed response (Fig. 2). Seasonal evolution in ERA5 indicates a stronger signal in early (late) winter for the El Niño (La Niña) winters. In LN, two out of nine models capture the observed late-winter response relatively well, and other models do not show any response or even the opposite direction (Fig. 3).

Major SSWs occur frequently during both El Niño and La Niña winters, compared to ENSO-neutral, in ERA5. Most models show more events during EN but they do not catch the LN response, implying that the QBOi models have some trouble in



750 reproducing observed SSWs statistics (Fig. 4). Major SSW frequencies in ERA5 show strong variation with QBO and ENSO phase. QBOi models are characterized by linear distributions between SSW frequencies and the polar vortex strength in NH winters (similar to ERA5) and overall the EN (LN) experiment displays high (relatively low) SSW frequencies (Fig. 5). SSW frequencies between QBO-W and QBO-E are indistinguishable in the models, indicating that polar vortex responses to the idealized ENSO forcing in the QBOi models are strong, while vortex responses to equatorial QBOs are fairly weak.

755 The APJ changes in response to the QBO are investigated (Figs. 6 and 7), focusing on the late winter when the subtropical route is strongest in the observations. In observations, the QBO westerly anomaly exhibits a distinct horseshoe-shaped pattern extending from the tropical lower stratosphere to the subtropical lower troposphere, indicating that the APJ shifts equatorward during the QBO-W winter compared to the QBO-E winter. However, most models underestimate or fail to reproduce the observed QBO-APJ connection. The observed QBO-APJ connection differs between El Niño and La Niña. In observations, as
760 the APJ strengthens over the Pacific sector in response to El Niño, the QBO subtropical wind anomalies become stronger near the APJ center during El Niño while they do not change much during La Niña as the APJ becomes slightly weaker. All models capture a stronger APJ in EN than in LN.

The positive equatorial Pacific signal in the GPCP dataset, which resembles an El Niño anomaly for W-E differences, is particularly strong and statistically significant in DJF, as shown by previous studies that highlight the issue of strong ENSO
765 events coinciding with the westerly phase (García-Franco et al., 2023). Although most of the models do not show such El-Niño-like precipitation anomaly patterns in either EN or LN experiments, some models (EC-EARTH, ECHAM5sh, WACCM and MIROC-ESM) show significant precipitation signals over the Indian Ocean and Australia (Fig. 8). The precipitation response to the QBO in these experiments depends on both the model, region and ENSO phase, as there is no consistent response between the experiments for each model (Fig. 9). For example, the simulated and observed QBO signals in the Niño
770 3.4 region disagree on the magnitude and sign. To explore the causes of model versus observation differences, the strength of the QBO impact on the TTL region was analyzed as it is considered to be important for the QBO teleconnection in the tropical route (Fig. 10). In particular, we verified whether the strength of the temperature anomaly could explain inter-model or inter-experiment differences in the precipitation signals. Overall, the QBOi models have too-weak wind amplitudes and too-weak temperature anomalies in the lower stratosphere, which could help explain the weak precipitation signals.

775 The QBO teleconnection to the Walker circulation in reanalyses is strongest under mean state conditions in boreal summer, followed by autumn, and weakest in winter. Because model diversity and biases in the simulated QBOs in each model may cause the simulated QBO teleconnection to vary, the strongest signal is identified for each model, defining the QBO across slightly different seasons and vertical levels. A distinct QBO signal with upper-level westerly and lower-level easterly anomalies is observed in the equatorial troposphere over the Maritime continent–Western Pacific in ERA5 during La Niña
780 years, and similar characteristics, but with a zonal shift, are also found in all the models (Fig. 11). The QBO signal in EN notably differs from that in LN, with longitudinal variations between ERA5 and the models (Fig. 12). Overall, the QBO, defined during JJA and SON, modulates the zonal circulation over the tropical Indian-Pacific Oceans, with its impact varying based on the ENSO phase (Fig. 13).



We now consider three issues about modelling QBO-ENSO complexity raised by these results: forced SSTs, the seasonality and variation of the Walker circulation, biases of the QBO and other diagnostics. AMIP-type experiments, where observed SSTs and other external forcing are used, have been used here to examine QBO-ENSO teleconnections. However, the responses of the climate system to ENSO forcing tend to be nonlinear with respect to ENSO intensity and asymmetric with respect to the polarity of ENSO (Domeisen et al., 2019; Rao and Ren, 2016b, c). This means that it is difficult to isolate physical meaningful mechanisms from such a nonlinearity system and gain scientific insights into QBO-ENSO teleconnections. Conducting idealized experiments that take into account the ENSO-QBO diversity could help us to further elucidate scientifically meaningful mechanisms in such a complex system. It is noted that the experimental design of QBOiENSO (Kawatani et al. in revision) is annually locked with monthly-mean anomalies from the climatology. For example, the precipitation responses to the QBO for the AMIP-type experiments with interannually varying SSTs (Serva et al., 2022; García-Franco et al., 2022) is different from those for the QBOi ENSO experiments with perpetual SSTs. The precipitation response to the QBO in the equatorial Pacific signal in the GPCP dataset shows a statistically significant, El-Niño-like anomaly pattern. Most of the models do not show such El-Niño-like precipitation anomaly patterns in the CTL, EN or LN experiments, while such patterns were seen in some of the QBOi models in the QBOi Experiment 1 (Serva et al., 2022). The lack of a robust and coherent QBO-related precipitation signal across experiments and models highlights significant spread in how convection and circulation respond to a QBO forcing. This raises the possibility that the QBO's downward impact on tropical precipitation is too sensitive to model physics, or is perhaps muted by the lack of SST feedbacks (García-Franco et al., 2023, Randall et al., 2024) to be clearly detected.

One of the most important points from this study is that the Walker circulation would potentially play an important role in tropical teleconnections as well as extratropical teleconnections. The QBO influences on the mean state and variations over Walker circulation are affected by El Niño La Niña. However, every ENSO event differs in both its magnitude and in its location in the tropical Pacific where maximum SST anomalies are observed, and El Niño events tend to exhibit a wider range in the longitudinal location of the maximum SST anomaly in the equatorial Pacific (Domeisen et al., 2019). Thus, we are interested in two distinct and documented El Niño patterns, Eastern Pacific (EP) versus Central Pacific (CP, or Modoki) El Niños, which make a large difference in the Hadley and Walker circulations and also have markedly different impacts on remote regions. Therefore, idealized experiments forced with ENSO SST patterns of different spatial configurations, strengths, and signs (e.g., Lopez-Parages et al., 2016) would be beneficial for us to better achieve the changing impact of ENSO events on the QBO teleconnections. Furthermore, although it may be beyond the scope of atmospheric-only models, we here emphasize that tropical convection is inherently coupled with the ocean. Long-term simulations from coupled global circulation models (CGCMs) would be a convenient tool for testing responses of QBO-ENSO teleconnections associated with internal variability of the ocean-atmosphere coupled system (García-Franco et al., 2023; Randall et al., 2023). We emphasize on the importance of seasonality about the combined effect of QBO-ENSO on the tropical troposphere. So far, many previous studies have examined QBO-ENSO teleconnections in one specific season, such a boreal winter month. This is because the Hadley circulation is the strongest and tropical tropopause highest in this season, such that the stratospheric signal might be



expected to be stronger in the troposphere. Our results suggest that QBO teleconnections with the Walker circulation exhibit seasonal variability and a distinct zonally asymmetric pattern. These findings emphasize the need for further investigation to elucidate the drivers of the seasonal dependence, the nature of the asymmetry and the underlying mechanisms governing these interactions.

In the extratropical stratosphere, the previous studies of QBOi models suggested that the systematic weakness of the QBO-polar vortex coupling in the models might arise from systematically weak QBO amplitudes at lower levels in the equatorial stratosphere, polar vortex biases in winter, inadequate representation of stratospheric-troposphere coupling, etc. (Bushell et al., 2022; Richter et al., 2022; Anstey et al., 2022c). In our QBOiENSO experiments, such systematic model biases were also found because most of the modes were the same as the previous studies. In the tropics, our results suggested that the systematic bias of the QBO impact on the tropical troposphere might arise from the systematically weak QBO amplitudes at lower levels, precipitation bias, and inadequate representation of the Walker circulation. Thus, the combination of several biases could be reason why we have not seen a consistent response of the QBO teleconnections across the models and experiments. Therefore, it is plausible that consistent response with observations will not improve without correcting such model biases. Currently, a project of QBOi Phase 2 is in progress to assess the impact of QBO biases by using zonal mean nudged toward observations in the QBO region. Bias-corrected QBO amplitude, achieved through nudging methods, may provide insights for improving the representation of QBO teleconnections to the extratropics and the tropical troposphere.

Data availability. The QBOi data archive was hosted by the Centre for Environmental Data Analysis (CEDA), UK, and processing was performed on the JASMIN infrastructure. The ERA5 reanalysis data can be obtained from the ECMWF website (<https://www.ecmwf.int/en/forecasts/datasets/browse-reanalysis-datasets>). The El Niño Monitoring Index (NINO.3) was derived from an ENSO monitoring site of the Japan Meteorological Agency (https://ds.data.jma.go.jp/tcc/tcc/products/el_nino/index.html). The GPCP v2.3 was downloaded from <https://doi.org/10.7289/V56971M6> (Adler et al., 2016).

Author contribution. YK, KH, JA, JHR designed the QBOi ENSO-QBO experiments. FS, HN, KY, TK, SW, YK, JGS, FMP, PAB, FL, CO, JHR, CCJ created model-experimented data and uploaded them to the CEDA. HN, JLGF, CHP, MR, FMP, FS and MT conducted the analyses, prepared the figures, and contributed to the manuscript. JA, JGS, SWS, NB, and SO contributed in the interpretation of results. The first draft of the manuscript was prepared by HN, who involved all authors for the final version.

Competing interests. The authors declare that they have no conflict of interest.



Acknowledgements. All authors acknowledge the CEDA, which kindly hosted the QBOi data archive. HN was supported by MEXT, JSPS KAKENHI (grant numbers: JP22H04493, JP24K07140); HN and KY were supported by JSPS KAKENHI JP24K00710. YK was supported by JSPS KAKENHI (JP22K18743) and the Environment Research and Technology Development Fund (JPMEERF20242001) of the Environmental Restoration and Conservation Agency provided by Ministry of the Environment of Japan. YK and SW were supported by JSPS KAKENHI (JP22H01303 and JP23K22574). SW was supported by MEXT-Program for the advanced studies of climate change projection (SENTAN) Grant Number JPMXD0722681344. The numerical simulations of MIROC models were performed using the Earth Simulator. CP and SS were supported by Korea Environment Industry & Technology Institute (KEITI) through 'Climate Change R&D Project for New Climate Regime', funded by Korea Ministry of Environment (MOE) (2022003560004). MR was supported by the 'Ayudas para la Formación de Profesorado Universitario' programme (FPU20/03517), FMP was supported by the EU/HORIZON-funded MSCA-IF-GF SD4SP project (GA 101065820), and JGS acknowledges funding from the Spanish DYNCAST project (CNS2022-135312). The ECHAM5sh simulations were performed thanks to an ECMWF Special Project awarded to FS.

860

References

- Adler, R. F., Huffman, G. J., Chang, A., Ferraro, R., Xie, P.-P., Janowiak, J., Rudolf, B., Schneider, U., Curtis, S., Bolvin, D., Gruber, A., Susskind, J., Arkin, P., and Nelkin, E.: The Version-2 Global Precipitation Climatology Project (GPCP) monthly precipitation analysis (1979–present), *J. Hydrometeor.*, 4, 1147–1167, [https://doi.org/10.1175/1525-7541\(2003\)004<1147:TVGPCP>2.0.CO;2](https://doi.org/10.1175/1525-7541(2003)004<1147:TVGPCP>2.0.CO;2), 2003.
- Adler, R., Wang, J. J., Sapiano, M., Huffman, G., Chiu, L., Xie, P. P., Ferraro, R., Schneider, U., Becker, A., Bolvin, D., Nelkin, E., Gu, G., and NOAA CDR Program: Global Precipitation Climatology Project (GPCP) Climate Data Record (CDR), Version 2.3 (Monthly), National Centers for Environmental Information, <https://doi.org/10.7289/V56971M6>, 2016.
- Anstey, J. A., and Shepherd, T. G.: High-latitude influence of the quasi-biennial oscillation, *Q. J. Roy. Meteor. Soc.*, 140, 1–21. <http://doi.wiley.com/10.1002/qj.2132>, 2014.
- Anstey, J. A., Butchart, N., Hamilton, K., and Osprey, S. M.: The SPARC quasi-biennial oscillation initiative, *Q. J. Roy. Meteor. Soc.*, 148, 1455–1458, <https://doi.org/10.1002/qj.3820>, 2022a.
- Anstey, J. A., Osprey, S. M., Alexander, J., Baldwin, M.P., Butchart, N., Gray, L., Kawatani, Y., and Richter, J. H.: Impacts, processes and projections of the quasi-biennial oscillation, *Nat. Rev. Earth Environ.*, 3, 588–603, <https://doi.org/10.1038/s43017-022-00323-7>, 2022b.
- Anstey, J. A., Simpson, I. R., Richter, J. H., Naoe, H., Taguchi, M., Serva, F., Gray, L. J., Butchart, N., Hamilton, K., Osprey, S., Bellprat, O., Braesicke, P., Bushell, A. C., Cagnazzo, C., Chen, C.-C., Chun, H.-Y., Garcia, R. R., Holt, L., Kawatani, Y.,



- Kerzenmacher, T., Kim, Y.-H., Lott, F., McLandress, C. Scinocca, J., Stockdale, T. N., Versick, S., Watanabe, S., Yoshida, K., Yukimoto S.: Teleconnections of the quasi-biennial oscillation in a multi-model ensemble of QBO-resolving models, *Q. J. Roy. Meteor. Soc.*, 148, 1568–1592, <https://doi.org/10.1002/qj.4048>, 2022c.
- Baldwin, M. P., Gray, L. J. Gray, Dunkerton, T. J., Hamilton, K., Haynes, P. H., Randel, W. J., Holton, J. R., Alexander, M. J., Hirota, I., Horinouchi, T., Jones, D. B. A., Kinnnersley, J. S., Marquardt, C., Sato, K., and Takahashi, M.: The quasi-biennial oscillation, *Rev. Geophys.*, 39, 179–229, <https://doi.org/10.1029/1999RG000073>, 2001.
- ratropical predictive skill, *Clim. Dyn.*, 31, 987–1000, <https://doi.org/10.1007/s00382-008-0379-5>, 2008.
- Bushell, A.C., Anstey, J. A., Butchart, N., Kawatani, Y., Osprey, S.M., Richter, J. H., Serva, F., Braesicke, P., Cagnazzo, C., Chen, C.-C., Chun, H.-Y., Garcia, R. R., Gray, L. J., Hamilton, K., Kerzenmacher, T., Kim, Y.-H., Lott, F., McLandress, C., Naoe, H., Scinocca, J., Smith, A. K., Stockdale, T. N., Versick, S., Watanabe, S., Yoshida, K., Yukimoto, S.: Evaluation of the Quasi-Biennial Oscillation in global climate models for the SPARC QBO-initiative, *Q. J. Roy. Meteor. Soc.*, 148, 1459–1489, <https://doi.org/10.1002/qj.3765>, 2022.
- Butchart, N., Anstey, J. A., Hamilton, K., Osprey, S., McLandress, C., Bushell, A.C., Kawatani, Y., Kim, Y.-H., Lott, F., Scinocca, J., Stockdale, T., Bellprat, O., Braesicke, P., Cagnazzo, C., Chen, C.-C., Chun, H.-Y., Dobrynin, M., Garcia, R. R., García-Serrano, J., Gray, L. J., Holt, L., Kerzenmacher, T., Naoe, H., Pohlmann, H., Richter, J. H., Scaife, A. A., Schenzinger, V., Serva, F., Versick, S., Watanabe, S., Yoshida, K., and Yukimoto, S.: Overview of experiment design and comparison of models participating in phase 1 of the SPARC Quasi-Biennial Oscillation initiative (QBOi), *Geosci. Model Dev.*, 11, 1009–1032, <https://doi.org/10.5194/gmd-11-1009-2018>, 2018.
- Butler, A., and Polvani, L.: El Niño, La Niña, and stratospheric sudden warmings: Reevaluation in light of the observational record, *Geophys. Res. Lett.*, 38, L13807, <https://doi.org/10.1029/2011GL048084>, 2011.
- Butler, A., Seidel, D. J., Hardiman, S. C., Butchart, N., Birner, T., and Match, A.: Defining sudden stratospheric warmings, *Bull. Amer. Meteor. Soc.*, 96, 1913–1928, <https://doi.org/10.1175/BAMS-D-13-00173.1>, 2015.
- Butler, A. H., Polvani, L. M.: El Niño, La Niña, and stratospheric sudden warmings: A reevaluation in light of the observational record, *Geophys. Res. Lett.*, 38, L13807, <https://doi.org/10.1029/2011GL048084>, 2011.
- Butler, A. H., Polvani, L. M., and Deser, C.: Separating the stratospheric and tropospheric pathways of El Niño–Southern Oscillation teleconnections, *Environ. Res. Lett.*, 9, 024014, DOI 10.1088/1748-9326/9/2/024014, 2014.
- Butler, A. H., Seidel, D. J., Hardiman, S. C., Butchart, N., Birner, T., and Match, A.: Defining sudden stratospheric warmings, *Bull. Am. Meteor. Soc.*, 96, 1913–1928, <https://doi.org/10.1175/BAMS-D-13-00173.1>, 2015.
- Calvo, N., Giorgetta, M. A., Garcia-Herrera, R., and Manzini, E.: Nonlinearity of the combined warm ENSO and QBO effects on the Northern Hemisphere polar vortex in MAECHAM5 simulations, *J. Geophys. Res.*, 114, D13109, <https://doi.org/10.1029/2008JD011445>, 2009.
- Camp, C. D., and Tung, K. K.: Stratospheric polar warming by ENSO in winter: A statistical study, *Geophys. Res. Lett.*, 34, L04809, <https://doi.org/10.1029/2006GL028521>, 2007.



- Capotondi, A., and Sardeshmukh, P. D.: Optimal precursors of different types of ENSO events, *Geophys. Res. Lett.*, 42, 9952–9960, <https://doi.org/10.1002/2015GL066171>, 2015.
- Charlton, A. J., and Polvani, L. M.: A new look at stratospheric sudden warmings. Part I: Climatology and modeling benchmarks, *J. Clim.*, 20, 449–469, <https://doi.org/10.1175/JCLI3996.1>, 2007.
- 915 Collimore, C. C., Martin, D. W., Hitchman, M. H., Huesmann, A., and Waliser, D. E.: On the relationship between the QBO and tropical deep convection, *J. Clim.*, 16, 2552–2568, [https://doi.org/10.1175/1520-0442\(2003\)016<2552:OTRBTQ>2.0.CO;2](https://doi.org/10.1175/1520-0442(2003)016<2552:OTRBTQ>2.0.CO;2), 2003.
- Crooks, S.A., and Gray, L.J.: Characterization of the 11-year solar signal using a multiple regression analysis of the ERA-40 dataset, *J. Clim.*, 18, 996–1015, <https://doi.org/10.1175/JCLI-3308.1>, 2005.
- 920 Domeisen, D. I., Garfinkel, C. I., and Butler, A. H.: The teleconnection of El Niño Southern Oscillation to the stratosphere, *Rev. Geophys.*, 57, 5–47, <https://doi.org/10.1029/2018rg000596>, 2019.
- Dommenget, D., Bayr, T., and Frauen, C.: Analysis of the nonlinearity in the pattern and time evolution of El Niño Southern Oscillation, *Clim. Dynam.*, 40, 2825–2847, <https://doi.org/10.1007/s00382-012-1475-0>, 2013.
- Elsbury, D., Serva, F., Caron, J., Back, S.-Y., Orbe, C., Richter, J., Anstey, J. A., Bretonniere, P.-A., Butchart, N., Chen, C.-
- 925 C., Garcia-Serrano, J., Glanville, A., Kawatani, Y., Kerzenmacher, T., Lott, F., Naoe, H., Osprey, S., Palmeiro, F. M., Son, S.-W., Taguchi, M., Versick, S., Watanabe, S., Yoshida, K.: QBOi El Niño Southern Oscillation experiments: Assessing relationships between ENSO, MJO, and QBO, *EGUsphere* [preprint], <https://doi.org/10.5194/egusphere-2024-3950>, *Weather and Climate Dynamics*, (submitted).
- Eyring, V., Bock, L., Lauer, A., Righi, M., Schlund, M., Andela, B., Arnone, E., Bellprat, O., Brötz, B., Caron, L.-P.,
- 930 Carvahais, N., Cionni, I., Cortesi, N., Crezee, B., Davin, E. L., Davini, P., Debeire, K., de Mora, L., Deser, C., Docquier, D., Earnshaw, P., Ehbrecht, C., Gier, B. K., Gonzalez-Reviriego, N., Goodman, P., Hagemann, S., Hardiman, S., Hassler, B., Hunter, A., Kadow, C., Kindermann, S., Koirala, S., Koldunov, N., Lejeune, Q., Lembo, V., Lovato, T., Lucarini, V., Massonnet, F., Müller, B., Pandde, A., Pérez-Zanón, N., Phillips, A., Predoi, V., Russell, J., Sellar, A., Serva, F., Stacke, T., Swaminathan, R., Torralba, V., Vegas-Regidor, J., von Hardenberg, J., Weigel, K., and Zimmermann, K.: Earth System Model
- 935 Evaluation Tool (ESMValTool) v2.0 ? an extended set of large-scale diagnostics for quasi-operational and comprehensive evaluation of Earth system models in CMIP, *Geosci. Model Dev.*, 13, 3383–3438, <https://doi.org/10.5194/gmd-13-3383-2020>, 2020.
- García-Franco, J. L., Gray, L. J., Osprey, S., Chadwick, R., and Martin, Z.: The tropical route of quasi-biennial oscillation (QBO) teleconnections in a climate model, *Weather Clim. Dynam.*, 3, 825–844. <https://doi.org/10.5194/wcd-3-825-2022>, 2022.
- 940 García-Franco, J. L., Gray, L. J., Osprey, S., Jaison, A. M., Chadwick, R., and Lin, J.: Understanding the mechanisms for tropical surface impacts of the quasi-biennial oscillation (QBO), *J. Geophys. Res.*, 128, e2023JD038474, <https://doi.org/10.1029/2023JD038474>, 2023.
- García-Serrano, J., Cassou, C., Douville, H., Giannini, A., and Doblas-Reyes, F. J.: Revisiting the ENSO teleconnection to the tropical North Atlantic, *J. Clim.*, 30, 6945–6957, <https://doi.org/10.1175/JCLI-D-16-0641.1>, 2017.



- 945 Garfinkel, C. I., and Hartmann, D. L.: Effects of the El Niño–Southern Oscillation and the Quasi-Biennial Oscillation on polar temperatures in the stratosphere, *J. Geophys. Res.*, 112, D19112, <https://doi.org/10.1029/2007JD008481>, 2007.
- Garfinkel, C. I., and Hartmann, D. L.: Different ENSO teleconnections and their effects on the stratospheric polar vortex, *J. Geophys. Res.*, 113, D18114, <https://doi.org/10.1029/2008JD009920>, 2008.
- Garfinkel, C. I., and Hartmann, D. L.: The influence of the quasi-biennial oscillation on the troposphere in winter in a hierarchy of models. Part I: Simplified dry GCMs, *J. Atmos. Sci.*, 68, 1273–1289, <https://doi.org/10.1175/2011JAS3665.1>, 2011a.
- 950 Garfinkel, C. I., and Hartmann, D. L.: Influence of the quasi-biennial oscillation on the troposphere in winter in a hierarchy of models. Part II: perpetual winter WACCM runs, *J. Atmos. Sci.*, 68, 2026–2041, <https://doi.org/10.1175/2011JAS3702.1>, 2011b.
- Garfinkel, C. I., Butler, A. H., Waugh, D. W., Hurwitz, M. M., and Polvani, L. M.: Why might stratospheric sudden warmings occur with similar frequency in El Niño and La Niña winters?, *J. Geophys. Res.*, 117, D19106, <https://doi.org/10.1029/2012JD017777>, 2012a.
- 955 Garfinkel, C. I., Shaw, T. A., Hartmann, D. L., and Waugh, D. W.: Does the Holton-Tan mechanism explain how the quasi-biennial oscillation modulates the Arctic polar vortex?, *J. Atmos. Sci.*, 69, 1713–1733, <https://journals.ametsoc.org/doi/10.1175/JAS-D-11-0209.1>, 2012b.
- 960 Garfinkel, C. I., Schwartz, C., Domeisen, D. I. V., Son, S.-W., Butler, A. H., and White, I. P.: Extratropical atmospheric predictability from the Quasi-Biennial Oscillation in subseasonal forecast models, *J. Geophys. Res.*, 123, 7855–7866, <https://doi.org/10.1029/2018JD028724>, 2018.
- Gray, L. J., Anstey, J. A., Kawatani, Y., Lu, H., Osprey, S., and Schenzinger, V.: Surface impacts of the Quasi Biennial Oscillation, *Atmos. Chem. Phys.*, 18, 8227–8247, <https://www.atmos-chemphys.net/18/8227/2018/>, 2018.
- 965 Gray, W. M., Scheaffer, J. D., and Knaff, J. A.: Influence of the stratospheric QBO on ENSO variability, *J. Meteor. Soc. Japan*, 70, 975–995, https://doi.org/10.2151/jmsj1965.70.5_975, 1992.
- Hansen, F., Matthes, K., and Wahl, S.: Tropospheric QBO-ENSO interactions and differences between Atlantic and Pacific, *J. Clim.*, 29, 1353–1368, <https://doi.org/10.1175/JCLI-D-15-0164.1>, 2016.
- Haynes, P., Hitchcock, P., Hitchman, M., Yoden, S., Hendon, H., Kiladis, G., Kodera, K., and Simpson, I.: The influence of the stratosphere on the tropical troposphere, *J. Meteorol. Soc. Japan. Ser. II*, 99, 803–845, <https://doi.org/10.2151/jmsj.2021-040>, 2021.
- 970 Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horanyi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., A. Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Holm, E., Janiskova, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thepaut, J.-N.: The ERA5 global reanalysis, *Quart. J. Roy. Meteor. Soc.*, 146, 1999–2049, <https://doi.org/10.1002/qj.3803>, 2020.



- Hitchman, M. H., Yoden, S., Haynes, P. H., Kumar, V., and Tegtmeier, S.: An observational history of the direct influence of the stratospheric Quasi-biennial Oscillation on the tropical and subtropical upper troposphere and lower stratosphere, *J. Meteorol. Soc. Japan. Ser. II*, 99, 239–267, <https://doi.org/10.2151/jmsj.2021-012>, 2021.
- Holm, S.: A simple sequentially rejective multiple test procedure, *Scand. J. Statist.*, 6, 65–70. <http://www.jstor.org/stable/4615733>, 1979.
- Holton, J. R., and Lindzen, R. S.: An updated theory for the quasi-biennial cycle of the tropical stratosphere, *J. Atmos. Sci.*, 29, 1076–1080, [https://doi.org/10.1175/1520-0469\(1972\)029<1076:AUTFTQ>2.0.CO;2](https://doi.org/10.1175/1520-0469(1972)029<1076:AUTFTQ>2.0.CO;2), 1972.
- 985 Holton, J. R., and Tan, H.-C.: The influence of the equatorial quasi-biennial oscillation on the global circulation at 50 mb, *J. Atmos. Sci.*, 37, 2200–2208, [https://doi.org/10.1175/1520-0469\(1980\)037<2200:TIOTEQ>2.0.CO;2](https://doi.org/10.1175/1520-0469(1980)037<2200:TIOTEQ>2.0.CO;2), 1980.
- Holton, J.R., and Tan, H.-C.: The quasi-biennial oscillation in the Northern Hemisphere lower stratosphere, *J. Meteorol. Soc. Japan, Ser. II*, 60, 140–148, https://doi.org/10.2151/jmsj1965.60.1_140, 1982.
- Iza, M., Calvo, N., and Manzini, E.: The stratospheric pathway of La Niña, *J. Clim.*, 29, 8899–8914, <https://doi.org/10.1175/JCLI-D-16-0230.1>, 2016.
- 990 Kawatani, Y., Hamilton, K., Watanabe, S., Anstey, J. A., Richter, J. H., Butchart, N., Orbe, C., Osprey, S. M., Naoe, H., Elsbury, D., Chen, C.-C., Garcia-Serrano, J., Glanville, A., Kerzenmacher, T., Lott, F., Palmerio, F. M., Park, M., Serva, F., Taguchi, M., Versick, S., and Yoshida, K.: QBOi El Niño Southern Oscillation experiments: Overview of experiment design and ENSO modulation of the QBO, *EGUsphere [preprint]*, <https://doi.org/10.5194/egusphere-2024-3270>, Weather and
- 995 Climate Dynamics, (in revision).
- Kumar, V., Yoden, S., and Hitchman, M. H.: QBO and ENSO effects on the mean meridional circulation, polar vortex, subtropical westerly jets, and wave patterns during boreal winter, *J. Geophys. Res.*, 127, e2022JD036691. <https://doi.org/10.1029/2022JD036691>, 2022.
- Liess, S. and Geller, M. A.: On the relationship between QBO and distribution of tropical deep convection, *J. Geophys. Res.*, 117, <http://doi.wiley.com/10.1029/2011JD016317>, 2012.
- 1000 Lindzen, R. S., and Holton, J. R.: A theory of the quasi-biennial oscillation, *J. Atmos. Sci.*, 25, 1095–1107, [https://doi.org/10.1175/1520-0469\(1968\)025<1095:ATOTQB>2.0.CO;2](https://doi.org/10.1175/1520-0469(1968)025<1095:ATOTQB>2.0.CO;2), 1968.
- Lopez-Parages, J., Rodriguez-Fonseca, B., Mohino, E., and Losada, T.: Multidecadal modulation of ENSO teleconnection with Europe in late winter: Analysis of CMIP5 models, *J. Clim.*, 29, 8067–8081. <https://doi.org/10.1175/JCLI-D-15-0596.1>,
- 1005 2016.
- Lott, F., Rani, R., McLandress, C., Podglagen, A., Bushell, A., Bramberger, M., Lee, H.-K., Alexander, J., Anstey, J., Chun, H.-Y., Hertzog, A., Butchart, N., Kim, Y.-H., Kawatani, Y., Legras, B., Manzini, E., Naoe, H., Osprey, S., Plougonven, R., Pohlmann, H., Richter, J. H., Scinocca, J., García-Serrano, J., Serva, F., Stockdale, T., Versick, S., Watanabe, S., Yoshida, K.: Comparison between non-orographic gravity-wave parameterizations used in QBOi models and StratoLe 2 constant-level
- 1010 balloons. *Quart. J. Roy. Meteor. Soc.*, 1–16, <https://doi.org/10.1002/qj.4793>, 2024.



- Lu, H., Hitchman, M., Gray, L., Anstey, J., and Osprey, S.: On the role of Rossby wave breaking in the quasi-biennial modulation of the stratospheric polar vortex during boreal winter, *Q. J. Roy. Meteor. Soc.*, 146, 1939–1959, <https://doi.org/10.1002/qj.3775>, 2020.
- Ma, T., Chen, W., An, X., Garfinkel, C. I., and Cai, Q.: Nonlinear effects of the stratospheric Quasi-Biennial Oscillation and ENSO on the North Atlantic winter atmospheric circulation, *J. Geophys. Res.*, 128, e2023JD039537, <https://doi.org/10.1029/2023JD039537>, 2023.
- Marshall, A. G., Hendon, H. H., Son, S.-W., and Lim, Y.: Impact of the quasi-biennial oscillation on predictability of the Madden–Julian oscillation, *Clim. Dyn.*, 49, 1365–1377, <https://doi.org/10.1007/s00382-016-3392-0>, 2017.
- Martin, Z., Son, S.-W., Butler, A., Hendon, H., Kim, H., Sobel, A., Yoden, S., and Zhang, C.: The influence of the quasi-biennial oscillation on the Madden–Julian oscillation, *Nat. Rev. Earth Environ.*, 2, 477–489, <https://www.nature.com/articles/s43017-021-00173-9>, 2021.
- Naoe, H., Kobayashi, C., Kobayashi, S., Kosaka, Y., and Shibata, K.: Representation of Quasi-Biennial Oscillation in JRA-3Q, *J. Meteor. Soc. Japan*, (EOR), <https://doi.org/10.2151/jmsj.2025-012>, 2025.
- Naoe, H., and Shibata, K.: Equatorial quasi-biennial oscillation influence on northern winter extratropical circulation, *J. Geophys. Res.*, 115, D19102, doi:10.1029/2009JD012952, 2010.
- Naoe, H., and Yoshida, K.: Influence of quasi-biennial oscillation on the boreal winter extratropical stratosphere in QBOi experiments, *Q. J. Roy. Meteor. Soc.*, 145, 2755–2771, <https://doi.org/10.1002/qj.3591>, 2019.
- Palmeiro, F. M., García-Serrano, J., Ruggieri, P., Batte, L., and Gualdi, S.: On the influence of ENSO on sudden stratospheric warmings, *J. Geophys. Res.*, 128, e2022JD037607, <https://doi.org/10.1029/2022JD037607>, 2023.
- Palmeiro, F., Iza, M., Barriopedro, D., Calvo, N., Garcia-Herrera, R.: The complex behavior of El Niño winter 2015–2016. *Geophys. Res. Lett.*, 44, 2902–2910. <https://doi.org/10.1002/2017GL072920>, 2017.
- Park, C.-H., Son, S.-W., Lim, Y., and Choi, J.: Quasi-biennial oscillation-related surface air temperature change over the western North Pacific in late winter, *Int. J. Clim.*, 30, 4351–4359, <https://doi.org/10.1002/joc.7470>, 2022.
- Park, C.-H., Son, S.-W.: Relationship between the QBO and surface air temperature in the Korean Peninsula, *Atmos. Korean Meteor. Soc.*, 32, 39–49, (In Korean), <https://doi.org/10.14191/Atmos.2022.32.1>, 2022.
- Plumb, R. A., and McEwan, A. D.: The instability of a forced standing wave in a viscous stratified fluid: A laboratory analogue of the quasi-biennial oscillation, *J. Atmos. Sci.*, 35, 1827–1839, [https://doi.org/10.1175/1520-0469\(1978\)035<1827:TIOAFS>2.0.CO;2](https://doi.org/10.1175/1520-0469(1978)035<1827:TIOAFS>2.0.CO;2), 1978.
- Randall, D. A., Tziperman, E., Branson, M. D., Richter, J. H., Kang, W.: The QBO-MJO connection: A possible role for the SST and ENSO, *J. Clim.*, 36, 6515–6531, <https://doi.org/10.1175/JCLI-D-23-0031.1>, 2023.
- Rao, J., Garfinkel, C. I., and White, I. P.: How does the quasi-biennial oscillation affect the boreal winter tropospheric circulation in CMIP5/6 models?, *J. Clim.*, 33, 8975–8996, <https://doi.org/10.1175/JCLI-D-20-0024.1> 2020.
- Rao, J., and Ren, R.: A decomposition of ENSO’s impacts on the northern winter stratosphere: Competing effect of SST forcing in the tropical Indian Ocean, *Clim. Dyn.*, 46, 3689–3707, <https://doi.org/10.1007/s00382-015-2797-5>, 2016a.



- 1045 Rao, J., and Ren, R.: Asymmetry and nonlinearity of the influence of ENSO on the northern winter stratosphere: 1. Observations, *J. Geophys. Res.*, 121, 9000–9016, <https://doi.org/10.1002/2015JD024520>, 2016b.
- Rao, J., and Ren, R.: Asymmetry and nonlinearity of the influence of ENSO on the northern winter stratosphere: 2. Model study with WACCM, *J. Geophys. Res.*, 121, 9017–9032, <https://doi.org/10.1002/2015JD024520>, 2016c.
- Richter, J.H., Butchart, N., Kawatani, Y., Bushell, A. C., Holt, L., Serva, F., Anstey, J., Simpson, I R., Osprey, S., Hamilton, K., Braesicke, P., Cagnazzo, C., Chen, C.-C., Garcia, R. R., Gray, L. J., Kerzenmacher, T., Lott, F., McLandress, C., Naoe, H., Scinocca, J., Stockdale, T. N., Versick, S., Watanabe, S., Yoshida, K., and Yukimoto, S.: Response of the quasi-biennial oscillation to a warming climate in global climate models, *Q. J. Roy. Meteor. Soc.*, 148, 1490–1518, <https://doi.org/10.1002/qj.3749>, 2022.
- 1050 Richter, J. H., Matthes, K., Calvo, N., and Gray, L. J.: Influence of the Quasi-Biennial Oscillation and El Niño–Southern Oscillation on the frequency of sudden stratospheric warmings, *J. Geophys. Res.*, 116, D20111. <https://doi.org/10.1029/2011JD015757>, 2011.
- Rodrigo, M., García-Serrano, J., and Bladé, I.: Quasi-Biennial Oscillation influence on tropical convection and El Niño variability, *Geophys. Res. Lett.*, [submitted, 2024GL112854], (in revision).
- Ruzmaikin, A., Feynman, J., Jiang, X., Yung, Y. L.: Extratropical signature of the quasi-biennial oscillation, *J. Geophys. Res.*, 110, D11111, <https://doi.org/10.1029/2004JD005382>, 2005.
- 1060 Scaife, A. A., Athanassiadou, M., Andrews, M., Arribas, A., Baldwin, M., Dunstone, N., Knight, J., MacLachlan, C., Manzini, E., Muller, W. A., Pohlmann, H., Smith, D., Stockdale, T., and Williams A.: Predictability of the quasi-biennial oscillation and its northern winter teleconnection on seasonal to decadal timescales, *Geophys. Res. Lett.* 41, 1752–1758, <https://doi.org/10.1029/2004JD005382>, 2014.
- 1065 Serva, F., Anstey, J. A., Bushell, A. C., Butchart, N., Cagnazzo, C., Gray, L., Kawatani, Y., Osprey, S. M., Richter, J. H., and Simpson, I. R.: The impact of the QBO on the region of the tropical tropopause in QBOi models: present-day simulations, *Q. J. Roy. Meteor. Soc.*, 148, 1945–1964, <https://doi.org/10.1002/qj.4287>, 2022.
- Simpson, I. R., Blackburn, M., and Haigh, J. D.: The role of eddies in driving the tropospheric response to stratospheric heating perturbations, *J. Atmos. Sci.*, 66, 1347–1365, <https://doi.org/10.1175/2008JAS2758.1>, 2009.
- 1070 Son, S.-W., Lim, Y., Yoo, C., Hendon, H. H., and Kim, J.: Stratospheric control of the Madden-Julian oscillation, *J. Clim.*, 30, 1909–1922, <http://journals.ametsoc.org/doi/10.1175/JCLI574>, 2017.
- Song, K. and Son, S.-W.: Revisiting the ENSO–SSW relationship, *J. Clim.*, 31, 2133–2143, <https://doi.org/10.1175/JCLI-D-17-0078.1>, 2018.
- Taguchi, M.: Wave driving in the tropical lower stratosphere as simulated by WACCM. Part II: ENSO-induced changes for northern winter, *J. Atmos. Sci.*, 67, 543–555. <https://doi.org/10.1175/2009JAS3144.1>, 2010a.
- 1075 Taguchi, M.: Observed connection of the stratospheric Quasi-Biennial Oscillation with El Niño–Southern Oscillation in radiosonde data, *J. Geophys. Res.*, 115, D18120. <https://doi.org/10.1029/2010JD014325>, 2010b.



- Taguchi, M., and Hartmann, D. L.: Increased occurrence of stratospheric sudden warmings during El Niño simulated by WACCM, *J. Clim.*, 19, 324–332, doi:10.1175/JCLI3655.1, 2006.
- 1080 Tegtmeier, S., Anstey, J., Davis, S., Ivanciu, I., Jia, Y., McPhee, D., Kedzierski, R. P.: Zonal asymmetry of the QBO temperature signal in the tropical tropopause region. *Geophys. Res. Lett.*, 47, e2020GL089533. <https://doi.org/10.1029/2020GL089533>, 2020.
- Tyrrell, N. L., Koskentausta, J. M., and Karpechko, A. Y.: Sudden stratospheric warmings during El Niño and La Niña: sensitivity to atmospheric model biases, *Weather Clim. Dynam.*, 3, 45–58, <https://doi.org/10.5194/wcd-3-45-2022>, 2022.
- 1085 Van Loon, H., and Labitzke, K.: The Southern Oscillation. Part V: The anomalies in the lower stratosphere of the Northern Hemisphere in winter and a comparison with the Quasi-Biennial Oscillation, *Mon. Weather Rev.*, 115, 357–369, [https://doi.org/10.1175/1520-0493\(1987\)115<0357:TSOPVT>2.0.CO;2](https://doi.org/10.1175/1520-0493(1987)115<0357:TSOPVT>2.0.CO;2), 1987.
- Walsh, A., Screen, J. A., Scaife, A. A., and Smith, D. M.: Non-linear response of the extratropics to tropical climate variability, *Geophys. Res. Lett.*, 49, e2022GL100416, <https://doi.org/10.1029/2022GL100416>, 2022.
- 1090 Wang, J., Kim, H.-M., and Chang, E. K. M.: Interannual modulation of Northern Hemisphere winter storm tracks by the QBO, *Geophys. Res. Lett.*, 45, 2786–2794, <http://doi.wiley.com/10.1002/4842017GL076929>, 2018.
- Watson, P. A. G., and Gray, L. J.: How does the quasi-biennial oscillation affect the stratospheric polar vortex?, *J. Atmos. Sci.*, 71, 391–409, <https://doi.org/10.1175/JAS-D-13-096.1>, 2014.
- Wei, K., Chen, W., and Huang, R.: Association of tropical Pacific sea surface temperatures with the stratospheric Holton-Tan oscillation in the Northern Hemisphere winter, *Geophys. Res. Lett.*, 34, L16814, doi:10.1029/2007GL030478, 2007.
- 1095 Weinberger, I., Garfinkel, C. I., White, I. P., Oman, L. D., Aquila, V., and Lim, Y.-K.: The salience of nonlinearities in the boreal winter response to ENSO: Arctic stratosphere and Europe, *Clim. Dyn.*, 53, 4591–4610, <https://doi.org/10.1007/s00382-019-04805-1>, 2019.
- White, I. P., Lu, H., Mitchell, N. J., and Phillips, T.: Dynamical response to the QBO in the northern winter stratosphere: signatures in wave forcing and eddy fluxes of potential vorticity, *J. Atmos. Sci.*, 72, 4487–4507, <https://doi.org/10.1175/JAS-D-14-0358.1>, 2015.
- 1100 Yamazaki, K., Nakamura, T., Ukita, J., and Hoshi, K.: A tropospheric pathway of the stratospheric quasibiennial oscillation (QBO) impact on the boreal winter polar vortex, *Atmos. Chem. Phys.*, 20, 5111–5127, <https://doi.org/10.5194/acp-20-5111-2020>, 2020.
- 1105 Yoo, C., and Son, S.-W.: Modulation of the boreal wintertime Madden-Julian oscillation by the stratospheric quasi-biennial oscillation, *Geophys. Res. Lett.*, 43, 1392–1398, doi:10.1002/2016GL067762, 2016.
- Yukimoto, S., Kawai, H., Koshiro, T., Oshima, N., Yoshida, K., Urakawa, S., Tsujino, H., Deushi, M., Tanaka, T.Y., Hosaka, M., Yabu, S., Yoshimura, H., Shindo, E., Mizuta, R., Obata, A., Adachi, Y. and Ishii, M.: The MRI Earth System Model version 2.0, MRI-ESM2.0: description and basic evaluation of the physical component, *J. Meteor. Soc. Japan*, 97, 931–965, <https://doi.org/10.2151/jmsj.2019-051>, 2019.
- 1110