

Sensitivity of Simulated Ammonia Fluxes in Rocky Mountain National Park to Measurement Time Resolution and Meteorological Inputs

Lillian E. Naimie¹, Da Pan¹, Amy P. Sullivan¹, John T. Walker², Aleksandra Djurkovic², Bret A. Schichtel^{3,4}, Jeffrey L. Collett, Jr.¹

¹Department of Atmospheric Science, Colorado State University, Fort Collins, CO 80523, USA

²United States Environmental Protection Agency, Office of Research and Development, Durham, NC 27709, USA

³Cooperative Institute for Research in the Atmosphere, Colorado State University, Fort Collins, CO 80523, USA

⁴US National Park Service, Air Resource Division, Lakewood, CO 80225-0287, USA

Correspondence to: Jeffrey L. Collett, Jr. (collett@colostate.edu)

Abstract. Gaseous ammonia (NH₃) is an important precursor for secondary aerosol formation and contributes to reactive nitrogen deposition. NH₃ dry deposition is ~~poorly~~~~rarely~~ quantified due to the complex bidirectional nature of NH₃ atmosphere-surface exchange and lack of high time-resolution in situ NH₃ concentration and meteorological measurements. To better quantify NH₃ dry deposition, measurements of NH₃ were made above a subalpine forest canopy in Rocky Mountain National Park (RMNP) and used ~~with~~ in situ micrometeorology to simulate bidirectional fluxes. NH₃ dry deposition was largest during the summer, with ~~47~~~~48~~% of annual net NH₃ dry deposition occurring in June, July, and August. ~~A net annual dry deposition estimated using measured 30-minute NH₃ concentrations and in situ meteorological data, accounted for 6% of total RMNP reactive inorganic N deposition.~~ Because in situ, high-time resolution concentration and meteorological data are often unavailable, the ~~impacts~~~~impact~~ on estimated deposition from ~~utilizing~~ more commonly available ~~biweekly NH₃ measurements and ERA5 meteorology were~~~~input data was~~ evaluated. Fluxes simulated with biweekly NH₃ concentrations, commonly available from NH₃ monitoring networks, underestimated NH₃ dry deposition by ~~45~~~~29~~%. These fluxes were strongly correlated with 30-minute fluxes integrated to a biweekly basis ($R^2 = 0.8889$) indicating that a correction factor could be applied to mitigate the observed bias. Application of an average NH₃ diel concentration pattern to the biweekly NH₃ concentration data removed the observed low bias. Annual NH₃ dry deposition from fluxes simulated with reanalysis meteorological inputs exceeded simulations using in situ meteorology measurements by ~~a factor of 2.59~~~~%~~.

1. Introduction

Gaseous ammonia (NH₃) is an important atmospheric constituent, with effects on atmospheric chemistry and the nitrogen cycle. Atmospheric deposition of reactive nitrogen (N_r) is linked to nitrogen oxides (NO_x) and NH₃ emissions. Emissions of NO_x and NH₃ have many potential fates including chemical transformation, dry deposition, particle formation, and wet deposition. Anthropogenic emissions of NO_x and NH₃ are produced predominantly by combustion and agriculture, respectively

(Walker et al., 2019a), although there are also NH₃ emissions from traffic, wastewater treatment, and wildfires (Tomsche et al., 2023; Walker et al., 2019b). Due to increased food demand and industrialization, anthropogenic N_r has been increasing annually (Galloway et al., 2008; Kanakidou et al., 2016). Excess reactive nitrogen deposition has well-documented adverse effects on ecosystem health including ~~lake~~-eutrophication, soil acidification, decreased biodiversity, and increased N in freshwater bodies (Baron, 2006; Bobbink, 1991; Boot et al., 2016; Holtgrieve et al., 2011; Pan et al., 2021; Zhan et al., 2017). As a result of effective NO_x emission controls, the balance of N_r wet deposition across the US has shifted from oxidized N-dominated to reduced N-dominated, and dry deposition of NH₃ at times dominates total N_r deposition (Driscoll et al., 2024; Li et al., 2016, Walker et al., 2019a). The National Emission Inventory (NEI) indicates that US NO_x emissions were reduced by 46% between 2013 and 2023, while NH₃ emissions increased by 13% (US EPA, 2023).

Critical loads, deposition levels below which harmful effects are not expected to occur, have been estimated for many ecosystems (e.g. Bowman et al., 2012; Schwede and Lear, 2014). In Rocky Mountain National Park (RMNP), a critical load of 1.5 kg N ha⁻¹ yr⁻¹, based on wet deposition of NO₃⁻ and NH₄⁺, has been established to avoid adverse effects on the ecosystem (Baron, 2006). The pre-industrial nitrogen load has been estimated at 0.2 kg N ha⁻¹ yr⁻¹ while the current ~~wet~~ deposition rate is as high as 3.65 kg N ha⁻¹ yr⁻¹, approximately 15x the natural background and significantly higher than the critical load (Benedict et al., 2013a; Burns, 2003; CDPHE, 2007). Although the RMNP N_r critical load only considers wet deposition of NO₃⁻ and NH₄⁺, dry deposition can also contribute significantly to total N_r deposition. NH₃ dry deposition in RMNP was estimated to be the third largest contributor to total N_r deposition, accounting for 18% of N_r deposition from November 2008 to November 2009 (Benedict et al., 2013a).

NH₃ dry deposition, however, remains a highly uncertain component of N_r deposition, and fluxes are rarely measured (Walker et al., 2019b). Previous studies in RMNP have estimated NH₃ dry deposition using unidirectional inferential models, where the NH₃ deposition velocity (V_d) was approximated as 70% of the HNO₃ deposition velocity (Beem et al., 2010; Benedict et al., 2013a; Benedict et al., 2013b) and NH₃ emission from the surface was ignored. In reality, NH₃ exchange between the atmosphere and surface is bidirectional, including deposition to and emission from the surface (Sutton et al., 1995). Several models have been developed to simulate the bidirectional exchange of NH₃ with the surface (Massad et al., 2010; Pleim et al., 2013; Zhang et al., 2010). Key model inputs include micrometeorology, soil and vegetation parameters, and atmospheric concentrations. In practice, fluxes can change quickly and even reverse direction with changing environmental conditions. Gaseous NH₃ is challenging and expensive to measure at high time resolution; lower-cost weekly or biweekly passive diffusion-based sampler measurements are more commonly utilized for long-term monitoring (Butler et al., 2016; Hu et al., 2021; Li et al., 2016; Schiferl et al., 2016). Previous efforts have used these low-cost measurements to quantify NH₃ dry deposition (Shen et al., 2016; Tanner et al., 2022; Walker et al., 2008). Detailed, high-time resolution meteorological observations at the location of interest are also desired when estimating dry deposition. Due to the frequent unavailability of such data, reanalysis meteorological data is often used as a substitute (Schrader et al., 2018; Wichink Kruit et al., 2012).

Schrader et al. (2018) investigated the impact of low time-resolution NH₃ concentrations on modeled fluxes. They found that using monthly NH₃ concentrations underestimates total NH₃ dry deposition. However, due to a linear relationship between

65 simulations using monthly NH_3 concentrations and those using hourly NH_3 concentrations, they were able to generate a site-specific correction to compensate for the use of low time-resolution concentration data. Simulations were done using a simplified parameterization of the bidirectional exchange model described in Massad et al. (2010) and the NH_3 concentrations were simulated using the LOTOS-EUROS model (Hendricks et al., 2016).

Understanding and managing these biases could unveil opportunities to estimate NH_3 deposition when high-time resolution, in situ concentration, and meteorological observations are unavailable. Using high-time resolution NH_3 concentration measurements, we provide the first estimate of NH_3 annual dry deposition to an RMNP forest canopy using a bidirectional exchange model driven by high-time resolution NH_3 concentration data and in situ micro-meteorological measurements. We use in situ data collected in RMNP to determine if site-specific correction factors suggested by Schrader et al. (2018) apply to real-world observations and whether correction factors can be employed to reduce biases associated with NH_3 simulations using lower-cost, low-time resolution NH_3 measurements such as those available from the U.S. Ammonia Monitoring network (AMoN) (Puchalski et al., 2011). We also tested if an average NH_3 diel pattern could be applied to reduce these biases and, if so, the length of measurements necessary to adequately describe the diel pattern. Finally, we examine biases introduced by substituting reanalysis meteorological data for high-time resolution in situ measurements.

2 Data and ~~methods~~Methods

80 2.1 Site ~~location~~Location

Study observations were collected in RMNP in northern Colorado. The park, established to preserve the natural landscape, including montane, subalpine, and alpine ecosystems, is predominantly above 3000 m where ecosystems developed under nutrient-deprived conditions and are therefore sensitive to excess inputs of nitrogen. Nitrogen deposition has been a historical problem in RMNP; with diatom changes documented starting in the 1950s and more recent effects including eutrophication and changes to plant species (Baron, 2006; Baron et al., 2000; Korb and Ranker, 2001).

The area east of RMNP (Fig. 1) includes a large urban corridor and extensive agricultural activity in the plains. The Front Range urban corridor, spanning from Denver to Fort Collins, is a major source of nitrogen oxide emissions (Benedict et al., 2013b). The northeast plains of Colorado are predominantly agricultural and include major sources of NH_3 emissions from both animal feeding operations and crop production. The spatial pattern seen for feedlots is broadly consistent with the spatial distribution of other agricultural activities. Pan et al. (2021) found that 40% of summertime dry deposition of NH_3 in RMNP was associated with transport from agricultural regions to the east.




Figure 1. A map of the study region. Animal units are shown as the number of permitted animals as of 2017, scaled by an animal unit factor relative to the species. Elevation data is from the US Geological Survey Global Multi-resolution Terrain Elevation Data 2010 (GMTED2010) at 7.5-arc-second spatial resolution (available at: <https://earthexplorer.usgs.gov/>).

Data was collected at two adjacent locations for this study, both near the base of Longs Peak in Rocky Mountain National Park: a National Ecological Observatory Network (NEON) tower site (40.275903, -105.54596) and a nearby National Park Service shelter (~500 m north of the NEON tower), from September 2021 through August 2022. The study location, denoted

100 with a star in Fig. 1, is 2750 m above sea level. The tower is surrounded by lower montane forest, comprised of predominantly evergreen needleleaf species, including ponderosa pine, juniper, and Douglas fir. There are also groves of quaking aspen located in the region. Meteorological transport to the site is generally bimodal. Prevailing downslope transport from the northwest occurs generally overnight and during the cooler months, when ammonia concentrations are typically low. The mountain-plains circulation generates daytime upslope transport, bringing air masses from the plains east of the park up into
105 RMNP. This pattern strengthens during warmer seasons. Periods of synoptically forced sustained upslope transport are also common, especially during spring and autumn (Gebhart et al., 2011). Downslope and upslope transport patterns are not due west and east at the study site because of ~~channelling~~channeling by local topography.

At RMNP, a diel pattern in ambient NH₃ concentrations has commonly been observed in past measurements. This pattern is primarily driven by changes in transport patterns that carry NH₃ emissions to the park (Benedict et al., 2013b; Juncosa
110 Calahorrano et al., 2024) and, sometimes, modified by changes in the atmosphere-surface exchange of NH₃, especially during NH₃ uptake and emission from dew formation and evaporation (Wentworth et al., 2016).

2.2 Micrometeorological ~~measurements~~Measurements

2.2.1 in situ ~~micrometeorology~~Micrometeorology

Meteorological and soil data were accessed from the RMNP NEON flux tower. The mean canopy height in the area
115 surrounding the tower is 19 m. ~~Temperature (mean=6 C), relative humidity (mean=40%), and annual days of precipitation are highly variable at the site due to its high elevation. Snowfall typically occurs between October and May.~~

Meteorological data accessed from the NEON site includes wind vectors, ~~friction~~frictional velocity, Obukhov length, soil temperature, short wave radiation, relative humidity, air density, air pressure, and air temperature above the tree canopy. Soil temperature was taken as the average across 5 collection sites within 200 m of the flux tower. ~~Leaf area index (LAI) is estimated at the site using remotely sensed data at 1 km resolution. The square kilometer of leaf area index values surrounding the tower site is shown in the supplementary information. A mean value of 0.8 was estimated using the landscape surrounding the site. The sensitivity to LAI can also be found in the supplementary information.~~ Additional information about each of the reported
120 NEON datasets can be found in the Site Management and Event Reporting documentation (available at: <https://doi.org/10.48443/9p2t-hj77>).

125 NEON meteorological data contained gaps due to power outages and scheduled instrument maintenance. Across the year of data, the gaps comprised 5.8% of the data (1021 data points). To quantify the annual deposition of NH₃ in RMNP, these gaps were filled using the average diel pattern of fluxes during the current biweekly NH₃ sampling period.

2.2.2 Reanalysis ~~meteorology data~~Meteorology Data

Detailed meteorological and soil data are not available at many locations where NH₃ dry deposition is of interest. Reanalysis
130 data, which combine short-range weather forecasts with assimilated observations, are a common source of meteorological data

Formatted: Subscript
Formatted: Subscript
Formatted: Subscript
Formatted: Subscript

that can be used in the absence of local observations. To probe the impact of using reanalysis data in place of in situ observations, a set of bidirectional flux simulations was conducted using ERA5 hourly reanalysis data (Hersbach et al., 2020). ERA5 hourly reanalysis data has a spatial resolution of 0.25° , or approximately 31 km. The parameters used from the ERA5 data are as follows: air temperature, air pressure, dewpoint temperature, turbulent surface stress, moisture flux, sensible heat flux, friction velocity, standard deviation of filtered subgrid orography, solar radiation, and soil temperature. Obukhov length (L) is not given in the ERA5 dataset and was calculated using [Eq. \(equation-5.7c\)](#) from Stull (1988), shown below. [Obukhov Length is the characteristic length scale of the atmosphere and is calculated from ERA5 data using surface sensible heat and moisture fluxes.](#)

$$L = \frac{-\overline{\theta'_v} u_*^3}{k g (w' \theta'_v)_s} = \frac{k g (t_{vst})}{T_s - T_v} \quad (1)$$

where k is the von Karman constant, g is gravitational acceleration, $\overline{\theta'_v}$ is the mean turbulent temperature scale, T_v is the virtual temperature near the surface, $w' \theta'_v$ is the surface flux of virtual potential temperature, and u_* is the friction velocity.

2.3 NH₃ data

2.3.1 Biweekly NH₃ measurements

Biweekly NH₃ ambient air concentration was measured using Radiello (<https://radiello.com/>) passive diffusion samplers. The Radiello sampling system includes a diffusive body and adsorbing cartridge, which is coated with phosphoric acid. NH₃ (g) diffuses across the exterior diffusive body and is collected on the adsorbing cartridge as ammonium (NH₄⁺) over two weeks. Collected ammonia (as NH₄⁺) is extracted from the cartridge into deionized water and analyzed using ion chromatography (IC) (Li et al., 2016). NH₃ passive samples were collected in duplicate ($\sigma = \pm 0.25 \mu\text{g m}^{-3}$) on top of the NEON tower (25.35 m-agl). [Across the study period, there were 27 sampling periods.](#) Passive NH₃ sampling methods have been shown to have a low bias when compared with other sampling methods, including University Research Glassware Denuders and Picarro Cavity Ringdown spectroscopy methods (Pan et al., 2020; Puchalski et al., 2011).

2.3.2 High temporal resolution NH₃ measurements

NH₃ (g) air concentration was also measured using an ion mobility spectrometer (IMS). Ion mobility spectroscopy separates ionized molecules based on their mobility through a carrier gas, under the influence of an electric field. The instrument used was the AirSentry II Point-of-Use IMS from Particle Measuring Systems (Boulder, CO). The instrument was in the [National Park Service \(NPS\)](#) shelter (located at 40.278129, -105.545635) 500 meters north of the NEON site with an inlet located approximately 2 m above natural grassland. The sampling inlet was $\frac{1}{4}$ " Teflon tubing, heated to 40 C to reduce NH₃ loss to the sampling tube. Inlet length was kept as short as possible to further prevent NH₃ loss. Particles were removed by a fiber

Formatted: Superscript

filter at the tip of the inlet. Due to the high altitude of the site location, the instrument was zeroed to account for pressure differences upon installation. Multi-point calibrations were conducted at the beginning and end of sampling. Calibration was confirmed using a known concentration ammonia gas sample split between the instrument and a phosphoric acid-coated denuder where the NH_3 collected by the denuder is extracted into deionized water and analyzed using ion chromatography.

Zero measurements were made periodically by overflowing the inlet with ultra-high purity clean air. The AirSentry samples at a 30-second frequency. During the study the AirSentry collected 919,000 data points. The limit of detection is 70 pptv.

2.3.3 NH_3 ~~data preparation~~Data Preparation

To investigate the effect of NH_3 (g) sampling time resolution, bidirectional fluxes were simulated with concentration data at: (i) 30-minute frequency (30-minute NH_3), (ii) with the 2-week integrated passive NH_3 (Biweekly Passive NH_3), and lastly with an average diel profile applied to each day within the 2-week passive period (Average Diel Pattern NH_3). The three NH_3 data products are shown in Fig. 2.




Figure 2. Three NH_3 concentration data sets are shown for the entire study period. The biweekly-passive- NH_3 is the mean of the NEON-tower-top-duplicate-measurements. 30-minute- NH_3 data was generated using the AirSentry- NH_3 concentration-sealed-to-match-biweekly-passive- NH_3 for each passive sampling period. Biweekly- NH_3 with diel profile applied was generated by applying an average diel profile from the AirSentry- NH_3 to each day of the biweekly-passive measurement. The two-week average across each concentration data product is the same.

The 30-minute NH_3 concentration data is generated using a combination of data from the AirSentry NH_3 located at the NPS shelter and passive NH_3 samples collected on the NEON tower. Data gaps, due to power outages and regular maintenance, were filled using the average diel pattern across the year of data collection. Data gaps accounted for about 3000 out of 5.8% of more than 900,000 points across the total data across the study period. To generate a 30-minute NH_3 data set above the tree canopy, the data was divided into biweekly periods which match the passive NH_3 collection periods. The average concentration from the AirSentry across each period was then scaled to match the biweekly passive NH_3 concentration. This preserves the

temporal variability of NH₃ concentrations while ensuring that the average air concentration across the sampling period is consistent with the passive NH₃ measurements atop the NEON tower which can differ from those above the adjacent grassland where the Air Sentry measurements are made.

The biweekly passive NH₃ with diel profile applied is generated using the annual average diel pattern of NH₃ from the AirSentry data. To determine if there are systematic differences between the NH₃ diel pattern at the two sites, raw and scaled AirSentry concentrations were compared to 4- and 6-hour University Research Glassware denuder measurements taken on the NEON tower. The NH₃ concentrations were well correlated between sites. This comparison is shown in Fig (S1). Each day of the biweekly passive period is assigned the average diel pattern, then the biweekly mean is scaled to match the biweekly passive concentration. This dataset was generated to investigate if the inclusion of a simple diel profile was sufficient to correct for the bias in bidirectional fluxes created by using low time-resolution NH₃ concentrations as shown by Schrader et al. (2018). These three concentration data sets will be used for bidirectional flux simulations of NH₃. For the rest of this work, the three NH₃ data sets will be referred to using the following nomenclature.

- 30-minute NH₃:** NH₃ concentration data at 30-minute frequency
- Biweekly NH₃:** Biweekly Passive NH₃ concentration data
- Average Diel Pattern NH₃:** Passive NH₃ concentration scaled using an average diel profile from the 30-minute NH₃ dataset

2.4 Additional ~~measurements~~Measurements

2.4.1 Wet ~~deposition data~~Deposition Data

Wet deposition data was obtained from the National Trends Network (NTN) (National Atmospheric Deposition Program, 2022) site at Beaver Meadows in RMNP ('CO19': located at 40.3639°N, -105.5810°E). The Beaver Meadows site location, at 2477 m elevation and located approximately 10 km north of the CASTNET site, is shown in Fig. 1.

2.4.2 Additional ~~gas~~Gas and ~~particle measurements~~Partiele Measurements

Additional air concentration data was obtained from the U.S. EPA Clean Air Status and Trends Network (CASTNET) site at the NPS shelter ('ROM206': located at 40.278129, -105.545635). Weekly filter pack concentrations of nitric acid (HNO₃) and sulfur dioxide (SO₂) were used to calculate the acid ratio (Eq. 10~~equation-8~~) in the bidirectional exchange simulations of NH₃ (U.S. EPA, 2024a).


Weekly dry deposition of HNO₃, NO₃⁻, and NH₄⁺ was generated by CASTNET (US EPA, 2024b) using the weekly filter pack concentrations and historical values of deposition velocity from the U.S. EPA Multi-Layer Model (MLM) (Meyers et al., 1998). The generation of deposition velocities was discontinued in 2019. Bowker et al. (2011) found that using historical values of deposition velocity from the U.S. EPA Multi-Layer Model did not significantly bias the annual mean of deposition.

One approach to estimating NH_3 deposition is to estimate the deposition velocity (V_d) as a fixed fraction (70%) of the deposition velocity of HNO_3 . This approach has been historically used to estimate the dry deposition velocity of NH_3 in RMNP (Beem et al., 2010; Benedict et al., 2013a; Benedict et al., 2013b).

$$V_d(\text{NH}_3) = 0.7 * V_d(\text{HNO}_3), \tag{2}$$

2.5 Bidirectional flux modellingFlux-Modeling of NH_3

Bidirectional NH_3 fluxes are simulated across the study period using the dry deposition inferential model described in Massad et al. (2010). The simulation framework (Fig. 3) accounts for the bidirectional nature of NH_3 fluxes and allows for deposition and emission. The model determines if the flux will be negative (deposition) or positive (emission) based on the relationship between the atmospheric concentration (χ_a) at a given reference height (z) and the canopy compensation point (χ_c). Canopy compensation point depends on the stomata resistance, cuticle resistance, and stomata compensation point.



225

χ_a	Atmospheric ammonia concentration	χ_c	Canopy compensation point
χ_{z0}	Surface compensation point (Z ₀)	χ_g	Ground layer/Soil compensation point
R_a	Aerodynamic resistance	χ_s	Stomata compensation point
R_b	Laminar boundary layer resistance	f_t	Total flux
R_{bg}	Ground laminar boundary layer resistance	f_g	Ground flux
R_w	Cuticle resistance	f_s	Stomata flux
R_{st}	Stomata resistance	f_w	Cuticle flux
R_{ac}	Aerodynamic resistance in the canopy		

Formatted Table

230

Figure 3. Dry deposition inferential model proposed in Massad et al. (2010). The table describes each model element. Arrows next to each flux show the allowed flux directions of the given pathway.

235

The relationship between resistances and compensation points is shown in Fig. 3. Aerodynamic (R_a) and laminar boundary layer resistance (R_b) capture the effects of turbulent and diffusive transfer from the atmosphere to the surface, respectively. R_a was calculated according to Thom (1975), where z is reference height (25.35 m), d is the displacement height (7.15 m), and z_0 is the roughness length (1.65 m). The stability functions are Ψ_H and Ψ_M for scalars and momentum, respectively, are empirical relationships dependent on Obukhov length (Thom 1975). Displacement and roughness lengths were provided from the RMNP NEON Tower (NEON, 2023).

240

$$R_a = (k \cdot u^*)^{-1} \cdot \left(\ln \left(\frac{z-d}{z_0} \right) - \Psi_H + \Psi_M \right), \quad (3)$$

R_b is modeled as described in Xiu and Pleim (2001), where γ_{air} is the kinematic diffusivity of air, and D_{NH_3} is the diffusivity of NH_3 .

$$R_b = \frac{5}{u^*} \cdot \left(\frac{\gamma_{air}}{D_{NH_3}} \right)^{2/3}, \quad (4)$$

245

In-canopy aerodynamic resistance (R_{ac}) captures the aerodynamic resistance from within the canopy layer and is the sum of aerodynamic resistance within the canopy (R_{ac}) and ground boundary layer resistance (R_{bg}). R_{ac} was calculated based on Nemitz et al. (2001) using Eq. (5) where α is a height dependent constant calculated using Eq. 16 and Eq. 17 from equations 15-17 of Massad et al. (2010).

$$R_{ac(d+z_0)} = \frac{\alpha(d+z_0)}{u^*} \quad (5)$$

250 ~~(2010)~~ Ground boundary layer resistance (R_{bg}) is based on Nemitz et al. (2001), where u is the wind speed at tower top
255 (25 canopy height ($h=11$ m)).

$$R_{bg} = \left(\frac{\gamma_{air}}{D_{NH_3}} - \ln \left(\frac{D_{NH_3}}{k \cdot \frac{u}{20} \cdot 0.1 \cdot h} \right) \right) \cdot \frac{1}{k \cdot \frac{u}{20}}, \quad (65)$$

Stomata resistance (R_{st}) captures the diffusion of NH_3 through plant stomata and is calculated as a minimum value related to the plant type proposed by Hicks et al. (1987). Further parameterization proposed by Nemitz et al. (2001) was used here to calculate R_{st} , where SR ($W\ m^{-2}$) is the solar radiation. The minimum value for R_{st} ($225\ s\ m^{-1}$) was determined using Table 1 of Zhang et al. (2003).

$$R_{st} = \min \left\{ 5000\ (s\ m^{-1}), 225\ (s\ m^{-1}) \cdot \left(1 + \left(\frac{180}{SR} \right) \right) \right\}, \quad (76)$$

Cuticle resistance (R_w) was calculated according to the proposed parameterization; for forest ecosystems ~~of~~ predominantly composed of Douglas Fir, as described in Massad et al. (2010). When relative humidity (RH) is below 100%, Eq. (8) equation
260 7 is used and when RH exceeds or is equal to 100%, Eq. (9) equation 8 is used.

$$R_w = 31.5 \cdot \frac{1}{AR} \cdot e^{(0.0318(100-RH))}, \quad (87)$$

$$R_w = \frac{31.5}{AR}, \quad (98)$$

In both equations, AR is the acid ratio which is calculated using the molar ratio of acids and bases in the atmosphere. The calculated acid ratio had a mean value of 1.3, a minimum of 0.22, and a maximum of 11.6. Acid ratios were the largest in the
265 winter months.

$$AR = \frac{2 \cdot [SO_2] + [HNO_3]}{[NH_3]}, \quad (109)$$

For this study period, the acid ratio was calculated using weekly CASTNET measurements of SO_2 and HNO_3 paired with our measurements of NH_3 .

~~Stomatal and ground~~ Stomata compensation points were calculated according to Massad et al. (2010). In the stomata compensation point (Eq. 11 equation 10), Γ_{st} is the emission potential of the stomata and is approximated as 294 based on vegetation samples from the area surrounding the NEON Tower. The sampling methods and determination of this value can be found in the supplementary information. Emission potentials are ratios that describe the potential for surface emission. Massad et al. (2010).
270

$$\chi_{st} = \frac{2.7457 \cdot 10^{15}}{T} \cdot e^{\left(\frac{-10378}{T} \right)} \cdot \Gamma_{st}, \quad (1140)$$

Soil compensation point was calculated according to Eq. (equations 3) through Eq. (5) of Stratton et al. (2018). In Eq. (12), equation 11, TAN is the concentration of total ~~ammoniacal~~ ammoniacal N (the sum of NH_3 and NH_4^+) in the soil aqueous phase ($mg\ kg^{-1}$), K_H is the Henry constant, and K_a is the equilibrium constant. TAN was estimated at 109.6 $mg\ kg^{-1}$ based on soil measurements in RMNP from Stratton et al. (2018). NH_3 flux simulations are very sensitive to TAN value. The supplementary information includes a test of the sensitivity of the flux results to TAN values within one standard deviation for
280 the measurements taken by Stratton et al. (2018).

$$\chi_g = \frac{K_H}{1+(10^{-pH})/(K_a)} \cdot TAN, \quad (12+1)$$

K_H and K_a were predicted using Eq. (13)equations-12 and Eq. (14)13 based on the models of Montes et al. (2009), where T is temperature.

$$K_H = \left(\frac{0.2138}{T} \right) \cdot 10^{(6.123-1825/T)}, \quad (13+2)$$

$$K_a = 10^{(0.05-2788/T)}, \quad (14+3)$$

Canopy compensation point, Eq. (15)-(equation-14 below,) was calculated using Eq. (equation-12) from Massad et al. (2010), where χ_a is the atmospheric NH_3 concentration.-

$$\chi_c = \frac{\chi_a \cdot (R_a \cdot R_b)^{-1} + \chi_{st} \cdot [(R_a \cdot R_{st})^{-1} + (R_b \cdot R_{st})^{-1} + (R_g \cdot R_{st})^{-1}] + \chi_g \cdot (R_b \cdot R_g)^{-1}}{(R_a \cdot R_b)^{-1} + (R_a \cdot R_{st})^{-1} + (R_a \cdot R_w)^{-1} + (R_b \cdot R_g)^{-1} + (R_b \cdot R_{st})^{-1} + (R_b \cdot R_w)^{-1} + (R_g \cdot R_{st})^{-1} + (R_g \cdot R_w)^{-1}},$$

$$(15) \chi_c = \frac{\chi_a \cdot (R_a \cdot R_b)^{-1} + \chi_{st} \cdot [(R_a \cdot R_{st})^{-1} + (R_b \cdot R_{st})^{-1} + (R_g \cdot R_{st})^{-1}] + \chi_g \cdot (R_b \cdot R_g)^{-1}}{(R_a \cdot R_b)^{-1} + (R_a \cdot R_{st})^{-1} + (R_a \cdot R_w)^{-1} + (R_b \cdot R_g)^{-1} + (R_b \cdot R_{st})^{-1} + (R_b \cdot R_w)^{-1} + (R_g \cdot R_{st})^{-1} + (R_g \cdot R_w)^{-1}}, \quad (14)$$

Compensation point at the a displacementreference height (d) above the roughness length (Z_0) is calculated using Eq. (16)equation-15 below as proposed in Massad et al. (2010). The surfacereference height is the same as the height at which NH_3 measurements were taken. The reference height compensation point (χ_{z0}) takes all other compensation points and resistances into account.

$$\chi_{z0} = \frac{\left(\frac{\chi_a}{R_a} + \frac{\chi_g}{R_g} + \frac{\chi_c}{R_b} \right)}{\left(\frac{1}{R_a} + \frac{1}{R_g} + \frac{1}{R_b} \right)}, \quad (16+5)$$

Finally, the total flux was calculated following Eq. (17equation-16) (Massad et al., 2010). NH_3 flux is defined in this framework as a difference between the roughnessreference height compensation point and the NH_3 concentration at that height, scaled by the aerodynamic resistance.

$$F_{NH_3} = \frac{\chi_{z0} - \chi_a}{R_a},$$

$$(17) \frac{\chi_{z0} - \chi_a}{R_a}, \quad (16)$$


Total exchange flux (F_{NH_3}) from the dry deposition inferential model gives the direction and magnitude of NH_3 fluxes.

3. Results and Discussion

3.1 Simulated bidirectional exchangeBidirectional Exchange of NH_3

Bidirectional fluxes were simulated using the 30-minute NH_3 concentration data set and in situ meteorological data as inputs to the Massad et al. (2010) model, described above. NH_3 concentration, surfacereference height compensation point, and fluxes have a strong seasonal cycle in RMNP (see Fig. 4). NH_3 flux direction is determined by the relative magnitudes of the NH_3

concentration and the ~~surface~~~~reference~~~~height~~ compensation point (Fig. 4a.). When NH₃ concentration exceeds the compensation point, NH₃ is deposited to the surface (a negative flux value). Both NH₃ concentrations and deposition fluxes
310 tend to be greatest during the summer, with ~~47~~~~48~~% of NH₃ modeled annual dry deposition occurring during June, July, and August. NH₃ fluxes also had the largest variability in the summer. Deposition in the spring closely follows, with ~~43~~~~33~~% of NH₃ modeled annual dry deposition occurring during March, April, and May. During all seasons there are periods of net emission from the surface (Fig. 4b.). ~~The largest periods of net emission occur in the spring. Daily~~~~These daily~~ NH₃ emission fluxes are most common in the winter ~~although where~~ they ~~are~~ ~~typically an order of magnitude~~ smaller than ~~typical~~ deposition
315 fluxes in the spring and summer.






Figure 4. Daily mean values of: (a.) Daily mean NH_3 concentration and surfacereference-height compensation point, and (b.) NH_3 flux.

Total modeled NH_3 flux can be broken down into stomata, ground, and cuticle fluxes. Figure 5 shows the distribution of simulated NH_3 fluxes for each of these components.

Deposition is driven primarily by stomata and cuticle fluxes, while ground emission fluxes are sometimes observed. Winter periods of net emission (see Fig. 4b.) are driven by the ground flux. One potential limitation of the model used for simulations is that it does not consider snow cover on the ground, which could alter winter fluxes in RMNP.






Figure 5. Total NH_3 simulated fluxes are separated into their component fluxes (stomata, ground, and cuticle). Simulated fluxes are shown for the entire study period. Boxes show the 25th and 75th percentile, and whiskers are determined at 1.5 times the interquartile range.

NH_3 concentrations at RMNP are impacted by emission and transport patterns, which can both increase daytime NH_3 concentrations. NH_3 emissions from agricultural sources have a strong diel pattern driven by volatilization during warmer

daytime temperatures. At RMNP, transport from these regions is driven on many days by the mountain-plains circulation, which begins in the late morning and transports polluted air masses westward and upslope to the park (Gebhart et al., 2011).

Previous studies have demonstrated that the upslope transport from sources in the Front Range has impacts on deposition and air concentrations in RMNP (Benedict et al., 2018; Pan et al., 2021). During this study, the largest χ_a values are also observed during upslope transport from source regions in the CO Front Range. These source regions likely disproportionately contribute to NH_3 dry deposition because the difference between χ_a and χ_{zo} drives the sign and magnitude of the NH_3 flux. On mornings following overnight dew formation, local volatilization from evaporating dew has also been shown to increase morning NH_3 concentrations (Wentworth et al., 2016). This phenomenon was observed in RMNP and corresponds to the increase in the NH_3 diel pattern around 10:00 observed in Fig. 6a. One limitation of the bidirectional flux model used is that NH_3 uptake and emission from dew are not simulated. NH_3 concentration, compensation point, and simulated fluxes each have a strong diel pattern, which peaks during the middle of the day (see Fig. 6). The peak value typically occurs close to 13:00. The soil temperature diel pattern contributes to a higher surface-reference-height compensation point during the middle of the day. The annual cycle of soil temperature also contributes to the higher surface-reference-height compensation points observed in summer. Although both NH_3 concentration and compensation point peak during the mid-day, we also observe peak fluxes during the middle of the day indicating that the influence of the diel pattern of NH_3 concentration is stronger than that from the compensation point diel pattern.






Figure 6. Diel pattern of: (a.) NH_3 concentration, (b.) simulated ~~surface~~~~reference-height~~ compensation point, and (c.) NH_3 fluxes are shown for the full study period in RMNP. Boxes show the 25th and 75th ~~percentile~~~~percentile~~, and whiskers are determined at 1.5 times the interquartile range.

To understand the relative importance of NH_3 deposition in RMNP, NH_3 flux simulation results are combined with NADP/NTN wet deposition fluxes and dry deposition fluxes for particulate ammonium (NH_4^+) and nitrate (NO_3^-) and gaseous HNO_3 derived from CASTNET concentration observations and MLM deposition velocities, to construct an updated seasonal and annual budget of inorganic N deposition at RMNP. This N_r deposition budget for all measured inorganic species is shown in Fig. 7a. Due to the lack of current measurements, wet and dry deposition of organic nitrogen are not included. Benedict et al. (2013b) reported annual organic nitrogen wet deposition of $0.6 \text{ kg N ha}^{-1} \text{ yr}^{-1}$ during their 2008-2009 study. NH_3 dry deposition is the net surface flux from the simulations using 30-minute NH_3 concentration. The inorganic annual N_r deposition budget totals $3.4 \text{ kg N ha}^{-1} \text{ yr}^{-1}$, with the largest contributions coming from NH_4^+ wet deposition ($1.34 \text{ kg N ha}^{-1} \text{ yr}^{-1}$), NH_3 net dry deposition ($0.1217 \text{ kg N ha}^{-1} \text{ yr}^{-1}$), NO_3^- wet deposition ($0.71 \text{ kg N ha}^{-1} \text{ yr}^{-1}$), and HNO_3 dry deposition ($0.33 \text{ kg N ha}^{-1} \text{ yr}^{-1}$). Overall, reduced N_r deposition comprises 59.60% of the total inorganic N deposition to RMNP. NH_3 dry deposition comprises 4.6% of total inorganic N_r deposition. Simulated NH_3 dry deposition ($0.1117 \text{ kg N ha}^{-1} \text{ yr}^{-1}$) is smaller than the value estimated by as the NH_3 concentration multiplied by 0.7 times the HNO_3 deposition velocity by Benedict et al. (2013b) during their 2008-2009 study ($0.66 \text{ kg N ha}^{-1} \text{ yr}^{-1}$). The previous value estimated NH_3 dry deposition velocity by scaling the HNO_3 dry deposition velocity by 0.7, instead of simulating the bidirectional exchange of NH_3 .




Figure 7. Reactive nitrogen deposition is shown for all species with measured concentrations or deposition for the full year of study. Wet deposition data is from the NADP NTN site at Beaver Meadows. NH_3 dry deposition is modeled using the bidirectional framework from Massad et. al (2010) and 30-minute NH_3 concentration data. Dry deposition of HNO_3 (g), NH_4^+ (p), and NO_3^- (p) are calculated from the nearby CASTNET site concentration data and deposition velocities from the U.S. EPA MLM. Panel (a.) has the annual deposition of all measured species. Panel (b.) has deposition of all measured N_r species grouped by month. ~~Reduced-N species are green. Oxidized-N species are blue.~~ Only one period of wet deposition was collected by the NADP NTN site during November 2021.

Speciated monthly dry deposition is plotted in Fig. 7b to probe the seasonality of N_r deposition in RMNP. Net dry deposition of NH_3 was largest during July and August. Total inorganic N_r deposition peaked during May, due to increased wet deposition. ~~Reduced~~ For all months except November and January, reduced N_r deposition exceeded oxidized N_r deposition in October, December, February, March, April, May, July, and August. Excluding November, where only one period of wet deposition

was recorded by the NADP NTN site, reduced N_e deposition had, with a fractional contribution ranging from 43.47 to 74.75%. In November and January, net NH_3 emission was estimated from the surface.

3.2 Impacts of biweekly NH_3 concentration data on simulated fluxes

The use of low time-resolution NH_3 concentrations for flux simulations can produce a low bias compared with fluxes simulated using higher time-resolution NH_3 concentrations. Here, we follow a similar method to that described by Schrader et al. (2018) and demonstrate that a site-specific correction can be generated to account for the bias introduced by lower time resolution NH_3 concentration data. Our methods differ from Schrader et al. (2018) in 3 major ways: (i) in situ data is used for both the higher frequency, 30-minute NH_3 concentration, and meteorology, (ii) biweekly passive NH_3 data is used instead of monthly NH_3 data, and (iii) Massad et al. (2010) is used as described instead of using a simplified parameterization. The results of the 30-minute NH_3 and Biweekly NH_3 bidirectional NH_3 flux simulations are compared to generate a site-specific factor to correct for any low bias in the lower time resolution flux calculations. Simulated fluxes at biweekly time resolution (Fig. 8) using the two NH_3 concentration data sets are well correlated ($R^2 = 0.8889$) and the NH_3 flux simulation using biweekly integrated NH_3 data can be corrected to match the control flux simulation using a linear fit (slope: 1.0307, y-intercept: -1.689468). As noted above, RMNP has few two-week periods of net NH_3 emission, and the efficacy of this method should be confirmed at a location with more extensive periods of net NH_3 emission. In particular, NH_3 fluxes above managed agricultural land could differ significantly from the pattern observed in RMNP. This study also focused on fluxes above a forest canopy, and results could differ for grassland ecosystems, which also occur in RMNP. To determine the efficacy in other locations, future investigations should select several sites with different land surface types and NH_3 concentrations to make biweekly and high-time resolution measurements for a year.




Figure 8. Bidirectional NH₃ flux simulated at 30-minute resolution is plotted for 30-minute NH₃ concentration data and biweekly integrated NH₃ concentration data. Fluxes are given as net flux over a two-week period. The least squares linear regression is plotted for the data.

Considering the net flux of NH₃ across the full study period, using the best available time resolution of 30 minutes, we find a total annual net NH₃ dry deposition flux of 0.1147 kg N ha⁻¹ yr⁻¹ (Fig. 9). The estimated NH₃ dry deposition drops by 4529% to 0.0612 kg N ha⁻¹ yr⁻¹ using biweekly vs. 30-min NH₃ concentration measurements. The annual NH₃ dry deposition flux increases to 0.78 kg N ha⁻¹ yr⁻¹ when simulating fluxes in a deposition-only unidirectional framework where the NH₃ deposition velocity is scaled as 0.7 times the nitric acid deposition velocity (generated by the US EPA MLM), an approach previously used for RMNP N deposition budgets (Beem et al., 2010; Benedict et al., 2013a; Benedict et al., 2013b).






Figure 9. Annual NH₃ dry deposition at the NEON Flux Tower in RMNP is shown for three bidirectional simulations using three sets of NH₃ concentration data (30-minute NH₃, Biweekly NH₃, and Average Diel Pattern NH₃) and one unidirectional simulation. Each simulation was run at 30-minute time steps with meteorological parameters from the NEON Flux Tower. The unidirectional NH₃ flux is calculated using simulation uses biweekly NH₃ concentrations, NH₃ and deposition velocities are calculated as 0.7 times the HNO₃ deposition velocity from based on the U.S. EPA MLM.

Bidirectional flux simulations using biweekly NH₃ data with an average diel pattern of NH₃ yield the same annual NH₃ dry deposition flux as the simulations run using 30-minute NH₃ concentration. This indicates that capturing daily variability in NH₃ concentration profiles is not critical to accurately simulating the annual NH₃ flux. Application of an annual averaged diel pattern misses the highest NH₃ concentrations (Fig. 10), however, across a full year of data the diel pattern effectively captures the net surface flux. Despite the scatter in Fig. 10a., fluxes simulated with an average diel pattern NH₃ data set are well

Formatted: Subscript

425 correlated with simulations using 30-minute NH_3 concentrations ($R^2=0.596$) and have a fit close to unity. The daily mean fluxes (Fig. 10b and Fig. 10c) of each simulation have similar seasonal patterns, with periods of net emission and deposition aligned between simulations.



430 **Figure 10.** NH_3 fluxes simulated with 30-minute NH_3 concentrations and annual average diel pattern NH_3 concentrations are shown for the full year of data. Panel (a.) directly compares 30-minute simulated fluxes for each data set. Panels (b.) and (c.) show the daily mean fluxes for simulations with 30-minute NH_3 concentration and average diel pattern NH_3 concentration respectively.


At RMNP, there is a large daily variability in concentration due especially to changes in upslope transport. When an air mass arrives from the Colorado Front Range and NE Colorado, NH_3 concentrations rise significantly due to the large emission sources upwind. For the comparison shown in Fig. 10, the diel pattern was determined using a full year of NH_3 concentration data. Fluxes were also simulated using diel patterns determined with only a month of data, to probe the necessary length of measurements to generate an effective diel pattern. Annual deposition from all flux simulations using a monthly diel pattern fell within 2% of the annual deposition using the annual average diel pattern. Therefore, in RMNP, one month of 30-minute

Formatted: Normal

measurements appears sufficient to generate a diel pattern thatwhich will effectively correct the net NH₃ surface flux. Other
440 locations may have larger and/or more complex variability in NH₃ diel pattern and may require longer periods of data collection
to establish an NH₃ diel pattern.

3.3 Impacts of reanalysis meteorological dataReanalysis Meteorological Data on simulatedSimulated NH₃ fluxes

Dry deposition inferential models require several meteorological and soil parameters, which may not be readily available for
many locations of interest. Reanalysis data can provide meteorological inputs for locations where required in situ
445 meteorological and soil measurements are unavailable. To examine the impact on flux simulation accuracy resulting from this
substitution at RMNP, the same simulations of NH₃ bidirectional fluxes were run using ERA5 meteorology and soil data. 30-
minute NH₃ simulations run with reanalysis data inputs are well correlated ($R^2 = 0.7680$) with 30-minute NH₃ simulations run
with in situ data inputs (see Fig. 11+1a) but overestimate the annual NH₃ deposition flux by a factor of 2.59%. From Fig.
11,11a, we find that the use of ERA5 reanalysis data in the simulation of NH₃ bidirectional fluxes introduces a low bias to the
450 flux magnitude in RMNP compared to in situ meteorological data, for both positive (emission) and negative (deposition)
fluxes. However, because the decrease to deposition fluxes is smaller than the decrease to emission fluxes, we observe anThe
annual overestimation from simulations using ERA5 is due in large part to missing periods of surface emission. Based on this
result, ERA5 reanalysis data should not be used to estimate NH₃ fluxes before additional sites and data have been considered
using in situ data.



455 **Figure 11. Bidirectional NH₃ flux simulated with ERA5 meteorology and NEON meteorology at 30-minute resolution using the 30-minute NH₃ concentration. The least squares linear regression is plotted for the data in red.**

The low bias for fluxes simulated using ERA5 reanalysis data is investigated further to explore what parametersparameter
differences influence this bias. Net NH₃ fluxes are simulated using Eq. (17Equation (12)), which relies on χ_{20} , NH₃

concentration, and aerodynamic resistance (R_a). We find that the simulations using reanalysis data generate surface reference height compensation points (χ_{z0}) which agree well with the simulations using that used in situ measurements (Slope=1.030.94, $R^2=0.98$).




Figure 12. Aerodynamic resistances are shown for simulations using in situ meteorological data from the NEON flux tower and reanalysis meteorological data from ERA5. The diel patterns are shown in panels (a.) and (b.) respectively. Panel (c.) directly compares simulated R_a values using NEON in situ and ERA5 reanalysis data.

Although the general diel pattern of R_a is well captured using reanalysis data, R_a magnitudes differ substantially between the two simulations (Fig. 12a and 12b), with the largest difference occurring overnight. Maximum R_a values from the reanalysis simulations are greater than an order of magnitude larger than those derived using in situ meteorology. A and a comparison of the two data sets shows (Fig. 12c) a typical enhancement of approximately a factor of four. Increased R_a values result in lower simulated NH_3 fluxes. The R_a bias is likely driven by differences in the friction velocity (u_*) and Obukhov Length (L), which are used to calculate simulate R_a . ERA5 data underestimates u_* by a factor of 5 when compared with the in situ NEON data (slope = 0.2). The in situ NEON data also sets a minimum u_* value (0.2 m s^{-1}), while the ERA5 data allows u_* values below 0.2 m s^{-1} . Comparisons of all meteorological parameters used can be found in the supplementary information. This discrepancy

Formatted: Subscript
Formatted: Subscript

in modeled R_a may be due to the gridded nature of reanalysis data, which represents a large area of variable land types and complex topography using only a single value (Hogrefe et al., 2023). ~~Obukhov Length is the characteristic length scale of the atmosphere and is calculated from ERA5 data using surface sensible heat and moisture fluxes.~~ Previous work has identified heat and moisture fluxes as large areas of uncertainty in ERA5 Reanalysis (Kong et al., 2022; Mayer et al., 2022). Two case studies were conducted to probe the relative importance of u_* and Obukhov Length. The case studies are described in the supplementary information. Differences in R_a were impacted by both u_* and L , accounting for 23% and 10%, respectively, of the discrepancy between in situ and ERA5 flux simulations. Comparisons of all meteorological parameters used can be found in the Supplement.

4. Conclusion

Fluxes of NH_3 (g) are best simulated using a bidirectional model, which uses rapidly changing meteorology paired with air concentrations and soil parameters to infer flux direction and magnitude. We use a bidirectional NH_3 flux model to simulate a year of NH_3 fluxes above a subalpine forest ecosystem in Rocky Mountain National Park. The net NH_3 dry deposition to the ecosystem is estimated at $0.1147 \text{ kg N ha}^{-1} \text{ yr}^{-1}$, comprising 46% of total inorganic reactive nitrogen deposition. This is significantly lower than previous estimates for RMNP, which did not consider the bi-directional nature of the exchange. Due to the observed low bias in passive NH_3 observations and the sensitivity of simulations to NH_3 concentrations, this is likely a low bound. The sensitivity of NH_3 flux modelling to χ_a was tested by scaling the input concentration by 9% to account for the error discussed in Puchalski et al. (2011). This resulted in an annual reduced-N deposition increase of 47%, indicating the importance of accurate NH_3 measurements for flux modelling. Additionally, since the highest NH_3 concentrations occur during upslope events, the sources contributing to these events likely have a disproportionate effect on inputs (wet and dry) econstitutes 60% of total N₂ deposition. One limitation of this model is the exclusion of snow cover, which could significantly change NH_3 fluxes in the winter, when RMNP has frequent snow events. Future works should investigate NH_3 fluxes above snow cover to better simulate the exchange of NH_3 in regions with snow.

Due to the cost and technical challenges of making continual, high-time resolution NH_3 concentration measurements, there is growing interest in using integrated biweekly passive NH_3 measurements, such as those from the NADP AMoN network, for flux simulations. Here, we establish that a site-specific correction can be used to correct a bias introduced by using lower time resolution passive NH_3 measurements over the studied forest canopy in RMNP. We also establish that an average NH_3 diel pattern can be used to interpolate 30-minute NH_3 concentration and correct for the bias introduced by passive NH_3 measurements. In RMNP, a month of measurements proved sufficient to determine the diel pattern used for flux simulations. The correction factor and diel pattern, however, likely vary by location due to differences in ecosystem characteristics and factors influencing NH_3 concentrations. Due to the potential regional differences and changes associated with land surface type, additional sites should be studied to assess the impact of measurement time resolution on NH_3 flux simulations. To

understand the seasonal variability in diel pattern and efficacy of diel pattern application for flux simulations, measurements should be conducted for a full year.

Local micrometeorological and soil measurements are also frequently unavailable, making the use of reanalysis data a desirable alternative for NH₃ flux simulations. In our location, the use of reanalysis data adds a bias that leads to overestimates of net NH₃ deposition. We found it was possible to apply a correction to address this bias, but this factor likely varies by location, in particular over different land surface types within a reanalysis grid cell. Future studies should explore the relationship between in situ measurements and reanalysis products above different land surface types, ~~above~~-varied topography, and in different regions.

Data Availability

The ammonia concentration data used in the study will be published after the manuscript is accepted. The NEON flux tower eddy covariance data bundle is available at: <https://data.neonscience.org/data-products/DP4.00200.001>. ERA5 reanalysis data is available at: <https://www.ecmwf.int/en/forecasts/dataset/ecmwf-reanalysis-v5>. CASTNET data are available at: <https://www.epa.gov/castnet/download-data>. NTN data are available at: <https://nadp.slh.wisc.edu/networks/national-trends-network/>.

Author Contributions

JC, BS, DP, and JW designed the measurement campaign. LN, AS, and DP made and processed Rocky Mountain National Park measurements. DP developed the model code. LN designed and ran the bidirectional exchange simulations. LN prepared the manuscript with contributions from all co-authors.

Competing interest

The authors declare that they have no conflict of interest.

Disclaimer

The results contain modified Copernicus Climate Change Service information 2020. Neither the European Commission nor ECMWF is responsible for any use that may be made of the Copernicus information or data it contains. The results make use of data collected by the CASTNET program from the U.S. Environmental Protection Agency. The views expressed are of the authors and do not necessarily reflect those of the U.S. EPA or any other organizations that the data used was obtained from.

Acknowledgments

The authors would also like to thank the NEON team for their support in collecting biweekly passive NH₃ data. The authors thank Nikolas Tafoya for his assistance in collecting measurements on the NEON Tower.

535 Financial support

This work was supported by the U.S. Environmental Protection Agency Regional Applied Research Efforts Program, Project #2237 and the National Park Service. This material is based in part upon work supported by the National Science Foundation through the National Ecological Observatory Network Program. The NEON Program is operated under a cooperative agreement by Batelle.

540 References

- Appel, K. W., Bash, J. O., Fahey, K. M., Foley, K. M., Gilliam, R. C., Hogrefe, C., Hutzell, W. T., Kang, D., Mathur, R., Murphy, B. N., Napelenok, S. L., Nolte, C. G., Pleim, J. E., Pouliot, G. A., Pye, H. O. T., Ran, L., Roselle, S. J., Sarwar, G., Schwede, D. B., Sidi, F. I., Spero, T. L., and Wong, D. C. (2021) The Community Multiscale Air Quality Model Version 5.3 and 5.3.1: System Updates and Evaluation, *Geosci. Model Dev.* doi: <https://doi.org/10.5194/gmd-14-2867-2021>
- 545 Baron, J. S. (2006), Hindcasting Nitrogen Deposition To Determine An Ecological Critical Load, *Ecological Applications*, 16(2), 433-439, doi:[https://doi.org/10.1890/1051-0761\(2006\)016\[0433:HNDTDA\]2.0.CO;2](https://doi.org/10.1890/1051-0761(2006)016[0433:HNDTDA]2.0.CO;2).
- Baron, J. S., H. M. Rueth, A. M. Wolfe, K. R. Nydick, E. J. Allstott, J. T. Minear, and B. Moraska (2000), Ecosystem Responses to Nitrogen Deposition in the Colorado Front Range, *Ecosystems*, 3(4), 352-368, doi:10.1007/s100210000032.
- Beem, K. B., et al. (2010), Deposition of reactive nitrogen during the Rocky Mountain Airborne Nitrogen and Sulfur (RoMANS) study, *Environmental Pollution*, 158(3), 862-872, doi:<https://doi.org/10.1016/j.envpol.2009.09.023>.
- 550 Benedict, K. B., C. M. Carrico, S. M. Kreidenweis, B. Schichtel, W. C. Malm, and J. L. Collett Jr (2013a), A seasonal nitrogen deposition budget for Rocky Mountain National Park, *Ecological Applications*, 23(5), 1156-1169, doi:<https://doi.org/10.1890/12-1624.1>.
- Benedict, K. B., D. Day, F. M. Schwandner, S. M. Kreidenweis, B. Schichtel, W. C. Malm, and J. L. Collett (2013b),
- 555 Observations of atmospheric reactive nitrogen species in Rocky Mountain National Park and across northern Colorado, *Atmospheric Environment*, 64, 66-76, doi:<https://doi.org/10.1016/j.atmosenv.2012.08.066>.
- Benedict, K. B., A. J. Prenni, A. P. Sullivan, A. R. Evansoski-Cole, E. V. Fischer, S. Callahan, B. C. Sive, Y. Zhou, B. A. Schichtel, and J. L. Collett, Jr. (2018), Impact of Front Range sources on reactive nitrogen concentrations and deposition in Rocky Mountain National Park, doi:<https://doi.org/10.7717/peerj.4759>.

560 Benish, S. E. and Bash, J. O. and Foley, K. M. and Appel, K. W. and Hogrefe, C. and Gilliam, R. and Pouliot, G. (2022) Long-term regional trends of nitrogen and sulfur deposition in the United States from 2002 to 2017, *Atmospheric Chemistry and Physics*, doi:<https://doi.org/10.5194/acp-22-12749-2022>

Bobbink, R. (1991), Effects of Nutrient Enrichment in Dutch Chalk Grassland, *Journal of Applied Ecology*, 28(1), 28-41, doi:[10.2307/2404111](https://doi.org/10.2307/2404111).

565 Boot, C. M., E. K. Hall, K. Denef, and J. S. Baron (2016), Long-term reactive nitrogen loading alters soil carbon and microbial community properties in a subalpine forest ecosystem, *Soil Biology and Biochemistry*, 92, 211-220, doi:<https://doi.org/10.1016/j.soilbio.2015.10.002>.

Bowker, G.E., Schwede, D.B., Lear, G.G. et al. Quality Assurance Decisions with Air Models: A Case Study of Imputation of Missing Input Data Using EPA's Multi-layer Model. *Water Air Soil Pollut* 222, 391-402 (2011).

570 <https://doi.org/10.1007/s11270-011-0832-7>

Bowman, W. D., J. Murgel, T. Blett, and E. Porter (2012), Nitrogen critical loads for alpine vegetation and soils in Rocky Mountain National Park, *Journal of Environmental Management*, 103, 165-171, doi:<https://doi.org/10.1016/j.jenvman.2012.03.002>.

Butler, T., F. Vermeulen, C. M. Lehmann, G. E. Likens, and M. Puchalski (2016), Increasing ammonia concentration trends

575 in large regions of the USA derived from the NADP/AMoN network, *Atmospheric Environment*, 146, 132-140, doi:<https://doi.org/10.1016/j.atmosenv.2016.06.033>.

Burns, D. A. (2003), Atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern Wyoming—a review and new analysis of past study results, *Atmospheric Environment*, 37(7), 921-932, doi:[https://doi.org/10.1016/S1352-2310\(02\)00993-7](https://doi.org/10.1016/S1352-2310(02)00993-7).

580 Colorado Department of Public Health and Environment (CDPHE), Air Pollution Control Division (2007), Nitrogen Deposition Reduction Plan. Colorado Department of Public Health and Environment. available at: <https://www.colorado.gov/pacific/cdphe/rocky-mountain-national-park-nitrogen-reduction-plan>

Driscoll, C., Milford, J. B., Henze, D. K., & Bell, M. D. (2024), Atmospheric reduced nitrogen: Sources, transformations, effects, and management, *Journal of the Air & Waste Management Association*, 74(6), 362-415,

585 doi:<https://doi.org/10.1080/10962247.2024.2342765>

Galloway, J. N., A. R. Townsend, J. W. Erisman, M. Bekunda, Z. Cai, J. R. Freney, L. A. Martinelli, S. P. Seitzinger, and M. A. Sutton (2008), Transformation of the Nitrogen Cycle: Recent Trends, Questions, and Potential Solutions, *Science*, 320(5878), 889-892, doi:[10.1126/science.1136674](https://doi.org/10.1126/science.1136674).

Gebhart, K. A., B. A. Schichtel, W. C. Malm, M. G. Barna, M. A. Rodriguez, and J. L. Collett (2011), Back-trajectory-based

590 source apportionment of airborne sulfur and nitrogen concentrations at Rocky Mountain National Park, Colorado, USA, *Atmospheric Environment*, 45(3), 621-633, doi:<https://doi.org/10.1016/j.atmosenv.2010.10.035>.

- Hendriks, C., R. Kranenburg, J. J. P. Kuenen, B. Van den Bril, V. Verguts, and M. Schaap (2016), Ammonia emission time profiles based on manure transport data improve ammonia modelling across north western Europe, *Atmospheric Environment*, 131, 83-96, doi:<https://doi.org/10.1016/j.atmosenv.2016.01.043>.
- 595 Hersbach, H. et al. (2020). The ERA5 Global Reanalysis. *Quarterly Journal of the Royal Meteorological Society*, 146, 1999-2049. <https://doi.org/10.1002/qj.3803>
- Hicks, B. B., D. D. Baldocchi, T. P. Meyers, R. P. Hosker, and D. R. Matt (1987), A preliminary multiple resistance routine for deriving dry deposition velocities from measured quantities, *Water, Air, and Soil Pollution*, 36(3), 311-330, doi:10.1007/BF00229675.
- 600 Hogrefe, C., Bash, J. O., Pleim, J. E., Schwede, D. B., Gilliam, R. C., Foley, K. M., Appel, K. W., and R. Mathur (2023), An analysis of CMAQ gas-phase dry deposition over North America through grid-scale and land-use-specific diagnostics in the context of AQMEII4, *Atmos. Chem. Phys.*, 23, 8119–8147, <https://doi.org/10.5194/acp-23-8119-2023>.
- Holtgrieve, G. W., et al. (2011), A Coherent Signature of Anthropogenic Nitrogen Deposition to Remote Watersheds of the Northern Hemisphere, *Science*, 334(6062), 1545-1548, doi:10.1126/science.1212267.
- 605 Hu, C., Griffiths, T. J., Frie, A., Baker, J. M., Wood, J. D., Millet, D. B., et al. (2021), A multiyear constraint on ammonia emissions and deposition within the US Corn Belt. *Geophysical Research Letters*, 48, e2020GL090865. <https://doi.org/10.1029/2020GL090865>
- Juncosa Calahorrano, J. F., Sullivan, A. P., Pollack, I. B., Roscioli, J. R., McCabe, M. E., Steinmann, K. M., Caulton, D. R., Li, E., Pierce, J. R., Naimie, L. E., Pan, D., Collett, J. L., Jr, & Fischer, E. V. (2024), Anatomy of Summertime Upslope Events in Northeastern Colorado: Ammonia (NH₃) Transport to the Rocky Mountains. *Environmental science & technology*, 58(38), 16922–16930. <https://doi.org/10.1021/acs.est.3c10902>
- 610 Kanakidou, M., S. Myriokefalitakis, N. Daskalakis, G. Fanourgakis, A. Nenes, A. R. Baker, K. Tsigaridis, and N. Mihalopoulos (2016), Past, Present, and Future Atmospheric Nitrogen Deposition, *Journal of the Atmospheric Sciences*, 73(5), 2039-2047, doi:<https://doi.org/10.1175/JAS-D-15-0278.1>.
- 615 Korb, J. E., and T. A. Ranker (2001), Changes in stand composition and structure between 1981 and 1996 in four Front Range plant communities in Colorado, *Plant Ecology*, 157(1), 1-11, doi:10.1023/A:1013772220131.
- Li, Y., B. A. Schichtel, J. T. Walker, D. B. Schwede, X. Chen, C. M. B. Lehmann, M. A. Puchalski, D. A. Gay, and J. L. Collett (2016), Increasing importance of deposition of reduced nitrogen in the United States, *Proceedings of the National Academy of Sciences*, 113(21), 5874-5879, doi:10.1073/pnas.1525736113.
- 620 Massad, R. S., E. Nemitz, and M. A. Sutton (2010), Review and parameterisation of bi-directional ammonia exchange between vegetation and the atmosphere, *Atmos. Chem. Phys.*, 10(21), 10359-10386, doi:10.5194/acp-10-10359-2010.
- Meyers, T. P., Finkelstein, P., Clarke, J., Ellestad, T. G., & Sims, P. F. (1998). A multilayer model for inferring dry deposition using standard meteorological measurements. *Journal of Geophysical Research*, 103(D17), 22645–22661.

Formatted: Check spelling and grammar, Subscript

Montes, F., C. A. Rotz, and H. Chaoui (2009). Process Modeling of Ammonia Volatilization from Ammonium Solution and
625 Manure Surfaces: A Review with Recommended Models. *Transactions of the ASABE*, 52(5), 1707-1720,
doi:<https://doi.org/10.13031/2013.29133>.

National Atmospheric Deposition Program (NRSP-3). 2022. NADP Program Office, Wisconsin State Laboratory of Hygiene,
465 Henry Mall, Madison, WI 53706.

NEON (National Ecological Observatory Network). Site management and event reporting (DP1.10111.001), RELEASE-2023.
630 <https://doi.org/10.48443/9p2t-hj77>. Dataset accessed from [https://data.neonscience.org/data-](https://data.neonscience.org/data-products/DP1.10111.001/RELEASE-2024)
products/DP1.10111.001/RELEASE-2024 on September 05, 2023.

Nemitz, E., C. Milford, and M. A. Sutton (2001), A two-layer canopy compensation point model for describing bi-directional
biosphere-atmosphere exchange of ammonia, *Quarterly Journal of the Royal Meteorological Society*, 127(573), 815-833,
doi:<https://doi.org/10.1002/qj.49712757306>.

635 Pan, D., et al. (2021), Ammonia Dry Deposition in an Alpine Ecosystem Traced to Agricultural Emission Hotspots,
Environmental Science & Technology, 55(12), 7776-7785, doi:10.1021/acs.est.0c05749.

Pan, Y., et al. (2020), Systematic low bias of passive samplers in characterizing nitrogen isotopic composition of atmospheric
ammonia, *Atmospheric Research*, 243, 105018, doi:<https://doi.org/10.1016/j.atmosres.2020.105018>.

Pleim, J. E., J. O. Bash, J. T. Walker, and E. J. Cooter (2013), Development and evaluation of an ammonia bidirectional flux
640 parameterization for air quality models, *Journal of Geophysical Research: Atmospheres*, 118(9), 3794-3806,
doi:<https://doi.org/10.1002/jgrd.50262>.

Puchalski, M. A., M. E. Sather, J. T. Walker, C. M. B. Lehmann, D. A. Gay, J. Mathew, and W. P. Robarge (2011), Passive
ammonia monitoring in the United States: Comparing three different sampling devices, *Journal of Environmental Monitoring*,
13(11), 3156-3167, doi:10.1039/C1EM10553A.

645 Schiferl, L. D., Heald, C. L., Van Damme, M., Clarisse, L., Clerbaux, C., Coheur, P.-F., Nowak, J. B., Neuman, J. A., Herndon,
S. C., Roscioli, J. R., and Eilerman, S. J. (2016), Interannual variability of ammonia concentrations over the United States:
sources and implications, *Atmos. Chem. Phys.*, 16, 12305–12328, doi:<https://doi.org/10.5194/acp-16-12305-2016>.

Schrader, F., M. Schaap, U. Zöll, R. Kranenburg, and C. Brümmer (2018), The hidden cost of using low-resolution
concentration data in the estimation of NH₃ dry deposition fluxes, *Scientific Reports*, 8(1), 969, doi:10.1038/s41598-017-
650 18021-6.

Schwede, D. B., and G. G. Lear (2014), A novel hybrid approach for estimating total deposition in the United States,
Atmospheric Environment, 92, 207-220, doi:<https://doi.org/10.1016/j.atmosenv.2014.04.008>.

Shen, J., D. Chen, M. Bai, J. Sun, T. Coates, S. K. Lam, and Y. Li (2016), Ammonia deposition in the neighbourhood of an
intensive cattle feedlot in Victoria, Australia, *Scientific Reports*, 6(1), 32793, doi:10.1038/srep32793.

655 Stratton, J. J., J. Ham, and T. Borch (2018), Ammonia Emissions from Subalpine Forest and Mountain Grassland Soils in
Rocky Mountain National Park, *Journal of Environmental Quality*, 47(4), 778-785,
doi:<https://doi.org/10.2134/jeq2018.01.0023>.

Formatted: Check spelling and grammar, Subscript

Stull R. B. (1988) An Introduction to Boundary Layer Meteorology; Kluwer Academic, Dordrecht, doi: <https://doi.org/10.1007/978-94-009-3027-8>.

660 Sutton, M. A., J. K. Schjorring, G. P. Wyers, J. H. Duyzer, P. Ineson, D. S. Powlson, D. Fowler, D. S. Jenkinson, J. L. Monteith, and M. H. Unsworth (1995), Plant—atmosphere exchange of ammonia, *Philosophical Transactions of the Royal Society of London. Series A: Physical and Engineering Sciences*, 351(1696), 261-278, doi:doi:10.1098/rsta.1995.0033.

Tanner, E., N. Buchmann, and W. Eugster (2022), Agricultural ammonia dry deposition and total nitrogen deposition to a Swiss mire, *Agriculture, Ecosystems & Environment*, 336, 108009, doi:<https://doi.org/10.1016/j.agee.2022.108009>.

665 Thom, A. S. (1975), *Momentum, mass and heat exchange of plant communities. Vol.1*, Edited by J.L. Monteith, London.

Tomsche, L., et al. (2023), Measurement report: Emission factors of NH₃ and NH_x for wildfires and agricultural fires in the United States, *Atmos. Chem. Phys.*, 23(4), 2331-2343, doi:10.5194/acp-23-2331-2023.

U.S. Environmental Protection Agency Clean Air Markets Division (2024 b), Clean Air Status and Trends Network (CASTNET), [Weekly Dry Deposition], Available at www.epa.gov/castnet, Date accessed: [02 20, 2024]

670 U.S. Environmental Protection Agency Clean Air Markets Division (2024b), Clean Air Status and Trends Network (CASTNET), [Weekly Filter Pack], Available at www.epa.gov/castnet, Date accessed: [02 20, 2024]

U.S. Environmental Protection Agency, National Emissions Inventory [Tier I CAPS] (2023), available via <https://www.epa.gov/air-emissions-inventories/national-emissions-inventory-nei>. Accessed June 05, 2024.

Walker, J. T., P. Spence, S. Kimbrough, and W. Robarge (2008), Inferential model estimates of ammonia dry deposition in the vicinity of a swine production facility, *Atmospheric Environment*, 42(14), 3407-3418, doi:<https://doi.org/10.1016/j.atmosenv.2007.06.004>.

675 Walker, J. T., et al. (2019a), Toward the improvement of total nitrogen deposition budgets in the United States, *Science of The Total Environment*, 691, 1328-1352, doi:<https://doi.org/10.1016/j.scitotenv.2019.07.058>.

Walker, J. T., M. D. Bell, D. Schwede, A. Cole, G. Beachley, G. Lear, and Z. Wu (2019b), Aspects of uncertainty in total reactive nitrogen deposition estimates for North American critical load applications, *Science of The Total Environment*, 690, 1005-1018, doi:<https://doi.org/10.1016/j.scitotenv.2019.06.337>.

Wentworth G. R., Murphy J. G., Benedict K. B., Bangs E. J., Collett Jr J. L. (2016), The role of dew as a night-time reservoir and morning source for atmospheric ammonia. *Atmospheric Chemistry and Physics*. 2016;16:7435–7449. doi: 10.5194/acp-16-7435-2016.

685 Wichink Kruit, R. J., M. Schaap, F. J. Sauter, M. C. van Zanten, and W. A. J. van Pul (2012), Modeling the distribution of ammonia across Europe including bi-directional surface—atmosphere exchange, *Biogeosciences*, 9(12), 5261-5277, doi:10.5194/bg-9-5261-2012.

Xiu, A., and J. E. Pleim (2001), Development of a Land Surface Model. Part I: Application in a Mesoscale Meteorological Model, *Journal of Applied Meteorology*, 40(2), 192-209, doi:[https://doi.org/10.1175/1520-0450\(2001\)040<0192:DOALSM>2.0.CO;2](https://doi.org/10.1175/1520-0450(2001)040<0192:DOALSM>2.0.CO;2).

690

Formatted: Check spelling and grammar, Subscript

Zhan, X., et al. (2017), Evidence for the Importance of Atmospheric Nitrogen Deposition to Eutrophic Lake Dianchi, China, *Environmental Science & Technology*, 51(12), 6699-6708, doi:10.1021/acs.est.6b06135.

Zhang, L., J. R. Brook., and R. Vet (2003), A revised parameterization for gaseous dry deposition in air-quality models, *Atmos. Chem. Phys.*, 3, 2067–2082, doi:https://doi.org/10.5194/acp-3-2067-2003.

- 695 Zhang, L., L. P. Wright, and W. A. H. Asman (2010), Bi-directional air-surface exchange of atmospheric ammonia: A review of measurements and a development of a big-leaf model for applications in regional-scale air-quality models, *Journal of Geophysical Research: Atmospheres*, 115(D20), doi:https://doi.org/10.1029/2009JD013589.