

5 **Sorting sudden stratospheric warmings with the downward tropospheric influence using ERA5 and CESM2-WACCM**

Rongzhao Lu¹ and Jian Rao^{1*}

¹Collaborative Innovation Center on Forecast and Evaluation of Meteorological Disasters / Key
Laboratory of Meteorological Disaster of Ministry of Education, Nanjing University of Information
10 Science and Technology, Nanjing 210044, China

Correspondence to: Jian Rao (raojian@nuist.edu.cn)

Abstract. Sudden Stratospheric Warming events (SSWs) can have a downward impact on the
troposphere, but the mechanism remains uncertain. This study focuses on classifying SSWs based on
their tropospheric responses and documenting associated dynamical characteristics.~~To aid further
15 research, this study explores the characteristics of SSWs that have different impacts on the troposphere.~~
Using the ERA5 data and CESM2-WACCM outputs, 52 SSWs are identified in ERA5 and 273 in
CESM2-WACCM, with 33 and 119 downward-propagating SSWs (DWs), respectively. The DWs are
classified into three types based on cold surges over Eurasia (EA), North America (NA), and both
(BOTH), respectively. Both DWs and non-downward-propagating SSWs (NDWs) weaken and deform
20 the polar vortex, but DWs induce stronger negative Northern Annular Mode (NAM) and North Atlantic
Oscillation (NAO) responses. For DWs, the anomalous high develops in the polar region, which deflects
to lower latitudes, consistent with the frequent appearance of the polar high and the midlatitude blockings.
The shape of the anomalous polar high varies with the DW type, and the extension and shift of the
anomalous high lead to different surface responses. The DWs are also accompanied by a southward shift
25 of the precipitation belt, especially over the oceanic and coastal regions. On average, the relatively
weaker tropospheric impact of NDWs may be partly explained by consistent with their weaker
stratospheric disturbance amplitude.~~NDWs show relatively weaker impact on the troposphere, which is
primarily related to the weaker stratospheric disturbance amplitude.~~ The three types of DWs differ in
spatiotemporal evolutions of the NAM and NAO pattern, different forcing by planetary waves, and
30 varying ratios between displacement and split. This study reveals the diversity of the DWs and
distinguishes their potential impacts on both continents in the Northern Hemisphere.

1 Introduction

In the northern winter stratosphere, the polar temperature increases dramatically in just a few days when the polar vortex deforms or even collapses, and meanwhile the circumpolar westerly winds decrease suddenly and even reverse the direction (Baldwin et al., 2021). This phenomenon, known as sudden stratospheric warming (SSW), is an important manifestation of stratosphere-troposphere coupling in spring and winter. The SSW occurrence is associated with strong, upward-propagating planetary waves from the troposphere (Sjoberg and Birner, 2012; Butler et al., 2015). Recent studies also found that stratospheric preconditioning might play a decisive role in inducing the SSW event, by determining the intensity of the interaction between planetary waves and the mean flow (White et al., 2019; Yang et al., 2023). Polvani and Waugh (2004) suggested that SSWs are caused by a sharp increase in the amplitude of planetary-scale waves (primarily wave 1 and 2) propagating upward from the troposphere and perturbing the stratospheric polar vortex. Modeling evidence shows that the wave 1 forces the displacement of the polar vortex from the North Pole (Lindgren and Sheshadri, 2020), while the wave 2 determines the extent of vortex elongation and deformation (Baldwin et al., 2021).

Stratospheric variability associated with SSWs affects the near surface weather and climate through stratosphere-troposphere coupling processes (Hitchcock and Simpson, 2014; Wu and Reichler, 2019). Several mechanisms have been proposed to explain the possible downward impact of the stratosphere on the troposphere, including wave-flow interaction theory (Kuroda and Kodera, 1999), balanced flow dynamics theory (Black, 2002; Haynes, 1991), planetary wave refraction theory (Schmitz and Grieger, 1980), non-local potential vorticity response (Ambaum and Hoskins, 2002), and isentropic atmospheric meridional mass circulation theory (Cai and Shin, 2014). The wave-mean flow interaction theory suggests that upward-propagating tropospheric wave forcing fluctuations affect stratospheric mean flows, which in turn affect the vertical propagation of planetary waves (Kuroda and Kodera, 1999; Hartmann, 2000). As a consequence, the stratospheric disturbances cause significant downward impact on tropospheric circulation variations and near surface climate anomalies (Colucci and Kelleher, 2015; Dall'Amico et al., 2010), which are proposed to be amplified by tropospheric eddy feedback (Kidston et

al., 2015).

SSWs can have a considerable impact on the troposphere and a sustained effect on surface weather for weeks or even months (Domeisen et al., 2020; Rao et al., 2021; Lu et al., 2023). The weakened polar vortex during SSW usually projects onto the negative phase of the Northern Annular Mode (NAM) and/or the North Atlantic Oscillation (NAO), which gradually propagates downward (Karpechko et al., 2017; Kunz and Greatbatch, 2013; White et al., 2019). The negative phase of the NAM is often accompanied by equatorward shifts in storm paths and tropospheric jets (Kidston et al., 2015), shifts in the centre of the East Asian jets, variation in the blocking frequency (Anstey et al., 2013), increased possibility of cold air outbreaks over Eurasia and North America (Baldwin and Dunkerton, 2001; Lehtonen and Karpechko, 2016; Lu et al., 2022; Yan et al., 2022), and likelihood of extreme rainfall (Karpechko and Manzini, 2012). The common characteristics of SSWs have been widely reported in literature (see Baldwin et al., 2021 and references therein). However, every SSW has its individual features and displays strong particularity (Karpechko et al., 2018; Rao et al., 2018, 2019, 2021; Lu et al., 2023, 2024). It is the differences between individual SSWs and the background conditions in observations that distinguish their influence on the troposphere. As a consequence, the tropospheric response signals vary in extent, area and scope. Possible factors explaining the SSW individuality especially in its influence include the SSW strength, the initial warming location or the warming center (Zhang and Chen, 2019; Yan et al., 2022), the warming duration time (Hitchcock and Simpson, 2014), details of the wave flux between the troposphere and stratosphere (Shi et al., 2024), and the geometry of the polar vortex (split or displacement) (Maycock and Hitchcock, 2015; Rao et al., 2020). Therefore, the SSWs can be classified into different types based on those metrics. Different SSW classifications are widely used in literature. For example, based on vortex geometry shape the SSWs are classified as split, displacement, and mixed types (Charlton and Polvani, 2007; Rao et al., 2019). Based on whether the stratosphere reflects planetary waves during the westerly recovery phase following the SSW onset, it can be divided into absorbing and reflecting types (Kodera et al., 2016). According to whether the event has had a significant impact on the troposphere, it can be grouped into downward (DW) or non-downward (NDW) types (Jucker, 2016; Runde et al., 2016; Karpechko et al., 2017; White et al., 2019; Chwat et al., 2022). Although the DW SSWs show dripping NAM signals from the troposphere to the stratosphere, the near surface response structure is still different among DWs, with cold extreme sometimes only appearing in Eurasia, sometimes only in North America, and sometimes in both continents. However, there is still not a widely accepted subclassification for DWs events based on

the coverage of the near-surface cold anomalies associated with SSW. This study is mainly concerned with two questions: (1) What causes the inter-case difference in the DW influence on the troposphere although by definition the NAM during DWs shows downward propagation to the lower troposphere?

95 (2) What can we learn from the flavor identification for DWs?

~~The primary objective of this paper is to classify SSWs based on their tropospheric responses and to document the associated dynamical characteristics. The primary objective of this paper is to classify SSWs based on different tropospheric responses and to conduct an investigation into the underlying mechanisms from the dynamic differences.~~ The organization of the paper is constructed as follows.

100 Following the introduction, Section 2 describes the data and methods used in this paper. Section 3 compares the tropospheric response characteristics of various DW events. Section 4 analyzes the dynamics related to various DW events. Finally, Section 5 provides a summary and discussion.

2 Data and methods

2.1 Reanalysis data

105 We use the European Centre for Medium-Range Weather Forecasts (ECMWF) fifth generation reanalysis (ERA5) dataset over the period from October 1940 to December 2022. The data prior to 1958 are used with an expectation of selecting more SSW samples. Both three-dimensional and two-dimensional data are used. The three-dimensional variables on the isobaric levels include the air temperature, the geopotential, the zonal and meridional winds, and the vertical velocity on p coordinate. The two-
110 dimensional variables at a single surface level include total precipitation and the temperature at two meters (t2m). The isobaric levels extend from 1000 hPa to 1 hPa, and the local horizontal resolution of the ERA5 is 0.25° latitude by 0.25° longitude (this dataset is collected at 1.5° latitude by 1.5° longitude for easy handling).

2.2 Model data

115 Considering that the SSW samples from the ERA5 reanalysis are not too large, three historical simulations (r1i1p1f1, r2i1p1f1, and r3i1p1f1) from the CESM2-WACCM are employed to increase the SSW number. The CESM2-WACCM model has been shown to simulate the SSW frequency, displacement versus split types, and the downward impact well (e.g., Liang et al., 2022). The historical simulation starts from 1850 and ends in 2014, which is one of the common experiments from CMIP6.

120 Daily outputs from CESM2-WACCM have a horizontal resolution of 0.9° latitude by 1.25° longitude at 8 standard pressure levels from 1000 to 10 hPa.

The daily climatology is computed as the long-term mean for each calendar day, and the raw daily climatology is smoothed using 31-day means. The daily anomalies refer to the detrended deviation relative to the smoothed daily climatology.

125 By examining the same classification framework for both ERA5 and CESM2-WACCM, this study aims to assess whether the identified SSW type distinctions are reproducible across sensitive to different data sources, rather than being dependent on a specific reanalysis dataset product.

2.3 DW and NDW event definitions

We use the definition from Charlton and Polvani (2007) to identify SSWs in the Northern Hemisphere.

130 Specifically, all days when a change in the zonal-mean zonal wind \bar{u} from westerlies to easterlies occurs at 60°N and 10 hPa within the period from 1 November to 31 March each year are selected. The first day when \bar{u} undergoes a transition from westerlies to easterlies is defined as the onset date of this SSW. Several SSW definitions considering the wind reversal at different latitudes and heights in the circumpolar region have been compared by Butler and Gerber (2018). They concluded that a small
135 modification in the wind position or the wind threshold used for the SSW definition does not lead to major changes in the statistics of the SSWs. Considering the zonal winds change radically even after the SSW onset, the onset date of the two SSW events must be more than 20 days of consecutive westerlies apart (roughly double the radiative timescales in the middle stratosphere). Further, \bar{u} must have recovered to westerlies at least for 10 consecutive days prior to 30 April, which can exclude final
140 warmings (e.g., Charlton and Polvani, 2007). Finally, 52 SSWs are selected from ERA5, and 273 SSWs are selected from CESM2-WACCM. The average occurrence frequency of SSW is approximately 0.63 per year in ERA5 and 0.55 per year in CESM2-WACCM, consistent with previous studies (Liu et al., 2019; Rao et al., 2021).

To classify the SSW as the DW or the NDW, the NAM index is calculated (Baldwin and Thompson,
145 2009). After removing the daily climatological mean from the geopotential heights, the anomaly data are area-weighted (i.e., multiplied by cosine of the latitude) over 20°-90°N. The anomaly data are normalized for each pressure level, and the empirical orthogonal function is performed to extract the leading mode (i.e., the NAM pattern). The anomaly field is projected onto the leading mode to compute the

corresponding time series (i.e., the NAM index). After the NAM index is obtained at each pressure level,
150 the index at 1000 hPa is used to represent the NAO index. The NAM index for each pressure level is
calculated separately, and the NAM at 150 and 850 hPa is used to identify DWs.

The definition of DWs used in this paper follows the method by Natarajan et al. (2019). Namely, a DW
is defined when the SSW satisfies the following three conditions: During the 45-day period from day 8
to day 52 relative to the SSW onset date, 1) the average NAM index is negative at both 1000 hPa and
155 150 hPa; 2) the percentage of days with a negative NAM index is greater than 70% at 150 hPa; and 3)
the percentage of days with a negative NAM index is greater than 50% at 1000 hPa. To reduce the effect
of topographic complexity over high-terrain regions, White et al. (2019) adjusted 1000 hPa to 850 hPa
for the first and third conditions, which are also adjusted in this study. Based on these criteria, 33 DWs
(63% of all SSWs) are identified in ERA5, compared to 119 DWs (44% of all SSWs) from CESM2-
160 WACCM.

2.4 Classification of DWs

It is assumed that following the DW event onset, continental cold anomalies can develop over Eurasia
and/or North America, implying an increase in cold air outbreaks after the DW SSW onset. To better
describe and distinguish the DWs, we further divide DWs based on the inland temperature anomalies
165 within 40 days after the onset of DWs. Considering that cold surges are more active over the Eurasia
(Europe + Asia) and North America (US + Canada) following the DWs, a comparison between the cold
anomalies over the two regions can further classify the DWs into three types: cold anomalies only
appearing over North America (NA), only appearing over Eurasia (EA), and over both regions (BOTH).
Regions focused in this study include Europe (40°-70°N, 0°-60°E), Asia (40°-70°N, 60°-140°E), United
170 States (30°-46°N, 70°-120°W), and Canada (46°-60°N, 60°-120°W). Within one region, the area-
averaged temperature anomalies over a 40-day period are considered to be associated with the downward
impact of the DW if the anomalies meet the following two criteria: 1) the percentage of days with cold
anomalies is greater than 50%; 2) the mean temperature anomaly over a 40-day period is less than -0.3°C.
This threshold can classify all DWs into the three types. For the mainland, as long as one district meets
175 the criteria, it is considered to be associated with the downward impact of the preexisting DW.

The classification of SSWs into split and displacement events is based on the method initially developed
by Mitchell et al. (2011) and adapted by Seviour et al. (2013) using the two-dimensional (2D) moment

analysis method. Maycock and Hitchcock (2015) modified this analysis method by using the geopotential height at 10 hPa instead of the potential vorticity used in Mitchell et al. (2011). Maycock and Hitchcock (2015) defined the edge of the polar vortex as the climatological mean geopotential height over 60°N and 10 hPa in winter, and this definition is adopted in this study. Maycock and Hitchcock (2015) employed the following thresholds: a displacement event is identified when the centroid latitude is less than 66.5°N for a minimum of seven consecutive days, while a split event is defined when the vortex aspect ratio is larger than 2.8 for at least seven consecutive days. Several modifications for some parameters in this study are as follows. We choose the thresholds as the most equatorward 5.7% of centroid latitudes and largest 5.2% of aspect ratios, yielding a threshold of 62.9°N for centroid latitude and 2.46 for aspect ratio, respectively (e.g., Mitchell et al., 2011; Seviour et al., 2013). Table 1 shows the onset date and specific categorization of each event in ERA5. The general statistics of SSWs from the three historical simulation members by CESM2-WACCM are shown in Table 2. In ERA5, types BOTH, EA, NA and NDW account for approximately 25%, 27%, 12% and 36% respectively. In CESM2-WACCM, the corresponding proportions are 14%, 19%, 11% and 56%, respectively.

Table 1. Statistics of the SSWs from 1940–2022. The second column (S/D) shows the SSW type based on the vortex shape (D=displacement, S=split). The third column shows the SSW type based on whether the stratospheric signal propagates downward (DW=downward, NDW=non-downward). The last column shows the subclassification of DWs.

onset date	S/D	DW/NDW	type	onset date	S/D	DW/NDW	type
1941-02-07	D	DW	BOTH	1981-03-04	D	NDW	—
1942-03-21	D	NDW	—	1981-12-04	D	NDW	—
1945-03-19	S	NDW	—	1984-02-24	D	DW	BOTH
1946-02-19	D	NDW	—	1985-01-01	S	DW	BOTH
1946-03-19	D	NDW	—	1987-01-23	D	DW	EA
1950-03-05	D	NDW	—	1987-12-08	S	NDW	—
1952-02-22	S	DW	BOTH	1988-03-14	S	NDW	—
1952-11-19	D	DW	EA	1989-02-21	S	NDW	—
1954-12-18	D	DW	EA	1998-12-15	D	NDW	—
1955-01-26	S	DW	BOTH	1999-02-26	S	DW	EA
1957-02-04	S	DW	EA	2000-03-20	D	NDW	—
1958-02-01	D	DW	NA	2001-02-11	S	DW	NA
1960-01-17	D	DW	BOTH	2001-12-30	D	NDW	—
1963-01-27	S	NDW	—	2002-02-17	D	DW	NA
1965-12-16	D	NDW	—	2003-01-18	S	NDW	—
1966-02-22	S	DW	EA	2004-01-05	D	DW	NA
1968-01-07	S	DW	EA	2006-01-21	D	DW	EA

1968-11-28	D	DW	BOTH	2007-02-24	D	NDW	—
1969-03-13	D	NDW	—	2008-02-22	D	DW	NA
1970-01-02	D	DW	BOTH	2009-01-24	S	DW	EA
1971-01-18	S	DW	BOTH	2010-02-09	S	DW	EA
1971-03-20	D	DW	BOTH	2010-03-24	D	DW	EA
1973-01-31	S	NDW	—	2013-01-06	S	DW	EA
1977-01-09	S	DW	BOTH	2018-02-12	S	DW	EA
1979-02-22	S	DW	BOTH	2019-01-01	S	DW	NA
1980-02-29	D	DW	BOTH	2021-01-05	D	DW	EA

Table 2. general statistics of SSWs from the three historical simulation members by CESM2-WACCM.

Member	SSW	DW			NDW
		BOTH	EA	NA	
r1i1p1f1	90	13	18	9	50
r2i1p1f1	87	11	15	10	51
r3i1p1f1	96	16	18	9	53
Total	273	40	51	28	154

2.5 Isentropic potential vorticity

200 In this paper, isentropic potential vorticity (IPV) is used. The vertical component of IPV is defined (Hoskins et al., 1985) as:

$$\text{IPV} = -\frac{g(f + \vec{k} \cdot \nabla_{\theta} \times \vec{V})}{\frac{\partial p}{\partial \theta}}, \quad (1)$$

where f is the Coriolis parameter, θ is the potential temperature, g is gravity acceleration, p is air pressure, \vec{k} is the vertical unit vector, and \vec{V} is horizontal wind vector at isentropic surface.

205 2.6 Eliassen-Palm (E-P) flux

The E-P flux and its divergence can characterize the propagation of quasi-geostrophic planetary waves and its interaction with the mean flows. The E-P flux and its divergence are employed to diagnose the dynamical processes during SSWs as follows (Edmon et al., 1980):

$$F_{\varphi} = -a(\cos \varphi) \overline{u'v'}, \quad (2)$$

$$210 \quad F_p = a(\cos \varphi) f \frac{\overline{v'\theta'}}{\theta_p}, \quad (3)$$

$$\nabla \cdot \mathbf{F} = \frac{1}{a \cos \varphi} \frac{\partial (F_{\varphi} \cos \varphi)}{\partial \varphi} + \frac{\partial F_p}{\partial p}, \quad (4)$$

where F_{φ} is the horizontal component of the E-P flux, F_p is the vertical component, and $\nabla \cdot \mathbf{F}$ is the

divergence of the E-P flux, and a , φ are the Earth's radius and latitude, respectively. The E-P flux vector characterizes the propagation direction of the planetary waves. The E-P flux divergence indicates the effect of the planetary waves on the mean flow. When the E-P flux is convergent (divergent), it indicates that there is easterly (westerly) forcing of the mean flow.

3 Comparison of tropospheric responses to three types of DWs

3.1 Spatiotemporal evolution of the NAM

The NAM index can be used to ~~characterize~~describe the downward propagation of stratospheric disturbance. The composite evolutions of the NAM index for the three types of DWs are compared in Fig. 1. Consistent with previous studies (Karpechko et al., 2017; Kunz and Greatbatch, 2013; White et al., 2019), negative NAM signals appear above 200 hPa within a few days around the SSW onset date. After the DW onset, the negative signal propagates downward into the troposphere (Fig. 1a1-c1), forming a typical dripping-paint pattern in the troposphere (e.g., Baldwin and Dunkerton, 2001). On day 20 and afterward, the upper stratosphere gradually recovers to the positive NAM, while the negative signal below 50 hPa returns to the positive NAM at different times for the three types of DWs. The negative NAM signal below 50 hPa can persist until day 60 and even afterward, suggesting that the downward influence of DWs is persistent, which is consistent with previous suggestions of enhanced tropospheric predictability following SSWs~~providing a possible predictability source for the troposphere~~ (Rao et al., 2021; Lu et al., 2023). Differences in the NAM evolutions are noticed for the three types of DWs in ERA5.

1) In the pre-SSW onset period, the NAM displays different behaviors. The positive NAM mainly develops around day -50 and day -30 with the most significant signals in the lower stratosphere and upper troposphere for the type BOTH. The positive NAM only develops in the upper stratosphere around day -30 for the type EA. In contrast, type NA exhibits characteristics markedly distinct from other types: the positive NAM dramatically develops from day -50 in the upper stratosphere to day -15 in the lower troposphere, forming a pronounced vertical structure extending from the upper stratosphere to the lower troposphere (Fig. 1c1)~~displaying a noticeable downward propagation for the type NA.~~

2) In the post-SSW onset period, the negative NAM is structured in different spatiotemporal dripping shapes. Around 20 days or more after the SSW onset, the NAM is reversed in the upper stratosphere,

while the reversion time in lower levels is later. The negative NAM at 50 hPa persists longer than any other level for all types of DWs and even NDWs. In contrast, the NAM sign reversion is earliest for NA out of the three DW types around day 50 at 50 hPa (Fig. 1c1), while the NAM sign change is much slower for BOTH and EA beyond day 60 (Fig. 1a1, b1). It is also seen that the NAM sign change for NDWs is also around day 50 at 50 hPa (Fig. 1d1).

3) ~~The process of converting positive NAM reversal from positive to negative NAM exhibits relatively minor differences. The positive NAM intensity before the SSW onset and the negative NAM intensity after the SSW onset are variously contrasted~~ for the DWs and NDWs. The NAM evolves from moderately positive to moderately negative for the type BOTH. It evolves from weakly positive to weakly but persistently negative for type EA in ERA5, and this signal transition is not significant in the model.


~~Further, it evolves from intensely positive to intensely but shortly negative for type NA.~~ The strong phase transition for NEA in ERA5 may be due to the small sample size and the influence of individual cases.

~~Further, it evolves from intensely positive to intensely but shortly negative for type NA. The positive NAM develops 40 days or more before the NDW, but the negative NAM intensity and the NAM contrast before and after the NDW onset are weak.~~

4) The near surface exhibits different behaviors in the NAM intermittent signals for the three types of DWs. Specifically, the negative NAM at the near surface is continuous for BOTH and EA types, while it is very short in the persistent time and shows a low significance level for NA. At the very beginning of the NA DWs, the NAM is still positive at the near surface due to the lagged downward impact of the stratosphere, with the negative NAM from day 10 to day 50. In contrast, the negative NAM signal fails to appear at the near surface for NDWs, only with significant positive NAM around day 30. The difference between BOTH and NDWs and between NA and EA is most significant in the troposphere and near surface, implying a diversity in the persistency and NAM intensity among SSW types.

Comparing CESM2-WACCM simulations (Fig. 1a2-d2) with ERA5, ~~the pre-SSW signals are not clearly discernible in CESM2-WACCM, yet the characteristic evolution~~ of the NAM ~~are~~ highly consistent.

The conclusions are nearly unchanged if the reanalysis is replaced by model simulations: the downward propagation of the NAM is more continuous during EA than NA, and the NAM is much shallower during the NDWs than during DWs. ~~The similar evolution patterns observed in ERA5 and CESM2-WACCM indicate that the characteristic differences are not confined to the fewer events in the observational data, lending confidence to the robustness of the conclusions.~~






Figure 1. The composite evolution of the NAM index for (a-c) three types of DWs and (d) NDW events.


275 (a) BOTH, (b) EA, (c) NA, (d) NDW. The reanalysis is numbered with “1” following the plot letters, and the model simulation is numbered with “2”. The units are in standard deviations. The dashed contours represent the difference between (a, d) BOTH and NDW (BOTH - NDW) and between (b, c) EA and NA (EA - NA), the contours show The black lines/solid contours represent/mark the statistical significance at the 95% level for the difference (contours) differences at the 95% confidence level. composite.

280

The NAO index is closely related to low-level circulation over the North Atlantic, which is significantly correlated with the probability of extremes, such as regional coldness, snowstorm, and strong winds (Thompson and Wallace, 2001; Scaife et al., 2014). Fig. 2 compares the probability density functions

(PDFs) of the 60-day average NAO index after the onset for each type of event. Comparisons with a
285 random sample of 2000 winters show that the probability distribution of the NAO index for DWs shifts
left toward negative, which indicates that DWs have a significant effect on the surface circulation by
projecting onto more negative NAO. In ERA5 (Fig. 2a), the NAO index after the DW onset for BOTH
is 2-3 times more likely to be less than -0.8 than for other cases, with the median roughly located at -0.7,
and the probability that the index is positive is almost zero. The median values of the NAO after EA and
290 NA are the same, roughly located at -0.5, but the difference is noticeable. Namely, the PDF of the NAO
for NA is more dispersed and the PDF peak is smaller than that for EA and BOTH. In contrast, the type
BOTH showed the largest composite mean of NAO (-0.762), followed by EA (mean NAO = -0.567) and
NA (mean NAO=-0.435). The PDF of NAO for NDWs is primarily concentrated between -0.2 and 0.4
(68.4%), with median and mean values near 0.1 and 0.088, which might indicate that the NAO nearly
295 has no preference during NDW events.

In the model simulation (Fig. 2b), the probability density function distribution of the NAO indices also
highly resembles that in ERA5 (Fig. 2a). Namely, the mean probability density function for BOTH and
EA is much more left-skewed than that for NA. Consistent with ERA5, the probability density functions
of BOTH and EA exhibit a higher degree of overlap. The probability density function for NA is
300 distributed more flattened than that for BOTH and EA. Moreover, due to the larger sample size, the peak
values can be seen more clearly for NA. The probability density function distribution is almost symmetric
on both sides of zero for NDWs in both ERA5 and CESM2-WACCM, although the peak value (~1.1) is
larger than that for the climatological probability density function (~0.8).



305 **Figure 2.** Probability density functions for the average NAO index during 2000 randomly averaged 60-day periods in winter (grey curve), the 60 days after SSW onset date for BOTH (black curve), EA (red curve), NA (blue curve), and NDW (orange curve) events from two datasets: (a) ERA5 and (b) historical simulations by CESM2-WACCM. The Kernel Density Estimation is used for the 60-day mean index to smooth the PDF curve.

310 **3.2 Comparison of near surface response**

315 Due to the definitions of the SSW types, the near-surface temperature impacts exhibit marked differences naturally. Figure 3 shows the composite t2m anomalies and their differences in the 40-day intervals before and after the SSW onset visually. To compare the possible different impact of DWs on the near surface, the composite t2m anomalies in the 40-day intervals before and after the SSW onset are shown in Fig. 3. In both ERA5 and CESM2-WACCM simulations (Fig. 3a-d), the cold anomalies appear over different regions during the SSW for DWs and the NDW. Specifically, in the pre-SSW period (day -40

to day 0), significant cold anomalies have well developed over northern Eurasia for BOTH and EA, while the cold anomalies are not detectable over Eurasia for NA (Fig. 3a-c). Although cold anomalies also appear over Eurasia, the anomaly magnitude is fairly weak for NDWs (Fig. 3d). The western coast of North America is covered with cold anomalies for BOTH and NDWs (Fig. 3a, d), while significant warm anomalies appear over most of North America for EA and NA (Fig. 3b, c). It is also found that significant warm anomalies develop over Europe for NA (Fig. 3c). For BOTH and EA, an anomalous high appears over the Arctic at 500 hPa, while for NA two anomalous high centers appear over North America and Europe, respectively (Fig. 3a-c). The tropospheric circulation anomalies are much weaker for NDWs than DWs (Fig. 3d). The general patterns of t2m and 500-hPa height anomalies are highly consistent between ERA5 and CESM2-WACCM simulations. _

In the post-SSW period, the t2m anomaly patterns differ among the DW types and the NDWs, consistent with the surface temperature t2m-based definition of the SSW events. DW events exhibit cold anomalies over their respective defining regions, while NDW events lack coherent surface temperature signals.

These characteristics are similarly represented in ERA5 and CESM2-WACCM. In the post-SSW period, the t2m anomaly pattern is contrastingly different for the three types of DWs and the NDW, consistent with the construction for t2m anomalies in ERA5 and CESM2-WACCM. For the type BOTH, the cold anomalies in the Eastern Hemisphere continue to strengthen and expand southward, covering most of Eurasia; the cold anomalies in the Western Hemisphere expand from western coasts of North America to most of the continent (Fig. 3e). The positive height anomaly center for the type BOTH is located over Greenland and the Bering Strait in ERA5 and over Greenland in model simulations. For the type EA, the cold anomalies persist over northern Eurasia and the cold center moves further eastward to Northeast Asia, while the warm anomalies over eastern Canada and southern Europe strengthen (Fig. 3f). The positive height anomaly center remains over the Arctic Ocean and Canada. For the type NA, warm anomalies over eastern Europe also intensify, while the warm anomalies over North America are replaced by anomalous coldness with the center over the central US (Fig. 3g). The positive height anomaly centers move from Canada and Europe to Greenland and Central Asia, respectively, although the composite anomalies are relatively weaker in CESM2-WACCM. For the NDW, the t2m and 500-hPa height anomalies over both lands are very scattered and less organized, although patches of warm anomalies are observed over Asia (Fig. 3h).





Figure 3. Composite 2-meter temperature (t_{2m}) anomalies (shadings; units: K) and 500-hPa geopotential height anomalies (contours; units: gpm) for (a, e) BOTH, (b, f) EA, (c, g) NA and (d, h) NDW events before the SSW onset (top row) and afterward (bottom row) from two datasets: (a1-h1) ERA5 and (a2-350 h2) historical simulations by CESM2-WACCM. The composite is based on the mean of 40-day intervals. The dots mark the composite anomalies at the 95% confidence level using the t -test.

To well depict the dynamic process of the near surface response to different types of SSWs, the composite evolution of t_{2m} anomalies over Eurasia and North America is shown in Fig. 4. In both datasets, for the type BOTH, cold anomalies develop following the SSW occurrence period from day 0 to day 40 over all 355 the four regions (Europe, Asia, US, and Canada), except that the negative t_{2m} anomalies at Canada are relatively weak and exhibit relatively large subseasonal variability (Fig. 4a). For the type EA, cold anomalies are persistent over northern Eurasia, larger in Asia than in Europe, while warm anomalies are

360 persistent over North America, larger in Canada than in US (Fig. 4b). For the type NA, the reversal of
temperature anomaly sign is observed over North America (US and Canada). The reversal of temperature
anomalies from positive to negative indicates outbreaks of cold air (Lehtonen and Karpechko, 2016).
Namely, cold air outbreaks increase in North America, while the anomalously cold state gradually
recovers to normal in Asia (Fig. 4c). For NDWs, the temperature anomalies are fairly weak throughout
the SSW onset except that Canada experiences a transition from anomalously warm state to cold (Fig.
365 4d).

Comparing ERA5 and historical simulations by CESM2-WACCM, the composite t2m anomaly
amplitude in model simulations is relatively weak and less significant in the pre-SSW period. Consistent
with ERA5, cold anomalies persist from pre-SSW to post-SSW periods over both continents for the
BOTH type in the model simulations. Further, the persistent cold anomalies over Asia from pre-SSW to
370 post-SSW periods are also present in model evidence for the EA type. The reversal of t2m anomalies
from positive to negative over North America and from negative to positive over Asia is also confirmed
by model simulations for the NA type. Both ERA5 and CESM2-WACCM simulations show that the t2m
anomalies over both continents are weak and insignificant most of the time during the NDWs (Fig. 4d1,
d2).



375

Figure 4. Evolution of the area-averaged t_{2m} anomalies (units: K) from day -40 to day 40 over four regions (I–IV), including Europe (40° - 70° N, 0° - 60° E), Asia (40° - 70° N, 60° - 140° E), United States (30° - 46° N, 70° - 120° W), and Canada (46° - 60° N, 60° - 120° W) for (a) BOTH; (b) EA; (c) NA and (d) NDW events from two datasets: (a1-d1) ERA5 and (a2-d2) historical simulations by CESM2-WACCM. The thickened part of the line denotes the composite at the 95% confidence level.


380

3.3 Analysis of isentropic potential vorticity

In order to further explore the process of cold air activity, an analysis of the isentropic potential vorticity (IPV) is investigated for the three types of DWs and the NDWs. According to Eq. (1), the IPV is

proportional to the absolute vorticity and static stability. In the case of equal vorticity, the cold air mass usually has a higher potential vorticity value due to its relatively larger static stability, which can be used to track the cold air activity on the isentropic surface (Hoskins et al., 1985; Lu and Ding, 2015). The 315K isentropic level has been used to track the cold air sources by detecting the movement of high IPV center (Jeong et al., 2006). The composite evolutions of IPV anomalies are shown in Fig. 5 for all types of events based on ERA5 and CESM2-WACCM simulations. For the type BOTH, a patch of high IPVs appear from North Pacific 30-20 days prior to the event and gradually move southeastwards in the following period, which finally reach the North America for both datasets (Fig. 5a). During day 10 to 20, anomalously high IPV air moves to 45°N and even more southward, with the anomaly amplitude gradually weakened. Significant negative IPV anomalies develop over Greenland, consistent with the local anomalous high (see Fig. 3e), which helps advect cold air to eastern North America. In contrast, there are two high IPV centers in Eurasia, one over Europe, and the other over Central Asia. The anomalously high IPV center over Europe has remained stationary and continues to intensify. The high IPV center over Central Asia is more stable from day -30 to -10 and redevelops from day 0 to 20. For the EA events, the high IPV center in the North Atlantic is weak and significant high IPV anomalies appear in Europe (Fig. 5b). The high IPV anomalies develop in North Atlantic soon after the SSW onset, and the Arctic is nearly covered by the positive IPV anomalies. Another positive IPV anomaly center is detected over North Asia since day -30 to -20, which diminishes from day -20 to -10 and redevelops from day -10 to 10. The high IPV patch moves southeastward to East Asia after day 10, indicating cold air outbreaks in local regions. For the NA events, the high IPV center first appears over North Pacific from day -20 to -10, which is still active from day -10 to 10 (Fig. 5c). The positive IPV anomaly intensity weakens from day 0 to 10 and then redevelops in the later period. Further, it is also noticed that a large patch of positive IPV anomalies form over North Atlantic after the SSW onset, which is nearly motionless. Similar to BOTH events, the blocking pattern denoted as a wide range of negative IPV anomalies over Greenland is observed, and the upstream winds can advect cold air to eastern North America. For the NDW events, the IPV anomalies are relatively weak, and the positive IPV anomalies are scattered and less organized over the land (Fig. 5d), consistent with the distribution of t2m anomalies. Weak positive IPV anomalies primarily appear over the oceans and are nearly motionless. It is also observed that a narrow band of positive IPV anomalies exists from the west to east across Canada during day 20

to day 30.



415

Figure 5. Composite isentropic potential vorticity (IPV) anomalies (shadings; units: PVU, 1 PVU = $10^{-6} \text{ m}^2 \text{ K kg}^{-1}$) at 315 K for (a) BOTH, (b) EA, (c) NA, and (d) NDW events from two datasets: (a1-d1) ERA5 and (a2-d2) historical simulations by CESM2-WACCM. The composite is based on the mean of 10-day intervals. The dots mark the composite anomalies at the 95% confidence level using the *t*-test.

420


3.4 Total precipitation

Previous studies have shown that SSWs not only influence the cold air outbreaks, but also affect the rainfall anomalies (King et al., 2019; Oehrlein et al., 2021). Negative NAM phases are usually accompanied by a possible equatorward shift of the storm track, and a possible intensification of the moving cyclone at low latitudes (Afargan-Gerstman and Domeisen, 2020; McAfee and Russell, 2008; Thompson and Wallace, 2001). The composite precipitation anomalies from day 0 to 40 are shown in Fig. 6 to compare the rainfall anomaly sensitivity to the SSW type. The most pronounced common features for the four conditions in both ERA5 and CESM2-WACCM simulations are the rainfall dipole

425

structure over North Atlantic - Europe. Positive rainfall anomalies form over Atlantic midlatitudes and the Mediterranean Sea, while negative rainfall anomalies form over Atlantic high-latitudes. This rainfall dipole is strong for BOTH and EA (Fig. 6a, b), while the intensity is relatively weak for NA and NDW (Fig. 6c, d). The positive rainfall anomaly in midlatitudes for NA is broken into two chains, one over the ocean, and the other biased toward the Eurasian land (Fig. 6c). Although the NAM fails to descend to the near surface, the rainfall dipole is biased farther northward for NDWs (Fig. 6d).

435 Comparing ERA5 and CESM2-WACCM simulations, the rainfall dipole is consistently present for the BOTH and EA types (Fig. 6a, b). The negative rainfall anomaly lobe in high latitudes is much weaker for the NA type than for the BOTH and EA types for both datasets, although the positive rainfall anomaly band in midlatitudes is still present (Fig. 6c). In ERA5, the dipole rainfall anomalies are nearly absent due to the non-downward propagation of NDW SSWs, while the rainfall dipole disappears in model simulations for the NDW events (Fig. 6d).



440

Figure 6. Composite precipitation anomalies (shadings; units: mm day^{-1}) for (a) BOTH, (b) EA, (c) NA and (d) NDW events after the SSW onset from two datasets: (a1-d1) ERA5 and (a2-d2) historical simulations by CESM2-WACCM. The composite is based on the mean of day 0 to 40. The dots mark the composite anomalies at the 99% confidence level using the t -test.

445 **4 Dynamic diagnostics**


In order to compare the large-scale atmospheric dynamics between different types of SSW events ~~and to better understand their downward influences~~, the composite circulation anomalies at 100 hPa and the sea level pressure anomalies are shown in Fig. 7. ~~These results are intended to document the type-dependent dynamical characteristics of the circulation, rather than to establish a mature causal physical mechanism for the observed differences that are far beyond the scope of the paper.~~ In the pre-SSW period, the circulation structure is differently organized for three types of DWs and the NDWs based on ERA5 and historical simulations by CESM2-WACCM. For the type BOTH, an anomalous high appears over Canada, and a low anomaly center forms over central northern Eurasia (Fig. 7a) with the polar vortex displaced toward North Asia. On the near surface, the negative NAM pattern has well developed, positive MSLP anomalies prevail over the Arctic, and negative anomalies develop over midlatitudes. The height anomaly distribution at 100 hPa for the type EA is similar to that for BOTH except that the height anomaly centers are further eastward situated (Fig. 7b). The anomalous high is situated at 60°E, while the anomalous low is situated over Northeastern Asia and Europe, implying that the polar vortex is likewise skewed towards Eurasia. Positive MSLP anomalies develop over the Arctic and northern Europe, while the negative MSLP anomalies form over northern Canada. For the type NA (Fig. 7c), the anomalous high at 100 hPa is centered over the Great Lakes, while the low covers most of northeast Asia. However, the near surface is covered by the anomalous low over most of the Arctic. ~~These characteristics are markedly different from EA (Fig. 7f). For NDWs, compared to DWs, the polar vortex at 100 hPa is not significantly disturbed, and no significant low-pressure systems developed at sea level (Fig. 7d, e). For NDWs, the polar vortex at 100 hPa is not significantly disturbed, although the near surface exhibits a pressure anomaly dipole, with the low over the Bering Strait—Alaska and the high over the northern Europe (Fig. 7d).~~ Comparing the three types of DWs, the precursor circulation anomalies for BOTH are more baroclinic from the lower stratosphere to the troposphere (Fig. 7a, b), while for NA and EA, the circulation anomalies are nearly barotropic in high latitudes (Fig. 7b, c). The height anomalies at 100 hPa are much weaker for NDWs (Fig. 7d) than for DWs. ~~The model results also suggest a weakening of the polar vortex at 100 hPa prior to DW events. At the surface, the spatial structure of the NAM, including the sign and location of the anomaly centers for different event types, broadly agree with those in the observational analysis. The circulation anomalies for the three types of DWs in the pre-SSW period are nearly barotropic in model simulations. The NAM pattern in model simulations is more clearly present~~

475 ~~in model simulations. For the type BOTH (Fig. 7a2), three anomalous low centers appear in the mid-~~
~~latitudes over East Asia, western North America, and the Atlantic Ocean, respectively. For the type EA~~
~~(Fig. 7b2), two anomalous high centers form in Europe and Canada, and anomalous low centers appear~~
~~in the North Pacific. For the type NA (Fig. 7c2), two anomalous high centers are located over the North~~
~~Pacific and North Atlantic, respectively. For NDW events (Fig. 7d2), the height anomalies are much~~
480 ~~weaker than the DWs, consistent with ERA5.~~ For the three types of DWs, during this period, negative
anomaly centres emerged over Eurasia (or East Asia or Europe), indicating that the polar vortex was
more inclined to shift towards Eurasia. This feature is consistent with the occurrence of cold anomalies
across Eurasia during this period for all three types of DWs. ~~This may be one reason why cold anomalies~~
~~occurred across Eurasia during this period for all three types of DWs.~~

485 In the post-SSW period, the Arctic is completely covered by high anomalies from the near surface to the
lower stratosphere with a nearly barotropic circulation structure (Fig. 7ge-jh). In the lower stratosphere,
the Arctic for all types of events is covered by the anomalous high, indicating the breakup of the
stratospheric polar vortex. The positive anomaly amplitude for the three types of DWs in both ERA5 and
model simulations is comparable (Fig. 7g-j), and their differences (Fig. 7k, l) are mainly featured by
490 midlatitude circulation anomalies. For BOTH, negative height anomalies are clearly present in
midlatitudes with three centers, one over North Atlantic, one over Canada, and one over East Asia at 100
hPa (Fig. 7e7g). Similarly, the MSLP anomalies are well structured in a negative NAM pattern with the
negative anomalies maximized over North Atlantic along midlatitudes. For EA, the negative height
anomalies are not as detectable as for BOTH, and only the negative center over East Asia and North
495 Atlantic is clearly observed at 100 hPa (Fig. 7f7h). On the near surface, the NAM structure is more clearly
present in the Atlantic sector than in the Pacific sector. The local anomalous high over North Pacific
disrupts the annular structure in the Eastern Hemisphere. For NA, the anticyclonic anomalies at 100 hPa
are further biased toward the Arctic Canada, while nominal negative height anomaly band in midlatitudes
for the negative NAM is only present over North Pacific (Fig. 7g7i). Similarly, the negative MSLP
500 anomaly band is concentrated from the Western Europe to East Asia and North Pacific (Fig. 7g7i),
different from the long anomalous low band in midlatitudes for BOTH and EA (Fig. 7i7j, 7l). For NDWs,
although positive height anomalies are present over the Arctic in the stratosphere at 100 hPa (Fig. 7h7j,
7k), the amplitude is nearly half or one third of the strength for DWs. The MSLP anomalies in Arctic are
very weak, while significantly negative MSLP anomalies over North Atlantic–Northern Eurasian are

505 observed. Similarly, centers of negative anomalies are also observed at 100 hPa over the regions where cold anomalies develop in the three types of DWs. The circulation anomaly distribution in the post-SSW period is consistent between ERA5 and model simulations. The negative NAM structure is more clearly present in model simulations, possibly due to a much larger sample size in model simulations than in ERA5.

510







Figure 7. Composite 100-hPa geopotential height anomalies (contours; units: gpm) and sea level pressure anomalies (shadings; units: hPa) for (a, **ge**) BOTH, (b, **hf**) EA, (c, **ig**) NA ~~and~~ and (d, **jh**) NDW, ~~as well~~ ~~as~~ ~~and~~ the composite difference between (e, **k**) BOTH and NDW, and between (f, **l**) EA and NA events before the SSW onset (top row) and afterward (bottom row) from two datasets: (a1-h1) ERA5 and (a2-h2) historical simulations by CESM2-WACCM. The composite is based on the mean of 40-day intervals. The dots mark the composite anomalies or differences at the 95% confidence level using the *t*-test.

515 To further analyze the wave dynamics and to better understand the difference between the three types of DWs and the NDWs, the E-P flux anomalies and the E-P flux divergence anomalies are shown in Fig. 8. The 10-day intervals before the SSW onset and afterward are examined for total planetary waves. For all types of SSWs, the upward propagation of planetary waves is enhanced before the event onset (Fig. 8a-d). Comparing the three types of DWs, the E-P flux anomalies are ~~very~~ comparable with the type BOTH a bit stronger, and the anomalous E-P flux convergence center is structured differently (Fig. 8a-~~ec~~, **f**). In both ERA5 and model simulations, the E-P flux convergence anomalies develop in the entire stratosphere for the type BOTH (Fig. 8a), while for EA and NA, the convergence anomalies are more concentrated around 50-150 hPa (Fig. 8b, c). The E-P flux convergence anomalies are weaker for the NDWs than all

types of DWs, likely due to the weak E-P flux itself (Fig. 8d, fd).



530

Figure 8. Composite E-P flux anomalies (vectors; units: F_y in 10^4 kg s^{-2} , F_z in 10^6 kg s^{-2}) and E-P flux divergence anomalies (shadings; units: m s^{-1}) by planetary waves (sum of wave 1 and wave 2) for (a, ~~ge~~) BOTH, (b, ~~hf~~) EA, (c, ~~ig~~) NA, ~~and and~~ (d, ~~jh~~) NDW, ~~as well as~~ and the composite difference between (e, k) BOTH and NDW, and between (f, l) EA and NA ~~events~~ before the SSW onset (top row) and afterward (bottom row) from two datasets: (a1-h1) ERA5 and (a2-h2) historical simulations by CESM2-WACCM. The composite is based on the mean of 10-day intervals. The black line represents the 90% confidence level for the composite E-P flux divergence anomalies or differences.

540 After the SSW onset, the composite 10-day intervals E-P flux anomalies are also not identical for the three types of SSWs (Fig. 8e-~~gg-i~~). The upward propagation ~~begins to weaken~~ for all types of DWs. However, the anomalous E-P flux convergence is still present in the lower stratosphere for type EA and NA ~~and their differences are mainly seen in the midlatitude troposphere (Fig. 8l)~~ all types of DWs. In contrast, weakening of the anomalous E-P flux convergence and the anomalous downward propagation of waves begins to form for the NDWs (Fig. 8j), ~~and the anomalous downward propagation of waves is also stronger for the NDW than the DW in ERA5, although model simulations only show E-P flux divergence anomalies in the polar stratosphere (Fig. 8h)~~. The relatively short lifetime and weak intensity of the wave forcing for the NDWs (Fig. 8k) likely explain the relatively weak intensity of the stratospheric disturbance ~~and even lacking impact on the near surface in the later period~~. It is also revealed that the E-P flux divergence (or convergence) anomalies in the upper stratosphere are much weaker than in the lower stratosphere and upper troposphere for all events (Fig. 8e-~~hg-j~~).




Figure 9. Same as in Figure 8, but for planetary wave 1 E-P flux.

The contribution of the wave 1 and wave 2 to the total E-P flux and its divergence is shown in Fig. 9 and Fig. 10, respectively. A decomposition of the waves for E-P flux before the SSW onset can easily reveal the dynamical difference among the three types of DWs. Specifically, in the pre-SSW onset, the upward propagation of planetary wave 1 is strengthened for all events in both ERA5 and model simulations, and the E-P flux convergence anomalies by wave 1 also appear for all events (Fig. 9a-d). Moreover, it is obviously observed that the E-P flux convergence anomalies by wave 1 are strongest for NA out of all groups (Fig. 9f) and weakest for NDWsother types (Fig. 9d-e, f).

In the post-SSW 10-day interval period, the upward propagation of wave 1 is suppressed, and the contribution of wave 1 to the deceleration of westerlies (and therefore weakening of the polar vortex) nearly disappears (Fig. 9e-hg-j). Namely, the E-P flux anomalies change the direction from upward to downward, and the E-P flux divergence anomalies are nearly zero. Wave forcing in midlatitudes is still present for DWs in ERA5, and the anomalous E-P flux convergence also persists in the lower stratosphere at midlatitudes (Fig. 9e-eg-i). For model simulations, although the upward flux still lasts, the convergence of the E-P flux has rapidly declined, ~~and NA even shows divergence of the flux.~~ Differences between NA and EA as well as between BOTH and NDWs are generally consistent in both datasets (Fig. 9k, l).




Figure 10. Same as in Figure 8, but for planetary wave 2 E-P flux.

575 Similarly, the dynamics difference for the three types of DWs can be easily seen from the contribution
of wave 2 to the total E-P flux and its convergence. In the pre-SSW periods, wave 2 cooperates with
wave 1 to enhance the upward propagation of planetary waves and therefore the anomalous E-P flux
convergence for BOTH and EA (Fig. 10a+, b+). Although the wave 2 is enhanced to propagate upward
to the stratosphere for NA and NDWs, ~~neither the composite E-P flux convergence of the E-P flux anomalies~~
580 ~~occurred nor undetectable~~ ~~the E-P flux is little converged,~~ ~~so~~ and the wave-driven change for the
circulation is not present (Fig. 10c+, d+). ~~For model simulations~~ In contrast, BOTH and EA show stronger
E-P flux convergence anomalies in the model than in ERA5. For NA and NDWs, ~~unlike in ERA5,~~
~~weak~~ the E-P flux divergence anomalies also occur in the lower stratosphere are stronger in ERA5
than in the model. The differences between EA and NA and those between BOTH and NDWs in the pre-
585 SSW period further confirm the diversity of the SSW dynamics (Fig. 10e, f).

This difference in the wave 2 forcing also exists in the post-SSW period for the three types of DWs (Fig.
10e-g-l). ERA5 shows that the vertical component of E-P flux anomalies by wave 2 reverses the sign for
BOTH and NA (Fig. 10g~~e~~, g~~i~~), while the upward propagation of wave 2 is still present for EA (Fig.
10f~~l~~10h+). As a consequence, the wave 2 forcing for the weakening of the polar vortex still persists after
590 the SSW onset for EA, while this forcing for BOTH and NA types by wave 2 has terminated. Changes
in the E-P flux anomalies and the divergence anomalies by wave 2 are not noticeable for NDWs (Fig.
10h~~l~~10j). For model simulations, the E-P flux anomalies in various types of events have rapidly decayed,
and the wave forcing has quickly disappeared. Differences between NA and EA as well as between BOTH
and NDWs are much larger in ERA5 than in model simulations (Fig. 10k, l).

595 The wave 1 and wave 2 can exert different effects on the stratospheric polar vortex (Baldwin et al., 2021;
Lindgren and Sheshadri, 2020; Yang et al., 2023). The former propels the polar vortex displaced away
from the Arctic, while the latter elongates and splits the vortex. Previous studies (Anstey et al., 2013;
Kidston et al., 2015; Lehtonen and Karpechko, 2016) found that 2–3 weeks before the displacement SSW,
it is anomalously warmer in the southeastern US and colder in Eurasia. These studies also found 2–3
600 weeks after the displacement SSW, the southeastern US is anomalously cold while Eurasia is unusually
warm. In the month around the split SSW, the probability of both continents (North America and Eurasia)
being cold at the same time is high. The different percentage of the displacement or split SSWs might
further account for the different circulation and t2m anomalies for different types of DWs.

5 Conclusions

605 SSWs show strong inter-case variability for their impact on the troposphere and near surface. In this paper, SSWs are classified into different groups based on whether there is downward impact and on where the downward impact occurs. Using a newly proposed method, this study further classifies 52 SSWs in ERA5 and 273 SSWs in CESM2-WACCM into four groups: DWs with near surface impact over both Eurasia and North America (BOTH), DWs with impact over Eurasia (EA), DWs with impact
610 over North America (NA), and NDWs. Finally, 33 DWs (13 + 14 + 6) and 19 NDWs were identified from ERA5, while 119 DWs (40 + 51 + 28) and 154 NDWs were selected from CESM2-WACCM simulations. Physical quantities in the CESM2-WACCM model tend to exhibit weaker amplitudes than those in ERA5, although it simulates the downward influence of the SSW effectively, successfully reproducing general features. The agreement between ERA5 and CESM2-WACCM does not imply that the underlying physical mechanisms are fully captured, but it does indicate that the identified type-dependent characteristics are reproducible/robust across distinct datasets/systems.
615 To well distinguish the potential impact diversity of SSWs, the tropospheric circulation response and the near surface behaviors following these four types of events are revisited in this study. The main findings in this study are as follows and summarized in Fig. 11.

620 1) The mean intensity of NDW events in terms of the NAM and the circulation anomalies in the stratosphere and troposphere are nearly half weaker than DWs events. Comparing the downward impact of three types of DWs, the persistency of the negative NAM pattern varies with the DW type. On average, the negative NAM signal in the lower stratosphere can last for >60 days for BOTH and EA, while it only lasts for ~40 days for NA events. Further, the dipping pattern at the near surface exhibits a continuous
625 negative NAM signal for BOTH and EA, while it is replaced by positive NAM at both the beginning and end of the SSW for NA.

2) An isentropic vorticity analysis reveals that anomalously high PV air can move southwards from the polar stratosphere and enter midlatitudes, thereby affecting the near ground. The cold anomalies over Eurasia and over North America have precursors in the upstream oceanic regions before the SSW onset.
630 Anomalously low PV appears over North Atlantic 10-30 days before the EA onset and anomalous high PV forms over North Atlantic 0-30 days afterward. In contrast, anomalously high PV appears 10-20 days before the NA onset to 0-10 days afterward over North Pacific – Alaska. The PV anomaly evolutions for BOTH have a joint characteristic of both EA and NA. Namely, anomalously high PV forms over North

Atlantic and North Pacific – Alaska as the anomalously low PV air enters the Arctic before the SSW
635 onset for BOTH, while anomalously high PV air sweeps the continents soon afterward. In contrast, the
t2m and 315-K PV anomalies are weaker and more scattered.


3) Adopting relative continental cold anomalies as the criterion for classifying SSWs, changes in
precipitation anomalies are not so sensitive to the DW type as t2m anomalies. Following the onset of all
types of SSWs, the rainfall band shows a southward shift especially over the Atlantic–Europe sector,
640 exhibiting a rainfall anomaly dipole in the Western Hemisphere. Namely, the precipitation in subtropical
Atlantic – Mediterranean Sea increases, while precipitation in high-latitude regions decreases. Previous
studies have found that the southward shift of the precipitation band is associated with an equatorward
shift in the storm track, and the intensification of moving cyclones at low latitudes (e.g., Thompson and
Wallace, 2001; Huang and Xie, 2015).

645 4) The dynamical processes during DWs and the NDW_s are various before and after the SSW onset. In
the pre-onset period, the negative NAM has been shaped at surface for BOTH and EA, while weak NAM
still dominates at surface for NA, although anomalous high begins to form over the Arctic. In the pre-
onset period, across all three types of DWs, the anomalously low centers are present over Eurasia, which
signifies that following the collapse of the polar vortex, its remnant tends to move towards Eurasia. For
650 BOTH, the anomalously low band stretching from the North Atlantic to North America represents the
portion of the polar vortex that also extends towards the America. In the post-onset period, anomalously
low centers exist above each cold anomaly region. This indicates that around the time of DWs, the polar
vortex either shifts or splits, and the movement of the collapsed polar vortex is a significant factor in the
region experiencing cold anomalies. The near surface NAM pattern is well organized with the midlatitude
655 negative pressure anomalies over sea larger than over lands for BOTH and EA, while the pressure
anomalies are larger over lands than over oceans for NA. For the NDW_s, the circulation anomaly
amplitude is only half or even one third of that for DWs.

5) The dynamic differences for the three types of DWs are also featured by the wave activities. Firstly,
the wave 1 forcing for them (BOTH, EA, and NA) shows a similar structure in the pre-onset period: the
660 upward propagation of wave 1 increases, and the anomalous E-P flux convergence denotes a dissipation
of wave 1 in the lower stratosphere. In the post-onset period, the enhancement of upward-travelling wave
1 terminates for BOTH and EA, but presents for NA. Secondly, the wave 2 forcing behaves in very
different spatiotemporal structures: the wave 2 is enhanced to propagate upward in the pre-onset period

for all types of DWs, but the wave dissipation in the stratosphere is only detected for BOTH and EA. In the post-onset period, the upward propagation of wave 2 is instantly suppressed for BOTH and NA, while the wave 2 forcing is still strong for EA.

665 6) Considering that displacement and split events have different impacts on the two continents (Anstey et al., 2013; Kidston et al., 2015; Lehtonen and Karpechko, 2016), a larger proportion of displacement SSW for NDWs and NA might explain their weak downward impact on near surface especially over
670 Eurasia.






Figure 11. Schematic charts comparing the three types of DWs and the NDW. Aspects of comparison include preceding wave forcings by wave 1 and 2, the circulation pattern in the stratosphere, the stratosphere-troposphere coupling, and the downward impact on near surface.

Compared with previous studies (Anstey et al., 2013; Domeisen et al., 2020), our study reveals the diversity of the DW events and distinguishes the potential impact on both continents in the Northern Hemisphere. The findings underscore the significant regional diversity in the tropospheric response to different SSW types, with distinct durations, intensities, and spatial patterns. These new SSW classifications and syntheses of their relevant dynamic features can improve our understanding of weather and climate variability associated with SSWs, though the specific mechanisms responsible for their differing impacts await further investigation. These new insights enhance our understanding of the mechanisms driving surface impacts and could improve our understanding of weather and climate variability associated with SSWs. This consideration is reasonable, because cold air outbreaks on both

continents are not in pace most of the time (Butler et al., 2017; Yu et al., 2024). White et al. (2019) found that at least 35 events should be present to get robust differences between DW and NDW events. The sample sizes from CESM2-WACCM generally meet the case requirement in this study. Using more samples from CMIP6 multimodel outputs, a deeper understanding of different types of DW events is possible, left for future investigation. Further, the distribution of various types of events (see Table 1) shows strong interdecadal variability in past decades. Whether this change is an internal climate variability or forced by global warming due to anthropogenic emission is still unknown, worth exploring in the future.

Data availability

The ERA5 reanalysis is available from the ECMWF (<https://cds.climate.copernicus.eu/cdsapp#!/dataset/reanalysis-era5-pressure-levels-monthly-means?tab=form>). The CESM2-WACCM historical simulations participating CMIP6 are available from the ESGF (<https://esgf-node.llnl.gov/projects/cmip6/>).

Author contributions

RL and JR designed this research. RL and JR analyzed the data. RL provided the data analysis methods. RL wrote the manuscript draft. JR reviewed and edited the manuscript.

Competing interests

The contact author has declared that none of the authors has any competing interests.

Acknowledgments

The authors express their gratitude to the National Natural Science Foundation of China for the funding support. The authors thank the High Performance Computing Center of Nanjing University of Information Science & Technology for their support of this work. The ECMWF is acknowledged by the authors for the accessible modern reanalysis data.

Funding

This research was supported by the National Natural Science Foundation of China (Grant nos.

42361144843, 42322503, 42175069, and 42288101) and the Qinglan project of Jiangsu.

References

- Afargan-Gerstman, H. and Domeisen, D. I. V.: Pacific modulation of the North Atlantic storm track response to sudden stratospheric warming events, *Geophys. Res. Lett.*, 47, e2019GL085007, 715 <https://doi.org/10.1029/2019GL085007>, 2020.
- Ambaum, M. H. P. and Hoskins, B. J.: The NAO troposphere–stratosphere connection, *J. Clim.*, 15, 1969-1978, [https://doi.org/10.1175/1520-0442\(2002\)015<1969:TNTSC>2.0.CO;2](https://doi.org/10.1175/1520-0442(2002)015<1969:TNTSC>2.0.CO;2), 2002. Anstey, J., Gray, L. J., Mitchell, D. M., Baldwin, M. P., and Charlton-Perez, A. J.: The influence of stratospheric vortex displacements and splits on surface climate, *J. Clim.*, 26, 2668-2682, 720 <https://doi.org/10.1175/jcli-d-12-00030.1>, 2013.
- Baldwin, M. P. and Dunkerton, T. J.: Stratospheric harbingers of anomalous weather regimes, *Science*, 294, 581-584, <https://doi.org/10.1126/science.1063315>, 2001.
- Baldwin, M. P. and Thompson, D. W. J.: A critical comparison of stratosphere-troposphere coupling indices, *Q. J. R. Meteorol. Soc.*, 135, 1661-1672, <https://doi.org/10.1002/qj.479>, 2009.
- 725 Baldwin, M. P., Ayarzagüena, B., Birner, T., Butchart, N., Butler, A. H., Charlton-Perez, A. J., Domeisen, D. I. V., Garfinkel, C. I., Garny, H., Gerber, E. P., Hegglin, M. I., Langematz, U., and Pedatella, N. M.: Sudden stratospheric warmings, *Rev. Geophys.*, 59, e2020RG000708, <https://doi.org/10.1029/2020RG000708>, 2021.
- Black, R. X.: Stratospheric forcing of surface climate in the arctic oscillation, *J. Clim.*, 15, 268-277, 730 [https://doi.org/10.1175/1520-0442\(2002\)015<0268:SFOSCI>2.0.CO;2](https://doi.org/10.1175/1520-0442(2002)015<0268:SFOSCI>2.0.CO;2), 2002.
- Butler, A. H. and Gerber, E. P.: Optimizing the definition of a sudden stratospheric warming, *J. Clim.*, 31, 2337-2344, <https://doi.org/10.1175/jcli-d-17-0648.1>, 2018.
- Butler, A. H., Sjoberg, J. P., Seidel, D. J., and Rosenlof, K. H.: A sudden stratospheric warming compendium, *Earth Syst. Sci. Data*, 9, 63-76, <https://doi.org/10.5194/essd-9-63-2017>, 2017.
- 735 Butler, A. H., Seidel, D. J., Hardiman, S. C., Butchart, N., Birner, T., and Match, A.: Defining sudden stratospheric warmings, *Bull. Am. Meteorol. Soc.*, 96, 1913-1928, <https://doi.org/10.1175/BAMS-D-13-00173.1>, 2015.
- Cai, M. and Shin, C.-S.: A total flow perspective of atmospheric mass and angular momentum circulations: Boreal winter mean state, *J. Atmos. Sci.*, 71, 2244-2263, <https://doi.org/10.1175/JAS->

- 740 [D-13-0175.1](#), 2014.
- Charlton, A. J. and Polvani, L. M.: A new look at stratospheric sudden warmings. Part I: Climatology and modeling benchmarks, *J. Clim.*, 20, 449-469, <https://doi.org/10.1175/JCLI3996.1>, 2007.
- Chwat, D., Garfinkel, C. I., Chen, W., and Rao, J.: Which sudden stratospheric warming events are most predictable?, *J. Geophys. Res.: Atmos.*, 127, e2022JD037521, 745 <https://doi.org/10.1029/2022JD037521>, 2022.
- Colucci, S. J. and Kelleher, M. E.: Diagnostic comparison of tropospheric blocking events with and without sudden stratospheric warming, *J. Atmos. Sci.*, 72, 2227-2240, <https://doi.org/10.1175/JAS-D-14-0160.1>, 2015.
- Dall'Amico, M., Stott, P. A., Scaife, A. A., Gray, L. J., Rosenlof, K. H., and Karpechko, A. Y.: Impact of 750 stratospheric variability on tropospheric climate change, *Clim. Dyn.*, 34, 399-417, <https://doi.org/10.1007/s00382-009-0580-1>, 2010.
- Domeisen, D. I. V., Grams, C. M., and Papritz, L.: The role of North Atlantic–European weather regimes in the surface impact of sudden stratospheric warming events, *Weather Clim. Dynam.*, 1, 373-388, <https://doi.org/10.5194/wcd-1-373-2020>, 2020.
- 755 Edmon, H. J., Hoskins, B. J., and McIntyre, M. E.: Eliassen–Palm cross sections for the troposphere, *J. Atmos. Sci.*, 37, 2600-2616, [https://doi.org/10.1175/1520-0469\(1980\)037<2600:EPCSFT>2.0.CO;2](https://doi.org/10.1175/1520-0469(1980)037<2600:EPCSFT>2.0.CO;2), 1980.
- Hartmann, D. L.: The key role of lower-level meridional shear in baroclinic wave life cycles, *J. Atmos. Sci.*, 57, 389-401, [https://doi.org/10.1175/1520-0469\(2000\)057<0389:TKROLL>2.0.CO;2](https://doi.org/10.1175/1520-0469(2000)057<0389:TKROLL>2.0.CO;2), 2000.
- 760 Haynes, P. H., McIntyre, M. E., Shepherd, T. G., Marks, C. J., and Shine, K. P.: On the “downward control” of extratropical diabatic circulations by eddy-induced mean zonal forces, *J. Atmos. Sci.*, 48, 651-678, [https://doi.org/10.1175/1520-0469\(1991\)048<0651:OTCOED>2.0.CO;2](https://doi.org/10.1175/1520-0469(1991)048<0651:OTCOED>2.0.CO;2), 1991.
- Hitchcock, P. and Simpson, I. R.: The downward influence of stratospheric sudden warmings, *J. Atmos. Sci.*, 71, 3856-3876, <https://doi.org/10.1175/JAS-D-14-0012.1>, 2014.
- 765 Hoskins, B. J., McIntyre, M. E., and Robertson, A. W.: On the use and significance of isentropic potential vorticity maps, *Q. J. R. Meteorol. Soc.*, 111, 877-946, <https://doi.org/10.1002/qj.49711147002>, 1985.
- Huang, P. and Xie, S.-P.: Mechanisms of change in ENSO-induced tropical Pacific rainfall variability in a warming climate, *Nat. Geosci.*, 8, 922-926, <https://doi.org/10.1038/ngeo2571>, 2015.
- Jeong, J.-H., Kim, B.-M., Ho, C.-H., Chen, D., and Lim, G.-H.: Stratospheric origin of cold surge

- 770 occurrence in East Asia, *Geophys. Res. Lett.*, 33, <https://doi.org/10.1029/2006GL026607>, 2006.
- Jucker, M.: Are sudden stratospheric warmings generic? Insights from an Idealized GCM, *J. Atmos. Sci.*, 73, 5061-5080, <https://doi.org/10.1175/jas-d-15-0353.1>, 2016.
- Karpechko, A. Y. and Manzini, E.: Stratospheric influence on tropospheric climate change in the Northern Hemisphere, *J. Geophys. Res.: Atmos.*, 117, <https://doi.org/10.1029/2011jd017036>, 2012.
- 775 Karpechko, A. Y., Hitchcock, P., Peters, D. H. W., and Schneidereit, A.: Predictability of downward propagation of major sudden stratospheric warmings, *Q. J. R. Meteorol. Soc.*, 143, 1459-1470, <https://doi.org/10.1002/qj.3017>, 2017.
- Kidston, J., Scaife, A. A., Hardiman, S. C., Mitchell, D. M., Butchart, N., Baldwin, M. P., and Gray, L. J.: Stratospheric influence on tropospheric jet streams, storm tracks and surface weather, *Nat. Geosci.*, 8, 433-440, <https://doi.org/10.1038/ngeo2424>, 2015.
- 780 King, A. D., Butler, A. H., Jucker, M., Earl, N. O., and Rudeva, I.: Observed relationships between sudden stratospheric warmings and european climate extremes, *J. Geophys. Res.: Atmos.*, 124, 13943-13961, <https://doi.org/10.1029/2019JD030480>, 2019.
- Kodera, K., Mukougawa, H., Maury, P., Ueda, M., and Claud, C.: Absorbing and reflecting sudden stratospheric warming events and their relationship with tropospheric circulation, *J. Geophys. Res.: Atmos.*, 121, 80-94, <https://doi.org/10.1002/2015jd023359>, 2016.
- 785 Kunz, T. and Greatbatch, R. J.: On the northern annular mode surface signal associated with stratospheric variability, *J. Atmos. Sci.*, 70, 2103-2118, <https://doi.org/10.1175/JAS-D-12-0158.1>, 2013.
- Kuroda, Y. and Kodera, K.: Role of planetary waves in the stratosphere-troposphere coupled variability in the northern hemisphere winter, *Geophys. Res. Lett.*, 26, 2375-2378, <https://doi.org/10.1029/1999g1900507>, 1999.
- Lehtonen, I. and Karpechko, A. Y.: Observed and modeled tropospheric cold anomalies associated with sudden stratospheric warmings, *J. Geophys. Res.: Atmos.*, 121, 1591-1610, <https://doi.org/10.1002/2015jd023860>, 2016.
- 795 Liang, Z., Rao, J., Guo, D., and Lu, Q.: Simulation and projection of the sudden stratospheric warming events in different scenarios by CESM2-WACCM, *Climate Dynamics*, 59, 3741-3761, <https://doi.org/10.1007/s00382-022-06293-2>, 2022.
- Lindgren, E. A. and Sheshadri, A.: The role of wave-wave interactions in sudden stratospheric warming formation, *Weather Clim. Dynam.*, 1, 93-109, <https://doi.org/10.5194/wcd-1-93-2020>, 2020.

- 800 Liu, S.-M., Chen, Y.-H., Rao, J., Cao, C., Li, S.-Y., Ma, M.-H., and Wang, Y.-B.: Parallel comparison of major sudden stratospheric warming events in CESM1-WACCM and CESM2-WACCM, *Atmosphere*, 10, 679, <https://doi.org/10.3390/atmos10110679>, 2019.
- Lu, C. and Ding, Y.: Analysis of isentropic potential vorticities for the relationship between stratospheric anomalies and the cooling process in China, *Sci. Bull.*, 60, 726-738, <https://doi.org/10.1007/s11434-015-0757-4>, 2015.
- 805
- Lu, Q., Rao, J., Shi, C., Ren, R., Liu, Y., and Liu, S.: Stratosphere-troposphere coupling during stratospheric extremes in the 2022/23 winter, *Weather Clim. Extremes.*, 42, 100627, <https://doi.org/10.1016/j.wace.2023.100627>, 2023.
- Lu, Q., Rao, J., Shi, C., Guo, D., Fu, G., Wang, J., and Liang, Z.: Possible influence of sudden stratospheric warmings on the atmospheric environment in the Beijing–Tianjin–Hebei region, *Atmos. Chem. Phys.*, 22, 13087-13102, <https://doi.org/10.5194/acp-22-13087-2022>, 2022.
- 810
- Maycock, A. C. and Hitchcock, P.: Do split and displacement sudden stratospheric warmings have different annular mode signatures?, *Geophys. Res. Lett.*, 42, <https://doi.org/10.1002/2015gl066754>, 2015.
- 815 McAfee, S. A. and Russell, J. L.: Northern annular mode impact on spring climate in the western United States, *Geophys. Res. Lett.*, 35, <https://doi.org/10.1029/2008GL034828>, 2008.
- Mitchell, D. M., Charlton-Perez, A. J., and Gray, L. J.: Characterizing the Variability and Extremes of the Stratospheric Polar Vortices Using 2D Moment Analysis, *J. Atmos. Sci.*, 68, 1194-1213, <https://doi.org/10.1175/2010JAS3555.1>, 2011.
- 820
- Oehrlein, J., Polvani, L. M., Sun, L., and Deser, C.: How well do we know the surface impact of sudden stratospheric warmings?, *Geophys. Res. Lett.*, 48, e2021GL095493, <https://doi.org/10.1029/2021GL095493>, 2021.
- Polvani, L. M. and Waugh, D. W.: Upward wave activity flux as a precursor to extreme stratospheric events and subsequent anomalous surface weather regimes, *J. Clim.*, 17, 3548-3554, [https://doi.org/10.1175/1520-0442\(2004\)017<3548:UWAFAA>2.0.CO;2](https://doi.org/10.1175/1520-0442(2004)017<3548:UWAFAA>2.0.CO;2), 2004.
- 825
- Rao, J. and Garfinkel, C. I. J. E. R. L.: CMIP5/6 models project little change in the statistical characteristics of sudden stratospheric warmings in the 21st century, *Environ. Res. Lett.*, 16, 034024, <https://doi.org/10.1088/1748-9326/abd4fe>, 2021.
- Rao, J., Garfinkel, C. I., and White, I. P.: Predicting the downward and surface influence of the february

- 830 2018 and january 2019 sudden stratospheric warming events in subseasonal to seasonal (S2S)
models, *J. Geophys. Res.: Atmos.*, 125, e2019JD031919, <https://doi.org/10.1029/2019JD031919>,
2020.
- Rao, J., Ren, R., Chen, H., Yu, Y., and Zhou, Y.: The stratospheric sudden warming event in february
2018 and its prediction by a climate system model, *J. Geophys. Res.: Atmos.*, 123, 13,332-313,345,
835 <https://doi.org/10.1029/2018JD028908>, 2018.
- Rao, J., Garfinkel, C. I., Wu, T., Lu, Y., Lu, Q., and Liang, Z.: The january 2021 sudden stratospheric
warming and its prediction in subseasonal to seasonal models, *J. Geophys. Res.: Atmos.*, 126,
e2021JD035057, <https://doi.org/10.1029/2021JD035057>, 2021.
- Rao, J., Ren, R., Chen, H., Liu, X., Yu, Y., Hu, J., and Zhou, Y.: Predictability of stratospheric sudden
840 warmings in the Beijing climate center forecast system with statistical error corrections, *J. Geophys.*
Res.: Atmos., 124, 8385-8400, <https://doi.org/10.1029/2019JD030900>, 2019.
- Runde, T., Dameris, M., Garny, H., and Kinnison, D. E.: Classification of stratospheric extreme events
according to their downward propagation to the troposphere, *Geophys. Res. Lett.*, 43, 6665-6672,
<https://doi.org/10.1002/2016GL069569>, 2016.
- 845 Scaife, A. A., Arribas, A., Blockley, E., Brookshaw, A., Clark, R. T., Dunstone, N., Eade, R., Fereday, D.,
Folland, C. K., Gordon, M., Hermanson, L., Knight, J. R., Lea, D. J., MacLachlan, C., Maidens, A.,
Martin, M., Peterson, A. K., Smith, D., Vellinga, M., Wallace, E., Waters, J., and Williams, A.:
Skillful long-range prediction of European and North American winters, *Geophys. Res. Lett.*, 41,
2514-2519, <https://doi.org/10.1002/2014GL059637>, 2014.
- 850 Schmitz, G. and Grieger, N.: Model calculations on the structure of planetary waves in the upper
troposphere and lower stratosphere as a function of the wind field in the upper stratosphere, *Tellus.*,
32, 207-214, <https://doi.org/10.3402/tellusa.v32i3.10576>, 1980.
- Seviour, W. J. M., Mitchell, D. M., and Gray, L. J.: A practical method to identify displaced and split
stratospheric polar vortex events, *Geophys. Res. Lett.*, 40, 5268-5273,
855 <https://doi.org/10.1002/grl.50927>, 2013.
- Shi, Y., Evtushevsky, O., Milinevsky, G., Wang, X., Klekociuk, A., Han, W., Grytsai, A., Wang, Y., Wang,
L., Novosyadlyj, B., and Andrienko, Y.: Impact of the 2018 major sudden stratospheric warming on
weather over the midlatitude regions of Eastern Europe and East Asia, *Atmos. Res.*, 297,
<https://doi.org/10.1016/j.atmosres.2023.107112>, 2024.

- 860 Sjoberg, J. P. and Birner, T.: Transient tropospheric forcing of sudden stratospheric warmings, *J. Atmos. Sci.*, 69, 3420-3432, <https://doi.org/10.1175/JAS-D-11-0195.1>, 2012.
- Thompson, D. W. J. and Wallace, J. M.: Regional climate impacts of the northern hemisphere annular mode, *Science*, 293, 85-89, <https://10.1126/science.1058958>, 2001.
- White, I., Garfinkel, C. I., Gerber, E. P., Jucker, M., Aquila, V., and Oman, L. D.: The downward influence
865 of sudden stratospheric warmings: Association with tropospheric precursors, *J. Clim.*, 32, 85-108, <https://doi.org/10.1175/JCLI-D-18-0053.1>, 2019.
- Wu, Z. and Reichler, T.: Surface control of the frequency of stratospheric sudden warming events, *J. Clim.*, 32, 4753-4766, <https://doi.org/10.1175/JCLI-D-18-0801.1>, 2019.
- Yan, H., Yuan, Y., Guirong, T., and Yucheng, Z.: Possible impact of sudden stratospheric warming on the
870 intraseasonal reversal of the temperature over East Asia in winter 2020/21, *Atmos. Res.*, 268, <https://doi.org/10.1016/j.atmosres.2022.106016>, 2022.
- Yang, P., Bao, M., and Ren, X.: Influence of preconditioned stratospheric state on the surface response to displacement and split sudden stratospheric warmings, *Geophys. Res. Lett.*, 50, e2023GL103992, <https://doi.org/10.1029/2023GL103992>, 2023.
- 875 Yu, Y., Ren, R., Li, Y., Yu, X., Yang, X., Liu, B., and Sun, M.: Continental cold-air-outbreaks under the varying stratosphere-troposphere coupling regimes during stratospheric Northern Annular Mode events, *Clim. Dyn.*, <https://doi.org/10.1007/s00382-024-07275-2>, 2024.
- Zhang, L. and Chen, Q.: Analysis of the variations in the strength and position of stratospheric sudden warming in the past three decades, *Atmos. Ocean. Sci. Lett.*, 12, 147-154,
880 <https://doi.org/10.1080/16742834.2019.1586267>, 2019.