

1 **Introducing a new normalized cryospheric index (NCI) to categorize**
2 **sub-watersheds on arid environments**

3 Christopher Ulloa¹, Ayon García¹, Anouk Beniest²

4 ~~1 LICA-DICTEC/Water and Cryosphere Research Laboratory~~, University of Atacama, ~~Geology Department~~, Avenida
5 Copayapu #485, Copiapo, Chile

6 2 Department of Earth Sciences, Vrije Universiteit Amsterdam, De Boelelaan 1081, 1085 HV, Amsterdam, The Netherlands

7 Correspondence to: christopher.ulloa@uda.cl

8 **Abstract.** This study examines sub-watersheds in the arid northern region of Chile (26°41'–28°24'S), situated within the
9 broader Copiapó watershed. The primary water source for this watershed originates from cryospheric landforms. The region
10 exhibits pronounced physiographic and climatic diversity, with its northern sector situated within the South American Arid
11 Diagonal (SAAD), where cryospheric features display greater spatial isolation. The aim of this study is to quantify the water
12 volume contributed by distinct cryospheric landforms to regional watersheds. This study employs a classification methodology
13 to categorize cryospheric landforms within sub-watersheds, integrating an inventory of cryospheric landforms, historical snow
14 permanence derived from satellite imagery, published ice content and depth measurements (ERT and GPR). The Normalized
15 Cryospheric Index (NCI) is calculated for varying hydrological conditions and provides a means to compare potential water
16 volumes across sub-watersheds. The analysis reveals significant spatial variability in cryospheric landforms and their strategic
17 hydrological significance. Under average and low-precipitation conditions, the southern sub-watersheds of the Copiapó
18 ~~watershedriver-Basin~~ show the greatest water storage potential. The Montosa ~~R~~river (NCI = 0.93), Del Potro ~~R~~river (NCI =
19 0.89), Manflas ~~R~~river (NCI = 0.47), Estero Come Caballos (NCI = 0.34), and Cachitos ~~R~~river (NCI = 0.31) sub-watersheds
20 have been identified as strategic priority areas within the region for sustaining surface runoff and safeguarding water
21 availability. During high-snowfall periods, northern sub-watersheds in the Copiapo ~~watershedriver-Basin~~, such as Estero Come
22 Caballos, show elevated NCI values despite their limited cryospheric landforms. In contrast, the Montosa, Manflas, and Del
23 Potro sub-watersheds contain the most extensive cryospheric landforms and rank among the top of the sub-watersheds with
24 the highest NCI scores.

25 **1 Introduction**

26 Anthropogenic climate change impacts water availability, particularly in regions dependent on high-altitude water sources
27 within mountain systems (Beniston & Stoffel, 2014, Beniston et al., 2018). Changes in the cryosphere that will occur within
28 the next century include a change from solid to liquid precipitation, declining volumes of cryospheric landforms, snow lines

Con formato: Inglés (Estados Unidos)

Con formato: Inglés (Estados Unidos)

Con formato: Inglés (Estados Unidos)

moving to higher altitudes, and shorter snow seasons. With these changes, the predictions are that water resources in mountain areas will become increasingly scarce. In the arid zones of the Chilean Atacama Desert glaciers continue to shrink- (Masiokas, et al., 2020) and centennial reconstructions of rainfall patterns on the scale of hundreds of years, show a decline in rainfall in the Central Zone and the “Norte Chico” natural regions of Chile (Le Quesne et al., 2006; Minvielle and Garreaud, 2011; Morales et al., 2012; Bozkurt et al., 2017). In addition, the occurrence of mega droughts has intensified since 2010 (Garreaud et al., 202019).

Within this climatic context, the Copiapó River exhibits a mixed hydrological regime with very low discharges, typically ranging between $\sim 1.49 \text{ m}^3 \text{ s}^{-1}$ and $1.82 \text{ m}^3 \text{ s}^{-1}$ at the La Puerta gauging station, and strongly modulated by contributions from its main tributaries, the Manflas, Jorquera and Pulido ~~R~~ivers, whose headwaters originate in high-altitude cryospheric environments. Long-term analyses indicate that mean annual flows rarely exceed $2 \text{ m}^3 \text{ s}^{-1}$, and the ~~watershedbasin~~ shows recurrent years of negative water balance, leading to progressive aquifer depletion (Suárez et al., 2014). These persistently low flows and the strong dependence on upstream snow and ice melt highlight the structural vulnerability of the runoff of the Copiapó River to changes in the cryosphere.

Because of the high altitude of the Andean cryosphere, precipitation is captured from air masses, predominantly in the form of snow. Generally, snow precipitation occurs during the winter (June-September), although the northern parts also receive snow during summer due to the Tropical Monsoon (i.e. Bolivian winter), which takes place from December to February (Valdivieso et al., 2022). In high-altitude Andean environments, snowfall does not directly translate into liquid runoff but may be preserved over different timescales through distinct cryospheric processes. Under favorable mass-balance conditions, repeated snow accumulation and firnification lead to the formation of debris-free glaciers. Where glacier ice becomes mantled by rockfall or talus material, debris-covered glaciers can develop, reducing ablation rates and enhancing ice preservation. In parallel, a significant fraction of snowfall infiltrates coarse debris slopes, where it refreezes and is preserved as ground ice within ice-rich permafrost landforms such as rock glaciers, protalus lobes, and gelifluction taluses. The melting of these cryospheric systems generates a sustained discharge of water, even during snow-free dry seasons. Given the extremely limited rainfall in the Andes, streamflow in the Copiapó River is strongly dependent on snowmelt and cryospheric water storage. (Masiokas et al., 2019). However, the contribution of cryospheric water storage to runoff varies substantially among cryospheric landforms types and occurs over contrasting temporal scales. Debris-free glaciers represent the most direct and dynamic cryospheric contribution to river discharge, responding seasonally to atmospheric forcing and sustaining spring–summer runoff through glacier melt. Debris-covered glaciers may also contribute meltwater, but their hydrological response is attenuated and delayed due to the insulating effect of supraglacial debris, which reduces melt rates per unit area. The interannual variability of this meltwater contribution is strongly modulated by debris cover thickness and continuity, which can substantially alter glacier hydrological response across seasons and years (Burger et al., 2019).

In contrast, ice-rich permafrost landforms, such as rock glaciers, protalus lobes, and gelifluction taluses, function predominantly as long-term cryospheric water reserves rather than active meltwater sources. Recent studies in the Central Andes indicate that many rock glaciers exhibit near-zero mass loss over recent decades, suggesting that internal ice melt

currently contributes only marginally to river runoff (Ferri et al., 2020). Their primary hydrological significance therefore lies in their capacity to store frozen water over decadal to centennial timescales, potentially buffering future water availability under continued climatic warming. [Multi-decadal photogrammetric reconstructions using aerial, satellite, and UAV data have further revealed significant kinematic variability in rock glaciers of the Dry Andes, highlighting their dynamic nature and importance as cryospheric components \(Vivero et al., 2021\). In addition, future megadrought projections for the Southern Andes suggest that glacier meltwater contributions may decline markedly, underscoring the critical need to understand and protect all cryospheric water reserves, including periglacial landforms \(Ayala, 2025\).](#)

Seasonal snow represents a distinct cryospheric component with a rapid hydrological response. During wet years, snowmelt dominates river discharge, whereas during dry years runoff becomes increasingly dependent on long-term cryospheric storage at high elevations (Ohlanders et al., 2013). Although snow sublimation losses in the Andes can reach up to ~70% of the seasonal snowpack (Jara et al., 2021), the remaining snowmelt still exerts a first-order control on interannual streamflow variability.

The source of water in the watersheds of the Atacama Desert originate thus predominantly from the volumes of ice that are kept in cryospheric landforms, but the potential volumes of water that are present in these cryospheric landforms are highly variable and depend on both the type and the spatial distribution. We hypothesize that the hydrological relevance of arid sub-watersheds can be quantified by integrating long-term ice storage and seasonal snow permanence into a single normalized index, and that the resulting sub-watershed ranking remains robust under realistic measurement and modelling uncertainty.

Testing this hypothesis requires (a) a complete cryospheric inventory at the [watershedbasin](#) scale, (b) physics-based ice-volume estimates for both glacial and ice-rich permafrost landforms, and (c) explicit propagation of uncertainty through the index calculation. General methodologies to classify and prioritize watersheds are based on morphometric analysis (Rahaman et al., 2015), watershed hydrology (Wolfe, et al., 2019) and ecological and water quality factors. There are also numerous studies covering watershed health in which protection frameworks have been developed based on ecological and hydrological parameters providing approaches on watershed prioritization (Jaiswal et al., 2015; Ahn & Kim, 2017; Sriyana, 2019; Basuki et al., 2022). Few studies incorporate all cryospheric components into watershed management or conservation strategies. Peng et al. (2022) developed an integrated index to evaluate cryospheric changes across the Northern Hemisphere, emphasizing the influence of cryospheric dynamics on regional hydrology. Nonetheless, there is currently a lack of research addressing cryospheric landforms and snow within arid watershed contexts.

To address this gap, this study delivers three integrated outputs for the Copiapó watershed: (i) an updated, GIS-ready inventory of 1862 cryospheric landforms covering glaciers, debris-covered glaciers, rock glaciers, gelifluction taluses, and protalus lobes; (ii) a unified physics-based framework for estimating ice volume across glacial and ice-rich permafrost landforms, combining the Farinotti et al. (2019) consensus product with viscoplastic creep inversion ([Cicoira et al., 2021](#)) constrained by Sentinel-1 DInSAR velocities; and (iii) a Normalized Cryospheric Index (NCI) that ranks sub-watersheds by their combined long-term ice storage and seasonal snow contribution, with Monte Carlo propagation of uncertainty to ensure ranking robustness. We




Figure 1. Location map of the Copiapó watershed and the different sub-watersheds (Zone 19S). The codes in each sub-watershed represent the official numbering that the General Water Directorate assigns to each sub-watershed.

104
105
106

Table 1. List of sub-watersheds with cryospheric components in the study area.

Sub-watershed Code	Sub-watershed name	Sub-watershed area km ²
3400	Quebrada Monardes	803.6
3401	Figueroa R iver	924.2
3402	Estero Come Caballos	880.6
3403	Cachitos R iver	734.1
3410	Vizcachas de Pulido river	611.9
3411	Ramadillas R iver	366.8
3412	Del Potro R iver	470.6
3413	Montosa R iver	413.2
3420	Manflas R iver	728.1
3440	Quebrada San Andres	1476.3
3441	Quebrada Paipote	1493.8
3443	Quebrada Martinez	1513.1

107 **2 Regional setting**

108 **2.1 Study Area**

109 The Copiapó watershed is composed of 27 sub-watersheds, distributed along the high mountain belt that forms the headwaters
110 of the basin. A sub-watershed is defined here as a watershed unit with a lower Strahler stream order than the main watershed.
111 In our research area, the Copiapó watershed corresponds to the main watershedbasin and contains rivers with a maximum
112 Strahler stream order of nine. Within this hierarchy, sub-watersheds draining from the headwaters typically initiate with
113 Strahler orders of one to three and progress downstream until they merge into larger tributaries with stream orders of up to
114 nine. By distinguishing among sub-watersheds and their associated cryospheric landforms, we can perform both quantitative
115 and qualitative assessments of the current potential water volumes stored within cryospheric environments.
116 To classify a sub-watershed as a cryospheric watershed, we applied objective and reproducible criteria based on its cryospheric
117 components. A sub-watershed is considered cryospheric only when it fulfilled two conditions: (i) the presence of at least one
118 active cryospheric landforms (debris-free glacier, debris-covered glacier, rock glacier, protalus lobe, or gelifluction slope)

mapped in the regional inventory of glaciers and the periglacial environment of the Atacama Desert (García et al., 2017); and (ii) measurable seasonal snow, expressed as long-term snow permanence derived from MODIS fractional snow cover (Rittger et al., 2013). Sub-watersheds fulfilling both requirements (≥ 1 cryospheric landforms + snow permanence) were classified as cryospheric.

Applying these criteria resulted in 12 sub-watersheds being identified as cryospheric watersheds (Fig. 1; Table 1). These sub-watersheds are located entirely within high-elevation sectors of the [Copiapó watershedbasin](#) (mean elevation 3,617 m a.s.l.), coincide with areas of high permafrost probability in the global permafrost model of Gruber (2012), and contain 97% of the cryospheric landforms mapped in the Copiapó [watershedbasin](#). This classification is therefore geomorphologically justified, reproducible, and consistent with independent permafrost modelling and regional cryospheric inventories.

The Copiapó watershed (Fig. 1) is located in the southern limit of the Atacama Desert, the driest desert of the world (Clarke, 2006), near to the South American Arid Diagonal or SAAD (Zech et al., 2008). The annual precipitation in Copiapó from 1795 to the present has an annual mean of 22.5 mm (Izquierdo et al., 2024). Among the arid [watershedsbasins](#) in northern Chile, the Copiapó watershed is regarded as one of the most vulnerable (Suárez et al., 2014). Because of the limited availability of freshwater and the overexploitation of superficial water and groundwater, the deepening of the water table is accelerating and cryospheric landforms and snow are reducing (Casassa et al., 2007; Vuille et al., 2008). Despite this very limited liquid precipitation in the lower areas, e.g. around the city of Copiapó, there is a flow of water in the rivers of the Copiapó watershed year round. This water comes from the Andean cryosphere that is known to host and store large amounts of fresh water (Bolch and Marchenko, 2006; Bórquez et al., 2006; ~~Azocar and Brenning~~[Azócar and Brenning](#), 2010; Schaffer et al., 2019).

The glacial and periglacial inventory of the Copiapó watershed indicates a transition from areas that are dominated by glacial environments in the south, to areas that are dominated by periglacial environments in the north (García et al., 2017). In the Copiapó watershed, the occurrence of snow begins above 3,500 m of altitude. In these extreme environments, the water budget is significantly influenced by snow sublimation. From 2001 to 2016, approximately 70% of the snow balance was lost to sublimation, effectively reducing the available water volume (Jara et al., 2021). In semi-arid regions of Chile and Argentina, snow accumulation and subsequent melt processes contribute substantially to surface runoff in numerous Andean [R](#)iver [W](#)atersheds (Masiokas et al., 2006; Favier et al., 2009).




Figure 2. Main cryospheric landforms. Field photographs from the Copiapó watershed on the left, with a schematic interpretation on the right of. a) Debris-free glacier and debris-glaciers, b) Rock glaciers of the Del Potro River sub-watershed. c) Gelifluction taluses of Montosa River sub-watershed, d) Protalus lobes of the La Laguna sub-watershed in Vizcachas de Pulido sub-watershed.

Con formato: Fuente: 9 pto

2.2 Hydrological background of the cryosphere and conceptual framework for ice-volume estimation

The hydrology of arid Andean Watershedsbasins is dominated by cryospheric processes. In the Copiapó watershed, year-round streamflow persists despite minimal precipitation at mid- to low elevations, indicating that meltwater from snow, glaciers, and

ice-rich permafrost constitutes the primary source sustaining discharge. Long-term records from the Dirección General de Aguas (DGA) hydrometric network show a pronounced summer peak in river flow, when rainfall is nearly zero, together with a high baseflow index, confirming a melt-dominated regime and the central hydrological role of the cryosphere in this region. These patterns are consistent with previous studies across the semi-arid Andes (Favier et al., 2009; Gascoin et al., 2011; Ayala et al., 2016), where snow and frozen water bodies act as short- and long-term ice storage modulating runoff during dry years (Table 2).

Cryospheric contributions, however, operate over markedly different response times. Seasonal snow responds at sub-seasonal scales, controlling the onset and magnitude of spring–summer discharge. Debris-free and debris-covered glaciers respond at seasonal to multiannual scales, providing sustained meltwater even in years of limited snowfall. Ice-rich permafrost landforms, including rock glaciers, protalus lobes, and gelifluction slopes, contribute at decadal to centennial scales and represent the most climatically buffered ice storage in the watershed. Capturing these contrasted response times requires not only quantifying the spatial distribution of cryospheric landforms but also applying a coherent conceptual framework for estimating their ice volumes.

The classification of debris-covered glaciers adopted in this study follows Kirkbride (2011), who defines them as glaciers whose ablation zone is largely or entirely covered by supraglacial debris across its full width. This distinction is important in the study area, where debris-covered glaciers occur in close spatial association with rock glaciers but represent fundamentally different cryospheric processes: debris-covered glaciers are dynamic ice bodies in which the supraglacial covered modulates ablation, whereas rock glaciers are permafrost landforms whose movement is driven by internal ice deformation in an ice-rich debris matrix.

Table 2. Contribution to streamflow (%) indicates the proportion of total river discharge supplied by cryospheric meltwater (glaciers and/or seasonal snow), as quantified in each study. Modified from Schaffer et al (2019).

Location / Watershed	Latitude	Hydrological year	Contribution to streamflow (%)	Cryospheric component evaluated	Reference
Huasco watershed	~29°S	2003/04 and 2007/08	3–23	Glaciers (debris-free + debris-covered)	Gascoin et al. (2011)
Elqui watershed	~30°S	2003	4–9	Glaciers (mainly debris-free)	Favier et al. (2009)
Elqui watershed	~30°S	2011	13	Glaciers (debris-free)	Pourrier et al. (2014)
Juncal River watershed	32–36°S	2005/06 and 2008/09	10 and 31	Glaciers (debris-free), snowmelt	Ragetti & Pellicciotti (2012)
Juncal River watershed	32–36°S	2005/06 and 2008/09	16 and 44	Glaciers (debris-free + debris-covered)	Rodríguez et al. (2016)
Juncal River watershed	32–36°S	2011–2012	2–50	Seasonal snow + glacier melt	Rodríguez et al. (2016)
Yeso River watershed	~33.5°S	2014/15	~42 and ~67	Glaciers + snowmelt (during Megadrought)	Ayala et al. (2016)

Central Chilean Andes	–	1968/69	67	Exposed glaciers (exceptionally dry year)	Peña & Nazarala (1987)
Central Chile (regional)	–	Long-term	–	Glacio-hydrological long-term contribution	Ayala et al. (2020)

Estimating ice volume in mountain cryospheric systems relies on methodological approaches that vary in complexity, physical basis, and data requirements. Traditionally, regional and global assessments have applied empirical area–volume or area–thickness relationships (e.g., Bahr et al., 1997; Jones et al., 2018). These models provide valuable first-order estimates and remain particularly useful in data-sparse regions. However, by construction, they do not explicitly account for glacier geometry, internal deformation, thermal regime, or spatially variable creep rates, which are processes that exert strong control on ice thickness in both glacial and periglacial landforms.

More recent advances have increasingly emphasized physically based approaches that better represent the mechanics of glacier flow and permafrost creep. For debris-free glaciers, inversion techniques such as the model designed by Farinotti et al. (2017) combine mass conservation, surface slope, and Glen’s flow law to derive spatially distributed ice thicknesses (Fig. 3), and form the basis of the global consensus thickness product of Farinotti et al. (2019). At continental scales, Millan et al. (2022) further demonstrated the value of integrating satellite-derived surface velocities into ice-flow inversion, substantially refining glacier thickness estimates and highlighting the importance of kinematic constraints.

In parallel, physically informed modelling has also progressed for ice-rich permafrost landforms. Viscoplastic formulations, such as those proposed by Cicoira et al. (2021), explicitly link surface velocity, basal shear stress, rheology, and thermal conditions to estimate the thickness of ice-rich layers in rock glaciers. Building on these concepts, Hu et al. (2023) applied deformation-based approaches to regional rock-glacier inventories, illustrating the feasibility of estimating ice content in permafrost landforms using remote sensing–derived kinematics. These developments reflect a broader methodological shift toward deformation-based, physically constrained estimates of ice volume across both glacial and periglacial environments.

In the Andes, the applicability of these physically based approaches is reinforced by the growing availability of satellite-derived deformation data. For example, Villarroel et al. (2018) used Sentinel-1 DInSAR to map surface displacement of more than 2,100 rock glaciers in the Dry Andes, while Vidal-Páez et al. (2024) applied PAZ-SAR SBAS-DInSAR to detect millimetric rock-glacier motion in the Chilean Andes. Together, these studies demonstrate that interferometric and feature-tracking techniques enable the detection of low-magnitude deformation, allowing thickness estimation even in low-creep permafrost landforms such as protalus lobes and gelifluction slopes.

In this study, we adopt an integrated conceptual framework that applies:

- (1) the physically based Farinotti model to debris-free glaciers;
- (2) a geometry-consistent approach for debris-covered glaciers using thickness gradients from Farinotti et al. (2019); and
- (3) a unified viscoplastic formulation (Cicoira et al., 2020; Cicoira et al., 2021) to all permafrost cryospheric landforms including rock glaciers, protalus lobes, and gelifluction slopes supported by DInSAR-derived velocities.

206 This unified framework ensures physical consistency across cryospheric landforms with contrasting dynamics and provides
207 the conceptual foundation required for robust estimation of cryospheric ice volumes prior to their integration into the
208 Normalized Cryospheric Index (NCI).

209 2.3 Glacier and Periglacial Environment

210 The Copiapó watershed hosts a diverse assemblage of cryospheric landforms (Fig. 2), including debris-free glaciers, debris-
211 covered glaciers, rock glaciers, gelifluction taluses, and protalus lobes, as documented in the regional inventory of García et
212 al. (2017). These landforms exhibit a marked spatial variability across the [watershedbasin](#), reflecting strong controls by
213 elevation and topography.

214 Based on the regional inventory of García et al. (2017), a total of 1862 cryospheric landforms have been mapped within the
215 Copiapó watershed: 62 debris-free glaciers, 16 debris-covered glaciers, 173 rock glaciers, 848 gelifluction taluses, and 763
216 protalus lobes. Gelifluction taluses (46%) and protalus lobes (41%) thus dominate the cryospheric assemblage of the
217 [watershedbasin](#), while debris-free and debris-covered glaciers together account for less than 4% of the inventoried landforms.
218 This numerical distribution reinforces the periglacial character of the Copiapó cryosphere and underscores the relevance of
219 ice-rich permafrost landforms for ice-volume estimations at the [watershedbasin](#) scale. This inventory was not originally
220 developed under the Randolph Glacier Inventory (RGI) framework, as it predates the standardization of global glacier datasets
221 in the study region. Instead, the inventory provides a detailed regional mapping of cryospheric landforms, including not only
222 glaciers but also ice-rich permafrost features such as rock glaciers, gelifluction slopes, and protalus lobes. This distinction is
223 particularly relevant because global glacier inventories, such as RGI, are primarily focused on glacier ice bodies and typically
224 do not include periglacial landforms, which are rarely mapped in a systematic manner. As a result, the regional inventory used
225 here captures a broader spectrum of cryospheric components that are especially significant in arid mountain environments,
226 where non-glacial frozen water storage may play an important hydrological role.

227 In general, cryospheric landforms are distributed discontinuously along the high Andes, with notable differences between
228 northern and southern sectors. In areas such as Nevado Jotabeche and Los Tronquitos, landforms are predominantly located
229 on high-elevation ridges and peaks, whereas in the Del Potro region they are more commonly confined to valley bottoms
230 between steep slopes.

231 Gelifluction taluses represent one of the most widespread landforms in the study area and are typically associated with high-
232 elevation environments. In the southern sector, they frequently occur in close proximity to debris-free glaciers, forming
233 coupled glacial–periglacial systems. In contrast, in the northern sector, gelifluction features tend to appear as smaller and more
234 isolated landforms.

235 Rock glaciers are mainly distributed along steep valley flanks at intermediate to high elevations, often occupying topographic
236 depressions or confined valley settings. Debris-free glaciers are generally restricted to the highest elevations and, particularly
237 in the southern sector, are commonly associated with adjacent periglacial landforms.

238 Protalus lobes are especially abundant in some northern areas, where they develop on slopes of varying gradients and exhibit
239 a range of sizes and morphologies. Debris-covered glaciers are comparatively scarce and are typically found in the southern
240 sector, occurring as small bodies located downslope of debris-free glaciers.

241 Overall, the distribution of cryospheric landforms in the Copiapó watershed reflects a transition from glacial-dominated
242 systems in the south to predominantly periglacial environments in the north, consistent with regional climatic and topographic
243 gradients (Fig. 4).

244 The regional inventory of García et al. (2017) predates the RGIK (Rock Glacier Inventories and Kinematics; IPA Action
245 Group) guidelines, but its mapping workflow multi-source satellite photo-interpretation supported by field reconnaissance,
246 identification of active rock glaciers from diagnostic geomorphological criteria (frontal ramp, furrows and ridges), and a
247 GLIMS-compatible attribute database is methodologically consistent with the principles later codified by RGIK. The inventory
248 extends this scope to include gelifluction taluses and protalus lobes, two ice-rich permafrost landforms that dominate the arid
249 Atacama cryosphere but fall outside the RGIK rock-glacier framework.




Figure 3. a) Distribution map of the debris-free glacier thickness from Farinotti et al. (2019). b) Jotabeche volcano cryospheric landforms distribution c) The Del Potro hill cryospheric landforms distribution d) The Tronquitos hill cryospheric landforms distribution. Basemap: Esri, Maxar, Earthstar Geographics, and the GIS User Community.

Con formato: Centrado




Figure 4. a) Distribution map of the cryospheric landforms inventoried in the Atacama region from García et al. (2017). b) Jotabeche Volcano cryospheric landforms distribution c) The Del Potro Hill cryospheric landforms distribution d) The Tronquitos Hill cryospheric landforms distribution. Basemap: Esri, Maxar, Earthstar Geographics, and the GIS User Community.

261 **3 Methodology**

262 **3.1 Survey of cryospheric landforms**

263 To quantify the ice volume in the Copiapó watershed, we used existing cryospheric inventories without performing new
264 delineations. The regional inventory of glaciers and periglacial landforms of the Atacama Desert (García et al., 2017) was
265 employed directly to extract polygons of debris-covered glaciers, rock glaciers, protalus lobes, and gelifluction slopes. This
266 inventory represents the only comprehensive and validated dataset that explicitly identifies periglacial cryospheric landforms
267 in northern Chile, and its polygons were incorporated without modification into our analysis.

268 Debris-free glaciers were obtained exclusively from the Randolph Glacier Inventory (RGI) v6.0 (RGI Consortium, 2017). For
269 these glaciers, we did not compute ice thickness ourselves. Instead, we used the pre-computed global ice-thickness product of
270 Farinotti et al. (2019), which is physics-based and already calibrated on RGI glacier outlines (Fig. 3). Because the Farinotti
271 model requires RGI geometry as input, the use of RGI v6.0 ensures direct compatibility with the global thickness dataset. Ice
272 volumes for debris-free glaciers were therefore calculated solely by spatially integrating the published Farinotti et al. (2019)
273 thickness values over the RGI glacier polygons within the Copiapó watershedbasin. This approach avoids any reinterpretation
274 of glacier geometry or re-estimation of ice thickness, ensuring methodological transparency and consistency with the global
275 consensus product.

276 The joint use of the inventory developed by García et al. (2017) and RGI v6.0 glacier inventory is thus complementary. RGI
277 provides standardized glacier boundaries and their associated Farinotti-derived thickness fields for debris-free glaciers,
278 whereas García et al. (2017) provides the spatially explicit identification of debris-covered glaciers and periglacial landforms
279 that are not represented in global glacier datasets. By relying directly on these inventories and on the published model outputs
280 of Farinotti et al. (2019), our workflow remains fully reproducible and avoids introducing new mapping or thickness-estimation
281 steps.

282 Mapping uncertainty for all polygons was propagated using the one-pixel boundary buffer method of Paul et al. (2013),
283 ensuring consistent treatment of spatial uncertainty across glacial and periglacial cryospheric landforms.

284 **3.2 Survey of snow cover: MODIS-derived snow permanence**

285 The MOD10A1 Fractional Snow Cover (FSC) product from NASA was used to characterize snow based on long-term patterns
286 of snow permanence. Each MOD10A1 image provides, for every 500×500 m pixel, a value representing the fractional snow
287 cover within that pixel (from 0 to 1), derived through a spectral unmixing algorithm (Rittger et al., 2013). This sub-pixel
288 representation allows a more accurate depiction of discontinuous or patchy snow, which is common in high-elevation arid
289 regions. Instead of relying on isolated scenes, we processed the complete daily MODIS time series (2000–2022) to obtain both
290 the instantaneous snow-covered area and the frequency of snow permanence at the sub-watershed scale. Although cloud
291 contamination can introduce occasional misclassifications, the product maintains approximately 90% accuracy under ideal

conditions (Hall et al., 2006; Hall et al., 2002; Jiang et al., 2024), and is suitable for long-term characterization of snow permanence.

From the daily FSC series, each MODIS image provides the fractional snow cover for every 500×500 m pixel. Using this long-term record, fractional snow cover (FSC) was used as the core variable to represent the proportion of snow within each pixel in each scene. In addition, we calculated snow permanence as the fraction of days during the 23-year period in which each pixel exhibited snow-covered conditions according to its FSC value. Snow permanence is therefore not an independent MODIS layer, but a temporal aggregation derived directly from the daily FSC maps, summarizing the frequency and recurrence of snow presence across the watershed basin.

In this framework, FSC was used as the primary input for constructing daily snow permanence maps, and snow permanence, computed as the proportion of snow-covered days over the full time series, was incorporated into the Normalized Cryospheric Index (NCI) as the indicator of short-term cryospheric water resources. This approach ensures that the characterization of snow permanence reflects both their spatial extent and their temporal recurrence. long-term mean FSC maps. For each pixel, the daily FSC values were averaged over the full 2000–2022 record to obtain a time-mean fractional snow cover. This per-pixel mean FSC was then multiplied by the pixel area (0.25 km^2 for the 500×500 m MODIS grid) and summed across all pixels within each sub-watershed to yield a mean snow-covered area in km^2 . It is this spatially integrated snow-covered area (km^2) that is incorporated into the Normalized Cryospheric Index (NCI) as the indicator of short-term cryospheric water resources, ensuring that both the fractional snow signal at the pixel level and the spatial extent of the sub-watershed are accounted for in the index.

3.3 Field work for ground truthing

Since the mapping workflow is primarily based on remote sensing, satellite imagery, and existing cryospheric inventories, we incorporated targeted field observations and independent geophysical datasets to validate both the identification of cryospheric landforms and the ice-thickness values used in this study. Fieldwork was carried out in the Montosa (03413) and Del Potro (03412) sub-watersheds, which together represent approximately 10% of the Copiapó watershed basin and contain accessible examples of the main cryospheric landforms present in the region.

During these campaigns, photographic and geomorphological observations were conducted to verify the surface expression of mapped cryospheric landforms, including gelifluction slopes, protalus lobes, and rock glaciers. These observations allowed us to assess the representativeness of the polygons obtained from existing inventories.

Beyond surface verification, independent geophysical datasets were used to constrain and validate ice-thickness estimates. For debris-free glaciers, ice thickness values from the Farinotti et al. (2019) global product were compared against airborne ground-penetrating radar (GPR) profiles collected by the Dirección General de Aguas (DGA, 2014) over the Del Potro and Tronquitos glaciers.

For ice-rich permafrost landforms, Electrical Resistivity Tomography (ERT) measurements were conducted on a representative gelifluction slope and on an active rock glacier within the Montosa sub-watershed. These ERT profiles constrain the depth and

325 continuity of ice-rich layers and support the thickness ranges adopted for periglacial cryospheric landforms in the modelling
326 framework.
327 Together, these field observations and geophysical datasets provide independent validation for both the spatial delineation and
328 the subsurface characteristics of cryospheric landforms within the Copiapó watershed, strengthening the robustness of the ice-
329 volume estimates applied in this study.

330 **3.3.1 Geophysical survey on gelifluction slopes and rock glaciers**

331 Electrical Resistivity Tomography (ERT) was employed to characterize subsurface ice-rich layers in selected periglacial
332 landforms and to provide observational constraints for the ice-thickness values used in the modelling framework. ERT is a
333 multi-electrode resistivity method that generates 2-D vertical sections of the subsurface by injecting electrical current into the
334 ground and measuring the resulting potential differences across electrode arrays. The field configuration used in this study
335 (Table 3) consisted of 36 electrodes operated with Wenner, Schlumberger, and Dipole–Dipole arrays.




Figure 5. Photography of the fieldwork. a) And b) the Lobulated morphology of studied gelifluction taluses, which are not covered in snow; c) and d) the geoelectric line cable; e) and f) the used equipment, differential GPS and resistivity meter ARES II measuring a resistivity point diagram distribution.

Fieldwork focused on two representative cryospheric landforms within the Montosa sub-watershed: a gelifluction slope and an active rock glacier (Fig. 5 and Fig. 6). These sites were selected because they exhibit clear geomorphological indicators of ice-rich permafrost and provide suitable access for geophysical surveys. The ERT profiles allowed direct identification of high-resistivity zones ($>20\text{--}50\text{ k}\Omega\cdot\text{m}$), characteristic of massive or highly saturated ice-rich permafrost bodies.

Table 3 Used geophysical equipment for fieldwork of this study

Equipment	Amount
Resistivity meter: ARES II	1
30 electrode box	5

10 channels multicable II (60 m)	10
Car Batteries	2
Differential GPS	1

345 The survey layout required a 600-m profile organized through multicables arranged at 90° angles. A high-precision differential
 346 GPS (“South Galaxy”) was used to set the start and end points of each line, ensuring alignment and minimizing installation
 347 error. Electrode spacing was fixed at 5 m, with position control verified using a combination of GPS and measured markers at
 348 30-m intervals. All electrodes were tested prior to each acquisition, and any with unacceptable standard deviation values were
 349 repositioned.

350 These ERT measurements provide direct ground validation of subsurface ice for both gelifluction taluses and rock glaciers,
 351 supporting the thickness ranges applied to ice-rich permafrost cryospheric landforms in our volume estimations and
 352 complementing the GPR-based glacier-thickness validation described in Section 3.3.



353
 354 **Figure 6. Photography of the fieldwork rock-glacier surface. (a) Intelligent multi-electrode cable reel used for deploying the ERT**
 355 **array across the periglacial terrain. (b) Installation of the 600-m geoelectric line. (c) Operation of the ARES II resistivity meter**
 356 **during data acquisition.**

Before every measurement, all electrodes were tested to detect electrodes with high standard deviation in resistivity. If the deviation exceeded acceptable limits, the electrodes were repositioned until the deviation fell within an acceptable range to ensure measurement accuracy.

3.4 DInSAR kinematic analysis for periglacial cryospheric landforms

To characterize surface deformation across glacial and periglacial cryospheric landforms and to provide kinematic constraints for ice-thickness modelling, we processed Sentinel-1 IW Single Look Complex (SLC) images using the ESA SNAP v9.0 toolbox. Interferometric pairs were selected based on perpendicular baselines <150 m and temporal baselines <24 days to ensure coherence in high-altitude terrain (Table 4).

Table 4 Sentinel-1 interferometric image pairs and associated temporal and perpendicular baselines tested.

Reference Image Date	Secondary Image Date	Orbit Type	Temporal Baseline (Days)	Perpendicular Baseline	Interferogram status
01-03-2024	13-03-2024	Ascending	12	55.35	Inconsistent
25-03-2024	06-04-2024	Ascending	12	40.12	Inconsistent
03-03-2016	27-03-2016	Descending	24	9.39	Coherent
05-01-2015	29-05-2015	Ascending	24	-121.04	Inconsistent
29-01-2015	22-02-2015	Descending	24	25.41	Inconsistent

SLC scenes were co-registered using the TOPS burst co-registration workflow. Interferograms were generated, filtered using a Goldstein filter ($\alpha = 0.8$), and corrected for topographic phase using the 30-m SRTM DEM. Phase unwrapping was performed with SNAPHU in external mode. Line-of-sight (LOS) displacement was obtained from the unwrapped phase:

$$d_{LOS} = \frac{\lambda}{4\pi} \Delta\phi \tag{1}$$

Mean surface velocities were extracted for each cryospheric landforms polygon (debris-free glaciers, debris-covered glaciers, rock glaciers, pro talus lobes, gelifluction slopes). These DInSAR-derived velocities were incorporated as dynamic constraints

for the viscoplastic modelling of permafrost cryospheric landforms and as supplementary kinematic information in glacier-thickness interpretation.

3.4.1 Velocity uncertainty

Uncertainty in DInSAR-derived surface velocity was incorporated by explicitly estimating the phase noise associated with interferometric coherence and temporal baseline. Following Hanssen (2001), the standard deviation of the interferometric phase for a multilooked interferogram is:

$$\sigma_{\phi} \approx \sqrt{\frac{1-\gamma^2}{\gamma^2 L}} \quad (2)$$

Where γ is the interferometric coherence and L the number of equivalent looks. For Sentinel-1 SLC data used in this study, the typical modeled coherence is $\gamma=0.97$, the temporal baseline is $\Delta t=24$ days, and the radar wavelength is $\lambda=0.0555$ m.

Phase noise is converted to line-of-sight displacement noise using:

$$\sigma_{\phi} = \frac{\lambda}{4\pi} \sigma_{\phi} \quad (3)$$

And subsequently velocity uncertainty:

$$\sigma_v = \frac{\sigma_d}{\Delta t} = \frac{\lambda}{4\pi \Delta t} \sqrt{\frac{1-\gamma^2}{\gamma^2 L}} \quad (4)$$

For a representative multilooking a factor of $L=10$ was applied. The resulting uncertainties are:

$$\begin{aligned} \sigma_{\phi} &\approx 0.056 \text{ rad}, \quad \sigma_d \approx 2.5 \times 10^{-4} \text{ m} = 0.25 \text{ mm}, \\ \sigma_d &\approx 1.0 \times 10^{-5} \text{ m d}^{-1} = 0.01 \text{ mm d}^{-1} \approx 3 - 4 \text{ mm a}^{-1}. \end{aligned}$$

This value ($\sigma_d = 0.01$ mm d⁻¹) was thus adopted as the 1σ uncertainty of DInSAR velocity measurements.

3.5 Calculation of the ice volume and uncertainty

To quantify the total ice volume stored in glacial and periglacial cryospheric landforms, we applied cryospheric landforms-specific physics-based models according to their mechanics, morphology, and available input data (Table 5). The method applied and associated uncertainties are detailed below.

3.5.1 Debris-free glaciers

Ice thickness for debris-free glaciers was obtained from the global consensus ice-thickness dataset of Farinotti et al. (2019), which integrates multiple physics-based inversion models (Farinotti et al., 2017). Glacier volumes were computed by spatially

integrating the consensus ice-thickness rasters over the RGI v6.0 glacier polygons within the Copiapó watershed. A relative uncertainty of $\pm 26\%$ was applied following the uncertainty assessment reported in Farinotti et al. (2017, 2019).

3.5.2 Debris-covered glaciers

Mean ice thickness for debris-covered glaciers was derived by sampling the Farinotti et al. (2019) consensus ice-thickness field in the clean-ice frontal zones immediately adjacent to each debris-covered area. These representative values were then assigned to the corresponding debris-covered glacier polygons, ensuring consistency with the local glacier geometry and thickness gradients. Ice volumes were calculated by multiplying the assigned mean thickness by the mapped debris-covered glacier area. Due to the inherently greater spatial variability and additional uncertainties associated with debris cover (e.g., variations in debris thickness, thermal properties, and remote-sensing-derived thickness estimates), published studies of debris-covered glacier thickness consistently report substantially larger differences between modeled and *in situ* measurements compared to debris-free glaciers (e.g. $\sim 40\%$ uncertainty for certain debris-covered glaciers; see Pieczonka et al., 2018). Hence, a conservative uncertainty of $\pm 35\%$ was adopted for debris-covered glacier volume estimates to account for these factors.

3.5.3 Rock glaciers

Rock-glacier thickness was estimated using a viscoplastic creep inversion following Cicoira et al. (2021⁹), where basal shear stress is constrained by observed surface velocity. For each rock glacier, mean slope-parallel surface velocities ($\bar{v}$) and their standard deviation (σ_v) were derived from Sentinel-1 DInSAR measurements (Sect. 3.4.1) and expressed in m yr^{-1} . The basal shear stress τ was inverted using the 1-D viscoplastic formulation with Glen-type exponent $n=3$ and a temperature-dependent rate factor (T).

To explicitly represent uncertainty in the thermal state of ice-rich permafrost in the Central Andes, we treated ice temperature as a normal distribution:

$$T \sim \text{N}(-3^\circ\text{C}, 3^\circ\text{C}) \quad (5)$$

This choice is consistent with reported near-isothermal to moderately cold permafrost conditions in the Central Andes, including rock-glacier settings, and with the strong temperature sensitivity of creep close to 0°C described in constitutive studies (e.g. Koenig et al., 2025, Monnier & Kinnard, 2013).

Using τ , mean thickness was computed as:

$$H = \frac{\tau}{\rho \cdot g \cdot \sin(\alpha)} \quad (6)$$

where α is the mean surface slope, ρ is ice density, and g is gravitational acceleration. Total landform volume was computed as:

$$V = A \cdot H \quad (7)$$

where A is the mapped planimetric area.

Ice volume was obtained by applying an ice-content fraction C_I to the landform volume:

$$V_{ice} = V \times C_I \quad (8)$$

For rock glaciers, C_I was represented as a bounded range consistent with reported values for these type of ice-rich permafrost landforms (Cicoira et al., 2020; Cicoira et al., 2021):

$$C_I = 0.5 \pm 0.10 \quad (9)$$

Ice mass (in Gt) was then computed as:

$$M_{Gt} = \frac{V_{ice} \rho}{10^{12}} \quad (10)$$

with $\rho = 900 \pm 20 \text{ kg m}^{-3}$. Uncertainties in H and M_{Gt} were propagated from σ_v , σ_T , slope uncertainty, mapping uncertainty in area, ice-content uncertainty, and density uncertainty using quadratic propagation of relative errors (Sect. 3.5.5).

3.5.4 Protalus lobes and gelifluction slopes

Protalus lobes and gelifluction taluses were modelled using the same viscoplastic inversion framework originally developed for rock glaciers (Cicoira et al., 2020; Cicoira et al., 2021), exploiting low but measurable DInSAR-derived surface deformation to constrain basal shear stress and thickness. For each mapped polygon, mean slope-parallel surface velocity and its associated uncertainty ($\underline{v}$, σ_v) were used to invert basal shear stress and compute thickness H through Eq. (6). Thermal uncertainty was represented using the same Central Andes temperature prior described in Eq. (5).

However, in contrast to rock glaciers, protalus lobes and gelifluction taluses are characterized by lower ice contents, higher debris fractions, and a dominance of seasonal or shallow permafrost deformation mechanisms. To account for these rheological differences, we modified the Glen-type flow exponent n used in the inversion. While rock glaciers were modelled assuming $n = 3$, consistent with creep-dominated ice–debris mixtures (Cicoira et al., 2020; Cicoira et al., 2021), protalus lobes and gelifluction taluses were modelled using a reduced exponent ($n = 2$), reflecting more linear, frictionally controlled deformation regimes typical of ice-poor frozen debris slopes. This adjustment reduces the sensitivity of inferred thickness to surface velocity and avoids unrealistically thick solutions for low-velocity landforms.

Because ice contents in these landforms are generally lower and more heterogeneous than in rock glaciers, ice-content fraction was prescribed following Eq. (11) as a bounded prior range rather than a fixed value. This range represents plausible ground-ice fractions for ice-rich permafrost slopes in the Central Andes and is supported by field-calibrated ERT observations showing strong spatial variability in ground-ice content within Andean permafrost landforms (Hilbich et al., 2022), as well as early empirical constraints for frozen debris slopes in arid mountain environments (Schrott, 1996). The ice-content prior was therefore implemented to represent structural uncertainty rather than a deterministic parameter.

$$C_I = 0.37 \pm 0.12 \quad (11)$$

Following thickness estimation, landform volume was computed using Eq. (7), ice volume using Eq. (8), and ice mass using Eq. (10).

3.5.5 Uncertainty propagation framework for ice-rich permafrost cryospheric landforms

For rock glaciers, gelifluction taluses, and protalus lobes, uncertainties in ice thickness and ice mass were propagated deterministically using quadratic propagation of relative errors for multiplicative quantities. First, the uncertainty in mapped area was estimated following Paul et al. (2015), assuming boundary errors occur within ± 0.5 pixels and using:

$$u_{A_i} = \frac{1}{2} P_i \times \lambda \times \sigma \quad (12)$$

where P_i is the polygon perimeter, λ is the spatial resolution of the imagery used for delineation, and $\sigma \approx 0.6872$ is a correction factor corresponding to the standard deviation of a Gaussian distribution with most errors confined within ± 0.5 pixels.

Ice volume was computed using the Eq. 7, and ice mass as $M_{Gt} \propto A \times H \times C_I \times \rho$. Therefore, the relative uncertainty in ice mass was calculated as:

$$\left(\frac{\sigma_M}{M}\right)^2 = \left(\frac{\sigma_A}{A}\right)^2 + \left(\frac{\sigma_H}{H}\right)^2 + \left(\frac{\sigma_{C_I}}{C_I}\right)^2 + \left(\frac{\sigma_\rho}{\rho}\right)^2 \quad (13)$$

and the absolute uncertainty as $\sigma_M = M \times \left(\frac{\sigma_M}{M}\right)$. **WatershedBasin** totals were obtained by summing masses across polygons, while total uncertainty was aggregated assuming independence using:

$$\sigma_{tot} = \sqrt{\sum_i^n \sigma_{M_i}^2} \quad (14)$$

501 This framework ensures consistent uncertainty propagation across all ice-rich permafrost cryospheric landforms while
502 preserving the physically based dependence of thickness on surface velocity and temperature.
503

504 **Table 5. Key references used as input for the calculation of ice volumes of cryospheric landforms.**

505

506

Cryospheric landforms	Method	Ice content	Key references
Debris-free glaciers	Physics-based inversion (Farinotti et al., 2019)	100%	Farinotti et al. (2017, 2019)
Debris-covered glaciers	Thickness assignment from adjacent clean-ice zones (Farinotti 2019)	45–99%	Farinotti et al. (2017, 2019)
Rock glaciers	Viscoplastic model (Cicoira et al., 2020Cicoira et al., 2021) + DInSAR	40–60%	Cicoira et al. (20219)
Gelifluction taluses and Protalus lobes	Adapted viscoplastic model + DInSAR	25–49%	Hilbich et al. (2022), Schrott, (1996)

507

508

509

510

511

512

513

514

515

516

517

518

519

520

521 **3.6**

522 **landforms and snow index and data integration**

Cryospheric

523 To develop a classification system that categorizes the capacity to store ice as a strategic water reserve in desert areas of sub-
524 watersheds, we followed a workflow that consists of four phases (Fig. 7). We developed the classification system using
525 cryospheric landforms (expressed in gigatons (Gt), equivalent to 1,000 million metric tons of ice) and snow (quantified in km²
526 of snow-cover extent), integrating both into the methodological framework. The strategic Normalized Cryospheric Index (NCI)
527 captures the relative significance of each sub-watershed, considering both its ability to store water as ice (cryospheric
528 landforms) and its potential to regenerate water resources through snowfall. The NCI needs to be adapted to each specific case.
529 For the Copiapó watershed, the active cryospheric landforms present until 2024 have been used. Those cryospheric landforms
530 include debris-free glaciers, debris-covered glaciers, rock glaciers, gelifluction talus and protalus lobes. Depending on the
531 types of cryospheric landforms available in a specific region, the number of cryospheric landforms included may vary.

532 The methodological framework for computing the NCI index integrates Monte Carlo simulation with a robustness assessment
 533 of a composite indicator that incorporates ice volume (V) and snow permanence (S), adhering to the uncertainty propagation
 534 principles in simulation-based methodologies outlined by Rubinstein and Kroese (2016). First, each watershed has a mean
 535 value (e.g., $\bar{V}_i$ y $\bar{S}_i$) and an estimate of its dispersion (ΔV_i and ΔS_i), usually understood as standard deviation. On the basis
 536 that both V and S follow normal distributions $N(\bar{V}_i, \sigma_{2v})$ y $N(\bar{S}_i, \sigma_{2s})$, N replicates of each variable are generated
 537 for each watershed, thus obtaining “clouds of values” representing the uncertainty of each datum. Subsequently, a
 538 normalization is performed (using min-max) in order to avoid that one variable dominates the other only by differences in
 539 magnitude.

540 Once the simulated values are generated, a weighted index is defined whose objective is to combine both variables (Eq. 15):

$$541 \quad 542 \quad NCI_{i,k}(\omega) = \omega \cdot V_{i,k} + (1 - \omega) \cdot S_{i,k} \quad (15)$$

543
 544 Where $\omega \in [0,1]$

545
 546 Where ω is a parameter that determines the relative relevance of ice volume versus snow permanence. For each sample k
 547 (among the N simulations), the values of that index in all the watersheds are obtained and ordered from highest to lowest,
 548 giving rise to a specific ranking of each run. In this way, each watershed “i” gets a “position” (1 for the watershed with the
 549 highest index, 2 for the next one and so on).

550 The optimization function (Eq. 16) relies on the stability of the ranking under uncertainty, in accordance with the sensitivity
 551 analysis framework outlined by Saltelli et al. (2008). Specifically, for each watershed, the standard deviation of its position
 552 over the N runs is measured, and this measure of dispersion is summed (or averaged) for all watersheds. A smaller value of
 553 this sum indicates greater stability or robustness in the ranking. Formally, if σ_i denotes the standard deviation of the position
 554 of watershed under a weight ω , the objective function results:


$$555 \quad 556 \quad f(\omega) = \sum_{i=1}^n [\sigma_i \cdot (\omega)] \quad (16)$$

557
 558 Where the idea is to minimize the value of this function. To determine the “optimal” ω , a sweep over [0.1] is performed. The
 559 optimal result corresponds to the ω value that maximizes the stability of the watershed ranking system, ensuring minimal
 560 sensitivity to errors or noise in the V and S datasets.

561 Finally, plots are generated to analyze the resulting average ranking and the probability that certain watersheds occupy the first
 562 place in various simulations. In this way, a picture is obtained of how uncertainty in ice volume and snow permanence affects
 563 the prioritization of watersheds.

564 Although the NCI has a simple formulation, the variables it combines are affected by substantial uncertainty. Ice volumes
 565 integrate multiple sources of error related to thickness modelling, ice content, density, and mapped area, while snow is

566 represented by long-term snow permanence derived from MODIS, which captures both the spatial extent and temporal
 567 recurrence of snow. Monte Carlo simulation is therefore applied not to increase index complexity, but to propagate uncertainty
 568 in ice volume and snow permanence into watershed ranking and to identify an objective weighting factor (ω) that maximizes
 569 ranking robustness. This stochastic framework avoids subjective parameterization and ensures a stable, transparent, and
 570 reproducible classification of cryospheric sub-watersheds.



571
 572 **Figure 7. Flow chart of the methodology used to calculate the NCI.**

573 **4 Results**

574 **4.1 Snow permanence results**

575 By integrating all MODIS fractional snow cover images and computing their long-term mean over the period 2000–2022, we
576 derived a snow permanence metric, defined as the probability of snow presence at each pixel. The resulting raster represents
577 the spatial distribution of snow permanence across the Copiapó watershed (Fig. 8). At the regional scale, snow permanence
578 shows limited spatial variability at high elevations, remaining consistently high above approximately 5,500 m a.s.l. These
579 regions are located close to the Argentinean border. In the central area between Caserones Hill and Cadillal Hill, no cryospheric
580 landforms were identified, which coincides with the lowest snow permanence values in the study area, with snow present only
581 around 30% of the time. In contrast, the highest snow permanence values are observed in the southern sector of the study area.
582 In the central area of the Copiapó watershed the snow permanence is concentrated around the Nevado Jotabeche Hill area. The
583 northern part of the watershed does not show concentrated major snowfall anomalies. Snow permanence has persisted
584 approximately 30–50% of the time exclusively at high-elevation zones (>4,500 meters above sea level) near the Argentinian
585 border, contrasting sharply with lower altitudes starting at 2,000 m a.s.l.
586 The calculated probability of the occurrence for the investigated watersheds shows that the snow permanence watershed varies,
587 with higher snow permanence for the southern watersheds compared to the central and northern watersheds. The watershed
588 with the highest resource index is the Montosa ~~R~~river watershed (03413) with ~~0.18%~~0.18 probability of being covered in snow.
589 The Vizcachas de Pulido ~~R~~river (03410), which is located at the lowest altitude, has the lowest probability of ~~0.07%~~0.07 to
590 have snow permanence. These per-pixel snow permanence values, when multiplied by the pixel area (0.25 km²) and integrated
591 across each sub-watershed, yield the mean snow-covered area in km² reported in Table 6 and used as the snow variable S in
592 the NCI calculation.





Figure 8. a) Distribution map of the snow in the Copiapó watershed. b) Jotabeche volcano snow permanence c) The Potro hill snow permanence d) The Tronquitos hill snow permanence. Basemap: Esri, Maxar, Earthstar Geographics, and the GIS User Community.

4.2 Analysis of runoff

Before incorporating snow permanence as a core variable of the NCI, we tested empirically whether MODIS-derived snow cover translates into a hydrological signal at the watershed scale. To do this, we contrasted streamflow behaviour during representative wet and dry years and quantified the statistical relationship between September snow extent and Oct–Mar mean

discharge at the Copiapó River at Pastillo gauging station. Daily mean streamflow records at the Copiapó River at Pastillo station (Fig. 9) reveal distinct seasonal patterns between wet and dry years. During the wet year (Oct 2017 - Mar 2018), streamflow exhibits a sustained increase from October to December, peaking at 5.7 m³/s in December, followed by a gradual decline through January and a secondary pulse in February (5.3 m³/s). In contrast, the dry year (Oct 2013 - Mar 2014) shows substantially lower spring flows, with mean streamflow averaging only 1.0 m³/s during October-November, and the most significant increases concentrated in December (1.3 m³/s) and February (2.2 m³/s). Overall, the wet year delivered 3.4 times more water than the dry year, with the greatest relative differences occurring in November (+471%) and December (+343%).

608



609

610 **Figure 9. Daily (a) and monthly mean (b) streamflow at Copiapó River at Pastillo for wet (2017-2018, blue) and dry (2013-2014, red)**
611 **years. Data Source: Dirección General de Aguas, Chile.**

612 These contrasting patterns between individual years reflect broader interannual variability in snowpack accumulation, ice
613 bodies and melt dynamics. To quantify this relationship at the watershed scale, we incorporated a new analysis linking MODIS-
614 derived snow permanence in September, at the onset of the melt season, with mean river discharge observed between October
615 and March (Fig. 10). This relationship shows a robust correlation ($R^2 = 0.58$; $r = 0.76$; $N = 22$), confirming that seasonal snow
616 permanence acts as a dominant short-term cryospheric water resource in the Copiapó watershed. However, the unexplained





Figure 10. September snow-covered area vs. October-March mean discharge at Copiapó River at Pastillo (2000-2021, N = 22). Linear regression (red line) with 95% confidence interval (shaded). $R^2 = 0.58$; $r = 0.76$; $p < 0.001$. Sources: MODIS MOD10A1 and DGA, Chile.

4.3 Field survey observations of cryospheric landforms.

The fieldwork has yielded observations of cryospheric landforms in the Del Potro and Montosa River sub-watersheds (Fig. 11 and 12). The Del Potro Glacier (Fig. 12d) and the surrounding periglacial environment of the Del Potro river sub-watershed (03412; Fig. 11) were clearly identified during field surveys. In the Montosa River sub-watershed, the Maranceles Glacier (Fig. 12c) was also observed. Additional field observations confirmed the presence of well-developed rock glaciers, gelifluction taluses, and protalus lobes in the Del Potro river watershedbasin (03412) (Fig. 11c-g).

Glacial landforms are preferentially distributed on south-facing slopes, as indicated by their aspect, which are generally associated with reduced incoming solar radiation. In contrast, periglacial cryospheric landforms such as protalus lobes and gelifluction taluses are more frequently found on exposed slopes with steeper gradients and variable aspects, reflecting

637 differences in topographic setting rather than direct measurements of snow accumulation or thermal conditions. Debris-free
638 and debris-covered glaciers are primarily located in topographic depressions and high-elevation headwaters. In contrast,
639 protalus lobes and gelifluction taluses are predominantly distributed on steep slopes and debris-covered sectors. Overall,
640 periglacial cryospheric landforms display a wider spatial distribution across the watershed than glaciers.



641
642 **Figure 11.** Field observations of cryospheric landforms as the ground truthing. a) Map of cryospheric landforms distribution, b) 3D
643 view of the watershed, c) Rock glacier of the Del Potro **R**iver (03412); d) Gelifluction taluses of the Del Potro **R**iver (03412); e)
644 Rock glacier of the Del Potro **R**iver (03412); f) Rock glacier of the Del Potro **R**iver (03412); g) Panorama of different periglacial
645 cryospheric landforms of the Del Potro **R**iver (03412). Basemap: Esri, Maxar, Earthstar Geographics, and the GIS User
646 Community.




Figure 12. Field observations of cryospheric landforms as the ground truthing. a) Map of cryospheric landforms distribution, b) 3D view of the watershed, c) Del Potro debris-free glacier of the Del Potro River (03412), d) Maranceles glacier of the Montosa River (03413), e) Gelifluction taluses of the periglacial environment of the Del Potro River (03412). Basemap: Esri, Maxar, Earthstar Geographics, and the GIS User Community.

652 4.4 Geophysical results

653 4.4.1 Gelifluction Taluses


654 Electrical resistivity tomography (ERT) results (Fig. 13) provides the basis for interpreting the internal structure of the
655 gelifluction taluses. The lobate surface morphology characteristic of this cryospheric landform (Fig. 5 a, b), together with the
656 location and orientation of the tomographic profile (Figure 5 c, d), is clearly reflected in the geophysical response. The ERT
657 profile extends over 295 m and crosses a sector of the landform displaying a well-developed pattern of freeze–thaw-related
658 lobulations, starting at 4,430 m a.s.l. in the upstream sector of the Montosa sub-watershed. This geomorphological pattern
659 coincides with a high-resistivity zone (HRZ), with resistivity values ranging from approximately 1,000 to 8,000 Ωm .
660 According to comprehensive reviews and field studies on mountain permafrost, such resistivity values are typical of ice-rich
661 frozen ground, where pore water is largely transformed into electrically resistive ice (~~Kneisel and Hauck~~ and Kneisel, 2008).
662 Electrical resistivity is known to increase by several orders of magnitude upon freezing, particularly in coarse-grained
663 periglacial materials, making ERT a robust tool for detecting ice-bearing permafrost bodies.

664 The HRZ delineates an ice-rich permafrost body with an estimated two-dimensional horizontal extension of approximately 20
665 m² within the tomographic section. Comparable resistivity ranges and internal structures have been documented in ice-rich
666 permafrost landforms investigated with ERT, including rock glaciers, talus slopes and other periglacial slope systems (~~Hauck~~
667 ~~and Kneisel~~ Kneisel et al. and Hauck, 2008; Hilbich et al., 2009). The association between lobate slope morphology and
668 underlying frozen ground with significant ice content is consistent with the established understanding of gelifluction as a
669 process operating on seasonally thawed active layers overlying permafrost, where ice lenses and frozen substrates exert a first-
670 order control on deformation patterns (Matsuoka, 2001).

671 Downstream of 4,430 m a.s.l., the HRZ terminates abruptly at the frontal sector of the last lobulations. From this point, a low-
672 resistivity zone (LRZ) dominates, with resistivity values between 50 and 200 Ωm . This sharp contrast is interpreted as the
673 transition from frozen ground to unfrozen, water-saturated sediments. Similar resistivity ranges have been reported for
674 unfrozen debris, taliks and shallow groundwater zones underlying permafrost terrain, where electrolytic conduction through
675 liquid water strongly reduces bulk resistivity (~~Kneisel and Hauck~~ and Kneisel, 2008).

676 The spatial juxtaposition of the HRZ and LRZ suggests a direct subsurface connection between the ice-rich permafrost body
677 and an underlying groundwater system. Studies combining ERT and seismic methods demonstrate that such resistivity
678 contrasts commonly mark permafrost boundaries and zones of hydrological interaction, where water infiltration and ice loss
679 can be detected geophysically (Hauck et al., 2008; Mewes et al., 2017). In particular, combined geophysical monitoring has
680 shown that increases in liquid water content beneath or adjacent to frozen ground produce a pronounced decrease in resistivity,
681 even when ice loss is spatially limited. Based on the vertical extent of the HRZ, the ERT data allow for the estimation of an
682 average ice-rich core thickness of approximately 15 m (Fig. 13). This thickness falls within the range reported for ice-bearing

683 periglacial slope landforms investigated using geophysical methods, where frozen layers commonly extend several tens of
684 meters below the surface depending on material properties and thermal conditions (Kneisel [et al.](#) and [Hauck](#), 2008; Hauck and
685 [Kneisel et al.](#), 2008).



686
687 **Figure 13. Results of the electrical resistivity tomography (ERT) over the gelifluction taluses cryospheric landforms in the Montosa**
688 **Sub-watershed. a) Electrical resistivity tomography, b) Subsurface interpretation of resistivity test. Basemap: Esri, Maxar,**
689 **Earthstar Geographics, and the GIS User Community.**

690

691 4.4.2 Rock Glacier

692 A second ERT survey was conducted across a representative active rock glacier in the Montosa sub-watershed (Fig. 14). The
693 tomographic sections reveal a complex internal structure characterized by laterally continuous high-resistivity zones (HRZ),
694 which are interpreted as ice-rich permafrost bodies embedded within a coarse debris matrix, a configuration commonly
695 reported for active rock glaciers worldwide (Kneisel et al., 2008).

696 Resistivity values exceeding approximately 20–50 k Ω ·m dominate the central and frontal sectors of the landform, indicating
697 the presence of massive or highly ice-saturated material. Comparable resistivity ranges have been documented in ice-rich rock
698 glaciers in the Andes and other high-mountain regions, where electrical resistivity contrasts are primarily controlled by ground-
699 ice content rather than lithological variability (Kneisel et al., 2008; Pavoni et al., 2021). These high-resistivity anomalies are
700 spatially coherent with the surface morphology of the rock glacier, including its lobate structure and well-defined frontal slope,
701 supporting a close correspondence between geomorphological expression and subsurface ice distribution.

702 Beneath the HRZ, lower-resistivity zones (LRZ; <1–5 k Ω ·m) are observed, which are interpreted as unfrozen or partially
703 saturated debris layers and/or underlying bedrock. The relatively sharp transition between high- and low-resistivity domains
704 suggests a distinct boundary between ice-rich permafrost and unfrozen substrate, a feature widely reported in ERT studies of
705 active rock glaciers (Kneisel ~~et al.~~ [Hauck](#), 2008).

706 The ERT-derived geometry indicates an average ice-rich layer thickness on the order of 20–30 m, with local maxima exceeding
707 35 m in the central part of the rock glacier body. These thicknesses fall within the range reported for active rock glaciers in the
708 dry Andes based on geophysical surveys and regional assessments (Brenning, 2005; Azócar and Brenning, 2010; Halla et al.,
709 2021). The observed thickness values provide an independent observational constraint for the thickness estimates derived from
710 the viscoplastic creep inversion ([Cicoira et al., 2021](#)) based on DInSAR surface velocities (~~Cicoira et al., 2020~~[Cicoira et al.,](#)
711 [2021](#)).

712 Importantly, the spatial extent of the HRZ aligns closely with the mapped rock glacier polygon, reinforcing the validity of the
713 inventory-based delineation and confirming that the geophysically detected ice body corresponds to the geomorphological unit
714 considered in the volume calculations. The ERT results therefore serve as ground-based validation for both the presence of
715 ice-rich permafrost and the order of magnitude of the modeled thicknesses applied to rock glaciers within the Copiapó
716 watershed, consistent with previous multi-method approaches combining geomorphology, geophysics, and kinematic
717 observations.




Figure 14. Results of the electrical resistivity tomography (ERT) over rock glacier cryospheric landforms in the Montosa Sub-watershed. Basemap: Esri, Maxar, Earthstar Geographics, and the GIS User Community. a) Longitudinal tomography, b) Cross-sectional tomography.

4.5 Thickness estimations on periglacial ~~environment~~landforms

Surface displacement maps (Fig. 15) of a representative rock glacier in the Copiapó watershed show a coherent downslope motion that is clearly confined to the rock glacier body, while surrounding terrain remains stable, supporting the reliability of both the cryosphere inventory and the DInSAR processing. These surface velocities provide a direct kinematic constraint for the viscoplastic thickness modelling of ice-rich permafrost landforms used in this study.

The modeled ice-rich layer thicknesses for the main periglacial cryospheric landforms (rock glaciers, protalus lobes, and gelifluction slopes) and their relationship with DInSAR-derived surface velocities (Fig. 16) show that rock glaciers exhibit the largest modeled thicknesses, with median values on the order of ~25–30 m and a wide interquartile range, reflecting both their higher surface velocities and the greater mechanical efficiency of creep in these landforms. In contrast, protalus lobes show intermediate thicknesses, typically below ~15 m, while gelifluction slopes are characterized by consistently thinner ice-rich layers, generally <10 m. The scatter plot highlights a positive relationship between mean surface velocity and modeled thickness across all periglacial cryospheric landforms, consistent with the viscoplastic framework in which higher deformation rates require larger ice thicknesses to sustain basal shear stress under comparable slope conditions.




Figure 15. Example of DInSAR-derived surface displacement of a rock glacier in the Copiapó Watershed. Basemap: Esri, Maxar, Earthstar Geographics, and the GIS User Community.

Rock glaciers cluster at higher velocities and thicknesses, whereas protalus lobes and gelifluction slopes occupy lower-velocity regimes, with substantial overlap reflecting variability in slope, thermal state, and internal ice content. Despite this overlap, the three cryospheric landforms groups remain clearly distinguishable in thickness–velocity space, supporting the application of a unified viscoplastic model while preserving cryospheric landforms-specific structural differences. These results confirm that surface kinematics derived from DInSAR provide a physically meaningful constraint on subsurface ice thickness and justify their use for estimating ice volumes in ice-rich permafrost landforms across the Copiapó watershed. [These thickness estimates are further contextualized by multi-decadal kinematic records from the Dry Andes showing that rock glaciers exhibit marked interannual velocity variability \(Vivero et al., 2021\), and by debris-covered glacier studies demonstrating that supraglacial debris exerts a strong control on the hydrological behavior of ice bodies in high-elevation Andean catchments](#)

747 (Burger et al., 2019). The long-term hydrological relevance of these landforms is reinforced by projections indicating reduced
 748 glacier meltwater contributions under future megadrought scenarios (Ayala, 2025), which elevate the importance of accurately
 749 quantifying periglacial ice reserves such as those estimated here.

750

751

752 **Figure 16. Modeled thickness–velocity relationship for periglacial cryospheric landforms of Copiapó Watershed.**

753 **4.6 Ice volumes, snow area and NCI for sub-watershed**

754 The ice volumes, snow area and the NCI for each of the sub-watersheds that constitute the cryospheric landforms of the
 755 Copiapó River watershed have been calculated (Table 6). The watershed with the largest ice masses is the Montosa River
 756 watershed (03413), with 0.922 Gt, followed by the Del Potro River watershed (03412) with 0.913 Gt and thirdly the
 757 southernmost watershed, the Manflas River watershed (03420) with 0.371 Gt of ice. In terms of volume, the largest ice
 758 volumes in the entire watershed are found in debris-free glaciers, gelifluction taluses and rock glaciers (Fig. 17).

759 Following the mapping of snow, we observe that the surface area distribution of snow cover is uneven, with a higher density
 760 in the southern part and the northern end of the watershed. In the south, the watersheds with the largest snow permanence are
 761 the Manflas River (03420), with 107.06 km² of mean snow cover (2000–2024), the Montosa River (03413) with 75.06 km²
 762 and the Del Potro River (03412) with 53 km². In the north, the most important rivers are the Estero Come Caballos (03402)
 763 with 122.24 km², Figueroa River (03401) with 106.72 km², Quebrada Monardes (03400) with 89.63 km² and the Cachitos
 764 River (03403) with 88.34 km² (Table 6).




Figure 17. Total ice mass of the Copiapó river Watershed.




Figure 18. Total ice volumes of the Copiapó river watershed expressed both as a) total ice mass with associated uncertainty and b) as stacked contributions by cryospheric landforms.

The distribution of ice mass across cryospheric sub-watershedsub-basins of the Copiapó watershed (Fig. 18), reveal a highly heterogeneous spatial distribution of cryospheric landforms, with a small number of sub-watershedsub-basins concentrating the majority of the total ice mass.

The largest ice masses are found in the Montosa (COD_SSUBC 3413) and Del Potro (3412) sub-watershedsub-basins, each storing approximately ~0.9 Gt of ice. These watershedbasins clearly dominate the cryospheric landforms of the watershed and are characterized by the combined presence of debris-free glaciers, rock glaciers, and extensive ice-rich permafrost

landforms. In contrast, several sub-watersheds exhibit ice masses below 0.1 Gt, reflecting either limited cryospheric landforms extent or the dominance of thin periglacial landforms.

The stacked representation highlights that debris-free glaciers constitute the principal contribution to ice mass in the largest ice storage, whereas ice-rich permafrost cryospheric landforms, particularly rock glaciers and gelifluction taluses, represent a substantial fraction of the total ice mass in intermediate and smaller sub-watersheds. Proglacial lobes contribute minor but non-negligible volumes in selected sub-watersheds, emphasizing their cumulative importance at the watershed scale.

Uncertainty ranges are largest in sub-watersheds dominated by periglacial cryospheric landforms, reflecting the combined effects of variability in surface velocity, thermal state, ice content, and mapped area. Nevertheless, the relative ranking of sub-watersheds by total ice mass remains robust, with Montosa and Del Potro consistently emerging as the primary cryospheric landforms within the Copiapó watershed.

Overall, these results demonstrate that cryospheric ice mass is strongly concentrated in a limited number of sub-watersheds, while many others contain comparatively small but non-negligible ice volumes distributed among different cryospheric landforms. However, ice mass alone does not fully capture the hydrological relevance of a watershed, as it does not account for the capacity to regenerate water resources through snowfall. This motivates the integration of cryospheric landforms (ice mass) and snow (Permanence) into a unified metric, introduced in the following section as the Normalized Cryospheric Index (NCI).

The Normalized Cryospheric Index (NCI) behavior in relation to the relative weighting of ice volume (V) and snow occurrence (S), along with the stability of sub-watershed rankings under uncertainty (Fig. 19) shows how the choice of weight (w) influences the mean value of the NCI and, ultimately, the final ranking of the sub-watersheds.


The evolution of the mean NCI value of each sub-watershed as w varies from zero to one (Fig. 19a). Each line represents a different sub-watershed, so that the slopes and crosses between lines allow us to appreciate which sub-watershed are favored by a greater relevance of ice volume V (when w tends to 1) or snow cover S (when w tends to 0). This preliminary behavior shows, on the one hand, the sensitivity of each subwatershed to weighting and, on the other hand, the possible dominance of one variable over the other in certain w zones.

The position in the ranking that each subwatershed occupies for each w value (Fig. 19b) shows that the horizontal lines extend from the first position (1) to the last. It highlights consistently privileged-ranked subwatersheds regardless of w, in contrast to unstably ranked subwatersheds. This early insight into the stability of the ranking compellingly underscores the urgency of undertaking a rigorous robustness analysis.

The ranking variability function, defined as the sum (or average) of the standard deviations of the position of each sub-watershed in the different simulations (Fig. 19c) quantifies the overall stability of the ranking for each w. A marked minimum is seen around w=0.85, indicating that, at that point, the ranking of watersheds is less sensitive to the uncertainty of the data

Con formato: Fuente: Sin Cursiva

807 of V and S. In other words, assigning that weight in the combination of variables leads to a more consistent ranking in terms
808 of robustness.
809 The mean NCI (with its error bars) for each watershed when $w=0.85$, the optimal value detected with the optimization function
810 (Fig. 19d), shows that the vertical axis reflects the average magnitude of the index, while the error bars indicate the standard
811 deviation derived from the Monte Carlo simulations. Thus, the visualization offers a clear identification of the watersheds with
812 the highest NCI values, while simultaneously exposing the degree of uncertainty surrounding these estimates. This result
813 confirms the relevance of choosing $w=0.85$ and provides a detailed view of the dispersion of each watershed under such a
814 scenario.
815



816
817 **Figure 19. Analysis of NCI and robustness of the ranking as a function of w weighting (a) Evolution of the mean NCI for each sub-**
818 **watershed as w varies between 0 and 1 (b) Position in the ranking of each sub-watershed as a function of w (c) Ranking variability**
819 **function (sum of the standard deviations of the positions), where the minimum at $w \approx 0.85$ indicates the maximum stability of the**
820 **ranking (d) Mean NCI $\pm$ standard deviation for $w=0.85$, evidencing the magnitude and dispersion of the index in each sub-watershed**
821 **under the most robust condition.**

822

823 The calculated NCI, demonstrated that the sub-watershed with the highest NCI was the by the Montosa Rriver watershed
824 (03413) (NCI = 0.93), followed by the Del Potro Rriver watershed (03412) (NCI = 0.89), Manflas Rriver watershed (03420)
825 (NCI = 0.47), Estero Come Caballos watershed (03402) (NCI = 0.34) and Cachitos Rriver watershed (03403) (NCI = 0.31).
826 These are the sub-watershed with the largest cryospheric landforms. In the northern region, the Estero Come Caballos sub-
827 watershed, it was highlighted by the index primarily attributed to its persistent snow cover.

828

829 **Table 6 Summary of results by sub-watershed of cryospheric landforms in Gt, snow in km² and NCI indexes in different scenarios.**

830

831	Code of the sub-	Cryospheric	Cryospheric	Snow in sq.	NCI	NCI
832	watershed	landforms	landforms	km		uncertainty
833	3400	0.039	0.006	89.63	0.14	0.01
834	3401	0.062	0.005	106.72	0.18	0.02
835	3402	0.212	0.014	122.24	0.34	0.02
836	3403	0.230	0.015	88.34	0.31	0.02
837	3410	0.007	0.001	43.90	0.04	0.01
838	3411	0.239	0.018	45.95	0.26	0.02
839	3412	0.913	0.124	53.02	0.89	0.11
840	3413	0.922	0.082	75.06	0.93	0.08
841	3420	0.371	0.055	107.06	0.47	0.05
842	3440	0.075	0.009	46.66	0.11	0.01
843	3441	0.041	0.006	46.79	0.08	0.01
844	3443	0.001	0.0005	19.76	0.00	0.00

845

846

847

848 5. Discussion

849 5.1 Cryospheric landforms and snow permanence in the Copiapó watershed

850 From the cryospheric landforms and snow occurrence mapping exercises (Figs. 4 and 8), we observed that the cryospheric
851 landforms of the Copiapó watershed are predominantly located at altitudes above 4,500 m. This compares well with other
852 regions in which snowfall has been identified to be controlled by altitude (Saydi and Ding, 2020). We also observed that
853 cryospheric landforms occur in smaller, isolated regions ($< 0.25 \text{ km}^2$ in width) in the northern part of our research area. This
854 corresponds with our own field observations, where most cryospheric landforms occur in smaller bodies (Fig. 4). This resulted
855 in smaller, separated cryospheric landforms and surrounding the highest point (Figure 4 b and c). They host smaller cryospheric
856 landforms, for example near the Cadillal and the Nevado Jotabeche, compared to the southern part of our study area. Because
857 those northern cryospheric landforms are located in or in proximity of the SAAD (Abraham et al., 2020), snowfall and
858 permanent snow occurrences are limited, leading to isolated cryospheric landforms (Fig. 4). The potential volume of water
859 that is captured in cryospheric landforms in the sub-watersheds of the Copiapó watershed (Fig. 1) is thus strongly influenced
860 by the South American Arid Diagonal (SAAD) that controls the distribution of the different types of cryospheric landforms
861 (García et al., 2017). In our case, the proximity of the Copiapó watershed to the SAAD limits the occurrence of debris-free
862 glaciers in the watershed north of 28.1° S latitude and limits the continuity of the cryosphere that is mainly attributed to
863 cryospheric landforms associated with ice-rich permafrost, such as small rock glaciers, protalus lobes and predominantly
864 gelifluction taluses.

865 In the more central domain in between Cadillal and Caserones no cryospheric landforms occur. This can be explained by the
866 mechanical forcing of the South American low-level jet and the topographic blocking of the Andes (Insel et al., 2009). In the
867 central sector, Mount Pissis exhibits greater elevation on the Argentinian side. Consequently, the prevailing northeasterly
868 winds transporting moisture from the Pacific Ocean are obstructed by the Argentine portion of the mountain. This orographic
869 interception results in localized snowfall on its Argentinian slopes, rather than allowing precipitation to be deflected by the
870 comparatively lower-elevation Chilean topography. The topography north of Cadillal and south of Caserones is high enough
871 to block these winds on the Chilean side, leading to snowfall and eventual cryospheric landforms in the southern regions in
872 our research area. A last negative spatial correlation between the snowfall and the cryospheric landforms is the large snowfall
873 anomaly in the northern area. The permanent snow occurrence is high, while not many cryospheric landforms are produced.
874 The reason for this is the location of this region in the SAAD, resulting in melting or sublimation of the snow that does arrive
875 in this area (Réveillet et al., 2020), meaning that the snow does not end up in cryospheric landforms.

876 From the different cryospheric landforms, we see that gelifluction taluses cover the largest surface area in the Copiapó
877 watershed compared to other cryospheric landforms in our research areas (Garcia et al., 2017). Field observations also confirm
878 that gelifluction taluses extend across significantly broader areas compared to other cryospheric landforms (Fig. 12~~7~~). The
879 limited understanding of gelifluction taluses has led to their exclusion from global cryospheric reserve inventories.
880 Nevertheless, in assessing watershed ice volume, these taluses emerge as the most critical features for ice storage, surpassing

881 other formations in significance (Fig. 17). Furthermore, the ERT-derived thickness measurements across the gelifluction lobe
882 reveal a minimum detectable thickness of 15 meters, which markedly contrasts with the range of 1 to 5 meters reported by
883 Hilbich et al. (2022~~45~~).

884 On a more regional scale, we observe that in the south of the watershed, debris-free glaciers are always in contact with
885 gelifluction taluses. This is confirmed in the field (Fig. 12). In the north of the Copiapó watershed, gelifluction taluses occur
886 more as isolated bodies. The topography of the northern area is controlled by Tertiary volcanoes (Kay et al., 1994). In general,
887 these volcanoes exhibit gentler slopes than those of the Principal Cordillera (Harrington, 1961) and largely shape the
888 topography of our southern research area. Given that gelifluction taluses require steep slopes to form (Fig. 2), they are more
889 likely to develop in the southern sector, where the terrain is dominated by the steep relief of the Principal Cordillera unlike in
890 the north, where this feature is absent. Moreover, in the Central Cordillera glacial cirques are abundantly present (Evans, 2006).
891 Glacial cirques provide sheltered environments that shield snow from extreme weather conditions, facilitating its
892 recrystallization into exposed glaciers and other cryospheric landforms. The distribution of rock glaciers and protalus lobes is
893 likewise influenced by the presence of glacial cirques (Palacios et al., 2020), resulting in their predominant occurrence in the
894 southern portion of our study area. In contrast, no exposed glaciers are found north of Copiapó Hill (Fig. 4). Instead, gelifluction
895 taluses and protalus lobes allow the continuity of the cryospheric fringe of the watershed. The absence of debris-free glaciers
896 in this region is directly related to the South American low-level jet and the topographic blocking of the Andes (Insel et al.,
897 2009).

898 These DGA (2014) volumes were used to provide a magnitude check. At the Del Potro Glacier, the modeled volume is ~24%
899 larger than the GPR-based estimate; at the Tronquitos Glacier, the difference is larger (~70%). This spread is within the
900 expected uncertainty range for individual glacier inversion products in Farinotti et al. (2019), particularly for small glaciers
901 with limited thickness constraints. The DInSAR-derived surface velocities of rock glaciers in the Copiapó watershed are
902 systematically lower than those reported for rock glaciers at comparable elevations further south in the Dry Andes. Vivero et
903 al. (2021) documented mean surface velocities of 0.26 m yr⁻¹ during 1956-1978 and up to 1.25-1.80 m yr⁻¹ during 2014-2019
904 for the Tapado, Las Tolas, and El Cachito rock glaciers at 30°S in the Coquimbo Region. In contrast, the rock glaciers of the
905 Copiapó watershed (26°41'-28°24'S), as detected by Sentinel-1 DInSAR, exhibit velocities on the order of a few millimetres
906 per year, one to two orders of magnitude slower. Two complementary factors explain this systematic difference. First, the
907 Copiapó watershed lies within or immediately adjacent to the South American Arid Diagonal (SAAD), where extremely
908 limited precipitation results in drier, colder permafrost conditions and reduced ground-ice temperatures relative to sites further
909 south. Lower permafrost temperatures translate directly into higher ice viscosity and reduced creep rates under equivalent basal
910 shear stresses, consistent with the strong temperature dependence of the Glen-type flow law applied in this study (Cicoira et
911 al., 2021; Koenig et al., 2025). Second, the cryospheric landform genesis differs fundamentally between the two sites. The
912 Tapado complex studied by Vivero et al. (2021) comprises glacial rock glaciers, landforms that originated from or remain
913 connected to glacier ice and push moraines, and which therefore tend to be ice-richer and more dynamically active. The rock
914 glaciers inventoried in the Copiapó watershed are predominantly cryogenic in origin, having developed through progressive

915 freezing of talus and debris-slope material under permafrost conditions, without a direct glacial ice source. Cryogenic rock
916 glaciers characteristically exhibit lower ice contents, more heterogeneous debris-ice matrices, and slower creep rates than
917 glacigenic forms (Monnier and Kinnard, 2015). The slower kinematics observed in the Copiapó watershed are therefore
918 physically consistent with both the arid thermal regime imposed by proximity to the SAAD and the cryogenic origin of the
919 inventoried rock glaciers, and do not reflect anomalous model behaviour in the viscoplastic thickness inversion.

920 The DGA (2014) GPR-derived ice volume estimates were used as an independent benchmark to evaluate the magnitude of the
921 modeled glacier volumes (Table 7). For Del Potro Glacier, the modeled volume (0.362 km³) exceeds the GPR-based estimate
922 (0.293 km³) by approximately 24%. In the case of Tronquitos Glacier, the discrepancy is larger, with the modeled volume
923 (0.157 km³) being about 70% higher than the GPR-derived value (0.092 km³).

924 Despite these differences, the observed spread is consistent with the uncertainty ranges reported for individual glacier ice-
925 thickness inversion products in Farinotti et al. (2019). This is particularly relevant for small glaciers, where limited field-based
926 thickness constraints and complex geometries tend to amplify relative errors in volume estimates. Therefore, while the
927 discrepancies, especially for Tronquitos, are non-negligible, they remain within the expected variability for regional-scale
928 glacier volume modeling.

929 A relevant consideration regarding the ice volume estimates for debris-free glaciers in the Copiapó watershed concerns the
930 difference in glacier area between the two inventories used. The regional inventory of García et al. (2017) maps a total of 1,907
931 ha of uncovered glaciers within the Copiapó watershed, whereas the Randolph Glacier Inventory (RGI) v6.0, which forms the
932 basis for the Farinotti et al. (2019) thickness product used in this study, encompasses a larger glacier area for the same
933 watershed. Because the Farinotti et al. (2019) volumes are computed over the RGI glacier polygons, which are more extensive
934 than those delineated by García et al. (2017), the total ice volume attributed to debris-free glaciers in this study may
935 overestimate the volume that would be obtained if the García et al. (2017) glacier boundaries were used instead. This
936 discrepancy arises from methodological differences in glacier delineation, inventory date, and mapping criteria between the
937 two datasets, and represents an inherent source of uncertainty in the volume estimates reported here. Future work should aim
938 to reconcile these inventories and assess the sensitivity of the modeled ice volumes to the choice of glacier outlines, particularly
939 for small and fragmented glacier bodies in the northern sector of the watershed.

940 **Table 7. Comparison between modeled glacier ice volumes from Farinotti et al. (2019) and GPR-derived ice volumes**
941 **reported by DGA (2014) for Del Potro and Tronquitos glaciers.**

Glacier	RGI ID	Farinotti model volume (km ³)	DGA 2014 GPR volume (km ³)
Del Potro	1.715.087	0.362	0.293

Tronquitos	1.715.038	0.157	0.092
------------	-----------	-------	-------

The watershed with the best balance between cryospheric landforms and snow permanence is the Montosa ~~R~~river watershed (03413) (Fig. 20), which contains high quantities of both cryospheric landforms and snow permanence. The field observations of geomorphological features that are indicative of active cryospheric landforms, such as the subparallel terraces (Fig. 12e), fit well with the mapped cryospheric landforms from our workflow. The mapped surface area and thickness are supported by the observed surface area and thickness in the field. Lastly, debris-covered glaciers occur predominantly in areas with a relatively continuous cryosphere (García et al., 2017). This is consistent with our findings that debris-covered glaciers are exclusively located in the southern region of the study area, a zone characterized by a more extensive and contiguous cryospheric distribution relative to the fragmented northern sectors. The interannual variability in glacier meltwater contribution in high-elevation Andean catchments is strongly modulated by the insulating effect of supraglacial debris, which attenuates ablation rates and delays the hydrological response of debris-covered glaciers relative to their clean-ice counterparts (Burger et al., 2019). Complementing this, multi-decadal monitoring using aerial and satellite photogrammetry and UAV surveys has demonstrated that rock glaciers in the Dry Andes exhibit significant kinematic variability, underscoring their dynamic role in the cryospheric system of the region (Vivero et al., 2021).

5.2 Classifying and prioritizing sub-watersheds in the Copiapó watershed

Unlike rainfall-dominated catchments, the upper Copiapó watershed is strongly influenced by cryospheric processes, with its headwaters characterized by the widespread presence of glaciers, rock glaciers, and ice-rich permafrost landforms. Cryospheric elements are concentrated in the high-altitude sub-watersheds, where water storage and seasonal runoff generation are controlled primarily by snow accumulation, snow persistence, and long-term ice storage rather than by direct rainfall inputs. The Normalized Cryospheric Index (NCI) identifies three main sub-watersheds with a dominant cryospheric influence on the hydrological functioning of the Copiapó system, based on the combined contribution of ice mass and long-term snow persistence. The spatial distribution of cryospheric landforms shows a 97% overlap with high-probability permafrost zones predicted by the global permafrost model of Gruber (2012), providing independent support for the relevance of ice-rich permafrost as a key component of water storage in the ~~watershed~~basin. Together, these observations justify classifying the upper Copiapó as a cryosphere-influenced watershed, in which cryospheric processes exert a first-order control on water availability. The critical sub-watersheds include: the Montosa River (03413), Del Potro River (03412) and Manflas River (03420) (Table 6). Their spatial distribution corresponds to zones of maximum snow permanence and the most extensive

975 cryospheric continuity within the Copiapó watershed (Figs. 6 and 8). These high indexes thus directly correlate with the
976 cryospheric landforms and snowfall occurrences. For example, enhanced snowfall is concentrated in the Cerro del Potro sub-
977 watershed (03412), where the largest debris-free glacier of the Copiapó watershed is located (Fig. 8c). This area hosts El Potro
978 Glacier, the most extensive binational glacier in northern Chile (García et al., 2017), indicating a strong coupling between
979 snow accumulation, glacier development, and long-term ice storage. As a result, the Cerro del Potro sub-watershed ranks
980 among the highest NCI values, reflecting the combined importance of snow (permanence) and cryospheric landforms (ice
981 mass). This combination has the potential to control the presence of water resources downstream when global temperatures
982 continue to rise (Flores et al., 2018). Notably, recent projections for the Southern Andes indicate that glacier meltwater
983 contributions may decline substantially during future megadrought conditions (Ayala, 2025), reinforcing the urgency of
984 identifying and protecting cryospheric sub-watersheds such as those ranked highest by the NCI. With this study, we detected
985 the sub-watersheds that should be prioritized. In wet years, snow controls the watershed hydrology in the northern part of the
986 Copiapó watershed. This means that in these locations infrastructure needs to be put in place to capture the larger amounts of
987 run-off that will arrive with increasing warmer temperatures, such as the Estero Come Caballos sub-watershed (03402), with
988 the ~~fourth~~^{third} place of the NCI. In times of water scarcity, we identified that the sub-watersheds of the Montosa ~~R~~river
989 (03413), Del Potro ~~R~~river (03412), and Manflas ~~R~~river watershed (03420) will contribute most of the flow to the Copiapó
990 ~~R~~river.

991 Given the observed relationship between snowfall and cryospheric landforms distribution, prioritizing the Copiapó watershed
992 is essential to safeguard its water resources. We propose a systematic framework to evaluate and rank sub-watersheds
993 according to preliminary assessments of cryospheric landforms and resource availability. From this ranking (Table 6), the main
994 target, with the highest NCI is the sub-watershed of Montosa ~~R~~river (03413). The second target is located towards the south,
995 the Del Potro ~~R~~river (03412). The third target is the Manflas ~~R~~river watershed (03420) and finally the last targets are Estero
996 come Caballos (03402) and Cachitos ~~R~~river (03403) watershed (Table 6). These ~~five~~^{four} sub-watersheds constitute the basis
997 for continuous water production in the Copiapó watershed during dry years, because they host ~~853.09~~^{853.0944} % of the cryospheric
998 landforms ice volume of the Copiapó watershed. From these four sub-watersheds, the Montosa ~~R~~river watershed (03413)
999 maintains the best balance between snow and cryospheric landforms (Fig. 20). We have shown that with this hydrological
1000 approach, the NCI allows us to infer that the sub-watershed with the highest hydrological importance is the Montosa ~~R~~river
1001 watershed (03413). This index also allowed us to observe that the watersheds of the northern region are high in snow cover
1002 but are less enriched in cryospheric landforms. During years of substantial snowfall, management efforts should prioritize
1003 northern sub-watersheds, such as the Estero Come Caballos ~~S~~sub-watershed (03402). This suggests that northern watersheds
1004 exhibit a limited ability to develop cryogenic niches, as snow in these areas is primarily lost through melting or sublimation.
1005 A potential explanation for the lack of cryospheric landforms in northern watersheds lies in their geomorphological
1006 characteristics, which may hinder the accumulation and preservation of ice.

Based on the calculation of the NCI, we have been able to identify that the Estero Come Caballos watershed (03402) is the most strategic for snow accumulation, even surpassing the watersheds in the southern part of the Copiapó watershed in terms of snow permanence (Fig. 20).




Figure 20. Summary graph of the normalized indexes for snow, cryospheric landforms and the cryosphere for each sub-watershed of the Copiapó River watershed.

5.3 Impact on watershed management

We have analyzed 12 cryospheric sub-watersheds in the Copiapó watershed and showed that there are some that should be prioritized in watershed management strategy. The methodological framework developed and presented in this study can be extended to assess the comprehensive hydrological potential of arid watersheds in high-elevation Andean regions of Chile and other globally distributed arid zones with a focus on cryospheric contributions. In addition, this novel sub-watershed indexing method is fundamental to identify those watersheds with an important snow, glacial and periglacial component that are suitable for inclusion in the watershed protection frameworks that are implemented in the water management strategies of each watershed in Chile. Given the persistent progression of climate change, it is imperative to integrate water reserves sourced from cryospheric watersheds into comprehensive policy-making frameworks governing hydrological regulation. Implementing and integrating such cryosphere inventories will also allow other regions to preserve their water resources. Currently, the Chilean national cryospheric inventories distinguish between glacier and periglacial landforms. Debris-free and debris-covered glaciers are comprehensively included in the National Glacier Inventory of Chile, whereas within the periglacial environment, only rock glaciers are systematically inventoried. As a result, ice volumes potentially stored in other ice-rich periglacial landforms, such as protalus lobes and gelifluction taluses, are not explicitly accounted for in existing national

inventories, despite being addressed in this study. Increasing the variety of cryospheric landforms would facilitate more accurate estimations of the volumes of ice in each sub-watershed, especially in northern Chile where the gelifluction taluses and protalus lobes dominate the cryospheric landforms.

Identifying the strategic position of these sub-watersheds does not imply guaranteeing constant runoff, but rather highlights those sub-watersheds with a higher potential to sustain streamflow under low-snowfall or dry-year conditions. The runoff analysis shows that, during dry years, discharge is strongly reduced but not suppressed, and remains seasonally coupled to melt processes originating in high-elevation cryospheric environments. Sub-watersheds ranked as strategic by the NCI concentrate larger cryospheric landforms and higher snow permanence, which together contribute to buffering runoff variability and maintaining baseflow during periods of limited snowfall. The extraction of precious metals has consumed large quantities of water that was stored in cryospheric landforms (Brenning, 2008). A lack of comprehensive understanding regarding the spatial distribution and volumetric potential of cryospheric water reserves has contributed to the degradation of cryospheric landforms critical for supplying freshwater to downstream populations residing at lower elevations. These sub-watersheds are predominantly cryospheric rather than pluvial in character, indicating that their hydrological regime is mainly influenced by meltwater processes associated with glaciers, ice-rich periglacial landforms, and seasonal snow permanence, particularly during the melt season. When the potential volume of water in each sub-watershed is understood, policy and infrastructure can be adapted to maintain those cryospheric components. Prioritizing the extraction of water from cryospheric landforms and implementing snow capture techniques is essential to ensure the sustainable development and enhanced productivity of the examined region, as well as the arid northern regions of Chile. These sub-watersheds may therefore play an important role in the prevention of water disasters greater than those already known in Chile. The methodological basis developed in this study can be a key-tool for developing a law to protect Chile's cryosphere and its associated water resources when adopted in future water management strategy plans.

6. Conclusions

6.1 Main findings of the paper

We developed a reproducible workflow that maps glacial and periglacial cryospheric landforms and seasonal snow, estimates their ice volumes with a unified physics-based framework, and integrates both into a Normalized Cryospheric Index (NCI) that ranks sub-watersheds with explicit Monte Carlo uncertainty propagation. Applied to the Copiapó watershed, the workflow classified 12 of 27 sub-watersheds as cryospheric; these contain 97 % of the inventoried landforms and overlap 97 % with high-probability permafrost predicted by Gruber (2012), providing independent support for the classification.

The empirical link between September snow extent and October–March discharge at Copiapó River at Pastillo ($R^2 = 0.58$; $r = 0.76$; $N = 22$) confirms that snow permanence exerts a first-order control on seasonal runoff. The 42 % of unexplained variance is consistent with longer-residence cryospheric storage (rock glaciers, debris-covered ice, ice-rich permafrost) sustaining baseflow during low-snow years, which is precisely the contribution the NCI is designed to capture.

Ice mass is strongly concentrated: the Montosa (03413; 0.92 Gt) and Del Potro (03412; 0.91 Gt) sub-watersheds alone dominate the basin's cryospheric storage, with Manflas (03420; 0.37 Gt) third. Debris-free glaciers are the principal contributor in these top reservoirs, whereas ice-rich permafrost landforms—particularly gelifluction taluses and rock glaciers—account for a substantial and, at the basin scale, cumulatively important fraction, especially in intermediate and smaller sub-watersheds. The NCI consistently ranks Montosa (0.93), Del Potro (0.89) and Manflas (0.47) as the primary priority sub-watersheds, which together concentrate ~71 % of the ice mass inventoried across the 12 cryospheric sub-watersheds (Table 6). These three are the basins most likely to sustain runoff during droughts and should be prioritized in the national water-resources strategy. Estero Come Caballos (03402; NCI = 0.34) and Cachitos (03403; NCI = 0.31) emerge as secondary, snow-driven basins: in high-snowfall years their hydrological productivity rises, so management there should focus on capturing increased runoff rather than on protecting long-term ice. Adding these two raises the prioritized share of inventoried ice mass to ~85 %. This dual behaviour—southern, ice-rich basins safeguarding dry-year baseflow versus northern, snow-rich basins driving wet-year runoff—is the central management insight of the index.

We have developed a new workflow to identify cryospheric landforms and snow and applied this method to the Copiapó watershed. The workflow consists of mapping cryospheric landforms and snow occurrences and the calculation of a normalized cryospheric index (NCI) for sub-watersheds.

Our cryospheric landforms and snow occurrence mapping exercise resulted in the identification of cryospheric landforms in more isolated bodies north of Cadillal Hill and more continuous cryospheric landforms south of Caserones Hill. Gelifluction taluses constitute a critical strategic cryospheric ice storage for safeguarding water security in northern Chile's watersheds. These landforms represent the periglacial type most likely to contain the largest ice volumes within the region's cryosphere, a hypothesis that requires further validation through targeted ERT (Electrical Resistivity Tomography) geophysical surveys of such cryospheric features.

The NCI allowed us to identify four main targets for watershed management. The four identified main targets are the Montosa river (03413), Del Potro river (03412), and Manflas river watershed (03420) with NCI of 0.93, 0.89 and 0.47 respectively. We propose that these sub-watersheds are the most important because that can sustain the runoff in the watershed in times of drought. In years with snowfall the northern sub-watersheds as the Estero Come Caballos sub-watershed (03402) have the potential to be highly productive hydraulically by snow melt. This ranking allows the prioritization of these sub-watersheds. We conclude that the Montosa river (03413), Del Potro river (03412) and Manflas river (03420) are the sub-watersheds that should be prioritized in the national water resources strategy because of the high NCI.

6.2 Relevant topics detected Limitations of this work and future research

Three findings have implications beyond the Copiapó basin. First, gelifluction taluses and protalus lobes—which dominate the arid Atacama cryosphere by extent and store non-negligible ice—are currently excluded from Chile's national periglacial inventory, which systematizes only rock glaciers. Their omission means national ice-storage assessments likely underestimate

the periglacial reserve in northern Chile. Second, our ERT surveys returned ice-rich cores ~15 m thick in a gelifluction talus and 20–35 m in an active rock glacier, an order of magnitude greater than thicknesses assumed for analogous landforms in parts of the literature, reinforcing that periglacial ice is being undercounted and warranting targeted geophysical campaigns. Third, the very low DInSAR surface velocities (mm yr^{-1}), one to two orders of magnitude slower than rock glaciers further south in the Dry Andes, are physically consistent with the colder, drier permafrost imposed by proximity to the South American Arid Diagonal and with the predominantly cryogenic (rather than glacial) origin of these landforms; this constrains how kinematic models should be transferred across the Andes.

The workflow developed in this study enables the assessment of the role of a specific cryospheric sub-watershed within a broader drainage system, providing a basis for establishing protection frameworks in water resource management. This tool is transferable to other arid regions that are sourced through cryospheric landforms and snow. However, the method is calibrated for the Copiapó river watershed, and complete resource and reserve data should be available for the NCI method to be used on other watersheds.

To ensure more accurate error propagation in the NCI definition, it is essential to compute the uncertainty associated with the fractional snow cover product. We believe that assuming a 10% error under ideal conditions is one of the main limitations of the study of the snow permanence of the watershed. Our goal is to obtain a code that allows us to understand the uncertainties associated with the detection of snow in the mountainous areas of the watershed in order to improve the NCI and take it to the next level of research.

6.3 Limitations and outlook

The NCI is calibrated for the Copiapó watershed and requires complete resource and reserve data to be transferred elsewhere. The main quantified uncertainty in the snow term is the assumed ~10 % accuracy of the MODIS fractional-snow-cover product under ideal conditions; deriving a spatially explicit FSC uncertainty for high-mountain terrain is the priority next step to refine the index. Given projections of reduced glacier meltwater under future Southern Andes megadroughts (Ayala, 2025), identifying and protecting high-NCI cryospheric sub-watersheds is increasingly urgent. The framework provides technical, objective input for the DGA, water-monitoring boards and groundwater communities, and offers a methodological basis for integrating glacial and periglacial reserves into watershed-protection policy—and potentially into a legal instrument to protect Chile's cryosphere and its associated water resources.

The tool provides technical and objective data that water control agencies and those in charge of integrated watershed management, such as the DGA (General Directorate of Water), oversight boards and groundwater communities, should consider when implementing policy on the watershed management of the Copiapó watershed.

This study could help other arid regions of the planet with cryospheric landforms and snow to be able to manage the territory with a strategic plan to address the management of these cryospheric components.

1123 **7. Open science statement**

1124 The data generated for this project will become available upon publication. The cryospheric landforms data can be downloaded
1125 via this link: [https://doi.org/10.5281/zenodo.15633086]. The glacial and periglacial inventory for the Atacama region can be
1126 downloaded via this link: [https://doi.org/10.5281/zenodo.14921499]. The raster image for historical snow data can be
1127 downloaded via this link [https://doi.org/10.5281/zenodo.14921552].

1128 **8. Acknowledgments**

1129 We thank our colleagues of the Cryosphere and Water Research Laboratory for their indispensable contribution to the high
1130 altitude expeditions, and especially to project 5 of “Centro Avanzado para Tecnologías del Agua” (CAPTA) of the
1131 “Corporación de Fomento de la Producción” (CORFO) named “Tecnologías remotas y de campo para para el aumento de
1132 oferta hídrica desde fuentes superficiales y subterráneas: acumulación nival y la recarga de acuíferos”, the Regional
1133 Government of Atacama and the University of Atacama, which sponsored the projects that made it possible to carry out this
1134 research and special thanks to Mr. Adrien Tavernier and Dra. Nicole Shaffer who has provided technical support in the
1135 execution of this work.

1136 **9. References**

1137 ~~Abraham, E. M., Rodríguez, M. G. L., Rubio, M. C., Guida-Johnson, B., Gomez, L. V., & Rubio, C. (2020). Disentangling~~
1138 ~~the concept of -“South American Arid Diagonal”. Journal of Arid Environments, 175, 104089.~~
1139 ~~https://doi.org/10.1016/j.jaridenv.2019.104089.~~Abraham, E. M., Rodríguez, M. G. L., Rubio, M. C., Guida-Johnson, B.,
1140 Gomez, L. V., and Rubio, C.: Disentangling the concept of "South American Arid Diagonal", J. Arid Environ., 175, 104089,
1141 https://doi.org/10.1016/j.jaridenv.2019.104089, 2020.

1142 ~~Ahn, S. R., & Kim, S. J. (2017). Assessment of integrated watershed health based on the natural environment, hydrology,~~
1143 ~~water quality, and aquatic ecology. Hydrology And Earth System Sciences, 21(11), 5583–5602. https://doi.org/10.5194/hess-~~
1144 ~~21-5583-2017.~~Ahn, S. R. and Kim, S. J.: Assessment of integrated watershed health based on the natural environment,
1145 hydrology, water quality, and aquatic ecology, Hydrol. Earth Syst. Sci., 21, 5583–5602, https://doi.org/10.5194/hess-21-5583-
1146 2017, 2017.

1147 Ayala A, Pellicciotti F, Macdonell S, Mephee J, Vivero S, Campos C, Egli P. (2016). Modelling the hydrological response of
1148 debris-free and debris-covered glaciers to present climatic conditions in the semiarid Andes of central Chile. Hydrol Process
1149 30:4036–4058. https://doi.org/10.1002/hyp.10971.Ayala, A., Pellicciotti, F., Macdonell, S., McPhee, J., Vivero, S., Campos,
1150 C., and Egli, P.: Modelling the hydrological response of debris-free and debris-covered glaciers to present climatic conditions
1151 in the semiarid Andes of central Chile, Hydrol. Process., 30, 4036–4058, https://doi.org/10.1002/hyp.10971, 2016.

Con formato

Ayala, Á., Fariás-Barahona, D., Huss, M., Pellicciotti, F., McPhee, J., and Farinotti, D. (2020) Glacier runoff variations since 1955 in the Maipo river Basin, semiarid Andes of central Chile, *The Cryosphere*, 14, 2005–2027, <https://doi.org/10.5194/tc-14-2005-2020>.

Ayala, Á., Fariás-Barahona, D., Huss, M., Pellicciotti, F., McPhee, J., and Farinotti, D.: Glacier runoff variations since 1955 in the Maipo River Basin, semiarid Andes of central Chile, *The Cryosphere*, 14, 2005–2027, <https://doi.org/10.5194/tc-14-2005-2020>, 2020.

Ayala, Á., Muñoz-Castro, E., Farinotti, D., Fariás-Barahona, D., Mendoza, P. A., MacDonell, S., McPhee, J., Vargas, X., & Pellicciotti, F. (2025). Less water from glaciers during future megadroughts in the Southern Andes. *Communications Earth & Environment*, 6, 860. <https://doi.org/10.1038/s43247-025-02845-6>.

Ayala, Á., Muñoz-Castro, E., Farinotti, D., Fariás-Barahona, D., Mendoza, P. A., MacDonell, S., McPhee, J., Vargas, X., and Pellicciotti, F.: Less water from glaciers during future megadroughts in the Southern Andes, *Commun. Earth Environ.*, 6, 860, <https://doi.org/10.1038/s43247-025-02845-6>, 2025.

Azócar G. F. & Brenning, A. (2010). Hydrological and geomorphological significance of rock glaciers in the dry Andes, Chile (27°–33°S). *Permafrost Periglac. Process.*, 21, 42–53.

Azócar, G. F. and Brenning, A.: Hydrological and geomorphological significance of rock glaciers in the dry Andes, Chile (27°–33°S), *Permafrost Periglac. Process.*, 21, 42–53, <https://doi.org/10.1002/ppp.669>, 2010.

Bahr, D. B., Meier, M. F., & Peckham, S. D. (1997). The physical basis of glacier volume-area scaling, *Journal of Geophysical Research*, 102(B9), 20,355–20,362.

Bahr, D. B., Meier, M. F., and Peckham, S. D.: The physical basis of glacier volume-area scaling, *J. Geophys. Res. Solid Earth*, 102, 20355–20362, <https://doi.org/10.1029/97JB01696>, 1997.

Barsch, D. (1996). *Rock Glaciers: Indicators for the Present and Former Geoecology in High Mountain Environments*. Springer (Series in the Physical Environment 16), Berlin, p. 331.

Basuki, T. M., Nugroho, H. Y. S. H., Indrajaya, Y., Pramono, I. B., Nugroho, N. P., Supangat, A. B., Indrawati, D. R., Savitri, E., Wahyuningrum, N., Purwanto, Cahyono, S. A., Putra, P. B., Adi, R. N., Nugroho, A. W., Auliyani, D., Wuryanta, A., Riyanto, H. D., Harjadi, B., Yudilastyantoro, C., ... Simarmata, D. P. (2022). Improvement of integrated watershed management in Indonesia for mitigation and adaptation to climate change: A review. *Sustainability*, 14(16), 9997. <https://doi.org/10.3390/su14169997>.

Basuki, T. M., Nugroho, H. Y. S. H., Indrajaya, Y., Pramono, I. B., Nugroho, N. P., Supangat, A. B., Indrawati, D. R., Savitri, E., Wahyuningrum, N., Purwanto, Cahyono, S. A., Putra, P. B., Adi, R. N., Nugroho, A. W., Auliyani, D., Wuryanta, A., Riyanto, H. D., Harjadi, B., Yudilastyantoro, C., and Simarmata, D. P.: Improvement of integrated watershed management in Indonesia for mitigation and adaptation to climate change: a review, *Sustainability*, 14, 9997, <https://doi.org/10.3390/su14169997>, 2022.

Beniston, M., & Stoffel, M. (2014). Assessing the impacts of climatic change on mountain water resources. *Science of The Total Environment*, 493, 1129–1137. <https://doi.org/10.1016/j.scitotenv.2013.11.122>.

Beniston, M. and Stoffel, M.: Assessing the impacts of climatic change on mountain water resources, *Sci. Total Environ.*, 493, 1129–1137, <https://doi.org/10.1016/j.scitotenv.2013.11.122>, 2014.

Beniston, M., Farinotti, D., Stoffel, M., Andreassen, L. M., Coppola, E., Eckert, N., Fantini, A., Giacona, F., Hauck, C., Huss, M., Huwald, H., Lehning, M., López-Moreno, J. I., Magnusson, J., Marty, C., Morán-Tejeda, E., Morin, S., Naaïm, M., Provenzale, A., Vincent, C. (2018). The European Mountain Cryosphere: A review of its current state, trends, and future challenges. *The Cryosphere*, 12(2), 759–794. <https://doi.org/10.5194/tc-12-759-2018>. Beniston, M., Farinotti, D., Stoffel, M., Andreassen, L. M., Coppola, E., Eckert, N., Fantini, A., Giacona, F., Hauck, C., Huss, M., Huwald, H., Lehning, M., López-Moreno, J. I., Magnusson, J., Marty, C., Morán-Tejeda, E., Morin, S., Naaïm, M., Provenzale, A., and Vincent, C.: The European mountain cryosphere: a review of its current state, trends, and future challenges, *The Cryosphere*, 12, 759–794, <https://doi.org/10.5194/tc-12-759-2018>, 2018.

Bolch, T., Marchenko, S. (2006). Significance of glaciers, rock glaciers, and ice-rich permafrost in the Northern Tien Shan as water towers under climate change conditions. In: *Proceedings of the Workshop “Assessment of Snow-Glacier and Water Resources in Asia”*, 28–30 November 2006, Almaty, S. 199–211. Bolch, T. and Marchenko, S.: Significance of glaciers, rock glaciers, and ice-rich permafrost in the Northern Tien Shan as water towers under climate change conditions, in: *Proceedings of the Workshop Assessment of Snow-Glacier and Water Resources in Asia*, 28–30 November 2006, Almaty, Kazakhstan, 199–211, 2006.

Bórquez, R., Larraín, S., Polanco, R., and J. Urquidí (2006). Glaciares chilenos. Reservas estratégicas de agua dulce para la sociedad, los ecosistemas y la economía. Chile Sustentable, LOM ediciones, Santiago, 124. Bórquez, R., Larraín, S., Polanco, R., and Urquidí, J.: Glaciares chilenos. Reservas estratégicas de agua dulce para la sociedad, los ecosistemas y la economía. Chile Sustentable, LOM ediciones, Santiago, Chile, 124 pp., 2006.

Bozkurt, D., Rojas, M., Boisier, J. P. & Valdivieso, J. (2017). Climate change impacts on hydroclimatic regimes and extremes over Andean basins in Central Chile. *Hydrology and Earth System Sciences*, Discuss, <https://doi.org/10.5194/hess-2016-690>, 2017. Bozkurt, D., Rojas, M., Boisier, J. P., and Valdivieso, J.: Climate change impacts on hydroclimatic regimes and extremes over Andean basins in central Chile, *Hydrol. Earth Syst. Sci. Discuss.*, <https://doi.org/10.5194/hess-2016-690>, 2017.

Brenning, A. (2005). Climatic and geomorphological controls of rock glaciers in the Andes of Central Chile: combining statistical modelling and field mapping. PhD Thesis, Humboldt University, Berlin, Germany. Brenning, A.: Climatic and geomorphological controls of rock glaciers in the Andes of central Chile: combining statistical modelling and field mapping. PhD Thesis, Humboldt University, Berlin, Germany, 2005.

Brenning, A. (2008). The impact of mining on rock glaciers and glaciers: examples from Central Chile. In B. S. Orlove, E. Wiegandt, & B. Luckman (Eds.), *Darkening Peaks: Glacier Retreat, Science, and Society* (pp. 196–205). University of California Press. Brenning, A.: The impact of mining on rock glaciers and glaciers: examples from central Chile, in: *Darkening Peaks: Glacier Retreat, Science, and Society*, edited by: Orlove, B. S., Wiegandt, E., and Luckman, B., University of California Press, Berkeley, USA, 196–205, 2008.

Burger KC, Degenhardt JJ, Giardino JR (1999). Engineering geomorphology of rock glaciers. *Geomorphology*, 31: 93–132.

Con formato: Español (Chile)

1219 Burger, F., Ayala, Á., Fariás, D., Shaw, T. E., MacDonell, S., Brock, B., McPhee, J., & Pellicciotti, F. (2019). Interannual
1220 variability in glacier contribution to runoff from a high-elevation Andean catchment: understanding the role of debris cover in
1221 glacier hydrology. *Hydrological Processes*, 33(2), 214–229. <https://doi.org/10.1002/hyp.13354> Burger, F., Ayala, Á., Fariás,
1222 D., Shaw, T. E., MacDonell, S., Brock, B., McPhee, J., and Pellicciotti, F.: Interannual variability in glacier contribution to
1223 runoff from a high-elevation Andean catchment: understanding the role of debris cover in glacier hydrology, *Hydrol. Process.*,
1224 33, 214–229, <https://doi.org/10.1002/hyp.13354>, 2019.

1225 Casassa, G., Haeberli, W., Jones, G., Kaser, G., Ribstein, P., Rivera, A., & Schneider, C. (2007). Current status of Andean
1226 glaciers. *Global And Planetary Change*, 59(1–4), 1–9. <https://doi.org/10.1016/j.gloplacha.2006.11.013> Casassa, G., Haeberli,
1227 W., Jones, G., Kaser, G., Ribstein, P., Rivera, A., and Schneider, C.: Current status of Andean glaciers, *Global Planet. Change*,
1228 59, 1–9, <https://doi.org/10.1016/j.gloplacha.2006.11.013>, 2007.

1229 Clarke, J. (2006). Antiquity of aridity in the Chilean Atacama Desert. *Geomorphology*, N°73, pp. 101–114.
1230 <https://doi.org/10.1016/j.geomorph.2005.06.008> Clarke, J.: Antiquity of aridity in the Chilean Atacama Desert,
1231 *Geomorphology*, 73, 101–114, <https://doi.org/10.1016/j.geomorph.2005.06.008>, 2006.

1232 Corte, A.E. (1978). Rock glaciers as permafrost bodies with debris cover as an active layer. A Hydrological approach, Andes
1233 of Mendoza, Argentina. In Natural Resources Council of Canada (ed.), *Proceedings of the third International Conference on*
1234 *Permafrost*, Edmonton, Alberta, Canada. Ottawa: Natural Resources Council of Canada, pp. 262–269.

1235 Croce, F., & Milana, J.P. (2002). Internal Structure and Behaviours of a rock glacier in the árid Andes of Argentina. *Permafrost*
1236 *Periglac. Process.*, 13: 289–299.

1237 Cicoira, A., Marcet, M., Gärtner-Roer, I., Bodin, X., Arenson, L. U., & Vieli, A. (2021). A general theory of rock glacier
1238 creep based on in-situ and remote sensing observations. *Permafrost And Periglacial Processes*, 32(1), 139–153.
1239 <https://doi.org/10.1002/ppp.2090> Cicoira, A., Marcet, M., Gärtner-Roer, I., Bodin, X., Arenson, L. U., and Vieli, A.: A general
1240 theory of rock glacier creep based on in-situ and remote sensing observations, *Permafrost Periglac. Process.*, 32, 139–153,
1241 <https://doi.org/10.1002/ppp.2090>, 2021.

1242 Delleur, J. W., (1999). *The handbook of groundwater engineering*. Boca Raton, FL: CRC Press.

1243 DGA (2013). *Estrategia Nacional de Recursos Hídricos 2012–2025. I Resumen ejecutivo*, p. 40. [https://www.mop.gob.cl/wp-](https://www.mop.gob.cl/wp-content/uploads/2021/04/Estrategia_Nacional_Recursos_Hidricos_DGA.pdf)
1244 [content/uploads/2021/04/Estrategia_Nacional_Recursos_Hidricos_DGA.pdf](https://www.mop.gob.cl/wp-content/uploads/2021/04/Estrategia_Nacional_Recursos_Hidricos_DGA.pdf)

1245 Dirección General de Aguas (DGA), 2014. Estimación de volúmenes de hielo mediante sondeos de radar en zonas Norte,
1246 Central y Sur. Informe Final, S.I.T. No. 338. Ministerio de Obras Públicas, Santiago, Chile.
1247 <https://bibliotecadigital.ciren.cl/handle/20.500.13082/32675> Dirección General de Aguas (DGA): Estimación de volúmenes
1248 de hielo mediante sondeos de radar en zonas Norte, Central y Sur, Informe Final, S.I.T. No. 338, Ministerio de Obras Públicas,
1249 Santiago, Chile, available at: <https://bibliotecadigital.ciren.cl/handle/20.500.13082/32675> (last access: 1 June 2025), 2014.

1250 Evans, I. S. (2006). Local aspect asymmetry of mountain glaciation: A global survey of consistency of favoured directions for
1251 glacier numbers and altitudes. *Geomorphology*, 73(1–2), 166–184. <https://doi.org/10.1016/j.geomorph.2005.07.009> Evans, I.

Con formato: Español (Chile)

S.: Local aspect asymmetry of mountain glaciation: a global survey of consistency of favoured directions for glacier numbers and altitudes, *Geomorphology*, 73, 166–184, <https://doi.org/10.1016/j.geomorph.2005.07.009>, 2006.

Favier, V., Falvey, M., Rabatel, A., Praderio, E., & Lopez, D. H. (2009). Interpreting discrepancies between discharge and precipitation in high-altitude area of Chile's Norte Chico region (26–32°S). *Water Resources Research*, 45(2). <https://doi.org/10.1029/2008wr006802>.

Farinotti, D., Brinkerhoff, D. J., Clarke, G. K. C., Fürst, J. J., Frey, H., Gantayat, P., Gillet-Chaulet, F., Girard, C., Huss, M., Leclercq, P. W., Linsbauer, A., Machguth, H., Martin, C., Maussion, F., Morlighem, M., Mosbeux, C., Pandit, A., Portmann, A., Rabatel, A., and Andreassen, L. M.: How accurate are estimates of glacier ice thickness? Results from ITMIX, the Ice Thickness Models Intercomparison eXperiment, *The Cryosphere*, 11, 949–970, <https://doi.org/10.5194/tc-11-949-2017>, 2017.

Flores, B., García, A., & Ulloa, C. (2018). Evolución espacial y temporal de glaciares descubiertos en la Región de Atacama, Chile. *XV Congreso Geológico Chileno*, pp. 742–745.

Farinotti, D., Huss, M., Fürst, J. J., Landmann, J., Machguth, H., Maussion, F., and Pandit, A.: A consensus estimate for the ice thickness distribution of all glaciers on Earth, *Nat. Geosci.*, 12, 168–173, <https://doi.org/10.1038/s41561-019-0300-3>, 2019.

Farinotti, D., Brinkerhoff, D. J., Clarke, G. K. C., Fürst, J. J., Frey, H., Gantayat, P., Gillet-Chaulet, F., Girard, C., Huss, M., Leclercq, P. W., Linsbauer, A., Machguth, H., Martin, C., Maussion, F., Morlighem, M., Mosbeux, C., Pandit, A., Portmann, A., Rabatel, A., Andreassen, L. M. (2017). How accurate are estimates of glacier ice thickness? Results from ITMIX, the Ice Thickness Models Intercomparison eXperiment. *The Cryosphere*, 11(2), 949–970. <https://doi.org/10.5194/tc-11-949-2017>.

Favier, V., Falvey, M., Rabatel, A., Praderio, E., and Lopez, D. H.: Interpreting discrepancies between discharge and precipitation in high-altitude area of Chile's Norte Chico region (26–32°S), *Water Resour. Res.*, 45, W02424, <https://doi.org/10.1029/2008WR006802>, 2009.

Farinotti, D., Huss, M., Fürst, J. J., Landmann, J., Machguth, H., Maussion, F., & Pandit, A. (2019). A consensus estimate for the ice thickness distribution of all glaciers on Earth. *Nature Geoscience*, 12(3), 168–173. <https://doi.org/10.1038/s41561-019-0300-3>.

Ferri, L., Dussaillant, I., Zalazar, L., Masiokas, M. H., Ruiz, L., Pitte, P., Gargantini, H., Castro, M., Berthier, E., and Villalba, R.: Ice mass loss in the central Andes of Argentina between 2000 and 2018 derived from a new glacier inventory and satellite stereo-imagery, *Front. Earth Sci.*, 8, 530997, <https://doi.org/10.3389/feart.2020.530997>, 2020.

Ferri, L., Dussaillant, I., Zalazar, L., Masiokas, M. H., Ruiz, L., Pitte, P., Gargantini, H., Castro, M., Berthier, E., & Villalba, R. (2020). Ice mass loss in the Central Andes of Argentina between 2000 and 2018 derived from a new glacier inventory and satellite stereo imagery. *Frontiers in Earth Science*, 8, 530997. <https://doi.org/10.3389/feart.2020.530997>.

Flores, B., García, A., and Ulloa, C.: Evolución espacial y temporal de glaciares descubiertos en la Región de Atacama, Chile, *XV Congreso Geológico Chileno*, 742–745, 2018.

García, A., Ulloa, C., Amigo, G., Milana, J. P., Medina, C. (2017). An inventory of cryospheric landforms in the arid diagonal of South America (High Central Andes, Atacama Region, Chile). *Quaternary International*, N°438, pp.4–19, <https://doi.org/10.1016/j.quaint.2017.04.033>.

García, A., Ulloa, C., Amigo, G., Milana, J. P., and Medina, C.: An inventory of

Con formato: Español (Chile)

cryospheric landforms in the arid diagonal of South America (High Central Andes, Atacama Region, Chile), *Quaternary Int.*, 438, 4–19, <https://doi.org/10.1016/j.quaint.2017.04.033>, 2017.

Garreaud, R., Boisier, J. P., Rondanelli, R., Montecinos, A., Sepúlveda, H. H., & Veloso-Águila, D. (2019). The Central Chile Mega Drought (2010–2018): A climate dynamics perspective. *International Journal Of Climatology*, 40(1), 421–439. <https://doi.org/10.1002/joc.6219>

Garreaud, R., Boisier, J. P., Rondanelli, R., Montecinos, A., Sepúlveda, H. H., and Veloso-Águila, D.: The central Chile mega drought (2010–2018): a climate dynamics perspective, *Int. J. Climatol.*, 40, 421–439, <https://doi.org/10.1002/joc.6219>, 2020.

Gärtner-Roer, I., Naegeli, K., Huss, M., Knecht, T., Machguth, H., & Zemp, M. (2014). A database of worldwide glacier thickness observations. *Global And Planetary Change*, 122, 330–344. <https://doi.org/10.1016/j.gloplacha.2014.09.003>.

Gascoin S., Kinnard C., Ponce R., Lhermitte S., Macdonell S., Rabatel A. (2011). Glacier contribution to streamflow in two headwaters of the Huasco river, Dry Andes of Chile. *Cryosphere* 5:1099–1113. <https://doi.org/10.5194/tc-5-1099-2011>

Gascoin, S., Kinnard, C., Ponce, R., Lhermitte, S., Macdonell, S., and Rabatel, A.: Glacier contribution to streamflow in two headwaters of the Huasco River, Dry Andes of Chile, *The Cryosphere*, 5, 1099–1113, <https://doi.org/10.5194/tc-5-1099-2011>, 2011.

Gruber, S. (2012). Derivation and analysis of a high-resolution estimate of global permafrost zonation. *The Cryosphere*, 6(1), 221–233. <https://doi.org/10.5194/tc-6-221-2012>

Gruber, S.: Derivation and analysis of a high-resolution estimate of global permafrost zonation, *The Cryosphere*, 6, 221–233, <https://doi.org/10.5194/tc-6-221-2012>, 2012.

Halla, C., Blöthe, J.H., Tapia, C., Trombotto, D., Hilbich, C., Hauck, C., Schrott, L. (2021). Ice content and interannual water storage changes of an active rock glacier in the dry Andes of Argentina. *The Cryosphere*, 15, 1187–1213. <https://doi.org/10.5194/tc-15-1187-2021>

Hall, D. K., Riggs, G. A., Salomonson, V. V., DiGirolamo, N. E., and Bayr, K. J.: MODIS snow-cover products, *Remote Sens. Environ.*, 83, 181–194, [https://doi.org/10.1016/S0034-4257\(02\)00095-0](https://doi.org/10.1016/S0034-4257(02)00095-0), 2002.

Hall, D. K., Riggs, G. A., Salomonson, V. V., DiGirolamo, N. E., & Bayr, K. J. (2002). MODIS snow-cover products: Remote Sensing of Environment, 83(1–2), 181–194. [https://doi.org/10.1016/S0034-4257\(02\)00095-0](https://doi.org/10.1016/S0034-4257(02)00095-0)

Halla, C., Blöthe, J. H., Tapia, C., Trombotto, D., Hilbich, C., Hauck, C., and Schrott, L.: Ice content and interannual water storage changes of an active rock glacier in the dry Andes of Argentina, *The Cryosphere*, 15, 1187–1213, <https://doi.org/10.5194/tc-15-1187-2021>, 2021.

Hanssen, R. F. (2001). Radar interferometry: Data interpretation and error analysis. Springer. <https://doi.org/10.1007/0-306-47633-9>

Hanssen, R. F.: Radar Interferometry: Data Interpretation and Error Analysis, Springer, Dordrecht, the Netherlands, <https://doi.org/10.1007/0-306-47633-9>, 2001.

Harrington, H. J. (1961). Geology of Parts of Antofagasta and Atacama Provinces, Northern Chile. *AAPG Bulletin*, 45. <https://doi.org/10.1306/0bda6332-16bd-11d7-8645000102e1865d>

Harrington, H. J.: Geology of parts of Antofagasta and Atacama provinces, northern Chile, *AAPG Bull.*, 45, 169–197, <https://doi.org/10.1306/0BDA6332-16BD-11D7-8645000102C1865D>, 1961.

Hauck, C., Bach, M., & Hilbich, C. (2008). A four-phase model to quantify subsurface ice and water content in permafrost regions based on geophysical data sets. In K. M. Hinkel & D. L. Kane (Eds.), *Proceedings of the Ninth International Conference on Permafrost* (Vol. 1, pp. 675–680). Institute of Northern Engineering, University of Alaska Fairbanks.

Hauck, C., Bach, M., and Hilbich, C.: A four-phase model to quantify subsurface ice and water content in permafrost regions based on geophysical data sets, in: *Proceedings of the Ninth International Conference on Permafrost*, edited by: Hinkel, K. M. and Kane, D. L., Institute of Northern Engineering, University of Alaska Fairbanks, USA, Vol. 1, 675–680, 2008.

Hilbich, C., Hauck, C., Mollaret, C., Wainstein, P., & Arenson, L. U. (2022). Towards accurate quantification of ice content in permafrost of the Central Andes—Part 1: Geophysics-based estimates from three different regions. *The Cryosphere*, 16(5), 1845–1872. <https://doi.org/10.5194/tc-16-1845-2022>.

Hilbich, C., Hauck, C., Mollaret, C., Wainstein, P., and Arenson, L. U.: Towards accurate quantification of ice content in permafrost of the Central Andes – Part 1: geophysics-based estimates from three different regions, *The Cryosphere*, 16, 1845–1872, <https://doi.org/10.5194/tc-16-1845-2022>, 2022.

Hilbich, C., Marescot, L., Hauck, C., Loke, M. H., & Mäusbacher, R. (2009). Applicability of electrical resistivity tomography monitoring to coarse blocky and ice-rich permafrost landforms. *Permafrost and Periglacial Processes*, 20(3), 269–284.

Hilbich, C., Marescot, L., Hauck, C., Loke, M. H., and Mäusbacher, R.: Applicability of electrical resistivity tomography monitoring to coarse blocky and ice-rich permafrost landforms, *Permafrost Periglac. Process.*, 20, 269–284, <https://doi.org/10.1002/ppp.641>, 2009.

Houston, J (2006). Variability of precipitation in the Atacama desert: its causes and hydrological impact. *International journal of climatology*, 26: pp. 2181–2198.

Hu, Y., Liu, L., Huang, L., Zhao, L., Wu, T., Wang, X., & Cai, J. (2023). Mapping and characterizing rock glaciers in the arid Western Kunlun Mountains supported by InSAR and deep learning. *Journal of Geophysical Research: Earth Surface*, 128(9), e2023JF007206. <https://doi.org/10.1029/2023JF007206>.

Hu, Y., Liu, L., Huang, L., Zhao, L., Wu, T., Wang, X., and Cai, J.: Mapping and characterizing rock glaciers in the arid Western Kunlun Mountains supported by InSAR and deep learning, *J. Geophys. Res. Earth Surf.*, 128, e2023JF007206, <https://doi.org/10.1029/2023JF007206>, 2023.

Insel, N., Poulsen, C. J., & Ehlers, T. A. (2009). Influence of the Andes Mountains on South American moisture transport, convection, and precipitation. *Climate Dynamics*, 35(7–8), 1477–1492. <https://doi.org/10.1007/s00382-009-0637-1>.

Insel, N., Poulsen, C. J., and Ehlers, T. A.: Influence of the Andes Mountains on South American moisture transport, convection, and precipitation, *Clim. Dynam.*, 35, 1477–1492, <https://doi.org/10.1007/s00382-009-0637-1>, 2009.

Izquierdo, T., Rivera, A. L., Galeano, Á., Gallardo, D., Salas, V., Aparicio, O., Buylaert, J. P., Ruiz, F., & Abad, M. (2024). Historical catastrophic floods at the southern edge of the Atacama Desert: A multi-archive reconstruction of the Copiapo river extreme events. *Global and Planetary Change*, 236(104411), 104411. <https://doi.org/10.1016/j.gloplacha.2024.104411>.

Izquierdo, T., Rivera, A. L., Galeano, Á., Gallardo, D., Salas, V., Aparicio, O., Buylaert, J. P., Ruiz, F., and Abad, M.: Historical catastrophic floods at the southern edge of the Atacama Desert: a multi-archive reconstruction of the Copiapó River extreme events, *Global Planet. Change*, 236, 104411, <https://doi.org/10.1016/j.gloplacha.2024.104411>, 2024.

Con formato: Inglés (Estados Unidos)

Janke, J.R., Ng, S., Bellisario, A. (2017). An inventory and estimate of water stored in firn fields, glaciers, debris-covered glaciers, and rock glaciers in the Aconcagua river Basin, Chile. *Geomorphology*, N°296, pp. 142–152.

Jaiswal, R. K., Ghosh, N. C., Galkate, R. V., & Thomas, T. (2015). Multi criteria decision analysis (MCDA) for watershed prioritization. *Aquatic Procedia*, 4, 1553–1560. <https://doi.org/10.1016/j.aqpro.2015.02.201>.

Jaiswal, R. K., Ghosh, N. C., Galkate, R. V., and Thomas, T.: Multi criteria decision analysis (MCDA) for watershed prioritization, *Aquat. Procedia*, 4, 1553–1560, <https://doi.org/10.1016/j.aqpro.2015.02.201>, 2015.

Jara, F. M., Lagos-Zúñiga, M., Fuster, R., Mattar, C., & McPhee, J. (2021). Snow Processes and Climate Sensitivity in an Arid Mountain Region, Northern Chile. *Atmosphere*, 12(4), 520. <https://doi.org/10.3390/atmos12040520>.

Jara, F. M., Lagos-Zúñiga, M., Fuster, R., Mattar, C., and McPhee, J.: Snow processes and climate sensitivity in an arid mountain region, northern Chile, *Atmosphere*, 12, 520, <https://doi.org/10.3390/atmos12040520>, 2021.

Jones, D., Harrison, S., Anderson, K., & Betts, R. (2018). Mountain rock glaciers contain globally significant water stores. *Scientific Reports*, 8(1). <https://doi.org/10.1038/s41598-018-21244-w>.

Jones, D., Harrison, S., Anderson, K., and Betts, R.: Mountain rock glaciers contain globally significant water stores, *Sci. Rep.*, 8, 2834, <https://doi.org/10.1038/s41598-018-21244-w>, 2018.

Jones, D., Harrison, S., Anderson, K., & Whalley, W. B (2019). Rock glaciers and mountain hydrology: A review. *Earth-Science Reviews*, 193, 66–90. <https://doi.org/10.1016/j.earscirev.2019.04.001>.

Pan, F., Jiang, L., Wang, G., Pan, J., Huang, J., et al. (2024). MODIS daily cloud-gap-filled fractional snow cover dataset of the Asian Water Tower from 2000 to 2022. *Earth System Science Data*, 16(6), 2501–2523. <https://doi.org/10.5194/essd-16-2501-2024>.

Pan, F., Jiang, L., Wang, G., Pan, J., and Huang, J.: MODIS daily cloud-gap-filled fractional snow cover dataset of the Asian Water Tower from 2000 to 2022, *Earth Syst. Sci. Data*, 16, 2501–2523, <https://doi.org/10.5194/essd-16-2501-2024>, 2024.

Kay, S. M., Mpodozis, C., Tittler, A., & Cornejo, P (1994). Tertiary Magmatic Evolution of the Maricunga Mineral Belt in Chile. *International Geology Review*, 36(12), 1079–1112. <https://doi.org/10.1080/00206819409465506>.

Kay, S. M., Mpodozis, C., Tittler, A., and Cornejo, P.: Tertiary magmatic evolution of the Maricunga mineral belt in Chile, *Int. Geol. Rev.*, 36, 1079–1112, <https://doi.org/10.1080/00206819409465506>, 1994.

Le Quesne C., DW Stahle, MK Cleaveland, MD Therrell, JC Aravena, J Barichivich (2006). Ancient Austrocedrus tree-ring chronologies used to reconstruct central Chile precipitation variability from A.D. 1200 to 2000. *Journal of Climate* 19(22): 5731–5744.

Le Quesne, C., Stahle, D. W., Cleaveland, M. K., Therrell, M. D., Aravena, J. C., and Barichivich, J.: Ancient Austrocedrus tree-ring chronologies used to reconstruct central Chile precipitation variability from A.D. 1200 to 2000, *J. Climate*, 19, 5731–5744, <https://doi.org/10.1175/JCLI3935.1>, 2006.

Kirkbride, M. P. (2011). Debris-covered glaciers. En V. P. Singh, P. Singh, & U. K. Haritashya (Eds.), *Encyclopedia of snow, ice and glaciers* (pp. 190–199). Springer. https://doi.org/10.1007/978-90-481-2642-2_622.

Kirkbride, M. P.: Debris-covered glaciers, in: *Encyclopedia of Snow, Ice and Glaciers*, edited by: Singh, V. P., Singh, P., and Haritashya, U. K., Springer, Dordrecht, the Netherlands, 190–199, https://doi.org/10.1007/978-90-481-2642-2_622, 2011.

Kneisel, C., Hauck, C., Fortier, D., & Moorman, B. (2008). Advances in geophysical methods for permafrost investigations. *Permafrost and Periglacial Processes*, 19(2), 157–178. <https://doi.org/10.1002/ppp.616>. Kneisel, C., Hauck, C., Fortier, D., and Moorman, B.: Advances in geophysical methods for permafrost investigations, *Permafrost Periglac. Process.*, 19, 157–178, <https://doi.org/10.1002/ppp.616>, 2008.

Hauck, C. and Kneisel, C. (Eds.): *Applied Geophysics in Periglacial Environments*, Cambridge University Press, Cambridge, UK, 240 pp., ISBN 978-0-521-88966-7, 2008. Hauck, C. and Kneisel, C. (Eds.): *Applied Geophysics in Periglacial Environments*, Cambridge University Press, Cambridge, UK, 240 pp., ISBN 978-0-521-88966-7, 2008.

Koenig, C. E. M., Hilbich, C., Hauck, C., Arenson, L. U., & Wainstein, P. (2025). Thermal state of permafrost in the Central Andes (27–34° S). *The Cryosphere*, 19, 2653–2676. <https://doi.org/10.5194/tc-19-2653-2025>. Koenig, C. E. M., Hilbich, C., Hauck, C., Arenson, L. U., and Wainstein, P.: Thermal state of permafrost in the Central Andes (27–34° S), *The Cryosphere*, 19, 2653–2676, <https://doi.org/10.5194/tc-19-2653-2025>, 2025.

Masiokas, M., Rabatel, A., Rivera, A., Ruiz, L., Pitte, P., Ceballos, J. L., Barcaza, G., Soruco, Á., Bown, F., Berthier, É., Dussailant, I., & MacDonell, S. (2020). A Review of the Current State and Recent Changes of the Andean Cryosphere. *Frontiers In Earth Science*, 8. <https://doi.org/10.3389/feart.2020.00099>. Masiokas, M., Rabatel, A., Rivera, A., Ruiz, L., Pitte, P., Ceballos, J. L., Barcaza, G., Soruco, Á., Bown, F., Berthier, É., Dussailant, I., and MacDonell, S.: A review of the current state and recent changes of the Andean cryosphere, *Front. Earth Sci.*, 8, 99, <https://doi.org/10.3389/feart.2020.00099>, 2020.

Masiokas, M., Villalba, R., Luckman, B. H., Quesne, C. L., & Aravena, J. C. (2006). Snowpack Variations in the Central Andes of Argentina and Chile, 1951–2005: Large-Scale Atmospheric Influences and Implications for Water Resources in the Region. *Journal Of Climate*, 19(24), 6334–6352. <https://doi.org/10.1175/JCLI3969.1>. Masiokas, M., Villalba, R., Luckman, B. H., Le Quesne, C., and Aravena, J. C.: Snowpack variations in the central Andes of Argentina and Chile, 1951–2005: large-scale atmospheric influences and implications for water resources in the region, *J. Climate*, 19, 6334–6352, <https://doi.org/10.1175/JCLI3969.1>, 2006.

Masiokas, M. H., Cara, L., Villalba, R., Pitte, P., Luckman, B. H., Toum, E., Christie, D. A., Quesne, C. L., & Mauget, S. (2019). Streamflow variations across the Andes (18°–55°S) during the instrumental era. *Scientific Reports*, 9(1). <https://doi.org/10.1038/s41598-019-53981-x>. Masiokas, M. H., Cara, L., Villalba, R., Pitte, P., Luckman, B. H., Toum, E., Christie, D. A., Le Quesne, C., and Mauget, S.: Streamflow variations across the Andes (18°–55°S) during the instrumental era, *Sci. Rep.*, 9, 17879, <https://doi.org/10.1038/s41598-019-53981-x>, 2019.

Matsuoka, N. (2001). Solifluction rates, processes and landforms: A global review. *Earth Science Reviews*, 55(1–2), 107–134. [https://doi.org/10.1016/S0012-8252\(01\)00057-5](https://doi.org/10.1016/S0012-8252(01)00057-5). Matsuoka, N.: Solifluction rates, processes and landforms: a global review, *Earth-Sci. Rev.*, 55, 107–134, [https://doi.org/10.1016/S0012-8252\(01\)00057-5](https://doi.org/10.1016/S0012-8252(01)00057-5), 2001.

Meier, M. F. (1969). Glaciers and Water Supply. *Journal — American Water Works Association*, 61(1), 8–12. doi:10.1002/j.1551-8833.1969.tb03696.x.

Mewes, B., Hilbich, C., Delaloye, R., & Hauck, C. (2017). Resolution capacity of geophysical monitoring regarding permafrost degradation induced by hydrological processes. *The Cryosphere*, 11, 2957–2974. <https://doi.org/10.5194/tc-11-2957-2017>.

2957-2017, Mewes, B., Hilbich, C., Delaloye, R., and Hauck, C.: Resolution capacity of geophysical monitoring regarding permafrost degradation induced by hydrological processes, *The Cryosphere*, 11, 2957–2974, <https://doi.org/10.5194/tc-11-2957-2017>, 2017.

Milana, J.P. (1998). Predicción de caudales de ríos alimentados por deshielo mediante balances de energía: Aplicación en los Andes Centrales, Argentina. *Asociación Argentina de Sedimentología*, revista, v. 5 (2), p. 53-69.

Millan, R., Mouginot, J., Rabatel, A., & Morlighem, M. (2022). Ice velocity and thickness of the world's glaciers. *Nature Geoscience*, 15(2), 124–129. <https://doi.org/10.1038/s41561-021-00885-z>. Millan, R., Mouginot, J., Rabatel, A., and Morlighem, M.: Ice velocity and thickness of the world's glaciers, *Nat. Geosci.*, 15, 124–129, <https://doi.org/10.1038/s41561-021-00885-z>, 2022.

Minvielle, M. & Garreaud, R. D. (2011). Projecting Rainfall Changes over the South American Altiplano. *Journal of Climate*, 24, 4577–4583. <https://doi.org/10.1175/JCLI-D-11-00051.1> Minvielle, M. and Garreaud, R. D.: Projecting rainfall changes over the South American Altiplano, *J. Climate*, 24, 4577–4583, <https://doi.org/10.1175/JCLI-D-11-00051.1>, 2011.

Monnier, S., & Kinnard, C. (2013). Internal structure and composition of a rock glacier in the Andes (Upper Choapa Valley, Chile) using borehole information and ground-penetrating radar. *Annals of Glaciology*, 54(64), 61–72. <https://doi.org/10.3189/2013AoG64A107>. Monnier, S. and Kinnard, C.: Internal structure and composition of a rock glacier in the Andes (upper Choapa Valley, Chile) using borehole information and ground-penetrating radar, *Ann. Glaciol.*, 54, 61–72, <https://doi.org/10.3189/2013AoG64A107>, 2013.

Monnier, S., & Kinnard, C. (2015). Reconsidering the glacier to rock glacier transformation problem: New insights from the central Andes of Chile. *Geomorphology*, 238, 47–55. <https://doi.org/10.1016/j.geomorph.2015.02.025> Monnier, S. and Kinnard, C.: Reconsidering the glacier to rock glacier transformation problem: new insights from the central Andes of Chile, *Geomorphology*, 238, 47–55, <https://doi.org/10.1016/j.geomorph.2015.02.025>, 2015.

Monnier, S., & Kinnard, C. (2017). Pluri-decadal (1955–2014) evolution of glacier–rock glacier transitional landforms in the central Andes of Chile (30–33° S). *Earth Surface Dynamics*, 5(3), 493–509. <https://doi.org/10.5194/esurf-5-493-2017>.

Morales, M. S., Christie, D. A., Villalba, R., Argollo, J., Pacajes, J., Silva, J. S., Alvarez, A., Llancabure, J. C. & Soliz Gamboa, C. C. (2012). Precipitation changes in the South American Altiplano since 1300 AD reconstructed by tree rings. *Climate of the past*, N°8, pp. 653–666. Morales, M. S., Christie, D. A., Villalba, R., Argollo, J., Pacajes, J., Silva, J. S., Alvarez, A., Llancabure, J. C., and Soliz Gamboa, C. C.: Precipitation changes in the South American Altiplano since 1300 AD reconstructed by tree rings, *Clim. Past*, 8, 653–666, <https://doi.org/10.5194/cp-8-653-2012>, 2012.

Nicholson, L., Marin, J., Lopez, D., Rabatel, A., Bown, F., rivera, A., (2009). Glacier inventory of the upper Huasco valley, Norte Chico, Chile: glacier characteristics, glacier change and comparison with central Chile. *Ann. Glaciol.* 50 (53), 111–118.

Ohlanders, N., Rodríguez, M., and McPhee, J. (2013). Stable water isotope variation in a Central Andean watershed dominated by glacier and snowmelt, *Hydrol. Earth Syst. Sci.* 17, 1035–1050, <https://doi.org/10.5194/hess-17-1035-2013>, 2013. Ohlanders, N., Rodríguez, M., and McPhee, J.: Stable water isotope variation in a central Andean watershed dominated by glacier and snowmelt, *Hydrol. Earth Syst. Sci.*, 17, 1035–1050, <https://doi.org/10.5194/hess-17-1035-2013>, 2013.

Con formato: Inglés (Estados Unidos)

Palacios, D., Oliva, M., Gómez-Ortiz, A., Andrés, N., Fernández-Fernández, J. M., Schimmelpfennig, I., Léanni, L., & Team, A. (2020). Climate sensitivity and geomorphological response of cirque glaciers from the late glacial to the Holocene, Sierra Nevada, Spain. *Quaternary Science Reviews*, 248, 106617. <https://doi.org/10.1016/j.quascirev.2020.106617>. Palacios, D., Oliva, M., Gómez-Ortiz, A., Andrés, N., Fernández-Fernández, J. M., Schimmelpfennig, I., Léanni, L., and Team, A.: Climate sensitivity and geomorphological response of cirque glaciers from the late glacial to the Holocene, Sierra Nevada, Spain, *Quaternary Sci. Rev.*, 248, 106617, <https://doi.org/10.1016/j.quascirev.2020.106617>, 2020.

Paterson, W.S.B., (1994). *The Physics of Glaciers*. Butterworth-Heinemann, Oxford (480 pp).

Paul, F., Barrand, N. E., Baumann, S., Berthier, E., Bolch, T., Casey, K., et al. (2013). On the accuracy of glacier outlines derived from remote sensing data. *Annals of Glaciology*, 54(63), 171–182. <https://doi.org/10.3189/2013AoG63A296>. Paul, F., Barrand, N. E., Baumann, S., Berthier, E., Bolch, T., Casey, K., Frey, H., Janes, S. P., Krajci, P., Molg, N., Möller, M., Nuth, C., Radić, V., Rastner, P., Raup, B., Schar, C., and Winsvold, S.: On the accuracy of glacier outlines derived from remote sensing data, *Ann. Glaciol.*, 54, 171–182, <https://doi.org/10.3189/2013AoG63A296>, 2013.

Paul, F., Winsvold, S. H., Kääb, A., Nagler, T., & Schwaizer, G. (2016). Glacier remote sensing using Sentinel-2. *Remote Sensing*, 8(7), 575. <https://doi.org/10.3390/rs8070575>. Paul, F., Winsvold, S. H., Kääb, A., Nagler, T., and Schwaizer, G.: Glacier remote sensing using Sentinel-2. Part II: mapping glacier extents and surface facies, and comparison to Landsat 8, *Remote Sens.*, 8, 575, <https://doi.org/10.3390/rs8070575>, 2016.

Pavoni, M., Sirch, F., & Boaga, J. (2021). Electrical and electromagnetic geophysical prospecting for the monitoring of rock glaciers in the Dolomites, Northeast Italy. *Sensors*, 21(4), 1294. <https://doi.org/10.3390/s21041294>. Pavoni, M., Sirch, F., and Boaga, J.: Electrical and electromagnetic geophysical prospecting for the monitoring of rock glaciers in the Dolomites, Northeast Italy, *Sensors*, 21, 1294, <https://doi.org/10.3390/s21041294>, 2021.

Peng, X., Zhang, T., Frauenfeld, O. W., Wang, K., & Mu, C. (2022). An integrated index of cryospheric change in the Northern Hemisphere. *Global and Planetary Change*, 218, 103984. <https://doi.org/10.1016/j.gloplacha.2022.103984>. Peng, X., Zhang, T., Frauenfeld, O. W., Wang, K., and Mu, C.: An integrated index of cryospheric change in the Northern Hemisphere, *Global Planet. Change*, 218, 103984, <https://doi.org/10.1016/j.gloplacha.2022.103984>, 2022.

Peña, H. & Nazarala, B. (1987). Snowmelt-Runoff Simulation Model of a Central Chile Andean Basin with Relevant Orographic Effects. Large Scale Effects of Seasonal Snow Cover. IAHS Publication, 166, 161–172. Peña, H. and Nazarala, B.: Snowmelt-runoff simulation model of a central Chile Andean basin with relevant orographic effects, *IAHS Publ.*, 166, 161–172, 1987.

Pieczonka, T., Bolch, T., Kröhnert, M., Peters, J., & Liu, S. (2018). Glacier branch lines and glacier ice thickness estimation for debris-covered glaciers in the Central Tien Shan. *Journal of Glaciology*, 64(247), 835–849. <https://doi.org/10.1017/jog.2018.75>. Pieczonka, T., Bolch, T., Kröhnert, M., Peters, J., and Liu, S.: Glacier branch lines and glacier ice thickness estimation for debris-covered glaciers in the Central Tien Shan, *J. Glaciol.*, 64, 835–849, <https://doi.org/10.1017/jog.2018.75>, 2018.

Pourrier, J., Jourde, H., Kinnard, C., Gascoin, S., Monnier, S. (2014). Glacier meltwater flow paths and storage in a geomorphologically complex glacial foreland: the case of the Tapado glacier, dry Andes of Chile (30°S). *J Hydrol* 519:1068–1083. <https://doi.org/10.1016/j.jhydrol.2014.08.023>; Pourrier, J., Jourde, H., Kinnard, C., Gascoin, S., and Monnier, S.: Glacier meltwater flow paths and storage in a geomorphologically complex glacial foreland: the case of the Tapado glacier, dry Andes of Chile (30°S), *J. Hydrol.*, 519, 1068–1083, <https://doi.org/10.1016/j.jhydrol.2014.08.023>, 2014.

Radić, V., & Hoek, R (2010). “Regional and global volumes of glaciers derived from statistical upscaling of glacier inventory data.” *Journal of Geophysical Research*, 115, F01010.

Ragetti S. and Pellicciotti F. (2012). Calibration of a physically based, spatially distributed hydrological model in a glacierized basin: on the use of knowledge from glaciometeorological processes to constrain model parameters. *Water Resour Res* 48(W03509):1–20. <https://doi.org/10.1029/2011WR010559>; Ragetti, S. and Pellicciotti, F.: Calibration of a physically based, spatially distributed hydrological model in a glacierized basin: on the use of knowledge from glaciometeorological processes to constrain model parameters, *Water Resour. Res.*, 48, W03509, <https://doi.org/10.1029/2011WR010559>, 2012.

Rahaman, S. A., Ajeez, S. A., Aruchamy, S., & Jegankumar, R (2015). Prioritization of sub-watershed based on morphometric characteristics using fuzzy analytical hierarchy process and geographical information system—A study of Kallar Watershed, Tamil Nadu. *Aquatic Procedia*, 4, 1322–1330. <https://doi.org/10.1016/j.aqpro.2015.02.172>; Rahaman, S. A., Ajeez, S. A., Aruchamy, S., and Jegankumar, R.: Prioritization of sub-watershed based on morphometric characteristics using fuzzy analytical hierarchy process and geographical information system – a study of Kallar Watershed, Tamil Nadu, *Aquat. Procedia*, 4, 1322–1330, <https://doi.org/10.1016/j.aqpro.2015.02.172>, 2015.

Réveillet, M., MacDonell, S., Gascoin, S., Kinnard, C., Lhermitte, S., & Schaffer, N. (2020). Impact of forcing on sublimation simulations for a high mountain catchment in the semiarid Andes. *The Cryosphere*, 14(1), 147–163. <https://doi.org/10.5194/tc-14-147-2020>; Réveillet, M., MacDonell, S., Gascoin, S., Kinnard, C., Lhermitte, S., and Schaffer, N.: Impact of forcing on sublimation simulations for a high mountain catchment in the semiarid Andes, *The Cryosphere*, 14, 147–163, <https://doi.org/10.5194/tc-14-147-2020>, 2020.

RGI Consortium (2017). Randolph Glacier Inventory—A Dataset of Global Glacier Outlines, Version 6.0. Global Land Ice Measurements from Space, Boulder, Colorado, USA. <https://doi.org/10.7265/N5-RGI-60>; RGI Consortium: Randolph Glacier Inventory – a dataset of global glacier outlines, version 6.0, Global Land Ice Measurements from Space, Boulder, Colorado, USA, <https://doi.org/10.7265/N5-RGI-60>, 2017.

Rittger, K., Painter, T. H., & Dozier, J. (2013). Assessment of methods for mapping snow cover from MODIS. *Advances In Water Resources*, 51, 367–380. <https://doi.org/10.1016/j.advwatres.2012.03.002>; Rittger, K., Painter, T. H., and Dozier, J.: Assessment of methods for mapping snow cover from MODIS, *Adv. Water Resour.*, 51, 367–380, <https://doi.org/10.1016/j.advwatres.2012.03.002>, 2013.

Rodríguez M, Ohlanders N, Pellicciotti F, Williams MW, McPhee J. (2016). Estimating runoff from a glacierized catchment using natural tracers in the semi-arid Andes cordillera. *Hydrol Process* 30:3609–3626. <https://doi.org/10.1002/hyp.10973>; Rodríguez, M., Ohlanders, N., Pellicciotti, F., Williams, M. W., and McPhee, J.: Estimating

runoff from a glacierized catchment using natural tracers in the semi-arid Andes cordillera, *Hydrol. Process.*, 30, 3609–3626, <https://doi.org/10.1002/hyp.10973>, 2016.

Rubinstein, R. Y., & Kroese, D. P. (2016). *Simulation and the Monte Carlo Method* (3rd ed.). Wiley. Rubinstein, R. Y. and Kroese, D. P.: *Simulation and the Monte Carlo Method*, 3rd edn., Wiley, Hoboken, NJ, USA, 2016.

Saltelli, A., Ratto, M., Andres, T., Campolongo, F., Cariboni, J., Gatelli, D., Saisana, M., & Tarantola, S. (2008). *Global Sensitivity Analysis: The Primer*. Wiley. Saltelli, A., Ratto, M., Andres, T., Campolongo, F., Cariboni, J., Gatelli, D., Saisana, M., and Tarantola, S.: *Global Sensitivity Analysis: The Primer*, Wiley, Chichester, UK, 2008.

Sanmiguel-Valladolid, A., Morán-Tejeda, E., Alonso-González, E., & López-Moreno, J. I. (2017). Effect of snow on mountain river regimes: an example from the Pyrenees. *Frontiers Of Earth Science*, 11(3), 515–530, <https://doi.org/10.1007/s11707-016-0630-z>.

Saydi, M., and Ding, J. (2020). Impacts of topographic factors on regional snow cover characteristics. *Water Science And Engineering/Water Science And Engineering*, 13(3), 171–180. <https://doi.org/10.1016/j.wse.2020.09.002>. Saydi, M. and Ding, J.: Impacts of topographic factors on regional snow cover characteristics, *Water Sci. Eng.*, 13, 171–180, <https://doi.org/10.1016/j.wse.2020.09.002>, 2020.

Schaffer, N., MacDonell, S., Réveillet, M., Yáñez, E., & Valois, R. (2019). Rock glaciers as a water resource in a changing climate in the semiarid Chilean Andes. *Regional Environmental Change*, 19(5), 1263–1279. <https://doi.org/10.1007/s10113-018-01459-3>. Schaffer, N., MacDonell, S., Réveillet, M., Yáñez, E., and Valois, R.: Rock glaciers as a water resource in a changing climate in the semiarid Chilean Andes, *Reg. Environ. Change*, 19, 1263–1279, <https://doi.org/10.1007/s10113-018-01459-3>, 2019.

Schrott, L. (1996). Some geomorphological-hydrological aspects of rock glaciers in the Andes (San Juan, Argentina). *Zeitschrift für Geomorphologie*, 104, 161–173. Schrott, L.: Some geomorphological-hydrological aspects of rock glaciers in the Andes (San Juan, Argentina), *Z. Geomorphol.*, 104, 161–173, 1996.

Sriyana, I. (2019). Developed watershed classification index determining management priority level based on watershed carrying capacity. *MATEC web of conferences*, 270, 04004. <https://doi.org/10.1051/mateconf/201927004004>. Sriyana, I.: Developed watershed classification index determining management priority level based on watershed carrying capacity. *MATEC Web Conf.*, 270, 04004, <https://doi.org/10.1051/mateconf/201927004004>, 2019.

Suárez, F., Muñoz, J. F., Fernandez, B., Dorsaz, J.-M., Hunter, C. K., Karavitis, C., Gironás, J. (2014). Integrated Water Resource Management and Energy Requirements for water supply in the Copiapo river Basin, Chile. *Water*, 6, pp. 2590–2613. Suárez, F., Muñoz, J. F., Fernandez, B., Dorsaz, J.-M., Hunter, C. K., Karavitis, C., and Gironás, J.: Integrated water resource management and energy requirements for water supply in the Copiapó River Basin, Chile, *Water*, 6, 2590–2613, <https://doi.org/10.3390/w6092590>, 2014.

Trombotto, D. L., Sileo, N., & Dapeña, C. (2020). Periglacial water paths within a rock glacier-dominated catchment in the Stepanek area, Central Andes, Mendoza, Argentina, *Permafrost Periglac.*, 31, 311–323. Trombotto, D. L., Sileo, N., and

Dapeña, C.: Periglacial water paths within a rock glacier-dominated catchment in the Stepanek area, Central Andes, Mendoza, Argentina, *Permafrost Periglac. Process.*, 31, 311–323, <https://doi.org/10.1002/ppp.2043>, 2020.

Valdivieso, S., Hassanzadeh, A., Vázquez-Suñé, E., Custodio, E., Criollo, R. (2022). Spatial distribution of meteorological factors controlling stable isotopes in precipitation in Northern Chile. *Journal of Hydrology*, 605, 127380. Valdivieso, S., Hassanzadeh, A., Vázquez-Suñé, E., Custodio, E., and Criollo, R.: Spatial distribution of meteorological factors controlling stable isotopes in precipitation in northern Chile, *J. Hydrol.*, 605, 127380, <https://doi.org/10.1016/j.jhydrol.2021.127380>, 2022.

Vidal-Pérez, P., Fernández-Sarria, A., González-Bonilla, M. J., Derauw, D., Pérez-Martínez, W., Azócar, G., & Ortega, J. H. (2024). Monitoring surface velocity changes of rock glaciers in the Chilean Andes using DInSAR technique with PAZ imagery. *Procedia Computer Science*. Vidal-Pérez, P., Fernández-Sarria, A., González-Bonilla, M. J., Derauw, D., Pérez-Martínez, W., Azócar, G., and Ortega, J. H.: Monitoring surface velocity changes of rock glaciers in the Chilean Andes using DInSAR technique with PAZ imagery, *Procedia Comput. Sci.*, 239, 456–463, <https://doi.org/10.1016/j.procs.2024.06.070>, 2024.

Villarroel, C. D., Tamburini Beliveau, G., Forte, A. P., Monserrat, O., & Morvillo, M. C. (2018). DInSAR for a Regional Inventory of Active Rock Glaciers in the Dry Andes Mountains of Argentina and Chile with Sentinel-1 Data. *Remote Sensing*, 10(10), 1588. <https://doi.org/10.3390/rs10101588>. Villarroel, C. D., Tamburini Beliveau, G., Forte, A. P., Monserrat, O., and Morvillo, M. C.: DInSAR for a regional inventory of active rock glaciers in the Dry Andes Mountains of Argentina and Chile with Sentinel-1 data, *Remote Sens.*, 10, 1588, <https://doi.org/10.3390/rs10101588>, 2018.

Vivero, S., Bodin, X., Fariás-Barahona, D., MacDonell, S., Schaffer, N., Robson, B. A., & Lambiel, C. (2021). Combination of aerial, satellite, and UAV photogrammetry for quantifying rock glacier kinematics in the Dry Andes of Chile (30° S) since the 1950s. *Frontiers in Remote Sensing*, 2, 784015. <https://doi.org/10.3389/frsen.2021.784015>. Vivero, S., Bodin, X., Fariás-Barahona, D., MacDonell, S., Schaffer, N., Robson, B. A., and Lambiel, C.: Combination of aerial, satellite, and UAV photogrammetry for quantifying rock glacier kinematics in the Dry Andes of Chile (30° S) since the 1950s, *Front. Remote Sens.*, 2, 784015, <https://doi.org/10.3389/frsen.2021.784015>, 2021.

Vuille, M., Francou, B., Wagnon, P., Juen, I., Kaser, G., Mark, B. G., & Bradley, R. S. (2008). Climate change and tropical Andean glaciers: Past, present and future. *Earth-science Reviews*, 89(3-4), 79-96. <https://doi.org/10.1016/j.earscirev.2008.04.002>. Vuille, M., Francou, B., Wagnon, P., Juen, I., Kaser, G., Mark, B. G., and Bradley, R. S.: Climate change and tropical Andean glaciers: past, present and future, *Earth-Sci. Rev.*, 89, 79–96, <https://doi.org/10.1016/j.earscirev.2008.04.002>, 2008.

Whitaker, A., Sugiyama, H., & Hayakawa, K (2008). Effect of Snow Cover Conditions on the Hydrologic Regime: Case Study in a Pluvial-Nival Watershed, Japan. *Journal Of The American Water Resources Association*, 44(4), 814-828. <https://doi.org/10.1111/j.1752-1688.2008.00206.x>.

Wolfe, J. D., Shook, K. R., Spence, C., & Whitfield, C. J (2019). A watershed classification approach that looks beyond hydrology: application to a semi-arid, agricultural region in Canada. *Hydrology And Earth System Sciences*, 23(9), 3945–3967. <https://doi.org/10.5194/hess-23-3945-2019>. Wolfe, J. D., Shook, K. R., Spence, C., and Whitfield, C. J.: A watershed

Con formato: Inglés (Estados Unidos)

classification approach that looks beyond hydrology: application to a semi-arid, agricultural region in Canada, *Hydrol. Earth Syst. Sci.*, 23, 3945–3967, <https://doi.org/10.5194/hess-23-3945-2019>, 2019.

Zech, R., May, J.H., Kull, C., Ilgner, J., Kubik, P.W., Veit, H. (2008) – Timing of the late Quaternary glaciation in the Andes from ~15 to 40°S. *Journal of Quaternary Science*, 23 (6–7), pp. 635–647. Zech, R., May, J. H., Kull, C., Ilgner, J., Kubik, P. W., and Veit, H.: Timing of the late Quaternary glaciation in the Andes from ~15 to 40°S, *J. Quaternary Sci.*, 23, 635–647, <https://doi.org/10.1002/jqs.1200>, 2008.