

1 **Introducing a new normalized cryospheric index (NCI) to categorize** 2 **sub-watersheds on arid environments**

3 Christopher Ulloa¹, Ayon García¹, Anouk Beniest²

4 1LICA-DICTEC/Water and Cryosphere Research Laboratory, University of Atacama, Avenida Copayapu #485, Copiapo,
5 Chile

6 2 Department of Earth Sciences, Vrije Universiteit Amsterdam, De Boelelaan 1081, 1085 HV, Amsterdam, The Netherlands

7 Correspondence to: christopher.ulloa@uda.cl

8 **Abstract.** This study examines sub-watersheds in the arid northern region of Chile (26°41'–28°24'S), situated within the
9 broader Copiapó watershed. The primary water source for this watershed originates from cryospheric landforms. The region
10 exhibits pronounced physiographic and climatic diversity, with its northern sector situated within the South American Arid
11 Diagonal (SAAD), where cryospheric features display greater spatial isolation. The aim of this study is to quantify the water
12 volume contributed by distinct cryospheric landforms to regional watersheds. This study employs a classification methodology
13 to categorize cryospheric landforms within sub-watersheds, integrating an inventory of cryospheric landforms, historical snow
14 permanence derived from satellite imagery, published ice content and depth measurements (ERT and GPR). The Normalized
15 Cryospheric Index (NCI) is calculated for varying hydrological conditions and provides a means to compare potential water
16 volumes across sub-watersheds. The analysis reveals significant spatial variability in cryospheric landforms and their strategic
17 hydrological significance. Under average and low-precipitation conditions, the southern sub-watersheds of the Copiapó river
18 Basin show the greatest water storage potential. The Montosa river (NCI = 0.93), Del Potro river (NCI = 0.89), Manflas river
19 (NCI = 0.47), Estero Come Caballo (NCI = 0.34), and Cachitos river (NCI = 0.31) sub-watersheds have been identified as
20 strategic priority areas within the region for sustaining surface runoff and safeguarding water availability. During high-snowfall
21 periods, northern sub-watersheds in the Copiapo river Basin, such as Estero Come Caballos, show elevated NCI values despite
22 their limited cryospheric landforms. In contrast, the Montosa, Manflas, and Del Potro sub-watersheds contain the most
23 extensive cryospheric landforms and rank among the top of the sub-watersheds with the highest NCI scores.

24 **1 Introduction**

25 Anthropogenic climate change impacts water availability, particularly in regions dependent on high-altitude water sources
26 within mountain systems (Beniston & Stoffel, 2014, Beniston et al., 2018). Changes in the cryosphere that will occur within
27 the next century include a change from solid to liquid precipitation, declining volumes of cryospheric landforms, snow lines
28 moving to higher altitudes, and shorter snow seasons. With these changes, the predictions are that water resources in mountain

29 areas will become increasingly scarce. In the arid zones of the Chilean Atacama Desert glaciers continue to shrink (Masiokas,
30 et al., 2020) and centennial reconstructions of rainfall patterns on the scale of hundreds of years, show a decline in rainfall in
31 the Central Zone and the “Norte Chico” natural regions of Chile (Le Quesne et al., 2006; Minvielle and Garreaud, 2011;
32 Morales et al., 2012; Bozkurt et al., 2017). In addition, the occurrence of mega droughts has intensified since 2010 (Garreaud
33 et al., 2019).

34 Within this climatic context, the Copiapó River exhibits a mixed hydrological regime with very low discharges, typically
35 ranging between $\sim 1.49 \text{ m}^3 \text{ s}^{-1}$ and $1.82 \text{ m}^3 \text{ s}^{-1}$ at the La Puerta gauging station, and strongly modulated by contributions from
36 its main tributaries, the Manflas, Jorquera and Pulido rivers, whose headwaters originate in high-altitude cryospheric
37 environments. Long-term analyses indicate that mean annual flows rarely exceed $2 \text{ m}^3 \text{ s}^{-1}$, and the basin shows recurrent years
38 of negative water balance, leading to progressive aquifer depletion (Suárez et al., 2014). These persistently low flows and the
39 strong dependence on upstream snow and ice melt highlight the structural vulnerability of the runoff of the Copiapó River to
40 changes in the cryosphere.

41 Because of the high altitude of the Andean cryosphere, precipitation is captured from air masses, predominantly in the form
42 of snow. Generally, snow precipitation occurs during the winter (June–September), although the northern parts also receive
43 snow during summer due to the Tropical Monsoon (i.e. Bolivian winter), which takes place from December to February
44 (Valdivieso et al., 2022). In high-altitude Andean environments, snowfall does not directly translate into liquid runoff but may
45 be preserved over different timescales through distinct cryospheric processes. Under favorable mass-balance conditions,
46 repeated snow accumulation and firnification lead to the formation of debris-free glaciers. Where glacier ice becomes mantled
47 by rockfall or talus material, debris-covered glaciers can develop, reducing ablation rates and enhancing ice preservation. In
48 parallel, a significant fraction of snowfall infiltrates coarse debris slopes, where it refreezes and is preserved as ground ice
49 within ice-rich permafrost landforms such as rock glaciers, protalus lobes, and gelifluction taluses. The melting of these
50 cryospheric systems generates a sustained discharge of water, even during snow-free dry seasons. Given the extremely limited
51 rainfall in the Andes, streamflow in the Copiapó River is strongly dependent on snowmelt and cryospheric water storage.
52 (Masiokas et al., 2019). However, the contribution of cryospheric water storage to runoff varies substantially among
53 cryospheric landforms types and occurs over contrasting temporal scales. Debris-free glaciers represent the most direct and
54 dynamic cryospheric contribution to river discharge, responding seasonally to atmospheric forcing and sustaining spring–
55 summer runoff through glacier melt. Debris-covered glaciers may also contribute meltwater, but their hydrological response
56 is attenuated and delayed due to the insulating effect of supraglacial debris, which reduces melt rates per unit area.

57 In contrast, ice-rich permafrost landforms, such as rock glaciers, protalus lobes, and gelifluction taluses, function
58 predominantly as long-term cryospheric water reserves rather than active meltwater sources. Recent studies in the Central
59 Andes indicate that many rock glaciers exhibit near-zero mass loss over recent decades, suggesting that internal ice melt
60 currently contributes only marginally to river runoff (Ferri et al., 2020). Their primary hydrological significance therefore lies
61 in their capacity to store frozen water over decadal to centennial timescales, potentially buffering future water availability
62 under continued climatic warming.

63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93

Seasonal snow represents a distinct cryospheric component with a rapid hydrological response. During wet years, snowmelt dominates river discharge, whereas during dry years runoff becomes increasingly dependent on long-term cryospheric storage at high elevations (Ohlanders et al., 2013). Although snow sublimation losses in the Andes can reach up to ~70% of the seasonal snowpack (Jara et al., 2021), the remaining snowmelt still exerts a first-order control on interannual streamflow variability.

The source of water in the watersheds of the Atacama Desert originate thus predominantly from the volumes of ice that are kept in cryospheric landforms, but the potential volumes of water that are present in these cryospheric landforms are highly variable and depend on both the type and the spatial distribution. We hypothesize that the hydrological relevance of arid sub-watersheds can be quantified by integrating long-term ice storage and seasonal snow permanence into a single normalized index, and that the resulting sub-watershed ranking remains robust under realistic measurement and modelling uncertainty. Testing this hypothesis requires (a) a complete cryospheric inventory at the basin scale, (b) physics-based ice-volume estimates for both glacial and ice-rich permafrost landforms, and (c) explicit propagation of uncertainty through the index calculation. General methodologies to classify and prioritize watersheds are based on morphometric analysis (Rahaman et al., 2015), watershed hydrology (Wolfe, et al., 2019) and ecological and water quality factors. There are also numerous studies covering watershed health in which protection frameworks have been developed based on ecological and hydrological parameters providing approaches on watershed prioritization (Jaiswal et al., 2015; Ahn & Kim, 2017; Sriyana, 2019; Basuki et al., 2022). Few studies incorporate all cryospheric components into watershed management or conservation strategies. Peng et al. (2022) developed an integrated index to evaluate cryospheric changes across the Northern Hemisphere, emphasizing the influence of cryospheric dynamics on regional hydrology. Nonetheless, there is currently a lack of research addressing cryospheric landforms and snow within arid watershed contexts.

To address this gap, this study delivers three integrated outputs for the Copiapó watershed: (i) an updated, GIS-ready inventory of 1862 cryospheric landforms covering glaciers, debris-covered glaciers, rock glaciers, gelifluction taluses, and protalus lobes; (ii) a unified physics-based framework for estimating ice volume across glacial and ice-rich permafrost landforms, combining the Farinotti et al. (2019) consensus product with viscoplastic creep inversion (Cicoira et al., 2020) constrained by Sentinel-1 DInSAR velocities; and (iii) a Normalized Cryospheric Index (NCI) that ranks sub-watersheds by their combined long-term ice storage and seasonal snow contribution, with Monte Carlo propagation of uncertainty to ensure ranking robustness. We apply this framework to the 12 cryospheric sub-watersheds of the Copiapó basin and identify those whose protection is critical for sustaining downstream water availability under continued climatic warming.




Figure 1. Location map of the Copiapó watershed and the different sub-watersheds (Zone 19S). The codes in each sub-watershed represent the official numbering that the General Water Directorate assigns to each sub-watershed.

99 **Table 1. List of sub-watersheds with cryospheric components in the study area.**

Sub-watershed Code	Sub-watershed name	Sub-watershed area km ²
3400	Quebrada Monardes	803.6
3401	Figueroa river	924.2
3402	Estero Come Caballos	880.6
3403	Cachitos river	734.1
3410	Vizcachas de Pulido river	611.9
3411	Ramadillas river	366.8
3412	Del Potro river	470.6
3413	Montosa river	413.2
3420	Manflas river	728.1
3440	Quebrada San Andres	1476.3
3441	Quebrada Paipote	1493.8
3443	Quebrada Martinez	1513.1

100 **2 Regional setting**

101 **2.1 Study Area**

102 The Copiapó watershed is composed of 27 sub-watersheds, distributed along the high mountain belt that forms the headwaters
103 of the basin. A sub-watershed is defined here as a watershed unit with a lower Strahler stream order than the main watershed.
104 In our research area, the Copiapó watershed corresponds to the main basin and contains rivers with a maximum Strahler stream
105 order of nine. Within this hierarchy, sub-watersheds draining from the headwaters typically initiate with Strahler orders of one
106 to three and progress downstream until they merge into larger tributaries with stream orders of up to nine. By distinguishing
107 among sub-watersheds and their associated cryospheric landforms, we can perform both quantitative and qualitative
108 assessments of the current potential water volumes stored within cryospheric environments.
109 To classify a sub-watershed as a cryospheric watershed, we applied objective and reproducible criteria based on its cryospheric
110 components. A sub-watershed is considered cryospheric only when it fulfilled two conditions: (i) the presence of at least one
111 active cryospheric landforms (debris-free glacier, debris-covered glacier, rock glacier, protalus lobe, or gelifluction slope)
112 mapped in the regional inventory of glaciers and the periglacial environment of the Atacama Desert (García et al., 2017); and

(ii) measurable seasonal snow, expressed as long-term snow permanence derived from MODIS fractional snow cover (Rittger et al., 2013). Sub-watersheds fulfilling both requirements (≥ 1 cryospheric landforms + snow permanence) were classified as cryospheric.

Applying these criteria resulted in 12 sub-watersheds being identified as cryospheric watersheds (Fig. 1; Table 1). These sub-watersheds are located entirely within high-elevation sectors of the basin (mean elevation 3,617 m a.s.l.), coincide with areas of high permafrost probability in the global permafrost model of Gruber (2012), and contain 97% of the cryospheric landforms mapped in the Copiapó basin. This classification is therefore geomorphologically justified, reproducible, and consistent with independent permafrost modelling and regional cryospheric inventories.

The Copiapó watershed (Fig. 1) is located in the southern limit of the Atacama Desert, the driest desert of the world (Clarke, 2006), near to the South American Arid Diagonal or SAAD (Zech et al., 2008). The annual precipitation in Copiapó from 1795 to the present has an annual mean of 22.5 mm (Izquierdo et al., 2024). Among the arid basins in northern Chile, the Copiapó watershed is regarded as one of the most vulnerable (Suárez et al., 2014). Because of the limited availability of freshwater and the overexploitation of superficial water and groundwater, the deepening of the water table is accelerating and cryospheric landforms and snow are reducing (Casassa et al., 2007; Vuille et al., 2008). Despite this very limited liquid precipitation in the lower areas, e.g. around the city of Copiapó, there is a flow of water in the rivers of the Copiapó watershed year round. This water comes from the Andean cryosphere that is known to host and store large amounts of fresh water (Bolch and Marchenko, 2006; Bórquez et al., 2006; Azocar and Brenning, 2010; Schaffer et al., 2019).

The glacial and periglacial inventory of the Copiapó watershed indicates a transition from areas that are dominated by glacial environments in the south, to areas that are dominated by periglacial environments in the north (García et al., 2017). In the Copiapó watershed, the occurrence of snow begins above 3,500 m of altitude. In these extreme environments, the water budget is significantly influenced by snow sublimation. From 2001 to 2016, approximately 70% of the snow balance was lost to sublimation, effectively reducing the available water volume (Jara et al., 2021). In semi-arid regions of Chile and Argentina, snow accumulation and subsequent melt processes contribute substantially to surface runoff in numerous Andean river watersheds (Masiokas et al., 2006; Favier et al., 2009).




Figure 2. Main cryospheric landforms. Field photographs from the Copiapó watershed on the left, with a schematic interpretation on the right of. a) Debris-free glacier and debris-glaciers, b) Rock glaciers of the Del Potro river sub-watershed. c) Gelifluction taluses of Montosa river sub-watershed, d) Protalus lobes of the La Laguna sub-watershed in Vizcachas de Pulido watershed.

2.2 Hydrological background of the cryosphere and conceptual framework for ice-volume estimation

The hydrology of arid Andean basins is dominated by cryospheric processes. In the Copiapó watershed, year-round streamflow persists despite minimal precipitation at mid- to low elevations, indicating that meltwater from snow, glaciers, and ice-rich permafrost constitutes the primary source sustaining discharge. Long-term records from the Dirección General de Aguas

(DGA) hydrometric network show a pronounced summer peak in river flow, when rainfall is nearly zero, together with a high baseflow index, confirming a melt-dominated regime and the central hydrological role of the cryosphere in this region. These patterns are consistent with previous studies across the semi-arid Andes (Favier et al., 2009; Gascoin et al., 2011; Ayala et al., 2016), where snow and frozen water bodies act as short- and long-term ice storage modulating runoff during dry years (Table 2).

Cryospheric contributions, however, operate over markedly different response times. Seasonal snow responds at sub-seasonal scales, controlling the onset and magnitude of spring–summer discharge. Debris-free and debris-covered glaciers respond at seasonal to multiannual scales, providing sustained meltwater even in years of limited snowfall. Ice-rich permafrost landforms, including rock glaciers, protalus lobes, and gelifluction slopes, contribute at decadal to centennial scales and represent the most climatically buffered ice storage in the watershed. Capturing these contrasted response times requires not only quantifying the spatial distribution of cryospheric landforms but also applying a coherent conceptual framework for estimating their ice volumes.

The classification of debris-covered glaciers adopted in this study follows Kirkbride (2011), who defines them as glaciers whose ablation zone is largely or entirely covered by supraglacial debris across its full width. This distinction is important in the study area, where debris-covered glaciers occur in close spatial association with rock glaciers but represent fundamentally different cryospheric processes: debris-covered glaciers are dynamic ice bodies in which the supraglacial covered modulates ablation, whereas rock glaciers are permafrost landforms whose movement is driven by internal ice deformation in an ice-rich debris matrix.

Table 2. Contribution to streamflow (%) indicates the proportion of total river discharge supplied by cryospheric meltwater (glaciers and/or seasonal snow), as quantified in each study. Modified from Schaffer et al (2019).

Location / Watershed	Latitude	Hydrological year	Contribution to streamflow (%)	Cryospheric component evaluated	Reference
Huasco watershed	~29°S	2003/04 and 2007/08	3–23	Glaciers (debris-free + debris-covered)	Gascoin et al. (2011)
Elqui watershed	~30°S	2003	4–9	Glaciers (mainly debris-free)	Favier et al. (2009)
Elqui watershed	~30°S	2011	13	Glaciers (debris-free)	Pourrier et al. (2014)
Juncal River watershed	32–36°S	2005/06 and 2008/09	10 and 31	Glaciers (debris-free), snowmelt	Ragetli & Pellicciotti (2012)
Juncal River watershed	32–36°S	2005/06 and 2008/09	16 and 44	Glaciers (debris-free + debris-covered)	Rodríguez et al. (2016)
Juncal River watershed	32–36°S	2011–2012	2–50	Seasonal snow + glacier melt	Rodríguez et al. (2016)
Yeso River watershed	~33.5°S	2014/15	~42 and ~67	Glaciers + snowmelt (during Megadrought)	Ayala et al. (2016)
Central Chilean Andes	–	1968/69	67	Exposed glaciers (exceptionally dry year)	Peña & Nazarala (1987)

Estimating ice volume in mountain cryospheric systems relies on methodological approaches that vary in complexity, physical basis, and data requirements. Traditionally, regional and global assessments have applied empirical area–volume or area–thickness relationships (e.g., Bahr et al., 1997; Jones et al., 2018). These models provide valuable first-order estimates and remain particularly useful in data-sparse regions. However, by construction, they do not explicitly account for glacier geometry, internal deformation, thermal regime, or spatially variable creep rates, which are processes that exert strong control on ice thickness in both glacial and periglacial landforms.

More recent advances have increasingly emphasized physically based approaches that better represent the mechanics of glacier flow and permafrost creep. For debris-free glaciers, inversion techniques such as the model designed by Farinotti et al. (2017) combine mass conservation, surface slope, and Glen’s flow law to derive spatially distributed ice thicknesses (Fig. 3), and form the basis of the global consensus thickness product of Farinotti et al. (2019). At continental scales, Millan et al. (2022) further demonstrated the value of integrating satellite-derived surface velocities into ice-flow inversion, substantially refining glacier thickness estimates and highlighting the importance of kinematic constraints.

In parallel, physically informed modelling has also progressed for ice-rich permafrost landforms. Viscoplastic formulations, such as those proposed by Cicoira et al. (2020), explicitly link surface velocity, basal shear stress, rheology, and thermal conditions to estimate the thickness of ice-rich layers in rock glaciers. Building on these concepts, Hu et al. (2023) applied deformation-based approaches to regional rock-glacier inventories, illustrating the feasibility of estimating ice content in permafrost landforms using remote sensing–derived kinematics. These developments reflect a broader methodological shift toward deformation-based, physically constrained estimates of ice volume across both glacial and periglacial environments.

In the Andes, the applicability of these physically based approaches is reinforced by the growing availability of satellite-derived deformation data. For example, Villarroel et al. (2018) used Sentinel-1 DInSAR to map surface displacement of more than 2,100 rock glaciers in the Dry Andes, while Vidal-Páez et al. (2024) applied PAZ-SAR SBAS-DInSAR to detect millimetric rock-glacier motion in the Chilean Andes. Together, these studies demonstrate that interferometric and feature-tracking techniques enable the detection of low-magnitude deformation, allowing thickness estimation even in low-creep permafrost landforms such as protalus lobes and gelifluction slopes.

In this study, we adopt an integrated conceptual framework that applies:

- (1) the physically based Farinotti model to debris-free glaciers;
- (2) a geometry-consistent approach for debris-covered glaciers using thickness gradients from Farinotti et al. (2019); and
- (3) a unified viscoplastic formulation (Cicoira et al., 2020) to all permafrost cryospheric landforms including rock glaciers, protalus lobes, and gelifluction slopes supported by DInSAR-derived velocities.

198 This unified framework ensures physical consistency across cryospheric landforms with contrasting dynamics and provides
199 the conceptual foundation required for robust estimation of cryospheric ice volumes prior to their integration into the
200 Normalized Cryospheric Index (NCI).

201 **2.3 Glacier and Periglacial Environment**

202 The Copiapó watershed hosts a diverse assemblage of cryospheric landforms (Fig. 2), including debris-free glaciers, debris-
203 covered glaciers, rock glaciers, gelifluction taluses, and protalus lobes, as documented in the regional inventory of García et
204 al. (2017). These landforms exhibit a marked spatial variability across the basin, reflecting strong controls by elevation and
205 topography.

206 Based on the regional inventory of García et al. (2017), a total of 1862 cryospheric landforms have been mapped within the
207 Copiapó watershed: 62 debris-free glaciers, 16 debris-covered glaciers, 173 rock glaciers, 848 gelifluction taluses, and 763
208 protalus lobes. Gelifluction taluses (46%) and protalus lobes (41%) thus dominate the cryospheric assemblage of the basin,
209 while debris-free and debris-covered glaciers together account for less than 4% of the inventoried landforms. This numerical
210 distribution reinforces the periglacial character of the Copiapó cryosphere and underscores the relevance of ice-rich permafrost
211 landforms for ice-volume estimations at the basin scale. This inventory was not originally developed under the Randolph
212 Glacier Inventory (RGI) framework, as it predates the standardization of global glacier datasets in the study region. Instead,
213 the inventory provides a detailed regional mapping of cryospheric landforms, including not only glaciers but also ice-rich
214 permafrost features such as rock glaciers, gelifluction slopes, and protalus lobes. This distinction is particularly relevant
215 because global glacier inventories, such as RGI, are primarily focused on glacier ice bodies and typically do not include
216 periglacial landforms, which are rarely mapped in a systematic manner. As a result, the regional inventory used here captures
217 a broader spectrum of cryospheric components that are especially significant in arid mountain environments, where non-glacial
218 frozen water storage may play an important hydrological role.

219 In general, cryospheric landforms are distributed discontinuously along the high Andes, with notable differences between
220 northern and southern sectors. In areas such as Nevado Jotabeche and Los Tronquitos, landforms are predominantly located
221 on high-elevation ridges and peaks, whereas in the Del Potro region they are more commonly confined to valley bottoms
222 between steep slopes.

223 Gelifluction taluses represent one of the most widespread landforms in the study area and are typically associated with high-
224 elevation environments. In the southern sector, they frequently occur in close proximity to debris-free glaciers, forming
225 coupled glacial–periglacial systems. In contrast, in the northern sector, gelifluction features tend to appear as smaller and more
226 isolated landforms.


227 Rock glaciers are mainly distributed along steep valley flanks at intermediate to high elevations, often occupying topographic
228 depressions or confined valley settings. Debris-free glaciers are generally restricted to the highest elevations and, particularly
229 in the southern sector, are commonly associated with adjacent periglacial landforms.

230 Protalus lobes are especially abundant in some northern areas, where they develop on slopes of varying gradients and exhibit
231 a range of sizes and morphologies. Debris-covered glaciers are comparatively scarce and are typically found in the southern
232 sector, occurring as small bodies located downslope of debris-free glaciers.

233 Overall, the distribution of cryospheric landforms in the Copiapó watershed reflects a transition from glacial-dominated
234 systems in the south to predominantly periglacial environments in the north, consistent with regional climatic and topographic
235 gradients (Fig. 4).

236 The regional inventory of García et al. (2017) predates the RGIK (Rock Glacier Inventories and Kinematics; IPA Action
237 Group) guidelines, but its mapping workflow multi-source satellite photo-interpretation supported by field reconnaissance,
238 identification of active rock glaciers from diagnostic geomorphological criteria (frontal ramp, furrows and ridges), and a
239 GLIMS-compatible attribute database is methodologically consistent with the principles later codified by RGIK. The inventory
240 extends this scope to include gelifluction taluses and protalus lobes, two ice-rich permafrost landforms that dominate the arid
241 Atacama cryosphere but fall outside the RGIK rock-glacier framework.

Debris-free glacier thickness (Farinotti et al., 2019)






Figure 4. a) Distribution map of the cryospheric landforms inventoried in the Atacama region from García et al. (2017). b) Jotabeche volcano cryospheric landforms distribution c) The Potro hill cryospheric landforms distribution d) The Tronquitos hill cryospheric landforms distribution. Basemap: Esri, Maxar, Earthstar Geographics, and the GIS User Community.

253 **3 Methodology**

254 **3.1 Survey of cryospheric landforms**

255 To quantify the ice volume in the Copiapó watershed, we used existing cryospheric inventories without performing new
256 delineations. The regional inventory of glaciers and periglacial landforms of the Atacama Desert (García et al., 2017) was
257 employed directly to extract polygons of debris-covered glaciers, rock glaciers, protalus lobes, and gelifluction slopes. This
258 inventory represents the only comprehensive and validated dataset that explicitly identifies periglacial cryospheric landforms
259 in northern Chile, and its polygons were incorporated without modification into our analysis.

260 Debris-free glaciers were obtained exclusively from the Randolph Glacier Inventory (RGI) v6.0 (RGI Consortium, 2017). For
261 these glaciers, we did not compute ice thickness ourselves. Instead, we used the pre-computed global ice-thickness product of
262 Farinotti et al. (2019), which is physics-based and already calibrated on RGI glacier outlines (Fig. 3). Because the Farinotti
263 model requires RGI geometry as input, the use of RGI v6.0 ensures direct compatibility with the global thickness dataset. Ice
264 volumes for debris-free glaciers were therefore calculated solely by spatially integrating the published Farinotti et al. (2019)
265 thickness values over the RGI glacier polygons within the Copiapó basin. This approach avoids any reinterpretation of glacier
266 geometry or re-estimation of ice thickness, ensuring methodological transparency and consistency with the global consensus
267 product.

268 The joint use of the inventory developed by García et al. (2017) and RGI v6.0 glacier inventory is thus complementary. RGI
269 provides standardized glacier boundaries and their associated Farinotti-derived thickness fields for debris-free glaciers,
270 whereas García et al. (2017) provides the spatially explicit identification of debris-covered glaciers and periglacial landforms
271 that are not represented in global glacier datasets. By relying directly on these inventories and on the published model outputs
272 of Farinotti et al. (2019), our workflow remains fully reproducible and avoids introducing new mapping or thickness-estimation
273 steps.

274 Mapping uncertainty for all polygons was propagated using the one-pixel boundary buffer method of Paul et al. (2013),
275 ensuring consistent treatment of spatial uncertainty across glacial and periglacial cryospheric landforms.

276 **3.2 Survey of snow cover: MODIS-derived snow permanence**

277 The MOD10A1 Fractional Snow Cover (FSC) product from NASA was used to characterize snow based on long-term patterns
278 of snow permanence. Each MOD10A1 image provides, for every 500×500 m pixel, a value representing the fractional snow
279 cover within that pixel (from 0 to 1), derived through a spectral unmixing algorithm (Rittger et al., 2013). This sub-pixel
280 representation allows a more accurate depiction of discontinuous or patchy snow, which is common in high-elevation arid
281 regions. Instead of relying on isolated scenes, we processed the complete daily MODIS time series (2000–2022) to obtain both
282 the instantaneous snow-covered area and the frequency of snow permanence at the sub-watershed scale. Although cloud
283 contamination can introduce occasional misclassifications, the product maintains approximately 90% accuracy under ideal
284 conditions (Hall et al., 2006; Jiang et al., 2024), and is suitable for long-term characterization of snow permanence.

285 From the daily FSC series, each MODIS image provides the fractional snow cover for every 500×500 m pixel. Using this
286 long-term record, fractional snow cover (FSC) was used as the core variable to represent the proportion of snow within each
287 pixel in each scene. In addition, we calculated snow permanence as the fraction of days during the 23-year period in which
288 each pixel exhibited snow-covered conditions according to its FSC value. Snow permanence is therefore not an independent
289 MODIS layer, but a temporal aggregation derived directly from the daily FSC maps, summarizing the frequency and recurrence
290 of snow presence across the basin.

291 In this framework, FSC was used as the primary input for constructing daily snow permanence maps, and snow permanence,
292 computed as the proportion of snow-covered days over the full time series, was incorporated into the Normalized Cryospheric
293 Index (NCI) as the indicator of short-term cryospheric water resources. This approach ensures that the characterization of snow
294 permanence reflects both their spatial extent and their temporal recurrence.
295

296 **3.3 Field work for ground truthing**

297 Since the mapping workflow is primarily based on remote sensing, satellite imagery, and existing cryospheric inventories, we
298 incorporated targeted field observations and independent geophysical datasets to validate both the identification of cryospheric
299 landforms and the ice-thickness values used in this study. Fieldwork was carried out in the Montosa (03413) and Del Potro
300 (03412) sub-watersheds, which together represent approximately 10% of the Copiapó basin and contain accessible examples
301 of the main cryospheric landforms present in the region.

302 During these campaigns, photographic and geomorphological observations were conducted to verify the surface expression of
303 mapped cryospheric landforms, including gelifluction slopes, protalus lobes, and rock glaciers. These observations allowed us
304 to assess the representativeness of the polygons obtained from existing inventories.

305 Beyond surface verification, independent geophysical datasets were used to constrain and validate ice-thickness estimates. For
306 debris-free glaciers, ice thickness values from the Farinotti et al. (2019) global product were compared against airborne ground-
307 penetrating radar (GPR) profiles collected by the Dirección General de Aguas (DGA, 2014) over the Del Potro and Tronquitos
308 glaciers.

309 For ice-rich permafrost landforms, Electrical Resistivity Tomography (ERT) measurements were conducted on a representative
310 gelifluction slope and on an active rock glacier within the Montosa sub-watershed. These ERT profiles constrain the depth and
311 continuity of ice-rich layers and support the thickness ranges adopted for periglacial cryospheric landforms in the modelling
312 framework.

313 Together, these field observations and geophysical datasets provide independent validation for both the spatial delineation and
314 the subsurface characteristics of cryospheric landforms within the Copiapó watershed, strengthening the robustness of the ice-
315 volume estimates applied in this study.

316 **3.3.1 Geophysical survey on gelifluction slopes and rock glaciers**

317 Electrical Resistivity Tomography (ERT) was employed to characterize subsurface ice-rich layers in selected periglacial
 318 landforms and to provide observational constraints for the ice-thickness values used in the modelling framework. ERT is a
 319 multi-electrode resistivity method that generates 2-D vertical sections of the subsurface by injecting electrical current into the
 320 ground and measuring the resulting potential differences across electrode arrays. The field configuration used in this study
 321 (Table 3) consisted of 36 electrodes operated with Wenner, Schlumberger, and Dipole–Dipole arrays.



322
 323 **Figure 5. Photography of the fieldwork. a) And b) the Lobulated morphology of studied gelifluction taluses, which are not covered**
 324 **in snow; c) and d) the geoelectric line cable; e) and f) the used equipment, differential GPS and resistivity meter ARES II measuring**
 325 **a resistivity point diagram distribution.**

326 Fieldwork focused on two representative cryospheric landforms within the Montosa sub-watershed: a gelifluction slope and
 327 an active rock glacier (Fig. 5 and Fig. 6). These sites were selected because they exhibit clear geomorphological indicators of

ice-rich permafrost and provide suitable access for geophysical surveys. The ERT profiles allowed direct identification of high-resistivity zones ($>20\text{--}50\text{ k}\Omega\cdot\text{m}$), characteristic of massive or highly saturated ice-rich permafrost bodies.

Table 3 Used geophysical equipment for fieldwork of this study

Equipment	Amount
Resistivity meter: ARES II	1
30 electrode box	5
10 channels multicable II (60 m)	10
Car Batteries	2
Differential GPS	1

The survey layout required a 600-m profile organized through multicables arranged at 90° angles. A high-precision differential GPS (“South Galaxy”) was used to set the start and end points of each line, ensuring alignment and minimizing installation error. Electrode spacing was fixed at 5 m, with position control verified using a combination of GPS and measured markers at 30-m intervals. All electrodes were tested prior to each acquisition, and any with unacceptable standard deviation values were repositioned.

These ERT measurements provide direct ground validation of subsurface ice for both gelifluction taluses and rock glaciers, supporting the thickness ranges applied to ice-rich permafrost cryospheric landforms in our volume estimations and complementing the GPR-based glacier-thickness validation described in Section 3.3.



Figure 6. Photography of the fieldwork rock-glacier surface. (a) Intelligent multi-electrode cable reel used for deploying the ERT array across the periglacial terrain. (b) Installation of the 600-m geoelectric line. (c) Operation of the ARES II resistivity meter during data acquisition.

Before every measurement, all electrodes were tested to detect electrodes with high standard deviation in resistivity. If the deviation exceeded acceptable limits, the electrodes were repositioned until the deviation fell within an acceptable range to ensure measurement accuracy.

3.4 DInSAR kinematic analysis for periglacial cryospheric landforms

To characterize surface deformation across glacial and periglacial cryospheric landforms and to provide kinematic constraints for ice-thickness modelling, we processed Sentinel-1 IW Single Look Complex (SLC) images using the ESA SNAP v9.0 toolbox. Interferometric pairs were selected based on perpendicular baselines <150 m and temporal baselines <24 days to ensure coherence in high-altitude terrain (Table 4).

Table 4 Sentinel-1 interferometric image pairs and associated temporal and perpendicular baselines tested.

Reference Image Date	Secondary Image Date	Orbit Type	Temporal Baseline (Days)	Perpendicular Baseline	Interferogram status
01-03-2024	13-03-2024	Ascending	12	55.35	Inconsistent
25-03-2024	06-04-2024	Ascending	12	40.12	Inconsistent
03-03-2016	27-03-2016	Descending	24	9.39	Coherent
05-01-2015	29-05-2015	Ascending	24	-121.04	Inconsistent
29-01-2015	22-02-2015	Descending	24	25.41	Inconsistent

SLC scenes were co-registered using the TOPS burst co-registration workflow. Interferograms were generated, filtered using a Goldstein filter ($\alpha = 0.8$), and corrected for topographic phase using the 30-m SRTM DEM. Phase unwrapping was performed with SNAPHU in external mode. Line-of-sight (LOS) displacement was obtained from the unwrapped phase:

$$d_{LOS} = \frac{\lambda}{4\pi} \Delta\phi \tag{1}$$

Mean surface velocities were extracted for each cryospheric landforms polygon (debris-free glaciers, debris-covered glaciers, rock glaciers, protalus lobes, gelifluction slopes). These DInSAR-derived velocities were incorporated as dynamic constraints

for the viscoplastic modelling of permafrost cryospheric landforms and as supplementary kinematic information in glacier-thickness interpretation.

3.4.1 Velocity uncertainty

Uncertainty in DInSAR-derived surface velocity was incorporated by explicitly estimating the phase noise associated with interferometric coherence and temporal baseline. Following Hanssen (2001), the standard deviation of the interferometric phase for a multilooked interferogram is:

$$\sigma_{\phi} \approx \sqrt{\frac{1-\gamma^2}{\gamma^2 L}} \quad (2)$$

Where γ is the interferometric coherence and L the number of equivalent looks. For Sentinel-1 SLC data used in this study, the typical modeled coherence is $\gamma=0.97$, the temporal baseline is $\Delta t=24$ days, and the radar wavelength is $\lambda=0.0555$ m.

Phase noise is converted to line-of-sight displacement noise using:

$$\sigma_{\phi} = \frac{\lambda}{4\pi} \sigma_{\phi} \quad (3)$$

And subsequently velocity uncertainty:

$$\sigma_v = \frac{\sigma_d}{\Delta t} = \frac{\lambda}{4\pi \Delta t} \sqrt{\frac{1-\gamma^2}{\gamma^2 L}} \quad (4)$$

For a representative multilooking a factor of $L=10$ was applied. The resulting uncertainties are:

$$\begin{aligned} \sigma_{\phi} &\approx 0.056 \text{ rad}, \quad \sigma_d \approx 2.5 \times 10^{-4} \text{ m} = 0.25 \text{ mm}, \\ \sigma_d &\approx 1.0 \times 10^{-5} \text{ m d}^{-1} = 0.01 \text{ mm d}^{-1} \approx 3 - 4 \text{ mm a}^{-1}. \end{aligned}$$

This value ($\sigma_d = 0.01 \text{ mm d}^{-1}$) was thus adopted as the 1σ uncertainty of DInSAR velocity measurements.

3.5 Calculation of the ice volume and uncertainty

To quantify the total ice volume stored in glacial and periglacial cryospheric landforms, we applied cryospheric landforms-specific physics-based models according to their mechanics, morphology, and available input data (Table 5). The method applied and associated uncertainties are detailed below.

3.5.1 Debris-free glaciers

Ice thickness for debris-free glaciers was obtained from the global consensus ice-thickness dataset of Farinotti et al. (2019), which integrates multiple physics-based inversion models (Farinotti et al., 2017). Glacier volumes were computed by spatially

401 integrating the consensus ice-thickness rasters over the RGI v6.0 glacier polygons within the Copiapó watershed. A relative
402 uncertainty of $\pm 26\%$ was applied following the uncertainty assessment reported in Farinotti et al. (2017, 2019).

403 3.5.2 Debris-covered glaciers

404 Mean ice thickness for debris-covered glaciers was derived by sampling the Farinotti et al. (2019) consensus ice-thickness
405 field in the clean-ice frontal zones immediately adjacent to each debris-covered area. These representative values were then
406 assigned to the corresponding debris-covered glacier polygons, ensuring consistency with the local glacier geometry and
407 thickness gradients. Ice volumes were calculated by multiplying the assigned mean thickness by the mapped debris-covered
408 glacier area. Due to the inherently greater spatial variability and additional uncertainties associated with debris cover (e.g.,
409 variations in debris thickness, thermal properties, and remote-sensing-derived thickness estimates), published studies of
410 debris-covered glacier thickness consistently report substantially larger differences between modeled and *in situ* measurements
411 compared to debris-free glaciers (e.g. $\sim 40\%$ uncertainty for certain debris-covered glaciers; see Pieczonka et al., 2018). Hence,
412 a conservative uncertainty of $\pm 35\%$ was adopted for debris-covered glacier volume estimates to account for these factors.

413 3.5.3 Rock glaciers

414 Rock-glacier thickness was estimated using a viscoplastic creep inversion following Cicoira et al. (2020), where basal shear
415 stress is constrained by observed surface velocity. For each rock glacier, mean slope-parallel surface velocities (v) and their
416 standard deviation (σ_v) were derived from Sentinel-1 DInSAR measurements (Sect. 3.4.1) and expressed in m yr^{-1} . The basal
417 shear stress τ was inverted using the 1-D viscoplastic formulation with Glen-type exponent $n = 3$ and a temperature-dependent
418 rate factor (T).

419
420 To explicitly represent uncertainty in the thermal state of ice-rich permafrost in the Central Andes, we treated ice temperature
421 as a normal distribution:

$$423 \quad T \sim N(-3^\circ\text{C}, 3^\circ\text{C}) \quad (5)$$

424
425 This choice is consistent with reported near-isothermal to moderately cold permafrost conditions in the Central Andes,
426 including rock-glacier settings, and with the strong temperature sensitivity of creep close to 0°C described in constitutive
427 studies (e.g. Koenig et al., 2025, Monnier & Kinnard, 2013).

428
429 Using τ , mean thickness was computed as:

$$430 \quad H = \frac{\tau}{\rho \cdot g \cdot \sin(\alpha)} \quad (6)$$

432 where α is the mean surface slope, ρ is ice density, and g is gravitational acceleration. Total landform volume was computed
433 as:

$$434 \quad V = A \cdot H \quad (7)$$

435 where A is the mapped planimetric area.

436 Ice volume was obtained by applying an ice-content fraction C_I to the landform volume:

$$437 \quad V_{ice} = V \times C_I \quad (8)$$

438 For rock glaciers, C_I was represented as a bounded range consistent with reported values for these type of ice-rich permafrost
439 landforms (Cicoira et al., 2020):

$$440 \quad C_I = 0.5 \pm 0.10 \quad (9)$$

441 Ice mass (in Gt) was then computed as:

$$442 \quad M_{Gt} = \frac{V_{ice} \rho}{10^{12}} \quad (10)$$

443 with $\rho = 900 \pm 20 \text{ kg m}^{-3}$. Uncertainties in H and M_{Gt} were propagated from σ_v , σ_T , slope uncertainty, mapping
444 uncertainty in area, ice-content uncertainty, and density uncertainty using quadratic propagation of relative errors (Sect. 3.5.5).

445 **3.5.4 Protalus lobes and gelifluction slopes**

446 Protalus lobes and gelifluction taluses were modelled using the same viscoplastic inversion framework originally developed
447 for rock glaciers (Cicoira et al., 2020), exploiting low but measurable DInSAR-derived surface deformation to constrain basal
448 shear stress and thickness. For each mapped polygon, mean slope-parallel surface velocity and its associated uncertainty (v ,
449 σ_v) were used to invert basal shear stress and compute thickness H through Eq. (6). Thermal uncertainty was represented using
450 the same Central Andes temperature prior described in Eq. (5).

451 However, in contrast to rock glaciers, protalus lobes and gelifluction taluses are characterized by lower ice contents, higher
452 debris fractions, and a dominance of seasonal or shallow permafrost deformation mechanisms. To account for these rheological
453 differences, we modified the Glen-type flow exponent n used in the inversion. While rock glaciers were modelled assuming n
454 $= 3$, consistent with creep-dominated ice–debris mixtures (Cicoira et al., 2020), protalus lobes and gelifluction taluses were
455 modelled using a reduced exponent ($n = 2$), reflecting more linear, frictionally controlled deformation regimes typical of ice-
456 poor frozen debris slopes. This adjustment reduces the sensitivity of inferred thickness to surface velocity and avoids
457 unrealistically thick solutions for low-velocity landforms.

Because ice contents in these landforms are generally lower and more heterogeneous than in rock glaciers, ice-content fraction was prescribed following Eq. (11) as a bounded prior range rather than a fixed value. This range represents plausible ground-ice fractions for ice-rich permafrost slopes in the Central Andes and is supported by field-calibrated ERT observations showing strong spatial variability in ground-ice content within Andean permafrost landforms (Hilbich et al., 2022), as well as early empirical constraints for frozen debris slopes in arid mountain environments (Schrott, 1996). The ice-content prior was therefore implemented to represent structural uncertainty rather than a deterministic parameter.

$$C_I = 0.37 \pm 0.12 \quad (11)$$

Following thickness estimation, landform volume was computed using Eq. (7), ice volume using Eq. (8), and ice mass using Eq. (10).

3.5.5 Uncertainty propagation framework for ice-rich permafrost cryospheric landforms

For rock glaciers, gelifluction taluses, and protalus lobes, uncertainties in ice thickness and ice mass were propagated deterministically using quadratic propagation of relative errors for multiplicative quantities. First, the uncertainty in mapped area was estimated following Paul et al. (2015), assuming boundary errors occur within ± 0.5 pixels and using:

$$u_{A_i} = \frac{1}{2} P_i \times \lambda \times \sigma \quad (12)$$

where P_i is the polygon perimeter, λ is the spatial resolution of the imagery used for delineation, and $\sigma \approx 0.6872$ is a correction factor corresponding to the standard deviation of a Gaussian distribution with most errors confined within ± 0.5 pixels.

Ice volume was computed using the Eq. 7, and ice mass as $M_{Gt} \propto A \times H \times C_I \times \rho$. Therefore, the relative uncertainty in ice mass was calculated as:

$$\left(\frac{\sigma_M}{M}\right)^2 = \left(\frac{\sigma_A}{A}\right)^2 + \left(\frac{\sigma_H}{H}\right)^2 + \left(\frac{\sigma_{C_I}}{C_I}\right)^2 + \left(\frac{\sigma_\rho}{\rho}\right)^2 \quad (13)$$

and the absolute uncertainty as $\sigma_M = M \times \left(\frac{\sigma_M}{M}\right)$. Basin totals were obtained by summing masses across polygons, while total uncertainty was aggregated assuming independence using:

$$\sigma_{tot} = \sqrt{\sum_i^n \sigma_{M_i}^2} \quad (14)$$

487 This framework ensures consistent uncertainty propagation across all ice-rich permafrost cryospheric landforms while
488 preserving the physically based dependence of thickness on surface velocity and temperature.

489

490 **Table 5. Key references used as input for the calculation of ice volumes of cryospheric landforms.**

491

492

493

494

495

496

497

498

499

500

501

502

503

504

505

506

Cryospheric landforms	Method	Ice content	Key references
Debris-free glaciers	Physics-based inversion (Farinotti et al., 2019)	100%	Farinotti et al. (2017, 2019)
Debris-covered glaciers	Thickness assignment from adjacent clean-ice zones (Farinotti 2019)	45–99%	Farinotti et al. (2017, 2019)
Rock glaciers	Viscoplastic model (Cicoira et al., 2020) + DInSAR	40–60%	Cicoira et al. (2020)
Gelifluction taluses and Protalus lobes	Adapted viscoplastic model + DInSAR	25–49%	Hilbich et al. (2022), Schrott, (1996)

507 **3.6 Cryospheric landforms and snow index and data integration**

508 To develop a classification system that categorizes the capacity to store ice as a strategic water reserve in desert areas of sub-
509 watersheds, we followed a workflow that consists of four phases (Fig. 7). We developed the classification system using
510 cryospheric landforms (expressed in gigatons (Gt), equivalent to 1,000 million metric tons of ice) and snow (quantified in km²
511 of snow-cover extent), integrating both into the methodological framework. The strategic Normalized Cryospheric Index (NCI)
512 captures the relative significance of each sub-watershed, considering both its ability to store water as ice (cryospheric
513 landforms) and its potential to regenerate water resources through snowfall. The NCI needs to be adapted to each specific case.
514 For the Copiapó watershed, the active cryospheric landforms present until 2024 have been used. Those cryospheric landforms
515 include debris-free glaciers, debris-covered glaciers, rock glaciers, gelifluction talus and protalus lobes. Depending on the
516 types of cryospheric landforms available in a specific region, the number of cryospheric landforms included may vary.
517 The methodological framework for computing the NCI index integrates Monte Carlo simulation with a robustness assessment
518 of a composite indicator that incorporates ice volume (V) and snow permanence (S), adhering to the uncertainty propagation

principles in simulation-based methodologies outlined by Rubinstein and Kroese (2016). First, each watershed has a mean value (e.g., $\bar{V}_i$ y $\bar{S}_i$) and an estimate of its dispersion (ΔV_i and ΔS_i), usually understood as standard deviation. On the basis that both V and S follow normal distributions $N(\bar{V}_i, \sigma_v)$ y $N(\bar{S}_i, \sigma_s)$, N replicates of each variable are generated for each watershed, thus obtaining “clouds of values” representing the uncertainty of each datum. Subsequently, a normalization is performed (using min-max) in order to avoid that one variable dominates the other only by differences in magnitude.

Once the simulated values are generated, a weighted index is defined whose objective is to combine both variables (Eq. 15):

$$NCI_{i,k}(\omega) = \omega \cdot V_{i,k} + (1 - \omega) \cdot S_{i,k} \quad (15)$$

Where $\omega \in [0,1]$

Where ω is a parameter that determines the relative relevance of ice volume versus snow permanence. For each sample k (among the N simulations), the values of that index in all the watersheds are obtained and ordered from highest to lowest, giving rise to a specific ranking of each run. In this way, each watershed “i” gets a “position” (1 for the watershed with the highest index, 2 for the next one and so on).

The optimization function (Eq. 16) relies on the stability of the ranking under uncertainty, in accordance with the sensitivity analysis framework outlined by Saltelli et al. (2008). Specifically, for each watershed, the standard deviation of its position over the N runs is measured, and this measure of dispersion is summed (or averaged) for all watersheds. A smaller value of this sum indicates greater stability or robustness in the ranking. Formally, if σ_i denotes the standard deviation of the position of watershed under a weight ω , the objective function results:

$$f(\omega) = \sum_{(i = 1 \text{ to } n)} [\sigma_i \cdot (\omega)] \quad (16)$$

Where the idea is to minimize the value of this function. To determine the “optimal” ω , a sweep over [0.1] is performed. The optimal result corresponds to the ω value that maximizes the stability of the watershed ranking system, ensuring minimal sensitivity to errors or noise in the V and S datasets.

Finally, plots are generated to analyze the resulting average ranking and the probability that certain watersheds occupy the first place in various simulations. In this way, a picture is obtained of how uncertainty in ice volume and snow permanence affects the prioritization of watersheds.

Although the NCI has a simple formulation, the variables it combines are affected by substantial uncertainty. Ice volumes integrate multiple sources of error related to thickness modelling, ice content, density, and mapped area, while snow is represented by long-term snow permanence derived from MODIS, which captures both the spatial extent and temporal

recurrence of snow. Monte Carlo simulation is therefore applied not to increase index complexity, but to propagate uncertainty in ice volume and snow permanence into watershed ranking and to identify an objective weighting factor (ω) that maximizes ranking robustness. This stochastic framework avoids subjective parameterization and ensures a stable, transparent, and reproducible classification of cryospheric sub-watersheds.





Figure 7. Flow chart of the methodology used to calculate the NCI.

558 **4 Results**

559 **4.1 Snow permanence results**

560 By integrating all MODIS fractional snow cover images and computing their long-term mean over the period 2000–2022, we
561 derived a snow permanence metric, defined as the probability of snow presence at each pixel. The resulting raster represents
562 the spatial distribution of snow permanence across the Copiapó watershed (Fig. 8). At the regional scale, snow permanence
563 shows limited spatial variability at high elevations, remaining consistently high above approximately 5,500 m a.s.l. These
564 regions are located close to the Argentinean border. In the central area between Caserones Hill and Cadillal Hill, no cryospheric
565 landforms were identified, which coincides with the lowest snow permanence values in the study area, with snow present only
566 around 30% of the time. In contrast, the highest snow permanence values are observed in the southern sector of the study area.
567 In the central area of the Copiapó watershed the snow permanence is concentrated around the Nevado Jotabeche Hill area. The
568 northern part of the watershed does not show concentrated major snowfall anomalies. Snow permanence has persisted
569 approximately 30–50% of the time exclusively at high-elevation zones (>4,500 meters above sea level) near the Argentinian
570 border, contrasting sharply with lower altitudes starting at 2,000 m a.s.l.

571 The calculated probability of the occurrence for the investigated watersheds shows that the snow permanence watershed varies,
572 with higher snow permanence for the southern watersheds compared to the central and northern watersheds. The watershed
573 with the highest resource index is the Montosa river watershed (03413) with 0.18% probability of being covered in snow. The
574 Vizcachas de Pulido river (03410), which is located at the lowest altitude, has the lowest probability of 0.07% to have snow
575 permanence.




4.2 Analysis of runoff

Before incorporating snow permanence as a core variable of the NCI, we tested empirically whether MODIS-derived snow cover translates into a hydrological signal at the watershed scale. To do this, we contrasted streamflow behaviour during representative wet and dry years and quantified the statistical relationship between September snow extent and Oct–Mar mean

584 discharge at the Copiapó River at Pastillo gauging station. Daily mean streamflow records at the Copiapó River at Pastillo
 585 station (Fig. 9) reveal distinct seasonal patterns between wet and dry years. During the wet year (Oct 2017 - Mar 2018),
 586 streamflow exhibits a sustained increase from October to December, peaking at 5.7 m³/s in December, followed by a gradual
 587 decline through January and a secondary pulse in February (5.3 m³/s). In contrast, the dry year (Oct 2013 - Mar 2014) shows
 588 substantially lower spring flows, with mean streamflow averaging only 1.0 m³/s during October-November, and the most
 589 significant increases concentrated in December (1.3 m³/s) and February (2.2 m³/s). Overall, the wet year delivered 3.4 times
 590 more water than the dry year, with the greatest relative differences occurring in November (+471%) and December (+343%).

591



592

593 **Figure 9. Daily (a) and monthly mean (b) streamflow at Copiapó River at Pastillo for wet (2017-2018, blue) and dry (2013-2014, red)**
 594 **years. Data Source: Dirección General de Aguas, Chile.**

595 These contrasting patterns between individual years reflect broader interannual variability in snowpack accumulation, ice
 596 bodies and melt dynamics. To quantify this relationship at the watershed scale, we incorporated a new analysis linking MODIS-
 597 derived snow permanence in September, at the onset of the melt season, with mean river discharge observed between October
 598 and March (Fig. 10). This relationship shows a robust correlation ($R^2 = 0.58$; $r = 0.76$; $N = 22$), confirming that seasonal snow
 599 permanence acts as a dominant short-term cryospheric water resource in the Copiapó watershed. However, the unexplained





Figure 10. September snow-covered area vs. October-March mean discharge at Copiapó River at Pastillo (2000-2021, N = 22). Linear regression (red line) with 95% confidence interval (shaded). $R^2 = 0.58$; $r = 0.76$; $p < 0.001$. Sources: MODIS MOD10A1 and DGA, Chile.

4.3 Field survey observations of cryospheric landforms.

The fieldwork has yielded observations of cryospheric landforms in the Del Potro and Montosa river sub-watersheds (Fig. 11 and 12). The Del Potro Glacier (Fig. 12d) and the surrounding periglacial environment of the Del Potro River sub-watershed (03412; Fig. 11) were clearly identified during field surveys. In the Montosa River sub-watershed, the Maranceles Glacier (Fig. 12c) was also observed. Additional field observations confirmed the presence of well-developed rock glaciers, gelifluction taluses, and protalus lobes in the Del Potro River basin (03412) (Fig. 11c–g). Glacial landforms are preferentially distributed on south-facing slopes, as indicated by their aspect, which are generally associated with reduced incoming solar radiation. In contrast, periglacial cryospheric landforms such as protalus lobes and gelifluction taluses are more frequently found on exposed slopes with steeper gradients and variable aspects, reflecting

620 differences in topographic setting rather than direct measurements of snow accumulation or thermal conditions. Debris-free
 621 and debris-covered glaciers are primarily located in topographic depressions and high-elevation headwaters. In contrast,
 622 protalus lobes and gelifluction taluses are predominantly distributed on steep slopes and debris-covered sectors. Overall,
 623 periglacial cryospheric landforms display a wider spatial distribution across the watershed than glaciers.



624
 625 **Figure 11. Field observations of cryospheric landforms as the ground truthing. a) Map of cryospheric landforms distribution, b) 3D**
 626 **view of the watershed, c) Rock glacier of the Del Potro river (03412); d) Gelifluction taluses of the Del Potro river (03412); e) Rock**
 627 **glacier of the Del Potro river (03412); f) Rock glacier of the Del Potro river (03412); g) Panorama of different periglacial cryospheric**
 628 **landforms of the Del Potro river (03412). Basemap: Esri, Maxar, Earthstar Geographics, and the GIS User Community.**




Figure 12. Field observations of cryospheric landforms as the ground truthing. a) Map of cryospheric landforms distribution, b) 3D view of the watershed, c) Del Potro debris-free glacier of the Del Potro river (03412), d) Maranceles glacier of the Montosa river (03413), e) Gelifluction taluses of the periglacial environment of the Del Potro river (03412). Basemap: Esri, Maxar, Earthstar Geographics, and the GIS User Community.

634 4.4 Geophysical results

635 4.4.1 Gelifluction Taluses


636 Electrical resistivity tomography (ERT) results (Fig. 13) provides the basis for interpreting the internal structure of the
637 gelifluction taluses. The lobate surface morphology characteristic of this cryospheric landform (Fig. 5 a, b), together with the
638 location and orientation of the tomographic profile (Figure 5 c, d), is clearly reflected in the geophysical response. The ERT
639 profile extends over 295 m and crosses a sector of the landform displaying a well-developed pattern of freeze–thaw-related
640 lobulations, starting at 4,430 m a.s.l. in the upstream sector of the Montosa sub-watershed. This geomorphological pattern
641 coincides with a high-resistivity zone (HRZ), with resistivity values ranging from approximately 1,000 to 8,000 Ωm .
642 According to comprehensive reviews and field studies on mountain permafrost, such resistivity values are typical of ice-rich
643 frozen ground, where pore water is largely transformed into electrically resistive ice (Kneisel and Hauck, 2008). Electrical
644 resistivity is known to increase by several orders of magnitude upon freezing, particularly in coarse-grained periglacial
645 materials, making ERT a robust tool for detecting ice-bearing permafrost bodies.

646 The HRZ delineates an ice-rich permafrost body with an estimated two-dimensional horizontal extension of approximately 20
647 m^2 within the tomographic section. Comparable resistivity ranges and internal structures have been documented in ice-rich
648 permafrost landforms investigated with ERT, including rock glaciers, talus slopes and other periglacial slope systems (Hauck
649 and Kneisel, 2008; Hilbich et al., 2009). The association between lobate slope morphology and underlying frozen ground with
650 significant ice content is consistent with the established understanding of gelifluction as a process operating on seasonally
651 thawed active layers overlying permafrost, where ice lenses and frozen substrates exert a first-order control on deformation
652 patterns (Matsuoka, 2001).

653 Downstream of 4,430 m a.s.l., the HRZ terminates abruptly at the frontal sector of the last lobulations. From this point, a low-
654 resistivity zone (LRZ) dominates, with resistivity values between 50 and 200 Ωm . This sharp contrast is interpreted as the
655 transition from frozen ground to unfrozen, water-saturated sediments. Similar resistivity ranges have been reported for
656 unfrozen debris, taliks and shallow groundwater zones underlying permafrost terrain, where electrolytic conduction through
657 liquid water strongly reduces bulk resistivity (Kneisel and Hauck, 2008).

658 The spatial juxtaposition of the HRZ and LRZ suggests a direct subsurface connection between the ice-rich permafrost body
659 and an underlying groundwater system. Studies combining ERT and seismic methods demonstrate that such resistivity
660 contrasts commonly mark permafrost boundaries and zones of hydrological interaction, where water infiltration and ice loss
661 can be detected geophysically (Hauck et al., 2008; Mewes et al., 2017). In particular, combined geophysical monitoring has
662 shown that increases in liquid water content beneath or adjacent to frozen ground produce a pronounced decrease in resistivity,
663 even when ice loss is spatially limited. Based on the vertical extent of the HRZ, the ERT data allow for the estimation of an
664 average ice-rich core thickness of approximately 15 m (Fig. 13). This thickness falls within the range reported for ice-bearing

665 periglacial slope landforms investigated using geophysical methods, where frozen layers commonly extend several tens of
666 meters below the surface depending on material properties and thermal conditions (Kneisel and Hauck, 2008; Hauck et al.,
667 2008).



668
669 **Figure 13. Results of the electrical resistivity tomography (ERT) over the gelifluction taluses cryospheric landforms in the Montosa**
670 **sub-watershed. a) Electrical resistivity tomography, b) Subsurface interpretation of resistivity test. Basemap: Esri, Maxar, Earthstar**
671 **Geographics, and the GIS User Community.**

673 4.4.2 Rock Glacier

674 A second ERT survey was conducted across a representative active rock glacier in the Montosa sub-watershed (Fig. 14). The
675 tomographic sections reveal a complex internal structure characterized by laterally continuous high-resistivity zones (HRZ),
676 which are interpreted as ice-rich permafrost bodies embedded within a coarse debris matrix, a configuration commonly
677 reported for active rock glaciers worldwide (Kneisel et al., 2008).

678 Resistivity values exceeding approximately 20–50 k Ω ·m dominate the central and frontal sectors of the landform, indicating
679 the presence of massive or highly ice-saturated material. Comparable resistivity ranges have been documented in ice-rich rock
680 glaciers in the Andes and other high-mountain regions, where electrical resistivity contrasts are primarily controlled by ground-
681 ice content rather than lithological variability (Kneisel et al., 2008; Pavoni et al., 2021). These high-resistivity anomalies are
682 spatially coherent with the surface morphology of the rock glacier, including its lobate structure and well-defined frontal slope,
683 supporting a close correspondence between geomorphological expression and subsurface ice distribution.

684 Beneath the HRZ, lower-resistivity zones (LRZ; <1–5 k Ω ·m) are observed, which are interpreted as unfrozen or partially
685 saturated debris layers and/or underlying bedrock. The relatively sharp transition between high- and low-resistivity domains
686 suggests a distinct boundary between ice-rich permafrost and unfrozen substrate, a feature widely reported in ERT studies of
687 active rock glaciers (Kneisel and Hauck, 2008).

688 The ERT-derived geometry indicates an average ice-rich layer thickness on the order of 20–30 m, with local maxima exceeding
689 35 m in the central part of the rock glacier body. These thicknesses fall within the range reported for active rock glaciers in the
690 dry Andes based on geophysical surveys and regional assessments (Brenning, 2005; Azócar and Brenning, 2010; Halla et al.,
691 2021). The observed thickness values provide an independent observational constraint for the thickness estimates derived from
692 the viscoplastic creep inversion based on DInSAR surface velocities (Cicoira et al., 2020).

693 Importantly, the spatial extent of the HRZ aligns closely with the mapped rock glacier polygon, reinforcing the validity of the
694 inventory-based delineation and confirming that the geophysically detected ice body corresponds to the geomorphological unit
695 considered in the volume calculations. The ERT results therefore serve as ground-based validation for both the presence of
696 ice-rich permafrost and the order of magnitude of the modeled thicknesses applied to rock glaciers within the Copiapó
697 watershed, consistent with previous multi-method approaches combining geomorphology, geophysics, and kinematic
698 observations.




Figure 14. Results of the electrical resistivity tomography (ERT) over rock glacier cryospheric landforms in the Montosa sub-watershed. Basemap: Esri, Maxar, Earthstar Geographics, and the GIS User Community. a) Longitudinal tomography, b) Cross-sectional tomography.

703 **4.5 Thickness estimations on periglacial environment**

704 Surface displacement maps (Fig. 15) of a representative rock glacier in the Copiapó watershed show a coherent downslope
705 motion that is clearly confined to the rock glacier body, while surrounding terrain remains stable, supporting the reliability of
706 both the cryosphere inventory and the DInSAR processing. These surface velocities provide a direct kinematic constraint for
707 the viscoplastic thickness modelling of ice-rich permafrost landforms used in this study.

708 The modeled ice-rich layer thicknesses for the main periglacial cryospheric landforms (rock glaciers, protalus lobes, and
709 gelifluction slopes) and their relationship with DInSAR-derived surface velocities (Fig. 16) show that rock glaciers exhibit the
710 largest modeled thicknesses, with median values on the order of ~25–30 m and a wide interquartile range, reflecting both their
711 higher surface velocities and the greater mechanical efficiency of creep in these landforms. In contrast, protalus lobes show
712 intermediate thicknesses, typically below ~15 m, while gelifluction slopes are characterized by consistently thinner ice-rich
713 layers, generally <10 m. The scatter plot highlights a positive relationship between mean surface velocity and modeled
714 thickness across all periglacial cryospheric landforms, consistent with the viscoplastic framework in which higher deformation
715 rates require larger ice thicknesses to sustain basal shear stress under comparable slope conditions.




Figure 15. Example of DInSAR-derived surface displacement of a rock glacier in the Copiapó watershed. Basemap: Esri, Maxar, Earthstar Geographics, and the GIS User Community.

Rock glaciers cluster at higher velocities and thicknesses, whereas protalus lobes and gelifluction slopes occupy lower-velocity regimes, with substantial overlap reflecting variability in slope, thermal state, and internal ice content. Despite this overlap, the three cryospheric landforms groups remain clearly distinguishable in thickness–velocity space, supporting the application of a unified viscoplastic model while preserving cryospheric landforms-specific structural differences. These results confirm that surface kinematics derived from DInSAR provide a physically meaningful constraint on subsurface ice thickness and justify their use for estimating ice volumes in ice-rich permafrost landforms across the Copiapó watershed.




Figure 16. Modeled thickness–velocity relationship for periglacial cryospheric landforms.

4.6 Ice volumes, snow area and NCI for sub-watershed

The ice volumes, snow area and the NCI for each of the sub-watersheds that constitute the cryospheric landforms of the Copiapó river watershed have been calculated (Table 6). The watershed with the largest ice masses is the Montosa river watershed (03413), with 0.922 Gt, followed by the Del Potro river watershed (03412) with 0.913 Gt and thirdly the southernmost watershed, the Manflas river watershed (03420) with 0.371 Gt of ice. In terms of volume, the largest ice volumes in the entire watershed are found in debris-free glaciers, gelifluction taluses and rock glaciers (Fig. 17).

Following the mapping of snow, we observe that the surface area distribution of snow cover is uneven, with a higher density in the southern part and the northern end of the watershed. In the south, the watersheds with the largest snow permanence are the Manflas river (03420), with 107.06 km² of mean snow cover (2000–2024), the Montosa river (03413) with 75.06 km² and the Del Potro river (03412) with 53 km². In the north, the most important rivers are the Estero Come Caballos (03402) with 122.24 km², Figueroa river (03401) with 106.72 km², Quebrada Monardes (03400) with 89.63 km² and the Cachitos river (03403) with 88.34 km² (Table 6).




Figure 17. Total ice mass of the Copiapó river watershed.




Figure 18. Total ice volumes of the Copiapó river watershed expressed both as a) total ice mass with associated uncertainty and b) as stacked contributions by cryospheric landforms.

The distribution of ice mass across cryospheric sub-sub-basins of the Copiapó watershed (Fig. 18), reveal a highly heterogeneous spatial distribution of cryospheric landforms, with a small number of sub-sub-basins concentrating the majority of the total ice mass.

The largest ice masses are found in the Montosa (COD_SSUBC 3413) and Del Potro (3412) sub-sub-basins, each storing approximately ~0.9 Gt of ice. These basins clearly dominate the cryospheric landforms of the watershed and are characterized by the combined presence of debris-free glaciers, rock glaciers, and extensive ice-rich permafrost landforms. In contrast,

751 several sub-sub-basins exhibit ice masses below 0.1 Gt, reflecting either limited cryospheric landforms extent or the dominance
752 of thin periglacial landforms.

753 The stacked representation highlights that debris-free glaciers constitute the principal contribution to ice mass in the largest
754 ice storage, whereas ice-rich permafrost cryospheric landforms, particularly rock glaciers and gelifluction taluses, represent a
755 substantial fraction of the total ice mass in intermediate and smaller sub-sub-basins. Protalus lobes contribute minor but non-
756 negligible volumes in selected basins, emphasizing their cumulative importance at the watershed scale.

757 Uncertainty ranges are largest in sub-sub-basins dominated by periglacial cryospheric landforms, reflecting the combined
758 effects of variability in surface velocity, thermal state, ice content, and mapped area. Nevertheless, the relative ranking of sub-
759 sub-basins by total ice mass remains robust, with Montosa and Del Potro consistently emerging as the primary cryospheric
760 landforms within the Copiapó watershed.

761 Overall, these results demonstrate that cryospheric ice mass is strongly concentrated in a limited number of sub-sub-basins,
762 while many others contain comparatively small but non-negligible ice volumes distributed among different cryospheric
763 landforms. However, ice mass alone does not fully capture the hydrological relevance of a watershed, as it does not account
764 for the capacity to regenerate water resources through snowfall. This motivates the integration of cryospheric landforms (ice
765 mass) and snow (Permanence) into a unified metric, introduced in the following section as the Normalized Cryospheric Index
766 (NCI).

767 The Normalized Cryospheric Index (NCI) behavior in relation to the relative weighting of ice volume (V) and snow occurrence
768 (S), along with the stability of sub-watershed rankings under uncertainty (Fig. 19) shows how the choice of weight (w)
769 influences the mean value of the NCI and, ultimately, the final ranking of the sub-watersheds.

770 The evolution of the mean NCI value of each sub-watershed as w varies from zero to one (Fig. 19a). Each line represents a
771 different sub-watershed, so that the slopes and crosses between lines allow us to appreciate which sub-watershed are favored
772 by a greater relevance of ice volume V (when w tends to 1) or snow cover S (when w tends to 0). This preliminary behavior
773 shows, on the one hand, the sensitivity of each subwatershed to weighting and, on the other hand, the possible dominance of
774 one variable over the other in certain w zones.


775 The position in the ranking that each subwatershed occupies for each w value (Fig. 19b) shows that the horizontal lines extend
776 from the first position (1) to the last. It highlights consistently privileged-ranked subwatersheds regardless of w , in contrast to
777 *unstably ranked* subwatersheds. This early insight into the stability of the ranking compellingly underscores the urgency of
778 undertaking a rigorous robustness analysis.

779 The ranking variability function, defined as the sum (or average) of the standard deviations of the position of each sub-
780 watershed in the different simulations (Fig. 19c) quantifies the overall stability of the ranking for each w . A marked minimum
781 is seen around $w=0.3$, indicating that, at that point, the ranking of watersheds is less sensitive to the uncertainty of the data of

782 V and S. In other words, assigning that weight in the combination of variables leads to a more consistent ranking in terms of
 783 robustness.

784 The mean NCI (with its error bars) for each watershed when $w=0.3$, the optimal value detected with the optimization function
 785 (Fig. 19d), shows that the vertical axis reflects the average magnitude of the index, while the error bars indicate the standard
 786 deviation derived from the Monte Carlo simulations. Thus, the visualization offers a clear identification of the watersheds with
 787 the highest NCI values, while simultaneously exposing the degree of uncertainty surrounding these estimates. This result
 788 confirms the relevance of choosing $w=0.3$ and provides a detailed view of the dispersion of each watershed under such a
 789 scenario.

790



791

792 **Figure 19. Analysis of NCI and robustness of the ranking as a function of w weighting (a) Evolution of the mean NCI for each sub-**
 793 **watershed as w varies between 0 and 1 (b) Position in the ranking of each sub-watershed as a function of w (c) Ranking variability**
 794 **function (sum of the standard deviations of the positions), where the minimum at $w \approx 0.3$ indicates the maximum stability of the**
 795 **ranking (d) Mean NCI $\pm$ standard deviation for $w=0.3$, evidencing the magnitude and dispersion of the index in each sub-watershed**
 796 **under the most robust condition.**

797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822

The calculated NCI, demonstrated that the sub-watershed with the highest NCI was the by the Montosa river watershed (03413) (NCI = 0.93), followed by the Del Potro river watershed (03412) (NCI = 0.89), Manflas river watershed (03420) (NCI = 0.47), Estero Come Caballos watershed (03402) (NCI = 0.34) and Cachitos river watershed (03403) (NCI = 0.31). These are the sub-watershed with the largest cryospheric landforms. In the northern region, the Estero Come Caballos sub-watershed, it was highlighted by the index primarily attributed to its persistent snow cover.

Table 6 Summary of results by sub-watershed of cryospheric landforms in Gt, snow in km² and indexes in different scenarios.

Code of the sub-watershed	Cryospheric landforms gigatons	Cryospheric landforms uncertainty	Snow in sq. km	NCI	NCI uncertainty
3400	0.039	0.006	89.63	0.14	0.01
3401	0.062	0.005	106.72	0.18	0.02
3402	0.212	0.014	122.24	0.34	0.02
3403	0.230	0.015	88.34	0.31	0.02
3410	0.007	0.001	43.90	0.04	0.01
3411	0.239	0.018	45.95	0.26	0.02
3412	0.913	0.124	53.02	0.89	0.11
3413	0.922	0.082	75.06	0.93	0.08
3420	0.371	0.055	107.06	0.47	0.05
3440	0.075	0.009	46.66	0.11	0.01
3441	0.041	0.006	46.79	0.08	0.01
3443	0.001	0.0005	19.76	0.00	0.00

5. Discussion

5.1 Cryospheric landforms and snow permanence in the Copiapó watershed

From the cryospheric landforms and snow occurrence mapping exercises (Figs. 4 and 8), we observed that the cryospheric landforms of the Copiapó watershed are predominantly located at altitudes above 4,500 m. This compares well with other regions in which snowfall has been identified to be controlled by altitude (Saydi and Ding, 2020). We also observed that cryospheric landforms occur in smaller, isolated regions ($< 0.25 \text{ km}^2$ in width) in the northern part of our research area. This corresponds with our own field observations, where most cryospheric landforms occur in smaller bodies (Fig. 4). This resulted in smaller, separated cryospheric landforms and surrounding the highest point (Figure 4 b and c). They host smaller cryospheric landforms, for example near the Cadillal and the Nevado Jotabeche, compared to the southern part of our study area. Because those northern cryospheric landforms are located in or in proximity of the SAAD (Abraham et al., 2020), snowfall and permanent snow occurrences are limited, leading to isolated cryospheric landforms (Fig. 4). The potential volume of water that is captured in cryospheric landforms in the sub-watersheds of the Copiapó watershed (Fig. 1) is thus strongly influenced by the South American Arid Diagonal (SAAD) that controls the distribution of the different types of cryospheric landforms (García et al., 2017). In our case, the proximity of the Copiapó watershed to the SAAD limits the occurrence of debris-free glaciers in the watershed north of 28.1° S latitude and limits the continuity of the cryosphere that is mainly attributed to cryospheric landforms associated with ice-rich permafrost, such as small rock glaciers, protalus lobes and predominantly gelifluction taluses.

In the more central domain in between Cadillal and Caserones no cryospheric landforms occur. This can be explained by the mechanical forcing of the South American low-level jet and the topographic blocking of the Andes (Insel et al., 2009). In the central sector, Mount Pissis exhibits greater elevation on the Argentinian side. Consequently, the prevailing northeasterly winds transporting moisture from the Pacific Ocean are obstructed by the Argentine portion of the mountain. This orographic interception results in localized snowfall on its Argentinian slopes, rather than allowing precipitation to be deflected by the comparatively lower-elevation Chilean topography. The topography north of Cadillal and south of Caserones is high enough to block these winds on the Chilean side, leading to snowfall and eventual cryospheric landforms in the southern regions in our research area. A last negative spatial correlation between the snowfall and the cryospheric landforms is the large snowfall anomaly in the northern area. The permanent snow occurrence is high, while not many cryospheric landforms are produced. The reason for this is the location of this region in the SAAD, resulting in melting or sublimation of the snow that does arrive in this area (Réveillet et al., 2020), meaning that the snow does not end up in cryospheric landforms.

From the different cryospheric landforms, we see that gelifluction taluses cover the largest surface area in the Copiapó watershed compared to other cryospheric landforms in our research areas (Garcia et al., 2017). Field observations also confirm that gelifluction taluses extend across significantly broader areas compared to other cryospheric landforms (Fig. 127). The limited understanding of gelifluction taluses has led to their exclusion from global cryospheric reserve inventories. Nevertheless, in assessing watershed ice volume, these taluses emerge as the most critical features for ice storage, surpassing

856 other formations in significance (Fig. 17). Furthermore, the ERT-derived thickness measurements across the gelifluction lobe
857 reveal a minimum detectable thickness of 15 meters, which markedly contrasts with the range of 1 to 5 meters reported by
858 Hilbich et al. (2015).

859 On a more regional scale, we observe that in the south of the watershed, debris-free glaciers are always in contact with
860 gelifluction taluses. This is confirmed in the field (Fig. 12). In the north of the Copiapó watershed, gelifluction taluses occur
861 more as isolated bodies. The topography of the northern area is controlled by Tertiary volcanoes (Kay et al., 1994). In general,
862 these volcanoes exhibit gentler slopes than those of the Principal Cordillera (Harrington, 1961) and largely shape the
863 topography of our southern research area. Given that gelifluction taluses require steep slopes to form (Fig. 2), they are more
864 likely to develop in the southern sector, where the terrain is dominated by the steep relief of the Principal Cordillera unlike in
865 the north, where this feature is absent. Moreover, in the Central Cordillera glacial cirques are abundantly present (Evans, 2006).
866 Glacial cirques provide sheltered environments that shield snow from extreme weather conditions, facilitating its
867 recrystallization into exposed glaciers and other cryospheric landforms. The distribution of rock glaciers and protalus lobes is
868 likewise influenced by the presence of glacial cirques (Palacios et al., 2020), resulting in their predominant occurrence in the
869 southern portion of our study area. In contrast, no exposed glaciers are found north of Copiapó Hill (Fig. 4). Instead, gelifluction
870 taluses and protalus lobes allow the continuity of the cryospheric fringe of the watershed. The absence of debris-free glaciers
871 in this region is directly related to the South American low-level jet and the topographic blocking of the Andes (Insel et al.,
872 2009).

873 These DGA (2014) volumes were used to provide a magnitude check. At the Del Potro Glacier, the modeled volume is ~24%
874 larger than the GPR-based estimate; at the Tronquitos Glacier, the difference is larger (~70%). This spread is within the
875 expected uncertainty range for individual glacier inversion products in Farinotti et al. (2019), particularly for small glaciers
876 with limited thickness constraints.

877 The DGA (2014) GPR-derived ice volume estimates were used as an independent benchmark to evaluate the magnitude of the
878 modeled glacier volumes (Table 7). For Del Potro Glacier, the modeled volume (0.362 km³) exceeds the GPR-based estimate
879 (0.293 km³) by approximately 24%. In the case of Tronquitos Glacier, the discrepancy is larger, with the modeled volume
880 (0.157 km³) being about 70% higher than the GPR-derived value (0.092 km³).

881 Despite these differences, the observed spread is consistent with the uncertainty ranges reported for individual glacier ice-
882 thickness inversion products in Farinotti et al. (2019). This is particularly relevant for small glaciers, where limited field-based
883 thickness constraints and complex geometries tend to amplify relative errors in volume estimates. Therefore, while the
884 discrepancies, especially for Tronquitos, are non-negligible, they remain within the expected variability for regional-scale
885 glacier volume modeling.

886

887 **Table 7. Comparison between modeled glacier ice volumes from Farinotti et al. (2019) and GPR-derived ice volumes**
 888 **reported by DGA (2014) for Del Potro and Tronquitos glaciers.**
 889

890	Glacier	RGI ID	Farinotti model volume (km³)	DGA 2014 GPR volume (km³)
891	Del Potro	1.715.087	0.362	
892				0.293
893	Tronquitos	1.715.038	0.157	0.092

896 The watershed with the best balance between cryospheric landforms and snow permanence is the Montosa river watershed
 897 (03413) (Fig. 20), which contains high quantities of both cryospheric landforms and snow permanence. The field observations
 898 of geomorphological features that are indicative of active cryospheric landforms, such as the subparallel terraces (Fig. 12e),
 899 fit well with the mapped cryospheric landforms from our workflow. The mapped surface area and thickness are supported by
 900 the observed surface area and thickness in the field. Lastly, debris-covered glaciers occur predominantly in areas with a
 901 relatively continuous cryosphere (García et al., 2017). This is consistent with our findings that debris-covered glaciers are
 902 exclusively located in the southern region of the study area, a zone characterized by a more extensive and contiguous
 903 cryospheric distribution relative to the fragmented northern sectors.

904 **5.2 Classifying and prioritizing sub-watersheds in the Copiapó watershed**

905 Unlike rainfall-dominated catchments, the upper Copiapó watershed is strongly influenced by cryospheric processes, with its
 906 headwaters characterized by the widespread presence of glaciers, rock glaciers, and ice-rich permafrost landforms. Cryospheric
 907 elements are concentrated in the high-altitude sub-watersheds, where water storage and seasonal runoff generation are
 908 controlled primarily by snow accumulation, snow persistence, and long-term ice storage rather than by direct rainfall inputs.
 909 The Normalized Cryospheric Index (NCI) identifies three main sub-watersheds with a dominant cryospheric influence on the
 910 hydrological functioning of the Copiapó system, based on the combined contribution of ice mass and long-term snow
 911 persistence. The spatial distribution of cryospheric landforms shows a 97% overlap with high-probability permafrost zones
 912 predicted by the global permafrost model of Gruber (2012), providing independent support for the relevance of ice-rich
 913 permafrost as a key component of water storage in the basin. Together, these observations justify classifying the upper Copiapó
 914 as a cryosphere-influenced watershed, in which cryospheric processes exert a first-order control on water availability. The
 915 critical sub-watersheds include: the Montosa River (03413), Del Potro River (03412) and Manflas River (03420) (Table 6).
 916 Their spatial distribution corresponds to zones of maximum snow permanence and the most extensive cryospheric continuity
 917 within the Copiapó watershed (Figs. 6 and 8). These high indexes thus directly correlate with the cryospheric landforms and
 918 snowfall occurrences. For example, enhanced snowfall is concentrated in the Cerro del Potro sub-watershed (03412), where
 919 the largest debris-free glacier of the Copiapó watershed is located (Fig. 8c). This area hosts El Potro Glacier, the most extensive

binational glacier in northern Chile (García et al., 2017), indicating a strong coupling between snow accumulation, glacier development, and long-term ice storage. As a result, the Cerro del Potro sub-watershed ranks among the highest NCI values, reflecting the combined importance of snow (permanence) and cryospheric landforms (ice mass). This combination has the potential to control the presence of water resources downstream when global temperatures continue to rise (Flores et al., 2018). With this study, we detected the sub-watersheds that should be prioritized. In wet years, snow controls the watershed hydrology in the northern part of the Copiapó watershed. This means that in these locations infrastructure needs to be put in place to capture the larger amounts of run-off that will arrive with increasing warmer temperatures, such as the Estero Come Caballos sub-watershed (03402), with the third place of the NCI. In times of water scarcity, we identified that the sub-watersheds of the Montosa river (03413), Del Potro river (03412), and Manflas river watershed (03420) will contribute most of the flow to the Copiapó river.

Given the observed relationship between snowfall and cryospheric landforms distribution, prioritizing the Copiapó watershed is essential to safeguard its water resources. We propose a systematic framework to evaluate and rank sub-watersheds according to preliminary assessments of cryospheric landforms and resource availability. From this ranking (Table 6), the main target, with the highest NCI is the sub-watershed of Montosa river (03413). The second target is located towards the south, the Del Potro river (03412). The third target is the Manflas river watershed (03420) and finally the last targets are Estero come Caballos (03402) and Cachitos river (03403) watershed (Table 6). These four sub-watersheds constitute the basis for continuous water production in the Copiapó watershed during dry years, because they host 83.44 % of the cryospheric landforms of the Copiapó watershed. From these four sub-watersheds, the Montosa river watershed (03413) maintains the best balance between snow and cryospheric landforms (Fig. 20). We have shown that with this hydrological approach, the NCI allows us to infer that the sub-watershed with the highest hydrological importance is the Montosa river watershed (03413). This index also allowed us to observe that the watersheds of the northern region are high in snow cover but are less enriched in cryospheric landforms. During years of substantial snowfall, management efforts should prioritize northern sub-watersheds, such as the Estero Come Caballos sub-watershed (03402). This suggests that northern watersheds exhibit a limited ability to develop cryogenic niches, as snow in these areas is primarily lost through melting or sublimation. A potential explanation for the lack of cryospheric landforms in northern watersheds lies in their geomorphological characteristics, which may hinder the accumulation and preservation of ice.

Based on the calculation of the NCI, we have been able to identify that the Estero Come Caballos watershed (03402) is the most strategic for snow accumulation, even surpassing the watersheds in the southern part of the Copiapó watershed in terms of snow permanence (Fig. 20).




Figure 20. Summary graph of the normalized indexes for snow, cryospheric landforms and the cryosphere for each sub-watershed of the Copiapó river watershed.

5.3 Impact on watershed management

We have analyzed 12 cryospheric sub-watersheds in the Copiapó watershed and showed that there are some that should be prioritized in watershed management strategy. The methodological framework developed and presented in this study can be extended to assess the comprehensive hydrological potential of arid watersheds in high-elevation Andean regions of Chile and other globally distributed arid zones with a focus on cryospheric contributions. In addition, this novel sub-watershed indexing method is fundamental to identify those watersheds with an important snow, glacial and periglacial component that are suitable for inclusion in the watershed protection frameworks that are implemented in the water management strategies of each watershed in Chile. Given the persistent progression of climate change, it is imperative to integrate water reserves sourced from cryospheric watersheds into comprehensive policy-making frameworks governing hydrological regulation.

Implementing and integrating such cryosphere inventories will also allow other regions to preserve their water resources. Currently, the Chilean national cryospheric inventories distinguish between glacier and periglacial landforms. Debris-free and debris-covered glaciers are comprehensively included in the National Glacier Inventory of Chile, whereas within the periglacial environment, only rock glaciers are systematically inventoried. As a result, ice volumes potentially stored in other ice-rich periglacial landforms, such as protalus lobes and gelifluction taluses, are not explicitly accounted for in existing national inventories, despite being addressed in this study. Increasing the variety of cryospheric landforms would facilitate more accurate estimations of the volumes of ice in each sub-watershed, especially in northern Chile where the gelifluction taluses and protalus lobes dominate the cryospheric landforms.

Identifying the strategic position of these sub-watersheds does not imply guaranteeing constant runoff, but rather highlights those sub-watersheds with a higher potential to sustain streamflow under low-snowfall or dry-year conditions. The runoff analysis shows that, during dry years, discharge is strongly reduced but not suppressed, and remains seasonally coupled to melt processes originating in high-elevation cryospheric environments. Sub-watersheds ranked as strategic by the NCI concentrate larger cryospheric landforms and higher snow permanence, which together contribute to buffering runoff variability and maintaining baseflow during periods of limited snowfall. The extraction of precious metals has consumed large quantities of water that was stored in cryospheric landforms (Brenning, 2008). A lack of comprehensive understanding regarding the spatial distribution and volumetric potential of cryospheric water reserves has contributed to the degradation of cryospheric landforms critical for supplying freshwater to downstream populations residing at lower elevations. These sub-watersheds are predominantly cryospheric rather than pluvial in character, indicating that their hydrological regime is mainly influenced by meltwater processes associated with glaciers, ice-rich periglacial landforms, and seasonal snow permanence, particularly during the melt season. When the potential volume of water in each sub-watershed is understood, policy and infrastructure can be adapted to maintain those cryospheric components. Prioritizing the extraction of water from cryospheric landforms and implementing snow capture techniques is essential to ensure the sustainable development and enhanced productivity of the examined region, as well as the arid northern regions of Chile. These sub-watersheds may therefore play an important role in the prevention of water disasters greater than those already known in Chile. The methodological basis developed in this study can be a key-tool for developing a law to protect Chile's cryosphere and its associated water resources when adopted in future water management strategy plans.

6. Conclusions

6.1 Main findings of the paper

We have developed a new workflow to identify cryospheric landforms and snow and applied this method to the Copiapó watershed. The workflow consists of mapping cryospheric landforms and snow occurrences and the calculation of a normalized cryospheric index (NCI) for sub-watersheds.

Our cryospheric landforms and snow occurrence mapping exercise resulted in the identification of cryospheric landforms in more isolated bodies north of Cadillal Hill and more continuous cryospheric landforms south of Caserones Hill.

Gelifluction taluses constitute a critical strategic cryospheric ice storage for safeguarding water security in northern Chile's watersheds. These landforms represent the periglacial type most likely to contain the largest ice volumes within the region's cryosphere, a hypothesis that requires further validation through targeted ERT (Electrical Resistivity Tomography) geophysical surveys of such cryospheric features.

The NCI allowed us to identify four main targets for watershed management. The four identified main targets are the Montosa river (03413), Del Potro river (03412), and Manflas river watershed (03420) with NCI of 0.93, 0.89 and 0.47 respectively. We propose that these sub-watersheds are the most important because that can sustain the runoff in the watershed in times of

drought. In years with snowfall the northern sub-watersheds as the Estero Come Caballos sub-watershed (03402) have the potential to be highly productive hydraulically by snow melt. This ranking allows the prioritization of these sub-watersheds. We conclude that the Montosa river (03413), Del Potro river (03412) and Manflas river (03420) are the sub-watersheds that should be prioritized in the national water resources strategy because of the high NCI.

6.2 Limitations of this work and future research

The workflow developed in this study enables the assessment of the role of a specific cryospheric sub-watershed within a broader drainage system, providing a basis for establishing protection frameworks in water resource management. This tool is transferable to other arid regions that are sourced through cryospheric landforms and snow. However, the method is calibrated for the Copiapó river watershed, and complete resource and reserve data should be available for the NCI method to be used on other watersheds.

To ensure more accurate error propagation in the NCI definition, it is essential to compute the uncertainty associated with the fractional snow cover product. We believe that assuming a 10% error under ideal conditions is one of the main limitations of the study of the snow permanence of the watershed. Our goal is to obtain a code that allows us to understand the uncertainties associated with the detection of snow in the mountainous areas of the watershed in order to improve the NCI and take it to the next level of research.

6.3 Broader impacts

The tool provides technical and objective data that water control agencies and those in charge of integrated watershed management, such as the DGA (General Directorate of Water), oversight boards and groundwater communities, should consider when implementing policy on the watershed management of the Copiapó watershed.

This study could help other arid regions of the planet with cryospheric landforms and snow to be able to manage the territory with a strategic plan to address the management of these cryospheric components.

7. Open science statement

The data generated for this project will become available upon publication. The cryospheric landforms data can be downloaded via this link: [<https://doi.org/10.5281/zenodo.15633086>]. The glacial and periglacial inventory for the Atacama region can be downloaded via this link: [<https://doi.org/10.5281/zenodo.14921499>]. The raster image for historical snow data can be downloaded via this link [<https://doi.org/10.5281/zenodo.14921552>].

8. Acknowledgments

We thank our colleagues of the Cryosphere and Water Research Laboratory for their indispensable contribution to the high altitude expeditions, and especially to project 5 of “Centro Avanzado para Tecnologías del Agua” (CAPTA) of the “Corporación de Fomento de la Producción” (CORFO) named “Tecnologías remotas y de campo para el aumento de oferta hídrica desde fuentes superficiales y subterráneas: acumulación nival y la recarga de acuíferos”, the Regional Government of Atacama and the University of Atacama, which sponsored the projects that made it possible to carry out this research and special thanks to Mr. Adrien Tavernier and Dra. Nicole Shaffer who has provided technical support in the execution of this work.

9. References

- Abraham, E. M., Rodríguez, M. G. L., Rubio, M. C., Guida-Johnson, B., Gomez, L. V., & Rubio, C (2020). Disentangling the concept of “South American Arid Diagonal”. *Journal of Arid Environments*, 175, 104089. <https://doi.org/10.1016/j.jaridenv.2019.104089>.
- Ahn, S. R., & Kim, S. J (2017). Assessment of integrated watershed health based on the natural environment, hydrology, water quality, and aquatic ecology. *Hydrology And Earth System Sciences*, 21(11), 5583-5602. <https://doi.org/10.5194/hess-21-5583-2017>.
- Ayala A, Pellicciotti F, Macdonell S, McPhee J, Vivero S, Campos C, Egli P (2016). Modelling the hydrological response of debris-free and debris-covered glaciers to present climatic conditions in the semiarid Andes of central Chile. *Hydrol Process* 30:4036–4058. <https://doi.org/10.1002/hyp.10971>.
- Ayala, Á., Fariás-Barahona, D., Huss, M., Pellicciotti, F., McPhee, J., and Farinotti, D (2020) Glacier runoff variations since 1955 in the Maipo river Basin, semiarid Andes of central Chile, *The Cryosphere*, 14, 2005–2027, <https://doi.org/10.5194/tc-14-2005-2020>, 2020.
- Azócar G. F. & Brenning, A. (2010). Hydrological and geomorphological significance of rock glaciers in the dry Andes, Chile (27°–33°S). *Permafr. Periglac. Proc.* 21, 42–53.
- Bahr, D. B., Meier, M. F., & Peckham, S. D (1997). “The physical basis of glacier volume-area scaling.” *Journal of Geophysical Research*, 102(B9), 20,355–20,362.
- Barsch, D (1996). *Rock Glaciers: Indicators for the Present and Former Geoecology in High Mountain Environments*. Springer (Series in the Physical Environment 16), Berlin, p. 331.
- Basuki, T. M., Nugroho, H. Y. S. H., Indrajaya, Y., Pramono, I. B., Nugroho, N. P., Supangat, A. B., Indrawati, D. R., Savitri, E., Wahyuningrum, N., Purwanto, Cahyono, S. A., Putra, P. B., Adi, R. N., Nugroho, A. W., Auliyani, D., Wuryanta, A., Riyanto, H. D., Harjadi, B., Yudilastyantoro, C., ... Simarmata, D. P (2022). Improvement of integrated watershed

management in Indonesia for mitigation and adaptation to climate change: A review. *Sustainability*, 14(16), 9997. <https://doi.org/10.3390/su14169997>.

Beniston, M., & Stoffel, M (2014). Assessing the impacts of climatic change on mountain water resources. *Science of The Total Environment*, 493, 1129-1137. <https://doi.org/10.1016/j.scitotenv.2013.11.122>.

Beniston, M., Farinotti, D., Stoffel, M., Andreassen, L. M., Coppola, E., Eckert, N., Fantini, A., Giacona, F., Hauck, C., Huss, M., Huwald, H., Lehning, M., López-Moreno, J. I., Magnusson, J., Marty, C., Morán-Tejeda, E., Morin, S., Naaïm, M., Provenzale, A., Vincent, C (2018). The European Mountain Cryosphere: A review of its current state, trends, and future challenges. *The Cryosphere*, 12(2), 759-794. <https://doi.org/10.5194/tc-12-759-2018>.

Bolch, T., Marchenko, S (2006). Significance of glaciers, rock glaciers, and ice-rich permafrost in the Northern Tien Shan as water towers under climate change conditions. In: *Proceedings of the Workshop ``Assessment of Snow-Glacier and Water Resources in Asia"*, 28-30 November 2006, Almaty, S. 199-211.

Bórquez, R., Larraín, S., Polanco, R., and J. Urquidi (2006). Glaciares chilenos. *Reservas estratégicas de agua dulce para la sociedad, los ecosistemas y la economía*. Chile Sustentable, LOM ediciones, Santiago, 124.

Bozkurt, D., Rojas, M., Boisier, J. P. & Valdivieso, J (2017). Climate change impacts on hydroclimatic regimes and extremes over Andean basins in Central Chile. *Hydrology and Earth System Sciences, Discuss*, <https://doi.org/10.5194/hess-2016-690>, 2017.

Brenning A (2005a). Climatic and geomorphological controls of rock glaciers in the Andes of Central Chile: combining statistical modelling and field mapping. PhD Thesis, Humboldt University, Berlin, Germany.

Brenning, A. (2008). The impact of mining on rock glaciers and glaciers: examples from Central Chile. In B. S. Orlove, E. Wiegandt, & B. Luckman (Eds.), *Darkening Peaks: Glacier Retreat, Science, and Society* (pp. 196–205). University of California Press.

Burger KC, Degenhardt JJ, Giardino JR (1999). Engineering geomorphology of rock glaciers. *Geomorphology*, 31: 93–132.

Casassa, G., Haeberli, W., Jones, G., Kaser, G., Ribstein, P., rivera, A., & Schneider, C (2007). Current status of Andean glaciers. *Global And Planetary Change*, 59(1-4), 1-9. <https://doi.org/10.1016/j.gloplacha.2006.11.013>.

Clarke, J (2006). Antiquity of aridity in the Chilean Atacama Desert. *Geomorphology*, N°73, pp. 101- 114. <http://dx.doi.org/10.1016/j.geomorph.2005.06.008>.

Corte, A.E (1978). Rock glaciers as permafrost bodies with debris cover as an active layer. A Hydrological approach, Andes of Mendoza, Argentina. In *Natural Resources Council of Canada (ed.), Proceedings of the third International Conference on Permafrost*, Edmonton, Alberta, Canada. Ottawa: Natural Resources Council of Canada, pp. 262-269.

Croce, F., & Milana, J.P (2002). Internal Structure and Behaviours of a rock glacier in the árid Andes of Argentina. *Permafrost Periglac. Process.*, 13: 289–299.

Cicoira, A., Marcer, M., Gärtner-Roer, I., Bodin, X., Arenson, L. U., & Vieli, A. (2020). A general theory of rock glacier creep based on in-situ and remote sensing observations. *Permafrost And Periglacial Processes*, 32(1), 139-153. <https://doi.org/10.1002/ppp.2090>.

1091 Delleur, J. W., (1999). The handbook of groundwater engineering. Boca Raton, FL: CRC Press.

1092 DGA (2013). Estrategia Nacional de Recursos Hídricos 2012-2025. I Resumen ejecutivo, p. 40. [https://www.mop.gob.cl/wp-](https://www.mop.gob.cl/wp-content/uploads/2021/04/Estrategia_Nacional_Recursos_Hidricos_DGA.pdf)

1093 [content/uploads/2021/04/Estrategia_Nacional_Recursos_Hidricos_DGA.pdf](https://www.mop.gob.cl/wp-content/uploads/2021/04/Estrategia_Nacional_Recursos_Hidricos_DGA.pdf)

1094 Evans, I. S (2006). Local aspect asymmetry of mountain glaciation: A global survey of consistency of favoured directions for

1095 glacier numbers and altitudes. *Geomorphology*, 73(1-2), 166-184. <https://doi.org/10.1016/j.geomorph.2005.07.009>.

1096 Favier, V., Falvey, M., Rabatel, A., Praderio, E., & Lopez, D. H (2009). Interpreting discrepancies between discharge and

1097 precipitation in high-altitude area of Chile's Norte Chico region (26–32°S). *Water Resources Research*, 45(2).

1098 <https://doi.org/10.1029/2008wr006802>.

1099 Flores, B., García, A., & Ulloa, C (2018). Evolución espacial y temporal de glaciares descubiertos en la Región de Atacama,

1100 Chile. XV Congreso Geológico Chileno, pp. 742-745.

1101 Farinotti, D., Brinkerhoff, D. J., Clarke, G. K. C., Fürst, J. J., Frey, H., Gantayat, P., Gillet-Chaulet, F., Girard, C., Huss, M.,

1102 Leclercq, P. W., Linsbauer, A., Machguth, H., Martin, C., Maussion, F., Morlighem, M., Mosbeux, C., Pandit, A., Portmann,

1103 A., Rabatel, A., Andreassen, L. M. (2017). How accurate are estimates of glacier ice thickness? Results from ITMIX, the Ice

1104 Thickness Models Intercomparison eXperiment. *The Cryosphere*, 11(2), 949–970. <https://doi.org/10.5194/tc-11-949-2017>.

1105 Farinotti, D., Huss, M., Fürst, J. J., Landmann, J., Machguth, H., Maussion, F., & Pandit, A. (2019). A consensus estimate for

1106 the ice thickness distribution of all glaciers on Earth. *Nature Geoscience*, 12(3), 168-173. [https://doi.org/10.1038/s41561-019-](https://doi.org/10.1038/s41561-019-0300-3)

1107 [0300-3](https://doi.org/10.1038/s41561-019-0300-3).

1108 Ferri, L., Dussailant, I., Zalazar, L., Masiokas, M. H., Ruiz, L., Pitte, P., Gargantini, H., Castro, M., Berthier, E., & Villalba,

1109 R. (2020). Ice mass loss in the Central Andes of Argentina between 2000 and 2018 derived from a new glacier inventory and

1110 satellite stereo-imagery. *Frontiers in Earth Science*, 8, 530997. <https://doi.org/10.3389/feart.2020.530997>.

1111 García, A., Ulloa, C., Amigo, G., Milana, J.P., Medina, C (2017). An inventory of cryospheric landforms in the arid diagonal

1112 of South America (High Central Andes, Atacama Region, Chile). *Quaternary International*, N°438, pp.4-19,

1113 <https://doi.org/10.1016/j.quaint.2017.04.033>.

1114 Garreaud, R., Boisier, J. P., Rondanelli, R., Montecinos, A., Sepúlveda, H. H., & Veloso-Águila, D (2019). The Central Chile

1115 Mega Drought (2010–2018): A climate dynamics perspective. *International Journal Of Climatology*, 40(1), 421-439.

1116 <https://doi.org/10.1002/joc.6219>

1117 Gärtner-Roer, I., Naegeli, K., Huss, M., Knecht, T., Machguth, H., & Zemp, M (2014). A database of worldwide glacier

1118 thickness observations. *Global And Planetary Change*, 122, 330-344. <https://doi.org/10.1016/j.gloplacha.2014.09.003>.

1119 Gascoin S, Kinnard C, Ponce R, Lhermitte S, Macdonell S, Rabatel A (2011). Glacier contribution to streamflow in two

1120 headwaters of the Huasco river, Dry Andes of Chile. *Cryosphere* 5:1099–1113. <https://doi.org/10.5194/tc-5-1099-2011>.

1121 Halla, C., Blöthe, J.H., Tapia, C., Trombotto, D., Hilbich, C., Hauck, C., Schrott, L (2021). Ice content and interannual water

1122 storage changes of an active rock glacier in the dry Andes of Argentina. *The Cryosphere*, 15, 1187-1213.

1123 <https://doi.org/10.5194/tc-15-1187-2021>.

1124 Hall, D. K., Riggs, G. A., Salomonson, V. V., DiGirolamo, N. E., & Bayr, K. J (2002). MODIS snow-cover products. Remote
 1125 Sensing of Environment, 83(1-2), 181-194. [https://doi.org/10.1016/S0034-4257\(02\)00095-0](https://doi.org/10.1016/S0034-4257(02)00095-0).

1126 Hanssen, R. F. (2001). Radar interferometry: Data interpretation and error analysis. Springer. <https://doi.org/10.1007/0-306-47633-9>.

1127

1128 Harrington, H. J (1961). Geology of Parts of Antofagasta and Atacama Provinces, Northern Chile. AAPG Bulletin, 45.
 1129 <https://doi.org/10.1306/0bda6332-16bd-11d7-8645000102c1865d>.

1130 Hauck, C., Bach, M., & Hilbich, C. (2008). A four-phase model to quantify subsurface ice and water content in permafrost
 1131 regions based on geophysical data sets. In K. M. Hinkel & D. L. Kane (Eds.), Proceedings of the Ninth International Conference
 1132 on Permafrost (Vol. 1, pp. 675–680). Institute of Northern Engineering, University of Alaska Fairbanks.

1133 Hilbich, C., Hauck, C., Mollaret, C., Wainstein, P., & Arenson, L. U (2022). Towards accurate quantification of ice content in
 1134 permafrost of the Central Andes – Part 1: Geophysics-based estimates from three different regions. The Cryosphere, 16(5),
 1135 1845-1872. <https://doi.org/10.5194/tc-16-1845-2022>.

1136 Hilbich, C., Marescot, L., Hauck, C., Loke, M. H., & Mäusbacher, R. (2009). Applicability of electrical resistivity tomography
 1137 monitoring to coarse blocky and ice-rich permafrost landforms. Permafrost and Periglacial Processes, 20(3), 269–284.

1138 Houston, J (2006). Variability of precipitation in the Atacama desert: its causes and hydrological impact. International journal
 1139 of climatology, 26: pp. 2181–2198.

1140 Insel, N., Poulsen, C. J., & Ehlers, T. A (2009). Influence of the Andes Mountains on South American moisture transport,
 1141 convection, and precipitation. Climate Dynamics, 35(7-8), 1477-1492. <https://doi.org/10.1007/s00382-009-0637-1>.

1142 Izquierdo, T., rivera, A.-L., Galeano, Á., Gallardo, D., Salas, V., Aparicio, O., Buylaert, J.-P., Ruiz, F., & Abad, M (2024).
 1143 Historical catastrophic floods at the southern edge of the Atacama Desert: A multi-archive reconstruction of the Copiapo river
 1144 extreme events. Global and Planetary Change, 236(104411), 104411. <https://doi.org/10.1016/j.gloplacha.2024.104411>.

1145 Janke, J.R., Ng, S., Bellisario, A (2017). An inventory and estimate of water stored in firn fields, glaciers, debris-covered
 1146 glaciers, and rock glaciers in the Aconcagua river Basin, Chile. Geomorphology, N°296, pp. 142-152.

1147 Jaiswal, R. K., Ghosh, N. C., Galkate, R. V., & Thomas, T (2015). Multi criteria decision analysis (MCDA) for watershed
 1148 prioritization. Aquatic Procedia, 4, 1553–1560. <https://doi.org/10.1016/j.aqpro.2015.02.201>.

1149 Jara, F. M., Lagos-Zúñiga, M., Fuster, R., Mattar, C., & McPhee, J (2021). Snow Processes and Climate Sensitivity in an Arid
 1150 Mountain Region, Northern Chile. Atmosphere, 12(4), 520. <https://doi.org/10.3390/atmos12040520>.

1151 Jones, D., Harrison, S., Anderson, K., & Betts, R (2018). Mountain rock glaciers contain globally significant water stores.
 1152 Scientific Reports, 8(1). <https://doi.org/10.1038/s41598-018-21244-w>.

1153 Jones, D., Harrison, S., Anderson, K., & Whalley, W. B (2019). Rock glaciers and mountain hydrology: A review. Earth-
 1154 Science Reviews, 193, 66-90. <https://doi.org/10.1016/j.earscirev.2019.04.001>.

1155 Jiang, Y., Hao, X., Li, Z., & Wang, J (2024). MODIS daily cloud-gap-filled fractional snow cover dataset of the Asian Water
 1156 Tower from 2000 to 2022. Earth System Science Data, 16(6), 2501-2523. <https://doi.org/10.5194/essd-16-2501-2024>.

1157 Kay, S. M., Mpodozis, C., Tittler, A., & Cornejo, P (1994). Tertiary Magmatic Evolution of the Maricunga Mineral Belt in
 1158 Chile. *International Geology Review*, 36(12), 1079-1112. <https://doi.org/10.1080/00206819409465506>.
 1159 Le Quesne C, DW Stahle, MK Cleaveland, MD Therrell, JC Ara-vena, J Barichivich (2006). Ancient Austrocedrus tree-
 1160 ringchronologies used to reconstruct central Chile precipitation variability from A.D. 1200 to 2000. *Journal of Climate* 19(22):
 1161 5731-5744.
 1162 Kirkbride, M. P. (2011). Debris-covered glaciers. En V. P. Singh, P. Singh, & U. K. Haritashya (Eds.), *Encyclopedia of snow,*
 1163 *ice and glaciers* (pp. 190–199). Springer. https://doi.org/10.1007/978-90-481-2642-2_622.
 1164 Kneisel, C., & Hauck, C. (2008). Advances in geophysical methods for permafrost investigations. *Permafrost and Periglacial*
 1165 *Processes*, 19(2), 157–178. <https://doi.org/10.1002/ppp.616>.
 1166 Koenig, L. S., Hauck, C., Azócar, G. F., et al. (2025). Thermal state of permafrost in the Central Andes (27–34° S). *The*
 1167 *Cryosphere*, 19, 2653–2680. <https://doi.org/10.5194/tc-19-2653-2025>.
 1168 Masiokas, M., Rabatel, A., rivera, A., Ruiz, L., Pitte, P., Ceballos, J. L., Barcaza, G., Soruco, Á., Bown, F., Berthier, É.,
 1169 Dussaillant, I., & MacDonell, S (2020). A Review of the Current State and Recent Changes of the Andean Cryosphere.
 1170 *Frontiers In Earth Science*, 8. <https://doi.org/10.3389/feart.2020.00099>.
 1171 Masiokas, M., Villalba, R., Luckman, B. H., Quesne, C. L., & Aravena, J. C (2006). Snowpack Variations in the Central Andes
 1172 of Argentina and Chile, 1951–2005: Large-Scale Atmospheric Influences and Implications for Water Resources in the Region.
 1173 *Journal Of Climate*, 19(24), 6334-6352. <https://doi.org/10.1175/jcli3969.1>.
 1174 Masiokas, M. H., Cara, L., Villalba, R., Pitte, P., Luckman, B. H., Toum, E., Christie, D. A., Quesne, C. L., & Mauget, S
 1175 (2019). Streamflow variations across the Andes (18°–55°S) during the instrumental era. *Scientific Reports*, 9(1).
 1176 <https://doi.org/10.1038/s41598-019-53981-x>.
 1177 Matsuoka, N. (2001). Solifluction rates, processes and landforms: A global review. *Earth-Science Reviews*, 55(1–2), 107–134.
 1178 [https://doi.org/10.1016/S0012-8252\(01\)00057-5](https://doi.org/10.1016/S0012-8252(01)00057-5).
 1179 Meier, M. F (1969). Glaciers and Water Supply. *Journal - American Water Works Association*, 61(1), 8–12.
 1180 doi:10.1002/j.1551-8833.1969.tb03696.x.
 1181 Mewes, B., Hilbich, C., Delaloye, R., & Hauck, C. (2017). Resolution capacity of geophysical monitoring regarding permafrost
 1182 degradation induced by hydrological processes. *The Cryosphere*, 11, 2957–2974. <https://doi.org/10.5194/tc-11-2957-2017>.
 1183 Milana, J.P (1998). Predicción de caudales de ríos alimentados por deshielo mediante balances de energía: Aplicación en los
 1184 Andes Centrales, Argentina. *Asociación Argentina de Sedimentología, revista*, v. 5 (2), p. 53-69.
 1185 Minvielle, M. & Garreaud, R. D (2011). Projecting Rainfall Changes over the South American Altiplano. *American*
 1186 *Meteorological Society*, N°24, pp. 4577-4583.
 1187 Monnier, S., & Kinnard, C. (2013). Internal structure and composition of a rock glacier in the Andes (Upper Choapa Valley,
 1188 Chile) using borehole information and ground-penetrating radar. *Annals of Glaciology*, 54(64), 61–72.
 1189 <https://doi.org/10.3189/2013AoG64A107>.

1190 Monnier, S., & Kinnard, C. (2017). Pluri-decadal (1955–2014) evolution of glacier–rock glacier transitional landforms in the
 1191 central Andes of Chile (30–33° S). *Earth Surface Dynamics*, 5(3), 493-509. <https://doi.org/10.5194/esurf-5-493-2017>.

1192 Morales, M. S., Christie, D. A., Villalba, R., Argollo, J., Pacajes, J., Silva, J. S., Alvarez, A., Llancabure, J. C. & Soliz Gamboa,
 1193 C. C (2012). Precipitation changes in the South American Altiplano since 1300 AD reconstructed by tree rings. *Climate of the*
 1194 *past*, N°8, pp. 653-666.

1195 Nicholson, L., Marin, J., Lopez, D., Rabatel, A., Bown, F., rivera, A., (2009). Glacier inventory of the upper Huasco valley,
 1196 Norte Chico, Chile: glacier characteristics, glacier change and comparison with central Chile. *Ann. Glaciol.* 50 (53), 111–118.

1197 Ohlanders, N., Rodriguez, M., and McPhee, J (2013). Stable water isotope variation in a Central Andean watershed dominated
 1198 by glacier and snowmelt, *Hydrol. Earth Syst. Sci.* 17, 1035–1050, <https://doi.org/10.5194/hess-17-1035-2013>, 2013.

1199 Palacios, D., Oliva, M., Gómez-Ortiz, A., Andrés, N., Fernández-Fernández, J. M., Schimmelpfennig, I., Léanni, L., & Team,
 1200 A (2020). Climate sensitivity and geomorphological response of cirque glaciers from the late glacial to the Holocene, Sierra
 1201 Nevada, Spain. *Quaternary Science Reviews*, 248, 106617. <https://doi.org/10.1016/j.quascirev.2020.106617>.

1202 Paterson, W.S.B., (1994). *The Physics of Glaciers*. Butterworth-Heinemann, Oxford (480 pp).

1203 Paul, F., Barrand, N. E., Baumann, S., Berthier, E., Bolch, T., Casey, K. & Wuite, J. (2013). On the accuracy of glacier outlines
 1204 derived from remote sensing data. *Annals of Glaciology*, 54(63), 171–182. <https://doi.org/10.3189/2013AoG63A296>

1205 Paul, F., Winsvold, S. H., Kääb, A., Nagler, T., & Schwaizer, G. (2015). Glacier remote sensing using Sentinel-2. *Remote*
 1206 *Sensing of Environment*, 162, 162–177. <https://doi.org/10.1016/j.rse.2015.06.037>.

1207 Pavoni, M., Sirch, F., & Boaga, J. (2021). Electrical and electromagnetic geophysical prospecting for the monitoring of rock
 1208 glaciers in the Dolomites, Northeast Italy. *Sensors*, 21(4), 1294. <https://doi.org/10.3390/s21041294>.

1209 Peng, X., Zhang, T., Frauenfeld, O. W., Wang, K., & Mu, C (2022). An integrated index of cryospheric change in the Northern
 1210 Hemisphere. *Global and Planetary Change*, 218, 103984. <https://doi.org/10.1016/j.gloplacha.2022.103984>.

1211 Peña, H. & Nazarala, B (1987). Snowmelt-Runoff Simulation Model of a Central Chile Andean Basin with Relevant
 1212 Orographic Effects. *Large Scale Effects of Seasonal Snow Cover*. IAHS Publication, 215 (166), 161-171.

1213 Pieczonka, T., Bolch, T., Kröhnert, M., Peters, J., & Liu, S. (2018b). Glacier branch lines and glacier ice thickness estimation
 1214 for debris-covered glaciers in the Central Tien Shan. *Cambridge Core*. <https://doi.org/10.1017/jog.2018.75>.

1215 Pourrier J, Jourde H, Kinnard C, Gascoin S, Monnier S (2014). Glacier meltwater flow paths and storage in a
 1216 geomorphologically complex glacial foreland: the case of the Tapado glacier, dry Andes of Chile (30°S). *J Hydrol* 519:1068–
 1217 1083. <https://doi.org/10.1016/j.jhydrol.2014.08.023>.

1218 Radić, V., & Hock, R (2010). “Regional and global volumes of glaciers derived from statistical upscaling of glacier inventory
 1219 data.” *Journal of Geophysical Research*, 115, F01010.

1220 Ragetli S. and Pellicciotti F (2012). Calibration of a physically based, spatially distributed hydrological model in a glacierized
 1221 basin: on the use of knowledge from glaciometeorological processes to constrain model parameters. *Water Resour Res*
 1222 48(W03509):1–20. <https://doi.org/10.1029/2011WR010559>.

1223 Rahaman, S. A., Ajeez, S. A., Aruchamy, S., & Jegankumar, R (2015). Prioritization of sub-watershed based on morphometric
 1224 characteristics using fuzzy analytical hierarchy process and geographical information system – A study of Kallar Watershed,
 1225 Tamil Nadu. *Aquatic Procedia*, 4, 1322-1330. <https://doi.org/10.1016/j.aqpro.2015.02.172>.
 1226 Réveillet, M., MacDonell, S., Gascoin, S., Kinnard, C., Lhermitte, S., & Schaffer, N (2020). Impact of forcing on sublimation
 1227 simulations for a high mountain catchment in the semiarid Andes. *The Cryosphere*, 14(1), 147-163. [https://doi.org/10.5194/tc-](https://doi.org/10.5194/tc-14-147-2020)
 1228 [14-147-2020](https://doi.org/10.5194/tc-14-147-2020).
 1229 Rittger, K., Painter, T. H., & Dozier, J (2013). Assessment of methods for mapping snow cover from MODIS. *Advances In*
 1230 *Water Resources*, 51, 367-380. <https://doi.org/10.1016/j.advwatres.2012.03.002>.
 1231 Rodriguez M, Ohlanders N, Pellicciotti F, Williams MW, McPhee J (2016). Estimating runoff from a glacierized catchment
 1232 using natural tracers in the semi-arid Andes cordillera. *Hydrol Process* 30:3609– 3626. <https://doi.org/10.1002/hyp.10973>
 1233 Sanmiguel-Valladolid, A., Morán-Tejeda, E., Alonso-González, E., & López-Moreno, J. I (2017). Effect of snow on mountain
 1234 river regimes: an example from the Pyrenees. *Frontiers Of Earth Science*, 11(3), 515-530. [https://doi.org/10.1007/s11707-016-](https://doi.org/10.1007/s11707-016-0630-z)
 1235 [0630-z](https://doi.org/10.1007/s11707-016-0630-z).
 1236 Saydi, M., and Ding, J (2020). Impacts of topographic factors on regional snow cover characteristics. *Water Science And*
 1237 *Engineering/Water Science And Engineering*, 13(3), 171-180. <https://doi.org/10.1016/j.wse.2020.09.002>.
 1238 Schaffer, N., MacDonell, S., Réveillet, M., Yáñez, E., & Valois, R (2019). Rock glaciers as a water resource in a changing
 1239 climate in the semiarid Chilean Andes. *Regional Environmental Change*, 19(5), 1263-1279. [https://doi.org/10.1007/s10113-](https://doi.org/10.1007/s10113-018-01459-3)
 1240 [018-01459-3](https://doi.org/10.1007/s10113-018-01459-3).
 1241 Schrott, L (1996). Some geomorphological-hydrological aspects of rock glaciers in the Andes (San Juan, Argentina).
 1242 *Zeitschrift für Geomorphologie*, 104, 161-173.
 1243 Sriyana, I (2019). Developed watershed classification index determining management priority level based on watershed
 1244 carrying capacity. *MATEC web of conferences*, 270, 04004. <https://doi.org/10.1051/mateconf/201927004004>.
 1245 Suárez, F., Muñoz, J. F., Fernandez, B., Dorsaz, J-M., Hunter, C. K., Karavitis, C., Gironás, J (2014). Integrated Water
 1246 Resource Management and Energy Requirements for water supply in the Copiapo river Basin, Chile. *Water*, 6, pp. 2590-2613.
 1247 Trombotto, D. L., Sileo, N., & Dapeña, C (2020). Periglacial water paths within a rock glacier-dominated catchment in the
 1248 Stepanek area, Central Andes, Mendoza, Argentina, *Permafrost Periglac.*, 31, 311–323.
 1249 Valdivieso, S., Hassanzadeh, A., Vázquez-Suñé, E., Custodio, E., Criollo, R (2022). Spatial distribution of meteorological
 1250 factors controlling stable isotopes in precipitation in Northern Chile. *Journal of Hydrology*, 605, 127380.
 1251 Vuille, M., Francou, B., Wagnon, P., Juen, I., Kaser, G., Mark, B. G., & Bradley, R. S (2008). Climate change and tropical
 1252 Andean glaciers: Past, present and future. *Earth-science Reviews*, 89(3-4), 79-96.
 1253 <https://doi.org/10.1016/j.earscirev.2008.04.002>.
 1254 Whitaker, A., Sugiyama, H., & Hayakawa, K (2008). Effect of Snow Cover Conditions on the Hydrologic Regime: Case Study
 1255 in a Pluvial-Nival Watershed, Japan1. *Journal Of The American Water Resources Association*, 44(4), 814-828.
 1256 <https://doi.org/10.1111/j.1752-1688.2008.00206.x>.

1257 Wolfe, J. D., Shook, K. R., Spence, C., & Whitfield, C. J (2019). A watershed classification approach that looks beyond
1258 hydrology: application to a semi-arid, agricultural region in Canada. Hydrology And Earth System Sciences, 23(9), 3945-
1259 3967. <https://doi.org/10.5194/hess-23-3945-2019>.
1260 Zech, R., May, J.H., Kull, C., Ilgner, J., Kubik, P.W., Veit, H (2008) - Timing of the late Quaternary
1261 glaciation in the Andes from ~15 to 40°S. Journal of Quaternary Science, 23 (6-7), pp. 635–647.
1262