

Direct assimilation of ground-based microwave radiometer observations with machine learning bias correction based on developments of RTTOV-gb v1.0 and WRFDA v4.5

Qing Zheng^{1,3}, Wei Sun^{2,3}, Zhiqian Liu⁴, Jiajia Mao⁵, Jieying He⁶, Jian Li^{2,3}, Xingwen Jiang^{1,3}

5 ¹Institute of Plateau Meteorology, China Meteorology Administration, Chengdu, 610213, China

²State Key Laboratory of Severe Weather, Chinese Academy of Meteorological Sciences, Beijing, 100081, China

³Institute of Tibetan Plateau Meteorology, China Meteorological Administration, Chengdu, 610213, China

⁴National Center for Atmospheric Research, Boulder, 80307-3000, USA

⁵Meteorological Observation Center, China Meteorological Administration, Beijing, 100081, China

10 ⁶National Space Science Center, Chinese Academy of Sciences, Beijing, 100190, China

Correspondence to: Wei Sun (sunwei@cma.gov.cn)

Abstract. The application of ground-based microwave radiometers (GMWRs), which provide high-quality and continuous vertical atmospheric observations, has traditionally focused on the indirect assimilation of retrieved profiles. This study

15 advanced this application by developing a direct assimilation capability for GMWR radiance observations within the Weather Research and Forecasting model data assimilation (WRFDA) system, along with a bias correction scheme based on the random forest technique. The proposed bias correction scheme effectively reduced the observation-minus-background (O–B) biases

20 and standard deviations by 0.83 K (97.1 %) and 1.63 K (64.6 %), respectively. A series of ten-day-long experiments demonstrated that assimilating GMWR radiances improves both the initial conditions and the forecasts, with additional benefits

25 from higher assimilation frequencies. In the initial conditions, hourly assimilation significantly enhanced low-level temperature and humidity fields, reducing the root-mean-square error (RMSE) for temperature and water vapor mixing ratio by 6.32 % below 1 km and 1.98 % below 5 km. These improvements extended to forecasts, where 2 m temperature and humidity showed sustained benefits for over 12 hours, and precipitation forecasts exhibited improvements to a certain extent.

The time-averaged Fractions Skill Score (FSS) for 3 h accumulated precipitation within the 24 h forecasts increased by 0.02–0.04 (3.9–10.2 %) for thresholds of 3–6 mm.

1 Introduction

Data assimilation (DA), a core component of numerical weather prediction (NWP), plays an important role in improving forecast accuracy by integrating observational data to refine initial conditions (Bauer et al., 2015; Gustafsson et al., 2018). Among various types of observations, microwave radiance data are crucial for DA due to their ability to penetrate the 30 atmosphere and their sensitivity to temperature, humidity, clouds, and precipitation. Correspondingly, satellite-borne

microwave radiance observations have been extensively studied and are considered among the most influential contributors to data assimilation systems (Geer et al., 2017; Kim et al., 2020; Candy and Migliorini, 2021).

Unlike satellite-borne microwave radiometers, ground-based microwave radiometers (GMWRs) offer unique advantages for DA, including high temporal resolution (minute-level) and greater sensitivity to the atmospheric boundary layer (ABL). Over 35 the past two decades, the assimilation of GMWRs has been increasingly studied, leading to improvements in the accuracy of NWP (Vandenberge and Ware, 2002; Otkin, 2010; Hartung et al., 2011; Otkin et al., 2011; Caumont et al., 2016; He et al., 2020; Qi et al., 2021, 2022; Lin et al., 2023). The assimilation of retrieved temperature and humidity profiles from GMWRs has shown improvements in forecasting fog, storms, and precipitation. However, the reliance on indirect assimilation methods introduces uncertainties and complicates error quantification, which limits their overall effectiveness in enhancing forecast 40 accuracy (Caumont et al., 2016; Martinet et al., 2017; Lin et al., 2023).

Direct assimilation of GMWR radiances, which bypasses the retrieval process, offers significant advantages by avoiding retrieval-related errors and improving the effective use of observations. This approach requires accurate observation operators and robust bias correction to address differences between radiance observations and model states. The direct assimilation of 45 satellite-borne radiance observations is relatively mature (Geer et al., 2008; Bauer et al., 2010; Geer et al., 2010; Eyre et al., 2020; Sun and Xu, 2021; Eyre et al., 2022) and utilizes fast radiative transfer models (RTMs) as observation-operators, such as the Radiative Transfer for Television and Infrared Observation Satellite (RTTOV) (Saunders et al., 2018). However, the unique characteristics of upward-looking GMWR observations, such as sensitivity to near-surface conditions, require 50 specialized RTMs and adaptation of existing techniques. Recent studies have developed fast RTMs suitable for GMWRs, which provide a foundation for constructing observation operators for assimilation of GMWR radiances (De Angelis et al., 2016; Cimini et al., 2019; Shi et al., 2025). The RTTOV-gb, a ground-based version of the RTTOV model, was used to simulate brightness temperature from GMWRs, demonstrating high accuracy (De Angelis et al., 2016, 2017; Cimini et al., 2019). Recent studies have demonstrated the potential of direct GMWR radiance assimilation using RTTOV-gb to improve temperature, 55 humidity, and precipitation forecasts (Cao et al., 2023; Vural et al., 2024).

Despite these advancements, previous studies have typically relied on limited GMWR networks or focused on specific case 60 studies. Additionally, research conducted in regions with relatively simple terrain may not fully address the complexities of areas like the Tibetan Plateau, where complex topography often leads to significant model biases (Yang et al., 2020; Wei et al., 2021). These biases make accurate bias correction essential for improving the effectiveness of direct assimilation, while traditional bias correction approaches developed for satellite-borne microwave radiance observations are not directly applicable to GMWRs.

To address these issues, this study integrates RTTOV-gb into the Weather Research and Forecasting Data Assimilation (WRFDA) system (Barker et al., 2012) to develop a direct assimilation module for GMWR radiances. A nonlinear bias correction scheme based on machine learning is also constructed using three months of observational data. The impact of direct 65 GMWR assimilation is then investigated through a series of ten-day experiments conducted in Southwest China, a region shaped by the influence of the Tibetan Plateau and characterized by complex terrain. The remainder of this paper is organized

65 as follows. Section 2 describes the data, the implementation of RTTOV-gb in WRFDA, and the model configuration. Section 3 evaluates the performance of the bias correction scheme. Section 4 presents the impacts of GMWR assimilation on the initial and forecast fields. The conclusions and discussion are presented in Sect. 5.

2 Methodology

2.1 Data

70 Two types of GMWR sensors were assimilated in this study, as shown in Fig. 1: the MP3000A and the Humidity And Temperature Profiler (HATPRO). Atmospheric radiance is measured as brightness temperatures in 14 channels for HATPRO and 22 channels for MP3000A (Table 1). For HATPRO, channels 1–7 are in the K-band, while channels 8–14 are in the V-band. For MP3000A, channels 1–8 are in the K-band, and channels 9–22 are in the V-band. The K-band channels correspond 75 to humidity-sensitive water vapor absorption lines, whereas the V-band channels correspond to temperature-sensitive oxygen absorption lines.

The Fengyun-4B (FY-4B) Advanced Geosynchronous Radiation Imager (AGRI) cloud mask (CLM) is used to identify GMWR-observed brightness temperatures under clear-sky conditions. The AGRI-based CLM product has a temporal resolution of 15 minutes and a horizontal resolution of 4 km, categorizing conditions as confidently cloudy, probably cloudy, probably clear, or confidently clear, with corresponding values of 0, 1, 2, and 3, respectively (Min et al., 2017). Due to its high 80 quality, this cloud mask product is widely applied in satellite data assimilation (Yin et al., 2020, 2021; Xu et al., 2023; Shen et al., 2024).

The National Centers for Environmental Prediction (NCEP) Final Operational Global Analysis data (FNL) ($0.25^\circ \times 0.25^\circ$, 6-hourly) were used to establish the initial and boundary conditions for regional NWP. Conventional observations from the Global Telecommunications System (GTS) were assimilated and evaluated, including land surface, marine surface, radiosonde, 85 and aircraft reports. The hourly precipitation analysis product from the China Meteorological Administration Multisource Precipitation Analysis System (Shen et al., 2014) was used for evaluation. This dataset has been widely used in precipitation studies (Xia et al., 2019; Su et al., 2020; Sun and Xu, 2021; Wang et al., 2021; Li et al., 2023; Zheng et al., 2024).

Table 1. Central frequency for GMWRs

Sensor	Frequencies for K-band (GHz)	Frequencies for V-band (GHz)
HATPRO	22.240;23.040;23.840;25.440; 26.240;27.840;31.400	51.260;52.280;53.860;54.940; 56.660;57.300;58.000
MP3000A	22.234;22.500;23.034;23.834; 25.000;26.234;28.000;30.000	51.248;51.760;52.280;52.804; 53.336;53.848;54.400;54.940; 55.500;56.020;56.660;57.288; 57.964;58.800

90 **2.2 Assimilation system and observation operator**

The WRFDA system, developed by the National Center for Atmospheric Research (NCAR), is designed for data assimilation and includes three-dimensional variational (3DVAR), four-dimensional variational (4DVAR), and hybrid data assimilation algorithms. In this study, version 4.5 of the WRFDA system with 3DVAR is used for the direct assimilation of GMWR radiances. The 3DVAR algorithm produces the analysis by minimizing a scalar objective cost function:

$$95 \quad J(\mathbf{x}) = \frac{1}{2}(\mathbf{x} - \mathbf{x}_b)^T \mathbf{B}^{-1}(\mathbf{x} - \mathbf{x}_b) + \frac{1}{2}(\mathbf{y} - \mathbf{H}(\mathbf{x}))^T \mathbf{R}^{-1}(\mathbf{y} - \mathbf{H}(\mathbf{x})), \quad (1)$$

where \mathbf{x} and \mathbf{x}_b represent the analysis and background fields of the model variables, \mathbf{y} is the vector of the observations, and \mathbf{B} and \mathbf{R} represent the background and observation error covariance matrices, respectively. The covariance matrices determine the weights assigned to the background and observations in the analysis, dictate how localized observation information is distributed vertically and horizontally in the model space, and maintains the balance among the model's control variables. \mathbf{H} is the non-linear observation operator, that transforms model variables to the observed quantities. The observation operator works slightly differently for different types of observations. For conventional observations (e.g., temperature), its primary role is to perform spatiotemporal interpolation of model grid values to the observation space. For unconventional observations (e.g., reflectivity and radiance), where the model state cannot be directly compared with the observations, the observation operator must also convert model variables into observed variables.

105 The static background error covariance for the variational experiments is estimated using the National Meteorological Center (NMC) method (Parrish and Derber, 1992), which uses the difference between WRF forecasts at lead times of 24 h and 12 h ($T + 24$ h minus $T + 12$ h) valid at the same time over a specified period. Control variables option 5 (CV5) is adopted for the background error covariance used in 3DVAR. CV5 is domain-dependent and therefore must be generated based on forecast or ensemble data over the same domain. It utilizes streamfunction, unbalanced velocity potential, unbalanced temperature, 110 unbalanced surface pressure, and pseudo relative humidity. In this study, the background error covariance matrix was generated using the Generalized Background Error Covariance Matrix Model (GEN_BE v2.0) (Descombes et al., 2015) based on one month of WRF forecasts. Observation-error correlations are typically assumed to be zero in WRFDA, resulting in a diagonal observation-error covariance matrix. Observation errors were specified based on the standard deviation of O–B.

115 RTMs serve as observation operators for assimilating radiance data by mapping model variables (e.g., temperature and water vapor) into radiance space. RTTOV, a fast RTM, is widely used for assimilating satellite radiance data. However, GMWR radiances are upward-looking microwave observations, differing from the downward-looking measurements of satellite-borne microwave radiometers. This difference in direction makes RTTOV difficult to apply in GMWR radiances assimilation. Fortunately, RTTOV-gb can simulate brightness temperatures from GMWRs, and serves as the observation operator in this study. The weighting function (WF) quantifies the contribution of emissions from each atmospheric layer, and the maximum 120 WF height indicates which atmospheric layer contributes most to the measured radiance (Carrier et al., 2008). According to Cui et al. (2020), WFs are calculated as the derivative of transmittance with respect to the natural logarithm of pressure. The

vertical distribution of WFs for HATPRO and MP3000A, calculated using RTTOV-gb, is shown in Fig. 2. The WFs reach their maximum at 1000 hPa and decrease monotonically with height. These results confirm that the lower atmosphere contributes most to the observed radiation across all channels, consistent with the findings of Shu et al. (2012).

125 It should be noted that RTTOV-gb is not included in the publicly available version of WRFDA. To address this limitation, a GMWR direct assimilation module was developed within WRFDA. A single-observation assimilation experiment was conducted to test the GMWR direct assimilation module. In this experiment, a pseudo radiance observation from the HATPRO sensor was assimilated in channel 6 (K band), with an assigned observation error of 1 K and an innovation of 2 K. Results confirm that the GMWR direct assimilation module performs correctly. The temperature and water vapor increments are
130 horizontally isotropic and show a maximum at lower atmospheric levels vertically (Fig. 2). It should also be noted that this experiment was conducted to verify the correct performance of the GMWR direct assimilation module and to provide valuable insights into the characteristics of GMWR assimilation. However, it is not representative of the subsequent multi-observation, multi-channel assimilation experiments.

135 2.3 Model configuration and experimental design

In this study, version 4.5 of the Weather Research and Forecasting (WRF) model (Skamarock et al., 2021) is used to simulate atmospheric evolution. The simulation employs a single domain (Fig. 1) with a horizontal resolution of 3 km, comprising $1,261 \times 811$ grid points and 51 vertical levels, with the top boundary at 10 hPa. The model physics configuration includes the Morrison two-moment microphysics scheme (Morrison et al., 2009), the Yonsei University PBL scheme (Hong et al., 2006),
140 the Rapid Radiative Transfer Model for General Circulation Models (RRTMG) shortwave and longwave radiation schemes (Iacono et al., 2008), and the unified Noah land-surface model (Chen and Dudhia, 2001). Cumulus parameterization was excluded due to the convection-permitting horizontal resolution of 3 km (Li et al., 2023; Moker et al., 2018).


Similar to previous studies (Jiang et al., 2017; Nie and Sun, 2023), the target region of Southwest China in this study is defined as the area within the rectangular domain 22° – 35° N, 93° – 110° E (Fig. 1). This region encompasses the Hengduan Mountains,
145 the Yunnan–Guizhou Plateau, and the Sichuan Basin, and is generally consistent with Chinese administrative divisions. Based on the model configuration described above, four parallel experiments were conducted to investigate the impact of GMWR assimilation (Table 2). Each experiment started at 12:00 UTC daily, incorporating 12 hours of data assimilation followed by a 24 h forecast. The primary differences among these experiments lie in the assimilated data and assimilation intervals. The CNTL experiment assimilated GTS data with a 6 h interval, while the GMWR_6H experiment added GMWR assimilation to
150 the CNTL setup, enabling an evaluation of GMWR assimilation's impact. The other two experiments, GMWR_3H and GMWR_1H, assimilated both GTS and GMWR data with 3 h and 1 h intervals, respectively, to assess the effects of observation frequency in GMWR assimilation.

The assimilation experiments were conducted under clear-sky conditions due to the uncertainties in the model and observation operators under cloudy or rainy conditions. All experiments were conducted over a ten-day period from 13 to 22 October 2023.

155 Among the available GMWR observations from August to October 2023, this period exhibited a notably higher frequency of clear-sky data, which was more favorable for demonstrating the role and potential of GMWR assimilation. Before implementing bias correction, clear-sky screening, first-guess departure check, and whitelist check were sequentially applied to improve measurement quality. Subsequently, a relative departure check was applied prior to minimization. For the 6 h, 3 h, and 1 h assimilation intervals, 34 (0.91%), 70 (1.42%), and 76 (0.72%) observations were rejected, respectively. The detailed

160 procedure prior to a single assimilation cycle is as follows:

- (1) Observation Selection: The observation nearest to the analysis time within ± 10 minutes is selected.
- (2) Clear-sky Screening: Clear-sky GMWR observations were screened using the AGRI-based CLM, with background-simulated cloud liquid water path equal to zero.
- (3) First-Guess Departure Check: Observations with (O–B) values greater than 20 K are excluded.
- 165 (4) Whitelist Check: Remove observations from stations identified as unreliable or displaying abnormal behavior.
- (5) Bias Correction: A machine learning bias correction scheme was applied (see Section 3.2).
- (6) Relative Departure Check: Applied when the absolute value of the O–B exceeds three times the standard deviation of the observational error, further rejecting questionable data.



170

Figure 1: Computation domain(shaded). The shaded denotes topography (units: m). The green rectangle denotes the target region of Southwest China. The blue empty circle denotes radiosonde. The 'x' and '+' symbols denote HATPRO and MP3000A, respectively.




Figure 2: Normalized weighting functions of (a) HATPRO and (b) MP3000A calculated using the RTTOV-gb. The (c, d) horizontal and (e, f) vertical analysis increments for (c, e) temperature and (d, f) water vapor mixing ratio in single-observation assimilation experiment. The vertical increments are cross-sections along the green lines shown in the horizontal increments. The colorbar tick labels for temperature and water vapor mixing ratio are expressed in scientific notation as 1×10^{-2} and 1×10^{-4} , respectively.

Table 2 Experimental design

Experiment	Assimilated Data	Assimilation Interval
CNTL	GTS	6-hour
GMWR_6H	GTS and GMWR	6-hour
GMWR_3H	GTS and GMWR	3-hour
GMWR_1H	GTS and GMWR	1-hour

180 3 Machine learning based bias correction for GMWR

3.1 Bias characteristics

Variational assimilation assumes that both observation and background errors follow an unbiased Gaussian distribution. However, due to instrument errors, limitations of the RTMs, and errors in the NWP model background, observed radiances (O) and simulated radiances (B) inherently contain errors (denoted as μ^o and μ^b), which may exhibit a biased distribution. Bias correction is a crucial process in radiance data assimilation, aiming to identify and remove these biases (Auligné et al., 2007;

Dee, 2005). In the real atmosphere, O and B are regarded as the true value (T) plus their respective deviations μ , as shown in Eq. (2):

$$\overline{O - B} = \overline{(O - T) - (B - T)} = \overline{\mu^o - \mu^b}, \quad (2)$$

It shows that the statistical expectation value of O–B can represent the systematic deviation ($\mu^o - \mu^b$). Therefore, it is critical
190 to evaluate the bias characteristics of O–B and correct them.

To estimate the bias and develop a bias correction scheme for GMWR direct assimilation, a long-term experiment was conducted from August to October 2023, yielding a three-month sample dataset. In this experiment, the WRF model was initialized every 6 hours using NCEP FNL data, and WRFDA operated hourly in monitoring mode (only calculating O–B). After a cloud check using the AGRI-based CLM and a gross check ($O - B < 20$ K), the bias of O–B for HATPRO and MP3000A
195 was estimated.

A comparative scatterplot analysis of observed and simulated brightness temperatures was conducted. For most channels, the scatter points are closely aligned along the diagonal and exhibit high correlation coefficients, indicating strong agreement between the simulations and observations. However, the scatter points for some channels forms two distinct clusters. To further investigate, representative channels from the K-band (water vapor absorption lines) and the V-band (temperature-sensitive
200 oxygen absorption lines) were selected. Figure 3 presents scatterplots for channel 1 (K-band) and channel 13 (V-band) of HATPRO, as well as channel 1 (K-band) and channel 14 (V-band) of MP3000A. Results for the remaining channels are shown in Figures A1 and A2. For HATPRO, more than 6,000 samples are analyzed. The O–B biases are 1.25 K for channel 1 and 2.14 K for channel 13, with standard deviations (STD) of 3.35 K and 2.82 K, respectively. Additionally, the scatter distribution for channel 13 is not centered, showing a cluster shifted to the right of the diagonal (Fig. 3b). For MP3000A, more than 2,000
205 samples are analyzed, with O–B biases of 3.06 K for channel 1 and –0.54 K for channel 14. The O–B STDs are 3.94 K and 3.08 K, respectively. Similar to the results for HATPRO channel 13 (V-band), the scatter points for MP3000A channel 14 (V-band) also shows a cluster offset from the diagonal, but to the left (Fig. 3d). Based on these results, significant O–B biases are detected in GMWR observations, with their characteristics varying across different sensors and channels. However, the correlation coefficients between observed and simulated brightness temperatures are high, at least 0.95, suggesting that these
210 biases can be effectively corrected.




Figure 3: Scatter plot of observed brightness temperature (Tb) versus simulated Tb, based on samples collected from August to October 2023. The top and bottom rows correspond to the HATPRO and MP3000A sensors, and the left and right columns represent the K-band and V-band, respectively. Each panel displays the number of samples (num), the O–B mean (bias), O–B standard deviation (STD), and the correlation coefficient (r) between observed and simulated Tb.

To further analyze the O–B bias characteristics at each station and investigate the reasons for the band shifting from the diagonal (Fig. 3b and d), the statistics for each station are presented in Fig. 4. For HATPRO, the O–B bias varies among stations. Stations near complex topography (e.g., 56312, 56137, 56029, and 55664) exhibit notable positive O–B biases in channels 8 to 13 (Fig. 4a), leading to a rightward shift of the band relative to the diagonal (Fig. 3b). These positive biases may result from biases in the background field over the topographic region, the limited applicability of RTTOV-gb coefficients, or calibration issues in the observations. Overall, the O–B STD at each station for the K-band is larger than that for the V-band (Fig. 4b). The correlation coefficients between observed and simulated brightness temperatures are high across all channels (typically above 0.90), although they are slightly lower for channels 4 to 9. For MP3000A, station 57461 exhibits a negative O–B bias in channels 9 to 14 (Fig. 4d), contributing to the band shifting to the left of the diagonal (Fig. 3d). Similar to the results for HATPRO, the O–B STD in the K-band is generally larger than that in the V-band (Fig. 4e), and the correlation coefficients are also overall higher, typically exceeding 0.9 (Fig. 4f).




Figure 4: Statistics at each station based on samples collected from August to October 2023. O–B (a) bias and (b) standard deviations (STD) for HATPRO; (c) correlation coefficient (r) between observed and simulated brightness temperatures for HATPRO; (d–f) same as (a–c) but for MP3000A. Some stations did not provide observations for specific channels; the corresponding missing data are displayed in grey in the figure.

3.2 Bias correction

Based on the results above, noticeable O–B biases were observed, varying across sensors, channels, and the geographical locations of stations. It is essential to remove these biases before assimilation. Static bias correction (Harris and Kelly, 2001) and variational bias correction (Dee, 2005) are commonly used in radiance data assimilation. O–B bias is commonly represented using multiple linear regression with several predictors. Compared to linear estimates, nonlinear approaches show improved performance in reducing systematic biases (Zhang et al., 2023, 2024). Following these works, this study employed a machine learning–based bias correction scheme using the Random Forest (RF) technique (Breiman, 2001).

According to Yin et al. (2020), the predictors include the 1,000–300 hPa thickness, 200–50 hPa thickness, model surface skin temperature (TS) and total precipitable water (PW). Considering that GMWRs are sensitive to the lower atmosphere, the predictors also include 1,000–700 hPa thickness, 700–500 hPa thickness, 500–300 hPa thickness, 2m temperature (T2), 2m water vapor mixing ratio (Q2), 10m zonal wind (U10), 10m meridional wind (V10), and surface pressure (PS). Finally, latitude,

longitude, and observed brightness temperatures (Tb) are included as predictors due to their potential importance (Zhang et al., 2023). The O–B biases vary across sensors and channels. Therefore, a separate model is trained for each type of instrument and channel. Biases also vary across the geographical locations of stations, potentially influenced by the large-scale topography of the Tibetan Plateau. As predictors, 2 m temperature, surface pressure, and latitude and longitude are important for explaining these biases.

There are two types of parameters in machine learning models: model parameters and hyperparameters. Model parameters are initialized and updated during the learning process. Hyperparameters, on the other hand, cannot be directly estimated from data. They must be configured before training because they define the model's architecture. Building an optimal machine learning model requires exploring a range of possibilities. The process of determining the ideal model architecture and hyperparameter configuration is known as hyperparameter tuning, which is a key component of developing an effective machine learning model (Yang and Shami, 2020).

The RF model has four key hyperparameters: the number of trees in the forest (*n_estimators*), the maximum depth of the tree (*max_depth*), the minimum number of samples required to split an internal node (*min_samples_split*), and the minimum number of samples required to be at a leaf node (*min_samples_leaf*). These hyperparameters were tuned using scikit-learn’s GridSearchCV (Pedregosa et al., 2011) with 5-fold cross-validation, which exhaustively searches over a predefined range of hyperparameters, training and evaluating the model for each configuration. The flowchart illustrating the training and evaluation process of the bias correction (BC) model is shown in Fig. 5. The three-month sample dataset (described in Section 3.1) was randomly split into a training set (70 %) and a test set (30 %). During training, GridSearchCV constructed a large grid of possible hyperparameter configurations, iteratively trained and evaluated the model for each, and calculated a score. Finally, the optimized model was trained using the configuration with the highest score.




Figure 5: Flowchart of the training and evaluation for bias correction model

265 To investigate the impact of hyperparameters on model training time and performance, the fit time and score of the RF model under various hyperparameter settings were analyzed (figure not shown). During hyperparameter tuning, *n_estimators* was varied between 10 and 150, *max_depth* was adjusted from 5 to 30, *min_samples_leaf* was tested with values between 1 and 3, and *min_samples_split* was tuned in the range of 2 to 6. Overall, the fit time and score demonstrated a positive correlation. As the *min_samples_leaf* and *min_samples_split* parameters increased, both fit time and score decreased monotonically.

270 Conversely, increasing the *max_depth* and *n_estimators* parameters resulted in a monotonic increase in both fit time and score. Notably, *max_depth* had the most significant impact on the score, while *n_estimators* primarily affected the fit time. For *n_estimators*, the score increased logarithmically, while the fit time grew linearly. These findings suggest that selecting a moderately small value for the *n_estimators* parameter can achieve better results while reducing computational time.

275 Using the above bias correction (BC) model, the corrected O–B values are obtained by subtracting the predicted O–B from the original observations. The effectiveness of the BC model is assessed based on the probability density function (PDF) of the O–B distributions in the test set (see Fig.6 and Appendix B). Similar to Fig. 3, channel 1 (K-band) and channel 13 (V-band) of HATPRO, as well as channel 1 (K-band) and channel 14 (V-band) of MP3000A, are selected for detailed analysis (Fig. 6). For HATPRO channels 1 and 13, the biases (STDs) are 1.24 K (3.38 K) and 2.21 K (2.90 K), respectively. For MP3000A channels 1 and 14, the biases (STDs) are 3.00 K (3.89 K) and –0.64 K (3.08 K), respectively. The differences 280 between the test set (Fig.6) and the full dataset (Fig. 3) are negligible, with maximum differences in bias and STD of 0.10 K and 0.08 K, respectively, highlighting the strong representativeness of the test set. The original PDF generally exhibit a unimodal shape, although their peaks deviate from zero. Moreover, some channels display bimodal features, often manifested as secondary peaks superimposed on the primary distribution—an issue that may affect 3D-Var, which typically assumes the errors to follow an unbiased Gaussian distribution.

285 After applying the BC model, both O–B bias and STD are reduced, and the distribution becomes more sharply concentrated around zero, accompanied by an increase in kurtosis. For instance, in MP3000A channel 1, the bias (STD) decreases from 1.24 K (3.38 K) to 0.03 K (1.44 K), respectively, while the kurtosis markedly increases from 1.53 to 9.44. It is noted that the PDF of corrected O–B approximate an unbiased distribution, and the secondary peaks are effectively suppressed. These results 290 confirm that the proposed BC scheme effectively reduces both the bias and STD in the O–B statistics, transforming the PDF from a bimodal to a unimodal distribution. The bimodal feature in the O–B PDF corresponds to the two distinct clusters observed in the scatter plots shown in Fig. 3. Specifically, when one cluster is concentrated along the diagonal and the other shifts to the right, a peak forms on the positive x-axis of the O–B PDF, resulting in a bimodal distribution. From the O–B PDF (Fig. 6), both instruments exhibit a positive bias with a unimodal distribution in the K-band. In contrast, the V-band displays a bimodal distribution: the second peak appears on the right for HATPRO and on the left for MP3000A. These results are 295 consistent with the scatter plots shown in Fig. 3. After BC, the O–B PDF changes from a bimodal to a unimodal distribution, indicating that the two clusters have been effectively merged.





Figure 6: Probability density function (PDF) of the O-B distributions, based on a test set randomly selected from 30% of the three-month sample dataset collected from August to October 2023. The top and bottom rows correspond to the HATPRO and MP3000A sensors, respectively, while the left and right columns represent the K-band and V-band. Each panel displays the number of samples (num), the mean (bias), standard deviation (STD), skewness, and kurtosis of the distributions.

300

305

Figure 7 illustrates the O-B bias and STD for each HATPRO channel. Before BC, the O-B bias ranged from 0 to 2 K, with the bias in the K-band (particularly channels 4 to 7) smaller than that in the V-band. After BC, the bias for each channel is approximately 0 K. In terms of the O-B STD, values ranged from 2 to 4 K without BC, with channels 4 to 7 exhibiting smaller values compared to other channels. After BC, the STD of O-B oscillates between 0.5 and 1.5 K. The application of this BC model significantly reduced both the O-B bias and STD, with reductions of 0.83 K (97.1 %) and 1.63 K (64.6 %), respectively. Meanwhile, the corrected O-B distributions display Gaussian characteristics centered around zero, indicating effective removal of systematic O-B biases.

310 Diagnosing the contributions of each predictor is crucial. Figure 7c illustrates the feature importance of several predictors for HATPRO. This BC model normalized the feature importance scores so that their sum equals 1. A higher score reflects a stronger correlation between predictors and O–B biases. Observed brightness temperature, total precipitable water, and surface pressure are significant contributors to BC for the K-band (water vapor channel). For the V-band (temperature channel), observed brightness temperature, latitude, and surface pressure are the most influential predictors. The contributions of 315 atmospheric thickness predictors are smaller compared to the other predictors; however, the 1,000–700 hPa thickness predictor has a relatively larger contribution among them. This may be because GMWR primarily observes radiation from the lower atmosphere. Notably, surface pressure plays a critical role in BC for V-band, which may account for the positive bias in O–B observed at plateau stations (Fig. 4a).



320 **Figure 7: (a) Bias and (b) standard deviation (STD) of O–B, based on a test set randomly selected from 30% of the three-month sample dataset collected from August to October 2023. (c) Feature importance of the predictors used in the bias correction (BC) model. The shaded regions and solid lines represent the range and mean feature importance for the K-band and V-band, respectively.**

4 Direct assimilation of GMWR radiance observations

4.1 Assimilation impacts on initial condition

325 The performance of GMWR assimilation in the observation space was evaluated. Figure 8 summarizes the bias and STD of the O–B and observation minus analysis (O–A) statistics, aggregated over time and across different channels. The bias of O–A was reduced compared to O–B, particularly in the V-band. Specifically, for channel 11 of GMWR_1H (1 h assimilation

interval), O–B was -0.40 K and O–A was -0.13 K. When GMWR observations were assimilated, the simulated brightness temperatures became closer to the observations, resulting in smaller STD. Moreover, as the frequency of GMWR observation assimilation increases, the bias and STD of the O–B gradually converge closer to zero. For channel 3, the O–B STD in GMWR_6H, GMWR_3H, and GMWR_1H significantly decreased from 1.03 K, 0.92 K, and 0.56 K to O–A STD values of 0.36 K, 0.34 K, and 0.34 K, respectively. Although the differences in O–A are less noticeable, the improvement of O–B suggests that increasing the frequency in cycling assimilation accumulates the impact of the GMWRs, producing a higher-quality first-guess field for the final cycle. O–A statistics were also computed based on the initial fields from the CNTL experiment. The O–A bias in CNTL is slightly larger than that in the GMWR assimilation experiments, with a more noticeable difference in the V band. Regarding the STD, the CNTL experiment shows higher values across all channels, with the largest difference approaching 1 K. These results suggest that assimilating GMWR data improves the consistency between the initial fields and the observed brightness temperatures. The assimilation of GMWR observations effectively influences the brightness temperatures, demonstrating the successful processing of GMWR data by the 3DVAR system.




Figure 8: Verification of the initial conditions against GMWR observations, based on the ten-day assimilation experiment conducted from 13 to 22 October 2023. (a) Bias and (b) standard deviation (STD) of the observation minus background (O–B) and observation minus analysis (O–A) for the GMWR assimilation in the target region of Southwest China (blue box in Fig. 1).

The above evaluation demonstrates the successful implementation of the newly introduced GMWR radiance direct assimilation in WRFDA. However, compared to brightness temperature simulations, greater attention should be given to the model state variables in the initial field, as they directly influence subsequent model forecasts. To this end, radiosonde observations in the

target region of Southwest China were used to evaluate the impact of GMWR assimilation. The root-mean-square error (RMSE) was calculated, and the RMSE differences between CNTL and other assimilation experiments are shown in Fig. 9. Results indicate that assimilating GMWR radiances enhances low-level temperature and humidity fields, with higher assimilation 350 frequencies offering the potential for additional improvements. GMWR assimilation has a neutral impact on atmospheric temperature above 1 km AGL, where the RMSE difference is minimal. However, it positively impacts lower atmospheric temperature, with the RMSE for temperature decreasing below 1 km AGL. Specifically, the average RMSE improvements below 1 km are 3.67 %, 5.28 %, and 6.32 % for GMWR_6H, GMWR_3H, and GMWR_1H, respectively. This indicates that increasing assimilation frequency enhances observational impacts and further improves the initial field. The improvement 355 becomes more pronounced with decreasing altitude, with 100 m RMSE improvements of 0.10 K (6.25 %), 0.13 K (7.90 %), and 0.19 K (11.34 %) in GMWR_6H, GMWR_3H, and GMWR_1H, respectively. For the water vapor mixing ratio (QVAPOR), GMWR assimilation demonstrates a positive impact extending into the middle troposphere, with average RMSE reductions below 5 km of 2.30 %, 2.20 %, and 1.98 % for GMWR_6H, GMWR_3H, and GMWR_1H, respectively. The impact of GMWR assimilation and the effect of assimilation frequency are more pronounced in the lower atmosphere, with 360 average RMSE reductions below 300 m of 3.01 % for GMWR_1H, compared to 2.43 % for GMWR_6H and 2.05 % for GMWR_3H.

It is noted that the GMWR assimilation has negative impacts on the wind fields. The RMSE for zonal and meridional winds exhibits a slight negative effect when GMWR is assimilated, with meridional winds even showing an increase in RMSE. These negative impacts on the wind field caused by GMWR assimilation may be attributed to two factors: (1) When assimilating 365 observed brightness temperature, the adjoint model of the observation operator directly adjusts temperature and humidity to optimize the simulation, while changes in the wind field are indirectly driven by these adjustments through the background error covariance. (2) GMWR assimilation primarily improves the lower atmosphere, while changes in the upper atmosphere are also governed by the background error covariance. The static background error covariance used here is climatological and isotropic, which does not fully align with evolving weather conditions, potentially resulting in ineffective wind field 370 improvements.




Figure 9: Verification of the initial conditions against radiosonde observations, based on the ten-day assimilation experiment conducted from 13 to 22 October 2023. Root mean square error (RMSE) of (a) temperature, (b) water vapor mixing ratio (QVAPOR), (c) zonal wind, and (d) meridional wind in the target region of Southwest China (blue box in Fig. 1).

375 Based on the evaluation against radiosonde observations, the assimilation of GMWR data improves the initial fields of temperature and humidity, aligning them more closely with observations, particularly in the lower atmosphere. Additionally, the initial fields are validated against surface station observations, including measurements of 2m temperature, 2m humidity, and 10m wind (Fig. 10). The RMSE differences indicate that GMWR assimilation effectively enhances the 2m temperature and humidity fields. Under 6-hourly GMWR assimilation, the temperature RMSE generally increased on the southern side of
 380 the basin, whereas other regions showed a positive effect with reduced RMSE values. Moreover, the temperature RMSE reduction in these positively affected areas further improved as the assimilation frequency increased, with overall differences ranging from -0.008 K (-0.3 \%) to -0.099 K (-4.1 \%). For humidity, GMWR assimilation shows a negative impact on relative humidity (RH) at a 6 h assimilation frequency. However, the RMSE over the plateau decreases as the assimilation frequency increases, with the RMSE difference shifting from positive to negative. In the GMWR_1H experiment, the RMSE is reduced
 385 by 0.276 (1.3 \%).

Unlike the temperature and humidity RMSEs, the improvement in the wind field RMSE does not exhibit a distinct spatial pattern. Compared to the CNTL experiment, the RMSE differences for zonal wind are -0.005 m s^{-1} (-0.3 \%), -0.017 m s^{-1} (-1.0 \%), and -0.019 m s^{-1} (-1.2 \%) in GMWR_6H, GMWR_3H, and GMWR_1H, respectively. Similarly, the RMSE differences for meridional wind are -0.008 m s^{-1} (-0.5 \%), -0.011 m s^{-1} (-0.7 \%), and -0.009 m s^{-1} (-0.5 \%) in GMWR_6H, GMWR_3H, and GMWR_1H, respectively. While the changes in wind RMSE are relatively small, the results indicate that assimilating GMWR data improves the initial field, with higher assimilation frequencies offering potential for further enhancement.





Figure 10: Verification of the initial conditions against surface station observations, based on the ten-day assimilation experiment conducted from 13 to 22 October 2023. The percentage in RMSE differences (scatter) of temperature (T), relative humidity (RH), zonal wind (U), and meridional wind (V) in the target region of Southwest China (blue box in Fig. 1). The grey solid line represents the topography height (m).

4.2 Assimilation impacts on forecast field

After presenting the improvements in the initial condition, this section investigated the impact of GMWR assimilation on the 400 24 h forecasts. The time series of RMSE for the CNTL experiment and RMSE differences (assimilation experiments minus the CNTL experiment) against surface station observations for 2 m temperature, 2 m relative humidity, and 10 m wind fields are shown in Fig. 11. In the CNTL experiment, the RMSE of temperature and relative humidity initially decreases and then increases with lead time, while the RMSE of the wind field exhibits the opposite trend, increasing at first and then decreasing.

The mean RMSEs over the 24-hour forecast period are 2.32 K for temperature, 16.26% for relative humidity, 1.92 m s^{-1} for zonal wind, and 2.08 m s^{-1} for meridional wind. Regarding assimilation impacts, the RMSE reduction for temperature gradually decreases, approaching zero at a lead time of 6 hours, with higher assimilation frequency (GMWR_1H) achieving a greater RMSE reduction. Similar results are observed for relative humidity, where the RMSE reduction also decreases and approaches zero at a lead time of 12 hours. GMWR_1H consistently demonstrates the largest RMSE reduction for relative humidity. However, it should be noted that the direct assimilation of GMWR data caused a negative impact on relative humidity at a lead time of 12 hours. The degradation of wind fields (Fig. 9) and the model's inherent nonlinearity may be responsible. For the wind field, no increase in the RMSE difference with lead time was observed, as previously described. However, the RMSE differences between the assimilation experiments and the CNTL experiment remain overall negative, indicating that GMWR assimilation improves wind forecasts. Additionally, GMWR_1H demonstrates the largest RMSE reduction in meridional wind, suggesting that increasing the frequency of GMWR assimilation may lead to further improvements. The quantitative statistics are presented in Table 3. The temperature RMSE differences between GMWR_6H and CNTL are -0.012 , -0.005 , and -0.004 K for lead time of 1–6 hours, 1–12 hours, and 1–24 hours, respectively. This gradual decrease in RMSE differences with increasing forecast time is also observed in other experiments and variables, indicating a weakening of the positive impact of GMWR assimilation as the forecast period extends. When the impact of GMWR assimilation is most pronounced (at a lead time of 1–6 hours), the temperature RMSE differences range from -0.012 K in GMWR_6H to -0.014 K in GMWR_3H, and -0.019 K in GMWR_1H. The temperature RMSE reduction increases with the frequency of GMWR assimilation, a trend also observed in relative humidity and wind, suggesting that increasing the assimilation frequency can further improve the short-term forecasts. Although these differences are small, the results reflect the potential for improved model forecasts with GMWR assimilation.



425 **Figure 11: Verification of the forecast against surface station observations, based on the ten-day assimilation experiment conducted from 13 to 22 October 2023. RMSE (black line) for the CNTL experiment and RMSE differences (colored lines) between the assimilation experiments and the CNTL experiment for (a) temperature, (b) relative humidity, (c) zonal wind, and (d) meridional wind.**

430

Table 3 RMSE difference against surface station observations.

EXP	Lead time (hour)	Temperature (K)	Relative Humidity (%)	Zonal wind (m s^{-1})	Meridional wind (m s^{-1})
GMWR_6H	1–6	-0.012	-0.046	-0.006	-0.004
GMWR_6H	1–12	-0.005	-0.027	-0.003	-0.006


minus CNTL	1–24	−0.004	0.046	−0.003	−0.003
GMWR_3	1–6	−0.014	−0.046	−0.005	−0.005
H	1–12	−0.010	−0.036	−0.006	−0.007
minus CNTL	1–24	−0.003	0.072	−0.005	−0.003
GMWR_1	1–6	−0.019	−0.127	−0.007	−0.013
H	1–12	−0.012	−0.087	−0.006	−0.013
minus CNTL	1–24	−0.005	0.065	−0.006	−0.009

435

Verification against surface station observations indicated that assimilating GMWR radiances improves near-surface forecasts, with higher assimilation frequencies offering potential for further enhancement. To further examine the impact of GMWR assimilation, Fig. 12 presents the forecast verification against radiosonde measurements. Unlike the RMSE differences in the initial condition (Fig. 9), the GMWR assimilation did not reduce RMSE for temperature or water vapor mixing ratio, indicating

440 a neutral impact of GMWR assimilation on forecasts. Similarly, the wind field verification results did not show noticeable improvements with GMWR assimilation. While the RMSE of zonal wind was reduced in the GMWR_1H experiment, the RMSE differences for the wind field in other experiments were close to or greater than zero, suggesting a neutral to slightly negative impact of GMWR assimilation on wind forecasting. According to the verifications against radiosonde data, limited improvements were found in the forecasts through GMWR assimilation. The limited improvement shown in this figure could

445 be related to the relatively long forecast lead times (12 and 24 hours), during which model errors tend to accumulate and weaken the benefits of improved initial conditions from GMWR assimilation. Verification against surface station observations indicates that the improvements were primarily confined to the first few hours, particularly for temperature and humidity. After 12 hours, the impact declined noticeably, with some cases even exhibiting negative effects (Fig. 11).



450 **Figure 12: Same as Fig. 9, but for forecast at lead time of 12 and 24 hours.**

To further explore the role of GMWR assimilation in precipitation forecasting, the fractions skill score (FSS) of 3 h accumulated precipitation forecasts was calculated. The radius of influence for the FSS was set to 18 km, equivalent to six times the grid spacing (Ha and Snyder, 2014; Zheng et al., 2024). Figure 13 presents the time series of FSS for the CNTL

455 experiment and FSS differences (assimilation experiments minus the CNTL experiment). The assimilation experiments were conducted during a period with a high frequency of clear-sky observations. Cloud cover and precipitation were sparse over the 10-day period, resulting in the absence of frequent heavy rainfall events. Consequently, the FSS was calculated using small precipitation thresholds. In the CNTL experiment, the FSS for 3 h accumulated precipitation shows an initial decline followed by a subsequent increase with lead time, with relatively low FSS values observed around the 9 h forecast period. Moreover,

460 the FSS generally decreases as the precipitation threshold increases. The time mean FSS values are 0.47, 0.45, 0.42, and 0.39 for thresholds of 3 mm, 4mm, 5mm, and 6 mm, respectively. Regarding the role of GMWR assimilation in precipitation forecasting, the results indicate that assimilating GMWR radiances enhances precipitation forecasts, with FSS differences increasing progressively at higher precipitation thresholds. Additionally, increasing assimilation frequency shows the potential to further enhance forecast performance. When assimilating GMWR data at a 1 h frequency, the time-averaged FSS

465 improvements for 3 h accumulated precipitation are 0.02 (3.9 %) for the 3 mm threshold, 0.02 (4.7 %) for 4 mm threshold, 0.03 (7.3 %) for 5 mm threshold, and 0.04 (10.2 %) for 6 mm threshold precipitation. For 3 h accumulated precipitation with a threshold of 6 mm, the time-averaged FSS improvements are 0.01, 0.02, and 0.03 for GMWR_6H, GMWR_3H, and GMWR_1H, respectively. These findings are consistent with the above verification against radiosonde and surface station observations, suggesting that GMWR assimilation can improve forecasts and that higher-frequency assimilation leads to

470 further enhancements.




Figure 13: The time series of FSS (black line) for CNTL experiment and FSS differences (colored lines) between the assimilation experiments and the CNTL experiment. These experiments were conducted from 13 to 22 October 2023. The FSS was calculated for 3 h accumulated precipitation for thresholds of (a) 3 mm, (b) 4 mm, (c) 5 mm, and (d) 6 mm.

475

5 Conclusions and Discussion

To investigate the impact of directly assimilating GMWRs in Southwest China, a GMWR assimilation module has been built in WRFDA-4.5, where RTTOV-gb is used as the observation operator. Based on this module, a three-month sample of O–B statistics was calculated to evaluate the bias and develop a BC model. Furthermore, 10-day assimilation experiments (Table 2) 480 were conducted using this GMWR assimilation module and BC model to investigate the impact of direct GMWR assimilation and the effects of assimilation frequency. The main findings are as follows:

1. Based on three months of hourly samples, noticeable O–B biases were observed, varying across sensors, channels, and geographical locations. The machine learning-based bias correction scheme, employing an RF model, effectively reduced these O–B systematic biases. After applying this BC model, both the bias and STD of the O–B were substantially reduced. 485 Specifically, the bias and STD decreased by 0.83 K (97.1 %) and 1.63 K (64.6 %), respectively. For some channels, the original O–B distribution exhibited a bimodal pattern, which was transformed into a unimodal distribution after BC. The corrected O–B distributions exhibited Gaussian characteristics centered around zero.

2.Assimilating GMWR enhances the accuracy of initial atmospheric conditions, with higher assimilation frequencies
490 amplifying the positive impact, particularly for temperature and humidity in the lower atmosphere. Evaluation against
radiosonde observations shows that the temperature RMSE below 1 km AGL decreases by 3.67 % to 6.32 % as the assimilation
frequency increases from 6 h to 1 h. For the water vapor mixing ratio, positive impacts extend up to 5 km AGL, with average
RMSE improvements ranging from 1.98 % to 2.30 %. Verification against surface station observations further supports these
findings, indicating that the RMSE for 2 m temperature decreases by up to 4.1 %, while the RMSE for 2 m relative humidity
495 decreases by up to 1.3 % at the 1 h assimilation frequency.

3.The assimilation of GMWR observations leads to improvements in forecasts, and increasing assimilation frequencies has the
potential to get further improvement. In the first 6 hours of the forecast, the temperature RMSE decrease by 0.012 K, 0.014 K,
and 0.019 K with 6 h, 3 h and 1 h assimilation frequency, respectively. Similar trends are observed for relative humidity, the
experiment with 1 h GMWR assimilation frequency showing the largest decrease in RMSE. GMWR assimilation also
500 improves precipitation forecasts, with further enhancements seen as assimilation frequency increases. For 1 h GMWR
assimilation, time-averaged FSS improvements reach 0.02 for both the 3 mm and 4mm, 0.03 for 5 mm, and 0.04 for 6 mm
thresholds.

In the three-month O–B statistics, the STD in the K-band is larger than that in the V-band, consistent with Vural et al. (2024)
and Cao et al. (2023). The O–B bias varies across sensors, channels, and geographical locations, with a notable positive bias
505 observed at high-altitude stations. This positive bias is potentially caused by large-scale topographical effects on the Tibetan
Plateau. In this region, model simulations may contain errors, and RTTOV-gb coefficients may be inapplicable. The RTTOV-
gb coefficients are based on global atmospheric profiles, which may differ significantly from the climatic conditions of plateau
regions, potentially affecting simulation accuracy.

A machine learning-based BC scheme using the RF technique was developed. The number and depth of trees are critical
510 hyperparameters that must be predetermined. Training time increases approximately linearly with the number of trees, while
performance exhibits a logarithmic-like saturation trend. In term of tree depth, both training time and performance increase
approximately logarithmically with depth. Thus, selecting a modest number (*n_estimators*) and depth (*max_depth*) of trees,
such as 50 and 15, can balance efficiency and accuracy. Feature importance analysis for BC predictors revealed observed
515 brightness temperature, atmospheric precipitable water, and surface pressure as key factors for correcting biases. The
importance of brightness temperatures aligns with findings in satellite data bias correction (Liu et al., 2022; Zhang et al., 2023).
Atmospheric precipitable water is essential for the K-band, a humidity-sensitive channel. Surface pressure plays a key role in
temperature channels, thereby accounting for the positive bias observed in plateau regions. Although atmospheric thickness
predictors contributed less overall, the 1,000–700 hPa thickness was relatively significant, likely due to GMWRs primarily
sensing radiation from the lower atmosphere.

520 The machine learning-based BC scheme effectively mitigated the bimodal distribution and systematic errors in the O–B
statistics. To assess its impact on the initial and forecast fields, a parallel experiment without bias correction was conducted,
based on the 1-hour assimilation interval experiment (GMWR_1H). For the initial fields, as verified against radiosonde

observations, the experiment without BC yielded only minor improvements in temperature and even degraded the water vapor field. As for the forecast fields, verification against surface station observations showed that the absence of BC led to a noticeable degradation in the forecast accuracy of 2 m temperature and relative humidity. These findings indicate that the machine learning-based BC scheme had a beneficial impact on both the initial conditions and the subsequent forecasts. Nevertheless, despite its demonstrated effectiveness, the scheme is subject to several limitations. Relying on offline O–B statistics, it implicitly assumes that all biases originate from the observations—an assumption that may not always hold and may, in some instances, mask model biases (Auligné et al., 2007; Eyre, 2016). Moreover, the offline scheme lacks anchoring observations, rendering the analysis fields more susceptible to model bias. Future efforts should consider bias correction strategies based on unbiased reference observations or adopt a constrained correction scheme, such as the constrained adaptive bias correction (Han and Bormann, 2016).

In this study, direct assimilation of GMWR radiance enhances both the initial conditions and the forecasts, showing a potential in improving ABL and precipitation simulations. Although the assimilation of GMWR radiance yields slight improvements in the forecast wind fields (Fig. 11), it exerts an overall negative impact on the wind fields in the initial conditions (Fig. 9). This discrepancy stems from the use of distinct observational datasets: the initial conditions are verified against radiosonde observations, while the forecasts are evaluated using surface station data. It should be noted that assimilating GMWR improves the wind field below 500 m AGL in the initial conditions. This improvement is consistent with the verification of the forecast, which demonstrate enhancements in the 2-meter wind fields. Regarding the degradation for wind field above 500m AGL, the background error covariance may contribute to this negative impact. On the one hand, it determines the response of the wind fields to temperature and humidity adjustments made by RTTOV-gb. On the other hand, since RTTOV-gb's adjustments are primarily concentrated in the ABL, the response above the ABL may be propagated through the background error covariance. As an initial study primarily focused on the direct variational assimilation of GMWR data, it is admitted that this study has some limitations. The GMWR assimilation was implemented using 3DVAR, based on RTTOV-gb and WRFDA, and only static background-error covariances were employed in this study. The background error covariance matrix plays an important role in variational data assimilation, but this type of covariance is climatological, spatially homogeneous, and isotropic. This may limit the impact of GMWR assimilation, and flow-dependent error covariances should be considered in future work.

Moreover, only clear-sky GMWRs were assimilated in this study. Since precipitation processes are often accompanied by extensive cloud cover, few clear-sky GMWRs were available. To better explore the potential of GMWR assimilation, experiments were conducted during periods with abundant clear-sky GMWRs (e.g., a ten-day period in October 2023), which coincided with minimal heavy precipitation. Studies on satellite all-sky assimilation have shown that incorporating cloud- and precipitation-affected data improves forecasts (Ma et al., 2022; Xian et al., 2019), highlighting the need for future research on all-sky assimilation of GMWRs. Under such conditions, assimilation experiments could be conducted during a different or longer period, given that assimilated GMWR observations would be relatively more abundant. It is noted that GMWRs exhibit higher sensitivity and provide more valuable observations of the lower troposphere and planetary boundary layer compared to satellite-based microwave radiometers (Shi et al., 2023). Building on this study, future research could explore the joint direct

assimilation of satellite-based and ground-based microwave radiometers. By leveraging their complementary observational capabilities, this approach has the potential to further enhance the accuracy of atmospheric analysis and improve forecasting across multiple layers of the atmosphere.

560 **Appendix A: Observed versus simulated Tb**





Figure A1: Scatter plot of observed brightness temperature (Tb) versus simulated Tb for HATPRO. Same as Fig. 3 but for additional channels.



565 Figure A2: Scatter plot of observed brightness temperature (Tb) versus simulated Tb for MP3000A. Same as Fig. 3 but for
additional channels.

Appendix B: PDF distributions of O-B




Figure B1: Probability density function (PDF) of the O-B distributions for HATPRO. Same as Fig. 6 but for additional




Figure B2: Probability density functions (PDF) of the O–B distributions for MP3000A. Same as Fig. 6 but for additional channels.

575 **Code availability**

The RTTOV-gb v1.0, WRF v4.5, WRFDA v4.5, along with the code for developing a direct assimilation module for GMWR radiances and training a machine learning-based GMWR bias correction model, are available on Zenodo (<https://doi.org/10.5281/zenodo.16731169>; Zheng et al., 2025a)

Data availability

580 The GMWR data and precipitation analysis product are provided by the Chinese Meteorological Administration and can be obtained via request from <https://www.cma.gov.cn/en/>. The AGRI CLM used are available at <https://satellite.nsrmc.org.cn/portalsite/default.aspx?currentculture=en-US>. The NCEP FNL data used are available at <https://rda.ucar.edu/datasets/d083003/>. The assimilated GTS data are available at <https://rda.ucar.edu/datasets/d337000/>. The model outputs for the single-observation assimilation experiment and the three-month sample dataset for the “Machine learning 585 based bias correction for GMWR” section is available on Zenodo (<https://doi.org/10.5281/zenodo.14586346>; Zheng et al., 2025b)

Author contribution

WS conceived the idea and designed the research. QZ performed the research and wrote the first draft of the manuscript. All authors discussed the results and contributed to writing and revisions.

590 **Competing interests**

The authors declare no conflicts of interest or competing financial interests.

Acknowledgements

595 The authors gratefully acknowledge Dr. Domenico Cimini for his help in usage of RTTOV-gb. They also thank the Numerical Weather Prediction Satellite Application Facility (NWP SAF) for providing the source code of RTTOV-gb v1.0, and the National Center for Atmospheric Research (NCAR) for providing the source code of WRF v4.5 and WRFDA v4.5. Additionally, the authors appreciate the National Centers for Environmental Prediction (NCEP) for providing the FNL data,

the China Meteorological Administration (CMA) for supplying the GMWR data and precipitation analysis product, and the National Supercomputing Center in Chengdu for their computational support.

Financial support

600 This study was jointly supported by the National Key Research and Development Program of China (2023YFC3007504), the National Natural Science Foundation of China (42475013 and U2442214), Xizang Science and Technology Major Program (XZ202402ZD0006), Sichuan Science and Technology Program (2024YFFK0110), Southwest Sichuan (Ya'an) Rainstorm Laboratory (CXBYSYSZD202401).

References

- 605 Auligné, T., McNally, A. P., and Dee, D. P.: Adaptive bias correction for satellite data in a numerical weather prediction system, *Quarterly Journal of the Royal Meteorological Society*, 133, 631–642, <https://doi.org/10.1002/qj.56>, 2007.
- Barker, D., Huang, X.-Y., Liu, Z., Auligné, T., Zhang, X., Rugg, S., Ajjaji, R., Bourgeois, A., Bray, J., Chen, Y., Demirtas, M., Guo, Y.-R., Henderson, T., Huang, W., Lin, H.-C., Michalakes, J., Rizvi, S., and Zhang, X.: The Weather Research and Forecasting Model's Community Variational/Ensemble Data Assimilation System: WRFDA, *Bulletin of the American Meteorological Society*, 93, 831–843, <https://doi.org/10.1175/BAMS-D-11-00167.1>, 2012.
- 610 Bauer, P., Geer, A. J., Lopez, P., and Salmond, D.: Direct 4D-Var assimilation of all-sky radiances. Part I: Implementation, *Quarterly Journal of the Royal Meteorological Society*, 136, 1868–1885, <https://doi.org/10.1002/qj.659>, 2010.
- Bauer, P., Thorpe, A., and Brunet, G.: The quiet revolution of numerical weather prediction, *Nature*, 525, 47–55, <https://doi.org/10.1038/nature14956>, 2015.
- 615 Bormann, N., H. Lawrence and J. Farnan, 2019: Global observing system experiments in the ECMWF assimilation system, ECMWF Tech. Memo, 839, 23pp.
- Breiman, L.: Random Forests, *Machine Learning*, 45, 5–32, <https://doi.org/10.1023/A:1010933404324>, 2001.
- Candy, B. and Migliorini, S.: The assimilation of microwave humidity sounder observations in all-sky conditions, *Quarterly Journal of the Royal Meteorological Society*, 147, 3049–3066, <https://doi.org/10.1002/qj.4115>, 2021.
- 620 Cao, Y., Shi, B., Zhao, X., Yang, T., and Min, J.: Direct Assimilation of Ground-Based Microwave Radiometer Clear-Sky Radiance Data and Its Impact on the Forecast of Heavy Rainfall, *Remote Sensing*, 15, 4314, <https://doi.org/10.3390/rs15174314>, 2023.
- Carrier, M. J., Zou, X., and Lapenta, W. M.: Comparing the Vertical Structures of Weighting Functions and Adjoint Sensitivity of Radiance and Verifying Mesoscale Forecasts Using AIRS Radiance Observations, *Monthly Weather Review*, 136, 1327–1348, <https://doi.org/10.1175/2007MWR2057.1>, 2008.
- 625 Caumont, O., Cimini, D., Löhnert, U., Alados-Arboledas, L., Bleisch, R., Buffa, F., Ferrario, M. E., Haefele, A., Huet, T., Madonna, F., and Pace, G.: Assimilation of humidity and temperature observations retrieved from ground-based microwave

- radiometers into a convective-scale NWP model, *Quarterly Journal of the Royal Meteorological Society*, 142, 2692–2704, <https://doi.org/10.1002/qj.2860>, 2016.
- 630 Chen, F. and Dudhia, J.: Coupling an Advanced Land Surface–Hydrology Model with the Penn State–NCAR MM5 Modeling System. Part I: Model Implementation and Sensitivity, *Monthly Weather Review*, 129, 569–585, [https://doi.org/10.1175/1520-0493\(2001\)129<0569:CAALSH>2.0.CO;2](https://doi.org/10.1175/1520-0493(2001)129<0569:CAALSH>2.0.CO;2), 2001.
- 635 Cimini, D., Hocking, J., De Angelis, F., Cersosimo, A., Di Paola, F., Gallucci, D., Gentile, S., Geraldi, E., Larosa, S., Nilo, S., Romano, F., Ricciardelli, E., Ripepi, E., Viggiano, M., Luini, L., Riva, C., Marzano, F. S., Martinet, P., Song, Y. Y., Ahn, M. H., and Rosenkranz, P. W.: RTTOV-gb v1.0 – updates on sensors, absorption models, uncertainty, and availability, *Geoscientific Model Development*, 12, 1833–1845, <https://doi.org/10.5194/gmd-12-1833-2019>, 2019.
- Cui, X., Z. Yao, Z. Zhao, Y. Zhai, Z. Sun, W. Cheng, and C. Gu: Use of Double Channel Differences for Reducing the Surface Emissivity Dependence of Microwave Atmospheric Temperature and Humidity Retrievals. *Earth and Space Science*, 7, e2019EA000854, <https://doi.org/10.1029/2019EA000854>, 2020.
- 640 De Angelis, F., Cimini, D., Hocking, J., Martinet, P., and Kneifel, S.: RTTOV-gb – adapting the fast radiative transfer model RTTOV for the assimilation of ground-based microwave radiometer observations, *Geoscientific Model Development*, 9, 2721–2739, <https://doi.org/10.5194/gmd-9-2721-2016>, 2016.
- 645 De Angelis, F., Cimini, D., Löhner, U., Caumont, O., Haefele, A., Pospichal, B., Martinet, P., Navas-Guzmán, F., Klein-Baltink, H., Dupont, J.-C., and Hocking, J.: Long-term observations minus background monitoring of ground-based brightness temperatures from a microwave radiometer network, *Atmospheric Measurement Techniques*, 10, 3947–3961, <https://doi.org/10.5194/amt-10-3947-2017>, 2017.
- Dee, D. P.: Bias and data assimilation, *Quarterly Journal of the Royal Meteorological Society*, 131, 3323–3343, <https://doi.org/10.1256/qj.05.137>, 2005.
- 650 Descombes, G., T. Auligné, F. Vandenberghe, D. M. Barker, and J. Barré: Generalized background error covariance matrix model (GEN_BE v2.0). *Geoscientific Model Development*, 8, 669–696, <https://doi.org/10.5194/gmd-8-669-2015>, 2015.
- Eyre, J. R.: Observation bias correction schemes in data assimilation systems: a theoretical study of some of their properties. *Quarterly Journal of the Royal Meteorological Society*, 142, 2284–2291, <https://doi.org/10.1002/qj.2819>, 2016.
- Eyre, J. R., English, S. J., and Forsythe, M.: Assimilation of satellite data in numerical weather prediction. Part I: The early years, *Quarterly Journal of the Royal Meteorological Society*, 146, 49–68, <https://doi.org/10.1002/qj.3654>, 2020.
- 655 Eyre, J. R., Bell, W., Cotton, J., English, S. J., Forsythe, M., Healy, S. B., and Pavelin, E. G.: Assimilation of satellite data in numerical weather prediction. Part II: Recent years, *Quarterly Journal of the Royal Meteorological Society*, 148, 521–556, <https://doi.org/10.1002/qj.4228>, 2022.
- 660 Geer, A. J., Bauer, P., and Lopez, P.: Lessons learnt from the operational 1D + 4D-Var assimilation of rain- and cloud-affected SSM/I observations at ECMWF, *Quarterly Journal of the Royal Meteorological Society*, 134, 1513–1525, <https://doi.org/10.1002/qj.304>, 2008.
- Geer, A. J., Bauer, P., and Lopez, P.: Direct 4D-Var assimilation of all-sky radiances. Part II: Assessment, *Quarterly Journal of the Royal Meteorological Society*, 136, 1886–1905, <https://doi.org/10.1002/qj.681>, 2010.

- 665 Geer, A. J., Baordo, F., Bormann, N., Chambon, P., English, S. J., Kazumori, M., Lawrence, H., Lean, P., Lonitz, K., and Lupu, C.: The growing impact of satellite observations sensitive to humidity, cloud and precipitation, *Quarterly Journal of the Royal Meteorological Society*, 143, 3189–3206, <https://doi.org/10.1002/qj.3172>, 2017.
- 670 Gustafsson, N., Janjić, T., Schraff, C., Leuenberger, D., Weissmann, M., Reich, H., Brousseau, P., Montmerle, T., Wattrelot, E., Bučánek, A., Mile, M., Hamdi, R., Lindskog, M., Barkmeijer, J., Dahlbom, M., Macpherson, B., Ballard, S., Inverarity, G., Carley, J., Alexander, C., Dowell, D., Liu, S., Ikuta, Y., and Fujita, T.: Survey of data assimilation methods for convective-scale numerical weather prediction at operational centres, *Quarterly Journal of the Royal Meteorological Society*, 144, 1218–1256, <https://doi.org/10.1002/qj.3179>, 2018.
- 675 Ha, S.-Y. and Snyder, C.: Influence of Surface Observations in Mesoscale Data Assimilation Using an Ensemble Kalman Filter, *Monthly Weather Review*, 142, 1489–1508, <https://doi.org/10.1175/MWR-D-13-00108.1>, 2014.
- 680 Harris, B. A. and Kelly, G.: A satellite radiance-bias correction scheme for data assimilation, *Quarterly Journal of the Royal Meteorological Society*, 127, 1453–1468, <https://doi.org/10.1002/qj.49712757418>, 2001.
- 685 Hartung, D. C., Otkin, J. A., Petersen, R. A., Turner, D. D., and Feltz, W. F.: Assimilation of Surface-Based Boundary Layer Profiler Observations during a Cool-Season Weather Event Using an Observing System Simulation Experiment. Part II: Forecast Assessment, *Monthly Weather Review*, 139, 2327–2346, <https://doi.org/10.1175/2011MWR3623.1>, 2011.
- 690 Han W. and Bormann N.: Constrained adaptive bias correction for satellite radiance assimilation in the ECMWF 4D-Var system, *ECMWF Technical Memoranda*, 783, <https://doi.org/10.21957/rex0omex>, 2016
- 695 He, W., Chen, H., and Li, J.: Influence of assimilating ground-based microwave radiometer data into the WRF model on precipitation, *Atmospheric and Oceanic Science Letters*, 13, 107–112, <https://doi.org/10.1080/16742834.2019.1709299>, 2020.
- 700 Hong, S.-Y., Noh, Y., and Dudhia, J.: A New Vertical Diffusion Package with an Explicit Treatment of Entrainment Processes, *Monthly Weather Review*, 134, 2318–2341, <https://doi.org/10.1175/MWR3199.1>, 2006.
- 705 Iacono, M. J., Delamere, J. S., Mlawer, E. J., Shephard, M. W., Clough, S. A., and Collins, W. D.: Radiative forcing by long-lived greenhouse gases: Calculations with the AER radiative transfer models, *Journal of Geophysical Research Atmospheres*, 113, <https://doi.org/10.1029/2008JD009944>, 2008.
- 710 Jiang, X., J. Shu, X. Wang, X. Huang, and Q. Wu: The Roles of Convection over the Western Maritime Continent and the Philippine Sea in Interannual Variability of Summer Rainfall over Southwest China. *Journal of Hydrometeorology*, 18, 2043–2056, <https://doi.org/10.1175/JHM-D-16-0292.1>, 2017.
- 715 Kim, M.-J., Jin, J., Akkraoui, A. E., McCarty, W., Todling, R., Gu, W., and Gelaro, R.: The Framework for Assimilating All-Sky GPM Microwave Imager Brightness Temperature Data in the NASA GEOS Data Assimilation System, *Monthly Weather Review*, 148, 2433–2455, <https://doi.org/10.1175/MWR-D-19-0100.1>, 2020.
- 720 Li, J., Jiang, X., Schiemann, R., Chen, H., Li, Y., and Heng, Z.: Prediction of the Diurnal Variation of Summertime Precipitation Over the Sichuan Basin by a Regional Model, *Journal of Geophysical Research: Atmospheres*, 128, e2021JD036247, <https://doi.org/10.1029/2021JD036247>, 2023.
- 725 Liu J. J., L. Xu, W. Cheng, B. Wang, X. Y. Gong, Z. R. Deng, Y. Y. Li, and D. Di, 2022. Bias Characteristics and Bias Correction of GIIRS Sounder onboard FY-4A Satellite for Data Assimilation. *Chinese Journal of Atmospheric Sciences (in Chinese)*, 46(2): 275–292. DOI: 10.3878/j.issn.1006-9895.2111.21034

- 700 Lin, H.-C., Sun, J., Weckwerth, T. M., Joseph, E., and Kay, J.: Assimilation of New York State Mesonet Surface and Profiler Data for the 21 June 2021 Convective Event, *Monthly Weather Review*, 151, 485–507, <https://doi.org/10.1175/MWR-D-22-0136.1>, 2023.
- Ma, S., Zhang, W., Cao, X., Zhao, Y., and Liu, B.: Assimilation of All-Sky Radiance from the FY-3 MWHS-2 with the Yinhe 4D-Var System, *Journal of Meteorological Research*, 36, 750–766, <https://doi.org/10.1007/s13351-022-1208-1>, 2022.
- 705 Martinet, P., D. Cimini, F. De Angelis, G. Canut, V. Unger, R. Guillot, D. Tzanos, and A. Paci: Combining ground-based microwave radiometer and the AROME convective scale model through 1DVAR retrievals in complex terrain: an Alpine valley case study. *Atmospheric Measurement Techniques*, 10, 3385–3402, <https://doi.org/10.5194/amt-10-3385-2017>, 2017.
- Min, M., Wu, C., Li, C., Liu, H., Xu, N., Wu, X., Chen, L., Wang, F., Sun, F., Qin, D., Wang, X., Li, B., Zheng, Z., Cao, G., and Dong, L.: Developing the science product algorithm testbed for Chinese next-generation geostationary meteorological 710 satellites: Fengyun-4 series, *J Meteorol Res*, 31, 708–719, <https://doi.org/10.1007/s13351-017-6161-z>, 2017.
- Moker, J. M., Castro, C. L., Arellano, A. F., Serra, Y. L., and Adams, D. K.: Convective-Permitting Hindcast Simulations during the North American Monsoon GPS Transect Experiment 2013: Establishing Baseline Model Performance without Data Assimilation, *Journal of Applied Meteorology and Climatology*, 57, 1683–1710, <https://doi.org/10.1175/JAMC-D-17-0136.1>, 2018.
- 715 Morrison, H., Thompson, G., and Tatarki, V.: Impact of Cloud Microphysics on the Development of Trailing Stratiform Precipitation in a Simulated Squall Line: Comparison of One- and Two-Moment Schemes, *Monthly Weather Review*, 137, 991–1007, <https://doi.org/10.1175/2008MWR2556.1>, 2009.
- Nie, Y., and J. Sun: Regional Persistent Extreme Precipitation Events over Southwest China under Different Low-Latitude Intraseasonal Oscillations during the Rainy Season. *Journal of Climate*, 36, 2873–2894, <https://doi.org/10.1175/JCLI-D-22-0310.1>, 2023.
- Otkin, J. A.: Clear and cloudy sky infrared brightness temperature assimilation using an ensemble Kalman filter, *Journal of Geophysical Research: Atmospheres*, 115, <https://doi.org/10.1029/2009JD013759>, 2010.
- 720 Otkin, J. A., Hartung, D. C., Turner, D. D., Petersen, R. A., Feltz, W. F., and Janzon, E.: Assimilation of Surface-Based Boundary Layer Profiler Observations during a Cool-Season Weather Event Using an Observing System Simulation Experiment. Part I: Analysis Impact, *Monthly Weather Review*, 139, 2309–2326, <https://doi.org/10.1175/2011MWR3622.1>, 2011.
- Parrish, D. F., and J. C. Derber: The National Meteorological Center's Spectral Statistical-Interpolation Analysis System. *Mon. Wea. Rev.*, 120, 1747–1763, [https://doi.org/10.1175/1520-0493\(1992\)120<1747:TNCSS>2.0.CO;2](https://doi.org/10.1175/1520-0493(1992)120<1747:TNCSS>2.0.CO;2). Pedregosa, F., and Coauthors, 2011: Scikit-learn: Machine Learning in Python. *J. Mach. Learn. Res.*, 12, 2825–2830, 1992.
- 730 Qi, Y., Fan, S., Mao, J., Li, B., Guo, C., and Zhang, S.: Impact of Assimilating Ground-Based Microwave Radiometer Data on the Precipitation Bifurcation Forecast: A Case Study in Beijing, *Atmosphere*, 12, 551, <https://doi.org/10.3390/atmos12050551>, 2021.
- Qi, Y., Fan, S., Li, B., Mao, J., and Lin, D.: Assimilation of Ground-Based Microwave Radiometer on Heavy Rainfall Forecast in Beijing, *Atmosphere*, 13, 74, <https://doi.org/10.3390/atmos13010074>, 2022.
- 735 Saunders, R., Hocking, J., Turner, E., Rayer, P., Rundle, D., Brunel, P., Vidot, J., Roquet, P., Matricardi, M., Geer, A., Bormann, N., and Lupu, C.: An update on the RTTOV fast radiative transfer model (currently at version 12), *Geoscientific Model Development*, 11, 2717–2737, <https://doi.org/10.5194/gmd-11-2717-2018>, 2018.

- 740 Shen, F., Shu, A., Liu, Z., Li, H., Jiang, L., Zhang, T., and Xu, D.: Assimilating FY-4A AGRI Radiances with a Channel-Sensitive Cloud Detection Scheme for the Analysis and Forecasting of Multiple Typhoons, *dqkxjz*, 41, 937–958, <https://doi.org/10.1007/s00376-023-3072-z>, 2024.
- 745 Shen, Y., Zhao, P., Pan, Y., and Yu, J.: A high spatiotemporal gauge-satellite merged precipitation analysis over China, *Journal of Geophysical Research: Atmospheres*, 119, 3063–3075, <https://doi.org/10.1002/2013JD020686>, 2014.
- 750 Shi, Y., Luo, Z., Chen, X., Zhang, Q., Liu, Y., and Liu, C.: Effects of joint assimilation of FY-4A AGRI and ground-based microwave radiometer on heavy rainfall prediction, *Front. Environ. Sci.*, 11, <https://doi.org/10.3389/fenvs.2023.1083517>, 2023.
- 755 Shi, Y.-N., J. Yang, W. Han, L. Han, J. Mao, W. Kan, and F. Weng: Development of a fast radiative transfer model for ground-based microwave radiometers (ARMS-gb v1.0): validation and comparison to RTTOV-gb. *Geoscientific Model Development*, 18, 1947–1964, <https://doi.org/10.5194/gmd-18-1947-2025>, 2025.
- 760 Shu, T.-T., Han, H.-M., Guo, L.-X., Lin, L.-K., Zhao, Z.-W., and Li, J.-M.: Profiling atmospheric temperature using temperature weighting functions, in: ISAPE2012, ISAPE2012, 1184–1187, <https://doi.org/10.1109/ISAPE.2012.6408989>, 2012.
- 765 Skamarock, C., Klemp, B., Dudhia, J., Gill, O., Liu, Z., Berner, J., Wang, W., Powers, G., Duda, G., Barker, D., and Huang, X.: A Description of the Advanced Research WRF Model Version 4.3, <https://doi.org/10.5065/1dfh-6p97>, 2021.
- 770 Su, J., Lü, H., Crow, W. T., Zhu, Y., and Cui, Y.: The Effect of Spatiotemporal Resolution Degradation on the Accuracy of IMERG Products over the Huai River Basin, <https://doi.org/10.1175/JHM-D-19-0158.1>, 2020.
- Sun, W. and Xu, Y.: Assimilation of FY-3D MWHS-2 Radiances with WRF Hybrid-3DVAR System for the Forecast of Heavy Rainfall Evolution Associated with Typhoon Ampil, <https://doi.org/10.1175/MWR-D-20-0127.1>, 2021.
- Vandenbergh, F. and Ware, R.: 4DIMENSIONAL VARIATIONAL ASSIMILATION OF GROUND-BASED MICROWAVE OBSERVATIONS DURING A WINTER FOG EVENT, 2002.
- 775 Vural, J., Merker, C., Löffler, M., Leuenberger, D., Schraff, C., Stiller, O., Schomburg, A., Knist, C., Haefele, A., and Hervo, M.: Improving the representation of the atmospheric boundary layer by direct assimilation of ground-based microwave radiometer observations, *Quarterly Journal of the Royal Meteorological Society*, 150, 1012–1028, <https://doi.org/10.1002/qj.4634>, 2024.
- 780 Wang, D., Wang, X., Liu, L., Wang, D., and Zeng, Z.: Urban Signatures in the Spatial Clustering of Precipitation Extremes over Mainland China, <https://doi.org/10.1175/JHM-D-20-0063.1>, 2021.
- Wei, J., Shi, Y., Ren, Y., Li, Q., Qiao, Z., Cao, J., Ayantobo, O. O., Yin, J., and Wang, G.: Application of Ground-Based Microwave Radiometer in Retrieving Meteorological Characteristics of Tibet Plateau, *Remote Sensing*, 13, 2527, <https://doi.org/10.3390/rs13132527>, 2021.
- 785 Xia, Y., Chen, J., Du, J., Zhi, X., Wang, J., and Li, X.: A Unified Scheme of Stochastic Physics and Bias Correction in an Ensemble Model to Reduce Both Random and Systematic Errors, <https://doi.org/10.1175/WAF-D-19-0032.1>, 2019.
- 790 Xian, Z., Chen, K., and Zhu, J.: All-Sky Assimilation of the MWHS-2 Observations and Evaluation the Impacts on the Analyses and Forecasts of Binary Typhoons, *Journal of Geophysical Research: Atmospheres*, 124, 6359–6378, <https://doi.org/10.1029/2018JD029658>, 2019.

- 775 Xu, L., Cheng, W., Deng, Z., Liu, J., Wang, B., Lu, B., Wang, S., and Dong, L.: Assimilation of the FY-4A AGRI Clear-Sky Radiance Data in a Regional Numerical Model and Its Impact on the Forecast of the “21·7” Henan Extremely Persistent Heavy Rainfall, *dqkxjz*, 40, 920–936, <https://doi.org/10.1007/s00376-022-1380-3>, 2023.
- Yang, L. and Shami, A.: On hyperparameter optimization of machine learning algorithms: Theory and practice, *Neurocomputing*, 415, 295–316, <https://doi.org/10.1016/j.neucom.2020.07.061>, 2020.
- 780 Yang, Z.-L., Zhao, L., He, Y., and Wang, B.: Perspectives for Tibetan Plateau data assimilation, *National Science Review*, 7, 495–499, <https://doi.org/10.1093/nsr/nwaa014>, 2020.
- Yin, R., Han, W., Gao, Z., and Di, D.: The evaluation of FY4A’s Geostationary Interferometric Infrared Sounder (GIIRS) long-wave temperature sounding channels using the GRAPES global 4D-Var, *Quarterly Journal of the Royal Meteorological Society*, 146, 1459–1476, <https://doi.org/10.1002/qj.3746>, 2020.
- 785 Yin, R., Han, W., Gao, Z., and Li, J.: Impact of High Temporal Resolution FY-4A Geostationary Interferometric Infrared Sounder (GIIRS) Radiance Measurements on Typhoon Forecasts: Maria (2018) Case With GRAPES Global 4D-Var Assimilation System, *Geophysical Research Letters*, 48, e2021GL093672, <https://doi.org/10.1029/2021GL093672>, 2021.
- Zhang, X., Xu, D., Li, X., and Shen, F.: Nonlinear Bias Correction of the FY-4A AGRI Infrared Radiance Data Based on the Random Forest, *Remote Sensing*, 15, 1809, <https://doi.org/10.3390/rs15071809>, 2023.
- 790 Zhang, X., D. Xu, F. Shen, and J. Min: Impacts of Offline Nonlinear Bias Correction Schemes Using the Machine Learning Technology on the All-Sky Assimilation of Cloud-Affected Infrared Radiances. *Journal of Advances in Modeling Earth Systems*, 16, e2024MS004281, <https://doi.org/10.1029/2024MS004281>, 2024.
- Zheng, Q., Sun, W., Li, J., Feng, Y., Heng, Z., and Jiang, X.: Impacts of Moisture Advection Scheme on Precipitation in the Steep Topography Region between the Tibetan Plateau and the Sichuan Basin, *Journal of Applied Meteorology and Climatology*, 63, 781–801, <https://doi.org/10.1175/JAMC-D-23-0111.1>, 2024.
- 795 Zheng, Q., Sun, W., Liu, Z., Mao, J., He, J., Li, J., and Jiang, X.: Codes related to the manuscript “Direct assimilation of ground-based microwave radiometer observations with machine learning bias correction based on developments of RTTOV-gb v1.0 and WRFDA v4.5”, Zenodo [code], <https://doi.org/10.5281/zenodo.16731169>, 2025a.
- Zheng, Q., Sun, W., Liu, Z., Mao, J., He, J., Li, J., and Jiang, X.: Data for manuscript “Direct assimilation of ground-based microwave radiometer observations with machine learning bias correction based on developments of RTTOV-gb v1.0 and WRFDA v4.5”, Zenodo [data set], <https://doi.org/10.5281/zenodo.14586346>, 2025b.