

Supplementary Materials for TECO-CNP Sv1.0: A coupled carbon-nitrogen-phosphorus model with data assimilation for subtropical forests

Fangxiu Wan, Chenyu Bian, Ensheng Weng, Yiqi Luo, Jianyang Xia

This PDF file includes:

Supplementary tables

Table S1. Carbon cycling observations from study site for model development.

Table S2. Nitrogen cycling observations from study site for model development.

Table S3. Phosphorus cycling observations from study site for model development.

Table S4. Reference codes and corresponding citations used in the main text.

Table S1 Carbon cycling observations from study site for model development.

Variables	Value	Unit	Sources	Measured Year
Evaluation				
Leaf C	0.22 ± 0.02	kg C m ⁻²	Zhou, 2020	2013-2017
Wood C	8.67 ± 0.56	kg C m ⁻²	Zhou, 2020	2013-2017
Root C	0.47 ± 0.03	kg kg ⁻¹	Measured data	2007
Root biomass	0.32 ± 0.11	kg m ⁻²	Zeng et al., 2008	2004-2005
Soil C	26.48 ± 10.99	kg C m ⁻²	Measured data	2017
Leaf C litterfall rate	0.23 ± 0.05	kg C m ⁻²	Measured data	2008-2018
Litterfall rate	0.44 ± 0.04	kg C m ⁻² a ⁻¹	Zhou, 2020	2011-2016
Soil respiration	0.99 ± 0.07	g C m ⁻² a ⁻¹	Zhou, 2020	2015-2017
Mean LAI	3.75 ± 0.15	m ² m ⁻²	Measured data	2021
Parametrization & Calibration				
LAI _{max}	5.2	m ² m ⁻²	Measured data	2021
LAI _{min}	2.8	m ² m ⁻²	Measured data	2021
Plant height	19.5±2.33	m	Measured data	2020
SLA	136.65 ± 15.87	cm ² g ⁻¹	Measured data	2020
V _{cmax}	26.2±0.97	μmol m ⁻² s ⁻¹	Measured data	2020
Fruit C	464.25 ± 19.56	g kg ⁻¹	Measured data	2007
J _{max}	38.01±1.75	μmol m ⁻² s ⁻¹	Measured data	2020

Table S2 Nitrogen cycling observations from study site for model development.

Variables	Value	Unit	Sources	Measured Year
Leaf N	6.1 ± 0.5	g C g N ⁻¹	Zhou, 2020	2013-2017
Wood N	31.42 ± 2.04	g C g N ⁻¹	Zhou, 2020	2013-2017
Root N	9.28 ± 2.18	g kg ⁻¹	Measured	2007
Fruit N	7.06 ± 0.25	g kg ⁻¹	Measured	2007
Soil N	1860.48 ± 550.90	gN m ⁻²	Measured	2017
Inorganic N	3.45 ± 0.59	gN m ⁻²	Zhou, 2020	2011-2013
Litterfall N rate	6.74 ± 0.68	g N m ⁻² a ⁻¹	Zhou, 2020	2013-2017
N net immobilization	13.14 ± 0.73	g m ² yr ⁻¹	Li, 2014	2014
Parametrization & Calibration				
CN _{leaf}	35.63 ± 4.23	g C gN ⁻¹	Measured	2020
CN _{wood}	275.99 ± 25.37	g C gN ⁻¹	Zhou, 2020	2013-2017
CN _{root}	50.56 ± 12.2	g C gN ⁻¹	Zhou, 2020	2013-2017
CN _{fruit}	65.76 ± 3.62	g C gN ⁻¹	Measured	2007
CN _{soil}	14.24 ± 7.26	g C gN ⁻¹	Measured	2017
N resorption rate	35.23 ± 19.42	%	Measured	2020

Table S3 Phosphorus cycling observations from study site for model development.

Variables	Value	Unit	Sources	Measured Year
Leaf P	0.19 ± 0.05	g C g P^{-1}	Measured data	2020
Leaf N:P	32.84 ± 9.12	g N g P^{-1}	Measured data	2020
Wood P	1.75 ± 0.11	g C g P^{-1}	Zhou, 2020	2013-2017
Root P	0.56 ± 0.17	g kg^{-1}	Measured data	2007
Fruit P	0.68 ± 0.21	g kg^{-1}	Measured data	2007
Soil P	181.59 ± 60.18	gP m^{-2}	Measured data	2017
Labile P	1.04 ± 0.56	gP m^{-2}	Measured data	2022
Secondary P	9.427 ± 0.521	gP m^{-2}	Wang 2022	2022
Litterfall P rate	0.79 ± 0.24	$\text{g P m}^{-2} \text{ a}^{-1}$	Zhou, 2020	2013-2017
Parametrization & Calibration				
CP_{leaf}	1144 ± 610.05	g C gP^{-1}	Measured	2020
CP_{wood}	4955.15 ± 448.27	g C gP^{-1}	Zhou, 2020	2013-2017
CP_{root}	837.84 ± 258.81	g C gP^{-1}	Zhou, 2020	2013-2017
CP_{fruit}	682.72 ± 212.79	g C gP^{-1}	Measured	2007
CP_{soil}	145.85 ± 77.46	g C gP^{-1}	Measured	2017
P resorption rate	54.53 ± 20.55	%	Measured data	2020

Table S4. Reference codes and corresponding citations used in the main text.

Reference Code	Citation
1	Wang et al., 2010
2	Xu et al., 2020
3	Yan, 2006
4	Zhu et al., 2016
5	Du et al., 2018
6	Goll et al., 2017
7	Schachtman et al., 1998
8	Olander & Vitousek, 2005
9	Bonan et al., 2014
10	Barracough and Tinker, 1981
11	Mollier et al., 2008