

Subauroral Contamination in POES/Metop TED Channels

Jan Maik Wissing¹, Olesya Yakovchuk², Stefan Bender³, and Christina Arras^{4,5}

¹German Aerospace Center (DLR), Institute for Solar-Terrestrial Physics, Neustrelitz, Germany

²Institute of Physics, University of Rostock, Rostock, Germany

³Instituto de Astrofísica de Andalucía, Granada, Spain

⁴GFZ Helmholtz-Centre for Geosciences, Potsdam, Germany

⁵Berlin University of Technology, Berlin, Germany

Correspondence: Jan Maik Wissing (jan.wissing@dlr.de)

Abstract. Particle measurements from the Polar Operational Environmental Satellites (POES) and the Meteorological Operational (Metop) satellite program, are widely used for various scientific applications. While most studies focus on the Medium Energy Proton and Electron Detector (MEPED), the low-energy (eV and keV) counterpart, the Total Energy Detector (TED), has received comparatively less attention. However, the recent rise in the altitudes considered in ionization and climate models has increased interest in low-energy particle measurements as inputs for atmospheric ionization models.

This study analyzes TED particle data (along with selected MEPED channels) from 2001 to 2025 and demonstrates that, in particular, the TED 0° proton channels –and, to a lesser extent, other TED channels– are contaminated by energetic electrons at $L < 6$. In some cases, the contaminated fluxes exceed typical auroral flux levels. The affected regions were cross-validated using auroral UV emissions and occurrences of GNSS derived $S4$ index to rule out the possibility that the observed fluxes correspond to real particle precipitation.

As correction approach, we provide a simple Kp- and channel-dependent latitude boundary that may serve as preliminary cut-off criterion for the contaminated regions. In a more advanced step, we identified the contamination characteristics of each particle channel on each satellite. The outcome is a list of problematic channels that should be neglected and a correction method based on background counts for the other channels. The corrected fluxes are in good agreement with UV emissions and the method is available in the auxiliary material.

1 Introduction

Accurate spaceborne particle measurements are essential for various scientific fields, particularly for modeling the atmospheric impact of precipitating particles. These impacts include direct ionization and the production of nitrogen oxides (NO_x) (Rusch et al., 1981) and hydroxyl (OH) (Solomon et al., 1981, 1983), which can lead to catalytic processes such as ozone depletion (Crutzen et al., 1975; Heath et al., 1977; McPeters and Jackman, 1985; Funke et al., 2011). In the case of TED protons, these effects are primarily confined to NO_x production in the lower thermosphere, between 130 and 150 km. Due to its link to ozone chemistry, particle precipitation may also influence the radiation budget and, consequently, atmospheric dynamics.

However, particle measurements are often biased by contamination from different energies or particle species, out-of-view
25 contamination, electric charging, and detector degradation. For the POES/Metop MEPED instrument, these issues have been
extensively discussed and quantified. For instance, MEPED electron channels are known to suffer from proton contamina-
tion (Evans and Greer, 2004b), which has been addressed using correction methods that subtract the spectral contribution of
interfering protons (Lam et al., 2010; Asikainen and Mursula, 2013; Peck et al., 2015; Nesse Tyssøy et al., 2016). Monte
Carlo simulations have also been employed to model the theoretical MEPED detector response to various particle species
30 and energies (Yando et al., 2011). These simulations indicate that MEPED proton detectors are affected by energetic electron
contamination, with the strongest impact on channel P1, decreasing in higher-energy channels. Another result of Yando et al.
(2011) is that the integral electron channels *MEPED* 0° e1, e2, and e3 (as well as their 90° equivalents) are differently sensitive
to high-energy electrons. In consequence a construction of differential channels by subtracting channels does not completely
remove the high energetic electron component (in particularly when subtracting e3). The MEPED electron channels might
35 also be affected by contamination from low-energy protons (see Yando et al., 2011). Additionally, proton detector degradation,
which becomes relevant after 2–3 years in orbit, has been quantified using orbit intersections of different satellites (Asikainen
et al., 2012). MEPED detector degradation has also been quantified by (Asikainen and Mursula, 2011; Ødegaard et al., 2016;
Sandanger et al., 2015).

In contrast, studies on the data quality of the POES/Metop TED instrument remain scarce. The instrument description
40 (Green, 2013) acknowledges energetic protons from the South Atlantic Anomaly (SAA) and energetic electrons from the
radiation belts as primary sources of TED background counts. A correction for TED energy flux based on background count
measurements is proposed in Green (2013), yet contamination from energetic electrons appears to persist. For example, Søråas
et al. (2018, Figures 4 and 5) observed a daily modulation in TED observations of sub-20 keV electrons between 50° and 62°
invariant latitude during the 17 March 2013 geomagnetic storm, which they attributed to the penetration of relativistic electrons.
45 Even more problematic is that no dedicated contamination analysis or correction has been conducted for the various TED
energy bands. Given the growing demand for accurate low-energy particle measurements to support high-altitude atmospheric
precipitation studies (e.g., Wissing and Kallenrode, 2009), this study aims to address that gap.

High-energy particle populations known to induce contamination are typically found within the radiation belts. For over 50
years, Earth's radiation belts have been understood to consist of two distinct zones—an inner and an outer belt-separated by the
50 slot region (van Allen, 1959, 1983). The inner radiation belt primarily contains electrons in the 100 keV range and energetic
protons exceeding 100 MeV, which are trapped by relatively strong magnetic fields. This belt typically spans $L = 1.2$ to $L = 2.5$
and remains relatively stable during quiet geomagnetic conditions (e.g., Selesnick et al., 2014). However, strong geomagnetic
storms can significantly alter the inner belt proton population (e.g., Selesnick et al., 2014), and in regions such as the SAA, the
inner boundary can extend to very low altitudes, where its downward extent is ultimately constrained by atmospheric losses.
55 The outer radiation belt, in contrast, consists of highly dynamic electron populations (0.1–10 MeV) modulated by geomagnetic
activity (e.g., Baker et al., 2013) with its intensity peaking between $L = 3.5$ and $L = 5.5$. In 2013, observations from the Van
Allen Probes revealed a long-lived relativistic electron storage ring—a third radiation belt—that persisted for four weeks (Baker
et al., 2013).

Contamination within the radiation belts has been documented for the POES/Metop MEPED instrument. For instance, Andersson et al. (2014) reported elevated MEPED electron count rates in the SAA that did not correspond to mesospheric OH enhancements, suggesting proton contamination as the likely cause. Similarly, Evans et al. (2008) identified anomalous increases in the POES $E > 16$ MeV proton channel at subauroral latitudes near the SAA, which they attributed to contamination from $E > 3$ MeV electrons.

During our analysis of TED data from POES and Metop satellites for the period 2001–2025, we identified very high fluxes –primarily in the proton bands– at subauroral latitudes. To investigate this phenomenon, we compared TED data with high-energy electron fluxes, TED background counts, and atmospheric measurements. Our findings indicate strong contamination, most likely from highly relativistic electrons, affecting the TED bands.

The objectives of this study are twofold: a) To establish a simple latitudinal cut-off to exclude the contamination-dominated region, thereby enabling the safe application of TED data. b) To develop a contamination correction method that can be applied to average TED fluxes, provided background counts are known, and that may also serve as an indicator of TED data quality in polar regions.

The structure of this paper is as follows:

Section 2 describes the data sets and processing methods. Section 3 presents the subauroral flux peak observations that motivated this investigation, along with their K_p dependence. Section 4 assesses the reliability of the measurements within the subauroral region. This section consists of four subsections: Subsection 4.1 provides an overview of the TED measurements. Subsection 4.2 explores energetic electron contamination as a potential cause of increased count rates at $L = 4$. Subsection 4.3 examines the differing effects of contamination on TED species and orientations. Subsection 4.4 compares observed particle fluxes with independent data sets, demonstrating that TED proton channels do not capture real flux in the subauroral region. Section 5 quantifies the contamination effect for each channel, with Subsection 5.1 proposing a correction method based on TED background counts. In Subsection 5.2 the correction method is applied to measurements of year 2013. Section 6 summarizes our findings. The Appendix discusses an issue in TED proton Level 1b netCDF (nc) files that, while unrelated to contamination, has not been documented elsewhere. The auxiliary data contains the suggested correction method.

2 Data Sets

This section describes the data sets and data processing methods of this study. Subsection 2.1 introduces the data handling of the total energy detector (TED) particle measurements. After that the Modified Apex coordinate system is described, followed by the K_p -binning which has been applied to part of the paper's data base. In the next subsections two independent data sets are introduced, which are compared to the particle fluxes. These data sets are the backscattered UV, see Subsection 2.4 and the S4 scintillation index, see Subsection 2.5. The aim is to falsify or verify particle precipitation in certain regions or during special periods with self-contained data.

90 2.1 Particle Data

The particle data originates from the Polar Operational Environmental Satellites (POES) and the Meteorological Operational satellite program (Metop). Both satellite series operate in sun-synchronous orbits at altitudes of approximately 820 km, with a revolution period of about 100 minutes and an inclination of approximately 98.5° . Although these satellites were initially placed in fixed local time orbits (morning-evening or day-night sector), their orbits have gradually drifted over time, enabling
95 near-complete magnetic local time coverage (while combining multiple years and satellites). Even though we do not think that it notably impacts our results, the coverage has a blind spot at 12 h MLT limited to southern hemisphere subauroral latitudes and another at 0 h MLT on the northern hemisphere. For details we refer to Figure 1 (lower right) in Yakovchuk and Wissing (2019).

The particle detectors aboard POES and Metop are identical and integrated within the Space Environment Monitor SEM-2
100 (Evans and Greer, 2006). The SEM-2 comprises two instruments: the Total Energy Detector (TED) for low-energy particles and the Medium Energy Proton and Electron Detector (MEPED) for medium-energy particles. TED is a cylindrical electrostatic analyzer (Fig. 1 of Green, 2013, shows a schematic of the TED configuration), while MEPED is a semiconductor detector with passive shielding.

This study focuses on the TED detector, which consists of a set of electrostatic analyzers (ESA). These analyzers utilize
105 curved plates with predefined radii, modulated by different voltages, to allow only particles of specific energies to pass through and reach a channeltron (multiplier) at the end of the curved plates. TED features eight ESAs that differentiate between low and high energies, along with two inlet openings at 0° (zenith) and 30° . The detector mounting and its orientation in relation to the satellite is shown in Evans and Greer (Fig. 2.1.1 of 2004a). Additionally, the curvature of the plates is oriented oppositely for protons and electrons. Particle flux measurements are conducted with the ESA plate voltage activated, while background
110 counts are recorded with the voltage deactivated. Although this study primarily focuses on the upward-looking (0°) detector, which is crucial for observing precipitating particles at high latitudes, it also demonstrates that the TED 30° detector is similarly affected. It is important to note that at low latitudes near the magnetic equator, the 0° detector predominantly measures trapped particles (for the MEPED instrument this is shown in Fig. 1 of Rodger et al., 2010).

This study utilizes data from POES satellites N15, N16, N17, N18, and N19, as well as Metop satellites M01, M02 and M03.
115 N16 is not used after 2006 as TED measurements are erroneous. In total the data covers the years 2001–2025. Details on the applied period and data averaging are given in the different sections.

Background counts are recorded but even though the TED data processing algorithm includes a background subtraction step (Green, 2013), this correction applies only to the total energy flux derived from a full energy sweep and not to individual bands. This holds true for both bin-format data prior to 2013 (Evans and Greer, 2004a) and netCDF-format data after 2013 (Green,
120 2013). The differential TED bands are -until now- always provided uncorrected. The primary distinction between raw data and processed TED bands in the processed-data files is a multiplication by the geometric factor. Note that whenever we are mentioning the TED particles in this paper, we refer to these differential channels.

Since our study investigates the contamination of the total energy detector, a few words on the instrument's shielding are in order. Unfortunately the housing is not well documented, except for the total weight of the instrument, its dimensions and images showing the aperture. We will discuss this aspect when looking at the background counts. One important aspect is that the detector is stacked and that the inner instruments seem to be better shielded than the outer ones.

Data from the MEPED e3 channels have been used in order to identify high energetic electron fluxes. These data have been assigned an error value if high-energy protons, which can cause contamination, are detected. This filtering was performed on a 2-second basis using the high-energy P7 proton channel from the omni-directional detector as a proxy, with measurements set to an error value when counts exceeded 2 per second. Consequently, MEPED electron data is excluded during strong proton events and within the South Atlantic Anomaly. The energy ranges for all particle channels are detailed in Table 1. The channel names may be abbreviated, e.g. *TED 0° proton band 11* as TED 0° p 11.

2.2 Coordinate System

This study employs the family of Magnetic Apex Coordinates (Van Zandt et al., 1972), which are derived from the International Geomagnetic Reference Field (IGRF) (Thébault et al., 2015). A key advantage of IGRF-based coordinate systems is that they account for the gradual movement of the geomagnetic poles, ensuring consistency in precipitation feature localization over extended time periods. This stability allows for multi-year spatial approximations of precipitation fluxes and facilitates the application of results to future datasets.

For ground-based locations and higher latitudes, Magnetic Apex Coordinates closely resemble the widely used Altitude Adjustment Corrected Geomagnetic (AACGM) coordinates. Specifically, poleward of 50°, the two systems are nearly identical (see Fig. 7 in Laundal and Richmond, 2017).

However, for positions above the ground, Magnetic Apex Coordinates provide a more accurate framework for describing charged particle behavior. These coordinates track the magnetic field lines from their footpoints upward, ensuring that charged particles remain on the same coordinates during their bounce motion. Thus, in Magnetic Apex Coordinates, the altitude of a satellite measurement does not alter the perceived location of a particle.

By default, the reference altitude for magnetic field-line footpoints is set to ground level. However, the Modified Apex Coordinates (Richmond, 1995) enable customization of the reference altitude. In this study, we use a reference altitude of 110 km, approximately where atmospheric interactions begin to dominate over magnetic field influences. Considering the grid resolution used, the effect of the reference altitude is negligible. Consequently, the coordinate system utilized is termed “Modified Apex 110 km Coordinates”, though we may refer to it simply as “APEX” in the text.

The conversion to Apex coordinates is done with an own C++ translation of the apex.f90 Fortran code as published on https://github.com/NCAR/apex_fortran/blob/master/apex.f90.

2.3 Kp-Binning of Particle Data

The planetary K-index (Kp) (Bartels et al., 1939) quantifies geomagnetic activity using a 3-hourly quasi-logarithmic scale ranging from 0 to 9. It is derived from observations at 13 geomagnetic observatories located between 44° and 60° in both

northern and southern geomagnetic latitudes. The Kp index is sensitive to multiple current systems, including the ring current, and thus provides a global measure of magnetospheric activity.

For this study, particle data has been categorized into seven Kp-level groups: 0–0.7, 1–1.7, 2–2.7, 3–3.7, 4–4.7, 5–5.7, and 6–9. This binning approach enables the examination of particle flux variations as a function of geomagnetic activity.

160 2.4 SSUSI Data Set

We use the electron energy flux data from the Special Sensor Ultraviolet Spectrographic Imager (SSUSI) instruments on board the Defense Meteorological Satellite Program (DMSP) Block-5D3 satellites F17 and F18 (Paxton et al., 1992, 2017, 2018). These satellites orbit at 850 km altitude in polar, sun-synchronous orbits, the equator crossing times of their ascending nodes are 17:34 LT (F17) and 20:00 LT (F18). The SSUSI detectors provide about 3000 km-wide spectrographic images of the
165 auroral zones with a 10 km×10 km pixel size at the nadir point. Of these spectra 5 UV channels are downlinked, including two colours for the LBH (Lyman-Birge-Hopfield) band emissions of N₂. From these two LBH colours the electron energies and energy fluxes are derived based on Strickland et al. (1983), additional discussion can be found in Knight et al. (2018). The electron energy flux data used here are provided within the Auroral-EDR (Environmental Data Record) data set at https://ssusi.jhuapl.edu/data_products (SSUSI, 2020). The data files contain the data point locations in geomagnetic latitude and
170 longitude in AACGM coordinates.

The data have been validated previously by comparing them to EISCAT ground-based observations (Bender et al., 2021). For the comparison here, for each DMSP/SSUSI orbit during 2013, the data points within 60°E ±4° geomagnetic longitude were selected. The electron energy fluxes within these sectors were then averaged in one-degree latitude bins to produce the time series as shown later in Sect. 4.4.2.

175 It should be mentioned that the SSUSI-derived fluxes are based on the hypothesis that the UV-aurora originates from a pure electron spectrum. There is an on-going discussion how pure proton or mixed auroral spectra would show up in the results of these pure-electron calculations. While model calculations of Knight et al. (2012, their Table 4) and Gelinias and Hecht (2016) agree that this retrieval technique should slightly over-estimate the atmospheric peak electron density under proton precipitation conditions, Knight (2021) did not see such an overestimation comparing SSUSI-derived atmospheric electron density with
180 ionosonde observations. Concerning our paper we may summarize that –even though the SSUSI algorithms threat all UV lines as effect of electron precipitation– particle precipitation from both, electrons and protons, should show up in the SSUSI data set.

2.5 S4 Index

The *S4 index* is a standard measure used to quantify ionospheric scintillation, referring to rapid fluctuations in the amplitude
185 and phase of radio signals as they pass through irregularities in the ionosphere. It is defined as the ratio of the standard deviation of signal intensity to the average signal intensity (Wu, 2020). For this study, we used *S4 index* values calculated from the signal-to-noise ratio (SNR) of GNSS radio occultation profiles. The S4 values, and thus the degree of disturbance of the profiles, refer to the altitude range of approximately 80 to 130 km above the Earth’s surface. We analyzed the atmPhs or conPhs data product




Figure 1. Spatial flux distribution of *TED* 0° proton band 14 for selected Kp levels (low Kp: left, medium Kp: right) averaged over the years 2001–2025.

provided by UCAR for nine different satellite missions (CHAMP, GRACE, Formosat-3/COSMIC-1, PAZ, Spire, TerraSAR-X, TanDEM-X, KompSat-5, Metop-B, Metop-C, and PlanetiQ) measured within the latitude belt between 55°S and 65°S from 2001 to 2024. Further, we concentrated on night-time measurements.

In total, 231,680 radio occultation profiles met our criteria. Using the method described by Arras and Wickert (2018), we calculated the maximum *S4 index* for each profile. Profiles that included a sporadic E signature were excluded from further analysis.

195 3 Observation of High Particle Fluxes in the Subauroral Zone

A distinct local maximum in TED proton flux, separate from both auroral precipitation and the South Atlantic Anomaly (SAA), was previously reported by Yakovchuk and Wissing (2019). This section examines which channels exhibit enhanced subauroral particle flux and how it correlates with geomagnetic activity.

Figure 1 presents the multi-year averaged spatial distribution of TED proton flux for the period 2001–2025, separated by geomagnetic activity levels (left: $Kp=0-0.7$, right: $Kp=4-4.7$). A localized increase in flux appears around 60° North and South APEX, with a stronger presence at longitudes that also intersect the SAA (most notably near 60°E APEX in the Southern Hemisphere).

At low Kp levels (Fig. 1, left), the flux distribution of all TED 0° proton channels exhibits a clear separation between the auroral oval and the subauroral peak, as indicated by the red dotted line. However, as geomagnetic activity increases (Fig. 1, right), the separation diminishes, with the auroral flux extending equatorward and merging with the subauroral peak. For Kp levels of 5 and above, the subauroral feature becomes indistinguishable from the auroral precipitation.




Figure 2. Cross-section at 60°E APEX for *TED 0° proton band 11* during varying levels of geomagnetic disturbance (based on years 2001–2025). The vertical lines indicate the minimum between the auroral and subauroral peaks. Note that the latitude axis is stitched together, as modified APEX latitudes at 110 km altitude are constrained to values greater than 18°.




Figure 3. The subauroral peak of the TED channels cannot be adequately corrected based on scaled background counts. Either it creates strong negative fluxes around the peak or the correction does not noteworthy change the fluxes.

The dependence of the subauroral peak on geomagnetic disturbance is further illustrated in Figure 2, which shows cross-sections of *TED 0° proton band 11* at 60°E APEX for different Kp levels. While the subauroral peak consistently appears between 56° and 60° APEX in both hemispheres, increasing geomagnetic activity broadens and intensifies the auroral oval, shifting its equatorial boundary closer to the subauroral feature. As a result, the two regions may merge at moderate Kp levels (e.g., Kp = 4–4.7 in Fig. 1, right, for *TED 0° proton band 14*). However, differences in latitudinal extent and flux intensity suggest that portions of the subauroral peak remain distinguishable at moderate Kp levels. At higher Kp values, the subauroral peak becomes fully embedded within the auroral precipitation, limiting its identification to periods of low geomagnetic disturbance.

It is worth noting that the northern subauroral peak is generally weaker than its southern counterpart and 60°E APEX does not intersect its strongest region. An hemispheric asymmetry similar to this is typically seen in radiation belt populations measured on the same altitude because the polar horns of the radiation belt reach lower altitudes in the southern hemisphere due to the weaker magnetic field strength. The different radiation belt particle fluxes are shown e.g. in Winant et al. (2023) who compare the van Allen probe measurements on POES/Metop altitudes against the AE-8 model (Vette, 1991).

Figure 3 shows the flux of a TED proton band against L-shell. We also added scaled background counts, with the scaling factor chosen large as possible but still preventing unphysical negative counts at L=2–3 and L=6. The peak at L=4 remains largely unchanged, indicating that a simple linear background correction cannot eliminate the subauroral maximum without compromising physical consistency. This suggests that the background-corrected total energy flux may exhibit similar issues, as seen in Søråas et al. (2018, their Figures 4 and 5).

	instrument	band	energy range	Separation of the contamination area at Kp				
				0–0.7	1–1.7	2–2.7	3–3.7	4–4.7
electrons	TED 0°	4	154–224 eV	-	-	-	-	-
		8	688–1000 eV	-	-	-	-	-
		11	2.115–3.075 keV	-	-	-	-	-
		14	6.503–9.457 keV	-	-	-	-	-
protons	TED 0°	4	154–224 eV	66°	64°	62°	60°	58°
		8	688–1000 eV	66°	64°	64°	62°	62°
		11	2.115–3.075 keV	66°	66°	64°	64°	62°
		14	6.503–9.457 keV	66°	64°	64°	62°	62°

Table 1. Channel and Kp combinations that exhibit enhanced subauroral flux, along with the location of the minimum between auroral and subauroral peaks, as identified visually from spatial maps such as Fig. 1. The latitude values apply to both the Northern and Southern Hemispheres.

225 Table 1 summarizes the TED 0° proton channels and Kp conditions under which enhanced subauroral fluxes were detected. The identification criterion required the presence of a local flux maximum exceeding the equatorial boundary of the auroral oval by a factor of at least 1.778 (equivalent to one-quarter of an order of magnitude in logarithmic flux levels) across different longitudes, ruling out statistical fluctuations. Enhanced subauroral flux is observed across all TED 0° proton channels, whereas no corresponding feature is detected in TED electron data.

230 To illustrate the potential impact of a misinterpreted subauroral peak, we estimate its contribution to atmospheric ionization. Wissing and Kallenrode (2009, see their Fig. 9, lower panel) reported that at 120 km altitude, auroral proton-induced ionization accounts for approximately 14% of total ionization, with electrons dominating the remaining contribution. If the observed subauroral TED flux is real, it would imply an additional ionization peak at this auroral proton level in the lower thermosphere (130–150 km). This highlights the importance of accurately characterizing the subauroral peak to refine atmospheric chemistry
235 models.

4 Is the Subauroral Peak a Real Flux or an Artifact from Contamination?

A critical question regarding the observed subauroral particle fluxes is the reliability of the measurements. As noted in the introduction, high particle fluxes are expected at low L-shells, particularly in the proton and electron radiation belts. Two primary hypotheses can explain the subauroral peak: a) The low-energy proton fluxes could represent a low-energy tail of the radiation belts. b) The observed fluxes result from contaminated detector counts due to high-energy proton or electron
240 contamination.

To investigate this, we analyze various particle measurements from the SEM-2 instrument.

4.1 Comparison of Onboard Particle Data




Figure 4. Particle fluxes and background sensor counts for the TED 0° (left panels) and the TED 30° detectors (right panels) in addition to the MEPED 0° or 90° e3 channel. The difference between 60°E and 160°E APEX is highlighted by the shaded area. Below L=6, the higher counts (or fluxes) consistently correspond to 60°E APEX longitude, while the lower values correspond to 160°E APEX. Elevated background counts in the $L \leq 2$ (SAA) region and in the L=3–6 range (subauroral peak) coincide with intense particle counts in TED proton channels and the MEPED 90° e3 channel.

Figure 4 presents the mean particle fluxes recorded by the TED 0° detectors (left panels) and the TED 30° detectors (right panels) for the year 2013 under geomagnetically quiet conditions ($K_p \leq 0.7$). The data are plotted against L-shell along two APEX longitudes: 60°E and 160°E. Below L=6, the upper boundaries of each color correspond to 60°E APEX, while the lower boundaries correspond to 160°E APEX. The year 2013 was chosen because NOAA's data format update allowed easy export of

TED background sensor counts per second (CPS). The figure also shows the MEPED e3 fluxes from the 0° or 90° instrument as indication for energetic electrons.

250 **General Trends in the Data:** Nearly all TED channels exhibit two local maxima: a) A peak at $L \leq 2$, associated with the South Atlantic Anomaly (SAA). b) A peak in the subauroral region ($L = 3-6$), with a maximum at $L \approx 4$ ($\sim 60^\circ$ N/S). This pattern is also reflected in the TED background counts (black for lower energy bands 1–8 and gray for higher energy bands 9–16).

The SAA and the Absence of Energetic Protons in the Subauroral Region: The SAA is well known for intense proton contamination, particularly affecting MEPED electron channels (Peck et al., 2015). Therefore we applied a filtering method for the MEPED electron channels, which uses the P6 channel to effectively remove heavily contaminated MEPED electron data from the SAA ($L \leq 2$). Indeed, all MEPED electron data of 60°E longitudinal cross section within the in the SAA are neglected.

However, the subauroral data remain unaffected by P6 filtering, which suggests that energetic protons are not responsible for the observed peak. This leaves two possible explanations: a) The TED proton fluxes represent a low-energy tail of the radiation belt. b) The measurements are contaminated by energetic electrons.

TED 0° Proton Fluxes: To gain further insight, we examine the flux characteristics in the subauroral region, starting with the 0° differential proton channels (Fig. 4, top left). All TED 0° proton channels exhibit a local maximum at $L=4$. The flux difference between 60°E APEX (peak) and 160°E APEX (minimum) roughly spans one order of magnitude across all TED 0° channels, albeit from different baseline levels. In contrast, background counts vary by only half an order of magnitude.

265 Notably, the shape of the background counts differs from the flux peaks. The background counts of TED 0° proton bands form a nearly plateau-like distribution, whereas the measured fluxes show a distinct peak at $L=4$. If the subauroral peak were purely due to contamination, a simple correction by subtracting background counts (even with a scaling factor) would not effectively remove the contamination.

TED 30° Proton Fluxes: A slightly different pattern emerges in the TED 30° proton channels (Fig. 4, top right). Here, the background count peak (as well as the flux) is lower than in the 0° channels. Additionally, the longitudinal variation (60°E vs. 160°E APEX) is less pronounced in lower-energy TED channels than in higher-energy ones. Interestingly, the expected subauroral peak in *TED 30° proton band 8* is not clearly observed at $L=4$, despite noticeable longitudinal differences.

TED 0° Electron Fluxes: The 0° electron fluxes (Fig. 4, bottom left) exhibit the smallest subauroral peaks among all detectors. In *TED 0° electron band 4*, the peak is absent, likely due to naturally high count rates that mitigate contamination. In 275 *TED 0° electron band 8*, a weak subauroral peak is visible only when comparing 60°E and 160°E APEX. The higher-energy electron channels (*TED 0° electron band 11* and 14) do show a subauroral peak, but it remains one to two orders of magnitude below auroral flux levels.

TED 30° Electron Fluxes: For the 30° electrons (Fig. 4, bottom right), background counts in the subauroral region are higher than for 0° electrons. *TED 30° electron band 4* remains unaffected, whereas the other TED electron channels show 280 similar 60°E–160°E differences, with subauroral peak fluxes surpassing the 0° case.

MEPED 0° and 90° e3 Fluxes: The MEPED electron instrument is known to have contamination issues. Thus at first we should assess if this is relevant in our context. According to Pettit et al. (Fig. 6 in 2019) the corrected *MEPED 0° e3* flux at

L=4–6 may be factor 2 smaller than the uncorrected flux. A similar correction approach including both viewing directions was done by Asikainen and Mursula (2013). Their Fig. 5 (left) also shows a factor 2 for strong contamination events (note that their graph shows the reciprocal value). Considering the corrected 90° equivalent Asikainen and Mursula (2013)[their Fig. 5, right] gives a factor 1.2 to 1.45 more than the uncorrected flux. The impact of a factor 1.2–1.45 on Figure 4 (as well as on Fig. 7, which follows in Section 5) is practically negligible.

As indicated by *MEPED 0° e3* (see Fig. 4, bottom left), the subauroral peak represents a local maximum in energetic electrons. But these fluxes are significantly substantially smaller than *MEPED 90° e3* (see other panels of Fig. 4). The latter shows relatively small fluxes in the polar region and a dominant subauroral peak between L=4 and L=5 (note that data in the SAA is mostly removed). Within the subauroral area (which -in geographic coordinates- stretches from approx. 300°E, 70°S to 40°E, 55°S, see Fig. 1, top left, in Yakovchuk and Wissing, 2019) the *MEPED 90°* detector measures trapped, drift loss cone and bounce loss cone particles (compare Fig. 1, middle panel, in Crack et al., 2025). As the trapped population typically exceeds the other populations, we may simplify that averaged *MEPED 90°* fluxes in the subauroral area are mostly representing the trapped population.

The shielding of the *MEPED* detector housing should eliminate out-of-view electron contamination only up to about 6 MeV (Evans and Greer, 2004b). Selesnick et al. (2020) showed that especially during periods of low pitch angle diffusion (most often occurring under low Kp conditions), most of the detector counts at L=4 in the 0° detector result from trapped or quasi-trapped (out-of-nominal field-of-view) electrons.

4.2 What might be the Reason for increased Subauroral Background Counts?

Energetic protons have already been excluded (see Section 4.1), but according to Li et al. (2001); Baker et al. (2013); Baker et al. (2019), highly relativistic electrons are common at L= 4 as this region belongs to the outer (electron) radiation belt. Additionally, the so-called impenetrable boundary of > 1 MeV electrons at about L=2.8 (Baker et al., 2014) fits to the background minimum seen in Fig. 4.

Energetic electron contamination can also explain the Kp dependence. On one hand there is the Dst-effect (McIlwain, 1966) which describes the outward radial motion of electron drift paths during increased geomagnetic activity in order to obey the third adiabatic invariant when the magnetic field strength drops. This is also connected with a reduction of electron energy. The Dst-effect itself is not connected to electron loss, it just temporarily displaces the electron populations, but particles may get lost to the interplanetary medium if their guiding field-lines cross the magnetopause (e.g. Kim et al., 2010). Loss processes are typically observed during high geomagnetic activity. Iles et al. (2002) states significant losses during storm main phase that may remove the majority of trapped electrons while conditions of recovery determine if there is an increase or decrease afterwards. Millan and Thorne (2007) provide a review summarizing loss processes and the connected wave activity during different geomagnetic activity. Further evidence is provided by a case study from Turner et al. (2012) showing that the radiation belt electron flux drops by orders of magnitude during geomagnetic storms. A quick radiation belt recovery, which was also noted by Turner et al. (2012), or increased radiation belt electrons fluxes after 53% of the investigated geomagnetic storms as observed by Reeves et al. (2003) do not contradict this statement as the increased fluxes are observed after the main event

and thus during lower Kp-values (or Dst-values, as seen in Fig. 2 of Reeves et al., 2003). The same is shown in Ross et al. (2021). Their Fig. 14 compares 4.2 MeV radiation belt electron measurements of the Van Allen Probes at different L-shells with Kp. Whenever the Kp rises to higher levels (say above 5), the electron flux decreases by orders of magnitude. This is most prominently seen at around $L \approx 4$ can be used as indication for reduced contamination at higher Kp.

In the context of our study, it is unnecessary to distinguish between the Dst-effect displacements and loss processes such as magnetopause shadowing or atmospheric interactions, as all lead to a (at least short-term) reduction in energetic electron populations, thereby resulting in reduced contamination during periods of high geomagnetic activity.

Comparing the location of the background peak with *MEPED 90° e3*, we conclude that the contaminating electrons should have slightly higher center energy than *MEPED 90° e3*. Observations from Peck et al. (2015, see their Fig. 5) constructing a virtual electron channel e4 with a higher energy center of 800 keV (based on the MEPED P6 proton channel) show a flux maximum at $L=4$, similar to Ross et al. (2021) for 4.2 MeV electrons, which agrees to the observed background count peak.

To conclude this subsection, the abundance of high-energy electrons is most likely responsible for the TED background counts and suggests that contamination in the TED proton channels (and to some smaller extent in the TED electron channels) is occurring. However, due to the absence of a correction algorithm, this does not provide a direct measure of the actual abundance of low-energy particles. In other words, it remains unclear whether the contamination is strong enough to fully account for the observed peaks in the low-energy channels.

4.3 Why do not all TED Channels show the same Background Counts?

The background counts, as well as the channels' difference between 60°E and 160°E APEX, are highest for TED 0° protons and TED 30° electrons, while TED 30° protons and TED 0° electrons show less impact. (Note that the different background levels are shown in Fig. 4 as well as in Fig. 8 where we are focussing on a single satellite.)

The shielding of the TED instrument is unfortunately completely undocumented. Using very rough estimations on the volume, mass, and material, we end up with an eventual tungsten shielding of probably less than 0.8 mm, which would allow penetration of electrons >2 MeV according to the ESTAR database¹ from NIST. However, this is very vague. According to private communication with Juan Rodriguez (NOAA), it might be better to assess contamination candidates by coincidence.

This, however, does not explain any differences between the detectors. In fact, combining a TED detector image (see <https://www.eoportal.org/satellite-missions/noaa-poes-series-5th-generation#sem-2-space-environment-monitor-2>) and the information on the size of the single detector openings, allowed us to determine the layout of the TED detector stack, being from side to side: 0° protons, 30° protons, 0° electrons, 30° electrons. Given that both inner detectors show fewer background counts, we conclude that this is an effect of better (lateral) shielding.

4.4 Comparison to Different Data Sets

The most relevant datasets for comparison are those from satellites with similar orbits. However, these datasets often suffer from the same issue – particle measurements in the radiation belts are problematic and frequently omitted. For instance, data from

¹<https://physics.nist.gov/PhysRefData/Star/Text/ESTAR.html>

the METEOR-3M satellite program (accessible at <https://ftp.sinp.msu.ru/meteor-3m/msgi/L1/> and <http://smdc.sinp.msu.ru/>)
350 are subject to such limitations.

Regarding different orbits, the TED data can be compared to the HOPE instrument onboard the Van Allen Probes. Yue et al. (2017, their Fig 2) shows statistical flux maps of low energetic 11 keV protons. During $Kp \leq 1$ the highest fluxes are observed between $L=5$ and the image boundary at $L=6$. It steadily decreases till $L=2$ or 3 where it reaches the minimum flux. During higher Kp the flux is increased from the outer boundary down to $L=4$ while $L=4$ does not represent a local maximum. This is
355 in contrast to the (contaminated) flux of the TED channels that show a maximum at $L=4$ (compare Fig. 4). However, it agrees with the SSUSI observations (discussed later in this Section) that show enhanced fluxes at $L=4$ (equivalent to -60° APEX) only during active periods. Higher proton energies of 55 keV from RBSP-A / RBSPICE as investigated by Li et al. (2023, their Fig. 1) show a similar pattern but are already relatively far from the TED energy range (<10 keV).

In this study we adopted a different comparison approach. If the peak observed in the low-energy 0° detectors is real, it
360 should manifest as particle precipitation in the atmosphere at the footpoints of the corresponding magnetic field lines (which coincide in APEX coordinates). To assess the atmospheric impact, we applied two independent methods. First, we examined the *S4 index*: a peak in TED proton flux at TED energies should result in an ionization peak at approximately 150 km altitude, which should be reflected in the *S4 index*. Second, particle interactions in the upper atmosphere should produce UV radiation, which can be detected using auroral imagers such as SSUSI. The benefit of this comparison is that these measurements are
365 independent of particle contamination issues.

4.4.1 Comparison to *S4 Index*

Figure 5 compares the *S4 index* (top panel), which serves as an indicator for ionization layer formation, with the particle flux from an affected TED channel, *TED 0° proton band 11* (middle and bottom panels).

The *S4* profiles in the first panel have been restricted to night-time² (20–04 local time). The night-time *S4 index* is elevated
370 between 80° – 190° E geographic longitude with a main peak at 100 – 120° E.

The second panel illustrates the particle flux for *TED 0° proton band 11* without any Kp filtering for the same period and night-time selection as the *S4 index* data set. The sinusoidal shape originates from the auroral oval in geographic coordinates, reaching latitudes closest to the equator at 90 – 150° E geographic. The highest flux values (above $10^{12} / (MeV s sr m^2)$, yellow) occur between 320° and 50° E geographic, corresponding to subauroral latitudes in APEX coordinates. Note that the SAA itself
375 is not shown here as it is located equatorward of 50° S. The lowest flux values appear between 250° and 300° E geographic, aligning with a region equatorward of the subauroral peak.

The bottom panel condenses the 2D information from the middle panel into a latitudinal mean, allowing direct comparison with the top panel. It shows three peaks, the highest one at 15° E which belongs to the subauroral TED flux but which does not appear at all in the *S4* data. The other two TED maxima at 70° E and -even higher- at 170° E belong to the auroral oval. The
380 peak at 170° E clearly appears in the *S4* data whereas the 70° E maximum is part of the main auroral *S4* maximum which also

²Including day-time *S4 index* would produce a second maximum at 270° E geographic –not associated with particle precipitation but being an artifact of the subsolar TEC maximum that expands into the observed domain only in the area of the SAA.




Figure 5. The top panel shows the GNSS-derived $S4$ index at 55–65°S, averaged over all available night-time (20–4 local time) profiles from 2001 to 2024 that do not exhibit a sporadic E-layer. The $S4$ index serves as an indicator of disturbances in the ionosphere. The middle panel presents the average particle flux in TED 0° proton band 11 for the same latitudinal band, covering the same years and the same filtering for night-time values only. The bottom panel condenses the middle panel’s data into a latitudinal mean, allowing for direct comparison with the $S4$ index.

contains other energy channels and of course the electron precipitation. Lower-energy particles create an auroral pattern that extends further poleward, leading to a single peak.

Note that Figure 5 is not very sensitive to high Kp particle precipitation. First, the temporal contribution is small and second all shown latitudes would be similarly affected, resulting in a mostly longitude-independent $S4$ offset.

385 The take-away message is, that in the region between 320° and 50°E geographic, where the TED channel registers a flux maximum, the $S4$ index does not indicate a corresponding ionization increase. Since particle precipitation at a specific energy should produce a distinct ionization and scintillation signature at a predictable altitude, the low $S4$ index suggests that the observed TED proton flux in this region is contamination from the horns of the outer radiation belt.




Figure 6. Temporal evolution of electron energy flux, proton energy flux, and their combined flux, compared to the energy flux derived from SSUSI auroral UV radiation as observed by two satellites, and the geomagnetic activity represented by Kp for the year 2013.

4.4.2 Comparison to SSUSI

390 The first 5 panels of Figure 6 present the precipitating energy flux for the year 2013 based on different instruments. For better comparison all energy ranges have been matched to the SSUSI data set, which is limited to an upper energy of about 20 keV.

Figure 6 (top panel) presents the total precipitating proton energy flux derived from spectrum fits of all TED 0° proton channels and *MEPED 0° P1*. The highest TED channel already covers 6.5–9.5 keV thus *MEPED 0° P1* is only used for the extrapolation up to 20 keV with very limited impact on the total precipitating flux. Two regions exhibit increased flux: one
395 centered around 70° S APEX latitude, indicative of auroral precipitation, and another at 60° S APEX latitude, corresponding to the subauroral peak. Periods of $K_p \geq 5$ are marked by blue lines in the upper part of the graph and also shown in the bottom panel. The subauroral peaks tend to follow periods of elevated K_p , and in some cases, the subauroral proton energy flux exceeds that of the auroral region. Two instances of strong subauroral peaks are highlighted for comparison.

The second panel displays the electron energy flux (154 eV to 20 keV) from the TED 0° electron channels and *MEPED*
400 *0° e1-e2*, a differential channel derived by subtracting e2 from e1. Based on results from Yando et al. (2011) this channel subtraction removes the impact of energetic electrons. As in the proton case the MEPED channel is used only from 9.5 keV to 20 keV in order to extrapolate the spectrum to the same energy range as the SSUSI data set.

Unlike the proton flux, the subauroral electron contribution is noticeably smaller, while the auroral electron flux is more pronounced. Additionally, auroral electron flux strongly depends on geomagnetic activity, intensifying and shifting equatorward
405 during high K_p conditions. The proton flux exhibits similar behavior, albeit at lower intensities.

The third panel combines proton and electron energy fluxes. The subauroral contribution is comparable to auroral levels, underscoring the need to determine whether the subauroral flux is real or an artifact.

Panels 4 and 5 present UV-derived energy fluxes from auroral emissions measured by DMSP-17 and DMSP-18, respectively. Both datasets exhibit a strong K_p dependence, with increased energy flux during geomagnetic disturbances and auroral
410 emissions extending to approximately 60° S APEX latitude. This aligns with the electron flux observations in the second panel. However, the marked subauroral peak does not appear in the UV data, even though the combined energy flux matches levels recognized as auroral emissions on some days. This suggests that the subauroral proton energy flux is overestimated.

Given the similarities in detector design, it is likely that the TED electron channels experience subauroral contamination as well. However, since the subauroral electron flux is lower, it may not be detectable in SSUSI data.

415 In summary, Figures 5 and 6 independently demonstrate –through the absence of ionization effects and auroral emissions– that the subauroral peak does not correspond to real precipitating low-energy particles. The only times when precipitation effects are recorded in this region coincide with geomagnetic storms, during which the auroral oval expands into this latitude range (see Fig. 6, second panel).

It is worth noting that Kavanagh et al. (2018) observed particle precipitation events at $L < 4$ (slot region) while comparing
420 electron measurements exceeding 640 keV (potentially above 1 MeV) with riometer measurements. These slot region filling events persist long after the storms that generate them and exhibit low K_p dependence. However, our observation methods, which focus on the upper atmosphere, do not capture deep atmospheric ionization (occurring in the lower mesosphere to




Figure 7. Correlation of *TED 0° proton band 11* and *MEPED 90° e3*.

stratopause, 50–55 km altitude). This suggests that while an energetic electron population is often present at low Kp, contributing to deep atmospheric ionization and contamination, the majority of TED proton channel fluxes during these periods are likely artifacts.

It is important to mention that disproving the existence of a subauroral low-energy proton peak during low Kp does not invalidate the possibility of subauroral proton precipitation in general. Kim et al. (2021) linked low-energy subauroral proton precipitation to EMIC wave activity. However, their study focused on substorm periods without high fluxes of electrons exceeding 100 keV.

430 5 Estimation of Contamination

Figure 7 illustrates the correlation between *TED 0° proton band 11* and *MEPED 90° e3* throughout 2013. It reveals two distinct particle populations. At low *MEPED 90° e3* flux levels (up to approximately $10^8 / (m^2 s sr MeV)$), the *TED 0° proton band 11* flux distribution remains largely independent of the electron flux. However, beyond this threshold, the peak fraction of the TED proton flux distribution (dark red area) increases with rising energetic electron flux. Notably, most data points in this second population originate from the subauroral region.

As demonstrated in Section 4, particularly in Subsection 4.4, the elevated TED fluxes in subauroral regions are not real. This strong correlation is a clear indication of contamination.

One might argue that the large difference in flux levels between these channels should prevent strong contamination; however, this is not the case. The secondary axis in Figure 7 presents integral fluxes across the full bandwidth of each channel. Within the correlated population, the peak fraction of the TED proton flux distribution follows a roughly one-to-one relationship with *MEPED 90° e3*, albeit with considerable spread. Additionally, *MEPED 90° e3* does not even peak at the same L-shell location, suggesting that a higher-energy electron channel –such as the virtual electron channel proposed by Peck et al. (2015)– might yield an even stronger correlation.

Another factor to consider is that both particle channels can record zero flux, indicating that the detectors' integration time may be insufficient to capture a meaningful flux level. Zeros in *MEPED 90° e3* are binned separately (see the left subfigure in Fig. 7), while *TED 0° proton band 11* registers zero flux at 16-second resolution in approximately 60% of cases, with this fraction decreasing above $3 \times 10^7 / (m^2 s sr MeV)$.

In conclusion, *TED 0° proton band 11* –and likely other TED channels– should be treated with caution when *MEPED 90° e3* flux level exceeds $10^8 / (m^2 s sr MeV)$. Due to the large spread in the data and the probably higher center-energy of the contaminating population, a simple subtraction of *MEPED 90° e3* scaled by its bandwidth would yield unreliable results and is therefore not recommended.

5.1 Correction Based on TED Background Counts

A natural approach to correct TED bands involves a subtraction of background counts, as applied by Green (2013) for total energy flux. However, Figure 4 (e.g., top left) and Figure 3 highlight two key challenges: (1) a proper conversion of background counts to physical flux is required, and (2) the non-linear response during high contamination must be accounted for. The background peaks observed in the South Atlantic Anomaly (SAA) and the subauroral region flatten at high values, a behavior not seen in uncorrected TED flux or the contaminating *MEPED 0° e3* peak. Consequently, applying only a simple linear scaling to match SSUSI observations at subauroral latitudes would introduce unphysical negative fluxes at L=2.5–3.

Different possible physical or technical sources of non-linearity come to mind. First, saturation effects may arise due to the longer integration time of the background detector (3.2 s) compared to 0.2 s for individual band measurements. However, the highest recorded background counts remain below 30 –far from the allowed maximum of 1,999,848 stated in Green (2013). Second, non-linearity could stem from data compression during satellite transmission (see Table 14 in Green, 2013). Yet, since we observe non-integer values, we trust that these represent decompressed counts (see Table 15 in Green, 2013, and notes in the netCDF files). Third, orbital motion during measurement could contribute: background counts are averaged over 16 instrument cycles (32 s total), during which the satellite travels about 270 km (or 1° latitude). This spatial averaging naturally smooths background peaks compared to individual band measurements. However, direct comparison with the old bin-format reveals that the saturation effect is caused by a different handling of the background data in the netCDF-files. In the netCDF-version all background data above an arbitrary level are cut and replaced by a linear interpolation of the closest surrounding background values. We probably can call this a bug as it unnecessarily hides the real (much higher) background values.




Figure 8. TED channels of NOAA15 plotted against their individual background channel.

470 The underlying assumption for an empirical contamination correction is that a given background count level corresponds to a predictable level of contamination in the particle bands. Since multiple flux measurements exist for each background count level, the lowest observed flux should best approximate the contamination component. We thus use the dataset from Figure 4 (without longitudinal restrictions and separately for each satellite as well as a similar data processing for M03 in year 2020, as it was not launched in 2013) to estimate the lower flux envelope for each background count value.

475 Figure 8 shows the particle populations at L=0–2.5 (grey, including the SAA), L=3–5 (black, subauroral area) and L>5
(colored) by plotting background counts against individual particle bands for NOAA15. Using *TED 0° proton band 11* (top
panel, third column) as typical example, the last group (in color, here blue) is centered around (10^0 , 3×10^{11}) and mostly
represents uncontaminated flux (meaning that the flux may be orders of magnitude higher than the contamination). The first
group (SAA, gray) forms a narrow band –on a wide range linear in log-log– extending to the highest background counts (about
480 2×10^1) where background channel reaches a (in this representation inverted) plateau, indicating the saturation-like cut-off. The
second group (black) corresponds to the subauroral region (L=3–5) and shows the same dependency on background counts as
the SAA population but does not reach the highest background counts. Also the flux variation for a specific background level
is typically higher (except for TED 0°/30° p 8).

Notably, contamination effects appear more pronounced in the outside detectors (TED 0° p and 30° e). This can be seen
485 clearly as the peak contamination within the subauroral population (gray) reaches much higher background values in 0° p than
in 30° p (same in 30° e compared to 0° e). This relation is observed in all satellites and is generally more pronounced than in
N15.

To estimate the lower contamination envelope we first need to define the boundaries. The upper boundary is the saturation
threshold. The saturation effect was evident in practically all SAA distributions³. In order to estimate the saturation level, we
490 binned the distribution (consisting of background count vs. particle flux pairs) into 20 logarithmically equidistant background
levels, derived the standard deviation of the logarithmic particle flux within these bins and selected the highest increase of the
standard deviation between neighboring bins as indication for the beginning saturation. The green vertical line (now called
saturation threshold) marks the background value that corresponds to the boundary between these bins. In many cases the
saturation threshold of 0° and 30° viewing angles were close to each other so that just one value has been selected. Only in a
495 few cases this method did not work out well and the saturation threshold has been set manually (integer values or 2.5).

Within each of the 20 logarithmically spaced background bins, the pairs were sorted by flux, and the logarithmic flux distance
to the following pair is determined. In order to suppress outliers, 1% of the data are omitted, depending on this distance. While
this filtering would also remove isolated data points within the distribution it turned out to be effective in removing outliers
below more populated areas. The pair with the lowest remaining flux value in each bin is returned and creates the data base
500 for the lower contamination envelope fit. However, a contamination should positively correlate with the background counts.
Therefore the minimum flux of all returned pairs is selected and its background count is used as lower boundary for the fit.
The we applied a linear fit to the logarithmic background counts $\log_{10}(b_g)$ and the logarithmic lower envelope channel flux
 $\log_{10}(\Phi_{LE})$ in form of $\log_{10}(\Phi_{LE}) = a \cdot \log_{10}(b_g) + b$. Fit parameters for the SAA (L=0–2.5) and subauroral region (L=3–
5) are presented in Table 2 and 3 and depicted in Figure 8 (for N15), Fig. 9 (for problematic channels) and Fig. 10 (for all
505 satellites).

³N17 TED 0° p 11/14 as well as N18 TED 0° p 11/14 had very few data points and did not allow to define a saturation threshold. We applied the thresholds
of the other viewing direction instead. M02 TED 30° p 11/14 are sufficiently populated without showing saturation effects. We handled it similarly as the
adopted threshold is too far above typical levels to be problematic.

☛	band	L=0–2.5		L=3–5		type	band	L=0–2.5		L=3–5		type	satu- ration
		a	b	a	b			a	b	a	b		
N15	0° p 4	1.3692	10.438	1.6471	10.1923	N	0° p 8	-	-	-	-	B	13.0573
N16		1.2691	10.5071	1.0656	10.5085	A		-	-	-	-	B	42.3749
N17		1.9867	10.0123	1.664	9.9264	A		-	-	-	-	B	9.9735
N18		1.2284	10.7717	1.3466	10.5727	N		-	-	-	-	B	11.9346
N19		1.2143	10.9064	1.622	10.562	N		-	-	-	-	B	12.7
M01		1.2413	10.7464	1.4671	10.5197	N		-	-	-	-	B	11.1117
M02		1.3515	10.5766	1.3922	10.5698	N		-	-	-	-	B	12.4286
M03		1.2408	10.7868	1.8257	10.3279	N		-	-	-	-	B	11.9792
N15	0° p 11	1.7854	9.6331	1.6871	9.6758	N	0° p 14	1.5908	9.2085	1.9199	8.8406	N	11.63
N16		-	-	-	-	C2		-	-	-	-	C2	0.2
N17		1.0452	10.292	0.3287	10.3134	A		1.054	9.5729	0.2457	9.6454	A	15.99
N18		1.9212	9.82	1.5581	9.9016	N		2.0183	9.1832	1.5781	9.2643	A	10.6284
N19		1.3347	10.2894	1.4744	10.1324	N		1.3957	9.5854	1.5225	9.4511	N	11.2012
M01		2.2212	8.7472	2.8392	8.5863	A		1.9873	8.2279	3.0604	7.5966	A	13.5737
M02		1.6974	9.7693	1.5476	9.9052	N		1.3228	9.4694	1.5713	9.229	N	11.374
M03		5.0627	5.6879	2.9747	7.7588	D2		5.3462	4.9232	3.2995	6.7386	D2	12.12
N15	30° p 4	1.4894	10.4026	1.7588	10.0822	N	30° p 8	-	-	-	-	B	12.752
N16		1.3923	10.6097	2.1465	9.7961	A		-	-	-	-	B	19.66
N17		1.6935	10.2283	1.8386	9.9347	N		-	-	-	-	B	11.0662
N18		1.1748	10.7141	1.5425	10.3959	N		-	-	-	-	B	14.2925
N19		1.1824	10.882	1.6277	10.3892	N		-	-	-	-	B	10.9012
M01		1.436	10.9302	2.0695	10.3583	A		-	-	-	-	B	10.9804
M02		1.2732	10.9672	1.8632	10.3523	N		-	-	-	-	B	11.8398
M03		1.0925	11.1182	2.2204	10.2314	A		-	-	-	-	B	12.153
N15	30° p 11	1.9747	9.2822	2.221	8.96	N	30° p 14	1.8418	8.7178	2.0934	8.4397	N	12.3852
N16		-	-	-	-	C2		-	-	-	-	C2	0.2034
N17		-	-	1.1097	9.2614	C1		-	-	1.3031	8.5724	C1	15.99
N18		-	-	-	-	C1		-	-	0.1464	9.0385	C1	10.6284
N19		1.5497	10.0123	0.9818	9.961	A		1.5522	9.329	1.3303	9.1878	N	7.9743
M01		0.4284	10.2786	0.9198	9.9783	N		0.538	9.4512	0.9033	9.3348	N	14.3504
M02		-	-	0.7271	9.5239	C1		-	-	1.5626	8.9905	C1	17.34
M03		-	-	-	-	C1		-	-	-	-	C1	2.6

Table 2. Fit parameters of the lower envelope of the contamination and the saturation threshold for the proton bands. When applying the (averaged) background flux b_g to $\log_{10}(\Phi_{LE}) = a \cdot \log_{10}(b_g) + b$ the resulting lower envelope flux Φ_{LE} is given in $1/(\text{MeV m}^2 \text{ s sr})$. The Table only states fits that may be used for a correction. “N” stands for “type normal”. Further information on the types is given in Fig. 9 and the related paragraph.

θ	band	L=0-2.5		L=3-5		type	band	L=0-2.5		L=3-5		type	saturation
		a	b	a	b			a	b	a	b		
N15	0° e 4	0.1757	12.7643	0.1479	12.7179	N	0° e 8	1.0405	10.4431	1.3256	10.2252	N	11.3735
N16		0.2582	12.3224	0.5066	12.1941	N		1.3427	10.4248	1.3278	10.2901	N	7.2533
N17		0.1507	12.7013	0.2385	12.3635	A		1.1995	10.2977	1.0513	10.1419	N	9.6104
N18		0.2434	12.226	0.2931	12.2106	N		1.3004	10.5618	0.6833	10.3584	A	12.5035
N19		0.2013	12.4409	0.2592	12.4548	N		1.059	10.528	1.3056	10.2366	N	10.2191
M01		0.9407	12.2016	0.2534	12.8199	A		1.1995	10.5339	1.2916	10.4423	N	11.4406
M02		0.4437	12.1087	0.2575	12.2763	N		1.2733	10.479	0.7848	10.316	N	10.77
M03		1.1911	11.533	0.4963	12.2266	A		1.2173	10.5204	2.1642	10.0818	A	11.2594
N15	0° e 11	3.0889	8.2427	3.2337	8.2487	N	0° e 14	1.4178	9.6304	1.8801	9.083	A	11.3193
N16		1.3126	10.077	1.364	9.9116	N		1.3113	9.7727	1.5205	9.5166	N	10.58
N17		2.5432	8.9685	0.9603	9.9276	A		2.1655	8.9129	1.2641	9.4011	A	11.6936
N18		2.2937	6.9668	3.116	6.2682	D1		2.7953	6.734	1.4694	7.9839	D1	14.85
N19		1.231	10.1163	1.6547	9.7326	A		1.0959	10.0199	1.5484	9.5294	A	8.7417
M01		1.2924	10.2585	1.6869	9.9044	A		1.2745	9.9674	1.8511	9.5145	A	10.6335
M02		3.4891	7.748	2.813	8.8181	A		2.5281	8.5939	2.649	8.6981	N	13.642
M03		3.6828	6.46	1.3791	9.0396	D2		9.4392	0.2084	1.3928	8.6598	D2	14.9028
N15	30° e 4	0.0899	12.9104	0.3497	12.6869	N	30° e 8	0.6108	11.0522	0.5996	11.032	N	10.4142
N16		0.1553	12.7563	0.4378	12.5086	N		1.1655	10.6201	1.2986	10.3105	A	11.45
N17		0.2087	12.9252	1.0532	11.9948	A		0.5132	10.9767	0.873	10.597	A	9.3792
N18		0.1898	12.1304	0.0856	11.9977	N		1.121	10.3719	0.5232	10.0557	A	10.13
N19		0.1071	12.6167	0.2328	12.5718	N		0.7693	10.646	0.9444	10.42	N	10.1744
M01		0.3425	12.7253	0.147	12.829	N		1.0557	10.5431	1.068	10.3681	N	9.303
M02		0.2131	12.5231	0.1355	12.5903	N		0.8635	10.6559	1.0241	10.4231	A	10.0025
M03		0.4474	12.1856	0.6044	12.1444	N		1.1192	10.541	1.7249	10.1694	A	8.7529
N15	30° e 11	1.7938	9.3485	2.009	9.282	N	30° e 14	1.6574	9.1646	1.9185	9.0407	N	11.73
N16		1.2646	10.0226	1.4302	9.7352	N		1.2905	9.6695	1.6476	9.1855	A	23.4004
N17		4.8817	3.74	1.3697	8.4371	D1		1.4431	7.6442	-	-	D1	15.8449
N18		0.5003	9.7502	3.984	5.8287	D1		3.2733	6.3603	2.5748	7.0328	D1	14.2552
N19		1.2582	9.8451	1.4969	9.6536	N		1.1985	9.5857	1.5305	9.2481	N	10.2177
M01		1.2233	10.1376	1.6744	9.7239	N		1.3658	9.6996	1.5543	9.5241	N	11.8745
M02		0.9888	9.6101	1.2065	9.4166	D2		0.3589	9.69	2.3486	7.9399	D2	10.2734
M03		1.2764	9.4402	0.8744	9.8616	D2		1.252	9.126	0.7579	9.6708	D2	15.0515

Table 3. Same as Table 2 for the electron bands.

N15 (see Fig. 8) seems to have relatively good particle detectors. Meaning that within most channels the contamination is proportional to the background values (below the saturation threshold). Also the contamination seems to be similar within the SAA and the subauroral area. Given the main expected contamination sources being protons in the SAA and electrons in the subauroral area the contamination is independent of particle type. This is indicated by the matching reddish solid and dashed lines labeled as lower envelope of the associated area. Also the saturation threshold is above typical values reached in the $L > 5$ region, which means that (at least at these K_p values) a cut at the saturation threshold is not problematic. We will call this case “type normal”. As a correction attempt all values below the saturation threshold may be subtracted by typical background contamination which should be the lower envelope with a certain factor taking into account statistical variations. In case that the TED flux does not show a positive correlation with background counts we omit the fit and no background is subtracted. All data above the saturation threshold should be discarded. However, *TED 0° proton band 8* shows that “type normal” does not apply to all channels.




Figure 9. Different types of problematic contaminaton (see text for details).

Figure 9 presents the different types of contamination that we found in examining all satellites: “type A” is similar to “type normal” but the contamination seems to depend on particle source. While the particle detector is equipped with a magnetic filter that might have a particle specific effect on the contamination. However it is not clear why this applies to some of the channels only. The affected channels are indicated in Tables 2 and 3. In case that a channel is also

identified as one of the following types, this supersedes the “type A” declaration. A correction for “type A” should be done as for “type normal” but considering the different kind of contamination.

525 “type B” applies to the proton band 8 on every satellite and both viewing directions. At least partly the distributions are not proportional to background counts. And we observed a different behavior in case of proton or electron contamination. Within the SAA region, enhanced flux is observed at relatively low background values but the neighboring channels 4 and 11 do not observe an enhancement. It is unclear what is going on here. A physical explanation might be that the satellite is negatively changed. But then the charging must always be on the same level because just this channel is affected. If we see charging effects, that would not impact particle precipitation. And charging should more be an issue for high flux regions as the auroral area than for high energy areas. Thus we recommend to not use p 8 on all satellites.

530 “type C1” seems to be particle specific as well. The subauroral flux seems not or not significantly affected by contamination. Within the SAA however the flux starts orders of magnitude smaller than anywhere else but then saturates almost instantly for increasing background counts. Consequently a correction can only be suggested for expected electron contamination. In case of proton contamination the data should be neglected.

535 “type C2” is very similar to “type C1” but electron contamination also seems to reduce the observed fluxes. As such any kind of contamination is uncorrectable. This type has only been observed in some channels of N16, which TED detector is officially declared as defect starting from 2006. We recommend to exclude these channels even before 2006.

540 In “type D” the saturation level is reached also outside typical contamination areas. For D1 no or just minor dependence on background flux below saturation level is observed while D2 shows obvious dependence on background. There is a smooth transition between D1 and D2. A correction might be done as in the “type normal” case but the correction will significantly change data outside of SAA and subauroral area as saturation level is reached there as well. Supposed that higher fluxes of the energetic particle populations (that are reflected in the background counts) are to some degree occurring simultaneously with increased fluxes of low energetic TED populations at higher L-shells, this correction will act as a high flux filter.

545 Figure 10 gives an overview on the lower envelopes of all TED channels on all satellites that are not excluded as uncorrectable type. *TED 0° electron band 8* shows the best overall agreement of the lower envelopes. Here, the background response seems to be almost identical for contaminating protons and electrons throughout all satellites. A very weak dependency on background counts is seen in both TED 0°/30° e 4 bands as indicated by the slow rise of the lower envelopes. The reason probably is that high flux values overshadow contamination effects. The strongest inter-satellite differences are typically connected with channels 11 and 14 (and the problematic p 8 bands) where the contamination effect gets stronger in relation to the lower flux values. Note that “type C1” occurs in p 30° 11/14 only while “type D” can be found in the 11/14 bands of p 0° and e 0°/30°. 550 The biggest differences to the other satellites are mostly found for N17, N18, M01 and M03 while no significant outliers have been observed for M02, N15 and N19. However, these differences are channel-specific –and may change over time. The plotted data is given in Tables 2 and 3.




Figure 10. Lower envelope of the contamination for all channels and satellites.

5.2 Application of the Contamination Correction:

Figure 11 demonstrates the effect of the suggested contamination correction under different geomagnetic conditions. The correction here has been done with a factor 1, meaning that if below saturation level only the lower contamination envelope has been subtracted. The applied correction method is available as Python code. In case that the background subtraction resulted in negative fluxes the channel has been removed and the particle spectrum fit has been done based on the remaining channels




Figure 11. Similar plot as Fig. 6 but with corrected fluxes. Note that the third panel (protons + electrons) contains data only in case that it is available for both particle species.

-if available. Period, location and color scale exactly matches to Fig. 6 and thus can be compared the contaminated fluxes and to SSUSI observations.

560 Starting with the proton precipitating flux, the moderately contaminated periods (outside the marked areas) are now practically cleaned from contamination, as. e.g. seen in early March, May or October. However, the proton correction does not come without the prize of missing data in cases of uncorrectable contamination. Given that this plot shows daily averages on 1° latitude resolution and on a 9° longitude patch the amount of data points per bin is not ideal for a correction attempt designed for average particle fluxes and average background counts. Within the marked heavily contaminated periods about
 565 half of the proton data is missing now. It should be said that this is just a minor problem for the atmospheric effects as the electron ionization is typically higher and mostly available after correction. Especially in July there are subauroral proton peaks remaining after the correction. Interestingly SSUSI on DMSP F18 (in contrast to F17) shows peaks in this period as well which are only slightly equatorward of this, so it remains unclear if the corrected fluxes are still contaminated here. Anyhow the overall proton picture is much closer to the SSUSI observations than with using uncorrected fluxes. The electron correction
 570 looks very convincing. Even during the heavily contaminated periods no elevated fluxes are seen any more. The proton plus

electron precipitating flux shows a much better agreement with the SSUSI observations. Please note that Figure 11 shows the worst-case scenario, with the strongest background counts occurring at 60°E APEX.

While the contamination is mostly removed now, this method does not replace an inter-comparison of the TED data. However, the corrected TED fluxes should be a better data base to use than the uncorrected fluxes.

575 Some uncertainty persists in the factor 1. It is not clear how strong natural fluctuations are impacting the results. It might be that at least part of the particle flux above the lower envelope is due to statistical variations that deviate from a direct relationship between background counts and contamination flux –which are not measured simultaneously or on the same location. While the factor 1 should be a good estimation for large ensembles, stronger statistical fluctuations when using just a few measurements might afford a factor >1 –with the cost of a reduced average (or negative fluxes). Within the scope of this study we can state that
580 the applied correction matches to the SSUSI observations in so far that the subauroral maximum is mostly removed while the auroral pattern persists with flux levels comparable to SSUSI observations. But the limitation of SSUSI's low flux resolution does not allow to quantify or the remaining subauroral contribution.

6 Conclusions

In this study, we investigate the subauroral flux maximum (located at approximately 60°N/S APEX, corresponding to L=4,
585 with a longitudinal focus between 0° and 100°E APEX, most prominent south of the South Atlantic Anomaly) observed in many TED channels. This subauroral maximum is predominantly visible in the vertical proton channels but is also present in nearly all other channels.

Two independent measurements of atmospheric impact contradict the presence of the observed subauroral peak of low-energy particle fluxes under low Kp conditions. Only at higher Kp levels does the expanding auroral oval contribute to low-energy precipitation in subauroral regions. It is important to note that our observational methods are limited to higher altitudes;
590 thus, precipitating energetic electrons with an ionization peak below 80km would not be detected.

Increased background counts indicate a likely instrumental issue, such as contamination, even though the absolute background levels remain low. The most probable source of elevated TED background counts is energetic electrons from the radiation belt. The increased background in the TED 0° proton and TED 30° electron channels may be attributed to the design
595 of the detector stack.

The vertical channels may be used to derive atmospheric ionization. However, failing to correct for the contaminated subauroral peak –such as in *TED 0° proton band 11* (which registers twice the typical auroral proton flux)– would result in subauroral thermospheric ionization rates reaching approximately 14% of the typical auroral electron ionization rates. This underscores the necessity of a correction method.

600 Table 1 presents a preliminary latitude-based cutoff approach, limited to channels with a clearly identifiable contamination peak. While this method leads to an underestimation of the precipitating flux (area), contamination appears to be the dominant source of subauroral signals and can be addressed without introducing a longitudinal dependence.

A detailed investigation of all particle channels and their individual contamination behavior (see Subsection 5.1) revealed that all channels are affected by a background count saturation level, above which the data should not be used. Lower envelopes
605 of the contamination, as a function of background counts, have also been identified and may be subtracted as a flux correction.

While many channels exhibit similar crosstalk responses to proton and electron particles, some channels react in a particle-specific manner. The 0° and 30° proton band 8 channels across all satellites show a distinctly different contamination pattern compared to neighboring channels, leading us to recommend excluding these channels from other studies (or at least using them with caution). The same applies to channels in which proton-induced background counts appear to suppress the measured
610 particle flux. Additionally, in some channels, the contamination threshold is reached even within the auroral region, meaning that applying a simple threshold-based cut would significantly impact the corrected flux.

The proposed channel- and satellite-specific corrections are summarized in Tables 2 and 3 and can be applied automatically using a provided Python routine (see the auxiliary material for this paper). These corrections are based on TED band flux, background counts, and L-shell, or alternatively allow for manual selection of protons or electrons as the contamination source.
615 Subsection 5.2 applies the Python routine to TED data from 2013, demonstrating (in Fig. 11) much better agreement with SSUSI observations. Limitations of the correction include: a) negative (i.e., missing) data due to over-correction, which primarily affects short-term binned proton data; and b) residual high subauroral proton fluxes, which may be either accurate (e.g., when compared to SSUSI F18) or inaccurate (e.g., when compared to SSUSI F17).

We should note that our proposed correction method is not limited to the recent netCDF-files. It should also work with
620 the older bin-files or a corrected netCDF algorithm without high-background cutting - even though a user should think about setting a maximum background flux (like the saturation threshold) that still allows for a reasonable subtractive correction, a certain proportion maybe. In case of the “type D” contamination a different background handling in the netCDF-files would be clearly beneficial.

Further studies may be valuable to assess the impact of fluctuations and determine whether a more aggressive correction
625 is warranted. Additionally, note that the correction functions are based solely on data from 2013 (and for N16, on 2005; for M03, on 2020); detector degradation or different voltage settings may alter the contamination effect in other years. Therefore, follow-up studies are encouraged.

Appendix A: Auxiliary Material: Python Code

The Python routine `TEDcorrection.py` contains the information from Tables 2 and 3 as well as the contamination type. It
630 provides the suggested correction functions based on TED particle flux, background counts and L-shell as used in Fig. 11.

Author contributions. Maik conducted most of the analysis and was the primary author of the paper. Olesya contributed to the analysis of TED data. Stefan provided the SSUSI dataset and associated tools. Christina facilitated the comparison with *S4 index*. All authors participated in discussions and refinement of the manuscript.

Competing interests. The authors declare that they have no conflict of interest.

635 *Acknowledgements.* This work was initially funded by the German Science Foundation (DFG project WI4417/2–1). The authors acknowledge the NOAA National Centers for Environmental Information (<https://ngdc.noaa.gov/stp/satellite/poes/dataaccess.html>) for the POES and Metop particle data used in this study. The Kp-index is provided from GFZ Potsdam (<https://kp.gfz-potsdam.de>). We thank Juan Rodriguez (NOAA) for a helpful comment on the energy range of the contaminating particles and Frank Heymann (DLR) for fruitful discussions on the background channels. The authors used ChatGPT for linguistic assistance in refining the manuscript.

640 References

- Andersson, M. E., Verronen, P. T., Rodger, C. J., Clilverd, M. A., and Wang, S.: Longitudinal hotspots in the mesospheric OH variations due to energetic electron precipitation, *Atmospheric Chemistry and Physics*, 14, 1095–1105, <https://doi.org/10.5194/acp-14-1095-2014>, 2014.
- Arras, C. and Wickert, J.: Estimation of ionospheric sporadic E intensities from GPS radio occultation measurements, *Journal of Atmospheric and Solar-Terrestrial Physics*, 171, 60–63, <https://doi.org/10.1016/j.jastp.2017.08.006>, vertical Coupling in the Atmosphere-Ionosphere System: Recent Progress, 2018.
- 645 Asikainen, T. and Mursula, K.: Recalibration of NOAA/MEPED energetic proton measurements, *J. Atmos. Sol. Terr. Phys.*, 73, 335–347, <https://doi.org/10.1016/j.jastp.2009.12.011>, 2011.
- Asikainen, T. and Mursula, K.: Correcting the NOAA/MEPED energetic electron fluxes for detector efficiency and proton contamination, *Journal of Geophysical Research: Space Physics*, 118, 6500–6510, <https://doi.org/10.1002/jgra.50584>, 2013.
- 650 Asikainen, T., Mursula, K., and Maliniemi, V.: Correction of detector noise and recalibration of NOAA/MEPED energetic proton fluxes, *J. Geophys. Res.*, 117, A09204, <https://doi.org/10.1029/2012JA017593>, 2012.
- Baker, D. N., Kanekal, S. G., Hoxie, V. C., Henderson, M. G., Li, X., Spence, H. E., Elkington, S. R., Friedel, R. H. W., Goldstein, J., Hudson, M. K., Reeves, G. D., Thorne, R. M., Kletzing, C. A., and Claudepierre, S. G.: A Long-Lived Relativistic Electron Storage Ring Embedded in Earth’s Outer Van Allen Belt, *Science*, 340, 186–190, <https://doi.org/10.1126/science.1233518>, 2013.
- 655 Baker, D. N., Jaynes, A. N., Hoxie, V. C., Thorne, R. M., Foster, J. C., Li, X., Fennell, J. F., Wygant, J. R., Kanekal, S. G., Erickson, P. J., Kurth, W., Li, W., Ma, Q., Schiller, Q., Blum, L., Malaspina, D. M., Gerrard, A., and Lanzerotti, L. J.: An impenetrable barrier to ultrarelativistic electrons in the Van Allen radiation belts, *Nature*, 515, 531–534, <https://doi.org/10.1038/nature13956>, 2014.
- Baker, D. N., Hoxie, V., Zhao, H., Jaynes, A. N., Kanekal, S., Li, X., and Elkington, S.: Multiyear Measurements of Radiation Belt Electrons: Acceleration, Transport, and Loss, *Journal of Geophysical Research (Space Physics)*, 124, 2588–2602, <https://doi.org/10.1029/2018JA026259>, 2019.
- 660 Bartels, J., Heck, N. H., and Johnston, H. F.: The three-hour-range index measuring geomagnetic activity, *Terrestrial Magnetism and Atmospheric Electricity*, 44, 411–454, <https://doi.org/10.1029/TE044i004p00411>, 1939.
- Bender, S., Espy, P. J., and Paxton, L. J.: Validation of SSUSI-derived auroral electron densities: comparisons to EISCAT data, *Ann. Geophys.*, 39, 899–910, <https://doi.org/10.5194/angeo-39-899-2021>, 2021.
- 665 Crack, M., Rodger, C. J., Clilverd, M. A., Hendry, A. T., and Sauvaud, J.-A.: Energetic Electron Precipitation From the Radiation Belts: Geomagnetic and Solar Wind Proxies for Precipitation Flux Magnitudes, *Journal of Geophysical Research: Space Physics*, 130, e2024JA033158, <https://doi.org/10.1029/2024JA033158>, e2024JA033158 2024JA033158, 2025.
- Crutzen, P. J., Isaksen, I. S. A., and Reid, G. C.: Solar Proton Events: Stratospheric Sources of Nitric Oxide, *Science*, 189, 457–459, <https://doi.org/10.1126/science.189.4201.457>, 1975.
- 670 Emmert, J. T., Richmond, A. D., and Drob, D. P.: A computationally compact representation of Magnetic-Apex and Quasi-Dipole coordinates with smooth base vectors, *Journal of Geophysical Research: Space Physics*, 115, <https://doi.org/10.1029/2010JA015326>, 2010.
- Evans, D., Garrett, H., Jun, I., Evans, R., and Chow, J.: Long-term observations of the trapped high-energy proton population ($L < 4$) by the NOAA Polar Orbiting Environmental Satellites (POES), *Advances in Space Research*, 41, 1261–1268, <https://doi.org/10.1016/j.asr.2007.11.028>, 2008.

- 675 Evans, D. S. and Greer, M. S.: Polar Orbiting Environmental Satellite Space Environment Monitor - 2, Instrument Descriptions and Archive Data Documentation, National Oceanic and Atmospheric Administration, NOAA Space Environ. Lab, Boulder, Colorado, USA, version 1.4b, including TED calibrations, 2004a.
- Evans, D. S. and Greer, M. S.: Polar Orbiting Environmental Satellite Space Environment Monitor - 2, Instrument Descriptions and Archive Data Documentation, National Oceanic and Atmospheric Administration, NOAA Space Environ. Lab, Boulder, Colorado, USA, version 680 2.0, 2004b.
- Evans, D. S. and Greer, M. S.: Polar Orbiting Environmental Satellite Space Environment Monitor - 2, Instrument Descriptions and Archive Data Documentation, National Oceanic and Atmospheric Administration, NOAA Space Environ. Lab, Boulder, Colorado, USA, version 2.0, 2006.
- Funke, B., Baumgaertner, A., Calisto, M., Egorova, T., Jackman, C. H., Kieser, J., Krivolutsky, A., López-Puertas, M., Marsh, D. R., Red- 685 dmann, T., Rozanov, E., Salmi, S.-M., Sinnhuber, M., Stiller, G. P., Verronen, P. T., Versick, S., von Clarmann, T., Vyushkova, T. Y., Wieters, N., and Wissing, J. M.: Composition changes after the "Halloween" solar proton event: the High Energy Particle Precipitation in the Atmosphere (HEPPA) model versus MIPAS data intercomparison study, *Atmospheric Chemistry and Physics*, 11, 9089–9139, <https://doi.org/10.5194/acp-11-9089-2011>, 2011.
- Gelinas, L. J. and Hecht, J. H.: Calibration/validation final report for the Special Sensor Ultraviolet Limb Imager (SSULI) and the Special 690 Sensor Ultraviolet Spectrographic Imager (SSUSI) on the Defense Meteorological Satellite Program (DMSP) F19 Spacecraft, Aerospace Report TOR-2016-00996, Space and Missile Systems Center, Air Force Space Command, 483 N. Aviation Blvd., El Segundo, CA, USA, attn: SMC/RS, 90245-2808, 2016.
- Green, J.: TED Data Processing - ALGORITHM THEORETICAL BASIS DOCUMENT - Version 1.0, Tech. rep., NOAA National Geo- physical Data Center, https://ngdc.noaa.gov/stp/satellite/poes/docs/NGDC/TEDprocessingATBD_V1.doc, version 1.0, 2013.
- 695 Heath, D. F., Krueger, A. J., and Crutzen, P. J.: Solar proton event - Influence on stratospheric ozone, *Science*, 197, 886–889, <https://doi.org/10.1126/science.197.4306.886>, 1977.
- Iles, R. H. A., Fazakerley, A. N., Johnstone, A. D., Meredith, N. P., and Bühler, P.: The relativistic electron response in the outer radiation belt during magnetic storms, *Annales Geophysicae*, 20, 957–965, <https://doi.org/10.5194/angeo-20-957-2002>, 2002.
- Kavanagh, A. J., Cobbett, N., and Kirsch, P.: Radiation Belt Slot Region Filling Events: Sustained Energetic Precipitation Into the Meso- 700 sphere, *Journal of Geophysical Research: Space Physics*, 123, 7999–8020, <https://doi.org/10.1029/2018JA025890>, 2018.
- Kim, H., Shiokawa, K., Park, J., Miyoshi, Y., Miyashita, Y., Stolle, C., Connor, H. K., Hwang, J., Buchert, S., Kwon, H.-J., Nakamura, S., Nakamura, K., Oyama, S.-I., Otsuka, Y., Nagatsuma, T., and Sakaguchi, K.: Isolated Proton Aurora Driven by EMIC Pc1 Wave: PWING, Swarm, and NOAA POES Multi-Instrument Observations, *Geophysical Research Letters*, 48, e2021GL095090, <https://doi.org/10.1029/2021GL095090>, e2021GL095090 2021GL095090, 2021.
- 705 Kim, K. C., Lee, D.-Y., Kim, H.-J., Lee, E. S., and Choi, C. R.: Numerical estimates of drift loss and Dst effect for outer radiation belt relativistic electrons with arbitrary pitch angle, *Journal of Geophysical Research: Space Physics*, 115, <https://doi.org/https://doi.org/10.1029/2009JA014523>, 2010.
- Knight, H. K.: Auroral ionospheric E region parameters obtained from satellite-based far-ultraviolet and ground-based ionosonde observations – effects of proton precipitation, *Annales Geophysicae*, 39, 105–118, <https://doi.org/10.5194/angeo-39-105-2021>, 2021.
- 710 Knight, H. K., Strickland, D. J., Correira, J., Hecht, J. H., and Straus, P. R.: An empirical determination of proton auroral far ultraviolet emission efficiencies using a new nonclimatological proton flux extrapolation method, *Journal of Geophysical Research: Space Physics*, 117, <https://doi.org/10.1029/2012JA017672>, 2012.

- 715 Knight, H. K., Galkin, I. A., Reinisch, B. W., and Zhang, Y.: Auroral Ionospheric E Region Parameters Obtained From Satellite-Based Far Ultraviolet and Ground-Based Ionosonde Observations: Data, Methods, and Comparisons, *J. Geophys. Res. Space Phys.*, 123, 6065–6089, <https://doi.org/10.1029/2017ja024822>, 2018.
- Lam, M. M., Horne, R. B., Meredith, N. P., Glauert, S. A., Moffat-Griffin, T., and Green, J. C.: Origin of energetic electron precipitation >30 keV into the atmosphere, *Journal of Geophysical Research: Space Physics*, 115, <https://doi.org/10.1029/2009JA014619>, 2010.
- Laundal, K. M. and Richmond, A. D.: Magnetic Coordinate Systems, *Space Sci. Rev.*, 206, 27–59, <https://doi.org/10.1007/s11214-016-0275-y>, 2017.
- 720 Li, J., Bortnik, J., Chu, X., Ma, D., Tian, S., Wang, C.-P., Manweiler, J. W., and Lanzerotti, L. J.: Modeling Ring Current Proton Fluxes Using Artificial Neural Network and Van Allen Probe Measurements, *Space Weather*, 21, e2022SW003257, <https://doi.org/10.1029/2022SW003257>, e2022SW003257 2022SW003257, 2023.
- Li, X., Baker, D. N., Kanekal, S. G., Looper, M., and Temerin, M.: Long term measurements of radiation belts by SAMPEX and their variations, *Geophysical Research Letters*, 28, 3827–3830, <https://doi.org/10.1029/2001GL013586>, 2001.
- 725 McIlwain, C. E.: Ring current effects on trapped particles, *Journal of Geophysical Research (1896-1977)*, 71, 3623–3628, <https://doi.org/10.1029/JZ071i015p03623>, 1966.
- McPeters, R. D. and Jackman, C. H.: The response of ozone to solar proton events during solar cycle 21: The observations, *Journal of Geophysical Research: Atmospheres*, 90, 7945–7954, <https://doi.org/10.1029/JD090iD05p07945>, 1985.
- Millan, R. and Thorne, R.: Review of radiation belt relativistic electron losses, *Journal of Atmospheric and Solar-Terrestrial Physics*, 69, 362–377, <https://doi.org/10.1016/j.jastp.2006.06.019>, global Aspects of Magnetosphere-Ionosphere Coupling, 2007.
- 730 Nesse Tyssøy, H., Sandanger, M. I., Ødegaard, L.-K. G., Stadsnes, J., Aasnes, A., and Zawedde, A. E.: Energetic electron precipitation into the middle atmosphere-Constructing the loss cone fluxes from MEPED POES, *Journal of Geophysical Research: Space Physics*, 121, 5693–5707, <https://doi.org/10.1002/2016JA022752>, 2016.
- Ødegaard, L.-K. G., Tyssøy, H. N., Sandanger, M. I. J., Stadsnes, J., and Søråas, F.: Space Weather impact on the degradation of NOAA POES MEPED proton detectors, *J. Space Weather Space Clim.*, 6, A26, <https://doi.org/10.1051/swsc/2016020>, 2016.
- 735 Paxton, L. J., Meng, C.-I., Fountain, G. H., Ogorzalek, B. S., Darlington, E. H., Gary, S. A., Goldsten, J. O., Kusnierkiewicz, D. Y., Lee, S. C., Linstrom, L. A., Maynard, J. J., Peacock, K., Persons, D. F., and Smith, B. E.: Special sensor ultraviolet spectrographic imager: an instrument description, in: *Instrumentation for Planetary and Terrestrial Atmospheric Remote Sensing*, edited by Chakrabarti, S. and Christensen, A. B., SPIE, <https://doi.org/10.1117/12.60595>, 1992.
- 740 Paxton, L. J., Schaefer, R. K., Zhang, Y., and Kil, H.: Far ultraviolet instrument technology, *J. Geophys. Res. Space Phys.*, 122, 2706–2733, <https://doi.org/10.1002/2016ja023578>, 2017.
- Paxton, L. J., Schaefer, R. K., Zhang, Y., Kil, H., and Hicks, J. E.: SSUSI and SSUSI-Lite: Providing Space Situational Awareness and Support for Over 25 Years, *Johns Hopkins APL Tech. Dig.*, 34, 388–400, 2018.
- 745 Peck, E. D., Randall, C. E., Green, J. C., Rodriguez, J. V., and Rodger, C. J.: POES MEPED differential flux retrievals and electron channel contamination correction, *Journal of Geophysical Research: Space Physics*, 120, 4596–4612, <https://doi.org/10.1002/2014JA020817>, 2015.
- Pettit, J. M., Randall, C. E., Peck, E. D., Marsh, D. R., van de Kamp, M., Fang, X., Harvey, V. L., Rodger, C. J., and Funke, B.: Atmospheric Effects of >30-keV Energetic Electron Precipitation in the Southern Hemisphere Winter During 2003, *Journal of Geophysical Research: Space Physics*, 124, 8138–8153, <https://doi.org/10.1029/2019JA026868>, 2019.

- 750 Reeves, G. D., McAdams, K. L., Friedel, R. H. W., and O'Brien, T. P.: Acceleration and loss of relativistic electrons during geomagnetic storms, *Geophysical Research Letters*, 30, <https://doi.org/10.1029/2002GL016513>, 2003.
- Richmond, A. D.: Ionospheric Electrodynamics Using Magnetic Apex Coordinates., *Journal of Geomagnetism and Geoelectricity*, 47, 191–212, <https://doi.org/10.5636/jgg.47.191>, 1995.
- Rodger, C. J., Clilverd, M. A., Green, J. C., and Lam, M. M.: Use of POES SEM-2 observations to examine radiation
755 belt dynamics and energetic electron precipitation into the atmosphere, *Journal of Geophysical Research: Space Physics*, 115, <https://doi.org/10.1029/2008JA014023>, 2010.
- Ross, J. P. J., Glauert, S. A., Horne, R. B., Watt, C. E. J., and Meredith, N. P.: On the Variability of EMIC Waves and the Consequences for the Relativistic Electron Radiation Belt Population, *Journal of Geophysical Research: Space Physics*, 126, e2021JA029754, <https://doi.org/10.1029/2021JA029754>, e2021JA029754 2021JA029754, 2021.
- 760 Rusch, D., Gérard, J.-C., Solomon, S., Crutzen, P., and Reid, G.: The effect of particle precipitation events on the neutral and ion chemistry of the middle atmosphere—I. Odd nitrogen, *Planetary and Space Science*, 29, 767–774, [https://doi.org/10.1016/0032-0633\(81\)90048-9](https://doi.org/10.1016/0032-0633(81)90048-9), 1981.
- Sandanger, M. I., Ødegaard, L.-K. G., Nesse Tyssøy, H., Stadsnes, J., Søråas, F., Oksavik, K., and Aarsnes, K.: In-flight calibration of NOAA POES proton detectors—Derivation of the MEPED correction factors, *Journal of Geophysical Research: Space Physics*, 120, 9578–9593,
765 <https://doi.org/10.1002/2015JA021388>, 2015.
- Selesnick, R. S., Baker, D. N., Jaynes, A. N., Li, X., Kanekal, S. G., Hudson, M. K., and Kress, B. T.: Observations of the inner radiation belt: CRAND and trapped solar protons, *Journal of Geophysical Research (Space Physics)*, 119, 6541–6552, <https://doi.org/10.1002/2014JA020188>, 2014.
- Selesnick, R. S., Tu, W., Yando, K., Millan, R. M., and Redmon, R. J.: POES/MEPED Angular Response Functions and
770 the Precipitating Radiation Belt Electron Flux, *Journal of Geophysical Research: Space Physics*, 125, e2020JA028240, <https://doi.org/10.1029/2020JA028240>, e2020JA028240 10.1029/2020JA028240, 2020.
- Solomon, S., Rusch, D., Gérard, J., Reid, G., and Crutzen, P.: The effect of particle precipitation events on the neutral and ion chemistry of the middle atmosphere: II. Odd hydrogen, *Planetary and Space Science*, 29, 885–893, [https://doi.org/10.1016/0032-0633\(81\)90078-7](https://doi.org/10.1016/0032-0633(81)90078-7), 1981.
- 775 Solomon, S., Reid, G. C., Rusch, D. W., and Thomas, R. J.: Mesospheric ozone depletion during the Solar Proton Event of July 13, 1982 Part II. Comparison between theory and measurements, *Geophysical Research Letters*, 10, 257–260, <https://doi.org/10.1029/GL010i004p00257>, 1983.
- Søråas, F., Sandanger, M. I., and Smith-Johnsen, C.: NOAA POES and MetOp particle observations during the 17 March 2013 storm, *Journal of Atmospheric and Solar-Terrestrial Physics*, 177, 115–124, <https://doi.org/10.1016/j.jastp.2017.09.004>, dynamics of the Sun-
780 Earth System: Recent Observations and Predictions, 2018.
- SSUSI: Data products, available at: https://ssusi.jhuapl.edu/data_products, last access: 21 September, 2020.
- Strickland, D. J., Jasperse, J. R., and Whalen, J. A.: Dependence of auroral FUV emissions on the incident electron spectrum and neutral atmosphere, *J. Geophys. Res. Space Phys.*, 88, 8051, <https://doi.org/10.1029/ja088ia10p08051>, 1983.
- Thébault, E., Finlay, C. C., Beggan, C. D., Alken, P., Aubert, J., Barrois, O., Bertrand, F., Bondar, T., Boness, A., Brocco, L., Canet, E.,
785 Chambodut, A., Chulliat, A., Coisson, P., Civet, F., Du, A., Fournier, A., Fratter, I., Gillet, N., Hamilton, B., Hamoudi, M., Hulot, G., Jager, T., Korte, M., Kuang, W., Lalanne, X., Langlais, B., Léger, J.-M., Lesur, V., Lowes, F. J., Macmillan, S., Mandea, M., Manoj, C., Maus, S., Olsen, N., Petrov, V., Ridley, V., Rother, M., Sabaka, T. J., Saturnino, D., Schachtschneider, R., Sirol, O., Tangborn, A.,

- Thomson, A., Tøffner-Clausen, L., Vigneron, P., Wardinski, I., and Zvereva, T.: International Geomagnetic Reference Field: the 12th generation, *Earth, Planets and Space*, 67, 79, <https://doi.org/10.1186/s40623-015-0228-9>, 2015.
- 790 Turner, D. L., Shprits, Y., Hartinger, M., and Angelopoulos, V.: Explaining sudden losses of outer radiation belt electrons during geomagnetic storms, *Nature Physics*, 8, 208–212, <https://doi.org/10.1038/nphys2185>, 2012.
- van Allen, J. A.: The Geomagnetically Trapped Corpuscular Radiation, *Journal of Geophysical Research*, 64, 1683–1689, <https://doi.org/10.1029/JZ064i011p01683>, 1959.
- van Allen, J. A.: *Origins of magnetospheric physics.*, University of Iowa Press, 1983.
- 795 van der Meeren, C., Laundal, K. M., Burrell, A. G., Lamarche, L., Starr, G., Reimer, A., Morschhauser, A., and Michaelis, I.: ApexPy 2.1.0, <https://doi.org/10.5281/zenodo.1214206>, 2024.
- Van Zandt, T., Clark, W., and Warnock, J.: Magnetic apex coordinates: A magnetic coordinate system for the ionospheric F2 layer, vol. 55, Environmental Research Laboratories, 1972.
- Vette, J. I.: *The AE-8 trapped electron model environment*, 1991.
- 800 Winant, A., Pierrard, V., and Botek, E.: Comparison of radiation belt electron fluxes simultaneously measured with PROBA-V/EPT and RBSP/MagEIS instruments, *Annales Geophysicae*, 41, 313–325, <https://doi.org/10.5194/angeo-41-313-2023>, 2023.
- Wissing, J. M. and Kallenrode, M.-B.: Atmospheric Ionization Module Osnabrück (AIMOS): A 3-D model to determine atmospheric ionization by energetic charged particles from different populations, *Journal of Geophysical Research: Space Physics*, 114, <https://doi.org/10.1029/2008JA013884>, 2009.
- 805 Wu, D. L.: Ionospheric S4 Scintillations from GNSS Radio Occultation (RO) at Slant Path, *Remote Sensing*, 12, <https://doi.org/10.3390/rs12152373>, 2020.
- Yakovchuk, O. and Wissing, J.: Magnetic local time asymmetries in precipitating electron and proton populations with and without substorm activity, *Annales Geophysicae*, 37, 1063–1077, <https://doi.org/10.5194/angeo-37-1063-2019>, 2019.
- Yando, K., Millan, R. M., Green, J. C., and Evans, D. S.: A Monte Carlo simulation of the NOAA POES Medium Energy Proton and Electron Detector instrument, *Journal of Geophysical Research: Space Physics*, 116, <https://doi.org/10.1029/2011JA016671>, 2011.
- 810 Yue, C., Bortnik, J., Chen, L., Ma, Q., Thorne, R. M., Reeves, G. D., and Spence, H. E.: Transitional behavior of different energy protons based on Van Allen Probes observations, *Geophysical Research Letters*, 44, 625–633, <https://doi.org/10.1002/2016GL071324>, 2017.