

Response and recovery of a *Sphagnum* peatland from long-term human-induced alkalinisation

Luke Oliver Andrews^{a,b}, Katarzyna Marcisz^a, Piotr Kołaczek^a, Leeli Amon^c, Siim Veski^c, Atko Heinsalu^c, Normunds Stivrins^{c,d,e}, Mariusz Bąk^a, Marco A. Aquino-Lopez^f, Anna Cwanek^g, Edyta Łokasⁱ, Monika Karpińska-Kołaczek^a, Sambor Czerwiński^{a,i}, Michał Słowiński^j, Mariusz Lamentowicz^a

Corresponding author: Luke Andrews, l.o.andrews@ljmu.ac.uk, ORCID ID: <https://orcid.org/0000-0002-8494-4791>

Manuscript in preparation for submission to: Biogeosciences

- a. Climate Change Ecology Research Unit, Adam Mickiewicz University, Poznań, Poland.
- b. School of Biological and Geographical Studies, Liverpool John Moores, Liverpool, UK (present address).
- c. Department of Geology, Tallinn University of Technology, Tallinn, Estonia.
- d. Department of Geology, Faculty of Science and Technology, University of Latvia, Riga, Latvia.
- e. Faculty of Forest and Environmental Sciences, Latvia University of Life Sciences and Technologies, Jelgava, Latvia
- f. Department of Geography, University of Cambridge, Cambridge, UK.
- g. Department of Mass Spectrometry, Institute of Nuclear Physics Polish Academy of Sciences, Krakow, Poland.
- h. Institute of Nuclear Physics, Polish Academy of Sciences, Warsaw, Poland.
- i. Department of Geomorphology and Quaternary Geology, University of Gdańsk, Bażyńskiego 4, 80-309, Gdansk, Poland
- j. Institute of Geography and Spatial Organization, Polish Academy of Sciences, Warsaw, Poland.

Keywords: Alkalinisation, Carbon, Vegetation, Palaeoecology, Multi-proxy, Peatland.

Abstract

Northern peatlands are significant terrestrial carbon stores but are increasingly threatened by human activities. Ombrotrophic peatlands, being naturally acidic, are particularly vulnerable to alkaline pollution. Despite their importance, the effects of alkalinisation on peatlands remain insufficiently studied. In Estonia, alkaline pollution from a cement industry and oil shale power plant emissions have degraded several peatlands since the 19th century. Although some sites have recovered in recent decades, more severely impacted areas remain in poor condition.

We investigated the effects of alkalinisation on Varudi peatland, a forested site in northeast Estonia, which was exposed to 125 years of alkaline emissions from a nearby cement factory. Using a multi-proxy, high-resolution palaeoecological approach combined with a precise and reliable age-depth model, we reconstructed changes in environmental, chemical, botanical, and hydrological conditions over the past millennium. Our findings revealed three successional phases: during the mid-13th century CE, land clearance and increased mineral deposition caused the site to transition from a bog to a poor fen phase between approximately 1250-1570 CE; and while the cement factory operated without efficient filters, the site became a pine-dominated fen between 1871-1995.

After the installation of filters in 1996, peatland pH returned to pre-disturbance levels, and some recovery was observed. However, the site remains degraded. Our results indicate that alkalinisation significantly disrupts peatland functioning, reducing carbon storage and altering vegetation communities. These effects can persist for decades even after the source of contamination is removed, underscoring the need for more comprehensive monitoring of peatlands impacted by alkaline pollution globally.

1. Introduction

Despite only covering c. 3% of the Earth's surface, northern peatlands contain >500 gigatonnes of carbon (Bridgham *et al.*, 2008; Yu *et al.*, 2010; Yu, 2012). Their capacity to accumulate and store carbon results from the waterlogged and acidic nature of their soils (Clymo *et al.*, 1998). These conditions preserve organic material, which accumulates and may be stored indefinitely (Harenda *et al.*, 2018). Since their initiation, peatlands have slowly removed carbon from the atmosphere, imparting a weak but persistent cooling effect upon global climate over millennial timescales (Frolking *et al.*, 2006).

Despite being recognised as a valuable tool for climate change mitigation, peatlands still receive little protection, regionally or nationally (Rawlins and Morris, 2010). As of 2018, approximately 10% of the remaining peatlands worldwide are in a degraded state (Leifeld and Menichetti, 2018), while in Europe this rises to 25% (Tanneberger *et al.*, 2021). Such disturbance can disrupt the fragile hydrological balance that maintains the carbon sink function of peatlands and may cause them to shift from sinks to sources of atmospheric carbon, exacerbating climate change (Leifeld and Menichetti, 2018).

Estonia is one of Europe's most peat-rich countries, with peatlands covering c. 22.5% of its land area (Orru and Orru, 2008). Due to this abundance, peat is a significant natural resource for Estonia, and has been heavily exploited, particularly after the Industrial Revolution (Paal *et al.*, 2010; Łuców *et al.*, 2022). The rise in anthropogenic pollution since this time has caused substantial changes in global geochemistry such that this era is informally termed 'the Anthropocene' (Fiałkiewicz-Kozieł *et al.*, 2018; Waters *et al.*, 2023). During this time in Estonia, emissions from industrial sources were characterised by high levels of calcium-rich particulate matter, with most of the emissions concentrated in the northeastern industrial region of the country (Liblik *et al.*, 1995; Karofeld, 1996). Ombrotrophic peat bogs, which are the dominant type of peatlands in Estonia, being naturally acidic and nutrient-poor ecosystems are particularly sensitive to alkaline atmospheric pollution (Paal *et al.*, 2010). These emissions caused significant changes in the geochemical and botanical composition of bogs adjacent to pollution sources, which resulted in dramatic increases in pore-water pH and losses of bog-specific vegetation (including *Sphagnum* mosses) near affected sites (Paal *et al.*, 2010; Vellak *et al.*, 2014). This was often followed by encroachment of *Pinus sylvestris* (Scots

pine) onto polluted sites alongside other species typical of nutrient-rich alkaline environments (Pensa *et al.*, 2004, 2007; Ots and Reisner, 2006; Kaasik *et al.*, 2008; Kask *et al.*, 2008).

By the 1990s, industrial emissions in Estonia began to fall, following a decline in power generation and improved filtration systems in factories (Liiv and Kaasik, 2004). Following these reductions, polluted peatland sites began to show signs of recovery, with acidic conditions and bog-specific vegetation returning (Karofeld, 1996; Kaasik *et al.*, 2008; Paal *et al.*, 2010). However, in more heavily polluted sites, this recovery has been slow, and the impact of past alkaline pollution persists to this day in some areas (Ots and Reisner, 2006). It remains unclear whether current levels of atmospheric pollution are sufficiently low to permit their full recovery in the future, or how long this process will take (Paal *et al.*, 2010). Despite growing concerns over alkaline pollution and its potential future effect on peatlands, particularly concerning their role as carbon reservoirs, research exploring the effects of alkalinisation upon peatland ecosystems and their subsequent recovery has been limited.

Atmospheric pollution remains a significant threat to peatland ecosystem functioning (Bobbink *et al.*, 1998; Turetsky and St Louis, 2006; Osborne *et al.*, 2024). The effects of alkaline pollution upon peatlands have been relatively overlooked relative to those of acid rain due to its effects being more localised (Vellak *et al.*, 2014; Sutton *et al.*, 2020). However, nearly two billion tonnes of alkaline residues are emitted into the atmosphere each year (Gomes *et al.*, 2016). Despite environmental standards curbing emissions in recent decades, in some areas these regulations are not consistently enforced or are merely declarative (Abril *et al.*, 2014; Ivanov *et al.*, 2018). Following global reductions in acid rain since the 1980s, the relative proportion of alkaline pollutants in airborne particulate matter has increased in the UK, much of Europe, North America and China since 1986 (Turetsky and St Louis, 2006; Sutton *et al.*, 2020). Additionally, climate change may exacerbate the effects of increased alkalinity in boreal regions, as permafrost thawing may cause the expansion of areas of open water, increasing surface runoff and infiltration in some regions (Walvoord and Kurylyk, 2016). This may allow for longer contact times between surface water and carbonate bedrock, accelerating weathering and raising the pH of surface waters which may then enter peatlands in the surrounding catchment (Schindler, 1997; Osterkamp *et al.*, 2000; Lehmann *et al.*, 2023).

In this study, we focus on the effects of alkalinisation resulting from over 125 years of intense emissions from a nearby cement factory upon Varudi soo (bog), a formerly ombrotrophic peatland in northeastern Estonia. By employing a high-resolution, multi-proxy palaeoecological framework, we reconstruct changes in the chemical, botanical, hydrological and environmental conditions of the site over the past millennium to address the following questions:

1. What is the current state of a heavily polluted raised bog almost 30 years after the reduction in alkaline pollution?
2. How has alkaline pollution altered the ecosystem functioning of the site and how does this compare with pre-disturbance conditions?
3. To what extent has this ecosystem function recovered 30 years after removing the point source of pollution?
4. Can we identify critical transitions that can be broadly applied to assess peatland condition and recovery following alkaline pollution?

2. Methods and materials

2.1. Study area

Varudi bog (59°26'19"N, 26°35'13"E) is located in Lääne-Virumaa, northeastern Estonia, consisting of fen-bog habitats. The site is approximately 10 km south of the coast of the Gulf of Finland and Baltic Klint (Figure 1) and spans c. 12.6 km². The site is primarily a forested *Sphagnum* bog interspersed by numerous bog pools and hummocks, with an overstory of *Pinus sylvestris*. Varudi peatland receives 478 mm of rainfall per year, has a mean annual temperature of 7.3 °C, and prevailing winds are from the southwest and south. The underlying bedrock is composed of Cambrian and Lower Ordovician siliciclastic sedimentary rocks and Middle Ordovician limestones (Sibul *et al.*, 2017) covered with a relatively thin layer of glacial and post-glacial sediments.




Figure 1: Map of study locations. A. Modern-day Estonia (in Gray). The red shaded square in map 1A indicates the area mapped in map 1B. B. Locations of sites relevant to this study (Kunda Cement Factory: Red, Aru-Lõuna Quarry: Yellow, Varudi peatland: Blue). C. Satellite image of Varudi bog (© Microsoft) showing the extent of peat cutting and drainage that has taken place over the past century. The red-shaded area indicates the area mapped in map 1D. D. Close-up of the location from where core VAR1 was recovered (red star). E. Coring location.




Figure 2. Photographs of the Kunda Cement factory and locale taken during the 1980s and 1990s, showing A. Cement dust emissions from the chimneys. B. Photograph of chimneys and factory surroundings. C. Photograph of the factory workings, showing substantial cement dust deposition in the surrounding area. D. Photograph showing cement dust deposition upon house near the factory. Photography by: A and D: Atko Heinsalu taken in the early 1990s; B: Estonian National Archives Photo Database (code EFA.204.0.268452) August 1994 Albert Truuväart. C: Estonian National Archives Photo Database (code EFA.204.0.168924) August 1989 Tiit Veermäe.

Varudi was selected for study due to its proximity (7.5 km NW) to the Kunda Nordic Tsement Factory (henceforth Kunda Cement Factory) (Figures 1 and 2), which has been operational since the 1870s. During the late 1970s, cement production peaked at c. 1.2 million tonnes per year (Figure 3). The dust emissions from the cement plant have fluctuated between 45,000 and 99,000 tonnes per year during the last decades of the 20th century, with the highest dust emissions recorded in 1991 (Ots and Mandre, 2012). In addition to emissions from the cement factory, the site has also received emissions from nearby alkaline generating industries and oil shale power plants, including the Balti power plant, located approximately 100 km west, and the Aru-Lõuna limestone quarry, situated 8 km to the east (Karofeld, 1996, Figure 1) which supplies raw materials for cement production at Kunda. However, significant emission reductions occurred after 1996, following the installation of pollution control filters, which

lowered emissions from 14,000 tonnes in 1996 to 530 tonnes in 2000 and 8 tonnes in 2020 (Figure 3).

The input of alkaline cement dust emissions has generated the drastic pH increase of bog's water. During 1996-1997, the pH levels taken from peat pools at Varudi varied from 7.6 to 8.5, whereas in natural conditions bog pools have a pH in the range of 3-4. Disturbances at the site are compounded by drainage and peat harvesting for horticulture, which continues to the present day, with harvesting affecting around 40% of the former centre of the site (Figure 1C). There is limited documentary information available related to the timing of drainage and peat cutting at Varudi, although historic maps (ETOMESTO, 2025; MAA-JA RUUMIAMET, 2025) indicate that most of the extensive drainage took place before 1977 and after 1944, probably relating to the large-scale drainage projects undertaken in Estonia in the 1960s under Soviet rule (Paavilainen and Päivänen, 1995). Peat extraction matching the areal extent of present-day works at the site by 1988. Prior to this, there is documentary evidence for drainage in the northern, eastern and western margins of the site having taken place around 1866 – 1911, likely indicative of the systematic forest drainage that took place in Estonia around this time (Paavilainen and Päivänen, 1995).




Figure 3. Cement annual production and emissions by the Cement Factory in Kunda. Data adapted from Trumm *et al.*, (2010), digitised from a figure at page 209 using WebPlotDigitiser: <https://automeris.io/WebPlotDigitizer/> Emissions data (available from 1985 to 2023) sourced from Raukas (1993); Pärtma (2023) and Heidelberg Materials Kunda AS (2023). Note the logarithmic scale for emission values.

The chemical composition of the cement dust emissions from Kunda consists primarily of CaO (12 – 17%), SiO₂ (6 – 9%), and several other trace metals, including lead (Pb, c. 60 mg kg⁻¹), cadmium (Cd, c. 0.9 mg kg⁻¹), and zinc (Zn, 129 mg kg⁻¹) (Mandre and Ots, 1999). The pH of the cement dust in the water solution ranged between pH 12.3 to 12.6 (Mandre and Korsjukov, 2007).

2.2. Coring method

In August 2022, an 86 cm peat core (VAR1) was extracted from Varudi peatland using a 1-meter-long Wardenaar peat corer (Wardenaar, 1987, coring location: 59°26'23"N, 26°35'55"E, Figure 1D). The core site was located near an actively harvested peat area but is within an intact section of the site in what was originally the bog's central raised dome. Vegetation was characterised by *Pinus sylvestris*, with the ground cover dominated by *Eriophorum* sp., *Menyanthes trifoliata*, *Vaccinium* sp. and scattered *Betula nana*. Both *Sphagnum* and brown mosses were also present. Despite its proximity to a drainage ditch, the coring location was representative of the overall condition of the site.

Following recovery, the core was wrapped in plastic and transported to the Faculty of Geographical and Geological Sciences at Adam Mickiewicz University, Poznań, Poland for analysis. The core was stored at 4 °C prior to sub-sampling.

2.2. Dating methods and age-depth model

Identifiable above-ground plant macrofossils were picked from 1 cm thick sub-samples taken at various depths throughout the core, following frameworks by Piotrowska *et al.* (2011) and Nilsson *et al.* (2001). *Sphagnum* stems, branches and leaves were preferentially used for dating where present. Where these were not available, above-ground remains of ericaceous plants (leaves, stems, seeds) were used instead. Initial samples were taken at 20 cm intervals throughout the core to establish a baseline chronology, which was then used to determine where additional samples would be selected. A total of 12 samples were sent for radiocarbon analysis. Each sample was pre-treated using the acid-base-acid approach and analysed by accelerator mass spectrometry (AMS) at the Poznań Radiocarbon Laboratory, Poland.

To provide a reliable chronology for recently accumulated peat, ²¹⁰Pb and ²³⁹⁺²⁴⁰Pu analyses were used on material from the upper 40 cm of the core at 1 cm contiguous resolution,

following methods outlined by Appleby (1998). Peaks in the activity of $^{239+240}\text{Pu}$, which are linked to nuclear fallout events (e.g., 1950s atmospheric nuclear tests), were also measured as independent time-markers to validate and supplement the age-depth model (Mroz *et al.*, 2017; Cwanek *et al.*, 2021). Samples were processed at the Polish Academy of Sciences' Institute of Nuclear Physics, Kraków, Poland using an AlphaAnalyst™ 7200 spectrometer (Mirion Technologies). Quality control and accuracy was ensured by measuring blanks and certified reference material (IAEA 447, IAEA 385) alongside the core samples (results are provided in Supplementary Table 1. The activity concentration of ^{210}Pb ($T_{1/2} = 22.3$ yr) was estimated by measurement of its decay product - ^{210}Po ($T_{1/2} = 138.4$ d), while $^{239+240}\text{Pu}$ was measured directly. Activity concentrations of ^{210}Pb and $^{239+240}\text{Pu}$ are reported in units of Bq kg^{-1} .

The age-depth model was constructed by integrating all ^{14}C , ^{210}Pb , and $^{239+240}\text{Pu}$ data within a Bayesian framework in R using the package 'rplum' (rplum package, Aquino-López *et al.*, 2018; Blaauw *et al.*, 2021; R working group, 2023). rplum generates maximum age probabilities at user-defined intervals (here every 1 cm), together with maximum and minimum ages based upon calculated 95% credible intervals. This method allows for the integration of ^{14}C dates with the ^{210}Pb dates without the need for re-modelling (Aquino-López *et al.*, 2018, 2020). Radiocarbon dates were calibrated in rplum using the INTCAL20 curve, with post-1950 samples using the BOMB1 curve for the Northern Hemisphere (Reimer *et al.*, 2020; Uno *et al.*, 2013). In this study, the resultant ages are expressed as calendar years (cal) CE, with 0 BP equal to 1950 cal CE.

2.3. Palaeoenvironmental proxies

2.3.1. Testate Amoebae

Samples were processed following a modified version of protocols by Hendon and Charman (1997). Samples were placed into 50 ml centrifuge tubes filled with deionised water and agitated for c. 10 minutes. These were sieved through a 300 μm mesh and the smaller fraction was retained. Sieved samples were centrifuged at >3000 rpm for 5 minutes and a sub-sample of the resultant material was transferred to a microscope slide for identification at 400x magnification. Samples were not heated or micro-sieved, to retain small but ecologically sensitive species, as recommended by Avel and Pensa (2013). A minimum count of 100 tests for each sample was considered statistically significant (Payne and Mitchell, 2008). Tests were

identified to the species level where possible, with reference to Siemensma (2023) and Mazei and Tsyganov (2006), and were later pooled into taxonomic groups defined by Amesbury *et al.* (2016). The relative abundance (%) of each taxa count was calculated for each sample. Water table depth and peat pore water pH were reconstructed using the pan-European tolerance down weighted with inverse de-shrinking transfer function model by Amesbury *et al.* (2016), based on a training set of 1302 samples spanning 35° of latitude and 55° of longitude. Reconstructions were performed using the 'Rioja' package in R (Juggins, 2019). Small species (<10-25 µm, broadly oval-shaped) not included in these groupings but present in the core from Varudi were grouped under the '*Cryptodifflugia oviformis*' group.

Stratigraphically Constrained Cluster Analysis (CONISS) was used to quantitatively define stratigraphic zones in the sub-fossil testate amoeba data (Grimm, 1987). This method is used to determine statistically significant zones, reflecting changes in testate amoebae community composition. The data were square root transformed prior to applying CONISS. The numbers of statistically significant zones were determined using Broken-Stick modelling.

We applied the framework outlined by Burge *et al.* (2023) to identify significant ecological transitions in the testate amoeba communities due to disturbance. Sub-fossil data were Hellinger transformed and analysed using the 'prcurve' function in the 'analogue' package in R (Simpson and Oksanen, 2016). Principal response curves (PrC), which reduce multi-dimensional community data to a single-dimensional curve, calculating the (dis)similarity between sample scores indicative of the difference between samples (De'ath, 1999; van Den Brink and Braak, 1999) provided the best fit for our data. A generalised additive model (GAM), an approach effective in capturing rapid and non-linear changes in palaeoecological studies, was applied to the PrC data to account for temporal autocorrelation (Auber *et al.*, 2017; Beck *et al.*, 2018; Burge *et al.*, 2023). Given the abrupt changes in our data, we also applied an adaptive spline GAM following Burge *et al.* (2023). However, this method does not account for temporal autocorrelation and thus remains incompatible with the GAM framework as outlined by Simpson (2018).

2.3.2. Plant macrofossils

Plant macrofossil analysis followed procedures adapted from Mauquoy *et al.* (2010). Samples of approximately 5 cm³ were sieved through a 200 µm test sieve and the larger fraction was

retained. Botanical composition was estimated as percentages under a low-powered microscope at between 10–100 times magnification, using a 10 x 10 grid eyepiece graticule to aid quantification of plant remains. Seeds, fruits, spindles, leaves, and wood were also identified at the species level where possible and counted as individual counts. Identification was aided by identification guides (Katz *et al.*, 1965, 1977; Grosse-Brauckmann, 1972, 1974; Tobolski, 2000; Mauquoy and van Geel, 2007; Bojnanský and Fargašová, 2007). *Sphagnum* remains were identified to sub-generic sections, with 100 leaves examined per sample, where possible, to calculate the relative abundance of each sub-section as a percentage of the total leaves identified.

2.3.3. Pollen, non-pollen palynomorphs and microscopic charcoal analyses

Past changes in vegetation cover at the landscape scale were assessed using pollen analysis (Seppä and Bennett, 2003). A total of 22 samples were prepared following the laboratory procedures outlined by Berglund and Ralska-Jasiewiczowa (1986). Each 1 cm thick sample, measuring 1 cm³, was sub-sampled at 5 cm intervals throughout the core. The samples were treated with 10% potassium hydroxide (KOH) to remove humic compounds before acetolysis. A *Lycopodium* tablet (batch no: 280521291, 18,407 spores per tablet; Manufacturer: Lund University) was added to each sample to calculate pollen concentrations, following methods by Stockmarr (1971). Samples were transferred to microscope slides and mounted in glycerine jelly for analysis. Pollen, spores, and selected non-pollen palynomorphs (NPPs) were identified and counted using a high-powered stereo microscope. Identification was based on established atlases and keys (Pollen: Moore *et al.*, 1991; Beug, 2009; NPPs: van Geel, 1978; van Geel and Aptroot, 2006; Miola, 2012). Although a target of 500 terrestrial pollen grains was aimed for per sample, this count was not always achievable due to low pollen concentrations in some core sections. The relative abundance of spores and NPPs was calculated as a proportion of the terrestrial pollen sum (TPS), which includes both arboreal (AP) and non-arboreal (NAP) pollen, excluding aquatic and wetland plant spores, Ericaceous pollen, and NPPs.

In addition to pollen counts, microscopic charcoal particles (> 10 - < 100 µm) were counted from pollen slides as past fire activity, both natural and anthropogenic (Finsinger and Tinner, 2005), while spheroidal carbonaceous particles (SCPs) counted as indicators of industrial activity (Patterson III *et al.*, 1987; Swindles, 2010). Microcharcoal concentrations per cm³

were calculated by dividing the number of particles counted with the number of *Lycopodium* spores and multiplying this by the total number of particles counted, and accumulation rates (reported as particles cm⁻³ yr⁻¹) were calculated by dividing the pollen concentration with the rPlum-derived age increments per cm slice.

2.3.4. Apparent rates of carbon accumulation

To measure apparent rates of carbon accumulation (aCAR), contiguous 1 cm-thick subsamples were taken throughout VAR1. The volume of each wet sample was determined by water displacement, and then the samples were dried in an oven at 105 °C until no further weight loss occurred. Dry bulk density was calculated by dividing the wet volume by the dry mass of each sample. Organic matter content (derived from LOI) was determined by ashing the samples at 550 °C for six hours, following the method of Chambers *et al.* (2011).

The carbon content of each sample was estimated indirectly by multiplying the LOI content by 0.52, based on the average ratio of organic carbon (OC) and LOI in ombrotrophic peat from multiple studies (Ball, 1964; Dean, 1974; Gorham, 1991; Clymo *et al.*, 1998). Carbon density was calculated by multiplying the dry bulk density (g/cm³) by the percentage of carbon content, as described by Chambers *et al.* (2011). Apparent carbon accumulation rates (aCAR) were then calculated by dividing the carbon density from each peat slice by the sedimentation rate, determined from the age-depth model (Young *et al.*, 2019, 2021).

2.3.5. μ XRF-Core Scanning (ITRAX)

To identify the section of the core affected by cement dust pollution, we followed methods similar to those used by Varvas and Punning (1993) to assess pollution histories from Estonian lake sediments. They identified rapid increases in micro-element concentrations associated with alkaline fly-ash emissions, accompanied by a decrease in organic matter, indicating the presence of particulate emissions from oil-shale combustion by power plants.

In this study, the concentrations of geochemical elements throughout the core were measured using an ITRAX μ XRF core scanner equipped with a molybdenum X-ray tube. Element concentrations were quantified as counts per second, based on the number of secondary fluorescence detected for each element over a given period. Measurements were

taken at 5 mm intervals (30 kV, 50 mA, exposure time: 30 seconds per step) at the Institute of Geography, University of Bremen, Germany. The scanner identified the activity of the following elements: Al, Si, P, S, Ar, K, Ca, Sc, Ti, V, Cr, Mn, Fe, Co, Ni, Cu, Zn, Ga, Ge, Se, Br, Rb, Sr, Y, Ba, Pt, Pb, Bi, and Fe.

The mean concentrations of each element were calculated for each 1 cm slice of the core. To account for variability in element concentrations due to sedimentation throughout the core, the results were normalised by dividing the total counts by the sum of the coherent and incoherent peaks, following the approach of Orme *et al.* (2015) as recommended by Longman *et al.* (2019). The chemical signature of cement dust was identified based on the composition of clinker emissions from the Kunda Cement factory, as detailed by Klöšeko *et al.* (2011) (Clinker emissions being the emissions produced by the production of clinker, the primary ingredient of cement, created by the high-temperature heating of limestone and other materials). Due to the uneven and unconsolidated nature of the top 6 cm of the core, this section was not scanned and was removed prior to analysis.

Principal Component Analysis (PCA) was used to summarise patterns of variation in the geochemical data, using the 'vegan' package in R (Oksanen *et al.*, 2019). To mitigate scaling effects, the data were standardised to z-scores. The analysis was conducted with varimax rotation in correlation mode, to explore correlations between elements and organic matter percentage (Silva-Sanchez *et al.*, 2014). The number of components to retain for analysis was determined using a Broken-Stick model.

3. Results.

3.1. Dating and age-depth model.

A total of 13 radiocarbon dates were analysed from core VAR1, four of which were post-bomb dates extending to a depth of 19.5 cm. The oldest dated section of the core produced a date of > 1000 yr. Radiocarbon and calibrated dates are shown in Table 1.

Table 1. Uncalibrated and calibrated radiocarbon dates from core VAR1, including depths and materials used for dating. PMC = Percent modern carbon

Code	Depth (cm)	Radiocarbon Age + error	Material dated	Calibrated dates + uncertainties
------	------------	-------------------------	----------------	----------------------------------

Poz-164520	5.5	104.51 ± 0.33 pMC	Seeds, ericaceous leaves	68.3% probability 1950 – 1957 (5.6%) 2010 – 2012 (62.7%) 95.4% probability 1956 – 1957 (10.6%) 2009 – 2012 (79.9%) 2012 – 2013 (5%)
Poz-164521	9.5	110.06 ± 0.67 pMC	Seeds, ericaceous leaves	68.3% probability 1957 – 1958 (7.5 %) 1997 – 1999 (50%) 1999 – 2000 (10.8%) 95.4% probability 1958 (11.2 %) 1996 – 2001 (85.2 %)
Poz-164662	15.5	110.36 ± 0.48 pMC	Seeds, ericaceous leaves	68.3% probability 1958 (7.3%) 1997 – 1999 (56.9%) 2000 (4.1%) 95.4% probability 1958 (11.9%) 1997 – 2000 (80.3%)
Poz-161987	19.5	118.8 ± 0.67 pMC	<i>Sphagnum</i> stems, leaves, branches	68.3% probability 1986 (68.3%) 95.4% probability 1959 (5.4%) 1960 (3.4%) 1986 - 1989 (86.7%)
Poz-162663	26.5	80±30	<i>Sphagnum</i> stems, leaves, branches	68.3% probability 1697 - 1724 (22.0%) 1814 - 1837 (20.1%) 1881 - 1913 (26.1%) 95.4% probability 1691 - 1729 (26.2%) 1809 - 1922 (69.3%)
Poz-164664	34.5	80±35	<i>Sphagnum</i> stems, leaves, branches	68.3% probability 1695 - 1725 (21.6%) 1813 - 1839 (19.6%) 1878 – 1916 (27.1%) 95.4% probability 1686 - 1725 (26.4%) 1805 - 1929 (69.0%)
Poz-161988	39.5	105±30	<i>Sphagnum</i> stems, leaves, branches	68.3% probability 1695 - 1725 (19.9%) 1813 - 1839 (17.5%) 1846 – 1853 (4%) 1869 - 1872 (1.4%) 1878 - 1916 (25.5%) 95.4% probability 1683 - 1737 (25.9%) 1803 - 1937 (69.5%)

Poz-163962	49.5	540±30	<i>Sphagnum</i> stems, leaves, branches	68.3% probability 1329 – 1338 (8.5%) 1397 - 1428 (59.8%) 95.4 % probability 1323 - 1357 (26%) 1392 - 1437 (69.5%)
Poz-161989	59.5	865±30	<i>Sphagnum</i> stems, leaves, branches	95.4% probability 1153 - 1263 (87.2%)
Poz-163963	69.5	875±30	<i>Sphagnum</i> stems, leaves, branches	95.4% probability 1126 - 1231 (79.2%) 1243 - 1258 (1.8%)
Poz-161990	79.5	910±30	<i>Sphagnum</i> stems, leaves, branches	68.3% probability 1047 - 1085 (28.6%) 1097 - 1103 (2.8%) 1126 - 1179 (30%) 1192 - 1205 (1.8%) 95.4% probability 1041 – 1214 (95.4%)
Poz-161992	85.5	985±30	<i>Sphagnum</i> stems, leaves, branches	68.3% probability 1022 - 1048 (26.8%) 1084 - 1128 (34.7%) 1140 - 1150 (6.8%) 95.4% probability 994 – 1055 (38.7%) 1076 – 1158 (56.7%)




Figure 4. Comparison between ^{210}Pb activity concentrations throughout core VAR1 and cement production. Values interpolated from a GAM fitted to annual cement production data from Kunda Cement Factory (Figure 2; Trumm *et al.*, (2010)) at the same age frequency as the VAR1 age depth model (Figure 5) using a GAMM. Age ranges are based upon the median ages from the age-depth model.

The ^{210}Pb activity profile throughout core VAR1 (Figure 4) declined throughout the core, although not in a typical monotonic pattern, with a substantial increase from 12.5 cm to 19.5 cm, indicating that the rate of ^{210}Pb accumulation was not constant throughout the core. This spike is due to the enrichment of ^{210}Pb from fly-ash fallout from the cement factory and other sources (Vaasma *et al.*, 2014; 2017). Comparison of the ^{210}Pb activities and cement production

rates applied to the age-depth model support this (Figure 4) with a significant correlation ($\tau_b = .487, p = <.0001$), indicating that ^{210}Pb accumulation in the core cannot be solely attributed to precipitation. From 32 cm core depth, ^{210}Pb activities achieve low levels, although they do not reach the background activity. Unsupported ^{210}Pb was calculated using linear regression of the last 5 samples, showing that the samples assumed background activities following the final measured samples.

The $^{239+240}\text{Pu}$ activity profile reflects the history of atmospheric deposition at the site (Figure 5). A clear peak at 28.5 cm corresponds to the onset of nuclear testing in 1945, followed by a second, larger peak at 22.5 cm, likely associated with the peak in global fallout from bomb testing in 1963, before the signing of the Partial Test Ban Treaty (Cwanek *et al.*, 2021). Smaller peaks at 16.5 cm suggest possibly the 1986 Chernobyl fallout (Ketterer *et al.*, 2004), and at 5.5 cm potentially originate from the 2011 Fukushima disaster (Bossew, 2013).




Figure 5. $^{239+240}\text{Pu}$ activity with depth in core VAR1. Selected median dates from the rPlum-derived age-depth model show good correspondence between known fallout events. Each peak may be tentatively related to a known nuclear fallout event, as shown on the right y-axis.

The age-depth relationship for core VAR1 (Figure 6) was calculated by aligning ^{210}Pb and $^{239+240}\text{Pu}$ data with the calibrated radiocarbon dates. Considering the enrichment of ^{210}Pb from industrial fallout, the assumption of a constant unsupported ^{210}Pb supply was violated.

Despite this, anchoring the model with the known 1963 peak in $^{239+240}\text{Pu}$ activity and adding a constant but uncertain reservoir effect of c. 15 years for radiocarbon dates improved the model alignment with the peaks in $^{239+240}\text{Pu}$. A radiocarbon reservoir effect is possible in *Sphagnum* peatlands, due to the recycling of 'old' gaseous carbon by mycorrhizal fungi associated with ericaceous plants near the peat surface. However, this does not usually occur when individual plant remains are dated, as was the case for our study (Piotrowska *et al.*, 2011). A more likely explanation could be the uptake of old carbon by vegetation of dissolved carbonates, particularly in the more recent samples (Madeja and Lafowski, 2008) where old carbon may derive from the buried cement dust.

The resulting age-depth model (Figure 6) indicates that peat accumulation rates were stable from the base of the core until around 1255 cal CE, after which accumulation slowed to around 0.22 yr cm^{-1} , remaining low until around 1940 cal CE, where there was a sharp acceleration to c. 2 yr cm^{-1}




Figure 6. Age-depth model for core VAR1, including calibrated radiocarbon dates and uncertainty (green) ^{210}Pb activity (Bq kg^{-1}) in blue and the calendar date for the 1963 nuclear treaty peak (orange). Uncertainties for the model are shown as the shaded area.

3.2. Peat physical and chemical properties

Figure 7 shows that activities of elements associated with clinker dust pollution began to increase around c. 1873 cal CE, marking the beginning of cement production in Kunda. There is evidence of increasing lithogenic dust deposition throughout the core, beginning as early as the mid-12th century. There is a clear negative correlation between the counts of lithogenic elements associated with cement dust (Ti, Pb, Ca, K, Fe, Cr) and organic matter content, most noticeably between c. 1942 - 2006 cal CE. Organic matter content falls sharply until c. 1988 cal CE, suggesting the accumulation of cement dust occurred during this period, mirroring the results of studies in NE Estonia lakes impacted by oil shale powerplants (Varvas and Punning, 1993; Punning *et al.*, 1996; Koff *et al.*, 2016).

The aCAR for the Varudi core is illustrated in Figure 7, which also shows the trends in LOI %, bulk density, and carbon accumulation throughout the core. Over the past millennium, the average aCAR for the entire core is $148.7 \text{ g OC m}^{-2} \text{ yr}^{-1}$. This is especially high in the upper section of the core, where carbon accumulation rates peak at $342.2 \pm 231.7 \text{ g OC m}^{-2} \text{ yr}^{-1}$. For most of the record (c. 1045 to 1910 cal CE), the mean aCAR is $72.9 \pm 28.9 \text{ g OC m}^{-2} \text{ yr}^{-1}$, aligning more closely with average values reported for Northern peatlands (Roulet *et al.*, 2007). The rate of carbon accumulation is relatively stable from the base of the core until c. 1280 cal CE, with aCAR averaging $94.5 \pm 19.8 \text{ g OC m}^{-2} \text{ yr}^{-1}$. After this point, aCAR falls to $51.3 \pm 19.7 \text{ g OC m}^{-2} \text{ yr}^{-1}$, remaining low until c. 1840 cal CE. After this date, aCAR increases significantly, reaching an average of $256.6 \pm 221.2 \text{ g OC m}^{-2} \text{ yr}^{-1}$.




Figure 7. Comparison of selected elements detected by the μ XRF core scanner, as indicated by the PCA analysis, for the detection of dust fallout from the Kunda Cement Factory. In addition to peat physical property parameters: Loss on ignition (LOI%), bulk density and apparent carbon accumulation rates (aCAR). μ XRF Data are presented as counts per second and normalised by dividing the sum of incoherent (inc) and coherent (coh) activities. The shaded area represents the section of core where most of the cement dust is concentrated. The uppermost aCAR sample was removed from the figure to aid interpretation.

The broken stick model shows that the two first components together explain a significant proportion (48.3%) of the total variance in the peat's chemical composition (PCA1: 25.5%, PCA2: 22.8%). These components highlighted elements associated with clinker dust deposition from the Kunda Cement Factory and are significantly negatively correlated with peat LOI content (Figure 8).




Figure 8. Biplot of first and second principal components showing factor loadings of individual elements scanned across core VAR1. Positively correlated variables point to the same side of the plot, while negatively correlated elements point to the opposite sides. Peat organic matter % (Derived from Loss on Ignition) content is shown in blue. The colour of the lines represents the sum factor loading of each variable for both axes, representing how strongly each variable contributes to the principal component.

3.3. Palaeoecological reconstructions

3.3.1. Plant Macrofossils

The results of plant macrofossil analysis are illustrated in Figure 9. The plant macrofossil data show three major phases of vegetation change, corresponding broadly with shifts observed in the testate amoeba and palynological records (Figures 10, 11 and 12).




Figure 9. Plant macrofossil stratigraphic diagram illustrating the changes in the botanical composition throughout core VAR1. Note the mixed data types used, percentages and total counts.

From c. 1000 cal CE, the site is dominated by *Sphagnum* mosses, particularly those of the subgenus *Sphagnum*, although by the beginning of the 12th century, *Sphagnum sub. Acutifolia* becomes the most abundant. By c. 1450 cal CE, there was an increasing abundance of monocots, likely from *Eriophorum* species and Cyperaceae, owing to the presence of spindles and fruits identified to these species. During this period, *Sphagnum* gradually declined, eventually disappearing entirely by the start of the next phase.

The latter portion of the record, starting c. 1970 cal CE, is characterised by a shift towards more ‘woody’ vegetation. Shrub-type taxa increase initially, followed by a rise in ligneous remains, particularly those of *Pinus sylvestris*. Mycorrhizal roots, bark fragments, and pine needles become significant components of the peat's botanical composition in the upper section. *Betula* is represented in the uppermost samples by its characteristic catkin scales and fruits, along with a *Betula nana* leaf recovered from the surface sample. This late phase is also characterised by a high percentage of unidentified organic material. At the top of the core, remains of the brown moss *Tomentypnum nitens*, characteristic of calcareous fens, were identified (Hájek *et al.*, 2021).

3.3.2. Testate amoeba sub-fossil communities

Reconstructed pH values were relatively stable throughout much of the record (4.1 ± 0.5). Around c. 1800 cal CE, pH began to rise, reaching a maximum of 5.4 ± 0.5 by c. 1998 cal CE. By c. 2006 cal CE until the top of the core, pH returned to pre-disturbance levels.

3.3.3. Pollen, spores, and non-pollen palynomorphs.

The results of the palynological analysis are illustrated in Figures 11 (Pollen and spores) and Figure 12 (Non-Pollen Palynomorphs- NPPs). The pollen sequence from the core VAR1 is dominated throughout by arboreal taxa, particularly *Pinus sylvestris* and *Betula* sp., with lesser contributions from shrubs such as *Calluna vulgaris* and *Vaccinium* type. The high dominance of these arboreal taxa and a low sampling resolution make detecting subtle shifts in human activity in the palynological record difficult to infer (Favre *et al.*, 2008). Other palynological studies from Estonia also report low variation in pollen assemblages and dominance of *P. sylvestris* and *Betula* across different regions, even during periods of significant land-use changes (e.g., Poska *et al.*, 2004; Veski *et al.*, 2005; Łuców *et al.*, 2022).




Figure 11. Stratigraphic plot illustrating the relative abundance of selected pollen taxa from core VAR1, as well as microcharcoal (expressed as the total number of counted fragments). Relative abundances of arboreal taxa are shaded green, disturbance indicators in pink, crops in orange, herbaceous taxa in brown and shrubs in teal. Grey shading represents 5x exaggeration. Only taxa with a minimum abundance of 2% are shown.




Figure 12. Stratigraphic plot illustrating the abundance of selected non-pollen palynomorphs from core VAR1, as well as spheroidal carbonaceous particles (SCPs) and *Pinus* stomata, both expressed as counts. Relative abundances of spores of aquatic taxa are shaded blue, algae: light green, fungi: red, invertebrate remains grey and testate amoebae in dark green. Grey shading represents 5x exaggeration. Only taxa with a minimum abundance of 2% are shown.

The oldest section of the core (c. 1000–1250 cal CE) provides little evidence of anthropogenic disturbance, indicating a largely forested landscape. This period is dominated by *Pinus sylvestris*, *Betula* sp. and *Picea abies*, suggesting a stable, predominantly woodland environment. After c. 1250 cal CE, there is a slight increase in pollen from grasses (Poaceae) from around 3% to c. 8% of the total assemblage until c. 1680 cal CE, as well as the appearance in low concentrations of plants indicative of disturbance, such as *Plantago lanceolata*, Chenopodiaceae and *Ranunculus acris* type and a general increase in *Secale* pollen which together may indicate agricultural activity, although wild populations of *Secale* are believed to have existed in Estonia (Veski, 1998; Poska et al., 2003). Despite this, arboreal pollen remains the dominant component of the record, with declines mainly affecting *Pinus sylvestris* (49 % at c. 1310 cal CE to 34% by c. 1680 cal CE) and *Picea abies* (16% at c. 1220 cal CE to 5% by c. 1680 cal CE). Tree species associated with early succession, such as *Betula*, remain largely unchanged, while *Alnus* increases during this period from c. 6% at the start of the record to c. 10% by c. 1680 cal CE. By c. 1680 cal CE, arboreal pollen has declined to its lowest relative abundance (around 82% of the total pollen sum). This decline is driven mainly by the changes in arboreal species, particularly declines in *P. sylvestris*, which becomes progressively less common through the core, a trend that continues until c. 1965 cal CE. By

c. 1680 cal CE a peak in cereals occurs, including *Hordeum*, although these are only present in low concentrations. The pollen record is stable after this period, with only slight (<1%) increases in *Calluna vulgaris* and *Vaccinium* type taxa occurring. These changes correspond with a rapid increase in microcharcoal accumulation rates, rising from 311 to 1027 particles $\text{cm}^{-3} \text{yr}^{-1}$ following the opening of the cement factory, peaking at 9837 particles $\text{cm}^{-3} \text{yr}^{-1}$ by c. 1970 cal CE. After this, *Pinus sylvestris* begins to recover, but the relative abundance of *Betula* sp. continues to rise. By c. 2010s cal CE, *Pinus sylvestris* becomes dominant once again, while *Betula* declines throughout the most recent samples, and microcharcoal accumulation reaches a maximum at c. 2010 cal CE, of 172237 particles $\text{cm}^{-3} \text{yr}^{-1}$.

The NPP record provides a more detailed picture of changes in the local environment than the pollen record. In the earliest portion of the record (c. 1000 - 1220 cal CE), fungi such as HdV-90 and HdV-13 (cf. *Entophlyctis lobata*) are common, and HdV-27 (*Bryophytomyces sphagni*) is present throughout. These fungi are typically associated with oligotrophic and ombrotrophic conditions, although HdV-90 can also thrive in more minerotrophic or poor-fen environments (van Geel, 1978; Kuhry, 1985).

Between c. 1220 - 1460 cal CE, HdV-13 increases in abundance. Around 1850 cal CE, taxa associated with the earlier section of the record start to decline, being replaced by the microalgae *Botryococcus* taxa, indicative of aquatic conditions in addition to several fungi including HdV-55A (*Sordaria* type), HdV-112 (*Cercophora* type). The most notable species in the upper section of the core during this period is HdV-201 (cf. *Xylomyces chlamydosporis*), a wood-inhabiting fungus linked to freshwater environments or pool vegetation (Goh *et al.*, 1997; Kuhry, 1997). This species is especially abundant c. 2006 cal CE, comprising around 34% of the assemblage. After c. 1950s cal CE, the diversity of NPPs declines. A key species identified in this section is *Botryococcus braunii*, a green alga typically found in environments with high levels of inorganic phosphorus, which peaks around 1985 cal CE (Órpez *et al.*, 2009) suggesting a shift toward more nutrient-enriched conditions. Overall, the record suggests a transition from nutrient-poor, *Sphagnum*-dominated peat towards an increasingly nutrient-enriched system, followed by a change towards decomposers of ligneous material.

3.3.4. Rate of change analysis

The principal response curves (PrC) explained 79% of the variance in the sub-fossil testate amoeba data from Varudi and are illustrated in Figure 13. The poor fit of GAM models to the PrCs prompted us to use adaptive splines with a GAM, which are recommended for data exhibiting abrupt changes (Burge *et al.*, 2023). However, adaptive spline GAMs cannot yet be used within the GAM framework as described by Simpson (2018). One outlier was removed due to the difference in testate amoebae composition owing to a high proportion of *Cryptodifflugia angusta* (33.5 cm).

The PrC revealed two separate periods of change at Varudi, as well as highlighting periods of rapid change associated with the onset of cement production in Kunda in 1871, and an earlier change indicated from the palynological record to coincide with increasing human activity in the landscape (Figure 11). Testate amoeba community composition was relatively consistent from c. 1040 cal CE until c. 1330 cal CE and again between c. 1645 to c. 1886 cal CE. Following c. 1998 cal CE, the direction of change in the PrC curve reverses, coinciding chronologically with the installation of filters in the cement factory in 1996. By the end of the record, testate amoeba community composition was characterised by the return of species such as *Cyclopyxis arcelloides*, *Cryptodifflugia oviformis*, and *Assulina muscorum*, which were not abundant since before c. 1970 cal CE.




Figure 13. Changes in principal response curve scores derived from sub-fossil testate amoebae data, modelled using an adaptive spline generalised additive model GAM (black line). Solid red lines indicate periods of rapid change, identified where the modelled confidence interval of the slope does not include zero. Vertical dashed lines denote key chronological events potentially influencing environmental conditions at Varudi: The onset of human activity recorded in the pollen data (c. 1250 cal CE, orange), the beginning of cement production in Kunda (1871, blue) and the installation of pollution control filters at the cement factory (1996, black).

4. Discussion

4.1. Palaeoecological assessment of ecosystem and functional changes before, during, and after intensive alkaline dust pollution

The palaeoecological record of the Varudi peatland reveals three distinct phases of botanical change occurring over the past 1,000 years: an ombrotrophic bog phase (c. 1000–1250 cal CE), a poor fen phase (c. 1250–1570 cal CE), and a wetland forest phase that continues to the present day. Unlike most research on peatland recovery, which has typically been conducted on historically drained, mined, or agricultural sites (Roderfeld, 1993; Graf *et al.*, 2008; Wagner *et al.*, 2008; Paal *et al.*, 2010), our study presents a unique pre-disturbance context for understanding the effects of atmospheric pollution upon Estonian peatlands.

The different phases of development exhibited by the site in contrast with the different periods of anthropogenic activity and disturbance are summarised in Figure 14. During the early ombrotrophic phase, low palynological richness, minimal microcharcoal counts and abundant peat organic matter indicate a largely forested landscape with limited human impact near the site, consistent with other palaeoecological records throughout Estonia during this period (Veski *et al.*, 2005). The dominance of *Sphagnum* mosses and mixotrophic testate amoeba species such as *Archerella flavum* and *Hyalosphenia papilio* reflect the relatively undisturbed and productive nature of the bog during this phase (Marcisz *et al.*, 2020; Łuców *et al.*, 2022). During this period, water table depths were high, and pH levels were low, consistent with natural conditions for ombrotrophic bogs in the Northern Hemisphere (Warner and Asada, 2006).




Figure 14: Summary of key trends in the Varudi peatland record. Shown are changes in reconstructed pH and water table depth, percentages of non-arboreal pollen (serving as a proxy for forest clearance related to agricultural activity), apparent carbon accumulation rates (aCAR), periods of significant change (as identified by principal response curves with Generalised Additive Models, GAMs), and a general timeline of environmental shifts. Dashed horizontal lines represent key historical events relevant to the Varudi record: the yellow line marks the approximate onset of human activity c. 1250 cal CE, possibly linked to the Livonian Crusades; the blue line indicates the establishment of the Kunda Cement Factory in 1871; the black line denotes the installation of effective filters at the Kunda Cement Factory; and the grey-shaded area highlights the section of the core with substantial cement dust, identified through LOI and μ XRF core scanning.

The beginning of agricultural activity around c. 1250 cal CE is broadly coincident with the German-Danish crusaders occupation of Estonia starting in 1227 CE (Veski *et al.*, 2005). From c. 1250 to 1570 cal CE, increased Poaceae, *Secale cereale*, and field weed pollen indicates a shift toward more open, meadow-like conditions (Poska and Saarse, 2002; Niinemets and Saarse, 2009), which may relate to the environmental effects resulting from economic changes resulting from the Teutonic order crusades, similar to those identified by Brown and Pluskowski (2011) in a peatland record from northern Poland.

Declines in *Pinus sylvestris* and *Picea abies*, alongside increases in early successional species such as *Betula* and *Alnus*, and minute increases in microcharcoal accumulation rates (Figure 11) suggest land clearance using slash-and-burn methods in the surrounding landscape (Jaats *et al.*, 2010). While arboreal pollen remained dominant, indicating continued forest cover around the Varudi peatland, mineral soil enrichment of the peat suggests overland flow or

aeolian deposition resulting from land clearance. Similar patterns of early human-induced soil erosion have been recorded in peatland records from central Estonia (Heinsalu and Veski, 2010). This is reflected by decreasing organic matter content and increasing concentrations of lithogenic elements such as Ca and Ti, patterns like those observed in Estonian lake sediment records following land clearance events (Koff *et al.*, 2016; Vandel and Vaasma, 2018). Ti, a conservative lithogenic element, indicates soil erosion from the surrounding catchment (Boyle, 2001; Boës *et al.*, 2011).

By c. 1570 cal CE, the site had transitioned into a wetter, more minerotrophic fen, indicated by the gradual replacement of *Sphagnum* by monocots such as *Carex* sp. and *Eriophorum vaginatum* in the plant macrofossil record and an increase in *Sphagnum* section *Cuspidata* remains and subsequent declines in *Sphagnum* subsection *Acutifolia* and subsection *Sphagnum* (Figure 9), and a shift towards wetter conditions during this period indicated by the testate amoebae reconstruction, possibly indicating permanent or seasonal standing water at this time (Figure 10). This interpretation is consistent with observations from contemporary bog pools, where *Sphagnum* is often rare or absent in shallow or temporary water bodies on a peatland's surface, while monocots such as *Eriophorum* species are abundant, as exemplified by the British National Vegetation Classification (NCV) bog pool community classification (M3) (Rodwell, 1998). Nutrient enrichment due to soil erosion resulting from forest clearance and agricultural activities in the landscape at this time likely contributed to this shift (Hölzer and Hölzer, 1998).

The establishment of the Kunda cement factory in 1871 brought significant ecological changes to the site. Substantial increases in Ti, Ca, Fe, and other elements associated with cement dust fallout occurred along with increased pH and declining organic matter content, indicating the chemical influence of cement dust pollution upon the peat (Figures 7 and 8). This pollution appears to have driven rapid changes in testate amoeba communities by c. 1890 cal CE, as seen in the PrC record (Figures 13 and 14). These results are consistent with previous palaeoecological studies on alkaline pollution impacts in Estonian peatlands and lakes (Varvas and Punning, 1993; Koff *et al.*, 2016; Vandel and Vaasma, 2018; Vellak *et al.*, 2014). The enrichment of the peat by cement dust is reported to have had a fertiliser-like effect on ligneous and ericaceous species, increasing tree cover, size, and stand density within the industrial region of Estonia at this time (Pensa *et al.*, 2004; Ots *et al.*, 2011).

The concurrent hydrological changes experienced by the site, characterised by persistent water table lowering after c. 1960, were likely driven by drainage and enhanced transpiration rates following tree and shrub expansion onto the site (Stelling *et al.*, 2023). The simultaneous compounding effect of hydrological change upon the site confounds interpretation of the effects of alkaline pollution at Varudi peatland, as it is likely that the changes that occurred at this time were driven by the combined effects of both disturbances.

From 1955 CE onward, cement production increased, reflected in peaks of Ca, Ti, and K, alongside rising bromine (Br) levels, coincident with increased microcharcoal accumulation rates (increasing from c. 11 to 58 particles yr⁻¹. The halogen Br is often linked to changes in storminess in peat palaeoecological studies due to being commonly sourced from marine aerosols (e.g., Shotyk, 1997; Turner *et al.*, 2014). Here, elevated Br levels correlate strongly with cement dust indicators, suggesting that increased Br in the core corresponds with enhanced plant decay and oxidation of abiotic organic matter, resulting in the formation of methyl bromide (Lee-Taylor and Holland, 2000; Keppler *et al.*, 2000). The increased presence of fungi in addition to microalgae species *Botryococcus braunii* in the palynological and plant macrofossil records further suggests that the soil decomposition rate increased during this period, coincident with periods of increased seasonal wetness (Barthelmes *et al.*, 2012; Defrenne *et al.*, 2023; Buttler *et al.*, 2023; Thorman *et al.*, 2003) (Figure 12).

Following the restoration of Estonia's independence in 1991, more rigorous environmental regulations resulted in a reduction of industrial emissions (Kask *et al.*, 2008). The effect of this decline is evident in our palaeoecological record, with declining pollution markers and pH returning to pre-disturbance levels after 1996 (Figures 7 and 10), as well as the apparent (albeit limited) recovery of testate amoeba communities shown by the PrC curve (Figure 13). Despite these improvements, the site has remained densely forested and enriched, with a persistently lowered water table. We find no evidence for botanical succession toward pre-disturbance conditions in the plant macrofossil record (Figure 9), while the presence of calciphilous *Tomentypnum nitens* at the core surface reflects the legacy of alkaline conditions at the site (Malmer *et al.*, 1992; Hájek *et al.*, 2006; Apolinarska *et al.*, 2024), likely resulting from the movement of mineral-rich water, enriched by the buried cement dust as it is transported upwards via capillary rise, indicating that these cement deposits continue to influence surface water chemistry and vegetation communities at the site, although *Sphagnum* has returned to the site since its disappearance, indicating a degree of

recovery. Identification and monitoring of the species present on the site could provide additional insight into the extent of this recovery.

4.2. Ecosystem Recovery After Reduction in Atmospheric Alkaline Pollution

Our study found evidence of recovery in PrC of the sub-fossil testate amoeba assemblages (Figure 13). The community composition of these environmentally sensitive proxies showed signs of turnover nearly 30 years after installing filters at the Kunda cement factory. However, the extent of recovery indicated by the testate amoebae is limited, as the community composition in the top sediment layer more closely resembles conditions from c. 1977 cal CE, the peak period of cement production at Varudi (Figures 3 and 13).

We used adaptive spline GAMs to analyse our data, which while suitable for datasets exhibiting abrupt changes such as the example from Varudi (Burge *et al.*, 2023), they cannot yet be incorporated into the Generalized Additive Mixed Model (GAMM) framework described by Simpson *et al.* (2018) and are more susceptible to issues such as temporal autocorrelation and boundary effects (Simpson, 2018; Burge *et al.*, 2023). While the model best fit our data, more evidence is needed to draw conclusive interpretations about recovery than these results alone. More recently accumulated samples may be more influenced by temporal autocorrelation (Burge *et al.*, 2023). Some testate amoeba species may infiltrate the sub-fossil community composition downcore, making near-surface communities appear more similar (Liu *et al.*, 2024). Another limitation of this study is that our results are based upon reconstructions taken from a single core and as such do not account for the spatial heterogeneity characteristic of peatlands. Although palaeoecological trends tend to be well replicated across multiple cores within a given site (e.g., Barber *et al.*, 1999; Hendon *et al.*, 2001), it may be that the trends reconstructed from VAR1 represent conditions specific to the sampling site, rather than the whole site (Bacon *et al.*, 2017). Paal *et al.* (2010) examined peatland vegetation communities in eleven bogs in northeastern Estonia at varying distances from sources of atmospheric alkaline pollution, finding evidence of *Sphagnum* re-establishment and the recovery of other bog-specific plant species at several sites as surface waters re-equilibrated to pre-pollution conditions, particularly within microforms that are less affected by contaminated groundwater. Therefore, recovery at Varudi may be spatially heterogenous, with some areas recovering more quickly depending upon their contact with polluted peat layers.

Studies investigating the recovery of vegetation communities following alkaline pollution show that ecosystem recovery may be limited due to the present, degraded condition of the ecosystem. For example, Vellak *et al.* (2014) studied bryophyte recovery following reductions in atmospheric fly-ash pollution across Estonia and northwestern Russian sites. They found that bryophyte species growing in more heavily affected sites tended to be adapted for growing in low light conditions, due to competing with the larger and more dense vascular plants that encroached on these sites, in response to the more enriched and alkaline conditions for which they are better adapted than bog-specific vegetation (Pärtel *et al.*, 2004; Zvereva *et al.*, 2008a, 2008b). Increased nutrient and litter supply and root exudates, coupled with the faster growth of vascular plants and trees may further hinder the reestablishment of bog and fen communities, delaying or preventing recovery (Konings *et al.*, 2019). Gunnarsson *et al.* (2002) demonstrated that nutrient enrichment, often accompanied by higher pH, can give vascular plants a competitive advantage over bog-specific vegetation (Limpens *et al.*, 2003; Dieleman *et al.*, 2015).

4.3. How has the pollution impacted apparent rates of carbon accumulation at the site?

Each of the phases in ecosystem development in the palaeoecological record corresponds with substantial changes in the apparent carbon accumulation rate (aCAR) throughout the Varudi record (Figure 7). Between c. 1000 to 1250 cal CE, when the site was in a relatively pristine, undisturbed condition, aCAR rates were higher ($32.0 \pm 10.7 \text{ g C m}^{-2} \text{ yr}^{-1}$) than the average for northern hemisphere ombrotrophic peatlands ($23 \pm 2 \text{ g C m}^{-2} \text{ yr}^{-1}$) (Korhola *et al.*, 1995; Yu *et al.*, 2009). Throughout the transition from bog to poor fen, from c. 1360 to 1570 cal CE, average aCAR rates most likely initially increased ($38.5 \pm 34.3 \text{ g C m}^{-2} \text{ yr}^{-1}$). However, they were highly variable, likely reflecting the incorporation of vascular plant roots from the overlying forested phase, introducing younger carbon into deeper sediments. Between c. 1630 and 1800 cal CE, rates fell as low as $19.0 \pm 5.4 \text{ g C m}^{-2} \text{ yr}^{-1}$, aligning with reported values for Finnish fens (Korhola *et al.*, 1995). Overall, minerotrophic fens typically exhibit lower carbon accumulation rates than ombrotrophic bogs (Loisel and Bunsen, 2020; Yang *et al.*, 2023).

Following the succession from poor fen to forested fen from c. 1871 cal CE to the present, there is an apparent increase in aCAR, especially in the most recently accumulated peat ($280.7 \pm 211.3 \text{ g C m}^{-2} \text{ yr}^{-1}$). Due to the incomplete decomposition of labile organic matter at the peat

surface, care must be taken when interpreting recently accumulated carbon from peat core records (Young *et al.*, 2019; 2021). This increase is likely due to high litter deposition from trees and shrubs, and the rapid decline in aCAR downcore suggests that this labile material is rapidly cycled back into the atmosphere rather than sequestered in the soil. Our results suggest that the current vegetation composition, dominated by trees and shrubs, is less effective for long-term carbon sequestration compared to earlier phases. While the water table remains reduced, the site shows limited potential for recovery to its original ombrotrophic condition due to ongoing drainage. At the same time, nutrient inputs and evapotranspiration from the overlying trees compound this limiting factor.

4.4. Indicators of critical thresholds to assess peatland condition and recovery due to alkaline pollution

By comparing our pH reconstruction with significant successional shifts at the site, we may infer the thresholds at which transitions in the steady state of the ecosystem occurred in response to changes in alkalinity at Varudi bog. Defining tipping points that may be broadly applied to other sites allows for predicting ecosystem shifts in response to global change, allowing for prediction of such shifts in the future (Munson *et al.*, 2018). To our knowledge, our study represents the first step towards defining thresholds for tipping points in peatland ecosystems in response to changes in alkalinity.

Around c. 1800 cal CE, the reconstructed pH for Varudi showed an increasing trend from the previous, relatively stable conditions, reaching a maximum of 5.4 ± 0.5 by c. 1990s cal CE (Figure 10). This value aligns with pH measurements for polluted sites in Estonia, supporting the reliability of our reconstruction (Paal *et al.*, 2010). We find that the transition from an ombrotrophic bog to fen-like conditions, driven by mineral soil enrichment resulting from land clearance for agriculture in the surrounding landscape, was associated with an increase of pH levels from c. pH 3.8 to pH 4.1 which represents (due to the logarithmic scale of pH) nearly a twofold increase in alkalinity. Likewise, the subsequent shift from fen-like to forested conditions corresponded with a small average increase in pH to c. 4.3, or c. 1.5 times more alkaline than the previous state and more than three times more alkaline than during the bog phase. Our results suggest that relatively small increases in pH (increases in pH of 0.2 – 0.3) may result in critical ecosystem transitions. However, we stress that the threshold values defined here are highly uncertain, owing to the limited predictive ability of the transfer

function models used for our pH reconstruction and the relatively significant uncertainties associated with transfer function models in general (Amesbury *et al.*, 2016). Higher pH values measured from bog pools at the site in 1996 – 1997 of 7.6 - 8.5 are higher than our estimates for this period, and by 2022 pH had only fallen to 6.3 - 6.8 (Pärtma, 2023), suggesting that our reconstruction underestimates pH levels experienced by the site by a wide margin, possibly due to the removal of high-pH samples from Amesbury *et al.*, (2016)'s dataset, causing the range of pH measurements to be low and biased towards more acidic conditions. This may warrant future re-evaluation of our data using transfer functions specifically developed for reconstructing pH in polluted peatlands. In addition, we cannot rule out the possibility that other impacts of cement deposition (e.g., reduced photosynthesis rates due to dust deposition) and, significantly- drainage; may have also played a role in driving botanical changes.

Contemporary field and lab experiments that manipulate the alkalinity of peatland soils directly may provide more precise values for ecosystem thresholds in response to alkaline pollution. However, to date few studies have done this for alkalinisation: one example is Kang *et al.* (2018) who conducted a series of field and laboratory pH manipulation experiments across seven peatlands in the UK, Japan, Indonesia and South Korea, finding that increases in pH resulted in significant changes in microbial composition, resulting in increased phenol oxidase activity and enhanced DOC releases. Another is the long-term (active since 2002) Whim Bog experiment near Edinburg, Scotland (Sheppard *et al.*, 2011; Levy *et al.*, 2019), where the effects of enrichment upon a peatland by gaseous ammonia (NH_3) and wet-deposited ammonium (NH_4^+) and nitrate (NO_3^-) are compared. Analysis by Sutton *et al.* (2020) indicates that vegetation declined three times more quickly when exposed to gaseous ammonia and three times for ammonium than for a similar dose of nitrate.

A significant limitation of these studies in defining tipping points is that the long timescale necessary for critical transitions to occur in some cases typically exceeds the lifespan of most observational and experimental studies (Taranu *et al.*, 2018; Lamentowicz *et al.*, 2019). Therefore, further palaeoecological work across a more extensive range of sites or with multiple cores from within a single site may also advance our understanding of ecosystem tipping points in response to alkalinisation. Furthermore, disentangling the effects of alkalinisation from those of drainage is challenging, as both factors may influence plant

community composition, carbon accumulation rates, overall ecosystem functioning (Word *et al.*, 2022) and possibly resilience to alkalinisation in peatlands. Since significant drainage and cement production appear to have begun at roughly the same period in the site's history, some of the ecological variability attributed here to cement dust pollution may instead be the result of drainage impacts. However, it is notable that despite reconstructed pH values increasing following the cessation of significant emissions from Kunda, water table depths have not recovered. This indicates that drainage is unlikely to have influenced the apparent modest ecosystem recovery observed at Varudi in recent decades.

5. Conclusion

This study is the first to investigate the long-term impacts of alkaline emissions on a peatland over centennial timescales. It establishes the first attempt to define threshold indicator values for ecosystem tipping points in response to alkalinisation. Our findings demonstrate that alkaline pollution, along with later drainage, has had a profound influence upon ecosystem development at Varudi for more than 750 years, with ecosystem succession following both low-level, sustained mineral soil enrichment due to agricultural activities and intensive fly-ash fallout sustained over 160 years. We find that an increase in pH of 0.2 to 0.3 (corresponding to, approximately, a two to threefold increase in alkalinity) is sufficient to trigger a critical ecosystem shift. These can lead to a long-term decline in carbon accumulation over long timescales, and such changes may be slow to recover or permanent, even if the point source of pollution is eliminated. Our results, while insightful, have limitations that underscore the need for additional experimental and palaeoecological research to assess peatland responses and resilience to alkalinisation across a range of spatial and temporal scales. This would allow a better understanding of the timescales required for peatland recovery and how these ecosystem transitions influence greenhouse gas dynamics from affected sites. Alkalinisation poses a growing threat to peatlands worldwide and is a developing challenge for the 21st century (Sutton *et al.*, 2020). As carbon-rich ecosystems, peatlands are important in future atmospheric greenhouse gas concentrations. Understanding how peatlands will respond to future alkalinisation is essential for predicting the role of peatlands in climate change mitigation.

6. Acknowledgements

899 The study was funded by the National Science Centre, Poland, grant no
900 2021/41/B/ST10/00060. We wish to thank the Estonian Research Council (projects PRG1993
901 and TK215) for providing additional financial support. Thanks to Eliise Kara, Merlin Liiv and
902 Triin Reitalu for their assistance in the field.

903 Data availability statement:

904 All relevant data files described here are available either in the supplementary materials or
905 via FigShare, accessible via the following link: <https://figshare.com/s/acfe140783b6f871f1f2>

906 Author contributions:

907

908 LOA prepared the manuscript with contributions from all authors. ML, MS and SV
909 conceptualised the study and secured funding. LOA, ML, MS, KM, LA and NS participated in
910 fieldwork. LA, SC, MB, ML, PK, MKK, EŁ, AC and MAL conducted formal analyses. SV and AH
911 provided historical data and photographs.

912 Declaration of interests:

913 The authors declare they have no known financial interests or personal relationships that
914 could influence the work represented in this manuscript.

7. References

- Abril, G.A., Wannaz, E.D., Mateos, A.C., Pignata, M.L.: Biomonitoring of airborne particulate matter emitted from a cement plant and comparison with dispersion modelling results. *Atmos. Environ.* **82**, 154–163. <https://doi.org/10.1016/j.atmosenv.2013.10.020>, 2014.
- Amesbury, M.J., Swindles, G.T., Bobrov, A., Charman, D.J., Holden, J., Lamentowicz, M., Mallon, G., Mazei, Y., Mitchell, E.A., Payne, R.J. and Roland, T.P.: 2016. Development of a new pan-European testate amoeba transfer function for reconstructing peatland palaeohydrology. *Quat. Sci. Rev.*, **152**, pp.132–151. <https://doi.org/10.1016/j.quascirev.2016.09.024>, 2016.
- Apolinarska, K., Pleskot, K., Aunina, L., Marzec, M., Szczepaniak, M., Kabaciński, M., Kiełczewski, R. and Gałka, M.: Late Holocene changes in the water table at an alkaline fen in Central Latvia: Their impacts on CaCO₃ deposition at the fen and relation to the hydroclimate patterns of the Eastern Baltic Region. *Quat. Sci. Rev.*, **334**, p.108717. <https://doi.org/10.1016/j.quascirev.2024.108717>, 2024.
- Appleby, P.G.: *Dating recent sediments by 210 Pb: problems and solutions* (No. STUK-A--145), 1998.
- Aquino-López, M.A., Blaauw, M., Christen, J.A. and Sanderson, N.K.: Bayesian analysis of 210 Pb dating. *J. of Agric. Biol. and Environ. Statistics*, **23**(3), pp.317–333. <https://doi.org/10.1007/s13253-018-0328-7>, 2018.
- Aquino-López, M.A., Ruiz-Fernández, A.C., Blaauw, M. and Sanchez-Cabeza, J.A.: Comparing classical and Bayesian 210Pb dating models in human-impacted aquatic environments. *Quat. Geochronol.*, **60**, p.101106. <https://doi.org/10.1016/j.quageo.2020.101106>, 2020.
- Auber, A., Travers-Trolet, M., Villanueva, M.C. and Ernande, B.: A new application of principal response curves for summarizing abrupt and cyclic shifts of communities over space. *Ecosphere*, **8**(12), p.e02023. <https://doi.org/10.1002/ecs2.2023>, 2017.
- Avel, E. and Pensa, M.: Preparation of testate amoebae samples affects water table depth reconstructions in peatland palaeoecological studies. *Estonian J. of Earth Sci.*, **62**(2), p.113. doi:10.3176/earth.2013.09, 2013.
- Bacon, K.L., Baird, A.J., Blundell, A., Bourgault, M.A., Chapman, P.J., Dargie, G., Dooling, G.P., Gee, C., Holden, J., Kelly, T.J. and McKendrick-Smith, K.A.: Questioning ten common assumptions about peatlands. *Mires and Peat*, **19**, pp.1–23, 2017.

- Ball, D.F.: Loss-on-ignition as an estimate of organic matter and organic carbon in non-calcareous soils. *J. of Soil Sci.*, 15(1), pp.84-92, 1964.
- Barber, K.E., Battarbee, R.W., Brooks, S.J., Eglinton, G., Haworth, E.Y., Oldfield, F., Stevenson, A.C., Thompson, R., Appleby, P.G., Austin, W.E.N. and Cameron, N.G.: Proxy records of climate change in the UK over the last two millennia: documented change and sedimentary records from lakes and bogs. *J. of the Geol. Soc.*, 156(2), pp.369-380. <https://doi.org/10.1144/gsjgs.156.2.0369>, 1999.
- Barthelmes, A., de Klerk, P., Prager, A., Theuerkauf, M., Unterseher, M. and Joosten, H.: Expanding NPP analysis to eutrophic and forested sites: Significance of NPPs in a Holocene wood peat section (NE Germany). *Rev. of Palaeobot. and Palynol.*, 186, pp.22-37. <https://doi.org/10.1016/j.revpalbo.2012.07.007>, 2012.
- Beck, K.K., Fletcher, M.S., Gadd, P.S., Heijnis, H., Saunders, K.M., Simpson, G.L. and Zawadzki, A.: Variance and rate-of-change as early warning signals for a critical transition in an aquatic ecosystem state: a test case from Tasmania, Australia. *J. of Geophysical Res.: Biogeosciences*, 123(2), pp.495-508. <https://doi.org/10.1002/2017JG004135>, 2018.
- Berglund, B.E. and Ralska-Jasiewiczowa, M.: Pollen analysis and pollen diagrams. *Handbook of Holocene palaeoecology and palaeohydrology*, 455, pp.484-486, 1986.
- Beug, H. J.: Hans-Jürgen Beug, Leitfaden der Pollenbestimmung für Mitteleuropa und angrenzende Gebiete. *Germania: Anzeiger der Römisch-Germanischen Kommission des Deutschen Archäologischen Instituts*, pp.699-702, 2009.
- Blaauw, M., Christen, J.A., Aquino-Lopez, M.A., Esquivel-Vazquez, J., Belding, T., Theiler, J., Gough, B., Karney, C. and Blaauw, M.M.: *Package 'rplum'*, 2021.
- Bobbink, R., Hornung, M. and Roelofs, J.G.: The effects of air-borne nitrogen pollutants on species diversity in natural and semi-natural European vegetation. *J. of Ecol.*, 86(5), pp.717-738. <https://doi.org/10.1046/j.1365-2745.1998.8650717.x>, 1998.
- Boës, X., Rydberg, J., Martinez-Cortizas, A., Bindler, R. and Renberg, I.: Evaluation of conservative lithogenic elements (Ti, Zr, Al, and Rb) to study anthropogenic element enrichments in lake sediments. *J. of Paleolimnology*, 46, pp.75-87. <https://doi.org/10.1007/s10933-011-9515-z>, 2011.
- Bojčanský, V. and Fargašová, A.: *Atlas of seeds and fruits of Central and East-European flora: the Carpathian Mountains region*. Springer Science & Business Media.
- Bossew, P., 2013. Anthropogenic radionuclides in environmental samples from Fukushima Prefecture. *Radiat Emerg Med*, 2(1), pp.69-75, 2007.

- Boyle, J.F.: 2001. Inorganic geochemical methods in palaeolimnology. *Tracking environmental change using lake sediments: physical and geochemical methods*, pp.83-141. https://doi.org/10.1007/0-306-47670-3_5, 2001.
- Bridgham, S.D., Pastor, J., Dewey, B., Weltzin, J.F. and Updegraff, K.: Rapid carbon response of peatlands to climate change. *Ecology*, 89(11), pp.3041-3048. <https://doi.org/10.1890/08-0279.1>, 2008.
- Brown, A. and Pluskowski, A., 2011. Detecting the environmental impact of the Baltic Crusades on a late-medieval (13th–15th century) frontier landscape: palynological analysis from Malbork Castle and hinterland, Northern Poland. *J. of Archaeol. Sci.*, 38(8), pp.1957-1966. <https://doi.org/10.1016/j.jas.2011.04.010>
- Burge, O.R., Richardson, S.J., Wood, J.R. and Wilmshurst, J.M.: A guide to assess distance from ecological baselines and change over time in palaeoecological records. *The Holocene*, 33(8), pp.905-917. <https://doi.org/10.1177/09596836231169986>, 2023.
- Buttler, A., Bragazza, L., Laggoun-Défarge, F., Gogo, S., Toussaint, M.L., Lamentowicz, M., Chojnicki, B.H., Słowiński, M., Słowińska, S., Zielińska, M. and Reczuga, M.: Ericoid shrub encroachment shifts aboveground–belowground linkages in three peatlands across Europe and Western Siberia. *Global Change Biology*, 29(23), pp.6772-6793. <https://doi.org/10.1111/gcb.16904>, 2023.
- Chambers, F.M., Beilman, D.W. and Yu, Z.: Methods for determining peat humification and for quantifying peat bulk density, organic matter and carbon content for palaeostudies of climate and peatland carbon dynamics. *Mires and Peat*, 7(7), pp.1-10, 2011
- Charman, D.J.: Biostratigraphic and palaeoenvironmental applications of testate amoebae. *Quat. Sci. Rev.*, 20(16-17), pp.1753-1764. [https://doi.org/10.1016/S0277-3791\(01\)00036-1](https://doi.org/10.1016/S0277-3791(01)00036-1), 2001.
- Clymo, R.S., Turunen, J. and Tolonen, K.: Carbon accumulation in peatland. *Oikos*, pp.368-388. <https://doi.org/10.2307/3547057>, 1998
- Cwanek, A., Łokas, E., Mitchell, E.A., Mazei, Y., Gaca, P. and Milton, J.A.: Temporal variability of Pu signatures in a 210Pb-dated Sphagnum peat profile from the Northern Ural, Russian Federation. *Chemosphere*, 281, p.130962. <https://doi.org/10.1016/j.chemosphere.2021.13096>, 2021.

1013 De'ath, G.: Principal curves: a new technique for indirect and direct gradient
 1014 analysis. *Ecology*, 80(7), pp.2237-2253. [https://doi.org/10.1890/0012-](https://doi.org/10.1890/0012-9658(1999)080[2237:PCANTF]2.0.CO;2)
 1015 [9658\(1999\)080\[2237:PCANTF\]2.0.CO;2](https://doi.org/10.1890/0012-9658(1999)080[2237:PCANTF]2.0.CO;2), 1999.

1016 Dean, W.E.: Determination of carbonate and organic matter in calcareous sediments and
 1017 sedimentary rocks by loss on ignition; comparison with other methods. *J. of*
 1018 *Sediment. Res.*, 44(1), pp.242-248. [https://doi.org/10.1306/74d729d2-2b21-11d7-](https://doi.org/10.1306/74d729d2-2b21-11d7-8648000102c1865d)
 1019 [8648000102c1865d](https://doi.org/10.1306/74d729d2-2b21-11d7-8648000102c1865d), 1974.

1020 Defrenne, C.E., Moore, J.A., Tucker, C.L., Lamit, L.J., Kane, E.S., Kolka, R.K., Chimner, R.A.,
 1021 Keller, J.K. and Lilleskov, E.A.: Peat loss collocates with a threshold in plant–
 1022 mycorrhizal associations in drained peatlands encroached by trees. *New*
 1023 *Phytologist*, 240(1), pp.412-425. <https://doi.org/10.1111/nph.18954>, 2023.

1024 Dieleman, C.M., Branfireun, B.A., McLaughlin, J.W. and Lindo, Z.: Climate change drives a
 1025 shift in peatland ecosystem plant community: implications for ecosystem function
 1026 and stability. *Glob. Change Biol.*, 21(1), pp.388-395. DOI: [10.1111/gcb.12643](https://doi.org/10.1111/gcb.12643), 2015.

1027 ETOMESTO: Online maps in real time - satellite and City: [http://www.etomesto.com/map-](http://www.etomesto.com/map-mir/?y=59.444377&x=26.592107)
 1028 [mir/?y=59.444377&x=26.592107](http://www.etomesto.com/map-mir/?y=59.444377&x=26.592107), last Access: 10 August 2025.

1029 Favre, E., Escarguel, G., Suc, J.P., Vidal, G. and Thévenod, L.: A contribution to deciphering
 1030 the meaning of AP/NAP with respect to vegetation cover. *Review of Palaeobot. and*
 1031 *Palynol.*, 148(1), pp.13-35. <https://doi.org/10.1016/j.revpalbo.2007.08.003>, 2008.

1032 Fiałkiewicz-Kozieł, B., De Vleeschouwer, F., Mattielli, N., Fagel, N., Palowski, B., Pazdur, A.
 1033 and Smieja-Król, B.: Record of Anthropocene pollution sources of lead in disturbed
 1034 peatlands from Southern Poland. *Atmos. Environm.*, 179, pp.61-68
 1035 <https://doi.org/10.1016/j.atmosenv.2018.02.002>, 2018.

1036 Finsinger, W. and Tinner, W.: Minimum count sums for charcoal concentration estimates in
 1037 pollen slides: accuracy and potential errors. *The Holocene*, 15(2), pp.293-297.
 1038 <https://doi.org/10.3389/fevo.2018.00149>, 2005.

1039 Frolking, S., Roulet, N., Fuglestad, J.: How northern peatlands influence the Earth's
 1040 radiative budget: Sustained methane emission versus sustained carbon
 1041 sequestration. *J. Geophys. Res. Biogeosciences* 111.
 1042 <https://doi.org/10.1029/2005JG000091>, 2006.

1043 Goh, T.K., Ho, W.H., Hyde, K.D. and Umali, T.E.: New records and species of *Sporoschisma*
 1044 and *Sporoschismopsis* from submerged wood in the tropics. *Mycological*

1045 *research*, 101(11), pp.1295-1307. <https://doi.org/10.1017/S0953756297003973>,
1046 1997.

1047 Gomes, H.I., Mayes, W.M., Rogerson, M., Stewart, D.I. and Burke, I.T.: Alkaline residues and
1048 the environment: a review of impacts, management practices and opportunities. *J.*
1049 *of Clean. Prod.*, 112, pp.3571-3582. <https://doi.org/10.1016/j.jclepro.2015.09.111>,
1050 2016.

1051 Gorham, E.: Northern peatlands: role in the carbon cycle and probable responses to climatic
1052 warming. *Ecolog. Appl.*, 1(2), pp.182-195. <https://doi.org/10.2307/1941811>, 1991.

1053 Graf, M.D., Rochefort, L. and Poulin, M.: Spontaneous revegetation of cutaway peatlands
1054 of North America. *Wetlands*, 28, pp.28-39. <https://doi.org/10.1672/06-136.1>, 2008.

1055 Grimm, E.C.: CONISS: a FORTRAN 77 program for stratigraphically constrained cluster
1056 analysis by the method of incremental sum of squares. *Computers & Geosci.*, 13(1),
1057 pp.13-35. [https://doi.org/10.1016/0098-3004\(87\)90022-7](https://doi.org/10.1016/0098-3004(87)90022-7), 1987.

1058 Grosse-Brauckmann, G.: *Über pflanzliche Makrofossilien mitteleuropäischer Torfe. I.*
1059 *Gewebereste krautiger Pflanzen und ihre Merkmale* (On plant macrofossils in central
1060 European peat. I. Remnants of vascular plant tissues and their characteristics).
1061 *Telma*, 2, 19 55 (in German), 1972.

1062 Grosse-Brauckmann, G.: *Über pflanzliche Makrofossilien mitteleuropäischer Torfe. II.*
1063 *Weitere Reste (Früchte und Samen, Moose u.a.) und ihre Bestimmungsmöglichkeiten*
1064 (On plant macrofossils in central European peat. II. Other remnants (e.g. fruits and
1065 seeds, mosses) and possibilities for their identification). *Telma*, 4, 51–117 (in
1066 German), 1974.

1067 Gunnarsson, U., Malmer, N. and Rydin, H.: Dynamics or constancy in Sphagnum dominated
1068 mire ecosystems? A 40-year study. *Ecography*, 25(6), pp.685-
1069 704. <https://doi.org/10.1034/j.1600-0587.2002.250605.x>, 2002.

1070 Hájek, M., Horsák, M., Hájková, P. and Dítě, D.: Habitat diversity of central European fens in
1071 relation to environmental gradients and an effort to standardise fen terminology in
1072 ecological studies. *Perspect. Plant Ecol. Evol. and Syst.*, 8(2), pp.97-114.
1073 <https://doi.org/10.1016/j.ppees.2006.08.002>, 2006.

1074 Hájek, M., Hájková, P., Apostolova, I., Sopotlieva, D., Goia, I., Dítě, D.: The vegetation of rich
1075 fens (*Sphagno warnstorffii*-*Tomentypnion nitentis*) at the southeastern margins of
1076 their European range. *Veg. Class. and Surv.* 2, 177–190.
1077 <https://doi.org/10.3897/vcs/2021/69118>, 2021.

1078 Harenda, K.M., Lamentowicz, M., Samson, M., Chojnicki, B.H.: The Role of Peatlands and
 1079 Their Carbon Storage Function in the Context of Climate Change, in: Zielinski, T.,
 1080 Sagan, I., Surosz, W. (Eds.), *Interdisciplinary Approaches for Sustainable Development*
 1081 *Goals: Economic Growth, Social Inclusion and Environmental Protection*. Springer
 1082 International Publishing, Cham, pp. 169–187. [https://doi.org/10.1007/978-3-319-](https://doi.org/10.1007/978-3-319-71788-3_12)
 1083 [71788-3_12](https://doi.org/10.1007/978-3-319-71788-3_12), 2018.

1084 Heidelberg Materials Kunda AS: Heidelberg Materials Kunda AS. Available at:
 1085 <https://www.kunda.heidelbergmaterials.ee/et/tegevusandmed> (Accessed: 24
 1086 February 2025), 2023

1087 Heinsalu, A., Veski, S.: Palaeoecological evidence of agricultural activity and human impact
 1088 on the environment at the ancient settlement centre of Keava, Estonia. *Estonian*
 1089 *Journal of Earth Sciences*, 59, pp.80-89. doi: 10.3176/earth.2010.1.06

1090 Hendon, D. and Charman, D.J., 1997. The preparation of testate amoebae (Protozoa:
 1091 Rhizopoda) samples from peat. *The Holocene*, 7(2), pp.199-205.
 1092 <https://doi.org/10.1177/095968369700700207>, 2010.

1093 Hendon, D., Charman, D.J. and Kent, M.: Palaeohydrological records derived from testate
 1094 amoebae analysis from peatlands in northern England: within-site variability,
 1095 between-site comparability and palaeoclimatic implications. *The Holocene*, 11(2),
 1096 pp.127-148. <https://doi.org/10.1191/095968301674575645>, 2001.

1097 Ivanov, Y.V., Kartashov, A.V., Ivanova, A.I., Ivanov, V.P., Marchenko, S.I., Nartov, D.I.,
 1098 Kuznetsov, V.V.: Long-term impact of cement plant emissions on the elemental
 1099 composition of both soils and pine stands and on the formation of Scots pine seeds.
 1100 *Environ. Pollut.* 243, 1383–1393. <https://doi.org/10.1016/j.envpol.2018.09.099>,
 1101 2018.

1102 Jalkanen, L., Mäkinen, A., Häsänen, E. and Juhanoja, J.: The effect of large anthropogenic
 1103 particulate emissions on atmospheric aerosols, deposition and bioindicators in the
 1104 eastern Gulf of Finland region. *Sci. of the Total Env.*, 262(1-2), pp.123-136.
 1105 [https://doi.org/10.1016/S0048-9697\(00\)00602-1](https://doi.org/10.1016/S0048-9697(00)00602-1), 2000.

1106 Joosten, H. and Clarke, D.: Wise use of mires and peatlands. *International mire conservation*
 1107 *group and international peat society*, 304, 2002.

1108 Joosten, H.: Wise use of mires: Background and principles, 239–250, 2003.

1109 Juggins, S. and Juggins, M.S.: Package ‘rioja’, 2019.

1110 Kaasik, A., Kont, R. and Lõhmus, A.: Modeling forest landscape futures: Full scale simulation
 1111 of realistic socioeconomic scenarios in Estonia. *Plos one*, 18(11), p.e0294650.
 1112 10.2495/GEO080171, 2023.

1113 Kaasik, M., Liblik, V. and Kaasik, H.: Long-term deposition patterns of airborne wastes in the
 1114 North-East of Estonia. *Oil Shale*, 16(4), pp.315-330. DOI: 10.3176/oil.1999.4.04,
 1115 1999.

1116 Kang, H., Kwon, M.J., Kim, S., Lee, S., Jones, T.G., Johncock, A.C., Haraguchi, A. and Freeman,
 1117 C.: Biologically driven DOC release from peatlands during recovery from
 1118 acidification. *Nature communications*, 9(1), p.3807.
 1119 <https://doi.org/10.1038/s41467-018-06259-1>, 2018.

1120 Karofeld, E.: The effects of alkaline fly ash precipitation on the Sphagnum mosses in
 1121 Niinsaare bog, NE Estonia. *Suoseura - Finnish Peatland Society Helsinki*, 1996.

1122 Kask, R., Ots, K., Mandre, M., Pikk, J.: Scots pine (*Pinus sylvestris* L.) wood properties in an
 1123 alkaline air pollution environment. *Trees* 22, 815–823.
 1124 <https://doi.org/10.1007/s00468-008-0242-7>, 2008.

1125 Katz, N.J., Katz, S.V. & Skobeyeva, E.I.: *Atlas Rastitel'nyh Oostatkov v Torfje (Atlas of Plant*
 1126 *Remains in Peats)*. Nedra, Moscow, 736pp. (in Russian), 1977.

1127 Katz, N.J., Katz, S.V. and Kipiani, M.G.: *Atlas and keys of fruits and seeds occurring in the*
 1128 *Quaternary deposits of the USSR*. Publishing House Nauka, Moscow, 1965.

1129 Keppler, F., Eiden, R., Niedan, V., Pracht, J. and Schöler, H.F.: Halocarbons produced by
 1130 natural oxidation processes during degradation of organic
 1131 matter. *Nature*, 403(6767), pp.298-301. <https://doi.org/10.1038/35002055>, 2000.

1132 Ketterer, M.E., Hafer, K.M. and Mietelski, J.W.: Resolving Chernobyl vs. global fallout
 1133 contributions in soils from Poland using Plutonium atom ratios measured by
 1134 inductively coupled plasma mass spectrometry. *J. of Env. Radioact.*, 73(2), pp.183-
 1135 201. <https://doi.org/10.1016/j.jenvrad.2003.09.001>, 2004.

1136 Klõšeiko, J., Kuznetsova, T., Tilk, M. and Mandre, M.: Short-term influence of Clinker dust
 1137 and wood ash on macronutrients and growth in Norway spruce (*Picea abies*) and
 1138 Scots pine (*Pinus sylvestris*) seedlings. *Commun in Soil Sci and Plant Analysis*, 45(16),
 1139 pp.2105-2117. <https://doi.org/10.1080/00103624.2014.929695>, 2014.

1140 Koff, T., Vandel, E., Marzecová, A., Avi, E. and Mikomägi, A.: Assessment of the effect of
 1141 anthropogenic pollution on the ecology of small shallow lakes using the

1142 palaeolimnological approach. *Estonian J. of Earth Sci.*, 65(4). Doi:
 1143 10.3176/earth.2016.19, 2016.

1144 Konings, W., Boyd, K. and Andersen, R.: Comparison of plant traits of sedges, shrubs and
 1145 Sphagnum mosses between sites undergoing forest-to-bog restoration and near-
 1146 natural open blanket bog: a pilot study. *Mires and Peat*, 23, pp.1-10, 2019.

1147 Korhola, A., Tolonen, K., Turunen, J. and Jungner, H.: Estimating long-term carbon
 1148 accumulation rates in boreal peatlands by radiocarbon dating. *Radiocarbon*, 37(2),
 1149 pp.575-584. 995;37(2) doi:10.1017/S0033822200031064, 1995.

1150 Kuhry, P.: Transgression of a raised bog across a coversand ridge originally covered with an
 1151 oak—lime forest: palaeoecological study of a middle holocene local vegetational
 1152 succession in the Amtsvien (Northwest Germany). *Rev. of Palaeobot. and*
 1153 *Palynol.*, 44(3-4), pp.303-353. [https://doi.org/10.1016/0034-6667\(85\)90023-5](https://doi.org/10.1016/0034-6667(85)90023-5),
 1154 1985.

1155 Kuhry, P.: The palaeoecology of a treed bog in western boreal Canada: a study based on
 1156 microfossils, macrofossils and physico-chemical properties. *Rev. of Palaeobot. and*
 1157 *Palynol.*, 96(1-2), pp.183-224. [https://doi.org/10.1016/S0034-6667\(96\)00018-8](https://doi.org/10.1016/S0034-6667(96)00018-8),
 1158 1997.

1159 Lamentowicz, M., Gałka, M., Marcisz, K., Słowiński, M., Kajukato-Drygalska, K., Dayras, M.D.
 1160 and Jassey, V.E. Unveiling tipping points in long-term ecological records from
 1161 Sphagnum-dominated peatlands. *Biol. Lett.*, 15(4), p.20190043.
 1162 <https://doi.org/10.1098/rsbl.2019.0043>, 2019.

1163 Lee-Taylor, J.M. and Holland, E.A.: Litter decomposition as a potential natural source of
 1164 methyl bromide. *J. of Geophys. Res.: Atmos.*, 105(D7), pp.8857-
 1165 8864. <https://doi.org/10.1029/1999JD901112>, 2000.

1166 Lehmann, N., Lantuit, H., Böttcher, M.E., Hartmann, J., Eulenburg, A., Thomas, H.: Alkalinity
 1167 generation from carbonate weathering in a silicate-dominated headwater
 1168 catchment at Iskorasfjellet, northern Norway. *Biogeosciences* 20, 3459–3479.
 1169 <https://doi.org/10.5194/bg-20-3459-2023>, 2023.

1170 Leifeld, J. and Menichetti, L.: The underappreciated potential of peatlands in global climate
 1171 change mitigation strategies. *Nature Commun.*, 9(1), p.1071.
 1172 <https://doi.org/10.1038/s41467-018-03406-6>, 2018.

1173 Levy, P., van Dijk, N., Gray, A., Sutton, M., Jones, M., Leeson, S., Dise, N., Leith, I. and
 1174 Sheppard, L., 2019. Response of a peat bog vegetation community to long-term

1175 experimental addition of nitrogen, *J. of Ecol.*, 107(3), pp.1167-1186.
 1176 <https://doi.org/10.1111/1365-2745.13107>

1177 Liblik, V., Rätsep, A. and Kundel, H.: Pollution sources and spreading of sulphur dioxide in
 1178 the North-Eastern Estonia, *Water, Air, and Soil Pollut.*, 85, pp.1903-1908.
 1179 <https://doi.org/10.1007/BF01186112>, 1995.

1180 Liiv, S. and Kaasik, M.: Trace metals in mosses in the Estonian oil shale processing region. *J.*
 1181 *of Atmos. Chemistry*, 49, pp.563-578. <https://doi.org/10.1007/s10874-004-1266-z>,
 1182 2004.

1183 Limpens, J., Berendse, F. and Klees, H.: N deposition affects N availability in interstitial water,
 1184 growth of Sphagnum and invasion of vascular plants in bog vegetation, *New*
 1185 *Phytologist*, 157(2), pp.339-347. [https://doi.org/10.1046/j.1469-](https://doi.org/10.1046/j.1469-8137.2003.00667.x)
 1186 [8137.2003.00667.x](https://doi.org/10.1046/j.1469-8137.2003.00667.x), 2003.

1187 Liu, B., Heinemeyer, A., Marchant, R. and Mills, R.T.: Exploring optimal sampling strategy of
 1188 testate amoebae as hydrological bioindicators in UK upland peatlands. *J. of Env.*
 1189 *Manag.*, 370, p.122959. <https://doi.org/10.1016/j.jenvman.2024.122959>, 2024.

1190 Loisel, J. and Bunsen, M.: Abrupt fen-bog transition across southern Patagonia: Timing,
 1191 causes, and impacts on carbon sequestration. *Front. in Ecol. and Evol.*, 8, p.273.
 1192 <https://doi.org/10.3389/fevo.2020.00273>, 2020.

1193 Longman, J., Veres, D. and Wennrich, V.: Utilisation of XRF core scanning on peat and other
 1194 highly organic sediments. *Quat. Int.*, 514, pp.85-96.
 1195 <https://doi.org/10.1016/j.quaint.2018.10.015>, 2019.

1196 Łuców, D., Küttim, M., Słowiński, M., Kołaczek, P., Karpińska-Kołaczek, M., Küttim, L., Salme,
 1197 M. and Lamentowicz, M.: Searching for an ecological baseline: Long-term ecology of
 1198 a post-extraction restored bog in Northern Estonia. *Quat. Int.*, 607, pp.65-78.
 1199 <https://doi.org/10.1016/j.quaint.2021.08.017>, 2022.

1200 MAA-JA RUUMIAMET: <https://xgis.maaamet.ee/xgis2/page/app/ajalooline>, last access: 10
 1201 August 2025.

1202 Madeja, J. and Latowski, D.: TOO OLD AMS RADIOCARBON DATES OBTAINED FROM MOSS
 1203 REMAINS FROM LAKE KWIECKO BOTTOM SEDIMENTS (N
 1204 POLAND). *Geochronometria: Journal on Methods & Applications of Absolute*
 1205 *Chronology*, 32. 10.2478/v10003-008-0029-2, 2008.

1206 Maimer, N., Horton, D.G. and Vitt, D.H.: Element concentrations in mosses and surface
 1207 waters of western Canadian mires relative to precipitation chemistry and

1208 hydrology. *Ecography*, 15(1), pp.114-128. <https://doi.org/10.1111/j.1600->
1209 [0587.1992.tb00015.x](https://doi.org/10.1111/j.1600-0587.1992.tb00015.x), 1992.

1210 Mandre, M. and Korsjukov, R.: The quality of stemwood of *Pinus sylvestris* in an alkalised
1211 environment. *Water, air, and soil pollution*, 182, pp.163-172.
1212 <https://doi.org/10.1007/s11270-006-9329-1>, 2007.

1213 Mandre, M. and Ots, K.: Growth and biomass partitioning of 6-year-old spruces under
1214 alkaline dust impact. *Water, Air, and Soil Poll.*, 114, pp.13-25.
1215 <https://doi.org/10.1023/A:1005048921852>, 1999.

1216 Marcisz, K., Jassey, V.E., Kosakyan, A., Krashevskaya, V., Lahr, D.J., Lara, E., Lamentowicz, Ł.,
1217 Lamentowicz, M., Macumber, A., Mazei, Y. and Mitchell, E.A.: Testate amoeba
1218 functional traits and their use in paleoecology. *Front. in Ecol. and Evol.*, 8, p.575966.
1219 <https://doi.org/10.3389/fevo.2020.575966>, 2020.

1220 Mauquoy, D and van Geel, B.: Mire and peat macros. *Encyclopedia Quaternary Science*, 3,
1221 pp.2315-2336, 2007.

1222 Mauquoy, D., Hughes, P.D.M. and Van Geel, B.: A protocol for plant macrofossil analysis of
1223 peat deposits. *Mires and Peat*, 7(6), pp.1-5, 2010.

1224 Mazei, Y and Tsyganov, A.: *Freshwater testate amoebae* [in Russian], KMK Scientific
1225 pressSBN: 5-87317-336-2, 2006.

1226 Mitchell, E.A., Charman, D.J. and Warner, B.G.: Testate amoebae analysis in ecological and
1227 paleoecological studies of wetlands: past, present and future. *Biodivers. and*
1228 *Conserv.* 17, pp.2115-2137. <https://doi.org/10.1007/s10531-007-9221-3>, 2008.

1229 Moore, P.D., Webb, J.A. and Collison, M.E.: *Pollen analysis* (pp. viii+-216), 1991.

1230 Mróz, T., Łokas, E., Kocurek, J. and Gąsiorek, M.: Atmospheric fallout radionuclides in
1231 peatland from Southern Poland. *J. of Envi. Radioact.*, 175, pp.25-33
1232 <https://doi.org/10.1016/j.jenvrad.2017.04.012>, 2017.

1233 Munson, S.M., Reed, S.C., Peñuelas, J., McDowell, N.G. and Sala, O.E.: Ecosystem thresholds,
1234 tipping points, and critical transitions. *New Phytologist*, 218(4), pp.1315-1317.
1235 <https://www.jstor.org/stable/90021529>, 2018.

1236 Niinemets, E. and Saarse, L.: Holocene vegetation and land-use dynamics of south-eastern
1237 Estonia. *Quat Int.*, 207(1-2), pp.104-116.
1238 <https://doi.org/10.1016/j.quaint.2008.11.015>, 2009.

1239 Oksanen, J., Blanchet, F.G., Kindt, R., Legendre, P., Minchin, P.R., O'hara, R.B., Simpson, G.L.,
 1240 Solymos, P., Stevens, M.H.H., Wagner, H. and Oksanen, M.J.: Package
 1241 'vegan'. *Community ecology package, version, 2(9)*, pp.1-295, 2019.
 1242 Orme, L.C., Davies, S.J. and Duller, G.A.T.: Reconstructed centennial variability of late
 1243 Holocene storminess from Cors Fochno, Wales, UK. *J. of Quat. Sci.*, 30(5), pp.478-
 1244 488. <https://doi.org/10.1002/jqs.2792>, 2015.
 1245 Órpez, R., Martínez, M.E., Hodaifa, G., El Yousfi, F., Jbari, N. and Sánchez, S.: Growth of the
 1246 microalga *Botryococcus braunii* in secondarily treated sewage. *Desalination*, 246(1-
 1247 3), pp.625-630. <https://doi.org/10.1016/j.desal.2008.07.016>, 2009.
 1248 Osborne, C., Gilbert-Parkes, S., Spiers, G., Lamit, L.J., Lilleskov, E.A., Basiliko, N., Watmough,
 1249 S., Andersen, R., Artz, R.E., Benscoter, B.W., Bragazza, L., Bräuer, S.L., Carson, M.A.,
 1250 Chen, X., Chimner, R.A., Clarkson, B.R., Enriquez, A.S., Grover, S.P., Harris, L.I.,
 1251 Hazard, C., Hribljan, J., Juutinen, S., Kane, E.S., Knorr, K.-H., Kolka, R., Laine, A.M.,
 1252 Larmola, T., McCalley, C.K., McLaughlin, J., Moore, T.R., Mykytczuk, N., Normand,
 1253 A.E., Olefeldt, D., Rich, V., Roulet, N., Rupp, D.L., Rutherford, J., Schadt, C.W.,
 1254 Sonnentag, O., Tedersoo, L., Trettin, C.C., Tuittila, E.-S., Turetsky, M., Urbanová, Z.,
 1255 Varner, R.K., Waldrop, M.P., Wang, M., Wang, Z., Wiedermann, M.M., Williams, S.T.,
 1256 Yavitt, J.B., Yu, Z.-G.: Global Patterns of Metal and Other Element Enrichment in Bog
 1257 and Fen Peatlands. *Arch. Environ. Contam. Toxicol.* 86, 125–139.
 1258 <https://doi.org/10.1007/s00244-024-01051-3>, 2024.
 1259 Osterkamp, T.E., Viereck, L., Shur, Y., Jorgenson, M.T., Racine, C., Doyle, A. and Boone, R.D.:
 1260 Observations of thermokarst and its impact on boreal forests in Alaska, USA. *Arctic,*
 1261 *Antarctic, and Alpine Res.*, 32(3), pp.303-315.
 1262 <https://doi.org/10.1080/15230430.2000.12003368>, 2000.
 1263 Ots, K. and Mandre, M.: Monitoring of heavy metals uptake and allocation in *Pinus sylvestris*
 1264 organs in alkalised soil. *Env. Monit. and Assess.*, 184, pp.4105-4117.
 1265 <https://doi.org/10.1007/s10661-011-2247-8>, 2012.
 1266 Ots, K. and Reisner, V.: Scots pine (*Pinus sylvestris* L.) and its habitat in Muraka bog under
 1267 the influence of wastes from the Narva power plants (North-East Estonia). *Proc.*
 1268 *Estonian Acad. Sci. Biol. Ecol*, 55(2), pp.137-148, 2006.
 1269 Ots, K., Tilk, M. and Aguraijuja, K.: The effect of oil shale ash and mixtures of wood ash and
 1270 oil shale ash on the above-and belowground biomass formation of Silver birch and

1271 Scots pine seedlings on a cutaway peatland. *Eco. Eng.* 108, pp.296-306.
 1272 <https://doi.org/10.1016/j.ecoleng.2017.09.002>, 2017.

1273 Paal, J., Vellak, K., Liira, J. and Karofeld, E.: Bog recovery in northeastern Estonia after the
 1274 reduction of atmospheric pollutant input. *Restor. Eco.*, 18, pp.387-400.
 1275 <https://doi.org/10.1111/j.1526-100X.2009.00608.x>, 2010.

1276 Paavilainen, E. and Päivänen, J.: Utilization of peatlands. In *Peatland Forestry: Ecology and*
 1277 *Principles* (pp. 15-29). Berlin, Heidelberg: Springer Berlin Heidelberg, 1995.

1278 Pärtel, M., Helm, A., Ingerpuu, N., Reier, Ü. and Tuvi, E.L., 2004. Conservation of Northern
 1279 European plant diversity: the correspondence with soil pH. *Biol. Conserv.*, 120(4),
 1280 pp.525-531. <https://doi.org/10.1016/j.biocon.2004.03.025>

1281 Pärtma, M. 2023. *The recovery of bogs in NE-Estonia bogs from alkaline air pollution. Case*
 1282 *study of Varudi, Sämi-Kurustiku and Uljaste bogs*. Master Thesis, Tallinn University.
 1283 97 pp. (in Estonian with English summary).

1284 Patterson III, W.A., Edwards, K.J., Maguire, D.J.: Microscopic charcoal as a fossil indicator of
 1285 fire. *Quat. Sci. Rev.* 6, 3–23. [https://doi.org/10.1016/0277-3791\(87\)90012-6](https://doi.org/10.1016/0277-3791(87)90012-6), 1987.

1286 Payne, R.J. and Mitchell, E.A.: How many is enough? Determining optimal count totals for
 1287 ecological and palaeoecological studies of testate amoebae. *J. of*
 1288 *Paleolimnology*, 42(4), pp.483-495. <https://doi.org/10.1007/s10933-008-9299-y>,
 1289 2009.

1290 Pensa, M., Jalkanen, R. and Liblik, V.: Variation in Scots pine needle longevity and nutrient
 1291 conservation in different habitats and latitudes. *Canadian J. of For. Res.* 37(9),
 1292 pp.1599-1604. <https://doi.org/10.1139/X07-012>, 2007.

1293 Pensa, M., Liblik, V. and Jalkanen, R. Temporal changes in the state of a pine stand in a bog
 1294 affected by air pollution in northeast Estonia. *Water, Air, and Soil Poll.*, 159, pp.87-
 1295 99. <https://doi.org/10.1023/B:WATE.0000049191.36830.a7>, 2004.

1296 Piotrowska, N., Blaauw, M., Mauquoy, D. and Chambers, F.M.: Constructing deposition
 1297 chronologies for peat deposits using radiocarbon dating. *Mires and Peat*, 7(10), pp.1-
 1298 14, 2011.

1299 Poska, A. and Saarse, L.: Vegetation development and introduction of agriculture to
 1300 Saaremaa Island, Estonia: the human response to shore displacement. *The*
 1301 *Holocene*, 12(5), pp.555-568. <https://doi.org/10.1191/0959683602hl567rp>, 2002.

1302 Poska, A.: *Human impact on vegetation of coastal Estonia during the Stone Age*. Doctoral
 1303 Dissertation, Uppsala Universitet, Sweden, 2003.

- R Core Team: R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria. <<https://www.R-project.org/>>., 2023.
- Raukas, A. (ed.), Environmental Impact Assessment for the Area of Influence of Reconstructed Kunda Cement Factory. Present Situation and Prediction of Potential Changes. Tallinn (manuscript), 1993.
- Rawlins, A. and Morris, J.: Social and economic aspects of peatland management in Northern Europe, with particular reference to the English case. *Geoderma*, 154(3-4), pp.242-251. <https://doi.org/10.1016/j.geoderma.2009.02.022>, 2010.
- Reimer, P.J., Austin, W.E., Bard, E., Bayliss, A., Blackwell, P.G., Ramsey, C.B., Butzin, M., Cheng, H., Edwards, R.L., Friedrich, M. and Grootes, P.M.: The IntCal20 Northern Hemisphere radiocarbon age calibration curve (0–55 cal kBP). *Radiocarbon*, 62(4), pp.725-757. <https://doi.org/10.1017/RDC.2020.41>, 2020.
- Roderfeld, H.: *Raised bog regeneration after peat harvesting in north-west Germany*, 1993.
- Rodwell, J.S. ed., *British plant communities: Volume 2, mires and heaths (Vol. 2)*. Cambridge University Press, 1998.
- Roulet, N.T., Lafleur, P.M., Richard, P.J., Moore, T.R., Humphreys, E.R. and Bubier, J.I.L.L.: Contemporary carbon balance and late Holocene carbon accumulation in a northern peatland. *Glob. Change Biol.*, 13(2), pp.397-411. <https://doi.org/10.1111/j.1365-2486.2006.01292.x>, 2007.
- Schindler, D.W.: Widespread effects of climatic warming on freshwater ecosystems in North America. *Hydrol. Process.*, 11(8), pp.1043-1067. [https://doi.org/10.1002/\(SICI\)1099-1085\(19970630\)11:8<1043::AID-HYP517>3.0.CO;2-5](https://doi.org/10.1002/(SICI)1099-1085(19970630)11:8<1043::AID-HYP517>3.0.CO;2-5), 1997.
- Seppä, H. and Bennett, K.D.: Quaternary pollen analysis: recent progress in palaeoecology and palaeoclimatology. *Prog. in Phys. Geog.*, 27(4), pp.548-579. <https://doi.org/10.1191/0309133303pp394oa>, 2003.
- Sheppard, L.J., Leith, I.D., Mizunuma, T., Neil Cape, J., Crossley, A., Leeson, S., Sutton, M.A., van Dijk, N. and Fowler, D.: Dry deposition of ammonia gas drives species change faster than wet deposition of ammonium ions: Evidence from a long-term field manipulation. *Glob. Change Biol.*, 17(12), pp.3589-3607. <https://doi.org/10.1111/j.1365-2486.2011.02478.x>, 2011.
- Shotyk, W.: Atmospheric deposition and mass balance of major and trace elements in two oceanic peat bog profiles, northern Scotland and the Shetland Islands. *Chem. Geol.* 138(1-2), pp.55-72. [https://doi.org/10.1016/S0009-2541\(96\)00172-6](https://doi.org/10.1016/S0009-2541(96)00172-6), 1997.

1337 Sibul, I., Plado, J. and Jõelet, A.: Ground-penetrating radar and electrical resistivity
 1338 tomography for mapping bedrock topography and fracture zones: a case study in
 1339 Viru-Nigula, NE Estonia. *Estonian J. of Earth Sci.*, 66(3), p.142.
 1340 <https://doi.org/10.3176/earth.2017.11>, 2017.

1341 Siemensma, F.: *Microworld – world of amoeboid organisms*. <https://arcella.nl> last accessed
 1342 20/10/2023

1343 Silva-Sánchez, N., Martínez Cortizas, A. and López-Merino, L.: Linking forest cover, soil
 1344 erosion and mire hydrology to late-Holocene human activity and climate in NW
 1345 Spain. *The Holocene*, 24(6), pp.714-725.
 1346 <https://doi.org/10.1177/0959683614526934>, 2014.

1347 Simpson, G.L.: Modelling palaeoecological time series using generalised additive
 1348 models. *Front. Ecol. Evol*, 6, p.149. <https://doi.org/10.3389/fevo.2018.00149>, 2018.

1349 Simpson, G.L., Oksanen, J. and Simpson, M.G.L.: Package ‘analogue’. *Analogue and*
 1350 *Weighted Averaging Methods for Palaeoecology*, 2016.

1351 Stelling, J.M., Slesak, R.A., Windmuller-Campione, M.A. and Grinde, A.: Effects of stand age,
 1352 tree species, and climate on water table fluctuations and estimated
 1353 evapotranspiration in managed peatland forests. *J. of Env. Management*, 339,
 1354 p.117783. <https://doi.org/10.1016/j.jenvman.2023.117783>, 2023.

1355 Stockmarr, J.: Tables with spores used in absolute pollen analysis. *Pollen et spores*, 13,
 1356 pp.615-621, 1971.

1357 Sutton, M.A., van Dijk, N., Levy, P.E., Jones, M.R., Leith, I.D., Sheppard, L.J., Leeson, S., Sim
 1358 Tang, Y., Stephens, A., Braban, C.F., Dragosits, U., Howard, C.M., Vieno, M., Fowler,
 1359 D., Corbett, P., Naikoo, M.I., Munzi, S., Ellis, C.J., Chatterjee, S., Steadman, C.E.,
 1360 Móring, A., Wolseley, P.A.: Alkaline air: changing perspectives on nitrogen and air
 1361 pollution in an ammonia-rich world. *Philos. Trans. R. Soc. Math. Phys. Eng. Sci.* 378,
 1362 20190315. <https://doi.org/10.1098/rsta.2019.0315>, 2020.

1363 Swindles, G.T.: Dating recent peat profiles using spheroidal carbonaceous particles (SCPs).
 1364 *Mires and peat*, 7, 2020.

1365 Szumigalski, A.R. and Bayley, S.E.: Decomposition along a bog to rich fen gradient in central
 1366 Alberta, Canada. *Canadian J. of Bot.*, 74(4), pp.573-581.
 1367 <https://doi.org/10.1139/b96-073>, 1996.

1368 Tanneberger, F., Moen, A., Barthelmes, A., Lewis, E., Miles, L., Sirin, A., Tegetmeyer, C. and
1369 Joosten, H.: Mires in Europe—Regional diversity, condition and protection. *Diversity*,
1370 13(8), p.381. <https://doi.org/10.3390/d13080381>, 2021.

1371 Taranu, Z.E., Carpenter, S.R., Frossard, V., Jenny, J.P., Thomas, Z., Vermaire, J.C. and Perga,
1372 M.E.: Can we detect ecosystem critical transitions and signals of changing resilience
1373 from paleo-ecological records?. *Ecosphere*, 9(10),
1374 p.e02438. <https://doi.org/10.1002/ecs2.2438>, 2018.

1375 Thormann, M.N., Currah, R.S. and Bayley, S.E.: Succession of microfungal assemblages in
1376 decomposing peatland plants. *Plant and Soil*, 250, pp.323-333.
1377 <https://doi.org/10.1023/A:1022845604385>, 2003.

1378 Tobolski, K. Przewodnik do oznaczania torfów i osadów jeziornych [Guide to the
1379 determination of peat and lake sediments, in Polish]. Vademecum Geobotanicum. T.
1380 2. Warszawa. Wydaw. Nauk. PWN, p.508, 2000.

1381 Trumm, U., Raju, T. and Kangur, P. Kunda tsement - 140: tsemendi tootmise ajalugu Kundas
1382 1870-2010. In *Nomine OÜ, Tallinn*. (in Estonian), 2010.

1383 Turetsky, M.R. and St. Louis, V.L. Disturbance in boreal peatlands. In *Boreal peatland*
1384 *ecosystems* (pp. 359-379). Berlin, Heidelberg: Springer Berlin Heidelberg, 2006.

1385 Turner, T.E., Swindles, G.T. and Roucoux, K.H. Late Holocene ecohydrological and carbon
1386 dynamics of a UK raised bog: impact of human activity and climate change. *Quat. Sci.*
1387 *Rev.*, 84, pp.65-85. <https://doi.org/10.1016/j.quascirev.2013.10.030>, 2014.

1388 Uno, K.T., Quade, J., Fisher, D.C., Wittemyer, G., Douglas-Hamilton, I., Andanje, S., Omondi,
1389 P., Litoroh, M. and Cerling, T.E. Bomb-curve radiocarbon measurement of recent
1390 biologic tissues and applications to wildlife forensics and stable isotope (paleo)
1391 ecology. *Proc. of the Nat. Acad. of Sci.*, 110(29), pp.11736-11741.
1392 <https://doi.org/10.1073/pnas.1302226110>, 2013.

1393 Vaasma, T., Kaasik, M., Loosaar, J., Kiisk, M. and Tkaczyk, A.H.: Long-term modelling of fly
1394 ash and radionuclide emissions as well as deposition fluxes due to the operation of
1395 large oil shale-fired power plants. *J. of Env. Radioact*, 178, pp.232-244.
1396 <https://doi.org/10.1016/j.jenvrad.2017.08.017>, 2017.

1397 Vaasma, T., Kiisk, M., Meriste, T. and Tkaczyk, A.H.: The enrichment of natural radionuclides
1398 in oil shale-fired power plants in Estonia—The impact of new circulating fluidized bed
1399 technology. *J. of Env. Radioact*, 129, pp.133-139.
1400 <https://doi.org/10.1016/j.jenvrad.2014.01.002>, 2014.

1401 Van den Brink, P.J. and Braak, C.J.T.: Principal response curves: Analysis of time-dependent
 1402 multivariate responses of biological community to stress. *Env. Toxicol. and Chemistry:*
 1403 *An Int. J.* 18(2), pp.138-148. <https://doi.org/10.1002/etc.5620180207>, 1999.
 1404 van Geel, B. and Aptroot, A.: Fossil ascomycetes in Quaternary deposits. *Nova*
 1405 *Hedwigia*, 82(3), pp.313-330. <https://hdl.handle.net/11245/1.255775>, 2006.
 1406 Van Geel, B.: A palaeoecological study of Holocene peat bog sections in Germany and the
 1407 Netherlands, based on the analysis of pollen, spores and macro-and microscopic
 1408 remains of fungi, algae, cormophytes and animals. *Rev of palaeobot and*
 1409 *palynol.*, 25(1), pp.1-120. [https://doi.org/10.1016/0034-6667\(78\)90040-4](https://doi.org/10.1016/0034-6667(78)90040-4), 1978.
 1410 Vandel, E. and Vaasma, T.: Development of small lakes in Estonia: paleolimnological
 1411 studies. *Dynamiques environnementales. J. Int. de géosciences et de*
 1412 *l'environnement*, (42), pp.368-375. <https://doi.org/10.4000/dynenviron.2533>, 2018.
 1413 Varvas, M. and Punning, J.M.: Use of the 210Pb method in studies of the development and
 1414 human-impact history of some Estonian lakes. *The Holocene*, 3(1), pp.34-44.
 1415 <https://doi.org/10.1177/095968369300300104>, 1993.
 1416 Vellak, K., Liira, J., Karofeld, E., Galanina, O., Noskova, M. and Paal, J.: Drastic turnover of
 1417 bryophyte vegetation on bog microforms initiated by air pollution in Northeastern
 1418 Estonia and bordering Russia. *Wetlands*, 34, pp.1097-1108.
 1419 <https://doi.org/10.1007/s13157-014-0569-3>, 2014.
 1420 Veski, S.: *Vegetation history, human impact and palaeogeography of West Estonia: Pollen*
 1421 *analytical studies of lake and bog sediments* (Doctoral dissertation, Societas
 1422 Upsaliensis pro geologia quaternaria [Kvartärgeologiska fören.]), 1998.
 1423 Veski, S., Koppel, K. and Poska, A.: Integrated palaeoecological and historical data in the
 1424 service of fine-resolution land use and ecological change assessment during the last
 1425 1000 years in Rõuge, southern Estonia. *J. of Biogeog* 32(8), pp.1473-1488.
 1426 <https://doi.org/10.1111/j.1365-2699.2005.01290.x>, 2005.
 1427 Wagner, K.I., Gallagher, S.K., Hayes, M., Lawrence, B.A. and Zedler, J.B.: Wetland restoration
 1428 in the new millennium: do research efforts match opportunities?. *Restor. Ecol.* 16(3),
 1429 pp.367-372. <https://doi.org/10.1111/j.1526-100X.2008.00433.x>, 2008.
 1430 Wardenaar, E.C.P.: A new hand tool for cutting peat profiles. *Canadian J. of Botany*, 65(8),
 1431 pp.1772-1773. <https://doi.org/10.1139/b87-243>, 1987.
 1432 Warner, B.G. and Asada, T.: Biological diversity of peatlands in Canada. *Aquatic Sci.*, 68,
 1433 pp.240-253. <https://doi.org/10.1007/s00027-006-0853-2>, 2006.

- Waters, C.N., Turner, S.D., Zalasiewicz, J., Head, M.J.: Candidate sites and other reference sections for the Global boundary Stratotype Section and Point of the Anthropocene series. *Anthropocene Rev.* <https://doi.org/10.1177/20530196221136422>, 2023.
- Word, C.S., McLaughlin, D.L., Strahm, B.D., Stewart, R.D., Varner, J.M., Wurster, F.C., Amestoy, T.J. and Link, N.T.: Peatland drainage alters soil structure and water retention properties: Implications for ecosystem function and management. *Hydrological Processes*, 36(3), p.e14533. <https://doi.org/10.1002/hyp.14533> , 2022.
- Yang, Q., Liu, Z. and Bai, E.: Comparison of carbon and nitrogen accumulation rate between bog and fen phases in a pristine peatland with the fen-bog transition. *Glob. Change Biol.* 29(22), pp.6350-6366. <https://doi.org/10.1111/gcb.16915>, 2023.
- Young, D.M., Baird, A.J., Charman, D.J., Evans, C.D., Gallego-Sala, A.V., Gill, P.J., Hughes, P.D., Morris, P.J. and Swindles, G.T.: Misinterpreting carbon accumulation rates in records from near-surface peat. *Sci. Rep.* 9(1), p.17939. <https://doi.org/10.1038/s41598-019-53879-8>, 2019.
- Young, D.M., Baird, A.J., Gallego-Sala, A.V. and Loisel, J.: A cautionary tale about using the apparent carbon accumulation rate (aCAR) obtained from peat cores. *Sci. Rep.* 11(1), p.9547. <https://doi.org/10.1038/s41598-021-88766-8> , 2021.
- Yu, Z., Loisel, J., Brosseau, D.P., Beilman, D.W. and Hunt, S.J.: Global peatland dynamics since the Last Glacial Maximum. *Geophys. Res. Lett.*, 37(13). <https://doi.org/10.1029/2010GL043584>, 2010.
- Yu, Z.C., Northern peatland carbon stocks and dynamics: a review. *Biogeosciences*, 9(10), pp.4071-4085. <https://doi.org/10.5194/bg-9-4071-2012>, 2012.
- Zvereva, E.L., Toivonen, E. and Kozlov, M.V.: Changes in species richness of vascular plants under the impact of air pollution: a global perspective. *Glob. Ecol. and Biogeog.* 17(3), pp.305-319. <https://doi.org/10.1111/j.1466-8238.2007.00366.x>, 2008a.
- Zvereva, V.E., Zvereva, E.L. and Kozlov, M.V.: Slow growth of *Empetrum nigrum* in industrial barrens: Combined effect of pollution and age of extant plants. *Env. Poll.* 156(2), pp.454-460. <https://doi.org/10.1016/j.envpol.2008.01.025>, 2008b.