

Controls over debris flow initiation in glacio-volcanic environments in the Southern Andes.

Ivo Fustos-Toribio¹, Daniel Basualto^{3*}, Ardy Gatica¹, Alvaro Bravo-Alarcón^{2,1}, José-Luis Palma⁴, Gabriel Fuentealba^{5,2}, Sergio A. Sepúlveda^{6,7}

¹Departamento de Ingeniería en Obras Civiles, Facultad de Ingeniería y Ciencias, Universidad de La Frontera. Francisco Salazar #01145, Temuco, Chile.

² Programa de Magíster en Ciencias de La Ingeniería, Universidad de La Frontera, Temuco, Chile.

³Departamento de Ingeniería Eléctrica, Facultad de Ingeniería y Ciencias, Universidad de La Frontera. Francisco Salazar #01145, Temuco, Chile.

⁴Departamento Ciencias de la Tierra, Facultad de Ciencias Químicas, Universidad de Concepción, Víctor Lamas 1290, Concepción, Chile.

⁵Ministerio del Interior, Temuco, Chile.

⁶ Departamento de Geología, Facultad de Ciencias Físicas y Matemáticas, Universidad de Chile, Santiago, Chile

⁷ Department of Earth Sciences, Faculty of Science, Simon Fraser University, Burnaby, Canada

Correspondence to: daniel.basualto@ufrontera.cl

Abstract. The Southern Andes is an active zone of mass wasting processes with unknown constraints for public policies. Several conditioning factors could have an impact on the generation of debris flows, being controlled by water accumulation. This study investigates the generation of the Ñisoleufu debris flow, an active area of debris flow generation in Southern Andes, reviewing the interplay between geomorphological, geotechnical and hydrometeorological controls in debris flow dynamics, focusing on the effects of soil properties, slope characteristics and precipitation events. Our results highlight significant changes in soil moisture content on critical days associated with debris flow events. We revealed that the combination of areas with high water accumulation capacity from local runoff and slopes that capture precipitation effectively were crucial in the generation of debris flows. Areas with granular volcanic soils acted as storage mediums for water, which, coupled with decreased shear strength, facilitated debris flow initiation. The thin and fine-grained layers of glacial deposits located beneath the volcanic soil, characterized by low hydraulic conductivity, created localized accumulation zones that reinforced the storage capacity of adjacent areas, particularly in pyroclastic volcanic deposits in the release zone. The hydraulic properties of the volcanic deposits suggest that water storage capacity and high hydraulic conductivity play a critical role in rainfall-induced debris flow initiation. Additionally, we observed that the debris flow of the Ñisoleufu event has evidence of reworked lapilli-sized particles (>5 mm), being consistent with the surface and shallow water movement that reduces the slope stability within the area. Analysis of ERA5-land dataset showed abrupt changes in soil moisture content at various depths and time periods, correlating with intense or prolonged rainfall events. These results underscore the role of geomorphological features in modulating soil

35 moisture and thereby affecting the stability and movement of debris flows. Our results provide a comprehensive understanding
36 of how geomorphology interacts with hydrological factors to influence debris flow behaviour in volcanic areas of the Southern
37 Andes for the first time. Overall, the research highlights the critical role of geomorphological and hydrological factors in debris
38 flow generation and dynamics. It emphasizes the need for incorporating detailed soil and slope characteristics into models for
39 predicting debris flow risks. By understanding the combined effects of water accumulation, soil properties, and slope dynamics,
40 this study contributes valuable insights into managing and mitigating debris flow hazards in vulnerable regions. These findings
41 enhance the predictive capacity for rainfall-induced debris flows and provide practical criteria for hazard assessment in post-
42 glacial volcanic terrains.
43

44 1 Introduction

45 Episodes of extreme rainfall have increased due to climate change, resulting in a greater frequency of debris flows (Jakob and
46 Lambert 2009; Lee 2017; Fustos et al., 2017; Dey and Sengupta 2018) and mainly related to fast changes of soil water content
47 (Fustos et al., 2021). Currently, mass wasting processes produce widespread damages, representing a significant threat to
48 human life (Sepúlveda and Petley 2015; Vega and Hidalgo 2016). Consequently, the need to forecast (Fustos et al., 2020a)
49 and mitigate (Fustos et al., 2021b) the effects of these events has become a high priority for governments facing increasing
50 episodes of rainfall-induced landslides linked with climate change. An accurate assessment of potential debris flows to regional
51 scale needs precise understanding of their triggering and controlling conditions. Therefore, we assessed the main conditioning
52 and triggering conditions of debris flows in Southern Andes in order to understand their evolution from stable slope to mass
53 wasting event.

54 Worldwide, the increasing frequency and intensity of such extreme precipitation events exacerbated by climate change to
55 regional scale (Stoffel et al., 2013; Pavlova et al., 2018) introduce severe threats to human life and property. Changing
56 precipitation patterns related to extreme precipitation, lead to conditions conducive for debris flows in wide areas in North
57 America (Bovis et al., 1999), Asia (Chang et al., 2017), Europe (Malet et al., 2005; Stoffel et al., 2013; Pavlova et al., 2018)
58 and South America (Sepúlveda et al., 2013; Sepúlveda et al., 2014; Fustos et al., 2022). Stand out heavy tropical storms, such
59 as Taiwan, increasing debris flow incidents (Chang et al., 2025) and compromising the safety of urban areas located near
60 mountainous terrains (Chen et al., 2015; Kang et al., 2017). The Wenchuan Earthquake in China exemplifies how seismic
61 activity can trigger extensive debris flows, resulting in not only immediate destruction but also long-lasting hazards due to the
62 formation of landslide-dammed lakes that threaten downstream communities (Cui et al., 2009; Wang and Yan 2015). Accurate
63 forecasting of debris flow events is vital for disaster preparedness and mitigation. Understanding the triggering conditions—
64 including rainfall intensity, groundwater levels, and geological features—is essential for developing conceptual models that
65 represent the regional conditions that could control a debris flow initiation. Debris flows are influenced by soil hydraulic
66 characteristics and the intensity/duration of rainfall events (Singh and Kumar, 2020), in which rainfall intensities serve as
67 crucial predictors in mountainous regions (Chang et al., 2017; Fustos-Toribio et al., 2022). Moreover, coarse-grained volcanic
68 soils exhibit transient increases in pore pressure during intense rainfall events (Huang et al., 2012). Conversely, fine-grained
69 soils with low infiltration rates do not experience significant changes in the pore pressure, generating failures due to decreased
70 soil shear strength (Dahal et al., 2008; Dahal et al., 2011). Hence, understanding soil composition and granulometric features
71 is pivotal in assessing debris flow susceptibility worldwide. Debris flows are mainly controlled by the geomorphological
72 features and the specific geological evolution of each region of the planet, highlighting the need for localized and context-
73 specific approaches for their study and management. One of the next frontier corresponds to constrain the debris flow
74 generation in glacial environments under changing precipitation events related to climate change.

75 Nowadays, understanding the impact of debris flows in glacial environments becomes critical in the Chilean southern Andes,
76 particularly due to the most part of the inhabitants live there. Changes of precipitation patterns related to climate change,

particularly fast and intense rainfall events, could amplify the frequency and magnitude of debris flows (Fustos et al., 2022). An increase of extreme hydrometeorological events affecting slopes in glacial settings is observed, whose mechanical properties and geomorphology have evolved since the Last Glacial Maximum (Fustos-Toribio et al., 2021b; Somos-Valenzuela et al., 2020; Ochoa-Cornejo et al. 2024). Considerable uncertainty remains about how the interaction between volcanic-derived soils over glacial landforms will respond to extreme hydrometeorological events. One of the current models for initiation of debris flows is related to slow deforming surfaces in hillslopes, mainly due to gravity and surface erosion during high precipitation events (Xie et al. 2020; Yi et al. 2021). Slow surface deformation could lead to extensional failures that could expand and deepen, generating landslides and evolving into debris flows, especially under water-saturated conditions or heavy rainfall (Gregoretti, 2000; Fustos et al., 2017; Wang et al., 2024). The capacity to oversee these extensional failures in remote areas close to roads is an open question yet, mainly in Southern Andes.

Historical debris flow events in the Southern Andes remain subject to uncertain conditioning factors. The transformation of landslides into debris flows typically occurs when sliding material incorporates water, significantly increasing its fluidity—as observed in the Villa Santa Lucía event in Chilean Patagonia (Somos-Valenzuela et al., 2020). The occurrence of debris flows in volcanic settings is of considerable scientific interest (Cheung & Giardino, 2023; Sepúlveda et al., 2008), largely due to the complex nature of volcanic soils and their marked textural variability (Thompson et al., 2023), which strongly influence water retention and infiltration dynamics. Recurrent debris flows in the Osorno volcano exemplify the role of intense rainfall events and the mobilization of autobrecciated lava blocks in initiating such processes (Fustos et al., 2022). Although numerical modeling has estimated total flow volumes (ranging from 4.7×10^5 to $5.5 \times 10^5 \text{ m}^3$) highlighting the high sensitivity of debris flow generation to the initial water content, a comprehensive conceptual model remains lacking. Over the past four decades, the Southern Andes has experienced significant volcanic activity (Galetto et al., 2023), resulting in widespread tephra deposition that has contributed to increased frequencies of debris flows (Korup et al., 2019). Nevertheless, critical knowledge gaps persist regarding the spatial and temporal variability of the textural and hydraulic properties of volcanic soils, particularly under extreme hydrometeorological conditions. Addressing these gaps is crucial to better constrain the hydraulic and geomechanical conditions that lead to debris flow initiation (Schmidt et al., 2001; Kuriakose et al., 2009). Such understanding is fundamental for effective territorial planning, risk mitigation, and the development of robust early warning systems.


In this paper, we analyse the precursory surface deformation related to tensile cracks leading to the Ñisoleufu debris flow event (May 31st, 2021). We utilised a multi-temporal InSAR approach with Sentinel-1 C-band SAR data, enabling us to create time series plots of deformation before the landslide and compare them with available daily precipitation records from nearby weather stations and satellite measurements. By combining remote sensing data, weather records, and soil laboratory analyses, we aim to provide valuable insights into the factors leading to such events and, consequently, improve hazard assessment and management in zones with soils derived from explosive volcanic events.

109 **2 Study area**

110 Debris flows are the most common manifestation of mass wasting triggered by precipitation in the Southern Andes due to the
111 soil heterogeneity and geological features, providing a unique opportunity to study the relationship between extreme rainfall
112 and debris flows (Figure 1 A-D). Recent extreme precipitation events have produced mass wasting hazard, especially in steep
113 zones near alluvial plains where human settlements are often established (Fustos et al., 2017; Fustos-Toribio et al., 2021). On
114 May 31st, 2021, a very fast debris flow was triggered due to extreme rainfall affecting houses and blocking the CH-201 route
115 in the Ñisoleufu zone, southern Chile, generating economic losses in a vulnerable rural area (Figure 1B; Figure 2). The deposit
116 mainly consisted of rock blocks and trunks covered by a thin layer of debris. Much of the debris flow fell into the adjacent
117 Calafquén lake (Figure 2), causing a small tsunami. The event was extremely rapid, based on the classification proposed by
118 Hung et al., (2014), with a speed estimated to be over 3 m/s. Stand out that the debris flow experienced reactivation events on
119 June 19, 2023, and again on June 28, 2024.

120 The debris flow was deposited in a flat area of the valley (slopes between 0-20°), flowing into Lake Calafquén. The presence
121 of landslide deposits and old flows in the vicinity of the lake (Figure 2A; geologic map [Ha]) suggests that this phenomenon
122 is common in the area.

123



124

125 **Figure 1** Rainfall-Induced mass wasting in Southern Andes. A) Regional map of Ice-sheet extension during 35 ka and 20 ka as
126 example and volcanoes emplaced in the area. B) Zoom to study area with Ñisoleufu in Northern Ice sheet sector showing the weather
127 stations. C) Zoom to Northern Patagonian area showing correlation between mass wasting events and moraine lines (blue line). D)
128 Zoom to Osorno volcano area showing high debris flow generation area discussed in Fustos et al., 2022.

129




Figure 2 A) geological map of area study based on Rodríguez et al. (1999). B) slope and elevation values (m.a.s.l.).

The Nisoleufu area exhibits a geological sequence (Figure 2A) starting at the base with the Futrono-Riñihue Batholith (CPgfr), followed by stratified volcano-sedimentary units (Pliv) and topped by glacial deposits (Plm2). Within this sequence, the Sierra Quinchilca volcano-sedimentary unit has been dated to less than 1 million years and is in nonconformity contact with the underlying rocks of the Futrono-Riñihue Batholith. Consolidated glacial deposits from the last glacial maximum period show variable depths, ranging from 1 metre to 10 centimetres, and lie in erosive unconformity over the volcano-sedimentary units (Pliv). The area also exhibits mass movement deposits (HrM and Ha) covering the older units on the north slope of Sierra Quinchilca (Pliv, Figure 2A; Rodríguez et al., 1999).

3 Methodology

To assess the triggering and conditioning factors in glacial environments in Southern Andes, we assessed in detail the Nisoleufu debris flow (Figure 3). To achieve this, we employed two complementary methodologies. The first methodology involved fieldwork, including soil sampling and subsequent laboratory analysis to evaluate the geotechnical features influencing debris flow initiation. The second methodology utilized numerical models to analyse meteorological conditions in the study area, such as precipitation patterns, variations in soil moisture, and landforms associated with erosion and deposition zones linked to the landslide. This endeavour involved a meticulous examination of the rheological and hydraulic attributes of the soil, facilitating an understanding of past and contemporary surface processes. Through this analytical framework, it became

feasible to discern causative factors contributing to the event, thereby enabling an informed evaluation of the potential risk posed by analogous occurrences in the future.




Figure 3 Methodological approach.

3.1 Geomorphological and geotechnical conditions

To analyse the geomorphological characteristics of the area, we utilised a 12.5 m resolution ALOS PALSAR DEM data, from which slope, aspect, and elevation maps were derived. A terrestrial and remote sensing survey was carried out to characterise the physiography of the hillslope before and after the debris flow. We employed the Normalized Difference Vegetation Index (NDVI) to delineate the area affected by the debris flow. We calculated the NDVI using two Sentinel-2 acquisitions on May 13th, 2021, and June 14th, 2021, corresponding to the Ñisoleufu debris-flow event on May 31st, 2021. Field campaigns were conducted one day and three months after the event (June 1st, 2021 and September 2021) to characterise post-event

geomorphological features. Field observations were determinant in the identification of the sediment budget of the channel, aiding the assessment of the event's characteristics (a channelised, stony debris flow) and the movement dynamics in the hillslope. On-channel deposits were assessed using cross-sections along the Ñisoleufu sector. Peak flow marks were documented, and erosion depths were estimated based on erosion marks and bedrock exposures over three cross-sections. Moreover, an exhaustive analysis of the geomorphological changes resulting from the event was conducted. A detailed survey of the stratigraphic column was conducted at three key points (Points c1, c2 and c4 in Figure 2B). Firstly, a stratigraphy sequence at the base of the debris flow was generated, obtaining the deposit sequence and assessing previous non-documented events. Secondly, a survey was conducted in the headscarp area where the event was initiated, evaluating the geological and geomorphological conditions that led to its generation. This approach allowed for the characterisation of the strata and structures in the affected area, seeking insights into the triggering mechanisms of the flow. Additionally, a detailed evaluation of the lateral erosion caused by the flow was carried out to understand the impact of local geomorphological changes as proxies for similar glacial-volcanic environments (Bucher et al., 2024). Finally, we assessed changes in vegetation that occurred as a result of the May 31st event and its subsequent reactivations (Figure 2B), allowing the estimation of the evolution of the landscape.

To analyse the geotechnical features that support a debris flow generation or another type of mass wasting antecedent, soil samples were collected in the generation zone. We determined soil density (ρ) following the UNE 103-301-94, soil moisture content (NCh 1515), specific weight (ASTM D854-14) standards. Granulometry analysis was carried out using sieve and hydrometer methods based on Kinde et al. (2024). The liquid limit was determined using AS 1289.3.9.1, and the plastic limit was evaluated following NCh 1517/2 standard, allowing to obtain the Plasticity Index. Finally, the soils were classified based on ASTM D2487-17. These comprehensive geotechnical analyses provided crucial insights into the soil properties and their potential role in the occurrence of the debris flow event.

3.2 Hydrometeorological conditions

To analyse the hydrometeorological conditions, we investigate the influence of rainfall in the study area, analysing hourly/daily data from four weather stations (Figure 1B) from the INIA agrometeorological and DMC networks (<https://agrometeorologia.cl/>) and the Climate Hazards Group InfraRed Precipitation with Station (CHIRPS) precipitation estimates (Funk et al., 2015). Soil moisture data from the ERA5-land product was utilised to complement the analysis considering the antecedent soil moisture at different depths before debris flow (Bordoni et al., 2023; Palazzolo et al., 2023), being considered suitable due to their accurate soil moisture data in hydrological cases (Muñoz-Sabater et al., 2021). The ERA5 product provides valuable and reliable information on soil moisture, enabling a more comprehensive understanding of the hydrological conditions that could trigger debris flows in the area. To understand the water transfer capacity along the soil, we measured the layer thicknesses from visual assessment of the soil profile in the scar (Figure 4). We followed the experimental design of De Pue et al., (2019) considering two samples per layer. One sample was used to measure the soil moisture (Θ (h)) and unsaturated hydraulic conductivity (K_u) using the evaporation method (HYPROP®, Meter Group),

meanwhile, saturated hydraulic conductivity K_s was measured using KSAT® equipment (Meter Group), using the falling head method (Dane et al., 2002). The hydraulic conductivity will provide valuable information about the layer's ability to allow water to flow through, which could have played a significant role in the initiation and propagation of the debris flow.

3.3 Remote sensing and precursory signals

We assessed precursory signals estimating surface deformation by the Stanford Method for Persistent Scatterers (StaMPS; Hooper et al., 2008; Hooper et al., 2012) using Sentinel-1 C-band SAR data. We downloaded 35 ascending orbit (track 164) and 18 descending orbit (track 83) Sentinel-1 images covering the period from November 2020 to June 2021, which includes seven months before the May 31st debris flow in the Ñisoleufu sector (Table 1). We also included two acquisitions after the debris flow in both orbits to analyse the slope's response to non-rainfall and rainfall periods. Initially, we analysed data until the end of July, but snow coverage led to coherence loss in the area, resulting in a low density of Persistent Scatterer points (PS points) per km^2 . Consequently, we evaluated different combinations of bands from Sentinel-2 images (based on bands 4, 3, 2) to assess the maximum amount of SAR images available before the snow period. This approach allowed us to optimise the data selection process and continue our analysis effectively.

The SAR data was processed using open-source Sentinel Application Platform (SNAP) packages through the snap2stamps routines, enabling us to generate single-master interferograms compatible with StaMPS. Further details on the snap2stamps routine in Fomelis et al. (2018) and Blasco et al. (2018). First, the initial selection of PS points is performed based on their noise characteristics, using the amplitude dispersion criterion, which is defined by $D_{Amp} = \sigma_{Amp}/m_{Amp}$, where σ_{Amp} and m_{Amp} are the standard deviation and mean of the amplitude in time, respectively (Ferretti et al., 2001). We selected a threshold value of 0.4 for D_{Amp} as a typical threshold value (Hooper et al., 2007), and subsequently, some initial parameters were modified according to the values proposed by Höser (2018). This allowed us to plot surface soil deformation using time series, which we then compared with daily precipitation records obtained from nearby weather stations and satellite measurements. Lastly, we used the GACOS correction (Yu et al., 2018) through the TRAIN toolbox (Bekaert et al., 2015) to reduce the atmospherical phase component. We complemented surface deformation with antecedent precipitation to establish precursory signals and their correlation with precipitation. We calculated the accumulated precipitation between the dates of the SAR acquisitions and the total accumulated precipitation for the entire period. We assessed daily precipitation and their temporal changes to understand the antecedent precipitation in the debris flow event. The four weather stations were identified within a reasonable radius for potential use as sources of meteorological data. However, upon examining the temporal coverage and continuity of their records, it was found that only the Pucón station had complete and operational data for the study period. Moreover, we merged this data with CHIRPS dataset for comparison with the time series of deformation to examine the relationship between precipitation and deformation.

Table 1 Data used in temporal analysis using SAR data.

	Tra ck	Fra me	First Image	Last Image	Total Images	Primary Image	Sub- Swath
DES	83	93	02 November 2020	18 June 2021	18	02 March 2021	IW2
ASC	164	1048	01 November 2020	17 June 2021	35	11 February 2021	IW3

226 **4 Results**

227 **4.1 Geomorphological and geotechnical conditions**

228 The extension of the area affected by the debris flow was evaluated, identifying and characterizing the triggering conditions
229 (Figure 5A). The results revealed significant differences in the Normalized Difference Vegetation Index (NDVI), which
230 facilitated the delimitation of the landslide area (Figure 5A). Low positive NDVI values (<0.20) are shown for the area affected
231 by the debris flow, encompassing 118,575 m², in contrast to the normally high NDVI values (0.6-0.8) of surrounding areas.
232 The debris flow moved along a complex and abrupt geometry, with slopes varying from 20° near the ridges (950 m. a.s.l.) and
233 the base of the slope (250 m a.s.l.), to almost vertical areas (~90°) in the intermediate zone (550 m a.s.l.) (Figure 4A **and** Figure
234 5D). This geomorphological configuration is typical of glacially eroded valleys (U-shaped valleys) in the southern Andes, a
235 recurrent phenomenon in the formation of valleys at this latitude (Muratli et al., 2010).

236 The initial landslide crown has an altitude of 950 m. a.s.l. and slopes of approximately 20 to 30° (column c1 in Figure 4B).
237 The flow release zone (inset C in Figure 4) has evidence of extensional failures, where c1 indicates that the first level S-1
238 corresponds to a very compact chaotic and polymictic till deposit (Plm2), with a greyish matrix containing a higher percentage
239 of clay than sand. Some clasts exceed 10 cm and are composed of volcanic fragments (Pliv), as well as intrusive material
240 (CPfgr). Towards the top of the glacial deposit, level S-2 is observed, a thin, grey fluvioglacial deposit (varves), approximately
241 40 cm thick, composed of a well-consolidated matrix of clay-rich (dark) and silt-rich (light) setting a couplet annual sediment
242 layer. (Figure 4C, Figure 4D **and** Figure 5D). Above the glacial deposit, level S-4 is identified, composed mainly of lapilli-
243 sized deposits (>5 mm), associated with pumice from the Neltume deposit of the Mocho-Choshuenco Volcanic Complex,
244 dating 10,200+-500 BP (Rawson et al., 2015; Moreno-Yaeger et al., 2024). Finally, level S-7 corresponds to the current soil
245 where native forest develops (Figure 4B).




Figure 4 Field photographs indicating soil deposits in scarp (B and C) and the deposits in the toe (inset D and E).

Stratigraphic analysis along the slope (Figure 5B–D) reveals a consistent sequence overlying the CPfgr basement, characterized by a basal glacial deposit (S-1) with large angular clasts, overlain by silt–clay layers (S-2). These low-permeability units are covered by interbedded paleosols and volcanic ash-falls. In column C3, the paleosol S-3 was initially considered older than the Neltume ashfall, but the presence of reworked pyroclasts in S-4 (Figure 4D) suggests a younger relative age. The upper sequence concludes with active soil formation (S-7; Figure 5D).




Figure 5 A) Normalized Difference Vegetation Index (NDVI) highlighting the erosive zone generated by the Nisoleufu debris flow. B) and C): Photos taken by a drone on 01 June 2021, highlighting the crown of the debris-flow (B), and the landslide deposit and the development of many waterfalls around the flow in red (C). D) Profile of the sequence where the flow occurred and stratigraphic columns. Photos: Carrasco and Ramirez (2021).

Further downslope (250 m a.s.l.), column C4 exhibits a distinct stratigraphy dominated by mass-wasting deposits. The basal level (S-5) consists of a polymictic unit with sub-rounded clasts and pumice fragments from the Neltume event, embedded in a clay-rich matrix derived from upstream glacial units. A similar, though finer, deposit (S-6) overlies S-5, and is capped by the same modern soil unit (S-7). This stratigraphic framework—composed of alternating low-permeability substrates and reworked tephra—is representative of Andean terrains between 39° and 42°S and is critical to understanding hydrological storage and slope instability (Figure 1).




Figure 6 Soil granulometric curve of S-3 and S-7.

Table 2 Physical properties of soils related to the column C3.

Soil type/Property	Normative	S-2	S-3	S-4	S-7
Moisture [w] (%)	NCh-1515	17.8	56.2	119.3	111.6
Density [ρ] (g/cm ³)	UNE-103-301-94	2.07	1.52	<1	1.06
Specific Gravity [G_s]	ASTM-D854-14	2.76	2.49	2.5	2.34
Liquid Limit [W_L] (%)	AS 1289.3.9.1	27.48	123.93	-	149.83
Plastic Limit [W_P] (%)	Nch 1517/2	16.07	91.3	-	114.13
Plasticity Index [PL]	NCh1517/2	11	33	-	36
Hydraulic Conductivity [k_u] (m/s)	Porchet and Laferrere (1935)	-	-	-	3.13E-4

From a geotechnical perspective, the first soil (S-2) has a liquid limit of 27.48 and a plastic limit of 16.07, resulting in a plasticity index of 11. This soil exhibits a low plasticity and is classified as a silt (CL) according to the Unified Soil Classification System (USCS). S-3 showed a liquid limit of 123.93 and a plastic limit of 91.3, resulting in a plasticity index of 33. This soil exhibits a moderate plasticity and is classified as a plastic silt (CH), according to the USCS. Meanwhile, S-7 has a liquid limit of 149.83 and a plastic limit of 114.13, resulting in a plasticity index of 36, showing a high plasticity and is

classified as a plastic organic soil (OH). The granulometric analysis (Figure 6; S-3 and S-7) confirms this classification, with silt and sand content between 70% and 90%. Due to the soil properties, S-4 (Neltume ashfall pyroclastic) has not been characterised in the laboratory due to the high fragility of the pyroclasts classified as lapilli ($\phi > 5$ mm) being too coarse to measure their limits.

4.2 Hydrometeorological conditions and precursory signals

The debris flow release zone (column c1 in Figure 4B and Figure 5B) provided accurate soil measurements using KSAT, enabling us to understand the factors governing water transport on the slope. Field mapping revealed a layered model that identified a highly stratified environment with distinct characteristics (Figure 4C; Figure 7). It was observed that the glacial deposits underlying layers of volcanic origin regulate water movement in the soil due to their low hydraulic conductivity (Figure 7). These glacial deposits, identified as moraines and varves, are associated with the Last Glacial Maximum (LGM), correlated with nearby deposits and natural terrain conditions.

Notably, volcanic soil deposits (Figure 6; granular texture description) and Neltume ashfall demonstrating high saturated hydraulic conductivity of 4.64×10^{-5} to 3.31×10^{-4} m/s (Figure 7). This characteristic is critical as it facilitates the movement of infiltrated water from the organic surface to the slope's interior. In contrast, Varves-type glacial deposits show low hydraulic conductivity, with values of 1.54×10^{-5} m/s, while the till reached a K_s of 2.65×10^{-5} m/s, both acting as partial barriers to water movement. Nevertheless, their storage capacity is significant, particularly when situated on moraines that, due to their varied granulometric distribution, retain water effectively. The hydraulic conductivity and the high infiltration rate derived from the Porchet test showed a moderately high infiltrated rate (112.70 mm/hr), computing a $k = 3.31 \times 10^{-4}$ (m/s) in the superior organic deposit layer (S-7; Table 2 and Figure 7).




Figure 7 Hydraulic properties of release zone in debris flow generation zone (column c1 in [Figure 1](#)).

Analysis using the ERA5 model reveals a well-defined annual cycle in soil moisture (Figure 8A). While the overall evolution of moisture content does not indicate an anomalous trend when compared to previous years, significant short-term variations in soil moisture were observed (Figure 8B) in response to extreme precipitation events (Figure 8C). The preliminary results showed that antecedent rainfall led to surface saturation, with moisture levels reaching full saturation within the first metre of soil depth, primarily due to accumulated rainfall in the days prior (Figure 8B–C). Notably, a substantial and rapid increase in soil moisture was also detected at greater depths—up to 2.89 meters—where a 50% change occurred shortly before the onset of debris flow events (Figure 8B). This marked fluctuation was concentrated at the interface between tephra and till-varves, suggesting critical implications for slope instability. Our findings underscore that, even in the absence of a long-term anomalous trend, the combination of saturated soil conditions and intense rainfall plays a decisive role in triggering debris flows.




Figure 8 Assessment of ERA5-land product over debris flow generation. A: Assessment of time series of soil moisture at different depths. B: Zoom to Soil moisture weeks previous to debris flow. C: Rainfall events accumulated at 24-hourly scale (ERA5-land).

The availability of a sufficient quantity of SAR images and the revisit times of Sentinel-1 enabled a well-distributed temporal analysis of slope behaviour before the debris-flow event on May 31, 2021. We measured surface deformation between +9 and -32 mm/year. The results of PS estimation suggest the occurrence of surface subsidence consistent change in soil water content (Figure 8B). Our analysis reveals a precursory deformation signal that comprised two distinct phases (Figure 9). The first phase, beginning in late January 2021 and extending through April, exhibited a high-deformation pattern associated with consistent precipitation (Figure 8C), which infiltrated the soil and increased soil moisture levels (Figure 8B). This timing coincides with a significant precipitation event in late January, suggesting that summer rainfall may have acted as an initial trigger, initiating a surface deformation that evolved progressively over time. A second deformation phase following a high-intensity rainfall event in mid-to-late May 2021, characterized by pre-event deformation patterns. Despite that most evident precipitation event occurred in late May (15-day before), our surface deformation estimations could suggests that the triggering process likely began approximately in summer period. This temporal relationship between precipitation and deformation further underscores the need for incorporating hydrological factors into models of surface stability and landslide risk assessment. Finally, the deformation measurements are interrupted a few days before the debris flow occurs. This data interruption is attributed to the formation of significant extensional fractures (Figure 4D-E), causing a loss of coherence in the data (red box).




Figure 9 PS points obtained with Sentinel-1 “ascending” orbit data before debris flow initiation. Precipitation time series using Pucón weather station. Deformation shows an subsidence in surface.

5 Discussion

To improve debris-flow hazard assessment, it is crucial to understand how multiple controlling factors converge to trigger these events. In the Ñisoleufu case, we studied the triggering conditions of debris flow in an area to understand precursory signals of mass wasting initiation in post glacial and volcanic environments. We considered a geomorphological, geotechnical, hydrometeorological and surface deformation approach to constraint the variability of the mass wasting processes in one representative area of the Southern Andes.

5.1 Geomorphological and geotechnical implications

The occurrence of mass wasting events, such as landslides and debris flows, following periods of intense rainfall has been extensively studied in the Southern Andes using local cases, mainly based on the rainfall control over the mass wasting generation (Fustos et al., 2020; Maragaño et al., 2023) without consider the soil features as noted recently (Vasquez-Antipan et al., 2025). Our results showed that the geomorphology plays a crucial role in the generation of debris flows, particularly

342 through its influence on water accumulation in catchment areas, such as micro-basins (Fustos-Toribio et al., 2021). We
343 considered two main geomorphological controls for debris flow initiation. First, stand out the steep slope ($>45^\circ$), which was a
344 significant contributing factor to the generation of the Ñisoleufu debris flow (Figure 2 **and** Figure 5B-D). Second, the northern
345 orientation of the slope could also be a relevant factor, as in the central-southern Chile domain (36° - 42° S), atmospheric moist
346 flux from extreme rainfall tends to flow in a northwest (NW) direction, with orographic forcing triggering being enhanced by
347 the Andean belt (Valenzuela and Garreaud, 2019; Vasquez-Antipan et al., 2025). The interplay between areas with high water
348 accumulation capacity from local runoff and the slope's propensity to capture precipitation was instrumental in the generation
349 of the debris flow. Stand out the slope that facilitated an efficient surface drainage of water from higher areas, leading to the
350 formation of local accumulation and infiltration zones that promote the build-up of subsurface pore-water pressures (Figure
351 6B) leading to the debris flow event, facilitating a rapid downslope movement of soil once it became saturated.

352

353 The geotechnical stability of soils in the Southern Andes, could be controlled by the interaction between explosive volcanic.
354 Mocho-Choshuenco volcanic complex (MCVC) and glacial processes and the resultant varved deposits (Figure 5) has been
355 modulated the slope stability around the 39° S. The Neltume ashfall deposit related to the eruption of the MCVC ($10,200\pm 500$
356 BP; Rawson et al., 2015) played a significant role in the formation of the S-4 (Figure 4B-C). The thickness of all soils varies
357 along the scar left by the debris flow, with the magnitude strongly influenced by the topographic gradient and/or erosion
358 processes that might have occurred and following reworking (Figure 4D). Moreover, glacial deposit like as varves, formed
359 from sedimentation during glacial periods, played a crucial role in the soil's physical and mechanical properties controlling the
360 area with low hydraulic conductivity (Figure 7) introducing a water barrier in the area to regional scale.

361 Moreover, field evidence suggests that the Ñisoleufu event is not an isolated case, as indicated by other remobilised events in
362 the area (Figure 2, geological map – alluvial deposit: Ha). The geotechnical properties of the remobilised materials are critical
363 for defining slope stability conditions. Granulometric analyses indicate that the deposits are primarily granular soils, such as
364 those associated with S-4, which are classified as frictional and are found overlying finer-grained cohesive soils, such as varves
365 (S-2). Other soils found in the Southern Andes, including S-3 and S-7, originate from the decomposition of volcanic glass and
366 glacial clays (Sanhueza et al., 2011; Vasquez et al., 2025), producing particles smaller than 0.1 mm (Figure 6).

367 Specifically, S-3 soils, derived from explosive eruptions of the Mocho-Choshuenco volcano, consist of non-cohesive volcanic
368 ash mixed with fine-grained sediments, forming a matrix with elevated plasticity and a high liquid limit (Vasquez et al., 2025).
369 These properties result from the introduction of fine material during the deposition. Moreover, S-7 soils, classified as organic
370 soils derived from volcanic deposits, exhibit notably high liquid limits due to the accumulation of organic matter. The organic
371 matter enhances the soil's water retention and promotes the formation of organic colloids, which may increase the liquid limit
372 (Deng et al., 2017; Fiantis et al., 2019). Our results are consistent with independent laboratory testing in the zone (Vásquez et
373 al., 2025), which shows that organic-rich paleosols were buried after the Last Glacial Maximum, approximately 5 km south of
374 the study area, and exhibit similar liquid limit values to those observed in S-7.

375 The spatial distribution of soil layers varies abruptly along the slope, as observed in columns C1 and C3 for S-1, S-2, and S-4,
376 indicating significant mass wasting and erosion processes near glacial lakes (Figure 5D). The frictional soils, such as those
377 related to S-4, generally exhibit high shear strength (Chen et al., 2021), and when combined with steep topography, may
378 contribute to the relative stability of post-glacial volcanic deposits (Walding et al., 2023; Ontiveros-Ortega et al., 2023).
379 However, under extreme precipitation events—such as those recorded in recent years in the Southern Andes—soil saturation
380 can substantially reduce the strength of even frictional soils, increasing the likelihood of failure (Fustos et al., 2017; Somos-
381 Valenzuela et al., 2020; Fustos et al., 2021). This mechanism aligns with the observed extensional failures that preceded the
382 initiation and reactivation of flows in June 2023 and 2024 (Figure 4C; Figure 10).
383 We propose that the event of Ñisoleufu, a classical case of the Southern Andes, was triggered by the soil saturation, reducing
384 the effective stress within the soil matrix leading to extensional failure in the volcanic deposits (Figure 4B; Figure 10),
385 highlighting the risks associated with this saturation, particularly in areas where explosive volcanic activity has previously
386 occurred, leading to increased susceptibility to debris flows and other forms of mass wasting (Korup et al., 2019). The
387 interaction between glacial deposits and volcanic materials creates a unique geotechnical environment where the stability of
388 slopes is contingent upon both the physical properties of the soil and the hydrological conditions present.


389 **5.2 Hydrometeorological conditions and precursory signals implications**

390 The volcanic origin and composition of the soils evidenced a high soil moisture variability along the year. ERA5-land data
391 reveals a sudden change in soil moisture content at various depths on the base of the debris flow initiation supporting the
392 surface deformation days before the landslide (Figure 9). The water role in the event proposes that accumulation of subsurface
393 water in granular soil could serve as a storage medium. The fine media of S-1 and S-2, with its low hydraulic conductivity,
394 acts as barrier layer, enhancing the storage capacity of S-4 in the crown of the debris flow. This is supported by the hydraulic
395 properties of S-3 (CH) and S-7 (OH), combined with Porchet's study on the current soil (S-7: 112.70 mm/hr), which indicates
396 a moderately high infiltration rate and suggests high hydraulic conductivity for the Ñisoleufu soils (Figure 7). The combination
397 of these factors contributes to the overall permeability of the volcanic soils, a main feature in the Southern Andes, potentially
398 influencing water movement and retention in the area.

399 The water storage is consistent with surface deformation data, reaching -85 mm/yr in the previous 30 days of the event, offering
400 insights of previous surface deformation to the occurrence of debris flow events in areas where glacial deposits and volcanic
401 deposits coexist (Figure 9). The atypical nature of these deformations suggests that the surface shows a slow movement related
402 to subsidence along LOS, followed by the debris-flow event on May 31, 2021 (159.8 mm/year). The constant slope deformation
403 supports the hypothesis of the development of extensional failure, ultimately resulting in the observed a small landslide and
404 subsequent release of water as debris flows (inset C in Figure 4; Figure 10).

405 The fast saturation of S-4 is exacerbated by increased water availability due to rainfall and by the presence of a bottom layer
406 with low hydraulic conductivity associated with fine soils (S-2), being a common denominator in the Southern Andes. S-2
407 retains water and further reduces the shear strength of the underlying granular soil. Additionally, these soils could act as

lubricants, reducing the overall slope shear strength. The interaction between glacial deposits and explosive volcanic eruption deposits in the cordilleran zone of the Southern Andes creates a scenario prone to slope deformation and mass removal, especially debris flow. Our results suggest that the combination of geological and climatic factors in this region generates ideal conditions for the occurrence of mass removal events under extreme precipitation events (Savi et al., 2016).



Our results showed a limited amount of PS in the study area similar to previous studies in the area (Vasquez-Antipan et al., 2025). The Southern Andes, and the Ñisoleufu area, is characterized by complex geomorphological features and varying precipitation patterns that could introduce uncertainty to the remote sensing measurements. The application of limited persistent scatterer data in assessing slope deformations offers a promising avenue for the development of a landslide early warning system (LEWS) in the Southern Andes. However, the investigation into this method requires a thorough understanding of potential limitations, particularly the extensive vegetation cover in the Southern Andes, which can obscure satellite signals and affect data accuracy. Vegetation serves as a significant barrier to radar signals, leading to incomplete datasets that might

obscure important geological signals indicative of slope movements (Maragaño-Carmona et al., 2023). Therefore, additional efforts must be considered to move forward to an operational scale.

Climate change is increasingly debris flow generation by altering precipitation patterns and soil moisture dynamics (Talebi et al., 2007). In the Southern Andes, volcanic soils with variable textures play a critical role in this process. Enhanced seasonal moisture variability, exacerbated by extreme precipitation, leads to fast soil saturation, especially where fine-grained soils form low-permeability layers above coarser materials (Figure 10). These stratified soil conditions promote subsurface water storage, increasing the slope instability under saturated conditions (Talebi et al., 2007). Fine volcanic over glacial deposits can act as lubricants, further weakening slope cohesion and promoting failure (Espinosa et al., 2024) during intense rainfall, as happened during the Ñisoleufu event (Figure 7B). This event highlights how short-duration storms, increasingly associated with climate change, can overwhelm the buffering capacity of mountainous terrain. The soil media S-7 and S-4, both composed of organic-rich and granular volcanic materials, played a critical role in this response. During the 2023 event, infiltrating rainwater rapidly percolated through these coarse upper layers until reaching the underlying varved glacial sediments (S-2), which have significantly lower permeability. This layering caused a perched water table, increasing the pore pressure and reducing shear strength, ultimately contributing to slope failure. These effects were captured in our remote sensing observations, which showed expanded saturated zones and local instability near the contact between volcanic and glacial deposits.

Our results suggest strong stratigraphic controls and extreme precipitation events in the soils derived from volcanic materials overlying denser glacial layers, acting as failure planes under saturated conditions (Figure 7; Figure 8B). Our conceptual model promotes water retention and changes in soil pore pressure, especially during extreme rainfall events such as 2021 (not observed previously). The conditioning factors are further exacerbated by mid-term climate trends, including the ongoing megadrought in the Southern Andes (Garreaud et al., 2019), which increased desiccation cracking, and weakened root cohesion. Such drought-induced degradation lowers slope resistance, making even moderate precipitation more hazardous. Therefore, our observations suggest that the Ñisoleufu debris flow event, may exemplify the climate-induced changes in both hydrological extremes and landscape memory (drought legacy) act in concert to reduce slope stability. As extreme precipitation becomes more frequent under future climate scenarios, similar failures are expected to occur across a broader area than observed in past events (Figure 2A). Our findings stress the urgent need for debris flow forecasting models to incorporate stratified soil behavior, seasonal soil moisture dynamics, and drought-related weakening—factors essential to anticipating the growing hazard posed by climate change (Iverson et al., 2010; Gariano and Guzzetti, 2016).

5.3 Regional implications and Future Scope

The Southern Volcanic Zone (37.5° - 41.5°S; Stern 2004) has been shaped by significant volcanic Holocene eruptions (Fontaine et al., 2021; Moreno-Yaeger et al., 2024; Singer et al., 2024), leading to the formation of soils primarily composed of lapilli and/or ashes that settled following the contours of the topography, resulting in layers with varying thicknesses (Stern,

2007). Old deposits in the zone show similarities to the current debris flow deposits (Figure 5B; Figure 10A), suggesting that the event triggered in 2021 is not an isolated occurrence in the area. This evidence points to a history of recurring debris flow events in the region. This condition is dominant in all the surroundings of the Mocho-Choshuenco volcanic complex (Rawson et al., 2015; Moreno-Yaeger et al., 2024).

The Ñisoleufu debris flow exhibited a characteristic pattern of mass-wasting processes in the Southern Andes, similar to the Petrohué event (Fustos et al., 2021) at Osorno Volcano (Figure 1D), which can be attributed to comparable climatic and volcanic conditions. The occurrence of debris flows in the Southern Andes, particularly in the border of the maximum extension of the Patagonian ice sheet suggest that debris flows are typically related to saturated soils that can transform into more fluidic mixtures. Our observations and previous studies (Davies et al., 2020; Fustos et al., 2021) proposes a strong correlation between debris flows and glacial moraines, particularly where the last glacial maximum shaped the relief (Figure 1B and Figure 1C). The interaction of past glacial dynamics with contemporary environmental processes related to new precipitation patterns provides a backdrop for increased debris flow activities in Southern Andes based on rainfall-induced mass wasting data from Fustos et al. (2022). Stand out regions at the borders of these past glaciers tend to exhibit increased susceptibility to debris flow generation due to the combination of steep slopes and the prevalence of loose moraine deposits (Figure 8B and C), which can be mobilized during significant precipitation events (Sepúlveda et al., 2014).

Our results show that precursory signals such as small progressive deformation of the surface (Figure 10) could suggest the indication of possible soil slides previous to initiation of the debris flow (Figure 10). This phenomenon is influenced by fine soil interspersed with glacial deposits, superimposed on moraines and volcanic deposits, common throughout the southern Andean region (Moreno-Yaeger et al., 2024; Singer et al., 2024). A simple geomorphological generalization highlights the necessity of investigating the interaction between glacial and volcanic deposits to understand better the interplay between media with constraints on soil hydraulic conductivity. Future studies should prioritise examining these dynamics to prevent and mitigate potential risks associated with similar landslide events across the broad area of the Southern Andes.

Debris flow generation in volcanic soils and their correlation with post-glacial eruptions, assumes a critical importance in comprehending the mass wasting hazards under the current climate crisis. Holocene eruptions, known for their Plinian events such as Mocho-Choshuenco, Carran-Los Venados, Calbuco, Chaiten in the Southern Andes (Singer et al., 2024), have yielded abundant tephra and volcanic soils possessing high hydraulic conductivity. These eruptions formed volcanic deposits over moraine and varve deposits (Figure 4). The volcanic soil acts as a high-rate infiltration layer meanwhile, the moraine crowned by impermeable varve layers in the base of the sequence plays a reservoir on (Figure 10). Therefore, the amalgamation of these deposits with intense precipitation events may expedite the loading of these reservoirs, heightening the probability of slope instability due to increased mechanical load and pore-stress changes (Bogaard & Greco, 2015). Early indications of this phenomenon are evident in occurrences such as Choclo 2023, and Volcán Osorno (every year), suggesting a potentially heightened recurrence in the future. To mitigate the associated risks effectively, it is imperative to enhance the zoning of areas

491 prone to debris flows in the Southern Andes, thus minimizing the impact on population and infrastructure within these
492 vulnerable zones.

493 On a regional scale, climate change is intensifying debris flow hazards (Gariano and Guzzetti, 2016). In the Southern Andes,
494 current changes in precipitation patterns will affect the stability of volcanic and glacial deposits through alterations in water
495 storage, as noted in this study (Figure 8B). The region's unique stratigraphy in South America, where volcanic soils overlay
496 glacial sediments, may become unstable during extreme rainfall events. Significant shifts in precipitation patterns, as predicted
497 by CMIP6 models, alter the spatial distribution of precipitation and their impact on soil moisture storage, with limited accurate
498 estimations (Salazar et al., 2023). We propose that future developments should carefully constrain areas with high
499 susceptibility to debris flow. Therefore, improved hazard debris flow delimitation and instrumental monitoring become critical
500 for reducing the impact of these hazards in the Southern Andes.

501 Our study contributes to understanding the relationship between volcanic and glacial deposits under extreme rainfall events
502 forcing in the Southern Andes. However, further research is necessary to improve the rainfall-induced mass wasting
503 susceptibility models due to an incomplete integration of critical soil properties. Several studies focus on rainfall thresholds as
504 primary trigger, oversimplifying the failure mechanisms. Our study proposes that additional assessment of the hydraulic and
505 mechanical influence of specific soil layers must be considered. Additionally, the pronounced spatial heterogeneity in soil
506 layer composition in Southern Andes, ranging from S-1 to S-7, and variability in thickness and permeability, further
507 complicates predictive accuracy to regional scale. Future developments must consider high-resolution subsurface mapping
508 introducing national scale models (Dinamarca et al., 2024), allowing better mass wasting risk models overlooking zones where
509 possible saturated soil could appear, leading to sudden failure. Moreover, surface deformation frequently observed as a slow
510 extensional signal prior to collapse introduces a limited amount of PS. Moreover, the hydrometeorological variability indicates
511 that better soil moisture models are necessary in the zone to improve the slope stability analysis. Together, these limitations
512 highlight the urgent need for multidisciplinary approaches that integrate geotechnical, geomorphological, and
513 hydrometeorological data into landslide hazard assessments.

514 Our study proposes considerations that must be assessed in landslide research in the Southern Andes, allowing a deep
515 understanding of the interplay between geomorphological, geotechnics, hydrometeorological and remote sensing analysis,
516 allowing a better landslide risk assessment and mitigation efforts in the region. We propose that the Ñisoleufu event could as
517 study case for researchers and authorities can better comprehend the specific conditions that lead to debris-flow occurrences
518 and implement appropriate measures to minimize the impact of such events in the future. Finally, these results suggest that
519 hazard assessment protocols should include hydrogeotechnical mapping of tephra-over-moraine systems, combined with
520 seasonal soil moisture dynamics and weather forecasts of extreme rainfall as early indicators of instability in these settings.

6 Conclusions

We studied the conditions that evolved in the generation of debris flows in the Southern Andes, an area modulated by the glacial and volcanic processes. The mass wasting could be influenced by complex interactions among geomorphological, geotechnical, and hydrometeorological factors (Figure 10). The geological environment of the Southern Andes, characterized by a mix of volcanic and glacial deposits, showcases unique soil properties affecting overall slope stability. The mechanisms leading to mass wasting, particularly the Ñisoleufu event, underscore the critical roles of soil saturation and effective stress reduction in triggering slope failures (Figure 8). The analysis of geomorphological factors revealed that slopes greater than 30 degrees significantly contribute to debris flow triggers, supporting the emphasis of rainfall's role in mass wasting (Figure 5). The orientation of these slopes, particularly aspects with more rainfall exposure, promotes precipitation accumulation from extreme weather patterns. Our findings corroborate the influence of gradual accumulation of subsurface water, aided by low hydraulic conductivity from underlying soil layers (Figure 7), can create critical saturation levels that reduce shear strength and lead to flow initiation. These processes, not considered in detail must be integrated in future assessment in detail into the future.

We conclude that recurrence of mass wasting is influenced by past volcanic eruptions and post-glacial conditions. A comprehensive understanding of the interplay between geological and hydrometeorological conditions is crucial for forecasting debris flow risks in the Southern Andes, particularly considering climate change, which may exacerbate extreme weather events. We highlight that while frictional soils can provide stability under dry conditions, they become increasingly vulnerable to failure under saturated conditions as was evidenced in the Ñisoleufu event. The data indicating surface deformation prior to debris flow events (Figure 9), demonstrating that precursory signals that can be leveraged for early warning systems. The velocity of surface movements preceding the Ñisoleufu event are correlated with increased soil moisture levels, emphasizing the utility of integrating remote sensing technologies to monitor these changes as proxy. Our findings could be extended to regional scale as a conceptual model of the landslides and debris flows in the Southern Andes, suggesting areas prone to such occurrences and should be monitored closely to develop effective mitigation strategies. Finally, our results demonstrate the value of integrating geomorphological, hydrometeorological, and hydrogeotechnical data to support debris-flow hazard assessments. In particular, the combination of tephra layers overlying low-permeability glacial deposits, together with rapid water infiltration on steep slopes during extreme rainfall events, defines a critical configuration that enhances susceptibility to failure. This integrated framework provides a robust basis for identifying high-risk areas and strengthening early warning strategies in the Southern Andes and comparable volcanic-glacial settings worldwide.

7 Code availability

All the codes used in this manuscript are reproducible from the main text. We can deliver the main script under any request.

Data availability

The datasets used in this study are available in the paper. The ALOS-PALSAR DEM is publicly accessible at <https://search.asf.alaska.edu/#/?dataset=ALOS> (last access: 30 September 2024). Additional information about the information or datasets can be obtained from Ivo Janos Fustos-Toribio, ivo.fustos@ufrontera.cl.

Author contributions

IF, DB, AB, GF and AG contributed to the conceptualisation and methodology of the research and performed the formal analysis, visualisation and validation. IF and DB were involved in the funding and supervision of the paper. IF and AB contributed with the supervision, review and editing of the paper. SS and JLP contributed to the discussion of the scientific results. All the authors provided input in terms of methodology and the review and editing of the paper.

Competing interests

The authors declare that they have no conflict of interest.

Disclaimer

Acknowledgements

This work was made possible thanks to the “Agencia Nacional de Investigación y Desarrollo (ANID)” of the Chilean Government; “Fondecyt Regular” grant 1230792); “Fondecyt post-doctoral” grant 3200387) and CIVUR-39° “Centro Interactivo Vulcanológico de La Araucanía” Project UFRO2193 of the Desarrollo de Actividades de Interés Nacional (ADAIN), Ministry of Education, Chilean Government. We appreciate the support of Mauricio Hermosilla for their support in geotechnical analysis.

Financial support

This research has been supported by ANID (grant nos. Fondecyt 3200387 and Fondecyt 1230792). Laboratory instruments were supported by FONDEF ID23i10118.

574 14 References

- 575 Bekaert, D. P. S., Walters, R. J., Wright, T. J., Hooper, A. J., and Parker, D. J.: Statistical comparison of InSAR tropospheric
576 correction techniques, *Remote Sensing of Environment*, 170, 40–47, doi: 10.1016/j.rse.2015.08.035, 2015.
- 577 Blasco, J. M. D. und Foumelis, M.: Automated SNAP Sentinel-1 DInSAR processing for StaMPS PSI with open source tools,
578 Zenodo, doi: 10.5281/ZENODO.1322353, 2018.
- 579 Bovis, M. J. and Jakob, M.: The role of debris supply conditions in predicting debris flow activity, *Earth Surf. Process.*
580 *Landforms*, 24, 1039–1054, [https://doi.org/10.1002/\(sici\)1096-9837\(199910\)24:11<1039::1039:aid-esp29>3.0.co;2-u](https://doi.org/10.1002/(sici)1096-9837(199910)24:11<1039::1039:aid-esp29>3.0.co;2-u), 1999.
- 581 Bogaard, T. A. und Greco, R.: Landslide hydrology: from hydrology to pore pressure, *WIREs Water*, 3, 439–459, doi:
582 10.1002/wat2.1126, 2015.
- 583 Bordoni, M., Vivaldi, V., Ciabatta, L., Brocca, L., und Meisina, C.: Temporal prediction of shallow landslides exploiting soil
584 saturation degree derived by ERA5-Land products, *Bull Eng Geol Environ*, 82, doi: 10.1007/s10064-023-03304-2, 2023.
- 585 Bucher, J., del Papa, C., Hernando, I. R., und Almada, G.: Upper-flow-regime deposits related to glacio-volcanic interactions
586 in Patagonia: Insights from the Pleistocene record in Southern Andes, *Sedimentology*, doi: 10.1111/sed.13216, 2024.
- 587 Chen, Y., Lin, H., Cao, R., und Zhang, C.: Slope Stability Analysis Considering Different Contributions of Shear Strength
588 Parameters, *Int. J. Geomech.*, 21, doi: 10.1061/(asce)gm.1943-5622.0001937, 2021.
- 589 Carrasco Felipe and Ramírez Paola: Caracterización de remociones en masa del 31 de mayo de 2021, en la ribera sur del Lago
590 Calafquén, sobre la Ruta CH-201, Comuna de Panguipulli, Región de Los Ríos. Informe inédito. INF-LOS RÍOS-09.2021.
591 2021. Subdirección Nacional De Geología, Sernageomin, Stgo, Chile. PP:23.
- 592 Chang, J.-M., Chen, H., Jou, B. J.-D., Tsou, N.-C., und Lin, G.-W.: Characteristics of rainfall intensity, duration, and kinetic
593 energy for landslide triggering in Taiwan, *Engineering Geology*, 231, 81–87, doi: 10.1016/j.enggeo.2017.10.006, 2017.
- 594 Chang, J.-M., Yang, C.-M., Chao, W.-A., Ku, C.-S., Huang, M.-W., Hsieh, T.-C., & Hung, C.-Y. (2025). Unraveling landslide
595 failure mechanisms with seismic signal analysis for enhanced pre-survey understanding. *Natural Hazards and Earth System*
596 *Sciences*, 25(2), 451–466. <https://doi.org/10.5194/nhess-25-451-2025>
- 597 Chen, Y.-W., Shyu, J. B. H., and Chang, C.-P.: Neotectonic characteristics along the eastern flank of the Central Range in the
598 active Taiwan orogen inferred from fluvial channel morphology, *Tectonics*, 34, 2249–2270,
599 <https://doi.org/10.1002/2014tc003795>, 2015.
- 600 Cheung, D. J. und Giardino, J. R.: Debris flow occurrence under changing climate and wildfire regimes: A southern California
601 perspective, *Geomorphology*, 422, 108538, doi: 10.1016/j.geomorph.2022.108538, 2023.
- 602 Dahal, R. K. und Hasegawa, S.: Representative rainfall thresholds for landslides in the Nepal Himalaya, *Geomorphology*, 100,
603 429–443, doi: 10.1016/j.geomorph.2008.01.014, 2008.
- 604 Dahal, R. K., Hasegawa, S., Yamanaka, M., und Bhandary, N. P.: Rainfall-induced landslides in the residual soil of andesitic
605 terrain, western Japan, *Journal of Nepal Geological Society*, 42, 137–152, doi: 10.3126/jngs.v42i0.31461, 2011.
- 606 Dane, H. Topp, G., 2002. *Methods of Soil Analysis Part-4 Physical Methods*.

607 Davies, B. J., Darvill, C. M., Lovell, H., Bendle, J. M., Dowdeswell, J. A., Fabel, D., García, J.-L., Geiger, A., Glasser, N. F.,
 608 Gheorghiu, D. M., Harrison, S., Hein, A. S., Kaplan, M. R., Martin, J. R. V., Mendelova, M., Palmer, A., Pelto, M., Rodés,
 609 Á., Sagredo, E. A., Smedley, R. K., Smellie, J. L., and Thorndycraft, V. R.: The evolution of the Patagonian Ice Sheet from
 610 35 ka to the present day (PATICE), *Earth-Science Reviews*, 204, 103152, <https://doi.org/10.1016/j.earscirev.2020.103152>,
 611 2020.
 612 Deng, Y., Cai, C., Xia, D., Ding, S., Chen, J., and Wang, T.: Soil Atterberg limits of different weathering profiles of the
 613 collapsing gullies in the hilly granitic region of southern China, *Solid Earth*, 8, 499–513, [https://doi.org/10.5194/se-8-499-](https://doi.org/10.5194/se-8-499-2017)
 614 2017, 2017.
 615 Dey, N. und Sengupta, A.: Effect of rainfall on the triggering of the devastating slope failure at Malin, India, *Nat Hazards*, 94,
 616 1391–1413, doi: 10.1007/s11069-018-3483-9, 2018.
 617 De Pue, J., Rezaei, M., Van Meirvenne, M., und Cornelis, W. M.: The relevance of measuring saturated hydraulic conductivity:
 618 Sensitivity analysis and functional evaluation, *Journal of Hydrology*, 576, 628–638, doi: 10.1016/j.jhydrol.2019.06.079, 2019.
 619 Dinamarca, D. I., Galleguillos, M., Seguel, O., and Faúndez Urbina, C.: CLSoilMaps: A national soil gridded database of
 620 physical and hydraulic soil properties for Chile, *Sci Data*, 10, <https://doi.org/10.1038/s41597-023-02536-x>, 2023.
 621 Ferretti, A., Prati, C., und Rocca, F.: Permanent scatterers in SAR interferometry, *IEEE Trans. Geosci. Remote Sensing*, 39,
 622 8–20, doi: 10.1109/36.898661, 2001.
 623 Fiantis, D., Ginting, F., Gusnidar, Nelson, M., and Minasny, B.: Volcanic Ash, Insecurity for the People but Securing Fertile
 624 Soil for the Future, *Sustainability*, 11, 3072, <https://doi.org/10.3390/su11113072>, 2019.
 625 Fontaine, C. M., Siani, G., Delpech, G., Michel, E., Villarosa, G., Manssouri, F., und Nouet, J.: Post-glacial tephrochronology
 626 record off the Chilean continental margin (~41° S), *Quaternary Science Reviews*, 261, 106928, doi:
 627 10.1016/j.quascirev.2021.106928, 2021.
 628 Fomelis, M., Delgado Blasco, J. M., Desnos, Y.-L., Engdahl, M., Fernandez, D., Veci, L., Lu, J., und Wong, C.: Esa Snap -
 629 Stamps Integrated Processing for Sentinel-1 Persistent Scatterer Interferometry, *IGARSS 2018 - 2018 IEEE International*
 630 *Geoscience and Remote Sensing Symposium*, doi: 10.1109/igarss.2018.8519545, 2018.
 631 Funk, C., Peterson, P., Landsfeld, M., Pedreros, D., Verdin, J., Shukla, S., Husak, G., Rowland, J., Harrison, L., Hoell, A., und
 632 Michaelsen, J.: The climate hazards infrared precipitation with stations—a new environmental record for monitoring extremes,
 633 *Sci Data*, 2, doi: 10.1038/sdata.2015.66, 2015.
 634 Fustos, I., Abarca-del-Rio, R., Ávila, A., und Orrego, R.: A simple logistic model to understand the occurrence of flood events
 635 into the Biobío River Basin in central Chile, *J Flood Risk Management*, 10, 17–29, doi: 10.1111/jfr3.12131, 2014.
 636 Fustos, I., Remy, D., Abarca-Del-Rio, R., und Muñoz, A.: Slow movements observed within situand remote-sensing techniques
 637 in the central zone of Chile, *International Journal of Remote Sensing*, 38, 7514–7530, doi: 10.1080/01431161.2017.1317944,
 638 2017.
 639 Fustos, I., Abarca-del-Rio, R., Moreno-Yaeger, P., und Somos-Valenzuela, M.: Rainfall-Induced Landslides forecast using
 640 local precipitation and global climate indexes, *Nat Hazards*, 102, 115–131, doi: 10.1007/s11069-020-03913-0, 2020a.

641 Fustos, I., Abarca-del-Río, R., Mardones, M., González, L., und Araya, L. R.: Rainfall-induced landslide identification using
642 numerical modelling: A southern Chile case, *Journal of South American Earth Sciences*, 101, 102587, doi:
643 10.1016/j.jsames.2020.102587, 2020b.

644 Fustos-Toribio, I. J., Morales-Vargas, B., Somos-Valenzuela, M., Moreno-Yaeger, P., Muñoz-Ramirez, R., Rodriguez
645 Araneda, I., und Chen, N.: Debris flow event on Osorno volcano, Chile, during summer 2017: new interpretations for chain
646 processes in the southern Andes, *Nat. Hazards Earth Syst. Sci.*, 21, 3015–3029, doi: 10.5194/nhess-21-3015-2021, 2021.

647 Fustos-Toribio, I., Manque-Roa, N., Vásquez Antipan, D., Hermosilla Sotomayor, M., und Letelier Gonzalez, V.: Rainfall-
648 induced landslide early warning system based on corrected mesoscale numerical models: an application for the southern Andes,
649 *Nat. Hazards Earth Syst. Sci.*, 22, 2169–2183, doi: 10.5194/nhess-22-2169-2022, 2022.

650 Galetto, F., Pritchard, M. E., Hornby, A. J., Gazel, E., and Mahowald, N. M.: Spatial and Temporal Quantification of Subaerial
651 Volcanism From 1980 to 2019: Solid Products, Masses, and Average Eruptive Rates, *Reviews of Geophysics*, 61,
652 <https://doi.org/10.1029/2022rg000783>, 2023.

653 Gariano, S. L. and Guzzetti, F.: Landslides in a changing climate, *Earth-Science Reviews*, 162, 227–252,
654 <https://doi.org/10.1016/j.earscirev.2016.08.011>, 2016.

655 Garreaud, R. D., Boisier, J. P., Rondanelli, R., Montecinos, A., Sepúlveda, H. H., and Veloso-Aguila, D.: The Central Chile
656 Mega Drought (2010–2018): A climate dynamics perspective, *Intl Journal of Climatology*, 40, 421–439,
657 <https://doi.org/10.1002/joc.6219>, 2019.

658 Gregoretti, C.: The initiation of debris flow at high slopes: experimental results, *Journal of Hydraulic Research*, 38, 83–88,
659 <https://doi.org/10.1080/00221680009498343>, 2000.

660 Hooper, A., Segall, P., und Zebker, H.: Persistent scatterer interferometric synthetic aperture radar for crustal deformation
661 analysis, with application to Volcán Alcedo, Galápagos, *J. Geophys. Res.*, 112, doi: 10.1029/2006jb004763, 2007.

662 Hooper, A.: A multi-temporal InSAR method incorporating both persistent scatterer and small baseline approaches,
663 *Geophysical Research Letters*, 35, doi: 10.1029/2008gl034654, 2008.

664 Hooper, A., Bekaert, D., Spaans, K., und Arian, M.: Recent advances in SAR interferometry time series analysis for
665 measuring crustal deformation, *Tectonophysics*, 514–517, 1–13, doi: 10.1016/j.tecto.2011.10.013, 2012.

666 Höser, T.: Analysing the Capabilities and Limitations of InSAR using Sentinel-1 Data for Landslide Detection and Monitoring,
667 Unpublished, doi: 10.13140/RG.2.2.35085.59362, 2018.

668 Huang, A.-B., Lee, J.-T., Ho, Y.-T., Chiu, Y.-F., und Cheng, S.-Y.: Stability monitoring of rainfall-induced deep landslides
669 through pore pressure profile measurements, *Soils and Foundations*, 52, 737–747, doi: 10.1016/j.sandf.2012.07.013, 2012.

670 Hungr, O., Leroueil, S., und Picarelli, L.: The Varnes classification of landslide types, an update, *Landslides*, 11, 167–194,
671 doi: 10.1007/s10346-013-0436-y, 2014.

672 Iverson, R., Logan, M., LaHusen, R., & Berti, M. (2010). The perfect debris flow? aggregated results from 28 large-scale
673 experiments. *Journal of Geophysical Research Atmospheres*, 115(F3). <https://doi.org/10.1029/2009jf001514>

674 Jakob, M. und Lambert, S.: Climate change effects on landslides along the southwest coast of British Columbia,
675 Geomorphology, 107, 275–284, doi: 10.1016/j.geomorph.2008.12.009, 2009.

676 Kang, D., Nam, D., Lee, S., YANG, W., You, K., & Kim, B. (2017). Comparison of impact forces generated by debris flows
677 using numerical analysis models., 1, 195–203. <https://doi.org/10.2495/wrm170191>

678 Kinde, M., Getahun, E., und Jothimani, M.: Geotechnical and slope stability analysis in the landslide-prone area: A case study
679 in Sawla – Laska road sector, Southern Ethiopia, Scientific African, 23, e02071, doi: 10.1016/j.sciaf.2024.e02071, 2024.

680 Korup, O., Seidemann, J., and Mohr, C. H.: Increased landslide activity on forested hillslopes following two recent volcanic
681 eruptions in Chile, Nat. Geosci., 12, 284–289, <https://doi.org/10.1038/s41561-019-0315-9>, 2019.

682 Kuriakose, S. L., van Beek, L. P. H., and van Westen, C. J.: Parameterizing a physically based shallow landslide model in a
683 data poor region, Earth Surf Processes Landf, 34, 867–881, <https://doi.org/10.1002/esp.1794>, 2009.

684 Lee, C.-T.: Landslide trends under extreme climate events, Terr. Atmos. Ocean. Sci., 28, 33–42, doi:
685 10.3319/tao.2016.05.28.01(cca), 2017.

686 Malet, J.-P., Laigle, D., Remaitre, A., and Maquaire, O.: Triggering conditions and mobility of debris flows associated to
687 complex earthflows, Geomorphology, 66, 215–235, <https://doi.org/10.1016/j.geomorph.2004.09.014>, 2005.

688 Maragaño-Carmona, G., Fustos Toribio, I. J., Descote, P.-Y., Robledo, L. F., Villalobos, D., und Gatica, G.: Rainfall-Induced
689 Landslide Assessment under Different Precipitation Thresholds Using Remote Sensing Data: A Central Andes Case, Water,
690 15, 2514, doi: 10.3390/w15142514, 2023.

691 Moreno-Yaeger, P., Singer, B. S., Edwards, B. R., Jicha, B. R., Nachlas, W. O., Kurz, M. D., E. Breunig, R., Fustos-Toribio,
692 I., Antipán, D. V., and Piergrossi, E.: Pleistocene to recent evolution of Mocho-Choshuenco volcano during growth and retreat
693 of the Patagonian Ice Sheet, Geological Society of America Bulletin, 136, 5262–5282, doi: 10.1130/b37514.1, 2024.

694 Muñoz-Sabater, J., Dutra, E., Agustí-Panareda, A., Albergel, C., Arduini, G., Balsamo, G., Boussetta, S., Choulga, M.,
695 Harrigan, S., Hersbach, H., Martens, B., Miralles, D. G., Piles, M., Rodríguez-Fernández, N. J., Zsoter, E., Buontempo, C.,
696 und Thépaut, J.-N.: ERA5-Land: a state-of-the-art global reanalysis dataset for land applications, Earth Syst. Sci. Data, 13,
697 4349–4383, doi: 10.5194/essd-13-4349-2021, 2021.

698 Muratli, J. M., Chase, Z., McManus, J., und Mix, A.: Ice-sheet control of continental erosion in central and southern Chile
699 (36°–41°S) over the last 30,000 years, Quaternary Science Reviews, 29, 3230–3239, doi: 10.1016/j.quascirev.2010.06.037,
700 2010.

701 Ochoa-Cornejo, F., Palma, S., Sepúlveda, S. A., Lara, M., Burgos, K., and Duhart, P.: Rock slides in paraglacial environments
702 in South America: three-dimensional modeling of glacier retreat and landslide inducing the 2017 Santa Lucía disaster in the
703 Chilean Patagonia, Landslides, 22, 1003–1025, <https://doi.org/10.1007/s10346-024-02419-1>, 2024.

704 Ontiveros-Ortega, A., Plaza, I., Calero, J., Moleon, J. A., und Ibañez, J. M.: High variability of interaction energy between
705 volcanic particles: implications for deposit stability, Nat Hazards, 117, 3103–3122, doi: 10.1007/s11069-023-05979-y, 2023.

706 Palazzolo, N., Peres, D. J., Creaco, E., and Cancelliere, A.: Using principal component analysis to incorporate multi-layer soil
707 moisture information in hydrometeorological thresholds for landslide prediction: an investigation based on ERA5-Land
708 reanalysis data, *Nat. Hazards Earth Syst. Sci.*, 23, 279–291, doi: 10.5194/nhess-23-279-2023, 2023.

709 Pavlova, I., Jomelli, V., Brunstein, D., Grancher, D., Martin, E., and Déqué, M.: Debris flow activity related to recent climate
710 conditions in the French Alps: A regional investigation, *Geomorphology*, 219, 248–259,
711 <https://doi.org/10.1016/j.geomorph.2014.04.025>, 2014. Pereyra, F. X. und Bouza, P.: Soils from the Patagonian Region, *World*
712 *Soils Book Series*, 101–121, doi: 10.1007/978-3-319-76853-3_7, 2018.

713 Porchet M. and Laferrere H. Détermination des caractéristiques hydrodynamiques des sols en place. Mémoire et notes
714 techniques. *Ann. Du Ministère de l’Agriculture, Tech. Rep. 2 (64)*, 1935.

715 Rawson, H., Naranjo, J. A., Smith, V. C., Fontijn, K., Pyle, D. M., Mather, T. A., und Moreno, H.: The frequency and
716 magnitude of post-glacial explosive eruptions at Volcán Mocho-Choshuenco, southern Chile, *Journal of Volcanology and*
717 *Geothermal Research*, 299, 103–129, doi: 10.1016/j.jvolgeores.2015.04.003, 2015.

718 Rodriguez, C., Perez, Y., Hugo Roa, Clayton, J., Antinao, J. L., & Duhart, P., Martin, M.: Geología del área de Panguipulli-
719 Riñihue, Región de Los Lagos. Servicio Nacional de Geología y Minería, Mapas Geológicos No: 10. 1999.

720 Salazar, Á., Thatcher, M., Goubanova, K., Bernal, P., Gutiérrez, J., and Squeo, F.: CMIP6 precipitation and temperature
721 projections for Chile, *Clim Dyn*, 62, 2475–2498, <https://doi.org/10.1007/s00382-023-07034-9>, 2023.

722 SANHUEZA, C., PALMA, J., VALENZUELA, P., ARANEDA, O., und CALDERÓN, K.: Evaluación del comportamiento
723 geotécnico de suelos volcánicos chilenos para su uso como material de filtro en la depuración de aguas residuales domésticas,
724 *Revista de la Construcción*, 10, 66–81, doi: 10.4067/s0718-915x2011000200007, 2011.

725 Savi, S., Schildgen, T. F., Tofelde, S., Wittmann, H., Scherler, D., Mey, J., Alonso, R. N., and Strecker, M. R.: Climatic
726 controls on debris-flow activity and sediment aggradation: The Del Medio fan, NW Argentina, *JGR Earth Surface*, 121, 2424–
727 2445, <https://doi.org/10.1002/2016jf003912>, 2016.

728 Sepúlveda, S. A. und Padilla, C.: Rain-induced debris and mudflow triggering factors assessment in the Santiago cordillera
729 foothills, Central Chile, *Nat Hazards*, 47, 201–215, doi: 10.1007/s11069-007-9210-6, 2008.

730 Sepúlveda, S. A.: LARGE VOLUME LANDSLIDES IN THE CENTRAL ANDES OF CHILE AND ARGENTINA (32°-
731 34°S) AND RELATED HAZARDS, *Italian Journal of Engineering Geology and Environment*, 287–294, doi:
732 10.4408/IJEGE.2013-06.B-26, 2013.

733 Sepúlveda, S. A., Moreiras, S. M., Lara, M., und Alfaro, A.: Debris flows in the Andean ranges of central Chile and Argentina
734 triggered by 2013 summer storms: characteristics and consequences, *Landslides*, 12, 115–133, doi: 10.1007/s10346-014-0539-
735 0, 2014.

736 Sepúlveda, S. A. und Petley, D. N.: Regional trends and controlling factors of fatal landslides in Latin America and the
737 Caribbean, *Nat. Hazards Earth Syst. Sci.*, 15, 1821–1833, doi: 10.5194/nhess-15-1821-2015, 2015.

738 Singer, B. S., Moreno-Yaeger, P., Townsend, M., Huber, C., Cuzzone, J., Edwards, B. R., Romero, M., Orellana-Salazar, Y.,
739 Marcott, S. A., Breunig, R. E., Ferrier, K. L., Scholz, K., Coonin, A. N., Alloway, B. V., Tremblay, M. M., Stevens, S., Fustos-

740 Toribio, I., Moreno, P. I., Vera, F., und Amigo, Á.: New perspectives on ice forcing in continental arc magma plumbing
 741 systems, *Journal of Volcanology and Geothermal Research*, 455, 108187, doi: 10.1016/j.jvolgeores.2024.108187, 2024.

742 Singh, K. und Kumar, V.: Rainfall Thresholds Triggering Landslides: A Review, *Lecture Notes in Civil Engineering*, 455–
 743 464, doi: 10.1007/978-3-030-51354-2_42, 2020.

744 Schmidt, K. M., Roering, J. J., Stock, J. D., Dietrich, W. E., Montgomery, D. R., and Schaub, T.: The variability of root
 745 cohesion as an influence on shallow landslide susceptibility in the Oregon Coast Range, *Can. Geotech. J.*, 38, 995–1024,
 746 <https://doi.org/10.1139/t01-031>, 2001.

747 Somos-Valenzuela, M. A., Oyarzún-Ulloa, J. E., Fustos-Toribio, I. J., Garrido-Urzua, N., und Chen, N.: The mudflow disaster
 748 at Villa Santa Lucía in Chilean Patagonia: understandings and insights derived from numerical simulation and postevent field
 749 surveys, *Nat. Hazards Earth Syst. Sci.*, 20, 2319–2333, doi: 10.5194/nhess-20-2319-2020, 2020.

750 Stern, C. R.: Active Andean volcanism: its geologic and tectonic setting, *Rev. geol. Chile*, 31, doi: 10.4067/s0716-
 751 02082004000200001, 2004.

752 Stern, C. R.: Holocene tephrochronology record of large explosive eruptions in the southernmost Patagonian Andes, *Bull*
 753 *Volcanol*, 70, 435–454, doi: 10.1007/s00445-007-0148-z, 2007.

754 Stoffel, M., Mendlik, T., Schneuwly-Bollschweiler, M., and Gobiet, A.: Possible impacts of climate change on debris-flow
 755 activity in the Swiss Alps, *Climatic Change*, 122, 141–155, <https://doi.org/10.1007/s10584-013-0993-z>, 2013.

756 Talebi, A., Uijlenhoet, R., and Troch, P. A.: Soil moisture storage and hillslope stability, *Nat. Hazards Earth Syst. Sci.*, 7, 523–
 757 534, <https://doi.org/10.5194/nhess-7-523-2007>, 2007.

758 Thompson, P. I. J., Dugmore, A. J., Newton, A. J., Cutler, N. A., und Streeter, R. T.: The influence of burial rate on variability
 759 in tephra thickness and grain size distribution in Iceland, *CATENA*, 225, 107025, doi: 10.1016/j.catena.2023.107025, 2023.

760 Valenzuela, R. A. and Garreaud, R. D.: Extreme Daily Rainfall in Central-Southern Chile and Its Relationship with Low-Level
 761 Horizontal Water Vapor Fluxes, *Journal of Hydrometeorology*, 20, 1829–1850, doi: 10.1175/jhm-d-19-0036.1, 2019.

762 Vásquez-Antipán, D., Fustos-Toribio, I., Riffo-López, J., Cortez-Díaz, A., Bravo, Á., und Moreno-Yaeger, P.: Landslide
 763 processes related to recurrent explosive eruptions in the Southern Andes of Chile (39° S), *Journal of South American Earth*
 764 *Sciences*, 157, 105469, <https://doi.org/10.1016/j.jsames.2025.105469>, 2025.

765 Vega, J. A. and Hidalgo, C. A.: Quantitative risk assessment of landslides triggered by earthquakes and rainfall based on direct
 766 costs of urban buildings, *Geomorphology*, 273, 217–235, doi: 10.1016/j.geomorph.2016.07.032, 2016.

767 Walding, N., Williams, R., Rowley, P., und Dowey, N.: Cohesional behaviours in pyroclastic material and the implications for
 768 deposit architecture, *Bull Volcanol*, 85, doi: 10.1007/s00445-023-01682-9, 2023.

769 Wang, J. and Yu, Y.: Debris Flow Hazard Zoning Based on Numerical Simulation in the Wenchuan Earthquake Meizoseismal
 770 Areas, *Proceedings of the 5th International Conference on Civil Engineering and Transportation 2015*,
 771 <https://doi.org/10.2991/iccet-15.2015.81>, 2015.

772 Wang, H., Cui, P., Li, Y., Tang, J., Wei, R., Yang, A., Zhou, L., Bazai, N. A., und Zhang, G.: Rock and ice avalanche-generated
773 catastrophic debris flow at Chamoli, 7 February 2021: New insights from the geomorphic perspective, *Geomorphology*, 452,
774 109110, doi: 10.1016/j.geomorph.2024.109110, 2024.

775 Xie, M., Zhao, W., Ju, N., He, C., Huang, H., und Cui, Q.: Landslide evolution assessment based on InSAR and real-time
776 monitoring of a large, reactivated landslide, Wenchuan, China, *Engineering Geology*, 277, 105781, doi:
777 10.1016/j.enggeo.2020.105781, 2020.

778 Yi, Z., Xingmin, M., Allesandro, N., Tom, D., Guan, C., Colm, J., Yuanxi, L., und Xiaojun, S.: Characterization of pre-failure
779 deformation and evolution of a large earthflow using InSAR monitoring and optical image interpretation, *Landslides*, 19, 35–
780 50, doi: 10.1007/s10346-021-01744-z, 2021.

781 Yu, C., Li, Z., Penna, N. T., und Crippa, P.: Generic Atmospheric Correction Model for Interferometric Synthetic Aperture
782 Radar Observations, *JGR Solid Earth*, 123, 9202–9222, doi: 10.1029/2017jb015305, 2018.