

1 **Tsunamigenic potential of unstable masses in the Gulf of Pozzuoli,**

2 **Campi Flegrei, Italy**

3

4 Filippo Zaniboni¹, Luigi Sabino², Cesare Angeli¹, Martina Zanetti¹ and Alberto Armigliato¹

5 ¹ Department of Physics and Astronomy “A. Righi”, Alma Mater Studiorum - University of Bologna, Bologna, 40127, Italy

6 ² Department of Mathematics and Physics, University of Campania “L. Vanvitelli”, Caserta, 81100, Italy

7 *Correspondence to:* Filippo Zaniboni (filippo.zaniboni@unibo.it)

8 **Abstract.** Campi Flegrei, one of the most monitored and studied volcanic areas in the world, has recently attracted significant
9 attention due to the reactivation of its peculiar activity, consisting ~~of~~ small earthquakes, geothermal phenomena and slow
10 subsidence/rapid uplift cycles, known as bradyseism. While much of the research and of the attention focuses on ~~potential~~
11 ~~eruptions or other volcanic-related activities~~, the potential hazard posed by gravitational instabilities has received little
12 consideration. The interaction of the destabilized masses with water can trigger tsunamis, potentially affecting the whole
13 coastline of the Gulf of Pozzuoli, which lies above the Campi Flegrei caldera. Moving from the limited available
14 geomorphological studies of the area, a set of four landslide-tsunami scenarios (one subaerial and three submarine sources)
15 are ~~reconstructed. These are simulated through a sequence of numerical codes, accounting for all the phases of the tsunami~~
16 process, providing insights into the distribution of tsunami energy and identifying the most affected coastal stretches.
17 Additionally, the study explores the influence of dispersion effects in the tsunami propagation and the occurrence of resonance
18 effects in some minor inlets of the Gulf, emphasizing the importance of accounting for complex and non-linear coastal
19 processes when treating ~~landslide-generated tsunamis~~.

20 **1 Introduction**

21 The recent strengthening of the Campi Flegrei activity has raised many concerns about the potential impacts of volcanic-related
22 manifestations on local population and infrastructures. The caldera, located in correspondence of an extremely densely
23 inhabited area in the surroundings of Naples (South Italy), is one of the most active and dangerous in the world. The study of
24 the hazard correlated to the Campi Flegrei activity, concerning the tephra dispersal in case of eruption, has been already treated
25 in the scientific literature (Selva et al., 2018). On this basis, a National Civil Protection plan has been realized in case of
26 emergency (<https://mappe.protezionecivile.gov.it/en/risks-maps-and-dashboards/national-planning-phlegraean-fields/>). Other
27 considered potential hazards linked to the eruption of the Campi Flegrei caldera are pyroclastic flows (Neri et al., 2015),
28 phreatic explosions (Mayer et al., 2016) and mud flows generated by the interaction between the ash ejected from the volcano
29 and the rain (Isaia et al., 2021). Earthquakes are frequent, though characterized by low magnitude. As an example, according

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

Formattato: Allineato a sinistra

ha eliminato: into

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha eliminato: volcanic manifestations

ha eliminato: and one subaerial

ha eliminato: here

ha eliminato: with

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

35 to the [September 2025 INGV - Osservatorio Vesuviano bulletin](https://www.ov.ingv.it/index.php/flegrei-stato-attuale) (<https://www.ov.ingv.it/index.php/flegrei-stato-attuale>), 423
36 events have been recorded during the month. Almost 88% of these had magnitude lower than 1, four exceeded $M_d = 2.0$ and
37 the maximum was 4.0.

38 The interaction of volcanic activity with the sea and the consequent generation of tsunamis received some attention as well:
39 the work by Paris et al. (2019) tested the effects of submarine volcanic explosions in the Gulf of Pozzuoli by means of
40 numerical scenarios and a Probabilistic Tsunami Hazard Analysis (PTHA) methodology. Grezio et al. (2020) attempted a
41 comprehensive approach including earthquakes, submarine landslides and volcanic explosions as potential sources for tsunami
42 generation affecting the Gulf of Naples, implementing them into a PTHA providing hazard curves for different localities along
43 the coasts.

44 Specifically, gravitational collapses are one of the least considered potential sources of natural hazards in the area. Volcanic
45 activity at Campi Flegrei can induce instability both in the short term, through seismic shaking, and in the long term, due to
46 slope steepening associated with the caldera uplift. The scenarios considered in Grezio et al. (2020) were purely synthetic,
47 without evaluation of geological and morphological evidence, of the sliding dynamics and of possible hydrodynamic effects
48 during the tsunami propagation. In occasion of the 30 June 2025 earthquake (magnitude 4.6), a rock mass detached from the
49 coastal ridge of Punta Pennata, in Bacoli, at the western end of the Gulf of Pozzuoli, provoking some local sea level oscillations
50 (<https://en.cronachedellacampania.it/2025/06/terremoto-crollata-una-parete-di-punta-pennata-a-bacoli/>), and putting in the
51 spotlight such type of phenomenon.

52 This work aims to fill these gaps, by assessing the tsunamigenic potential of local submarine and subaerial landslides,
53 evaluating the tsunami distribution patterns and the respective impact on the coasts, at the scale of the Gulf of Pozzuoli basin.
54 The masses generating the waves are selected based on a worst-case credible approach (see e.g. Tonini et al., 2011), which
55 relies upon the present morphology and upon the existing knowledge of the mass transport processes in the area. The tsunami
56 hazard is then assessed through a set of numerical codes, already tested and applied in many other landslide-tsunami cases.
57 This investigation has been preliminarily performed and reported in Sabino (2024): here it is extended including the role of
58 dispersion in the tsunami propagation, since it can have significant influence in the impact of the waves on the coasts.

59 1.1 Campi Flegrei volcano


60 Campi Flegrei is a complex volcanic system with a history spanning at least the last 78,000 years (Perrotta et al., 2006),
61 characterized by intense unrest episodes involving ground deformation and seismicity (de Natale et al., 2006), together with
62 explosive eruptions and variable vent locations (Bevilacqua et al., 2015). Seismic studies have revealed a large magmatic sill
63 at depth, potentially connected to the surface through deep fractures (Zollo et al., 2008). The heart of Campi Flegrei is the vast
64 caldera, an almost circular structure with a diameter of about 10 km (marked by the red dashed line in Figure 1) involving the
65 western districts of Naples in its subaerial expression, with a total potentially affected population of more than 360 thousand,
66 living in the cities of Pozzuoli, Bagnoli and Bacoli. The submarine part of the caldera is covered by the Gulf of Pozzuoli, a
67 small, shallow water sub-basin of the wider Gulf of Naples. The landscape of this area has been shaped by several volcanic

ha eliminato: December
ha eliminato: 4
ha eliminato: 273
ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman), Non Evidenziato
ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)
ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman), Non Evidenziato
ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)
ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)
ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)
ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman), Non Evidenziato
ha eliminato: Over 93%
ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman), Non Evidenziato
ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)
ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)
ha eliminato: 3.
ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman), Non Evidenziato
ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)
ha eliminato: In the specific
ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)
ha eliminato: ology
ha eliminato: to bridge
ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)
ha eliminato: ; Zaniboni and Armigliato, 2025
ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)
ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

77 events: the caldera was formed by a catastrophic volcanic explosion that occurred approximately 39,000 years ago. This event,
78 an eruption of 100-200 km³ of rock called "Ignimbrite Campana" (Rosi et al., 1983; Perrotta et al. 2006), shaped the region's
79 topography, creating a unique landscape characterized by hills, fumaroles, and hot springs. At the centre of the caldera rises
80 the Solfatara crater, a focal point for the volcanic and geothermal activity in the area. Historical reports from Roman times
81 reveal a general trend of slow subsidence (rate of about 1-2 cm/yr), alternated with occasional episodes of faster uplift (Di Vito
82 et al., 2016; De Vivo et al., 2020), a peculiar behaviour that took the name of "bradyseism". Soil subsidence can lead to coastal
83 flooding, as testified by the submerged park of Baia, where Roman houses and constructions are still visible now at 6-8 m
84 depth. The last significant eruption of Campi Flegrei caldera occurred in 1538, with 0.03 km³ of erupted products that gave
85 rise to the hill of Monte Nuovo in one single night (De Vivo et al., 2001). After that, the floor of the caldera underwent slow
86 and regular subsidence. In more recent times, some episodes of uplift interrupted this phase: 74 cm in 1950-52, 159 cm in
87 1970-72 and 178 cm in 1982-84 (Del Gaudio et al., 2010), resulting in a maximum rise of about 3.5 meters of the ground in
88 the city of Pozzuoli, with shallow micro-seismicity recorded in response to fluid movement episodes.

89 At present, bradyseism, which resumed in mid-2023, is still ongoing. The GNSS network has measured, since January 2024,
90 an uplift of more than 33 cm in some stations (see <https://www.ov.ingv.it/index.php/monitoraggio-e-infrastrutture/bollettini-web>). Seismic activity has been intensifying during this period, although most of the events are characterized by low
92 magnitudes (about 90% of the events had magnitudes less than 1.0, see <https://www.ov.ingv.it/index.php/monitoraggio-e-infrastrutture/bollettini-tutti/bollett-mensili-cf>), with maximum depths around 4 km, predominantly concentrated within the
94 first 2 km (Danesi et al., 2024).

ha eliminato: has
ha eliminato: since half of 2023
ha eliminato: from
ha eliminato: and
ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)
ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)
ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)
Commentato [AA1]: uplift?
ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)
ha eliminato: in such
ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)
ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)
ha eliminato: of
ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)




101
102 **Figure 1.** Morphological map of the Gulf of Naples (South Italy). The red-dashed circle delimits approximately the Campi Flegrei
103 caldera, involving a subaerial part (west of Naples) and a submarine portion (Gulf of Pozzuoli).

104 **2 Data and methods**

105 The investigation on the tsunami hazard in the ~~Gulf of Pozzuoli~~ associated to the Campi Flegrei activity, triggering potential
106 instabilities interacting with water, is here performed through numerical methods, which in turn implement approaches that
107 are described in this section. An initial necessary premise concerns the importance of considering the dispersive effects in
108 landslide-tsunami simulations, a task that is usually underestimated but whose effects could be relevant in the analysis of the
109 tsunami impact on the coast.

110 **2.1 Landslide-tsunamis and dispersive effects**

111 Tsunamis are oscillations induced by a perturbation of the equilibrium state involving the whole water body, which can be
112 produced by sudden changes in the sea bottom (earthquakes, submarine landslides, underwater volcanic explosions) or upon
113 the sea surface (subaerial landslides, atmospheric perturbations, cosmogenic tsunamis).



101
102 **ha eliminato:** 1
103 **ha formattato:** Tipo di carattere: (Predefinito) +Titoli (Times New Roman)
104 **ha formattato:** Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

105 **ha eliminato:** Starting from the second half of 2023, the
106 bradyseism has resumed, with seismic activity intensifying, although
107 most of the events were characterized by low magnitudes (about 90%
108 of the events had magnitudes less than 1.0, see
109 <https://www.ov.ingv.it/index.php/monitoraggio-e-infrastrutture/bollettini-tutti/bollett-mensili-cf>), with maximum
110 depths of around 4 km, predominantly concentrated within the first 2
111 km (Danesi et al., 2024). The sequence culminated with the seismic
112 events on September 27th, 2023 ($M_d = 4.2$) and October 2nd, 2023
113 ($M_d = 4.0$), located respectively in the area between Bagnoli and
Pozzuoli and in the Pisciarelli-Solfatara area. 1

101 **ha eliminato:** g

102 **ha eliminato:**

130 In general, in hydrodynamics, each wave is subject to the dispersion relation, linking the phase velocity c to the wave number
 131 k (or to its inverse, the wavelength λ): the smaller is the second (i.e., the bigger is the wavelength), the higher is the first. In
 132 short, longer oscillations are faster than shorter ones (for a general overview about this topic see for example Saito, 2019).
 133 A tsunami can be considered mostly as a “packet” of waves, meaning that it is not a monochromatic wave, but it results from
 134 the superposition of different components, each characterized by a specific wavelength, then with its own velocity. When λ is
 135 much bigger than the water depth h of the involved basin, the hydrodynamic equations can be significantly simplified by
 136 means of the shallow-water (SW) approach, in which tsunamis are treated as “long-waves”. This is generally valid for waves
 137 generated by earthquakes, since the source has dimensions that is usually much bigger than the typical depth of the sea. When
 138 the tsunami trigger is provided by other phenomena, such as landslides, SW is known to neglect important hydrodynamic
 139 effects, such as dispersion, that can affect significantly the wave propagation and impact on the coast. It is generally agreed
 140 that SW is considered proper when the ratio λ/h is bigger than 20, while for lower values (i.e., for “shorter” waves) a more
 141 sophisticated and higher-order version of the hydrodynamic equations should be considered, as for example the Boussinesq
 142 approximation (hereafter accounted for as non-hydrostatic approach, NH).
 143 A quantification of the dispersive effects on the tsunami propagation was attempted in Glimsdal et al. (2013). A rearrangement
 144 of the considerations found in that work leads to the estimation of the ratio between distance and initial signal wavelength for
 145 which the dispersion becomes relevant and cannot be neglected anymore. Calling this (non-dimensional) dispersion distance
 146 as D , the expression is:

$$147 D = 0.025 \cdot \left(\frac{\lambda}{h}\right)^2 \quad (1)$$

148 For “long waves”, then, $\lambda/h \approx 20$ and $D \approx 10$; dispersive effects manifest at least at a distance ten times the initial
 149 wavelength. Table 1 reports some typical values for wavelengths and water basin depths found respectively for: earthquake-
 150 tsunamis (first line), for which generally SW approach is applicable; submarine landslide tsunamis (second line) for which SW
 151 is valid only on limited domains; subaerial landslide tsunamis (third line), which can be considered in general as deep-water
 152 (DW) waves, for which the NH approach is more suitable.

153 In general, we can see that dispersion affects every tsunami, also the “longer” ones, but it manifests only on very long distances,
 154 where probably the perturbation itself has already been damped by other propagation effects. Neglecting dispersion means that
 155 the different components of the tsunami travel all together and impact the coast at the same time; conversely, when considering
 156 it, the longer components travel much faster than the shorter ones. This provokes an “unpacking” of the tsunami, that is much
 157 more evident and effective with increasing distance from the source: in general, then, neglecting dispersion causes an
 158 overestimation of the tsunami effects.

159
 160 **Table 1. Examples of waves and of computing of respective dispersion distance. λ : initial wavelength; h : water depth; D : dispersion**
 161 **distance, computed by Eq. (1); d : distance for which dispersion effects become predominant ($d = D \cdot \lambda$).**

λ (km)	h (m)	λ/h	Wave type	D	d (km)
----------------	---------	-------------	-----------	-----	----------

100	4000	25	SW	15.6	1562.5
5	500	10	Weakly SW	2.5	12.5
1	200	5	DW	0.625	0.625

163 **2.2 Overview on numerical techniques**

164 Landslide-generated tsunamis are highly complex processes. Assessing their impact on the coast through numerical
 165 simulations requires numerous approximations and assumptions. The approach adopted here divides the entire process into
 166 distinct phases: i) the landslide motion, ii) the transfer of energy from the mass to the water, iii) tsunami propagation, and iv)
 167 the wave's impact on the coast. Each phase is treated independently, and back-interactions are not considered; for example,
 168 the mass-water interaction, where the wave's propagation can alter the water column and influence the dynamics of the
 169 landslide on the seafloor up to 20% (Harbitz et al., 2006), is neglected in this study, since those effects do not affect
 170 considerably the tsunami propagation. The phases are addressed sequentially, with the output of one stage serving as the input
 171 for the next.

172 *Landslide motion.* The movement of the mass along the seafloor triggers the tsunami. Unlike earthquake-generated tsunamis,
 173 where the source process can be considered instantaneous relative to the wave's propagation (at least as a first approximation),
 174 landslide-tsunamis exhibit a generation phase that evolves in comparable time with the ensuing waves. Thus, it is essential to
 175 accurately describe the mass dynamics and the evolution of shape changes. This is achieved using the numerical code UBO-
 176 BLOCK (see detailed description in Tinti et al., 1997), which divides the sliding mass into interacting portions. The equation
 177 governing the centre of mass (CoM) motion is determined by the forces acting on the mass: gravity, buoyancy, basal friction,
 178 surface drag, and an internal interaction force accounting for energy dissipation due to deformation. The last element is crucial,
 179 since it accounts for shape changes of the mass during its motion, a factor that influences also the wave generation and that
 180 can't be neglected; it is found, in fact, that rigid block approximation for landslide collapses produces an overestimation of the
 181 generated tsunamis (see for example the review in Yavari-Ramshe and Ataje-Ashtiani, 2016). The resulting time-series of
 182 geometric and dynamic quantities describing the landslide motion is then generated. Applications of UBO-BLOCK to
 183 landslide-tsunami events can be retrieved in Tryantafillou et al. (2020), Zaniboni et al. (2021), Gallotti et al. (2021), Gasperini
 184 et al. (2022), Gallotti et al. (2023), Zaniboni et al. (2024) and in the references therein.

185 *Tsunamigenic impulse.* The next step involves assessing the perturbation to the water column caused by the sliding motion
 186 along the seafloor. This disturbance provides the dynamic forcing term for the wave propagation equations, which are not
 187 instantaneous but evolve over time. The change in the seafloor due to the landslide is interpolated onto the tsunami grid nodes,
 188 with a filter applied to local sea depths that suppresses high frequencies. These tasks are handled by the intermediate code
 189 UBO-TSUIMP (details in Tinti et al., 2006).

190 *Tsunami propagation and coastal impact.* The final two phases are simulated using the JAGURS code, which solves the fluid
 191 dynamics equations using finite difference methods. Tsunami propagation can be modelled using either the shallow-water

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha eliminato: with

ha eliminato: no

ha eliminato: (

ha eliminato:)

ha eliminato: involve a finite time

ha eliminato:

ha eliminato: for the

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

199 (SW) equations or a more sophisticated approach that accounts for vertical variations in hydrodynamic quantities,
200 implementing for example the Boussinesq model (NH). The non-hydrostatic method allows the code to capture dispersion
201 effects, which are crucial for landslide-generated tsunamis, as described in Section 1.1. JAGURS supports simulations over
202 computational grids with varying resolutions (using a nested grid approach) and can model coastal inundation. It is also
203 optimized for parallel computing with MPI and OpenMPI coding, enabling the simulation of dispersive effects over large
204 computational domains (see Baba et al., 2015, for further description). While JAGURS is widely used for earthquake-generated
205 tsunamis (Ren et al., 2021; Ehara et al., 2023), for landslide-generated events it requires some adaptations, due to the nature
206 of the phenomenon itself. As previously mentioned, the tsunamigenic impulse for landslide-tsunamis is not instantaneous:
207 then, it must be provided to the code as a sequence of single impulses, one for each time step describing the landslide motion
208 along the sea bottom and producing the perturbation.

209 2.3 Computational grid

210 The numerical codes applied here require as input a regularly spaced computational grid, whose definition and assembling
211 needs a compromise among different aspects: the detailed description of the morphology, mainly for the coastal areas, requires
212 a huge number of nodes, that on the contrary needs to be limited basing on the computational resources available. Additionally,
213 landslide-tsunamis are known to affect limited domains, due to the dispersive effects characterizing their oscillations that
214 produce a rapid damping of their amplitude. In the specific case studied here, the morphology of the seabed suggests that the
215 mass instabilities induced by the Campi Flegrei activity have typically small volumes, producing waves that will presumably
216 travel modest distances, with reduced consequences on the coasts. Under all these considerations, the selected tsunami
217 computational domain covers the Pozzuoli Gulf for an area of 12 km x 11 km approximately, with a spatial step of 20 m and
218 a total number of nodes of about 340 thousand. Raw data have been retrieved from the database MaGIC for the coastline and
219 the bathymetry (DPC), and from the database Tinitaly (Tarquini et al., 2023) as concerns the topography.

220 The morphology, that can be inferred from the general map in Figure 1 and with more detail in Figure 2, shows some
221 remarkable features as concerns landslide-tsunami triggering and propagation: i) the underwater slopes are in general quite
222 gentle, inhibiting high initial acceleration of the sliding masses, one of the most important factors in tsunami genesis (e.g.,
223 Lovholt et al., 2015). Only the areas close to the Gulf's mouth opposite sides, i.e., Capo Miseno on the west and Nisida Bank
224 on the east (see Figure 2 for toponyms), show steep submarine gradients. ii) the coastal slope is generally flat, favoring water
225 ingressions and tsunami penetration; only the two abovementioned areas (Capo Miseno and Nisida) are characterized by steep
226 coastal profiles, that can also produce rapidly moving collapses into the water. iii) the sea is rather shallow within the Gulf,
227 with maximum depth reaching about 100 m, while it deepens eastward and southward. This will have consequences on the
228 wave propagation in the basin, since, as we have seen, the sea depth deeply influences the tsunami behaviour during the
229 propagation. iv) finally, but not less importantly, at this level of detail it is possible to represent the main piers and harbour
230 structures (well visible at Pozzuoli and Bagnoli, Figure 2), which influence the wave propagation with local effects, such as
231 for example multiple reflections and resonance.

ha eliminato: (see Table 2)

ha eliminato: represented in

ha eliminato: 2


ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

235 **2.4 Mass instabilities in the Gulf of Pozzuoli and landslide scenarios.** The evaluation of the landslide-tsunami hazard in the
236 Gulf of Pozzuoli requires, as a first step, to identify the sources, i.e., masses that have the potential to generate waves, both in
237 submarine and in the subaerial environments. This investigation is here performed through a scenario approach, meaning that
238 those collapses are hypothesized starting from geological and morphological evidence, studying their signatures (scars, slopes,
239 deposits) and the ensuing tsunamis evaluated through the numerical routine previously illustrated. The assumption is that such
240 scenarios represent the range of possible and credible worst events occurring in the area, following the approach called Worst-
241 case Credible Tsunami Hazard Assessment (WCTHA, see Tonini et al., 2011 for description and application), which is suitable
242 to phenomena like landslide-tsunamis, where neither recurrence time are defined nor extensive catalogues of events are
243 available, contrary to the most adopted approach in tsunami science (PTHA, see e.g. Behrens et al., 2021). In the specific case
244 of the Gulf of Pozzuoli, we will see that, lacking detailed bathymetric surveys of the seabed specifically devoted to map and
245 describe such occurrences, the WCTHA is not fully applicable: evidence of past collapses is scarce and a general pattern of
246 the potential mass movements is not defined. Moreover, in a context like the Campi Flegrei, bradiseismic activity, morphology
247 and trigger can change rapidly in time, defining new threshold for “worst-case credible” scenarios. The scenarios that have
248 been individuated, then, must not be considered as detailed reconstruction of past events, but as representative of categories of
249 collapse events that potentially occurred, whose potential hazard is here evaluated through numerical simulations. In the
250 following, as a first step, the search for mass collapse traces in the Gulf of Pozzuoli within the scientific literature is reported.
251 Recent magnetic surveys in the larger Gulf of Naples evidenced the presence of consistent deposits on the seabed, testifying
252 an intense mass transport activity in the whole area (de Ritis et al., 2024), that could have probably generated tsunamis affecting
253 the area of interest of this study. The focus, though, is here centred on local sources, i.e., potential landslides triggered by the
254 seismic shaking related to the Campi Flegrei caldera, then along the slopes (submarine and subaerial) in the Gulf of Pozzuoli
255 area. As already mentioned, this is a small, shallow water basin, whose detailed stratigraphic and morphological features have
256 been revealed by high-resolution seismic and bathymetric surveys (Aiello et al., 2012; Somma et al., 2016). The area is
257 characterized by numerous seismic units, both volcanic and sedimentary, and by a complex tectonic setting related to the
258 intense Campi Flegrei volcanic activity. The northern part of the basin shows an inner shelf with maximum depth of 50 m,
259 deepening to 100 m with gentle slopes and delimited by a belt of submarine volcanic edifices: starting from East, they are
260 called Miseno Bank, Pentapalummo Bank and Nisida Bank (see Figure 2). The morphological investigations have revealed
261 the presence of deposits in the deeper and flat part of the Gulf, especially around Miseno Bank and in the central part of the
262 basin (areas in light brown, Figure 2; Aiello et al., 2012). The seismic sections reported in the same publication evidence the
263 presence of buried deposits with peculiar characteristics, which have been characterized and denoted as paleo-landslides.
264 Figure 2 reports the profiles along which they have been found: one lies along the transect L69_07 (in red), extending in the
265 E-W direction and showing a potential deposit close to Nisida; the second refers to the line L74_07 (in magenta), running in
266 the N-S direction from Pozzuoli. The morphological evidence close to Miseno Bank and the two buried deposits found in the
267 seismic profiles will be taken as the basis for the submarine landslide scenarios adopted for this investigation.

ha eliminato: ¶
ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman), Grassetto
ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)
ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)
ha eliminato: the
ha eliminato: , Zaniboni and Armigliato, 2025

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)
ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)
ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)
ha eliminato: the collapse pieces of
ha eliminato: are few
ha eliminato: the
ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)
ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)
ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)
ha eliminato: ;
ha eliminato: m
ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)
ha eliminato: have
ha eliminato: to
ha eliminato: while
ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)
ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)
ha eliminato: s
ha eliminato: which
ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)
ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)
ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)



282

283 **Figure 2.** Morphological map of the Gulf of Pozzuoli. Black labels denote the main coastal toponyms, blue ones the submarine
 284 volcanic edifices. The brown contours mark the recognized depositional areas; the red and magenta lines report the seismic line
 285 called L69_07 and L74_07 respectively. Such features are retrieved from Aiello et al. (2012).

286

287 Regarding subaerial sliding, a comprehensive geodatabase (CAFLAG) documents 2302 landslide events from 1828 to 2017,
 288 most of which consists of rock falls affecting volcanic slopes and rainfall-induced slides in pyroclastic deposits (Esposito and
 289 Matano, 2023). Landslide hazards affect over 15% of the subaerial Campi Flegrei area, with varying risk levels among towns
 290 (Calcaterra and di Martire, 2022). The events of interest here are the ones potentially interacting with water: the candidates are
 291 then restricted to collapses along the coastline that are not simple rockfalls (surely generating high waves but only locally,
 292 dissipating quickly, as in the case of June 30th, 2025, previously cited), but more complex body collapses involving a coherent
 293 mass impacting the water and moving along the seafloor.
 294 Based on all these considerations, a set of four landslide scenarios has been arranged for the numerical simulations. Table 2
 295 summarizes their geomorphological characteristics (volume, area, maximum thickness and initial elevation), while Figure 3
 296 reports the position of the initial masses for each case. Three of them are submarine and cover different positions over the Gulf
 297 of Pozzuoli, while the fourth is subaerial.
 298 Scenario 1 (blue contour in Figure 3) is located just south of Capo Miseno, at the western end of the Gulf, and has been
 299 reconstructed starting from the deposit shown in brown in Figure 2. The hypothesis is that the mass detached from the seafloor
 300 depression between Capo Miseno and the Miseno Bank, with the constraint to obtain a volume comparable to the observed
 301 deposit at the toe of the slope: since an accurate estimate of this does not exist, a conservative approach has been adopted

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha eliminato: As concerns

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha eliminato: such

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman), Apice

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

304 basing on the hypothesized boundary and on a reasonable thickness distribution, basing on the seafloor morphology. The
 305 volume obtained for this scenario is of almost 3 million m³.

306 Scenario 2 (in magenta, Figure 3) lies in the central part of the Gulf and has been built based on the buried paleo-landslide
 307 recognized in the seismic profile just south of Pozzuoli (magenta line, Figure 2). Its morphology is quite arbitrary, since scarce
 308 information exist on this hypothesis. Adopting again a conservative approach, this has been taken similar to Scenario 1, with
 309 volume and initial area slightly smaller, though its shape is more elongated in the sliding direction.

310 Scenario 3 (green boundary, Figure 3) is placed in the eastern part of the basin, and recalls a buried paleo-landslide as well,
 311 relative to another seismic transect (E-W from Nisida, see Figure 2, red line). This scenario shows a slightly larger volume
 312 (around 4 million m³), a slightly larger initial thickness and moves from shallower water: all these elements suggest a likely
 313 higher capability of triggering relevant waves.

314 Scenario 4 (in black, Figure 3) is the only subaerial scenario, and is located along the coastal cliffs of Capo Miseno, at the
 315 western end of the Gulf of Pozzuoli. It has been chosen by morphological considerations and recalling the 30 June 2025
 316 rockfall, assuming the large coastal subaerial scar of the eastern flank of Capo Miseno - still visible now - originated from a
 317 single, sudden collapse. The resulting scenario is quite different from the other cases: volume much smaller (around half
 318 million m³) and much larger initial thickness (maximum of more than 50 m).

319 As mentioned previously, accurate characterisation of geotechnical properties and rheological behaviour of these masses are
 320 not available. Since these factors can significantly influence both the landslide dynamics and the ensuing tsunami, this study
 321 adopts a set of reasonable assumptions, common to all involved cases. The slides are considered predominantly translational,
 322 allowing for elongation during motion and descent toward the deeper, central part of the basin. An exception is Scenario 1,
 323 where the presence of a hypothesized depositional area acts as a constraint on the model parameters regulating the dynamics.
 324 Although these assumptions are simplified and approximate, and each case would deserve more accurate reconstruction and
 325 characterization, they are considered acceptable within the context of WCTHA and for the purposes of this work, aiming at
 326 providing a first glance of the potential hazard of landslide-tsunamis in the area.

327
 328 **Table 2. Morphological characteristics of the Gulf of Pozzuoli landslide scenarios.**

329 Scenario	330 Environment	331 Volume (10 ⁶ m ³)	332 Area (km ²)	333 Maximum Thickness (m)	334 Initial maximum elevation (m)
332 1	333 Submarine	334 2.98	335 0.77	336 9.47	49
333 2	334 Submarine	335 2.30	336 0.56	337 10.11	37
334 3	335 Submarine	336 4.12	337 0.63	338 12.73	27
335 4	336 Subaerial	337 0.58	338 0.03	339 56.77	126

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)


ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

337
338



339

340 **Figure 3. Location of the initial bodies for the four landslide scenarios hypothesized: in blue, Scenario 1, between Capo Miseno and**
341 **Miseno Bank; in magenta, Scenario 2, south of Pozzuoli; in green, Scenario 3, offshore Bagnoli; in black, Scenario 4, along the cliffs**
342 **of Capo Miseno.**

343

344 3 Results

345 3.1 Landslide simulations

346 The code UBO-BLOCK has been applied to each of the scenarios described [in the previous section](#). The software requires as
347 input the initial landslide configuration (the undisturbed sliding surface and the upper surface of the mass), the predefined
348 trajectory of the CoM and the lateral boundaries of the surface swept by the sliding motion. In this way it is possible to obtain
349 the time history of the landslide shape changes and of its dynamics, representing the input for the computation of the
350 tsunamigenic impulse.

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha eliminato: Table 2. Parameters of the tsunami computational domain covering the Gulf of Pozzuoli: [\[1\]](#)
 N_x , number of nodes in the west-east direction; N_y , number of nodes in the north-south direction; $\Delta x, \Delta y$ spatial grid step along the x and y directions respectively; N_{tot} , total number of nodes; h_{max} maximum water depth in the computational grid. [\[2\]](#)

... [1]

ha eliminato:

ha eliminato: The evaluation of the landslide-tsunami hazard in the Gulf of Pozzuoli requires, as a first step, to identify the sources, i.e. masses that have the potential to generate waves, both in the submarine and in the subaerial environments. This investigation is here performed through a scenario approach, meaning that the sources are hypothesized starting from geological and morphological evidence, studying their signatures (scars, slopes, deposits) and the ensuing tsunamis evaluated through the numerical routine previously illustrated. The assumption is that such scenarios represent the range of possible and credible worst events occurring in the area, following the approach called Worst-case Credible Tsunami Hazard Assessment (WCTHA, Zaniboni and Armigliato, 2025), which is suitable to phenomena like landslide-tsunamis, where neither recurrence time are defined nor extensive catalogues of events are available, contrary to the most adopted approach in tsunami science (PTHA, see e.g. Behrens et al., 2021). [\[3\]](#)

In the following, as a first step, the search for mass collapse traces in the Gulf of Pozzuoli within the scientific literature is reported; from this a set of numerical scenarios is defined and the numerical routine previously described applied, providing a quantification of the hazard associated to the generated tsunamis. [\[4\]](#)

3.1 Mass instabilities in the Gulf of Pozzuoli and landslide scenarios [\[5\]](#)
Recent magnetic surveys in the Gulf of Naples evidenced the presence of consistent deposits on the seabed, testifying an intense mass transport activity in the whole area (de Ritis et al., 2024), that could have probably generated tsunamis affecting also the Gulf of Pozzuoli. The focus, though, is here centred on local sources, i.e. potential landslides triggered by the seismic shaking related to the Campi Flegrei caldera, then along the slopes (submarine and subaerial) in the Gulf of Pozzuoli area. [\[6\]](#)

As already mentioned, this is a small, shallow water basin, whose detailed stratigraphic and morphological features have been revealed by high-resolution seismic and bathymetric surveys (Aiello et al., 2012; Aiello et al., 2016; Somma et al., 2016). The area is characterized by numerous seismic units, both volcanic and sedimentary, and by a complex tectonic setting related to the intense Campi Flegrei volcanic activity. The northern part of the basin shows an inner shelf with maximum depth of 50 m, deepening to 100 m with gentle slopes and delimited by a belt of submarine volcanic edifices: starting from East, they are called Miseno Bank, Pentapalummo Bank and Nisida Bank (see Figure 3A). The [\[7\]](#)

... [2]

ha eliminato: [\[8\]](#)


... [3]

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman), Colore carattere: Automatico, Inglese (Regno Unito)

ha eliminato: previously

467 3.1 Scenario 1

468 The first scenario is submarine, with a volume around 3 million m³, and is placed south of Capo Miseno. Figure 4A shows the
 469 initial thickness distribution (yellow-red scale), and the predefined trajectory for the CoM, that has been determined based on
 470 the position of the final deposit found from the geomorphological survey (dashed-blue boundary). Figure 4B reports a section
 471 of the landslide, taken along the CoM trajectory: as previously noticed, the slopes are quite gentle, with an average value of
 472 1.5° for this case. In the simulation, the sliding mass settles between 80 and 100 m sea depth (red line, Figure 4B), reaching a
 473 maximum velocity of around 5 m/s after 200 seconds (Figure 4C), and decelerating quickly due to low seafloor gradient.
 474 Figure 4C reports the Froude number (Fr) evolution in time as well: this quantity is computed as the ratio between the landslide
 475 velocity and the tsunami phase velocity in the shallow-water approximation, depending on the water depth in a first instance.
 476 When Fr approaches one, the velocities of the two quantities are similar, and the mass-wave system reaches resonance
 477 condition: for supercritical values ($Fr > 1$) the mass moves faster than the wave: this occurs mainly in shallow water; in the
 478 subcritical regime ($Fr < 1$), typical of submarine slides, the wave travels much faster than the slide, which is then unable to
 479 "feed" the main tsunami front. In other words, Fr measures the efficiency of the energy transfer from the sliding mass to the
 480 wave. As already mentioned, in this investigation the wave-mass feedback (i.e. the effect of the generated tsunami on the slide
 481 motion) is not accounted for. For Scenario 1, Fr attains maximum values around 0.2 corresponding to the velocity peak,
 482 accounting then for a poor efficiency in the tsunami generation process.



484

485

ha eliminato: 2

ha eliminato: 5A

ha eliminato: 5B

ha eliminato: 5B

ha eliminato: 5C

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha eliminato:


ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)


Formattato: Allineato al centro



492 **Figure 4.** Panel A) Map of the initial sliding mass for Scenario 1, with initial thickness shown by the yellow-red scale. The black line
 493 marks the CoM predefined trajectory, defined to fit the observed deposit (dashed-blue boundary). The area swept by the sliding
 494 motion is highlighted in green. Panel B) Landslide profile along the CoM trajectory: in black the undisturbed sliding surface; in
 495 blue the initial mass; in red the simulated deposit. Panel C) Sliding velocity (red for average, black for each CoM) and Froude
 496 Number (in blue) time histories.
 497

3.1.2 Scenario 2

498 The second scenario is still submarine and is placed in the central part of the Gulf of Pozzuoli, about 1 km offshore from the
 499 piers of the homonymous city (Figure 5A). The hypothesized initial mass has been placed along the steeper slope connecting
 500 the shallow-water shelf to the deeper sea. The volume is similar to Scenario 1 (see Table 2), and the predefined sliding direction
 501 (black line in Figure 5A) mainly extends in the north-south direction. The simulation shows the deposit reaching the sub-
 502 horizontal seafloor at about 100 m depth (Figure 5B), with acceleration and deceleration phases almost symmetric, around the
 503 peak velocity value of 9 m/s (Figure 5C). The maximum value of Fr is here higher than the previous case (more than 0.3),
 504 remaining however largely subcritical and then limiting the build-up of the wave front.
 505



506 **Figure 5.** Panel A) Map of the initial sliding mass for Scenario 2, with initial thickness shown by the yellow-red scale. The black line
 507 marks the CoM predefined trajectory. The area swept by the sliding motion is highlighted in green. Panel B) Landslide profile along
 508 the CoM trajectory: in black the undisturbed sliding surface; in blue the initial mass; in red the simulated deposit. Panel C) Sliding
 509 velocity (red for average, black for each CoM) and Froude Number (in blue) time histories.
 510

3.1.3 Scenario 3

511 The last submarine scenario is located on the eastern side of the basin, just north of the small peninsula of Nisida and few
 512 hundreds of meters offshore Bagnoli (Figure 6A). As for the previous cases, the initial thickness is shown by the area in the
 513 yellow-red scale, showing that the mass is not distributed uniformly, but with the thicker part placed in deeper water. The
 514

ha eliminato: 5

ha eliminato: y: in red the average, in black the values for each CoM....

ha eliminato: 2

ha eliminato: 6A


ha eliminato: 6A

ha eliminato: 6

ha eliminato: 6

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)



ha eliminato:

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

Formattato: Allineato al centro


ha eliminato: 6

ha eliminato: Sliding velocity time history: in red the average, in black the values for each CoM

ha eliminato: 2

ha eliminato: 7A

529 sliding motion follows the main bathymetric gradient, south-westward (black line, Figure 6A), stopping at about 100 m depth,
 530 where the slope is quite horizontal (Figure 6B). The dynamics is characterized by a shorter acceleration phase, reaching the
 531 maximum velocity of almost 9 m/s after around 2 minutes (Figure 6C), followed by a longer deceleration taking 4 minutes
 532 before stopping. Similarly to scenario 2, Fr attains maximum values of 0.3, meaning poor efficiency in tsunami generation.



533
 534 Figure 6. Panel A) Map of the initial sliding mass for Scenario 3, with initial thickness shown by the yellow-red scale. The black line
 535 marks the CoM predefined trajectory. The area swept by the sliding motion is highlighted in green. Panel B) Landslide profile along
 536 the CoM trajectory: in black the undisturbed sliding surface; in blue the initial mass; in red the simulated deposit. Panel C) Sliding
 537 velocity (red for average, black for each CoM) and Froude Number (in blue) time histories.

538 3.4 Scenario 4

539 This scenario is the only subaerial one and presents very different morphological features compared to the previously illustrated
 540 cases: much smaller volume and initial area (see Table 2), much larger initial thickness, and an aspect ratio (length/width) very
 541 different from the submarine cases, as visible from Figure 7A. The initial mass has been hypothesized “filling” the subaerial,
 542 coastal scar still visible now along the eastern flank of Capo Miseno. Though purely theoretical, this type of collapse is not
 543 unusual in the whole area of the Gulf of Naples, as testified by the June 30th 2025 event at Punta Pennata, located close to this
 544 area; this scenario can be considered an endmember for this category of landslides.


545 The simulation shows that the deposit reaches a sub-horizontal area at 70-80 m depth (Figure 7B), with dynamics again
 546 different from the previous cases: a sudden acceleration brings the sliding mass at about 23 m/s within 10 seconds (Figure 7C),
 547 due to the very steep slope characterizing the first part of the trajectory (black line in Figure 7B); then the submarine shelf in
 548 very shallow water provokes an abrupt deceleration, down to 5 m/s, before a second acceleration due to the increasing slope
 549 between 20 and 60 m b.s.l., up to 10 m/s. Finally, the slide stops about 2 minutes after the onset. The Froude number crossed
 550 the critical value after around 20 seconds, when the mass is decelerating and moving along the shallow water platform offshore

ha eliminato: 7A

ha eliminato: 7B

ha eliminato: 7C

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)



ha eliminato:

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman), Inglese americano

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

Formattato: Allineato al centro

ha eliminato: 7

ha eliminato: Sliding velocity time history: in red the average, in black the values for each CoM

ha eliminato: 2

ha eliminato: 8A

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman), Apice

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)


ha eliminato: 8

Formattato: Normale

ha eliminato: 8C

ha eliminato: 8

564 Capo Miseno: this configuration turns out to be very efficient in generating the tsunami, since the waves are moving slower
 565 than in deep water and they are coupled with the mass motion on the seafloor.




566
 567 **Figure 7** Panel A) Map of the initial sliding mass for Scenario 4, with initial thickness shown by the yellow-red scale. The black line
 568 marks the CoM predefined trajectory. The area swept by the sliding motion is highlighted in green. Panel B) Landslide profile along
 569 the CoM trajectory: in black the undisturbed sliding surface; in blue the initial mass; in red the simulated deposit. Panel C) Sliding
 570 velocity (red for average, black for each CoM) and Froude Number (in blue) time histories.

571 3.2 Tsunami simulations

572 As illustrated previously, tsunamis can be significantly affected, during the propagation, by the hydrodynamic effect of
 573 dispersion, due to the different phase velocity characterizing its components. This phenomenon is particularly evident for short
 574 oscillations, which are more often induced by landslides. It is possible to estimate the distance at which such effects become
 575 predominant through Eq. (1), applying it to each of the studied scenarios. The dispersion can be quantified through the ratio
 576 λ/h_s with λ tsunami wavelength and h_s sea depth that can be determined as follows:

- 577 - λ_s , the initial wavelength of the tsunami, can be assessed in a first approximation as twice the longitudinal length of
 578 the slide, called b_s . This assumption, adopted in Heidarzadeh et al. (2023), though quite simplistic and rough, provides
 579 a first reasonable indication for this quantity.
- 580 - Since the focus here goes to the minimum value for the ratio λ/h_s delimiting the validity of SW approach, a value
 581 representing the maximum depth h_s for the tsunami propagation inside the Gulf of Pozzuoli is assumed: 105 m. The
 582 waves that are in the SW regime for this value, will satisfy this requirement also for shallower water.

583 Table 3 reports the estimations obtained for the landslide scenarios here adopted: the three submarine cases are in the SW
 584 regime, with the dispersion manifesting only for distances larger than the Gulf of Pozzuoli dimension. The subaerial case
 585 (Scenario 4) generates waves that are suddenly dominated by the dispersive effects: for this case the use of SW for the
 586 propagation on the whole Gulf of Pozzuoli seems not suitable.




594
595
596
597

Table 2. Computation of the dispersion distance for the four scenarios here studied (b : landslide length; λ : initial wavelength, assumed to coincide with b ; h : water depth (fixed at 105 m); D : dispersion distance, computed by Eq. (1); d : distance for which dispersion effects become predominant).

Scenario	Environment	b (m)	λ/h	D	d (km)
1	Submarine	1090	20.7	10.7	11.7
2	Submarine	1260	24.0	14.4	18.1
3	Submarine	1010	19.2	9.2	9.3
4	Subaerial	170	3.24	0.26	0.05

598



599


Figure 8. Maximum water amplitude on each node of the computational grid for the four scenarios considered: each of them are simulated through the shallow-water (SW) and the Boussinesq (NH) approach. The coloured boundaries report the initial position of the respective landslide source.

600 The simulations of the tsunamis generated by the sliding scenarios were performed using the JAGURS software, employing
601 both the SW and the non-hydrostatic (NH) approaches, with the non-linear version of the equations in both cases. The
602 computational grid resolution (20 m) is considered sufficient to investigate most of the tsunami features of interest in this
603 work; grid nesting (meaning higher resolution grid in specific, target areas) is thus not implemented here. This strategy allows
604 to investigate the suitability of the considerations previously made on the effects of dispersion on the tsunami propagation.

615 Figure 8 illustrates the maximum water amplitude on each point of the computational domain for each scenario, comparing
616 the two approaches: SW (upper row of plots) and NH (lower row). For the submarine scenarios (1, 2 and 3) the differences are
617 negligible; for Case 4 (last column) significant differences are evident, with the NH approach showing much more localized
618 effects compared to SW. This confirms the earlier hypothesis that dispersion effects cannot be neglected for sources of this
619 type.

620 From a hazard assessment perspective, the submarine scenarios here treated generate relatively small tsunamis in the Gulf of
621 Pozzuoli. In Cases 1 and 2, the maximum wave amplitudes are on the order of some tens of centimetres. In Case 3, the
622 maximum amplitude exceeds 50 cm along the coastal stretch between Bagnoli and Nisida, which, while not catastrophic, could
623 still cause damages to small boats and generate currents in smaller sub-basins. In contrast, Case 4 produces more significant
624 waves, especially at the local scale. Although the NH simulations show a rapid damping, localized amplifications can be
625 observed in more distant coastal stretches, such as around Pozzuoli (northern coast) and the Nisida peninsula (on the east).

626 Figure 9 depicts the maximum tsunami amplitude along the coast vs the cumulative coastal distance, measured from the eastern
627 extreme of the computational domain and represented with the black labels, in the left plot. The SW-NH simulations are almost
628 indistinguishable for the three submarine scenarios (right plot in Figure 9), with the respective waves amplitude that are almost
629 superimposed. Cases 1 and 2 show limited effects on the coast, while Case 3 generates maximum amplitudes of over 0.5 m
630 between Pozzuoli and Nisida. As previously observed, on the contrary, for the subaerial case (Scenario 4) the dispersive effects
631 play a key role, lowering considerably the maximum amplitude at the coast starting from the Pozzuoli coastal stretch (around
632 20 km of cumulative distance along the coast), with values almost halved at the opposite side of the Gulf. Conversely, the two
633 approaches produce similar waves for coastal stretches closer to the source, since for these the tsunami travels in shallower
634 water and the dispersive effects are then less intense. As already observed, this scenario produces the most impacting tsunami,
635 with peak value close to 5 m in Capo Miseno and local amplifications at Pozzuoli and Nisida with amplitudes of almost 2 m.



636

642 **Figure 9.** The map on the left reports the initial boundaries of the landslide scenarios. The cumulative distance along the coast is
643 measured from the eastern extreme of the computational domain. The plot on the right depicts the maximum water amplitude along
644 the coast for the four scenarios, with comparison between NH (continuous lines) and SW (dashed lines).

ha eliminato: 10


645 Figure 10 depicts the sketches of the first minutes of propagation for each scenario: all simulations show the typical feature
646 characterizing landslide-tsunamis, the almost circular shape of the tsunami signal, mimicking a point-like source. This polar
647 symmetry is lost when the wave interacts with shallow water and non-linear effects become predominant. A positive front
648 (yellow-red scale, meaning sea level increase) propagates in the same direction of the sliding motion, while a negative one
649 (cyan-blue scale) moves in the opposite side. For cases 2 and 3, the first manifestation of the tsunami at the coastal stretch
650 close to the source - presumably affected by the larger waves - is a negative signal, meaning that the water withdraws and is
651 followed later by a sea level increase: in terms of early warning this is undoubtedly an advantage, since it can act as a precursor
652 of an upcoming flooding. For case 1 the situation is different, since the slide motion does not have a direction opposite to the
653 dryland: a positive front enters the Gulf of Pozzuoli, while a negative one moves west of Capo Miseno, to the coastal stretch
654 out of the basin. As to case 4, the sliding motion starts in the subaerial environment, resulting into an always positive tsunami
655 front. Notice also the sequence of high frequency waves characterizing this scenario, especially evident in the 3- and 4-minutes
656 sketches (last row of plots), reflecting the smaller spatial dimensions of the tsunami source.

ha eliminato: 1

657

ha eliminato: an

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)



ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha eliminato: 1

ha eliminato: 11


ha eliminato: 12

ha eliminato: 12

661

662 **Figure 10.** Propagation sketches at 1 minute intervals of the four landslide-tsunami scenarios investigated (NH approach). The
 663 yellow-red scale marks the positive values (sea level increase), the cyan-blue scale is for negative ones (sea level sinking). The coloured
 664 contours represent the respective initial landslide boundary.

665 In all cases represented in Figure 10, the tsunami affects most of the Gulf of Pozzuoli coasts within 4 minutes. Figure 11 reports
 666 the travel time for each point of the computational domain, providing precious insights from the warning point of view:
 667 independently from the initial position, the waves take approximately 3 to 6 minutes to affect all the coasts of the basin. Case
 668 2 is the fastest, also due to its position at the centre of the Gulf. The north-western coastal stretch, on the contrary, is the one
 669 reached latest (5 to 6 minutes) in every scenario: the waves are slowed down by the shallow-water shelf that in this area is
 670 particularly large, compared to the other areas (as confirmed by the morphology, Figure 4). It is worth to specify that the code
 671 JAGURS registers the first positive signal for each computational cell: this explains the anomalous pattern of the tsunami
 672 opposite to the slide direction, particularly evident in Case 1 (westward, upper left panel, Figure 11) and Case 2 (northward,
 673 upper right panel).



678
679 **Figure 14.** Travel time, in seconds, of the tsunamis generated by the four landslide scenarios here hypothesized. The position of the
680 initial mass is marked by the respective coloured boundaries.

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha eliminato: 2

681 4 Discussion

682 The numerical simulations illustrated in the previous section provide some precious insights about the tsunami hazard pattern
683 within the Gulf of Pozzuoli. Submarine collapses of the size adopted in this study generate waves that do not represent a threat
684 for coastal population, but which can damage harbour facilities and small boats, which are present in a great number within
685 the Gulf of Pozzuoli. Conversely, the subaerial scenario produces high waves especially in the near field (almost 5 meters
686 high), which rapidly attenuate with distance thanks to dispersive effects. Some distant coastal stretches are affected by local
687 amplifications, with maximum amplitudes reaching almost 2 m (Pozzuoli, Nisida), highlighting the need to investigate this
688 type of events. In most cases, the tsunami reaches the shoreline in a few minutes with a positive signal, meaning that it manifests
689 as a water level increase. Only in limited coastal stretches, and not for all scenarios, the first signal is negative, i.e., the sea
690 withdraws for some minutes, providing a crucial precursor of an incoming wave in terms of early warning. In a few words,
691 these events can occur totally unannounced, reflecting the definition of “surprise tsunamis” given in Ward (2001). In the
692 following, some specific issues arising from the approach adopted and the simulation results are discussed.

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

694 Landslide-tsunami modelling. As shown, the scenarios considered involve heterogeneous contexts, occurring both in subaerial
695 and submarine environment. The use of the same modelling approach for both type of events deserves some discussion. The
696 interaction of a subaerial slide impacting water encompasses all the three phases (solid-liquid-air), resulting in a highly three-
697 dimensional, nonlinear process that is particularly difficult to simulate, especially near the source area. To address this
698 complexity, the phenomenon is typically divided into two main phases: i) the “splash-zone”, where such unsteady, complex
699 interactions dominate. The signals generated in this area, though intense and characterized by large amplitude, have high
700 frequency and dissipate quickly (Abadie et al., 2008). Consequently, their effects are confined to a limited area, generally
701 comparable in size to the slide run-out (Walder et al., 2006); ii) the propagation phase, beyond the splash-zone, where a
702 coherent, longer-period signal travels outward. Indeed, the tsunamigenic physical mechanism - whole water column uplift due
703 to the passage of the mass on the seabed – is the same. These oscillations exhibit peculiar characteristics but are broadly
704 analogous to submarine-generated tsunamis, apart from dispersive effects that can be captured by proper modelling. A
705 comprehensive review about this topic is provided by Yavari-Ramshe and Ataje-Ashtiani (2016). In this study, the adopted
706 modelling strategy deliberately neglects the splash-zone effects, assuming they are confined to a very narrow coastal stretch
707 at Capo Miseno. Consequently, the tsunami hazard assessment for the broader Gulf of Pozzuoli could be considered reliable.
708 This assumption is further supported by previous applications of the same numerical routine to other subaerial landslide-
709 generated tsunamis, such as the 1783 Scilla landslide-tsunami (see Zaniboni et al., 2016; Zaniboni et al., 2019).
710 Landslide scenarios. The scenario approach here adopted is a consequence of the lack of knowledge about the underwater
711 landslide bodies in the Gulf of Pozzuoli, and it is based on a worst-case methodology (WCTHA, Tonini et al., 2011).
712 Geophysical and bathymetric surveys have evidenced the presence of some ancient collapses, buried by the sediments, in the
713 deeper part of the basin at the toe of the slopes, but a general pattern of mass transport processes in the Gulf of Pozzuoli, with
714 recurrence time and volume estimation, does not exist. The sources adopted have been reconstructed based on the pieces of
715 evidence found in the scientific literature and on geomorphological considerations (margin slopes, existing scars, basin depth),
716 assuming that they are representative of the maximum credible occurrences expected in the area with the present morphology
717 and knowledge of the area. As a result, three submarine masses have been reconstructed, with similar volumes (few million
718 m³), thickness (a maximum of about ten meters) and detachment depth (in shallow water, between 30 and 50 m). The fourth
719 case is a coastal subaerial collapse and is characterized by a very different morphology: it can be considered as the endmember
720 of this type of landslide, since no direct evidence exists of bigger collapses interacting with the sea. Specific characteristics of
721 each sliding scenario could not be retrieved from the available geological and geophysical evidence: then, reasonable and
722 standard assumptions have been made for their rheology, hypothesizing for all of them a moderate translational behaviour.
723 These scenarios cover the whole extent of the Gulf, providing then a general idea of the impact expected from the ensuing
724 tsunamis. However, larger collapses cannot be ruled out, especially in case of intensification of the Campi Flegrei volcanic
725 activity providing possible triggers and, in the long term, slope oversteepening.
726 Dispersion effects. For tsunamis of non-seismic origin, the effect of dispersion should be always taken into consideration, since
727 it can change consistently the propagation pattern with respect to the classic SW approach. The discrepancy grows with the

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman), Corsivo

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha eliminato: s

ha eliminato: , Zaniboni and Armigliato, 2025

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha eliminato: doesn't

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha eliminato: can't

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

732 distance from the source, depending on the wavelength of the initial signal and on the depth of the basin where the perturbation
733 propagates. Eq. (1) provided rough estimates of this distance for the scenarios considered (reported in Table 2), suggesting
734 that for the submarine cases the dispersion is negligible within the Gulf of Pozzuoli domain. Numerical simulations confirmed
735 this hypothesis: the application of the code with (NH) and without (SW) dispersion produced almost identical tsunamis,
736 proving that the simpler and faster approach is sufficient to assess properly the tsunami hazard for these cases. For the subaerial
737 case, on the contrary, the difference is marked, as evidenced in Figure 9 by the maximum amplitudes along the coast. The
738 modelling effort, then, should consider the morphological features of the source generating the tsunami, keeping in mind that
739 subaerial masses collapsing into the sea usually generate shorter perturbations. In these cases, dispersive effects can become
740 relevant even for brief distances, and the application of the SW approach could produce an overestimation of the tsunami
741 impact on the coasts.

742 *Coastal and non-linear phenomena.* From the propagation plots (Figure 10) it is possible to infer some peculiar features of the
743 tsunami close to the coast, where the interaction with shallow water and minor basins can induce non-linear effects. For
744 example, for case 2 a positive signal propagating in the Bacoli Bay - a minor inlet just north of Capo Miseno on the western
745 part of the Gulf (see also Figure 12 for location) - can be noticed, evident especially in the $t = 3$ and 4 minutes sketches. This
746 is visible, while less marked, also for the other scenarios, and suggests the possibility of the excitation of the normal modes of
747 this sub-basin by the tsunami: this phenomenon, known as resonance, can occur for every basin affected by an external
748 perturbation, and is for example the mechanism at the basis of the generation of the meteotsunamis (Vilibic et al., 2016). The
749 morphology of the basin determines the periods of the resonant modes typical of the basin. Rabinovich (2009) obtained a set
750 of simple expressions allowing to estimate them for basic geometries. For example, for an open rectangular basin of length L
751 and uniform depth h , the period T of the fundamental mode is:

$$752 \quad T = \frac{4L}{\sqrt{gh}} \quad (2)$$

753 where g is the gravitational acceleration. Focusing on the Bacoli Bay case, Figure 12 reports the marigrams obtained from two
754 virtual tide gauges placed inside the inlet (tg#1, in red) and at its mouth (tg#2, in black), depicting the two respective time
755 histories in the four scenarios considered. The comparison of the signals shows clearly, for Cases 1, 2 and 3, that inside the
756 bay the perturbations behave as standing waves, characterized by regular oscillations lasting for at least 30 minutes (final
757 simulation time) with an approximate period of 200 s and evident amplifications if compared to the oscillations out of the basin
758 (in black). The tsunami generated by Case 3, in particular, is amplified five times with respect to the incoming signal. Assuming
759 for the Bacoli Bay a simplified rectangular geometry ($L \approx 1000 \text{ m}$, $h \approx 5 \text{ m}$) and applying Eq. (2), one can estimate the
760 fundamental mode as $T \approx 200 \text{ s}$, in full agreement with the features deduced from the virtual tide gauges. Moreover, the
761 subaerial scenario (Case 4) is less subject to the amplification when entering the inlet compared to the other cases, due to the
762 shorter period characterizing its oscillations.

763

ha eliminato: 4

ha eliminato: 10

ha eliminato: 11

ha eliminato: 13

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha eliminato: 2

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha eliminato: 3


ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha eliminato: 10

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)



771

772 **Figure 12.** Virtual tide gauges at the entrance (in black) and inside (in red) the Bacoli Bay. On the left, their location; on the right,
 773 the comparison between the marigrams for the scenarios here simulated.

774 5 Conclusions

775 The Gulf of Pozzuoli covers a significant portion of the Campi Flegrei caldera, a region of significant geological and volcanic
 776 activity. Intense seismic and volcanic processes have the potential to destabilize large sliding bodies both in the subaerial and
 777 in the submarine environment. When these masses interact with the water, they can generate tsunamis that may impact the
 778 entire basin's coastline, posing potential risks to local communities, coastal infrastructures, and marine activities.

779 Despite the region being the object of extensive geological and geophysical investigations, a comprehensive understanding of
 780 mass transport processes remains limited. A worst-case, scenario-based approach is then adopted, analysing four representative
 781 cases, based on the limited geological evidence and on morphological considerations: three submarine bodies, with similar
 782 volume (few million m³, occurring in shallow water) and one subaerial slide (smaller mass, occurring on the coastline). These
 783 scenarios are not intended to reproduce actual occurrences, while they aim at quantifying the tsunami hazard associated to
 784 events of this entity, providing precious insights into the potential tsunami generation mechanisms and their subsequent impact
 785 on coastal areas, as an important basis for possible risk mitigation strategies.

786 The sliding dynamics and the resulting tsunamis are simulated using numerical techniques that account for key hydrodynamic
 787 phenomena such as dispersion, nonlinear coastal effects, and resonance.

788 The simulations results indicate that submarine landslides generally produce waves of limited amplitude on the coast, with a
 789 maximum height of 0.5 meters, as observed for Case 3. While these tsunamis do not represent a major threat to coastal
 790 communities, they could still cause localized damages, for example to small boats moored in the harbours across the Gulf of

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha eliminato: 3

ha eliminato: , though not reproducing

ha eliminato: e

ha eliminato: providing

ha eliminato: case

796 Pozzuoli. Furthermore, resonance effects in smaller basins, such as harbours, can amplify an incoming wave, preventing its
797 dissipation and resulting in standing wave affecting the coast for long time: simulation outcomes show that in the Bacoli Bay
798 - placed in the western sector of the gulf – the incoming signal is amplified up to five times, with a sequence of regular
799 oscillations with a period of 200 s, affecting the basin for at least 30 minutes, which reflects the normal modes typical of the
800 inlet. This can repeat in every coastal basin, each one characterized by its own geometry and fundamental mode and is worthy
801 of specific and detailed investigation, implementing also nested grids at higher resolution. In contrast, the subaerial slide
802 scenario results in significantly larger waves, exceeding 4 meters close to the source. The tsunami amplitude damps rapidly
803 with distance, due to dispersion, but in some coastal stretches, Pozzuoli on the north and Nisida on the east, it reaches almost
804 2 meters.

805 The investigation here presented highlights the complex interplay between geological processes, hydrodynamic phenomena,
806 and coastal hazard in the Gulf of Pozzuoli. The results emphasize the need for detailed study and monitoring of potential
807 unstable masses both in the submarine and in the subaerial realm, determining their main features (geotechnical parameters,
808 possible rheology) which can influence deeply the tsunamigenesis. Coastal subaerial events, in particular, can give rise to
809 large tsunamis threatening the whole Gulf of Pozzuoli, enhancing the need, for risk management strategies to mitigate the
810 potential impact of tsunamis in this active volcanic region.

811 **Code availability**

812 The numerical codes UBO-BLOCK and UBO-TSUIMP for landslide dynamics and tsunamigenic impulse computation
813 respectively are available upon request to the authors; the code JAGURS can be freely downloaded from the link:
814 <https://github.com/jagurs-admin/jagurs>

815 **Data availability**

816 The computational grids have been obtained from the elaboration of raw datasets available online, which have been
817 interpolated and readjusted. They are available under request to the authors.

818 **Author contribution**

819 FZ and AA conceptualized the investigation; FZ, CA and MZ prepared the computational grids and the landslide scenario
820 datasets; FZ and LS performed the simulations; FZ prepared the manuscript, with the contribution from all co-authors; FZ and
821 AA supervised the whole manuscript realization.

ha eliminato: ,
ha eliminato: especially in c
ha eliminato: environment where they
ha eliminato:
ha eliminato: and

ha eliminato: nder
ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)
ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

828 **Competing interests**

829 The authors declare that they have no conflict of interest.

830 **Acknowledgements**

831 The authors are grateful to Prof. Jacopo Selva, University of Naples, Federico II, for the productive discussion about the
832 potential tsunami sources in the Gulf of Pozzuoli related to the Campi Flegrei activity, and two anonymous reviewers for their
833 constructive and insightful comments, improving considerably the manuscript.

834 

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

835 **References**

836 Abadie, S., Morichon, D., Grilli, S., Glockner, S.: VOF/Navier-Stokes numerical modeling of surface waves generated by
837 subaerial landslides, La Houille Blanche 1:21–26, doi:10.1051/lhb:2008001, 2016.

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman), Inglese americano

838 Aiello, G., Marsella, E. and Fiore, V. D.: New seismo-stratigraphic and marine magnetic data of the Gulf of Pozzuoli (Naples
839 Bay, Tyrrhenian Sea, Italy): inferences for the tectonic and magmatic events of the Phleorean Fields volcanic complex
840 (Campania), Mar Geophys Res, 33, 97–125, <https://doi.org/10.1007/s11001-012-9150-8>, 2012.

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

841 Baba, T., Ando, K., Matsuoka, D., Hyodo, M., Hori, T., Takahashi, N., Obayashi, R., Imato, Y., Kitamura, D., Uehara, H.,
842 Kato, T. and Saka, R.: Large-scale, highspeed tsunami prediction for the Great Nankai Trough Earthquake on the K computer,
843 Intern Jour of High Per Comp App, 30, doi: 10.1177/1094342015584090, 2015.

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

844 Behrens, J., Løvholt, F., Jalayer, F., Lorito, S., Salgado-Gálvez, M. A., Sørensen, M., Abadie, S., Aguirre-Ayerbe, I., Aniel-
845 Quiroga, I., Babeyko, A., Baiguera, M., Basili, R., Belliazz, S., Grezio, A., Johnson, K., Murphy, S., Paris, R., Raflana, I.,

ha eliminato: Aiello, G., Giordano, L. and Giordano, F.: High-resolution seismic stratigraphy of the Gulf of Pozzuoli (Naples Bay) and relationships with submarine volcanic setting of the Phleorean Fields volcanic complex, Rend. Fis. Acc. Lincei, 27, 775–801, <https://doi.org/10.1007/s12210-016-0573-z>, 2016.

846 De Risi, R., Rossetto, T., Selva, J., Taromi, M., Del Zoppo, M., Armigliato, A., Bureš, V., Cech, P., Cecioni, C.,
847 Christodoulides, P., Davies, G., Dias, F., Bayraktar, H. B., González, M., Gritsevich, M., Guillias, S., Harbitz, C. B., Kânoğlu,
848 U., Macías, J., Papadopoulos, G. A., Polet, J., Romano, F., Salamon, A., Scala, A., Stepinac, M., Tappin, D. R., Thio, H. K.,
849 Tonini, R., Triantafyllou, I., Ulrich, T., Varini, E., Volpe, M. and Vyhmeister, E.: Probabilistic Tsunami Hazard and Risk
850 Analysis: A Review of Research Gaps, Front. Earth Sci, 9:628772, doi: 10.3389/feart.2021.628772, 2021.

851 Bevilacqua, A., Isaia, R., Neri, A., Vitale, S., Aspinall, W. P., Bisson, M., Flandoli, F., Baxter, P. J., Bertagnini, A., Esposti
852 Ongaro, T., Iannuzzi, E., Pistolesi, M. and Rosi, M.: Quantifying volcanic hazard at Campi Flegrei caldera (Italy) with
853 uncertainty assessment: 1. Vent opening maps, Journal of Geophysical Research: Solid Earth, 120(4), 2309–2329.
854 <https://doi.org/10.1002/2014JB011775>, 2015.

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

860 Calcaterra, D. and Di Martire, D.: Landslide Hazard and Risk in the Campi Flegrei Caldera, Italy. In: Orsi, G., D'Antonio, M.,
 861 Civetta, L. (eds) Campi Flegrei. Active Volcanoes of the World, Springer, Berlin, Heidelberg, https://doi.org/10.1007/978-3-642-37060-1_13, 2022.

863 Danesi, S., Pino, N. A., Carlino, S. and Kilburn, C. R.: Evolution in unrest processes at Campi Flegrei caldera as inferred from
 864 local seismicity, *Earth and Planetary Science Letters*, 626, 118530, doi: <https://doi.org/10.1016/j.epsl.2023.118530>, 2024.

865 De Natale, G., Troise, C., Pingue, F., Mastrolorenzo, G., Pappalardo, L., Battaglia, M. and Boschi, E.: The Campi Flegrei
 866 caldera: unrest mechanisms and hazards. In Troise, C., De Natale, G., Kilburn C. R. J. (eds), *Mechanisms of Activity and*
 867 *Unrest at Large Calderas*, Geological Society, London, Special Publications, <https://doi.org/10.1144/GSL.SP.2006.269.01.03>,
 868 2006.

869 De Ritis, R., Cocchi, L., Passaro, S. and Chiappini, M.: Giant landslide, hidden caldera structure, magnetic anomalies and
 870 tectonics in southern Tyrrhenian Sea (Italy), *Geomorphology*, 466, 109445, <https://doi.org/10.1016/j.geomorph.2024.109445>,
 871 2024.

872 De Vivo, B., Rolandi, G., Gans, P. B., Calvert, A., Bohrson, W. A., Spera, F. J. and Belkin, H. E.: New constraints on the
 873 pyroclastic eruptive history of the Campanian volcanic Plain (Italy), *Mineralogy and Petrology*, 73, 47-65,
 874 <https://doi.org/10.1007/s007100170010>, 2001.

875 De Vivo, B., Belkin, H. E. and Rolandi, G.: Introduction to Vesuvius, Campi Flegrei, and Campanian Volcanism. In: De Vivo,
 876 B., Belkin, H. E., Rolandi, G. (eds) *Vesuvius, Campi Flegrei, and Campanian Volcanism*, Elsevier,
 877 <https://doi.org/10.1016/B978-0-12-816454-9.00001-8>, 2020.

878 Del Gaudio, C., Aquini, I., Ricciardi G. P., Ricco, C. and Scandone R.: Unrest episodes at Campi Flegrei: A reconstruction of
 879 vertical ground movements during 1905–2009, *Journal of Volcanology and Geothermal Research*, 195,10,
 880 <https://doi.org/10.1016/j.jvolgeores.2010.05.014>, 2010.

881 Di Vito, M., Acocella, V., Aiello, G., Barra, D., Battaglia, M., Carandente A., Del Gaudio C., de Vita, S., Ricciardi, G. P.,
 882 Ricco, C., Scandone, R. and Terrasi, F.: Magma transfer at Campi Flegrei caldera (Italy) before the 1538 AD eruption, *Sci
 883 Rep*, 6, 32245, <https://doi.org/10.1038/srep32245>, 2016.

884 DPC (Dipartimento della Protezione Civile - Civil Protection Department, Italy), MaGIC - Marine Geohazards along the Italian
 885 Coasts, <https://github.com/pcm-dpc/MaGIC>, last access 18 May 2023.

886 Ehara, A., Salmanidou, D. M., Heidarzadeh, M. and Guillas, S.: Multi-level emulation of tsunami simulations over Cilacap,
 887 South Java, Indonesia, *Computational Geosciences*, 27(1), 127-142, <https://doi.org/10.1007/s10596-022-10183-1>, 2023.

888 Esposito, G. and Matano, F.: A geodatabase of historical landslide events occurring in the highly urbanized volcanic area of
 889 Campi Flegrei, Italy, *Earth Syst. Sci. Data*, 15, 1133–1149, <https://doi.org/10.5194/essd-15-1133-2023>, 2023.

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman), Italiano

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

890 Gallotti, G., Zaniboni, F., Arcangeli, D., Angeli, C., Armigliato, A., Cocchi, L., Muccini, F., Zanetti, M., Tinti, S. and Ventura,
 891 G.: The tsunamigenic potential of landslide-generated tsunamis on the Vavilov seamount, *Journal of Volcanology and*
 892 *Geothermal Research*, 434, 107745, <https://doi.org/10.1016/j.jvolgeores.2023.107745>, 2023.

893 Gallotti, G., Zaniboni, F., Pagnoni, G., Romagnoli, C., Gamberi, F., Marani, M. and Tinti, S.: Tsunamis from prospected mass
 894 failure on the Marsili submarine volcano flanks and hints for tsunami hazard evaluation, *Bulletin of Volcanology*, 83, 2,
 895 <https://doi.org/10.1007/s00445-020-01425-0>, 2021.

896 Gasperini, L., Zaniboni, F., Armigliato, A., Tinti, S., Pagnoni, G., Özeren, M. S., Ligi, M., Natali, F. and Polonia, A.: Tsunami
 897 potential source in the eastern Sea of Marmara (NW Turkey), along the North Anatolian Fault system, *Landslides*, 19, 2295–
 898 2310, doi: 10.1007/s10346-022-01929-0, 2022.

899 Glimsdal, S., Pedersen, G., Harbitz, C. B. and Løvholt, F.: Dispersion of tsunamis: does it really matter?, *Nat. Hazards Earth*
 900 *Syst. Sci.*, 13, 1507–1526, doi: 10.5194/nhess-13-1507-2013, 2013.

901 Grezio, A., Cinti, F. R., Costa, A., Faenza, L., Perfetti, P., Pierdominici, S., Pondrelli, S., Sandri, L., Tierz, P., Tonini, R. and
 902 Selva, J.: Multisource Bayesian probabilistic tsunami hazard analysis for the Gulf of Naples (Italy), *Journal of Geophysical*
 903 *Research: Oceans*, 125, 2, <https://doi.org/10.1029/2019JC015373>, 2020.

904 [Harbitz, C.B., Løvholt, F., Pedersen, G. and Masson, D.G.: Mechanisms of tsunami generation by submarine landslides: a](#)
 905 [short review, Norwegian Journal of Geology, Vol. 86, pp. 255–264, Trondheim 2006, ISSN 029-196X, 2006.](#)

906 Heidarzadeh, M., Gusman, A.R. and Mulia, I.E.: The landslide source of the eastern Mediterranean tsunami on 6 February
 907 2023 following the M_w 7.8 Kahramanmaraş (Türkiye) inland earthquake, *Geosci. Lett.*, 10, 50, <https://doi.org/10.1186/s40562-023-00304-8>, 2023.

909 Isaia, R., Di Giuseppe, M. G., Natale, J., Tramparulo, F. D. A., Troiano, A. and Vitale, S.: Volcano-tectonic setting of the
 910 Pisciarelli fumarole field, Campi Flegrei caldera, southern Italy: insights into fluid circulation patterns and hazard scenarios,
 911 *Tectonics*, 40, 5, <https://doi.org/10.1029/2020TC006227>, 2021.

912 Løvholt, F., Pedersen, G., Harbitz, C. B., Glimsdal, S. and Kim, J.: On the characteristics of landslide tsunamis, *Phil. Trans. R. Soc. A*, 373: 20140376, doi: 10.1098/rsta.2014.0376, 2015.

914 Mayer, K., Scheu, B., Montanaro, C., Yilmaz, T. I., Isaia, R., Aßbichler, D. and Dingwell, D. B.: Hydrothermal alteration of
 915 surficial rocks at Solfatara (Campi Flegrei): Petrophysical properties and implications for phreatic eruption processes, *Journal*
 916 *of Volcanology and Geothermal Research*, 320, 128–143, <https://doi.org/10.1016/j.jvolgeores.2016.04.020>, 2016.

917 Neri, A., Bevilacqua, A., Esposti Ongaro, T., Isaia, R., Aspinall, W. P., Bisson, M., Flandoli, F., Baxter P. J., Bertagnini, A.,
 918 Iannuzzi, E., Orsucci, S., Pistolesi, M., Rosi, M. and Vitale, S.: Quantifying volcanic hazard at Campi Flegrei caldera (Italy)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

919 with uncertainty assessment: 2. Pyroclastic density current invasion maps, Journal of Geophysical Research: Solid Earth, 120,
 920 4, 2330-2349, <https://doi.org/10.1002/2014JB011776>, 2015.

921 Paris, R., Ulvrová, M., Selva, J., Brizuela, B., Costa, A., Grezio, A., Lorito, S. and Tonini, R.: Probabilistic hazard analysis for
 922 tsunamis generated by subaqueous volcanic explosions in the Campi Flegrei caldera, Italy. Journal of Volcanology and
 923 Geothermal Research, 379, 106-116, <https://doi.org/10.1016/j.jvolgeores.2019.05.010>, 2019.

924 Perrotta, A., Scarpati, C., Luongo, G. and Morra, V.: The Campi Flegrei caldera boundary in the city of Naples. In: De Vivo,
 925 B. (ed.), Developments in Volcanology, Elsevier, Volume 9, 85-96, [https://doi.org/10.1016/S1871-644X\(06\)80019-7](https://doi.org/10.1016/S1871-644X(06)80019-7), 2006.

926 Rabinovich, B. A.: Seiches and Harbor Oscillations, Handbook of Coastal and Ocean Engineering, 93-236, 2009.

927 Ren, Z., Hou, J., Wang, P. and Wang, Y.: Tsunami resonance and standing waves in Hangzhou Bay, Physics of Fluids, 33, 8,
 928 <https://doi.org/10.1063/5.0059383>, 2021.

929 Rosi, M., Sbrana, A. and Principe, C.: The Phlegraean Fields: structural evolution, volcanic history and eruptive
 930 mechanisms, Journal of Volcanology and Geothermal Research, 17, 1-4, 273-288, [https://doi.org/10.1016/0377-0273\(83\)90072-0](https://doi.org/10.1016/0377-0273(83)90072-0), 1983.

932 Sabino, L., Analisi della pericolosità da tsunami generati da frana nell'area del Golfo di Pozzuoli e dei Campi Flegrei, Second
 933 Level Degree in Physics of the Earth System, discussed on 14 march 2024, 2024 (in italiano).

934 Saito, T.: Tsunami Generation and Propagation, 265 - Springer Geophysics, <https://doi.org/10.1007/978-4-431-56850-6>,
 935 2019.

936 Selva, J., Amato, A., Armigliato, A., Basili, R., Bernardi, F., Brizuela, B., Cerminara, M., de' Micheli Vitturi, M., Di Bucci,
 937 D., Di Manna, P., Esposti Ongaro, T., Lacanna, G., Lorito, S., Løvholt, F., Mangione, D., Panunzi, E., Piatanesi, A., Ricciardi,
 938 A., Ripepe, M., Romano, F., Santini, M., Scalzo, A., Tonini, R., Volpe, M. and Zaniboni, F.: Tsunami risk management for
 939 crustal earthquakes and non-seismic sources in Italy, Riv. Nuovo Cim., 44, 69-144, <https://doi.org/10.1007/s40766-021-00016-9>, 2021.

941 Somma, R., Iuliano, S., Matano, F., Molisso, F., Passaro, S., Sacchi, M., Troise, C. and De Natale, G.: High-resolution morpho-
 942 bathymetry of Pozzuoli Bay, southern Italy, Journal of Maps, 12, 2, 222-230,
 943 <https://doi.org/10.1080/17445647.2014.1001800>, 2016.

944 Tarquini, S., Isola, I., Favalli, M., Battistini, A. and Dotta, G.: TINITALY, a digital elevation model of Italy with a 10 meters
 945 cell size (Version 1.1), Istituto Nazionale di Geofisica e Vulcanologia (INGV), <https://doi.org/10.13127/tinitaly/1.1>, 2023.

946 Tinti, S., Bortolucci, E. and Vannini, C.: A block-based theoretical model suited to gravitational sliding, Natural Hazards, 16,
 947 1-28, <https://doi.org/10.1023/A:1007934804464>, 1997.

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

948 Tinti, S., Pagnoni, G. and Zaniboni, F.: The landslides and tsunamis of 30th December 2002 in Stromboli analysed through
 949 numerical simulations, *Bulletin of Volcanology*, 68, 462-479, doi: 10.1007/s00445-005-0022-9, 2006.

950 [Tonini, R., Armigliato, A., Pagnoni, G., Zaniboni, F., Tinti, S.: Tsunami hazard for the city of Catania, eastern Sicily, Italy,](#)
 951 [assessed by means of Worst-case Credible Tsunami Scenario Analysis \(WCTSA\)](#), *Nat. Hazards Earth Syst. Sci.*, 11, 1217-
 952 1232, [www.nat-hazards-earth-syst-sci.net/11/1217/2011/](#), doi: 10.5194/nhess-11-1217-2011, 2011.

953 Triantafyllou, I., Zaniboni, F., Armigliato, A., Tinti, S. and Papadopoulos, G. A.: The Large Earthquake (~M7) and Its Associated Tsunami of 8 November
 954 1905 in Mt. Athos, Northern Greece, *Pure Appl. Geophys.*, 177, 1267-1293, <https://doi.org/10.1007/s0024-019-02363-5>,
 955 2020.

956 Vilibić, I., Šepić, J., Rabinovich, A. B. and Monserrat, S.: Modern approaches in meteotsunami research and early warning,
 957 *Frontiers in Marine Science*, 3, 57, <https://doi.org/10.3389/fmars.2016.00057>, 2016.

958 Walder, J.S., Watts, P., Waythomas, C.F.: Case study: mapping tsunami hazards associated with debris flow into a reservoir.
 959 *J Hydraul Eng* 132:1-11. doi:10.1061/(ASCE)0733-9429(2006)132:1(1), 2006.

960 [Yavari-Ramshe, S. and Ataie-Ashtiani, B.: Numerical modeling of subaerial and submarine landslide-generated tsunami](#)
 961 [waves—recent advances and future challenges. *Landslides* \(2016\) 13:1325–1368. DOI: 10.1007/s10346-016-0734-2,](#)
 962 [2016.](#)

963 Zaniboni, F. and Armigliato, A.: Worst-case tsunami approach applied to Catania (eastern Sicily). In Sørensen, M.,
 964 Behrens, J., Jalayer, F., Løvholt, F., Lorito, S., Rafliana, I., Salgado, M., Selva, J. (eds), *Probabilistic Tsunami Hazard and*
 965 *Risk Analysis*, Springer Nature, Switzerland, 2025 (in publication).

966 [Zaniboni, F., Armigliato, A. and Tinti, S.: A numerical investigation of the 1783 landslide-induced catastrophic tsunami in](#)
 967 [Scilla, Italy. *Nat Hazards* \(2016\) 84:S455–S470, doi: 10.1007/s11069-016-2461-3, 2016.](#)

968 Zaniboni, F., Pagnoni, G., Paparo, M. A., Gauchery, T., Rovere, M., Argnani, A., Armigliato, A. and Tinti S.: Tsunamis From
 969 Submarine Collapses Along the Eastern Slope of the Gela Basin Strait of Sicily, *Front. Earth Sci.* 8, 602171, doi:
 10.3389/feart.2020.602171, 2021.

970 [Zaniboni, F., Pagnoni, G., Gallotti, G., Paparo, M.A., Armigliato, A. and Tinti, S.: Assessment of the 1783 Scilla landslide-](#)
 971 [tsunami's effects on the Calabrian and Sicilian coasts through numerical modelling. *Nat. Hazards Earth Syst. Sci.*, 19, 1585–](#)
 972 [1600, DOI: https://doi.org/10.5194/nhess-19-1585-2019, 2019.](#)

973 Zaniboni, F., Pagnoni, G., Gallotti, G., Tinti, S. and Armigliato, A.: Landslide-tsunamis along the flanks of Mount Epomeo,
 974 Ischia: propagation patterns and coastal hazard for the Campania Coasts, Italy. In Marotta, E., D'Auria, L., Zaniboni, F., Nave,
 975 R. (eds), *Volcanic Island: from Hazard Assessment to Risk Mitigation*, Geological Society, London, Special Publications, 519,
 976 <https://doi.org/10.1144/SP519-2020-128>, 2024.

ha eliminato: 1

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha eliminato: 1

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman), Inglese (Regno Unito)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

979 Zollo, A., Maercklin, N., Vassallo, M., Dello Iacono, D., Virieux, J. and Gasparini, P.: Seismic reflections reveal a massive
980 melt layer feeding Campi Flegrei caldera, Geophysical Research Letters, 35, 12, <https://doi.org/10.1029/2008GL034242>,
981 2008.

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)

ha formattato: Tipo di carattere: (Predefinito) +Titoli (Times New Roman)