



Differentiation of primary and secondary marine organic aerosol with machine learning

Baihua Chen¹, Lu Lei², Emmanuel Chevassus², Wei Xu^{1*}, Ling Zhen¹, Haobin Zhong¹, Lin Wang¹, Chunshui Lin³, Ru-Jin Huang³, Darius Ceburnis², Colin O'Dowd², Jurgita Ovadnevaite^{2*}

¹State Key Laboratory of Advanced Environmental Technology, Institute of Urban Environment, Chinese Academy of Sciences, Xiamen, China.

²School of Physics, Centre for Climate & Air Pollution Studies, Ryan Institute, University of Galway, Galway, Ireland.

³State Key Laboratory of Loess and Quaternary Geology and Key Laboratory of Aerosol Chemistry and Physics, Institute of Earth Environment, Chinese Academy of Sciences, Xi'an, China.

Correspondence to: Wei Xu (wxu@iue.ac.cn) and Jurgita Ovadnevaite (jurgita.ovadnevaite@universityofgalway.ie)

Abstract. Marine primary organic aerosols (POA) are important components of the marine climate system, regulating solar radiation budget and cloud dynamics. Despite their importance, there is a lack of extensive long-term observations of POA properties, introducing great uncertainty in their parameterization in models. The lack of information originated from the complexity of POA chemical composition, very few long-term high-resolution measurements of clean marine air, and the difficulty in performing source apportionment techniques over a long-term period. In this study, we utilize a comprehensive high-resolution time-of-flight aerosol mass spectrometer dataset spanning a decade (2009-2018) and introduce a machine learning approach to differentiate and quantify the contribution of marine POA from marine secondary organic aerosol (SOA). Results indicate that marine POA concentrations peak during summer months and reach lowest levels in winter. On average, marine POA constitutes 51% (ranging from 21% to 76%) of the marine organic aerosol annually and up to 63% (48% to 75%) in summer. With the differentiated POA and SOA, we found diverse impacts of POA and SOA on aerosol hygroscopicity and mixing state. Increase in POA reduces the hygroscopicity and leads to external state of mixing, while the increase in SOA sustains the relatively high hygroscopicity and leads to internal mixing. This study provides observational dataset for marine POA and SOA and their diverse impacts on aerosol hygroscopicity, emphasizing a better appreciation of marine POA and SOA to improve the climate projections.

1 Introduction

Marine aerosols constitute a large portion of the global aerosol budget and are pivotal in regulating the Earth's climate system (Fitzgerald, 1991; O'Dowd and Leeuw, 2007). It has been known for quite some time that marine aerosols contain a significant amount of organic matter (Blanchard, 1964). Cavalli et al. (2004) and O'Dowd et al. (2004) show the great contribution of organic matter in the Northeast Atlantic marine aerosol during periods of high biological activities. These marine organic aerosol originates from two sources: (1) bubble bursting that scavenges surface-active organic matter and other biogenic



materials (bacteria, viruses and detritus), producing primary organic aerosol (POA) (Barger and Garrett, 1970; Blanchard, 1964; Blanchard and Woodcock, 1957); (2) oxidation of marine volatile organic compounds (VOCs) such as dimethyl sulfide, aliphatic amines, isoprene and monoterpenes, which can form secondary organic aerosol (SOA) (Bates et al., 1992; Bonsang et al., 1988; Charlson et al., 1987; Facchini et al., 2008; Meskhidze and Nenes, 2006; Wohl et al., 2023; Zheng et al., 2020).

- 35 Marine POA is crucial for regulating cloud properties, acting as cloud condensation nuclei (CCN) or ice nuclei (IN). Ovadnevaite et al. (2011a) documented a marine POA plume with a peak POA concentration of up to $3.8 \mu\text{g m}^{-3}$, which is comparable to levels found in European continental air. Ovadnevaite et al. (2011b) further highlighted that marine POA has low hygroscopicity but high CCN potential. Additionally, sea spray tank experiments have demonstrated a significant correlation between seawater nanophytoplankton cell abundances and sea spray CCN number fluxes (Sellegri et al., 2021).
- 40 Incorporating marine POA into global models necessitates a comprehensive understanding of source strength and environmental response of POA. O'Dowd et al. (2008) proposed an integrated organic-inorganic sea spray source function that accounted for a size-dependent contribution of POA to total sea spray aerosol. Further parameterization efforts have considered factors such as chlorophyll-a concentration and wind speed (Gantt et al., 2011; Gantt et al., 2012; Rinaldi et al., 2013), which have been integrated into global chemical transport models. However, the influence of marine biota on the
- 45 chemical composition and cloud activation properties of POA remains a contentious topic. O'Dowd et al. (2015) observed significant changes in the CCN activities of sea spray aerosol during a phytoplankton plume, whereas Quinn et al. (2014) and Bates et al. (2020) reported no substantial alterations in CCN activity.

In summary, the source intensity, chemical composition, mixing state, and cloud condensation activation potential of marine POA remain poorly understood (Gantt and Meskhidze, 2013) with large discrepancies between modelled and measured POA

50 (Gantt et al., 2015). A major challenge in improving POA parameterization and modelling is the lack of long-term datasets, which are critical for both understanding the environmental drivers of POA emissions, and developing emission schemes for regional or global chemical transport models. The majority of the available data, such as that in Rinaldi et al. (2013), are derived from filter measurements that require extended sampling durations and result in low time resolution (days to weeks). Although filter-based methods can distinguish POA from SOA by using chemical molecular fingerprints (O'Dowd et al.,

55 2004), they suffer from low temporal resolution, limiting their ability to capture dynamic changes in aerosol composition (minutes to hours).

The deployment of the Aerosol Mass Spectrometer (AMS) in both coastal and remote marine atmospheres has provided an opportunity to improve the POA parameterization and refine model predictions (Choi et al., 2017; Huang et al., 2018; Ovadnevaite et al., 2014; Saliba et al., 2020; Sanchez et al., 2020; Schmale et al., 2013; Willis et al., 2017). The AMS enables

60 near-real-time measurements of aerosol chemical composition (DeCarlo et al., 2004, 2006), including organic aerosol (OA), non-sea salt sulfate (nss-SO₄), ammonium (NH₄), nitrate, methanesulphonic acid (Ovadnevaite et al., 2014) and sea salt (Ovadnevaite et al., 2012). While POA and SOA can be differentiated using their mass spectra fingerprints through positive



matrix factorization (PMF), this method faces challenges in high time resolution long-term data sets due to the required workforce and computational cost (Chevassus et al., 2024).

65 In this study, we developed a machine learning (ML) model to differentiate the contributions of marine POA and SOA from the measured total marine OA, using long-term marine aerosol measurements obtained by an AMS at the Mace Head Atmospheric Research Station. This model effectively identified and quantified contributions from the marine POA from SOA. The impacts of POA and SOA on aerosol hygroscopicity were then investigated.

2 Methods

70 2.1 Data and instrumentation

Aerosol measurements were conducted at the Mace Head Atmospheric Research Station (53°19' N, 9°54' W) on the west coast of Ireland from 2009 to 2018. The station, regularly exposed to clean marine air masses from the North Atlantic, has been a representative site for studying clean marine aerosols for several decades (O'Dowd et al., 2014).

We employed a High-Resolution Time-of-Flight AMS (DeCarlo et al., 2004) at Mace Head (Ovadnevaite et al., 2014) to
75 measure the PM₁ (particulate matter with diameter smaller than 1 µm) chemical composition including organic aerosol (OA), non-sea-salt sulfate (nss-SO₄), sea salt, methanesulfonic acid (MSA) (Ovadnevaite et al., 2014), ammonium (NH₄), and nitrate (NO₃). Additionally, black carbon (BC) was measured using a Multi-Angle Absorption Photometer (MAAP) to trace anthropogenic emissions. Meteorological conditions were also recorded, including air temperature, pressure, precipitation, relative humidity, wind speed, and wind direction.

80 The humidified tandem differential mobility analyzer (HTDMA) (Swietlicki et al., 2000) was used to measure aerosol hygroscopic growth at a fixed relative humidity of 90% for aerosol with selected dried sizes of 35, 50, 75, 110, and 165 nm. The growth factors measured by HTDMA were inverted using a piecewise linear function (Gysel et al., 2009) and converted to hygroscopicity parameter κ (Kreidenweis et al., 2008), assuming the surface tension of water-air interface.

Data from the AMS and MAAP data were averaged to a 10-minute resolution, while the meteorological records were initially
85 recorded hourly and later downsampled to 10-minute intervals using linear interpolation to enlarge the dataset's availability. Any gaps in the AMS or MAAP data that contain invalid measurements were removed. Hourly boundary layer heights were obtained from ERA5 (Hersbach et al., 2020) and downsampled to 10-minute resolution using linear interpolation.

2.2 Clean sector criteria and machine learning strategy

To differentiate between marine POA and SOA in MHD, this study employs an ML model to predict the mass concentration
90 of marine SOA. Subsequently, POA concentrations are obtained by subtracting SOA from the measured total OA. To ensure the high quality of the SOA production period, we implemented multiple screening criteria to minimize the influence of



anthropogenic and marine primary signatures. First, we applied clean sector criteria, limiting BC concentration to less than 15 ng m⁻³ and selecting wind direction between 190 and 300 °(O'Dowd et al., 2014), to exclude continental outflows and ship plumes. The clean sector criteria, which have been established and applied in various MHD research studies (O'Dowd et al., 2014; Ovadnevaite et al., 2014; Xu et al., 2022), were employed to isolate marine air masses from anthropogenic influences.

We then applied additional filtering processes to reduce the impact of POA production. We further refined the data by keeping instances with wind speed below 6 m/s and sea salt mass concentration under 0.03 µg m⁻³ to minimize the concentration of marine POA. Finally, we retained only those data points where nss-SO₄ was the dominant component (nss-SO₄/OA > 4) to ensure a predominantly secondary source. We presumed that in these selected data, OA was predominantly SOA, with the contribution of POA represented by a minor and constant background concentration (POA_{bg}).

Subsequently, we employed a support vector regression (SVR)(Awad et al., 2015) trained on these rigorously selected SOA production periods. SVR was chosen for its generalizability in handling small datasets and its resistance to overfitting. Unlike tree-based models like random forest (Breiman, 2001), SVR model can predict continuous values. The hyperparameters, including the penalty coefficient (C) and gamma (γ) of the Radial Basis Functions (RBF) kernel were tuned via grid research. The model targeted OA concentration, using nss-SO₄, MSA, NH₄, and meteorological parameters (temperature, relative humidity, boundary layer height, wind direction, and pressure) as predictors. We also included hours of the day to capture diurnal variations. Predictors not directly linked to secondary production, e.g., sea salt, NO₃, and BC, are excluded to avoid over-fitting and ensure generalizability, even though including these might have enhanced the model performance in the training dataset. Wind speed was used to select the SOA production period, therefore, it was not suitable as a predictor. A summary of the variables employed as predictors is shown in Table 1.

Table 1. Predictors used for SOA ML model.

Predictors	Acronyms	Source	Rationale
Hour of the day	hour		to capture any diurnal pattern
nss-SO ₄	SO ₄	AMS measurement	Secondary aerosol marker
MSA	MSA	AMS measurement	Secondary aerosol marker
NH ₄	NH ₄	AMS measurement	Secondary aerosol marker
Temperature	temp	Meteorological records	Known to influence secondary processes
Rain	rain	Meteorological records	Related to wet removal
wind direction	wddir	Meteorological records	Related to sources
Boundary layer height	blh	ERA5 reanalysis	Related to concentration

The selected SOA production periods were split by the year 2015: data before 2015 were used to train the mode. The training dataset covered a significant fraction of the variabilities of predictors, as illustrated in Fig. S1. The hyperparameters of the



115 SOA-SVR model were optimized using grid search and 5-fold cross validation. The data after 2015, which was unseen for the training process, was used to challenge the model's generalizability. Overall, there are 1700 hours of SOA production period, and 477 hours (27.8%) after 2015.

120 A schematic diagram of the proposed methodology is shown in Fig. 1. Initially, the clean marine dataset was extracted by applying the clean sector criteria, followed by additional filtering processes to minimize the influence of POA. To assess the representativeness of the selected data as secondary sources, the Fuzzy C-Means (FCM) clustering method was utilized. Subsequently, data were divided into training, validation, and test sets for ML, with SOA (including minor POA_{bg}) as the predictive variable. To mitigate experimental uncertainties, cross-validation and Monte Carlo simulations were performed. The study further investigated the magnitude of the POA_{bg} values. Finally, ML method was applied to predict SOA concentrations, enabling the differentiation of POA concentrations within the total marine OA.




Figure 1. The proposed data processing and model construction workflow. POA_{bg} represents the assumed constant POA background concentration. ML (chem + met + time) represents the machine learning model that uses chemical composition, meteorological and time parameters as predictors.

2.3 Fuzzy-C Means clustering

FCM is a clustering algorithm that enables the grouping of data points into multiple clusters with varying degrees of membership. Unlike traditional hard clustering techniques, where each data point is assigned to a single cluster, FCM assigns membership levels to each data point, indicating the degree to which it belongs to each cluster. This soft clustering approach



is particularly useful when dealing with complex datasets where boundaries between clusters are not well-defined or overlap significantly. By optimizing an objective function that minimizes the weighted sum of squared errors, FCM iteratively updates the cluster centers and membership degrees, providing a flexible and robust means of uncovering underlying patterns and structures within data.

3 Results and Discussions

3.1 Performance of ML model in SOA production period

We first examined the aerosol composition in the selected secondary marine aerosol dataset. In these SOA production period, nss-SO₄ constituted $68.9 \pm 8.7\%$ of the PM₁ mass, followed by OA at $12.1 \pm 3.4\%$, and MSA at $6.8 \pm 4.0\%$. The average concentration of sea salt and wind speed were $0.015 \pm 0.009 \mu\text{g m}^{-3}$ and $4.6 \pm 1.2 \text{ m/s}$, respectively. The MSA to nss-SO₄ ratio was 0.10 ± 0.06 , aligning with previous findings at the same site (Ovadnevaite et al., 2014). BC concentrations remained well below the 15 ng m^{-3} threshold, averaging at $6.0 \pm 3.7 \text{ ng m}^{-3}$. Taken together, this chemical composition indicates the selected data mainly originated from secondary sources.

To ensure the representativeness of the selected training data as secondary sources, we applied FCM method on the clean marine dataset to identify the characteristics of chemical and meteorological parameters from typical sources. Compared to more conventionally used k-means clustering, FCM allows data instances to belong to multiple clusters with varying degrees of membership (or probability). The membership is used to determine how strongly each data instance belongs to each cluster. The FCM clustering, which is independent of the selection of SOA production periods, provides further validation and examination of the data selection for model training. We selected those with any cluster membership higher than 80% to show the clustering center of each cluster. As shown in Fig. S2a, the 2nd factor, which is featured by low sea-salt, low wind speed and high temperature, is most likely to be of the secondary origin. This factor also showed high MSA and nss-SO₄, supporting the selection criteria for SOA production period. Indeed, the highest possibility of the selected training data is found to be the 2nd factor (Fig. S2b), reaffirming the SOA production characteristics.

Subsequently, we employed the SVR model, leveraging the clean marine dataset, to predict OA (SOA + POA_{bg}) concentrations. Cross-validation yields Pearson's R of 0.97 for training and validation datasets, demonstrating the model's accuracy in predicting OA concentration using the selected predictors. The slopes between estimated SOA + POA_{bg} and measured OA were 0.97 for both the train and validation data, indicating robust pattern recognition across most concentration ranges. The model also exhibited great generalizability and performed consistently well on an unseen dataset (Fig. 2c), with a Pearson's R value of 0.94 and a slope of 0.98 between observed OA and estimated SOA + POA_{bg}, reaffirming the model's efficacy in modeling the complex dynamics of SOA.




Figure 2. Observed OA versus predicted (SOA + POA_{bg}) for (a) training, (b) validation and (c) test datasets. Data density is illustrated using a color gradient with darker colour indicating lower data density. Black lines denote the 1:1 correspondence lines, blue lines represent regression lines.

Permutation importance analysis highlighted nss-SO₄ and MSA as the most influential variables, followed by NH₄ (Fig. S3a). Partial dependent plots (Fig. S3b) indicate a nonlinear relationship between SOA and increasing levels of nss-SO₄ and MSA. While these plots do not imply a causal relationship, they highlight the complexity of the interactions and underscore the importance of employing a machine learning model to effectively capture such intricate patterns. Various meteorological parameters were also found to influence SOA concentration, especially relative humidity and precipitation, but to a lower extent.

In this study, one of the major assumptions of this approach is to assume that the OA in the selected secondary marine data for training the model is dominated by SOA. To evaluate the potential influence of POA contributions in those secondary marine data cases on entire clean marine dataset, we conducted a sensitivity analysis. We assumed that POA_{bg} constitutes 5% to 30% of OA. As shown in Fig. S4, compared to the original assumption, the monthly averaged concentration of POA systematically increased throughout most of the year, except during winter, when alternative assumption predicted POA concentrations lower than 0, which is, of course, non-physical. Then, we applied the model on the entire clean marine dataset and tried different fixed POA_{bg} values iteratively. As shown in Fig. S5, the POA_{bg} of 0.01 μg m⁻³ yields the least nonphysical predictions (either POA or SOA lower than 0). Therefore the POA_{bg} of 0.01 μg m⁻³ was used in the following calculation. Based on this strategy, SOA concentrations were predicted, enabling the estimation of POA concentrations. Furthermore, if POA production period is defined as periods with POA concentrations exceeding 0.1 μg m⁻³ for more than 12 hours, there were more than 60 such POA production periods during the 10-year period (Fig. S6). Detecting these POA production periods allows for detailed characterization, potentially enhancing its parameterization.



Note that the measurement itself contains relatively large uncertainty, compared to summertime measurement. Indeed, we manually tuned the POA_{bg} value to minimize the negative values. This is supported by previous studies that marine is a large organic pool (Quinn et al., 2014). Given the inherent uncertainties in aerosol measurements, as well-documented in previous study (Ovadnevaite et al., 2014), we further quantified the uncertainties associated with ML model using Monte Carlo simulations. To do this, we performed a robustness test by randomly validating the model 1000 times, each time excluding 20% of the data from the training set. The lower and upper limits of the estimated POA seasonality are shown in Fig. S7, which is similar to the original model. The Monte Carlo ensembles demonstrated negligible differences in the contribution of POA, indicating stable model performance across different scenarios.

To validate the ML-based POA concentrations, we further compared it with PMF-based POA concentrations from Chevassus et al. (2024). The PMF-based source apportionment was conducted for about one month. The Pearson's correlation coefficient between ML-based POA and PMF-based POA was about 0.91, indicating strong agreement between the two methods (Fig. 3). Compared to the conventional AMS-based OA source apportionment techniques, e.g. PMF, this ML approach requires significantly fewer computational resources and is less dependent on detailed knowledge of the mass spectra signatures of marine POA and SOA. For example, the model performed equally well even after removing MSA as a predictor (Fig. S8). Given that MSA can only be resolved by high-resolution AMS, this suggests that our approach could be extended to Aerosol Chemical Species Monitor (ACSM) data, which is more affordable and widely used but has lower mass resolution. This would enable broader applications of our method, offering a more comprehensive understanding of marine POA over global oceans. Finally, although secondary production of OA and $nss-SO_4$ rely on similar meteorological conditions, it should be noted that many marine VOC species do not share the same sources and oxidization pathways as DMS and its derivatives. For example, some VOCs are produced by different organisms or abiotically from sea surface microlayer (Ciuraru et al., 2015; Mungall et al., 2017), which could introduce additional uncertainty in the SOA quantification based on high $nss-SO_4$ periods, which adds to the uncertainty for POA attribution as well.




Figure 3. The comparison between machine learning-derived POA and HR-PMF POA. The HR-PMF POA data is taken from Chevassus et al. (2024). The blue line is regression line and grey area represents 95% confidence interval. The black line represents 1:1 line. Pearson's correlation coefficient and the equation of regression line are shown in the top left.

210 3.2 Case study and long-term seasonality

The evaluation of the model's performance has shown a clear relationship between SOA and the predictor variables in the chosen SOA production period. Assuming there is little and constant contribution from POA_{bg} , the estimated SOA concentration should be similar to the measured OA. We then evaluated the model's performance in well-defined cases over finer time scales. During a typical SOA production period from 11st to 14th August 2011 (Fig. 4a), which was not included in the training dataset because of the slightly elevated sea salt above the threshold of $0.03 \mu g m^{-3}$, the measured OA closely followed the variation of nss-SO₄ and MSA, with Pearson's R values of over 0.80 and 0.84, respectively. These high correlations indicate a predominant secondary source of OA. The model's estimates of SOA concentrations were very close to the measured total OA (Fig. 4b), with an OA/SOA ratio of 1.03 ± 0.04 . This further suggests that the ML model is able to predict the variability of SOA.

220 In contrast, during a well-documented marine POA plume, from 13rd to 18th August 2009 (Fig. 4c), the OA dominated the PM₁ concentration and showed little correlation with nss-SO₄ or MSA (Pearson's R of 0.05 and 0.08, respectively). Notably, the



estimated SOA deviates significantly from the observed OA (Fig. 4d). In this instance, the SOA estimated by the ML model accounted for only about 20% of the total OA during the plume, underscoring the significant contribution of marine POA. The different performance, presented in Fig. 4, is expected. In Fig. 4a, the model is mostly influenced by the nss-SO₄, a marker for secondary species, leading to a great agreement between modelled SOA and observed OA. Conversely, Fig. 4b shows a significant discrepancy between modelled SOA and observed OA. The difference is largely attributed to the contribution of POA. This case was reported by Ovadnevaite et al. (2011a), in which the marine POA was identified using HR-ToF-AMS mass spectra, and the SOA during the period was not quantified.




Figure 4. Case study of (a-b) SOA production period and (c-d) POA production period. (a,c) time series of PM₁ chemical species and (b,c) measured OA versus estimated SOA, the colour presents data density with darker colour indicating lower data density, the black lines represent 1:1 lines.

The cases of SOA and POA production periods indicate the model's capability to predict the SOA and POA levels. As illustrated in Fig. 5, both POA and SOA reached their peak in June and dropped to their lowest during the winter months. Typically, median concentrations of SOA were higher than those of POA across most months, except for May. However, POA concentrations spiked periodically, highlighted by outliers. The mean relative contribution of POA to total OA, detailed in Fig. 5b, shows the lowest contribution in winter and the largest from May to July, peaking at approximately 50% and slightly later in summer than SOA. The pattern corresponds with the enhanced marine activities in later spring and early summer of the North Atlantic, involving extensive phytoplankton proliferation and other marine organisms that release organic matter into the atmosphere through wave breaking and bubble bursting.



It is important to note that the total marine OA concentrations during winter at Mace Head are very low, introducing substantial uncertainty in OA separation during this season. The minimal POA concentration observed in winter suggests a distinct relationship between total OA, secondary species, and environmental factors. This relationship closely mirrors the dynamics observed in SOA. As shown in Fig. S9, the estimated SOA closely aligns with measured OA throughout the winter, pointing to low contribution from POA. Conversely, during the summer, numerous data points shown in Fig. S9 deviate to the right of the 1:1 lines, indicating a substantial contribution from POA.




Figure 5. The seasonality of POA and SOA. (a) boxplot of mass concentration of POA (dark green) and SOA (purple), the horizontal lines represent median, the boxes represent 25th and 75th quantile, and the whiskers represent 1.5 inter-quarter ranges. Note that outliers are not fully shown to ease to visualization. (b) The contribution of POA to total OA, the line represents the monthly median, and the shaded area represents the 25th and 75th quantile.

The differentiation of POA from SOA is further substantiated by additional correlation analysis. As shown in Fig. S10, the correlation between OA and nss-SO_4 across the entire clean marine dataset is relatively low, at approximately 0.17, suggesting that nss-SO_4 explains less than 3% of the variability of marine OA. Upon decoupling the OA into POA and SOA, we observed distinct correlation patterns: the correlation between SOA and nss-SO_4 increased to 0.88, reflecting a strong linkage. Whereas for POA, it decreased to 0.08. This stark contrast underscores the different sources and atmospheric behaviors of marine POA and SOA. This different correlation analysis provides a clear delineation of how POA and SOA contribute to marine OA and emphasizes the capacity of advanced modeling techniques and long-term observations to unravel complex atmospheric processes.

The potential sources of POA and SOA were investigated using the Potential Source Contribution Function (PSCF) combined with air mass backward trajectories (Mansour et al., 2020). As shown in Fig. 6, POA likely originates from the Northeast



Atlantic polar marine regions (Fig. 6a), which are recognized as biologically active waters. In contrast, SOA sources are traced to tropical marine regions (Fig. 6b). The identified POA sources align with previous studies suggesting that regional marine biological activity is a key driver of POA production (O'Dowd et al., 2004; Sellegri et al., 2021). For SOA, enhanced photochemical reactions in lower-latitude waters likely promote the formation of secondary species. The distinct source regions of marine POA and SOA underscore the need for models to incorporate specific parameterisation schemes that account for these spatial and mechanistic differences.




Figure 6. The potential source contribution function analysis of marine POA (a) and SOA (b). Redish colour represents higher probability.

In contrast to prior studies that relied on filter-based measurements with limited temporal resolution, this study introduces a ML framework to systematically differentiate and quantify marine POA and SOA. While seasonal variations in POA/SOA have been reported previously, our decade-long dataset—the most extensive of its kind to date—provides unprecedented resolution to constrain and develop POA and SOA parameterization for climate models. Furthermore, the distinct source regions identified for POA (polar marine zones) and SOA (tropical waters) underscore their divergent formation mechanisms. Current model estimated of global emissions of POA span from 6.9 - 76 Tg year⁻¹ for <1 μm emissions and global source of SOA were thought to be substantially smaller than marine POA. Our measurements found similar contribution of POA and SOA. While it has to be noted that some fractional of SOA were transformed via atmospheric aging from POA, which is difficult to quantified. Recent study found the photochemical reactions in the sea-air interface produces substantial VOC as the precursors of marine SOA (Brüggemann et al., 2018), the complex sources of SOA highlights the need for field observational data to challenge the models. The combined sensitivity to marine biological activities and photochemistry of marine OA was also supported by Sanchez et al. (2020), which found high correlation with downward shortwave flux and net primary production, while they did not separate POA and SOA.



3.3 Impact of marine organic aerosol to aerosol hygroscopicity and mixing state

- 285 The long-term marine POA and SOA time series now enable an assessment of their influence on aerosol hygroscopicity—a relationship previously uncertain. As shown in Fig. 7, increasing POA concentrations from 0.1 to 1 $\mu\text{g m}^{-3}$ reduces the hygroscopicity parameter (κ) across all particle sizes, dropping values from ~ 0.5 to below 0.25. This finding contrasts with earlier experimental work suggesting POA production has negligible effects on aerosol hygroscopicity and CCN activity (Quinn et al., 2014), but aligns with our prior case study highlighting the inherently low hygroscopicity of POA
- 290 (Ovadnevaite et al., 2011b). Quinn et al. (2014) investigated the impact of marine POA on its cloud condensation nuclei using a sea sweep devices, they found no significant difference of POA contribution at different oceanic regions with diverse ranges of chlorophyll-a, and they attributed the POA to the ocean carbon pool. However, based on our field measurements, it is unlikely that ocean carbon pool induces such large variations of the observed POA, underscoring the importance of oceanic biological activities.
- 295 As for the SOA, the increasing SOA concentrations from 0.1 to 0.3 $\mu\text{g m}^{-3}$ only slightly reduce κ_{HTDMA} values (from 0.5 to 0.45) for particles between 50 and 165 nm, with no significant change observed for 35 nm particles. This muted response may arise from co-varying increases in secondary species such as nss-SO_4^{2-} or MSA, which help maintain hygroscopicity at relatively high levels. This is consistent with our recent high-temporal online measurement, which shows a simultaneous increase in SOA and nss-SO_4^{2-} during particle growth (Xu et al., 2024; Zheng et al., 2020). These results underscore the distinct
- 300 roles of POA and SOA in modulating aerosol water uptake and cloud-forming potential, emphasizing the need to explicitly represent OA composition and sources in climate models.




Figure 7. The impact of POA (a) and SOA (b) on aerosol hygroscopicity parameter (K_{HTDMA}) at different sizes. The analysis was limited to April to August to minimize the seasonal variations. The black lines represent medians, dark shaded areas represent 25th to 75th percentiles, darker shaded areas represent 10th to 90th percentiles.

The influence of POA and SOA on aerosol mixing state was further investigated using the spread factor, calculated from growth-factor probability density functions (Xu et al., 2020). A spread factor of 0 indicates a theoretically *internal* mixture, while higher values reflect increasing *external* mixing. Based on established thresholds (Swietlicki et al., 2008; Xu et al., 2019), a spread factor of ≤ 0.05 is classified as internal mixture, whereas values ≥ 0.2 signify external mixing. POA and SOA exhibit divergent impacts on aerosol mixing state. As shown in Fig. 8a, increasing the POA contribution from 0% to 100% elevates the spread factor, suggesting POA production promotes external mixing. Conversely, SOA accumulation drives the



system toward a more internally mixed state. This aligns with aerosol aging processes, where particles tend to homogenize over time. Accurately representing the hygroscopicity and mixing-state dynamics of POA and SOA is critical for assessing their climatic impacts, as these properties directly influence aerosol-cloud interactions and radiative forcing.




Figure 8. The impact of percentage contribution of POA (a) and SOA (b) to aerosol mixing state (spread factor) at different sizes. The analysis was limited to April to August to minimize the seasonal variations. The black lines represent medians, dark shaded areas represent 25th to 75th percentiles, darker shaded areas represent 10th to 90th percentiles.



4 Conclusion

320 Quantifying marine POA and SOA traditionally relies on PMF applied to aerosol mass spectra, which is challenging for long-term data. This study presents a data-driven ML framework to identify and quantify marine POA by leveraging temporal data patterns rather than chemical mass signatures. The ML model, trained on rigorously selected SOA-dominated periods, was applied to a multi-year aerosol dataset, enabling the identification of numerous POA production events. At Mace Head, marine POA constitutes ~50% of total marine organic aerosol (OA), increasing to 63% during late spring and early summer. Unlike
325 PMF, this ML approach proves particularly effective for disentangling OA components in complex, long-term environments where high-resolution AMS data are unavailable.

Combined with aerosol hygroscopicity measurements, our analysis reveals distinct climatic impacts: marine POA significantly reduces aerosol hygroscopicity and promotes external mixing, whereas SOA exhibits weaker effects. These findings underscore the need to accurately quantify marine POA abundance and its influence on cloud-relevant properties. A key
330 limitation lies in the selection of SOA-dominated periods for model training; future work should optimize ML performance for smaller or less curated datasets. Additionally, validating these marine POA results with global oceanic measurements is essential to refine POA parameterizations in climate and chemical transport models.

Supplement

The supplement related to this article is available.

335 Acknowledgement

This study is financial supported by National Key R&D Program of China (2023YFC3705503), the Guiding Project of Seizing the Commanding Heights of City Self-purifying Technology (No. IUE-CERAE-202403), Natural Science Foundation of China (No. 42307591), State Environmental Protection Key Laboratory of Formation and Prevention of Urban Air Pollution Complex, European Union's Horizon Europe Research and Innovation programme under HORIZON-CL5-2022-D1-02 (grant
340 no. 101081430-PARIS), EPA-Ireland and the Department of Environment, Climate and Communications.

Data availability

The AMS datasets used and/or analyzed during the current study are available on reasonable request from the corresponding author Jurgita Ovadnevaite (jurgita.ovadnevaite@universityofgalway.ie). Aerosol hygroscopicity dataset is available via mendeley data (<http://doi.org/10.17632/3dx6pnx869.1>). ERA5 data is available from the CDS
340 (<https://cds.climate.copernicus.eu/#!/home>).



Code availability

The code is available upon request to the corresponding author (wxu@iue.ac.cn) and Jurgita Ovadnevaite (jurgita.ovadnevaite@universityofgalway.ie).

Author contribution

350 WX, LL and JO conceived the study, WX, BC, LZ, EC, CL, LW and HZ analyzed the data and constructed the models. JO and DC conducted the AMS and BC measurements. WX, RH and CO wrote the paper with the input from all the authors.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

355 References

- Awad, M., Khanna, R., Awad, M., and Khanna, R.: Support vector regression, Efficient learning machines: Theories, concepts, and applications for engineers and system designers, 67–80, doi:<https://dl.acm.org/doi/10.5555/2785649>, 2015.
- Barger, W. and Garrett, W.: Surface active organic material in the marine atmosphere, *Journal of Geophysical Research*, 75, 4561–4566, doi:<https://doi.org/10.1029/JC075i024p04561>, 1970.
- 360 Bates, T. S., Calhoun, J. A., and Quinn, P. K.: Variations in the methanesulfonate to sulfate molar ratio in submicrometer marine aerosol particles over the south pacific ocean, *Journal of Geophysical Research: Atmospheres*, 97, 9859–9865, doi: <https://doi.org/10.1029/92JD00411>, 1992.
- Bates, T. S., Quinn, P. K., Coffman, D. J., Johnson, J. E., Upchurch, L., Saliba, G., Lewis, S., Graff, J., Russell, L. M., and Behrenfeld, M. J.: Variability in Marine Plankton Ecosystems are not Observed in Freshly Emitted Sea Spray Aerosol over
- 365 the North Atlantic Ocean, *Geophysical Research Letters*, doi:<https://doi.org/10.1029/2019gl085938>, 2020.
- Blanchard, D. C.: Sea-to-air transport of surface active material, *Science*, 146, 396–397, 1964.
- Blanchard, Dc. and Woodcock, A.: Bubble formation and modification in the sea and its meteorological significance, *Tellus*, 9, 145–158, doi: <https://doi.org/10.3402/tellusa.v9i2.9094> ,1957.
- Bonsang, B., Kanakidou, M., Lambert, G., and Monfray, P.: The marine source of c 2-c 6 aliphatic hydrocarbons, *Journal of*
- 370 *Atmospheric Chemistry*, 6, 3–20, doi: <https://doi.org/10.1007/BF00048328>, 1988.
- Breiman, L.: Random Forests, *Machine Learning*, 45, 5–32, doi:<https://doi.org/10.1023/a:1010933404324>, 2001.
- Brüggemann, M., Hayeck, N., and George, C.: Interfacial photochemistry at the ocean surface is a global source of organic vapors and aerosols, *Nature Communications*, 9, 2101, doi:<https://doi.org/10.1038/s41467-018-04528-7>, 2018.



- 375 Cavalli, F., Facchini, M., Decesari, S., Mircea, M., Emblico, L., Fuzzi, S., Ceburnis, D., Yoon, Y., O'Dowd, C., Putaud, J.-P.,
et al.: Advances in characterization of size-resolved organic matter in marine aerosol over the north atlantic, *Journal of*
Geophysical Research: Atmospheres, 109, doi:<https://doi.org/10.1029/2004JD005137>, 2004.
- Charlson, R. J., Lovelock, J. E., Andreae, M. O., and Warren, S. G.: Oceanic phytoplankton, atmospheric sulphur, cloud
albedo and climate, *Nature*, 326, 655–661, doi:<https://doi.org/10.1038/326655a0>, 1987.
- Chevassus, E., Fossum, K. N., Ceburnis, D., Lei, L., Lin, C., Xu, W., O'Dowd, C. D., and Ovadnevaite, J.: Marine organic
380 aerosols at mace head: Effects from phytoplankton and source region variability, *EGUsphere*, 2024, 1–33,
doi:<https://doi.org/10.5194/egusphere-2024-2890>, 2024.
- Choi, Y., Rhee, T. S., Collett, J. L., Park, T., Park, S.-M., Seo, B.-K., Park, G., Park, K., and Lee, T.: Aerosol concentrations
and composition in the North Pacific marine boundary layer, *Atmospheric Environment*, 171, 165–172, doi:
<https://doi.org/10.1016/j.atmosenv.2017.09.047>, 2017.
- 385 Ciuraru, R., Fine, L., Pinxteren, M. van, D'Anna, B., Herrmann, H., and George, C.: Photosensitized production of
functionalized and unsaturated organic compounds at the air-sea interface, *Scientific Reports*, 5, 12741,
doi:<https://doi.org/10.1038/srep12741>, 2015.
- DeCarlo, P. F., Slowik, J. G., Worsnop, D. R., Davidovits, P., and Jimenez, J. L.: Particle Morphology and Density
Characterization by Combined Mobility and Aerodynamic Diameter Measurements. Part 1: Theory, *Aerosol Science and*
390 *Technology*, 38, 1185–1205, doi:<https://doi.org/10.1080/027868290903907>, 2004.
- DeCarlo, P. F., Kimmel, J. R., Trimborn, A., Northway, M. J., Jayne, J. T., Aiken, A. C., Gonin, M., Fuhrer, K., Horvath, T.,
Docherty, K. S., Worsnop, D. R., and Jimenez, J. L.: Field-Deployable, High-Resolution, Time-of-Flight Aerosol Mass
Spectrometer, *Analytical Chemistry*, 78, 8281–8289, doi:<https://doi.org/10.1021/ac061249n> 2006.
- Facchini, M. C., Decesari, S., Rinaldi, M., Carbone, C., Finessi, E., Mircea, M., Fuzzi, S., Moretti, F., Tagliavini, E.,
395 Ceburnis, D., and O'Dowd, C. D.: Important Source of Marine Secondary Organic Aerosol from Biogenic Amines,
Environmental Science & Technology, 42, 9116–9121, doi:<https://doi.org/10.1021/es8018385>, 2008.
- Fitzgerald, J. W.: Marine aerosols: A review, *Atmospheric Environment. Part A. General Topics*, 25, 533–545, 1991.
- Gantt, B. and Meskhidze, N.: The physical and chemical characteristics of marine primary organic aerosol: a review,
Atmospheric Chemistry and Physics, 13, 3979–3996, doi:<https://doi.org/10.5194/acp-13-3979-2013>, 2013.
- 400 Gantt, B., Meskhidze, N., Facchini, M. C., Rinaldi, M., Ceburnis, D., and O'Dowd, C.: Wind speed dependent size-resolved
parameterization for the organic mass fraction of sea spray aerosol, *Atmospheric Chemistry and Physics*, 11, 8777–8790,
doi:<https://doi.org/10.5194/acp-11-8777-2011> 2011.
- Gantt, B., Johnson, M. S., Meskhidze, N., Sciare, J., Ovadnevaite, J., Ceburnis, D., and O'Dowd, C. D.: Model evaluation of
marine primary organic aerosol emission schemes, *Atmospheric Chemistry and Physics*, 12, 8553–8566,
405 doi:<https://doi.org/10.5194/acp-12-8553-2012>, 2012.
- Gysel, M., McFiggans, G., and Coe, H.: Inversion of tandem differential mobility analyser (TDMA) measurements, *Journal*
of Aerosol Science, 40, 134–151, doi:<https://doi.org/10.1016/j.jaerosci.2008.07.013>, 2009.



- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., Rosnay, P. de, Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J.-N.: The ERA5 global reanalysis, *Quarterly Journal of the Royal Meteorological Society*, 146, 1999–2049, doi:<https://doi.org/10.1002/qj.3803>, 2020.
- Huang, S., Wu, Z., Poulain, L., Pinxteren, M. van, Merkel, M., Assmann, D., Herrmann, H., and Wiedensohler, A.: Source apportionment of the organic aerosol over the atlantic ocean from 53° n to 53° s: Significant contributions from marine emissions and long-range transport, *Atmospheric Chemistry and Physics*, 18, 18043–18062, doi: <https://doi.org/10.5194/acp-18-18043-2018>, 2018.
- Kreidenweis, S. M., Petters, M. D., and DeMott, P. J.: Single-parameter estimates of aerosol water content, *Environmental Research Letters*, 3, 035002, doi: <https://doi.org/10.1088/1748-9326/3/3/035002>, 2008.
- Mansour, K., Decesari, S., Facchini, M. C., Belosi, F., Paglione, M., Sandrini, S., Bellacicco, M., Marullo, S., Santoleri, R., Ovadnevaite, J., Ceburnis, D., O'Dowd, C., Roberts, G., Sanchez, K., and Rinaldi, M.: Linking Marine Biological Activity to Aerosol Chemical Composition and Cloud-Relevant Properties Over the North Atlantic Ocean, *Journal of Geophysical Research: Atmospheres*, 125, doi:<https://doi.org/10.1029/2019jd032246>, 2020.
- Meskhidze, N. and Nenes, A.: Phytoplankton and Cloudiness in the Southern Ocean, *Science*, 314, 1419–1423, doi:<https://doi.org/10.1126/science.1131779>, 2006.
- Mungall, E. L., Abbatt, J. P. D., Wentzell, J. J. B., Lee, A. K. Y., Thomas, J. L., Blais, M., Gosselin, M., Miller, L. A., Papakyriakou, T., Willis, M. D., and Liggio, J.: Microlayer source of oxygenated volatile organic compounds in the summertime marine Arctic boundary layer, *Proceedings of the National Academy of Sciences*, 114, 6203–6208, doi:<https://doi.org/10.1073/pnas.1620571114>, 2017.
- O'Dowd, C. and Leeuw, G. de: Marine aerosol production: a review of the current knowledge, *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 365, 1753–1774, doi:<https://doi.org/10.1098/rsta.2007.2043>, 2007.
- O'Dowd, C., Facchini, M. C., Cavalli, F., Ceburnis, D., Mircea, M., Decesari, S., Fuzzi, S., Yoon, Y. J., and Putaud, J.-P.: Biogenically driven organic contribution to marine aerosol, *Nature*, 431, 676–680, doi:<https://doi.org/10.1038/nature02959>, 2004.
- O'Dowd, C., Ceburnis, D., Ovadnevaite, J., Vaishya, A., Rinaldi, M., and Facchini, M. C.: Do anthropogenic, continental or coastal aerosol sources impact on a marine aerosol signature at Mace Head?, *Atmospheric Chemistry and Physics*, 14, 10687–10704, doi: <https://doi.org/10.5194/acp-14-10687-2014>, 2014.
- O'Dowd, C. D., Langmann, B., Varghese, S., Scannell, C., Ceburnis, D., and Facchini, M. C.: A combined organic-inorganic sea-spray source function, *Geophysical Research Letters*, 35, doi:<https://doi.org/10.1029/2007gl030331>, 2008.



- O'Dowd, C. D., Ceburnis, D., Ovadnevaite, J., Bialek, J., Stengel, D. B., Zacharias, M., Nitschke, U., Connan, S., Rinaldi, M., Fuzzi, S., Decesari, S., Facchini, M. C., Marullo, S., Santolero, R., Dell'Anno, A., Corinaldesi, C., Tangherlini, M., and Danovaro, R.: Connecting marine productivity to sea-spray via nanoscale biological processes: Phytoplankton Dance or Death Disco?, *Scientific Reports*, 5, 1 11, doi:<https://doi.org/10.1038/srep14883>, 2015.
- 445 Ovadnevaite, J., O'Dowd, C., Dall'Osto, M., Ceburnis, D., Worsnop, D. R., and Berresheim, H.: Detecting high contributions of primary organic matter to marine aerosol: A case study, *Geophysical Research Letters*, 38, 2011a.
- Ovadnevaite, J., Ceburnis, D., Martucci, G., Bialek, J., Monahan, C., Rinaldi, M., Facchini, M. C., Berresheim, H., Worsnop, D. R., and O'Dowd, C.: Primary marine organic aerosol: A dichotomy of low hygroscopicity and high CCN activity, *Geophysical Research Letters*, 38, L21806, doi:<https://doi.org/10.1029/2011GL048869>, 2011b.
- 450 Ovadnevaite, J., Ceburnis, D., Canagaratna, M., Berresheim, H., Bialek, J., Martucci, G., Worsnop, D. R., and O'Dowd, C.: On the effect of wind speed on submicron sea salt mass concentrations and source fluxes, *Journal of Geophysical Research: Atmospheres*, 117, D16201, doi:<https://doi.org/10.1029/2011JD017379>, 2012.
- Ovadnevaite, J., Ceburnis, D., Leinert, S., Dall'Osto, M., Canagaratna, M., O'Doherty, S., Berresheim, H., and O'Dowd, C.: Submicron NE Atlantic marine aerosol chemical composition and abundance: Seasonal trends and air mass categorization, *Journal of Geophysical Research: Atmospheres*, 119, 11, 850–11, 863, doi:<https://doi.org/10.1002/2013JD021330>, 2014.
- 455 Quinn, P. K., Bates, T. S., Schulz, K. S., Coffman, D. J., Frossard, A. A., Russell, L. M., Keene, W. C., and Kieber, D. J.: Contribution of sea surface carbon pool to organic matter enrichment in sea spray aerosol, *Nature Geoscience*, 7, 228, 2014.
- Rinaldi, M., Fuzzi, S., Decesari, S., Marullo, S., Santolero, R., Provenza, A., Hardenberg, J. von, Ceburnis, D., Vaishya, A., O'Dowd, C. D., and Facchini, M. C.: Is chlorophyll- a the best surrogate for organic matter enrichment in submicron primary
- 460 marine aerosol?, *Journal of Geophysical Research: Atmospheres*, 118, 4964–4973, doi:<https://doi.org/10.1002/jgrd.50417>, 2013.
- Saliba, G., Chen, C., Lewis, S., Russell, L. M., Quinn, P. K., Bates, T. S., Bell, T. G., Lawler, M. J., Saltzman, E. S., Sanchez, K. J., Moore, R., Shook, M., Rivellini, L., Lee, A., Baetge, N., Carlson, C. A., and Behrenfeld, M. J.: Seasonal Differences and Variability of Concentrations, Chemical Composition, and Cloud Condensation Nuclei of Marine Aerosol
- 465 over the North Atlantic, *Journal of Geophysical Research: Atmospheres*, doi:<https://doi.org/10.1029/2020jd033145>, 2020.
- Sanchez, K. J., Zhang, B., Liu, H., Saliba, G., Chen, C.-L., Lewis, S. L., Russell, L. M., Shook, M. A., Crosbie, E. C., Ziemba, L. D., Brown, M. D., Shingler, T. J., Robinson, C. E., Wiggins, E. B., Thornhill, K. L., Winstead, E. L., Jordan, C., Quinn, P. K., Bates, T. S., Porter, J., Bell, T. G., Saltzman, E. S., Behrenfeld, M. J., and Moore, R. H.: Linking marine phytoplankton emissions, meteorological processes and downwind particle properties with FLEXPART, *Atmospheric*
- 470 *Chemistry and Physics Discussions*, doi:<https://doi.org/10.5194/acp-2020-702>, 2020.
- Schmale, J., Schneider, J., Nemitz, E., Tang, Y. S., Dragosits, U., Blackall, T. D., Trathan, P. N., Phillips, G. J., Sutton, M., and Braban, C. F.: Sub-antarctic marine aerosol: Dominant contributions from biogenic sources, *Atmospheric Chemistry and Physics*, 13, 8669–8694, doi:<https://doi.org/10.5194/acp-13-8669-2013>, 2013.



- Sellegri, K., Nicosia, A., Freney, E., Uitz, J., Thyssen, M., Grégori, G., Engel, A., Zäncker, B., Haëntjens, N., Mas, S.,
475 Picard, D., Saint-Macary, A., Peltola, M., Rose, C., Trueblood, J., Lefevre, D., D'Anna, B., Desboeufs, K., Meskhidze, N.,
Guieu, C., and Law, C. S.: Surface ocean microbiota determine cloud precursors, *Scientific Reports*, 11, 281,
doi:<https://doi.org/10.1038/s41598-020-78097-5>, 2021.
- Swietlicki, E., Zhou, J., Covert, D. S., Hameri, K., Busch, B., Vakeva, M., Dusek, U., Berg, O. H., Wiedensohler, A., Aalto,
P. P., Makela, J., Martinsson, B. G., Papaspiropoulos, G., Menten, B., Frank, G., and Stratmann, F.: Hygroscopic properties
480 of aerosol particles in the north- eastern Atlantic during ACE-2, *Tellus B: Chemical and Physical Meteorology*, 52, 201–227,
doi:<https://doi.org/10.1034/j.1600-0889.2000.00036.x>, 2000.
- Swietlicki, E., Hansson, H. C., Hämeri, K., Svenningsson, B., Maßling, A., McFiggans, G., McMurry, P. H., Petaja, T.,
Tunved, P., Gysel, M., Topping, D. O., Weingartner, E., Baltensperger, U., Rissler, J., Wiedensohler, A., and Kulmala, M.:
Hygroscopic properties of submicrometer atmospheric aerosol particles measured with H-TDMA instruments in various
485 environmentsa review, *Tellus B: Chemical and Physical Meteorology*, 60, 432–469, doi:<https://doi.org/10.1111/j.1600-0889.2008.00350.x>, 2008.
- Willis, M. D., Köllner, F., Burkart, J., Bozem, H., Thomas, J. L., Schneider, J., Aliabadi, A. A., Hoor, P. M., Schulz, H.,
Herber, A. B., Leaitch, W. R., and Abbatt, J. P. D.: Evidence for marine biogenic influence on summertime Arctic aerosol,
Geophysical Research Letters, 44, 6460–6470, doi:<https://doi.org/10.1002/2017gl073359>, 2017.
- 490 Wohl, C., Li, Q., Cuevas, C. A., Fernandez, R. P., Yang, M., Saiz-Lopez, A., and Simó, R.: Marine biogenic emissions of
benzene and toluene and their contribution to secondary organic aerosols over the polar oceans, *Science Advances*, 9,
eadd9031, doi:<https://doi.org/10.1126/sciadv.add9031>, 2023.
- Xu, W., Ovadnevaite, J., Fossum, K. N., Lin, C., Huang, R.-J., O'Dowd, C., and Ceburnis, D.: Aerosol hygroscopicity and
its link to chemical composition in coastal atmosphere of mace head: Marine and continental air masses, *Atmospheric*
495 *Chemistry and Physics*, 2019, 1–25, doi:<https://doi.org/10.5194/acp-2019-839>, 2019.
- Xu, W., Ovadnevaite, J., Fossum, K. N., Lin, C., Huang, R.-J., O'Dowd, C., and Ceburnis, D.: Aerosol hygroscopicity and
its link to chemical composition in the coastal atmosphere of mace head: Marine and continental air masses, *Atmospheric*
Chemistry and Physics, 20, 3777–3791, doi:<https://doi.org/10.5194/acp-20-3777-2020>, 2020.
- Xu, W., Ovadnevaite, J., Fossum, K. N., Lin, C., Huang, R.-J., Ceburnis, D., and O'Dowd, C.: Sea spray as an obscured
500 source for marine cloud nuclei, *Nature Geoscience*, 15, doi:<https://doi.org/10.1038/s41561-022-00917-2>, 2022.
- Xu, W., Ovadnevaite, J., Fossum, K. N., Huang, R.-J., Huang, D. D., Zhong, H., Gu, Y., Lin, C., Huang, C., O'Dowd, C.,
and Ceburnis, D.: Condensation of organic-inorganic vapours governs the production of ultrafine secondary marine cloud
nuclei, *Communications Earth & Environment*, 5, 359, doi:<https://doi.org/10.1038/s43247-024-01519-z>, 2024.
- Zheng, G., Kuang, C., Uin, J., Watson, T., and Wang, J.: Large contribution of organics to condensational growth and
505 formation of cloud condensation nuclei (CCN) in remote marine boundary layer, doi: <https://doi.org/10.5194/acp-20-12515-2020>, 2020.