

**Contribution of dark inorganic carbon fixation to bacterial carbon demand in
the oligotrophic Southeastern Mediterranean Sea**

Tom Reich ^{1,2*}, Natalia Belkin ¹, Guy Sisma-Ventura ¹, Hagar Hauzer ¹, Maxim
Rubin-Blum ^{1,2}, Ilana Berman-Frank ^{2*}, and Eyal Rahav ^{1,3,4 *}

¹ National Institute of Oceanography, Israel Oceanographic and Limnological
Research, Haifa, Israel.

² Department of Marine Biology, Leon H. Charney School of Marine Sciences,
University of Haifa, Haifa, Israel.

³ Department of Earth and Environmental Science, Ben-Gurion University of the
Negev, Beer Sheva, Israel.

⁴ Institute of Marine Science, University of California, Santa Cruz, CA, USA.

*Correspondence: TR- treich02@campus.haifa.ac.il; IBF- iberman2@univ.haifa.ac.il;
ER- eyal.rahav@ocean.org.il

Abstract

Photosynthetically derived organic matter sinking to depth from the illuminated layers
is often not sufficient to meet the energy demands of microbes in the dark ocean. This
‘mismatch’ is especially notable in the warm and oligotrophic eastern Mediterranean
Sea where the annual photosynthesis is one of the lowest in the world’s oceans, yet its
aphotic zone is considered a hotspot for microbial activity and biomass. Here, we
investigated the role of photic and aphotic dark inorganic carbon fixation rates (DCF)

and its contribution to bacterial carbon demand in the southeastern Mediterranean Sea during the mixed and stratified periods. Our results demonstrate that DCF rates are measurable throughout the water column (0-1750 m) and are on the same order of magnitude as photosynthesis (34 vs. 45 g C m⁻² y⁻¹, respectively). Using a carbon mass balance that considers photosynthesis, DCF and bacterial production (and hence respiration) we show that chemoautotrophy provides ~35% of the ‘missing carbon’ supply needed for microbial growth and activity in the aphotic layer, while other sources of dissolved organic carbon remain to be elucidated. These findings underscore the need for further research into the factors affecting DCF, its role in global carbon budgets, and its potential to enhance atmospheric inorganic carbon sequestration.

1 Introduction

The oceans aphotic layers contain the world’s largest reservoir of dissolved inorganic carbon (DIC) (Baltar et al., 2010; Burd et al., 2010; Reinthaler et al., 2010), and harbor ~65% of all prokaryotes (Whitman et al., 1998). Aphotic prokaryotes typically rely on utilization of organic matter (and carbon), fixed by photoautotrophs via photosynthesis and exported from the euphotic zone, to sustain their growth and accumulate biomass (del Giorgio and Duarte, 2002). Current estimates reveal, however, a discrepancy between the supply of particulate organic carbon from photosynthesis and the bacterial organic carbon demand (BCD) in the aphotic zones (Ducklow, 2000; Karl et al., 1988; Smith and Azam, 1992). This mismatch suggests that there are other source/s of carbon that are being utilized by aphotic microorganisms (Baltar et al., 2009; Herndl and Reinthaler, 2013). One such source, that has been far less investigated, involves the fixation of DIC by chemo-autotrophic microbes and its assimilation into new biomass

(Baltar and Herndl, 2019). This could subsequently provide bioavailable DOC to other microbial populations at depth (Baltar et al., 2010).

DIC uptake by heterotrophic bacterioplankton is generally attributed to anaplerotic reactions (Dijkhuizen and Harder, 1984; Erb, 2011) which are metabolic pathways that replenish intermediates enzymes in the citric acid cycle by fixing CO₂, but other microorganisms such as nitrifying bacteria can also fix DIC (Alonso-Sáez et al., 2010). Genomic studies on deep-sea microbial communities identified several genes and metabolic pathways that enable some microbes to thrive as chemoautotrophs on inorganic substrates (Berg et al., 2007; Hallam et al., 2006). Measurements of CO₂ fixation by chemoautotrophs and heterotrophic bacterioplankton are scarce, yet substantial dark DIC fixation (DCF) rates have been reported in various oceanic settings and water masses (Swan et al. 2011; Zhou et al. 2017; La Cono et al. 2018; Alothman et al. 2023) and maybe more common than previously thought (Hansman et al., 2009; Herndl et al., 2005).

The deep waters of the southeast Mediterranean Sea are characterized by higher concentrations of inorganic nutrients compared to the photic zone (e.g., ~6 μmol NO₃+NO₂ kg⁻¹ and ~0.2 PO₄ μmol kg⁻¹ (Ben-Ezra et al., 2021; Sisma-Ventura et al., 2021) and low bioavailable dissolved organic carbon (Martínez-Pérez et al., 2017; Santinelli, 2015; Santinelli et al., 2010). Despite these characteristics, the southeast Mediterranean Sea's aphotic waters are considered a hotspot for bacterial activity compared to other oceanic regimes at similar depths (Luna et al., 2012; Rahav et al., 2019). Nutrient addition bioassays and water mixing simulations suggest that aphotic prokaryotes are primarily carbon-limited (Hazan et al., 2018; Rahav et al., 2019).

Here, we report on both photic and aphotic DCF and heterotrophic bacterial production rates from 6 cruises held between 2021-2023 in the southeastern Mediterranean (bottom

depth 1500-1750 m) during the mixed (winter) and stratified (summer) periods. Our results demonstrate that DCF rates cannot be neglected (contrary to past convention, Nielsen 1952) and are within the same order of magnitude as photosynthesis or heterotrophic bacterial production (BP). We also show that DCF substantially contributes to bacterial carbon demand (BCD), therefore providing, some of the ‘missing carbon’ supply needed for microbial growth and activity in the aphotic layer of the southeast Mediterranean Sea.

2 Material and methods

2.2 Sample collection - Seawater was collected during six seasonal cruises in the Levantine Basin, southeast Mediterranean Sea, on-board the R/V *Bat-Galim* between 2021-2023. Three cruises were held during the stratified period and three during the winter mixing. The mixed layer depth was calculated using a temperature difference of $\Delta 0.3$ °C (Mena et al., 2019). Two ‘deep’ stations were sampled in each cruise; one located at the edge of the continental shelf (H05 33.00 Lat, 34.50 Lon, bottom depth ~1500 m, 50 Km from the coast) and the other at the edge of Israel’s exclusive economic zone (H06 33.15 Lat, 34.16 Lon, bottom depth 1750 m, 90 Km from shoreline). Seawater was sampled at discrete depths throughout the water column, from the surface (~0.5 m) to the bottom (1500-1750 m) using Niskin bottles. Sampling depths were chosen in real-time based on reads of Conductivity Temperature Depth (CTD) (Seabird 19 Plus), chlorophyll fluorescence (Turner designs, Cyclops-7) and PAR (Sea Bird). The raw hydrological data can be freely downloaded from <https://isramar.ocean.org.il/isramar2009/>. Measurements included DIC ($\text{NaH}^{14}\text{CO}_3$) uptake under ambient light (hereafter ‘light primary productivity, LPP) or under full dark conditions (DCF), bacterial productivity (BP) and nutrient quantification.

Nitrite and ammonium concentrations – Samples for nitrite (NO_2^-) and ammonium (NH_4^+) concentrations were collected only in the 2023 cruises. The samples were pre-filtered ($0.45\ \mu\text{m}$), placed in acid-washed plastic vials, and were kept frozen at $-20\ ^\circ\text{C}$ until analysis. Nutrients were measured with a Seal Analytical AA-3 system. The limits of detection for NO_2^- and NH_4^+ were $0.06\ \mu\text{M}$ and $0.09\ \mu\text{M}$, respectively.

2.3 LPP and DCF - Seawater was collected in triplicates into transparent (for LPP measurements) or dark (for DCF) Nalgene bottles (45-250 ml) and spiked with $\text{NaH}^{14}\text{CO}_3$ (Perkin Elmer, specific activity $56\ \text{mCi mmol}^{-1}$) following Nielsen, (1952). The bottles were maintained in on-deck incubators covered with a gradient of neutral mesh simulating the irradiance intensity (no change in spectrum) at 100%, 50%, 10%, 1%, and 0.1% of surface light intensities or under complete dark conditions (Belkin et al., 2022; Reich et al., 2024). Incubators were kept at constant ambient surface temperatures ($\sim 19\text{-}20\ ^\circ\text{C}$ in winter and $\sim 28\text{-}29\ ^\circ\text{C}$ in the summer cruises). We acknowledge that temperature differences between surface and deeper depths may alter the LPP or DCF rates measurements, especially during the summer when the water column is stratified. While *in situ* measurements may offer more precise rate estimates, they are generally impractical during research cruises that involve sampling at multiple locations and times throughout the day and night. Nevertheless, preliminary comparisons between the incubation setup used here **versus** *in situ* incubations using a mooring line showed negligible differences in primary productivity, falling within the expected range of measurement variability (see also Reich et al., 2022). All the incubation bottles were spiked at sunrise and terminated after 24 h (Reich et al., 2022; Robinson et al., 2009) by filtering the particulate matter onto GF/F filters using low vacuum pressure ($<50\ \text{mmHg}$). Next, the excess ^{14}C -bicarbonate was removed by fuming with $50\ \mu\text{l}$ of 37% hydrochloric acid overnight. Finally, 5 ml scintillation

cocktail (ULTIMA-GOLD) was added, and the disintegrations per minute (DPM) from the particulate matter concentrated on the filters were counted using a TRI-CARB 4810 TR (Packard) liquid scintillation counter. Blank seawater spiked with $\text{NaH}^{14}\text{CO}_3$ was filtered immediately without incubation and the reads were subtracted from the sample's DPM. The blank DPM reads were usually negligible (<5% of the sample's DPM). Aliquots (50 μl) from random spiked samples were placed onto new GFF filters, added with 50 μl ethanolamine and scintillation liquid, and counted immediately without incubation to account for the 'added activity' of the radiolabeled working solution used. LPP was calculated as the difference between the DPM retrieved from the samples incubated under ambient light ('total primary production) and the 'dark' bottles. Dark or light dissolved inorganic carbon fixation was calculated based on the Bermuda Atlantic Time-series Study (BATS) protocol (<https://bios.asu.edu/bats/bats-data>). More details can be found in Reich et al. (2024).

2.3 Bacterial production- Triplicate samples per depth (1.7 ml) were incubated in the dark with 10 nmol/L ^3H -leucine L^{-1} (Perkin Elmer, specific activity 123 Ci mmol^{-1}) for 4-6 h under ambient temperature (Simon et al., 1990). The incubations were terminated with 100 μl of trichloroacetic acid (100%), processed as described by Smith and Azam (1992), and counted using a TRI-CARB 4810 TR (Packard) liquid scintillation counter. Killed control samples containing ^3H -leucine L^{-1} and trichloroacetic acid (without incubation) were also measured and these control sample's DPMs were subtracted from the sample's reads. A conversion factor of 3 kg C mol^{-1} per mole leucine incorporated was used, assuming an isotopic dilution of 2.0 (Simon and Azam 1989).

2.4 Molecular analyses and statistics - DNA was extracted from water samples with the PowerWater kit (Qiagen, USA), using the FastPrep-24TM Classic (MP Biomedicals,

USA) bead-beating to disrupt the cells (2 cycles at 5.5 m sec⁻¹, with a 5 min interval). The V4 region (~ 300 bp) of the 16S rRNA gene was amplified from the DNA (~50 ng) using the 515Fc/806Rc primers amended with relevant tags (Apprill et al., 2015; Parada et al., 2016). PCR conditions were as follows: initial denaturation at 94 °C for 45 s, 30 cycles of denaturation (94 °C for 15 sec), annealing (15 cycles at 50 °C and 15 cycles at 60 °C for 20 sec) and extension (72 °C for 30 s). Two annealing temperatures were used to account for the melting temperature of both forward (58.5-65.5 °C), and reverse (46.9-54.5 °C), primers.

Demultiplexed paired-end reads were processed in QIIME2 V2022.2 environment (Bolyen et al., 2019). Reads were truncated based on quality plots, checked for chimeras, merged and grouped into amplicon sequence variants (ASVs) with DADA2 (Callahan et al., 2016), as implemented in QIIME2. The amplicons were classified with Scikit-Learn classifier that was trained on Silva database v138 (16S rRNA, (Glöckner et al., 2017)). Mitochondrial and chloroplast sequences were removed from the 16S rRNA amplicon dataset. Downstream analyses were performed in R v4.1.1 (R Core Team, 2021), using packages Phyloseq (McMurdie and Holmes, 2013) and Ampvis2 (Andersen et al., 2018). Indicator species analyses were performed using Indic species package v1.7.9 (De Ca'ceres et al., 2009). Amplicon reads were deposited to the NCBI SRA archive under project number PRJNA1215023.

2.5 Bacterial respiration (BR), bacterial carbon demand (BCD) and zooplankton respiration (ZR) - BR was calculated based on the following equation and assuming an average open-ocean bacterial growth efficiency (BGE) of 20% (Herndl and Reinthaler, 2013) similar to previous direct measurements from the Mediterranean Sea ranging from 0.21-0.29 (Zweifel et al., 1993).

$$BGE = \frac{BP}{BP + BR}$$

BCD was then calculated as the sum of BP and BR (Gasol et al., 1998). Zooplankton respiration (ZR) and excretion were compiled from Belkin et al. (2022).

3 Results and discussion

3.1 Dark and light inorganic carbon fixation rates – As expected, LPP was restricted to the photic layer with highest rates usually measured at the surface (~0.5 m) that gradually decreased to reach minimum rates at the bottom of the photic layer (~180 m) (Fig 1A). Relatively low LPP values were measured during the stratified summer (~0.1-0.8 $\mu\text{g C L}^{-1} \text{ d}^{-1}$), whereas higher rates were measured during the winter mixing period (~0.2-7.4 $\mu\text{g C L}^{-1} \text{ d}^{-1}$) (Fig 1A). This resulted in ~10-fold higher integrated rates measured during the mixed period compared to those measured during the stratified period (Table 1), in accordance with studies from the area (Psarra et al., 2005; Reich et al., 2022; Sisma-Ventura et al., 2022b). In contrast with LPP, DCF was not restricted to the photic layer and ranged from 0 to ~0.4 $\mu\text{g C L}^{-1} \text{ d}^{-1}$ throughout the water column (Fig 1B, Fig 2A), without significant differences in the absolute rates between the photic and aphotic zones (t-test, $p > 0.05$, Fig 2B). The observed decrease in DCF rates with depth (Figure 2A) during the summer cruises may be partly attributed to a decline in the abundance of chemoautotrophs with depth. For example, Agogu   et al., (2008) reported a decline in archaeal *amoA* gene copy numbers with depth in the eastern North Atlantic. Normalizing DCF rates to chemoautotrophic microbial cell abundance (or gene copy) could reveal a different vertical pattern. Another possible explanation for the decline in DCF rates with depth may be related to the weakening flux of sinking organic matter with depth that limits the substrates that fuel DCF (discussion below). The integrated photic DCF was typically lower than the rates reported in the central and

western Mediterranean Sea (La Cono et al., 2018). The aphotic DCF rates were ~3.5 fold higher during the mixed than during the stratified period (Table 1, Fig 2A).




Figure 1: Spatial and temporal variability in rates of LPP (a), DCF (b), BP (c) and the contribution of DCF to bacterial carbon demand (BCD) (d) at the offshore SE Mediterranean Sea (Lat. 33.15 N, Lon. 34.16 E) between 2021-2023. BCD was calculated assuming a bacterial gross efficiency of 0.20 (Gasol et al., 1998).

Table 1: Integrated rates and contribution of DCF to metabolic processes in the photic (0-180) and aphotic (>180 m) depths of the pelagic southeast Mediterranean Sea. The values represent the minimum and maximum ranges observed across the cruises, with the averages and their corresponding standard deviations provided in parentheses. BDL = Below detection limit.

Variable	Season	Photic (0-180 m)	Aphotic (180-1750 m)
LPP (mg C m ⁻² d ⁻¹)	Mixed	158-649 (368 ± 205)	BDL
	Stratified	4-69 (32 ± 26)	BDL
DCF (mg C m ⁻² d ⁻¹)	Mixed	6-27 (15 ± 8)	17-342 (152 ± 127)

	Stratified	7-19 (14 ± 5)	8-127 (59 ± 48)
	Mixed	6-58 (28 ± 21)	12-65 (33 ± 22)
BP (mg C m ⁻² d ⁻¹)	Stratified	9-55 (30 ± 20)	7-123 (81 ± 55)
	Mixed	23-221 (109 ± 88)	49-594 (213 ± 200)
DCF contribution to BCD (%) *	Stratified	8-31 (18 ± 9)	8-42 (23 ± 13)
	Mixed	1-15 (6 ± 5)	---
DCF contribution to total PP (%)	Stratified	12-81 (40 ± 32)	---

* Assuming bacterial gross efficiency of 0.2 (Gasol et al., 1998) and that the available DOC for bacteria is 20% of the total primary productivity at the photic layer (Teira et al., 2003).

The higher aphotic DCF in the mixed versus the stratified periods may be related to more bioavailable carbon that is transported from the photic layer as marine snow and supplies organic carbon to heterotrophic activity in the winter (coinciding higher LPP). However, given the oligotrophic nature of the southeast Mediterranean Sea (Berman-Frank and Rahav, 2012; Reich et al., 2022), most of the organic carbon (both particulate and dissolved originating from LPP) is recycled within the photic layer. Only a small fraction fluxes down to the aphotic zone and has been recorded in sediment traps (Alkalay et al., 2024).

3.2 Bacterial productivity in relation to DCF and BCD - Another possible mechanism that may, potentially, explain higher DCF rates during the winter versus the summer is anaplerosis. The extent of anaplerotic reactions is primarily driven by the availability of labile organic carbon to heterotrophs (Dijkhuizen and Harder 1984). Therefore, assuming anaplerosis drives DCF, we expect it will be positively coupled to BP.

Yet, our results do not support the likelihood of significant anaplerosis reactions, predominantly evident from the spatiotemporal distribution of aphotic BP (Fig 1C) differing considerably from that of the DCF (Fig 1B) and does not correlate with it (Fig 3A). In fact, BP seems to be coupled with LPP at the photic layer reaching $\sim 0.4 \mu\text{g C L}^{-1} \text{ d}^{-1}$ (not shown). Excluding some sporadic measurements, aphotic BP rates were usually of similar magnitude and typically $< 0.1 \mu\text{g C L}^{-1} \text{ d}^{-1}$ (Fig 1C). Moreover, the highest integrated aphotic BP was measured during the summers of 2021 and 2022 and not during the winter cruises when generally higher DCF was recorded (Table 1, Fig 2A).




Figure 2: Dark carbon fixation rates in the southeastern Mediterranean Sea. Averaged vertical distribution of DCF in the offshore southeast Mediterranean Sea during the mixed (white) and stratified (gray) periods (a), and a box plot showing the DCF rates at the photic (0-180 m) and aphotic (>180 m) water depths (b).

Despite the lack of a clear positive relationship between DCF and BP, DCF may contribute to bacterial carbon demand (BCD) in the aphotic zone. Thus, we use a

244 literature standard, a bacterial growth efficiency of 0.20 (Gasol et al., 1998) to calculate
 245 BR and BCD (see the ‘material and methods’ section for more details). This calculation
 246 yielded bacterial respiration (BR) ranging from 29-494 mg C m⁻² d⁻¹ (average 209±172
 247 mg C m⁻² d⁻¹), and the concurrent BCD ranges from 36 to 648 mg C m⁻² d⁻¹ (average
 248 262±121 mg C m⁻² d⁻¹). Under these circumstances, exudation of DOC from primary
 249 productivity at the photic layer estimated as 20% of the rates (Teira et al., 2003) equals
 250 to ~1-130 mg C m⁻² d⁻¹.

251 This new DOC, that originated from the photic zone therefore cannot support the
 252 aphotic BCD in our system in all of our observations. However, if we consider the
 253 contribution of DOC produced by aphotic DCF, part of the missing carbon may be
 254 accounted for. Thus, when considering aphotic DCF in addition to the sequestered DOC
 255 from the photic layer, the ‘abnormally high’ aphotic BCD could be explained in full
 256 (≥100%) in ~35% of the observations (Fig 1D). In the other 65% of the observations
 257 the missing carbon sources needed to support the aphotic BCD remains an enigma. We
 258 note that these calculations are based on global averages and assumptions and therefore
 259 may be subject to some uncertainties. For example, BGE can vary between seasons and
 260 sites (del Giorgio and Cole, 1998). In the Mediterranean Sea, long-term measurements
 261 of BGE ranged from 0.21 (similar to our calculations and the global average used by
 262 Herndl and Reinthaler, 2013) to 0.29 (Zweifel et al., 1993). If the 0.29 value is used,
 263 the contribution of DCF to the aphotic BCD increases to ~45% of the observations
 264 rather than ~35% when using BGE of 0.2. Similarly, if we apply an exported DOC
 265 estimate of ~4% from the photic zone, as reported for the Ionian Sea/western
 266 Mediterranean (Moutin and Raimbault, 2002), the relative contribution of DCF to
 267 aphotic BCD would be even higher than in our current calculations, which assume
 268 ~20% DOC export (Teira et al., 2003). These uncertainties warrant future investigation.

Yet, even when using conservative estimates for BGE and DOC export as done here, the contribution of DCF to aphotic BCD remains substantial.

Evidence suggests that dissolved methane may be more abundant in oxygenated environments than previously thought (Grossart et al., 2011). Methane can potentially be one of the missing energy sources for marine microbes and support high BCD (Brankovits et al., 2017) as observed at the aphotic southeast Mediterranean Sea (Fig 1D). In agreement, methanotrophs were found in aphotic cold seeps at the southeast Mediterranean Sea (Sisma-Ventura et al., 2022a), as well as across the aphotic water column in our samples (see discussion below).

3.3 Interannual variability in aphotic DCF - Interannual variability in DCF, but not in LPP or BP, was observed with higher rates recorded in March 2021-March 2022 and lower rates observed in August 2022-August 2023 (Fig 1B). Inorganic nutrients such as PO_4^{3-} or $\text{NO}_3^- + \text{NO}_2^-$ are unlikely to explain this variability as their ambient levels were similar between periods (<https://isramar.ocean.org.il/isramar2009/>). Alternatively, we surmise that differences in the bioavailability and concentration of sinking organic particles, possibly attributed by the BiOS (Bimodal Oscillating System) oscillation circulation of deep water between the Adriatic and Ionian seas, could potentially explain the higher aphotic chemoautotrophic activity in March 2021-March 2022 versus August 2022-August 2023. This mechanism is known to influence the bioavailability of organic nutrients in the deep Mediterranean Sea by modulating deep-water circulation and ventilation patterns (Civitarese et al., 2010). These shifts affect the transport and residence time of organic matter (Civitarese et al., 2023), thereby potentially altering availability of organic nutrients to aphotic microbial populations, including to chemoautotrophs. Supporting this hypothesis are recent studies from the northern Red Sea and South China Sea showing that DCF is limited by labile organic

nutrients such as phosphonates and even carbon-rich molecules (Reich et al., 2024; Zhou et al., 2017). Aphotic free-living chemoautotrophs are likely to encounter an increasingly refractory pool of dissolved organic matter for metabolism that may result in lower DCF rates, as shown in exported material through the water column (Santinelli et al., 2013). Particle-attached chemoautotrophs may have access to higher concentrations of organic substrates. Therefore we surmise these microbes would preferentially have a particle-attached lifestyle in the deep ocean. The patchy nature of particulate matter sinking and lateral transport during wintertime (Alkalay et al., 2024) and aggregate concentrations (Bar-Zeev et al., 2012) in the deep southeast Mediterranean Sea could also potentially explain the interannual variability in DCF between periods. Understanding how chemoautotrophs transform labile dissolved organic matter into refractory dissolved organic matter, which is an essential process in the ‘microbial carbon pump’, is crucial as it influences the efficiency of the biological pump (Jiao et al., 2010).

Oxidizing reduced inorganic compounds as electron donors (e.g., NO_2^- or NH_4^+) may provide chemoautotrophic prokaryotes sufficient energy to fix DIC (Hügler and Sievert, 2011). We therefore measured the vertical distribution of NO_2^- and NH_4^+ (only in the March and August 2023 cruises) and examined if these chemical species are coupled or uncoupled with DCF at the aphotic zone. Our results show a negative-linear relationship between DCF and NO_2^- (Fig 3B) and NH_4^+ (Fig 3C), suggesting nitrification. This is because chemoautotrophs consume NO_2^- and NH_4^+ during nitrification to yield energy to fix DIC in the aphotic zone (REF), thereby reducing nutrient standing stocks in the water. We surmise that this ‘depletion’ may, theoretically, explain the observed negative correlation between nutrient levels and chemoautotrophic activity. In agreement, both ammonia and nitrite oxidizers were

found in the aphotic zone of all cruises (DNA level, discussion below), further highlighting their potential role as contributors to DCF in the southeast Mediterranean Sea. Nitrification measurements along with metagenomic tools, DCF (and BP) in aphotic water should be included in future dedicated studies to better refute or reinforce that oxidation of NO_2^- or NH_4^+ may provide chemoautotrophic prokaryotes the energy to fix DIC.




Figure 3: The relationship between aphotic DCF and BP (a), NO_2^- (b) and NH_4^+ (c). Note that NO_2^- and NH_4^+ was measured only during the 2023 cruises. The 95% confidence interval is shown in gray.

*3.4 Potential chemoautotrophs based on microbial community structure - Analyses of 16S rRNA gene amplicons suggest that diverse bacteria and archaea may drive DCF in the aphotic southeast Mediterranean Sea (Fig 4A). Microbes found in our collected genetic material primarily include the order Nitrosopumilales ammonia-oxidizing archaea, which become dominant below DCM (up to ~30% read abundance near the bottom), corresponding to previous estimates based on *in-situ* fluorescent hybridization (De Corte et al., 2009). Nitrite-oxidizing Nitrospirales comprised ~1% read abundance at depths below 300 m. Among these lineages, analyses of indicator species identified seasonal variation in abundance of the orders Nitrosopumilales and Nitrososphaeria that were more prominent in the stratified period than in wintertime (p-value<0.05)*

when the water column is mixed, while the deep-sea community in general exhibited only mild seasonal changes (Fig 4B). These ammonia oxidizers may thus drive ammonia depletion during summertime at the southeast Mediterranean Sea.

Additionally, we identified consistent occurrence of UBA10353 (Arenicellales, including the UBA868 group) and SAR324 gammaproteobacterial groups at depths below the deep chlorophyll maxima (Fig 4A). Members from these ubiquitous clades are mixotrophs that can fix inorganic carbon conserving energy from sulfur oxidation (Baltar et al., 2023; Jaffe et al., 2024; Swan et al., 2011). We did not find the SUP05/Arctic96BD-19 gammaproteobacterial sulfur oxidizers that occur in productive dark oxygenated waters (Swan et al., 2011). We note that Methylococcales methane oxidizers were typical in the near-bottom water layer, as found previously in the southeast Mediterranean Sea basin (Sisma-Ventura et al., 2022a; Techtmann et al., 2015), suggesting methylotrophy as a potential mechanism of 1-carbon molecule acquisition in the dark southeast Mediterranean Sea. While this DNA-based community analysis provides insight into the potential contributors to DCF, it does not reflect a direct link as we cannot determine which of the identified microbes are responsible for the measured DCF rates. We stress that future studies should examine the link between DCF and specific microbial groups such as archaea by targeting RNA-level expression and functional genes (e.g., *amoA* and *amoB*), as demonstrated by (Agogu   et al., 2008)




Figure 4: (a) Read abundance of the 40 most abundant taxa (order level) from different depths offshore the Southeastern Mediterranean Sea. Surface samples are within the range of 1-5 m depth; near surface are between 20 and deep chlorophyll maximum (DCM); below DCM corresponds to 180-240 m depths; near bottom samples were taken circa 5 meters above the seafloor. Potential DCF microbes are shown in blue, Methylococcales methanotrophs are marked in magenta, and photosynthetic Synechococcales are shown in green for reference. (b) A principal component analysis

showing the differences in the structure of microbial populations (40 most abundant taxa) based on 16S rRNA gene read mapping.

4 Conclusions

Based on the conceptual model in Figure 5 that summarizes the annual microbial carbon exchanges in the southeast Mediterranean sea's offshore area, DOC supplied by LPP is negligible and cannot explain the 'high' BCD in the area, especially in the aphotic zone that is considered a 'microbial hotspot' with relatively high bacterial activity per cell (Hazan et al., 2018; Rahav et al., 2019). Our observations suggest that DCF may provide a substantial amount of the missing carbon, at least in the southeast Mediterranean Sea, while source/s for the remaining missing carbon are currently unknown and warrant more research.




Figure 5: A schematic illustration showing microbial carbon exchange in the southeast Mediterranean Sea. Values shown are the annual averages of LPP, DCF and BP. DOC exudation from LPP was assumed to be 20% of the rates, BR was calculated from BP and assuming BGE=0.20, Zooplankton's respiration (ZR) and excretion were compiled from Belkin et al. (2022). The numbers in brackets show the integrated values over 1750 m and expressed as $\text{g C m}^{-2} \text{ y}^{-1}$.

Regardless of the yet missing DOC sources, our results demonstrate the pivotal role DCF plays in compensating metabolic imbalances in carbon sources at the aphotic southeast Mediterranean Sea. Similarly, DCF was shown to be a significant process supporting microbial respiration and/or activity in aphotic layers (Baltar et al., 2009; Herndl et al., 2005; Yakimov et al., 2011), as well as in hydrothermal vents (Mattes et al., 2013) and cold seeps (Nakagawa et al., 2007). Note that, while our estimates of DCF contribution to aphotic BCD are based on widely accepted assumptions, they are subject to some uncertainties, particularly regarding BGE that may be changed on both spatial and temporal scales, as well as the fraction of DOC exported from the photic zone that may also change between seasons and water provinces (see discussion below). These uncertainties underscore the need for more precise and region-specific measurements of BGE and DOC fluxes to better constrain the role of DCF in deep ocean carbon cycling.

Another potential uncertainty in measuring aphotic metabolic rates such as DCF or BP lies in the unclear effects of hydrostatic pressure on the activity of bulk microbial communities (Riebesell et al., 2009; Tamburini et al., 2013). Laboratory-based manipulations that do not account for *in situ* pressure conditions may alter DCF rates, potentially misrepresenting the actual contribution of chemoautotrophs to aphotic

BCD. This highlights the urgent need for more detailed investigations into how hydrostatic pressure influences microbial activity in the deep ocean.

Additionally, despite the contribution of DCF to the DOC pool (taking into account the uncertainties associated with it discussed above), as well as the other sources, very little of fixed carbon as particulate organic matter (POC) ends up in sediment traps located above the seabed (2-6%). This suggests that most of the fixed carbon arriving from DCF (as well as LPP other potential sources) is recycled in the water column and does not reach the seabed. The rapid microbial recycling of nutrients was mostly investigated in the photic layer of the southeast Mediterranean Sea (e.g., PO₄, Thingstad et al., 2005) and little is known about the processes, which are often cryptic (e.g., NO₂⁻ oxygenation), occurring in the aphotic layers.

Understanding the ‘dark end’ of the biological pump in oligotrophic oceans, which plays an important (yet variable) role in oceanic carbon cycling and sequestration, will require a multidisciplinary approach that take into account all the uncertainties discussed above in light of our (and others) observations of DCF, especially in the context of ongoing and significant changes in the marine environment. Our study supports the need for adding DCF measurements to global carbon budgets as also mentioned by Baltar and Herndl (2010).

Data availability statement

The raw physiological data used to generate figures 1-3 can be downloaded from the PANGAEA repository: Reich, Tom (2025): Primal, chemo and bacterial productivity coupled with inorganic nutrient concentrations from an off shore, outgoing transect cruises: <https://doi.org/10.1594/PANGAEA.975231>. Hydrological data can be

downloaded from <https://isramar.ocean.org.il/isramar2009/>. Genomic data can be
downloaded from the NCBI SRA archive project number PRJNA1215023.

Author contribution

TR: Formal experimental work and data analyses; Investigation; Data Curation;
Visualization; Writing – Original Draft Preparation. NB: Resources; Writing –
Review & Editing. GSV: Investigation; Writing – Review & Editing. HH:
Investigation. MRB: Investigation; Writing – Review. IBF: Conceptualization;
Funding Acquisition; Supervision; data interpretation, Writing – Review & Editing.
ER: Conceptualization; Formal Analysis; Funding Acquisition; Supervision;
Visualization; Writing – Review & Editing.

Competing interests

The authors declare that they have no conflict of interest.

Acknowledgements

We would like to thank the University of Haifa for supporting T. Reich, and the
Helmholtz funded International Laboratory; The Eastern Mediterranean Sea Centre-
An Early-Warning Model-System for our Future Oceans: EMS Future Ocean
Research (EMS FORE).

This study is in partial fulfillment of the PhD requirements for T. Reich at the Leon
H. Charney School of Marine Sciences, at the University of Haifa.

450 **Statements and Declarations**

451 The authors declare that they have no conflict of interest.

452

453 **Funding**

454 This work was supported by the German-Israeli Foundation for Scientific Research
455 and Development 2016021

456

457 **References**

- 458 Agogu , H., Brink, M., Dinasquet, J., and Herndl, G. J.: Major gradients in putatively
459 nitrifying and non-nitrifying Archaea in the deep North Atlantic, *Nature*, 456, 788–
460 792, <https://doi.org/10.1038/nature07535>, 2008.
- 461 Alkalay, R., Weinstein, Y., Herut, B., Ozer, T., Zlatkin, O., Bar, T., Berman-Frank, I.,
462 and Katz, T.: Temporal Pattern and Profile of a Coastal-Deep Sea Conveyor at a
463 Marginal Deep Oligotrophic Sea, *J Geophys Res Oceans*, 129,
464 <https://doi.org/10.1029/2023JC020441>, 2024.
- 465 Alonso-S  ez, L., Galand, P. E., Casamayor, E. O., Pedr  s-Ali  , C., and Bertilsson, S.:
466 High bicarbonate assimilation in the dark by Arctic bacteria, *ISME Journal*, 4, 1581–
467 1590, <https://doi.org/10.1038/ismej.2010.69>, 2010.
- 468 Alothman, A., L  pez-Sandoval, D., Duarte, C. M., and Agust  , S.: Bacterioplankton
469 dark CO₂ fixation in oligotrophic waters, *Biogeosciences Discussions*,
470 <https://doi.org/10.5194/bg-2023-65>, 2023.
- 471 Andersen, K. S., Kirkegaard, R. H., Karst, S. M., and Albertsen, M.: ampvis2: an R
472 package to analyse and visualise 16S rRNA amplicon data,
473 <https://doi.org/10.1101/299537>, 11 April 2018.
- 474 Apprill, A., McNally, S., Parsons, R., and Weber, L.: Minor revision to V4 region
475 SSU rRNA 806R gene primer greatly increases detection of SAR11 bacterioplankton,
476 *Aquatic Microbial Ecology*, 75, 129–137, <https://doi.org/10.3354/ame01753>, 2015.
- 477 Baltar, F. and Herndl, G. J.: Ideas and perspectives: Is dark carbon fixation relevant
478 for oceanic primary production estimates?, *Biogeosciences*, 16, 3793–3799,
479 <https://doi.org/10.5194/bg-16-3793-2019>, 2019.
- 480 Baltar, F., Ar  stegui, J., Gasol, J. M., Sintes, E., and Herndl, G. J.: Evidence of
481 prokaryotic metabolism on suspended particulate organic matter in the dark waters of
482 the subtropical North Atlantic, *Limnol Oceanogr*, 54, 182–193,
483 <https://doi.org/10.4319/lo.2009.54.1.0182>, 2009.
- 484 Baltar, F., Ar  stegui, J., Sintes, E., Gasol, J. M., Reinthaler, T., and Herndl, G. J.:
485 Significance of non-sinking particulate organic carbon and dark CO₂ fixation to
486 heterotrophic carbon demand in the mesopelagic northeast Atlantic, *Geophys Res*
487 *Lett*, 37, <https://doi.org/https://doi.org/10.1029/2010GL043105>, 2010.
- 488 Baltar, F., Mart  nez-P  rez, C., Amano, C., Vial, M., Robaina-Est  vez, S., Reinthaler,
489 T., Herndl, G. J., Zhao, Z., Logares, R., Morales, S. E., and Gonz  lez, J. M.: A
490 ubiquitous gammaproteobacterial clade dominates expression of sulfur oxidation
491 genes across the mesopelagic ocean, *Nat Microbiol*, 8, 1137–1148,
492 <https://doi.org/10.1038/s41564-023-01374-2>, 2023.
- 493 Bar-Zeev, E., Berman-Frank, I., Girshevitz, O., and Berman, T.: Revised paradigm of
494 aquatic biofilm formation facilitated by microgel transparent exopolymer particles,
495 *Proc Natl Acad Sci U S A*, 109, 9119–9124,
496 <https://doi.org/10.1073/pnas.1203708109>, 2012.

497 Bar-Zeev, E., Passow, U., Romero-Vargas Castrillón, S., and Elimelech, M.:
 498 Transparent exopolymer particles: From aquatic environments and engineered
 499 systems to membrane biofouling, <https://doi.org/10.1021/es5041738>, 20 January
 500 2015.

501 Belkin, N., Guy-haim, T., Rubin-blum, M., Lazar, A., and Sisma-, G.: Influence of
 502 cyclonic and anti-cyclonic eddies on plankton biomass , activity and diversity in the
 503 southeastern Mediterranean Sea, 1–56, 2022.

504 Ben-Ezra, T., Krom, M. D., Tsemel, A., Berman-Frank, I., Herut, B., Lehahn, Y.,
 505 Rahav, E., Reich, T., Thingstad, T. F., and Sher, D.: Deep-Sea Research Part I
 506 Seasonal nutrient dynamics in the P depleted Eastern Mediterranean Sea, Deep-Sea
 507 Research Part I, 176, 103607, <https://doi.org/10.1016/j.dsr.2021.103607>, 2021.

508 Berg, I. A., Kockelkorn, D., Buckel, W., and Fuchs, G.: A 3-hydroxypropionate/4-
 509 hydroxybutyrate autotrophic carbon dioxide assimilation pathway in archaea, *Science*
 510 (1979), 318, 1782–1786, <https://doi.org/10.1126/science.1149976>, 2007.

511 Berman-Frank, I. and Rahav, E.: Dinitrogen fixation as a source for new production in
 512 the Mediterranean Sea: A review, *Life in the Mediterranean Sea: A Look at Habitat*
 513 *Changes*, 199–226, 2012.

514 Bolyen, E., Rideout, J. R., Dillon, M. R., Bokulich, N. A., Abnet, C. C., Al-Ghalith,
 515 G. A., Alexander, H., Alm, E. J., Arumugam, M., Asnicar, F., Bai, Y., Bisanz, J. E.,
 516 Bittinger, K., Brejnrod, A., Brislawn, C. J., Brown, C. T., Callahan, B. J., Caraballo-
 517 Rodríguez, A. M., Chase, J., Cope, E. K., Da Silva, R., Diener, C., Dorrestein, P. C.,
 518 Douglas, G. M., Durall, D. M., Duvallet, C., Edwardson, C. F., Ernst, M., Estaki, M.,
 519 Fouquier, J., Gauglitz, J. M., Gibbons, S. M., Gibson, D. L., Gonzalez, A., Gorlick,
 520 K., Guo, J., Hillmann, B., Holmes, S., Holste, H., Huttenhower, C., Huttley, G. A.,
 521 Janssen, S., Jarmusch, A. K., Jiang, L., Kaehler, B. D., Kang, K. Bin, Keefe, C. R.,
 522 Keim, P., Kelley, S. T., Knights, D., Koester, I., Kosciulek, T., Kreps, J., Langille, M.
 523 G. I., Lee, J., Ley, R., Liu, Y. X., Loftfield, E., Lozupone, C., Maher, M., Marotz, C.,
 524 Martin, B. D., McDonald, D., McIver, L. J., Melnik, A. V., Metcalf, J. L., Morgan, S.
 525 C., Morton, J. T., Naimey, A. T., Navas-Molina, J. A., Nothias, L. F., Orchanian, S.
 526 B., Pearson, T., Peoples, S. L., Petras, D., Preuss, M. L., Priesse, E., Rasmussen, L.
 527 B., Rivers, A., Robeson, M. S., Rosenthal, P., Segata, N., Shaffer, M., Shiffer, A.,
 528 Sinha, R., Song, S. J., Spear, J. R., Swofford, A. D., Thompson, L. R., Torres, P. J.,
 529 Trinh, P., Tripathi, A., Turnbaugh, P. J., Ul-Hasan, S., van der Hooft, J. J. J., Vargas,
 530 F., Vázquez-Baeza, Y., Vogtmann, E., von Hippel, M., et al.: Reproducible,
 531 interactive, scalable and extensible microbiome data science using QIIME 2,
 532 <https://doi.org/10.1038/s41587-019-0209-9>, 1 August 2019.

533 Brankovits, D., Pohlman, J. W., Niemann, H., Leigh, M. B., Leewis, M. C., Becker,
 534 K. W., Iliffe, T. M., Alvarez, F., Lehmann, M. F., and Phillips, B.: Methane-and
 535 dissolved organic carbon-fueled microbial loop supports a tropical subterranean
 536 estuary ecosystem, *Nat Commun*, 8, <https://doi.org/10.1038/s41467-017-01776-x>,
 537 2017.

538 Burd, A. B., Hansell, D. A., Steinberg, D. K., Anderson, T. R., Aristegui, J., Baltar,
 539 F., Beaupré, S. R., Buesseler, K. O., DeHairs, F., Jackson, G. A., Kadko, D. C.,

540 Koppelman, R., Lampitt, R. S., Nagata, T., Reinthaler, T., Robinson, C., Robison, B.
 541 H., Tamburini, C., and Tanaka, T.: Assessing the apparent imbalance between
 542 geochemical and biochemical indicators of meso- and bathypelagic biological
 543 activity: What the @\$#! is wrong with present calculations of carbon budgets?, *Deep*
 544 *Sea Res 2 Top Stud Oceanogr*, 57, 1557–1571,
 545 <https://doi.org/10.1016/j.dsr2.2010.02.022>, 2010.

546 De Ca'ceres, M., Ca'ceres, C., and Legendre, P.: Associations between species and
 547 groups of sites: indices and statistical inference, *Ecology*, 3566–3574 pp., 2009.

548 Callahan, B. J., McMurdie, P. J., Rosen, M. J., Han, A. W., Johnson, A. J. A., and
 549 Holmes, S. P.: DADA2: High-resolution sample inference from Illumina amplicon
 550 data, *Nat Methods*, 13, 581–583, <https://doi.org/10.1038/nmeth.3869>, 2016.

551 Civitarese, G., Gačić, M., Lipizer, M., and Eusebi Borzelli, G. L.: On the impact of
 552 the Bimodal Oscillating System (BiOS) on the biogeochemistry and biology of the
 553 Adriatic and Ionian Seas (Eastern Mediterranean), *Biogeosciences*, 7, 3987–3997,
 554 <https://doi.org/10.5194/bg-7-3987-2010>, 2010.

555 Civitarese, G., Gačić, M., Batistić, M., Bensi, M., Cardin, V., Dulčić, J., Garić, R.,
 556 and Menna, M.: The BiOS mechanism: History, theory, implications,
 557 <https://doi.org/10.1016/j.pocean.2023.103056>, 1 August 2023.

558 La Cono, V., Ruggeri, G., Azzaro, M., Crisafi, F., Decembrini, F., Denaro, R., La
 559 Spada, G., Maimone, G., Monticelli, L. S., Smedile, F., Giuliano, L., and Yakimov,
 560 M. M.: Contribution of bicarbonate assimilation to carbon pool dynamics in the deep
 561 Mediterranean Sea and cultivation of actively nitrifying and CO₂-fixing bathypelagic
 562 prokaryotic consortia, *Front Microbiol*, 9, <https://doi.org/10.3389/fmicb.2018.00003>,
 563 2018.

564 De Corte, D., Yokokawa, T., Varela, M. M., Agogué, H., and Herndl, G. J.: Spatial
 565 distribution of Bacteria and Archaea and amoA gene copy numbers throughout the
 566 water column of the Eastern Mediterranean Sea, *ISME Journal*, 3, 147–158,
 567 <https://doi.org/10.1038/ismej.2008.94>, 2009.

568 Dijkhuizen, L. and Harder, W.: Current views on the regulation of autotrophic carbon
 569 dioxide fixation via the Calvin cycle in bacteria, *Antonie Van Leeuwenhoek*, 50, 473–
 570 487, 1984.

571 Ducklow, H. W.: Bacterial Production and Biomass in the Oceans., *Microbial*
 572 *Ecology of the Oceans.*, 85–119, 2000.

573 Erb, T. J.: Carboxylases in natural and synthetic microbial pathways,
 574 <https://doi.org/10.1128/AEM.05702-11>, December 2011.

575 Gasol, J. M., Doval, M. D., Pinhassi, J., Calderón-Paz, J. I., Guixa-Boixareu, N.,
 576 Vaqué, D., and Pedrós-Alió, C.: Diel variations in bacterial heterotrophic activity and
 577 growth in the northwestern Mediterranean Sea, *Mar Ecol Prog Ser*, 164, 107–124,
 578 <https://doi.org/10.3354/meps164107>, 1998.

del Giorgio, P. A. and Cole, J. J.: Bacterial growth efficiency in natural aquatic systems, *Annu Rev Ecol Syst*, 29, 503–541, <https://doi.org/10.1146/annurev.ecolsys.29.1.503>, 1998.

del Giorgio, P. A. and Duarte, C. M.: Respiration in the open ocean, *Nature*, 420, 379–384, <https://doi.org/10.1038/nature01165>, 2002.

Glöckner, F. O., Yilmaz, P., Quast, C., Gerken, J., Beccati, A., Ciuprina, A., Bruns, G., Yarza, P., Peplies, J., Westram, R., and Ludwig, W.: 25 years of serving the community with ribosomal RNA gene reference databases and tools, <https://doi.org/10.1016/j.jbiotec.2017.06.1198>, 10 November 2017.

Grossart, H. P., Frindte, K., Dziallas, C., Eckert, W., and Tang, K. W.: Microbial methane production in oxygenated water column of an oligotrophic lake, *Proc Natl Acad Sci U S A*, 108, 19657–19661, <https://doi.org/10.1073/pnas.1110716108>, 2011.

Hallam, S. J., Mincer, T. J., Schleper, C., Preston, C. M., Roberts, K., Richardson, P. M., and DeLong, E. F.: Pathways of carbon assimilation and ammonia oxidation suggested by environmental genomic analyses of marine Crenarchaeota, *PLoS Biol*, 4, 520–536, <https://doi.org/10.1371/journal.pbio.0040095>, 2006.

Hansman, R. L., Griffin, S., Watson, J. T., Druffel, E. R. M., Ingalls, A. E., Pearson, A., and Aluwihare, L. I.: The radiocarbon signature of microorganisms in the mesopelagic ocean, *Proceedings of the National Academy of Sciences*, 106, 6513–6518, <https://doi.org/10.1073/pnas.0810871106>, 2009.

Hazan, O., Silverman, J., Sisma-Ventura, G., Ozer, T., Gertman, I., Shoham-Frider, E., Kress, N., and Rahav, E.: Mesopelagic prokaryotes alter surface phytoplankton production during simulated deep mixing experiments in Eastern Mediterranean Sea waters, *Front Mar Sci*, 5, 1–11, <https://doi.org/10.3389/fmars.2018.00001>, 2018.

Herndl, G. J. and Reinthaler, T.: Microbial control of the dark end of the biological pump, <https://doi.org/10.1038/ngeo1921>, September 2013.

Herndl, G. J., Reinthaler, T., Teira, E., Van Aken, H., Veth, C., Pernthaler, A., and Pernthaler, J.: Contribution of Archaea to total prokaryotic production in the deep atlantic ocean, *Appl Environ Microbiol*, 71, 2303–2309, <https://doi.org/10.1128/AEM.71.5.2303-2309.2005>, 2005.

Hügler, M. and Sievert, S. M.: Beyond the Calvin cycle: Autotrophic carbon fixation in the ocean, *Ann Rev Mar Sci*, 3, 261–289, <https://doi.org/10.1146/annurev-marine-120709-142712>, 2011.

Jaffe, A. L., Salcedo, R. S. R., and Dekas, A. E.: Abundant and metabolically flexible lineages within the SAR324 and gammaproteobacteria dominate the potential for rubisco-mediated carbon fixation in the dark ocean, <https://doi.org/10.1101/2024.05.09.593449>, 10 May 2024.

Jiao, N., Herndl, G. J., Hansell, D. A., Benner, R., Kattner, G., Wilhelm, S. W., Kirchman, D. L., Weinbauer, M. G., Luo, T., Chen, F., and Azam, F.: Microbial production of recalcitrant dissolved organic matter: Long-term carbon storage in the global ocean, <https://doi.org/10.1038/nrmicro2386>, August 2010.

620 Karl, D. M., Knauer, G. A., and Martin, J. H.: Downward flux of particulate organic
621 matter in the ocean: a particle decomposition paradox, *Nature*, 332, 438–441,
622 <https://doi.org/10.1038/332438a0>, 1988.

623 Li Zweifel, U., Norrman, B., and Hagström, Å.: Consumption of dissolved organic
624 carbon by marine bacteria and demand for inorganic nutrients, *Source: Marine*
625 *Ecology Progress Series*, 23–32 pp., 1993.

626 Luna, G. M., Bianchelli, S., Decembrini, F., De Domenico, E., Danovaro, R., and
627 Dell’Anno, A.: The dark portion of the Mediterranean Sea is a bioreactor of organic
628 matter cycling, *Global Biogeochem Cycles*, 26,
629 <https://doi.org/https://doi.org/10.1029/2011GB004168>, 2012.

630 Martínez-Pérez, A. M., Álvarez-Salgado, X. A., Aristegui, J., and Nieto-Cid, M.:
631 Deep-ocean dissolved organic matter reactivity along the Mediterranean Sea: Does
632 size matter, *Sci Rep*, 7, <https://doi.org/10.1038/s41598-017-05941-6>, 2017.

633 Mattes, T. E., Nunn, B. L., Marshall, K. T., Proskurowski, G., Kelley, D. S., Kawka,
634 O. E., Goodlett, D. R., Hansell, D. A., and Morris, R. M.: Sulfur oxidizers dominate
635 carbon fixation at a biogeochemical hot spot in the dark ocean, *ISME Journal*, 7,
636 2349–2360, <https://doi.org/10.1038/ismej.2013.113>, 2013.

637 McMurdie, P. J. and Holmes, S.: Phyloseq: An R Package for Reproducible
638 Interactive Analysis and Graphics of Microbiome Census Data, *PLoS One*, 8,
639 <https://doi.org/10.1371/journal.pone.0061217>, 2013.

640 Mena, C., Reglero, P., Hidalgo, M., Sintés, E., Santiago, R., Martín, M., Moyà, G.,
641 and Balbín, R.: Phytoplankton Community Structure Is Driven by Stratification in the
642 Oligotrophic Mediterranean Sea, *Front Microbiol*, 10,
643 <https://doi.org/10.3389/fmicb.2019.01698>, 2019.

644 Moutin, T. and Raimbault, P.: Primary production, carbon export and nutrients
645 availability in western and eastern Mediterranean Sea in early summer 1996 (MINOS
646 cruise), *Journal of Marine Systems*, 33, 273–288, 2002.

647 Nakagawa, T., Mori, K., Kato, C., Takahashi, R., and Tokuyama, T.: Distribution of
648 Cold-Adapted Ammonia-Oxidizing Microorganisms in the Deep-Ocean of the
649 Northeastern Japan Sea, *Microbes Environ*, 365–372 pp., 2007.

650 Nielsen, E. S.: The use of radio-active carbon (C14) for measuring organic production
651 in the sea, *ICES Journal of Marine Science*, 18, 117–140, 1952.

652 Parada, A. E., Needham, D. M., and Fuhrman, J. A.: Every base matters: Assessing
653 small subunit rRNA primers for marine microbiomes with mock communities, time
654 series and global field samples, *Environ Microbiol*, 18, 1403–1414,
655 <https://doi.org/10.1111/1462-2920.13023>, 2016.

656 Psarra, S., Zohary, T., Krom, M. D., Mantoura, R. F. C., Polychronaki, T., Stambler,
657 N., Tanaka, T., Tselepides, A., and Frede Thingstad, T.: Phytoplankton response to a
658 Lagrangian phosphate addition in the Levantine Sea (Eastern Mediterranean), *Deep*
659 *Sea Res 2 Top Stud Oceanogr*, 52, 2944–2960,
660 <https://doi.org/10.1016/j.dsr2.2005.08.015>, 2005.

661 Rahav, E., Silverman, J., Raveh, O., Hazan, O., Rubin-Blum, M., Zeri, C., Gogou, A.,
662 Kralj, M., Pavlidou, A., and Kress, N.: The deep water of Eastern Mediterranean Sea
663 is a hotspot for bacterial activity, *Deep Sea Res 2 Top Stud Oceanogr*, 164, 135–143,
664 <https://doi.org/10.1016/j.dsr2.2019.03.004>, 2019.

665 Reich, T., Ben-Ezra, T., Belkin, N., Tsemel, A., Aharonovich, D., Roth-Rosenberg,
666 D., Givati, S., Bialik, M., Herut, B., Berman-Frank, I., Frada, M., Krom, M. D.,
667 Lehahn, Y., Rahav, E., and Sher, D.: A year in the life of the Eastern Mediterranean:
668 Monthly dynamics of phytoplankton and bacterioplankton in an ultra-oligotrophic
669 sea, *Deep Sea Research Part I: Oceanographic Research Papers*, 182, 103720,
670 <https://doi.org/https://doi.org/10.1016/j.dsr.2022.103720>, 2022.

671 Reich, T., Belkin, N., Sisma-Ventura, G., Berman-Frank, I., and Rahav, E.:
672 Significant dark inorganic carbon fixation in the euphotic zone of an oligotrophic sea,
673 *Limnol Oceanogr*, 69, 1129–1142, <https://doi.org/10.1002/lno.12560>, 2024.

674 Reinthaler, T., van Aken, H. M., and Herndl, G. J.: Major contribution of autotrophy
675 to microbial carbon cycling in the deep North Atlantic's interior, *Deep Sea Res 2 Top*
676 *Stud Oceanogr*, 57, 1572–1580, <https://doi.org/10.1016/j.dsr2.2010.02.023>, 2010.

677 Riebesell, U., Kö, A., and Oschlies, A.: Sensitivities of marine carbon fluxes to ocean
678 change, 2009.

679 Robinson, C., Tilstone, G. H., Rees, A. P., Smyth, T. J., Fishwick, J. R., Tarran, G.
680 A., Luz, B., Barkan, E., and David, E.: Comparison of in vitro and in situ plankton
681 production determinations, *Aquatic Microbial Ecology*, 54, 13–34,
682 <https://doi.org/10.3354/ame01250>, 2009.

683 Santinelli, C.: DOC in the Mediterranean Sea, in: *Biogeochemistry of marine*
684 *dissolved organic matter*, Elsevier, 579–608, 2015.

685 Santinelli, C., Nannicini, L., and Seritti, A.: DOC dynamics in the meso and
686 bathypelagic layers of the Mediterranean Sea, *Deep Sea Res 2 Top Stud Oceanogr*,
687 57, 1446–1459, <https://doi.org/10.1016/j.dsr2.2010.02.014>, 2010.

688 Simon, M., Alldredge, A., and Azam, F.: Bacterial carbon dynamics on marine snow,
689 *Mar Ecol Prog Ser*, 65, 205–211, 1990.

690 Sisma-Ventura, G., Kress, N., Silverman, J., Gertner, Y., Ozer, T., Biton, E., Lazar,
691 A., Gertman, I., Rahav, E., and Herut, B.: Post-eastern Mediterranean Transient
692 Oxygen Decline in the Deep Waters of the Southeast Mediterranean Sea Supports
693 Weakening of Ventilation Rates, *Front Mar Sci*, 7,
694 <https://doi.org/10.3389/fmars.2020.598686>, 2021.

695 Sisma-Ventura, G., Bialik, O. M., Makovsky, Y., Rahav, E., Ozer, T., Kanari, M.,
696 Marmen, S., Belkin, N., Guy-Haim, T., Antler, G., Herut, B., and Rubin-Blum, M.:
697 Cold seeps alter the near-bottom biogeochemistry in the ultraoligotrophic
698 Southeastern Mediterranean Sea, *Deep Sea Research Part I: Oceanographic Research*
699 *Papers*, 183, 103744, <https://doi.org/https://doi.org/10.1016/j.dsr.2022.103744>, 2022a.

700 Sisma-Ventura, G., Belkin, N., Rubin-Blum, M., Jacobson, Y., Hauzer, H., Bar-Zeev,
701 E., and Rahav, E.: Discharge of polyphosphonate-based antiscalants via desalination

brine: impact on seabed nutrient flux and microbial activity, *Environ Sci Technol*,
<https://doi.org/10.1021/acs.est.2c04652>, 2022b.

Smith, D. and Azam, F.: A simple, economical method for measuring bacterial
protein synthesis rates in seawater using, *Marine microbial food webs*, 6, 107–114,
1992.

Swan, B. K., Martinez-Garcia, M., Preston, C. M., Sczyrba, A., Woyke, T., Lamy, D.,
Reinthal, T., Poulton, N. J., Masland, E. D. P., Gomez, M. L., Sieracki, M. E.,
DeLong, E. F., Herndl, G. J., and Stepanauskas, R.: Potential for
chemolithoautotrophy among ubiquitous bacteria lineages in the dark ocean, *Science*
(1979), 333, 1296–1300, <https://doi.org/10.1126/science.1203690>, 2011.

Tamburini, C., Boutrif, M., Garel, M., Colwell, R. R., and Deming, J. W.: Prokaryotic
responses to hydrostatic pressure in the ocean - a review,
<https://doi.org/10.1111/1462-2920.12084>, May 2013.

Techtmann, S. M., Fortney, J. L., Ayers, K. A., Joyner, D. C., Linley, T. D., Pfiffner,
S. M., and Hazen, T. C.: The unique chemistry of Eastern Mediterranean water
masses selects for distinct microbial communities by depth, *PLoS One*, 10,
<https://doi.org/10.1371/journal.pone.0120605>, 2015.

Teira, E., Pazó, M., Quevedo, M., Fuentes, M., Niell, F., and Fernández, E.: Rates of
dissolved organic carbon production and bacterial activity in the eastern North
Atlantic Subtropical Gyre during summer, *Mar Ecol Prog Ser*, 249, 53–67,
<https://doi.org/10.3354/meps249053>, 2003.

Thingstad, T. F., Krom, M. D., Mantoura, R. F. C., Flaten, G. A. F., Groom, S., Herut,
B., Kress, N., Law, C. S., Pasternak, A., and Pitta, P.: Nature of phosphorus limitation
in the ultraoligotrophic eastern Mediterranean, *Science* (1979), 309, 1068–1071,
2005.

Whitman, W. B., Coleman, D. C., and Wiebe, W. J.: Perspective Prokaryotes: The
unseen majority, 6578–6583 pp., 1998.

Yakimov, M. M., Cono, V. La, Smedile, F., Deluca, T. H., Juárez, S., Ciordia, S.,
Fernández, M., Albar, J. P., Ferrer, M., Golyshin, P. N., and Giuliano, L.:
Contribution of crenarchaeal autotrophic ammonia oxidizers to the dark primary
production in Tyrrhenian deep waters (Central Mediterranean Sea), *ISME Journal*, 5,
945–961, <https://doi.org/10.1038/ismej.2010.197>, 2011.

Zhou, W., Li, Y., Liu, X., He, S., and Huang, J. C.: Comparison of microbial
communities in different sulfur-based autotrophic denitrification reactors, *Appl*
Microbiol Biotechnol, 101, 447–453, <https://doi.org/10.1007/s00253-016-7912-y>,
2017.