

Supplement to Technical note: Apportionment of Southeast Asian Biomass Burning and Urban Influence via In Situ Trace Gas Enhancement Ratios

Joshua P. DiGangi¹, Glenn S. Diskin¹, Subin Yoon^{2,*}, Sergio L. Alvarez², James H. Flynn², Claire E. Robinson^{1,3,†}, Michael A. Shook¹, K. Lee Thornhill^{1,3}, Edward L. Winstead^{1,3}, Luke D. Ziemba¹, Maria O. Cambaliza^{4,5}, James B. Simpas^{4,5}, Miguel Ricardo A. Hilario⁶, Armin Sorooshian⁶

¹NASA Langley Research Center, Hampton, VA, 23681, USA

²University of Houston, Houston, TX, 77204, USA

³Analytical Mechanics Associates, Inc., Hampton, VA, 23666, USA

⁴Air Quality Dynamics Laboratory, Manila Observatory, Quezon City, 1101, Philippines

⁵Department of Physics, Ateneo de Manila University, Quezon City, 1101, Philippines

⁶University of Arizona, Tucson, AZ, 85721, USA

*Now at: Scripps Institution of Oceanography, University of California San Diego, La Jolla, CA, 92037, USA

†deceased

Correspondence to: Joshua P. DiGangi (joshua.p.digangi@nasa.gov)




Figure S1: Example of hysteresis in identifying urban influence. Dashed lines represent hysteresis cutoffs.




Figure S2: Highlighted data from the high ozone flag (in red). Scatterplots of (a) CH₄ vs CO, (b) O₃ vs H₂O(v), (c) O₃ vs CO, (d) O₃ vs CH₄.