

Track-changes document for egusphere-2025-1464

Valentin Bruch

October 24, 2025

Normal tracking of changes is not suitable since the document has been restructured. We produced a track-change document for each of the two new parts by the following procedure:

1. Start from the submitted version 1.
2. Remove sections or text that is not contained in the respective part of version 2. Removed parts are marked with [...]. In some cases, the text was removed because the markup of changes made rewritten paragraphs unreadable.
3. Rearrange the sections to match the structure of the respective part of version 2. Old section numbers are indicated in the section titles.
4. Highlight changes between the new version and the obtained LaTeX-document using latexdiff.

Since individual paragraphs were not rearranged before using latexdiff, some identical or similar paragraphs are marked as new in the tracked changes.

The thus obtained documents follow below.

German methane fluxes ~~in 2021~~ estimated ~~with an~~ ~~ensemble-enhanced top-down~~ using ~~ICON-ART~~ – Part 1: ~~Ensemble-enhanced~~ scaling inversion ~~based on the ICON-ART~~ ~~model~~

Valentin Bruch¹, Thomas Rösch¹, Diego Jiménez de la Cuesta Otero¹, Beatrice Ellerhoff¹,
Buhalgem Mamtimin¹, Niklas Becker¹, Anne-Marlene Blechschmidt¹, Jochen Förstner¹, and Andrea
K. Kaiser-Weiss¹

¹Deutscher Wetterdienst, Frankfurter Str. 135, 63067 Offenbach

Correspondence: Valentin Bruch (valentin.bruch@dwd.de) and Andrea K. Kaiser-Weiss (andrea.kaiser-weiss@dwd.de)

Abstract. ~~A reliable~~ ~~This two-part study explores the~~ quantification of greenhouse gas emissions ~~is important for climate~~
~~change mitigation strategies using atmospheric observations in order to validate national emission inventories.~~ Inverse methods
~~based on observations and atmospheric transport simulations~~ can support emission quantification ~~down to at~~ the national scale
~~based on observations and atmospheric transport simulations~~, yet, they are often limited by the ~~observing systems~~ ~~observation~~
5 ~~coverage~~, transport model uncertainties, and inversion methodologies. Here, we ~~present~~ ~~introduce~~ a system for ~~observation-based,~~
~~regional methane flux estimation, which has the potential for long-term operational support of national emission reporting.~~
~~We regional estimation of methane fluxes and~~ apply this to Central Europe ~~in 2021 with~~ ~~with a~~ focus on Germany, where we
distinguish emissions from different anthropogenic sectors. ~~We evaluate the robustness of the method using sensitivity tests~~
~~with in-situ observations from the Integrated Carbon Observation System (ICOS).~~ Using synthetic observation experiments,
10 ~~we estimate the impact of transport errors on the flux estimates.~~ The atmospheric transport is calculated ~~with~~ ~~employing~~ the
numerical weather prediction model ~~ICON-ART~~ ~~ICON with its module ART~~ at 6.5 km resolution, sampling the meteorological
uncertainty with a 12-member transport ensemble. ~~We use a priori fluxes from national reporting to facilitate the validation of~~
~~reported fluxes.~~ ~~The same transport ensemble is used to generate pseudo-observations with a simulated transport uncertainty.~~
Posterior fluxes are estimated with a ~~modified~~ synthesis inversion method ~~, relying on observations from the Integrated Carbon~~
15 ~~Observation System (ICOS).~~ ~~for three different approximations of the model–observation error covariance matrix.~~ Compared
to the a priori, we find a significant increase in methane emissions in Germany and in the Benelux. ~~We estimate German~~
~~methane emissions (32 ± 19) % higher than the anthropogenic emissions in the national inventory, and attribute this difference~~
~~mainly to the agricultural sector, although separation from Land Use, Land Use Change and Forestry (LULUCF) as well as~~
~~natural fluxes requires further research.~~ ~~We find that using ensemble-estimated transport uncertainties can significantly reduce~~
20 ~~the random error of emission estimates.~~ ~~The combination of an ensemble-enhanced numerical weather prediction model for~~
~~atmospheric transport and good observation coverage paves the way to sector-specific~~ ~~Our results highlight the importance of~~
~~analyzing biases in flux inversions for reliable,~~ observation-based ~~national~~ emission estimates.

1 Introduction

~~Reducing-Quantifying~~ greenhouse gas (GHG) emissions is ~~erucial for mitigating current anthropogenic global warming-~~

25 ~~UNFCCC (essential for effective mitigation of anthropogenic climate change. Atmospheric GHG inversions provide such~~
~~quantification by connecting the observed atmospheric composition to surface fluxes using transport models. This so-called~~
~~“top-down” approach is complementary to “bottom-up” emission estimates, which are based on activity data and emission~~
~~factors (IPCC et al., 2019). Top-down emission estimates can be used to validate national bottom-up GHG inventories reported~~
~~to the United Nations Framework Convention on Climate Change)-compliant national inventories and -or process models~~
30 ~~quantify anthropogenic GHG emissions for the purpose of monitoring the effectiveness of mitigation as planned, e. g., in~~
~~the Paris Agreement. [...]~~ (UNFCCC) (Manning et al., 2003, 2011; Henne et al., 2016). Such national-scale estimates are
~~typically limited by the observation coverage (Petrescu et al., 2023) and uncertainties in atmospheric transport modeling (Gerbig et al., 2008).~~
~~This motivates estimating methane emissions in the comparably well-observed Central Europe using a high-resolution transport~~
~~model and applying methods from numerical weather prediction (NWP) to estimate the transport uncertainty.~~

35 Regional top-down estimates of long-lived GHG can be based on different types of transport models. Lagrangian models
calculate trajectories from selected locations by moving with air parcels transported by the wind. They have been widely used
for inversions of trace gases like halocarbons, nitrous oxide and methane (CH_4) in European regions, see e.g., Stohl et al.
(2009); Ganesan et al. (2015); Henne et al. (2016). In contrast, Eulerian models – such as ICON–ART – continuously transport
trace gas concentrations through three-dimensional grid boxes. Although they are computationally more expensive for cases
40 where a relatively small number of trajectories would suffice, they become superior when the amount of data grows and, as
Engelen et al. (2002) pointed out, open the road for data assimilation methods as used in ~~numerical weather prediction-NWP.~~
~~Among the Eulerian models, also NWP models have been used for regional flux inversions of CO_2 (Lauvaux et al., 2013) and~~
 ~~CH_4 (Steiner et al., 2024b).~~ Regardless whether Lagrangian or Eulerian or even combined approaches (Rigby et al., 2011) are
applied, the top-down estimation requires solving an inverse problem (Enting, 2002). Eulerian transport model based inversions
45 may employ emission ensembles, as in Steiner et al. (2024b) with a localized Kalman filter, and other data assimilation methods
(see, e.g., Meirink et al., 2008). Alternatively, the method of synthesis inversion scales a set of a priori emission categories
(Kaminski et al., 2001). ~~Note that Meirink et al. (2008) compare 4D-Var and synthesis inversion methods.~~

In this work, we ~~present a modular system for regional-~~introduce a system for national-scale top-down ~~estimates of CH_4~~
~~fluxes designed to validate national inventories, including the discrimination of economic sectors such as agriculture and~~
50 ~~industry. [...]~~estimation of CH_4 emissions based on modeling experience from NWP. We analyze the benefit of constraining
the transport uncertainty using a meteorological ensemble as proposed by Ghosh et al. (2021) and Steiner et al. (2024a). A
synthesis inversion method is used to estimate emissions with a focus on Germany based on high-resolution a priori emissions
from national reporting and in situ observations of atmospheric CH_4 concentrations.

In the present Part 1 of this two-part study, we describe our new inversion system and evaluate its performance. Section 2 introduces
55 the method with a detailed description of the uncertainty estimation. The description of the inversion system is completed by
the input data described in Sect. 3. In Sect. 4, we analyze the performance using synthetic observation experiments and test the

sensitivity to tuning parameters with real observations. We conclude in Sect. 5 and refer to Part 2 (Bruch et al., 2025a) for a discussion of the emission estimates obtained using real observations.

2 Method

60 ~~In the employed offline scaling inversion, we categorize the a priori fluxes and scale each flux category~~

~~We use a synthesis inversion method (Kaminski et al., 2001) that scales the CH₄ fluxes to optimize the agreement between model prediction of model predictions and observations. We start by defining flux categories which~~ In this method, the fluxes are initially grouped into a manageable set of flux categories. Here, these are 46 categories that subdivide the fluxes by region and ~~sector. emission sector.~~ With the Eulerian transport model, the ~~concentrations concentration~~ from each flux category ~~are~~ is calculated separately at all grid cells and time points. At the location and time of the observations, the model writes out the predicted concentrations from the flux category contributions and their sum is compared to the observed concentration. ~~Our inversion makes~~ The inversion then minimizes the mismatch between model prediction and observations by scaling each of the flux categories by one number – the scaling factor – making use of the linear relation between ~~surface~~ fluxes and concentrations in the atmosphere. ~~The categorized fluxes are scaled to minimize the mismatch between model prediction and observed concentrations.~~ Thus, the inversion result consists of one scaling factor for each flux category. ~~The~~ By multiplying the a priori fluxes ~~multiplied by with~~ the scaling factors ~~yield we obtain~~ the a posteriori fluxes. This scaling method cannot provide a correction where a priori fluxes are zero (Kountouris et al., 2018). However, this is less of a problem for CH₄, as inventories can collect where methane-emitting activities are normally located, but emission factors which translate the activities into bottom-up emissions are uncertain (Dammers et al., 2024).

75 The described method relies on high quality model predictions as well as accurate concentration observations. ~~Furthermore, we need to estimate the model uncertainty and error correlations to assess whether deviations between model and observations contain information on the fluxes.~~ To match these requirements, we have carefully chosen the configuration of the transport model (Sect. 2.1) ~~. Also, selected Sect. 2.1) and consider the specific difficulties in modeling strong plumes (2.2).~~ Selected observational data are employed to remedy model boundary effects and therefore improve the overall model predictions (Sect. 2.2 Sect. 2.3). In ~~this section, we further Sect. 2.4, we~~ introduce the Bayesian inversion framework ~~(Sect. 2.3), the model uncertainties (Sect. 2.4), and our strategy for dealing with strong plumes (Sect. 2.5).~~ To assess whether deviations between model and observations contain information on the fluxes, we estimate the model uncertainty and error correlations. We compare three different methods for estimating these uncertainties and correlations (Sections 2.5 and 2.6). Furthermore, we define the time window and a priori uncertainties of the inversion (Sections 2.7 and 2.8). A summary of the method and data streams will be provided in Sect. 3.5.

2.1 Transport ~~model~~ simulation

2.1.1 Transport model

The atmospheric transport is simulated using the ~~numerical-weather-prediction-NWP~~ model ICON (Zängl et al., 2015) ~~with the ART module (Rieger et al., 2015; Schröter et al., 2018).~~ in a configuration close to operational NWP at Germany's Meteorological Service (DWD), extended with the module for Aerosol and Reactive Trace gases (ART) (Rieger et al., 2015; Schröter et al., 2018). The model is run in limited area mode for a domain covering large parts of the European continent (latitudes 34° N to 70° N, longitudes 21° W to 59° E, see Fig. 1) with a horizontal resolution of 6.5 km (ICON grid R3B8) and 74 vertical levels up to a maximal height of 22.77 km. ~~The~~ The ICON model simulates the meteorology and the tracer transport. Re-initialization of the meteorological fields every 24 h with operationally produced analysis fields ensures that the meteorology stays close to reality.

The surface CH₄ fluxes are provided to the transport model using the online emission module (Jähn et al., 2020; Steiner et al., 2024b). We do not simulate any chemical reactions, because the typical lifetime of CH₄ in the atmosphere is much longer than the time that an air parcel typically spends in our modeling domain.

For long living tracers like methane, the correct treatment of the lateral boundary concentrations is of importance. Therefore, we extended the model by implementing lateral boundary nudging for ART tracers in order to obtain smooth fields and avoid strong ~~gradients~~ spatial gradients. The nudging is limited to a boundary zone of width < 250 km. Further, so-called meteogram output has been implemented for ART tracers, providing model output in the vicinity of observation locations with high temporal resolution.

2.1.2 Meteorological ensemble

For improved uncertainty estimates, we run a meteorological ensemble of 12 members. Each ensemble member uses ~~slightly different but equally likely parametrizations and different~~ meteorological initial and ~~boundary conditions. The construction of this ensemble follows the operational weather prediction at DWD (Schraff et al., 2016).~~ lateral boundary conditions from the operational ensemble data assimilation used for global NWP at DWD (Schraff et al., 2016; Reinert et al., 2025). Since our meteorological input fields and the transport model setup are taken from operational NWP at DWD, the ensemble provides a reasonable estimate for the meteorological uncertainty in our model, including uncertainties in the simulated wind field and atmospheric stability.

In the following, we ~~therefore~~ distinguish a so-called deterministic model run providing the best estimate of the modeled CH₄ concentration, and the ensemble runs providing 12 different CH₄ concentrations to estimate the uncertainty. The ensemble will only be used to estimate model uncertainties and error covariances (see Sect. 2.5), and to generate pseudo-observations (Sect. 3.4).

2.2 **Implementation of CH₄ fluxes (Sect. 4.1)**

2.1.1 Definition of flux categories




Figure 1. Model domain, colored to distinguish 35 patches defining regional flux categories. Observation sites (dots) are colored by the choice of model equivalent height (see Table C1). Dark blue at the domain boundary indicates regions for which emissions are not categorized and therefore not modified in the inversion. Other colors only distinguish neighboring patches. In white hatched regions, natural fluxes are also categorized and scaled. A white ellipse marks the Upper Silesian Coal Basin, in which fugitive emissions define their own flux category. In Germany, the map shows the six regions used for the agricultural sector. For other sectors in Germany, we use four regions: south (yellow and light green), west (dark blue), north (light green), and east (dark green and yellow).

Estimating CH_4 fluxes in $> 10^5$ grid cells based on 50 observation sites seems impossible without reducing the number of degrees of freedom of the fluxes. Here, we reduce the degrees of freedom drastically by parametrizing the fluxes using only 46 basis vectors. A basis vector in this parametrization is a flux category that contains all fluxes from one region, possibly limited to specific emission sectors. For example, we define all anthropogenic emissions from Denmark as one flux category. We thereby assume that the distribution of anthropogenic emissions within Denmark is correct in the a priori and only allow the inversion to adjust the total emissions from Denmark.

For the inversion, we define the flux categories based on sector and region with the primary aim of providing an accurate estimate for emissions from Germany, resolving federated states where possible, to address the user requirements of potential stakeholders.

2.1.2 Definition of flux categories (Sect. 4.1.1)

When distinguishing emission sectors, we stay close to the national reporting by using definitions from the gridded aggregated nomenclature for reporting (GNFR, Veldeman et al., 2013). For the agricultural sector (GNFR sectors K+L), which contributes roughly two thirds of all German CH_4 emissions, we distinguish six regions within Germany as depicted in Fig. 1. We furthermore try to distinguish the sectors waste (GNFR sector J) and public power (GNFR A) from the For the sum of all remaining sectors (“other”, GNFR B–I). However, we will only present results in which these sectors are combined (see discussion in Sect. 6.3 and Appendix I). For the sectors waste, public power, and other other sectors – excluding natural and LULUCF fluxes – we distinguish four regions, i.e., the federated states south: Baden-Wuerttemberg and Bavaria, west:

Table 1. Overview of sectors distinguished in the inversion and number of flux categories. We distinguish the focus region, well-observed regions near the focus region, and regions in large distance from the focus region (“remote”). The latter are split in very large flux categories with low a priori uncertainty. Natural plus LULUCF fluxes are separated from other anthropogenic emissions only in regions where the natural fluxes are strong and in Germany. One extra category in the well-observed regions is the Upper Silesian Coal Basin (marked* in the last column). See Fig. 1 for the definition of flux categories on the map.

Classification	Countries and regions	Sectors	# of areas	# of flux categories
focus region	Germany	agriculture, LULUCF + natural, other	6 agr., 4 other, 1 LULUCF	11
focus region	Netherlands	agriculture, other	1	2
well observed	Sweden, Norway	LULUCF + natural, anthropogenic	2	4
well observed	DK, PL, CZ, AU, SK, HU, SV, HR, BA, CH, FR, BE, LU, UK, IE, northern IT, North Sea	anthropogenic (excl. LULUCF)	16	17*
remote	Finland, north-western Russia	LULUCF + natural, anthropogenic	2	4
remote	other	anthropogenic (excl. LULUCF)	8	8

North Rhine-Westphalia, Hesse, Rhineland-Palatinate and Saarland, north: Lower Saxony, Bremen, Hamburg and Schleswig-
135 Holstein, as well as east: Mecklenburg-Western Pomerania, Brandenburg, Berlin, Saxony, Saxony-Anhalt and Thuringia. Natural plus LULUCF fluxes in Germany are treated as a single flux category.

Outside Germany, we do not distinguish sectoral emissions, with one exception. Agriculture emissions in the Netherlands form their own category, as we found that they strongly influence the CH₄ concentrations in Germany, caused by the proximity and high emission rates in the Netherlands. ~~We define~~ We define further categories by area for anthropogenic emissions
140 excluding LULUCF such that a comparably high resolution is obtained in regions near Germany with high observation coverage. These area-defined flux categories follow borders as feasible for the inversion. Areas with small expected influence on inversion results for Germany are combined in large categories, such as Spain plus Portugal, Türkiye plus Greece, and large areas east of Poland. All area-defined categories are shown in Fig. 1 ~~and an overview of the sector resolution is given in Table 1.~~ ~~We mitigate and analyze the plume problem (Sect. 2.2) in our inversion system by defining a separate flux category for the strongest CH₄ plume in Central Europe. These are fugitive emissions from the Upper Silesian Coal Basin with yearly emissions of 567 kt in our prior (white ellipsis in Fig. 1).~~

We treat natural plus LULUCF fluxes separately and categorize them only in Germany, Scandinavia, and the north-eastern part of our domain (hatched regions in Fig. 1). This is motivated by strong CH₄ emissions from wetlands in summer in Scandinavia and northern Russia in our prior (Segers and Houweling, 2020). Uncategorized fluxes – whether natural or anthropogenic – are not scaled in the inversion, but still included in the transport simulation such that no fluxes are discarded. To avoid strong spatial gradients in the concentration fields, the boundaries between different area-defined categories are smoothened as visualized in Fig. 1.

We furthermore define a separate flux category for the strongest CH₄ plume in Central Europe to mitigate the plume localization problem described below (Sect. 2.2). These are fugitive emissions from the Upper Silesian Coal Basin with yearly emissions of 567 kt in our prior (white ellipse in Fig. 1).

2.1.2 Tracer assignment in the transport model (Sect. 4.1.2)

In the transport simulation, we consider not only the categorized fluxes, but also the CH₄ from lateral boundaries and from uncategorized emissions. Overall, we simulate the transport of 58-50 tracer fields in the deterministic model run:¹

- (i) ~~AH~~ Sum of all anthropogenic emissions excluding LULUCF. This constitutes a single, common tracer.
- (ii) ~~AH~~ Sum of all natural plus LULUCF fluxes. This constitutes another single, common tracer, which summed with (i) covers all a priori emissions in the domain.
- (iii) Far field. The far field contains the CH₄ from initial and lateral boundary conditions.

The sum of (i)–(iii) is the total a priori CH₄ concentration. The a posteriori concentration is not computed directly. Instead, we treat the deviation of the posterior concentration from the prior as a perturbation. To compute this perturbation, we simulate the transport of each flux category:

- (iv) **Flux categories**. For each ~~flux category of the 46 flux categories~~ an own tracer field is defined. To avoid the accumulation of categorized CH₄ beyond the time scale on which we consider the modeled transport reliable, we set an artificial decay rate of these concentrations. After emission, the concentration in these tracer fields decays exponentially with a mean lifetime of five days. ~~In combination with (i) and (ii) this technical feature~~ This technical feature constitutes a localization in time similar to the commonly used localization in space (e.g., Steiner et al., 2024b) and allows a waning of sectoral and regional attribution over a few days ~~—no CH₄ is lost, though, as full CH₄ transport is modeled without decay in (i) and (ii).~~ This regulates that any attribution of a CH₄ anomaly to a certain region or sector is only attempted if the emission was fresh or a few days ago. Furthermore, this allows us to save computing time by limiting the transport of these flux category tracer fields to altitudes below 8 km. ~~In total, 49 categories for anthropogenic emission excluding LULUCF are covered—including 18 sector-resolving categories in Germany, two sector categories in the Netherlands and one plume category—and complemented by five natural plus LULUCF flux categories. The artificial decay rate~~

¹Technically, the simulation includes 58 tracers in an attempt to split up the sector “other” in Germany in three sectors. Since we do not use these additional data here, we describe the setup for the 50 tracers we actually used.

affects the posterior concentration and the sensitivity of the inversion to changes in the emissions. However, assuming that the typical time between emission and observation is short compared to the artificial lifetime and in the presence of transport model errors, we expect that this feature of our inversion system leads to more robust results.

180 (v) ~~Far-field. The far-field contains the CH₄ from initial and lateral boundary conditions.~~

(vi) **Auxiliary field for plume detection.** For the purpose of investigating the model uncertainty due to the plume from the Upper Silesian Coal Basin, an auxiliary tracer is added (see Sect. 2.6.1). This tracer is never added to the total CH₄ concentration but only serves as an indicator for the plume location.

185 ~~In the post-processing of the model data, the categorized concentrations from (iii) are subtracted from the combined tracers (i) or (ii).~~

2.2 Plume localization problem (Sect. 2.2)

~~Plumes caused by~~ In our transport simulation and inversion, we address the specific challenge posed by plumes from high emissions in a small area require special treatment to avoid a potential bias in the inversion small areas. The inversion may be biased for such plumes due to the so-called double penalty issue (Vanderbecken et al., 2023). In cases where our model 190 falsely predicts that the plume reaches an observation site, the inversion will reduce the emissions to improve the agreement with the observation. In the opposite case, when the model fails to predict that a plume reaches the observation, the inversion will not change the plume emission amount but will wrongly increase emissions in other areas instead. This can cause a systematic underestimation of fluxes from localized plumes. ~~In the posterior- R inversion, this problem is mitigated because the underestimation of emissions is penalized by a lower model uncertainty.~~ To avoid biases in the inversion results, we suggest to 195 treat strong plumes separately, with their own flux categories. This allows us to quantify the problem (see Sect. 5.6 Sect. 4.2) and to limit the plume penalty influence on other flux categories.

2.3 ~~The need for a far-field~~ Far-field correction

For cases where the model predicts almost no influence from our categorized emissions (i.e., clean air cases), deviations between model and observations point to the need for correcting the CH₄ advected across the lateral boundaries – here referred 200 to as “far field”.² For our regional inversion problem, it is essential to separate the CH₄ emitted within the domain from the far field, in order to avoid model biases which would confound the aspired flux scaling (see, e.g., Chen et al., 2019, for CO₂). To minimize potential biases arising from imperfect boundary conditions, we construct a correction field which is added to the modeled far-field concentration in the whole domain after the transport simulation. We require this correction field to be smooth on large length and time scales, chosen in our case as spatial and temporal scales 320 km (horizontal), 1 km (vertical), 205 and 16 h (time). We construct this far-field correction using a Kalman smoother as described in detail in Appendix A. This construction uses only clean-air observations with a cumulated signal of all flux categories of ≤ 20 ppb and a total signal from emissions within our domain of ≤ 50 ppb.

²Technically, the far field also includes the initial CH₄ concentration. But this is hardly relevant due to our generous spin-up period of 17 days.




Figure 2. Statistical evaluation of the far-field correction at the observation coordinates when using synthetic observations (light blue area) or real observations (dark red line). Considering all data points used in the inversion, histograms of the far-field correction show (a) the range of the correction and (b) its temporal variation. For each station, month, and realization of pseudo-observations, we compute the root mean square (RMS) and the mean (or bias). Histograms combining these values for all stations and months are shown in (c) and (d).

The Figure 2 shows a statistical overview of the far-field correction when using real observations (red line) or pseudo-observations (shaded area). The considered pseudo-observations are generated from the ensemble members of the transport simulation and represent the case where simulated emissions and boundary conditions are perfect, i.e., equal to the truth. The far-field correction range is usually limited to ± 10 ppb (see Fig. 2 for details). Section 2.2 comprises the mathematical derivation, parameters, and a statistical overview of the when using real observation data and ± 5 ppb in the synthetic observation experiments (Fig. 2 a) with variations of a few ppb per day (Fig. 2 b). The broad distribution of the root mean square (RMS) for different observation sites and months in Fig. 2 (c) indicates significant differences among the stations when using real observations.

Figure 2 (d) shows that the correction has a small bias towards positive corrections even when using synthetic observations with unbiased fluxes and boundary conditions. This is partially due to the pseudo-observations, which are biased by $+0.5$ ppb compared to the simulated concentrations due to details of the transport model configuration. The other part of the bias hints to a more general problem. We construct the far-field correction using observations for which the model predicts clean air, i.e., a low signal from the emissions. Since the transport model is not perfect, this introduces a sampling bias: We select more observations for which the model underestimates the concentrations and thereby increase the bias to 1.2 ppb. In response to this bias, the far-field correction increases the simulated concentrations by 1.0 ppb.

The sampling bias will likely also occur when working with real observations. But the estimated correction bias of 0.6 ppb due to the sampling is small compared to the accuracy of the Copernicus Atmosphere Monitoring Service (CAMS) inversion-optimized data product used for our boundary conditions (Segers et al., 2023) (see Sect. 3.1). We therefore do not expect a significant impact on the emission estimates.

2.4 Inversion General approach of the inversion framework

We use a Bayesian inversion to optimize the agreement of model and observations ~~by scaling the flux categories. This is formulated in the~~. We define a vector of scaling factors – in our application $s \in \mathbb{R}^{46}$ – ~~consisting of one prefactor for each flux category. This low-dimensional parametrization of the fluxes leads to the~~ optimization problem

$$s^{\text{post}} = \arg \min_s \left\{ \frac{1}{2} (y - Hs - x^{\text{ff}})^{\top} R^{-1} (y - Hs - x^{\text{ff}}) + \frac{1}{2} (s - s^{\text{prior}})^{\top} B^{-1} (s - s^{\text{prior}}) \right\} \quad (1)$$

for the posterior scaling factors s^{post} . Here, the first term ~~penalized~~ penalizes the deviation from the ~~concentration observations observed concentrations~~, and the second term penalizes the deviation from the prior fluxes. In the first term, the vector y of observed concentrations is compared to the model prediction, which consists of the ~~transported fluxes contribution~~ Hs of fluxes within the model domain and the modeled far field x^{ff} ~~including the far-field correction. All model predictions (x^{ff} and Hs) are already~~ projected to the observation space. The ~~transported contribution of~~ fluxes Hs ~~depend depends~~ linearly on the vector ~~s of scaling factors for the flux categories, which is optimized.~~ The difference between modeled and observed values is weighted by the error covariance matrix R describing the combined uncertainty of the transport model and the observations. With the second term we constrain the deviation of s by defining from a priori scaling factors s^{prior} (~~$s_i^{\text{prior}} = 1$ for all i~~ $s_k^{\text{prior}} = 1$ for all k) with an error covariance matrix B characterizing the a priori uncertainty (see Sect. 2.8).

In Eq. (1), ~~the fluxes are parametrized by the vector s of scaling factors. The flux categories thereby define the low-dimensional space in which the inversion can optimize the fluxes. The~~ the model observation operator H connects the space of scaling factors (vectors s^{prior} , s^{post}) to the observation space (vectors y , x^{ff}). Computing H requires the transport model which distinguishes the flux categories. The setup is designed for optimizing a low-dimensional vector s^{post} of scaling factors ($\sim 10^2$ degrees of freedom) using a large number of observations ($\sim 10^4$), but an extension to more degrees of freedom and/or more observations is possible.

2.5 Computation of model uncertainties Approximations for the error covariance matrix R'

~~The inversion in Eq. (1) requires an estimate of the model–observation mismatch uncertainties described by~~ The definition of the error covariance matrix R in Eq. (1) is crucial for the inversion. R describes the combined uncertainties and correlations of observations and model predictions. In our case, the ~~measurement error is negligible in comparison to the model error, and the non-negligible representativity error is part of~~ observation uncertainty (usually $\lesssim 1$ ppb, ICOS RI (2020)) is small compared to the ensemble-estimated transport uncertainty (typically 5 ppb to 10 ppb). We therefore focus on the model uncertainty.

Many works have used diagonal R matrices (e.g. Bergamaschi et al., 2010; Petrescu et al., 2023; Steiner et al., 2024b) and others found non-diagonal approximations for R (Ghosh et al., 2021; Steiner et al., 2024a). Here, we use the diagonal R for comparison to two different ways of constructing a non-diagonal R matrix from our transport ensemble. We therefore compare three ways of constructing R :

Diagonal R : This baseline scenario considers a diagonal R matrix and discards all information from the transport ensemble.

Prior R : In a standard ensemble approach, we construct R using the transport ensemble with a priori fluxes.

Posterior R : We extend the standard approach by estimating R using the posterior fluxes in the transport ensemble.

The construction of the different R matrices consists of two steps that are described below. First, we construct a matrix R' that estimates the dominant uncertainties and correlations using one of the three methods. Second, we obtain R from R' by inflating and adding additional uncertainties to mitigating some known issues of the inversion (Sect. 2.6).

2.5.1 Diagonal R

In the baseline scenario of a diagonal R matrix, all observation and model uncertainties are assumed to be uncorrelated. However, it is known that model predictions for observations separated by only one hour usually have correlated errors. To avoid underestimating the overall uncertainty without introducing correlations in R , we assume high uncertainties of each observation. We simply refer to this as “model error” below. We estimate this uncertainty and the cross-correlations of model equivalents at the observation locations using an Following Steiner et al. (2024b), we assume that the signal from CH_4 emissions within our domain will generally increase the model uncertainty in the predicted CH_4 concentration. This motivates defining $R'_{ii} = \sigma_{\text{const}}^2 + (\beta H s^{\text{prior}}_i)^2$ where $\sigma_{\text{const}} = 10$ ppb and $\beta = 0.5$ are scalar tuning factors. Index i labels observation data points that are typically distinguished by location, time, and sampling height. The diagonal R scenario uses crude approximations because the selection of observations is designed for an inversion that can handle correlations. However, we will obtain qualitative insights from the comparison to the other approximations for R .

2.5.2 Prior R

This approximation of R is based on an ensemble of $M = 12$ different transport ~~model realizations, realizations~~. The potential of using a small transport ensemble for estimating model uncertainties was demonstrated by Steiner et al. (2024a). We can use the covariance of the ensemble members to estimate the transport uncertainty. ~~The main contribution R' of the uncertainty matrix is~~ We define

$$R'_{ij} = C_{ij} \frac{1}{M-1} \sum_{m=1}^M (x_i^m - \bar{x}_i)(x_j^m - \bar{x}_j) + \delta_{ij} \sigma_{\text{const}}^2, \quad (2)$$

where x_i^m is the prediction of ensemble member m for observation y_i assuming a priori fluxes, $\bar{x}_i = \frac{1}{M} \sum_m x_i^m$ is the ensemble mean, and $\sigma_{\text{const}} = 10$ ppb is a constant uncertainty added to each observation accounting for any representativity error. With this uncorrelated uncertainty σ_{const} , we account for additional uncertainties, such as representativity errors inherent to a simulation at finite resolution. Indices i, j label observation data points that are typically distinguished by location, time, and sampling height. By C_{ij} we denote a localization in space and time such that $C_{ii} = 1$ and $C_{ij} = 0$ for any observations i and j that we expect to be uncorrelated because of their temporal or spatial separation. In the application to Germany, we choose C_{ij} to be a Gaussian localization matrix with standard deviations 6 h (time), 319 km (horizontal), and 400 m (vertical). We use the notation $\delta_{ij} = 1$ if $i = j$ and $\delta_{ij} = 0$ if $i \neq j$.

2.5.3 **Uncertainty inflation** Posterior R

The transport ensemble may not necessarily include the full uncertainty of the transport model. The posterior R approximation is a variation of the prior R approximation. In Eq. (2), we use model predictions for the concentrations x_i^m . Instead of using the prior concentrations as in the prior R construction, we can define x_i^m as the posterior concentrations and thereby allow x_i^m to change as the inversion changes the fluxes. This leads to a self-consistent estimate of R' in the inversion. Consequently, Eq. (2) remains valid but x_i^m , and the localization C_{ij} further reduces the simulated uncertainty by suppressing correlations. This motivates an inflation of the uncertainty to avoid overconfidence in the model prediction. We inflate the uncertainty by a factor $f_i > 1$ depending on the observation site of observation i , leading to the matrix $f_i f_j R'_{ij}$. In the application below, we choose $f_i = 2$ except for some stations with known difficulties, for which $f_i = 3$ (see Table C1). R' , and R become functions of the scaling factors s . Since R is estimated using posterior scaling factors, we call this method the posterior R inversion as opposed to the prior R estimate. To compute the posterior concentration $x_i^m(s)$ for each ensemble member without prohibitive computational effort, we use an approximation described in Appendix B1.

As opposed to the diagonal R and prior R inversion with fixed R , the posterior R inversion does not allow for a closed form solution of Eq. (1). To solve the minimization problem in Eq. (1) numerically, we used SciPy's "trust-exact" implementation of a trust-region method (Virtanen et al., 2020; Moré and Sorensen, 1983; Conn et al., 2000). Within each iteration, the incomplete LU decomposition (Li et al., 1999; Li and Shao, 2011) of the sparse matrix $R(s)$ is the most computationally expensive task when the number of observations is large.

2.6 **Additional uncertainties and final error covariance matrix R**

The previously derived approximations for the error covariance matrices R' describe our knowledge of the transport uncertainty and the observation uncertainty. In the next four steps, we increase uncertainties and include other possible sources of uncertainty to obtain approximations for R that are suitable for the inversion.

2.6.1 **Mitigating the plume localization problem**

To reduce the bias which we predicted for strong plumes in Sect. 2.2, we increase the uncertainty for all observations that are likely affected by a plume. The transport ensemble will already lead to an increased uncertainty when the model cannot predict reliably whether a plume hits an observation site. But with an ensemble of only 12 members, this will not cover all cases where model and observations deviate. We therefore introduce an auxiliary tracer that contains emissions from the Upper Silesian Coal Basin, spatially smoothened on a length scale of 0.4° (one standard deviation of a Gaussian filter). Denoting the concentration of this tracer at observation i by ρ_i , we increase the uncertainties to $R^{\text{step 1}}_{ij} = R'_{ij} + 0.25\rho_i^2\delta_{ij}$.

2.6.2 **Dynamic uncertainty inflation**

To avoid potential biases through site-specific small-scale features not captured in the model, we aim to base our inversion on many observations. To this end, we limit the influence of individual data points on the inversion result by inflat-

ing the uncertainty further in the case of a very large disagreement between model and observation. This is achieved by replacing f_i with $f'_i \geq f_i$ to increase the uncertainty an uncertainty inflation of individual observations until the deviation $\mu = y - H s^{\text{prior}} - x^{\text{ff}}$ between model and observations is at most three standard deviations of the resulting error covariance matrix $R_{ij} = f'_i f'_j R_{ij}^{\text{step 2}} = g_i g_j R_{ij}^{\text{step 1}}$, i.e., $f'_i = \max\{f_i, \frac{|\mu_i|}{3\sqrt{R_{ii}}}\}$, $g_j = \max\{1, \frac{|\mu_j|}{3\sqrt{R_{jj}^{\text{step 1}}}}\}$. This is justified because large deviations between model and observations, $|\mu_i| > 3\sqrt{R_{ii}}|\mu_j| > 3\sqrt{R_{jj}^{\text{step 1}}}$, are likely caused by local pollution or modeling problems that are not captured appropriately in our uncertainty estimate. This correction makes sure that inversion results will be based on many observations and no single measurement can have an extreme impact. At the same time, this method it is less sensitive to tuning parameters than discarding outliers completely.

2.6.3 Flux dependence of model uncertainties

In Eq. (2), we estimate the model uncertainty utilizing the model concentration x_i^m , which itself depends on the fluxes. In principle, one can compute x_i^m using either the a priori fluxes or the a posteriori fluxes. In the latter case, the error covariance matrix

2.6.3 Static uncertainty inflation

The transport ensemble in the prior R depends on the scaling factors s , and the optimization problem Eq. (1) can only be solved numerically (see Appendix D) and posterior R construction may not necessarily include the full uncertainty of the transport model, and the localization C_{ij} further reduces the simulated uncertainty by suppressing correlations. This motivates another inflation of the uncertainty to avoid overconfidence in the model prediction. We inflate the uncertainty by a factor $f_i > 1$ depending on the observation site of observation i , leading to $R_{ij}^{\text{step 3}} = f_i f_j R_{ij}^{\text{step 2}}$. We choose $f_i = 2$ except for some stations with known difficulties, for which $f_i = 3$ (see Table C1). To keep the methods for constructing R comparable, we apply this inflation also to the diagonal R matrix.

2.6.4 Far-field uncertainty

We furthermore account for the uncertainty in the far-field correction, although the effect of this additional uncertainty is small. We define $R_{ij} = R_{ij}^{\text{step 4}} = R_{ij}^{\text{step 3}} + 0.5|c_i c_j| \tilde{C}_{ij}$ where c_i denotes the smooth correction field introduced in Sect. 2.3 at observation i and $\tilde{C}_{ij}$ is the Gaussian localization matrix constructed by the length and time scales of the far-field correction (see Appendix A).

2.6.5 χ^2 analysis

To assess whether the estimated uncertainties are reasonable, one can compute the χ^2/N_{dof} value (Pearson, 1900). This value compares the a priori model–observation mismatch to the uncertainty assumed for this mismatch (see Appendix D for details). We will refer to these two variants of the inversion as the prior. A value of $\chi^2/N_{\text{dof}} > 1$ indicates that uncertainties are underestimated, whereas values smaller than one indicate the opposite. When comparing the observations to the far-field-corrected

Table 2. Median of χ^2/N_{dof} for different configurations. χ^2/N_{dof} for the prior R inversion also serves as an approximation for the posterior R inversion. Synthetic observations are generated using the ensemble simulation, assuming that the a priori fluxes and the CH_4 concentration on lateral boundaries are known exactly.

Observations	Far-field correction	χ^2/N_{dof} , diagonal R	χ^2/N_{dof} , prior R
real	yes	0.18	0.16
real	no	0.21	0.18
synthetic	yes	0.05	0.03
synthetic	no	0.06	0.03

model, we find $\chi^2/N_{\text{dof}} \approx 0.16$ for the prior R and ~~posterior- R inversion~~-inversion when using real observations (see Table 2). In an idealized setup, this indicates that the uncertainties of the model-data mismatch are overestimated by a factor 2.5. This
350 implies that our uncertainty inflation by a factor $f_i = 2$ for most observations seems unnecessary in the idealized setup. However, our data can contain unknown biases in transport and boundary conditions, and simplifying assumptions about the representativity of the low-dimensional state space of the inversion. We contain these potential issues of unknown error components by inflating the uncertainties.

In the synthetic experiments, the idealized transport uncertainty and perfect a priori emissions lead to even lower χ^2 , which
355 is expected because not all uncertainties are contained in the pseudo-observations of these synthetic experiments. Computing ~~posterior- R inversion~~ exactly requires that each ensemble member distinguishes the various flux categories similar to the deterministic modelrun. To enable the posterior- R inversion without significantly increased computational effort in the ensemble run, we approximate the posterior-based concentration x_i^m in the ensemble members as described in Appendix B1. The posterior- χ^2/N_{dof} for the posterior R inversion generally yields slightly higher emission estimates because larger scaling
360 factors lead to higher model uncertainties and thereby reduce the first term in Eq. (1). To provide robust inversion results with cautionary uncertainty estimates, we compute the prior- χ^2/N_{dof} is more difficult, but the result is expected to be similar to the prior R inversion and posterior-. The tuning parameters of the diagonal R inversion separately and interpret the difference of the two methods as methodological uncertainty. matrix were chosen such that the posterior uncertainties are similar to the prior R inversion, which also leads to similar χ^2/N_{dof} (see Table 2).

365 2.7 Inversion time window and temporal aggregation (Sect. 4.3)

We apply the inversion separately for each month in 2021. In each monthly time window, we start from fixed We simulate the transport for the whole year 2021 without any interruption. The inversion is then applied to each month separately by selecting only observations within one month. The scaling factors of the months are treated as independent, each month starting with the same a priori scaling factors $s_i^{\text{prior}} = 1$ and use observations within the time window to compute a posteriori scaling factors
370 as explained in Sect. 2.4. ($s_k^{\text{prior}} = 1$ for all k) and the same a priori scaling uncertainties (B matrix). However, when The continuous transport simulation over the whole year implies that the initial CH_4 concentration is hardly relevant after the

spin-up. At the beginning of each month, the modeled CH_4 concentration already consists of the far field – the contribution of the lateral boundaries – and the contribution of the fluxes, which will be adjusted by the inversion.

In summary, we correct the contribution of the lateral boundaries on the time scale of 16 h by the far-field correction, and the fluxes on the time scale of one month defined by the inversion time window. The inversion results consist of one vector $s^{\text{post}} \in \mathbb{R}^{46}$ of scaling factors and the corresponding error covariance matrix for each month. When aggregating results for the whole year, we treat the uncertainties of the prior or posterior fluxes of different months as correlated because these likely include systematic uncertainties and biases which we cannot fully separate from the statistical uncertainty. We therefore aggregate by adding up absolute emissions and their uncertainties linearly.

2.8 Prior uncertainties(Sect. 4.4)

In each inversion time window, we consider ~~uncorrelated~~ a priori scaling factors with a two standard deviation (2σ) uncertainty of ~~80%~~0.8 for most flux categories, corresponding to a 95 % confidence interval of ~~$\pm 80\%$~~ ± 0.8 . Throughout this paper, uncertainties will denote two standard deviations or 95 % confidence intervals. Categories resolving emission sectors have a higher prior 2σ uncertainty of ~~$\pm 100\%$~~ 1.0, and within Germany categories describing the same sector have an a priori uncertainty correlation of ~~50%~~0.5 (e.g., uncertainties of agriculture emissions in the German states of Bavaria and Baden-Wuerttemberg are assumed to be correlated). ~~Importantly, the previously defined sectors waste, public power and “other” are combined and denoted “non-agr.” in the following. The $\pm 100\%$ uncertainty applies to the combination. All other categories are treated as uncorrelated in the a priori.~~ For the Upper Silesian Coal Basin as well as regions with low observation density outside of our primary focus in Central Europe ~~and with low observation density(marked “remote” in Table 1)~~, the 2σ uncertainty is set to ~~$\pm 50\%$. Figure 4(b) shows these a priori uncertainties on a map. The sensitivity of our results to these choices and many more tuning parameters is tested in Appendix E.~~0.5.

3 Input data and processing

We apply the method to estimate CH_4 fluxes in the year 2021 in Germany and in the surrounding European domain, relying on input data for the transport simulation~~and~~ CH_4 concentration on the lateral boundary (Sect. 3.1), a priori fluxes (Sect. 3.2), and observations (Sect. 3.3).

3.1 Initial and lateral boundary conditions

The meteorological initial and lateral boundary conditions used to drive our transport model are taken from the archive of DWD’s operational ~~numerical weather prediction (NWP)~~NWP, which also employs the ICON model. As we do not assimilate meteorological data in our application, we re-initialize the meteorological fields every night at 0 UTC, using the analysis fields from the operational NWP data assimilation. Lateral boundary conditions for the meteorological fields are taken from the NWP short term forecasts with hourly resolution.

For the CH₄ concentrations, we use initial and lateral boundary concentrations from the ~~global-CAMS~~ CAMS global inversion-optimized dataset (Segers and Houweling, 2020), version v22r2, in the variant based on surface air-sample data for the inversion. The CAMS data have a resolution of 1° × 1° and are interpolated onto our model grid. In contrast to the meteorological fields, the CH₄ concentrations are only transported and never re-initialized. Each transport ensemble member uses slightly different initial and lateral boundary conditions for meteorological fields (see Sect. 2.1.2), but equal CH₄ concentrations on the lateral boundaries.

3.2 A priori CH₄ fluxes

For the inversion, we employ a priori CH₄ fluxes that were compiled from six datasets of anthropogenic and natural fluxes, as detailed in Table 3. We ensured mass conservation when interpolating to our model grid. We generally distinguish between anthropogenic emissions excluding LULUCF, and natural fluxes plus LULUCF. ~~Anthropogenic fluxes excluding LULUCF are split further into 12 GNFR sectors (gridded-aggregated-NFR, nomenclature for reporting, Veldeman et al. (2013)), but we only use this separation in selected regions of interest. Since the input datasets for anthropogenic emissions are based on reporting to the UNFCCC, these distinguish between GNFR sectors following the reporting conventions (Veldeman et al., 2013). For the inversion, we combine these sectors and only distinguish between agriculture and the sum of all other sectors as described in Sect. 2.1.1.~~ Natural plus LULUCF fluxes of CH₄ are mostly dominated by wetland emissions, for which we do not distinguish between natural and anthropogenic origin.

~~The datasets in Table 3 were chosen such that our~~ For Germany, we obtained a priori fluxes ~~are based on the national reporting to the UNFCCC. For Germany, the~~ directly from the national inventory agencies. ~~The~~ a priori LULUCF fluxes obtained from the Thünen Institute cover the emissions from mineral and organic soils. Notably, this excludes emissions from artificial water bodies in Germany – such as ponds – amounting to 160 kt or 8.5 % of the total German emissions in the national reporting, though these numbers are associated with large uncertainties (UBA, 2024, Table 399). These emissions are missing in our a priori estimate, leading to a low bias in the a priori.

3.3 Observations and **applied** pre-processing

We compare our model predictions to the high quality ground-based in situ observations of CH₄ concentrations collected in the European Obspack (ICOS RI et al., 2024), which includes the ICOS stations among others. These observations are assumed to be representative for a larger area (Storm et al., 2023). Table C1 lists all 53 available stations and Fig. 1 shows 50 stations that were used for the inversion. For tower observations, we ~~only consider sampling levels at least use up to three sampling heights per station, preferring the highest three sampling heights and discarding observations below~~ 50 m above ground level to reduce the influence of very local emissions. ~~For towers providing more than three sampling heights, we consider the three highest levels. Due to~~ Due to significant model–observation mismatch, we exclude the IPR, FKL and LMP stations. For LUT, BIR and HUN we only consider some seasons, specified in Table C1.

The model data are interpolated horizontally and vertically to the station sampling locations. The vertical sampling locations in model coordinates ~~is~~ are derived from the station sampling ~~height~~ heights and the modeled station ~~elevation~~ elevations.

Table 3. Input data for a priori CH₄ fluxes. The second column lists where these fluxes were considered. Here, “Germany” refers to all model grid cells that lie fully within the German borders. The national reporting distinguishes emissions by GNFR sectors of which A–M include all anthropogenic emissions excluding land use, land use change and forestry (LULUCF).

Data provider	Domain	Fluxes	Original grid	Time profile	Remarks
Umweltbundesamt (UBA)	Germany	GNFR sectors A–L (excluding LULUCF) <u>A–M</u>	native (ICON)	constant	based <u>Based</u> on reporting to the UNFCCC (UBA, 2023), spatially distributed using the Gridding Emission Tool for ArcGIS (GRETA 1.2.01) (Feigenspan et al., 2024)
Thünen Institute	Germany	organic and mineral soils <u>(part of LULUCF)</u>	100 m × 100 m	constant	emissions <u>Emissions</u> from organic and mineral soils, including wetlands but excluding artificial ponds (approx. 160 kt CH ₄ per year) (Fuß and Akubia, 2024)
CAMS-REG-ANT, v7.0	model domain excl. Germany	GNFR sectors A–L (excluding LULUCF) <u>A–M</u>	0.05° × 0.1°	constant	based <u>Based</u> on data reported to the UNFCCC for countries in Western and Central Europe (incl. Finland and the Baltic states) (Kuenen et al., 2021, 2022)
CAMS inversion optimized, v22r2	model domain excl. Germany, excl. oceans	wetlands	1° × 1°	monthly averages	variant <u>Variant</u> using surface air-sample data for the inversion (Segers and Houweling, 2020); Fluxes in model grid cells located over the ocean are set to zero.
Rocher-Ros et al. (2023), version 1.1	full model domain	rivers and streams	0.25° × 0.25°	monthly averages	
Weber et al. (2019)	oceans (full model domain)	oceans	0.25° × 0.25°	constant	

435 depending on the station characteristics (column “mountain” in Table C1). For high mountain stations, the modeled station elevation is given by the real station elevation above mean sea level. For stations on smaller mountains, we consider the arithmetic mean between real station elevation and model topography as proposed by Brunner et al. (2012) and Henne et al. (2016), and for all other stations the modeled station elevation is set to the model topography.

To make use of observations which are likely well represented by the model, we filter the observations based on the local
440 time of day, wind speed, and model–data mismatch. Table 4 lists how the root mean square error (RMSE) of the model output

Table 4. Average root mean square error (RMSE in ppb), mean absolute bias of the model prediction minus observation (in ppb), and number of available data points after each processing step (1–6) for synthetic (left) and real observations (right). Each row adds a processing step to all previous steps and improves the RMSE. Three numbers for steps 7 and 8 distinguish diagonal R , prior R , and posterior R inversion. Step 7 (uncertainty weighting) is not a processing step in the inversion since it uses only the diagonal of the uncertainty matrix R , but it underscores the importance of accurate uncertainty estimation. Step 8 refers to the result of ~~our~~the inversion. RMSE and absolute bias are computed separately for each station, sampling height and month. The obtained values are weighted by the number of data points and averaged. By taking the mean of multiple RMSEs for different stations, sampling heights and months, we obtain lower numbers than for the RMSE of the combined dataset, which would average squared values and thereby would give higher weight to large deviations between model and observations.

Step	<u>Synthetic observations (ppb)</u>			<u>Real observations (ppb)</u>		
	<u>RMSE</u>	<u>Absolute bias</u>	<u>Data points</u>	<u>Mean-RMSE</u>	<u>Mean-absolute bias</u>	<u>Data points</u>
1 horizontal and vertical interpolation	<u>–</u>	<u>–</u>	<u>–</u>	27.6 ppb	9.6 ppb	$6.02 \cdot 10^5$
2 time average (3 h)	<u>11.1</u>	<u>0.9</u>	<u>$6.02 \cdot 10^5$</u>	25.8 ppb	9.6 ppb	$6.02 \cdot 10^5$
3 time window 11 h–17 h / 23 h–5 h	<u>10.2</u>	<u>1.1</u>	<u>$1.48 \cdot 10^5$</u>	23.5 ppb	9.8 ppb	$1.48 \cdot 10^5$
4 minimal wind speed 2 m s^{-1}	<u>9.6</u>	<u>1.0</u>	<u>$1.30 \cdot 10^5$</u>	22.4 ppb	9.7 ppb	$1.30 \cdot 10^5$
5 extreme mismatch-exclusion <u>exclude extreme deviations</u>	<u>9.6</u>	<u>1.0</u>	<u>$1.30 \cdot 10^5$</u>	21.5 ppb	9.4 ppb	$1.29 \cdot 10^5$
6 far-field correction	<u>9.0</u>	<u>0.9</u>	<u>$1.30 \cdot 10^5$</u>	19.4 ppb	7.2 ppb	$1.29 \cdot 10^5$
7 weight by inverse uncertainty	<u>7.1, 6.9, 6.9</u>	<u>0.7, 0.8, 0.8</u>	<u>$1.30 \cdot 10^5$</u>	<u>14.4, 16.6, 16.6</u> ppb	<u>5.7, 6.6, 6.6</u> ppb	$1.29 \cdot 10^5$
8 inversion (posterior fluxes)	14.1 ppb <u>6.9, 6.8, 6.8</u>	<u>0.6, 0.8, 0.6</u>	<u>$1.30 \cdot 10^5$</u>	<u>12.4, 14.2, 14.0</u>	<u>2.5, 3.4, 3.0</u> ppb	$1.29 \cdot 10^5$

changes during these pre-processing steps. We start by ~~smoothing~~smoothing both observations and modeled concentrations in a time window of approximately $\pm 1.5 \text{ h}$ around each observation time as depicted in Fig. 3. This allows for some uncertainty in the timing of modeled tracer transport. The resulting correlation of neighboring time steps is automatically considered in the ensemble-based uncertainty estimate.

445 In the next steps, we filter the data by time in order to keep only observations expected to be representative for large regions. Observations within the planetary boundary layer are most representative in the afternoon hours whereas measurements at high mountains ~~have less local influence~~are less influenced by very local fluxes at night time (Bergamaschi et al., 2015). ~~We therefore use~~. Inversions therefore commonly use afternoon observations for flat land stations and night times at mountain




Figure 3. Weighting function for time interpolation of model and observations. For example, an interpolated model point at 16:30 UTC averages over all model output between 15:30 UTC and 17:30 UTC with full weight and another 1 h with linearly decreasing relative weight. The model yields instantaneous values every 15 min, whereas observations are provided as hourly averages, three of which contribute to the observational time average. Reference times are those times for which observations are available.

sites (Bergamaschi et al., 2015; Steiner et al., 2024b). We use the time windows 23 h to 5 h (local mean time) for stations on
450 high mountains and 11 h to 17 h for all other stations.

We furthermore exclude times with no wind to avoid a strong influence of local emissions that are not resolved in the model, motivated by Ganesan et al. (2015). All data points for which the model predicts a wind speed of $< 2 \text{ m s}^{-1}$ are excluded, which improves the overall agreement of model and observations as shown in Table 4 (step 4). Figure 4 shows that the RMSE indeed increases significantly at low wind speeds. This increase is partially captured by an increase of the ensemble spread,
455 supporting the idea of an uncertainty estimate depending on wind speed as proposed by Bergamaschi et al. (2022).

In the last filtering step – step 5 in Table 4 – we exclude data points with extreme mismatch between ~~model and observations of more than 200 ppb~~, far-field corrected a priori and observations, where $|y - Hs - x^{\text{ff}}| > 200 \text{ ppb}$. Data points where ~~the observations are more than 20 ppb below the model-predicted far field are~~ $y - x^{\text{ff}} < -20 \text{ ppb}$ are also discarded. Since no strong sinks of CH_4 are expected, the contribution of CH_4 from the lateral boundaries should not exceed the observations.
460 Thus, an observation below the model-predicted far field indicates an error in this far field. Steps 6–8 in Table 4 complete our processing chain by applying the far-field correction (Sect. 2.2 Sect. 2.3), indicating the relevance of the model uncertainty (Sect. 2.4 Sections 2.5 and 2.6), and finally yielding using the inversion results.

4 Application to Germany and neighboring areas for the year 2021

3.4 Synthetic observation experiments

465 To test our setup and analyze biases, we use synthetic experiments in which observation data are replaced by model-generated pseudo-observations. These synthetic experiments use exactly the same setup and the same observation coordinates. Only the observation values are replaced by the simulation result of one of our 12 ensemble members. We thus obtain 12 separate datasets of pseudo-observations, in which a transport error is simulated by using the transport ensemble members. The true fluxes assumed for these synthetic experiments are identical to the prior fluxes. This allows us to estimate a bias and a random




Figure 4. Dependency of RMSE dependence and proxies for the model uncertainty on wind speed (left axis). All data points from step 3 in Table 4 were ordered by the model-predicted wind speed and split into 100 bins, each containing approximately 1500 data points. The blue line indicates the cumulative fraction of observations (right axis). The figure shows the RMSE difference of model and observation (black line) and, the mean ensemble spread multiplied by factor 4 (magenta-red line), and the mean a priori concentration due to categorized emissions (green line) for each of these bins. The ensemble spread is the standard deviation of the model prediction in the 12 ensemble members. It is a main contribution to our uncertainty estimate for the model–data mismatch in the prior R and posterior R inversion. The signal of categorized emissions is used to estimate the uncertainty for the diagonal R matrix. Much of the larger RMSE at low wind speed is well captured by the ensemble spread inflated by factor 4-4 and by the mean a priori emission signal. In the inversion, we discard data points with wind speeds below 2 m s^{-1} (gray vertical line).

470 error in the posterior scaling factor. We will repeat this procedure with modified true fluxes in Sect. 4.3. An analysis of the sensitivity to random changes in the true fluxes is included in Part 2 (Bruch et al., 2025a).

3.5 Summary and overview

We can now summarize the inversion method following the required data streams in Fig. 5. After collecting the input data for the transport simulation (Sections 3.1 and 3.2, top of Fig. 5), we prepare the inversion by categorizing the fluxes (Sect. 2.1.1). The transport is simulated separately for the deterministic and ensemble run (Sect. 2.1.1, white ellipses in Fig. 5). Using observation data from the ICOS carbon portal and the simulation output, we compute model equivalents and filter these to ensure a high quality of the model predictions (Sect. 3.3). The data from the deterministic run are used to construct a far-field correction to mitigate uncertainties in the boundary conditions (Sect. 2.3). The ensemble data are used to construct the uncertainty matrix $R(s)$ as required for the prior R and posterior R inversion (Sect. 2.5.2). The far-field corrected data and the R matrix serve as input for the Bayesian inversion (Sect. 2.4). By combining the resulting posterior scaling factors with the categorized fluxes, we obtain posterior flux estimates.

[...]




Figure 5. Overview of the inversion system including input data sources. Arrows indicate data streams. Dashed lines indicate data streams with small or negligible impact on the inversion results. Colored areas group the input data (top), the deterministic model run and data processing (left), and the ensemble model run including processing of the resulting data (right). Colored text boxes distinguish gridded fluxes (green), data in observation space (blue, matrices in purple), and data in the space of scaling factors (red). Observation data are included when working in observation space (not explicitly marked). At the end of the processing chain (bottom), the three methods for estimating R lead to different scaling factors from which we can compute national emission estimates.




Figure 6. Mean (a, c) and standard deviation (b, d) of monthly flux estimates relative to the prior in synthetic experiments for diagonal R (blue), prior R (orange), and posterior R inversion (green). Each bar represents the posterior fluxes for 144 inversions, obtained from 12 datasets of pseudo-observations, each covering 12 monthly time windows. Black horizontal lines indicate the 2σ statistical uncertainty estimate. Panels (a, c) show the bias as the relative deviation of the mean posterior from the prior, which is equal to the synthetic truth. The standard deviation (b, d) among the 144 emission estimates indicates the random error expected in each monthly inversion. Colored lines in (b, d) show the mean posterior 1σ uncertainty, which is similar for all three methods.

4 Results and discussion

In this section, we examine the presented inversion system using synthetic experiments and sensitivity tests. We start by considering synthetic observation experiments in which the synthetic truth is equal to the a priori fluxes. Figure 6 shows a statistical evaluation of inversion results for this case, which we analyze for multiple aspects.

4.1 Simulated transport error (Sect. 5.6)

4.1 Random error

In Fig. 6, we see the bias (panels a, c) and random error (b, d) of the inversion results for selected countries or emission sources relative to the a priori emissions, distinguishing the three methods for constructing R . The random error is estimated by the standard deviation obtained from 144 inversions and indicates the precision or reliability of these results for a single month. The comparison of the three methods shows that the prior R and posterior R method lead to a very similar random error, which is considerably lower than for the diagonal R in all considered regions. This leads to the conclusion that using

495 a transport ensemble to estimate uncertainties and their correlations can significantly reduce the random error in emission estimates, independent of the far-field correction.

Since the diagonal R construction uses different tuning parameters than the prior R and posterior R inversion, we need to make sure that the chosen configurations are comparable. This is achieved by aiming for a similar posterior uncertainty in all methods for constructing R . Thin lines in Fig. 6 (b, d) show the posterior 1σ uncertainties to validate the similarity.

500 By comparing emission estimates without (panels a, b) and with the far-field correction (c, d), one can identify that the far-field correction changes the bias and slightly reduces the random error. Both effects are very similar for all three choices of R . Since the far-field correction pulls the simulated prior concentrations towards the observations, we can expect that it brings the emission estimates closer to the prior. But we can see in Fig. 6 (b, d) that the resulting reduction in random error is only weak.

505 4.2 Inversion bias

The bias shown in Fig. 6 (a, c) clearly depends on the far-field correction. The pseudo-observations without far-field correction have a bias of $+0.5$ ppb. The far-field correction reverts this to a negative bias of -0.5 ppb due to a sampling bias as explained in Sect. 2.3. Ideally, we would therefore expect a small positive bias in Fig. 6 (a) and an equally strong negative bias in panel (c). But the bias differs depending on how R is constructed.

510 For the diagonal R inversion, we see overall a positive bias for most regions. This approximation for R assumes a large uncertainty if the model predicts a strong signal from emissions. For an imperfect transport model, this implies that the model will tend to have a higher uncertainty when it overestimates the concentration and a lower uncertainty when it underestimates the real emission signal. As the model is more confident when observations are higher than the model prediction, it will tend to overestimate the emissions.

515 For the prior R approximation, we find a negative bias in the emission estimates in many regions. This may be due to the plume bias problem introduced in Sect. 2.2. For the Upper Silesian Coal Basin as a very strong and localized source, all methods show the expected negative bias. Notably, a considerable negative bias is also found for the Netherlands as a small country with high emission rates.

In the posterior R approximation, the negative bias for plumes is reduced, but also all other emission estimates are higher compared to the prior R inversion. To understand this, we recall that a transport error in our model only leads to an error in the predicted CH_4 concentration if the concentration field contains spatial gradients. Such gradients are caused by emissions. Stronger emissions directly cause higher uncertainty estimates in the meteorological ensemble. In the posterior R inversion, the inversion can adjust the emissions of the transport ensemble and thereby change the uncertainties. As we optimize the agreement of model and observations relative to the uncertainties, the system will prefer larger uncertainties. Thus, the inversion will tend to overestimate emissions to reach higher uncertainties. This counteracts the negative plume bias, but it may also lead to a positive bias.

By combining bias and random error, we obtain the RMSE. For Germany, the monthly results with far-field correction show an RMSE between 2.4% (posterior R) and 4.3% (diagonal R). For yearly totals, this reduces to 1.2% for posterior R and




Figure 7. Mean (a, c) and standard deviation (b, d) of monthly flux estimates relative to the prior in synthetic experiments with 20 % increased anthropogenic emissions in the synthetic truth for diagonal R (blue), prior R (orange), and posterior R inversion (green). In (a, c), the a priori has value 1.0 and a black vertical line shows the synthetic truth. Bars connect the prior to the posterior. Like in Fig. 6, each bar represents the posterior fluxes for 144 inversions, combining 12 months with 12 datasets of pseudo-observations. Horizontal lines show 2σ statistical uncertainties and colored lines in (b, d) indicate the posterior 1σ uncertainty.

1.8 % for diagonal R , while the prior R inversion is dominated by the bias and has an RMSE of 2.9 %. This indicates that the simulated transport error in our synthetic experiments leads to an error of approximately 2 % on the German yearly total emission estimate. Overall, the posterior R inversion shows the best performance as it has a lower random error and only a small bias.

4.3 Sensitivity tests (Sect. 5.4) to increased true emissions

To test the sensitivity of the inversion to true fluxes, we repeat the synthetic experiments with an identical setup but different pseudo-observations. For these new pseudo-observations, we increase all anthropogenic emissions by 20 %. The a priori emissions remain unchanged and are thus lower than the synthetic truth. The inversion results are summarized in Fig. 7, which is analogous to Fig. 6.

Figure 7 (a) and (c) show the mean posterior (bars) compared to the synthetic truth (black vertical line). Without the far-field correction, the inversion is too sensitive in many regions, as it increases the emissions beyond the synthetic truth. This leads to an overestimation, which is likely due to the artificial lifetime of the flux category tracers (see Sect. 2.1.2). With the far-field correction (panel c), the deviation of the posterior from the prior is damped and we obtain a low bias compared to the truth, as expected when the a priori emissions are underestimated. The random error (b, d) remains similar to the case with perfect prior emissions, albeit a small increase can be seen (compare Fig. 6). Like for the perfect prior emissions, the best performance with the lowest RMSE is found for the posterior R inversion.

4.4 Sensitivity to bias and noise in observations




Figure 8. Total posterior emissions in 2021 of selected countries and German sectors for synthetic experiments with perfect transport. Markers show the average of the emission estimates obtained from the prior R and posterior R inversion. Thin horizontal lines indicate the synthetic truth. Vertical lines show uncertainties (95% confidence intervals).

We now turn from the focus on the transport error to uncertainties in the observations. To this end, we consider different pseudo-observations without any transport error that follow scenarios defined in Fig. 8. To avoid the transport error, we generate these pseudo-observations based on the deterministic model run. For simplicity, we only consider the average of prior R and posterior R inversion.

550 In the first scenarios, we shift all pseudo-observations by -5 ppb (case 01 in Fig. 8) and $+5$ ppb (case 02). This bias is mostly compensated by the far-field correction with monthly averages of ± 2.75 ppb to ± 3.8 ppb, the sign depends on the scenario. Because of this correction, the effect on the estimated German total emissions remains well within the posterior uncertainty. This is in stark contrast to the same scenarios without the far-field correction (cases 03 and 04) and demonstrates the benefits of the far-field correction.

555 We furthermore test the effect of correlated and uncorrelated Gaussian noise added to the observations (cases 10–12), finding that the effect on the posterior emissions is small compared to the posterior uncertainties. The correlated Gaussian noise is a three-dimensional Gaussian random field in flat (longitude, latitude, time) coordinates with a lower cutoff for fluctuations on scales $\leq 2.5^\circ$ (horizontal) and ≤ 12 days (time) such that it acts as a slowly varying random bias. The RMS of the noise is normalized to 5 ppb. For the last three test cases (20–22), we scale either the natural and LULUCF fluxes or all other emissions in the synthetic truth while leaving the a priori emissions unchanged. Overall, the emission estimates follow the change in the synthetic truth well as already found in Sect. 4.3.

4.5 Sensitivity to inversion parameters

Our inversion method has various tuning parameters. Above we have described the inversion and its results synthetic experiments for one choice of these parameters. We analyzed the sensitivity of analyze the sensitivity to these parameters by repeating the

565 inversion ~~48 times with~~ 50 times with real observations and modified parameters. Table E1 lists these test cases with their ID, parameters, and influence on the inversion results. An overview of the national emission estimates for each test case is provided in Fig. E1. Here, we summarize the main results and refer to Table E1 for details. We use the average of the prior R and posterior R inversion results and focus on the influence of the parameters on the emission estimates, leaving the discussion of the inversion results for Part 2 (Bruch et al., 2025a).

570 4.5.1 Comparison to observations

~~The filtering steps listed in Table 4~~ Before comparing model and observations, we apply multiple filtering steps that influence the inversion results ~~significantly~~ considerably. Most prominently, selecting nighttime observations for high mountain stations and afternoon hours for other stations strongly affects the inversion and improves the model representativeness (case 201 in Table E1). This is one of only ~~four~~ five sensitivity tests with posterior fluxes deviating from the reference case by $\gtrsim 30\%$ of the posterior uncertainty, which we call a strong change in inversion results. Other filtering parameters such as the number of sampling heights used per station (case 202) and the minimal wind speed (cases 203–205) affect the inversion results noticeably, although changes are small compared to the uncertainties. ~~Neglecting extreme outliers has only a small effect (cases 206, 207), but limiting~~ Limiting the influence of outliers with model–observation mismatch $|\mu_i| > 3\sqrt{R'_{ii}}$ by increasing their uncertainty (see Sect. 2.6.2) has a considerable impact (cases 208, 209). Completely neglecting extreme outliers – defined
580 by $|y - Hs - x^{\text{ff}}| > 200 \text{ ppb}$ or $y - x^{\text{ff}} < -20 \text{ ppb}$ – has only a small effect (cases 206, 207).

The choice of observation sites is analyzed in cases 601 and 602, which select subsets of stations with good observation coverage over the full year. When using only 27 stations (case 602), the results change strongly compared to the reference case with 50 stations, also because some regions are hardly observed in case ~~602 (compare also Fig. 6 with Fig. A2).~~ 602. Varying the elevation of high mountain stations has only little impact on the inversion results (case 100). The effect of time-averaging
585 over 3 h (as chosen in step 2 of Sect. 3.3) is noticeable in the results, but small compared to the uncertainties (case 101).

4.5.2 Uncertainty

The diagonal R inversion deviates from the reference case by one third of the posterior uncertainty (case 311). Also the construction of the error covariance matrix R following ~~Sections 2.4 and 4.2~~ Sections 2.5 and 2.6 contains numerous tuning parameters. Key parameters are the overall uncertainty inflation factors f_i (Sect. 2.6.3, cases 302 and 303 in Table E1) and the
590 uncorrelated additive uncertainty σ_{const} (see Eq. (2)) of each data point (cases 309, 310). Variations of these parameters change the inversion results considerably. The tuning parameter σ_{const} illustrates the importance of hidden patterns in the considered data. Increasing to $\sigma_{\text{const}} = 20 \text{ ppb}$ effectively reduces the weight of observations with a small ensemble-estimated transport uncertainty. As observations with strong emission signals and high transport uncertainty become more relevant, the emission estimate for Germany is increased by 5 % (case 310 in Fig. E1).

595 Other important parameters are the correlation scales in the localization matrix C for the ensemble-based uncertainty estimate (see Sect. 2.5.2). The overall effect of these scales on the posterior scaling factors is small (cases 304–308), but these parameters also influence the posterior uncertainties. The sensitivity tests indicate that 12 ensemble members are sufficient

to estimate the uncertainties and correlations even without a strong localization. In general, we expect that a larger transport ensemble will yield better statistical estimates for uncertainties and their correlations. This reduces the need for a localization which suppresses spurious correlations. The considered additional plume localization uncertainty (see ~~Sect. 2.2~~Sect. 2.6.1, cases 300 and 301) arising from the Upper Silesian Coal Basin seems negligible when considering the full domain. However, the additional plume localization uncertainty reduces the negative bias for the plume emissions (~~see Sect. 5.6~~), that was discussed in Sect. 2.2.

4.5.3 Far-field correction

The synthetic experiments already showed that the far-field correction ~~explained in Sect. 2.2~~strongly introduced in Sect. 2.3 influences the results ~~– Removing considerably~~ (see Figs. 6 and 7). When using real observations, removing the correction field leads to ~~striking strong~~ changes in the inversion results ~~– including unrealistic negative~~ (case 400), albeit the results remain within the posterior uncertainty bounds. Without the correction, the scaling factors for some natural ~~flux categories (case 400)~~ fluxes in Scandinavia even become negative for some months – a clearly unrealistic result that underlines the importance of the far-field correction. However, changing various tuning parameters of the far-field correction within a reasonable range has much smaller effects. The selection of data points used for the far-field correction (cases 409, 410) and the overall correction strength (cases 401, 402) have modest influence, whereas correlation scales in the correction play a minor role (cases 403–408). The additional uncertainty added to R due to the far-field correction (see Sect. 2.6.4) has little influence on the inversion results (cases 412–414). ~~We draw the conclusions that (i) the far-field correction is important for the inversion and (ii) the inversion extracts most information from signals that are hardly affected by the precise form of the far-field correction, especially from strong emission signals ($\gg 20$ ppb).~~

4.5.4 A priori error covariance matrix

Modifying the a priori uncertainty or correlations of the scaling factors (B in Eq. (1)) changes the results quantitatively, but not qualitatively (cases 500–502). ~~We notice that a smaller a priori uncertainty (case 500) narrows the ability to discriminate sector emissions, because the sector attribution tends to follow the a priori uncertainties.~~ A coarser spatial resolution in Germany (case 504) and different choices of sectors (cases 503, 506) yield aggregated German sector emissions that agree well with the reference case.

4.5.5 Inversion time windows

In the reference case, we considered each month independently. Increasing the inversion time windows to three months has a considerable influence on the results (case 702). As the inversion time window increases, the overall weight of the observations in the inversion also increases. Thus, posterior uncertainties are reduced and the deviations between posterior and prior are amplified.

5 Discussion

[...]

630 5 Conclusions

We presented a novel system for regional flux inversion designed to validate national CH₄ emission reporting. Applying this method to Central Europe in 2021 with a focus on Germany. This study introduced a new flux inversion system that explores the potential of a transport ensemble from NWP for observation-based regional estimation of methane emissions. In experiments with pseudo-observations and simulated transport error, we found a significant increase in emissions from Germany and the Benelux. Careful estimation of posterior uncertainties revealed that total German posterior emissions are $(32 \pm 19)\%$ higher than the anthropogenic emissions reported to the UNFCCC (submission 2024). This increase is most likely due to emissions from the agriculture sector, possibly with contributions from LULUCF and natural sources. that using a transport ensemble can substantially reduce the random error of the flux estimates compared to a simple baseline scenario (“diagonal R ”). This is in line with findings by Ghosh et al. (2021) and by Steiner et al. (2024a), who estimated CH₄ emissions in Europe using an ensemble Kalman smoother. But in contrast to Ghosh et al. (2021), who studied CO₂ at urban scale using an ensemble transform Kalman filter, we identified no significant improvement in the bias of the emission estimates. Instead, our results indicate systematic biases depending on the emissions characteristics. Most notably, localized sources causing strong plumes can be underestimation by 10% by our synthesis inversion. To benefit from the transport ensemble and to reduce such biases, we proposed to use the posterior concentrations in the ensemble when constructing R . This posterior R inversion showed the best performance in the synthetic experiments. Our results were confirmed by an exhaustive range of sensitivity tests and by validation with independent observation sites. Synthetic experiments with known truth verified the ability to distinguish emission sectors in Germany. Overall, we expect an error of 2% for the total German CH₄ emissions in 2021 in our inversion system due to random transport errors.

Methodological comparison to other regional inversion systems highlights the advantages of our method for distinguishing emission sectors and fitness for purpose for validation of national emission estimates. The qualitative gap between UNFCCC reporting and our estimates for Germany and the Benelux is consistent with earlier works (Petrescu et al., 2023; Bergamaschi et al., 2022, 2023). We complement these studies by providing an emission estimate for the German agriculture sector that can be directly compared to the national reporting, revealing a significant mismatch.

When applying our regional inversion system to real observations, we face the challenge of uncertain CH₄ concentrations at the lateral boundaries. Different approaches exist to correct biased boundary conditions. In some cases, selected measurements can provide a baseline (Lauvaux et al., 2013). At national or continental scale, a coarse discretization of the boundaries allows optimization along with the emissions (Ganesan et al., 2015; Steiner et al., 2024b). Here, we followed a different path by adding a smooth correction field for the simulated concentrations. This allowed us to use different time scales for the inversion and the far-field correction. The far-field correction causes a small bias towards the prior fluxes, but without the correction we expect errors from wrongly projecting any boundary bias onto the fluxes. We demonstrated the potential of the

far-field correction using biased pseudo-observations and analyzed its importance in sensitivity tests, for which we repeated the inversion with different tuning parameters. These tests with real observations show that switch on the far-field correction changes the results considerably within the uncertainty ranges, but the specific choices made in constructing the correction field have only minor or moderate effects. Also other tested changes in tuning parameters only lead to variations of the full-year flux estimates well within the uncertainty ranges, indicating that we found robust settings for our application. This establishes a basis for applying our system to validate the German emission inventory in Part 2 (Bruch et al., 2025a).

In this study we only presented the first application of an extensible, novel inversion system. Future developments may include the integration of satellite data. The presented novel inversion system leverages the potential of the incorporation of temporal profiles, a more comprehensive treatment of boundary conditions and flux uncertainties using ensemble methods, and an extension of the state space. The close connections to operational numerical weather prediction — especially in the underlying transport simulation — and ICON-ART model and the ensemble modeling capabilities from operational NWP for national scale estimation of CH₄ fluxes. It is tailored to the modular design establish the potential for long-term operational support of national emissions reporting. validation of national inventories by using high-resolution a priori emission estimates from national reporting and allowing for distinguishing emission sectors, as will be discussed in detail in Part 2. With synthetic experiments and sensitivity tests we demonstrated the suitability for estimating national CH₄ emissions.

Data availability. A collection of model data, inversion results, and data for reproducing most figures in this work is available at <https://doi.org/10.5281/zenodo.17414768> (Bruch et al., 2025b).

Appendix A: Far-field Formal definition of far-field correction

This appendix provides details for the far-field correction introduced in Sect. 2.2. Sect. 2.3. We correct the computed far field by a smooth field that may only vary on temporal scales $\gtrsim 16$ h and horizontal scales $\gtrsim 320$ km. This correction field is determined using all data points where the cumulated signal of all flux categories is at most 20 ppb, the total concentration due to all fluxes in the domain – including natural and uncategorized fluxes – is at most 50 ppb, and natural plus LULUCF fluxes contribute at most 20 ppb. These criteria aim to select only measurements of sufficiently clean air for the far-field correction. Figure 2 shows that this correction lies typically in the range ± 10 ppb and has a bias of few ppb towards higher concentrations.




Figure B1. Statistical evaluation of the far-field correction. Considering all data points used in the inversion (steps 5–8 in Table 4), histograms of the far-field correction (a) and its time gradient (b) show that the correction is usually in the range

± 10 ppb and can vary by a few ppb per day. For each station and month, we compute the RMS and the mean (or bias). Histograms combining these values for all stations and months are shown in (c) and (d). The broad distribution of the RMS values indicates significant differences among the stations. The bias shows a tendency towards positive values, which implies that the far-field correction tends to increase the modeled CH_4 concentration. This increase in model concentrations leads to a decrease in emission estimates.

The far-field correction is realized as a Kalman smoother on the selected data points. For simplicity, we only provide the definition of the correction at the observation coordinates. Consider the vector of all model predictions x , which is aligned with the observation vector y . By P we denote the projector selecting those data points that shall be used to determine the far-field correction. We aim to find a correction vector c aligned with x and y that minimizes

$$\arg \min_c \left\{ \frac{1}{2} (x + c - y)^\top P^\top (P \tilde{R} P^\top)^{-1} P (x + c - y) + \frac{1}{2} c^\top P^\top (P \tilde{C} P^\top)^{-1} P c \right\}, \quad (\text{A1})$$

where $\tilde{R} = 16I$ is a diagonal matrix and $\tilde{C}$ ~~is an unnormalized (i.e. $\tilde{C}_{ii} = 1$ for all i)~~ a Gaussian localization matrix with standard deviations 16 h (time), 319 km (horizontal) and 1 km (vertical), normalize to $\tilde{C}_{ii} = 1$ for all i . The matrix $\tilde{C}$ ensures that the correction field c is smooth on these scales. For the under-determined Eq. (A1) we use the solution

$$c = \tilde{C} P^\top \left[P (\tilde{C} + \tilde{R}) P^\top \right]^{-1} P (y - x). \quad (\text{A2})$$

This only defines c at the observations, but we can generalize Eq. (A2) to arbitrary locations and times by including these coordinates in $\tilde{C}$. Formally, this then defines a smooth field.

To prove that Eq. (A2) ~~solves~~ is one possible – albeit not unique – solution of Eq. (A1), we use that Eq. (A1) is a quadratic form and compute its gradient with respect to c :

$$0 \stackrel{!}{=} \underline{2} P^\top (P \tilde{R} P^\top)^{-1} P (x + c - y) + \underline{2} P^\top (P \tilde{C} P^\top)^{-1} P c. \quad (\text{A3})$$

~~This can be solved by requiring~~

$$\begin{aligned} & \underline{0 \stackrel{!}{=} (P \tilde{R} P^\top)^{-1} P (x + c - y) + (P \tilde{C} P^\top)^{-1} P c} \\ & \underline{= \left[(P \tilde{R} P^\top)^{-1} + (P \tilde{C} P^\top)^{-1} \right] P c + (P \tilde{R} P^\top)^{-1} P (x - y)} \\ & \underline{\implies P c = \left[1 + P \tilde{R} P^\top (P \tilde{C} P^\top)^{-1} \right]^{-1} P (y - x)} \\ & \underline{= P \tilde{C} P^\top \left[P (\tilde{C} + \tilde{R}) P^\top \right]^{-1} P (y - x).} \end{aligned}$$

Since $PP^\top$ has full rank, this implies that

$$0 \stackrel{!}{=} \left[(P\tilde{R}P^\top)^{-1} + (P\tilde{C}P^\top)^{-1} \right] Pc + (P\tilde{R}P^\top)^{-1} P(x - y) \quad (\text{A4})$$

$$\Rightarrow Pc = \left[1 + P\tilde{R}P^\top (P\tilde{C}P^\top)^{-1} \right]^{-1} P(y - x) \quad (\text{A5})$$

$$= P\tilde{C}P^\top \left[P(\tilde{C} + \tilde{R})P^\top \right]^{-1} P(y - x). \quad (\text{A6})$$

It follows that Eq. (A2) is a solution of Eq. (A1) that is independent of the non-selected data points. One can furthermore show that Eq. (A2) is optimal in the sense that it minimizes $c^\top \tilde{C}^{-1} c$ under constraint that c is a solution of Eq. (A1). Thus, this solution is as close as possible to zero under the constraint of smoothness (quantified by $\tilde{C}$). By defining $\xi = \left[P(\tilde{C} + \tilde{R})P^\top \right]^{-1} P(y - x)$

and introducing Lagrange multipliers λ , we obtain

$$f(c, \lambda) = c^\top \tilde{C}^{-1} c + \lambda^\top (Pc - P\tilde{C}P^\top \xi), \quad \frac{\partial f}{\partial c_i} = 0, \quad \frac{\partial f}{\partial \lambda_j} = 0, \quad (\text{A7})$$

$$c = -\tilde{C}P^\top \lambda \quad \text{from } \partial_{c_i} f(c, \lambda) = 0, \quad (\text{A8})$$

$$Pc = P\tilde{C}P^\top \xi \quad \text{from } \partial_{\lambda_j} f(c, \lambda) = 0. \quad (\text{A9})$$

Since $P\tilde{C}P^\top$ has full rank, combining Eqs. (A8) and (A9) implies that $\lambda = -\xi$ and thereby $c = \tilde{C}P^\top \xi$ is the unique solution of the optimization problem $\arg \min_c f(c, 0)$ under the constraint that $Pc = P\tilde{C}P^\top \xi$.

Appendix B: ~~Posterior-based model uncertainty estimate~~ Posterior R with reduced ensemble

[...]

B1 Reduced ensemble

When using a priori scaling factors to estimate the model uncertainty in R , we need only the total concentration $x_i^m(s^{\text{prior}})$ for each ensemble member m and each observation i , where s^{prior} is known. Thus, only a single tracer field is required in the ensemble transport simulation. To ~~fully~~ compute $x_i^m(s)$ ~~as function of s , the tracer for arbitrary $s \in \mathbb{R}^{46}$, the flux~~ categories need to be distinguished for each ensemble member, resulting in > 40 tracer fields in the ensemble simulation. To avoid wasting numerical resources, we chose to approximate $x_i^m(s)$ by only a few tracer fields, using additional information from the deterministic model run which distinguishes all tracer fields.

From the deterministic model run, we know the operator H mapping scaling factors s to a model prediction $HS + x^{\text{ff}}$ for the concentrations. For ensemble member m , we would ideally know H^m and $x^{\text{ff}, m}$ yielding to compute a model prediction $H^m s + x^{\text{ff}, m}$. ~~To avoid calculating the full matrix~~ In lack of computational resources to compute H^m , we for every ensemble member, we combine information from the deterministic run (H) and selected tracers for the ensemble run to approximate

H^m . We group the flux categories into groups $\{g\}$ and denote by P_g the projector of scaling vectors s on the subspace spanned by the flux categories in group g . ~~Using the~~ In the ensemble members, we compute the total concentration from group g , ~~$x_i^{mg} = x_i^m(P_g s^{\text{prior}})$, we~~ $x_i^{mg} = H^m P_g s^{\text{prior}}$. We distribute the 46 flux categories to only three groups and thereby reduce the computational effort considerably. To estimate the full dependence on the scaling factors in the ensemble, we approximate:

$$x_i^m(s) \approx \sum_g \frac{(H P_g s)_i}{(H P_g s^{\text{prior}})_i} x_i^{mg} + x_i^{\text{ff},m}. \quad (\text{B1})$$

Thus, we compute the transport ensemble for a few tracer groups and estimate $x^m(s)$ for arbitrary s by using the ratios of tracer fields within the tracer groups from the deterministic run. Using the approximation in Eq. (B1), we estimate the posterior model uncertainties with only five tracer fields in an ensemble of 12 transport simulations:

1. far field (initial and lateral boundary conditions)
2. total anthropogenic fluxes
3. total natural fluxes
- 750 4. total anthropogenic fluxes from Germany with lifetime five days
5. total anthropogenic fluxes from outside Germany with lifetime five days

Appendix C: ~~Extended data tables and figures~~ Observation sites

~~[...]~~

Table C1. Observation stations from the European Obspack (ICOS RI et al., 2024). Column 6 (“mountain”) characterizes the stations as high mountains, small mountains, and other stations. This serves as a reference for computing the station height in the model and for the daily time window. We indicate the sampling heights used in the inversion (column 7) and mark those sampling heights with an asterisk that have good observation coverage in each month (used in sensitivity test 602). Column 8 indicates times in which the station was excluded due to modeling problems. Column 9 (“inflation”) defines the factor f_i of the static uncertainty inflation (see Sect. 2.6.3).

755

Code	Name	Coun- try	ICOS class	Elevation (m)	Mountain	Sampling heights (m)	Limitations	Infla- tion
BIK	Białystok	PL	–	183	no	90, 180, 300		2
BIR	Birkenes	NO	2	219	no	75	excl. Apr–Aug	3
BIS	Biscarrosse	FR	–	73	small	47*		2
BRM	Beromunster	CH	–	797	no	72, 132, 212		2
BSD	Bilsdale	UK	–	382	no	108, 248		2
CBW	Cabauw	NL	1	0	no	67, 127*, 207*		2
CMN	Monte Cimone	IT	2	2165	high	8		2
CRA	Centre de Recherches Atmosphériques	FR	–	600	no	60*		2
CRP	Carnsore Point	IE	–	9	no	14		2
ERS	Ersa	FR	–	533	small	40		3
FKL	Finokalia	GR	–	250	small	–	excluded	–
GAT	Gartow	DE	1	70	no	132*, 216*, 341*		2
HEI	Heidelberg	DE	–	113	no	30*		3
HEL	Helgoland	DE	2	43	no	110*		2
HPB	Hohenpeissenberg	DE	1	934	small	50, 93*, 131*		2
HTM	Hyltemossa	SE	1	115	no	70, 150		2
HUN	Hegyhátsál	HU	2	248	no	82, 115	incl. Mar–Oct	3
IPR	Ispira	IT	2	210	no	–	excluded	–
JFJ	Jungfraujoch	CH	1	3571.8	high	13.9		2
JUE	Jülich	DE	2	98	no	120*		3
KAS	Kasprowy Wierch	PL	–	1987	high	7*		2
KIT	Karlsruhe	DE	1	110	no	60*, 100*, 200*		2
KRE	Křešín u Pacova	CZ	1	534	no	50, 125, 250		2
LHW	Laegern-Hochwacht	CH	–	840	small	32		3
LIN	Lindenberg	DE	1	73	no	98		2
LMP	Lampedusa	IT	2	45	no	–	excluded	–

Code	Name	Coun-try	ICOS class	Elevation (m)	Mountain	Sampling heights (m)	Limitations	Infla-tion
LMU	La Muela	ES	–	571	no	79	excl. Nov–Dec	2
LUT	Lutjewad	NL	2	1	no	60		2
MHD	Mace Head	IE	–	5	no	24*		2
MLH	Malin Head	IE	–	22	no	47		2
NOR	Norunda	SE	1	46	no	58*, 100*		2
OHP	Observatoire de Haute Provence	FR	–	650	no	50, 100		2
OPE	Observatoire pérenne de l’environnement	FR	1	390	no	50*, 120*		2
OXK	Ochsenkopf	DE	1	1022	small	90, 163		2
PAL	Pallas	FI	1	565	no	12*		2
PDM	Pic du Midi	FR	–	2877	high	28		2
PRS	Plateau Rosa	IT	2	3480	high	10		2
PUI	Puijo	FI	2	232	small	84*		2
PUY	Puy de Dôme	FR	2	1465	small	10*		2
RGL	Ridge Hill	UK	2	207	no	90*		2
ROC	Roc’h Trédudon	FR	–	362	no	25, 80, 140		2
SAC	Saclay	FR	1	160	no	60*, 100*		2
SMR	Hyytiälä	FI	1	181	no	67.2*, 125*		2
SSL	Schauinsland	DE	2	1205	small	12, 35		2
STE	Steinkimmen	DE	1	29	no	127*, 187*, 252*		2
SVB	Svartberget	SE	1	269	no	85*, 150*		2
TAC	Tacolneston	UK	–	64	no	54*, 100*, 185*		2
TOH	Torfhaus	DE	2	801	small	76*, 110*, 147*		2
TRN	Trainou	FR	2	131	no	50*, 100*, 180*		2
UTO	Utö - Baltic sea	FI	2	8	no	57*		2
WAO	Weybourne	UK	2	17	no	10*		2
WES	Westerland	DE	2	12	no	14		2
ZSF	Zugspitze	DE	2	2666	high	3*		2

Chi-square-

Appendix D: χ^2 analysis

In this appendix, we provide the mathematical details for the χ^2/N_{dof} analysis ~~used in Sect. 4.2.~~ (see, e.g., Greenwood and Nikulin, 1996) ~~used in Sect. 2.6.5.~~ The aim of this analysis is to quantify whether the data used in the inversion agree with the assumed uncertain-

ties. ~~The inversion formally relies on~~ We restrict this analysis to the prior R and diagonal R inversion, for which the matrix R is constant. These inversions formally rely on the assumption of Gaussian probability distributions of the a priori scaling factors (error covariance matrix B) and the model–observation mismatch (R). ~~For the~~

We start from the probability density of observations y under the assumption that s describes the true emissions:

$$765 \quad P(y|s) \propto \exp \left[-\frac{1}{2} (y - Hs - x^{\text{ff}})^\top R^{-1} (y - Hs - x^{\text{ff}}) \right]. \quad (\text{D1})$$

Like in the inversion, R describes uncertainties in the transport, in the corrected far-field contribution x^{ff} , and in the observations y . By a change of variables we obtain the probability for the a priori model–observation mismatch $\mu = y - Hs^{\text{prior}} - x^{\text{ff}}$ this yields the probability $\mu^{\text{prior}} = y - Hs^{\text{prior}} - x^{\text{ff}}$: $P(\mu^{\text{prior}}|s) d\mu = P(y|s)|_{y=Hs^{\text{prior}}+x^{\text{ff}}+\mu^{\text{prior}}} dy$.

To estimate whether a given μ^{prior} is realistic, we need to integrate out the scaling factors s to obtain $P(\mu^{\text{prior}})$. We denote the
770 integral over the vector space of scaling factors s with probability measure dP_s by $\int_s \bullet dP_s = \int_s P(s) \bullet d^n s$ for $s \in \mathbb{R}^n$. Using the above definitions in Eq. (D1), we obtain³ (Berchet et al., 2015)

$$\begin{aligned} P(\mu = y - Hs^{\text{prior}} - x^{\text{ff}}) \\ = \int_s P(y = Hs + \mu + x^{\text{ff}} | s) dP_s \end{aligned} \quad (\text{D2})$$

$$\propto \int_s \exp \left[-\frac{1}{2} (y - Hs - x^{\text{ff}})^\top R^{-1} (y - Hs - x^{\text{ff}}) - \frac{1}{2} (s - s^{\text{prior}})^\top B^{-1} (s - s^{\text{prior}}) \right]_{y=Hs^{\text{prior}}+x^{\text{ff}}+\mu^{\text{prior}}} d^n s \quad (\text{D3})$$

$$775 \quad \stackrel{\tau = s - s^{\text{prior}}}{=} \int_\tau \exp \left[-\frac{1}{2} (H\tau + \mu^{\text{prior}} - H\tau)^\top R^{-1} (H\tau + \mu^{\text{prior}} - H\tau) - \frac{1}{2} \tau^\top B^{-1} \tau \right] d^{N_{\text{dof}}} \tau \quad (\text{D4})$$

$$\propto \exp \left[-\frac{1}{2} \mu^{\text{prior}^\top} (R^{-1} - R^{-1} H (B^{-1} + H^\top R^{-1} H)^{-1} H) H^\top R^{-1} \mu^{\text{prior}} \right] \quad (\text{D5})$$

$$=: \exp \left(-\frac{1}{2} \mu^{\text{prior}^\top} Q \mu^{\text{prior}} \right). \quad (\text{D6})$$

~~Such~~ This result is a high-dimensional Gaussian probability distributions share various useful properties. Here, we use that when sampling distribution, $\mu^{\text{prior}} \sim \mathcal{N}(0, Q^{-1})$. When drawing a random vector μ from a probability distribution $P(\mu)$ as
780 in Eq. (D6), it is very likely to find μ such that $\chi^2 \equiv \mu^\top Q \mu \approx N_{\text{dof}}$ where N_{dof} denotes the number of degrees of freedom, which is the dimension of vector μ . In our case, $N_{\text{dof}} \sim 10^4$ is the number of observation data points used per one-month time window. In the limit of large N_{dof} , one can approximate ~~$P(\chi^2) \sim \mathcal{N}(N_{\text{dof}}, 2N_{\text{dof}})$ the probability distribution for χ^2 by~~ $\chi^2 \sim \mathcal{N}(N_{\text{dof}}, 2N_{\text{dof}})$ (Gaussian distribution with mean N_{dof} and variance $2N_{\text{dof}}$) (Abramowitz and Stegun, 1964, Sect. 26.4). Thus, in an idealized setup we expect that $\chi^2/N_{\text{dof}} = 1 \pm 0.03$ (95% confidence interval). Values $\gtrsim 1.05$ ~~imply that uncertainties~~
785 ~~were underestimated~~ hint at underestimated uncertainties and $\chi^2/N_{\text{dof}} \lesssim 0.95$ indicates that uncertainties were too high. However, in reality we may have biases and other problems not fully described errors such that the assumption of a Gaussian

³In Eq. (D5), we first solve the Gaussian integral to obtain $\exp \left\{ -\frac{1}{2} \mu^{\text{prior}^\top} [R^{-1} - R^{-1} H (B^{-1} + H^\top R^{-1} H)^{-1} H^\top R^{-1}] \mu^{\text{prior}} \right\}$ and then use that $(R + H B H^\top) [R^{-1} - R^{-1} H (B^{-1} + H^\top R^{-1} H)^{-1} H^\top R^{-1}] = I$.

uncertainty in the model–observation mismatch becomes invalid and $\chi^2/N_{\text{dof}} < 1$ does not necessarily imply that uncertainties ~~should~~can simply be reduced.

Appendix E: Sensitivity tests

790 Table E1 provides an overview of the sensitivity tests. For this table, we quantify the impact of a parameter variation on the inversion results by the following, heuristic metric: Consider a fixed region, sector and inversion time window with posterior fluxes F , defined as the average of the prior R and posterior R inversion result. The normalized deviation from the reference inversion is defined as $\Delta = \frac{2|F - F^{\text{ref.}}|}{F^{\text{ref. upper}} - F^{\text{ref. lower}}}$, where $F^{\text{ref. upper}}$ and $F^{\text{ref. lower}}$ denote the bounds of the posterior uncertainty range. The overall impact is computed as the arithmetic mean of Δ over the (usually monthly) time windows and a selection of
795 regions and sectors. In the regions UK+Ireland, France, Italy, Poland, Austria+Czechia, Netherlands, Belgium+Luxembourg, Switzerland, and Denmark we consider only total fluxes without distinguishing sectors. In Germany we include Δ for the total fluxes in four different regions (north, east, south, west) and additionally for national total fluxes distinguishing the three sectors agriculture, natural plus LULUCF, and other sectors. Effectively, this counts all fluxes in Germany twice and gives them more weight in the impact metric for Table E1.

800

Table E1. Sensitivity tests for estimating the robustness of the inversion results with respect to tuning parameters. Modified numbers are marked in bold font. The impact column quantifies the deviation of the inversion results relative to the uncertainties and shall qualitatively indicate the relevance of the modified parameters (see explanation in the text). An impact of 100 % means that the average deviation from the reference case is as large as the posterior uncertainty. Overall, we see that most tests have an impact of $\lesssim 15\%$, implying that the effect on the inversion results is small compared to the uncertainty in the reference case. See also Fig. E1 for the posterior emissions in the sensitivity tests.

ID	Test case	Explanation	Impact
0	reference	<u>reference case</u> as explained in Seet. 4 <u>Sections 2 and 3 and discussed in Part 2 (Bruch et al., 2025a), uses 129 117 observations in 2021</u>	
Model equivalent calculation (see Sect. 3.3)			
100	station elevation for mountain stations	treat all mountain stations like small mountains when computing model heights, as proposed by Brunner et al. (2012); Henne et al. (2016); Bergamaschi et al. (2022), <u>uses 127 087 observations</u>	5.3 %
101	no <u>additional</u> time averaging	average only over 1 h instead of over like in the observations, instead of averaging 3 h	13 %
Filtering observations (see Sect. 3.3)			
200	fewer hours of day	use time window 12 h–16 h (0 h–4 h for high mountains), <u>85 674 observations (reference case uses 11 h–17 h / 23 h–5 h)</u>	11 %
201	all hours of day	no filtering by time of day, increase uncertainty inflation <u>(factors f_i in Sect. 2.6.3) by factor 1.5, uses 508 594 observations</u>	38 %
202	one sampling height per station	use only highest sampling height of each station <u>instead of up to 3 highest levels, 80 132 observations</u>	16 %
203	no filtering based on wind	include data points with low wind speed, <u>147 019 observations</u>	12 %
204	low min. wind speed	minimum wind speed: 1.11 ms^{-1} <u>(reference: 2 ms^{-1}), 140 650 obs.</u>	9.4 %
205	high min. wind speed	minimum wind speed: 3.0 ms^{-1} <u>(reference: 2 ms^{-1}), 112 275 obs.</u>	11 %
206	low max. model-obs. mismatch	discard when absolute deviation exceeds 120 ppb, or model far field minus observation exceeds 12 ppb <u>$y - Hs - x^{\text{ff}} > 120 \text{ ppb}$ or $y - x^{\text{ff}} < -12 \text{ ppb}$, 127 055 obs. (reference case: 200 ppb / -20 ppb)</u>	3.5 %
207	high max. model-obs. mismatch	discard when absolute deviation exceeds 300 ppb, or model far field minus observation exceeds 30 ppb <u>$y - Hs - x^{\text{ff}} > 300 \text{ ppb}$ or $y - x^{\text{ff}} < -30 \text{ ppb}$, 129 706 obs.</u>	1.3 %
208	low max. data point influence	increase uncertainty if $\mu_i > 2.5 \sqrt{R_{ii}}$ in Sect. 2.4.1 <u>$\mu_i > 2.5 \sqrt{R_{ii}^{\text{step 1}}}$ in Sect. 2.6.2 (reference value: 3)</u>	11 %
209	high max. data point influence	increase uncertainty if $\mu_i > 4 \sqrt{R_{ii}}$ in Sect. 2.4.1 <u>$\mu_i > 4 \sqrt{R_{ii}^{\text{step 1}}}$ in Sect. 2.6.2 (reference value: 3)</u>	15 %

ID	Test case	Explanation	Impact
Uncertainty / error covariance matrix R (see Sections 2.5 and 2.6)			
300	no plume uncertainty	no extra uncertainty due to point-like emissions <u>localized emissions</u> (Sect. 2.6.1)	0.27 %
301	high plume uncertainty	uncertainty due to point-like emissions: use $R'_{ij} + 0.5\rho_i^2\delta_{ij}$ in Sect. 4.2 <u>extra uncertainty: $R_{ij}^{\text{step 1}} = R'_{ij} + 0.5\rho_i^2\delta_{ij}$ in Sect. 2.6.1 (reference: 0.25)</u>	0.56 %
302	low uncertainty inflation	uncertainty inflation by (1.5, 2.25) <u>instead of (2, 3)</u> $f_i = 1.5$ or 2.25 <u>instead of 2 or 3 in Sect. 2.6.3</u>	8.6 %
303	high uncertainty inflation	uncertainty inflation by (3, 4.5) <u>instead of (2, 3)</u> $f_i = 3$ or 4.5 <u>instead of 2 or 3 in Sect. 2.6.3</u>	13 %
304	small horizontal error correlation scale	scale 191 km instead of 319 km <u>in localization matrix C_{ij} (Sect. 2.5.2)</u>	6.0 %
305	large horizontal error correlation scale	scale 510 km instead of 319 km <u>in localization matrix C_{ij} (Sect. 2.5.2)</u>	8.3 %
306	small vertical error correlation scale	scale 400 m instead of 1 km <u>in localization matrix C_{ij} (Sect. 2.5.2)</u>	2.3 %
307	short error correlation time scale	scale 4 h instead of 6 h <u>in localization matrix C_{ij} (Sect. 2.5.2)</u>	2.5 %
308	long error correlation time scale	scale 10 h instead of 6 h <u>in localization matrix C_{ij} (Sect. 2.5.2)</u>	2.8 %
309	low uncorrelated uncertainty	$\sigma_{\text{const}} = 5$ ppb instead of 10 ppb in Eq. (2)	21 %
310	high uncorrelated uncertainty	$\sigma_{\text{const}} = 20$ ppb instead of 10 ppb in Eq. (2)	22 %
311	<u>diagonal R without ensemble</u>	<u>see Sect. 2.5.1</u>	<u>33 %</u>
height	Far-field correction (see Sect. 2.3 and Appendix A)		
400	no far-field correction		97 <u>35</u> %
401	weak far-field correction	$\tilde{R} = 100I$ instead of $16I$ in Eq. (A1)	16 %
402	strong far-field correction	$\tilde{R} = 2.78I$ instead of $16I$ in Eq. (A1)	9.2 %
403	small horiz. far-field correction scale	scale 191 km instead of 319 km <u>in localization matrix $\tilde{C}_{ij}$ in Appendix A</u>	6.8 %
404	large horiz. far-field correction scale	scale 510 km instead of 319 km <u>in localization matrix $\tilde{C}_{ij}$ in Appendix A</u>	4.5 %
405	short far-field correction time scale	time scale 10 h instead of 16 h <u>in localization matrix $\tilde{C}_{ij}$ in Appendix A</u>	3.7 %
406	long far-field correction time scale	time scale 28 h instead of 16 h <u>in localization matrix $\tilde{C}_{ij}$ in Appendix A</u>	3.8 %
407	long <u>extra-long</u> far-field correction time scale	time scale 48 h instead of 16 h <u>in localization matrix $\tilde{C}_{ij}$ in Appendix A</u>	7.1 %
408	low vertical far-field correction scale	scale 400 m <u>instead of 1 km in localization matrix $\tilde{C}_{ij}$ in Appendix A</u>	0.92 %
409	strict far-field observation selection	max. signal 10 ppb, max. due to natural fluxes 10 ppb <u>construct far-field correction based on observations with cumulated signal from categorized fluxes ≤ 10 ppb (reference: 20 ppb) and from natural fluxes ≤ 10 ppb (reference: 20 ppb)</u>	20 %

ID	Test case	Explanation	Impact
410	loose far-field observation selection	max. signal 30 ppb, max. due to natural fluxes 30 ppb, max. uncategorized 80 ppb far-field correction uses observations with cumulated signal from categorized fluxes ≤ 30 ppb (ref.: ≤ 20 ppb), from natural fluxes ≤ 30 ppb (ref.: 20 ppb), and from all emissions within the domain ≤ 80 ppb (ref.: 50 ppb)	14 %
411	unrestricted iterative far-field correction	max. signal 50 ppb, no other selection criteria; $\tilde{C}$ far-field correction uses all observations with cumulated signal from categorized fluxes ≤ 50 ppb; $\tilde{C}_{ij}$ uses localization scales 10 h, 191 km; iterate far-field correction and inversion are iterated 3 times, the correction always uses the posterior concentrations from the previous iteration. This aggressively suppresses large scale signals (biases) in the observations.	30 %
412	low correction uncertainty	use $R_{ij} + 0.25 c_i c_j \tilde{C}_{ij}$ in Sect. 4.2 $R_{ij}^{\text{step } 4} = R_{ij}^{\text{step } 3} + 0.25 c_i c_j \tilde{C}_{ij}$ in Sect. 2.6.4 (reference value: 0.5)	2.5 %
413	high correction uncertainty	use $R_{ij} + 1.0 c_i c_j \tilde{C}_{ij}$ in Sect. 4.2 $R_{ij}^{\text{step } 4} = R_{ij}^{\text{step } 3} + 1.0 c_i c_j \tilde{C}_{ij}$ in Sect. 2.6.4 (reference value: 0.5)	4.2 %
414	uncorrelated correction uncertainty	use $R_{ij} + 2c_i^2 \delta_{ij}$ instead of $R_{ij} + 0.5 c_i c_j \tilde{C}_{ij}$ in Sect. 4.2 $R_{ij}^{\text{step } 4} = R_{ij}^{\text{step } 3} + 2c_i^2 \delta_{ij}$ in Sect. 2.6.4	3.6 %
A priori scaling factor error covariance matrix B (see Sect. 2.8)			
500	low prior uncertainty	1σ prior uncertainty for most areas 0.25, set to 0.25 (ref.: 0.4) for well-observed areas, 0.2 (ref.: 0.25) for remote and plume 0.2, sector-resolved 0.33 categories, 0.33 (ref.: 0.5) for sector-resolving categories	14 %
501	high prior uncertainty in Germany	1σ prior uncertainty such that <u>national</u> total sector emissions in Germany have uncertainty 0.6 1σ uncertainty 60% for each distinguished sector (reference: approx. 40%)	8.6 %
502	uncorrelated prior, B is diagonal	1σ prior uncert. uncertainty in sector categories in Germany: 0.75 <u>0.75</u> ; uncertainty on national total: 35% for agriculture, 39% for other anthropogenic)	6.35.6 %
503	no sector distinction in prior	four regions in Germany with uncorrelated 1σ prior uncertainty of 0.4 <u>0.4</u>	7.7 %
504	low spatial resolution in Germany	two initially uncorrelated regions in Germany (<u>south-west and north-east</u>), each distinguishing sectors like in the reference case	15 %
506	distinguish 5 sectors in Germany	see Appendix I split “non-agr.” into sectors waste, public power, and other emissions	2.1 %
Station selection			
601	stations covering require full-year coverage	require ≥ 10 days each month use coverage each month ; 35 of 50 stations, 105 701 obs.	13 %




Figure E1. Posterior emissions and uncertainties of selected countries and German sectors for all sensitivity tests. Thin horizontal lines indicate the posterior of the reference case 0. Markers show the average of prior R and posterior R inversion. Vertical lines show uncertainties (95% confidence intervals) that are not extended by excluding stations and cover the uncertainty range of prior R and posterior R inversion. The individual tests are listed in Table E1. One can clearly see the strongest deviations for cases 400 (no far-field correction), whereas for all other test cases only lead to changes, the emission estimates for the shown countries remain within the uncertainty range of the reference case.

ID	Test case	Explanation	Impact
602	<u>stations covering require good full-year coverage</u>	<u>require ≥ 20 days each month use coverage each month; 27 of 50 stations as detailed, 82912 observations (discussed in Fig. A2 of Part 2)</u>	33 %
Inversion time windows (see Sect. 2.7)			
701	2 month inversion window	<u>uncertainties are not adjusted to the longer window</u>	12 %
702	3 month inversion window	<u>uncertainties are not adjusted to the longer window</u>	18 %

Author contributions. VB and TR conceptualized the inversion method. VB implemented the inversion method and wrote the original draft together with AKW. TR configured the transport model. TR and BE interpolated the a priori flux data which BE collected. DJCO organized data streams of CH₄ concentrations and observations. JF, BM, AMB, DJCO, TR and VB contributed to testing and tuning the transport model. NB contributed to the model–observation comparison. AKW supervised and coordinated the project. All authors reviewed and edited the manuscript.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. In our simulations we use modified Copernicus Atmosphere Monitoring Service information and ECCAD products for initial and lateral boundary conditions, and for a priori fluxes. We thank Stefan Feigenspan, Christian Mielke, Theo Wernicke, John Akubia and Roland Fuß for helpful discussions and providing a priori emission fields. We thank Roland Potthast, Frank-Thomas Koch, Christoph Gerbig, Dominik Brunner, Michael Steiner, David Ho, Thomas Kaminski, Hannes Imhof and our partners in the ITMS project for very helpful and inspiring discussions. We also wish to thank Peter Bergamaschi, Aurélie Colomb, Martine De Mazière, Lukas Emmenegger, Dagmar Kubistin, Irene Lehner, Kari Lehtinen, Markus Leuenberger, Cathrine Lund Myhre, Michal V. Marek, Simon O’Doherty, Stephen M. Platt, Christian Plaß-Dülmer, Francesco Apadula, Sabrina Arnold, Pierre-Eric Blanc, Dominik Brunner, Huilin Chen, Lukasz Chmura, Lukasz Chmura, Sébastien Conil, Cédric Couret, Paolo Cristofanelli, Grant Forster, Arnoud Frumau, Christoph Gerbig, François Gheusi, Samuel Hammer, Laszlo Haszpra, Juha Hatakka, Michal Heliasz, Stephan Henne, Arjan Hensen, Antje Hoheisel, Tobias Kneuer, Eric Larmanou, Tuomas Laurila, Ari Leskinen, Ingeborg Levin, Matthias Lindauer, Morgan Lopez, Ivan Mammarella, Giovanni Manca, Andrew Manning, Damien Martin, Frank Meinhardt, Meelis Mölder, Jennifer Müller-Williams, Steffen Manfred Noe, Jarosław Nęcki, Mikael Ottosson-Löfvenius, Carole Philippon, Joseph Pitt, Michel Ramonet, Pedro Rivas-Soriano, Bert Scheeren, Marcus Schumacher, Mahesh Kumar Sha, Gerard Spain, Martin Steinbacher, Lise Lotte Sørensen, Alex Vermeulen, Gabriela Vítková, Irène Xueref-Remy, Alcide di Sarra, Franz Conen, Victor Kazan, Yves-Alain Roulet, Tobias Biermann, Marc Delmotte, Daniela Heltai, Ove Hermansen, Kateřina Komínková, Olivier Laurent, Janne Levula, Chris Lunder, Per Marklund, Josep-Anton Morguí, Jean-Marc Pichon, Martina Schmidt, Damiano Sferlazzo, Paul Smith, Kieran Stanley, Pamela Trisolino and Giulia Zazzeri for providing the atmospheric observations for the stations listed in Table C1. VB, DJCO, NB and AMB acknowledge funding by the German Federal Ministry for Education and Research (BMBF) in the ITMS project (grant 01LK2102B) as well as BE (grant 01LK2104A). Map plots were made with Natural Earth.

References

- Abramowitz, M. and Stegun, I. A.: Handbook of Mathematical Functions With Formulas, Graphs, and Mathematical Tables, Applied Mathematics Series, United States Department of Commerce, National Bureau of Standards, Washington, D.C., 1964.
- Berchet, A., Pison, I., Chevallier, F., Bousquet, P., Bonne, J.-L., and Paris, J.-D.: Objectified quantification of uncertainties in Bayesian
835 atmospheric inversions, *Geosci. Model Dev.*, 8, 1525–1546, <https://doi.org/10.5194/gmd-8-1525-2015>, 2015.
- Bergamaschi, P., Krol, M., Meirink, J. F., Dentener, F., Segers, A., van Aardenne, J., Monni, S., Vermeulen, A. T., Schmidt, M., Ramonet, M., Yver, C., Meinhardt, F., Nisbet, E. G., Fisher, R. E., O’Doherty, S., and Dlugokencky, E. J.: Inverse modeling of European CH₄ emissions 2001–2006, *J. Geophys. Res. Atmos.*, 115, <https://doi.org/10.1029/2010JD014180>, 2010.
- Bergamaschi, P., Corazza, M., Karstens, U., Athanassiadou, M., Thompson, R. L., Pison, I., Manning, A. J., Bousquet, P., Segers, A.,
840 Vermeulen, A. T., Janssens-Maenhout, G., Schmidt, M., Ramonet, M., Meinhardt, F., Aalto, T., Haszpra, L., Moncrieff, J., Popa, M. E., Lowry, D., Steinbacher, M., Jordan, A., O’Doherty, S., Piacentino, S., and Dlugokencky, E.: Top-down estimates of European CH₄ and N₂O emissions based on four different inverse models, *Atmos. Chem. Phys.*, 15, 715–736, <https://doi.org/10.5194/acp-15-715-2015>, 2015.
- Bergamaschi, P., Karstens, U., Manning, A. J., Saunio, M., Tsuruta, A., Berchet, A., Vermeulen, A. T., Arnold, T., Janssens-Maenhout, G., Hammer, S., Levin, I., Schmidt, M., Ramonet, M., Lopez, M., Lavric, J., Aalto, T., Chen, H., Feist, D. G., Gerbig, C., Haszpra, L.,
845 Hermansen, O., Manca, G., Moncrieff, J., Meinhardt, F., Necki, J., Galkowski, M., O’Doherty, S., Paramonova, N., Scheeren, H. A., Steinbacher, M., and Dlugokencky, E.: Inverse modelling of European CH₄ emissions during 2006–2012 using different inverse models and reassessed atmospheric observations, *Atmos. Chem. Phys.*, 18, 901–920, <https://doi.org/10.5194/acp-18-901-2018>, 2018.
- Bergamaschi, P., Segers, A., Brunner, D., Haussaire, J.-M., Henne, S., Ramonet, M., Arnold, T., Biermann, T., Chen, H., Conil, S., Delmotte, M., Forster, G., Frumau, A., Kubistin, D., Lan, X., Leuenberger, M., Lindauer, M., Lopez, M., Manca, G., Müller-Williams, J., O’Doherty, S., Scheeren, B., Steinbacher, M., Trisolino, P., Vítková, G., and Yver Kwok, C.: High-resolution inverse modelling of European CH₄ emissions using the novel FLEXPART-COSMO TM5 4DVAR inverse modelling system, *Atmos. Chem. Phys.*, 22, 13 243–13 268, <https://doi.org/10.5194/acp-22-13243-2022>, 2022.
- Bruch, V., Rösch, T., Jiménez de la Cuesta Otero, D., Ellerhoff, B., Mamtimin, B., Becker, N., Blechschmidt, A.-M., Förstner, J., and Kaiser-Weiss, A. K.: German methane fluxes estimated top-down using ICON–ART – Part 2: Inversion results for 2021, submitted to ACP,
855 2025a.
- Bruch, V., Rösch, T., Jiménez de la Cuesta Otero, D., Ellerhoff, B., Mamtimin, B., Becker, N., Blechschmidt, A.-M., Förstner, J., and Kaiser-Weiss, A. K.: German methane fluxes in 2021 estimated with an ensemble-enhanced scaling inversion based on the ICON–ART model, Zenodo [data set], <https://doi.org/10.5281/zenodo.15083479>, 2025b.
- Brunner, D., Henne, S., Keller, C. A., Vollmer, M. K., Reimann, S., and Buchmann, B.: Estimating European Halocarbon Emissions Using Lagrangian Backward Transport Modeling and in Situ Measurements at the Jungfrauoch High-Alpine Site, in: *Lagrangian Modeling of the Atmosphere*, pp. 207–222, John Wiley & Sons, Ltd., Chichester, England, UK, ISBN 978-1-11870457-8, <https://doi.org/10.1029/2012GM001258>, 2012.
- Chen, H. W., Zhang, F., Lauvaux, T., Davis, K. J., Feng, S., Butler, M. P., and Alley, R. B.: Characterization of Regional-Scale CO₂ Transport Uncertainties in an Ensemble with Flow-Dependent Transport Errors, *Geophys. Res. Lett.*, 46, 4049–4058,
865 <https://doi.org/10.1029/2018GL081341>, 2019.

- Conn, A. R., Gould, N. I. M., and Toint, Ph. L.: Trust Region Methods, MOS-SIAM Series on Optimization, Society for Industrial and Applied Mathematics (SIAM, 3600 Market Street, Floor 6, Philadelphia, PA 19104), Philadelphia, PA, USA, ISBN 978-0-89871985-7, 2000.
- Dammers, E., Tokaya, J., Mielke, C., Hausmann, K., Griffin, D., McLinden, C., Eskes, H., and Timmermans, R.: Can TROPOMI NO₂ satellite data be used to track the drop in and resurgence of NO_x emissions in Germany between 2019–2021 using the multi-source plume method (MSPM)?, *Geosci. Model Dev.*, 17, 4983–5007, <https://doi.org/10.5194/gmd-17-4983-2024>, 2024.
- Engelen, R. J., Denning, A. S., and Gurney, K. R.: On error estimation in atmospheric CO₂ inversions, *J. Geophys. Res. Atmos.*, 107, ACL10–1–ACL10–13, <https://doi.org/10.1029/2002JD002195>, 2002.
- Enting, I. G.: Inverse Problems in Atmospheric Constituent Transport, Cambridge University Press, Cambridge, England, UK, ISBN 978-0-52181210-8, <https://doi.org/10.1017/CBO9780511535741>, 2002.
- Feigenspan, S., Wernicke, T., and Mielke, C.: Personal communication, 2024.
- Fuß, R. and Akubia, J.: Personal communication, 2024.
- Ganesan, A. L., Manning, A. J., Grant, A., Young, D., Oram, D. E., Sturges, W. T., Moncrieff, J. B., and O’Doherty, S.: Quantifying methane and nitrous oxide emissions from the UK and Ireland using a national-scale monitoring network, *Atmos. Chem. Phys.*, 15, 6393–6406, <https://doi.org/10.5194/acp-15-6393-2015>, 2015.
- Gerbig, C., Körner, S., and Lin, J. C.: Vertical mixing in atmospheric tracer transport models: error characterization and propagation, *Atmos. Chem. Phys.*, 8, 591–602, <https://doi.org/10.5194/acp-8-591-2008>, 2008.
- Ghosh, S., Mueller, K., Prasad, K., and Whetstone, J.: Accounting for Transport Error in Inversions: An Urban Synthetic Data Experiment, *Earth Space Sci.*, 8, e2020EA001 272, <https://doi.org/10.1029/2020EA001272>, 2021.
- Greenwood, P. E. and Nikulin, M. S.: A Guide to Chi-Squared Testing, Wiley Series in Probability and Statistics, Wiley, ISBN 9780471557791, 1996.
- Henne, S., Brunner, D., Oney, B., Leuenberger, M., Eugster, W., Bamberger, I., Meinhardt, F., Steinbacher, M., and Emmenegger, L.: Validation of the Swiss methane emission inventory by atmospheric observations and inverse modelling, *Atmos. Chem. Phys.*, 16, 3683–3710, <https://doi.org/10.5194/acp-16-3683-2016>, 2016.
- ICOS RI: ICOS Atmosphere Station Specifications V2.0 (editor: O. Laurent), <https://doi.org/10.18160/GK28-2188>, 2020.
- ICOS RI, Bergamaschi, P., Colomb, A., De Mazière, M., Emmenegger, L., Kubistin, D., Lehner, I., Lehtinen, K., Leuenberger, M., Lund Myhre, C., Marek, M. V., O’Doherty, S., Platt, S. M., Plaß-Dülmer, C., Apadula, F., Arnold, S., Blanc, P.-E., Brunner, D., Chen, H., Chmura, L., Chmura, Ł., Conil, S., Couret, C., Cristofanelli, P., Forster, G., Frumau, A., Gerbig, C., Gheusi, F., Hammer, S., Haszpra, L., Hatakka, J., Heliasz, M., Henne, S., Hensen, A., Hoheisel, A., Kneuer, T., Larmanou, E., Laurila, T., Leskinen, A., Levin, I., Lindauer, M., Lopez, M., Mammarella, I., Manca, G., Manning, A., Martin, D., Meinhardt, F., Mölder, M., Müller-Williams, J., Noe, S. M., Nęcki, J., Ottosson-Löfvenius, M., Philippon, C., Pitt, J., Ramonet, M., Rivas-Soriano, P., Scheeren, B., Schumacher, M., Sha, M. K., Spain, G., Steinbacher, M., Sørensen, L. L., Vermeulen, A., Vítková, G., Xueref-Remy, I., di Sarra, A., Conen, F., Kazan, V., Roulet, Y.-A., Biermann, T., Delmotte, M., Heltai, D., Hermansen, O., Komínková, K., Laurent, O., Levula, J., Lunder, C., Marklund, P., Morguí, J.-A., Pichon, J.-M., Schmidt, M., Sferlazzo, D., Smith, P., Stanley, K., Trisolino, P., Zazzeri, G., ICOS Carbon Portal, ICOS Atmosphere Thematic Centre, ICOS Flask And Calibration Laboratory, and ICOS Central Radiocarbon Laboratory: European Obspack compilation of atmospheric methane data from ICOS and non-ICOS European stations for the period 1984–2024; obspack_ch4_466_GVeu_v9.2_20240502, <https://doi.org/10.18160/9B66-SQM1>, 2024.

- IPCC, Calvo Buendia, E., Tanabe, K., Kranjc, A., Baasansuren, J., Fukuda, M., Ngarize, S., Osako, A., Pyrozhenko, Y., Shermanau, P., and Federici, S., eds.: 2019 Refinement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories, vol. 1, The Intergovernmental Panel on Climate Change (IPCC), <https://www.ipcc-nggip.iges.or.jp/public/2019rf/index.html>, 2019.
- Jähn, M., Kuhlmann, G., Mu, Q., Haussaire, J.-M., Ochsner, D., Osterried, K., Clément, V., and Brunner, D.: An online emission module for atmospheric chemistry transport models: implementation in COSMO-GHG v5.6a and COSMO-ART v5.1-3.1, *Geosci. Model Dev.*, 13, 2379–2392, <https://doi.org/10.5194/gmd-13-2379-2020>, 2020.
- Kaminski, T., Rayner, P. J., Heimann, M., and Enting, I. G.: On aggregation errors in atmospheric transport inversions, *J. Geophys. Res. Atmos.*, 106, 4703–4715, <https://doi.org/10.1029/2000JD900581>, 2001.
- Kountouris, P., Gerbig, C., Rödenbeck, C., Karstens, U., Koch, T. F., and Heimann, M.: Technical Note: Atmospheric CO₂ inversions on the mesoscale using data-driven prior uncertainties: methodology and system evaluation, *Atmos. Chem. Phys.*, 18, 3027–3045, <https://doi.org/10.5194/acp-18-3027-2018>, 2018.
- Kuenen, J., Dellaert, S., Visschedijk, A., Jalkanen, J.-P., Super, I., and Denier van der Gon, H.: Copernicus Atmosphere Monitoring Service regional emissions version 4.2 (CAMS-REG-v4.2), Copernicus Atmosphere Monitoring Service (CAMS) [publisher], ECCAD [distributor], <https://doi.org/10.24380/0vzb-a387>, 2021.
- Kuenen, J., Dellaert, S., Visschedijk, A., Jalkanen, J.-P., Super, I., and Denier van der Gon, H.: CAMS-REG-v4: a state-of-the-art high-resolution European emission inventory for air quality modelling, *Earth Syst. Sci. Data*, 14, 491–515, <https://doi.org/10.5194/essd-14-491-2022>, 2022.
- Lauvaux, T., Miles, N. L., Richardson, S. J., Deng, A., Stauffer, D. R., Davis, K. J., Jacobson, G., Rella, C., Calonder, G.-P., and DeCola, P. L.: Urban Emissions of CO₂ from Davos, Switzerland: The First Real-Time Monitoring System Using an Atmospheric Inversion Technique, *Journal of Applied Meteorology and Climatology*, 52, 2654–2668, <https://doi.org/10.1175/JAMC-D-13-038.1>, 2013.
- Li, X. S. and Shao, M.: A Supernodal Approach to Incomplete LU Factorization with Partial Pivoting, *ACM Trans. Math. Softw.*, 37, <https://doi.org/10.1145/1916461.1916467>, 2011.
- Li, X. S., Demmel, J. W., Gilbert, J. R., Grigori, L., Shao, M., and Yamazaki, I.: SuperLU Users' Guide, june 2018 edn., 1999.
- Manning, A. J., Ryall, D. B., Derwent, R. G., Simmonds, P. G., and O'Doherty, S.: Estimating European emissions of ozone-depleting and greenhouse gases using observations and a modeling back-attribution technique, *J. Geophys. Res. Atmos.*, 108, <https://doi.org/10.1029/2002JD002312>, 2003.
- Manning, A. J., O'Doherty, S., Jones, A. R., Simmonds, P. G., and Derwent, R. G.: Estimating UK methane and nitrous oxide emissions from 1990 to 2007 using an inversion modeling approach, *J. Geophys. Res. Atmos.*, 116, <https://doi.org/10.1029/2010JD014763>, 2011.
- Meirink, J. F., Bergamaschi, P., and Krol, M. C.: Four-dimensional variational data assimilation for inverse modelling of atmospheric methane emissions: method and comparison with synthesis inversion, *Atmos. Chem. Phys.*, 8, 6341–6353, <https://doi.org/10.5194/acp-8-6341-2008>, 2008.
- Moré, J. J. and Sorensen, D. C.: Computing a Trust Region Step, *SIAM Journal on Scientific and Statistical Computing*, 4, 553–572, <https://doi.org/10.1137/0904038>, 1983.
- Pearson, K.: X. On the criterion that a given system of deviations from the probable in the case of a correlated system of variables is such that it can be reasonably supposed to have arisen from random sampling, *Philosophical Magazine Series 5*, 50, 157–175, <https://doi.org/10.1080/14786440009463897>, 1900.
- Petrescu, A. M. R., Qiu, C., McGrath, M. J., Peylin, P., Peters, G. P., Ciais, P., Thompson, R. L., Tsuruta, A., Brunner, D., Kuhnert, M., Matthews, B., Palmer, P. I., Tarasova, O., Regnier, P., Lauerwald, R., Bastviken, D., Höglund-Isaksson, L., Winiwarter, W., Etiope, G.,

- Aalto, T., Balsamo, G., Bastrikov, V., Berchet, A., Brockmann, P., Ciotoli, G., Conchedda, G., Crippa, M., Dentener, F., Groot Zwaaftink, C. D., Guizzardi, D., Günther, D., Haussaire, J.-M., Houweling, S., Janssens-Maenhout, G., Kouyate, M., Leip, A., Leppänen, A., Lugato, E., Maisonnier, M., Manning, A. J., Markkanen, T., McNorton, J., Muntean, M., Oreggioni, G. D., Patra, P. K., Perugini, L., Pison, I., Raivonen, M. T., Saunio, M., Segers, A. J., Smith, P., Solazzo, E., Tian, H., Tubiello, F. N., Vesala, T., van der Werf, G. R., Wilson, C., and Zaehle, S.: The consolidated European synthesis of CH₄ and N₂O emissions for the European Union and United Kingdom: 1990–2019, *Earth Syst. Sci. Data*, 15, 1197–1268, <https://doi.org/10.5194/essd-15-1197-2023>, 2023.
- Reinert, D., Prill, F., Frank, H., Denhard, M., Baldauf, M., Schraff, C., Gebhardt, C., Marsigli, C., Förstner, J., Zängl, G., Schlemmer, L., Blahak, U., and Welzbacher, C.: DWD Database Reference for the Global and Regional ICON and ICON-EPS Forecasting System, https://www.dwd.de/DWD/forschung/nwv/fepub/icon_database_main.pdf, 2025.
- Rieger, D., Bangert, M., Bischoff-Gauss, I., Förstner, J., Lundgren, K., Reinert, D., Schröter, J., Vogel, H., Zängl, G., Ruhnke, R., and Vogel, B.: ICON–ART 1.0 – a new online-coupled model system from the global to regional scale, *Geosci. Model Dev.*, 8, 1659–1676, <https://doi.org/10.5194/gmd-8-1659-2015>, 2015.
- Rigby, M., Manning, A. J., and Prinn, R. G.: Inversion of long-lived trace gas emissions using combined Eulerian and Lagrangian chemical transport models, *Atmos. Chem. Phys.*, 11, 9887–9898, <https://doi.org/10.5194/acp-11-9887-2011>, 2011.
- Rocher-Ros, G., Stanley, E. H., Loken, L. C., Casson, N. J., Raymond, P. A., Liu, S., Amatulli, G., and Sponseller, R. A.: Global methane emissions from rivers and streams, *Nature*, 621, 530–535, <https://doi.org/10.1038/s41586-023-06344-6>, 2023.
- Schraff, C., Reich, H., Rhodin, A., Schomburg, A., Stephan, K., Perriñez, A., and Potthast, R.: Kilometre-scale ensemble data assimilation for the COSMO model (KENDA), *Q. J. R. Meteorolog. Soc.*, 142, 1453–1472, <https://doi.org/10.1002/qj.2748>, 2016.
- Schröter, J., Rieger, D., Stassen, C., Vogel, H., Weimer, M., Werchner, S., Förstner, J., Prill, F., Reinert, D., Zängl, G., Giorgetta, M., Ruhnke, R., Vogel, B., and Braesicke, P.: ICON-ART 2.1: a flexible tracer framework and its application for composition studies in numerical weather forecasting and climate simulations, *Geosci. Model Dev.*, 11, 4043–4068, <https://doi.org/10.5194/gmd-11-4043-2018>, 2018.
- Segers, A. and Houweling, S.: CAMS global inversion-optimised greenhouse gas fluxes and concentrations, v22r2, Copernicus Atmosphere Monitoring Service [data set], <https://ads.atmosphere.copernicus.eu/datasets/cams-global-greenhouse-gas-inversion>, (last accessed: 18 April 2024), 2020.
- Segers, A., Nanni, R., and Houweling, S.: Evaluation and Quality Control document for observation-based CH₄ flux estimates for the period 1979–2022, ECMWF Copernicus Report, <https://atmosphere.copernicus.eu/supplementary-services#ptab-3-1-content>, 2023.
- Steiner, M., Cantarello, L., Henne, S., and Brunner, D.: Flow-dependent observation errors for greenhouse gas inversions in an ensemble Kalman smoother, *Atmos. Chem. Phys.*, 24, 12 447–12 463, <https://doi.org/10.5194/acp-24-12447-2024>, 2024a.
- Steiner, M., Peters, W., Lujikx, I., Henne, S., Chen, H., Hammer, S., and Brunner, D.: European CH₄ inversions with ICON-ART coupled to the CarbonTracker Data Assimilation Shell, *Atmos. Chem. Phys.*, 24, 2759–2782, <https://doi.org/10.5194/acp-24-2759-2024>, 2024b.
- Stohl, A., Seibert, P., Arduini, J., Eckhardt, S., Fraser, P., Grealley, B. R., Lunder, C., Maione, M., Mühle, J., O’Doherty, S., Prinn, R. G., Reimann, S., Saito, T., Schmidbauer, N., Simmonds, P. G., Vollmer, M. K., Weiss, R. F., and Yokouchi, Y.: An analytical inversion method for determining regional and global emissions of greenhouse gases: Sensitivity studies and application to halocarbons, *Atmos. Chem. Phys.*, 9, 1597–1620, <https://doi.org/10.5194/acp-9-1597-2009>, 2009.
- Storm, I., Karstens, U., D’Onofrio, C., Vermeulen, A., and Peters, W.: A view of the European carbon flux landscape through the lens of the ICOS atmospheric observation network, *Atmos. Chem. Phys.*, 23, 4993–5008, <https://doi.org/10.5194/acp-23-4993-2023>, 2023.
- UBA: Submission under the United Nations Framework Convention on Climate Change 2023, German Environment Agency, <https://doi.org/10.60810/openumwelt-2570>, 2023.

- UBA: Submission under the United Nations Framework Convention on Climate Change 2024, German Environment Agency,
 980 <https://doi.org/10.60810/openumwelt-7441>, 2024.
- Vanderbecken, P. J., Dumont Le Brazidec, J., Farchi, A., Bocquet, M., Roustan, Y., Potier, E., and Broquet, G.: Accounting for meteorological biases in simulated plumes using smarter metrics, *Atmos. Meas. Tech.*, 16, 1745–1766, <https://doi.org/10.5194/amt-16-1745-2023>, 2023.
- Veldeman, N., van der Maas, W., van Aardenne, J., Goodwin, J., Mareckova, K., Adams, M., Ruysenaars, P., Wankmüller, R., and Pye, S.: 7. Spatial mapping of emissions, in: EMEP/EEA air pollutant emission inventory guidebook 2013,
 985 European Environment Agency, <https://www.eea.europa.eu/publications/emep-eea-guidebook-2013/part-a-general-guidance-chapters/7-spatial-mapping-of-emissions/view>, 2013.
- Virtanen, P., Gommers, R., Oliphant, T. E., Haberland, M., Reddy, T., Cournapeau, D., Burovski, E., Peterson, P., Weckesser, W., Bright, J., van der Walt, S. J., Brett, M., Wilson, J., Millman, K. J., Mayorov, N., Nelson, A. R. J., Jones, E., Kern, R., Larson, E., Carey, C. J., Polat, I., Feng, Y., Moore, E. W., VanderPlas, J., Laxalde, D., Perktold, J., Cimrman, R., Henriksen, I., Quintero, E. A., Harris, C. R., Archibald,
 990 A. M., Ribeiro, A. H., Pedregosa, F., and van Mulbregt, P.: SciPy 1.0: fundamental algorithms for scientific computing in Python, *Nat. Methods*, 17, 261–272, <https://doi.org/10.1038/s41592-019-0686-2>, 2020.
- Weber, T., Wiseman, N. A., and Kock, A.: Global ocean methane emissions dominated by shallow coastal waters, *Nat. Commun.*, 10, 1–10, <https://doi.org/10.1038/s41467-019-12541-7>, 2019.
- Zängl, G., Reinert, D., Rípodas, P., and Baldauf, M.: The ICON (ICOsahedral Non-hydrostatic) modelling framework of DWD and MPI-M:
 995 Description of the non-hydrostatic dynamical core, *Quart. J. Roy. Meteorol. Soc.*, 141, 563–579, <https://doi.org/10.1002/qj.2378>, 2015.

German methane fluxes ~~in 2021~~ estimated ~~with an ensemble-enhanced scaling inversion based on the top-down using ICON-ART model~~ Part 2: Inversion results for 2021

Valentin Bruch¹, Thomas Rösch¹, Diego Jiménez de la Cuesta Otero¹, Beatrice Ellerhoff¹,
Buhalgem Mamtimin¹, Niklas Becker¹, Anne-Marlene Blechschmidt¹, Jochen Förstner¹, and Andrea
K. Kaiser-Weiss¹

¹Deutscher Wetterdienst, Frankfurter Str. 135, 63067 Offenbach

Correspondence: Valentin Bruch (valentin.bruch@dwd.de) and Andrea K. Kaiser-Weiss (andrea.kaiser-weiss@dwd.de)

Abstract. A reliable quantification of greenhouse gas emissions is important for climate change mitigation strategies. Inverse methods based on observations and atmospheric transport simulations can support emission quantification ~~down-to-at~~ the national scale, yet, they are often limited by the observing systems, transport model uncertainties, and inversion methodologies. ~~Here, we present~~ This two-part study introduces a system for observation-based, regional methane flux estimation, ~~which has the potential for long-term operational support of national emission reporting. We apply this to Central Europe in 2021 with focus on Germany, where we distinguish emissions from different anthropogenic sectors.~~ In the present Part 2, we apply this system to estimate German methane emissions in 2021. The ~~atmospheric transport is calculated with the~~ numerical weather prediction model ~~ICON-ART at 6.5 km resolution, sampling the meteorological uncertainty with a 12-member~~ ICON with its ART module for trace gases is used to simulate the atmospheric transport while estimating uncertainties using a transport ensemble. We use a priori fluxes from national reporting to facilitate the validation of reported fluxes. Posterior fluxes are estimated with a modified synthesis inversion method introduced in Part 1, relying on ~~observations from the Integrated Carbon Observation System (ICOS).~~ in-situ observations. Compared to the a priori, we find a significant increase in methane emissions in Germany and in the Benelux. We estimate German methane emissions (32 ± 19) % higher than the anthropogenic emissions in the national inventory, and attribute our inversion method attributes this difference mainly to the agricultural sector, although separation from Land Use, Land Use Change and Forestry (LULUCF) as well as natural fluxes requires further research. The combination of an ensemble-enhanced numerical weather prediction model for atmospheric transport and good observation coverage paves the way to sector-specific, observation-based national emission estimates.

1 Introduction

Reducing greenhouse gas (GHG) emissions is crucial for mitigating current anthropogenic global warming. UNFCCC (United Nations Framework Convention on Climate Change) compliant national inventories and/or process models quantify anthropogenic GHG emissions for the purpose of monitoring the effectiveness of mitigation as planned, e.g., in the Paris Agreement. In addition to so-called “bottom-up” methods, atmospheric GHG concentration observations are used in “top-down” flux es-

timations. The latter are complementary, as they are sensitive to the total fluxes (i.e., anthropogenic and natural) and provide options for independent validation of a priori fluxes provided by inventories (IPCC et al., 2019). The usefulness of top-down estimates has been demonstrated, e.g., for the United Kingdom (Manning et al., 2011), Switzerland (Henne et al., 2016), Europe (Petrescu et al., 2023) and globally (Deng et al., 2022; Petrescu et al., 2024).

Although research foundations for top-down methods have been developed in recent decades (see Janssens-Maenhout et al. (2020) and references therein), applications remain limited due to sparse observation coverage and representativeness, and most critically, due to transport model uncertainties (Engelen et al., 2002; Gerbig et al., 2008). The latter is a well-known issue not solved yet (Munassar et al., 2023). Inversions using satellite observations (e.g. Estrada et al., 2024) benefit from larger spatial observation coverage, but the uncertainties of the observations are larger compared to in situ data and the influence on the inversion results was found smaller where in situ coverage is good (Thompson et al., 2025). The benefits of increased model resolution (Agustí-Panareda et al., 2019; Bergamaschi et al., 2022) can be reaped with regional high resolution modeling and ensembles can cover parts of the meteorological uncertainty (Steiner et al., 2024a). At short time scales, the regional model uncertainties will constitute the main uncertainty, while at longer time scales, the boundary conditions become critical for tracer transport (Chen et al., 2019).

~~Regional top-down estimates of long-lived GHG can be based on different types of transport models. Lagrangian models calculate trajectories from selected locations by moving with air parcels transported by the wind. They have been widely used for inversions of trace gases like halocarbons, nitrous oxide and methane (CH₄) in European regions, see e.g., Stohl et al. (2009); Ganesan et al. (2015). In contrast, Eulerian models—such as ICON-ART—continuously transport trace gas concentrations through three-dimensional grid-boxes. Although they are computationally more expensive for cases where a relatively small number of trajectories would suffice, they become superior when the amount of data grows and, as Engelen et al. (2002) pointed out, open the road for data assimilation methods as used in numerical weather prediction. Regardless whether Lagrangian or Eulerian or even combined approaches (Rigby et al., 2011) are applied, the top-down estimation requires solving an inverse problem (Enting, 2002). Eulerian transport model-based inversions may employ emission ensembles, as in Steiner et al. (2024b) with a localized Kalman filter, and other data assimilation methods. Alternatively, the method of synthesis inversion scales a set of a priori emission categories (Kaminski et al., 2001). Note that Meirink et al. (2008b) compare 4D-Var and synthesis inversion methods.~~

In this work, ~~we present~~ we present first results of a modular system for regional top-down estimates of CH₄ fluxes designed to validate national inventories, including the discrimination of economic sectors such as agriculture and industry. We apply this method focusing on German inventories (provided by Umweltbundesamt and Thünen Institute) for the year 2021 using in situ observations collected by ICOS (ICOS RI, 2024). Atmospheric transport is simulated using the numerical weather prediction model ICON (Zängl et al., 2015) extended with the module for Aerosol and Reactive Trace gases (ART) (Rieger et al., 2015; Schröter et al., 2018) with a spatial resolution of 6.5 km. The model is combined with a synthesis inversion approach (Kaminski et al., 2001) which is developed further to make use of the ensemble-estimated transport uncertainty. For minimizing transport errors, we rely on the operational numerical weather prediction at Germany's Meteorological Service (DWD) for meteorological initial conditions, lateral boundaries and transport ensemble calculations. Further, we use the Copernicus

~~Atmospheric~~ Atmosphere Monitoring Service (CAMS) for boundary conditions of methane, and compensate possible biases on the boundaries by deriving a correction field. Benefiting from the numerical weather prediction model and spatially highly resolved a priori fluxes from the inventory agencies, we explore the basis for future operational top-down validation of national emission reporting, with special emphasis on further use in Germany.

In Sect. 2, we ~~explain the transport model and the inversion methodology, while leaving some technical details for the appendices. Section 3 contains an overview of the utilized initial and boundary data as well as the a priori fluxes. Further, we describe the pre-processing of the observations used. Section 4 is dedicated to the details of the method concerning its application for Germany. Section 5 summarize the methodology which is introduced in detail in Part 1 of this work (Bruch et al., 2025a). Section 3 contains the results for our example year 2021, together with the results of the validation and sensitivity tests. The potential of the method is demonstrated by model performance tests using pseudo-observations with known true emissions. In Sect. 6 validation tests and an analysis of the ability to distinguish emission sectors. In Sect. 4 we discuss limitations and capabilities of the method and compare to other studies, followed by a conclusion in Sect. 7. Sect. 5.~~

2 Method

This section is a non-technical summary of the detailed method description in Part 1 (Bruch et al., 2025a).

[...]

2.1 ~~Transport model~~ Parametrization of fluxes

~~The atmospheric transport is simulated using the numerical weather prediction model ICON (Zängl et al., 2015) with the ART module (Rieger et al., 2015; Schröter et al., 2018). The model is run in limited area mode for a domain covering large parts of~~ We aim to validate the national reporting of German CH₄ emissions to the UNFCCC. A simple way to address this validation problem is the ~~European continent (latitudes 34° N to 70° N, longitudes 21° W to 59° E, see Fig. 1) with a horizontal resolution of 6.5 km (ICON grid R3B8) and 74 vertical levels up to a maximal height of 22.77 km. following question: By which single number should we multiply all reported German CH₄ emissions based on the information from observed CH₄ concentrations? The surface CH₄ fluxes are provided to the transport model using the online emission module (Jähn et al., 2020; Steiner et al., 2024b)~~ We can extend this question and estimate different scaling factors for different regions and different emission sectors. In this work, we estimate scaling factors for 46 categories of CH₄ fluxes for each month in 2021. The spatial definition of these flux categories is shown in Fig. 1. In Germany, we distinguish 11 flux categories, consisting of six regions for the agriculture sector, one flux category for land use, land use change and forestry (LULUCF) plus natural fluxes, and four regions for the sum of all remaining emissions. In summary, the state space of our inversion is defined by the flux categories and consists of only 46 numbers.

~~For long living tracers like methane, the correct treatment of the lateral boundary concentrations is of importance. Therefore, we extended the model by implementing lateral boundary nudging for ART tracers in order to obtain smooth fields and avoid~~




Figure 1. Overview of the model domain indicating flux categories (colored areas) and observation sites (white dots), modified from Part I (Bruch et al., 2025a). Each connected area of equal color defines one flux category for anthropogenic emissions, except in Germany and the Netherlands, where the categories are split up further to distinguish agriculture emissions from other sectors. In white hatched regions, natural fluxes form additional flux categories because large natural fluxes are expected. Close to the eastern and western domain boundary (dark blue), emissions are not adjusted by the inversion. Fugitive emissions from the Upper Silesian Coal Basin (white ellipse) define their own flux category.

~~strong gradients. Further, so-called meteogram output has been implemented for ART tracers, providing model output in the vicinity of observation locations with high temporal resolution.~~

~~For improved uncertainty estimates, we run a meteorological~~

2.2 A priori fluxes

For the a priori fluxes outside Germany, we combine CAMS-REG (Kuenen et al., 2021, 2022) for anthropogenic emissions with wetland emissions from the CAMS global inversion-optimized dataset (Segers and Houweling, 2020), version v22r2. For Germany, we use emissions obtained from the inventory agencies, that is, the Umweltbundesamt (German Environmental Agency, Feigensperger et al., 2018), the Thünen Institute (Fuß and Akubia, 2024). Moreover, we consider emissions from rivers and streams (Rocher-Ros et al., 2023), as well as oceans (Weber et al., 2019).

2.3 Transport simulation

To connect surface fluxes and observations, we need to simulate atmospheric transport. This simulation is done using the numerical weather prediction model ICON (Zängl et al., 2015) with the module for Aerosol and Reactive Trace gases (ART) (Rieger et al., 2015; Schröter et al., 2018) at a horizontal resolution of 6.5 km. Initial and lateral boundary conditions for the CH_4 concentrations are taken from the CAMS global inversion-optimized dataset (Segers and Houweling, 2020), version v22r2. To mitigate a possible bias in the lateral boundary conditions, we construct a smooth correction field that is added

to all model predictions of the boundary contributions. This far-field correction is constructed based on observations for which the model predicts clean air with small influence of emissions from within our domain. We estimate transport uncertainties and their correlations using an ensemble of 12 members. Each ensemble member uses slightly different but equally likely parametrizations and meteorological initial and boundary conditions. The construction of this ensemble follows the operational with slightly different meteorology, derived from the operational numerical weather prediction at DWD (Schraff et al., 2016). In the following, we therefore distinguish a so-called deterministic model run providing the best estimate of the modeled CH_4 concentration, and the ensemble runs providing 12 different CH_4 concentrations to estimate the uncertainty.

[...]

2.4 Observations

We use CH_4 concentration observations from the European Obspack (ICOS RI et al., 2024) as provided on the Integrated Carbon Observation System (ICOS) carbon portal. The hourly observations are filtered by time of day and wind speed to use only observations that can be predicted well by the transport model. We use night time observations (23 h to 5 h local mean time) for high mountain stations and afternoon hours (11 h to 17 h local mean time) for all other sites, discarding observations at wind speeds below 2 m s^{-1} .

2.5 Inversion (Sect. 2.3)

We use a Bayesian inversion to optimize the agreement of model and observations by scaling

2.5 Bayesian Inversion

To estimate the scaling factors of the flux categories, this is formulated in, we use a Bayesian inversion. Denoting the scaling factors as a vector $s \in \mathbb{R}^{46}$, the inversion is formulated as the optimization problem

$$s^{\text{post}} = \arg \min_s \left\{ \frac{1}{2} ([y - Hs - x^{\text{ff}} H'(s)]^\top R^{-1} [y - Hs - x^{\text{ff}} H'(s)] + \frac{1}{2} (s - s^{\text{prior}})^\top B^{-1} (s - s^{\text{prior}}) \right\}. \quad (1)$$

for the posterior scaling factors s^{post} . Here, the first term penalized the deviation from the concentration observations, and the second term penalizes the deviation from the prior fluxes. In the first term, the vector y of observed concentrations is compared to denotes a vector of all observations and $H'(s)$ is the model prediction, which consists of the transported fluxes Hs and the modeled far field x^{ff} . The transported fluxes Hs depend linearly on the vector s of scaling factors for the flux categories, which is optimized. The difference between modeled and observed values is weighted by for these observations, which includes the previously mentioned far-field correction. R is the error covariance matrix R describing the combined uncertainty of the transport model and of the observations. With the second term we constrain s by defining a priori scaling factors s^{prior} ($s_i^{\text{prior}} = 1$ for all i) with an model-observation mismatch and B is the error covariance matrix B characterizing of the a priori uncertainty.

In Eq. (1), the fluxes are parametrized by the vector s of scaling factors. The flux categories thereby define the low-dimensional space in which the inversion can optimize the fluxes. The model-observation operator H connects the space of scaling factors

135 (vectors scaling factors s^{prior} , s^{post}) to the observation space (vectors y , x^{ff}). Computing H requires the transport model which distinguishes the flux categories. The setup is designed for optimizing a low-dimensional vector s^{post} of scaling factors ($\sim 10^2$ degrees of freedom) using a large number of observations ($\sim 10^4$), but an extension to more degrees of freedom and/or more observations is possible.

[...]

140 3 Input data and processing (Sect. 3)

[...]

3 Application to Germany and neighboring areas for the year 2021 (Sect. 4)

[...]

2.1 Prior uncertainties (Sect. 4.4)

145 In each inversion time window, we consider uncorrelated a priori scaling factors with a two standard deviation (2σ) uncertainty of 80% for most flux categories, corresponding to a 95% confidence interval of $\pm 80\%$. Throughout this paper, uncertainties will denote. Since s describes prefactors to the a priori emissions, we initially set $s_k^{\text{prior}} = 1$ for all k . In B we assume an a priori uncertainty of $2\sigma = 0.8$ (two standard deviations) for the scaling factors of most regions. This gives the inversion enough freedom to adjust the scaling factors. In large distance from Germany, the a priori uncertainty is reduced to $2\sigma = 0.5$ (see Fig. 2 b), and for emission sectors in Germany and the Netherlands we use $2\sigma = 1.0$.

150 The construction of R based on the transport ensemble is discussed in detail in Part 1 (Bruch et al., 2025a). In Eq. (1), R can be estimated using either a priori or 95% confidence intervals. Categories resolving emission sectors have a higher prior 2σ uncertainty of $\pm 100\%$, and within Germany categories describing the same sector have an a priori uncertainty correlation of 50% (e. g., uncertainties of agriculture emissions in the German states of Bavaria and Baden-Wuerttemberg are assumed to be correlated). Importantly, the previously defined sectors waste, public power and “other” are combined and denoted “non-agr.” in the following. The $\pm 100\%$ uncertainty applies to the combination. For the Upper Silesian Coal Basin as well as regions outside of our primary focus in Central Europe and with low observation density, the 2σ uncertainty is set to $\pm 50\%$. Figure 2(b) shows these a priori uncertainties on a map. The sensitivity of our results to these choices and many more tuning parameters is tested in Appendix E.

160 2.1 Posterior uncertainty estimates (Sect. 4.5)

Our inversion setup necessarily makes idealized assumptions on uncertainties of a priori and a posteriori fluxes. Based on these assumptions, the inversion yields posterior statistical uncertainties as part of the posterior error covariance matrix. However, these uncertainties neglect possible unknown biases and other systematic errors which we need to anticipate when working

with real data. We therefore use an enhanced notion of posterior uncertainty that combines statistical and methodological uncertainties. The main contribution to the posterior uncertainties is the statistical uncertainty, which we provide as two standard deviations (95 % confidence interval). Additionally, we combine the two variants of inversion (prior- R and posterior- R ; see Sect. 2.4.2) by taking the arithmetic mean of the two separate inversion results, arriving at the combined scaling factors. Assuming that the difference This defines two slightly different methods that are introduced in Sect. 2.5 of Part 1 as “prior R ” and “posterior R ” inversion. Here, we only consider the average of the two results and the union of the two variants indicates an additional, methodological uncertainty, our combined uncertainty range includes the uncertainty ranges of both individual results posterior uncertainty ranges.

Furthermore, each observation site may have a bias due to very local pollution or topography that is not modeled properly. We therefore only consider an inversion result reliable if it is robust with respect to the choice of used observations. To represent this in the uncertainty estimate

175 2.6 Posterior uncertainties

To estimate the uncertainties of posterior fluxes conservatively, we repeat the inversions ~~50 times for both prior- R and posterior- R , excluding each station once~~. The final posterior uncertainty range is spanned by the inversion 50×2 times with each of the 50 observation sites excluded once for each of the two approximations for R . The lower and upper bounds of the 100 uncertainty ranges, each describing a resulting hundred 2σ uncertainty ranges form our posterior 95 % confidence interval. This method of compensating for possible methodological weaknesses in a cautionary uncertainty estimate implicates that the posterior uncertainty may not necessarily be smaller than the prior uncertainty. Such cases we interpret as no information gain from the observation-based inversion with respect to the prior. This ensures that a result that is only based on a single observation site will not be considered significant.

2.7 Inversion time window ~~and temporal aggregation (Sect. 4.3)~~

185 ~~We apply the inversion~~ The scaling factors are estimated separately for each month in 2021. ~~In each monthly time window, we start from fixed a priori scaling factors $s_i^{\text{prior}} = 1$ and use observations within the time window to compute a posteriori scaling factors as explained in Sect. 2.3. However, when aggregating results for the whole year, we treat the uncertainties of the prior or posterior fluxes of different months as correlated because these likely~~ 2021 by using only observations from the selected month. The results for different months are thus independent. But since the posterior uncertainty estimates include systematic 190 ~~uncertainties and biases which we cannot fully separate from the statistical uncertainty. We therefore aggregate by adding up absolute emissions and their uncertainties linearly.~~ we assume that uncertainties from different months are correlated.

3 Results ~~(Sect. 5)~~




Figure 2. Full-year averages of (a) a priori fluxes, (b) a priori uncertainty on scaling factors, (c) a posteriori scaling factors, and (d) a posteriori uncertainty on scaling factors. Multiplying the a priori emissions (a) with the scaling factors (c) yields the a posteriori emissions. (b) and (d) show half of the 95 % confidence interval of the fluxes relative to the a priori fluxes, i.e., a 2σ uncertainty of $\pm 50\%$ on the a priori appears as 0.5 (i.e., 50%) 0.5 on the color scale. The direct comparison indicates the uncertainty reduction. The smooth boundaries between two regions with separate scaling factors appear as darker lines because these scaling factors are assumed to be initially uncorrelated.

~~This section presents the inversion results for Germany and the considered European regions, along with examinations of the seasonal cycle, validation of the results, and sensitivity tests. All uncertainty estimates are presented as 95 % confidence intervals as detailed above.~~

3.1 Resulting scaling factors

Figure 2 presents an overview of (a) the a priori CH_4 fluxes accumulated over the year 2021, (c) the resulting scaling factors averaged over 2021, and the respective uncertainties (b, d), ~~all accumulated over the year 2021.~~ The a posteriori scaling factors (Fig. 2 c) show the correction to the a priori emissions obtained in the inversion. A considerable increase in emissions is found for Germany and the Benelux. Lower emissions compared to the a priori are predicted for Scandinavia (see discussion in Sect. 4.3). The scaling factors should be considered jointly with their uncertainties. The comparison of Fig. 2 (b) and (d)




Figure 3. National CH₄ emission estimates comparing reported (NIR), prior, and posterior fluxes for 2021 with horizontal lines indicating 95% confidence intervals. Countries are grouped by the expected robustness of their inversion results. Some neighboring countries are combined to obtain more accurate results. For Germany, the inversions can resolve the agricultural sector, though the separation against natural and LULUCF fluxes is difficult. All other anthropogenic sectors are combined in the category “~~other excl. LULUCF~~” “~~other excl. LULUCF~~”. The inclusion of two inversion methods (~~prior-R~~ “prior R” and ~~posterior-R~~ “posterior R”, markers) ~~yields~~ provides an estimate of the methodological uncertainty. Accumulated ~~fluxes-emissions~~ from national inventory reports (NIR) to the UNFCCC submitted 2024 (including LULUCF emissions) are shown for reference (light blue bars, UNFCCC, 2024). For France (Citepa, 2024) and the United Kingdom (Department for Energy Security and Net Zero, 2024), the light blue bars show emission data from the respective inventory agencies excluding overseas territories and crown dependencies. Posterior uncertainties that are asymmetric with respect to flux estimates such as in Switzerland indicate the strong influence of a single observation site.

shows a substantial uncertainty reduction for Germany and most of the surrounding countries, for which we chose a high a priori uncertainty.

For a more detailed comparison of a priori and a posteriori emissions and uncertainties, we consider selected national emission estimates in Fig. 3.

Reliable inversion results are expected for countries or regions with sufficient observation coverage, strong emission signals, representation in the respective flux categories, and only moderate issues due to complex topography. These criteria are met for Germany, the Netherlands and the United Kingdom plus Ireland as grouped in Fig. 3. For Germany (first ~~line-entry~~ in Fig. 3),

the total posterior CH₄ emissions (red bar) are $(32 \pm 19)\%$ higher than the anthropogenic emissions ~~(including LULUCF)~~
210 including LULUCF reported to the UNFCCC in 2024 (light blue bar). The direct comparison to the reporting neglects the
unreported natural fluxes, but for Germany these are expected to be small because all relevant soil emissions are included in
the LULUCF sector. The inversion significantly increases emission estimates from the agriculture sector while the combined
other sectors remain nearly unchanged. Note, however, that the uncertainty in the sector attribution is large (horizontal lines,
see further discussion in Sections 3.4.2 and 4.3).

215 For the Netherlands, we also find significantly higher emissions than in the inventory. Compared to Germany, the attribution
to sectors has an even larger uncertainty, associated with fewer observations that could distinguish the sectors. Nevertheless,
the total emissions from the Netherlands are comparably well constrained by the observations. For the United Kingdom and
Ireland – which we combine to obtain more accurate results – the inversion yields a strong uncertainty reduction while hardly
changing the total emissions, indicating a good agreement of observations and national inventory.

220 In most countries, the observations do not cover the whole country, or the inversion results rely on few observations. In
Fig. 3 (gray-shaded part) we provide emission estimates also for countries or regions affected by this issue, though these have
a large posterior uncertainty. Another issue arises from the definition of the flux categories, which do not necessarily follow
country borders (see Fig. 1). In France, Belgium, and Switzerland, the inversion scales flux categories ~~overlapping that overlap~~
multiple countries¹. This implies that national emission estimates derived for these countries have an additional uncertainty and
225 artificial correlations with neighboring countries. However, this is of no concern for our application for Germany. The national
emission estimates are computed from the gridded posterior fluxes and precisely follow the country borders as shown in Fig. 2.
The scaling factors and uncertainties of all flux categories are listed in Fig. A1 for completeness.

3.2 Seasonal cycle

Although the national emission estimates are given for the full year, a closer examination of the seasonal cycle ~~yields~~ provides
230 additional insights. ~~The posterior fluxes are computed independently for each month and region.~~ Figure 4 shows the monthly
emission rates for the countries considered in Fig. 3. While the seasonal cycle is strikingly different depending on the region,
we find some recurrent features. For Germany, Poland, the Netherlands, and Austria plus Czechia (panel (a) in Fig. 4), the
posterior emission rates have their minimum in May. A local minimum between April and June is also found for northern
France and Belgium plus Luxembourg, see panel (b). In most countries, this minimum is followed by a local maximum in July
235 or August, which is most prominent in the Netherlands and Austria plus Czechia (panel (a)).

The differences between the regions become larger in autumn and winter. In September, posterior emission rates reach their
maximum in Germany and Italy, and their minimum in (northern) France. France and Belgium plus Luxembourg have their
highest emission rates in winter, when Switzerland and Spain plus Portugal have their minimum. For some regions – most
notably Italy and the United Kingdom plus Ireland – no clear pattern is found in the seasonal cycle for 2021 (panel (c) in
240 Fig. 4).

¹Technically, the issue also affects Italy because Corsica is combined with parts of Italy in one flux category. But the a priori emissions from Corsica are
so low that the effect on the national emission estimate is negligible.




Figure 4. Monthly posterior emission rates for selected countries or regions. Colored areas show the posterior uncertainties, and dotted lines with small markers indicate prior emission rates. In the prior, only the natural and LULUCF fluxes are time-dependent. The panels show (a) countries with minimum in May, (b) countries with a maximum in winter, and (c) other countries and regions. For France and Germany, selected regions are shown additionally (white markers). “DE, northwest” includes Rhineland-Palatinate, Saarland, Hesse, North Rhine-Westphalia, Lower Saxony, Schleswig-Holstein, Bremen and Hamburg.

The seasonal cycle in the inversion results may be partially influenced by the observation coverage because many stations lack data covering the whole year. To avoid this effect, we repeated the inversion using only stations which provide data for at least 20 days of each month. The seasonal cycle in these results does not change significantly, see [Fig. A2: supplementary Fig. A2](#). We further note that there is a seasonal cycle in the observations (East et al., 2024), which is captured well by the far field in the model though (see Fig. A3). [This “far field” is defined as CH₄ transported into our domain from the lateral boundaries.](#) A possible bias in the lateral boundary conditions [determining the far-field could also could](#) influence the seasonal cycle in the estimated fluxes. Moreover, the different meteorology in summer and winter – especially influencing the planetary boundary layer and vertical mixing (Seidel et al., 2012) – can lead to a seasonal bias in our transport model (Bessagnet et al., 2016; Canepa and Buitjes, 2017). This highlights the need for careful interpretation of the seasonal cycle, as meteorological differences could introduce biases that mask true emission patterns. Another potential contribution to the seasonal cycle could arise from neglecting the OH sink of CH₄ in our limited domain [-\(Logan et al., 1981\).](#)

3.3 Validation

A straightforward validation of the inversion results is possible using independent validation stations. Having excluded each station once in separate inversion runs, we can use every station as an independent validation site in the respective inver-




Figure 5. Statistics of the relative (a) and absolute (b) improvement of the model–observation mismatch by the inversion at independent validation stations. Each station and month is considered separately in its own (prior- R and posterior- R) inversion, with the validation station excluded from the inversion to remain independent. The histograms show (a) $1 - r^{\text{post}}/r^{\text{prior}}$ and (b) $r^{\text{prior}} - r^{\text{post}}$ where r^{post} and r^{prior} refer to the RMSE of the model–observation comparison in the case of posterior scaling and prior scaling, respectively. Each time series contributing to the histogram is weighted by the number of its data points. We consider all data points within the daily time window without filtering for wind speed, extreme mismatch, uncertainty weighting or model–observation mismatch and without the far-field correction (i.e., after step 3 introduced in Table 2) Part 1 (Bruch et al., 2025a) to keep the comparison as close as possible to the original data. Positive values indicate an improvement in the model prediction due to the inversion.

255 sion run. Figure 5 shows histograms of the ~~RMSE statistics evaluating root mean square error (RMSE) statistics obtained from~~ the model–data mismatch before and after the inversion. The validation stations agree on average significantly better with observations when using a posteriori emissions compared to the a priori. A comparison of the same histograms for the different methods of estimating uncertainties introduced in Part 1 (Bruch et al., 2025a) shows no significant differences (see supplementary Fig. A4).

260 3.4 Sensitivity tests (Sect. 5.4) Potential for detecting emissions

[...]

3.5 ~~Potential for detecting emissions (Sect. 5.5)~~

In this section, we complement the uncertainty estimates of our inversion results by separate measures for the sensitivity of the posterior to true emissions. The potential for detecting emissions from different sources can be identified using the posterior error covariance matrix B_{post} . However, the real error reduction is also influenced by the far-field correction and the filtering of
265 observations, ~~which is as detailed in Part 1 (Bruch et al., 2025a).~~ These aspects are not fully captured in B_{post} . We therefore use experiments with a “synthetic”, i.e., ~~define defined~~ truth and pseudo-observations to test the full inversion system.

3.4.1 National emission estimates

We first aim to verify that the inversion yields meaningful posterior emission estimates and uncertainties given a perfect trans-
270 port model. To this end, we generate 100 random vectors of scaling factors following the probability distribution assumed in the a priori uncertainty. ~~These scaling factors define the~~ Each vector of scaling factors defines a synthetic truth, and the model




Figure 6. RMSE and mean uncertainty of CH₄ emission estimates in synthetic experiments for selected countries, regions, and [German emission](#) sectors. Each of the 100 synthetic experiments [generates-uses](#) random true emissions. The vertical axis shows the root mean square (RMS) deviation of the posterior from these true emissions, relative to the RMS deviation of the prior from the truth. Lower values indicate that the inversion improves the emission estimate. The horizontal axis shows the posterior uncertainty relative to the prior uncertainty. [Therefore, the bottom left indicates best performance.](#) The disk size indicates the [amount-magnitude](#) of the prior emissions.

prediction for the observations obtained using these scaling factors defines our pseudo-observations. We further add uncorrelated Gaussian noise of standard deviation 2ppb to these pseudo-observations. Since the pseudo-observations are inferred from the model data, there is no transport error in these synthetic experiments. [This construction of pseudo-observations clearly underestimates the true error in the model–observation comparison, but it allows us to test the interplay of far-field correction and inversion in a controlled setup. Synthetic experiments with a simulated transport uncertainty are discussed in Part 1 \(Bruch et al., 2025a\).](#)

The quality of the model prediction [for-this-synthetic-truth](#) is shown in Fig. 6 for selected countries and German sectors. By comparing to the synthetic truth, we find the prior and posterior error. Their ratio (vertical axis in Fig. 6) shows a significant improvement by the inversion for all considered regions and German sectors, with the exception of German natural and LULUCF fluxes. The uncertainty reduction of the inversion (horizontal axis) provides a realistic estimate of the real error reduction (vertical axis) for the case of high quality observations, ideal transport modeling [and-perfect-far-field, and perfect lateral boundary conditions.](#) In some cases (Netherlands, Switzerland, Belgium, and Luxembourg), the real error reduction is significantly better than the uncertainty reduction suggests. This is no surprise because in this synthetic setup the transport




Figure 7. Averaging kernel matrices of German sector emissions (a, c) and scaling factors (b, d). The kernel is estimated using either the posterior covariance matrix (a, b) or 100 synthetic experiments with random truth (c, d). The small matrices on the bottom indicate what we aim for (posterior equals truth). The value 0.96 in the first row (“total”), second column (“agriculture”) of panel (a) means that if in reality all German agriculture emissions were 1 kt higher than in our prior, then we would expect an increase in the posterior total German emissions by 0.96 kt. Similarly, the value 0.67 in the same cell of panel (b) means that increasing real agriculture emissions by 10% should increase our posterior total emissions by 6.7%. All matrices are averaged over the whole year. Red lines separate the individual sectors from their sum (“total”). By “non-agr.” we denote anthropogenic emissions excluding agriculture and LULUCF.

error as the main source of uncertainty is switched off. Overall, the synthetic experiments confirm the potential for a strong uncertainty reduction in Central Europe.

3.4.2 Distinguishing sectors in Germany

Within Germany, we distinguish agriculture from other emissions. The discrimination of emission sectors works in the same way as we distinguish emissions from different areas. Each sector has a specific spatial distribution of emissions, which we assume to be correct in the a priori. The predicted CH₄ concentration at the observation sites will therefore depend on how the individual sectors are scaled. In the inversion, the sector emissions are scaled to find optimal agreement of model prediction and observations.

The ability to distinguish sectors can be described by averaging kernel matrices which estimate the dependence of the posterior on the true emissions, $A_{ij}^{\text{emis}} = \partial e_i^{\text{post}} / \partial e_j^{\text{truth}}$ where e_i denotes emissions from sector i . Since the true emissions e^{truth} are generally unknown, the averaging kernels A_{ij}^{emis} can only be estimated. Figure 7 shows such estimates for A^{emis} (panels a, c) and the averaging kernel for scaling factors, $A_{ij}^{\text{scaling factors}} = \partial s_i^{\text{post}} / \partial s_j^{\text{truth}}$ (panels b, d). Assuming a perfect transport model

and perfect far field, the averaging kernel matrix can be estimated by $A^{\text{emis}} \approx I - B_{\text{post. emis}} B_{\text{prior emis}}^{-1}$ (Rodgers, 2000) using the prior and posterior covariance matrices of the emissions from the ~~prior- R~~ “prior R ” inversion (see Appendix B1). I denotes the identity matrix. Figure 7 (a) shows this averaging kernel estimate for German sector emissions, extended by a row and column
 300 for the total German emissions.

The first row of Fig. 7 (a) indicates that the total German posterior emissions follow changes in every sector with high accuracy (88 % to 96 %). The diagonal of Fig. 7 (a) signifies that changes in the agriculture will be detected very well and also the attribution to the sum of all other anthropogenic sectors excluding LULUCF (“non-agr.”) will be mostly correct. However, LULUCF plus natural fluxes will in large parts be falsely attributed to the agriculture (second row, last column). Note that
 305 ideally, the first row and the diagonal elements would be close to 100 % (color-coded in the small matrix bottom left). The averaging kernel $A^{\text{scaling factors}}$ in Fig. 7(b) shows that the influence of LULUCF and natural emissions on the posterior scaling factor for agriculture emissions remains low (second row, last column). But if all emissions are scaled by the same factor (first column), the changes will be mostly attributed to the agriculture sector (as explained in Appendix C). This effect is expected because the agriculture sector has the highest absolute a priori uncertainty, which makes changes in agriculture more likely than changes in any other sector. A formal derivation of this argument is presented in Appendix C.

The averaging kernel matrices in Fig. 7 (a) and (b) are estimated based on the ~~prior- R~~ “prior R ” inversion while neglecting the far-field correction. We complement these by a statistical estimate of the averaging kernels using 100 synthetic experiments with random truth (see Appendix B2), shown in Fig. 7 (c) and (d). Here, the far-field correction is applied as implemented in our processing chain. While these statistical estimates reproduce all qualitative features in the averaging kernels, the matrix
 315 entries estimated using synthetic experiments are generally lower. This is likely due to the far-field correction and indicates that deviations from the prior emissions may be underestimated by our inversion. Importantly, both presented strategies for estimating the averaging kernels assume a perfect transport model. The real sensitivity of the posterior to the true emissions is therefore expected to be lower.

3.5 Simulated transport error (Sect. 5.6)

320 [...]

4 Discussion (Sect. 6)

Our inversion system combines precise in situ observations, accurate a priori fluxes from national reporting, the ICON–ART transport model at 6.5 km resolution, and an ensemble-estimated transport uncertainty. We further rely on CAMS boundary conditions and high-resolution meteorological fields from operational numerical weather prediction. This yields in general a
 325 good agreement between the model prediction and filtered observations, allowing us robust emission estimates for ~~countries~~ with well-observed emissions countries, such as Germany. We compare top-down CH_4 emission estimates to the reported German inventory and its agriculture sector with enough accuracy to lay the technical foundations for a future long-term observation-based national inventory verification. This section discusses our main results (Sect. 4.1), including a comparison

with other studies (Sect. 4.2). We elaborate the limitations of our approach (Sect. 4.3) and its potential for the development of
330 observation-based national inventory verification to inform climate policy (Sect. 4.4).

4.1 Key findings

Firstly, we find that our top-down CH₄ emission estimates are significantly higher than reported for Germany. Secondly, we identify the agriculture sector and possibly LULUCF and natural fluxes as the likely main source of this discrepancy. Thirdly, we ~~estimated the transport uncertainty using a meteorological ensemble. The same ensemble allowed us to simulate the effect~~
335 ~~of a transport error on the inversion results, which we estimate to be 2% for the total German emissions.~~

~~Fourthly, our study points to the relevant tuning parameters. recall from Part 1 (Bruch et al., 2025a) that the transport error simulated in the meteorological ensemble leads to an uncertainty of 2% on the total German CH₄ emissions. The far-field correction significantly impacts the inversion results, but the specific choices made in its setup have only minor or moderate effects. Switching on the far-field correction lets us err slightly on the side of caution, i.e., our results will be biased~~
340 ~~towards the prior flux estimates. Without far-field correction, on the other hand, we expect errors from wrongly projecting any boundary bias onto the fluxes. For the station observation filtering (e.g., time window selection, outlier identification), we found a robust setting for our application because of sufficient observation coverage. However, this may become a problem for other applications in less well-observed circumstances. Tuning the model-observation uncertainty parameters proved important, and thus we chose them cautiously to avoid overfitting.~~

4.2 Comparison to other methods

Our Eulerian approach with sectoral segregation differs from other studies on CH₄ inversions for single countries, e.g., Henne et al. (2016) for Switzerland and Ganesan et al. (2015) for the United Kingdom that use Lagrangian transport models. The latter both qualitatively attribute deviations from the inventory reporting to the agriculture sector by comparing the spatial and/or temporal patterns in the posterior fluxes to sectoral a priori fluxes. A similar strategy for sectoral segregation based on
350 ~~spatial flux patterns~~ a known spatial distribution of fluxes is followed by Varon et al. (2022) and analyzed by Cusworth et al. (2021). For deriving sector estimates, some inversions assume a spatial correlation of gridded emissions within each sector (Rödenbeck et al., 2003; Meirink et al., 2008b; Bergamaschi et al., 2010). Based on the same assumption, Steiner et al. (2024b) and Tenkanen et al. (2025) construct ensembles of perturbed a priori fluxes to distinguish natural and anthropogenic fluxes utilizing the CarbonTracker Data Assimilation Shell (van der Laan-Luijkx et al., 2017). Notably, Tenkanen et al. (2025) avoid
355 the lateral boundary problem by simulating transport globally with nested zoom in Europe to estimate Finnish CH₄ emissions on a coarse resolution of $1^\circ \times 1^\circ$. In the present work, we take the next step by validating sectoral emissions reported to UNFCCC and analyzing possible false attributions, making use of a significantly higher model resolution.

Our results are qualitatively in line with the discrepancy of top-down estimates and UNFCCC reporting for Germany and the Benelux found in different regional inversions for the years 2018 and earlier (Petrescu et al., 2023; Bergamaschi et al.,
360 2022, 2018; Steiner et al., 2024b). Furthermore, it appears as a robust feature in our results that emissions from the UK plus Ireland agree well with reported emissions, in line with Bergamaschi et al. (2022) for the year 2018. For the French emissions,

our inversion shows a tendency towards slightly higher emissions similar to Steiner et al. (2024b), whereas other inversions suggest significantly higher emissions (Petrescu et al., 2023; Bergamaschi et al., 2022).

4.3 Limitations

365 ~~Though simulating. Although we simulate~~ emissions and transport in a large domain, we can only provide reliable emission estimates for selected countries (compare Fig. 3). Regions without notable uncertainty reduction and regions with known modeling difficulties do not benefit from our model setup. In Scandinavia, we find strong wetland emissions with insufficiently modeled fine-scale spatial and temporal variability. Combined with only small signals from non-LULUCF anthropogenic emissions, this leads to a low signal-to-noise ratio, which prevents conclusive results for Scandinavia. Furthermore, the synthesis inversion
370 may be prone to underestimating large localized sources due to transport errors ~~— an issue we address in Part 1 (Bruch et al., 2025a). We quantified this effect (see Sect. 5.6) and reduced the bias by introducing the posterior- R inversion as well as an additional model uncertainty.~~

Another limitation comes from the challenges for the regional flux inversion caused by biases in the lateral boundary conditions, ~~motivating our~~. ~~The uncertainty in lateral boundary concentrations motivates the~~ far-field ~~correction. The correction~~
375 ~~effectively reduces the influence of observations with low signal from categorized emissions, while leaving observations with strong signals mostly unchanged. This makes the estimation of well-observed emissions more robust. However, due to the selection of observations for the correction that is discussed in Part 1 (Bruch et al., 2025a). We expect that the~~ far-field ~~correction, this correction is more likely to increase the far field rather than decreasing it, leading to~~ ~~correction leads to~~ ~~more robust estimates for well-observed emissions, but it may also cause~~ a bias towards ~~the prior and towards~~ lower emission
380 estimates, ~~as was proven with synthetic experiments. We expect similar difficulties as long as the bias correction of the lateral boundaries is based on the same observations as the flux estimation. This highlights the difficulty of determining fluxes in a limited area when concentrations at the lateral boundaries cannot be directly inferred from dedicated observations.~~

In our highly resolved transport simulation, every flux category is numerically expensive. Aiming to validate reported German emissions, we could reduce the state space of the inversion to only 46 scaling factors with monthly time resolution. This
385 substantially limits the spatial and temporal variations that can be represented in the inversion. This approach is justified if the a priori fluxes already provide a realistic spatial distribution of all major CH₄ sources within each flux category. While this may be the case in Germany and neighboring countries, the constant scaling factors for large flux categories in more distant regions may be oversimplified and could lead to less accurate results in these regions. Moreover, adjusting ~~only a~~ few degrees of freedom may not be sufficient to obtain realistic flux estimates in regions with limited or highly uncertain information on a
390 priori fluxes, such as Scandinavia. ~~The scaling method would further fail to correct zero prior fluxes (Kountouris et al., 2018). However, this is less of a problem for CH₄, as inventories can collect where CH₄-emitting activities are normally located, but emission factors which translate the activities into emissions are generally not well known (Dammers et al., 2024).~~

When constructing the state space, we unevenly distributed the 46 degrees of freedom on our model domain – using 11 degrees of freedom for Germany and only four for mainland France plus Belgium and Luxembourg. But the choice of flux
395 categories affects the results and can lead to biases depending on the location of the observations (Kaminski et al., 2001). In

our application, this effect is small because of the good observation coverage in Germany. ~~Sensitivity tests with 5, 9 and 19 degrees of freedom in Germany (cases 503, 504, 506 in Table E1) indicate robust results for Germany and thereby confirm the potential of the synthesis inversion when focusing on a well-observed region.~~ This is checked in Part 1 (Bruch et al., 2025a) using sensitivity tests.

400 We exploit the sectoral discrimination of emission in a well-observed region as a key feature of our inversion method. This relies heavily on an accurate spatial distribution and completeness of the a priori fluxes, which appears to be sufficient for the major emitting sectors in Germany. Furthermore, the sector discrimination relies on resolving comparably small spatial scales, which poses a challenge to the transport modeling. A general problem in sector attribution is that sectors with large absolute uncertainty – such as agriculture – may be falsely blamed for any change in total emissions when the observations
405 do not clearly distinguish the sectors (see Appendix C). By quantifying this effect in the averaging kernels (see Fig. 7), we confirmed that in Germany agriculture can be distinguished from other anthropogenic emissions excluding LULUCF. Small sectors like natural plus LULUCF fluxes could not be reliably distinguished from large sectors such as agriculture, and we therefore combined smaller sectors like waste and public power into the larger category “non-agr.”.

4.4 Implications for future research

410 We chose the synthesis inversion for the first application of our modular inversion system, but designed this framework to be expandable to other inversion methods. For instance, most of the steps in the inversion can be applied with only minor adjustments when replacing the flux categories by an ensemble of randomly perturbed surface fluxes, similar to Steiner et al. (2024b), or by grid cell clusters as used by Estrada et al. (2024). Such applications with a larger state space are limited by the computational effort of the transport simulation, which is much higher than the computational effort of the inversion itself.
415 Similar to the inversion method, the far-field correction can be replaced by a different strategy for mitigating a boundary bias. For example, one could construct the far field based on an ensemble of boundary concentrations.

Further possibilities of extension ~~concern~~ involve other observation types, including satellite data. Our Eulerian system allows in principle the handling of large observation datasets without prohibitive computational effort, albeit changes in the construction and handling of R may be required when reaching $\gtrsim 10^5$ observations per time window. This potential is leveraged
420 by many inversion systems that use Eulerian transport simulations (e.g., Varon et al., 2022; Meirink et al., 2008a; Bergamaschi et al., 2013). The increasing availability of satellite data is especially interesting for constraining concentrations and emissions in ~~less-observed regions~~ regions with few or no ground-based observations, such as near the boundaries of our domain, which is an aspect to be addressed in future studies.

We identified potentials and risks in separating sectors based on ~~highly-resolved spatial flux patterns~~ the spatially highly
425 resolved distribution of fluxes. Extending this by temporal profiles for a priori fluxes offers ~~an a yet~~ untapped potential for ~~improvement.~~ future improvement of our system. Moreover, our inversion ~~would benefit from~~ could benefit from an a priori emission ~~ensembles~~ ensemble reflecting the uncertainty in the spatial and temporal distribution of the fluxes. ~~Significant improvements may become possible by distinguishing sectors at the observations~~ It remains to be explored whether improvements

in distinguishing sectors can be achieved in our system using co-tracers such as ethane for fossil CH₄ emissions (Ramsden et al., 2022; Mead et al., 2024) or by distinguishing carbon isotopes (Basu et al., 2022; Thanwerdas et al., 2024; Chandra et al., 2024).

5 Conclusions

We presented first results from a novel system for regional flux inversion designed to validate national CH₄ emission reporting. Applying this method to Central Europe in 2021 with a focus on Germany, we found ~~a significant increase in~~ significantly higher emissions from Germany and the Benelux ~~compared to the reporting~~. Careful estimation of posterior uncertainties revealed ~~that for the investigated year that the~~ total German posterior emissions are (32 ± 19) % higher than the respective anthropogenic emissions reported to the UNFCCC (submission 2024). ~~This increase is most likely due~~ With our inversion method the difference is attributed to emissions from the agriculture sector, possibly with contributions from ~~LULUCF~~ the LULUCF sector and natural sources. Our results were confirmed by validation with independent observation sites and by an exhaustive range of sensitivity tests ~~and by validation with independent observation sites~~ presented in Part 1 (Bruch et al., 2025a). Synthetic experiments with known truth ~~verified the~~ revealed the method's ability to distinguish ~~emission~~ the agricultural from the non-agricultural sectors in Germany, whereas disentangling possible influences from natural and LULUCF sources requires further work and possibly more observations.

~~Methodological~~ A methodological comparison to other regional inversion systems highlights the advantages of our method for the purpose of distinguishing emission sectors and ~~fitness for purpose for validation of its suitability for validating~~ national emission estimates. The qualitative gap between UNFCCC reporting and our estimates for Germany and the Benelux is consistent with earlier works (Petrescu et al., 2023; Bergamaschi et al., 2022, 2018; Steiner et al., 2024b). We complement these studies by providing an emission estimate for the German agriculture sector that can be directly compared to the national reporting, revealing a significant mismatch.

In this study we ~~only~~ presented the first application of an extensible, novel inversion system. Future developments may include the integration of satellite data, the incorporation of temporal profiles, a more comprehensive treatment of boundary conditions and flux uncertainties using ensemble methods, and an extension of the state space. The close connections to operational numerical weather prediction – especially in the underlying transport simulation – and the modular design establish the potential for long-term operational support of national emissions reporting.

Data availability. A collection of model data, inversion results, and data for reproducing most figures in this work is available at <https://doi.org/10.5281/zenodo.17414768> (Bruch et al., 2025b).

Appendix A: ~~Extended data tables and~~ Supplementary figures

~~{...}~~




Figure A1. Prior and posterior emissions (a) and scaling factors (b) for all flux categories, ordered by prior emissions. Horizontal lines indicate 95% confidence intervals. See Fig. 1 for the geographical definition of the flux categories and Fig. 2 for the resulting map of scaling factors. (a) If no sector is explicitly specified, the flux categories contain all anthropogenic fluxes excluding LULUCF. For flux categories marked with an asterisk, the inversion does not reduce the absolute uncertainty. Thus, reliable information is only gained by our inversion for flux categories without asterisk (see [Sect. 4.5](#) [Sect. 2.6](#)). Red color of the category names indicates a statistically significant increase of emissions. (b) Scaling factors are the raw results of our inversion, though here they are already combined for the whole year. The posterior scaling factor is defined as the [mean-center](#) of the methodological uncertainty range indicated by brown boxes.




Figure A2. (a–c) Seasonal cycle when using only observations from stations that were active during the whole year. We select those stations and sampling heights, for which we used at least two data points per day on at least 20 days of each month in 2021 in our main inversion. This yields selects 27 stations shown in (d) with $8.3 \cdot 10^4$ data points for the inversion (step 5 in Table 2), compared to 50 stations with $1.29 \cdot 10^5$ data points in the reference case (compare Fig. 4). Colored areas show the posterior uncertainties (95% confidence intervals), which were computed without excluding individual stations from the inversion and are therefore smaller than in Fig. 4. Prior emission rates are shown as dotted lines with small markers.




Figure A3. Seasonal cycle in observations at stations with elevation below 500m above sea level (a, b) and above 1000 m (c, d), supplementary to the discussion in Sect. 3.2. Thin blue lines represent the 10% quantile of each month, station, and sampling height for (a, c) observations and (b, d) model predictions (prior). The 10% quantile is chosen to minimize the effect of local pollution. Thick black lines indicate the mean of all selected stations and sampling heights. Thick red lines in (b) and (d) show the 10% quantile of the modeled far-field concentration. The flatland stations show a pronounced seasonal cycle with minimum in summer for both model and observations. This cycle is dominated by the contribution of the far field. The mountain stations have a weaker seasonal cycle.




Figure A4. Statistics of the relative (a) and absolute (b) improvement of the model–observation mismatch at independent validation stations for different choices of the error covariance matrix R discussed in Part 1 (Bruch et al., 2025a). The figure is analogous to Fig. 5, where the visualization and the data selection is explained. Here, we distinguish three inversion methods that differ in how R is constructed, as introduced in Sect. 2.5 of Part 1. No clear advantage of one method over the others can be seen. The diagonal R inversion has the lowest posterior RMSE at validation sites, followed by the posterior R and prior R inversion, but the differences are not statistically significant.

Appendix B: Averaging kernel matrices

As introduced in Sect. 3.4.2, the averaging kernel matrices A^{emis} and $A^{\text{scaling factors}}$ estimate the change in the posterior when
 460 changing the truth, $A^{\text{emis}} = \partial e^{\text{post}} / \partial e^{\text{truth}}$ where e denotes the vector of emissions. Here, we summarize how these matrices are
 estimated using either the ~~prior and posterior~~ error covariance matrices B and B_{post} , or the statistics from inversion runs with
 synthetic truth.

B1 Analytic estimate using error covariance matrices

We first estimate the sensitivity of the posterior scaling factor to the true emissions under the assumption that the transport
 465 model, far field, ~~and the flux pattern observations, and the a priori spatial distribution~~ within each flux category are perfect. Un-
 der these idealized assumptions, the model–observation mismatch ~~is $\mu = y - H s^{\text{prior}} - x^{\text{ff}} = H(s^{\text{truth}} - s^{\text{prior}})$ for given scaling~~
~~factors s is $\mu(s) = y - H s - x^{\text{ff}} = H(s^{\text{truth}} - s)$~~ where s^{truth} denotes the true scaling factors. Our ~~prior- R~~ “~~prior R~~ ” inversion
 will now maximize

$$P(s) \propto \exp \left[-\frac{1}{2} (s - s^{\text{truth}})^{\top} H^{\top} R^{-1} H (s - s^{\text{truth}}) - \frac{1}{2} (s - s^{\text{prior}})^{\top} B^{-1} (s - s^{\text{prior}}) \right] \quad (\text{B1})$$

$$470 \quad \propto \exp \left[-\frac{1}{2} (s - s^{\text{post}})^{\top} B_{\text{post}}^{-1} (s - s^{\text{post}}) \right]. \quad (\text{B2})$$

This yields $s^{\text{post}} = s^{\text{prior}} + A(s^{\text{truth}} - s^{\text{prior}})$ with the averaging kernel $A = I - B_{\text{post}} B^{-1}$ and ~~the posterior error covariance matrix~~
 $B_{\text{post}}^{-1} = H^{\top} R^{-1} H + B^{-1}$ (Rodgers, 2000). Knowing B and B_{post} , we can compute the averaging kernel A to estimate how the
 posterior scaling factors depend on the true scaling factors.

B2 Statistical estimate using synthetic experiments

475 ~~We aim to~~ ~~In the statistical approach, we~~ estimate the sensitivity of posterior scaling factors $\xi := s^{\text{post}} - s^{\text{prior}}$ to changes in
 the synthetic truth $\zeta := s^{\text{truth}} - s^{\text{prior}}$ ~~using 100 synthetic experiments with random synthetic truth s^{truth}~~ . Given a sample of N
 realizations $\{\xi^n\}_n$ and $\{\zeta^n\}_n$, we aim to find the scaling factor averaging kernel matrix A that solves

$$A = \arg \min_{A'} \sum_{n=1}^N \|\xi^n - A' \zeta^n\|^2. \quad (\text{B3})$$

For $\|x\|^2 = \sum_i x_i^2$, differentiation by A'_{ij} yields $0 = \sum_{n=1}^N \zeta_j^n (\xi^n - A \zeta^n)_i$ for all i, j and thereby

$$480 \quad A = \Xi Z^{-1}, \quad \Xi_{ij} = \sum_{n=1}^N \xi_i^n \zeta_j^n, \quad Z_{ij} = \sum_{n=1}^N \zeta_i^n \zeta_j^n. \quad (\text{B4})$$

Equation (B4) was used to produce panels (c) and (d) of Fig. 7.

Appendix C: Relevance of absolute prior uncertainty in sector attribution

When observations can detect a change in total emissions but cannot distinguish between different emission sectors, the sector-
 resolving inversion will change the sectoral distribution based on the prior uncertainties. To understand this problem quali-

tatively, we consider the worst case: We assume that fluxes from all sectors are uncorrelated in the prior but 100% spatially correlated such that they cannot be distinguished in the inversion. The a priori probability density for an emission vector e of sector emissions e_i is

$$P(e) \propto \exp \left[-\frac{1}{2} \sum_i (e_i - e_i^{\text{prior}})^2 \sigma_i^{-2} \right], \quad (\text{C1})$$

where σ_i denotes the a priori standard deviation of e_i . The inversion will yield a result for estimate the total emissions $e_{\text{tot}}^{\text{post}}$ that maximizes the probability $P(e)$ when including information from the observations such that the a posteriori probability density $P(e|y)$ is maximized. But by assumption, these observations do not distinguish between sectors such that Eq. (C1) remains valid the a posteriori probability density fulfills $P(e|y) \propto P(e)$ as long as $\sum_i e_i$ is fixed. We thus obtain the posterior emissions of the sectors by maximizing Eq. (C1) with the constraint $\sum_i e_i = e_{\text{tot}}^{\text{post}}$. By introducing a Lagrange multiplier, one can show² that this yields implies

$$e_i - e_i^{\text{prior}} = \alpha \sigma_i^2, \quad \alpha = \frac{e_{\text{tot}}^{\text{post}} - e_{\text{tot}}^{\text{prior}}}{\sum_i \sigma_i^2}. \quad (\text{C2})$$

This shows that sectors with larger absolute a priori uncertainty are disproportionally stronger corrected. Applied to our emission estimates for Germany, this implies that if the observations were unsuitable for distinguishing sectors, the inversion would attribute up to 95 % of the changes in total fluxes to the agriculture sector, which is responsible for 69 % of the total a priori emissions. Fortunately, this worst case scenario is not realistic because the observations do contain information on the different sectors as indicated e.g. by Figs. 6 and 7. But a tendency remains to correct the agriculture stronger than the other sectors.

Appendix D: Attempt to distinguish five sectors in Germany

Our setup for the transport simulation was designed to separte-separate five sectors in Germany: agriculture, natural plus LULUCF, waste, public power, and the sum of all other sectors (“other”). We test the separation of try to distinguish these sectors in sensitivity-test a separate inversion run, in which each of these sectors is scaled separately (sensitivity tests 506 (Table E1 and Fig. E1) and in Part 1 (Bruch et al., 2025a)). This inversion uses 19 separate scaling factors in Germany instead of 11. We find no notable changes in the posterior emissions compared to our reference setup, in which we combined waste, public power, and other into one larger sector “non-agr.”. However, the uncertainties and the averaging kernels change considerably. We assume an a priori 2σ uncertainty of $\pm 100\%$ for each sector-resolving flux category. Thus, splitting the total fluxes in more uncorrelated flux categories reduces the a priori uncertainty of the total fluxes.

Figure D1 shows the averaging kernel matrices (introduced in Sect. 3.4.2 and Appendix B) for the inversion when separating five sectors. These matrices indicate that waste, public power, and “other” cannot be distinguished: The corresponding columns Fig. D1(a) are approximately equal. Thus, trying to distinguish these sectors does not yield provide any additional information. By comparing the row and column for “non-agr.” to Fig. 7, we identify drawbacks of the attempt to distinguish smaller sectors.

²We define $L(e, \lambda) = -\frac{1}{2} \sum_i (e_i - e_i^{\text{prior}})^2 \sigma_i^{-2} + \lambda (e_{\text{tot}}^{\text{post}} - \sum_i e_i)$ and require $\frac{\partial L}{\partial e_i} = 0, \frac{\partial L}{\partial \lambda} = 0$.




Figure D1. Averaging kernel matrices of German sector emissions (a) and the corresponding scaling factors (b) when trying to distinguish sectors waste, public power and other, in-estimated using the same-representation-as-in-posterior error covariance matrix. Small matrices at the bottom indicate the ideal result. See Fig. 7 (a)–(b) for an explanation of the representation. Panel (a), third row, shows that increasing true emissions in any sector is expected to cause higher posterior agriculture emissions with a false attribution of 46 % to 70 %. The same row in panel (b) shows that when looking at relative changes in the emissions, the influence of the false attribution on the agriculture sector is not very large.

When trying to distinguish five sectors, the false attribution of emissions to the agriculture sectors is more severe than when distinguishing only three sectors (48 % compared to 28 %). Consequently, the expected error reduction in the combined non-agriculture sectors (excluding natural plus LULUCF) is better when considering only three sectors. Qualitatively, this is what we expect from Appendix C for cases where the observations are insufficient to distinguish the considered sectors.

Author contributions. VB and TR conceptualized the inversion method. VB implemented the inversion method and wrote the original draft together with AKW. TR configured the transport model. TR and BE interpolated the a priori flux data which BE collected. DJCO organized data streams of CH₄ concentrations and observations. JF, BM, AMB, DJCO, TR and VB contributed to testing and tuning the transport model. NB contributed to the model–observation comparison. AKW supervised and coordinated the project. All authors reviewed and edited the manuscript.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. In our simulations we use modified Copernicus Atmosphere Monitoring Service information and ECCAD products for initial and lateral boundary conditions, and for a priori fluxes. We thank Stefan Feigenspan, Christian Mielke, Theo Wernicke, John Akubia and Roland Fuß for helpful discussions and providing a priori emission fields. We thank Roland Potthast, Frank-Thomas Koch, Christoph

Gerbig, Dominik Brunner, Michael Steiner, David Ho, Thomas Kaminski, Hannes Imhof and our partners in the ITMS project for very helpful and inspiring discussions. We also wish to thank Peter Bergamaschi, Aurélie Colomb, Martine De Mazière, Lukas Emmenegger, Dagmar Kubistin, Irene Lehner, Kari Lehtinen, Markus Leuenberger, Cathrine Lund Myhre, Michal V. Marek, Simon O'Doherty, Stephen

530 M. Platt, Christian Plaß-Dülmer, Francesco Apadula, Sabrina Arnold, Pierre-Eric Blanc, Dominik Brunner, Huilin Chen, Lukasz Chmura, Łukasz Chmura, Sébastien Conil, Cédric Couret, Paolo Cristofanelli, Grant Forster, Arnoud Frumau, Christoph Gerbig, François Gheusi, Samuel Hammer, Laszlo Haszpra, Juha Hatakka, Michal Heliasz, Stephan Henne, Arjan Hensen, Antje Hoheisel, Tobias Kneuer, Eric Larmanou, Tuomas Laurila, Ari Leskinen, Ingeborg Levin, Matthias Lindauer, Morgan Lopez, Ivan Mammarella, Giovanni Manca, Andrew Manning, Damien Martin, Frank Meinhardt, Meelis Mölder, Jennifer Müller-Williams, Steffen Manfred Noe, Jarosław Nęcki, Mikael

535 Ottosson-Löfvenius, Carole Philippon, Joseph Pitt, Michel Ramonet, Pedro Rivas-Soriano, Bert Scheeren, Marcus Schumacher, Mahesh Kumar Sha, Gerard Spain, Martin Steinbacher, Lise Lotte Sørensen, Alex Vermeulen, Gabriela Vítková, Irène Xueref-Remy, Alcide di Sarra, Franz Conen, Victor Kazan, Yves-Alain Roulet, Tobias Biermann, Marc Delmotte, Daniela Heltai, Ove Hermansen, Kateřina Komínková, Olivier Laurent, Janne Levula, Chris Lunder, Per Marklund, Josep-Anton Morguí, Jean-Marc Pichon, Martina Schmidt, Damiano Sferlazzo, Paul Smith, Kieran Stanley, Pamela Trisolino and Giulia Zazzeri for providing the atmospheric observations~~for the stations listed in Table A1.~~

540 VB, DJCO, NB and AMB acknowledge funding by the German Federal Ministry for Education and Research (BMBF) in the ITMS project (grant 01LK2102B) as well as BE (grant 01LK2104A). Map plots were made with Natural Earth.

References

- Agustí-Panareda, A., Diamantakis, M., Massart, S., Chevallier, F., Muñoz-Sabater, J., Barré, J., Curcoll, R., Engelen, R., Langerock, B., Law, R. M., Loh, Z., Morguí, J. A., Parrington, M., Peuch, V.-H., Ramonet, M., Roehl, C., Vermeulen, A. T., Warneke, T., and Wunch, D.:
545 Modelling CO₂ weather – why horizontal resolution matters, *Atmos. Chem. Phys.*, 19, 7347–7376, [https://doi.org/10.5194/acp-19-7347-](https://doi.org/10.5194/acp-19-7347-2019)
2019, 2019.
- Basu, S., Lan, X., Dlugokencky, E., Michel, S., Schwietzke, S., Miller, J. B., Bruhwiler, L., Oh, Y., Tans, P. P., Apadula, F., Gatti, L. V., Jordan, A., Necki, J., Sasakawa, M., Morimoto, S., Di Iorio, T., Lee, H., Arduini, J., and Manca, G.: Estimating emissions of methane consistent with atmospheric measurements of methane and $\delta^{13}\text{C}$ of methane, *Atmos. Chem. Phys.*, 22, 15 351–15 377, [https://doi.org/10.5194/acp-](https://doi.org/10.5194/acp-22-15351-2022)
550 22-15351-2022, 2022.
- Bergamaschi, P., Krol, M., Meirink, J. F., Dentener, F., Segers, A., van Aardenne, J., Monni, S., Vermeulen, A. T., Schmidt, M., Ramonet, M., Yver, C., Meinhardt, F., Nisbet, E. G., Fisher, R. E., O'Doherty, S., and Dlugokencky, E. J.: Inverse modeling of European CH₄ emissions 2001–2006, *J. Geophys. Res. Atmos.*, 115, <https://doi.org/10.1029/2010JD014180>, 2010.
- Bergamaschi, P., Houweling, S., Segers, A., Krol, M., Frankenberg, C., Scheepmaker, R. A., Dlugokencky, E., Wofsy, S. C., Kort, E. A.,
555 Sweeney, C., Schuck, T., Brenninkmeijer, C., Chen, H., Beck, V., and Gerbig, C.: Atmospheric CH₄ in the first decade of the 21st century: Inverse modeling analysis using SCIAMACHY satellite retrievals and NOAA surface measurements, *J. Geophys. Res. Atmos.*, 118, 7350–7369, <https://doi.org/10.1002/jgrd.50480>, 2013.
- Bergamaschi, P., Karstens, U., Manning, A. J., Saunio, M., Tsuruta, A., Berchet, A., Vermeulen, A. T., Arnold, T., Janssens-Maenhout, G., Hammer, S., Levin, I., Schmidt, M., Ramonet, M., Lopez, M., Lavric, J., Aalto, T., Chen, H., Feist, D. G., Gerbig, C., Haszpra, L.,
560 Hermansen, O., Manca, G., Moncrieff, J., Meinhardt, F., Necki, J., Galkowski, M., O'Doherty, S., Paramonova, N., Scheeren, H. A., Steinbacher, M., and Dlugokencky, E.: Inverse modelling of European CH₄ emissions during 2006–2012 using different inverse models and reassessed atmospheric observations, *Atmos. Chem. Phys.*, 18, 901–920, <https://doi.org/10.5194/acp-18-901-2018>, 2018.
- Bergamaschi, P., Segers, A., Brunner, D., Haussaire, J.-M., Henne, S., Ramonet, M., Arnold, T., Biermann, T., Chen, H., Conil, S., Delmotte, M., Forster, G., Frumau, A., Kubistin, D., Lan, X., Leuenberger, M., Lindauer, M., Lopez, M., Manca, G., Müller-Williams, J., O'Doherty, S., Scheeren, B., Steinbacher, M., Trisolino, P., Vítková, G., and Yver Kwok, C.: High-resolution inverse modelling of
565 European CH₄ emissions using the novel FLEXPART-COSMO TM5 4DVAR inverse modelling system, *Atmos. Chem. Phys.*, 22, 13 243–13 268, <https://doi.org/10.5194/acp-22-13243-2022>, 2022.
- Bessagnet, B., Pirovano, G., Mircea, M., Cuvelier, C., Aulinger, A., Calori, G., Ciarelli, G., Manders, A., Stern, R., Tsyro, S., García Vivanco, M., Thunis, P., Pay, M.-T., Colette, A., Couvidat, F., Meleux, F., Rouil, L., Ung, A., Aksoyoglu, S., Baldasano, J. M., Bieser, J., Briganti, G., Cappelletti, A., D'Isidoro, M., Finardi, S., Kranenburg, R., Silibello, C., Carnevale, C., Aas, W., Dupont, J.-C., Fagerli, H., Gonzalez, L., Menut, L., Prévôt, A. S. H., Roberts, P., and White, L.: Presentation of the EURODELTA III intercomparison exercise – evaluation of the chemistry transport models' performance on criteria pollutants and joint analysis with meteorology, *Atmos. Chem. Phys.*, 16, 12 667–12 701, <https://doi.org/10.5194/acp-16-12667-2016>, 2016.
- Bruch, V., Rösch, T., Jiménez de la Cuesta Otero, D., Ellerhoff, B., Mamtimin, B., Becker, N., Blechschmidt, A.-M., Förstner, J., and Kaiser-
575 Weiss, A. K.: German methane fluxes estimated top-down using ICON-ART – Part 1: Ensemble-enhanced scaling inversion, submitted to ACP, 2025a.

- Bruch, V., Rösch, T., Jiménez de la Cuesta Otero, D., Ellerhoff, B., Mantimin, B., Becker, N., Blechschmidt, A.-M., Förstner, J., and Kaiser-Weiss, A. K.: German methane fluxes in 2021 estimated with an ensemble-enhanced scaling inversion based on the ICON-ART model, Zenodo [data set], <https://doi.org/10.5281/zenodo.15083479>, 2025b.
- 580 Canepa, E. and Builtjes, P. J. H.: Thoughts on Earth System Modeling: From global to regional scale, *Earth-Sci. Rev.*, 171, 456–462, <https://doi.org/10.1016/j.earscirev.2017.06.017>, 2017.
- Chandra, N., Patra, P. K., Fujita, R., Höglund-Isaksson, L., Umezawa, T., Goto, D., Morimoto, S., Vaughn, B. H., and Röckmann, T.: Methane emissions decreased in fossil fuel exploitation and sustainably increased in microbial source sectors during 1990–2020, *Commun. Earth Environ.*, 5, 1–15, <https://doi.org/10.1038/s43247-024-01286-x>, 2024.
- 585 Chen, H. W., Zhang, F., Lauvaux, T., Davis, K. J., Feng, S., Butler, M. P., and Alley, R. B.: Characterization of Regional-Scale CO₂ Transport Uncertainties in an Ensemble with Flow-Dependent Transport Errors, *Geophys. Res. Lett.*, 46, 4049–4058, <https://doi.org/10.1029/2018GL081341>, 2019.
- Citepa: Format Secten, <https://www.citepa.org/fr/secten/>, (last accessed: 18 March 2025), 2024.
- Cusworth, D. H., Bloom, A. A., Ma, S., Miller, C. E., Bowman, K., Yin, Y., Maasakkers, J. D., Zhang, Y., Scarpelli, T. R., Qu, Z., Jacob, D. J.,
 590 and Worden, J. R.: A Bayesian framework for deriving sector-based methane emissions from top-down fluxes, *Commun. Earth Environ.*, 2, 1–8, <https://doi.org/10.1038/s43247-021-00312-6>, 2021.
- Dammers, E., Tokaya, J., Mielke, C., Hausmann, K., Griffin, D., McLinden, C., Eskes, H., and Timmermans, R.: Can TROPOMI NO₂ satellite data be used to track the drop in and resurgence of NO_x emissions in Germany between 2019–2021 using the multi-source plume method (MSPM)?, *Geosci. Model Dev.*, 17, 4983–5007, <https://doi.org/10.5194/gmd-17-4983-2024>, 2024.
- 595 Deng, Z., Ciais, P., Tzompa-Sosa, Z. A., Saunio, M., Qiu, C., Tan, C., Sun, T., Ke, P., Cui, Y., Tanaka, K., Lin, X., Thompson, R. L., Tian, H., Yao, Y., Huang, Y., Lauerwald, R., Jain, A. K., Xu, X., Bastos, A., Sitch, S., Palmer, P. I., Lauvaux, T., D’Aspremont, A., Giron, C., Benoit, A., Poulter, B., Chang, J., Petrescu, A. M. R., Davis, S. J., Liu, Z., Grassi, G., Albergel, C., Tubiello, F. N., Perugini, L., Peters, W., and Chevallier, F.: Comparing national greenhouse gas budgets reported in UNFCCC inventories against atmospheric inversions, *Earth Syst. Sci. Data*, 14, 1639–1675, <https://doi.org/10.5194/essd-14-1639-2022>, 2022.
- 600 Department for Energy Security and Net Zero: Final UK greenhouse gas emissions national statistics: 1990 to 2022, <https://www.gov.uk/government/statistics/final-uk-greenhouse-gas-emissions-national-statistics-1990-to-2022>, (last accessed: 17 January 2025), 2024.
- East, J. D., Jacob, D. J., Balasus, N., Bloom, A. A., Bruhwiler, L., Chen, Z., Kaplan, J. O., Mickley, L. J., Mooring, T. A., Penn, E., Poulter, B., Sulprizio, M. P., Worden, J. R., Yantosca, R. M., and Zhang, Z.: Interpreting the Seasonality of Atmospheric Methane, *Geophys. Res. Lett.*, 51, e2024GL108494, <https://doi.org/10.1029/2024GL108494>, 2024.
- 605 Engelen, R. J., Denning, A. S., and Gurney, K. R.: On error estimation in atmospheric CO₂ inversions, *J. Geophys. Res. Atmos.*, 107, ACL10–1–ACL10–13, <https://doi.org/10.1029/2002JD002195>, 2002.
- Enting, I. G.: *Inverse Problems in Atmospheric Constituent Transport*, Cambridge University Press, Cambridge, England, UK, ISBN 978-0-52181210-8, <https://doi.org/10.1017/CBO9780511535741>, 2002.
- Estrada, L. A., Varon, D. J., Sulprizio, M., Nesser, H., Chen, Z., Balasus, N., Hancock, S. E., He, M., East, J. D., Mooring, T. A., Oort Alonso, A., Maasakkers, J. D., Aben, I., Baray, S., Bowman, K. W., Worden, J. R., Cardoso-Saldaña, F. J., Reidy, E., and Jacob, D. J.: Integrated
 610 Methane Inversion (IMI) 2.0: an improved research and stakeholder tool for monitoring total methane emissions with high resolution worldwide using TROPOMI satellite observations, *EGUsphere* [preprint], pp. 1–31, <https://doi.org/10.5194/egusphere-2024-2700>, 2024.
- Feigenspan, S., Wernicke, T., and Mielke, C.: Personal communication, 2024.
- Fuß, R. and Akubia, J.: Personal communication, 2024.

- 615 Ganesan, A. L., Manning, A. J., Grant, A., Young, D., Oram, D. E., Sturges, W. T., Moncrieff, J. B., and O'Doherty, S.: Quantifying methane and nitrous oxide emissions from the UK and Ireland using a national-scale monitoring network, *Atmos. Chem. Phys.*, 15, 6393–6406, <https://doi.org/10.5194/acp-15-6393-2015>, 2015.
- Gerbig, C., Körner, S., and Lin, J. C.: Vertical mixing in atmospheric tracer transport models: error characterization and propagation, *Atmos. Chem. Phys.*, 8, 591–602, <https://doi.org/10.5194/acp-8-591-2008>, 2008.
- 620 Henne, S., Brunner, D., Oney, B., Leuenberger, M., Eugster, W., Bamberger, I., Meinhardt, F., Steinbacher, M., and Emmenegger, L.: Validation of the Swiss methane emission inventory by atmospheric observations and inverse modelling, *Atmos. Chem. Phys.*, 16, 3683–3710, <https://doi.org/10.5194/acp-16-3683-2016>, 2016.
- ICOS RI: ICOS Handbook 2024, <https://doi.org/10.18160/28AV-80QR>, 2024.
- ICOS RI, Bergamaschi, P., Colomb, A., De Mazière, M., Emmenegger, L., Kubistin, D., Lehner, I., Lehtinen, K., Leuenberger, M.,
 625 Lund Myhre, C., Marek, M. V., O'Doherty, S., Platt, S. M., Plaß-Dülmer, C., Apadula, F., Arnold, S., Blanc, P.-E., Brunner, D., Chen, H., Chmura, L., Chmura, L., Conil, S., Couret, C., Cristofanelli, P., Forster, G., Frumau, A., Gerbig, C., Gheusi, F., Hammer, S., Haszpra, L., Hatakka, J., Heliasz, M., Henne, S., Hensen, A., Hoheisel, A., Kneuer, T., Larmanou, E., Laurila, T., Leskinen, A., Levin, I., Lindauer, M., Lopez, M., Mammarella, I., Manca, G., Manning, A., Martin, D., Meinhardt, F., Mölder, M., Müller-Williams, J., Noe, S. M., Nęcki, J., Ottosson-Löfvenius, M., Philippon, C., Pitt, J., Ramonet, M., Rivas-Soriano, P., Scheeren, B., Schumacher, M., Sha, M. K., Spain, G.,
 630 Steinbacher, M., Sørensen, L. L., Vermeulen, A., Vítková, G., Xueref-Remy, I., di Sarra, A., Conen, F., Kazan, V., Roulet, Y.-A., Biermann, T., Delmotte, M., Heltai, D., Hermansen, O., Komínková, K., Laurent, O., Levula, J., Lunder, C., Marklund, P., Morguá, J.-A., Pichon, J.-M., Schmidt, M., Sferlazzo, D., Smith, P., Stanley, K., Trisolino, P., Zazzeri, G., ICOS Carbon Portal, ICOS Atmosphere Thematic Centre, ICOS Flask And Calibration Laboratory, and ICOS Central Radiocarbon Laboratory: European Obspack compilation of atmospheric methane data from ICOS and non-ICOS European stations for the period 1984–2024; `obspace_ch4_466_GVeu_v9.2_20240502`,
 635 <https://doi.org/10.18160/9B66-SQM1>, 2024.
- IPCC, Calvo Buendia, E., Tanabe, K., Kranjc, A., Baasansuren, J., Fukuda, M., Ngarize, S., Osako, A., Pyrozhenko, Y., Shermanau, P., and Federici, S., eds.: 2019 Refinement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories, vol. 1, The Intergovernmental Panel on Climate Change (IPCC), <https://www.ipcc-nggip.iges.or.jp/public/2019rf/index.html>, 2019.
- Janssens-Maenhout, G., Pinty, B., Dowell, M., Zunker, H., Andersson, E., Balsamo, G., Bézy, J.-L., Brunhes, T., Bösch, H., Bojkov, B.,
 640 Brunner, D., Buchwitz, M., Crisp, D., Ciais, P., Counet, P., Dee, D., van der Gon, H. D., Dolman, H., Drinkwater, M. R., Dubovik, O., Engelen, R., Fehr, T., Fernandez, V., Heimann, M., Holmlund, K., Houweling, S., Husband, R., Juvyns, O., Kentarchos, A., Landgraf, J., Lang, R., Löschner, A., Marshall, J., Meijer, Y., Nakajima, M., Palmer, P. I., Peylin, P., Rayner, P., Scholze, M., Sierk, B., Tamminen, J., and Veefkind, P.: Toward an Operational Anthropogenic CO₂ Emissions Monitoring and Verification Support Capacity, *Bull. Am. Meteorol. Soc.*, 101, E1439–E1451, <https://doi.org/10.1175/BAMS-D-19-0017.1>, 2020.
- 645 Jähn, M., Kuhlmann, G., Mu, Q., Haussaire, J.-M., Ochsner, D., Osterried, K., Clément, V., and Brunner, D.: An online emission module for atmospheric chemistry transport models: implementation in COSMO-GHG v5.6a and COSMO-ART v5.1-3.1, *Geosci. Model Dev.*, 13, 2379–2392, <https://doi.org/10.5194/gmd-13-2379-2020>, 2020.
- Kaminski, T., Rayner, P. J., Heimann, M., and Enting, I. G.: On aggregation errors in atmospheric transport inversions, *J. Geophys. Res. Atmos.*, 106, 4703–4715, <https://doi.org/10.1029/2000JD900581>, 2001.
- 650 Kountouris, P., Gerbig, C., Rödenbeck, C., Karstens, U., Koch, T. F., and Heimann, M.: Technical Note: Atmospheric CO₂ inversions on the mesoscale using data-driven prior uncertainties: methodology and system evaluation, *Atmos. Chem. Phys.*, 18, 3027–3045, <https://doi.org/10.5194/acp-18-3027-2018>, 2018.

- Kuenen, J., Dellaert, S., Visschedijk, A., Jalkanen, J.-P., Super, I., and Denier van der Gon, H.: Copernicus Atmosphere Monitoring Service regional emissions version 4.2 (CAMs-REG-v4.2), Copernicus Atmosphere Monitoring Service (CAMs) [publisher], ECCAD [distributor], <https://doi.org/10.24380/0vzb-a387>, 2021.
- Kuenen, J., Dellaert, S., Visschedijk, A., Jalkanen, J.-P., Super, I., and Denier van der Gon, H.: CAMs-REG-v4: a state-of-the-art high-resolution European emission inventory for air quality modelling, *Earth Syst. Sci. Data*, 14, 491–515, <https://doi.org/10.5194/essd-14-491-2022>, 2022.
- Logan, J. A., Prather, M. J., Wofsy, S. C., and McElroy, M. B.: Tropospheric chemistry: A global perspective, *J. Geophys. Res. Oceans*, 86, 7210–7254, <https://doi.org/10.1029/JC086iC08p07210>, 1981.
- Manning, A. J., O'Doherty, S., Jones, A. R., Simmonds, P. G., and Derwent, R. G.: Estimating UK methane and nitrous oxide emissions from 1990 to 2007 using an inversion modeling approach, *J. Geophys. Res. Atmos.*, 116, <https://doi.org/10.1029/2010JD014763>, 2011.
- Mead, G. J., Herman, D. I., Giorgetta, F. R., Malarich, N. A., Baumann, E., Washburn, B. R., Newbury, N. R., Coddington, I., and Cossel, K. C.: Apportionment and Inventory Optimization of Agriculture and Energy Sector Methane Emissions Using Multi-Month Trace Gas Measurements in Northern Colorado, *Geophys. Res. Lett.*, 51, e2023GL105973, <https://doi.org/10.1029/2023GL105973>, 2024.
- Meirink, J. F., Bergamaschi, P., Frankenberg, C., D'Amelio, M. T. S., Dlugokencky, E. J., Gatti, L. V., Houweling, S., Miller, J. B., Röckmann, T., Villani, M. G., and Krol, M. C.: Four-dimensional variational data assimilation for inverse modeling of atmospheric methane emissions: Analysis of SCIAMACHY observations, *J. Geophys. Res. Atmos.*, 113, <https://doi.org/10.1029/2007JD009740>, 2008a.
- Meirink, J. F., Bergamaschi, P., and Krol, M. C.: Four-dimensional variational data assimilation for inverse modelling of atmospheric methane emissions: method and comparison with synthesis inversion, *Atmos. Chem. Phys.*, 8, 6341–6353, <https://doi.org/10.5194/acp-8-6341-2008>, 2008b.
- Munassar, S., Monteil, G., Scholze, M., Karstens, U., Rödenbeck, C., Koch, F.-T., Totsche, K. U., and Gerbig, C.: Why do inverse models disagree? A case study with two European CO₂ inversions, *Atmos. Chem. Phys.*, 23, 2813–2828, <https://doi.org/10.5194/acp-23-2813-2023>, 2023.
- Petrescu, A. M. R., Qiu, C., McGrath, M. J., Peylin, P., Peters, G. P., Ciais, P., Thompson, R. L., Tsuruta, A., Brunner, D., Kuhnert, M., Matthews, B., Palmer, P. I., Tarasova, O., Regnier, P., Lauerwald, R., Bastviken, D., Höglund-Isaksson, L., Winiwarter, W., Etiope, G., Aalto, T., Balsamo, G., Bastrikov, V., Berchet, A., Brockmann, P., Ciotoli, G., Conchedda, G., Crippa, M., Dentener, F., Groot Zwaafink, C. D., Guizzardi, D., Günther, D., Haussaire, J.-M., Houweling, S., Janssens-Maenhout, G., Kouyate, M., Leip, A., Leppänen, A., Lugato, E., Maisonnier, M., Manning, A. J., Markkanen, T., McNorton, J., Muntean, M., Oreggioni, G. D., Patra, P. K., Perugini, L., Pison, I., Raivonen, M. T., Saunois, M., Segers, A. J., Smith, P., Solazzo, E., Tian, H., Tubiello, F. N., Vesala, T., van der Werf, G. R., Wilson, C., and Zaehle, S.: The consolidated European synthesis of CH₄ and N₂O emissions for the European Union and United Kingdom: 1990–2019, *Earth Syst. Sci. Data*, 15, 1197–1268, <https://doi.org/10.5194/essd-15-1197-2023>, 2023.
- Petrescu, A. M. R., Peters, G. P., Engelen, R., Houweling, S., Brunner, D., Tsuruta, A., Matthews, B., Patra, P. K., Belikov, D., Thompson, R. L., Höglund-Isaksson, L., Zhang, W., Segers, A. J., Etiope, G., Ciotoli, G., Peylin, P., Chevallier, F., Aalto, T., Andrew, R. M., Bastviken, D., Berchet, A., Broquet, G., Conchedda, G., Dellaert, S. N. C., Denier van der Gon, H., Gütschow, J., Haussaire, J.-M., Lauerwald, R., Markkanen, T., van Peet, J. C. A., Pison, I., Regnier, P., Solum, E., Scholze, M., Tenkanen, M., Tubiello, F. N., van der Werf, G. R., and Worden, J. R.: Comparison of observation- and inventory-based methane emissions for eight large global emitters, *Earth Syst. Sci. Data*, 16, 4325–4350, <https://doi.org/10.5194/essd-16-4325-2024>, 2024.
- Ramsden, A. E., Ganesan, A. L., Western, L. M., Rigby, M., Manning, A. J., Foulds, A., France, J. L., Barker, P., Levy, P., Say, D., Wisher, A., Arnold, T., Rennick, C., Stanley, K. M., Young, D., and O'Doherty, S.: Quantifying fossil fuel methane emissions using observations

- of atmospheric ethane and an uncertain emission ratio, *Atmos. Chem. Phys.*, 22, 3911–3929, <https://doi.org/10.5194/acp-22-3911-2022>, 2022.
- Rieger, D., Bangert, M., Bischoff-Gauss, I., Förstner, J., Lundgren, K., Reinert, D., Schröter, J., Vogel, H., Zängl, G., Ruhnke, R., and Vogel, B.: ICON–ART 1.0 – a new online-coupled model system from the global to regional scale, *Geosci. Model Dev.*, 8, 1659–1676, <https://doi.org/10.5194/gmd-8-1659-2015>, 2015.
- Rigby, M., Manning, A. J., and Prinn, R. G.: Inversion of long-lived trace gas emissions using combined Eulerian and Lagrangian chemical transport models, *Atmos. Chem. Phys.*, 11, 9887–9898, <https://doi.org/10.5194/acp-11-9887-2011>, 2011.
- Rocher-Ros, G., Stanley, E. H., Loken, L. C., Casson, N. J., Raymond, P. A., Liu, S., Amatulli, G., and Sponseller, R. A.: Global methane emissions from rivers and streams, *Nature*, 621, 530–535, <https://doi.org/10.1038/s41586-023-06344-6>, 2023.
- Rodgers, C. D.: *Inverse Methods for Atmospheric Sounding*, vol. 2 of *Series on Atmospheric, Oceanic and Planetary Physics*, World Scientific Publishing Company, Singapore, ISBN 978-981-02-2740-1, <https://doi.org/10.1142/3171>, 2000.
- Rödenbeck, C., Houweling, S., Gloor, M., and Heimann, M.: CO₂ flux history 1982–2001 inferred from atmospheric data using a global inversion of atmospheric transport, *Atmos. Chem. Phys.*, 3, 1919–1964, <https://doi.org/10.5194/acp-3-1919-2003>, 2003.
- Schraff, C., Reich, H., Rhodin, A., Schomburg, A., Stephan, K., Periañez, A., and Potthast, R.: Kilometre-scale ensemble data assimilation for the COSMO model (KENDA), *Q. J. R. Meteorolog. Soc.*, 142, 1453–1472, <https://doi.org/10.1002/qj.2748>, 2016.
- Schröter, J., Rieger, D., Stassen, C., Vogel, H., Weimer, M., Werchner, S., Förstner, J., Prill, F., Reinert, D., Zängl, G., Giorgetta, M., Ruhnke, R., Vogel, B., and Braesicke, P.: ICON-ART 2.1: a flexible tracer framework and its application for composition studies in numerical weather forecasting and climate simulations, *Geosci. Model Dev.*, 11, 4043–4068, <https://doi.org/10.5194/gmd-11-4043-2018>, 2018.
- Segers, A. and Houweling, S.: CAMS global inversion-optimised greenhouse gas fluxes and concentrations, v22r2, Copernicus Atmosphere Monitoring Service [data set], <https://ads.atmosphere.copernicus.eu/datasets/cams-global-greenhouse-gas-inversion>, (last accessed: 18 April 2024), 2020.
- Seidel, D. J., Zhang, Y., Beljaars, A., Golaz, J.-C., Jacobson, A. R., and Medeiros, B.: Climatology of the planetary boundary layer over the continental United States and Europe, *J. Geophys. Res. Atmos.*, 117, <https://doi.org/10.1029/2012JD018143>, 2012.
- Steiner, M., Cantarello, L., Henne, S., and Brunner, D.: Flow-dependent observation errors for greenhouse gas inversions in an ensemble Kalman smoother, *Atmos. Chem. Phys.*, 24, 12 447–12 463, <https://doi.org/10.5194/acp-24-12447-2024>, 2024a.
- Steiner, M., Peters, W., Luijkx, I., Henne, S., Chen, H., Hammer, S., and Brunner, D.: European CH₄ inversions with ICON-ART coupled to the CarbonTracker Data Assimilation Shell, *Atmos. Chem. Phys.*, 24, 2759–2782, <https://doi.org/10.5194/acp-24-2759-2024>, 2024b.
- Stohl, A., Seibert, P., Arduini, J., Eckhardt, S., Fraser, P., Grealley, B. R., Lunder, C., Maione, M., Mühle, J., O’Doherty, S., Prinn, R. G., Reimann, S., Saito, T., Schmidbauer, N., Simmonds, P. G., Vollmer, M. K., Weiss, R. F., and Yokouchi, Y.: An analytical inversion method for determining regional and global emissions of greenhouse gases: Sensitivity studies and application to halocarbons, *Atmos. Chem. Phys.*, 9, 1597–1620, <https://doi.org/10.5194/acp-9-1597-2009>, 2009.
- Tenkanen, M. K., Tsuruta, A., Denier van der Gon, H., Höglund-Isaksson, L., Leppänen, A., Markkanen, T., Petrescu, A. M. R., Raivonen, M., Aaltonen, H., and Aalto, T.: Partitioning anthropogenic and natural methane emissions in Finland during 2000–2021 by combining bottom-up and top-down estimates, *Atmos. Chem. Phys.*, 25, 2181–2206, <https://doi.org/10.5194/acp-25-2181-2025>, 2025.
- Thanwerdas, J., Saunio, M., Berchet, A., Pison, I., and Bousquet, P.: Investigation of the renewed methane growth post-2007 with high-resolution 3-D variational inverse modeling and isotopic constraints, *Atmos. Chem. Phys.*, 24, 2129–2167, <https://doi.org/10.5194/acp-24-2129-2024>, 2024.

- Thompson, R. L., Krishnankutty, N., Pisso, I., Schneider, P., Stebel, K., Sasakawa, M., Stohl, A., and Platt, S.: Efficient use of a Lagrangian Particle Dispersion Model for atmospheric inversions using satellite observations of column mixing ratios, *EGUsphere* [preprint], pp. 1–28, <https://doi.org/10.5194/egusphere-2025-147>, 2025.
- UNFCCC: National Inventory Submissions 2024, <https://unfccc.int/ghg-inventories-annex-i-parties/2024>, (last accessed: 18 March 2025), 2024.
- van der Laan-Luijkx, I. T., van der Velde, I. R., van der Veen, E., Tsuruta, A., Stanislawska, K., Babenhauserheide, A., Zhang, H. F., Liu, Y., He, W., Chen, H., Masarie, K. A., Krol, M. C., and Peters, W.: The CarbonTracker Data Assimilation Shell (CTDAS) v1.0: implementation and global carbon balance 2001–2015, *Geosci. Model Dev.*, 10, 2785–2800, <https://doi.org/10.5194/gmd-10-2785-2017>, 2017.
- Varon, D. J., Jacob, D. J., Sulprizio, M., Estrada, L. A., Downs, W. B., Shen, L., Hancock, S. E., Nesser, H., Qu, Z., Penn, E., Chen, Z., Lu, X., Lorente, A., Tewari, A., and Randles, C. A.: Integrated Methane Inversion (IMI 1.0): a user-friendly, cloud-based facility for inferring high-resolution methane emissions from TROPOMI satellite observations, *Geosci. Model Dev.*, 15, 5787–5805, <https://doi.org/10.5194/gmd-15-5787-2022>, 2022.
- Weber, T., Wiseman, N. A., and Kock, A.: Global ocean methane emissions dominated by shallow coastal waters, *Nat. Commun.*, 10, 1–10, <https://doi.org/10.1038/s41467-019-12541-7>, 2019.
- Zängl, G., Reinert, D., Rípodas, P., and Baldauf, M.: The ICON (ICOsahedral Non-hydrostatic) modelling framework of DWD and MPI-M: Description of the non-hydrostatic dynamical core, *Quart. J. Roy. Meteorol. Soc.*, 141, 563–579, <https://doi.org/10.1002/qj.2378>, 2015.