

Process-based diagnostics using atmospheric budget analysis and nudging technique to identify sources of model systematic errors

Chihiro Matsukawa^{1,2}, José M. Rodríguez², and Sean F. Milton³

¹Japan Meteorological Agency, Tokyo, Japan

²Met Office, Exeter, UK

³University of Leeds, Leeds, UK

Correspondence: Chihiro Matsukawa (chi_matsukawa@met.kishou.go.jp)

Abstract. Identifying sources of model systematic errors is a fundamental step to successfully reduce them in general circulation models by improving the representation of relevant physical processes. In this study, we examine model error sources in the Met Office Unified Model at numerical weather prediction timescale by the combined use of two diagnostics: 1) the relaxation or "nudging" in which wind and/or temperature fields are relaxed back towards analyses throughout the simulations, and 2) atmospheric zonal-mean zonal momentum and thermal budgets. The budget analysis quantifies resolved processes and subsequently estimates unresolved processes as a residual, corresponding to model dynamics and physics, respectively. This correspondence is demonstrated by a direct comparison between the budgets and the model tendencies. A systematic error addressed in this paper is the Northern Hemisphere mid-latitude zonal wind bias in the lower stratosphere in boreal winter, characterized by an initial easterly bias that subsequently develops as a westerly bias. The momentum and thermal budget analysis for control and nudging experiments indicates that a mechanical forcing predominantly from parametrized gravity wave drag causes the easterly error and an overly strong temperature gradient around the tropopause is one of the main sources of the westerly error through the Coriolis forcing. The relevant warm bias over the tropical tropopause is mainly attributed to the budget residual term that corresponds to a thermal forcing dominated by radiative processes. This is consistent with the experimental result that temperature nudging over the tropical tropopause significantly reduces the westerly wind bias.

Copyright statement. The works published in this journal are distributed under the Creative Commons Attribution 4.0 License. This licence does not affect the Crown copyright work, which is re-usable under the Open Government Licence (OGL). The Creative Commons Attribution 4.0 License and the OGL are interoperable and do not conflict with, reduce or limit each other.

1 Introduction

A model systematic error, or bias, is a deviation of the mean model states from the corresponding mean observed states, affecting not only local regions but also other regions remotely through associated teleconnections. Continuous enhancements incorporated into general circulation models (GCMs), such as increasing horizontal and vertical resolutions, improving representations of dynamical and physical processes, and incorporating new physical components, have massively contributed to a signif-

same diagnostic as Klinker and Sardeshmukh (1992) and indicated systematic deficiencies in parametrized tendencies. They demonstrated that the incorporation of the new parametrizations associated with subgrid-scale orography leads to better initial momentum balances as well as reduced systematic errors in the general circulation. As an analogous diagnostic, van Niekerk et al. (2016) used an angular momentum budget analysis to examine a sensitivity of the resolved and parametrized surface drag to changes in horizontal resolution and parametrization. Their approach was to use the nudging framework to constrain the contribution from the angular momentum flux convergence (AMFC), and to determine the contributions from the resolved mountain torque and parametrized surface torque by balancing the AMFC. They found that a parametrized orographic torque in their model was excessive at lower resolutions.

The present study is aimed at understanding mechanisms of forecast error growth in global Met Office Unified Model (hereafter MetUM) at NWP timescales, and finding possible sources of the model errors. We focus on the Northern Hemisphere (NH) mid-latitude zonal wind errors from the upper troposphere to the stratosphere in boreal winter. Many studies have been dedicated to understanding mechanisms of the model systematic errors in the different version of the MetUM, particularly for a temperature bias around the tropopause (Hardiman et al., 2015; Bland et al., 2021). Hardiman et al. (2015) have executed several sensitivity experiments to examine the tropical tropopause warm bias that indicate that microphysical and radiative processes influence the temperature. Bland et al. (2021) have investigated a connection between a cold bias and a moist bias in the extratropical lowermost stratosphere through long-wave radiative effects. The approach adopted in this study is to analyse the atmospheric zonal-mean zonal momentum and thermal budgets in initialized NWP hindcasts and various nudging experiments. We use the globally nudged simulation to provide a best possible estimate of the truth to validate individual components of the momentum and thermal budgets in the MetUM. The budgets are examined to identify which component has a dominant contribution to initial and subsequent error growth through their comparison between the non-nudged control experiment and the globally nudged experiment. In addition, we apply the budget analysis to the regional nudging experiments to examine the impact of regional forcing on the general circulation patterns as well as the specific model errors. A combined application of the momentum and thermal budget analysis enables us to obtain information on wind circulation and temperature interactions and possible error compensations.

This paper is organized as follows. Section 2 describes the model and nudging framework used in this study, experiments carried out, and the details of the budget analysis. Section 3 presents a general view of the model systematic errors and highlights the NH mid-latitude zonal wind errors in boreal winter. Section 4 shows zonal-mean zonal momentum and thermal budgets in control and global nudging experiments and demonstrates a correspondence between budget components and model tendencies. In Sect. 5, we address the NH mid-latitude lower-stratospheric zonal wind errors and investigate sources of the errors using the budget analysis. Section 5 also examines remote and indirect impacts of regional nudging and its momentum budget. Finally, Sect. 6 summarizes the main conclusions and discusses further potential applications of the diagnostics and possible model error sources.

In this section, we describe details of the global MetUM used in this study, the nudging framework in the MetUM, the experimental design, and the zonal-mean zonal momentum and thermal budget analysis.

2.1 Model description

The MetUM is a numerical model which has been developed for use in regional and global simulations across weather to
 95 climate timescales (Cullen, 1993; Senior et al., 2011; Brown et al., 2012). Its scientific configuration used in this study is a
 global atmosphere and land configuration with a version of Global Atmosphere Land 9 (hereafter GAL9, in prep.; containing
 incremental upgrades from the former versions (e.g., Walters et al., 2019)), which is uncoupled with ocean and sea-ice dynam-
 ical model. The dynamical core, ENDGame, adopts a semi-implicit semi-Lagrangian formulation to solve the non-hydrostatic,
 fully compressible deep-atmosphere equations of motion (Wood et al., 2014). The atmospheric prognostic variables are zonal,
 100 meridional, and vertical wind velocity, dry virtual potential temperature, Exner pressure, dry density, and moist prognostic
 variables such as mixing ratio of moisture variables (water vapour, cloud water, and cloud ice) and cloud prognostic fields.
 Physical processes which are not represented or not resolved in the dynamical core, such as friction, condensation and evap-
 oration, radiative heating and cooling, and too small-scale phenomena to be resolved at the grid scale, are accounted for by
 parametrizations. Parametrizations employed by the MetUM include shortwave and longwave radiation (Edwards and Slingo,
 105 1996), microphysics (Wilson and Ballard, 1999), gravity wave drag consisting of non-orographic gravity wave (Scaife et al.,
 2002) and sub-grid scale orographic drag (Appendix in Vosper (2015) for details), convection (Gregory and Rowntree, 1990),
 turbulent mixing represented by the boundary-layer scheme (Lock et al., 2000), and large-scale cloud (Wilson et al., 2008a, b).
 Processes at the land surface and in the subsurface soil are represented by a community land surface model, the Joint UK
 Land Environment Simulator (JULES; Best et al., 2011). Contributions of the model dynamics and the individual physics
 110 parametrizations to the time evolution of prognostic variables can be diagnosed using increments per model time step, or
 tendencies that are equivalent to the increments per unit time.

Different applications of the MetUM across a wide range of temporal and spatial scales employ essentially the same model
 configuration, such as dynamical core and physics parametrizations. The MetUM used in this study has a horizontal resolution
 of N320 grid (0.5625° longitude $\times$ 0.375° latitude; approximately 40 km in the midlatitudes) with 70 vertical levels extending
 115 to 80 km altitude. The forecast model time step is 12 minutes. The horizontal resolution is lower than that of the Met Office's
 operational global deterministic NWP model (N1280 grid; approximately 10 km horizontal resolution in the midlatitudes).
 Under the across-scale approach, we adopt a moderate horizontal resolution to investigate large-scale model systematic errors,
 which provides benefits with regard to computational resources. However, since representations of the dynamical core and the
 physics parametrizations may have a sensitivity to model horizontal resolutions, extending our analysis to different resolutions
 120 of the MetUM may also be required.

2.2 Nudging

The nudging technique with Newtonian relaxation is a method that relaxes predicted variables of GCMs back towards given meteorological fields by adding an unphysical relaxation term to the prognostic equations (Jeuken et al., 1996). The nudging process incorporated into the MetUM at the very end of each model time step is written as follows (Telford et al., 2008; van Niekerk et al., 2016):

$$X_F = X_M + \frac{\Delta t}{\tau} \frac{\delta t}{\tau} (X_A - X_M) \quad (1)$$

where X is the prognostic model variable at the current model time step, Δt is the time interval of the model integration (i.e., 12 minutes in this study), and τ is the relaxation timescale of nudging. Subscripts F , M , and A denote the variables after nudging, those after dynamics and physics calculations just before nudging, and those used as a nudging forcing, respectively. In this study, the 6-hourly operational MetUM analysis at N1280 grid is regridded onto the model resolution (i.e., N320 grid) and then used as a forcing. Since the analysis data are available every 6 hours, they are linearly interpolated into each model time step. The choice of the relaxation timescale is arbitrary. The shorter the relaxation timescale is, the more strongly the model prognostic variables are relaxed towards forcing data (i.e. MetUM analysis data). Too long relaxation timescale is ineffective, and too short relaxation timescale could make the model unstable (Telford et al., 2008). We select relaxation timescale used in this study to be 6 hours which corresponds to the temporal spacing of the MetUM analysis. Notice that nudging is applied throughout model integration regardless of the relaxation timescale.

Additional increments due to nudging, which are expressed by the second term of the right-hand side of Eq. (1), are calculated as a difference of the variables between forcing data X_A and the constrained model predictions X_M within the nudged run multiplied by the relaxation coefficient. Therefore, the nudging increments are comparable to the forecast errors at a particular timescale depending on the relaxation coefficient least those growing over the timescale of the relaxation. Since the nudging forcing can alter the time evolution of nudged prognostic variables alongside the model dynamics and physics, the nudging tendencies that are equivalent to the increments per unit time and are comparable with the model tendencies due to the dynamics and physics.

The prognostic model variables X which can be constrained by nudging are zonal wind, meridional wind, and potential temperature. Other prognostic variables, such as vertical wind velocity and mixing ratio of moisture variables, are allowed to evolve freely and respond to the forced variables. Nudged domains and model levels can be prescribed arbitrarily and are accompanied by interfacing transition zones and layers to ensure a smooth transition between the nudged and free-running parts of the simulation. The transition zones and layers are smoothed using the hyperbolic tangent function over 10 degrees in the horizontal and the linear function over two model levels in the vertical.

2.3 Experimental design

To examine model error growth in boreal winter at NWP timescale, deterministic 15-day forecasts are started at 00:00 UTC between 16 November 2018 and 27 February 2019 and evaluated over the period from the December 2018 to the February 2019

Table 1. Control and nudging experiments executed. U, V, and Θ in nudged variables indicate zonal wind velocity, meridional wind velocity, and potential temperature, respectively.

Experiments	Nudged domains	Nudged model levels	Nudged variables
CNTL	N/A	N/A	N/A
GLN	Globe (0° – 360° , 90° S– 90° N)	Whole model levels	U, V, Θ
GLNT	Globe (0° – 360° , 90° S– 90° N)	Whole model levels	Θ
NHTrpT	NH Tropics (0° – 360° , 0° – 30° N)	Whole model levels	Θ
NHTrpTrpT	NH Tropics (0° – 360° , 0° – 30° N)	Around the tropopause (approx. 200–70 hPa)	Θ
NHHLT	NH High Latitude (0° – 360° , 60° N– 90° N)	Whole model levels	Θ
NHHLTTrpT	NH High Latitude (0° – 360° , 60° N– 90° N)	Around the tropopause (approx. 300–100 hPa)	Θ

(90 cases in total). A series of the operational global MetUM analyses produced by the data assimilation system, Hybrid-4DVar (Clayton et al., 2013) based on GA6.1 configuration (Walters et al., 2017) operational in 2018/2019, is spatially interpolated to the model resolution and used as the initial conditions. Lower boundary conditions are given by the sea surface temperature and sea ice concentration of the Operational Sea Surface Temperature and Ice Analysis (OSTIA; Donlon et al., 2012) products fixed at the field on initial dates throughout the 15-day simulations.

The momentum and thermal budgets described below in Sect. 2.4 and model tendencies due to dynamics and individual parametrizations (and nudging forcing) are calculated using the experimental data. The budgets are analysed at the same horizontal resolution as the model on 28 pressure levels (i.e., 1000, 950, 925, 850, 700, 600, 500, 400, 300, 250, 200, 150, 100, 70, 50, 30, 20, 10, 7, 5, 3, 2, 1, 0.7, 0.5, 0.3, 0.2, 0.1 hPa) with masking of the levels below the ground. On the other hand, model tendencies are evaluated at the 70 model native levels. Temporal-mean variables required in the analysis of the budgets and the model tendencies are calculated from the fields at each time step in the model.

Control and various nudging experiments executed in this study are summarized in Table 1. We apply momentum and thermal budget analysis (see details in Sect. 2.4) to the control experiment referred to as CNTL and the global nudging experiment referred to as GLN. ~~Considering~~We consider individual budget components of GLN as the best possible estimate of truth instead of analysis data, ~~which is unable to provide the temporal-mean variables required in this study, we use GLN to validate the budget components in CNTL and identify which components contribute to a particular error growth. This is why the reasonably short relaxation timescale is selected in the nudging experiments.~~ Nudging frameworks which constrain a part of the variables could bring a better understanding of feedbacks among variables and disentangle compensating errors specifically between horizontal wind and temperature (Wehrli et al., 2018). In addition, regional nudging, in which the model variables are relaxed back towards the analyses over subdomains where there might be significant systematic errors, potentially provide experimental insights into origins of the errors. We performed another global nudging experiment in which only temperature is constrained over the globe, referred to as GLNT, and nudging sensitivity experiments in which temperature is constrained over

175 different domains at various altitudes (i.e., NHTrpT, NHTrpTrpT, NHHLT, and NHHLTrpT; see Table 1 in detail) to test how much wind biases in CNTL could be influenced by temperature biases and diagnose the remote impact of nudging. Note that the tropopause temperature nudging in NHTrpTrpT and NHHLTrpT is applied in a different range of pressure levels depending on their latitudinal positions.

2.4 ~~Zonal~~ Atmospheric zonal momentum and thermal budgets

180 The framework of the atmospheric zonal-mean momentum and thermal budgets is a well-established diagnostic tool for examining the contribution of resolved and unresolved processes to the large-scale structure of the atmosphere. Based on the primitive equations, the zonal-mean zonal momentum and thermal budget equations in spherical and pressure coordinates are derived as below (Hartmann, 1976; Andrews, 1987):

$$\begin{aligned}
 \frac{\partial [\bar{u}]}{\partial t} = & - \underbrace{\left(\frac{[\bar{v}]}{a \cos \phi} \frac{\partial [\bar{u}] \cos \phi}{\partial \phi} + [\bar{\omega}] \frac{\partial [\bar{u}]}{\partial p} \right)}_{\text{Mean Stationary Flow Advection}} \\
 & - \underbrace{\left(\frac{1}{a \cos^2 \phi} \frac{\partial [\bar{u}^* \bar{v}^*] \cos^2 \phi}{\partial \phi} + \frac{\partial [\bar{u}^* \bar{\omega}^*]}{\partial p} \right)}_{\text{Stationary Eddy Component}} \\
 & - \underbrace{\left(\frac{1}{a \cos^2 \phi} \frac{\partial [\bar{u}' v'] \cos^2 \phi}{\partial \phi} + \frac{\partial [\bar{u}' \omega']}{\partial p} \right)}_{\text{Transient Eddy Component}} \\
 & + \underbrace{f [\bar{v}]}_{\text{Coriolis}} + \underbrace{[\bar{F}_u]}_{\text{Residual}}
 \end{aligned} \tag{2}$$

$$\begin{aligned}
 \frac{\partial [\bar{T}]}{\partial t} = & - \underbrace{\left(\frac{[\bar{v}]}{a} \frac{\partial [\bar{T}]}{\partial \phi} + [\bar{\omega}] \frac{\partial [\bar{T}]}{\partial p} - \frac{R_d}{p c_p} [\bar{\omega}] [\bar{T}] \right)}_{\text{Mean Stationary Flow Component}} \\
 & - \underbrace{\left(\frac{1}{a \cos \phi} \frac{\partial [\bar{v}^* \bar{T}^*] \cos \phi}{\partial \phi} + \frac{\partial [\bar{\omega}^* \bar{T}^*]}{\partial p} - \frac{R_d}{p c_p} [\bar{\omega}^* \bar{T}^*] \right)}_{\text{Stationary Eddy Component}} \\
 & - \underbrace{\left(\frac{1}{a \cos \phi} \frac{\partial [\bar{v}' T'] \cos \phi}{\partial \phi} + \frac{\partial [\bar{\omega}' T']}{\partial p} - \frac{R_d}{p c_p} [\bar{\omega}' T'] \right)}_{\text{Transient Eddy Component}} \\
 & + \underbrace{\frac{[\bar{Q}]}{c_p}}_{\text{Residual}}
 \end{aligned} \tag{3}$$

where u and v are zonal and meridional components of wind velocity, ω is vertical pressure velocity, a is the mean radius of Earth, ϕ is latitude, p is pressure, f is the Coriolis parameter, T is temperature, R_d is the gas constant, c_p is the specific heat

190 of air at constant pressure, F_u is a mechanical forcing term of the zonal momentum equation, and Q is a diabatic heating rate. Overbars and primes denote the temporal mean and departure from the temporal mean, respectively, and square brackets and asterisks denote the zonal mean and departure from the zonal mean, respectively. The temporally and zonally averaged flow (e.g., $[\bar{u}]$) is referred to as mean stationary flow, and the departure from the temporal mean (e.g., $u' = u - \bar{u}$) is referred to as transient eddy, and the departure of the temporally averaged variable from its zonal mean (e.g., $\bar{u}^* = \bar{u} - [\bar{u}]$) is referred to as
195 stationary eddy.

The right-hand side of Eq. (2), in order, represents an advection of mean zonal wind by mean stationary flows, a convergence of stationary eddy momentum fluxes, that of transient eddy momentum fluxes, Coriolis forcing, and a residual term. The transient and stationary eddy fluxes are evaluated, for instance, as $\overline{u'v'} = \overline{uv} - \bar{u}\bar{v}$ and $[\bar{u}^*\bar{v}^*] = [\bar{u}\bar{v}] - [\bar{u}][\bar{v}]$, respectively. In Eq. (3), the right-hand side, from left to right, shows a warming/cooling due to advection of mean temperature by mean
200 stationary flow and adiabatic heating with mean vertical motions, a convergence and an energy conversion of stationary eddy heat flux, those of transient eddy heat flux, and a residual term. The terms other than the residual term are interpreted as contributions from resolved dynamical components and are evaluated directly from ~~forecast or analysis data~~ the forecast data specifically u , v , ω , and T fields. As in the manner in Martineau et al. (2016), a second-order centered difference scheme is used for calculating meridional and vertical derivatives, while meridional derivatives over the poles and vertical derivatives at
205 the top or bottom of the pressure levels require a first-order difference with their adjacent grids.

The last term of right-hand side of Eq. (2, 3), determined as a residual of the other terms including the left-hand side and referred to as residual term, represents contributions of the processes which are not represented by the resolved processes. In the partial differential equations of Eq. (2, 3), the residual term corresponds to a source or sink of the momentum and thermodynamic equations which consists of the frictional forcing and the diabatic heating due to radiative forcing, latent heat releases,
210 and latent and sensible heat fluxes at the surface. On the other hand, the residual ~~terms in the discretized equation adopted in the numerical model~~ term diagnosed using spatially discretized data could include effects of subgrid processes, which are too-small scale to be resolved in a grid scale, on a grid-box mean field as well as the frictional or diabatic forcing. ~~These unresolved processes are being parametrized in the model.~~ Ideally, the ~~total parametrized forcing should be~~ residual term should be qualitatively equal to the ~~residual term (Holopainen, 1987). In the case of nudging experiments, forcing due to nudging~~
215 ~~should also constitute a part of the residual term. However, in our study, they do not necessarily coincide because our residual term could contain other effects including imbalances caused by nudging increments (in the case of nudging experiments) and computational errors through the interpolation from model levels to pressure levels and the numerical methods employed to evaluate the budgets (Martineau et al., 2016).~~ total parametrized forcing calculated in the model (Holopainen, 1987), although this is not necessarily the case as described in Sect. 2.5. It is noteworthy that "residual" in this study, which is equivalent
220 to estimated unresolved tendencies, is defined differently from "residual" termed in some previous studies (e.g., Klinker and Sardeshmukh, 1992; Milton and Wilson, 1996), which is the difference between estimated unresolved tendencies in the analysis and parametrized physics tendencies. ~~In our framework, the physics tendencies parametrized in the model are not used for the budget calculation at all.~~

There are several options of how to define and compute temporal-mean fields denoted by overbars in Eq. (2, 3), and it
 225 **definitely** changes what the temporal derivative term on the left-hand side means. In this study, overbars represent the temporal
 averaging over the given forecast length in each case, not over the 3 months of the experimental period. Thus, a temporal-mean
 field over the forecast length (e.g., $\overline{u}$) varies among experiments and also among cases. The temporal average over the forecast
 length is computed using the fields at each model time step, which is the reason why globally nudged simulation data instead
 of analysis data is used to calculate individual budget components for validations because the analysis data has only 6-hourly
 230 instantaneous fields. This definition of temporal mean allows the budget equation to describe a time evolution over a particular
 forecast length in a single case, which makes it straightforward to diagnose forecast error growth at NWP timescale. The left-
 hand side of the Eq. (2, 3) indicating a total tendency of the variable is evaluated as a temporal change from initial conditions
 to model states at a given forecast lead time.

The momentum and thermal budget analysis is applied to each simulated case individually, and then averaged over the cases
 235 if mean budgets are examined. In this context, the terms "mean flow" and "transient eddy" in this paper should be interpreted
 cautiously because the methodology is different from that commonly used in climatological studies (e.g., Peixoto and Oort,
 1992). The partition of the nonlinear advective term among three components including mean stationary flow, stationary eddies,
 and transient eddies becomes slightly more complex and critically depends on the forecast lead time in this study. For example,
 in the momentum budget from an initial condition to a forecast at Day 1, if a migratory cyclone or anticyclone is simulated to
 240 be quasi-stationary throughout that time period, it is identified as the stationary eddy rather than the transient eddy. Therefore,
 the contribution of the transient eddy flux becomes smaller at a shorter forecast lead time, and the mean stationary flow and
 the stationary eddy components are responsible for the remaining part. Although the Coriolis forcing term involves the mean
 stationary flow and constitutes a part of the resolved processes, it can be independently evaluated regardless of the forecast
 length because of its linearity. The partition should depend not only on the definition of the temporal mean but also on the
 245 horizontal and vertical resolution of the model (and data) and the configuration of the model dynamics and physics.

2.5 Atmospheric budgets and model tendencies

Whilst the model dynamics and physics tendencies themselves, calculated in the model online, are not used for the offline
 budget calculation in our framework, the diagnosed resolved processes and the residual term correspond to the model dynamics
 and physics tendencies, respectively. A comparison of two diagnostics, atmospheric budgets and model tendencies, is helpful
 250 in understanding their physical meaning. The budget analysis diagnosed using forecast data on pressure levels quantifies
 contributions to the total tendencies from the resolved processes and the unresolved processes. On the other hand, model
 tendencies of prognostic variables are calculated by dynamical core and physics parametrizations in numerical models at each
 model time step during their time integration at each model level. Individual components of both atmospheric budgets and
 model tendencies represent contributions of the processes in the respective framework to the total tendency of the corresponding
 255 variable. Therefore, the atmospheric budget diagnostics can be compared with the model tendencies to demonstrate a qualitative
 similarity between the unresolved term in the budget equation and the model tendencies of the physics parametrizations and
 between the resolved term and the tendencies of the model dynamics.

The correspondence between the budget diagnostics and the model tendencies is expected from the derivation of the budget equation, but they are not necessarily exactly equivalent. The residual term could contain effects from imbalances caused by nudging increments (in the case of nudging experiments) and computational errors through the interpolation of forecast fields from model levels to pressure levels and the numerical methods employed to evaluate the budgets (Martineau et al., 2016) as well as the unresolved processes. A discrepancy between the primitive equations used in the budget diagnostics and the governing equations of the MetUM could also generate a difference between the resolved processes and the model dynamics tendencies and therefore between the residual term and the parametrization tendencies to some extent. In the case of nudging experiments, forcing due to nudging, which is associated with the very short timescale model error in the MetUM, should also constitute a part of the residual term.

2.6 Validation of atmospheric budgets

To validate the budget components in CNTL and identify which components contribute to a particular forecast error growth, we use the nudging technique to create quasi-analysis data and calculate the best possible estimate of truth of atmospheric budgets. An error of a given forecast variable against analysis in a single case, defined as its difference between forecast and analysis, is proportional to an error of total tendency as follows:

$$\begin{aligned} x_e(t_i; \Delta t) &\equiv x_f(t_i; \Delta t) - x_a(t_i + \Delta t) \\ &= \left\{ \overline{\left(\frac{\partial x}{\partial t} \right)_{f, t_i}} - \overline{\left(\frac{\partial x}{\partial t} \right)_{a, t_i}} \right\} \Delta t \end{aligned} \quad (4)$$

where x is a given variable, t_i is an initialized time in a given single case, and Δt is a forecast length. The subscripts e , f , and a denote error, forecast, and analysis, respectively. The overbars are the same as defined in Eq. (2, 3). In this study, the total tendency of the analysis averaged over a given forecast length is approximated by that of GLN, and we refer to a difference in budget components between CNTL and GLN as "quasi-error". As a result, the validation of the momentum and thermal budgets can decompose a quasi-error in the total tendencies into individual budget components.

The atmospheric budgets of GLN should be interpreted with caution because they have an error against the analysis. Firstly, nudged variables in nudging experiments still have errors against the MetUM analysis. The nudging in some cases is not able to relax sufficiently towards the forcing data with a moderate relaxation timescale especially if the error grows rapidly at a very short timescale. In addition, the vertical velocity, required by the budget analysis but not directly nudged, is driven by the horizontal wind and temperature nudging forcing. Therefore, there should be a non-negligible error in the individual budget components in GLN, making it difficult to precisely evaluate the residual term as a consequence. Secondly, since the 6-hourly MetUM analyses used as a forcing data of the nudging are linearly interpolated into each model time step as part of the nudged simulation methodology, the temporal interpolation leads to an underestimation of the intra-6-hour variability, and therefore results in underestimated transient eddy components and consequently overestimated residual terms. Although these deficiencies might affect a quasi-error in budget components of CNTL against GLN to some extent, it is reasonable to use GLN as the best estimate of truth.

290 The interpretation of differences in the residual term among experiments is slightly complex. It is attributed, by definition,
to differences in the other resolved terms because the residual term in each experiment is determined by the sum of the
other terms to close the budget. From another perspective, the residual term is expected to correspond to the sum of the
physics parametrization tendencies and the nudging forcing. The quasi-error in the residual term against GLN could provide
some implications for deficiencies in the parametrizations through a comparison between the residual term and the individual
295 parametrization tendencies.

Unlike the budget diagnostics, a difference in individual model tendencies between CNTL and GLN does not show a
quasi-error in the model tendencies. The physics parametrization schemes themselves used in CNTL and GLN are identical and
have substantial deficiencies in principle. The parametrization tendencies in GLN are calculated using the deficient scheme with
nudged grid-box mean fields which are much closer to the analysis than CNTL. The difference in parametrization tendencies
300 between two experiments represents a difference in the responses of the parametrization scheme to different grid-box mean
fields.

3 Model systematic errors

Prior to analysing the momentum and thermal budgets, we present a general view of the model systematic errors in zonal
wind and temperature. Figure 1 exhibits mean errors in zonal-mean zonal wind and temperature in CNTL against the MetUM
305 analysis in December-January-February (DJF) 2018/2019 at specific forecast lead times. The CNTL simulations up to Day
15 show a westerly wind bias in the NH and Southern Hemisphere (SH) midlatitudes from the upper troposphere to the
stratosphere, a westerly wind bias in the tropical upper troposphere, a warm temperature bias around the tropical tropopause, a
cold temperature bias around the tropopause over the poles, and a cold temperature bias in the stratosphere. Some of the biases
exist even at Day 1 and evolve steadily in the same location, such as the westerly bias in the tropical upper troposphere, the
310 warm bias around the tropical tropopause, and the cold bias in the stratosphere. The NH mid-latitude westerly wind bias in the
stratosphere appears to shift poleward gradually as a forecast lead time gets longer. Most of the pronounced errors in Fig. 1 are
also discernible in 20-years low-resolution climate simulations with a minor sensitivity to horizontal resolution (not shown).
These errors develop quickly in the NWP timescale and remain in the climate timescale.

The westerly wind bias in the NH mid-latitude stratosphere is consistent with the warm temperature bias in the tropical
315 tropopause and the cold temperature bias over the poles through thermal wind relation. Although causality is unclear, an
increase in the zonal wind error with altitude balances with an excessive meridional temperature gradient arising from the
temperature biases. To investigate these biases, we focus ~~in~~ on three features highlighting the pronounced zonal-mean system-
atic errors: A) zonal wind in the NH mid-latitude stratosphere averaged from 30°N to 60°N at 50 hPa, B) temperature around
the NH tropical to subtropical tropopause averaged from the equator to 30°N at 70 hPa, and C) temperature around the NH
320 high-latitude tropopause averaged from 60°N to 90°N at 200 hPa. These features are illustrated by the black horizontal lines
A, B, and C in Fig. 1(e, f). Figure 2 presents the zonal wind and temperature forecast error growth over these domains in the
individual cases up to Day 7 (Fig. 2(a–c)) and time evolution of the zonal wind error averaged over all the cases (Fig. 2(d)).




Figure 1. Zonal-mean errors of the CNTL forecast at a forecast lead time of (a, b) Day 1, (c, d) Day 5, and ~~Day 15~~ (from top to bottom, e, f) Day 15 averaged over the 90 cases from 1 Dec. 2018 to 28 Feb. 2019 in (a, c, e) zonal wind velocity [m s^{-1}], and (b, d, f) temperature [K]. Contour indicates the forecasts, and colour indicates mean errors against the MetUM analysis. The black horizontal lines A, B, and C are inserted for later reference (Figs. 2, 12, 13, and 14; see the text in detail).

The forecasts at Day 0 obviously have no errors against the analysis because the forecasts are started from the corresponding MetUM analysis. Timeseries of the CNTL forecasts reveal the noticeable drifts of the wind and temperature in most of the cases. Zonal wind tends to increase relative to the analysis in the NH mid-latitude lower stratosphere (Fig. 2(a)). Meanwhile, temperature increases around the tropical tropopause (Fig. 2(b)) and decreases near the NH polar tropopause (Fig. 2(c)) relative to the analysis. In particular, these temperature errors develop quickly and systematically in almost all the cases and are




Figure 2. (a–c) Timeseries of the MetUM analysis and the CNTL and GLN forecasts of the individual cases up to Day 7 for (a) area-averaged zonal wind velocity [m s^{-1}] over 30°N – 60°N at 50 hPa, (b) area-averaged temperature [K] over 0° – 30°N at 70 hPa, and (c) area-averaged temperature [K] over 60°N – 90°N at 200 hPa. These features are depicted by the line A, B, and C in Fig. 1, respectively. (d) Time evolution of NH mid-latitude (30°N – 60°N) 50 hPa zonal wind errors [m s^{-1}] against the MetUM analysis, shown in (a), averaged for DJF 2018–2019. Error bars indicate a 95 % confidence interval estimated using the bootstrap method with 10,000 resamples.

statistically significant at the 95 % confidence level throughout the forecast lead time up to Day 15. The rapid error growth and consistent drifts suggest that these errors are relevant to the model physics rather than changes in the atmospheric circulation (Martin et al., 2010; Ma et al., 2014; Martin et al., 2021). By contrast, the NH mid-latitude zonal wind shows an easterly bias at Day 1, and it is replaced by a westerly wind bias at Day 3 and thereafter (Fig. 2(d)). This characteristic time evolution of the zonal wind error implies that the sources of the initial easterly bias and the subsequent westerly bias are different.

As expected, timeseries of the GLN forecasts are apparently almost identical with those of the MetUM analysis. This illustrates that the model variables are properly relaxed towards the forcing data, the MetUM analysis. However, there are systematic errors with a small magnitude especially in the NH mid-latitude stratospheric zonal wind (Fig. 2(a, d)). The small initial easterly wind bias in GLN continues uniformly throughout the forecasts up to Day 15, even though the zonal wind itself is nudged. Zonal wind errors in GLN vary among cases but does not depend on the forecast lead time if their valid time is the same (see below in Fig. 12(a, b)). In the nudging framework, how strongly model variables are constrained depends on the relaxation parameter (i.e., $1/\tau$ in Eq. (1)). If the error grows rapidly at a very short timescale, the nudging is not able to relax sufficiently towards the forcing data with a moderate relaxation parameter. Although this consistent error in GLN seemingly looks strange, this is reasonable because the forcing data, the MetUM analysis, which model prognostic variables relaxed




Figure 3. Spatial distributions of zonal wind errors [m s^{-1}] at 50 hPa at a forecast lead time of (a, b) Day 1 and Day 5 (from top to bottom, c, d) Day 5 averaged over the 90 cases from 1 Dec. 2018 to 28 Feb. 2019 in (a, c) CNTL, and (b, d) GLN. Contour indicates the forecasts, and colour indicates mean errors against the MetUM analysis.

towards, does not depend on forecast lead time. As a result, GLN shows the same forecast fields and therefore the same error fields against analysis at specific valid time regardless of the forecast lead time.

Spatial distributions of the model biases could also reveal error characteristics. Global spatial maps of the zonal wind bias at 50 hPa in CNTL and GLN at Day 1 and Day 5 are shown in Fig. 3. The easterly wind bias over the Himalayas, the Tibetan Plateau, and its downstream region are the main contributors to the zonally averaged wind error in the lower stratosphere (Fig. 3(a)). It becomes less obvious relative to other regions as forecast lead time gets longer (Fig. 3(c)). Whilst the nudging of zonal wind as well as meridional wind and temperature in GLN makes the error against the analysis much smaller in the spatial-distribution perspective, the easterly wind bias remains in GLN especially over the Himalayas and the Tibetan Plateau (Fig. 3(b, d)). This local easterly wind bias is probably associated with the modelled orography and/or the orographic gravity wave drag parametrization.

4 Zonal-mean zonal momentum and thermal budget

~~Atmospheric~~The atmospheric zonal momentum and thermal budget analysis specifies to what extent the individual components in the budget equation contribute to the total tendency of the corresponding variable. In this section, we will establish a correspondence between these budget components and the model tendencies due to dynamics and physics. To identify




Figure 4. Latitude-height cross sections of zonal-mean (a, d) total tendency of zonal wind, and budget contributions from (b, e) the resolved processes (i.e., sum of the mean stationary flow advection term, the stationary eddy momentum flux term, the transient eddy momentum flux term, and the Coriolis term shown in Fig. 5) and (c, f) the residual term of the zonal-mean zonal momentum budget equation [$\text{m s}^{-1} \text{ day}^{-1}$] written in Eq. (2) from initial conditions to T+120h averaged over the 90 cases (aa-c) in CNTL, and (bd-f) their differences between CNTL and GLN. Contour in (a), for reference, indicates the zonal wind forecast at T+120h which is identical with that in the middle panel of Fig. 1(ac). Note that the colour scales in (aa-c) and (bd-f) are different.

which component has a dominant contribution to the forecast error growth, budget analysis in GLN is used for validating the equivalent budgets in CNTL. Differences between CNTL and GLN in the rest of the paper denoted by "CNTL minus GLN" will be interpreted as errors in CNTL against GLN a quasi-error in budget components of CNTL against GLN is evaluated.

4.1 Zonal momentum budgets

As written in Eq. (2), The atmospheric zonal-mean zonal momentum equation indicates that the total tendency of zonal-mean zonal wind is separated into contributions from the mean stationary flow advection, the stationary and transient eddy momentum flux convergence, the Coriolis forcing, and the residual term. A zonal-mean zonal momentum budget at Day 0–5 is selected here to examine the evolving errors. Figure 4 shows vertical distributions of the mean total tendency from initial conditions to T+120h and contributions from the resolved processes and the residual term averaged over the 90 cases, and their differences between CNTL and quasi-error of CNTL against GLN. The total forecast tendency in CNTL averaged over all the cases shows a considerably similar pattern to the mean error (the middle panel of Fig. 1(ac)). This is because the analysis




Figure 5. Latitude-height cross sections of ~~contributions to a breakdown of~~ the resolved processes (shown in Fig. 4(b, e) ~~from~~) to the individual terms of the zonal-mean zonal momentum budget equation [$\text{m s}^{-1} \text{ day}^{-1}$] written in Eq. (2) (i.e., from left to right, (a, e) the mean stationary flow advection term, (b, f) the stationary eddy momentum flux term, (c, g) the transient eddy momentum flux term, and (d, h) the Coriolis term) from initial conditions to T+120h averaged over the 90 cases (aa–d) in CNTL, and (be–h) their differences between CNTL and GLN. Note that the colour scales in (aa–d) and (be–h) are different.

tendency averaged over a long experimental period becomes vanishingly small compared with the forecast tendency so that the error is approximately proportional to the total forecast tendency (Rodwell and Palmer, 2007). Therefore, the difference in the total tendency between CNTL and GLN, shown in Fig. 4(bd), is almost the same as the total tendency in CNTL, shown in Fig. 4(a), whilst the color scales are different. Figure 5 shows the individual terms that constitute the resolved processes shown in Fig. 4(b, e). The mean stationary flow advection in the troposphere contributes to a decrease in zonal-mean zonal wind in the tropics with the southerly wind (i.e., $-\frac{[\bar{v}]}{a \cos \phi} \frac{\partial [\bar{u}] \cos \phi}{\partial \phi} < 0$ because $[\bar{v}] > 0$ and $\frac{\partial [\bar{u}] \cos \phi}{\partial \phi} > 0$), a decrease in zonal-mean zonal wind in the mid-latitude with the northerly wind (i.e., $-\frac{[\bar{v}]}{a \cos \phi} \frac{\partial [\bar{u}] \cos \phi}{\partial \phi} < 0$ because $[\bar{v}] < 0$ and $\frac{\partial [\bar{u}] \cos \phi}{\partial \phi} < 0$), and an increase in zonal-mean zonal wind in the subtropics with the descents (i.e., $-\frac{[\bar{\omega}]}{a \cos \phi} \frac{\partial [\bar{u}] \cos \phi}{\partial \phi} > 0$ because $[\bar{\omega}] > 0$ and $\frac{\partial [\bar{u}] \cos \phi}{\partial \phi} < 0$) (Fig. 5(a)). This illustrates meridional and vertical momentum transports by the upper and descending branch of the Hadley and Ferrel cells. The momentum budget from the middle troposphere to the stratosphere is characterized by the balance among the mean stationary flow advection, the convergence of the eddy momentum fluxes, and the Coriolis forcing. ~~Whilst the transient eddy is a deviation from temporal mean over the medium-range timescale of 5 days, the transient eddy momentum flux plays a substantial role in the budget in the middle-to-upper troposphere.~~ (Fig. 5(a–d)).




Figure 6. Latitude-height cross sections of zonal-mean (a, e) total tendency of zonal wind, and their breakdown into model tendencies of model (b, f) dynamics, (c, g) physics, and (d, h) nudging [$\text{m s}^{-1} \text{ day}^{-1}$] from initial conditions to T+120h averaged over the 90 cases (aa–d) in CNTL, and (be–h) their differences between CNTL and GLN. Zonal wind tendency due to nudging in CNTL is completely null. Note that the colour scales in (aa–d) and (be–h) are different.




Figure 7. Latitude-height cross sections of contributions to a breakdown of the model physics zonal-wind tendency (shown in Fig. 6(ac)) from to the individual parametrizations, (a) convection, (b) boundary layer, and (c) gravity wave drag [$\text{m s}^{-1} \text{ day}^{-1}$] in CNTL from initial conditions to T+120h averaged over the 90 cases.

380 We now establish a ~~correspondence~~ qualitative similarity between the budget terms and the model tendencies in CNTL. Figure 6 exhibits tendencies due to the model dynamics, physics, and the nudging forcing, ~~calculated directly from model simulations~~. Model tendencies due to individual parametrizations constituting model physics (i.e., convection, boundary layer, and gravity wave drag) are shown in Fig. 7. A comparison of Fig. 4(aa-c) with Fig. 6(aa-c) indicates that the resolved processes (Fig. 4(b)) and the residual term (Fig. 4(c)) in the momentum budget correspond to the dynamics tendency (Fig. 6(b)) and the physics parametrizations tendency (Fig. 6(c)) in the model, respectively. Although the residual term is relatively small from the upper troposphere to the stratosphere, there is a recognizable westward forcing around the NH mid-latitude lower stratosphere. The residual term (Fig. 4(ac)) near the surface balances with the Coriolis forcing (Fig. 5(ad)). A relatively large deceleration by the residual term is shown in the upper stratosphere to the mesosphere (i.e., a deceleration of the westerly wind in winter hemisphere and a deceleration of the easterly wind in summer hemisphere). ~~These forcings~~ Most of the residual term can
390 ~~mostly~~ be accounted for by unresolved momentum transports which are being parametrized by the gravity wave drag scheme, the boundary layer scheme, and the convection scheme shown in Fig. 7.

The ~~momentum budget in GLN is considered as a best possible estimate of the truth and used as the reference to validate the individual budget components. However, it should be interpreted cautiously in several respects. Firstly, nudged variables in nudging experiments still have errors against the MetUM analysis as shown by Fig. 2(d) and Fig. 3(b). In addition, the vertical velocity, required by the budget analysis but not directly nudged, is driven by the horizontal wind and temperature nudging forcing. Therefore, there should be a non-negligible error in the individual budget components in GLN, making it difficult to precisely evaluate the residual term as a consequence. Secondly, since the 6-hourly MetUM analyses used as a forcing data of the nudging are linearly interpolated into each model time step as part of the nudged simulation methodology, the temporal interpolation leads to an underestimation of the intra-6-hour variability, and therefore results in underestimated transient eddy components and consequently overestimated residual terms. These possible caveats are discussed in Sect. 6.~~
395 ~~components and consequently overestimated residual terms. These possible caveats are discussed in Sect. 6.~~

The ~~error in the total tendency against GLN~~ quasi-error in the total tendency is dominated by the large compensation between Coriolis forcing (Fig. 5(bh)) and the residual term (Fig. 4(bf)), particularly in the troposphere and the stratosphere. Whilst the Coriolis term in GLN may have a non-negligible error as described ~~above~~ in Sect. 2.6, a verification of zonal-mean meridional wind in CNTL and GLN against the MetUM analysis demonstrates that the error of the Coriolis term in GLN is smaller than
405 that in CNTL (not shown). This suggests that the difference of the Coriolis forcing between CNTL and GLN ~~can be a proxy of~~ could approximate the error in Coriolis term in CNTL against truth. The westerly wind bias in the midlatitudes over the upper troposphere to the middle stratosphere is attributable to the ~~error~~ quasi-error in Coriolis forcing around the tropopause and the residual term in the middle stratosphere, which is examined further in Sect. 5.

The ~~interpretation of differences in the residual term among experiments is more complex. It is attributed, in theory, to differences in the other resolved terms, because the residual term in each experiment is determined by the sum of the other terms to close the budget. From another perspective, the residual term corresponds to the sum of the physics parametrization tendencies and the nudging forcing as shown by the comparison between Fig. 4 and Fig. 6. The error in the residual term against GLN suggests deficiencies in the model physics parametrizations. As the nudging increment in this study is~~ increments are comparable to the forecast errors at ~~a least those growing over the~~ 6-hour timescale, a ~~resemblance between the error~~




Figure 8. Latitude-height cross sections of zonal-mean (a, d) total tendency of temperature, and budget contributions from (b, e) the resolved processes (i.e., sum of the mean stationary flow term, the stationary eddy heat flux term, and the transient eddy heat flux term shown in Fig. 9) and (c, f) the residual term of the zonal-mean thermal budget equation [K day^{-1}] written in Eq. (3) from initial conditions to T+120h averaged over the 90 cases (aa-c) in CNTL, and (bd-f) their differences between CNTL and GLN. Contour in (a), for reference, indicates the temperature forecast at T+120h which is identical with that in the middle panel of Fig. 1(bd). Note that the colour scales in (aa-c) and (bd-f) are different.

415 similarity between the quasi-error in the residual term (Fig. 4(bf)) and the reversed sign of the nudging forcing in GLN (Fig. 6(bh)) is consistent with the finding of the previous studies that fastest growing errors at short timescales are likely associated with the model physics (Martin et al., 2010; Ma et al., 2014; Martin et al., 2021). On the other hand, the error in the Little difference in the physics parameterization tendencies between CNTL and GLN, shown in Fig. 6(g), demonstrates that the tendency diagnostics using GLN are less useful to estimate correct unresolved physics tendency to be parameterized in the

420 model and indicates that the physics schemes themselves have substantial deficiencies. The quasi-error in the Coriolis forcing may be a response to the incorrect parametrized mechanical or thermal forcing.

4.2 Thermal budgets

In the same manner as the zonal momentum budget analysis, the total tendencies of zonal-mean temperature can be diagnosed using the thermal budget analysis as written in Eq. (3). Figures 8 and 9 exhibit latitude-height distributions of the mean total

425 forecast tendency and its thermal budget from initial conditions to Day 5, and their differences between CNTL and quasi-error




Figure 9. Latitude-height cross sections of contributions to a breakdown of the resolved processes (shown in Fig. 8**(b, e)**from) to the individual terms of the zonal-mean thermal budget equation [K day^{-1}] written in Eq. (3) (i.e., from left to right, (a, d) the mean stationary flow term, (b, e) the stationary eddy heat flux term, and (c, f) the transient eddy heat flux term) from initial conditions to T+120h averaged over the 90 cases (a-c) in CNTL, and (b-d-f) their differences between CNTL and GLN. Note that the colour scales in (a-c) and (b-d-f) are different.

against GLN. In CNTL shown in Fig. 9(a-c), the mean stationary flow component has a vertically consistent structure in the troposphere -(Fig. 9(a)). Separating it into the meridional advection, the vertical advection, and the adiabatic heating with mean vertical motions reveals that a contribution of the meridional advection is considerably small (not shown). In general, the vertical advection (the second term of the mean stationary flow component in Eq. (3)) and the adiabatic cooling (warming) with ascents (descents) (the third term of the mean stationary flow component in Eq. (3)) compensate throughout the troposphere because the vertical pressure derivative of temperature is positive. As a result, the adiabatic heating with vertical motions dominates over the vertical temperature advection. The mean stationary flow component illustrates the Hadley circulation leading to the cooling with ascents in the SH tropics and the warming with descents in the NH tropics. The cooling by the residual term in the middle to the upper troposphere in the NH high latitude (Fig. 8(a-c)) counteracts a warming by the transient and stationary eddy heat flux convergence (Fig. 9(a,b, c)). In the stratospheric winter hemisphere, the mean stationary flow component, dominated by the adiabatic heating with vertical motions, and the eddy heat flux component offset each other, and the cooling by the residual term balances with the warming due to the resolved dynamical processes. By contrast, in the stratospheric summer hemisphere where the eddy components are quiescent -(Fig. 9(b, c)), the warming by the residual term




Figure 10. Latitude-height cross sections of zonal-mean [\(a, e\)](#) total tendency of temperature, and [their breakdown into model](#) tendencies of [model](#) [\(b, f\)](#) dynamics, [\(c, g\)](#) physics, and [\(d, h\)](#) nudging [K day^{-1}] from initial conditions to T+120h averaged over the 90 cases [\(aa-d\)](#) in CNTL, and [\(be-h\)](#) their differences between CNTL and GLN. Temperature tendency due to nudging in CNTL is completely null. Note that the colour scales in [\(aa-d\)](#) and [\(be-h\)](#) are different.

counterbalances the cooling by the mean flow which constitutes a part of the stratospheric meridional circulations [-\(Figs. 8\(c\) and 9\(a\)\)](#).

As with the momentum budget, the residual term in each experiment corresponds to the parametrized forcing shown in Fig. 10(c). Figure 11 presents contributions to the model physics tendency from the individual parametrizations and shows that the cooling and the warming in the stratosphere are attributable to dominant longwave and shortwave radiative processes in winter and summer hemisphere, respectively [-\(Fig. 11\(a, b\)\)](#). The vertical and latitudinal distribution of the residual term is more complex in the troposphere. Latent heat release and subgrid heat transport as well as radiative processes play a substantial role, and these processes interact and balance each other. The warming in the tropics in summer hemisphere appears to correspond with latent heat release and subgrid heat transport represented by model physics (e.g., convection, cloud, microphysics, vertical mixing, etc.). The cooling in the upper troposphere is associated with a dominant longwave radiation process partly compensated by a shortwave radiative heating.

Errors The quasi-errors in the individual budget terms against GLN, presented in Figs. 8(bd-f) and 9(bd-f), show that there are large discrepancies in the mean stationary flow component [\(Fig. 9\(d\)\)](#) and the residual term [\(Fig. 8\(f\)\)](#) relative to GLN in the troposphere to the middle stratosphere. A set of negative and positive errors around the equator throughout the troposphere




Figure 11. Latitude-height cross sections of contributions to a breakdown of the model physics temperature tendency (shown in Fig. 10(ac)) from to the individual parametrizations, (a) shortwave radiation, (b) longwave radiation, (c) convection, (d) microphysics, (e) cloud, (f) boundary layer, and (g) gravity wave drag [K day⁻¹] in CNTL from initial conditions to T+120h averaged over the 90 cases.

suggests a meridional displacement of the large-scale ascents in summer hemisphere. ~~Unlike the momentum budget, the error~~ The quasi-error in the residual term shows a qualitatively similar distribution with that in the total tendency especially from the upper troposphere to the middle stratosphere (Fig. 8(b))-d, f), which is not the case in the zonal momentum budget (Fig. 4(d, f)). This suggests that the unresolved thermodynamic processes ~~cause the total forecast tendency error over the tropics to subtropics.~~ are the main sources of the temperature error from the upper troposphere to the middle stratosphere, and an error generated by deficient unresolved processes at a very short-range timescale is not completely compensated for by model dynamics responses even at Day 5. The similarity between the nudging forcing with the reversed sign (Fig. 10(h)) and the quasi-error in the residual term ((Fig. 8(f)) suggests that initial growing error at the very short range timescale, which the nudging would reduce in GLN, is associated with model physics parameterizations. There are likely to be systematic deficiencies in the relevant physics parametrizations (e.g., a deficient cooling or an excessive warming around the tropical tropopause, a deficient warming or an excessive cooling in the middle stratosphere, etc.).

Another interesting characteristic in CNTL is an underestimated warming by the stationary eddy component (mainly the meridional convergence of the stationary eddy heat flux) in the NH high latitude from the upper troposphere to the lower stratosphere compared with GLN (Fig. 9(b, e)). This indicates that the cold bias in CNTL arises from the resolved dynamical processes as well as from the residual term. The warming due to the eddy heat flux convergence in the lower stratosphere

appears to be substantially affected by the sudden stratospheric warming (SSW) event that has occurred in the end of December 2018 (Rao et al., 2019). This is revealed by the timeseries of the thermal budget at a certain pressure level in the stratosphere
470 (for example in Fig. 14).

5 NH mid-latitude wind bias

Given the study of the large-scale structure of the model biases and the momentum and thermal budgets, we investigate specific model biases, the NH mid-latitude wind bias and the relevant temperature biases, in detail. To identify sources of the NH mid-latitude stratospheric zonal wind errors and the relevant temperature errors shown in Fig. 2, we analyse the zonal momentum
475 and thermal budgets, as well as the model dynamics, physics and nudging tendencies over the domains A, B, and C (see Sect. 3). Finally, temperature-only nudging experiments provide experimental insights on impacts of thermal forcing on the NH mid-latitude wind bias.

5.1 Zonal momentum and thermal budgets

Zonal momentum budgets and model tendencies averaged over the NH midlatitude are examined. Figure 12 shows timeseries
480 of the zonal momentum budget and the model tendencies due to the dynamics, physics, and nudging forcing in the individual forecasts up to Day 1 (from initial conditions to T+24h) and Day 5 (from initial conditions to T+120h) averaged over the NH mid-latitude band (30°N–60°N) at 50 hPa (see the black line A in Fig. 1(e)). The zonal momentum budget and physics parametrization tendencies up to Day 1 as well as Day 5 are shown in Fig. 12 to clarify differences in contributions to the total tendency between Day 1 with initial easterly bias and Day 5 with subsequent westerly bias. Note that the total tendencies ~~in~~
485 ~~the physics parametrizations (depicted in Fig. 12(g–j))~~ are not identical with those depicted in the zonal momentum budget (Fig. 12(c–f)) because the ~~physics parametrization tendencies are evaluated model tendencies~~ at the specific model level closest to the pressure level of 50 hPa are selected whilst the momentum budget is analysed in pressure coordinates. In 1-day CNTL forecasts, the stationary eddy momentum flux and the Coriolis forcing contribute to an acceleration of the westerly wind, and the residual term decelerates and partly offsets the acceleration (Fig. 12(c)). Decelerations by the residual term are quantitatively
490 consistent with those by the physics parametrization tendency dominated by the gravity wave drag (Fig. 12(c, g)). Unlike the consistent deceleration by the residual term in CNTL (Fig. 12(c)), that in GLN shows a temporal variation with accelerations and decelerations day by day (Fig. 12(c, e)). In almost all the cases, whilst the Coriolis term provides excessive westerly tendencies relative to GLN, the residual term has excessive easterly tendencies against GLN (Fig. 12(e)). The other resolved processes including the mean stationary flow advection and the eddy components show smaller ~~errors~~ quasi-errors against
495 GLN. The error compensation between the Coriolis term and the residual term results in a net westerly ~~error~~ quasi-error in CNTL against GLN (Fig. 12(e)), even though CNTL has the easterly wind bias according to the verification against the MetUM analysis (Fig. 12(a)). This suggests that the validation of the budget in CNTL against GLN could mislead the total tendency error in the case that GLN has relatively large errors against analysis in comparison to CNTL, for example at a short timescale. The ~~difference~~ quasi-error in the residual term ~~between CNTL and GLN is attributed to~~ can be accounted for by the nudging




Figure 12. Timeseries area-averaged over the NH mid-latitude (30°N–60°N) at 50 hPa depicted by the line A in Fig. 1(c) of (a, b) zonal wind errors against the MetUM analyses [m s^{-1}], (c, d) zonal momentum budget in CNTL [$\text{m s}^{-1} \text{ day}^{-1}$], (e, f) differences in the zonal momentum budget (CNTL minus GLN) [$\text{m s}^{-1} \text{ day}^{-1}$], (g, h) individual model physics, dynamics, and nudging tendencies in CNTL [$\text{m s}^{-1} \text{ day}^{-1}$], and (i, j) differences in the individual model physics, dynamics, and nudging tendencies (CNTL minus GLN) [$\text{m s}^{-1} \text{ day}^{-1}$]. Left column shows Day 1 forecasts: (a) the errors at a T+24h forecast lead time, and (c, e, g, i) temporal changes from initial conditions to T+24h. Right column shows Day 5 forecasts: (b) the errors at a T+120h forecast lead time, and (d, f, h, j) temporal changes from initial conditions to T+120h. Note that the nudging tendency in (g, h) is completely null, and physics parametrization tendencies of the convection scheme and the boundary layer scheme are omitted in (g–j) because they are negligibly small in this case.

500 forcing (Fig. 12(e, i)), which accelerates the NH mid-latitude lower stratosphere wind in GLN (Fig. 12(i)). It is probable that direct mechanical forcing predominantly from the parametrized gravity wave drag is a source of the easterly wind bias at the very initial stage of the forecasts. On the other hand, the westerly wind errors growing thereafter could be caused by dynamical responses possibly to diabatic heating and temperature biases. The easterly forcing due to the gravity wave drag becomes slightly stronger in GLN (Fig. 12(i)), implying that the scheme ~~in GLN~~ responds to the westerly nudging forcing. Another
505 interesting point is that there is little difference in the zonal momentum budget and in the model tendencies between up to Day 1 and up to Day 5. This implies that budget analysis at a short timescale is also beneficial to understand contributions to the total tendency at a medium-range timescale except for the partition among the advective terms even if characteristics of the error are different among timescales. ~~The nudging forcing effectively corrects the model error against analysis at shorter timescales (depending on the relaxation timescale and nudging frequency) rather than at longer timescales. It is for this reason that, even~~
510 ~~though the model~~ Although the model zonal wind error is positive at lead time Day 5 (Fig. 12(b)), the nudging tendency at that lead time is still positive in GLN (Fig. 12(j)). This is because the nudging forcing effectively corrects the model error against analysis at shorter timescales (depending on the relaxation timescale and nudging frequency) rather than at longer timescales.

To investigate the relevant temperature bias, thermal budgets and model temperature tendencies averaged over the NH tropics around the tropopause are examined. Figure 13 presents timeseries of the thermal budget and the temperature tendencies due
515 to dynamics, physics, and nudging averaged over the NH tropics to subtropics band (0° – 30° N) at 70 hPa (see the black line B in Fig. 1(f)). Figure 13(a, b) reveals that CNTL has positive temperature errors at a 1-day forecast lead time in almost all the cases and at a 5-day forecast lead time in all the cases, indicating a robustness of the warm bias around the tropical tropopause. The dominant warming by the residual term leads to positive total tendencies, although the resolved processes contribute to a cooling and partly offset the residual warming (Fig. 13(c, d)). ~~The variability of the total tendency can mainly be accounted for~~
520 ~~by that of the mean stationary flow component. Errors~~ The quasi-errors in the individual terms against GLN show a substantial agreement between the total tendencies and the residual term (Fig. 13(e, f)). This agreement indicates that the residual term plays a key role in the error of the total tendency at both Day 1 and Day 5. The residual term corresponds to the diabatic warming parametrized in the longwave and shortwave radiation schemes (Fig. 13(g, h)), and the other physical processes have negligible impacts in this case. A comparison between the residual term (Fig. 13(e, f)) and the tendency due to nudging (Fig.
525 13(i, j)) demonstrates that the direct nudging forcing accounts for the ~~difference~~ quasi-error in the residual term ~~between CNTL and GLN~~. It is likely that deficiencies in the radiation schemes or model variables ingested into the schemes cause the tropical tropopause warm bias, although further ~~investigations are~~ investigation is needed.

The temperature bias over the NH high latitude around the tropopause, which is another relevant temperature bias, is also addressed. Figure 14 shows the thermal budget and the model tendencies averaged over the NH high-latitude band (60° N–
530 90° N) at 200 hPa (see the black line C in Fig. 1(f)). Compared with the NH tropics region shown in Fig. 13, the stationary eddy component plays an important role in the budget especially from the beginning of the December to the mid-January (Fig. 14(c, d)). This might respond to a dynamical situation prior to and during the SSW. The warming by the stationary eddy heat flux component is underestimated in CNTL compared with GLN throughout the experimental period (Fig. 14(e, f)). The magnitude of the temperature nudging forcing in this domain (Fig. 14(i, j)) is smaller than that in the previous domain (Fig. 13(i, j)). In




Figure 13. Same as Fig. 12, but area-averaged over the NH tropics to subtropics (0° – 30° N) at 70 hPa depicted by the line B in Fig. 1(f) of (a, b) temperature errors against the MetUM analyses [K], (c, d) thermal budget in CNTL [K day^{-1}], (e, f) differences in the thermal budget (CNTL minus GLN) [K day^{-1}], (g, h) individual model physics, dynamics, and nudging tendencies in CNTL [K day^{-1}], and (i, j) differences in the individual model physics, dynamics, and nudging tendencies (CNTL minus GLN) [K day^{-1}]. Note that physics parametrization tendencies of the convection scheme, the cloud scheme, the microphysics scheme, and the gravity wave drag scheme are omitted in (g–j) because they are negligibly small in this case.

535 contrast to the NH tropics where the residual term is consistently overestimated in CNTL relative to GLN (Fig. 13(e, f)), the residual term shows the positive and negative ~~error~~-quasi-error in the NH high latitude (Fig. 14(e, f)). These suggest that the




Figure 14. Same as Fig. 13, but area-averaged over the NH high-latitude region (60°N – 90°N) at 200 hPa depicted by the line C in Fig. 1(f). Note that physics parametrization tendencies of the convection scheme, the cloud scheme, the microphysics scheme, and the gravity wave drag scheme are omitted in (g–j) because they are negligibly small in this case.

NH high-latitude temperature bias arises from dynamical responses as well as model physics. The initial temperature error in this domain (Fig. 14(a)) is also less robust than in the NH tropics (Fig. 13(a)).

Given the diagnostics of the error evolution and the momentum and thermal budget analysis, a possible mechanism of the forecast error growth is summarized. The main source of the NH mid-latitude lower-stratospheric westerly wind bias in CNTL is probably the temperature biases around the tropopause especially in the tropics. The mechanical forcing itself has an




Figure 15. Same as Fig. 2(d), but for various nudging experiments summarized in Table 1. CNTL and GLN are identical with those in Fig. 2(d). Dotted lines with open circles and smaller caps of error bars indicate nudging sensitivity experiments in which variables are relaxed only around the tropopause.

erroneous excessive easterly tendency (and/or a deficient westerly forcing) at least at N320 horizontal resolution which causes the NH mid-latitude easterly wind bias at an initial stage of the forecasts. The overly strong temperature gradient causes the subsequently developing westerly wind bias through the excessive eastward Coriolis forcing. There are compensating errors between an excessive westerly tendency caused by thermal forcing originating probably from the radiative processes and an excessive easterly tendency predominantly from the gravity wave drag. Possible deficiencies in the parametrizations in the MetUM need to be investigated further and are discussed in Sect. 6.

5.2 Nudging sensitivity experiments

A partial nudging technique is adopted to attempt to disentangle compensating errors and understand remote impacts of temperature nudging forcing particularly on the NH mid-latitude lower-stratospheric wind bias, which corroborates findings of the possible error mechanism summarized above. The thermal forcing due to the temperature nudging can, in some sense, be interpreted as an additional artificial diabatic heating required to correct model biases. Figure 15 shows time evolutions of NH mid-latitude zonal wind error at 50 hPa in the various nudging experiments outlined by Table 1. Many interesting experimental results are found. The westerly wind bias in CNTL is eliminated by the global temperature nudging, and conversely an easterly wind bias emerges and evolves gradually and monotonically up to Day 15. The temperature error around the tropical

tropopause is probably one of the main drivers of the NH mid-latitude wind bias in the lower stratosphere. Note that, however, these experimental results do not determine genuine sources of the wind errors because errors that the nudging directly mitigates are contaminated at a longer forecast lead time by those which are remotely generated by dynamical responses. Impacts of the temperature nudging are significantly different between the whole column nudging and the tropopause nudging over the NH high latitude. There is a difference in the timescales on which temperature nudging over the tropics and at the high latitudes act. The temperature nudging ~~at-over~~ the tropics reduces substantially the wind bias at shorter timescales, whereas the impact of temperature nudging ~~at-over~~ the high latitudes is relatively substantial at Day 9 and afterwards. This suggests a difference in the timescales of the temperature error growth in CNTL in these regions. The temperature error around the tropical tropopause evolves more quickly than that in the NH high-latitude tropopause (cf. Figs. 13(a) and 14(a)), and therefore the impact of nudging also appears more rapidly.

Zonal momentum budget analysis could reveal the contribution from the individual terms and help understand the differences in impact timescales of the temperature nudging over the tropics and the high latitudes. Changes in the momentum budget of each experiment relative to that of CNTL are presented for Day 5 and Day 10 in Fig. 16 and Fig. 17, respectively. Day 5 represents a case on which the impact on the wind error by tropical-tropopause temperature nudging is dominant, while Day 10 is a case on which the effects by high-latitude temperature nudging are more significant. Impacts outside the nudged domain shown by green dashed boxes are interpreted as a remote response to errors in the target nudged domain. Momentum budgets up to Day 5 show that changes of the total tendencies from CNTL in the NH mid-latitude lower stratosphere are dominated by changes of the Coriolis forcing in all the temperature nudging experiments shown in Fig. 16. Temperature nudging alters the meridional circulation through geopotential gradients, which is clearly demonstrated by a meridional momentum budget analysis (~~not shown~~), and consequently the Coriolis forcing in the zonal momentum budgets ~~-(Fig. 16(a5-d5))~~. The stationary eddy momentum flux convergence also contributes to the deceleration on the poleward side of where the Coriolis forcing decelerates the westerly wind ~~-(Fig. 16(a3-d3))~~. In NHTrpT and NHTrpTrpT, the residual term also has a substantial contribution to the deceleration in the NH midlatitude (Fig. 16(~~a~~, ~~ba6~~, ~~b6~~)). ~~The Note that temperature is relaxed towards the analysis, but zonal wind is not relaxed at all in the temperature nudging experiments. Therefore, the zonal wind tendencies due to nudging are completely zero, and the~~ difference in the residual term of ~~NHTrpT and NHTrpTrpT cannot be fully accounted for by changes in the physics parametrization tendencies (not shown). Hence, zonal momentum budgets is attributed to the effects other than the nudging. Hence, a different responses of parameterizations to temperature nudged grid-box mean fields or an~~ imbalance caused by the temperature nudging may lead to the deceleration in the residual term. In the momentum budgets up to Day 10 in the NH tropics temperature nudging experiments shown in Fig. 17(a, b), the eddy momentum flux convergence and the Coriolis forcing weakens the total deceleration in the lower and higher latitude domain, respectively, in the NH lower stratosphere ~~-(Fig. 17(a3-a5, b3-b5))~~.

There is a difference in the impacts of nudging between the NH tropics nudging and the NH high-latitude nudging with respect to a latitudinal spreading of the impacts and their vertical distribution. The temperature nudging over the NH tropics affects not only the tropics but also the mid- to high-latitude regions, and its impact gradually extends from the low-latitude region to the high-latitudes in both hemispheres. On the other hand, the impact of temperature nudging over the NH high latitude




Figure 16. Latitude-height cross sections of differences in zonal-mean (a1–d1) total tendency of zonal wind, and budget contributions from the the individual terms of the zonal-mean zonal momentum budget equation (i.e., from left to right, (a2–d2) the mean stationary flow advection term, (a3–d3) the stationary eddy momentum flux term, (a4–d4) the transient eddy momentum flux term, (a5–d5) the Coriolis term, and (a6–d6) the residual term) [$\text{m s}^{-1} \text{ day}^{-1}$] written in Eq. (2) from initial conditions to T+120h averaged over the 90 cases in (a) NHTrpT minus CNTL, (b) NHTrpTrpT minus CNTL, (c) NHHLT minus CNTL, and (d) NHHLTrpT minus CNTL. Green dashed boxes indicate the temperature-nudging domain including interfacing transition zones and layers. The black horizontal line A, the same as that in Fig. 1(e), are inserted for reference. Note that the colour scale is different from that in Fig. 4.

is confined to the extratropics. The NH tropics nudging shows vertically tripolar structure in the total tendency around the tropics. The total deceleration in the lower stratosphere is confined to below the middle stratosphere and overlaid by acceleration in the upper stratosphere. On the other hand, the NH high-latitude nudging decelerates zonal wind consistently from the upper troposphere to the stratosphere. This might also be associated with the different vertical structure of the temperature error
595 between the tropics and the high latitudes shown in Fig. 1(b, d, f).




Figure 17. Same as Fig. 16, but for momentum budgets from initial conditions to T+240h.

6 Conclusions

To understand mechanisms of model systematic errors in the global MetUM and identify their sources, we have applied the momentum and thermal budget analysis and the nudging technique to NWP simulations in boreal winter. The model systematic errors addressed in this paper include a westerly wind bias in the NH midlatitude from the upper troposphere to the stratosphere, a warm temperature bias around the tropical tropopause, and a cold temperature bias around the tropopause over the poles. The warm and cold temperature errors near the tropopause develop quickly and systematically in almost all the cases, indicating their robustness. On the other hand, the NH mid-latitude wind in the lower stratosphere has an initial easterly bias especially over the Himalayas and the Tibetan Plateau and a subsequent westerly bias in the zonal-mean zonal wind emerging at Day 3 and evolving thereafter. The momentum and thermal budget analysis is a practical diagnostic method that specifies to what extent the individual components in the budget equation contribute to the total tendency of wind or temperature. The residual term corresponds to the effects of unresolved processes parametrized in GCMs on the total tendency. Simulations in which global horizontal wind and temperature are relaxed towards the MetUM analysis can be considered as a proxy of the analysis with some caveats, and their momentum and thermal budgets are used as the reference budgets to ~~validate those in CNTL.~~

~~The residual term in GLN includes the artificial nudging forcing which counteracts the model errors.~~ evaluate "quasi-error"
610 in the budget components. The zonal momentum budget analysis reveals that the westerly wind bias in the midlatitudes over the upper troposphere to the middle stratosphere is attributable to the dominant ~~error~~ quasi-error in the Coriolis forcing around the tropopause and the ~~error in~~ quasi-error in the residual term over the middle stratosphere. It is probable that an excessive mechanical forcing due to the orographically induced gravity wave drag is one of the sources of the easterly wind bias at the very initial stage of the forecasts. On the other hand, with respect to the thermal budget, the ~~error~~ quasi-error in the residual
615 term ~~against GLN~~ dominates that in the total temperature tendency from the upper troposphere to the middle stratosphere. Hence, the temperature bias is probably caused by systematic deficiencies in the unresolved processes represented by physics parametrizations. The NH high-latitude cold bias appears to be associated with the resolved dynamical processes as well as unresolved physical processes. This mechanism suggests error compensations in the zonal momentum budget.

The momentum and thermal budgets analysis to examine forecast and analysis time evolutions in NWP timescales is ex-
620 pected to be a powerful diagnostic tool not only for identifying sources of model systematic errors but also for understanding impacts of scientific changes or artificial forcing (e.g., nudging) among experiments. From zonal-mean perspective, one possible mechanism of the wind forecast error growth is that an overly strong temperature gradient with the warm (and cold) bias developing quickly near the tropopause leads to an acceleration of the southerly wind through an increased geopotential gradient, consequently resulting in an acceleration of the westerly wind by the Coriolis forcing. The fact that temperature
625 nudging over the NH tropical tropopause can significantly reduce the NH mid-latitude stratospheric westerly wind bias supports the idea that the tropical tropopause warm bias is one of the main sources of the mid-latitude wind bias. The excessive deceleration due to the gravity wave drag should partly compensate the ~~developing~~ westerly wind errors developing at the longer forecast lead time. Although this paper focuses on the zonal-mean budgets, it will also be profitable to analyse the longitudinally varying budgets (e.g., Yang et al., 2013) because mean total tendencies closely linked to model systematic errors
630 are zonally asymmetric and strongly localized. It is also important to explore a sensitivity of the budgets and model tendencies to the model horizontal resolution. Whilst the Eulerian framework is used in this study, the quasi-Lagrangian framework, such as the Transformed Eulerian Mean (TEM; Andrews and McIntyre, 1976) or the Mass-weighted Isentropic zonal Means (MIM; e.g., Iwasaki, 1989, 1992; Tanaka et al., 2004) equation, could also be useful for understanding large-scale atmospheric circulations.

635 Momentum and thermal budgets in the global nudging experiment used in this study should be interpreted cautiously because of errors even in the directly constrained variables and possible underestimations of the eddy components. These errors could be avoided or reduced using the budgets in analysis or reanalysis data instead of the global nudging experiment data as in main previous studies. However, it is required that the temporal, horizontal, and vertical resolutions of the analysis data are identical with those of the forecast data to compare their budgets fairly. It could be quite challenging to assess the budgets in
640 reanalyses and validate the budgets in models because of different resolutions or interpolations to get the resolutions aligned. One promising way ~~specially in for~~ operational meteorological centres to avoid this issue is applying the budget analysis with a different computation and/or definition of temporal mean to an objective analysis created by data assimilation system. If the budget components calculated using the fields at each model time step are well reproduced using 6-hourly instantaneous fields,

the analysis data potentially provide a better reference than the global nudged simulation. Defining temporal mean differently
645 as the average over the experimental period, a series of analysis and first guess, which is a 6-hour forecast from the previous
objective analysis, is available for diagnosing the budgets in the so-called analysis world and forecast world, although this is for
the purpose of identifying sources of initial model errors rather than understanding model error growth. This should also allow
us to reduce an initial spinup generated by discrepancies in the configuration between data assimilation systems and forecast
models.

650 We now discuss possible ~~physical~~ causes of the biases presented above [at the physical process level and potential further
investigations](#). The excessively strong deceleration in the NH midlatitude in winter is probably attributed to the orographic grav-
ity wave drag scheme. The vertical integrated momentum budget analysis (Milton and Wilson, 1996) or the angular momentum
budget analysis (Brown, 2004; van Niekerk et al., 2016) may help to identify if the gravity wave excitation process or the ver-
tical propagation and deposition process needs to be modified. Since there might be deficiencies in parametrizations employed
655 by the MetUM with specific horizontal resolutions, a horizontal resolution sensitivity also needs to be investigated. [The effects
of the gravity wave drag scheme on atmospheric circulations depend on the seasons and the hemispheres because of different
atmospheric stability and wind fields near the land surface and the asymmetric orography between the hemispheres. Therefore,
investigations of model systematic errors for the seasons \(e.g., boreal summer\) could be informative.](#) With respect to the warm
bias around the tropical tropopause, it is possible that there are deficiencies in the radiation schemes or relevant variables includ-
660 ing cloud fields. ~~Many studies have been dedicated to understanding mechanisms of the model systematic errors in the different
version of the MetUM, particularly for a temperature bias around the tropopause (Hardiman et al., 2015; Bland et al., 2021).
Hardiman et al. (2015) have executed several sensitivity experiments to examine the tropical tropopause warm bias that indicate
that microphysical and radiative processes influence the temperature. This is essentially consistent with the conclusion of the
previous studies (e.g., Hardiman et al., 2015).~~ Further investigations using the diagnostics framework in this study combined
665 with other diagnostic methods are a promising approach to establishing how to improve the model representations.

Code and data availability. Due to intellectual property right restrictions, we cannot provide either the source code or the documentation
papers for the Met Office Unified Model (MetUM). The MetUM is available for use under licence. For further information on how to
apply for a licence, see <https://www.metoffice.gov.uk/research/approach/collaboration/unified-model/partnership> (Met Office, 2024). JULES
is available under licence free of charge. For further information on how to gain permission to use JULES for research purposes, see
670 <https://jules.jchmr.org/> (JULES, 2024). Model data used in this study are archived at the Met Office and are available to research collaborators
upon request.

Author contributions. CM conducted the experiments, performed the analyses, and prepared the paper with contributions from all authors.
SFM initiated the study, designed the experiments and the analyses, and gave important guidance and feedback. JMR designed and conducted
the experiments, and gave important guidance and feedback.

675 *Competing interests.* The contact author has declared that none of the authors has any competing interests.

Acknowledgements. This study was carried out in a collaboration between the Met Office and Japan Meteorological Agency while C. Matsukawa was visiting the Met Office. J. M. Rodríguez was funded by the Met Office Climate Science for Service Partnership (CSSP) China project under the International Science Partnerships Fund (ISPF). Insightful and supportive comments from two anonymous reviewers are gratefully acknowledged.

- Andrews, D.: Middle Atmosphere Dynamics, Academic Press, 1987.
- Andrews, D. and McIntyre, M. E.: Planetary waves in horizontal and vertical shear: The generalized Eliassen-Palm relation and the mean zonal acceleration, *Journal of Atmospheric Sciences*, 33, 2031–2048, 1976.
- Bauer, P., Thorpe, A., and Brunet, G.: The quiet revolution of numerical weather prediction, *Nature*, 525, 47–55, <https://doi.org/10.1038/nature14956>, 2015.
- Best, M. J., Pryor, M., Clark, D., Rooney, G. G., Essery, R., Ménard, C., Edwards, J., Hendry, M., Porson, A., Gedney, N., et al.: The Joint UK Land Environment Simulator (JULES), model description–Part 1: energy and water fluxes, *Geoscientific Model Development*, 4, 677–699, 2011.
- Bland, J., Gray, S., Methven, J., and Forbes, R.: Characterising extratropical near-tropopause analysis humidity biases and their radiative effects on temperature forecasts, *Quarterly Journal of the Royal Meteorological Society*, 147, 3878–3898, <https://doi.org/10.1002/qj.4150>, 2021.
- Brown, A., Milton, S., Cullen, M., Golding, B., Mitchell, J., and Shelly, A.: Unified modeling and prediction of weather and climate: A 25-year journey, *Bulletin of the American Meteorological Society*, 93, 1865–1877, <https://doi.org/10.1175/BAMS-D-12-00018.1>, 2012.
- Brown, A. R.: Resolution dependence of orographic torques, *Quarterly Journal of the Royal Meteorological Society: A journal of the atmospheric sciences, applied meteorology and physical oceanography*, 130, 3029–3046, 2004.
- Chagnon, J., Gray, S., and Methven, J.: Diabatic processes modifying potential vorticity in a North Atlantic cyclone, *Quarterly Journal of the Royal Meteorological Society*, 139, 1270–1282, 2013.
- Clayton, A. M., Lorenc, A. C., and Barker, D. M.: Operational implementation of a hybrid ensemble/4D-Var global data assimilation system at the Met Office, *Quarterly Journal of the Royal Meteorological Society*, 139, 1445–1461, 2013.
- Cullen, M. J.: The unified forecast/climate model, *Meteorological magazine*, 122, 81–94, 1993.
- Donlon, C. J., Martin, M., Stark, J., Roberts-Jones, J., Fiedler, E., and Wimmer, W.: The operational sea surface temperature and sea ice analysis (OSTIA) system, *Remote Sensing of Environment*, 116, 140–158, 2012.
- Duykerke, P. G., de Roode, S. R., van Zanten, M. C., Calvo, J., Cuxart, J., Cheinet, S., Chlond, A., Grenier, H., Jonker, P. J., Köhler, M., et al.: Observations and numerical simulations of the diurnal cycle of the EUROCS stratocumulus case, *Quarterly Journal of the Royal Meteorological Society: A journal of the atmospheric sciences, applied meteorology and physical oceanography*, 130, 3269–3296, 2004.
- Edwards, J. and Slingo, A.: Studies with a flexible new radiation code. I: Choosing a configuration for a large-scale model, *Quarterly Journal of the Royal Meteorological Society*, 122, 689–719, 1996.
- Elvidge, A. D., Sandu, I., Wedi, N., Vosper, S. B., Zadra, A., Boussetta, S., Bouysse, F., van Niekerk, A., Tolstykh, M. A., and Ujiie, M.: Uncertainty in the representation of orography in weather and climate models and implications for parameterized drag, *Journal of Advances in Modeling Earth Systems*, 11, 2567–2585, 2019.
- Frasconi, A., Reynolds, C., Wedi, N., Bouallègue, Z. B., Caltabiano, A. C. V., Casati, B., Christophersen, J. A., Coelho, C. A., De Falco, C., Doyle, J. D., et al.: Systematic errors in weather and climate models: Challenges and opportunities in complex coupled modeling systems, *Bulletin of the American Meteorological Society*, 104, E1687–E1693, 2023.
- Gregory, D. and Rowntree, P.: A mass flux convection scheme with representation of cloud ensemble characteristics and stability-dependent closure, *Monthly Weather Review*, 118, 1483–1506, 1990.

- Hamilton, K.: Diagnostic study of the momentum balance in the Northern Hemisphere winter stratosphere, *Monthly weather review*, 111, 1434–1441, [https://doi.org/10.1175/1520-0493\(1983\)111%3C1434:DSOTMB%3E2.0.CO;2](https://doi.org/10.1175/1520-0493(1983)111%3C1434:DSOTMB%3E2.0.CO;2), 1983.
- Hardiman, S. C., Boutle, I. A., Bushell, A. C., Butchart, N., Cullen, M. J., Field, P. R., Furtado, K., Manners, J. C., Milton, S. F., Morcrette, C., et al.: Processes controlling tropical tropopause temperature and stratospheric water vapor in climate models, *Journal of Climate*, 28, 6516–6535, <https://doi.org/10.1175/JCLI-D-15-0075.1>, 2015.
- Hartmann, D. L.: The dynamical climatology of the stratosphere in the Southern Hemisphere during late winter 1973, *Journal of the Atmospheric Sciences*, 33, 1789–1802, [https://doi.org/10.1175/1520-0469\(1976\)033%3C1789:TDCOTS%3E2.0.CO;2](https://doi.org/10.1175/1520-0469(1976)033%3C1789:TDCOTS%3E2.0.CO;2), 1976.
- Holopainen, E.: Recent estimates of diabatic forcing on the planetary scale: A review, in: *Proc. of the ECMWF Workshop on Diabatic Forcing*, pp. 29–57, 1987.
- Hoskins, B., Fonseca, R., Blackburn, M., and Jung, T.: Relaxing the tropics to an ‘observed’ state: Analysis using a simple baroclinic model, *Quarterly Journal of the Royal Meteorological Society*, 138, 1618–1626, <https://doi.org/10.1002/qj.1881>, 2012.
- Iwasaki, T.: A diagnostic formulation for wave-mean flow interactions and Lagrangian-mean circulation with a hybrid vertical coordinate of pressure and isentropes, *Journal of the Meteorological Society of Japan. Ser. II*, 67, 293–312, 1989.
- Iwasaki, T.: General circulation diagnosis in the pressure-isentrope hybrid vertical coordinate, *Journal of the Meteorological Society of Japan. Ser. II*, 70, 673–687, 1992.
- Jeuken, A., Siegmund, P., Heijboer, L., Feichter, J., and Bengtsson, L.: On the potential of assimilating meteorological analyses in a global climate model for the purpose of model validation, *Journal of Geophysical Research: Atmospheres*, 101, 16 939–16 950, <https://doi.org/10.1029/96JD01218>, 1996.
- Karmalkar, A. V., Sexton, D. M., Murphy, J. M., Booth, B. B., Rostron, J. W., and McNeill, D. J.: Finding plausible and diverse variants of a climate model. Part II: development and validation of methodology, *Climate dynamics*, 53, 847–877, 2019.
- Klinker, E.: Investigation of systematic errors by relaxation experiments, *Quarterly Journal of the Royal Meteorological Society*, 116, 573–594, <https://doi.org/10.1002/qj.49711649304>, 1990.
- Klinker, E. and Sardeshmukh, P. D.: The diagnosis of mechanical dissipation in the atmosphere from large-scale balance requirements, *Journal of Atmospheric Sciences*, 49, 608–627, [https://doi.org/10.1175/1520-0469\(1992\)049%3C0608:TDOMDI%3E2.0.CO;2](https://doi.org/10.1175/1520-0469(1992)049%3C0608:TDOMDI%3E2.0.CO;2), 1992.
- Lenderink, G., Siebesma, A. P., Cheinet, S., Irons, S., Jones, C. G., Marquet, P., üLLER, F. M., Olmeda, D., Calvo, J., Sanchez, E., et al.: The diurnal cycle of shallow cumulus clouds over land: A single-column model intercomparison study, *Quarterly Journal of the Royal Meteorological Society: A journal of the atmospheric sciences, applied meteorology and physical oceanography*, 130, 3339–3364, 2004.
- Lock, A., Brown, A., Bush, M., Martin, G., and Smith, R.: A new boundary layer mixing scheme. Part I: Scheme description and single-column model tests, *Monthly weather review*, 128, 3187–3199, 2000.
- Ma, H.-Y., Xie, S., Klein, S., Williams, K., Boyle, J., Bony, S., Douville, H., Fermepin, S., Medeiros, B., Tyteca, S., et al.: On the correspondence between mean forecast errors and climate errors in CMIP5 models, *Journal of Climate*, 27, 1781–1798, <https://doi.org/10.1175/JCLI-D-13-00474.1>, 2014.
- Martin, G., Milton, S., Senior, C., Brooks, M., Ineson, S., Reichler, T., and Kim, J.: Analysis and reduction of systematic errors through a seamless approach to modeling weather and climate, *Journal of Climate*, 23, 5933–5957, <https://doi.org/10.1175/2010JCLI3541.1>, 2010.
- Martin, G. M., Levine, R. C., Rodriguez, J. M., and Vellinga, M.: Understanding the development of systematic errors in the Asian summer monsoon, *Geoscientific Model Development*, 14, 1007–1035, <https://doi.org/10.5194/gmd-14-1007-2021>, 2021.
- Martineau, P., Son, S.-W., and Taguchi, M.: Dynamical consistency of reanalysis datasets in the extratropical stratosphere, *Journal of Climate*, 29, 3057–3074, <https://doi.org/10.1175/JCLI-D-15-0469.1>, 2016.

- Milton, S. and Wilson, C.: The impact of parameterized subgrid-scale orographic forcing on systematic errors in a global NWP model, *Monthly weather review*, 124, 2023–2045, [https://doi.org/10.1175/1520-0493\(1996\)124%3C2023:TIOPSS%3E2.0.CO;2](https://doi.org/10.1175/1520-0493(1996)124%3C2023:TIOPSS%3E2.0.CO;2), 1996.
- Palmer, T., Shutts, G., and Swinbank, R.: Alleviation of a systematic westerly bias in general circulation and numerical weather prediction models through an orographic gravity wave drag parametrization, *Quarterly Journal of the Royal Meteorological Society*, 112, 1001–1039, <https://doi.org/10.1002/qj.49711247406>, 1986.
- Peixoto, J. P. and Oort, A. H.: *Physics of Climate*, vol. 520, American Institute of Physics, 1992.
- Phillips, T. J., Potter, G. L., Williamson, D. L., Cederwall, R. T., Boyle, J. S., Fiorino, M., Hnilo, J. J., Olson, J. G., Xie, S., and Yio, J. J.: Evaluating parameterizations in general circulation models: Climate simulation meets weather prediction, *Bulletin of the American Meteorological Society*, 85, 1903–1916, <https://doi.org/10.1175/BAMS-85-12-1903>, 2004.
- Rao, J., Garfinkel, C. I., Chen, H., and White, I. P.: The 2019 new year stratospheric sudden warming and its real-time predictions in multiple S2S models, *Journal of Geophysical Research: Atmospheres*, 124, 11 155–11 174, <https://doi.org/10.1029/2019JD030826>, 2019.
- Rodríguez, J. M. and Milton, S. F.: East Asian summer atmospheric moisture transport and its response to interannual variability of the West Pacific subtropical high: An evaluation of the Met Office Unified Model, *Atmosphere*, 10, 457, <https://doi.org/10.3390/atmos10080457>, 2019.
- Rodwell, M. and Palmer, T.: Using numerical weather prediction to assess climate models, *Quarterly Journal of the Royal Meteorological Society: A journal of the atmospheric sciences, applied meteorology and physical oceanography*, 133, 129–146, <https://doi.org/10.1002/qj.23>, 2007.
- Saffin, L., Methven, J., and Gray, S.: The non-conservation of potential vorticity by a dynamical core compared with the effects of parametrized physical processes, *Quarterly Journal of the Royal Meteorological Society*, 142, 1265–1275, 2016.
- Sánchez, C., Methven, J., Gray, S., and Cullen, M.: Linking rapid forecast error growth to diabatic processes, *Quarterly Journal of the Royal Meteorological Society*, 146, 3548–3569, 2020.
- Scaife, A., Butchart, N., Warner, C., and Swinbank, R.: Impact of a spectral gravity wave parameterization on the stratosphere in the Met Office Unified Model, *Journal of the atmospheric sciences*, 59, 1473–1489, 2002.
- Senior, C., Arribas, A., Brown, A., Cullen, M., Johns, T., Martin, G., Milton, S., Webster, S., and Williams, K.: Synergies between numerical weather prediction and general circulation climate models, *The development of atmospheric general circulation models*, edited by: Donner, L., Schubert, W., and Somerville, R., Cambridge University Press, Cambridge, UK, 2011.
- Sexton, D., Karmalkar, A., Murphy, J., Williams, K., Boutle, I., Morcrette, C., Stirling, A., and Vosper, S.: Finding plausible and diverse variants of a climate model. Part I: Establishing the relationship between errors at weather and climate time scales, *Climate dynamics*, 53, 989–1022, 2019.
- Smith, A. K. and Lyjak, L. V.: An observational estimate of gravity wave drag from the momentum balance in the middle atmosphere, *Journal of Geophysical Research: Atmospheres*, 90, 2233–2241, <https://doi.org/10.1029/JD090iD01p02233>, 1985.
- Svensson, G., Holtlag, A., Kumar, V., Mauritsen, T., Steeneveld, G., Angevine, W., Bazile, E., Beljaars, A., De Bruijn, E., Cheng, A., et al.: Evaluation of the diurnal cycle in the atmospheric boundary layer over land as represented by a variety of single-column models: The second GABLS experiment, *Boundary-Layer Meteorology*, 140, 177–206, 2011.
- Tanaka, D., Iwasaki, T., Uno, S., Ujiie, M., and Miyazaki, K.: Eliassen–Palm flux diagnosis based on isentropic representation, *Journal of the atmospheric sciences*, 61, 2370–2383, 2004.
- Telford, P., Braesicke, P., Morgenstern, O., and Pyle, J.: Description and assessment of a nudged version of the new dynamics Unified Model, *Atmospheric Chemistry and Physics*, 8, 1701–1712, <https://doi.org/10.5194/acp-8-1701-2008>, 2008.

- van Niekerk, A., Shepherd, T. G., Vosper, S. B., and Webster, S.: Sensitivity of resolved and parametrized surface drag to changes in resolution and parametrization, *Quarterly Journal of the Royal Meteorological Society*, 142, 2300–2313, <https://doi.org/10.1002/qj.2821>, 2016.
- 795 van Niekerk, A., Sandu, I., Zadra, A., Bazile, E., Kanehama, T., Köhler, M., Koo, M.-S., Choi, H.-J., Kuroki, Y., Toy, M. D., et al.: Constraining Orographic Drag Effects (COORDE): A model comparison of resolved and parametrized orographic drag, *Journal of Advances in Modeling Earth Systems*, 12, e2020MS002160, 2020.
- Vosper, S.: Mountain waves and wakes generated by South Georgia: Implications for drag parametrization, *Quarterly Journal of the Royal Meteorological Society*, 141, 2813–2827, 2015.
- 800 Walters, D., Boutle, I., Brooks, M., Melvin, T., Stratton, R., Vosper, S., Wells, H., Williams, K., Wood, N., Allen, T., et al.: The Met Office unified model global atmosphere 6.0/6.1 and JULES global land 6.0/6.1 configurations, *Geoscientific Model Development*, 10, 1487–1520, 2017.
- Walters, D., Baran, A. J., Boutle, I., Brooks, M., Earnshaw, P., Edwards, J., Furtado, K., Hill, P., Lock, A., Manners, J., et al.: The Met Office Unified Model global atmosphere 7.0/7.1 and JULES global land 7.0 configurations, *Geoscientific Model Development*, 12, 1909–1963, 2019.
- 805 Wehrli, K., Guillod, B. P., Hauser, M., Leclair, M., and Seneviratne, S. I.: Assessing the dynamic versus thermodynamic origin of climate model biases, *Geophysical research letters*, 45, 8471–8479, <https://doi.org/10.1029/2018GL079220>, 2018.
- Williams, K., van Niekerk, A., Best, M., Lock, A., Brooke, J., Carvalho, M., Derbyshire, S., Dunstan, T., Rumbold, H., Sandu, I., et al.: Addressing the causes of large-scale circulation error in the Met Office Unified Model, *Quarterly Journal of the Royal Meteorological Society*, 146, 2597–2613, 2020.
- 810 Wilson, D. R. and Ballard, S. P.: A microphysically based precipitation scheme for the UK Meteorological Office Unified Model, *Quarterly Journal of the Royal Meteorological Society*, 125, 1607–1636, 1999.
- Wilson, D. R., Bushell, A. C., Kerr-Munslow, A. M., Price, J. D., and Morcrette, C. J.: PC2: A prognostic cloud fraction and condensation scheme. I: Scheme description, *Quarterly Journal of the Royal Meteorological Society: A journal of the atmospheric sciences, applied meteorology and physical oceanography*, 134, 2093–2107, 2008a.
- 815 Wilson, D. R., Bushell, A. C., Kerr-Munslow, A. M., Price, J. D., Morcrette, C. J., and Bodas-Salcedo, A.: PC2: A prognostic cloud fraction and condensation scheme. II: Climate model simulations, *Quarterly Journal of the Royal Meteorological Society: A journal of the atmospheric sciences, applied meteorology and physical oceanography*, 134, 2109–2125, 2008b.
- Wood, N., Staniforth, A., White, A., Allen, T., Diamantakis, M., Gross, M., Melvin, T., Smith, C., Vosper, S., Zerroukat, M., et al.: An inherently mass-conserving semi-implicit semi-Lagrangian discretization of the deep-atmosphere global non-hydrostatic equations, *Quarterly*
- 820 *Journal of the Royal Meteorological Society*, 140, 1505–1520, <https://doi.org/10.1002/qj.2235>, 2014.
- Yanai, M., Esbensen, S., and Chu, J.-H.: Determination of bulk properties of tropical cloud clusters from large-scale heat and moisture budgets, *Journal of Atmospheric Sciences*, 30, 611–627, [https://doi.org/10.1175/1520-0469\(1973\)030%3C0611:DOBPOT%3E2.0.CO;2](https://doi.org/10.1175/1520-0469(1973)030%3C0611:DOBPOT%3E2.0.CO;2), 1973.
- Yang, W., Seager, R., and Cane, M. A.: Zonal momentum balance in the tropical atmospheric circulation during the global monsoon mature
- 825 months, *Journal of the atmospheric sciences*, 70, 583–599, <https://doi.org/10.1175/JAS-D-12-0140.1>, 2013.