

# Implementation and evaluation of sea level operators in OceanVar2.0: an open-source oceanographic three-dimensional variational data assimilation system

Paolo Oddo<sup>1,2</sup>, Mario Adani<sup>2</sup>, Francesco Carere<sup>2</sup>, Andrea Cipollone<sup>2</sup>, Anna Chiara Goglio<sup>2</sup>, Eric Jansen<sup>2</sup>,  
5 Ali Aydogdu<sup>2</sup>, Francesca Mele<sup>2</sup>, Italo Epicoco<sup>2</sup>, Jenny Pistoia<sup>2</sup>, Emanuela Clementi<sup>2</sup>, Nadia Pinardi<sup>1,2</sup>,  
Simona Masina<sup>2</sup>

<sup>1</sup>Bologna University, Department of Physics and Astronomy, Italy.

<sup>2</sup>CMCC Foundation - Euro-Mediterranean Center on Climate Change, Italy.

10

*Correspondence to:* Paolo Oddo (paolo.oddo@unibo.it)

## Abstract

This study presents the development and sensitivity analysis of the sea level operator within the OceanVar software which implements an oceanographic incremental three-dimensional variational data assimilation scheme. In OceanVar, the background error covariance matrix is decomposed into a sequence of physically based linear operators, allowing for individual analysis of specific error matrix components. The key development of OceanVar2.0 is the full integration of both dynamic height and barotropic model formulations as a flexible option for handling sea level covariance. The comparison of the two formulations of the sea level operator which provides correlations between Sea Level Anomaly, temperature and salinity increments is presented. The sensitivity experiments were performed in the Mediterranean Sea and the quality of the analysis assessed by comparing background estimates with observations for the period January-December 2021. The results confirm the methodological advantage of the barotropic model operator, which successfully overcomes the physical and operational limitations associated with choosing an appropriate level-of-no-motion for the dynamic height formulation. Furthermore, we present a method to assimilate along track satellite altimetry considering a forecasting model with tides.

## 1. Introduction

25 Understanding the past state of the ocean and predicting its future behaviour is critical for sustainable development and for climate change mitigation and adaptation strategies. Oceans are a key component of the Earth's climate system, and they require specific data assimilation schemes due to the sparsity of data in the ocean interior. There are different methodologies for ocean data assimilation, each with its own strengths and weaknesses. Within inverse problem theory, the two most used approaches are the variational and the Kalman filter (Carrassi *et al.*, 2018). Schemes based on Monte-Carlo algorithms, such 30 as the Particle filter, have been proven to be successful on low-dimensional systems and have become feasible for high-

dimensional geophysical systems only recently (Van Leeuwen *et al.*, 2019). The choice of data assimilation method depends on factors such as the type of the observations, the desired forecast horizon, and the available computational resources.

Recent machine learning (ML) advancements offer potential optimizations for ocean data assimilation (e.g., Barthélémy *et al.*, 2022; Beauchamp *et al.*, 2023). ML can refine the representation of the errors and reveal complex relationships, improving accuracy. To fully leverage ML and new data streams, modular and flexible data assimilation software is essential. As research progresses, these advancements will significantly enhance our ability to understand and predict ocean behaviour.

The data assimilation community has made substantial strides in developing widely used software tools and specific implementations. General frameworks such as PDAF (Nerger and Hiller, 2013) and DART (Data Assimilation Research Testbed, Anderson *et al.*, 2009) have contributed significantly to the field. Additionally, model-specific implementations such as ROMS-4DVAR (Moore *et al.*, 2011) have been instrumental in advancing ocean data assimilation. Each of these tools offers unique capabilities, and the increasing complexity of data assimilation problems—particularly with the adoption of ML—drives the continuous development and innovation of both existing and new tools in the community, as evidenced by recent research (Martin *et al.*, 2025).

OceanVar is the data assimilation software used in this study. It was first introduced by Dobricic and Pinardi (2008, hereafter DP08) and implements an incremental three-dimensional variational method. OceanVar features a modular design that allows for flexibility in incorporating diverse data sources and error covariance representations. This adaptability has made it suitable for a wide range of applications and research needs. The software has been extensively used in several operational and reanalysis systems, as evidenced by numerous publications (Dobricic *et al.*, 2007, Storto *et al.*, 2016, Escudier *et al.*, 2021, Lima *et al.*, 2021, Ciliberti *et al.*, 2022, Coppini *et al.*, 2023). It has been used to test new hybrid formulations (Oddo *et al.*, 2016; Storto *et al.*, 2018) and to implement new observational operators for different data types, including Lagrangian trajectories (Nilsson *et al.*, 2012), sea-ice variables (Cipollone *et al.* 2023), daytime SST from SEVIRI (Storto and Oddo, 2019), and AI-based operators (Storto *et al.*, 2021, Broccoli and Cipollone, 2025). Furthermore, it has been instrumental in improving error covariance models (Dobricic *et al.*, 2015) and advancing specific applications, such as biogeochemical modelling (Teruzzi *et al.*, 2014; Teruzzi *et al.*, 2018), and new interdisciplinary uses like assessing the impact on underwater acoustic predictions (Storto *et al.*, 2020).

The extensive use of the code in diverse applications has led to the creation of different versions, some of which are inconsistent or incompatible. This core issue drives the development presented here. For this study, we used OceanVar2.0 (hereafter OceanVar2 for improved readability), a new version of the underlying OceanVar framework that unifies the various developments into a single, consistent, and fully parallelized software package.

In DP08 and all subsequent developments, horizontal covariance was approximated using the recursive filter (Lorenc, 1992; Hayden and Purser, 1995). The filter is conceptually simple, typically requiring only a few iterations to approximate the Gaussian function, and its application on a horizontal grid can be split into two independent directions (Purser *et al.*, 2003). However, in cases of spatially or temporally varying correlation radii, the computational advantage of the recursive filter may be questioned (Purser *et al.*, 2022). To provide the system with greater flexibility in terms of horizontal correlation radii,

65 OceanVar2 models the horizontal correlation by a repeated application of the Laplacian operator, which is also the solution of the horizontal diffusion equation (e.g., Derber and Rosati, 1989; Weaver and Courtier, 2001).  
 Satellite altimetry and satellite-derived sea surface temperature are crucial in bridging gaps left by in-situ observations (Le Traon, *et al.*, 2025). Satellites help to cover almost the entire ocean surface and satellite altimetry contains information of the subsurface thermohaline structure that is key to obtain best estimates of the ocean variability at depth. Satellite altimetric sea  
 70 level data are available from 1992, and altimeters have increased in coverage in the past ten years. The effective integration of satellite altimetry and sea surface temperature data into model corrections requires advanced extrapolation algorithms. While this necessity was first emphasized in foundational work (De Mey and Robinson, 1987), it continues to drive development in the field, as evidenced by cutting-edge solutions using machine learning (Zavala-Romero *et al.*, 2025). In this work we demonstrate the capability of OceanVar2 to effectively assimilate along-track satellite altimetry with the use of a barotropic  
 75 model operator and multivariate sea level, temperature and salinity statistics. Although the barotropic model was conceptually defined in DP08, its implementation was not carried forward or maintained in subsequent mainline versions. This work presents its full integration and optimization within the OceanVar2 framework. We present, for the first time, a detailed, direct comparison of the barotropic model defined by DP08 with the dynamic height operator. Additionally, OceanVar2 is applied to the latest version of the Mediterranean Sea forecasting System (Clementi *et al.*, 2023) that considers tidal forcing. Tides are  
 80 becoming an essential component of the resolved variability of the ocean general circulation, and they can no longer be neglected in numerical ocean circulation models (Arbic, 2022). Satellite altimeters sample tides along their track as well as the mesoscales. Using OceanVar2 we present a preliminary solution to the problem of assimilation in presence of tidal components both in model and observations.

The manuscript is organized as follows. After the introduction, Section 2 provides a general overview of the variational  
 85 formulation and the characteristics of OceanVar2. Section 3 presents the background and the observational error covariance matrix formulation and their specific operators. Section 4 describes the experimental set-up and the altimetry assimilation methods. Section 5 discusses the results. Section 6 provides an overview of the computational performance of the code. Finally, Section 7 provides the summary and conclusions.

## 2. Formulation of the Variational Assimilation Scheme

90 The standard cost function in three-dimensional variational data assimilation is defined as:

$$J(\mathbf{x}) = \frac{1}{2}(\mathbf{x} - \mathbf{x}_b)^T \mathbf{B}^{-1}(\mathbf{x} - \mathbf{x}_b) + \frac{1}{2}(\mathbf{y} - \mathbf{h}[\mathbf{x}])^T \mathbf{R}^{-1}(\mathbf{y} - \mathbf{h}[\mathbf{x}]) \quad (1)$$

where,  $\mathbf{x}$  is the analysis state vector containing the ocean model prognostic variables,  $\mathbf{x}_b$  is the background state vector,  $\mathbf{B}$  is the background error covariance matrix,  $\mathbf{h}$  is the non-linear observational operator,  $\mathbf{y}$  are the observations,  $\mathbf{R}$  is the observational error covariance matrix and  $T$  indicates the matrix transpose. Equation (1) is linearized around the background state (e.g.,  
 95 Lorenc, 1997) and expressed in terms of the increments  $\delta\mathbf{x} = \mathbf{x} - \mathbf{x}_b$ :

$$J(\delta\mathbf{x}) = \frac{1}{2} \delta\mathbf{x}^T \mathbf{B}^{-1} \delta\mathbf{x} + \frac{1}{2} (\mathbf{d} - \mathbf{H}\delta\mathbf{x})^T \mathbf{R}^{-1} (\mathbf{d} - \mathbf{H}\delta\mathbf{x}), \quad (2)$$

where  $\mathbf{d} = \mathbf{y} - \mathbf{h}[\mathbf{x}_b]$  are the misfits (or innovations), and  $\mathbf{H}$  is the Jacobian matrix of  $\mathbf{h}$  at  $\mathbf{x} = \mathbf{x}_b$ . The analysis at a particular time is defined as  $\mathbf{x}_a = \mathbf{x}_b + \delta\mathbf{x}$ , where  $J$  attains its minimum.

Existence and uniqueness of  $\mathbf{x}_a$  is guaranteed because  $J$  is quadratic with  $\mathbf{R}$  and  $\mathbf{B}$  positive definite matrices. The minimum 100 can be found by forcing the gradient of the cost function to zero. The gradient of equation (2) is:

$$\nabla J(\delta\mathbf{x}) = (\mathbf{B}^{-1} + \mathbf{H}^T \mathbf{R}^{-1} \mathbf{H}) \delta\mathbf{x} - \mathbf{H}^T \mathbf{R}^{-1} \mathbf{d}$$

Following DP08, the OceanVar scheme assumes that the  $\mathbf{B}$  matrix can be decomposed as:

$$\mathbf{B} = \mathbf{V}\mathbf{V}^T$$

and the cost function may equivalently be minimized using a new control variable  $\mathbf{v}$  (e.g., Lorenc, 1997) defined using the 105 transformation matrix  $\mathbf{V}^+$ :

$$\mathbf{v} = \mathbf{V}^+ \delta\mathbf{x},$$

where the superscript “+” indicates the generalized inverse. The vector  $\mathbf{v}$  is defined on the control space, and the increment vector  $\delta\mathbf{x}$  on the physical space. The cost function (2) now obtains the form:

$$J(\mathbf{v}) = \frac{1}{2} \mathbf{v}^T \mathbf{v} + \frac{1}{2} (\mathbf{H}\mathbf{v} - \mathbf{d})^T \mathbf{R}^{-1} (\mathbf{H}\mathbf{v} - \mathbf{d}).$$

110 The misfit in OceanVar is estimated using the FGAT (First Guess at Appropriate Time) method. The valid time of the increment using the FGAT algorithm has been discussed and investigated in literature (see Massart *et al.*, 2010). The purpose of FGAT is to ensure that each observation ( $y_i$ ) falling within the assimilation window  $[t_n; t_{n+1}]$  is compared against the model background ( $\mathbf{x}_b$ ) at the specific time of the observation ( $t_i$ ), rather than only at the final analysis time ( $t_{n+1}$ ). The innovations ( $d_i$ ) are thus computed as:

$$115 \quad \mathbf{d}_i = (\mathbf{y}_i - \mathbf{h}[\mathbf{x}_b(t_i)]),$$

where  $\mathbf{x}_b(t_i)$  is the model background interpolated to the observational time  $t_i$ . The assimilation system then minimizes all these innovations to compute a single analysis increment ( $\delta\mathbf{x}$ ) which is applied at  $t_{n+1}$ . Formally our analysis is defined as the instantaneous field:

$$\mathbf{x}_a(t_{n+1}) = \mathbf{x}_b(t_{n+1}) + \delta\mathbf{x},$$

120 where  $\mathbf{x}_b(t_{n+1})$  is the instantaneous background field simulated by the nonlinear ocean model starting from  $t_n$ .

### 3. Modelling the background error covariance matrix

The transformation matrix  $\mathbf{V}$  is modelled at each minimization iteration as a sequence of linear operators (e.g., Weaver *et al.*, 2003). In this way,  $\mathbf{V}$  successively transforms increments in the control space towards final increments in the physical space. In OceanVar and OceanVar2 the matrix  $\mathbf{V}$  is defined in the following way:

$$125 \quad \mathbf{V} = \mathbf{V}_D \mathbf{V}_{u,v} \mathbf{V}_\eta \mathbf{V}_H \mathbf{V}_V \quad (3)$$

From right to left,  $\mathbf{V}_v$  defines the vertical error covariance;  $\mathbf{V}_H$  the horizontal error correlation;  $\mathbf{V}_\eta$  is the sea level operator containing correlation between temperature, salinity and sea surface elevation; and  $\mathbf{V}_{u,v}$  forces a geostrophic balance between temperature, salinity and the velocity components. Finally,  $\mathbf{V}_D$  is a divergence-damping operator avoiding spurious currents close to the coast in the presence of complex coastlines, as defined by DP08.

130 The vertical transformation operator  $\mathbf{V}_v$  has the form:

$$\mathbf{V}_v = \mathbf{S}_c \mathbf{\Lambda}_c^{1/2}$$

where columns of  $\mathbf{S}_c$  contain eigenvectors and  $\mathbf{\Lambda}_c$  is a diagonal matrix with eigenvalues of multivariate Empirical Orthogonal Functions (EOFs). In the OceanVar and OceanVar2 code, the EOFs can be defined pointwise (Coppini *et al.*, 2023) or by regions (DP08).

135 Differently from DP08, to account for horizontal correlations,  $\mathbf{V}_H$  is considered as the discretized form of the diffusive operator:

$$\mathbf{V}_H = \nabla_H \cdot (k_c \nabla_H C)$$

where  $\nabla_H$  is the horizontal differential operator,  $k_c$  is the spatially variable diffusivity coefficient corresponding to horizontal correlation lengths, and  $C$  is a generic increment. Assuming a gaussian solution (Weaver and Courtier, 2001), the relation between  $k_c$  and the horizontal correlation radius is:

140 
$$R_H = \sqrt{2k_c \Delta t}$$

Where  $R_H$ , in meters, is the horizontal correlation radius, and  $\Delta t = 1$  second is the pseudo time-step used to integrate the diffusion equation. This filter was implemented using a dimensional splitting approach. The operator is discretized with a Euler-Backward implicit scheme solved by means of LU decomposition of a tridiagonal matrix, following Hoffman *et al.* (2001). The tridiagonal matrix algorithm is chosen for its simplicity and the resulting ease of implementing its adjoint operator, 145 its memory efficiency, and its ability to avoid external libraries implementing direct methods, thus forming a compact implementation which simplifies the porting of OceanVar2.

The  $\mathbf{V}_\eta$  operator, the focus of this study, incorporates correlations between sea surface elevation and the subsurface thermohaline structure. Section 3.1 is dedicated to the detailed description of the two formulations implemented in OceanVar2. The present formulation of OceanVar also allows for the computation of corrections for the velocity field: the  $\mathbf{V}_{u,v}$  operator 150 computes the velocity corrections assuming geostrophic balance. This assumption results in an operator with small computational cost. However, this assumption is not valid at the equator and may produce velocity vectors orthogonal to the coast. When imposing the zero-boundary condition for the velocity component perpendicular to the coast, the divergence component of the velocity field may become unrealistically large. Therefore, the divergence damping operator  $\mathbf{V}_D$  in (3) is implemented to damp velocity divergence near coasts, while the vorticity remains unchanged. Details on the implementation 155 of the divergence damping operator are provided in DP08.

As highlighted in DP08, the sequence of operator multiplication is critical and determined through a combination of physical reasoning and iterative experimentation. Initially, all increments are projected onto multivariate EOFs for sea level, temperature, and salinity, as these effectively capture ocean stratification (Sanchez de la Lama *et al.*, 2016) and the relationship

with sea level for assimilation purposes (De Mey and Robinson, 1987). Next, the increments are distributed horizontally.

160 Following this, adjustments due to the sea level operator are computed based on the vertically projected temperature and salinity increments. Subsequently, increments in horizontal velocities are derived, and the process concludes with the application of the divergence damping filter.

### 3.1 The sea level operator and its relationship with the velocity terms

165 Dobricic *et al.* (2007) found that the vertical EOFs computed from the covariance between temperature, salinity and sea level could produce corrections that are not geostrophically balanced and proved that the enforcement of the geostrophic relationship for the sea level in the error covariance matrix has a significant positive impact on the accuracy of the analysis. Thus, in OceanVar2 the sea surface height increments from EOF projections are overwritten using a sea level operator. Two different sea level operators, with different levels of complexity, are implemented in OceanVar2.

170 The first is the commonly used Dynamic Height operator ( $\mathbf{V}_\eta = \mathbf{V}_{DH}$ ), which is defined as:

$$\mathbf{V}_{DH} = \delta\eta^{DH} = -\frac{1}{\rho_0} \int_{-D}^0 [-\alpha\delta T + \beta\delta S] dz \quad (4)$$

Where  $D$  is a uniform reference level, corresponding to a level-of-no-motion,  $\delta T$  and  $\delta S$  are temperature and salinity increments respectively,  $\alpha$  and  $\beta$  the expansion and contraction coefficients. The correlation between Sea Level Anomaly (SLA) from altimetry measurements and the dynamic height anomaly computed from in-situ measurements is high for regions

175 deeper than 1000 m (Dhomps *et al.*, 2011), at synoptic, seasonal and interannual time scales.

OceanVar2 allows the application of a more complex linear  $\mathbf{V}_\eta$ . This operator is based on the formulation published in DP08 and derives from the steady state results of a linear barotropic model forced by buoyancy anomalies induced by the temperature and salinity increments. While the methodology was previously described in the literature, the barotropic model code itself was fully re-introduced and parallelized to function efficiently within the new OceanVar2 framework, with its general 180 mathematical structure remaining unchanged. Its inclusion as a fully integrated option is a new feature of this version. The barotropic model equations, discretized in time by the semi-implicit scheme (Kwizak and Robert, 1971), are:

$$\frac{U^{n+1} - U^{n-1}}{2\Delta t} - fV^n = -gH \frac{\partial\eta^*}{\partial x} - \int_{-H}^0 \left[ \int_{-z}^0 \frac{\partial(\delta b)}{\partial x} dz' \right] dz + \gamma \nabla^2 U^* \quad (5)$$

$$\frac{V^{n+1} - V^{n-1}}{2\Delta t} + fU^n = -gH \frac{\partial\eta^*}{\partial y} - \int_{-H}^0 \left[ \int_{-z}^0 \frac{\partial(\delta b)}{\partial y} dz' \right] dz + \gamma \nabla^2 V^* \quad (6)$$

$$\frac{\eta^{n+1} - \eta^{n-1}}{2\Delta t} + \left( \frac{\partial U^*}{\partial x} + \frac{\partial V^*}{\partial y} \right) = 0$$

185 where  $U$  and  $V$  are vertically integrated velocity components,  $f$  is the Coriolis parameter,  $g$  is the acceleration due to gravity,  $H$  is the bottom depth,  $\eta$  is the surface elevation,  $\delta b$  is the buoyancy anomaly, and  $\gamma$  is the horizontal viscosity coefficient. The superscripts indicate the time step relative to  $n$ , and the superscript “ $*$ ” indicates the weighted average between two

timesteps. A more detailed description of the barotropic model and its discretization can be found in DP08. In the present version the barotropic model assumes closed lateral boundaries.

190 In the previous equations (5) and (6), the buoyancy forcing term is defined as:

$$\delta b = g \left( \frac{\delta \rho}{\rho_0} \right)$$

and the density perturbation  $\delta \rho$  is estimated by the linear equation of state:

$$\delta \rho = \alpha \delta T - \beta \delta S \quad (7)$$

Expansion and contraction coefficients ( $\alpha$  and  $\beta$ ) in eqs. (4) and (7) can be assumed space independent or spatially variable

195 and estimated by linearizing the equation of state around a user defined background temperature ( $T_b$ ) and salinity field ( $S_b$ ):

$$\alpha = \frac{\partial \rho}{\partial T} \Big|_{T=T_b, S=S_b}$$

$$\beta = \frac{\partial \rho}{\partial S} \Big|_{T=T_b, S=S_b}$$

In the latter case the coefficients are read from an external input file.

In OceanVar2 the sea level operators produce the final sea surface height increments, replacing the increment produced by the

200 cross-covariance between temperature, salinity and sea level provided by the EOFs (DP08, Storto *et al.*, 2018).

The choice on the sea level operator has consequences on the velocity operator  $\mathbf{V}_{u,v}$ .  $\mathbf{V}_{u,v}$  computes the velocity correction assuming geostrophic balance under the Boussinesq and incompressibility approximations:

$$fu_g(z) = - \frac{1}{\rho_0} \frac{\partial p}{\partial y}$$

$$fv_g(z) = + \frac{1}{\rho_0} \frac{\partial p}{\partial x}$$

205 Decomposing the pressure  $p$  at any level  $z$  as:

$$p(z) = p_{atm} + g \rho_0 \left( \eta + \int_{-z}^0 \frac{\delta \rho}{\rho_0} dz \right)$$

where  $p_{atm}$  is the atmospheric pressure,  $g$  is the effective gravity,  $\eta$  is the free surface elevation and  $\delta \rho$  is the density departure from a reference state  $\rho_0$ . Neglecting the atmospheric pressure, and rewriting the hydrostatic term as buoyancy term, the geostrophic velocities become:

$$fu_g(z) = - \left( g \frac{\partial \eta}{\partial y} + \int_{-z}^0 \frac{\partial \delta b}{\partial y} dz \right) \quad (8)$$

$$fv_g(z) = + \left( g \frac{\partial \eta}{\partial x} + \int_{-z}^0 \frac{\partial \delta b}{\partial x} dz \right) \quad (9)$$

When adopting the barotropic model as  $\mathbf{V}_\eta$ , the sea surface height in eqs. (8) and (9) is replaced by the increments deriving from the solution of the barotropic model and the lower limit of the baroclinic integral reaches  $-H(x,y)$ , the ocean floor. On the other hand, when  $\mathbf{V}_\eta = \mathbf{V}_{DH}$  the horizontal pressure-gradient force must vanish at the level-of-no-motion,  $D$ , in (4). In eqs.

215 (8) and (9)  $\eta$  is then substituted with  $\delta\eta^{DH}$  and the velocity increments are computed only up to depth D. Thus, in case of the barotropic model, velocity corrections are provided for the entire water column, while in case of dynamic height, velocity corrections are provided from the surface to the level-of-no-motion.

### 3.2. The observational error covariance matrix and quality checks

220 The observation error  $\epsilon^0$  is defined as the difference between the observation vector  $y$  and the observation counterpart in the true state  $x^t$  (Cohn, 1997)

$$y = h^*[x^t] + \epsilon^0 \rightarrow \epsilon^0 = (y - y^t) - (h^*[x^t] - y^t) = \epsilon^m + \epsilon^r,$$

225 where  $y^t$  is the *true* (unknown) observation value,  $h^*$  is an observation operator,  $\epsilon^m$  labels the instrumental or measurement error (distance of the actual value from the true state) and  $\epsilon^r$  gathers the different components of the representativeness errors, due to inaccuracies in  $h^*$  and the sampling error of the observations with respect to the true signal. Under the assumption of unbiased error  $\langle \epsilon^0 \rangle = 0$  and that  $\epsilon^m$  and  $\epsilon^r$  are uncorrelated, the error covariance matrix  $\mathbf{R}$  can be constructed as the sum of two terms that can be estimated independently:

$$\mathbf{R} = \langle \epsilon^0 \epsilon^0 \rangle \approx \langle \epsilon^m \epsilon^m \rangle + \langle \epsilon^r \epsilon^r \rangle = \mathbf{R}^m + \mathbf{R}^r.$$

230 If the errors associated to different observations are uncorrelated, the two matrices simplify greatly in diagonal ones. This hypothesis is valid for most of the current global/regional observational datasets and it is generally correct when observations are sampled relatively far in time (say few hours to avoid cross-correlation term in  $\mathbf{R}^m$ ) or are sparse with respect to the model grid resolution (to not include off-diagonal elements in  $\mathbf{R}^r$ ). Observation errors are a function of observation type in OceanVar2. OceanVar2 contains various procedures for the quality control and preprocessing of observations. A background quality check is included to reject observations that are too far from the model estimate. This quality check uses a threshold on the squared misfit. While OceanVar2 supports approximating the misfit distribution with a Huber norm PDF (Storto, 2016) to mitigate the 235 impact of outliers, this feature was not required for the analysis presented in this work.

240 To ensure that the assumption of spatially and temporally uncorrelated observation error is satisfied, data thinning and superrobbing procedures are employed. Horizontal thinning rejects observations that are too close in space, and in cases where multiple data from the same instrument fall into the same model grid cell, only the observation closest to the analysis time is retained. Along the vertical dimension, a superrobbing procedure takes the average of all data falling within the same model layer. Additionally, two important rejection criteria can be activated: coastal rejection prevents the assimilation of altimetric and in-situ coastal observations to avoid inconsistencies between observed and modelled coastal processes, and a bathymetry-based rejection prevents the assimilation of data in shallow areas.

## 4. Experimental Design

245 The experiments are designed to determine the best configuration of OceanVar2 for the assimilation of satellite altimetry together with ARGO floats and XBT in the Mediterranean Sea. The set-up of the ocean model used in this study is a simplified

version of the physical component of the Mediterranean Forecasting System of the Copernicus Marine Service (Clementi *et al.*, 2023). The model is implemented over the entire Mediterranean basin (Fig.1) with a horizontal grid resolution of 1/24° (approx. 4 km) and 141 non-uniformly distributed vertical levels. The ocean model code is based on the Nucleus for European Modelling of the Ocean (NEMO) v4.2 (Madec *et al.*, 2023) and includes the representation of tides. Atmospheric forcings are 250 calculated interactively with the operational fields of the European Centre for Medium-Range Weather Forecasts (ECMWF). The only difference to the Copernicus operational system is the omission of the surface wave model coupling.

Starting from an operational analysis, we performed a 1-year simulation followed by one year of daily assimilation cycles of 255 in-situ (XBT and ARGO floats) and satellite SLA data for the whole year of 2021. Figure 1 shows the positions of the assimilated in-situ and SLA data. The SLA data coverage in the bottom panel refers to the full 21-day repeat cycle of the satellite orbits. In our experimental set-up we perform daily assimilation cycles starting at midnight every day and we assimilate all observations available in the 24 hours before the analysis time. The figure also highlights the data used during a single daily cycle (11 August 2021), specifically showing the in-situ and SLA track subset daily coverage.

#### 4.1. Correcting the misfits for tides

A fundamental aspect to consider when assimilating SLA is the possible presence of tides in the modelled solution and in the 260 observed data. Discrepancies between modelled and observed tides can, as a first approximation, be attributed to inaccuracies in the bathymetry of the model, the bottom and/or the coastal frictional dynamics. If the difference between observed and modelled estimates is due to tides, then this part of the misfit is primarily composed of external gravity waves. In the present OceanVar2 formulation, this high-frequency signal would be incorrectly projected into baroclinic increments by the covariance 265 matrix of the background error (eq. 3). Since tidal errors are not dynamically linked to the slow baroclinic ocean state, using the standard **B** matrix leads to spurious analysis increments. It is therefore essential to filter out the tidal signal from both the observed and modelled SLA. This work offers a solution to the assimilation of satellite altimetry in a model with tides, showing that a filtering procedure can be accurate enough and that no additional adjustment is required in the analysis.

The Copernicus along-track sea level anomalies are provided together with an estimate of the tidal signal along the tracks so 270 tides can be filtered easily from the observations. To remove the tidal signal from the model background field, the tidal amplitude and phase for the eight components included in the Mediterranean Sea model (M2, S2, K1, O1, N2, P1, Q1 and K2) have been derived from a simulation output by harmonic analysis of the hourly sea level field. Following Cao *et al.* (2015), six months of hourly data were used for the harmonic analysis. The harmonic analysis was performed using the Pawlowicz *et al.* (2002) algorithm, based on the Foreman method (Foreman 1977; Foreman 1978) at each model grid point. Knowing the tidal constants, it is possible to estimate the model tidal sea level at the exact time and location of the altimetry data and remove 275 this component from the model outputs.

During the model simulation, misfits between model estimates and observations are computed and before entering OceanVar2, the misfits are updated, removing the tidal signal from both observations and model results. In Fig.2 an example of an SLA satellite track is provided with model estimates and the satellite observations, the position of the track is shown in Fig.1. In the

upper panel of Fig.2, the full signal from the model simulation and the observations is drawn as a function of latitude along  
280 the track. The middle panel of Fig. 2 shows the signals without tides, in addition to a debiasing procedure described by Dobricic  
et al. (2012). Dobricic *et al.* (2012) show that this method is the best for considering differences between the large-scale steric  
signal and the mean dynamic topography between observations and model. The average difference along the track is removed  
if the track is continuous, or for individual segments if the track is discontinuous due to the presence of land. Finally, in the  
285 bottom panel of Fig.2 the two tidal components for the observational and modelled SLA are shown indicating the large-scale  
signal of tides in the open ocean.

#### 4.2. Sensitivity experiments

In addition to a free-run (non-assimilative model simulation), we present results from two sets of six 1-year assimilative  
experiments each, comparing them against each other and against observations. All the experiments assimilate ARGO floats  
and XBT temperature and salinity data in the whole domain including the Atlantic part, while SLA data are assimilated only  
290 within the Mediterranean Basin (see Fig.1). In every experiment the vertical component of the background error covariance  
matrix is modelled using 25 tri-variate EOFs (temperature, salinity and SLA) computed following Dobricic *et al.* (2006) for  
every model grid point. The EOFs are computed from a 30-year timeseries of the Mediterranean Sea reanalysis (Escudier *et  
al.*, 2021). The horizontal correlation radius was set to a constant value of 27 kilometres, determined through sensitivity  
experiments. The diffusive filter was iterated five times to model the horizontal covariance. To account for coastal effects, the  
295 correlation radius was linearly decreased starting from about 30 kilometres offshore to the minimum grid resolution near the  
coast. Additionally, a Neumann boundary condition was applied at the coast, setting the normal derivative of the field to zero.  
Observations are rejected if they are less than 15 km from the coast and if the misfits are larger than fixed thresholds: 5°C for  
300 temperature, 2 psu for salinity; and 30 cm for SLA. The observational error covariance matrix is assumed diagonal. The results  
presented in Fig. 2 demonstrate that our model accurately reproduces along-track SLA tidal gradients, with the difference  
between the modelled and observed tidal signal being nearly constant. We effectively remove this along-track bias by  
subtracting the mean residual along each satellite track. This critical step ensures that the resulting misfit is very similar to the  
one computed without tides. Based on these considerations, we decided not to modify the SLA observational error, maintaining  
consistency with previous work (Escudier *et al.*, 2021). Consequently, all the SLA data have an associated error of 3 cm  
305 regardless of the satellite and the geographic distribution. The observational errors for in-situ observations were tuned via the  
Desroziers' method (Desroziers *et al.*, 2005) and vary monthly. We first prescribed the observation errors used in the previous  
system. The assimilation system was then run repeatedly, using the innovations and residuals to apply the Desroziers' formula  
to obtain an estimate for the observation error **R**. This process was iterated until the errors converged (e.g., Escudier *et al.*,  
2021). The resulting vertical error profiles are as follows: temperature and salinity observational errors peak at the surface with  
values of 0.45 °C and 0.14 psu, from 75 to 325m depth they decrease linearly to values of 0.2 °C and 0.05 psu, starting from  
310 750m they have constant values of 0.1 °C and 0.02 psu respectively.

In all the experiments, the state vector contains the following model state variables:

$$\mathbf{x} = [T, S, u, v, \eta]^T$$

where  $T$  is the three-dimensional temperature field,  $S$  the three-dimensional salinity field,  $u$  and  $v$  are the total horizontal velocity components and  $\eta$  the two-dimensional sea surface height.

315 Two sets of experiments are performed to compare the operational stability of mass-field-only  $(T, S, \eta)$  correction against the maximal impact of full-state variable  $(T, S, u, v, \eta)$  correction. Each of the two sets consists of six individual experiments (Exp-1 through Exp-6) with different OceanVar2 sea level operators and choices of free parameters. In Exp-1, which is used as a reference experiment, the barotropic model is used as the sea level operator with constant (in space and time)  $\alpha$  and  $\beta$  in eq. (7) and SLA data are rejected when falling in areas shallower than 100m. In the second experiment (Exp-2), consistently with  
 320 Adani *et al.* (2011), we rejected SLA data falling in areas shallower than 150m. Experiment 3 is similar to Exp-2, but we test the sensitivity to variable expansion and contraction coefficients in eq. (7). The coefficients are computed by linearizing the equation of state around a monthly mean climatology. In all these three experiments we integrated the barotropic model for 3 days with a time-step of 3600 sec and then used the average of the last day as an approximation of the steady state solution. The integration of the barotropic model is fully implicit and the turbulent viscosity is equal to  $650 \text{ m}^2/\text{s}$ .

325 Experiments 4, 5 and 6 use the dynamic height as the sea level operator. The difference between them is the choice of the level-of-no-motion. In Exp-4 we used a level-of-no-motion equal to 150m, thus Exp-2 and Exp-4 differ only for the sea level operator. This shallow depth was specifically chosen to test the upper limit of observation inclusion, ensuring that the assimilation scheme uses the maximum possible Sea Level Anomaly (SLA) data coverage, even in areas shallower than the generally accepted level-of-no-motion. In Exp-5 the level-of-no-motion is 350m in agreement with the Mean Dynamic  
 330 Topography (Rio *et al.*, 2014) used. Finally, in Exp-6 the level-of-no-motion is set at 1000 m which is the traditional choice for the operational setting of the Mediterranean Sea forecasting system (Coppini *et al.* 2023). In all these experiments, the depth of the level-of-no-motion naturally coincides with the minimum depth of SLA observations inclusion in the data assimilation scheme.

The configuration of the six individual experiments (Exp-1 to Exp-6) is identical across both sets. The two sets of experiments  
 335 are designed to test the critical operational trade-off between numerical stability and maximal observational impact. In the first set of experiments, we prioritize stability by applying the analysis increments only to the assimilated variables ( $T, S$  and  $\eta$ ), even though the state vector contains the velocity field. This forces the model dynamics to adjust the velocities indirectly. In the second set of experiments, which tests the maximal assimilation scope, also velocity corrections are applied directly in the analysis step. Throughout the remainder of the manuscript, experiments marked with an asterisk refer to those with velocity  
 340 corrections. The first sets of experiments are summarized in Table 1. The second set of experiments has the same naming convention as the first set, with the sole difference being the application of the velocity correction in the analysis definition.

345

	Exp-1	Exp-2	Exp-3	Exp-4	Exp-5	Exp-6
<b>Sea level operator</b>	BM	BM	BM	DH	DH	DH
$\alpha/\beta$	Const.	Const.	MC	Const.	Const.	Const.
<b>Reference Level for DH (m)</b>	N/A	N/A	N/A	150	350	1000
<b>SLA Min depth rejection (m)</b>	100	150	150	150	350	1000

**Table 1** Sensitivity experiments configurations. First row indicates the sea level operator used: BM= Barotropic Model; DH=Dynamic height. Second row indicates the choice for expansion and contraction coefficients: constant in space and time (Const) and spatially and temporally variable as computed from monthly climatology (MC). Third row indicates the reference level for the lower integral limit of the dynamic height operator and thus the level-of-no-motion for the  $\mathbf{V}_{u,v}$  part of the model background error covariance matrix. Fourth row indicates the minimum depth used as criterium to reject SLA data.

#### 4.3. Performance Metrics

To fully assess the performances of the experiments listed in Table 1, the mean squared error (e.g., Murphy, 1988) is decomposed and the single components are analysed:

$$MB = \bar{m} - \bar{o}, \quad (10)$$

$$SDE = \sigma_m - \sigma_o, \quad (11)$$

$$CC = \frac{1}{\sigma_o \sigma_m} \frac{1}{N} \sum_{i=1}^N (m_i - \bar{m})(o_i - \bar{o}), \quad (12)$$

where MB is the mean bias error, SDE is the standard deviation error and CC is the cross correlation between the modelled and observed fields. The  $i$ -th modelled and observed variable is denoted by  $m_i$  and  $o_i$ , respectively;  $\bar{m}$  and  $\bar{o}$  are the respective averages (horizontal and temporal); while  $\sigma_m$  and  $\sigma_o$  are the respective standard deviations. In addition, the unbiased root mean squared error ( $uRMSE$ ) is computed:

$$uRMSE = \sqrt{\frac{1}{N} \sum_{i=1}^N [(m_i - \bar{m}) - (o_i - \bar{o})]^2}. \quad (13)$$

It is important to note that the model results and observations used here are the same as those used to calculate misfits within the assimilation cycle. However, while not all misfits are utilized in the assimilation process, all available observations are included in the error statistics. This ensures that all experiments are evaluated based on the same reference dataset of observations. Furthermore, to evaluate model performance even in very shallow regions, the observational dataset used in the misfits, and thus in the calculation of the error statistics, includes sea level anomaly (SLA) data covering regions up to 10m

370 depth. This allows for the assessment of model skill in very shallow regions where data are not assimilated in any of the presented experiments.

Given that the  $CC$  is always positive for all the experiments, the misfit statistics for the experiments listed in Table 1 are also analysed in terms of relative improvement with respect to the free-run or Exp-1 according to the following metrics definition:

$$S_{uRMSE_{Exp\#}} = \left( \frac{uRMSE_{Ref} - uRMSE_{Exp\#}}{uRMSE_{Ref}} \right) \times 100, \quad (14)$$

$$S_{CC_{Exp\#}} = \left( \frac{CC_{Exp\#} - CC_{Ref}}{CC_{Ref}} \right) \times 100, \quad (15)$$

375 where  $\#$  indicates the different experiments listed in Table 1 and “ $Ref$ ” denotes the free-run or Exp-1.  $S_{CC}$  is only calculated using Exp-1 as the reference. The exclusion of the free-run as a reference is necessary because the very small  $CC$  values would lead to excessively large percentage values that hinder meaningful comparison between the different experiments. In the manuscript the relative performance statistics are presented only for the SLA data given that a similar comparison for the  
380 temperature and salinity did not provide additional insights.

## 5. Results

Figure 3 shows the temperature and salinity increments resulting from different assimilation scheme setups (Table 1), all starting with the same set of SLA misfits. These increments are presented before analysing the skills of the various experiments. The differences between the experiments are generally small, and of the order of 10%. The largest  
385 differences are due to the different number of SLA data assimilated, a result of the constraint imposed by the level-of-no-motion used or the different minimum depth rejection criterion adopted. We note that when the same data are assimilated, thus in areas deeper than 1000m, very similar increments in SLA are generated by the OceanVar2 regardless of the schemes adopted. However, the schemes differ on how these increments are projected into temperature and salinity increments. Note that the ordinate axes are strongly stretched in the figure to highlight the first 150m depth where most of the corrections are  
390 confined. Comparing Exp-2 and 3, which differ only in the use of spatially and temporally variable expansion and contraction coefficients, we observed small but noticeable differences in the temperature and salinity increments, particularly in the amplitude of near-surface maxima. The choice of sea level operator substantially influenced the results. When using Dynamic Height with a level-of-no-motion set at 1000 meters (Exp-6), temperature and salinity increments were comparable to those obtained with the barotropic model. The primary cause of the observed differences appears to be the constraint imposed on  
395 assimilated data by the level-of-no-motion. Reducing the level-of-no-motion (Exp-4 and 5) allowed for the assimilation of more sea level anomaly data but resulted in considerably different temperature and salinity increments within the first 100 meters compared to the barotropic model.

## 5.1. Barotropic sea level operator and free-run comparison

400 The performance of Exp-1 is compared with free-run. In Fig.4 the statistics for the SLA are shown. Every point in the time-series represents 5-day window statistics. That is, the overbars and the standard deviations in eqs. (10), (11), (12) and (13) are computed over a 5-day time window.

The free-run has an error of about 5 cm, slightly growing during the second half of the year. The model with assimilation underwent a 10-20 day adaptation period, after which the  $uRMSE$  of the misfit stabilizes around 3 cm. A consistent 405 improvement is noted in the  $CC$ . No seasonal cycle is observed in the  $CC$  of Exp-1, whereas the free-run exhibits a distinct summer minimum in the correlations. The  $SDE$  in the free-run is on average negative and it is characterized by 5-day oscillations. In Exp-1, the  $SDE$  stabilized around values of 0.25 cm, indicating an overestimation of the observed variability.

The SLA yearly averaged statistics were clustered according to ocean bathymetry and are plotted in Fig.5. In areas with 410 bathymetry between 150 and 2500 m, the free-run exhibited an almost constant  $uRMSE$ . However, the error increases in shallower and deeper regions, reaching the maximum in areas deeper than 3500 m. The  $uRMSE$  of the Exp-1 was more stable and approximately half of the corresponding free-run statistics. Regardless of the region considered, Exp-1 has better statistics than the free-run, indicating the effectiveness of the assimilation procedure. For the  $CC$  differences between free-run and Exp-1 results are also evident. In the free-run  $CC$  decreases with depth, while for the Exp-1 we observe an opposite tendency. In terms of  $SDE$ , the largest improvements with respect to the free-run are in shallow areas. The free-run underestimates the 415 variance in areas with bathymetry shallower than 1000m, this is particularly evident in areas shallower than 150m. The model with data assimilation tends to overestimate the observed variability particularly in shallow areas.

In Fig.6 the vertical profiles of  $uRMSE$ ,  $MB$ ,  $CC$  and  $SDE$  for salinity (upper panels) and temperature (bottom panels) for the free-run and the Exp-1 are shown. Statistics are computed against all available ARGO and XBT profiles. The vertical structure 420 of the salinity  $uRMSE$  is similar between the free-run and the Exp-1. The salinity errors are characterized by a near surface maximum which is reduced in the assimilative run. The  $MB$  in the free-run has a subsurface minimum at 100 m depth, while Exp-1 misfits have almost homogenous values through all the water column. The  $CC$  resembles the vertical distribution of the  $uRMSE$ , with values approaching the unity in both the experiments below 500 m depth. Finally the salinity  $SDE$  confirms the large improvement arising from the assimilation procedure. In the near surface layer, the free and the assimilative run have opposite behaviours, with the free-run overestimating the observed variability while the assimilative run underestimating it.

425 Below 100 m depth the  $SDE$  salinity values of Exp-1 are noticeably reduced with respect to the free-run.

The temperature  $uRMSE$  and  $CC$  are characterized by a strong, summer intensified (not shown), subsurface maximum/minimum due to the model difficulties in reproducing the correct stratification. A second  $uRMSE$  local maximum (CC minimum) is present around 300 m probably related to the misrepresentation of the Levantine Intermediate Water (LIW) advection in the different Mediterranean regions. A third temperature error relative maximum is present between 1000 and 430 1500 m depth. The assimilation corrects all the errors by approximately 30-50% down to 500 m, less below this depth.

Temperature  $MB$  is largely improved by assimilation. The free-run tends to overestimate the observed temperature variability, and the  $SDE$  has a marked vertical structure. In general, the assimilation seems capable of correcting most of the model errors except in the upper thermocline/mixed layer depth. Analysis of the corresponding time-series (not shown) indicates a clear summer maximum in all the error statistics in proximity of the mixed layer depth. This behaviour is shared between all the 435 different experiments, but it is clearly reduced in the assimilative runs. However, the persistence of this error maximum suggests a limitation in the current formulation of the Background Error Covariance. Specifically, the static, climatological nature of the EOFs used to model the vertical error component struggles to fully capture the rapidly evolving stratification and strong vertical gradients characteristic of the summer mixed layer. Future development will prioritize replacing these EOFs with more dynamic and stratification-aware operators to address this deficiency.

#### 440 5.2 Sensitivity experiments to the sea level operator

Figure 7 illustrates the performance of the six different data assimilation experiments (without velocity corrections) in terms of correlation coefficient ( $CC$ ) and improvements in root-mean-square error ( $S\_uRMSE$ ) relative to the model free-run. The top panels display the time evolution of these improvements, highlighting both short-term fluctuations and overall trends. The bottom panels present the time-averaged  $S\_uRMSE$  and  $CC$  clustered by bathymetric depth ranges, revealing how the 445 effectiveness of each experiment varies with depth. All assimilative experiments outperformed the model free-run. Experiments using the dynamic height as the sea level operator, with levels of no motion set at 150 meters (Exp-4) or 1000 meters (Exp-6), generally performed worse than the other experiments. For Exp-6, the differences in both  $uRMSE$  and  $CC$  were particularly noticeable in regions shallower than 1000 meters, where SLA data were not assimilated. However, even in these regions, Exp-6 substantially outperformed the model free-run, suggesting that corrections applied in deeper areas 450 effectively propagated into shallower regions. In Exp-4, the deterioration in results compared to other experiments was primarily confined to deeper regions. In areas shallower than 150 m Exp-1 outperforms the other experiments, however the performances of the experiments are similar indicating that the coastal areas are strongly constrained by the open ocean dynamics. A clear dependence of the  $CC$  on model bathymetry was evident in all experiments with the most significant improvements (with respect to the free-run) observed in areas deeper than 3500 meters, where the model free-run exhibited

455 the smallest  $CC$ . In shallow regions, Exp-6 generally provided the smallest improvement in  $CC$  compared to the other experiments. The results demonstrate that certain experiments achieve substantial improvements in deep-ocean regions, while others show more consistent performance across all depths. These results highlight the challenges associated with choosing an appropriate level-of-no-motion in data assimilation of SLA. The requirement to define a spatially constant level-of-no-motion, which is inherent to the dynamic height operator, demonstrates the critical limitations imposed by its theoretical assumptions, 460 and our experimental results explicitly highlight these deficiencies. This constraint imposes a critical methodological compromise. For instance, maintaining a conservative, deep level-of-no-motion (e.g., 1000m in Exp-6) ensures physical consistency but mandates the exclusion of a large fraction of the available SLA observations. Conversely, adopting a shallower level-of-no-motion (e.g., 150m in Exp-4) maximizes observation coverage, yet the resulting analysis is compromised due to

the violation of the theoretical assumptions in deeper water, ultimately leading to degraded accuracy. The choice of the level-  
465 of-no-motion can significantly impact the accuracy of model analysis, especially in complex regions with varying bathymetry and ocean dynamics, thereby confirming the methodological advantage of the barotropic model operator which successfully assimilates all available SLA data without imposing a constant level-of-no-motion constraint.

Figure 8 presents the relative performance of five data assimilation experiments (Exp-2 through Exp-6) compared to a baseline assimilative experiment (Exp-1), now used as the reference. The top panels illustrate the time series of the relative change in  
470 root-mean-square error ( $S_{uRMSE}$ ) and correlation coefficient ( $S_{CC}$ ). The bottom panels depict the time-averaged  $S_{uRMSE}$  and  $S_{CC}$  changes, categorized by bathymetric depth ranges. In contrast to the previous figure, where improvements were relative to a model free-run, this figure demonstrates the relative performance of each experiment against the initial data assimilation run. Negative values indicate a decrease in performance (higher  $uRMSE$  or lower  $CC$ ) compared to Exp-1, while positive values indicate improvement. This comparison highlights the incremental benefits or drawbacks of different  
475 experimental configurations in relation to a specific data assimilation configuration.

All the experimental configurations employing the barotropic model perform similarly to each other. In terms of time-series comparison, Exp-4 (with dynamic height as sea level operator and level-of-no-motion equal to 150m) has performance worse than all the other experiments. Among the experiments using the dynamic height operator, Exp-5 generally has better results both in terms of  $uRMSE$  and  $CC$ . The analysis per bathymetric class highlights the differences among the experiments. None  
480 of the experiments outperform Exp-1 in regions shallower than 150m both in terms of  $uRMSE$  and  $CC$ . In deeper areas we see that Exp-4 and Exp-5 produce, in general, worse results, and the worsening is amplified as the depth increases and the level-of-no-motion decreases; 1000m depth is a clear boundary for the effectiveness of Exp-6. Employing the barotropic model as a sea level operator yields consistent results, with small sensitivity to the minimum depth used in the rejection criterion or to the choice of constant or variable expansion/contraction coefficients. This confirms the difficulty of establishing a constant  
485 level-of-no-motion and highlights the benefit of using the barotropic model as a balancing mechanism.

### 5.3. Sensitivity experiments to velocity corrections

The second set of experiments (Exp-1\* to Exp-6\*) was carried out including velocity corrections in the analysis estimates. This second set represents the maximal scope of multivariate assimilation, providing a benchmark against the mass-field-only correction of the first set. The OceanVar2 configuration used in Exp-4\* generated velocity increments that led to numerical  
490 instabilities in the ocean model, preventing this simulation from completing. This critical instability confirms the operational risk associated with introducing direct momentum corrections and highlights the sensitivity of Exp-4. In contrast, the other experiments in this set ran without such issues, underscoring the challenges associated with dynamic height methods. Consequently, our focus is on the stable experiments. Across all the stable configurations, the successful experiments in the second set, which included velocity corrections, demonstrated improved performance compared to their counterparts in the  
495 first set, without velocity corrections. The extent of improvement varies depending on the specific experiment and the region analysed.

Figure 9 shows the time series and Fig.10 presents temporally averaged statistics by bathymetric class for the second set of experiments. In terms of time series (Fig.9), the error components exhibit the same characteristics as those discussed for Exp-1 (Fig.4 and 5). The  $uRMSE$  exhibits a summer minimum in all experiments. Exp-6\* performs significantly worse than the others throughout the year. All experiments using the barotropic model have similar  $uRMSE$  values, with Exp-1\* generally appearing slightly better than the others. The correlation coefficient ( $CC$ ) increases throughout the experiment's length. For this statistic as well, Exp-6\* is the worst, showing consistently lower values than the other experiments. Even for the SDE, which is generally reduced compared to the previously studied experiments without velocity corrections, the relative performance of the different experiments is confirmed.

Figure 10 presents the temporally averaged statistics clustered according to the bathymetry. All the stable experiments benefited greatly from the inclusion of the velocity corrections. The overall reduction in error confirms the importance of the velocity increments when the system remains stable. Exp-6\* confirms its poor performance in areas shallower than 1500m. However, by also correcting the velocities, its statistics in deep areas are now similar or slightly better than those obtained from experiments using the barotropic model as operator in the background error covariance matrix. Exp-1\* is now the best among those analysed for all the bathymetric classes shallower than 500m.

In terms of correlation coefficient, the results previously obtained by analysing  $uRMSE$  seem to be confirmed. For shallow areas ( $<1000m$ ), Exp-6\* is significantly worse than the others. In all other bathymetric classes, while confirming the previous findings, the differences between the experiments are less pronounced. A different behaviour is observed when analysing the standard deviation of the error. Exp-1\* remains the configuration that shows significantly lower error values than the others in almost all bathymetric classes. Notably, Exp-6\* is the one that benefits the most from velocity corrections, demonstrating that for this specific configuration, the direct momentum increments overcome other methodological weaknesses.

## 6. Computational performance and parallelization

To improve computational performance, OceanVar2 adopts a domain-decomposition scheme. This scheme leverages the computing power of a parallel computer by partitioning the computational domain into subdomains. Each process executes the necessary operations to update its portion of the global domain, sharing communications with neighbouring processes for lateral boundary treatments using Message Passing Interface (MPI) calls (The MPI forum, 1993).

Rigorous testing has been conducted to guarantee bit-for-bit (BFB) reproducibility for the entire data assimilation system, excluding the cost function minimizer, across runs with different MPI processes as well as runs with the same amount of MPI processes but different partitioning of the structured geographic grid. The quasi-Newton L-BFGS minimizer (Byrd *et al.*, 1995), employed for numerical minimization of the cost function, necessitates global matrix-vector multiplication, which precludes BFB reproducibility when domain decomposition is utilized. Divergences between executions stem from the non-associativity of floating-point operations, particularly floating-point summation within the minimizer. To specifically test and ensure that the parallelization of the rest of the assimilation system is fully reproducible, OceanVar2 offers the flexibility to

execute the minimizer serially (by aggregating variables from all domains and forcing a fixed order of operations) while the  
530 remaining code is parallelized using MPI domain decomposition. Extensive testing has demonstrated that using this specific serial option for the minimizer ensures BFB reproducibility. Moreover, even when the minimizer is executed in parallel, differences arising from various domain decompositions are statistically insignificant. Possible future work includes the introduction of a different minimizer suited for MPI parallelization.

Neglecting the differences arising from the parallel execution of the minimizer, the computational performance of the different  
535 experiments was evaluated in terms of minimizer iterations and code scalability. Figure 11 compares the number of iterations

required for the minimizer to converge in the various OceanVar2 experiments. The results are presented as a probability distribution, with statistics calculated based on the year of assimilation testing. Consistently, all experiments using the dynamic height operator converged with fewer iterations than those employing the barotropic model. The choice of the level-of-no-motion had a small effect, where the median increased from 24 to 25 iterations when using 350m or 150m instead of 1000m.

540 However, the primary insight is that the choice of the free surface operator formulation and the level-of-no-motion have negligible impact on the overall convergence speed. The median number of required iterations remains stable across all the experiments, varying only from 24 to 26 iterations. These differences between the median values appear insignificant when considered in the context of the wide day-to-day variability in the required convergence steps observed across the assimilation year, which spans approximately 12 to 45 iterations due to fluctuation in data availability and associated misfit values. The  
545 observed increase in the median number of iterations for the barotropic model schemes is likely due to the barotropic model's inherent physical complexity, which results in a more intricate optimization landscape for the minimizer to navigate.

To assess scalability, we limited the comparison to Exp-2 and Exp-4, as they had the same number of assimilated observations. We tested OceanVar2's performance with increasing numbers of cores. For a fixed number of cores, we performed experiments using eight different sets of observations and various decomposition strategies. (e.g., with 16 cores, we tested 4x4 and 8x2).

550 The results are shown in Fig.12. The model grid consisted of 1307x380x141 points along the x, y, and z directions, respectively. Analysis not shown here confirms that, in these experiments, the computational load across subdomains does not show significant differences based on geometric peculiarities, such as coastal proximity. Instead, the computational load of a subdomain, and thus the total execution time, was primarily determined by the overall number of assimilated observations. This finding is consistent with the global nature of the variational solver and the number of iterations required for convergence.

555 Notably, subdomains completely covered by land still required execution time comparable to the other domains. This occurred as a direct result of the land-sea mask not being used to optimize the domain decomposition, since the decomposition was based solely on the model grid geometry. The top panel of Fig.12 shows the CPU time per core as a function of the number of cores for Exp-2 and Exp-4. Consistently with Fig.11, the solid lines, representing the average CPU time, indicate that the difference between the dynamic height scheme (Exp-4) and the barotropic model (Exp-2) operators is minimal across all core  
560 counts, especially considering the large variability in run times, evidenced by the shaded areas indicating the maximum and minimum CPU time. More importantly, the speedup achieved is limited; increasing the cores from 1 to 36 yields a speedup of

approximately 8, corresponding to a parallel efficiency of about 22%. Furthermore, both experiments reach a performance plateau around 36 cores, with run time generally deteriorating at higher core counts due to increased communication overhead. To understand this limited parallel efficiency, the bottom panel of Fig.12 presents the scalability (expressed as a percentage of efficiency relative to the serial code) for the different **B** matrix operators. The diffusive filter operator acts as the main bottleneck. Its efficiency decreases rapidly at higher core counts and then exhibits a negligible further performances gain, stabilizing beyond 36 cores. This filter, which utilizes an implicit scheme, was implemented using a dimensional splitting approach. It is solved by means of LU decomposition of a tridiagonal matrix. We acknowledge that the implementation and parallelization strategy employed is suboptimal. This design choice simultaneously precludes optimal parallelization and results in the diffusive filter's extreme sensitivity to the geometry of the domain decomposition, which is the primary cause of the wide run-time variability shown in the top panel. This sequential process requires all processors associated with subsequent columns of subdomains to wait and then repeats along the second dimension (rows of subdomains). Although this crude, initial implementation constrains the overall parallel efficiency, the resulting total execution time remains within acceptable limits, rendering the software suitable for our current purposes. Future development will prioritize a more scalable parallelization strategy for this critical routine. Examining the remaining key operators reveals a strong general trend: while efficiency is high up to 36 cores, all remaining components experience a reduction in scaling performance thereafter. The barotropic model operator is the least affected by this change in scaling efficiency observed past 36 cores, maintaining the highest efficiency among the **B** operators.

580

## 7. Conclusions

This study describes recent developments of the OceanVar software which implements an incremental variational ocean data assimilation scheme. Key innovations compared to the previous schemes (DP08, Storto *et al.*, 2011, 2014) include the implementation and evaluation of two alternative solutions for the sea level operator, encompassing both barotropic model and 585 dynamic height operator. Furthermore, a diffusive operator has been adopted to model Gaussian horizontal covariances, replacing the recursive filter used in previous code versions. Finally, the geostrophic velocity operator is utilized for total velocity corrections, deviating from the DP08 approach and applied to both dynamic height and barotropic sea level operators. Furthermore, a method for filtering the tidal components of the background model fields is applied and tested allowing the 590 assimilation of SLA without tides, together with in-situ temperature and salinity data to produce analyses. These new and old features in OceanVar2 have been tested and compared for a regional implementation of the assimilation scheme in the Mediterranean Sea.

It has emerged that the barotropic operator is the only operator capable of consistently assimilating sea level anomaly data in shallow and deep ocean regions. Variable *alpha* and *beta* parameters in the linear equation of state (7) yielded minor differences in our experiments, however this assumption is likely not to be valid in global models.

595 The dynamic height operator, though easy to implement, has clear limitations. Requiring the definition of a spatially independent level-of-no-motion, it does not provide an optimal solution in domains with highly variable bottom topography and dynamics. For the Mediterranean Sea, a level-of-no-motion equal to 1000m is appropriate, as demonstrated by the quality of the corrections obtained with OceanVar2. However, this represents a significant limitation, as it excludes the assimilation of SLA observations in shallower areas. Decreasing the level-of-no-motion depth reveals the limitations of this approach. For  
600 shallower levels, the benefits of assimilating more data are offset by the loss of the quality of the corrections in deeper areas. The results are corroborated by the numerical instabilities arising when velocity corrections are applied in experiments with a level-of-no-motion shallower than 350m. The emergence of this critical instability provides a clear empirical demonstration that violating the inherent physical consistency assumptions used to construct the **B** matrix can lead to unstable solutions. Consequently, the optimal choice of operator depends on balancing the need for maximal observational impact with the  
605 requirement for numerical stability within the specific operational framework.

Although the barotropic model is inherently more complex and thus computationally more expensive than the dynamic height operator, its impact on the total run time is minor. This is because the semi-implicit scheme used to discretize the barotropic equations allows for large time-steps, significantly limiting the computational demand. This small computational overhead ensures that the barotropic model operator does not become the dominant factor in execution time.

610 The adopted solutions simplify the application of OceanVar2 in complex areas of the world's oceans. To our knowledge OceanVar2 is the only data assimilation software employing a barotropic model in its model background error covariance matrix. It is important to note that the current implementation of the barotropic model uses closed lateral boundary conditions. Its applicability is therefore limited to basins with a geometry that allows this approximation. In the future the barotropic model will be implemented also with open boundary conditions.

615 Regarding parallel performance, the diffusive filter overhead is currently the dominant computing factor, constraining the overall scalability. We acknowledge that the first implementation of the diffusive filter used in OceanVar2 limits the system's parallel efficiency. However, we chose this direction, replacing the previously used recursive filter, due to the superior flexibility and adaptability the diffusive filter offers for modeling horizontal error correlations.

The mathematical and algorithmic core of the system, first presented in DP08 and subsequently used in various operational  
620 forecasting frameworks, has been largely documented in several scientific works. Based on extensive testing, the revised code is stable and robust, with its performance results presented throughout this work. The present version, OceanVar2, is also open to the community.


## 625 Code availability

The OceanVar2.0 code is publicly available under a GPLv3 licence (<https://www.gnu.org/licenses/gpl-3.0.txt>) at <https://github.com/CMCC-Foundation/OceanVar2> (this manuscript) together with a user guide on compiling and running the code (Adani *et al.*, 2025, [https://github.com/CMCC-Foundation/OceanVar2/blob/main/doc/OceanVar\\_User\\_Manual.pdf](https://github.com/CMCC-Foundation/OceanVar2/blob/main/doc/OceanVar_User_Manual.pdf)). The code used in this manuscript is permanently archived at <https://doi.org/10.5281/zenodo.15593468> (Oddo *et al.*, 2025). A 630 test case can be downloaded at <https://github.com/CMCC-Foundation/MedFS831>. The ocean model used is based on the NEMO source code (version 4.2.0) is accessible on Zenodo. <https://doi.org/10.5281/zenodo.6334656> (Madec *et al.*, 2022).

### Author contribution

635 PO is the main author; he is the lead developer of OceanVar2. MA played a central role in the discussion; he led the developments; he wrote the code, and he performed all the experiments. FC contributed to the MPI parallelization, BFB reproducibility and debugging of the OceanVar2 code. AC contributed to the writing of the manuscript, and in incorporating some routines from previous code version. ACG computed the tidal constants used in all the experiments. EJ, AA, FM and IE were involved in the discussions and the definition of the development strategy. JP computed the EOFs used in all the 640 experiments. EC and SM played a central role in the discussion. NP contributed to the writing of the manuscript; she was active co-leading the scientific development.


## Figures




**Fig.1** Top panel: Model domain and bathymetry. Green and yellow dots indicate the position of the assimilated ARGO floats and XBT respectively. Red triangles indicate the subset of in-situ data assimilated during the single daily cycle of 11 August 2021. Bottom panel: an example of 21-day altimetry data. Satellite tracks in red and green are used in Figure 2 and 6 respectively. The solid, bold black tracks highlight the satellite altimetry data available during the single daily assimilation cycle of 11 August 2021. Three isobaths are drawn in both panels: 150, 350 and 1000m.

645

650




**Fig.2** An example of Sea Level Anomaly data along the red track of Figure 1. Blue lines indicate model free-run results, while  
655 black lines indicate observational data. In the upper panel the full signals are plotted. The dashed line indicates where SLA are  
in regions shallower than 1000m. The middle panel shows the SLA after the removal of the tidal signals, separately in the  
model and observations, and the along track averaged difference. The bottom panel shows the along track observational and  
model tidal signals.



660 **Fig.3** SLA, Temperature (red/blue) and Salinity (magenta/green) increments obtained from the different OceanVar2 configurations listed in Table1, starting from the same misfits. The SLA track used is drawn in green in Fig.1. For each experiment in the top panel there are the SLA increments where black dots indicate assimilated data, green dots indicate


rejected based on the coastal distance criteria, red dots indicate data rejected due to the level-of-no-motion or minimum depth (in case of the barotropic model).

665




**Fig.4** Time series of SLA error statistics for the entire year 2021. Blue ('free-run') and green ('EXP-1') lines indicate the statistics for the free-run results and the misfit of Exp-1 respectively. Top panel: unbiased root-mean-square error. Middle panel: correlation coefficient, in the bottom panel the standard deviation error is plotted.

670




**Fig.5** SLA statistics as a function of the ocean depth. Blue and green bars indicate the free-run and Exp-1 results respectively. In the top panel the number of observations used is also provided with dark bars.




675


**Fig.6** Vertical profiles of yearly averaged misfit statistics for salinity (top panels) and temperature (bottom panels). From left to right: *uRMSE*, *ME*, *CC* and *SDE*. Blue and green lines indicate free-run and Exp-1 results respectively.




680 **Fig.7**  $S_uRMSE$  and  $CC$  indices for the SLA misfit errors for the different experiments.  $S_uRMSE$  is computed w.r.t the free-run. Top two panels: time-series. Bottom panels: average indices for bathymetry classes.



685 **Fig.8**  $S_{uRMSE}$  and  $S_{CC}$  indices for % of improvement of the SLA errors for the different experiments w.r.t Exp-1. Top two  
panels: time-series. Bottom panels: average indices for bathymetry classes.




**Fig.9** Second set of experiments SLA time series error statistics. Top panel: unbiased root-mean-square error. Middle panel: correlation coefficient. Bottom panel: standard deviation error. Colour code is provided in the middle panel legend.



690

**Fig.10** SLA statistics as a function of the ocean depth. Top panel: unbiased root-mean-square error. Middle panel: correlation coefficient. Bottom panel: standard deviation error. Colour code is provided in the top panel legend.

695



700 **Fig.11** Probability density function (Y axis) of the number of iteration (X axis) needed for the OceanVar2 minimization algorithm to converge. For each experiment, the lighter and darker shaded areas correspond to the 90% and 50% of the events respectively. The vertical lines are the medians of each distribution. Exp-1 to Exp-6 from top to bottom.




Fig.12 Parallel performance and routine-level scalability. Top panel: CPU time per core (minutes) as a function of the number of cores for Exp-2 and Exp-4. Solid lines represent the average CPU time, while shaded areas indicate the maximum and minimum CPU time observed for each core count. Bottom panel: Scalability analysis of individual Background Error

705 Covariance matrix components. The Y-axis is the percentage of reduction time with respect to the serial (single core) execution. Solid lines indicate the minimum execution time and dashed lines indicate the maximum execution time observed across different days and domain decompositions for both the tangent linear and adjoint routines. The solid black line indicates the linear reduction with the number of cores, and coloured lines show the performance of individual routines (component colours are defined in the figure legend).

## References

- Adani, M., A. Aydogdu, A. Cipollone, E. Jansen, F. Carere, P. Oddo. 2025. OceanVar User Manual. [https://github.com/CMCC-Foundation/OceanVar2/blob/main/doc/OceanVar\\_User\\_Manual.pdf](https://github.com/CMCC-Foundation/OceanVar2/blob/main/doc/OceanVar_User_Manual.pdf)
- Adani, M., S. Dobricic, and N. Pinardi, 2011: Quality Assessment of a 1985–2007 Mediterranean Sea Reanalysis. *J. Atmos. Oceanic Technol.*, 28, 569–589, <https://doi.org/10.1175/2010JTECHO798.1>.
- Anderson, J. L., T. Hoar, K. Raeder, H. Liu, N. Collins, R. Torn and A. Arellano, 2009 The Data Assimilation Research Testbed: A Community Facility. *Bulletin of the American Meteorological Society*, 90, 1283-1296, doi:10.1175/2009BAMS2618.1
- Arbic, B.K., 2022: Incorporating tides and internal gravity waves within global ocean general circulation models: A review. *Progress in Oceanography* 206, 102824.
- Barthélémy, S., Brajard, J., Bertino, L. and Counillon, F., 2022. Super-resolution data assimilation. *Ocean Dynamics*, 72(8), pp.661-678. Vancouver
- Beauchamp, M., Febvre, Q., Georghenthum, H., and Fablet, R.: 4DVarNet-SSH: end-to-end learning of variational interpolation schemes for nadir and wide-swath satellite altimetry, *Geosci. Model Dev.*, 16, 2119–2147, <https://doi.org/10.5194/gmd-16-2119-2023>, 2023.
- Broccoli M. & Cipollone A. 2025: Assimilation of diurnal satellite retrieval of sea surface temperature with convolutional neural network, *Mach. Learn.: Earth* 1 015007, <https://doi.org/10.1088/3049-4753/adfb7e>
- Byrd, R.H., Lu, P., Nocedal, J., Zhu, C., 1995. A limited memory algorithm for bound constrained optimization. *SIAM Journal on Scientific Computing* 16, 1190–1208.
- Cao A.-Z., Wang D.-S., Lv X.-Q; Harmonic Analysis in the Simulation of Multiple Constituents: Determination of the Optimum Length of Time Series; *Journal of Atmospheric and Oceanic Technology*, 32(5), 1112–1118; 2015
- Carrassi, A., Bocquet, M., Bertino, L., and Evensen, G.: Data assimilation in the geosciences: An overview of methods, issues, and perspectives, *WIREs Climate Change*, 9, e535, <https://doi.org/10.1002/wcc.535>, 2018.
- Cipollone, A., Banerjee, D. S., Iovino, D., Aydogdu, A., & Masina, S. Bivariate sea-ice assimilation for global-ocean analysis–reanalysis, *Ocean Sci.*, 19, 1375–1392, <https://doi.org/10.5194/os-19-1375-2023>. 2023.
- Clementi, E., Drudi, M., Aydogdu, A., Moulin, A., Grandi, A., Mariani, A., Goglio, A. C., Pistoia, J., Miraglio, P., Lecci, R., Palermo, F., Coppini, G., Masina, S., & Pinardi, N. (2023). Mediterranean Sea Physical Analysis and Forecast (CMS MED-Physics, EAS8 system) (Version 1) [Data set]. Copernicus Marine Service (CMS). [https://doi.org/10.25423/CMCC/MEDSEA\\_ANALYSISFORECAST\\_PHY\\_006\\_013\\_EAS8](https://doi.org/10.25423/CMCC/MEDSEA_ANALYSISFORECAST_PHY_006_013_EAS8)
- Ciliberti, S.A.; Jansen, E.; Coppini, G.; Peneva, E.; Azevedo, D.; Causio, S.; Stefanizzi, L.; Creti, S.; Lecci, R.; Lima, L.; et al. The Black Sea Physics Analysis and Forecasting System within the Framework of the Copernicus Marine Service. *J. Mar. Sci. Eng.* 2022, 10, 48. <https://doi.org/10.3390/jmse10010048>
- Cohn, S. E., 1997: An introduction to estimation theory. *J. Meteor. Soc. Japan*, 75, 257–288

745 • Coppini, G., Clementi, E., Cossarini, G., Salon, S., Korres, G., Ravdas, M., Lecci, R., Pistoia, J., Goglio, A. C., Drudi, M., Grandi, A., Aydogdu, A., Escudier, R., Cipollone, A., Lyubartsev, V., Mariani, A., Creti, S., Palermo, F., Scuro, M., Masina, S., Pinardi, N., Navarra, A., Delrosso, D., Teruzzi, A., Di Biagio, V., Bolzon, G., Feudale, L., Coidessa, G., Amadio, C., Brosich, A., Miró, A., Alvarez, E., Lazzari, P., Solidoro, C., Oikonomou, C., and Zacharioudaki, A.: The Mediterranean Forecasting System – Part 1: Evolution and performance, *Ocean Sci.*, 19, 1483–1516, <https://doi.org/10.5194/os-19-1483-2023>, 2023.

750 • De Mey, P. and A.R. Robinson, 1987: Assimilation of altimetric fields in a limited area quasigeostrophic model. *J. Phys. Oceangr.*, 17, 2280-2293.

755 • Derber, J., Rosati, A., 1989. A global oceanic data assimilation system. *Journal of Physical Oceanography* 19, 1333–1347.

760 • Desroziers, G., Berre, L., Chapnik, B. and Poli, P. (2005), Diagnosis of observation, background and analysis-error statistics in observation space. *Q.J.R. Meteorol. Soc.*, 131: 3385-3396. <https://doi.org/10.1256/qj.05.108>

765 • Dhomps, A.-L., S. Guinehut, P.-Y. Le Traon, and G. Larnicol, 2011. A global comparison of Argo and satellite altimetry observations, *Ocean Sci.*, 7, 175–183, doi:10.5194/os-7-175-2011

770 • Dobricic, S., Pinardi, N., Adani, M., Tonani, M., Fratianni, C., Bonazzi, A., Fernandez, V., 2007. Daily oceanographic analyses by the Mediterranean basin scale assimilation system. *Ocean Sciences* 3, 149–157.

775 • Dobricic, S. and Pinardi, N.: An oceanographic three-dimensional variational data assimilation scheme, *Ocean Model.*, 22, 89–105, 2008.

• Dobricic, S., Dufau, C., Oddo, P., Pinardi, N., Pujol, I., and Rio, M.-H.: Assimilation of SLA along track observations in the Mediterranean with an oceanographic model forced by atmospheric pressure, *Ocean Sci.*, 8, 787–795, <https://doi.org/10.5194/os-8-787-2012>, 2012.

• Dobricic, S., Pinardi, N., Adani, M., Bonazzi, A., Fratianni, C. and Tonani, M. (2006). Mediterranean Forecasting System: An improved assimilation scheme for sea-level anomaly and its validation. *Q. J. R. Meteorol. Soc.* 131: 3627–3642, doi: 10.1256/qj.05.100.

• Dobricic, S., Wikle, C. K., Milliff, R. F., Pinardi, N., Berliner, L. M., 2015. Assimilation of oceanographic observations with estimates of vertical background-error covariances by a Bayesian hierarchical model *Quarterly Journal of the Royal Meteorological Society*, 141, 182-194, doi: <http://dx.doi.org/10.1002/qj.2348>

• Escudier, R., Clementi, E., Cipollone, A., Pistoia, J., Drudi, M., Grandi, A., Lyubartsev, V., Lecci, R., Aydogdu, A., Delrosso, D., et al.: A high resolution reanalysis for the Mediterranean Sea, *Frontiers in Earth Science*, 9, 702285, 2021.

• Foreman M.G.G.; *Manual for tidal heights analysis and prediction*. Pacific Marine Science Report 77-10, Institute of Ocean Sciences, Patricia Bay, Sidney, BC, 97pp; 1977

- Foreman M.G.G.; Manual for tidal currents analysis and prediction. Pacific Marine Science Report 78-6, Institute of Ocean Sciences, Patricia Bay, Sidney, BC, 57pp; 1978
- Hayden, C.M., Purser, R.J., 1995. Recursive filter objective analysis of meteorological fields: applications to NESDIS operational processing. *Journal of Applied Meteorology* 34, 3–15.
- 780 • Hoffman, J.D., Hoffman, J.D., & Frankel, S. (2001). Numerical Methods for Engineers and Scientists (2nd ed.). CRC Press. <https://doi.org/10.1201/9781315274508>
- Kwizak, M., Robert, A.J., 1971. A semi-implicit scheme for grid point atmospheric models of the primitive equations. *Monthly Weather Review* 99, 32–36.
- 785 • Le Traon, P.-Y., Dibarboire, G., Lellouche, J.-M., Pujol, M.-I., Benkiran, M., Drevillon, M., Drillet, Y., Faugère, Y., and Remy, E.: Satellite altimetry and operational oceanography: from Jason-1 to SWOT, *Ocean Sci.*, 21, 1329–1347, <https://doi.org/10.5194/os-21-1329-2025>, 2025.
- Lima, L., Ciliberti, S.A., Aydoğdu, A., Masina, S., Escudier, R., Cipollone, A., Azevedo, D., Causio, S., Peneva, E., Lecci, R., Clementi, E., Jansen, E., Ilicak, M., Creti, S., Stefanizzi L., Palermo, F. and Coppini G., 2021. Climate Signals in the Black Sea From a Multidecadal Eddy-Resolving Reanalysis. *Front. Mar. Sci.* 8:710973. <https://doi.org/10.3389/fmars.2021.710973>.
- 790 • Lorenc, A.C., 1992. Iterative analysis using covariance functions and filters. *Quarterly Journal of the Royal Meteorological Society* 118, 569–591.
- Lorenc, A.C., 1997. Development of an operational variational assimilation scheme. *Journal of the Meteorological Society of Japan* 75, 339346.
- 795 • Madec, G. and the NEMO System Team, 2023. NEMO Ocean Engine Reference Manual, Zenodo, <https://doi.org/10.5281/zenodo.8167700>
- Madec, G., and the NEMO System Team, 2022. NEMO ocean engine. In *Scientific Notes of IPSL Climate Modelling Center (v4.2, Number 27)*. Zenodo. <https://doi.org/10.5281/zenodo.6334656>
- 800 • Martin, M. J., Hoteit, I., Bertino, L., and Moore, A. M.: Data assimilation schemes for ocean forecasting: state of the art, in: *Ocean prediction: present status and state of the art (OPSR)*, edited by: Álvarez Fanjul, E., Ciliberti, S. A., Pearlman, J., Wilmer-Becker, K., and Behera, S., Copernicus Publications, State Planet, 5-opsr, 9, <https://doi.org/10.5194/sp-5-opsr-9-2025>, 2025.
- 805 • Massart, S., B. Pajot, A. Piacentini, and O. Pannekoucke, 2010: On the Merits of Using a 3D-FGAT Assimilation Scheme with an Outer Loop for Atmospheric Situations Governed by Transport. *Mon. Wea. Rev.*, 138, 4509–4522, <https://doi.org/10.1175/2010MWR3237.1>.
- Moore, A.M., H.G. Arango, G. Broquet, B.S. Powell, A.T. Weaver, and J. Zavala-Garay, 2011: The Regional Ocean Modeling System (ROMS) 4-dimensional variational data assimilation systems, Part I - System overview and formulation, *Progress in Oceanography*, 91, 34-49.

810 • Murphy, A. H., 1988: Skill Scores Based on the Mean Square Error and Their Relationships to the Correlation Coefficient. *Mon. Wea. Rev.*, 116, 2417–2424.

815 • Nerger, L., Hiller, W., Software for ensemble-based data assimilation systems—Implementation strategies and scalability, *Computers & Geosciences*, Volume 55, 2013, Pages 110-118, ISSN 0098-3004, <https://doi.org/10.1016/j.cageo.2012.03.026>.

820 • Nilsson, J. A. U., Dobricic, S., Pinardi, N., Poulain, P.-M., and Pettenuzzo, D., 2012. Variational assimilation of Lagrangian trajectories in the Mediterranean ocean Forecasting System. *Ocean Sci.*, 8, 249-259, doi:10.5194/os-8-249-2012

825 • Oddo, P. Storto, A. Dobricic, S. Russo, A. Lewis, C. Onken, R. Coelho, E., A hybrid variational-ensemble data assimilation scheme with systematic error correction for limited-area ocean models, *Ocean Science*, 12, 2016, 5, 1137-1153, doi:10.5194/os-12-1137-2016

830 • Oddo, P., Adani, M., Carere, F., Cipollone, A., Goglio, A. C., Jansen, E., Aydogdu, A., Mele, F., Epicoco, I., Pistoia, J., Clementi, E., Pinardi, N., & Masina, S. (2025). OceanVar2.0 (2.0). Zenodo. <https://doi.org/10.5281/zenodo.15593468>

835 • Pawlowicz R., Beardsley B., and Lentz S.; Classical Tidal Harmonic Analysis Including Error Estimates in MATLAB using T TIDE; *Computers and Geosciences*, 28, 929-937; 2002

840 • Purser, R.J., Wu, W.-S., Parish, D.F., Roberts, N.M., 2003. Numerical aspects of the application of recursive filters to variational statistical analysis. Part II: spatially inhomogeneous and anisotropic covariances. *Monthly Weather Review* 131, 1524–1535.

845 • Purser, R.J., Rancic, M. and Pondeca, M.S.D., 2022. The Multigrid Beta Function Approach for Modeling of Background Error Covariance in the Real-Time Mesoscale Analysis (RTMA). *Monthly Weather Review*, 150(4), pp.715-732. Vancouver

850 • Rio, M.-H., Pascual, A., Poulain, P.-M., Menna, M., Barceló, B., and Tintoré, J.: Computation of a new mean dynamic topography for the Mediterranean Sea from model outputs, altimeter measurements and oceanographic in situ data, *Ocean Sci.*, 10, 731–744, <https://doi.org/10.5194/os-10-731-2014>, 2014.

855 • Sanchez de la Lama, M., J. H. La Casce, and H. K. Fuhr, 2016: The vertical structure of ocean eddies. *Dyn. Stat. Climate Syst.*, 1, 1–16, <https://doi.org/10.1093/climsys/dzw001>.

860 • Storto, A., Masina, S. and Navarra, A. (2016), Evaluation of the CMCC eddy-permitting global ocean physical reanalysis system (C-GLORS, 1982–2012) and its assimilation components. *Q.J.R. Meteorol. Soc.*, 142: 738–758. <https://doi.org/10.1002/qj.2673>

865 • Storto, A., S. Dobricic, S. Masina, and P. Di Pietro, 2011: Assimilating Along-Track Altimetric Observations through Local Hydrostatic Adjustment in a Global Ocean Variational Assimilation System. *Mon. Wea. Rev.*, 139, 738–754, <https://doi.org/10.1175/2010MWR3350.1>.

- Storto, A., Variational quality control of hydrographic profile data with non-Gaussian errors for global ocean variational data assimilation systems, *Ocean Modelling*, Volume 104, 2016, Pages 226-241, ISSN 1463-5003, <https://doi.org/10.1016/j.ocemod.2016.06.011>.

845

- Storto, A., Oddo, P., Cipollone, A., Mirouze, I., Lemieux-Dudon, B. Extending an oceanographic variational scheme to allow for affordable hybrid and four-dimensional data assimilation. *Ocean Modelling*, 128 (2018), pp. 67-86, 10.1016/j.ocemod.2018.06.005
- Storto A, Oddo P. Optimal Assimilation of Daytime SST Retrievals from SEVIRI in a Regional Ocean Prediction System. *Remote Sensing*. 2019; 11(23):2776. <https://doi.org/10.3390/rs11232776>

850

- Storto, A., Falchetti, S., Oddo, P., Jiang, Y.-M., & Tesei, A. (2020). Assessing the impact of different ocean analysis schemes on oceanic and underwater acoustic predictions. *Journal of Geophysical Research: Oceans*, 125, e2019JC015636. <https://doi.org/10.1029/2019JC015636>
- Storto, A., G. De Magistris, S. Falchetti, and P. Oddo, 2021: A Neural Network-Based Observation Operator for Coupled Ocean-Acoustic Variational Data Assimilation. *Mon. Wea. Rev.*, 149, 1967–1985, <https://doi.org/10.1175/MWR-D-20-0320.1>.

855

- Teruzzi, A., S. Dobricic, Solidoro, C., Cossarini, G. A 3-D variational assimilation scheme in coupled transport-biogeochemical models: forecast of Mediterranean biogeochemical properties. *J. Geophys. Res.: Oceans*, 119 (1) (2014), pp. 200-217 URL <https://doi.org/10.1002/2013JC009277>.
- Teruzzi, A., Bolzon, G., Salon, S., Lazzari, P., Solidoro, C., Cossarini, G. Assimilation of coastal and open sea biogeochemical data to improve phytoplankton simulation in the Mediterranean Sea. *Ocean Model.*, 132 (2018), pp. 46-60
- The MPI Forum, CORPORATE (November 15–19, 1993). "MPI: A Message Passing Interface". Proceedings of the 1993 ACM/IEEE conference on Supercomputing. *Supercomputing '93*. Portland, Oregon, USA: ACM. pp. 878–883. doi:10.1145/169627.169855

860

- Van Leeuwen, P.J., Künsch, H.R., Nerger, L., Potthast, R. and Reich, S., 2019. Particle filters for high-dimensional geoscience applications: A review. *Quarterly Journal of the Royal Meteorological Society*, 145(723), pp.2335-2365.
- Weaver A., Courtier P. Correlation modeling on the sphere using a generalized diffusion equation Q. *J. R. Meteorol. Soc.*, 122 (2001), pp. 535-561.
- Weaver, A. T., Vialard, J. and Anderson, D. L. T. 2003 Three- and four-dimensional variational assimilation with an ocean general circulation model of the tropical Pacific Ocean. Part 1: formulation, internal diagnostics and consistency checks. *Mon. Weather Rev.*, 131, 1360–1378
- Zavala-Romero, O., Bozec, A., Chassagnet, E. P., and Miranda, J. R.: Convolutional neural networks for sea surface data assimilation in operational ocean models: test case in the Gulf of Mexico, *Ocean Sci.*, 21, 113–132, <https://doi.org/10.5194/os-21-113-2025>, 2025.

865

870