

1 **Toward a typology of river functioning: a**  
2 **comprehensive study of the POM-particulate organic**  
3 **matter composition at the multi-rivers scale**

4  
5 **Férechiche** Florian Férechiche,<sup>1</sup> **Liénart** Camilla- Liénart,<sup>1</sup> **Charlier** Karine- Charlier,<sup>1</sup> **Deborde** Jonathan-  
6 Deborde,<sup>2,3</sup> **Giraud** Mélanie- Giraud,<sup>4</sup> **Kerhervé** Philippe- Kerhervé,<sup>5</sup> **Polsenaere** Pierre- Polsenaere,<sup>1,3</sup>  
7 **Savoye** Nicolas Savoye.<sup>1\*</sup>

8 <sup>1</sup> Univ. Bordeaux, CNRS, EPHE, Bordeaux INP, UMR 5805 EPOC, F-33600 Pessac, France

9 <sup>2</sup> Univ. Pau & Pays Adour, CNRS, E2S UPPA – MIRA, UMR 5254 IPREM, F-64000 Pau, F-64600 Anglet, France

10 <sup>3</sup> Ifremer, COAST, F-17390 La Tremblade, France

11 <sup>4</sup> MNHN, CRESCO, Station Marine de Dinard, F-35800 Dinard, France

12 <sup>5</sup> Univ. Perpignan, CNRS, UMR 5110 CEFREM, F-66860 Perpignan, France

13 \*Correspondence to: Nicolas Savoye ([nicolas.savoye@u-bordeaux.fr](mailto:nicolas.savoye@u-bordeaux.fr)) (+33)556223916

14 Université de Bordeaux - UMR 5805 EPOC - Station marine d'Arcachon - 2 rue du Professeur Jolyet - 33120  
15 Arcachon

16  
17  
18 Keywords: river-estuary interface; particulate organic matter; stable isotopes; multi-ecosystems  
19 study.

20 **Abstract**

21 In riverine systems, particulate organic matter (POM) originates from various sources, each  
22 having its proper dynamics related to production, decomposition, transport and burial. ~~There~~  
23 ~~is resulting in~~ a significant ~~amount of~~ spatiotemporal heterogeneity in the POM pool. The  
24 current study, based on C and N elemental and isotopic ratios, applies Bayesian mixing models  
25 associated with statistical multivariate analyses to 1) quantify and examine relationships  
26 between POM composition and environmental forcings, and 2) draw a typology of river  
27 functioning based on POM composition and its seasonal dynamics.

28 Twenty-three ~~temperate~~ rivers ~~of temperate climate~~ accounting for a large diversity of  
29 environmental conditions were sampled fortnightly to monthly for one to seven years at the ~~ir~~  
30 River-Estuary Interface (REI). Phytoplankton and labile terrestrial material ~~were~~  
31 ~~present occurred~~ in all rivers, ~~contrary to whereas~~ sewage and refractory terrestrial material ~~that~~  
32 were present in only a few. At the twenty-three ~~studied~~ rivers scale, ~~POM sources are strongly~~  
33 ~~related to watershed characteristics~~, phytoplankton ~~dominance was being~~ associated with  
34 agricultural surfaces ~~and, while~~ labile terrestrial material ~~was linked to organic-rich leached~~ soil  
35 ~~and refractory terrestrial matter to steep catchments with little soil organic carbon content and~~  
36 ~~erosion rate. Overall, Seasonal dynamics were primarily driven by phytoplankton growth, river~~  
37 ~~discharge (labile terrestrial material), and sediment resuspension (refractory terrestrial~~  
38 ~~material). variations of phytoplankton, labile and refractory terrestrial material were mainly~~  
39 ~~related to drivers of phytoplankton growth, river flow, and sediment resuspension, respectively.~~

40 A statistical regionalization defined four ~~types of~~ river ~~dynamic types~~: (1) systems whose  
41 POM is dominated by labile terrestrial material ~~yeaaer-round all year long~~; (2) systems whose  
42 POM is composed of labile and refractory terrestrial material, in addition to phytoplankton,  
43 ~~with showing~~ variable seasonality ~~according to rivers~~; systems whose POM is composed of  
44 phytoplankton and labile terrestrial material (3) without and (4) with pronounced seasonality.

45 This work offers a comprehensive understanding of POM composition, ~~spatio-temporal~~  
46 dynamics and ~~drivers controlling factors~~ at the REI in temperate climates, complementing  
47 similar work dedicated to coastal systems. Future work dedicated to estuaries is called to get a  
48 comprehensive understanding of POM composition, dynamics and drivers along the Land-  
49 Ocean Aquatic Continuum.

50 ~~This study examines particulate organic matter (POM) composition and dynamics in 23~~  
51 ~~temperate rivers. Carbon and nitrogen isotope analysis revealed four river types based on~~  
52 ~~dominant POM sources (phytoplankton, terrestrial material). Watershed characteristics~~  
53 ~~influence POM composition while seasonal variations in river flow and sediment resuspension~~  
54 ~~drive POM dynamics. This study improves the understanding of river systems and calls for~~  
55 ~~further studies exploring downstream estuarine functioning.~~

56 **1. Introduction**

57 The River-Estuary Interface (REI) is a crucial biogeochemical interface for understanding the  
58 transition between continental and coastal systems, beginning at estuaries, because of its key  
59 location within the Land-Ocean Aquatic Continuum (LOAC) (Bate et al., 2002). Indeed, rivers  
60 then estuaries are important filters for matters received from land, transporting and transforming  
61 organic matter and nutrients along their courses (Bouwman et al., 2013; Dürr et al., 2011;  
62 Middelburg and Herman, 2007). These processes are fundamental in understanding global  
63 biogeochemical cycles (Regnier et al., 2013), as these matters directly fuel coastal ocean trophic  
64 networks (Dagg et al., 2004). However, in a Human-impacted world, anthropogenic activities  
65 and disturbances can modify natural matter fluxes. For example, damming rivers directly  
66 impacts nutrient flows (Wang et al., 2022) and sediment transportation (Kang et al., 2021).  
67 Indirectly, land use in river basins can lead to changes in the river matter quality (Lambert et  
68 al., 2017).

69 In aquatic systems, particulate organic matter (POM), i.e., non-mineral particles, is composed  
70 of different sources that originate from different compartments: phytoplankton, macrophytes  
71 from the aquatic systems, as well as ~~soil particles and~~ plant litter, ~~soil and~~ ~~petrogenic particles~~  
72 from terrestrial compartments and even treated and untreated anthropogenic organic matter (Ke  
73 et al., 2019; Sun et al., 2021; Zhang et al., 2021). Depending on its composition, POM exhibits  
74 different levels of lability, i.e., different levels of biogeochemical reactivity and bioavailability.  
75 For instance, phytoplankton is usually considered ~~mainly highly~~ labile, and thus highly  
76 biogeochemically reactive and bioavailable for primary consumers, while terrestrial POM is  
77 usually considered ~~mainly more and more~~ refractory ~~through degradation -processes and~~, thus  
78 lightly biogeochemically reactive and poorly bioavailable for the food webs (Brett et al., 2017;  
79 David et al., 2005; Etcheber et al., 2007). In other words, the determination and quantification  
80 of POM composition (i.e., the relative proportion of each source composing the POM) allow a  
81 better understanding of biogeochemical cycles and trophic ecology in aquatic systems (e.g.,  
82 Grunicke et al., 2023; Minaudo et al., 2016). Nevertheless, POM composition and concentration  
83 are not only involved in biogeochemical and biological processes (e.g., primary production,  
84 remineralization, feeding) but also undergo other processes inside and at the interface of the  
85 aquatic compartment (Canuel and Hardison, 2016). River ~~hydrodynamics hydrology~~ is ~~one of~~  
86 ~~the main drivers of a key factor controlling~~ POM composition and concentration. ~~The~~  
87 ~~geological and soil characteristics of each catchment, together with climatic conditions, shape~~  
88 ~~the erosional processes,~~ leading to great variabilities in ~~hydrodynamics,~~ terrestrial material  
89 quality and quantity and phytoplankton growth conditions (Dalzell et al., 2007; Hilton et al.,  
90 2010; Lebreton et al., 2016), possibly leading This variability leads to changes shifts in POM  
91 source origins (Arellano et al., 2019; Barros et al., 2010). ~~Additionaly~~~~Additionally~~~~lso, changes~~  
92 ~~in anthropic pressures~~~~anthropogenic disturbances~~ can change directly or indirectly affect  
93 seasonal as well as long-term patterns of POM ~~seasonal~~ composition and concentration and

94 ~~their seasonal variations patterns, like a~~ For instance, a decrease in nutrient load ~~affects~~  
95 ~~phytoplankton production and biomass (Minaudo et al., 2015), or affect year-round a river's~~  
96 ~~biogeochemistry by altering stable controlling factors (e.g., agricultural surfaces altering the~~  
97 ~~soil properties and erosion and consequently soil particle export to rivers, or damming altering~~  
98 ~~the river hydromorphology and consequently particle dynamics and export (Kang et al., 2021;~~  
99 ~~Zhang et al., 2021), etc.~~

100 This dependency of POM composition and concentration ~~on to~~ physical, biogeochemical and  
101 biological processes and their responses to environmental conditions and characteristics (Bonin  
102 et al., 2019; Falkowski et al., 1998; Field et al., 1998; Galeron et al., 2017; Goñi et al., 2009;  
103 Lebreton et al., 2016) may lead to distinguishing different types ~~esology~~ of rivers, i.e., the  
104 likeliness of rivers to carry preferential sources. For instance, highly turbid systems are more  
105 likely to carry refractory materials (Savoye et al., 2012), while eutrophicated rivers carry high  
106 biomass of phytoplankton (Hounshell et al., 2022; Minaudo et al., 2015), and contrasting  
107 processes can lead to a mixture between different detrital sources, as soil matter vs. fresh  
108 terrestrial plants (Ogrinc et al., 2008). However, to date, no study clearly determined typologies  
109 of river ~~dynamicss~~ based on POM composition and its seasonal variability.

110 To distinguish POM sources and quantify their contribution to POM composition, different  
111 tools such as elemental and isotopic ratios, pigments or specific compounds like fatty acids or  
112 alkanes can be used (e.g., Chevalier et al., 2015; Liénart et al., 2020, 2017; Savoye et al., 2012).  
113 Elemental and isotopic ratios are usually considered robust and allow the quantification of POM  
114 composition in this kind of study (e.g., Liénart et al., 2016; Onstad et al., 2000; Wang et al.,  
115 2021). Indeed, they usually allow the discrimination of, e.g., riverine phytoplankton, terrestrial  
116 POM and wastewater POM (Ke et al., 2019) and they can be used for running mixing models  
117 that quantify the proportion of the different sources ~~in to~~ a POM mixture (Parnell et al., 2013).  
118 However, studies using mixing models for quantifying POM composition in river systems are  
119 still ~~quite~~ scarce (e.g., Ferchiche et al., 2025, 2024; Kelso and Baker, 2022, 2020; Zhang et al.,  
120 2021).

121 Within the scope of better understanding the role of the LOAC in modifying matter fluxes and  
122 quality, the present study gathered published data and results from 23 rivers at the river-estuary  
123 interface with the aim of 1) quantifying the POM composition ~~of~~ each river, 2) describing the  
124 seasonal variations of this composition, 3) determining the drivers of the seasonal variability  
125 within each river and the spatial variability among the 23 rivers, and then 4) determining a  
126 typology of river ~~dynamicss~~ according to their POM composition and dynamics. This study is  
127 the first ~~that~~ precisely quantify POM composition in numerous and various temperate river  
128 systems ~~in a world region (here, the Western Europe)~~ and classify river types according to POM  
129 composition and dynamics.


## 130 2. Materials and methods

131 Twenty-three temperate rivers were studied at their river-estuary interface (i.e., right upstream  
132 of the tidal influence). All the data come from published studies or national open databases. To  
133 minimize the heterogeneity of the datasets in terms of sampling strategy, we have considered  
134 for this study the datasets only when 1) C/N ratio along with isotopic ratio of carbon and/or  
135 nitrogen were available, 2) particulate matter characteristics like, suspended particulate matter  
136 (SPM), particulate organic carbon (POC), particulate nitrogen (PN) or; chlorophyll *a* (chl *a*)  
137 were also available, 3) datasets exhibited at least a monthly temporal resolution for one full  
138 year. When needed, published datasets were completed and harmonized thanks to national  
139 databases.

140 **2.1. Study sites**

141 The studied rivers and associated watersheds are all located in France (except the upper basin  
142 of the Rhône River) and distributed in all regions of the mainland. Three, fifteen and five of  
143 these rivers flow into the English Channel, the Atlantic Ocean and the Mediterranean Sea (Fig.  
144 1). Even if located in a somewhat restricted area (Western Europe) at the global scale, they  
145 encompass large gradients of environmental characteristics (Tab. 1) for a temperate climate.  
146 For instance, the Loire River is one of the largest in Europe (length: 1006 km; watershed:  
147 117,356 km<sup>2</sup>), while the littlest studied river is a very small stream of the Arcachon lagoon  
148 (length: 3 km; watershed: 18 km<sup>2</sup>). They encompass large gradients of river flow (annual mean:  
149 0.3 m<sup>3</sup>/s – 1572 m<sup>3</sup>/s), turbidity (SPM annual mean: 2.7 mg/l – 40.9 mg/l) and trophic status  
150 (from oligotrophic to eutrophic rivers; chlorophyll *a* annual mean: 0.4 µg/l – 57.1 µg/l). At last,  
151 they undergo a gradient of anthropic pressures as illustrated by the proportion of artificial  
152 surfaces (0.1 % – 5.6 %) and agricultural areas (0 % – 86 %) in the watersheds (Fig. 1).

153



154

155 **Figure 1** Studied rivers (thick blue lines), sampling locations (black stars) and watersheds (thin  
 156 black lines), including the main land uses (red, yellow and green *colorscolours*). 1: Seine; 2:  
 157 Orne; 3: Rance; 4: Elorn; 5: Aulne; 6: Loire; 7: Sèvre niortaise; 8: Charente; 9: Seudre; 10:  
 158 Canal du Porge; 11: Cirès; 12: Milieu; 13: Lanton; 14: Renet; 15: Tagon; 16: Leyre; 17: Canal  
 159 des Landes; 18: Adour; 19: Têt; 20: Aude; 21: Orb; 22: Hérault; 23: Rhône.

160 **Table 1** Overview of river samplings and characteristics. Values are given as annual mean over the study period for river flow, temperature,  
 161 suspended particulate matter (SPM) and chlorophyll *a* (chl *a*). Id: identification number; Number: number of sampling dates. River types were  
 162 defined within the scope of the present study (see section 3.4).

River	Id	River type	Sampled period	Sampling Periodicity	Num ber	Latitud e	Longitu de	River length (km)	Catchment area (km <sup>2</sup> )	River flow (m <sup>3</sup> /s)	Water temperat ure (°c)	SPM (mg/l)	Chl <i>a</i> (µg/l)	References
Seine	1	IV	06/2014 to 06/2015	monthly	13	49.3067	1.2425	774	79000	496	15.0	21	2.8	Liénart et al., 2017, 2018
Orne	2	III	06/2014 to 06/2015	monthly	13	49.1797	-0.3491	169	2932	16	14.5	11	1.8	Liénart et al., 2017, 2018
Rance	3	IV	06/2014 to 05/2015	monthly	12	48.4916	-2.0014	103	1195	1.37	15.1	21	57.1	Liénart et al., 2017, 2018
Elorn	4	IV	01/2014 to 06/2015	monthly	17	48.4505	-4.2483	56	385	6	12.3	16	3.0	Liénart et al., 2017, 2018
Aulne	5	IV	01/2014 to 06/2015	monthly	17	48.2127	-4.0944	144	1875	30	14.4	7	3.3	Liénart et al., 2017, 2018
Loire	6	IV	10/2009 to 07/2012	bi-monthly	67	47.3920	-0.8604	1006	117356	630	14.1	19	18.7	Ferchiche et al., 2024
Sèvre	7	IV	03/2014 to 03/2015	monthly	13	46.3153	-1.0039	158	3650	3.72	15.7	13	3.8	Liénart et al., 2017, 2018
Charente	8	III	03/2014 to 03/2015	monthly	13	45.8680	-0.7131	381	9855	68	15.1	13	1.3	Liénart et al., 2017, 2018
Seudre	9	I	03/2014 to 09/2015	monthly	15	45.6740	-0.9331	68	855	1.81	14.3	17	0.5	Liénart et al., 2017, 2018
Porge	10	III	01/2008 to 02/2009	monthly	14	44.7898	-1.1612	57	222	3.48	13.3	12	5.0	Polsenaere et al., 2013
Cirès	11	I	02/2008 to 02/2009	monthly	13	44.7598	-1.1107	12	45	0.58	12.2	5	0.4	Polsenaere et al., 2013
Renet	12	I	02/2008 to 02/2009	bi-monthly	23	44.7144	-1.0441	3	18	0.56	12.9	10	0.6	Polsenaere et al., 2013
Lanton	13	I	02/2008 to 02/2009	monthly	13	44.7002	-1.0244	15	36	0.26	12.5	11	1.2	Polsenaere et al., 2013
Milieu	14	I	02/2008 to 02/2009	monthly	13	44.6973	-1.0225	7	21	0.58	12.7	7	0.4	Polsenaere et al., 2013
Tagon	15	I	02/2008 to 02/2009	bi-monthly	26	44.6590	-0.9891	10	30	0.64	12.6	13	1.3	Polsenaere et al., 2013
Leyre	16	I	01/2008 to 03/2010 and 02/2014 to 02/2015	bi-monthly or monthly	59	44.6263	-0.9961	116	1700	17	13.0	11	0.9	Dubois et al., 2012 / Polsenaere et al., 2013 / Liénart et al., 2017, 2018
Landes	17	III	02/2008 to 02/2009	monthly	12	44.6169	-1.1091	14	117	0.49	14.1	3	1.1	Polsenaere et al., 2013
Adour	18	III	04/2013 to 06/2014 and 05/2017 to	monthly	24	43.4988	-1.2949	308	16912	516	14.0	48	2.4	Liénart et al., 2016 / Deborde, 2019
Têt	19	II	01/2006 to 05/2010	monthly	52	42.7137	2.9935	115	1369	23	15.7	8	NA	Higuera et al., 2014
Hérault	20	II	01/2006 to 05/2010	monthly	52	43.3594	3.4354	148	2582	53	16.0	7	NA	Higuera et al., 2014
Orb	21	II	01/2006 to 05/2010	monthly	52	43.2850	3.2813	136	1585	23	15.7	8	NA	Higuera et al., 2014
Aude	22	II	01/2006 to 05/2010	monthly	52	43.2442	3.1527	223	5327	40	14.2	31	NA	Higuera et al., 2014
Rhône	23	II	12/2003 to 01/2011	monthly	105	43.6787	4.6212	812	95590	1572	15.9	41	1.9	Harmelin-Vivien et al., 2010 / Cathalot et al., 2013 / Higuera et al., 2014

163

164 **2.2.Data origin**

165 Regarding the core parameters (C/N ratio,  $\delta^{13}\text{C}$ ,  $\delta^{15}\text{N}$ , water temperature, SPM, POC, PN, chl  
166 *a*), most of the data sets come from published studies (Canton et al., 2012; Cathalot et al., 2013;  
167 Dubois et al., 2012; Ferchiche et al., 2024; Harmelin-Vivien et al., 2010; Higueras et al., 2014;  
168 Liénart et al., 2016, 2017, 2018; Polsenaere et al., 2013), while most of additional parameters  
169 come from national databases (Tab. A1). When not available in the cited studies, concentrations  
170 of SPM,  $\text{NO}_3^-$ ,  $\text{NH}_4^+$  and  $\text{PO}_4^{3-}$ , pH and water temperature were retrieved from the *Naiades*  
171 database (<https://naiades.eaufrance.fr/>, consulted the 07/10/2023). Note that these parameters  
172 were not necessarily measured or sampled exactly at the same location or date for *Naiades* ~~than~~  
173 ~~was in~~ the cited studies. In that case, the location was chosen as close as possible to the study  
174 location and data values were time-interpolated to match the study date. Meteorological  
175 variables (air temperature, zonal and meridional wind, irradiance; ~~used to qualify~~  
176 ~~photosynthetic favourable conditions or wind-induced resuspension~~) come from Météo France,  
177 the French meteorological service. Wind data ~~were~~ received originally as direction and speed.  
178 To remove the angular bias, they were combined using scalar products to get zonal and  
179 meridional wind speeds, which range between minus and plus infinity (see Lheureux et al.,  
180 2022, for more details). River flows (~~used to qualify the hydrodynamics forcing~~) were retrieved  
181 from the *Banque Hydro* database (<https://www.hydro.eaufrance.fr/>, consulted the 07/10/2023)  
182 or from Polsenaere et al. (2013) for the small streams.

183 ~~Catchment properties were retrieved when available for the 23 rivers. Land use proportions~~  
184 originate from the ~~national~~ Corine Land Cover database  
185 (<https://www.statistiques.developpement-durable.gouv.fr/corine-land-cover-0>, consulted the  
186 10/01/2024). Soil organic carbon data originate from the ~~SoilTrEC~~ database  
187 (<https://esdac.jrc.ec.europa.eu/content/predicted-distribution-soc-content-europe-based-lucas-biosoil-and-czo-context-eu-funded-1>, consulted the 10/01/2024). Net erosion soil data  
188 originates from the WaTEM/SEDEM database (<https://esdac.jrc.ec.europa.eu/content/estimate-net-erosion-and-sediment-transport-using-watemseDEM-european-union>, consulted the  
189 10/01/2024). Strahler numbers originate from the CARTHAGE database  
190 (<https://www.sandre.eaufrance.fr/atlas/srv/api/records/c1d89cc3-c530-4b0d-b0ae-06f5ebf7997d>, consulted the 15/08/2025). Useful reserve values come from the GSF database  
191 (Le Bas, 2025). ~~Geological and soil types come from the GISSOL database (gathered by great~~  
192 ~~geological and soil types; INRA, 2025). Wastewater treatment capacities originate from the Eau~~  
193 ~~France~~ ~~WFS~~ ~~services~~  
194 (<https://services.sandre.eaufrance.fr/geo/odp?REQUEST=getCapabilities&service=WFS&VERSION=2.0.0, couche sa:SysTraitementEauxUsees>, consulted the 15/08/2025). All these  
195 ~~catchment data were pre-processed on a Geographical Information System to extract~~  
196 ~~information for each catchment surface, then averaged or weighted (depending on continuous~~  
197 ~~or semi-quantitative data) to characterise each system by with a value.~~

202

203 It should be noted that a complete study was already dedicated to the Loire River and reported  
204 as a companion article (Ferchiche et al., 2024). Consequently, the results are not reported in the  
205 present study but are used for multi-system comparisons (Fig. 5 and 7, and corresponding text).

### 2.3.Determination of sources signatures

207 To run mixing models for quantifying POM composition, it is previously needed to 1)  
208 determine sources of POM, and 2) associate elemental and isotopic signatures to these sources.

In riverine systems, phytoplankton autochthonous (mainly phytoplankton) and allochthonous (terrestrial POM resuspended sediment, terrestrial fresh litter or rock-derived soil) are the main sources that are usually considered as fueling the POM (e.g., Ferchiche et al., 2024; Pradhan et al., 2016; Sarma et al., 2014). Nevertheless, sewage POM may also contribute (Higueras et al., 2014). Consequently, phytoplankton, labile and refractory terrestrial POM and sewage POM were considered as potential sources in this study.

Phytoplankton cannot be easily picked up from bulk particles to measure its elemental and isotopic ratios. Therefore, the method developed and used by Savoye et al. (2012), Liénart et al. (2017) and Ferchiche et al. (2024) was applied here. It consists of determining the elemental and isotopic ratios from a subset of the bulk dataset. Briefly, phytoplankton-dominated POM is characterized by a low POC/chl  $a$  ratio ( $\leq 200$  or even  $\leq 100$  g/g; Savoye et al., 2003 and references therein). Thus, elemental and isotopic ratios of samples exhibiting a low POC/chl  $a$  ratio can be considered as good estimates of phytoplankton elemental and isotopic ratios. When the POC/chl  $a$  ratio is not available, samples exhibiting a high PN/SPM ratio can be used. Additional constraints may be used to minimize potential overlap between phytoplankton and terrestrial elemental and isotopic signatures. Phytoplankton elemental and especially isotopic ratios may deeply vary over time and space depending on primary production intensity and potential limiting factors, nutrient origin, etc. (e.g., Miller et al., 2013; Savoye et al., 2003). When existing, this variability has to be taken into account to avoid using elemental and isotopic signatures that are not valid at the time or location of the sampling. This could be performed by using regressions between elemental and/or isotopic ratios and environmental variables (see Ferchiche et al., 2024; Liénart et al., 2017; Savoye et al., 2012). At last, when no samples exhibit a low POC/chl  $a$  ratio, samples exhibiting the lowest (even if high) POC/chl  $a$  ratios can be used, but the data should be firstly be corrected from the contribution of the terrestrial POM using Equations 1-3.

$$\delta^{13}\text{C}_{\text{sample}} = ([\text{POC}]_{\text{phytoplankton}} \times \delta^{13}\text{C}_{\text{phytoplankton}} + [\text{POC}]_{\text{terrestrial}} \times \delta^{13}\text{C}_{\text{terrestrial}}) / [\text{POC}]_{\text{sample}}$$

(eq. 1)

$$236 \quad [\text{POC}]_{\text{phytoplankton}} = [\text{chl } a]_{\text{sample}} \times (\text{POC}/\text{chl } a)_{\text{mean}} \quad (\text{eq. 2})$$

$$237 \quad [\text{POC}]_{\text{terrestrial}} = [\text{POC}]_{\text{sample}} - [\text{POC}]_{\text{phytoplankton}} \quad (\text{eq. 3})$$

238 where  $(\text{POC}/\text{chl } a)_{\text{mean}}$  is the mean POC/chl  $a$  ratio of the samples used to determine  
239 phytoplankton signatures. Similar equations are used for the N/C ratio,  $\delta^{15}\text{N}$  and C/N ratio, but  
240 using PN instead of POC for  $\delta^{15}\text{N}$  and C/N ratio.

241 Elemental and isotopic signatures of terrestrial POM can be estimated by directly measuring  
242 elemental and isotopic ratios in ~~terrestrial materials~~ a sample like soil, ~~rocks or~~ ~~and~~ vascular  
243 plants (e.g., Sarma et al., 2014). However, this does not take into account the reworking of this  
244 material within the river system, which can affect these signatures (Hou et al., 2021). Thus,  
245 similarly to phytoplankton, elemental and isotopic signatures of terrestrial POM can be  
246 estimated using subsets of bulk data, following the approach of Savoye et al. (2012), Liénart et  
247 al. (2017) and Ferchiche et al. (2025, 2024). Labile ~~t~~Terrestrial POM is usually characteris~~zed~~  
248 by high POC/chl  $a$  and C/N ratios and low POC/SPM ratios (Etcheber et al., 2007; Savoye et  
249 al., 2003 and references therein). However, during its decay in aquatic systems, terrestrial POM  
250 is colonis~~zed~~ by bacteria (low C/N ratio), resulting in a consortium terrestrial POM + bacteria  
251 of lower C/N ratio than the original terrestrial POM (Etcheber et al., 2007; Savoye et al., 2012).  
252 Finally, one can discriminate two kinds of terrestrial POM: refractory terrestrial POM,  
253 characteris~~zed~~ by high POC/chl  $a$  and C/N ratios and very low POC/SPM ratio, and quite labile  
254 terrestrial POM characteris~~zed~~ by high POC/chl  $a$  ratio, intermediate C/N ratios and low  
255 POC/SPM ratio (Etcheber et al., 2007; Savoye et al., 2012). Thus, subsets of high POC/chl  $a$   
256 ratio can be selected to determine the elemental and isotopic signatures of terrestrial POM. The  
257 C/N ratio can be used to discriminate labile from refractory terrestrial POM. When no samples  
258 exhibit a high POC/chl  $a$  ratio, samples exhibiting the highest (even if quite low) POC/chl  $a$   
259 ratio can be used, but the data should ~~be~~ firstly be corrected ~~from~~ the contribution of the  
260 phytoplankton POM using Equations 1-3.

261 Elemental and isotopic ratios of riverine POM can exhibit a departure from a simple  
262 phytoplankton-terrestrial POM mixing. In the present study, this was the case in only two rivers.  
263 For the Têt River, the elemental and isotopic signature of anthropogenic POM was available in  
264 Higueras et al. (2014). It consisted of analyses of POM sampled in the wastewater treatment  
265 plant (WWTP) ~~the~~ closest to the sampling site. For the Orb River, the signatures were estimated  
266 using the sample exhibiting the lowest  $\delta^{15}\text{N}$ , typical of anthropogenic POM (Ke et al., 2019).

267 The estimation of POM-source signatures was performed independently for each river, except  
268 for some of the tributaries of the Arcachon Lagoon (rivers 11 to 15), where data sets were  
269 gathered, thanks to very similar characteristics (same  $\delta^{13}\text{C}$  of dissolved inorganic carbon;  
270 Polsenaere et al., 2013), to get a larger subset of data for estimating elemental and isotopic  
271 signatures more accurately. All criteria used for defining the above-described subsets are  
272 reported in Table 2.

273 Table 2 Elemental and isotopic signatures of POM sources and criteria used to choose the data  
274 subset to determine them. When the signature did not vary over time, average  $\pm$  standard

275 deviation are reported. When the signature did vary over time, minimum and maximum values,  
 276 standard deviations, as well as equations are reported. The types of mixing models performed  
 277 for each river are also indicated (carbon mixing models were performed using  $\delta^{13}\text{C}$  and N/C  
 278 ratio, or only  $\delta^{13}\text{C}$ ; nitrogen mixing models were performed using  $\delta^{15}\text{N}$  and C/N ratio; mixed  
 279 mixing models were performed using  $\delta^{13}\text{C}$ ,  $\delta^{15}\text{N}$  and N/C ratio). POC% (or PN%) = Particulate  
 280 Organic Carbon (or Particulate Nitrogen) to Suspended Particulate Matter ratio (%); C/N =  
 281 POC/PN ratio (mol/mol); chla = chlorophyll *a* ( $\mu\text{g/l}$ ); phaeo = phaeopigments ( $\mu\text{g/l}$ ); conduc =  
 282 conductivity ( $\mu\text{S}$ ); temp = water temperature ( $^{\circ}\text{C}$ ); - Q7 = mean of past seven days river flow;  
 283  $\text{NO}_3^-$  = nitrate ( $\text{mg}(\text{NO}_3^-)/\text{l}$ ).

River	Source discriminants			WWTP's POM	Model performed			Labile terrestrial matter				Refractory terrestrial matter			
	Labile terrestrial matter	Refractory terrestrial matter	Phytoplankton		Carbon	Nitrogen	Mixed	$\delta^{13}\text{C}$	$\delta^{15}\text{N}$	C/N	N/C	$\delta^{13}\text{C}$	$\delta^{15}\text{N}$	C/N	N/C
Seine	C/N > 10		POC/chla < 200		X	X		-28.5 ± 0.3	6.6 ± 0.9	10.6 ± 0.3	0.093 ± 0.002				
Orne	C/N > 11		POC/chla < 500		X	X		-28.4 ± 0.3	5.8 ± 1.0	12.4 ± 0.4	0.082 ± 0.003				
Rance	POC/chla > 200 and chla < 10		POC/chla < 150		X	X		-26.8 ± 0.2	6.1 ± 0.7	8.8 ± 0.4	0.113 ± 0.007				
Elorn	C/N > 12		POC/chla < 200		X	X		-28.4 ± 0.7	5.8 ± 0.9	13.0 ± 0.8	0.077 ± 0.005				
Aulne	C/N > 11		POC/chla < 200 and C/N < 9		X	X		-28.9 ± 0.8	5.8 ± 0.8	12.1 ± 1.1	0.08 ± 0.008				
Loire	POC/chla > 500		POC/chla < 200		X	X		-28.1 ± 0.1	5.9 ± 0.3	10.3 ± 0.2	0.097 ± 0.002				
Sèvre Niortaise	C/N > 14		POC/chla < 300		X			-28.0 ± 0.4			0.057 ± 0.040				
Charente	C/N > 12		POC/chla < 300		X	X		-29.0 ± 0.4	4.7 ± 0.2	14.5 ± 0.5	0.069 ± 0.002				
Seudre	POC/chla > 2000 and C/N > 12		POC/chla < 1000		X			-28.5 ± 0.1							
Porge	C/N > 15		$\delta^{13}\text{C}$ : POC/chla < 100 ; N/C : mean of Cirès to Landes		X			-26.5 ± 1.1			0.050 ± 0.007				
Cirès / Renet / Milieu / Lanton / Tagon	C/N > 15 and chla < 1		POC/chla < 1000 and POC% > 10		X			-28.5 ± 0.5			0.053 ± 0.013				
Leyre	C/N > 15 and chla < 1		POC/Chla < 1000. $\delta^{13}\text{C} < 28.59$ and POC% > 10		X			-28.3 ± 0.5			0.06 ± 0.005				
Landes	C/N > 12		POC/Chla < 600. $\delta^{13}\text{C} < -29.1$		X			-29.1 ± 0.4			0.075 ± 0.002				
Adour	POC/chla > 3000		POC/chla < 200		X			-26.0 ± 0.9			0.099 ± 0.008				
Têt	C/N > 11.5	POC% < 4.25	PN% > 2, $\delta^{13}\text{C} < 26$ and $\delta^{15}\text{N} > 5$	Measured		X		-26.0 ± 0.2	3.7 ± 0.6	12.2 ± 0.5	0.082 ± 0.002	-26.0 ± 0.6	6.7 ± 1.4	5.8 ± 1.4	0.180 ± 0.045
Aude	C/N > 12	Q7 > 70	PN% > 1 or 2 and C/N < 6		X			-28.1 ± 0.6	6.3 ± 0.1	15.3 ± 1.6	0.066 ± 0.007	-28.0 ± 0.7	4.7 ± 0.4	7.3 ± 1.0	0.139 ± 0.018
Orb	C/N > 10		PN% > 2, $\delta^{15}\text{N} > 4.06$	Lower $\delta^{15}\text{N}$	X			-27.1 ± 0.4	3.7 ± 0.4	10.5 ± 0.3	0.095 ± 0.350				
Hérault	C/N > 12	Q > 45	PN% > 2		X			-27.7 ± 0.2	6.1 ± 0.7	13.7 ± 1.2	0.073 ± 0.007	-27.8 ± 0.4	4.7 ± 0.6	8.2 ± 1.5	0.124 ± 0.019
Rhône	C/N > 12	POC% < 1.25	C/N < 6.68 and $\delta^{15}\text{N} > 3.92$		X			-26.4 ± 1.3	5.2 ± 1.0	17.0 ± 3.2	0.061 ± 0.012	-25.9 ± 0.4	3.1 ± 0.8	8.8 ± 3.1	0.119 ± 0.032

284

285

286 **Table 2 (continued)**

River	Phytoplankton				WWTP's POM					
	$\delta^{13}\text{C}$ + equations	$\delta^{15}\text{N}$ + equations	C/N	N/C + equations	$\delta^{13}\text{C}$	$\delta^{15}\text{N}$	C/N	N/C		
Seine	-32.8 ± 1.1	8.4 ± 1.7	7.4 ± 0.7	0.136 ± 0.012						
Orne	-31.4 ± 0.8	4.3 ± 0.8	6.6 ± 1.3	0.141 ± 0.010						
Rance	[-31.4;-25;6] ± 1.7	$5.7 \times 10^{-4} \times [\text{chl}a + \text{phaeo}]^2 - 0.04 \times [\text{chl}a + \text{phaeo}] - 30.6$	[4.7;11.4] ± 0.7	$-0.28 \times [\text{NO}_3^-] + 12.7$	6.2 ± 0.4	0.161 ± 0.010				
Elorn	-27.4 ± 0.3	6.9 ± 0.5	10.0 ± 0.9	0.101 ± 0.007						
Aulne	-28.1 ± 0.2	8.6 ± 0.2	8.2 ± 0.2	0.122 ± 0.003						
Loire	[-30.6;-25.0] ± 0.9	$5 \times 10^{-4} \times [\text{chl}a + \text{phaeo}]^2 - 0.02 \times [\text{chl}a + \text{phaeo}] - 0.39 \times [\text{chl}a + \text{phaeo}] - 27.9$	[3.0;10.4] ± 1.2	$4.2 \times 10^{-4} \times [\text{chl}a]^2 - 0.08 \times [\text{chl}a] + 8.2$	7.2 ± 0.6	0.140 ± 0.011				
Sèvre Niortaise	[-35.7;-29.2] ± 1.0	$-258 \times \exp([\text{chl}a + \text{phaeo}]^2 / 16055) - 0.15 \times [\text{temp}] + 229$			[0.106;0.145] ± 0.006	$2.9 \times 10^{-3} \times [\text{chl}a + \text{phaeo}] + \alpha_1$				
Charente	-30.8 ± 0.03	7.5 ± 1.6	6.6 ± 0.3	0.152 ± 0.006						
Seudre	-33.3 ± 0.1									
Porge	-33.6 ± 0.4			0.128 ± 0.008						
Cirès / Renet / Milieu / Lanton / Tagon	-34.9 ± 0.4			0.133 ± 0.006						
Leyre	-30.1 ± 0.3			0.140 ± 0.016						
Landes	-29.9 ± 0.3			0.112 ± 0.010						
Adour	-28.2 ± 0.6			0.111 ± 0.010						
Têt	[-29.7;-27.8] ± 0.6	$-5.2 \times 10^{-3} \times [\text{temp}]^2 + 0.08 \times [\text{temp}] - 27.5$	[5.3;13.3] ± 1.8	$5.53 \times [\text{temp}] - 5.5$	5.6 ± 0.7	0.181 ± 0.021	-26.3 ± 0.1	-0.7 ± 0.1	6.3 ± 0.3	0.160 ± 0.017
Aude	[-32.6;-27.8] ± 0.6	$-0.21 \times [\text{temp}] - 26.5$	[5.2;10.6] ± 1.6	$-1.13 \times \delta^{13}\text{C} - 26.2$	5.0 ± 0.8	0.205 ± 0.033				
Orb	[-30.7;-23.4] ± 0.6	$-0.19 \times [\text{temp}] - 26.0$	[4.9;8.4] ± 0.6	$8.44 - (3.63 \times (\text{conduc} - 505)) / (\text{conduc} - 111)$	4.8 ± 0.9	0.213 ± 0.039	-27.1 ± 0.4	1.9 ± 1.9	3.7 ± 3.7	0.270 ± 0.270
Hérault	[-31.5;-27.5] ± 1.0	$-0.19 \times [\text{temp}] - 26.0$	[6.3;10;9] ± 1.3	$3.6 \times 10^{-2} \times [\text{temp}]^2 + 1.15 \times [\text{temp}] + 14.6$	5.0 ± 0.7	0.203 ± 0.031				
Rhône	-27.8 ± 1.2		5.6 ± 0.8		5.5 ± 0.8	0.180 ± 0.030				

287

288 **2.4. Quantification of POM composition**

289 POM composition was quantified using a Bayesian mixing model ('*simmr*' R package version  
 290 0.4.5, Govan and Parnell, 2023), which solves the equations system based on bulk and source  
 291 POM elemental and isotopic signatures. Mixing models were computed for each sampling date  
 292 of each river (Tab. 1), using carbon ( $\delta^{13}\text{C}$  and N/C ratio, Eq. 4, 7, 8), nitrogen ( $\delta^{15}\text{N}$  and C/N  
 293 ratio, Eq. 5, 6, 8), and/or a combination of three ( $\delta^{13}\text{C}$ ,  $\delta^{15}\text{N}$  and N/C ratio, Eq. 4, 5, 7, 8) tracers.  
 294 From the three mixing models performed for each sampling date and river (carbon, nitrogen or  
 295 mixed), one model was selected as the best estimation of bulk POM data. It should be noted  
 296 that N/C and C/N ratios give information on the mixing of C and N, respectively (Perdue and  
 297 Koprivnjak, 2007). We used at least the same number of equations as unknowns (sources) to  
 298 avoid running underdetermined models that result in large uncertainty in model outputs (Phillips  
 299 et al., 2014). Equations of the models were:

300  $\delta^{13}\text{C}_{\text{mixture}} = x_1 \delta^{13}\text{C}_{\text{source 1}} + x_2 \delta^{13}\text{C}_{\text{source 2}} + x_3 \delta^{13}\text{C}_{\text{source 3}} + x_4 \delta^{13}\text{C}_{\text{source 4}}$  (Eq. 4)

301  $\delta^{15}\text{N}_{\text{mixture}} = x_1 \delta^{15}\text{N}_{\text{source 1}} + x_2 \delta^{15}\text{N}_{\text{source 2}} + x_3 \delta^{15}\text{N}_{\text{source 3}} + x_4 \delta^{15}\text{N}_{\text{source 4}}$  (Eq. 5)

302  $\text{C/N}_{\text{mixture}} = x_1 \text{C/N}_{\text{source 1}} + x_2 \text{C/N}_{\text{source 2}} + x_3 \text{C/N}_{\text{source 3}} + x_4 \text{C/N}_{\text{source 4}}$  (Eq. 6)

303  $\text{N/C}_{\text{mixture}} = x_1 \text{N/C}_{\text{source 1}} + x_2 \text{N/C}_{\text{source 2}} + x_3 \text{N/C}_{\text{source 3}} + x_4 \text{N/C}_{\text{source 4}}$  (Eq. 7)

304  $x_1 + x_2 + x_3 + x_4 = 1$  (Eq. 8)

305 As there was no *a priori* knowledge of sources contributions to the POM mixture, the models  
 306 were set with an uninformative prior (1, 1, 1, 1) following a Dirichlet distribution (all sources  
 307 have an equal probability to contribute to the mix; Phillips et al., 2014). Model runs were set  
 308 following the recommendations of Phillips et al. (2014). Models outputs were evaluated with  
 309 Gelman-Rubin diagnostic (verification of chain convergence) and predictive distributions to  
 310 ensure the good fit of the models to the observed data. Models outputs are given as medians.  
 311 Absolute uncertainties for the models varied from 1 to 18 % (range of average for each river)  
 312 with an overall average of 8 % (all models).

## 313 2.5. Forcings at local and multi-systems scales

314 Environmental forcings driving POM composition were determined using redundancy analysis  
 315 (RDA; ‘dudi.pca’ and ‘pcaiv’ functions; R package {ade4} version 1.7-19). RDA summarises  
 316 multiple linear regressions between the response variable (POM composition: mixing model  
 317 outputs) and a set of explanatory variables (environmental forcings) to assess causality links  
 318 (Legendre et al., 2011). RDAs were performed at single-river and multi-river scales. Regarding  
 319 the multi-rivers scale, the annual mean POM composition of each river was used to determine  
 320 the drivers of spatial (i.e., between-rivers) variations of POM composition.

321 The proxies of the environmental forcings were chosen to directly or indirectly reflect the  
 322 forcings that affect the processes that occurring in the river and the adjacent ecosystems (e.g.,  
 323 primary production, soil leaching or WWTP's discharge) and influencing POM source inputs  
 324 and isotopic values. To homogenise the data sets for running the single-river RDAs, the same  
 325 combination of twelve parameters (see Table A2) proxies for environmental forcings was used  
 326 for each river. They are linked to primary production (chlorophyll *a*, phaeopigments,  
 327 temperature, pH, ammonium, nitrate, phosphate, irradiance), upstream and lateral, natural  
 328 and/or anthropogenic inputs (river flow, rain, SPM, ammonium, nitrate, phosphate), and  
 329 resuspension (SPM, zonal and meridional wind energy). SPM, chlorophyll *a*, phaeopigments,  
 330 temperature, daily river flow, pH, ammonium, nitrates, phosphates, irradiance, zonal and  
 331 meridional wind. For the multi-river RDA, environmental proxies were selected to reflect  
 332 processes occurring at large spatial scales and in the river basin. Hence, a new combination of  
 333 sixteen thirty-nine proxies was Forty parameters (See Fig. A6) were used. They are linked to:  
 334 river flow, water quality (conductivity, nitrates), climate setting (river flow, latitude,  
 335 longitude, air temperature, precipitations, zonal, and meridional wind and total wind energy),

336 artificial, hydromorphology (river length, basin surface area, slope, Strahler number), land use  
337 coverage (agricultural, artificial, forest and natural, -and wetlands and water bodies surface)  
338 areas, net soil erosion, soil properties (organic carbon content, net erosion in the soil,  
339 granulometry, useful reserves), soil type (podzol, brown, organic and hydromorphic soil),  
340 geological type (alluvial, calcareous, clayey, detrital, sandy, loamy, crystalline and  
341 metamorphic, volcanic and other/organic), and urban pressure (WWTP capacities, WWTP  
342 capacities to river flow ratio), river length, basin surface area, latitude, longitude. From this  
343 initial list of proxies, some were removed to limit the auto-correlation (use of the Variance  
344 Inflation Factor, Borcard et al., 2011) and to improve the adjusted R<sup>2</sup> of each RDA analysis  
345 (Tab. A2 and Fig. A6).

## 346 2.6. Typology of systemsriver dynamics

347 Rivers were classified based on POM composition and theirits temporal dynamics by  
348 performing a regionalisation analysis as in Liénart et al. (2018) (Fig. A1). This method, based  
349 on multivariate cluster analysis (Souissi et al., 2000), allows to consider the temporal (seasonal)  
350 variations specific to each river in addition to the spatial (between-rivers) component. The  
351 regionalisation analysis was based on POM composition data (i.e., proportions of sources)  
352 computed for each river and each month. When the sampling was fortnightly, averages were  
353 performed to get one value per month. When more than twelve months were available (bi-  
354 monthly sampling or more years)one year was sampled, a standard year of twelve months was  
355 chosen (averaged by month if fortnight dates). (Souissi et al., 2000). Nevertheless, to check if  
356 the choice of one year over the other ones would modify the typology, another regionalisation  
357 was performed using all available years for all rivers. Also, in order to check if the over-  
358 representation of the small rivers and streams fuelling the Arcachon Bay would bias the  
359 typology, a third regionalisation was performed, reducing the numbers of these rivers from  
360 8 to 3 (and especially from 6 to 1 regarding rivers of Type I). The results (Fig. 5, Fig. A7) are  
361 very similar, indicating the robustness of the method.

362 A contingency matrix (rivers, sources, months) was created from monthly values of source  
363 contributions (i.e., mixing model outputs). For each month, a dendrogram was performed, and  
364 ten cut-off levels were considered. Then, for each cut-off level, similarities between stations  
365 were identified within the twelve-monthly dendrograms. Ultimately, global similarities  
366 between rivers were computed using a fuzzy cluster that returns probabilities of membership of  
367 each river to each cluster type. The best number of river types, i.e., river dynamics typology,  
368 was determined considering the best Dunn coefficient (Dunn, 1974) and Silhouette score  
369 (Rousseeuw, 1987).

## 370 3. Results


371 Hereafter, four rivers (Rance, Charente, Milieu and Hérault Rivers) were selected and  
372 considered as representative of each type of studied river (see section 3.4). Thus, most of the

373 results are illustrated using these four rivers. Graphs of all the other rivers are reported in the  
374 supplementary material.

### 375 **3.1. Contrast~~inged~~ seasonalities in river characteristics**

376 As stated in section 2.1, the 23 studied rivers encompassed large gradients of environmental  
377 characteristics, as illustrated by the lowest and highest annual means of river flow (0.3 and 1572  
378 m<sup>3</sup>/s; Lanton and Rhône Rivers), water temperature (12.3 to 17.1 °C; Cirès and Têt Rivers),  
379 SPM (2.7 and 40.9 mg/l; Cirès and Rhône River), POC (0.3 and 5.1 mg/l; Hérault and Loire  
380 Rivers) and chlorophyll *a* (0.4 to 57.1 µg/l; Cirès and Rance Rivers) concentrations as well as  
381 POC/chl *a* (199 and 6444 g/g; Loire and Leyre Rivers) and C/N (5.9 and 20.3 mol/mol; Têt and  
382 Lanton Rivers) ratios; this was less contrast~~inged~~ among rivers for δ<sup>13</sup>C (-30.2 and -26.2 ‰;  
383 Sèvre and Têt Rivers) and especially δ<sup>15</sup>N (4.0 and 8.0 ‰; Leyre and Rance Rivers) (Fig. 2,  
384 A2).

385 As generally observed in rivers from mid-latitude, ~~the~~ studied rivers exhibited clear seasonal  
386 patterns in water temperature with lower and higher values in winter and summer, respectively.  
387 However, such clear seasonal patterns were not always recorded for all the parameters, as there  
388 were contrast~~inged~~ patterns of seasonal variability among rivers. Indeed, the seasonal  
389 variability of river flow was quite smooth (e.g., the Rance and Charente Rivers) with a higher  
390 flow in winter/spring and lower flow in summer/fall for some rivers, whereas it was highly  
391 pulsed for some others with constant low levels marked by short and strong floods (e.g., 53m<sup>3</sup>/s  
392 in mean but 1169m<sup>3</sup>/s in flood time for the Hérault River) (Fig. 2). Overall, one can distinguish  
393 rivers that are characterized by high concentrations of chlorophyll *a* and clear seasonal patterns  
394 of most parameters (e.g., 53 µg/l of chlorophyll *a* in mean ranging from 3 to 135 µg/l in the  
395 Rance River) from rivers characterized by low concentrations of chlorophyll *a*, high POC/chl  
396 *a* and low seasonal variability for most of the parameters (e.g., 1.1 µg/l of chlorophyll *a* in mean  
397 ranging from 0.7 to 1.7 µg/l in the Milieu River) and from rivers that are characterized by high  
398 seasonal variability of most parameters but without a clear seasonal pattern (e.g., Hérault  
399 River). Other rivers exhibited intermediate behaviour (e.g., Charente River) (Fig. 2, A2).  
400 Usually, Rance-like rivers exhibited high concentrations of chlorophyll *a* in spring/summer  
401 associated with POC/chl *a* ratio lower than 200 g/g, C/N ratio lower than 8 mol/mol and low  
402 δ<sup>13</sup>C (down to -31 ‰ or -33 ‰; e.g., Seine River, Fig. A2). In contrast, Milieu-like rivers  
403 exhibited high POC/chl *a* (> ~700 g/g) and C/N ratio (> 15 mol/mol) and quite constant δ<sup>13</sup>C  
404 (~-29 – -28 ‰) all year round (e.g., Cirès and Renet Rivers). These rivers are tributaries of the  
405 Arcachon Lagoon. Hérault-like rivers flowing into the Mediterranean Sea exhibited highly and  
406 suddenly variable C/N ratios (4 – 17 mol/mol), δ<sup>13</sup>C (~-33 – -26 ‰) and δ<sup>15</sup>N (~2 – 12 ‰) (e.g.,  
407 Aude and Orb Rivers; Fig. A2).




409 Figure 2 Temporal variations of matter characteristics for representative rivers along the studied  
410 periods for  $\delta^{13}\text{C}$  (left axis; black dotted line) and  $\delta^{15}\text{N}$  (right axis; blue line) (first column); C/N  
411 (left axis; black dotted line) and POC/chl *a* (right axis; blue line) ratios (second column);  
412 SPM(left axis; black dotted line), POC (right axis; blue line) and chl *a* (right axis; blue dotted  
413 line) concentrations (third column) and river flow (left axis; black dotted line) and temperature  
414 (right axis; blue line) (fourth column).

### 415 **3.2. Elemental and isotopic signatures of POM sources**

416 Elemental and isotopic signatures of phytoplankton were estimated for each of the twenty-three  
417 rivers (Tab. 2, Fig. 3 and A3). Most of them (all of them for the C/N ratio) were found to be  
418 constant over time. Their annual mean values varied between -34.9 ‰ (some tributaries of the  
419 Arcachon Lagoon) and -27.4 ‰ (Elorn River) for  $\delta^{13}\text{C}$ , between 4.3 ‰ (Elorn River) and 8.6  
420 ‰ (Aulne River) for  $\delta^{15}\text{N}$  and between 4.8 mol/mol (Orb River) and 10.0 mol/mol (Elorn River)  
421 for the C/N ratio. Some of them varied over time along with pigment concentration and ratio or  
422 with temperature for  $\delta^{13}\text{C}$ , and with pigment concentration (chlorophyll *a* and/or  
423 phaeopigments), nitrate concentration, temperature,  $\delta^{13}\text{C}$  or conductivity for  $\delta^{15}\text{N}$  (Tab. 2). The  
424 range of temporal variability was usually 4-6 ‰ for  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$ . Overall, phytoplankton  
425 signatures are comprised between -35.6 and -23.8 ‰ for the  $\delta^{13}\text{C}$  and between 3.0 and 13.2 ‰  
426 for the  $\delta^{15}\text{N}$ .

427 All other signatures were found to be constant over time (Tab. 2 and A2, Fig. 3 and A3) but  
428 may differ between rivers. Signatures mean annual values of labile terrestrial POM were  
429 comprised between -29.1 and -26.0 ‰ for the  $\delta^{13}\text{C}$ , between 3.7 and 6.6 ‰ for the  $\delta^{15}\text{N}$  and  
430 between 8.8 and 17.0 mol/mol for the C/N ratio. Signatures mean annual values of refractory  
431 terrestrial POM were comprised between -28.0 and -25.9 ‰ for the  $\delta^{13}\text{C}$ , between 3.1 and 6.7  
432 ‰ for the  $\delta^{15}\text{N}$  and between 5.8 and 8.8 mol/mol for the C/N ratio. Signatures mean annual  
433 values of sewage POM were -27.1 and -26.3 ‰ for  $\delta^{13}\text{C}$ , 1.9 and -0.7 ‰ for  $\delta^{15}\text{N}$  and 3.7 and  
434 6.3 mol/mol for C/N ratio for Orb and Têt Rivers, respectively.





436 Figure 3  $\delta^{13}\text{C}$ ,  $\delta^{15}\text{N}$ , N/C or C/N values of bulk POM (black crosses) and sources. The latter are  
437 presented as closed circles (average) and bars (standard deviation) when the signatures were  
438 constant over time and by colored area when at least one of the proxies was variable over time  
439 (see Table 2). This colored area corresponds to the dispersion of the values, including their  
440 uncertainties.

### 441 **3.3. Dynamics of particulate organic matter composition**

442 Particulate organic matter composition resulting from mixing models outputs is presented  
443 hereafter, for each river, as the relative contribution of each source to the POM pool (Fig. 4).  
444 Among rivers whose POM is composed of only two sources (terrestrial POM and  
445 phytoplankton), one can distinguish rivers with terrestrial-dominated POM (e.g., Milieu River:  
446 terrestrial POM accounted for  $94 \pm 3\%$  of the mixture) to rivers of intermediate POM  
447 composition (e.g., Charente and Rance Rivers where phytoplankton accounted for  $34 \pm 10\%$   
448 and  $62 \pm 10\%$  of the mixture, respectively). All these rivers flow in-into the English Channel  
449 and the Atlantic Ocean. The rivers whose POM is composed of three or four sources flow in  
450 into the Mediterranean Sea. In these rivers, terrestrial POM is present as refractory and labile  
451 materials. The contribution of labile terrestrial POM ranged between  $16 \pm 15\%$  (Têt River) and  
452  $46 \pm 21\%$  (Orb River), and of refractory terrestrial POM between  $21 \pm 9\%$  (Rhône River) and  
453  $39 \pm 15\%$  (Aude River). The contribution of phytoplankton ranged between  $34 \pm 15\%$  (Aude  
454 River) and  $51 \pm 30\%$  (Hérault River) for the Mediterranean rivers. The fourth source of POM  
455 was the WWTP's POM. It was identified as a source in the Orb and Têt Rivers and accounted  
456 for  $15 \pm 6\%$  and  $10 \pm 7\%$  in these two rivers, respectively. Regarding temporal variations of  
457 POM composition, some rivers exhibited clear seasonal patterns, whereas others revealed a  
458 homogeneous composition over the annual cycle (Fig. 4). The rivers where POM was highly  
459 dominated by terrestrial POM (Seudre, Cirès, Renet, Lanton, Milieu, Tagon, Leyre Rivers)  
460 showed almost no seasonal variability. In contrast, some rivers like the Rance, the Elorn or the  
461 Aulne River showed a clear seasonal pattern with the dominance of terrestrial material in winter  
462 and phytoplankton in summer. At last, other rivers exhibited less clear (e.g., Landes, Porge,  
463 Charente Rivers) or even no clear seasonal pattern but a quite stochastic variability over the  
464 annual cycle (e.g., Sèvre, Adour, Aude, Orb).

465 It should be noted that the above is valid for carbon and mixed as well as nitrogen models (cf.  
466 Tab. 2; Fig. 4 and A4).






468

469 Figure 4 Temporal dynamic (rectangle graphs) and (inter-)annual mean (pie charts) of POC  
470 source proportions. Sources are phytoplankton (green), labile terrestrial material (brown),  
471 refractory terrestrial material (yellow) and anthropogenic POM (orange).

472

### 473 3.4. Typology of river dynamics

474 Four types of river dynamics were determined by the regionalisation analysis based on river  
 475 POM compositions and their temporal dynamics (Fig. 5). The seven rivers (Renet, Cirès,  
 476 Lanton, Milieu, Seudre, Tagon and Leyre River), mainly belonging to Type I, are were  
 477 characterised by terrestrial-dominated POM and no/low seasonality. Six of them are small  
 478 streams/rivers flowing to the Arcachon Lagoon. The five rivers (Aude, Hérault, Têt, Rhône and  
 479 Orb River), mainly belonging to Type II, are were characterised by the co-occurrence of labile  
 480 and refractory terrestrial POM and large temporal variability, but, except for the Hérault River,  
 481 without a clear seasonal pattern. They all flow to the Mediterranean Sea. The five rivers (Porge,  
 482 Adour, Charente, Orne and Landes River), mainly belonging to Type III, are were composed  
 483 of phytoplankton and terrestrial POM, and exhibited moderate seasonality. Type III is clearly  
 484 an intermediary between Type I and Type IV. These five rivers flow to the Atlantic Ocean or  
 485 the English Channel. Among the seven rivers flowing to the Arcachon Lagoon, the two that  
 486 mainly belong to Type III are man-managed streams and flow through lakes, contrary to the six  
 487 other ones, which mainly belong to Type I and are natural streams that do not flow through  
 488 lakes. Finally, the six rivers (Rance, Elorn, Aulne, Loire, Seine and Sèvre River) mainly  
 489 belonging to Type IV are were composed of phytoplankton and terrestrial POM, and exhibited  
 490 high seasonality. These six rivers flow to the Atlantic Ocean or the English Channel. It should  
 491 be noted that the regionalisations performed using all sampled years for all rivers (Fig. A7)  
 492 resulted in the same typology and in the same type for each river, whatever the sampling year.  
 493 The only exception is the Leyre River, which switched from Type III in 2008 to Type I in 2009.








494

495 Figure 5 Typology of river dynamicss following a hierarchical cluster analysis on POM source  
496 proportions. The percentages of membership for each type attributed to each river are shown.

### 497 **3.5. Environmental forcings driving POM composition**

498 One redundancy analysis was performed for each river to relate environmental parameters,  
499 considered as proxies of drivers, to the POM composition, i.e., to assess the drivers of the  
500 temporal variability of POM composition for each river (Fig. 6 and A5). It should be kept in  
501 mind that the POC or PN concentration of each source was used for these analyses and not the  
502 relative proportion of the sources. In type-I rivers, i.e., rivers characteriszed by terrestrial-  
503 dominated POM and no/low seasonality, terrestrial POM is usually linked to river flow and/or  
504 SPM concentration (e.g., Milieu River on Fig. 6, Leyre and Tagon Rivers in Fig. A5). However,  
505 this feature is not always clear since the POM of these rivers is always dominated by terrestrial  
506 material, almost regardless of whatever the environmental conditions are. In type-II rivers, i.e.,  
507 rivers characteriszed by the co-occurrence of labile and refractory terrestrial POM and large  
508 temporal variability, phytoplankton POM is usually positively linked to temperature and  
509 negatively linked to river flow, whereas labile and refractory terrestrial POM are bothis  
510 positively linked to SPM and/or river flow. InterestinglyPrecisely, labile terrestrial POM is  
511 usually better linked to river flow and refractory terrestrial POM to SPM (e.g., Hérault River in  
512 Fig. 6 and Rhône River in Fig. A5). In the Têt River, anthropogenic POM was linked to nitrate  
513 concentration (Fig. A5). In rivers characteriszed by phytoplankton and terrestrial-POM  
514 composition with moderate (Type III) or high (Type IV) seasonality, terrestrial POM was  
515 almost always positively linked to river flow and/or SPM concentration, while phytoplankton  
516 was usually linked with chlorophyll *a* concentration (e.g., Charente and Rance Rivers on Fig.  
517 6, Landes and Sseine Rivers on Fig. A5).

**Milieu (I)**Adj.  $R^2 0.76$ **Hérault (II)**Adj.  $R^2 0.40$ **Charente C (III)**Adj.  $R^2 0.50$ **Rance C (IV)**Adj.  $R^2 0.35$ 



519

520 **Figure 6** Redundancy analyses (correlation circles) of rivers standing for each type of river.  
521 Black arrows represent explained variables (concentration of POC sources) and red arrows  
522 represent explaining variables (environmental variables). River types are recalled (Roman  
523 numerals). POC<sub>lt</sub>-LTPOC = Labile terrestrial POC; RTPOCPOC<sub>lt</sub> = Refractory terrestrial POC;  
524 POC<sub>φ</sub>POC = Phytoplankton POC; Chl *a* = chlorophyll *a*; Phaeo. = phaeopigments; M. wind =  
525 meridional wind; Z. wind = zonal wind; R. flow = river flow; Temp. = temperature; Irrad. =  
526 Irradiance; NH<sub>4</sub><sup>+</sup> = ammonium; NO<sub>3</sub><sup>-</sup> = nitrate; PO<sub>4</sub><sup>3-</sup> = phosphates-; Adj. R<sup>2</sup> = adjusted R<sup>2</sup>.

527

528 At last, another RDA was performed, gathering all rivers to relate environmental parameters to  
529 the mean annual POM composition at the multi-rivers scale (Fig. 7). As anthropogenic POM  
530 was only detected in two rivers (Orb, Têt), it was not included in the multi-rivers analysis to  
531 avoid analysis bias. At this scale, phytoplankton is strongly positively linked-correlated to  
532 agricultural surfaces and conductivity, labile terrestrial material to soil erosion rate and soil  
533 organic carbon content and podzol coverage, and refractory terrestrial material to river flow and  
534 water temperature catchment slope; refractory terrestrial material is also and negatively  
535 correlated to soil useful reserves of water (all correlations are significant; Fig. A6). Note that  
536 the phytoplankton and labile terrestrial matter, as well as their related environmental variables,  
537 are negatively correlated.





Figure 7 Multi-rivers redundancy analysis. Black arrows represent explained variables (relative proportions), red arrows represent explaining variables (environmental variables) and numbers are river identifiers (cf. Fig.1). R. flow = river flow; %OC soil = percentages of organic carbon in soil; Soil erosion = soil erosion rate; Rain. = precipitations;  $\text{NO}_3^-$  = nitrates concentration; Useful reserve = Useful reserve in soil; Conduc. = conductivity; %Agri. Surf. = Proportion of agricultural surface; %Podzol = Proportion of podzol coverage; Slope = Catchment slope; Adj.  $R^2$  = adjusted  $R^2$ .

## 4. Discussion

### 4.1. Bulk data-POM and source signatures in temperate rivers

550 Over the 23 studied rivers,  $\delta^{13}\text{C}$ ,  $\delta^{15}\text{N}$ , and C/N ratios of bulk POM ranged between -35.2 and  
551 -24.5 ‰, -0.3 and 12.6 ‰, and 3 and 23.4 mol/mol, respectively. This corresponds to usual  
552 values recorded for riverine POM over temperate systems, except for the lowest C/N ratios  
553 (Ferchiche et al., 2024; Kendall et al., 2001; Ogrinc et al., 2008).

554 In the present study, isotopic and elemental signatures of terrestrial POM and phytoplankton  
555 were determined from subsets of the bulk data sets following the approaches of Savoye et al.  
556 (2012), Liénart et al. (2017) and Ferchiche et al. (2025, 2024). It has the double advantage of  
557 1) taking into account the reworking of terrestrial POM within the river and thus discriminating  
558 labile from refractory terrestrial POM, and 2) taking into account the variability of  
559 phytoplankton signature over time, due to differences in growth conditions (see below). Labile  
560 terrestrial POM mainly appears during high river flow (Fig. 6 and A5; Savoye et al., 2012) and  
561 is usually composed of riparian litter (e.g., Veyssy et al., 1998). In the studied rivers,  $\delta^{13}\text{C}$ ,  $\delta^{15}\text{N}$   
562 and C/N ratio of labile terrestrial POM ranged between  $-28.9 \pm 0.8$  ‰ and  $-26 \pm 0.9$  ‰,  $3.7 \pm$   
563  $0.6$  ‰ and  $6.6 \pm 0.9$  ‰, and  $8.8 \pm 0.4$  and  $17 \pm 3.2$  mol/mol, respectively. These values are very  
564 similar to values found in other temperate systems like the Gironde Estuary ( $\delta^{13}\text{C} = -28.7 \pm 0.9$   
565 ‰; Savoye et al., 2012), the Sava River ( $\delta^{13}\text{C} = -28 \pm 5$  ‰;  $\delta^{15}\text{N} = 5 \pm 2$  ‰; C/N =  $33 \pm 15$   
566 mol/mol; Ogrinc et al., 2008) or Taiwanese rivers ( $\delta^{13}\text{C} = -26.6 \pm 1.8$  ‰; C/N =  $31.1 \pm 23.4$   
567 mol/mol; Hilton et al., 2010) and very similar to direct measurement of C3 plants ( $\delta^{13}\text{C} = -28.1$   
568  $\pm 2.5$  ‰; O’Leary, 1981 and references therein;  $\delta^{13}\text{C} = -28 \pm 1.3$  ‰;  $\delta^{15}\text{N} = 0.8 \pm 2.9$  ‰; C/N  
569 =  $39.6 \pm 25.7$  mol/mol; Dubois et al., 2012;  $\delta^{13}\text{C} = -27.9 \pm 0.1$  ‰; Fernandez et al., 2003).  
570 Refractory terrestrial POM is terrestrial POM that has undergone large reworking within river  
571 water, river sediment or even the estuarine-maximum turbidity zone (e.g., Etcheber et al., 2007;  
572 Veyssy et al., 1998). In the studied rivers where it was found,  $\delta^{13}\text{C}$ ,  $\delta^{15}\text{N}$  and C/N ratios of  
573 refractory terrestrial POM ranged between  $-28 \pm 0.7$  ‰ and  $-25.9 \pm 0.4$  ‰,  $3.2 \pm 0.8$  ‰ and  $6.7 \pm$   
574  $1.4$  ‰, and  $5.8 \pm 1.4$  and  $8.8 \pm 3.1$  mol/mol, respectively. These values are very similar to  
575 the large gradient of refractory POM origins those found in other temperate systems like the  
576 Gironde Estuary (France) (resuspended sediment,  $\delta^{13}\text{C} = -25.2 \pm 0.3$  ‰;  $\delta^{15}\text{N} = 5.5 \pm 0.4$  ‰;  
577 C/N =  $8.5 \pm 0.8$  mol/mol; Savoye et al., 2012), Taiwanese rivers (petrogenic POM,  $\delta^{13}\text{C} = -23.6 \pm 1.1$  ‰;  
578 C/N =  $6.5 \pm 1.6$  mol/mol; Hilton et al., 2010) and in the Pearl River (China) (soil,  
579  $\delta^{13}\text{C}$ : between  $-28.3 \pm 0.8$  ‰ and  $-21.7 \pm 0.7$  ‰; C/N: between  $8.9 \pm 1.1$  and  $17.9 \pm 3.6$  mol/mol;  
580 Yu et al., 2010).

581 Isotopic signatures of phytoplankton vary depending on biogeochemical conditions and  
582 processes like nutrient availability and utilis~~z~~ation, growth rate and limitation (e.g., Fry, 1996;  
583 Liénart et al., 2017; Lowe et al., 2014; Miller et al., 2013; Savoye et al., 2003; Sigman et al.,  
584 2009; Yan et al., 2022) and can be estimated using measured environmental parameters  
585 (Ferchiche et al., 2024, 2025; Liénart et al., 2017; Savoye et al., 2012). For the seven rivers  
586 where phytoplankton isotopic signatures were found variable over time, phytoplankton  $\delta^{13}\text{C}$  or

587  $\delta^{15}\text{N}$  were correlated to: concentrations and ratio of chlorophyll *a* and phaeopigments, water  
588 temperature, nitrate concentration and/or conductivity (Tab. 2). Chlorophyll *a* and  
589 phaeopigments concentrations are direct proxies of phytoplankton fresh and degraded  
590 biomasses and are related to phytoplankton growth and decay, two processes that increase  
591 phytoplankton  $\delta^{13}\text{C}$  (Golubkov et al., 2020; Michener and Kaufman, 2007 and references  
592 therein). Similar processes may explain phytoplankton- $\delta^{15}\text{N}$  increase with chlorophyll *a*  
593 increase. An increase in water temperature accelerates bio-mediated carbon remineralisation  
594 processes, bringing a lower  $\delta^{13}\text{C}$  value than  $\text{CO}_2$  coming from water-atmosphere equilibration  
595 and rock-leaching  $\text{CO}_2$  (Polsenaere et al., 2013 and references therein). Consequently,  
596 phytoplankton  $\delta^{13}\text{C}$  decreases as phytoplankton uses remineralised  $\text{CO}_2$  and thus as water  
597 temperature increases. Phytoplankton  $\delta^{15}\text{N}$  depends on N-nutrient origin and availability  
598 (Savoye et al., 2003 and references therein). Especially, it increases with nutrient concentration  
599 decrease (Sigman et al., 2009) as reported for the Rance River (Tab. 2). Water conductivity  
600 could be considered as a proxy of water mass and thus of nitrate origin. This may explain the  
601 relationship between phytoplankton  $\delta^{15}\text{N}$  and water conductivity in the Orb River (Tab. 2).

602 In the studied rivers, phytoplankton  $\delta^{13}\text{C}$ ,  $\delta^{15}\text{N}$  and C/N ratio ranged between  $-34.9 \pm 0.4$  and  $-23.8 \pm 0.6 \text{‰}$ ,  $4.3 \pm 0.8$  and  $13.2 \pm 1.8 \text{‰}$ , and  $4.8 \pm 0.9$  and  $10 \pm 0.9 \text{ mol/mol}$ , respectively.  
603 This is similar to values reported for the Loire River, another French river ( $-30.6 \leq \delta^{13}\text{C} \leq -25.0$   
604  $\text{‰}$ ;  $3.0 \leq \delta^{15}\text{N} \leq 10.4 \text{‰}$ ; C/N =  $7.2 \pm 0.6 \text{ mol/mol}$ : Ferchiche et al., 2024), but narrower ranges  
605 can be found in the literature. In the Sava River (Eastern Europe), phytoplankton signature was  
606  $-30.4 \pm 2.1 \text{‰}$ ,  $5.0 \pm 1.5 \text{‰}$  and  $6.5 \pm 1.5 \text{ mol/mol}$  for  $\delta^{13}\text{C}$ ,  $\delta^{15}\text{N}$  and C/N ratio, respectively  
607 (Ogrinc et al., 2008), similar to that of Indian ( $\delta^{13}\text{C} = -30.6 \pm 1.7 \text{‰}$ ,  $\delta^{15}\text{N} = 7.0 \pm 2.3 \text{‰}$ ;  
608 Gawade et al., 2018) and Texan ( $\delta^{13}\text{C} = -31.4 \text{‰}$ ; Lebreton et al., 2016) rivers. Lower  $\delta^{13}\text{C}$   
609 values ( $\leq -32 \text{‰}$ ) were also found (Finlay et al., 2010; Hellings et al., 1999; Sato et al., 2006;  
610 Savoye et al., 2012). However, values of elemental and isotopic ratios for riverine  
611 phytoplankton are scarce in the literature. Indeed, it is not easy to estimate phytoplankton  
612 signature since it cannot be separated from other particles. Thus, literature estimates may not  
613 be perfectly representative of the variability of phytoplankton isotopic signatures.  
614

## 615 **4.2. Watershed characteristics~~s~~ drive ~~on~~—spatial dynamics of POM 616 composition**

617 At the annual scale, we observed deep variations between studied rivers ~~concerning regarding~~  
618 the mean POC proportion of the different sources ( $5 \leq \text{phytoplankton} \leq 80 \text{ \%}$ ;  $17 \leq \text{labile}$   
619  $\text{terrestrial POC} \leq 95 \text{ \%}$ ;  $0 \leq \text{refractory terrestrial POC} \leq 39 \text{ \%}$ ).

620 Interestingly, phytoplankton proportions ~~were~~ highly correlated to the proportion of  
621 agriculture surface areas ~~and conductivity and in a less extent to river nitrate concentration (Fig.~~  
622 ~~7). (Fig. 7;  $R^2 = 0.59$  or even 0.72 when Seudre River is not included in the statistics) and nitrate~~

623 ~~concentration (Fig. 7;  $R^2 = 0.41$  when Seudre River is not included in the statistics).~~ Such  
624 relationship between agriculture surface ~~area~~ and phytoplankton is well-known, as agricultural  
625 activities increase nutrient inputs ~~s~~ to river bodies (Khan and Mohammad, 2014), leading to better  
626 conditions for phytoplankton growth (Dodds and Smith, 2016; Minaudo et al., 2015).

627 ~~Also interestingly, the proportions of labile terrestrial matter were strongly positively linked~~  
628 ~~to soil erosion and soil organic carbon content, soil erosion, and the podzol~~  
629 ~~coverage and sandy rock coverage (Fig. 7, Fig. A6), indicating a strong relationship between~~  
630 ~~terrestrial matter in rivers and soil nature with undecomposed and fresh detrital matter~~  
631 (McCorkle et al., 2016). They are also negatively correlated to phytoplankton proportions and

632 ~~their related environmental parameters. Rivers which POM is dominated by labile terrestrial~~  
633 ~~POM (mainly rivers of type I) flow through catchments dominated by sandy rocks and podzol.~~  
634 ~~This kind of soil is submitted to soil erosion and releases large amounts of colored~~  
635 ~~dissolved organic carbon, favouring the input of terrestrial material (soil erosion) and~~  
636 ~~disfavouring Podzol coverage can explain the dominance of labile terrestrial matter as much as~~  
637 ~~the inability for phytoplankton to grow in the river water because of the turbidity due to the~~  
638 ~~dissolved organic carbon—a significant biomass, this soil type being highly favourable to~~  
639 ~~leaching particulate and coloured dissolved components in the river and drastically limiting the~~  
640 ~~river euphotic zone (Canton et al., 2012; Polsenaere et al., 2013).~~

641 ~~The dominance proportions of refractory terrestrial matter are correlated to the catchment slope~~  
642 ~~and negatively correlated to linked to poor useful reserve of water (Fig. 7, Fig. A6). Rivers for~~  
643 ~~which POM- is partly composed of refractory terrestrial POM (most of the rivers of type II)~~  
644 ~~flow through catchments of and high slope inform of specific catchment settings, where more~~  
645 ~~mountainous surfaces, which are associated with shallower, poorer topsoil and more~~  
646 ~~outcropping bedrock. It favours more reactive and abrupt transfer of water to the river, leading~~  
647 ~~to enhanced episodes of sediment resuspension, as well as permitting a rock-derived POM~~  
648 ~~weathering (Copard et al., 2018; Higueras et al., 2014; Yaalon, 1997).~~

#### 649 **4.3. Temporal dynamics of POM composition and river ~~dynamics~~**

650 **typology**

651 ~~If average quantitative difference between rivers can be input to differences in the catchment~~  
652 ~~characteristics (see section 4.2). In aquatic systems seasonally, phytoplankton likely appears~~  
653 ~~during spring and summer in favorable conditions, related to low discharge, high-temperature~~  
654 ~~conditions and enough nutrients to support its growth, while in winter, high turbidity and low-~~  
655 ~~temperature conditions limit its presence (Turner et al., 2022). Quantitative differences between~~  
656 ~~rivers can be due to differences in nutrient availability, either because of anthropic mitigation~~  
657 ~~(Minaudo et al., 2015), competition with other nutrient users (Desey et al., 2012; Minaudo et~~  
658 ~~al., 2016) or sewage inputs (Codiga et al., 2022) in addition to agricultural inputs (see section~~  
659 ~~4.2). Terrestrial material likely appears during winter conditions, related to floods that transport~~

660 great amounts of terrestrial material (Dalzell et al., 2007). Such a seasonal dichotomy between  
661 phytoplankton and terrestrial POM was clearly visible for most of the studied rivers (Fig. 4),  
662 especially for type-IV and type-III rivers, but even for some of those highly dominated by the  
663 labile terrestrial POM (e.g., Milieu and Tagon Rivers; Type-I rivers). This was illustrated by  
664 the relationships between phytoplankton POM and chlorophyll *a* concentration and/or  
665 temperature (as proxies of favourable conditions for phytoplankton production) on the one  
666 hand, and between labile terrestrial POM and river flow and/or SPM concentration on the other  
667 hand (Fig. 6 and A5). This dichotomy in POM composition was also reported in other similar  
668 studies (e.g., Kelso and Baker, 2020; Lu et al., 2016). In rivers where refractory terrestrial POM  
669 was present in addition to the labile one (type-II rivers), ~~it was interesting to see that both~~  
670 ~~terrestrial sources were linked to river flows and SPM concentrations. More precisely, it was~~  
671 ~~interesting to see that~~ the refractory terrestrial POM was more related to SPM concentration  
672 than river flow and inversely for the labile terrestrial POM. This indicates that labile and  
673 refractory terrestrial POM were preferentially associated with direct river input and sediment  
674 resuspension, respectively. The origin of the refractory terrestrial POM may be  
675 fossil/bedrock/petrogenic OM (e.g., Copard et al., 2022; Hilton et al., 2010; Sun et al., 2021)  
676 brought by river flow (in quantity undetectable in the bulk POM using our tools), especially in  
677 Type-II rivers which watersheds are characterized by high slopes (Fig. 7). This POM can be  
678 and then accumulated in the downstream sediments and be resuspended (in quantity calculable  
679 in the bulk POM using our tools). Refractory terrestrial POM may also come from, and/or labile  
680 terrestrial POM brought by the river flow and then accumulated and reworked/decayed until  
681 refractory POM in the sediment (e.g., Etcheber et al., 2007; Savoye et al., 2012), which can be  
682 resuspended.

683 Sewage POM was detected in two of the studied rivers, ~~but~~ with different associated temporal  
684 dynamics. In the Têt River, because the former WWTP was dysfunctional, a new one replaced  
685 it in late 2007 (<https://www.assainissement.developpement-durable.gouv.fr/pages/data/fiche-060966136002>, last visit 10/09/24). This explains the shift in sewage POM between the two  
686 studied periods (2006-2007 versus 2008-2010 without anthropogenic POM). In the Orb River,  
687 sewage POM was detected throughout the studied periods. The WWTP is located only a few  
688 kilomet~~reers~~ upstream of the sampling site and is large enough (220~~0~~-000 inhabitant equivalent)  
689 compared to the river flow (annual mean: 23m<sup>3</sup>/s) to make the sewage POM detectable in the  
690 bulk POM using  $\delta^{15}\text{N}$ . Such a result is quite common for urban rivers (e.g., Kelso and Baker,  
691 2020).

#### 693 4.4. Originality of the study

694 ~~The originality of the present study firstly lies in its approach. Even if C and N stable isotopes~~  
695 ~~have been used for decades to investigate POM origins within river waters, the quantification~~  
696 ~~of POM composition (i.e., the relative proportion of each source composing the POM) using~~  
697 ~~mixing models, and especially Bayesian mixing models, is not so common. In addition, most~~

698 of the previous studies either use literature data for phytoplankton isotopic signature (e.g.,  
699 Zhang et al., 2021) or use lake or autochthonous POM as a proxy of phytoplankton (e.g., Kelso  
700 and Baker, 2020). Also, most of these studies use direct measurements of soil or plants to assess  
701 the isotopic signature of terrestrial POM whereas this material is able to rework within the water  
702 column or the sediment, which changes its elemental and isotopic values (e.g., Savoye et al.,  
703 2012). These approaches do not consider that isotopic signatures of phytoplankton and  
704 terrestrial material may change over time. In the present study, we used the approach developed  
705 by Savoye et al. (2012) in an estuary, Liénart et al. (2017) in coastal systems and Ferchiche et  
706 al. (2025, 2024) in a river to assess the elemental and isotopic signatures from subsets of bulk  
707 POM and when needed, empirical equations. This approach has the great advantage of 1) using  
708 signatures dedicated to the sampling area and 2) taking into account the potential variability of  
709 these signatures over time, i.e., depending on the environmental conditions for phytoplankton  
710 growth and taking into account its decay for phytoplankton and terrestrial POM. Especially, we  
711 were able to discriminate labile from refractory terrestrial POM in some rivers, as Savoye et al.  
712 (2012) in an estuary. Another great originality of the present study lies in the multi-systems  
713 approach: studying 23 rivers in a single study allowed the detection of four types of river  
714 functioning regarding the POM composition and its temporal dynamics. It also highlights the  
715 great influence of land use (agriculture) and characteristics (erosion, organic carbon content)  
716 on the POM composition of rivers. At last, using  $\delta^{13}\text{C}$ ,  $\delta^{15}\text{N}$  and C/N ratio all together allowed  
717 either to perform mixing models with up to four end-members or to study separately POC and  
718 PN composition. It showed that POC and PN display very similar compositions and dynamics  
719 in rivers.

## 720 **5. Synthesis, originality of the study and perspectives**

721 The present study proposes a comprehensive estimation of POM composition and its spatial  
722 and seasonal variability in temperate rivers. Thanks to the inclusion of twenty three rivers,  
723 encompassing large gradients of environmental conditions under a temperate climate, a river  
724 typology is proposed based on the POM composition and its temporal dynamics. In type I  
725 rivers, POM is dominated by labile terrestrial material all year long. This material is mainly  
726 associated with suspended particulate matter. Phytoplankton slightly contributes, especially  
727 during summertime. Type II rivers are mainly characterized by the presence of both labile and  
728 refractory terrestrial material, in addition to phytoplankton. The temporal variability between  
729 these sources is high but the seasonality is not always pronounced even if phytoplankton and  
730 terrestrial POM can dominate the POM composition during summer and winter, respectively.  
731 Nevertheless, labile terrestrial POM is mainly related to river flow and refractory terrestrial  
732 POM to SPM, indicating the sedimentary origin of the latter. In type III rivers, POM is  
733 composed of phytoplankton and labile terrestrial material. The seasonality of POM composition  
734 is not very pronounced even if the contribution of labile terrestrial POM is deeply related to  
735 river flows. Type III is an intermediary between type I and type IV. In type IV rivers, POM is

736 ~~also composed of phytoplankton and labile terrestrial material but the seasonality of POM~~  
737 ~~composition is very pronounced with a clear balance between high phytoplankton contribution~~  
738 ~~in summer and high terrestrial contribution in winter. Labile terrestrial POM is deeply related~~  
739 ~~to river flow. Beyond this typology, the main difference in POM composition between the~~  
740 ~~studied rivers is that the phytoplankton contribution to the POM composition is related to the~~  
741 ~~proportion of agricultural surface in the watershed and the contribution of labile terrestrial POM~~  
742 ~~is related to soil erosion and organic carbon content in the watershed. The present study~~  
743 ~~provides a comprehensive assessment of POM composition and its spatial and seasonal~~  
744 ~~variability in temperate rivers. By including twenty-three rivers spanning a wide range of~~  
745 ~~environmental conditions under a temperate climate, a river-dynamics typology is proposed~~  
746 ~~based on POM composition and its temporal patterns. In type-I rivers, POM is dominated by~~  
747 ~~labile terrestrial material throughout the year. This material is mainly associated with suspended~~  
748 ~~particulate matter. Phytoplankton makes a slight contribution, especially during summer. Type-~~  
749 ~~II rivers are characterised by the presence of both labile and refractory terrestrial material, along~~  
750 ~~with phytoplankton. The variability between these sources over time is high, but seasonality is~~  
751 ~~not always evident, although phytoplankton and terrestrial POM can dominate the POM~~  
752 ~~composition during summer and winter, respectively. Nonetheless, if both terrestrial sources~~  
753 ~~are primarily linked to river flow and SPM, a better coupling of refractory terrestrial POM with~~  
754 ~~SPM indicates that this material is probably stored in sediments and resuspended, whatever its~~  
755 ~~origin (soil, litter, petrogenic POM). In type-III rivers, POM consists of phytoplankton and~~  
756 ~~labile terrestrial material. The seasonality of POM composition is not very pronounced, though~~  
757 ~~the contribution of labile terrestrial POM is closely related to river flow. Type III is an~~  
758 ~~intermediate between type I and type IV. In type-IV rivers, POM is also composed of~~  
759 ~~phytoplankton and labile terrestrial material, but the seasonality is highly marked, with a clear~~  
760 ~~shift from high phytoplankton contribution in summer to high terrestrial contribution in winter.~~  
761 ~~Labile terrestrial POM remains closely associated with river flow. Beyond this typology, the~~  
762 ~~main differences in POM composition between the studied rivers is related to catchment~~  
763 ~~inherent properties. The contribution of phytoplankton is correlated with the proportion of~~  
764 ~~agricultural coverage, while the contribution of labile terrestrial POM is linked to leached OM-~~  
765 ~~rich thick soil features and the refractory terrestrial POM to thin OM-poor soils with high rock-~~  
766 ~~derived features.~~

767 The originality of the present study lies firstly in its approach. Even if C and N stable isotopes  
768 have been used for decades to investigate POM origins within river waters, the quantification  
769 of POM composition (i.e., the relative proportion of each source composing the POM) using  
770 mixing models, especially Bayesian mixing models, is not so common. Most previous studies  
771 either use literature data for phytoplankton isotopic signature (e.g., (Zhang et al., 2021) or use  
772 lake or autochthonous POM as a proxy of phytoplankton (e.g., (Kelso and Baker, 2020). Also,  
773 most of these studies use direct measurements of soil or plants to assess the isotopic signature

774 of terrestrial POM, although this material may rework within the water column or sediment,  
775 changing its elemental and isotopic values (e.g., (Savoye et al., 2012). These approaches do not  
776 consider the temporal variability of phytoplankton and terrestrial material isotopic signatures.  
777 In the present study, we used the approach developed by (Savoye et al., (2012) in an estuary,  
778 (Liénart et al., (2017) in coastal systems, and (Ferchiche et al., (2024, 2025) in a river to assess  
779 elemental and isotopic signatures from subsets of bulk POM and, when needed, empirical  
780 equations. This approach has the advantage of 1) using signatures dedicated to the sampling  
781 area and 2) taking into account the potential variability of these signatures over time, i.e.,  
782 depending on environmental conditions for phytoplankton growth and its decay for  
783 phytoplankton and terrestrial POM. Especially, we discriminated labile from refractory  
784 terrestrial POM in some rivers, as (Savoye et al., (2012) did in an estuary. Another great  
785 originality of the present study lies in the multi-systems approach: studying 23 rivers in a single  
786 study allowed the detection of four types of river functioning regarding the POM composition  
787 and its temporal dynamics, which has not been performed before. It also highlights the great  
788 influence of land use (agriculture) and characteristics (erosion, organic carbon content, type of  
789 soil) on the POM composition of rivers. At last, the multi-parameter use of  $\delta^{13}\text{C}$ ,  $\delta^{15}\text{N}$ , and C/N  
790 ratio allowed either to perform mixing models with up to four end-members or to study POC  
791 and PN composition separately. It showed that POC and PN display very similar compositions  
792 and dynamics in rivers.

793 The originality of this study mainly lies in 1) the approach used to determine the elemental and  
794 isotopic signatures of POM sources, which allowed to discriminate labile from refractory  
795 terrestrial POM and to take into account, when any, the variability of the signatures over time,  
796 and 2) determining a typology of temperate rivers based on the POM composition and its  
797 temporal dynamics.

798 Overall, this study, which focuses on the River-Estuary Interface, brings meaningful  
799 information for the comprehension of C and N cycles along the LOAC and especially the  
800 behaviour, dynamics and drivers of POM that leaves the river and enters the estuary.

801 From a methodological perspective, such a study could be strengthened by the use of non-  
802 exchangeable  $\delta^2\text{H}$  as an additional tool to even better distinguish and quantify more sources in  
803 mixing models. This tool has been recently shown to be powerful for such purposes (Ferchiche  
804 et al., 2025). From a fundamental perspective, aggregating more datasets from other temperate  
805 rivers would allow testing the robustness of this typology and probably detecting additional  
806 types, but also datasets from polar and tropical rivers to perform an even more comprehensive  
807 study at a global climate scale. Clearly, the approach developed in the present study is  
808 transferable in to other other regions of the planet and used at broader space-spatial scales.  
809 In addition, a similar study dedicated to the estuarine systems would even increase our  
810 comprehensive understanding of the origin and fate of POM along the Land-Ocean Aquatic

811 Continuum by complementing the present study dedicated to the River-Estuary Interface and  
812 those of Liénart et al.<sup>5</sup> (2017, 2018) dedicated to the coastal systems. <sup>6</sup>  
813

814

## Appendix

### 815 Toward a typology of river functioning: a 816 comprehensive study of the particulate organic 817 matterPOM composition at the multi-rivers scale

818

819 Ferchiche F.<sup>1</sup>, Liénart C.<sup>1</sup>, Charlier K.<sup>1</sup>, Deborde J.<sup>2,3</sup>, Giraud M.<sup>4</sup>, Kerhervé P.<sup>5</sup>, Polsenaere P.<sup>1,3</sup>, Savoye  
820 N.<sup>1\*</sup>

821 <sup>1</sup> Univ. Bordeaux, CNRS, EPHE, Bordeaux INP, UMR 5805 EPOC, F-33600 Pessac, France

822 <sup>2</sup> Univ. Pau & Pays Adour, CNRS, E2S UPPA – MIRA, UMR 5254 IPREM, F-64000 Pau, F-64600 Anglet, France

823 <sup>3</sup> Ifremer, COAST, F-17390 La Tremblade, France

824 <sup>4</sup> MNHN, CRESCO, Station Marine de Dinard, F-35800 Dinard, France

825 <sup>5</sup> Univ. Perpignan, CNRS, UMR 5110 CEFREM, F-66860 Perpignan, France

826

827 \*Corresponding author

828 [nicolas.savoye@u-bordeaux.fr](mailto:nicolas.savoye@u-bordeaux.fr)

829 Station marine - 2 rue du Professeur Jolyet 33120 ARCACHON

830

831

832

833

834

835

836

837

838

839

840

841

842

843 Keywords : River-Estuary Interface; particulate organic matter; isotopes; multi-ecosystems  
844 study

846 Table A1 Summary of data availability and origin (X means that the data were available; \* means that the data were retrieved from the Naïades  
 847 database; \*\* means that the data were retrieved from the Météo France database). n = number of sampling dates; SPM = Suspended Particulate  
 848 Matter; POC or PN = Particulate Organic Carbon or Nitrogen; Chl *a* = Chlorophyll *a*; Phaeo = Phaeopigments.

River	Samplings							Parameters																	
	Dates	Periodicity	n	Latitude	Longitude	$\delta^{13}\text{C}$	$\delta^{15}\text{N}$	C/N ratio	SPM	POC	PN	Chl <i>a</i>	Phaeo	Water temperature	pH	Conductivity	$\text{NH}_4^+$	$\text{NO}_2^-$	$\text{NO}_3^-$	$\text{PO}_4^{3-}$	Flow	Rainfall	Air temperature	Air wind direction	Wind intensity
Seine	06/2014 to 06/2015	monthly	13	49.306667	1.242500	X	X	X	X*	X	X	X	X	X*	X*	X*	X*	X*	X*	X*	X	X**	X**	X**	
Orne	06/2014 to 06/2015	monthly	13	49.179722	-0.349167	X	X	X	X*	X	X	X	X	X*	X*	X*	X*	X*	X*	X*	X	X**	X**	X**	
Rance	06/2014 to 05/2015	monthly	12	48.491667	-2.001389	X	X	X	X*	X	X	X	X	X*	X*	X*	X*	X*	X*	X*	X	X**	X**	X**	
Elorn	01/2014 to 06/2015	monthly	17	48.450556	-4.248333	X	X	X	X*	X	X	X	X	X*	X*	X*	X*	X*	X*	X*	X	X**	X**	X**	
Aulne	01/2014 to 06/2015	monthly	17	48.212778	-4.094444	X	X	X	X*	X	X	X	X	X*	X*	X*	X*	X*	X*	X*	X	X**	X**	X**	
Loire	10/2009 to 07/2012	bi-monthly	67	47.392095	-0.860351	X	X	X	X*	X	X	X	X	X	X	X	X	X	X	X	X	X**	X**	X**	
Sevre niortaise	03/2014 to 03/2015	monthly	13	46.315348	-1.003891	X	X	X	X*	X	X	X	X	X*	X*	X*	X*	X*	X*	X*	X	X**	X**		
Charente	03/2014 to 03/2015	monthly	13	45.868056	-0.713056	X	X	X	X*	X	X	X	X	X*	X*	X*	X*	X*	X*	X*	X	X**	X**	X**	
Seudre	03/2014 to 09/2015	monthly	15	45.674027	-0.933123	X	X	X	X*	X	X	X	X	X*	X*	X*	X*	X*	X*	X*	X	X**	X**	X**	
Porge	01/2008 to 02/2009	monthly	14	44.789868	-1.161181	X		X	X	X	X	X	X								X*	X**	X**	X**	
Renet	02/2008 to 02/2009	bi-monthly	23	44.714466	-1.044013	X		X	X	X	X	X	X								X*	X**	X**	X**	
Milieu	02/2008 to 02/2009	monthly	13	44.697326	-1.022532	X		X	X	X	X	X	X								X*	X**	X**	X**	
Cirès	02/2008 to 02/2009	monthly	13	44.759820	-1.110657	X		X	X	X	X	X	X								X*	X**	X**	X**	
Lanton	02/2008 to 02/2009	monthly	13	44.700283	-1.024385	X		X	X	X	X	X	X								X*	X**	X**	X**	
Tagon	02/2008 to 02/2009	bi-monthly	26	44.659049	-0.989050	X		X	X	X	X	X	X								X*	X**	X**	X**	
Landes	02/2008 to 02/2009	monthly	12	44.616912	-1.109066	X		X	X	X	X	X	X								X*	X**	X**	X**	
Leyre	02/2008 to 02/2015	bi-monthly or monthly	59	44.626389	-0.996111	X	X	X	X*	X	X	X	X	X	X*	X*	X*	X*	X*	X*	X	X**	X**	X**	
Adour	04/2013 to 05/2018	monthly	24	43.498880	-1.294899	X	X	X	X	X	X	X	X	X		X	X	X			X	X**	X**	X**	
Têt	01/2006 to 05/2010	monthly	52	42.713704	2.993488	X	X	X	X	X	X				X	X	X	X*	X*	X*	X	X**	X**	X**	
Aude	01/2006 to 05/2010	monthly	52	43.244281	3.152733	X	X	X	X	X	X				X	X	X	X*	X*	X*	X	X**	X**	X**	
Orb	01/2006 to 05/2010	monthly	52	43.285004	3.281278	X	X	X	X	X	X				X	X	X	X*	X*	X*	X	X**	X**		
Hérault	01/2006 to 05/2010	monthly	52	43.359415	3.435398	X	X	X	X	X	X				X	X	X	X*	X*	X*	X	X**	X**	X**	
Rhône	12/2003 to 01/2011	monthly	105	43.678724	4.621188	X	X	X	X	X	X	X	X	X	X	X	X	X*	X*	X*	X	X**	X**	X**	

850 Table A2 Summary of parameters kept (informative) to perform the local seasonal RDAs, opposite to those not considered (because non-  
 851 informative, not available, or auto-correlated (see section 2.5).

852

River	SPM	Chlorophyll <i>a</i>	Phaeopigment s	Temperature	River flow	pH	Ammonium	Nitrates	Phosphates	Irradiance	Zonal wind	Meridional wind
Seine	X	X	X	X	X	X	X	-	-	-	-	-
Orne	-	X	-	X	X	X	-	X	X	-	X	-
Rance	X	X	-	X	X	-	-	X	-	-	-	-
Elorn	-	X	X	-	X	-	X	-	X	-	-	-
Aulne	X	X	-	-	X	X	-	X	-	-	-	-
Sèvre	X	X	X	-	-	X	-	-	-	-	-	-
Charente	-	X	X	-	X	X	-	-	-	-	-	X
Seudre	X	X	X	X	X	-	-	-	-	X	X	X
Porge	X	X	X	-	-						X	X
Cirès	X	X	X	X	X						-	-
Milieu	X	X	X	-	X						-	X
Lanton	X	X	X	X	-						-	X
Renet	X	X	X	X	-						-	-
Tagon	X	X	X	X	X						-	-
Leyre	X	X	X	X	-	X	-				-	-
Landes	X	X	X	-	X						-	X
Adour	X	-	X	X	X		X	-			-	-
Têt	X			X	X	X	-	X	-		-	X
Aude	X			X	X	-	X	-	X	X	-	-
Orb	X			X	X	X	-	X	X		-	-
Hérault	X			X	X	-	-	-	X	-	-	X
Rhône	X			-	X	-				-	-	-

X = Kept

= No data

- = Non kept

= Auto-correlated

853

854

855

856

857

858

859

860

861

862

863

864

865

866

867

868

869

870








Figure A1 Diagram detailing the regionalization method, adapted from Souissi et al. (2000).










875

876



877


878




879

880 Figure A2 Temporal variations of matter characteristics for representative rivers along the studied periods for  $\delta^{13}\text{C}$  (left axis; black dotted line) and  
 881  $\delta^{15}\text{N}$  (right axis; blue line) (first column); C/N (left axis; black dotted line) and POC/chl *a* (right axis; blue line) ratios (second column); SPM (left  
 882 axis; black dotted line), POC (right axis; blue line) and chlorophyll *a* (right axis; blue dotted line) concentrations (third column) and river flow (left  
 883 axis; black dotted line) and temperature (right axis; blue line) (fourth column).









887 Figure A3  $\delta^{13}\text{C}$ ,  $\delta^{15}\text{N}$ , N/C and/or C/N values of bulk POM (black crosses) and sources. The latter are presented as closed circles (average) and  
888 bars (standard deviation) when the signatures were constant over time and by colored area when at least one of the proxies was variable over time  
889 (see Table 2). This colored area corresponds to the dispersion of the values, including their uncertainties.













890  
891  
892  
893  
894




895 Figure A4 Temporal dynamic (rectangle graphs) and (inter-)annual mean (pie charts) of PN  
896 source proportions. Sources are phytoplankton (green) and labile terrestrial material (brown).

897  
898  
899




**Tagon (I)**Adj.  $R^2$  0.70**Lanton (I)**Adj.  $R^2$  0.93**Leyre (I)**Adj.  $R^2$  0.41**Cirès (I)**Adj.  $R^2$  0.74**Renet (I)**Adj.  $R^2$  0.97**Seudre (I)**Adj.  $R^2$  0.72**Porge (II)**Adj.  $R^2$  0.38**Adour (II)**Adj.  $R^2$  0.53**Landes (II)**Adj.  $R^2$  0.61**Orne C (II)**Adj.  $R^2$  0.72**Orne N (II)**Adj.  $R^2$  0.74**Charente N (II)**Adj.  $R^2$  0.43




Seine C (IV)

Adj.  $R^2$  0.95




Aulne C (IV)

Adj.  $R^2$  0.92




Elorn C (IV)

Adj.  $R^2$  0.24




Seine N (IV)

Adj.  $R^2$  0.80




Aulne N (IV)

Adj.  $R^2$  0.74




Elorn N (IV)

Adj.  $R^2$  0.04




Rance N (IV)

Adj.  $R^2$  0.56



Sèvre (IV)

Adj.  $R^2$  0.01



906

907

908


909

910


911

912


Aude (axis 1-2) (III)

Adj.  $R^2 0.36$ 


Têt (axis 1-2) (III)

Adj.  $R^2 0.58$ 


Rhône (III)

Adj.  $R^2 0.65$ 


Aude (axis 1-3) (III)

Adj.  $R^2 0.43$ 

Têt (axis 1-3) (III)


Adj.  $R^2 0.57$ 

Orb (III)


Adj.  $R^2 0.68$ 

913


Aude (axis 1-2) (III)

Adj.  $R^2 0.43$ 


Têt (axis 1-2) (III)

Adj.  $R^2 0.57$ 


Rhône (III)

Adj.  $R^2 0.65$ 


Aude (axis 1-3) (III)

Adj.  $R^2 0.43$ 

Têt (axis 1-3) (III)

Adj.  $R^2 0.57$ 

Orb (III)

Adj.  $R^2 0.68$ 


914

915

916  
917  
918  
919

920 **Figure A5** Redundancy analyses (correlation circles) of rivers standing for each type of river.  
921 Black arrows represent explained variables (concentration of POC or PN sources) and red  
922 arrows represent explaining variables (environmental variables). River types are recalled  
923 (Roman numerals). LT<sub>POC</sub> or PN = Labile terrestrial POC or PN; RT<sub>POC</sub> = Refractory terrestrial  
924 POC; φ<sub>POC</sub> or PN = Phytoplankton POC or PN; Anth. POC = Anthropogenic POM; SPM =  
925 Suspended particulate matter; Chl a = chlorophyll a; Phaeo. = phaeopigments; M. wind =  
926 meridional wind; Z. wind = zonal wind; R. flow = river flow; Temp. = temperature; Irrad. =  
927 Irradiance; NH<sub>4</sub><sup>+</sup> = ammonium; NO<sub>3</sub><sup>-</sup> = nitrate; PO<sub>4</sub><sup>3-</sup> = phosphates; Adj. R<sup>2</sup> = adjusted R<sup>2</sup>.

928




929

930 [Figure A6 Correlogram of multi-system RDA parameters, including source proportions and](#)  
 931 [accompanying parameters. Descriptions of environmental parameters can be retrieved in](#)  
 932 [section 2.5. Temperature = Water temperature; SPM = Suspended particulate matter; OC in soil](#)  
 933 [= Organic carbon proportions in soil; WWTP population equivalent = sewage treatment](#)  
 934 [capacities; WWTP ...River flow = sewage treatment capacities to average river flow ratio.](#)

935

936

937



938

939 [Figure A7 Typology of river dynamics following a hierarchical cluster analysis on POM](#)  
 940 [source proportions. The percentages of membership for each type attributed to each river are](#)  
 941 [shown. Left panel: considering all sampling years for all rivers. Right panel: considering only](#)  
 942 [one stream of type I fueling the Bay of Arcachon \(Milieu River\).](#)

943

944

945

946

## 947 **Author contributions**

948 FF: Formal analysis, Investigation, **Visualization****Visualisation**, Writing – original draft. CL:  
949 Formal analysis, **Conceptualization****Conceptualisation**, Supervision, Writing – original draft,  
950 Writing – review & editing. KC: Investigation. JD: Investigation, **Visualization****Visualisation**,  
951 Writing – review & editing. MG: Investigation. PK: Investigation. PP: Investigation,  
952 **Visualization****Visualisation**, Writing – review & editing.

953 NS: **Conceptualization****Conceptualisation**, Formal analysis, Investigation, Methodology,  
954 Supervision, Writing – original draft, Writing – review & editing.

955

## 956 **Competing interest**

957 The authors declare that they have no known competing financial interests or personal  
958 relationships that could have appeared to influence the work reported in this paper.

959

## 960 **Data availability**

961 All POM and environmental data used in this article are stored in Figshare, accessible for the  
962 review process through this private link : <https://figshare.com/s/a7101028e6ab5452c4db>.

963

## 964 **References**

965 Arellano, A. R., Bianchi, T. S., Osburn, C. L., D'Sa, E. J., Ward, N. D., Oviedo-Vargas, D.,  
966 Joshi, I. D., Ko, D. S., Shields, M. R., Kurian, G., and Green, J.: Mechanisms of Organic  
967 Matter Export in Estuaries with Contrasting Carbon Sources, *Journal of Geophysical  
968 Research: Biogeosciences*, 124, 3168–3188, <https://doi.org/10.1029/2018JG004868>, 2019.

969 Barros, G. V., Martinelli, L. A., Oliveira Novais, T. M., Ometto, J. P. H. B., and Zuppi, G.  
970 M.: Stable isotopes of bulk organic matter to trace carbon and nitrogen dynamics in an  
971 estuarine ecosystem in Babitonga Bay (Santa Catarina, Brazil), *Science of The Total  
972 Environment*, 408, 2226–2232, <https://doi.org/10.1016/j.scitotenv.2010.01.060>, 2010.

973 Bate, G. C., Whitfield, A. K., Adams, J. B., Huizinga, P., and Wooldridge, T. H.: The  
974 importance of the river-estuary interface (REI) zone in estuaries, *Water SA*, 28, 271–280,  
975 <https://doi.org/10.4314/wsa.v28i3.4894>, 2002.

976 Bonin, P., Prime, A.-H., Galeron, M.-A., Guasco, S., and Rontani, J.-F.: Enhanced biotic  
977 degradation of terrestrial POM in an estuarine salinity gradient: interactive effects of organic  
978 matter pools and changes of bacterial communities, *Aquatic Microbial Ecology*, 83, 147–159,  
979 <https://doi.org/10.3354/ame01908>, 2019.

980 Borcard, D., Legendre, P., and Gillet, F.: Numerical Ecology with R, Springer, 315 pp., 2011.

981 Bouwman, L., Beusen, A., Glibert, P. M., Overbeek, C., Pawlowski, M., Herrera, J., Mulsow,  
982 S., Yu, R., and Zhou, M.: Mariculture: significant and expanding cause of coastal nutrient  
983 enrichment, *Environ. Res. Lett.*, 8, 044026, <https://doi.org/10.1088/1748-9326/8/4/044026>,  
984 2013.

985 Brett, M. T., Bunn, S. E., Chandra, S., Galloway, A. W. E., Guo, F., Kainz, M. J., Kankaala,  
986 P., Lau, D. C. P., Moulton, T. P., Power, M. E., Rasmussen, J. B., Taipale, S. J., Thorp, J. H.,  
987 and Wehr, J. D.: How important are terrestrial organic carbon inputs for secondary production  
988 in freshwater ecosystems?, *Freshwater Biology*, 62, 833–853,  
989 <https://doi.org/10.1111/fwb.12909>, 2017.

990 Canton, M., Anschutz, P., Coynel, A., Polsenaere, P., Auby, I., and Poirier, D.: Nutrient  
991 export to an Eastern Atlantic coastal zone: first modeling and nitrogen mass balance,  
992 *Biogeochemistry*, 107, 361–377, <https://doi.org/10.1007/s10533-010-9558-7>, 2012.

993 Canuel, E. A. and Hardison, A. K.: Sources, Ages, and Alteration of Organic Matter in  
994 Estuaries, *Annual Review of Marine Science*, 8, 409–434, <https://doi.org/10.1146/annurev-marine-122414-034058>, 2016.

995 Cathalot, C., Rabouille, C., Tisnérat-Laborde, N., Toussaint, F., Kerhervé, P., Buscail, R.,  
996 Loftis, K., Sun, M.-Y., Tronczynski, J., Azoury, S., Lansard, B., Treignier, C., Pastor, L., and  
997 Tesi, T.: The fate of river organic carbon in coastal areas: A study in the Rhône River delta  
998 using multiple isotopic ( $\delta^{13}\text{C}$ ,  $\Delta^{14}\text{C}$ ) and organic tracers, *Geochimica et Cosmochimica  
999 Acta*, 118, 33–55, <https://doi.org/10.1016/j.gca.2013.05.001>, 2013.

1000 Chevalier, N., Savoye, N., Dubois, S., Lama, M. L., David, V., Lecroart, P., le Ménach, K.,  
1001 and Budzinski, H.: Precise indices based on *n*-alkane distribution for quantifying sources of  
1002 sedimentary organic matter in coastal systems, *Organic Geochemistry*, 88, 69–77,  
1003 <https://doi.org/10.1016/j.orggeochem.2015.07.006>, 2015.

1004 Copard, Y., Eyrolle, F., Radakovitch, O., Poirel, A., Raimbault, P., Gairoard, S., and Di-  
1005 Giovanni, C.: Badlands as a hot spot of petrogenic contribution to riverine particulate organic  
1006 carbon to the Gulf of Lion (NW Mediterranean Sea), *Earth Surface Processes and Landforms*,  
1007 43, 2495–2509, <https://doi.org/10.1002/esp.4409>, 2018.

1008 Copard, Y., Eyrolle, F., Grosbois, C., Lepage, H., Ducros, L., Morereau, A., Bodereau, N.,  
1009 Cossonnet, C., and Desmet, M.: The unravelling of radiocarbon composition of organic  
1010 carbon in river sediments to document past anthropogenic impacts on river systems, *Science  
1011 of The Total Environment*, 806, 150890, <https://doi.org/10.1016/j.scitotenv.2021.150890>,  
1012 2022.

1013 Dagg, M., Benner, R., Lohrenz, S., and Lawrence, D.: Transformation of dissolved and  
1014 particulate materials on continental shelves influenced by large rivers: plume processes,  
1015 *Continental Shelf Research*, 24, 833–858, <https://doi.org/10.1016/j.csr.2004.02.003>, 2004.

1016 Dalzell, B. J., Filley, T. R., and Harbor, J. M.: The role of hydrology in annual organic carbon  
1017 loads and terrestrial organic matter export from a midwestern agricultural watershed,  
1018 *Geochimica et Cosmochimica Acta*, 71, 1448–1462,  
1019 <https://doi.org/10.1016/j.gca.2006.12.009>, 2007.

1021 David, V., Sautour, B., Chardy, P., and Leconte, M.: Long-term changes of the zooplankton  
1022 variability in a turbid environment: The Gironde estuary (France), *Estuarine, Coastal and*  
1023 *Shelf Science*, 64, 171–184, <https://doi.org/10.1016/j.ecss.2005.01.014>, 2005.

1024 Deborde, J.: *Dynamique des sels nutritifs et de la matière organique dans le système*  
1025 *fluvioestuarien de l'Adour/Golfe de Gascogne*, Université de Pau et des Pays de l'adour,  
1026 2019.

1027 Dodds, W. K. and Smith, V. H.: Nitrogen, phosphorus, and eutrophication in streams, *Inland*  
1028 *Waters*, 6, 155–164, <https://doi.org/10.5268/IW-6.2.909>, 2016.

1029 Dubois, S., Savoye, N., Grémare, A., Plus, M., Charlier, K., Beltoise, A., and Blanchet, H.:  
1030 Origin and composition of sediment organic matter in a coastal semi-enclosed ecosystem: An  
1031 elemental and isotopic study at the ecosystem space scale, *Journal of Marine Systems*, 94, 64–  
1032 73, <https://doi.org/10.1016/j.jmarsys.2011.10.009>, 2012.

1033 Dunn†, J. C.: Well-Separated Clusters and Optimal Fuzzy Partitions, *Journal of Cybernetics*,  
1034 4, 95–104, <https://doi.org/10.1080/01969727408546059>, 1974.

1035 Dürr, H. H., Laruelle, G. G., van Kempen, C. M., Slomp, C. P., Meybeck, M., and  
1036 Middelkoop, H.: Worldwide Typology of Nearshore Coastal Systems: Defining the Estuarine  
1037 Filter of River Inputs to the Oceans, *Estuaries and Coasts*, 34, 441–458,  
1038 <https://doi.org/10.1007/s12237-011-9381-y>, 2011.

1039 Etcheber, H., Taillez, A., Abril, G., Garnier, J., Servais, P., Moatar, F., and Commarieu, M.-  
1040 V.: Particulate organic carbon in the estuarine turbidity maxima of the Gironde, Loire and  
1041 Seine estuaries: origin and lability, *Hydrobiologia*, 588, 245–259,  
1042 <https://doi.org/10.1007/s10750-007-0667-9>, 2007.

1043 Falkowski, P. G., Barber, R. T., and Smetacek, V.: Biogeochemical Controls and Feedbacks  
1044 on Ocean Primary Production, *Science*, 281, 200–206,  
1045 <https://doi.org/10.1126/science.281.5374.200>, 1998.

1046 Ferchiche, F., Liénart, C., Charlier, K., Coynel, A., Gorse-Labadie, L., and Savoye, N.:  
1047 Quantifying particulate organic matter: source composition and fluxes at the river-estuary  
1048 interface, *Front. Freshw. Sci.*, 2, <https://doi.org/10.3389/ffwsc.2024.1437431>, 2024.

1049 Ferchiche, F., Liénart, C., Savoye, N., and Wassenaar, L. I.: Unlocking the potential of  
1050 hydrogen isotopes ( $\delta^{2\text{H}}$ ) in tracing riverine particulate organic matter sources and dynamics,  
1051 *Aquat Sci*, 87, 5, <https://doi.org/10.1007/s00027-024-01127-1>, 2025.

1052 Fernandez, I., Mahieu, N., and Cadisch, G.: Carbon isotopic fractionation during  
1053 decomposition of plant materials of different quality, *Global Biogeochemical Cycles*, 17,  
1054 <https://doi.org/10.1029/2001GB001834>, 2003.

1055 Field, C. B., Behrenfeld, M. J., Randerson, J. T., and Falkowski, P.: Primary Production of the  
1056 Biosphere: Integrating Terrestrial and Oceanic Components, *Science*, 281, 237–240,  
1057 <https://doi.org/10.1126/science.281.5374.237>, 1998.

1058 Finlay, J. C., Doucett, R. R., and McNEELY, C.: Tracing energy flow in stream food webs  
1059 using stable isotopes of hydrogen, *Freshwater Biology*, 55, 941–951,  
1060 <https://doi.org/10.1111/j.1365-2427.2009.02327.x>, 2010.

1061 Fry, B.: *13C/12C fractionation by marine diatoms*, *Marine Ecology Progress Series*, 134,  
1062 283–294, <https://doi.org/10.3354/meps134283>, 1996.

1063 Galeron, M.-A., Radakovitch, O., Charrière, B., Vaultier, F., and Rontani, J.-F.: *Autoxidation*  
1064 as a major player in the fate of terrestrial particulate organic matter in seawater, *Journal of*  
1065 *Geophysical Research: Biogeosciences*, 122, 1203–1215,  
1066 <https://doi.org/10.1002/2016JG003708>, 2017.

1067 Gawade, L., Krishna, M. S., Sarma, V. V. S. S., Hemalatha, K. P. J., and Venkateshwara Rao,  
1068 Y.: *Spatio-temporal variability in the sources of particulate organic carbon and nitrogen in a*  
1069 *tropical Godavari estuary*, *Estuarine, Coastal and Shelf Science*, 215, 20–29,  
1070 <https://doi.org/10.1016/j.ecss.2018.10.004>, 2018.

1071 Golubkov, M. S., Nikulina, V. N., Tiunov, A. V., and Golubkov, S. M.: *Stable C and N*  
1072 *Isotope Composition of Suspended Particulate Organic Matter in the Neva Estuary: The Role*  
1073 *of Abiotic Factors, Productivity, and Phytoplankton Taxonomic Composition*, *Journal of*  
1074 *Marine Science and Engineering*, 8, 959, <https://doi.org/10.3390/jmse8120959>, 2020.

1075 Goñi, M. A., Voulgaris, G., and Kim, Y. H.: *Composition and fluxes of particulate organic*  
1076 *matter in a temperate estuary (Winyah Bay, South Carolina, USA) under contrasting physical*  
1077 *forcings*, *Estuarine, Coastal and Shelf Science*, 85, 273–291,  
1078 <https://doi.org/10.1016/j.ecss.2009.08.013>, 2009.

1079 Govan, E. and Parnell, A.: *simmr: A Stable Isotope Mixing Model*, 2024.

1080 Grunicke, F., Wagner, A., von Elert, E., Weitere, M., and Berendonk, T.: *Riparian detritus vs.*  
1081 *stream detritus: food quality determines fitness of juveniles of the highly endangered*  
1082 *freshwater pearl mussels (Margaritifera margaritifera)*, *Hydrobiologia*, 850, 729–746,  
1083 <https://doi.org/10.1007/s10750-022-05120-3>, 2023.

1084 Harmelin-Vivien, M., Dierking, J., Bănaru, D., Fontaine, M. F., and Arlhac, D.: *Seasonal*  
1085 *variation in stable C and N isotope ratios of the Rhone River inputs to the Mediterranean Sea*  
1086 *(2004–2005)*, *Biogeochemistry*, 100, 139–150, <https://doi.org/10.1007/s10533-010-9411-z>,  
1087 2010.

1088 Hellings, L., Dehairs, F., Tackx, M., Keppens, E., and Baeyens, W.: *Origin and fate of*  
1089 *organic carbon in the freshwater part of the Scheldt Estuary as traced by stable carbon isotope*  
1090 *composition*, *Biogeochemistry*, 47, 167–186, <https://doi.org/10.1023/A:1006143827118>,  
1091 1999.

1092 Higueras, M., Kerhervé, P., Sanchez-Vidal, A., Calafat, A., Ludwig, W., Verdoit-Jarraya, M.,  
1093 Heussner, S., and Canals, M.: *Biogeochemical characterization of the riverine particulate*  
1094 *organic matter transferred to the NW Mediterranean Sea*, *Biogeosciences*, 11, 157–172,  
1095 <https://doi.org/10.5194/bg-11-157-2014>, 2014.

1096 Hilton, R. G., Galy, A., Hovius, N., Horng, M.-J., and Chen, H.: *The isotopic composition of*  
1097 *particulate organic carbon in mountain rivers of Taiwan*, *Geochimica et Cosmochimica Acta*,  
1098 74, 3164–3181, <https://doi.org/10.1016/j.gca.2010.03.004>, 2010.

1099 Hou, P., Eglinton, T. I., Yu, M., Montluçon, D. B., Haghipour, N., Zhang, H., Jin, G., and  
1100 Zhao, M.: *Degradation and Aging of Terrestrial Organic Carbon within Estuaries*:

1101 Biogeochemical and Environmental Implications, *Environmental Science and Technology*,  
1102 55, 10852–10861, <https://doi.org/10.1021/acs.est.1c02742>, 2021.

1103 Hounshell, A. G., Fegley, S. R., Hall, N. S., Osburn, C. L., and Paerl, H. W.: Riverine  
1104 Discharge and Phytoplankton Biomass Control Dissolved and Particulate Organic Matter  
1105 Dynamics over Spatial and Temporal Scales in the Neuse River Estuary, North Carolina,  
1106 *Estuaries and Coasts*, 45, 96–113, <https://doi.org/10/gj64px>, 2022.

1107 INRA: Base de Données Géographique des Sols de France à 1/1 000 000 version 3.2.8.0,  
1108 10/09/1998, <https://doi.org/10.15454/BPN57S>, 2025.

1109 Kang, S., Kim, J.-H., Joe, Y. J., Jang, K., Nam, S.-I., and Shin, K.-H.: Long-term  
1110 environmental changes in the Geum Estuary (South Korea): Implications of river  
1111 impoundments, *Marine Pollution Bulletin*, 168, 112383, <https://doi.org/10/gnpb2t>, 2021.

1112 Ke, Z., Tan, Y., Huang, L., Liu, J., Xiang, C., Zhao, C., and Zhang, J.: Significantly depleted  
1113 15N in suspended particulate organic matter indicating a strong influence of sewage loading  
1114 in Daya Bay, China, *Science of The Total Environment*, 650, 759–768,  
1115 <https://doi.org/10.1016/j.scitotenv.2018.09.076>, 2019.

1116 Kelso, J. E. and Baker, M. A.: Organic Matter Is a Mixture of Terrestrial, Autochthonous, and  
1117 Wastewater Effluent in an Urban River, *Frontiers in Environmental Science*, 7,  
1118 <https://doi.org/10.3389/fenvs.2019.00202>, 2020.

1119 Kelso, J. E. and Baker, M. A.: Organic matter sources and composition in four watersheds  
1120 with mixed land cover, *Hydrobiologia*, 849, 2663–2682, <https://doi.org/10.1007/s10750-022-04884-y>, 2022.

1122 Kendall, C., Silva, S. R., and Kelly, V. J.: Carbon and nitrogen isotopic compositions of  
1123 particulate organic matter in four large river systems across the United States, *Hydrological  
1124 Processes*, 15, 1301–1346, <https://doi.org/10.1002/hyp.216>, 2001.

1125 Khan, M. N. and Mohammad, F.: Eutrophication: Challenges and Solutions, in:  
1126 Eutrophication: Causes, Consequences and Control: Volume 2, edited by: Ansari, A. A. and  
1127 Gill, S. S., Springer Netherlands, Dordrecht, 1–15, [https://doi.org/10.1007/978-94-007-7814-6\\_1](https://doi.org/10.1007/978-94-007-7814-6_1), 2014.

1129 Lambert, T., Bouillon, S., Darchambeau, F., Morana, C., Roland, F. A. E., Descy, J.-P., and  
1130 Borges, A. V.: Effects of human land use on the terrestrial and aquatic sources of fluvial  
1131 organic matter in a temperate river basin (The Meuse River, Belgium), *Biogeochemistry*, 136,  
1132 191–211, <https://doi.org/10.1007/s10533-017-0387-9>, 2017.

1133 Le Bas, C.: Carte de la Réserve Utile en eau issue de la Base de Données Géographique des  
1134 Sols de France, <https://doi.org/10.15454/JPB9RB>, 2025.

1135 Lebreton, B., Beseres Pollack, J., Blomberg, B., Palmer, T. A., Adams, L., Guillou, G., and  
1136 Montagna, P. A.: Origin, composition and quality of suspended particulate organic matter in  
1137 relation to freshwater inflow in a South Texas estuary, *Estuarine, Coastal and Shelf Science*,  
1138 170, 70–82, <https://doi.org/10.1016/j.ecss.2015.12.024>, 2016.

1139 Legendre, P., Oksanen, J., and ter Braak, C. J. F.: Testing the significance of canonical axes in  
1140 redundancy analysis, *Methods in Ecology and Evolution*, 2, 269–277,  
1141 <https://doi.org/10.1111/j.2041-210X.2010.00078.x>, 2011.

1142 Lheureux, A., David, V., Del Amo, Y., Soudant, D., Auby, I., Ganthy, F., Blanchet, H.,  
1143 Cordier, M.-A., Costes, L., Ferreira, S., Mornet, L., Nowaczyk, A., Parra, M., D'Amico, F.,  
1144 Gouriou, L., Meteigner, C., Oger-Jeanneret, H., Rigouin, L., Rumebe, M., Tournaire, M.-P.,  
1145 Trut, F., Trut, G., and Savoye, N.: Bi-decadal changes in nutrient concentrations and ratios in  
1146 marine coastal ecosystems: The case of the Arcachon bay, France, *Progress in Oceanography*,  
1147 201, 102740, <https://doi.org/10.1016/j.pocean.2022.102740>, 2022.

1148 Liénart, C., Susperregui, N., Rouaud, V., Cavalheiro, J., David, V., Del Amo, Y., Duran, R.,  
1149 Lauga, B., Monperrus, M., Pigot, T., Bichon, S., Charlier, K., and Savoye, N.: Dynamics of  
1150 particulate organic matter in a coastal system characterized by the occurrence of marine  
1151 mucilage – A stable isotope study, *Journal of Sea Research*, 116, 12–22,  
1152 <https://doi.org/10.1016/j.seares.2016.08.001>, 2016.

1153 Liénart, C., Savoye, N., Bozec, Y., Breton, E., Conan, P., David, V., Feunteun, E., Grangeré,  
1154 K., Kerhervé, P., Lebreton, B., Lefebvre, S., L'Helguen, S., Mousseau, L., Raimbault, P.,  
1155 Richard, P., Riera, P., Sauriau, P.-G., Schaal, G., Aubert, F., Aubin, S., Bichon, S., Boinet, C.,  
1156 Bourasseau, L., Bréret, M., Caparros, J., Cariou, T., Charlier, K., Claquin, P., Cornille, V.,  
1157 Corre, A.-M., Costes, L., Crispi, O., Crouvoisier, M., Czamanski, M., Del Amo, Y.,  
1158 Derriennic, H., Dindinaud, F., Durozier, M., Hanquiez, V., Nowaczyk, A., Devesa, J.,  
1159 Ferreira, S., Fornier, M., Garcia, F., Garcia, N., Geslin, S., Grossteffan, E., Gueux, A.,  
1160 Guillaudeau, J., Guillou, G., Joly, O., Lachaussée, N., Lafont, M., Lamoureux, J., Lecuyer, E.,  
1161 Lehodey, J.-P., Lemeille, D., Leroux, C., Macé, E., Maria, E., Pineau, P., Petit, F., Pujo-Pay,  
1162 M., Rimelin-Maury, P., and Sultan, E.: Dynamics of particulate organic matter composition in  
1163 coastal systems: A spatio-temporal study at multi-systems scale, *Progress in Oceanography*,  
1164 156, 221–239, <https://doi.org/10.1016/j.pocean.2017.08.001>, 2017.

1165 Liénart, C., Savoye, N., David, V., Ramond, P., Rodriguez Tress, P., Hanquiez, V., Marieu,  
1166 V., Aubert, F., Aubin, S., Bichon, S., Boinet, C., Bourasseau, L., Bozec, Y., Bréret, M.,  
1167 Breton, E., Caparros, J., Cariou, T., Claquin, P., Conan, P., Corre, A.-M., Costes, L.,  
1168 Crouvoisier, M., Del Amo, Y., Derriennic, H., Dindinaud, F., Duran, R., Durozier, M.,  
1169 Devesa, J., Ferreira, S., Feunteun, E., Garcia, N., Geslin, S., Grossteffan, E., Gueux, A.,  
1170 Guillaudeau, J., Guillou, G., Jolly, O., Lachaussée, N., Lafont, M., Lagadec, V., Lamoureux,  
1171 J., Lauga, B., Lebreton, B., Lecuyer, E., Lehodey, J.-P., Leroux, C., L'Helguen, S., Macé, E.,  
1172 Maria, E., Mousseau, L., Nowaczyk, A., Pineau, P., Petit, F., Pujo-Pay, M., Raimbault, P.,  
1173 Rimelin-Maury, P., Rouaud, V., Sauriau, P.-G., Sultan, E., and Susperregui, N.: Dynamics  
1174 of particulate organic matter composition in coastal systems: Forcing of spatio-temporal  
1175 variability at multi-systems scale, *Progress in Oceanography*, 162, 271–289,  
1176 <https://doi.org/10.1016/j.pocean.2018.02.026>, 2018.

1177 Liénart, C., Savoye, N., Conan, P., David, V., Barbier, P., Bichon, S., Charlier, K., Costes, L.,  
1178 Derriennic, H., Ferreira, S., Gueux, A., Hubas, C., Maria, E., and Meziane, T.: Relationship  
1179 between bacterial compartment and particulate organic matter (POM) in coastal systems: An  
1180 assessment using fatty acids and stable isotopes, *Estuarine, Coastal and Shelf Science*, 239,  
1181 106720, <https://doi.org/10.1016/j.ecss.2020.106720>, 2020.

1182 Lowe, A. T., Galloway, A. W. E., Yeung, J. S., Dethier, M. N., and Duggins, D. O.: Broad  
1183 sampling and diverse biomarkers allow characterization of nearshore particulate organic  
1184 matter, *Oikos*, 123, 1341–1354, <https://doi.org/10.1111/oik.01392>, 2014.

1185 Lu, L., Cheng, H., Pu, X., Wang, J., Cheng, Q., and Liu, X.: Identifying organic matter  
1186 sources using isotopic ratios in a watershed impacted by intensive agricultural activities in  
1187 Northeast China, *Agriculture, Ecosystems & Environment*, 222, 48–59,  
1188 <https://doi.org/10.1016/j.agee.2015.12.033>, 2016.

1189 McCorkle, E. P., Berhe, A. A., Hunsaker, C. T., Johnson, D. W., McFarlane, K. J., Fogel, M.  
1190 L., and Hart, S. C.: Tracing the source of soil organic matter eroded from temperate forest  
1191 catchments using carbon and nitrogen isotopes, *Chemical Geology*, 445, 172–184,  
1192 <https://doi.org/10.1016/j.chemgeo.2016.04.025>, 2016.

1193 Michener, R. H. and Kaufman, L.: Stable Isotope Ratios as Tracers in Marine Food Webs: An  
1194 Update, 1st ed., edited by: Michener, R. and Lajtha, K., Wiley,  
1195 <https://doi.org/10.1002/9780470691854.ch9>, 2007.

1196 Middelburg, J. J. and Herman, P. M. J.: Organic matter processing in tidal estuaries, *Marine  
1197 Chemistry*, 106, 127–147, <https://doi.org/10.1016/j.marchem.2006.02.007>, 2007.

1198 Miller, R. J., Page, H. M., and Brzezinski, M. A.:  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  of particulate organic  
1199 matter in the Santa Barbara Channel: drivers and implications for trophic inference, *Marine  
1200 Ecology Progress Series*, 474, 53–66, <https://doi.org/10.3354/meps10098>, 2013.

1201 Minaudo, C., Meybeck, M., Moatar, F., Gassama, N., and Curie, F.: Eutrophication mitigation  
1202 in rivers: 30 years of trends in spatial and seasonal patterns of biogeochemistry of the Loire  
1203 River (1980–2012), *Biogeosciences*, 12, 2549–2563, <https://doi.org/10.5194/bg-12-2549-2015>, 2015.

1205 Minaudo, C., Moatar, F., Coynel, A., Etcheber, H., Gassama, N., and Curie, F.: Using recent  
1206 high-frequency surveys to reconstitute 35 years of organic carbon variations in a eutrophic  
1207 lowland river, *Environ Monit Assess*, 188, 41, <https://doi.org/10.1007/s10661-015-5054-9>,  
1208 2016.

1209 Ogrinc, N., Markovics, R., Kanduč, T., Walter, L. M., and Hamilton, S. K.: Sources and  
1210 transport of carbon and nitrogen in the River Sava watershed, a major tributary of the River  
1211 Danube, *Applied Geochemistry*, 23, 3685–3698,  
1212 <https://doi.org/10.1016/j.apgeochem.2008.09.003>, 2008.

1213 O’Leary, M. H.: Carbon isotope fractionation in plants, *Phytochemistry*, 20, 553–567,  
1214 [https://doi.org/10.1016/0031-9422\(81\)85134-5](https://doi.org/10.1016/0031-9422(81)85134-5), 1981.

1215 Onstad, G. D., Canfield, D. E., Quay, P. D., and Hedges, J. I.: Sources of particulate organic  
1216 matter in rivers from the continental usa: lignin phenol and stable carbon isotope  
1217 compositions, *Geochimica et Cosmochimica Acta*, 64, 3539–3546,  
1218 [https://doi.org/10.1016/S0016-7037\(00\)00451-8](https://doi.org/10.1016/S0016-7037(00)00451-8), 2000.

1219 Parnell, A. C., Phillips, D. L., Bearhop, S., Semmens, B. X., Ward, E. J., Moore, J. W.,  
1220 Jackson, A. L., Grey, J., Kelly, D. J., and Inger, R.: Bayesian stable isotope mixing models,  
1221 *Environmetrics*, 24, 387–399, <https://doi.org/10.1002/env.2221>, 2013.

1222 Perdue, E. M. and Koprivnjak, J.-F.: Using the C/N ratio to estimate terrigenous inputs of  
1223 organic matter to aquatic environments, *Estuarine, Coastal and Shelf Science*, 73, 65–72,  
1224 <https://doi.org/10.1016/j.ecss.2006.12.021>, 2007.

1225 Phillips, D. L., Inger, R., Bearhop, S., Jackson, A. L., Moore, J. W., Parnell, A. C., Semmens,  
1226 B. X., and Ward, E. J.: Best practices for use of stable isotope mixing models in food-web  
1227 studies, *Can. J. Zool.*, 92, 823–835, <https://doi.org/10.1139/cjz-2014-0127>, 2014.

1228 Polsenaere, P., Savoye, N., Etcheber, H., Canton, M., Poirier, D., Bouillon, S., and Abril, G.:  
1229 Export and degassing of terrestrial carbon through watercourses draining a temperate  
1230 podzolized catchment, *Aquat Sci*, 75, 299–319, <https://doi.org/10.1007/s00027-012-0275-2>,  
1231 2013.

1232 Pradhan, U. K., Wu, Y., Wang, X., Zhang, J., and Zhang, G.: Signals of typhoon induced  
1233 hydrologic alteration in particulate organic matter from largest tropical river system of Hainan  
1234 Island, South China Sea, *Journal of Hydrology*, 534, 553–566,  
1235 <https://doi.org/10.1016/j.jhydrol.2016.01.046>, 2016.

1236 Regnier, P., Friedlingstein, P., Ciais, P., Mackenzie, F. T., Gruber, N., Janssens, I. A.,  
1237 Laruelle, G. G., Lauerwald, R., Luyssaert, S., Andersson, A. J., Arndt, S., Arnosti, C., Borges,  
1238 A. V., Dale, A. W., Gallego-Sala, A., Goddérus, Y., Goossens, N., Hartmann, J., Heinze, C.,  
1239 Ilyina, T., Joos, F., LaRowe, D. E., Leifeld, J., Meysman, F. J. R., Munhoven, G., Raymond,  
1240 P. A., Spahni, R., Suntharalingam, P., and Thullner, M.: Anthropogenic perturbation of the  
1241 carbon fluxes from land to ocean, *Nature Geosci*, 6, 597–607,  
1242 <https://doi.org/10.1038/ngeo1830>, 2013.

1243 Rousseeuw, P. J.: Silhouettes: A graphical aid to the interpretation and validation of cluster  
1244 analysis, *Journal of Computational and Applied Mathematics*, 20, 53–65,  
1245 [https://doi.org/10.1016/0377-0427\(87\)90125-7](https://doi.org/10.1016/0377-0427(87)90125-7), 1987.

1246 Sarma, V. V. S. S., Krishna, M. S., Prasad, V. R., Kumar, B. S. K., Naidu, S. A., Rao, G. D.,  
1247 Viswanadham, R., Sridevi, T., Kumar, P. P., and Reddy, N. P. C.: Distribution and sources of  
1248 particulate organic matter in the Indian monsoonal estuaries during monsoon, *Journal of  
1249 Geophysical Research: Biogeosciences*, 119, 2095–2111,  
1250 <https://doi.org/10.1002/2014JG002721>, 2014.

1251 Sato, T., Miyajima, T., Ogawa, H., Umezawa, Y., and Koike, I.: Temporal variability of  
1252 stable carbon and nitrogen isotopic composition of size-fractionated particulate organic matter  
1253 in the hypertrophic Sumida River Estuary of Tokyo Bay, Japan, *Estuarine, Coastal and Shelf  
1254 Science*, 68, 245–258, <https://doi.org/10.1016/j.ecss.2006.02.007>, 2006.

1255 Savoye, N., Aminot, A., Tréguer, P., Fontugne, M., Naulet, N., and Kérouel, R.: Dynamics of  
1256 particulate organic matter d15N and d13C during spring phytoplankton blooms in a  
1257 macrotidal ecosystem (Bay of Seine, France), *Marine Ecology Progress Series*, 255, 27–41,  
1258 <https://doi.org/10.3354/meps255027>, 2003.

1259 Savoye, N., David, V., Morisseau, F., Etcheber, H., Abril, G., Billy, I., Charlier, K., Oggian,  
1260 G., Derriennic, H., and Sautour, B.: Origin and composition of particulate organic matter in a  
1261 macrotidal turbid estuary: The Gironde Estuary, France, *Estuarine, Coastal and Shelf Science*,  
1262 108, 16–28, <https://doi.org/10.1016/j.ecss.2011.12.005>, 2012.

1263 Sigman, D. M., DiFiore, P. J., Hain, M. P., Deutsch, C., Wang, Y., Karl, D. M., Knapp, A. N.,  
1264 Lehmann, M. F., and Pantoja, S.: The dual isotopes of deep nitrate as a constraint on the cycle  
1265 and budget of oceanic fixed nitrogen, Deep Sea Research Part I: Oceanographic Research  
1266 Papers, 56, 1419–1439, <https://doi.org/10.1016/j.dsr.2009.04.007>, 2009.

1267 Souissi, S., Yahia-Kéfi, O. D., and Yahia, M. N. D.: Spatial characterization of nutrient  
1268 dynamics in the Bay of Tunis (south-western Mediterranean) using multivariate analyses:  
1269 consequences for phyto- and zooplankton distribution, Journal of Plankton Research, 22,  
1270 2039–2059, <https://doi.org/10.1093/plankt/22.11.2039>, 2000.

1271 Sun, X., Fan, D., Cheng, P., Hu, L., Sun, X., Guo, Z., and Yang, Z.: Source, transport and fate  
1272 of terrestrial organic carbon from Yangtze River during a large flood event: Insights from  
1273 multiple-isotopes ( $\delta^{13}\text{C}$ ,  $\delta^{15}\text{N}$ ,  $\Delta^{14}\text{C}$ ) and geochemical tracers, *Geochimica et*  
1274 *Cosmochimica Acta*, 308, 217–236, <https://doi.org/10/gn8qs5>, 2021.

1275 Turner, R. E., Milan, C. S., Swenson, E. M., and Lee, J. M.: Peak chlorophyll a concentrations  
1276 in the lower Mississippi River from 1997 to 2018, *Limnology and Oceanography*, n/a,  
1277 <https://doi.org/10/gpjm83>, 2022.

1278 Veyssi, E., Etcheber, H., Lin, R. G., Buat-Menard, P., and Maneux, E.: Seasonal variation  
1279 and origin of Particulate Organic Carbon in the lower Garonne River at La Reole  
1280 (southwestern France), *Hydrobiologia*, 391, 113–126,  
1281 <https://doi.org/10.1023/A:1003520907962>, 1998.

1282 Wang, X., Chen, Y., Yuan, Q., Xing, X., Hu, B., Gan, J., Zheng, Y., and Liu, Y.: Effect of  
1283 river damming on nutrient transport and transformation and its countermeasures, *Frontiers in*  
1284 *Marine Science*, 9, 2022.

1285 Wang, Y., Song, J., Duan, L., Yuan, H., Li, X., Li, N., Zhang, Q., Liu, J., and Ren, C.:  
1286 Combining sterols with stable carbon isotope as indicators for assessing the organic matter  
1287 sources and primary productivity evolution in the coastal areas of the East China Sea,  
1288 *Continental Shelf Research*, 223, 104446, <https://doi.org/10.1016/j.csr.2021.104446>, 2021.

1289 Yaalon, D. H.: Soils in the Mediterranean region: what makes them different?, *CATENA*, 28,  
1290 157–169, [https://doi.org/10.1016/S0341-8162\(96\)00035-5](https://doi.org/10.1016/S0341-8162(96)00035-5), 1997.

1291 Yan, X., Yang, J.-Y. T., Xu, M. N., Wang, H., Dai, M., and Kao, S.-J.: Nitrogen isotope  
1292 constraint on the zonation of multiple transformations between dissolved and particulate  
1293 organic nitrogen in the Changjiang plume, *Science of The Total Environment*, 818, 151678,  
1294 <https://doi.org/10.1016/j.scitotenv.2021.151678>, 2022.

1295 Yu, F., Zong, Y., Lloyd, J. M., Huang, G., Leng, M. J., Kendrick, C., Lamb, A. L., and Yim,  
1296 W. W.-S.: Bulk organic  $\delta^{13}\text{C}$  and C/N as indicators for sediment sources in the Pearl River  
1297 delta and estuary, southern China, *Estuarine, Coastal and Shelf Science*, 87, 618–630,  
1298 <https://doi.org/10.1016/j.ecss.2010.02.018>, 2010.

1299 Zhang, Y., Meng, X., Bai, Y., Wang, X., Xia, P., Yang, G., Zhu, Z., and Zhang, H.: Sources  
1300 and features of particulate organic matter in tropical small mountainous rivers (SW China)  
1301 under the effects of anthropogenic activities, *Ecological Indicators*, 125, 107471,  
1302 <https://doi.org/10.1016/j.ecolind.2021.107471>, 2021.

1303

