

Hydrodynamic and Primary Production Effects on Seasonal DO Variability in the Danube River

Jan Maier¹, Anna-Neva Visser¹, Christina M. Schubert¹, Simon T. Wander¹, Johannes A. C. Barth¹

¹Applied Geology, GeoZentrum Nordbayern, Friedrich-Alexander-University Erlangen-Nürnberg, Erlangen 91054, Germany

Correspondence to: Jan Maier (jan.m.maier@fau.de)

Abstract

Dissolved oxygen (DO) is a fundamental indicator for water quality and ecosystem health, particularly in the context of anthropogenic impacts and climate change. This study presents the first large-scale dataset of DO concentrations combined with its stable oxygen isotope ratios (expressed as $\delta^{18}\text{O}_{\text{DO}}$), particulate organic carbon concentrations (POC) and respiration/ photosynthesis (R/P ratios) from five seasonal campaigns along the entire Danube River in 2023 and 2024. Our findings reveal pronounced seasonal DO dynamics driven by temperature, biological activity and hydrodynamic conditions. During spring and summer, enhanced photosynthesis increased DO up to 0.40 mmol/L, with $\delta^{18}\text{O}_{\text{DO}}$ values down to +12.1 ‰ and POC up to 0.25 mmol/L in two highly productive river sections. Low R/P ratios of up to 0.1 further indicated strong net autotrophic conditions. Strong correlations between $\delta^{18}\text{O}_{\text{DO}}$ and POC further confirm influence of primary producers (i.e., photosynthetic organisms) in a river section where a reduced slope led to slower flow and lower turbulence. Notably, $\delta^{18}\text{O}_{\text{DO}}$ values were lower than those expected for atmospheric equilibrium ($+24.6 \text{ ‰} \pm 0.4 \text{ ‰}$), a pattern rarely documented in large river systems. In contrast, tributary inflows from the Tisa and Sava Rivers diluted biomass and organic material inputs and led to declines in DO and POC. By late summer, intensified respiration reversed photosynthetic signals, led to the lowest DO concentrations down to 0.16 mmol/L and raised $\delta^{18}\text{O}_{\text{DO}}$ up to +23.7 ‰, particularly in the Sava River. In fall, DO levels partially recovered despite continued respiration, as also indicated by the highest observed $\delta^{18}\text{O}_{\text{DO}}$ values of 25.9 ‰ and the highest R/P ratios of the entire season, that reaching up to 8.9. In winter, oxygen input from the atmosphere became dominant with minimal biological influences. Overall, this study provides new insights into dynamics between oxygen sources and sinks across the river continuum over seasons. These new insights underscore the need for continuous DO monitoring, particularly in late summer when oxygen levels can become critically low. Understanding these interactions can help to establish efficient aqueous ecosystem management and conservation strategies in the face of land use and climate change.

1. Introduction

Freshwater ecosystems are increasingly threatened by complex global stressors (Borgwardt et al., 2019; Vörösmarty et al., 2013). Beyond climate change, anthropogenically-induced alterations of river morphology (Belletti et al., 2015), disrupted flow

30 regimes (Acreman and Dunbar, 2004; Poff and Zimmerman, 2010), nutrient input (Fowler et al., 2013; Seitzinger et al., 2006; Sutton et al., 2011), and pollution from urban and industrial sources (Nyenje et al., 2010; Qing et al., 2015; Suthar et al., 2009; Xia et al., 2016) continue to degrade water quality and stability of aquatic habitats. Land use changes further intensify these pressures, thus often jeopardizing river basin health. Such influences can also amplify the impacts of environmental stressors (Honisch et al., 2002; Hua, 2017). Consequently, between 2015 and 2021, only 37 % of all European surface water bodies
35 achieved a 'good' ecological status and only 29 % a 'good' chemical status (EEA, 2021). These numbers highlight the urgency to address environmental challenges for terrestrial aqueous systems. Particularly excessive agricultural nutrient inputs from mostly agriculture land use, primarily phosphorus and nitrogen, threaten aquatic life and can trigger algal blooms that lead to eutrophication, oxygen depletion and ultimately a decline in biodiversity (Carpenter et al., 1998; Dudgeon et al., 2006; Grizzetti et al., 2017).

40 Among many consequences of these stressors, disruptions of dissolved oxygen (DO) sources and sinks can impact freshwater habitats and influence biodiversity, biogeochemical cycles and overall environmental health (Franklin, 2014; Killgore and Hoover, 2001; North et al., 2014). Primary producers, such as phytoplankton, serve as a key source of DO via photosynthesis, whereas respiration by heterotrophic organisms provide major sinks that consume DO (Heddam, 2014; Wetzel, 2011). However, DO concentrations are not solely regulated by biological activity. They are also influenced by atmospheric
45 exchange and environmental factors such as temperature, light availability and nutrient levels (Benson et al., 1979; Odum, 1956; Stumm and Morgan, 1995). In addition, chemical processes, such as redox reactions and the mineralization of organic carbon under stratified or eutrophic conditions, can also serve as important DO sinks, particularly in hypolimnetic lake environments, anoxic groundwaters and rivers with elevated groundwater input (Jurgens et al., 2009; Piatka et al., 2021; Rosecrans et al., 2017).

50 In the context of fluvial systems, a better understanding and prediction of DO distributions and their controlling processes are essential to assess aquatic health and to prevent or at least manage potential anoxic events. This is especially important for the Danube River, Europe's second-longest waterway. While the river provides important services for agriculture and energy production, these activities also impose significant pressures on its natural system. Additionally, the Danube serves as a crucial ecological corridor that promotes biodiversity across central and eastern Europe (Habersack et al., 2016; ICPDR, 2015; Sommerwerk et al., 2009). Although the ecological quality improved in the last 30 years, the Danube River still faces
55 ongoing threats, particularly from organic pollution downstream of major cities and after confluences of key tributaries (Mănoiu and Crăciun, 2021; Wachs, 1997).

Despite the critical role of DO in freshwater ecosystems, research has primarily focused on DO concentration patterns as an indicator of water quality, while the relative contributions of biological and atmospheric inputs often remain poorly
60 quantified. In particular, photosynthesis and community respiration drive DO dynamics in fundamentally different ways, yet their individual effects cannot be fully distinguished through concentration measurements alone. Traditional approaches lack a framework to separate DO sources and sinks, thus leaving significant gaps in our understanding of aqueous DO cycling. Stable isotope measurements of DO provide a powerful tool to overcome these limitations by distinguishing between three

key processes that govern DO dynamics in aqueous environments: photosynthesis, respiration and atmospheric exchange.

65 Aquatic photosynthesis, by splitting water molecules, transfers a typically ^{16}O -enriched signature into the DO pool (Guy et al., 1993; Limburg et al., 1999). However, under low DO concentrations, concurrent DO consumption can lead to ^{18}O -enrichments even during photosynthesis (Eisenstadt et al., 2010). Similarly, aerobic respiration preferentially consumes ^{16}O , leaving the remaining DO enriched in ^{18}O (Mader et al., 2017). This study is among few to apply DO isotopes to river systems and contributes to a still limited body of research on DO dynamics in aqueous systems (e.g., Parker et al., 2010; Quay et al., 1995; Tobias et al., 2007; Wassenaar et al., 2010).

With increasing environmental pressures, a comprehensive understanding of DO sources and sinks in the Danube is essential for assessing and managing river health. To address this issue, we conducted the first large-scale study of the DO budget of the entire Danube main channel and its key tributaries (e.g., Inn, Tisa and Sava) (Figure 1), based on five sampling campaigns between 2023 and 2024. By integrating spatial high-resolution DO measurements with $\delta^{18}\text{O}_{\text{DO}}$ analyses, particulate organic carbon (POC), and ratios between respiration and photosynthesis (R/P), we were able to reveal DO sources and sinks along the river continuum together with its seasonal dynamics. This approach could identify critical periods and regions characterized by elevated or depleted DO levels and also disentangled variable contributions by photosynthesis, respiration and atmospheric exchange. Together, these findings contribute to a more comprehensive understanding of the Danube River's functioning.

80 **2. Material and methods**

2.1 Study area

The Danube has a total length of 2,857 km and a mean annual discharge of 6486 m³/s (Sommerwerk et al., 2009). Its catchment area of 807,827 km² hosts a population of approximately 83 million people in 10 different countries and serves as a vital freshwater resource for Central and Eastern Europe (Habersack et al., 2016; ICPDR, 2015; Sommerwerk et al., 2009).

85 In 2023 and 2024, five sampling campaigns were conducted on the main river and its major tributaries (e.g., Inn, Tisa and Sava), covering the following key seasons: summer (July 2023), fall (late October to early November 2023), winter (February 2024), spring (April 2024) and late summer (late August to early September 2024). During each sampling campaign, between 54 to 89 sampling locations along the entire mainstream were surveyed (Figure 1). The coordinates of the sampling sites were recorded in the field by Google Maps and confirmed by a Garmin eTrex HC-series GPS device. Elevation data were determined with a barometric altimeter via the mobile Elevation App from Mapnitude Company Limited (<https://mapnitude.com/elevation>). Discharge data were provided by the International Commission of the Danube River (ICPDR, <https://www.danubehis.org>, (ICPDR, 2025; last access: 11 March 2025)).




Figure 1: a) Overview map of Europe and b) detailed view of the Danube River Basin (red outline), including the Danube River (thick blue line) and its major tributaries (thin blue lines), data provided by the ICPDR (data source ICPDR, 2025, last access: 11 March 2025). Yellow dots represent the sampling locations of late summer 2024; red squares show main cities along the Danube. Sampled tributaries include the Brigach, Breg, Lech, Isar, Inn, Enns, March, Váh, Dráva, Tisa, Sava and Siret. The map was created using QGIS v 3.28.3 with raster data from natural earth data (version 3.2.0, last access: 11 December 2024: <https://www.naturalearthdata.com/downloads/10m-raster-data/10m-cross-blend-hypso/>) and shapefiles from geoBoundaries (last access: 11 December 2024: <https://www.geoboundaries.org/globalDownloads.html>).

2.2 Field methods

Samples were collected using a weighted narrow mouth 2-L polyethylene sampling bottle that was submerged between 1 and 2 meters below the water surface to obtain well-mixed water and to minimize influences by rain and evaporation. Sampling took place either from the center (via bridges or passenger boats) or occasionally from the riverbank. In each case, well-mixed water samples were obtained from flowing section of the river. To ensure that the samples represented the entire river section, two cross-sectional profiles were sampled during each sampling campaign and in all cases, they confirmed homogeneously mixed and representative water of each location.

In situ measurements of temperature (T), DO, and oxygen saturation (DO %) were performed with a multiparameter instrument (HQ40d, HACH™, Loveland, CO, USA). This instrument was calibrated daily. Measurement standard deviations were ± 0.1 mg/L for DO, ± 0.42 % for DO %, and ± 0.07 °C for T.

For $\delta^{18}\text{O}_{\text{DO}}$ analyses, samples were filtered through $0.45 \mu\text{m}$ pore size nylon syringe filters (Sartorius™) into pre-poisoned 12-mL vials (Labco LTD. Lampeter Exetainer™). These vials contained 10 μL of a saturated HgCl_2 solution to

inhibit microbial activity after sampling. Vials were filled completely and sealed with screw caps and butyl septa for efficient sealing against atmospheric influences. Triplicate samples were stored in the dark at 4 °C. To evaluate potential diel effects on DO, nighttime samples were collected in the main river channel at two sites during the campaign of late summer 2024 and reveal minor differences in DO concentration (< 0.01 mg/L) and $\delta^{18}\text{O}_{\text{DO}}$ (<1‰). For water isotopes (expressed as $\delta^{18}\text{O}_{\text{H}_2\text{O}}$), necessary for R/P calculations, water samples were collected in identical 12-mL exetainers however without poisoning by HgCl_2 .

For the determination of POC, 500 mL of unfiltered water were collected in acid-washed high-density polyethylene Nalgene bottles. Before sampling, the bottles were thoroughly rinsed three times with sample water. Preparation for POC analysis involved filtering the collected water through pre-weighed glass fiber filters (GF-5, pore size 0.4 μm ; Macherey-Nagel, Düren, Germany). To eliminate residual organic carbon, these filters were pre-heated at 400 °C for 4 h and stored under sterile conditions until sampling.

2.3 Laboratory methods

$\delta^{18}\text{O}_{\text{DO}}$ measurements were performed using a modified automated equilibration system (Gasbench II, ThermoFisher Scientific™) connected in continuous flow mode to a DELTA V Advantage isotope ratio mass spectrometer (IRMS, ThermoFisher Scientific™). The analytical approach was based on methods described by Barth et al. (2004) and Wassenaar and Koehler (1999). Briefly, a 3 mL pure helium headspace was established in the sample vial, and dissolved gases were extracted by shaking these sample vials on an orbital shaker at 250 rotations per minute for 30 minutes. The extracted O_2 in the headspace was separated from nitrogen (N_2) and other trace gases using a gas chromatography column (CP-Molsieve 5 Å, 25 m length, 0.53 mm outer diameter, 0.05 mm inner diameter; Agilent™, Santa Clara, CA, USA) before introduction into the isotope ratio mass spectrometer (IRMS) for analysis. Results are reported as averages of triplicate measurements, with an external reproducibility better than $\pm 0.2 \text{ ‰}$ (1σ).

Values of $\delta^{18}\text{O}_{\text{H}_2\text{O}}$ were determined by infrared spectroscopy (IRIS) with a Picarro™ analyzer (L 1102-i WS-CRDS, Santa Clara, CA, USA). The analysis was conducted following the protocol outlined by van Geldern and Barth (2012).

All stable isotope values are reported in the standard δ -notation relative to the Vienna Standard Mean Ocean Water (VSMOW) and are calculated as:

$$\delta = (R_{\text{sample}} / R_{\text{reference}} - 1) \quad (1)$$

and then multiplied by 1000 to express them in per mille (‰). R represents the molar ratio of the heavy to light isotopes ($^{18}\text{O}/^{16}\text{O}$ for oxygen; $^2\text{H}/^1\text{H}$ and $^{18}\text{O}/^{16}\text{O}$ for H_2O) in the sample and the reference (Coplen, 2011). The ratio of VSMOW is $2005.20 \pm 0.43 \text{ ppm}$ (Baertschi, 1976). For $\delta^{18}\text{O}_{\text{H}_2\text{O}}$, external reproducibility was better than $\pm 0.1 \text{ ‰}$ (1σ).

For POC determination, filters previously loaded with particulate material were freeze-dried for 60 minutes under vacuum conditions (<10 mbar) using a freeze-dryer (Lyovac GT 2 GT 2-E, FinnAqua, Gemini BV, Apeldoorn, Netherlands).

145 The dried filters were then pulverized for 60 s using a ball mill (CryoMill, Retsch, Verder, Vleuten, Netherlands). To ensure complete removal of potential carbonate residues, the powdered filter material was fumigated with a concentrated HCl in a desiccator for 24 h. After fumigation, aliquots of the prepared filter material were weighed into tin capsules (5×9 mm, IVA Analysentechnik GmbH & Co. KG, Meerbusch, Germany). Carbon contents and $^{13}\text{C}/^{12}\text{C}$ isotope ratios of the samples was then determined with an elemental analyzer (Costech ECS 4010, NC Technologies, Bussero, Italy) coupled in helium
150 continuous flow mode to an IRMS (Delta V plus, ThermoFisher, Bremen, Germany).

2.4 Isotope calculations

Ratios of respiration to photosynthesis (R/P) were determined according to Quay et al. (1995) with:

$$\frac{R}{P} = \frac{\left(\frac{^{18}}{^{16}}\text{O}_w * \alpha_p - \frac{^{18}}{^{16}}\text{O}_g \right)}{\left(\frac{^{18}}{^{16}}\text{O} * \alpha_r - \frac{^{18}}{^{16}}\text{O}_g \right)} \quad (2)$$

where $\frac{^{18}}{^{16}}\text{O}_w$ and $\frac{^{18}}{^{16}}\text{O}$ represent the isotope ratios of oxygen in water and in DO, respectively. The photosynthesis fractionation factor (α_p) was assumed to be 1.000 ± 0.003 (Russ et al., 2004; Stevens et al., 1975) and the respiration fractionation factor
155 (α_r) is commonly assumed with a value of 0.982 for community respiration (Quay et al., 1995).

The parameter $\frac{^{18}}{^{16}}\text{O}_g$, described in equation 3, accounts for the isotope ratio of net air-water fluxes and is calculated as:

$$\frac{^{18}}{^{16}}\text{O}_g = \frac{\alpha_g * \left(\frac{^{18}}{^{16}}\text{O}_a * \alpha_s - \frac{\text{O}_2}{\text{O}_{2s}} * \frac{^{18}}{^{16}}\text{O} \right)}{\left(1 - \frac{\text{O}_2}{\text{O}_{2s}} \right)} \quad (3)$$

With α_g being the fractionation factor for gas transfer velocities (0.9972 at 20 °C; Knox et al., 1992) and α_s the fractionation factor for oxygen dissolution in water (1.0007 at 28 °C) as calculated by Benson and Krause, (1984). The atmospheric oxygen isotopic ratio $\left(\frac{^{18}}{^{16}}\text{O}_a \right)$ is known with a value of + 23.9 ‰ (Dordoni et al., 2022). The $\frac{\text{O}_2}{\text{O}_{2s}}$ ratio represents the concentration of
160 DO in the sample relative to the maximum temperature-dependent equilibrium concentration after Henry's law. Multiplying this ratio by 100 is equal to DO saturation (DO %) as measured in the field.

This model assumes isotopic steady-state conditions, constant α_r , and simplified gas exchange. Although it cannot resolve short-term variability such as diel cycles, these effects appear minor in the entire Danube mainstream and renders the model suitable for this broad-scale seasonal and spatial assessment.

165 2.5 Statistical analyses

All statistical analyses were conducted in R (v.4.3.2; R Core Team, 2023) using the `lm()` function to create the linear model and `anova()` for variance analysis. The coefficient of determination (R^2) was reported to evaluate the proportion of variance in

170 $\delta^{18}\text{O}_{\text{DO}}$ explained by DO concentration (Figure 6a, b) and POC concentration (Figure 6c, d). To assess the effect of DO/POC concentration on $\delta^{18}\text{O}_{\text{DO}}$, we performed a one-way analysis of variance (ANOVA; Fisher, 1932) based on the following linear model:

$$\delta^{18}\text{O}_{\text{DO}} = \beta_0 + \beta_1 \text{DO/POC} + \varepsilon \quad (4)$$

where β_0 is the intercept, β_1 the regression coefficient, and ε the error term. ANOVA was used to determine whether DO/POC concentration explains a significant proportion of the variance in $\delta^{18}\text{O}_{\text{DO}}$. The p-value from the F-test was used to assess statistical significance with a commonly accepted threshold value α of 0.05.

3. Results

175 To investigate seasonal variations in DO, $\delta^{18}\text{O}_{\text{DO}}$ and POC concentrations, data were analyzed along the Danube River for the five different sampling campaigns in spring, summer, late summer, fall and winter. DO concentrations showed clear seasonal fluctuations and variations, that ranged from 0.16 mmol/L to 0.40 mmol/L (5.1 and 12.8 mg/L) (Figure 2a-d). The highest mean DO concentrations occurred during winter ($0.36 \text{ mmol/L} \pm 0.01$), while the lowest mean occurred in late summer ($0.25 \text{ mmol/L} \pm 0.03$). Spatial variability was most pronounced in summer 2023, late summer 2024 and spring 2024 (Figure 2a, d).
180 In contrast, concentrations during fall 2023 and winter 2024 showed little variation over the entire course of the river and ranged from ca. 0.30 to 0.35 mmol/L (Figure 2b, c). As indicated by the red arrows in Figure 2a, three DO maxima occurred during summer with two being the most pronounced: first in the mid Danube ($\sim 1220 \text{ km}$; 0.35 mmol/L) and second in the lower Danube ($\sim 440 \text{ km}$; 0.40 mmol/L). Similar but less pronounced patterns were also observed in spring and late summer. However, in these seasons, the downstream maximum did not exceed the upstream one and blue arrows mark the DO minima.

185





Figure 2: Dissolved oxygen (DO) concentrations (mmol/L) in the Danube River at various distances from the river mouth. Standard error bars are within the symbol size. Circles represent Danube River samples, triangles denote tributary samples, and dotted lines indicate the river km where tributaries enter. Vertical solid lines represent the Iron Gate I and II dams. The red arrows show DO maxima and blue arrows DO minima.

Stable isotope values measured for DO ($\delta^{18}\text{O}_{\text{DO}}$) ranged from +25.9 ‰ to +12.1 ‰ in the entire dataset (Figure 3a-d). Here, the equilibrium with atmospheric oxygen (+24.6 ‰ \pm 0.4 ‰) serves as the boundary value between photosynthesis and respiration (Dordoni et al., 2022). Values below this threshold refer to photosynthesis, while values above it refer to respiration. Consequently, samples from summer 2023, late summer 2024, and spring 2024 predominantly showed photosynthesis-driven signals with $\delta^{18}\text{O}_{\text{DO}}$ values lower than +24.6 ‰ (Figure 3a, d). Conversely, samples from fall 2023 and winter 2024 often fell into the respiration range, even though they were close to the equilibrium (Figure 3b, c). Variability in $\delta^{18}\text{O}_{\text{DO}}$ was highest during summer (+19.4 \pm 3.2 ‰), late summer (+22.4 \pm 2.3 ‰) and spring (+21.9 \pm 1.4 ‰). On the other hand, fall and winter exhibited lower variabilities with +24.3 \pm 1.1 ‰ and +24.7 \pm 0.4 ‰. As shown by the red arrows in Figure 3a, four minima in $\delta^{18}\text{O}_{\text{DO}}$ were observed during summer: one near the source region and three others at the same

200 locations as the DO maxima. Among these, the two most pronounced occurred in the mid Danube (+14.8 ‰) and the lower Danube (+12.1 ‰). In late summer, two $\delta^{18}\text{O}_{\text{DO}}$ minima occurred, with the upstream minimum being more pronounced than the downstream one. In spring, three smaller minima of $\delta^{18}\text{O}_{\text{DO}}$ appeared across the upper, middle and lower Danube.



205 **Figure 3: Stable isotopes of dissolved oxygen ($\delta^{18}\text{O}_{\text{DO}}$) in the Danube River at various distances from the river mouth: Standard error bars are within the symbol size. Circles represent Danube River samples, triangles denote tributary samples, and dotted lines indicate the river km where tributaries enter. Vertical solid lines represent the Iron Gate I and II dams, and the horizontal solid line at +24.6 ‰ is the equilibrium for atmospheric oxygen. Values >+24.6 ‰ indicate respiration and <+24.6 ‰ photosynthesis. The red arrows show $\delta^{18}\text{O}_{\text{DO}}$ minima.**

POC concentrations ranged from 0.01 to 0.26 mmol/L (0.1 to 3.1 mg/L) (Figure 4a-d) with pronounced seasonal fluctuations. Maxima in POC occurred in summer 2023, late summer 2024 and spring 2024, aligning well with elevated DO concentrations and inversely related $\delta^{18}\text{O}_{\text{DO}}$ minima. In contrast, POC levels in fall 2023 and winter 2024 were less variable and mostly remained below 0.10 mmol/L. As highlighted by the red arrows in Figure 4a, three maxima in POC concentrations

were also observed during summer, with two being the most pronounced: the first one in the mid Danube (~1210 km; 0.25 mmol/L) and the second in the lower Danube (~500 km; 0.24 mmol/L). Samples from late summer showed a scattered pattern, with slightly elevated values in the upper Danube, a weaker increase in the mid Danube and generally lower values downstream. Spring samples showed two peaks in the middle and lower Danube, where the downstream peak was smaller than the upstream one.




Figure 4: Particular organic carbon (POC) concentrations (mmol/L) in the Danube River at various distances from the river mouth. Standard error bars are within the symbol size. Circles represent Danube River samples, triangles denote tributary samples, and dotted lines indicate the river km where tributaries enter. Vertical solid lines represent the Iron Gate I and II dams. The red arrows indicate POC maxima.

R/P ratios were not plotted here over the entire range of the river but showed similar trends as shown for DO, $\delta^{18}\text{O}_{\text{DO}}$ and POC (Figure S1). However, a cross plot between $\delta^{18}\text{O}_{\text{DO}}$ and R/P ratio showed a non-linear relationship of both parameters (Figure 5). R/P ratios ranged from 0.1 in summer to 8.9 in fall. Spring and summer samples were more dominated by photosynthesis, with R/P ratios smaller than 1, while winter and fall samples were more dominated by respiration, with R/P ratios larger than 1. Late summer samples exhibited a mixed signal, with most values in a range of overlapping photosynthesis and respiration and R/P values mostly larger than 1.




Figure 5: Cross plots of $\delta^{18}\text{O}_{\text{DO}}$ versus respiration/photosynthesis (R/P) ratios of the Danube River. The horizontal dotted line at +24.6 ‰ marks the equilibrium for atmospheric oxygen. Values with $>+24.6$ ‰ indicate respiration and $<+24.6$ ‰ photosynthesis. The vertical dotted line at R/P = 1 denotes the transition between photosynthesis and respiration. R/P values <1 indicate dominance by photosynthesis, and R/P >1 dominance by respiration.

Correlation plots between $\delta^{18}\text{O}_{\text{DO}}$ and DO concentrations are shown for spring and summer (Figure 6a), and for late summer, fall and winter (Figure 6b). Here the strongest and statistically significant correlations were observed in spring ($R^2 = 0.52$, $p < 0.05$) and summer ($R^2 = 0.75$, $p < 0.05$). In spring, higher DO coincided with lower $\delta^{18}\text{O}_{\text{DO}}$, while in summer, this relationship was even stronger. In contrast, correlations weakened in late summer ($R^2 = 0.14$, $p < 0.05$) and fall ($R^2 = 0.33$, $p < 0.05$), even though they were still statistically significant. This suggests that while $\delta^{18}\text{O}_{\text{DO}}$ and DO are still related, other factors such as respiration, chemical processes or elevated groundwater input likely contributed to $\delta^{18}\text{O}_{\text{DO}}$ variability. In winter ($R^2 = 0.04$, $p > 0.05$), no significant correlation was observed between both parameters.

To further investigate potential biological influences on $\delta^{18}\text{O}_{\text{DO}}$, it was correlated with POC concentrations at selected areas of high DO contents (ca. 1600 to 200 km) that were found during spring and summer (Figure 6c) and for late summer, fall, and winter (Figure 6d). The strongest and statistically significant correlations were observed in spring ($R^2 = 0.60$, $p < 0.05$) and summer ($R^2 = 0.88$, $p < 0.05$). In spring, this correlation was moderately strong, and increasing POC concentrations

coincided with decreasing $\delta^{18}\text{O}_{\text{DO}}$ values. However, no direct biological influence on $\delta^{18}\text{O}_{\text{DO}}$ values could be identified in late summer ($R^2 = 0.16$), fall ($R^2 = 0.06$), and winter ($R^2 = 0.13$), as reflected by weak, non-significant correlations ($p > 0.05$).




Figure 6: Correlation plots for all Danube sampling locations of $\delta^{18}\text{O}_{\text{DO}}$ and dissolved oxygen (DO) in summer and spring (a), in late summer, fall, and winter (b). High productivity areas (~1600 to ~200 km) related correlation plots of $\delta^{18}\text{O}_{\text{DO}}$ and particulate organic carbon (POC) for the same periods are presented to highlight potential biological inputs (c, d). The negative correlation indicates that as POC increases as $\delta^{18}\text{O}_{\text{DO}}$ values decrease. This pattern is consistent with biological oxygen production ($< +24.6$ ‰) and consumption ($> +24.6$ ‰), as oxygen produced by algal activity is usually enriched in ^{16}O when compared to the atmosphere. These results support the role of algae-derived organic matter in influencing the $\delta^{18}\text{O}$ signature of dissolved oxygen.

4. Discussion

4.1 Seasonal Dynamics

Seasonal variations in DO and $\delta^{18}\text{O}_{\text{DO}}$ along the Danube River are primarily influenced by temperature, biological activity, and atmospheric exchange. During winter, colder water temperatures decreased to 6.5 °C and thereby enhanced O_2 solubility (Figure S2c) (Rettich et al., 2000). Additionally, higher river discharge and associated turbulence likely accelerated atmospheric oxygen exchange and contribute to the highest DO concentrations measured throughout the year (Figure 2c; Figure S3, S4c) (Vautier et al., 2020). The associated $\delta^{18}\text{O}_{\text{DO}}$ data support this interpretation because values remained consistently close to those expected for atmospheric equilibrium (± 24.6 ‰ ± 0.4 ‰) (Figure 3c; Figure S5). Although

260 respiration must have occurred as well, its overall impact on the DO budget was likely minimal and largely masked by continuous atmospheric equilibration (Figure 3c, 6b). At first glance, associated R/P ratios might contradict this interpretation with values larger than 1 (Figure 5, Figure S1c). However, according to Dordoni et al. (2022), true dominance of respiration only becomes evident when R/P ratios exceed 2. Therefore, winter conditions reflect a mixed signal, with minor respiration influences and dominant atmospheric exchange.

265 With rising temperatures and longer daylight in spring, biological activity became more prominent and led to moderate to high DO concentrations (Figure 2d; Figure S2d, S3) (Wetzel, 2011). The associated $\delta^{18}\text{O}_{\text{DO}}$ values indicated enhanced photosynthetic activity, as they gradually shifted from atmospheric equilibrium toward less positive values (Figure 3d; Figure S5). Such a photosynthesis increase was further confirmed by slight rises in POC concentrations when compared to winter. While some of this POC may have originated from external sources, such as soil material washed into the river
270 (Aramaki et al., 2010; Reddy et al., 2021), the concurrent buildup of POC, DO, and $\delta^{18}\text{O}_{\text{DO}}$ suggests an accumulation of algae biomass driven by in-river photosynthetic activity (Figures 2d, 3d, 4d). A moderate but statistically significant correlation between DO concentration and their $\delta^{18}\text{O}_{\text{DO}}$ values ($R^2 = 0.52$, $p < 0.05$) further supported direct influences of photosynthesis on DO, thus highlighting the link between oxygen production and its photosynthetic source. Although photosynthesis predominantly influenced DO in spring, atmospheric exchange likely remained a significant contributor. This process was
275 likely supported by cool temperatures that remained below 18 °C and elevated discharge rates that also increased turbulent flow (Figures S2d, S3d). Moreover, the interplay between biological and physical processes was also reflected by R/P ratios (Figure 5, Figure S1d). Most of these ratios remained below 1 and indicated shifts towards photosynthesis. However, occasional mixed signals, with R/P ratios around and larger than 1, suggest that both atmospheric exchange and respiration processes also influenced the river's DO budget.

280 The intensification of biological activity in summer was most evident by $\delta^{18}\text{O}_{\text{DO}}$ values that decreased to +12.1 ‰ (Figure 3a; Figure S5). This distinct shift marks the time of peak photosynthetic activity in the Danube and is particularly notable because such pronounced deviations from atmospheric equilibrium are typically observed in standing waters (e.g., Quay et al., 1995; Wassenaar, 2012), or smaller streams (e.g., Parker et al., 2010; Wassenaar et al., 2010), but rarely in larger river systems (Quay et al., 1995) and support our finding that rivers are capable of strong photosynthetic DO input. The strong
285 photosynthetic signal was further reflected in the increases of POC concentrations, that indicate an accumulation of organic matter. Overall, the simultaneous increases of POC and DO, along with parallel decreases in $\delta^{18}\text{O}_{\text{DO}}$ and characteristic fluctuations in R/P ratios that also consistently fell below 1 in summer, underscore intensified photosynthetic activity (Figures 2a, 3a, 4a, 5; Figure S1a). The strong linear correlation and statistical significance between $\delta^{18}\text{O}_{\text{DO}}$ and DO further confirm this relationship ($R^2 = 0.76$, $p < 0.05$; Figure 6a). These indicators reflect the preferential input of ^{16}O via photosynthetic water
290 splitting, a process that proceeds without isotopic fractionation (Mader et al., 2017). Interestingly, despite such strong biological DO inputs, DO concentrations were often lower in summer when compared to winter and spring (Figure S3). These declines in DO concentrations can be attributed to higher water temperatures (Figure S2), which reduce O_2 solubility (Rettich et al., 2000). Another factor that may have reduced DO could have been the simultaneous decomposition of POC by respiration.

If so, and while this process likely contributed to DO depletion, its impact on POC levels must have been minor. In fact, POC
295 input appeared to outperform its consumption, as indicated by rising concentrations (Figure 4a). However, it is possible that
short term fluctuations in POC were not fully captured by the sampling frequency and implies that elevated POC levels do not
necessarily contradict intensified decomposition activity under warmer conditions. Although DO levels were higher in winter,
photosynthetic DO production likely played an important ecological role during warmer periods when atmospheric O₂
exchange was less effective. However, such photosynthetic DO increases may also associate with declines during nighttime
300 due to ongoing respiration. In riparian zones and in headwaters as this process may have been more important than in the main
river, where our limited night measurements indicated only minor diel variations. Notably, recent research has shown that
 $\delta^{18}\text{O}_{\text{DO}}$ responds more sensitively to metabolic shifts than DO concentrations, as it can detect transitions between
photosynthesis and respiration before they become evident in net DO concentrations (Dordoni et al., 2024). Such observations
justify the additional application of stable isotopes.

305 Late summer samples differed from those of mid-summer, despite similar water temperatures (Figure S2a). During
this later period, DO concentrations reached their lowest levels, particularly in the Sava River, where oxygen levels decreased
to 0.16 mmol/L. After its confluence with the Danube, DO levels showed a slight recovery but remained depleted at around
0.20 mmol/L (see blue arrow in Figure 2a). Given that healthy aquatic systems should maintain DO levels above 0.156 mmol/L
(WHO, Atlas Scientific), these late summer values raise concerns about critical oxygen depletion. This concern is further
310 supported by a shift in R/P ratios from a photosynthesis-dominated signal (R/P<1) to a mixed signal (R/P>1) (Figure 5; Figure
S1a), thus indicating that respiration rates had begun to exceed those of photosynthesis. In parallel, DO saturations increasingly
fell below 100%, thus suggesting net oxygen production and further supporting the interpretation of enhanced respiration.
Additionally, compared to mid-summer, the weakening correlation between DO and $\delta^{18}\text{O}_{\text{DO}}$ ($R^2 = 0.14$) suggests a reduced
influence of photosynthesis and a relatively greater contribution of respiration or atmospheric exchange, both of which can
315 introduce more variable isotopic signatures and alter the relationship (Figure 6b). A combination of declining primary
production, enhanced respiration, and potential organic matter decomposition likely contributed to the observed decrease in
DO concentrations.

In fall, declining water temperatures increased O₂ solubility and led to a moderate rise in DO concentrations when
compared to late summer (Figure 2b; Figure S2b) (Rettich et al., 2000). These temperature decreases also contributed to a
320 more homogeneous DO distribution along the entire river system, which showed a large-scale seasonal effect that contrasted
with the localized areas of high DO production in summer. This seems plausible because such temperature shifts by changing
seasons are able to affect an entire river system. As a result, $\delta^{18}\text{O}_{\text{DO}}$ values shifted towards more positive values and approached
those of atmospheric exchange (Figure 3b; Figure S5). On the other hand, reduced light intensity and a decline in
photosynthetic activity during fall also affected the entire river system (Aruga, 1965; Collins and Boylen, 1982). With the
325 decline in photosynthesis, respiration became the dominant process, as also reflected by R/P ratios that were mostly above 2,
especially in the lower section of the Danube (Figure 5; Figure S1b). This transition from a photosynthesis to a respiration-
dominated river was likely further enhanced by accumulation of organic material over the summer, which provided fresh

biomass for decomposition by respiration in fall. (DeNicola, 1996; Uehlinger et al., 2000). Although in fall increased respiration was evident in the R/P ratios, $\delta^{18}\text{O}_{\text{DO}}$ levels remained close to atmospheric equilibrium (Figure 3b; Figure S5). This pattern suggests that during this time the Danube system is buffered by atmospheric O_2 exchange which serves as a crucial DO source preventing DO depletion despite the dominance of respiration.

Overall, seasonal fluctuations in DO and $\delta^{18}\text{O}_{\text{DO}}$ along the Danube River were primarily influenced by temperature, associated atmospheric exchange and biological activity. In spring and summer, photosynthesis was an important process that increased DO and lowered $\delta^{18}\text{O}_{\text{DO}}$, while in late summer, enhanced respiration became more important predominant and caused lower DO levels and higher $\delta^{18}\text{O}_{\text{DO}}$, particularly evident in the Sava tributary. With cooler temperatures in fall, DO levels recovered, although respiration remained an important driver. In winter, oxygen solubility was the dominant control, with minimal biological influence.

4.2 Areas of increased primary production

Beyond the seasonal dynamics, our data revealed three areas of increased DO production. They occurred during the warm season, with one area located in the middle and the other in the lower section of the river. These areas were characterized by elevated DO and POC concentrations, while $\delta^{18}\text{O}_{\text{DO}}$ showed corresponding decreases (Figures 2a, d; 3a, d; 4a, d; Figure S6a-c). Especially the pronounced decline in $\delta^{18}\text{O}_{\text{DO}}$ indicated enhanced photosynthesis, which has already been identified as the primary driver of oxygen production during the warmer months (Figure 6a, b). The strong photosynthetic impact is further supported by correlations between $\delta^{18}\text{O}_{\text{DO}}$ values and POC concentrations (Figure 6c). This relationship indicates the importance of autotrophic organisms as key DO sources in these highly productive river sections during spring and summer and is supported by previous studies on the Danube (Hein et al., 1999; Riedler P. and Schagerl M., 1998).

As the warm season progressed, the correlation between $\delta^{18}\text{O}_{\text{DO}}$ and POC strengthened, thus reflecting the increasing role of autotrophic production, with a moderate correlation in spring ($R^2 = 0.60$) that became even stronger in summer ($R^2 = 0.88$). These positive correlations indicate that a substantial portion of the organic material must have originated from primary producers and directly contributed to the formation of these maxima. This pattern aligns well with the composition of the Danube's phytoplankton community, which is primarily composed of diatoms, alongside notable co-occurrence of Cyanobacteria, Chlorophyta and Cryptophyta. In particular, the downstream shift from benthic to planktonic diatoms, observed by (Liška et al., (2021), likely supported enhanced primary production in the middle and lower Danube and supports the isotope signals. Although the same areas of elevated productivity did not change in late summer (Figure 2a, 3a, 4a), the previously strong correlation disappeared during this period (Figure 6d). This lack of correlation may result from the gradual depletion of DO and POC during this transition period, thereby suggesting a weakening of autotrophic activity. At the same time, the increasing influence of allochthonous POC sources (i.e., material transported into the river) may have further decoupled the relationship between $\delta^{18}\text{O}_{\text{DO}}$ and POC. Despite occasionally elevated DO levels, this seasonal shift likely reflects a decline in primary production as environmental conditions changed and relative contribution of autotrophic material to the overall organic matter pool decreased.

The link between primary producers and DO observed in this study is further supported by previous research on the Danube. For instance, phytoplankton data from Literáthy et al. (2002) showed that the highest biomass occurred in the middle section of the river, thus aligning with the area identified by our study and supports the connection between increased primary production and DO-rich zones. Furthermore, Dokulil, (2015) highlighted findings from several chlorophyll- α studies in the Danube that indicated peak algae growth during summer with significant temporal variations in both the intensity and timing of algae blooms across several years (e.g., 1988, 1998 and 2001). The agreement between these findings and our data suggests that algal blooms occur regularly in this section of the Danube, thus contributing to and reinforcing seasonal variations.

One key factor that influences these areas of increased productivity is the decreasing flow gradient of the Danube (Figure S6d). According to Habersack et al. (2016), the riverbed slope decreases in its middle section from a steeper gradient of about 0.4 % in the upper Danube to a much flatter gradient of about 0.1 %. The observed decrease in slope already begins in the region of the first lower productivity area and becomes increasingly pronounced towards the second and third high productivity area in the middle and lower Danube. This reduction in slope decreases the velocity of the river and creates flow conditions that favor primary producers. In addition, slower flow velocities enhance sedimentation, which in turn decreases turbidity and light penetration and further supports photosynthetic activity. These conditions also align with findings of Dokulil (2006, 2015), who identified the middle reach of the Danube as an optimal zone for primary production due to moderate flow velocities and increased light availability. Moreover, the Danube had low to moderate discharge levels during summer that could further intensify these effects (Figure S4a). While not at its lowest, this discharge still enables longer water residence times that increase light exposure for algae and create favorable growth conditions (Kamjunke et al., 2021; Weitere and Arndt, 2002).

Another factor that may have contributed to the observed productivity patterns is the proximity of Budapest (Figure S6a-d). Located upstream of the productivity maximum in the mid Danube, it could influence primary production through nutrient inputs. Urban areas such as Budapest are known sources of nutrient pollution due to wastewater discharge (Nyenje et al., 2010; Xia et al., 2016), while agricultural activities further contribute to nutrient loading. However, our analyses were unable to reveal links between nutrient concentrations, POC, DO levels or $\delta^{18}\text{O}_{\text{DO}}$ values. For instance, nitrate concentrations ranged around a mean value of $0.10 \text{ mmol/L} \pm 0.0025$ (standard error) throughout the year and along the entire river (Figure S7a-d), while phosphate levels remained consistently below the detection limit. These results align well with previous findings published by Liška et al. (2021), who reported stable nitrate concentrations in the Danube over the past decades without noticeable peaks from potential point sources. One possible explanation is that primary producers rapidly absorb and utilize available nutrients and prevent their accumulation in measurable concentrations (Joint et al., 2001; Wetzel, 2011). This mechanism could explain the observed productivity patterns, despite minimal variance in nutrient levels and may particularly account for undetectable phosphate concentrations, because this nutrient is rapidly taken up by primary producers. To further investigate this possibility, more detailed studies on aquatic biomass and nutrient uptake processes would be required.

Following the upstream productivity maximum near Budapest, DO concentrations declined abruptly, together with decreases in POC and increases in $\delta^{18}\text{O}_{\text{DO}}$ values (Figures 2a, d; 3a, d; 4a, d). Previous studies have also documented reductions

395 in phytoplankton biomass and chlorophyll- α concentrations in this region (Dokulil, 2006, 2015; Dokulil and Kaiblinger, 2008; Literáthy et al., 2002). These patterns suggest that external factors, particularly the inflows of the Tisa and Sava rivers, play a role. Both tributaries introduce large volumes of water and contribute substantial dilution to the main course of the Danube. Although the Tisa and Sava rivers could not be sampled during all campaigns, data from late-summer and spring indicate that they contain lower DO concentrations (Figure 2a, d), higher $\delta^{18}\text{O}_{\text{DO}}$ values, and reduced POC levels compared to the main
400 stem of the Danube (Figure 3a, d; 4a, d). This observation supports dilution as a plausible mechanism for the observed decrease. Further evidence arises from previous studies, which estimated that this confluence with the Danube becomes diluted by approximately 27 % during average discharge conditions (Dokulil, 2015). Discharge data from summer and spring confirm this effect and show a substantial increase in total river discharge following the confluence with these tributaries (Figure S4a, d). Additionally, the inflow of the Morava River (not analyzed in this study) likely added further dilution effects (Dokulil,
405 2015). Moreover, a similar dilution effect likely occurred between the smaller upstream increase and the mid Danube productivity maximum. In this case it was influenced by the March and Váh Rivers. Although not sampled during summer, both tributaries contribute substantial discharge and likely carry water with lower DO and POC and higher $\delta^{18}\text{O}_{\text{DO}}$, thereby contributing to the observed decline.

Beyond these hydrological factors, land use including agriculture and urbanization across the basin may act as key
410 drivers on DO dynamics. Although nitrate levels remain relatively low, and indicate no critical eutrophication risk, these inputs still support the observed productivity maxima. To better trace and understand sources and transformation of land use and associated nutrient influences on DO dynamics, further investigations with detailed GIS mapping and other tracers such as nitrogen isotopes may be necessary.

Further downstream, the Danube featured a second and even more intense productivity zone with similar patterns to
415 the upstream peak area. While this area shows increased chlorophyll- α and suspended solids (Dokulil, 2015; Dokulil and Kaiblinger, 2008), phytoplankton biomass does not rise accordingly (Literáthy et al., 2002), thus suggesting an unconventional productivity pattern. On the other hand, a pronounced decrease in $\delta^{18}\text{O}_{\text{DO}}$ indicated active photosynthesis despite potential phytoplankton growth limitations (Wetzel, 2011). Notably, this region exhibited the lowest $\delta^{18}\text{O}_{\text{DO}}$ values in the study, along with R/P ratios close to 0 in summer, thus underscoring the significance of photosynthetic activity (Figure 5; Figure S1a).
420 These findings align with those of Dokulil (2006, 2015), who reported that despite significant increases in chlorophyll- α , primary production remained low due to poor light availability in the water column that in turn have been caused by elevated turbidity. Similar to the upstream peak, the emergence of this second productivity zone coincided with a decrease of the river gradient (0.05 – 0.01 %) that likely fosters favorable hydrodynamic conditions for algae (Habersack et al., 2016).

The spatial patterns observed also coincide with the above-discussed seasonality. They enhance spatial DO dynamics
425 and play a crucial role, as the second most productive area is clearly visible in summer, but weakens and even disappears by late summer (Figure 2a, 3a). Initially, the peak was driven by primary production, but as the season progressed, it declined rapidly. This decrease was also reflected by the absence of a pronounced $\delta^{18}\text{O}_{\text{DO}}$ peak and a shift in R/P ratios toward respiration in late summer (Figure 5b; Figure S1a) and highlights growing influences of heterotrophic processes, particularly in the lower

section of the river. Additionally, the decreasing correlation between $\delta^{18}\text{O}_{\text{DO}}$ and POC concentration in late summer suggests
430 a shift towards respiration (Figure 6d). While a strong correlation in summer indicated active primary production, this link
weakened as phytoplankton started to decompose, thus leading to organic matter accumulation and increased microbial
respiration. This transition further supports the interpretation that the second peak was initially fueled by photosynthesis during
slow flow but shifted toward organic matter degradation as the season progressed. Therefore, areas with shallow gradients not
only promote algal growth but also facilitate biomass accumulation and subsequent respiration, making them key zones of
435 enhanced biogeochemical activity in the river.

Connecting these findings with previous research underscores the critical role of primary producers in shaping oxygen
dynamics and in influencing organic material distribution across the Danube River. The persistence of intensive algae growth
also highlights the need for stricter compliance with EU regulations on water quality. Since DO serves as a key parameter for
assessing ecosystem health, further monitoring is essential to mitigate human-induced impacts and maintain balanced aquatic
440 conditions in the Danube system.

5. Conclusions

Our study demonstrated that seasonal variations in temperature, biological activity and hydrodynamic conditions
drive complex seasonal and spatial DO dynamics in the Danube River. In spring and summer, enhanced photosynthesis raised
POC and DO levels, while lowering associated $\delta^{18}\text{O}_{\text{DO}}$ values. This precise source identification is novel in the Danube River
445 and serves as a crucial indicator of ecosystem health. Photosynthetic effects were particularly evident in two areas of higher
productivity where reduced slopes resulted in slower flow velocities and lower turbulence. In these zones, where river
conditions approached those of standing waters, primary producers played a critical role, as shown by strong correlations
between $\delta^{18}\text{O}_{\text{DO}}$ and POC. These findings underscore the importance of autotrophic DO production in the Danube and suggest
that with rising temperatures and resulting reduced DO solubility the ecosystem may increasingly rely on this internal form of
450 oxygen input. However, tributary inflows from the Tisa and Sava Rivers diluted biomass and organic material inputs and
contributed to declines in DO and POC levels downstream.

Although the Danube River's water quality has improved over last three decades, vulnerable sections such as the Sava
and Lower Danube continue to face the risk of further DO depletion. This risk is especially pronounced during late summer
and early fall, when photosynthetic oxygen production declines and respiration intensifies while the river is often still too
455 warm to dissolve sufficient atmospheric O_2 . With ongoing climate warming, these potential DO deficits are expected to worsen,
a trend likely to extend to other temperate large rivers systems.

By integrating high-resolution measurements of DO, $\delta^{18}\text{O}_{\text{DO}}$, POC and R/P ratios, our study identified new arrays of
biogeochemical processes that regulate DO. Notably, DO concentrations alone are often insufficient to identify their sources,
but when combined with $\delta^{18}\text{O}_{\text{DO}}$ values, relative contributions of different sources (photosynthetic and atmospheric input) and
460 sinks (respiration) become clearer. Such knowledge is crucial for future river management and may help to plan ecological
conservation amid increasing environmental pressures. We therefore recommend high-resolution monitoring of DO and its

sources and sinks, with a particular focus on vulnerable regions with critically low DO in late summer. The potential to identify DO sources and sinks with the application of stable isotopes in other aquatic systems, offers a powerful tool for understanding and conserving diverse aquatic ecosystems.

465 **Data availability**

We already submitted our data sets from this manuscript to PANGAEA in March 2025, until its publication all data are available upon request.

Author contributions

JM: sampling, conceptualization, formal analysis, investigation, methodology, visualization, and writing (original draft
470 preparation). ANV: conceptualization, formal analysis, investigation, methodology, visualization, and writing (original draft preparation). CMS: investigation, methodology, visualization, and writing (original draft preparation). STW: investigation, methodology, visualization, and writing (original draft preparation). JACB: conceptualization, resources, supervision, and writing (original draft preparation).

Competing interests

475 The contact author has declared that none of the authors has any competing interests.

Acknowledgements

This research has been funded by the DALIA project (Danube Region Water Lighthouse Action) – project no. 101094070 that supported work by Friedrich-Alexander-University Erlangen-Nürnberg. Discharge visualization used in this manuscript was produced by the International Commission for the Protection of the Danube River (ICPDR). We acknowledge and appreciate
480 their contribution to the availability and accessibility of valuable data for our research. We acknowledge Open Ai's ChatGPT for its support in statistical computations and creating R-based visualizations. We also express our gratitude to Christian Hanke, Irene Wein, Anja Schuster and Robert van Geldern for providing technical and analytical support. We further thank SE-Tours GmbH and nicko Cruises Schiffreisen GmbH for allowing us to join their cruises at partially reduced prices to conduct our sampling from their ships. Their cooperation has greatly facilitated our research.

485 **Financial support**

This research has been supported by the DALIA project – project no. 101094070.

References

- Acreman, M. and Dunbar, M. J.: Defining environmental river flow requirements—a review, *Hydrol Earth Syst Sci*, 8 (5), 861–876, 2004.
- ICPDR-Database: <https://www.danubehis.org>, last access: 11 March 2025.
- Aramaki, T., Nagao, S., Nakamura, Y.-H., Uchida, M., and Shibata, Y.: The Effects of Rainfall on Carbon Isotopes of POC in the Teshio River, Northern Japan, *Radiocarbon*, 52(2), 808–814, 2010.
- Aruga, Y.: Ecological Studies of Photosynthesis and Matter Production of Phytoplankton II. Photosynthesis of Algae in Relation to Light Intensity and Temperature, Tokyo, 78, 360–365, 1965.
- Baertschi, P.: Absolute ^{18}O content of standard mean ocean water, *Earth Planet Sci Lett*, 31(3), 341–344, 1976.
- Barth, J. A. C., Tait, A., and Bolshaw, M.: Automated analyses of $^{18}\text{O}/^{16}\text{O}$ ratios in dissolved oxygen from 12-mL water samples, *Limnol Oceanogr Methods*, 2(2), 35–41, <https://doi.org/10.4319/lom.2004.2.35>, 2004.
- Belletti, B., Rinaldi, M., Buijse, A. D., Gurnell, A. M., and Mosselman, E.: A review of assessment methods for river hydromorphology, *Environ Earth Sci*, 73(5), 2079–2100, <https://doi.org/10.1007/s12665-014-3558-1>, 2015.
- Benson, B. B. and Krause, D.: The concentration and isotopic fractionation of oxygen dissolved in freshwater and seawater in equilibrium with the atmosphere, *Limnol Oceanogr*, 29(3), 620–632, <https://doi.org/10.4319/lo.1984.29.3.0620>, 1984.
- Benson, B. B., Krause, D., and Peterson, M. A.: The Solubility and Isotopic Fractionation of Gases in Dilute Aqueous Solution. I. Oxygen, *J Solution Chem*, 8(9), 1979.
- Borgwardt, F., Robinson, L., Trauner, D., Teixeira, H., Nogueira, A. J. A., Lillebø, A. I., Piet, G., Kuemmerlen, M., O’Higgins, T., McDonald, H., Arevalo-Torres, J., Barbosa, A. L., Iglesias-Campos, A., Hein, T., and Culhane, F.: Exploring variability in environmental impact risk from human activities across aquatic ecosystems, *Science of the Total Environment*, 652, 1396–1408, <https://doi.org/10.1016/j.scitotenv.2018.10.339>, 2019.
- Carpenter, S. R., Caraco, N. F., Correll, D. L., Howarth, R. W., Sharpley, A. N., and Smith, V. H.: Nonpoint pollution of surface waters with phosphorus and nitrogen, *Ecological Applications*, 8(3), 559–568, [https://doi.org/10.1890/1051-0761\(1998\)008\[0559:NPOSWW\]2.0.CO;2](https://doi.org/10.1890/1051-0761(1998)008[0559:NPOSWW]2.0.CO;2), 1998.
- Collins, C. D. and Boylen, C. W.: Physiological responses of *Anabaena Variabilis* (Cyanophyceae) to instantaneous exposure to various combination of light intensity and temperature, *J Phycol*, 18(2), 206–211, <https://doi.org/10.1111/j.1529-8817.1982.tb03175.x>, 1982.
- Coplen, T. B.: Guidelines and recommended terms for expression of stable-isotope-ratio and gas-ratio measurement results, *Rapid Communications in Mass Spectrometry*, 25(17), 2538–2560, <https://doi.org/10.1002/rcm.5129>, 2011.
- DeNicola, D. M.: Periphyton responses to temperature, in: Stevenson, R.J., Bothwell, M.L., Lowe, R.L. (Eds.), *Algal Ecology: Freshwater Benthic Ecosystems*, Academic Press, San Diego, 149–181, 1996.

- Dokulil, M. T.: Assessment of potamoplankton and primary productivity in the river Danube: A review. In Proceedings 36th International Conference of IAD. Austrian Committee Danube Research/IAD. Available from http://www.oen-iad.org/conference/docs/1_introduutory/dokulil.pdf (pp. 1-5), 2006.
- Dokulil, M. T.: Phytoplankton of the River Danube: Composition, Seasonality and Long-Term Dynamics, The Danube River Basin, 411–428 pp., https://doi.org/10.1007/698_2014_293, 2015.
- Dokulil, M. T. and Kaiblinger, C. M.: Phytoplankton. In: Liska I, Wagner F, Slobodnik J (eds) Joint danube survey 2, Final scientific report. ICPDR. <http://www.icpdr.org/main/activities-projects/joint-danube-survey-2>, 2008.
- Dordoni, M., Seewald, M., Rinke, K., Schmidmeier, J., and Barth, J. A. C.: Novel evaluations of sources and sinks of dissolved oxygen via stable isotopes in lentic water bodies, *Science of the Total Environment*, 838, <https://doi.org/10.1016/j.scitotenv.2022.156541>, 2022.
- Dordoni, M., Tittel, J., Rosenlöcher, Y., Rinke, K., and Barth, J. A. C.: Metabolic activity of *Planktothrix rubescens* and its consequences on oxygen dynamics in laboratory experiment: A stable isotope study, *J Phycol*, 60, 642–653, <https://doi.org/10.1111/jpy.13455>, 2024.
- Dudgeon, D., Arthington, A. H., Gessner, M. O., Kawabata, Z. I., Knowler, D. J., Lévêque, C., Naiman, R. J., Prieur-Richard, A. H., Soto, D., Stiassny, M. L. J., and Sullivan, C. A.: Freshwater biodiversity: Importance, threats, status and conservation challenges, *Biol Rev Camb Philos Soc*, 81(2), 163–182, <https://doi.org/10.1017/S1464793105006950>, 2006.
- EEA [European Environment Agency]: Water resources across Europe-confronting water stress: an updated assessment. EEA Report 12/2021, <https://doi.org/10.2800/320975>, 2021.
- Eisenstadt, D., Barkan, E., Luz, B., and Kaplan, A.: Enrichment of oxygen heavy isotopes during photosynthesis in phytoplankton, *Photosynth Res*, 103, 97–103, <https://doi.org/10.1007/s11120-009-9518-z>, 2010.
- Fisher, R. A.: Statistical Methods for Research Workers (4th ed.). Oliver & Boyd, 1932.
- Fowler, D., Coyle, M., Skiba, U., Sutton, M. A., Cape, J. N., Reis, S., Sheppard, L. J., Jenkins, A., Grizzetti, B., Galloway, J. N., Vitousek, P., Leach, A., Bouwman, A. F., Butterbach-Bahl, K., Dentener, F., Stevenson, D., Amann, M., and Voss, M.: The global nitrogen cycle in the Twentyfirst century, *Philosophical Transactions of the Royal Society B: Biological Sciences*, 368(1621), <https://doi.org/10.1098/rstb.2013.0164>, 2013.
- Franklin, P. A.: Dissolved oxygen criteria for freshwater fish in New Zealand: A revised approach, *N Z J Mar Freshwater Res*, 48(1), 112–126, <https://doi.org/10.1080/00288330.2013.827123>, 2014.
- van Geldern, R. and Barth, J. A. C.: Optimization of instrument setup and post-run corrections for oxygen and hydrogen stable isotope measurements of water by isotope ratio infrared spectroscopy (IRIS), *Limnol Oceanogr Methods*, 10(12), 1024–1036, <https://doi.org/10.4319/lom.2012.10.1024>, 2012.
- Grizzetti, B., Pistocchi, A., Liqueste, C., Udias, A., Bouraoui, F., and Van De Bund, W.: Human pressures and ecological status of European rivers, *Sci Rep*, 7(1), <https://doi.org/10.1038/s41598-017-00324-3>, 2017.
- Guy, R. D., Fogel, M. L., and Berry, J. A.: Photosynthetic Fractionation of the Stable Isotopes of Oxygen and Carbon, *Plant Physiol*, 101(1), 37–47, 1993.

- Habersack, H., Hein, T., Stanica, A., Liska, I., Mair, R., Jäger, E., Hauer, C., and Bradley, C.: Challenges of river basin management: Current status of, and prospects for, the River Danube from a river engineering perspective, *Science of the Total Environment*, 543, 828–845, <https://doi.org/10.1016/j.scitotenv.2015.10.123>, 2016.
- Heddam, S.: Modelling hourly dissolved oxygen concentration (DO) using dynamic evolving neural-fuzzy inference system (DENFIS)-based approach: Case study of Klamath River at Miller Island Boat Ramp, OR, USA, *Environmental Science and Pollution Research*, 21(15), 9212–9227, <https://doi.org/10.1007/s11356-014-2842-7>, 2014.
- Hein, T., Baranyi, C., Heiler, G., Holarek, C., Riedler, P., and Schiemer, F.: Hydrology as a major factor determining plankton development in two floodplain segments and the River Danube, Austria, *Large Rivers*, 3, 439–452, 1999.
- Honisch, M., Hellmeier, C., and Weiss, K.: Response of surface and subsurface water quality to land use changes, *Geoderma*, 105(3–4), 277–298, [https://doi.org/https://doi.org/10.1016/S0016-7061\(01\)00108-2](https://doi.org/https://doi.org/10.1016/S0016-7061(01)00108-2), 2002.
- Hua, A. K.: Land Use Land Cover Changes in Detection of Water Quality: A Study Based on Remote Sensing and Multivariate Statistics, *J Environ Public Health*, 2017(1), <https://doi.org/10.1155/2017/7515130>, 2017.
- ICPDR: The Danube River Basin District Management Plan — Update 2015. International Commission for the Protection of the Danube River, 2015.
- Joint, I., Rees, A. P., Malcolm, E., and Woodward, S.: Primary production and nutrient assimilation in the Iberian upwelling in August 1998, *Prog Oceanogr*, 51(2–4), 303–320, 2001.
- Jurgens, B. C., McMahon, P. B., Chapelle, F. H., and Eberts, S. M.: An Excel workbook for identifying redox processes in ground water (No. 2009-1004), <https://doi.org/10.3133/ofr20091004>, 2009.
- Kamjunke, N., Rode, M., Baborowski, M., Kunz, J. V., Zehner, J., Borchardt, D., and Weitere, M.: High irradiation and low discharge promote the dominant role of phytoplankton in riverine nutrient dynamics, *Limnol Oceanogr*, 66(7), 2648–2660, <https://doi.org/10.1002/lno.11778>, 2021.
- Killgore, K. J. and Hoover, J. J.: Effects of Hypoxia on Fish Assemblages in a Vegetated Waterbody, *J. Aquat. Plant Manage*, 39(1), 40–44, 2001.
- Knox, M., Quay, P. D., and Wilbur, D.: Kinetic isotopic fractionation during air-water gas transfer of O₂, N₂, CH₄, and H₂, *J Geophys Res*, 97(C12), <https://doi.org/10.1029/92jc00949>, 1992.
- Limburg, J., Vrettos, J. S., Liable-Sands, L. M., Rheingold, A. L., Crabtree, R. H., and Brudvig, G. W.: A functional model for OO bond formation by the O₂-evolving complex in photosystem II, *Science* (1979), 283(5407), 1524–1527., 1999.
- Liška, I., Wagner, F., Sengl, M., Deutsch, K., Slobodník, J., and Paunovic, M.: Joint Danube survey 4 scientific report: A shared analysis of the Danube river. Vienna: International Commission for the Protection of the Danube River-ICPDR, 2021.
- Literáthy, P., Koller-Kreiml, V., and Liška, I.: Final Report of the Joined Danube Survey. International Commission for the Protection of the Danube River—ICPDR, Vienna, Austria, 262 pp., 2002.
- Mader, M., Schmidt, C., van Geldern, R., and Barth, J. A. C.: Dissolved oxygen in water and its stable isotope effects: A review, *Chem Geol*, 473, 10–21, <https://doi.org/10.1016/j.chemgeo.2017.10.003>, 2017.

- Mănoiu, V. M. and Crăciun, A. I.: Danube river water quality trends: A qualitative review based on the open access web of science database, *Ecohydrology and Hydrobiology*, 21(4), 613–628, <https://doi.org/10.1016/j.ecohyd.2021.08.002>, 2021.
- North, R. P., North, R. L., Livingstone, D. M., Köster, O., and Kipfer, R.: Long-term changes in hypoxia and soluble reactive phosphorus in the hypolimnion of a large temperate lake: Consequences of a climate regime shift, *Glob Chang Biol*, 20(3), 811–823, <https://doi.org/10.1111/gcb.12371>, 2014.
- Nyenje, P. M., Foppen, J. W., Uhlenbrook, S., Kulabako, R., and Muwanga, A.: Eutrophication and nutrient release in urban areas of sub-Saharan Africa - A review, *Science of the Total Environment*, 408(3), 447–455, <https://doi.org/10.1016/j.scitotenv.2009.10.020>, 2010.
- Odum, H. T.: Primary Production in Flowing Waters, *Limnol Oceanogr*, 1(2), 102–117, <https://doi.org/10.4319/lo.1956.1.2.0102>, 1956.
- Parker, S. R., Gammons, C. H., Poulson, S. R., DeGrandpre, M. D., Weyer, C. L., Smith, M. G., Babcock, J. N., and Oba, Y.: Diel behavior of stable isotopes of dissolved oxygen and dissolved inorganic carbon in rivers over a range of trophic conditions, and in a mesocosm experiment, *Chem Geol*, 269(1–2), 22–32, <https://doi.org/10.1016/j.chemgeo.2009.06.016>, 2010.
- Piatka, D. R., Wild, R., Hartmann, J., Kaule, R., Kaule, L., Gilfedder, B., Peiffer, S., Geist, J., Beierkuhnlein, C., and Barth, J. A. C.: Transfer and transformations of oxygen in rivers as catchment reflectors of continental landscapes: A review, <https://doi.org/10.1016/j.earscirev.2021.103729>, 1 September 2021.
- Poff, N. L. and Zimmerman, J. K. H.: Ecological responses to altered flow regimes: A literature review to inform the science and management of environmental flows, *Freshw Biol*, 55(1), 194–205, <https://doi.org/10.1111/j.1365-2427.2009.02272.x>, 2010.
- Qing, X., Yutong, Z., and Shenggao, L.: Assessment of heavy metal pollution and human health risk in urban soils of steel industrial city (Anshan), Liaoning, Northeast China, *Ecotoxicol Environ Saf*, 120, 377–385, <https://doi.org/10.1016/j.ecoenv.2015.06.019>, 2015.
- Quay, P. D., Wilbur, D. O., Richey, J. E., Devol, A. H., Benner, R., and Forsberg, B. R.: The 18O : 16O of dissolved oxygen in rivers and lakes in the Amazon Basin: Determining the ratio of respiration to photosynthesis rates in freshwaters, *Limnol. Oceanogr*, 40 (4), 718–729, 1995.
- R Core Team: R: A Language and Environment for Statistical Computing, [online] Available from: <https://www.r-project.org/>, 2023.
- Reddy, S. K. K., Gupta, H., Badimela, U., Reddy, D. V., Kurakalva, R. M., and Kumar, D.: Export of particulate organic carbon by the mountainous tropical rivers of Western Ghats, India: Variations and controls, *Science of the Total Environment*, 751, <https://doi.org/10.1016/j.scitotenv.2020.142115>, 2021.
- Rettich, T. R., Battino, R., and Wilhelm, E.: Solubility of gases in liquids. 22. High-precision determination of Henry's law constants of oxygen in liquid water from T = 274 K to T = 328 K, *Journal of Chemical Thermodynamics*, 32(9), 1145–1156, <https://doi.org/10.1006/jcht.1999.0581>, 2000.

- Riedler P. and Schagerl M.: Pelagic primary production and related parameters in the River Danube near Vienna (Austria),
 620 Archiv für Hydrobiologie, Supplementbände, 115(2), 139–151, 1998.
- Rosecrans, C. Z., Nolan, B. T., and Gronberg, J. A. M.: Prediction and visualization of redox conditions in the groundwater of
 Central Valley, California, J Hydrol (Amst), 546, 341–356, <https://doi.org/10.1016/j.jhydrol.2017.01.014>, 2017.
- Russ, M. E., Ostrom, N. E., Gandhi, H., Ostrom, P. H., and Urban, N. R.: Temporal and spatial variations in R:P ratios in Lake
 Superior, an oligotrophic freshwater environment, J Geophys Res Oceans, 109(C10), <https://doi.org/10.1029/2003JC001890>,
 625 2004.
- Seitzinger, S., Harrison, J. A., Böhlke, J. K., Bouwman, A. F., Lowrance, R., Peterson, B., Tobias, C., and Van Drecht, G.:
 Denitrification across landscapes and waterscapes: A synthesis, Ecological Applications, 16(6), 2064–2090,
[https://doi.org/10.1890/1051-0761\(2006\)016\[2064:DALAWA\]2.0.CO;2](https://doi.org/10.1890/1051-0761(2006)016[2064:DALAWA]2.0.CO;2), 2006.
- Sommerwerk, N., Hein, T., Schneider-Jacoby, M., Baumgartner, C., Ostojić, A., Siber, R., Bloesch, J., Paunović, M., and
 630 Tockner, K.: The Danube River Basin, in: Rivers of Europe, Elsevier, 59–112, [https://doi.org/10.1016/B978-0-12-369449-](https://doi.org/10.1016/B978-0-12-369449-2.00003-5)
 2.00003-5, 2009.
- Stevens, C. L. R., Schultz, D., Van Baalen, C., and Parker, P. L.: Oxygen Isotope Fractionation during Photosynthesis in a
 Blue-Green and a Green Algal, Plant Physiol, 56(1), 126–129, 1975.
- Stumm, W. and Morgan, J. J.: Aquatic Chemistry: Chemical Equilibria and Rates in Natural Waters, 3rd ed., Wiley-
 635 Interscience, New York, 455–464 pp., 1995.
- Suthar, S., Nema, A. K., Chabukdhara, M., and Gupta, S. K.: Assessment of metals in water and sediments of Hindon River,
 India: Impact of industrial and urban discharges, J Hazard Mater, 171(1–3), 1088–1095,
<https://doi.org/10.1016/j.jhazmat.2009.06.109>, 2009.
- Sutton, M., Howard, C. M., Erisman, J. W., Billen, G., Bleeker, A., Grennfelt, P., Van Grinsven, H., and Grizzetti, B.: The
 640 European Nitrogen Assessment: Sources, Effects and Policy Perspectives, Cambridge University Press, 2011.
- Tobias, C. R., Böhlke, J. K., and Harvey, J. W.: The oxygen-18 isotope approach for measuring aquatic metabolism in high-
 productivity waters, Limnol Oceanogr, 52(4), 1439–1453, <https://doi.org/10.4319/lo.2007.52.4.1439>, 2007.
- Uehlinger, U., König, C., and Reichert, P.: Variability of photosynthesis-irradiance curves and ecosystem respiration in a small
 river, Freshw Biol, 44(3), 493–507, <https://doi.org/10.1046/j.1365-2427.2000.00602.x>, 2000.
- 645 Vautier, C., Abhervé, R., Labasque, T., Laverman, A. M., Guillou, A., Chatton, E., Dupont, P., Aquilina, L., and de Dreuz, J. R.: Mapping gas exchanges in headwater streams with membrane inlet mass spectrometry, J Hydrol (Amst), 581,
<https://doi.org/10.1016/j.jhydrol.2019.124398>, 2020.
- Vörösmarty, C. J., Pahl-Wostl, C., Bunn, S. E., and Lawford, R.: Global water, the anthropocene and the transformation of a
 science, Curr Opin Environ Sustain, 5(6), 539–550, <https://doi.org/10.1016/j.cosust.2013.10.005>, 2013.
- 650 Wachs, B.: Zustand und Qualität der Donau. Verantwortung für einen europäischen Strom, Schriftenreihe des Bundesamtes
 für Wasserwirtschaft, 28–51, 1997.

- Wassenaar, L. I.: Dissolved oxygen status of Lake Winnipeg: Spatio-temporal and isotopic ($\delta^{18}\text{O}$ - O_2) patterns, *J Great Lakes Res*, 38(3), 123–134, <https://doi.org/10.1016/j.jglr.2010.12.011>, 2012.
- Wassenaar, L. I. and Koehler, G.: An on-line technique for the determination of the $\delta^{18}\text{O}$ and $\delta^{17}\text{O}$ of gaseous and dissolved oxygen, *Anal Chem*, 71(21), 4965–4968, <https://doi.org/10.1021/ac9903961>, 1999.
- Wassenaar, L. I., Venkiteswaran, J. J., Schiff, S. L., and Koehler, G.: Aquatic community metabolism response to municipal effluent inputs in rivers quantified using diel $\delta^{18}\text{O}$ values of dissolved oxygen, *Canadian Journal of Fisheries and Aquatic Sciences*, 67(8), 1232–1246, <https://doi.org/10.1139/F10-057>, 2010.
- Weitere, M. and Arndt, H.: Top-down effects on pelagic heterotrophic nanoflagellates (HNF) in a large river (River Rhine): Do losses to the benthos play a role?, *Freshw Biol*, 47(8), 1437–1450, <https://doi.org/10.1046/j.1365-2427.2002.00875.x>, 2002.
- Wetzel, R. G.: *Limnology: Lake and River Ecosystems*, 3rd edition., Academic Press, San Diego, California, 2011.
- Xia, Y., Ti, C., She, D., and Yan, X.: Linking river nutrient concentrations to land use and rainfall in a paddy agriculture–urban area gradient watershed in southeast China, *Science of the Total Environment*, 566, 1094–1105, <https://doi.org/10.1016/j.scitotenv.2016.05.134>, 2016.