

# Evolution of flood generating processes under climate change in France

Yves Tramblay <sup>1</sup>, Guillaume Thirel <sup>2,3</sup>, Laurent Strohmer <sup>2,4</sup>, Guillaume Evin <sup>5</sup>, Lola Corre <sup>6</sup>, Louis Heraut <sup>7</sup>, Eric Sauquet <sup>7</sup>

<sup>1</sup> Espace-Dev, Univ. Montpellier, IRD, Montpellier, France

<sup>2</sup> University of Paris-Saclay, INRAE, HYCAR, Antony, France

<sup>3</sup> Univ Toulouse, CNES/IRD/CNRS/INRAE, CESBIO, Toulouse, France

<sup>4</sup> Department of Ecohydrology and Biogeochemistry, Leibniz Institute of Freshwater Ecology and Inland Fisheries (IGB), Berlin, Germany

<sup>5</sup> Univ. Grenoble Alpes, INRAE, CNRS, IRD, Grenoble INP, IGE, Grenoble, France

<sup>6</sup> CNRM, Université de Toulouse, Météo-France CNRS, Toulouse, France

<sup>7</sup> INRAE, UR RiverLy, Villeurbanne, France

Correspondance to: Yves Tramblay ([yves.tramblay@ird.fr](mailto:yves.tramblay@ird.fr))

Revised manuscript, July 2025

## Abstract

The impact of climate change on floods varies spatially, and often the observed trends in flood characteristics can be explained by differentiated changes in flood-generating processes. This study explores changes in flood magnitude and flood-generating processes under different climate change scenarios for a large number of basins in France. It is based on an unprecedented exercise to model the impacts of climate change on hydrology, using a semi-distributed model (GRSD) applied to 3727 basins with 22 Euro-CORDEX bias-corrected climate projections using two greenhouse gas emission scenarios (RCP4.5 and RCP8.5). Annual maxima of daily simulated streamflow were extracted for the period 1975-2100, resulting in a set of over 10 million flood events, and a trend analysis was carried out on both flood magnitudes and flood generating processes. ~~Increasing~~The trends in flood magnitude ~~are contrasted, with increasing~~ trends in flood magnitudes are only found in the northern regions of France, although multi-model convergence rarely exceeds 60 %. The highest increases are observed for the 20-year floods and under the RCP8.5 scenario. A classification of floods according to their generating process revealed that floods linked to soil saturation represent more than half of all floods in France. The relative change in the importance of the different flood-generating processes is not spatially homogeneous and varies by region. ~~The proportion of floods linked to soil saturation excess is increasing in the Northeast, while decreasing in the southern Mediterranean regions. In these Mediterranean regions, the proportion of floods linked to infiltration excess related to extreme rainfall is increasing. The proportion of floods linked to soil saturation excess is decreasing while the proportion of floods linked to infiltration excess related to extreme rainfall is increasing, particularly in the southern Mediterranean regions.~~ Both the frequency and magnitude of floods linked to snowmelt processes are decreasing in mountainous areas. On the contrary, the most extreme floods associated with rainfall on dry soils tend to increase, in line with the increase of rainfall intensity. Overall, trends in antecedent soil moisture conditions are as important as trends in intense rainfall to explain flood hazard trends in the different climate projections. This study shows how important it is to decipher the changes in the different flood generating processes in order to better understand their evolution in different hydroclimatic regions.

## 1- Introduction

The impact of climate change on floods is uncertain, or unknown, in many regions of the world (Intergovernmental Panel On Climate Change (IPCC), 2023). In particular, it has been shown that competing changes in flood-generating processes can modulate, or even offset, changes in flood hazards (Ivancic and Shaw, 2015; Sharma et al., 2018; Trambly et al., 2019; Brunner et al., 2021; Zhang et al., 2022). For example, in the context of an increase in intense rainfall events in some temperate and arid regions, the concomitant drop in soil moisture, resulting in lower runoff coefficients, could result in a lack of trend in flood magnitude (Vasko and Nathan, 2019; Ho et al., 2022; Trambly et al., 2023; Scussolini et al., 2023). Similarly, the large decline in the frequency of snowmelt-induced floods in many regions may compensate for the increasing proportion of floods caused by rainstorms, resulting in the absence of trends in overall flood hazards in numerous regions worldwide where snowmelt is prevalent (Zhang et al., 2022). To comprehend the evolution of floods, it is essential to analyze not only their severity but also to consider in detail whether the processes that underpin them are also changing (Tarasova et al., 2019; Kemter et al., 2020; Blöschl, 2022a, b; Jiang et al., 2022; Tarasova et al., 2023; Kemter et al., 2023; McMillan et al., 2025). This is particularly salient

given that trends in extreme rainfall do not generally translate into the same trends in floods (Ivancic and Shaw, 2015; Sharma et al., 2018; Wasko et al., 2023). Numerous studies have focused on understanding and categorizing floods according to the hydroclimatic mechanisms behind their generation (Tarasova et al., 2019; Stein et al., 2020; Kemter et al., 2023). We can distinguish studies that use seasonality to categorize the various causal mechanisms based on their temporal proximity (Berghuijs et al., 2016, 2019; Trambly et al., 2021) from other studies that use different hydrological criteria, predominantly thresholds on precipitation intensity, snowmelt, or soil moisture levels, to differentiate between different types of floods (Froidevaux et al., 2015; Tarasova et al., 2018; Kim et al., 2019; Tarasova et al., 2020; Stein et al., 2020, 2021; Trambly et al., 2022; Tarasova et al., 2023; Trambly et al., 2023).

To provide future flood projections, it is necessary to use hydrological models to translate climate projections into hydrological projections, i.e. providing simulated streamflow values for the future. Such a modeling chain, which is generally constituted by greenhouse gas concentration scenarios, global (GCM) and regional (RCM) climate models, bias-correction methods, and hydrological models, necessarily suffers from uncertainties (Clark et al., 2016). The evolution of our society is inherently unpredictable, and thus, climate scenarios describe its evolution through greenhouse gas concentrations (Meinshausen et al., 2020). These concentrations, in turn, influence climate model outputs, which are subject to uncertainties related to process representation and climate variability (Knutti and Sedláček, 2013). Additionally, bias-correction methods and hydrological models are affected by process representation uncertainties (Teutschbein and Seibert, 2012; Maraun et al., 2017). Furthermore, their application in an extrapolation mode may compromise their temporal transferability. Indeed, these models and methods are generally calibrated, optimized, and evaluated over well-known past periods. With climate projections, they are used under potentially very different conditions, with unprecedented air temperatures, and possibly very different precipitation regimes, which may alter their transferability. This issue has been extensively documented in the context of hydrological modeling (Brigode et al., 2013; Thirel et al., 2015b; Dakhlaoui et al., 2017), and various protocols have been proposed to assess the robustness of these models (Klemeš, 1986; Thirel et al., 2015a). It is important to note that models of varying complexity exist, depending on their spatial discretization and the processes included in the equations, as well as the use of automatic parameter optimization (Hrachowitz and Clark, 2017). For large-scale studies, at the regional or even the global scale, most often the hydrological models are not calibrated for all stations but sometimes only in a subset of basins (Alfieri et al., 2015; Roudier et al., 2016; Do et al., 2020; Di Sante et al., 2021; Zhang et al., 2022). Nevertheless, all types of models can suffer from robustness issues (Santos et al., 2025). However, it has been shown that the relative uncertainty associated with hydrological modeling is typically lower for high flows, which are the focus of this study, in comparison to low flows (Vidal et al., 2016; Lemaitre-Basset et al., 2021).

This study analyzes a large ensemble of hydrological projections based on climate projections for a diverse set of basins in France, which represents a range of varied physiographic properties. The aim is to investigate the future evolution of flood events, both in terms of magnitude and flood-generating processes. In a multi-model context, it is important to analyze both the convergence of the scenarios for the different models and the signal of change given by these different models. The methodology enables a multi-factorial analysis of floods and the processes underlying their generation, leading to a better understanding of which factors

most influence flood trends. By distinguishing the hydro-climatic drivers regionally, we identify those that may either increase or decrease future flood risks.

## 2- Data and Methods

### 2.1 Hydrological model

The GRSD model, a semi-distributed rainfall-runoff model based on the lumped GR4J model structure (Perrin et al., 2003) and the CemaNeige snow model (Valéry et al., 2014), has been chosen for this study. The rationale behind this choice is its capacity to be applied to a very large range of hydro-meteorological conditions and numerous basins across the world, as shown by the GR4J or GRSD applications in Brazil (Kuana et al., 2024), Nepal (Nepal et al., 2017), Iran (Jahanshahi et al., 2025), Haiti (Bathelemy et al., 2024), Chile (Abbenante et al., 2024), Australia (Stephens et al., 2019), and France (De Lavenne et al., 2019; Lemaitre-Basset et al., 2024), as well as its large number of applications in the context of climate change (Chauveau et al., 2013; Cornelissen et al., 2013; Tian et al., 2013; Fabre et al., 2016; Stephens et al., 2018; Givati et al., 2019; Thirel et al., 2019; Tarek et al., 2021; Wasko et al., 2023; Poncet et al., 2024). The GRSD model calibration and simulation were applied within the airGR and airGRiwrn open-source R packages (Coron et al., 2017; Dorchies, 2022).

GR4J is a four-parameter rainfall-runoff model consisting of two conceptual stores: a production store and a routing store, whose flows are routed to the river after transformation by a unit hydrograph. While the GR4J model is applied on a sub-catchment spatial scale, and the simulated streamflow is routed from upstream to downstream with a lag function (de Lavenne et al., 2019), the CemaNeige snow model uses an additional sub-division of each sub-catchment into five zones of equal area to better describe the process heterogeneities linked to the topography (Valéry et al., 2014). GRSD provides daily streamflow simulation covering 3727 simulation points in France, and different variables have been extracted: the total, solid, and liquid precipitation, the soil water index (SWI, i.e. the ratio between water content and the storage capacity of the production store), and the snowmelt discharge (SMD, i.e. the water flow resulting from snowpack melting). [The GRSD simulations used in the present study are described in Sauquet et al., \(2024, 2025\).](#) The GRSD model has been calibrated over the 1976-2019 period, using the Kling-Gupta Efficiency criterion (Gupta et al., 2009) applied on squared-root transformed streamflow, against the observed streamflow of 611 gauging stations in France considered as nearly natural (Strohmenger et al., 2023), using the SAFRAN (Vidal et al., 2010) reanalysis as meteorological input. To simulate streamflow over the remaining - ungauged - stations, parameters were transferred from neighbor gauged catchments to the target ungauged catchments following Oudin et al. (2008), and these parameter sets were used to produce pseudo-observed streamflow on the ungauged stations, on which the GRSD model was subsequently calibrated (Sauquet et al., 2024, 2025). [The validation of the GRSD model is presented in Sauquet et al., 2024, 2025 and Héraut et al., 2024. In addition, the comparison of the observed and simulated 2-year and 20-year floods is presented in the supplementary materials, figure S7, showing a relative bias of -3% and mean absolute relative error of 16% for 2-year floods, and 20% for 20-year floods.](#)

### 2.2 Climate simulations

191  
192 The GRSD model was thereafter applied using as inputs climate projections from the  
193 Explore2-Climat-2022-ADAMONT dataset (Corre et al., 2025), available on the French  
194 national climate data portal (*DRIAS - les futurs du climat* - [www.drias-climat.fr](http://www.drias-climat.fr)), and described  
195 in Sauquet et al. (2025). This ensemble is derived from the 12-km resolution EURO-CORDEX  
196 ensemble (Jacob et al., 2018; Vautard et al., 2021), which consists of regional climate model  
197 simulations that downscale global climate model simulations over Europe from the CMIP5 (5th  
198 phase of the Coupled Model Intercomparison Project, (Taylor et al., 2012)). For this study, we  
199 use the 11 GCM/RCM pairs for which both emission scenarios (RCP4.5 and RCP8.5) are  
200 available (Table T1 in supplementary materials). The outputs of the climate models were  
201 statistically adjusted over mainland France, against the 8- km resolution SAFRAN reanalysis  
202 (Vidal et al. 2010). The bias correction method consists of a quantile mapping relying on  
203 seasonal weather regimes (ADAMONT method, Verfaillie et al., 2017). In the following, we  
204 use the daily rainfall data and the variables used to compute evapotranspiration (temperature,  
205 relative humidity, wind, radiation) from these bias adjusted simulations. These data are  
206 available from 1975-2100 at the daily time scale with an 8-km spatial resolution and have been  
207 aggregated at the catchment scale.

208  
209 Several studies describe the future climate changes in France based on these projections  
210 (Marson et al., 2024). Overall, they show a shift towards hotter and drier summers, as well as  
211 an expansion of temperate climate in mountainous regions (Strohmenger et al., 2024). Despite  
212 variations in the magnitude of warming depending on the scenario, time horizon, or level of  
213 warming considered, all studies identify common features. These include spatial contrasts,  
214 with stronger warming projected in the south, east, and mountainous regions compared to the  
215 Atlantic coast, showing a gradient of about 1°C. Seasonal contrasts are also consistent, with  
216 warming more pronounced in summer than in winter (with also a difference of 1°C). Regarding  
217 precipitation, all studies highlight the limited agreement among models on the annual average,  
218 which is linked to the influence of internal variability and to France's geographical position,  
219 which places it in a transition zone between Northern Europe, where precipitation is projected  
220 to increase, and the Mediterranean region, where it is projected to decrease (Terray and Boé,  
221 2013; Coppola et al., 2021). On a seasonal scale, however, projections are more consistent.  
222 In winter, median changes show an increase in rainfall (particularly in the northern regions)  
223 alongside a notable decrease in snowfall, especially in the low- and mid-mountain ranges. In  
224 summer, projections indicate a decrease in rainfall (particularly in the southern half). These  
225 changes in precipitation combined with increased evaporation due to warming, lead to a  
226 decrease in water resources availability and, consequently, a marked drying of the soil. In  
227 terms of intense rainfall (characterized by the annual maximum daily rainfall), the climate  
228 models show large dispersion regarding the direction of future changes (Tramblay et al.,  
229 2024), except for the long-term horizon under the RCP8.5 scenario or for high levels of  
230 warming (e.g., +4°C on average across France), where they tend to agree on an increase,  
231 particularly in the northern part of France

232  
233 **2.3 Extraction of flood events**  
234

235 From this dataset, we focused on the annual maximum of daily streamflow that has been  
236 extracted for each simulation point. For each station, a time series of 124 values, between  
237 1976 and 2099 considering hydrological years starting in September, is available and  
238 described thereafter as the Annual Maximum Flood (AMF) series, referred to as 'floods' to

Code de champ modifié

improve readability. This sampling results in a total of 10,167,256 flood events. To estimate the base flow contribution, that is the proportion of direct runoff from rainfall and/or snowmelt, we applied a base flow filter (Lyne and Hollick, 1979) to compute the direct streamflow fraction from the full daily times series, to be able to compute the base flow contribution for each flood event. To extract the characteristics of the events in terms of rainfall and snowmelt, we computed the antecedent cumulative rainfall (in mm) and snowmelt (in mm) for each AMF value, and this aggregation stops if a day has rainfall below 1 mm or if a maximum of seven days is reached. Thus, this duration estimated for each event is extracted and hereafter referred to as the time of concentration, which may differ for each event. This time window of seven days is chosen given the size of the basins considered in the present study (from 64 to 111 570 km<sup>2</sup>) and also to be consistent with previous studies using similar approaches (Ivancic and Shaw, 2015; Stein et al., 2020). For each simulated flood event, we extracted: the concentration time (in days), the antecedent soil moisture (i.e. SWI, between zero and one) one day before the ~~flood date~~concentration time, ~~the concentration time (in days)~~, the total and maximum daily rainfall (mm) during the event, the fraction of the total flood streamflow being direct streamflow (%), and the total snowmelt discharge (mm) during the event.

## 2.4 Classification of flood events

The flood event classification was determined using a decision tree that had previously been applied in Europe and the USA (Kemter et al., 2020, 2023) to estimate the importance of five different flood-generating processes in each catchment, namely long rain, short rain, snowmelt, rain and snowmelt, and soil moisture excess. This classification is based on the catchment and climatic conditions occurring during the period defined by the time of concentration before the day of the flood peak. The rationale behind the choice of this classification over others (e.g. Tarasova et al., 2019) is that it does not rely on fixed quantities for the different variables, and notably precipitation extremes, compared to other classification schemes (Stein et al., 2020; Trambly et al., 2022, 2023). Indeed, in a climate change context, these quantities could vary over time, notably those related to extreme rainfall that are inherently difficult to estimate precisely using climate simulations at these spatial and temporal resolutions (8 km, daily).

The flood event classification from Kemter et al. (2023) is adapted using the following sequential rules:

1. If the snowmelt is larger than the rainfall, the event is classified as 'snowmelt'.
2. If the snowmelt is larger than 25 % of the rainfall, the event is classified as 'rain and snowmelt'.
3. If antecedent soil moisture before the event is larger than the soil moisture threshold, the event is classified as 'soil water excess'. These events correspond to soil saturation excess.
4. If during the event, the daily maximum rainfall exceeds 75 % of the total rainfall, the event is classified as 'short rain'. These events are representing soil infiltration excess caused by a short and intense rainfall event.
5. Else, the event is classified as 'long rain'.

In the present study, we estimated the soil moisture threshold for each basin to identify flood events related to soil moisture excess. Considering specific soil moisture thresholds for each basin is undoubtedly more realistic than using a single soil saturation threshold for all basins,

as done in other studies. The approach used herein has been applied previously (Tarasova et al., 2018; Trambly et al., 2022) and is data-based, since there is no method for estimating these thresholds a priori from catchment characteristics. It relies on the extraction of all streamflow events above the 10th percentile of the whole streamflow time series between 1975 and 2005, fitting an exponential function to the soil moisture/runoff relationship, and then to identify the inflection point in the soil moisture/runoff relationship (i.e., the max slope of the curve). Here, we apply the Pruned Exact Linear Time method (Killick et al., 2012) to detect a change point (i.e., the soil moisture threshold, figure S1) in the exponential fit.

## 2.5 Trend detection

To check for trends, the entire annual time series between 1976 and 2009 was considered. The rationale for using trend detection over the whole record is that floods in many regions are strongly impacted by multidecadal variability, notably in Europe (Hodgkins et al., 2017; Lun et al., 2020; Blöschl et al., 2020), making it difficult to detect climate change response using the standard approach of comparing historical and future time windows of about 30 years. Two approaches are applied to detect these trends. First, a quantile regression (Koenker and Bassett, 1978), is applied to check the presence of trends in the 2-year flood and in the 20-year flood. The significance of these trends at the 5 % level is then assessed using a bootstrap method (Efron, 1979). In this case, the magnitude of the trend is given by the slope of the regression. We also applied a variant of the Mann-Kendall test adjusted for serial correlation in the data (Hamed and Ramachandra Rao, 1998), and the Sen slope (Sen, 1968) method to estimate the magnitude of the trends. The Mann-Kendall test was applied to the different flood event characteristics (event rainfall, antecedent soil moisture, direct runoff fraction, snowmelt contribution), as well as the relative frequencies of the different flood event types. In that latter case, 20-year sliding windows were used to compute these frequencies, using a similar approach as in Tarasova et al. (2023). Given that some processes are not relevant to all basins (e.g. snowmelt), we applied the trend detection test only if a minimum of 20 occurrences for the given flood process is present in the data sample over the 1975-2100 period. To present the trend detection results, we considered the multi-model index of agreement (MIA) (Trambly and Somot, 2018) that describes the model convergence towards an increase (1) or a decrease (-1) for a given indicator. The objective is to identify robust changes, where all model projections converge towards the same result. Finally, the trends in the flood quantiles (Q50 and Q95) and the trends in the five flood-generating processes have been clustered for the two RCP separately, using the Ward linkage method, commonly used in hierarchical clustering, together with silhouette plots to identify the optimal number of clusters (Kaufman and Rousseeuw, 1990). This, in order to define regions with similar flood changes.


Code de champ modifié

## 3- Results and discussion

### 3.1 Changes in flood characteristics



335 Flood trends are analyzed here using the quantile regression technique, which distinguishes  
 336 the change signal for AMF corresponding to 2-year (Q50) and 20-year (Q95) return periods.  
 337 The results presented in Figure 1 show the convergence of the different simulations under the  
 338 RCP4.5 and RCP8.5 scenarios with the MIA index. The figure shows a contrasting signal, with  
 339 a relatively good convergence of the models towards an increase in Q50 and Q95 in the  
 340 northern half of France and an absence of consensus for trends or even a decrease in some  
 341 areas for the southern regions, that is more marked under RCP8.5. Overall, convergence at  
 342 the country level is rather weak, with the median MIA value around 0.5 (Figure S2).  
 343 Convergence between models is slightly greater for Q50 rather than Q95, expressing a larger  
 344 uncertainty for rarest floods. In terms of magnitude (Figure 2), trends are more marked for the  
 345 rarest floods (median change +28 % with RCP4.5, +34 % with RCP8.5), than for the most  
 346 frequent floods (median change +20 % with RCP4.5, +27 % with RCP8.5). Overall, the  
 347 projected changes between Q50 and Q95 are more correlated under the RCP8.5 than under  
 348 the RCP4.5. This north/south contrast in the projections is consistent with previous findings  
 349 using observational data (Blöschl et al., 2019; Tramblay et al., 2019).  
 350



351  
 352 Figure 1: Maps of the multi-model index of agreement for the Q50 (2-year flood) and Q95  
 353 (20-year flood) for the 3727 simulation points, for the RCP4.5 and RCP8.5. The multi-model  
 354 index of agreement (MIA) is equal to 1 if all models project a significant increase, and equal  
 355 to -1 if all models project a significant decrease.  
 356  
 357






Figure 2: Comparison of the mean multi-model magnitude of significant trends on 2-year (Q50) and 20-year (Q95) floods under the RCP4.5 and RCP8.5.

This analysis of flood trends is complemented by trends in various indicators calculated during floods: maximum daily rainfall during the episode, initial soil moisture conditions, fraction of flood discharge from direct runoff, and contribution of snowmelt to flooding. The trends are shown for the RCP4.5 and RCP8.5 scenarios in Figure 3, with a general increase in maximum daily rainfall during floods, though less marked in southern France. The trends in antecedent soil moisture are more contrasted, with a good convergence of models towards an increase, more marked for RCP8.5, in the northern half of France and particularly in the eastern region. On the opposite, antecedent soil moisture decreases in the mountainous regions of the Alps and Pyrenees, and around the Mediterranean Sea. For the fraction of direct runoff (that is the total runoff minus the base flow for the day of the flood event), there is a slight trend towards an increase in RCP4.5, and a noticeable trend towards an increase in RCP8.5, in the western regions. Finally, the contribution of snowmelt to flood flows is decreasing wherever this component influences floods. Thus, it is interesting to note that in a context of increasing rainfall intensity, outside Mediterranean regions, we observe an increase in flood magnitude, that is associated either with an increase in initial soil saturation conditions, in eastern France, or with an increase in the fraction of direct runoff, in western France. It is also worth noticing that the spatial pattern of antecedent soil moisture trends seen in Figure 3 seems to mimic flood trends more closely than extreme rainfall trends, as observed in other regions of the world -(Wasko and Nathan, 2019).




Figure 3: Multi-model index of agreement of the trends in maximum rainfall during floods, antecedent soil moisture conditions, direct streamflow fraction, and snowmelt fraction, under the RCP4.5 and RCP8.5 with ADAMONT

### 3.2 Classification of flood-generating processes

The floods extracted from the various simulations were classified according to the different categories; snowmelt, rain and snowmelt, soil water excess, short rain, and long rain. The result of this classification from the multimodel ensemble during the historical period 1975-2005 is presented in Figure 4, which shows the relative contribution of these different flood types for each station. Events linked to soil saturation are predominant, accounting for more than 50 % of floods in France. Long precipitation events are responsible for almost 25 % of floods and are mainly located in the north-west and south of France. It should be noted that most of these events also correspond to soil saturation processes, although in this case, the soil saturation occurs not before but during these events. Indeed, Figure S3 shows that during these long rain events, the largest floods are only associated with SWI values above 0.6, and that the modal value of SWI maxima is close to 0.75 and 0.8, corresponding to the value of the soil saturation threshold most frequently found in French basins (Figure S1). And, on average for all basins, the threshold that splits between long rain and soil water excess floods are exceeded 69% of the time during long rain floods. Snowmelt-related events, caused by snowmelt or a mixture of rain and snowmelt, account for around 11 % of the total number of floods, mainly for mountainous areas and central-eastern regions. Lastly, short rain events account for less than 10 % of the events, with a spatial distribution very similar to that of long rain. In terms of magnitude, when all the standardized floods for the different processes are grouped, short rain events show the highest magnitude, followed by soil water excess and long rain events, which have similar magnitudes (Figure 5a). This result is consistent with Tarasova et al. (2023) who found a more pronounced tail for the distribution of floods caused by intense rainfall on dry soils.




Figure 4: Relative proportion of the different flood-generating processes for each basin during the historical period 1975-2005. The average contribution of each process to the total number of floods is given in the titles of the sub-plots in percentages.

As these results are obtained with an ensemble of bias-corrected climate projections, the results have also been extracted for each GCM/RCM couple independently to check the consistency between results. Over the historical period, the classifications obtained with the different climate models show a very high degree of consistency in the proportions of the different classes between models (Figure S4). This result was quite expected given that the RCM simulations are bias-corrected using the same reference. The results of this classification are also very consistent with previous studies in Europe applying similar classification schemes but with different datasets and a different methodology. For instance, Berghuijs et al. (2019) have found that 49 % of floods are driven by soil water excess in Western Europe, and about 22 % of floods are driven by maximum annual rainfall. Tarasova et al. (2023) obtained similar results in their pan-European study using slightly different classes, with 11 % of rain on dry soils, 67 % of rain on wet soils, and 21 % of rain-on-snow events in Atlantic regions, and for the Mediterranean region, with 25 % of rain on dry soils, 59 % of rain on wet soils, and 15 % of rain-on-snow events. In both regions, the proportion of snowmelt-only driven floods was equivalent to 1 % only.




Figure 5: Distribution of floods all basins together, for the different types of floods during the historical period 1975-2005, and for 2070-2100 under RCP4.5 and RCP8.5. The annual flood ( $AMF$ ) for each station have been standardized by the mean annual flood ( $\overline{AMF}$ )

a mis en forme : Police :Italique

### 3.3 Changes in flood-generating processes

The direction of changes in the contribution of the various flood-generating processes in RCP4.5 and RCP8.5 is shown in Figure 6. Given the low frequency of snowmelt-only events, it is difficult to draw robust conclusions about changes in these events. However, there is a general decrease in snowmelt or rain-and-snowmelt-induced floods in both scenarios. This applies both to the mountain ranges of the Alps and the Pyrenees and to regions located from the center of France to the northeast. There is also an increase in the contribution of events linked to soil saturation, especially in the northeast, that is more widespread spatially for RCP4.5 than for RCP8.5. Floods induced by short rains and long rains are increasing in areas where soil water excess events are decreasing, most importantly in the southern regions and the Alps. It should be noted that the change in rain and snowmelt events shows a very similar spatial distribution compared to the trends for events linked to soil water excess. Indeed, the spatial correlations between changes in soil water excess events on the one hand, and rain and snowmelt events on the other hand, are significant (-0.6 for RCP4.5 and -0.67 for RCP8.5). This means that where the proportion of events linked to the combination of rain and snowmelt is reducing, these events tend to be replaced by floods associated with saturated soils. Similarly, there are significant correlations between changes in the proportion of soil water excess events and the proportion of short rain or long rain events (correlations of -0.31 between changes in soil water excess and changes in short rain, -0.4 with changes in long rain under RCP4.5, and respectively -0.54 and -0.45 under RCP8.5 - yet the magnitude of these correlations remains small). We can therefore see that in regions where the proportion of floods related to saturated soils is decreasing, the proportion of floods associated with short rain or long rain is increasing, particularly in southern regions. This result shows that these shifts, previously observed in historical records (Jiang et al., 2022; Tarasova et al., 2023; Trambly et al., 2023), are likely to amplify under the two emission scenarios considered herein.




Figure 6: Multi-model index of agreement for the trends in the contributions of the different flood-generating processes, under the RCP4.5 and RCP8.5

The trends in the flood quantiles (Q50 and Q95) and the trends in the five flood generating processes have been clustered for the two RCP separately, using the Ward linkage method, commonly used in hierarchical clustering, together with silhouette plots to identify the optimal number of clusters (Kaufman and Rousseeuw, 1990). The classification of the trends in flood quantiles and the five flood generating processes is classification results in four spatial clusters as shown in Figure 7. The first cluster (Mountain) Cluster-1 includes mountainous basins in the Alps and Pyrenees, where Q50 and Q95 show mostly declining trends, or an absence of trends, linked to a sharp decrease in snowmelt-related events and an increase in events caused by short rain and long rain. The second cluster (Mediterranean) Cluster-2 includes basins located in southern regions, where there is no trend or a decrease in floods, associated with a drop in events linked to soil saturation and an increase in events caused by short rain and long rain. The third and fourth clusters (Atlantic and Continental) Cluster-3 and -4 gather the stations in the northern half of France, where flooding trends are on the rise, but with a sharp decrease in snowmelt-related flooding in the east (Continental cluster Cluster-4), while in the west (Atlantic cluster Cluster-3) there is a slight increase of the proportion-frequency of intense rainfall (short rain) on dry soils, associated with an increase in direct runoff, as already shown in the previous sections. The spatial organization of the different clusters is very similar under the two scenarios RCP4.5 and RCP8.5, with one notable difference concerning cluster-2, the corresponding to Mediterranean cluster events in the South of France, which has a much more marked northward extension under the RCP8.5 scenario. Overall, it should be noted that this spatial distribution is strongly reminiscent of the different climatic zones observed in France (Strohmenger et al., 2024), with the western and coastal regions under a temperate oceanic climate, the center and east of France with a more continental climate, the southern part corresponding to areas with a Mediterranean climate, and finally the mountain regions of the Alps and the Pyrenees.

Code de champ modifié

Code de champ modifié




Figure 7: Regional clusters of the trends in flood magnitudes and flood-generating processes

### 3.4 Regional drivers of change for flood hazard in future projections

The magnitude of floods will change differently, depending on their generating processes. When aggregated over France, both the frequency and the magnitude of snowmelt-related events is projected to decrease (Figures 5b and 5c, and figure S5 and S6 in the supplementary materials), while soil water excess events increase slightly. It is mainly the magnitude of the rarest and most intense short rain and long rain events that are increasing. Marked differences between future and historical distributions are mainly projected for events associated with non-exceedance probabilities greater than 0.95 (i.e. floods corresponding to a 20-year return period and beyond). However, this overall assessment at the country level is difficult to assess and hides regional differences depending on the flood-generating processes. If we look by region and by flood process (supplementary figures S5 and S6), we observe that this increase in the magnitude of the rarest floods (i.e. the distribution tails), only affects the northern regions of France, the Atlantic and continental clusters, and events linked to short rains and long rains, which are flood events linked to rainfall on dry soils, and also the events related to soil water excess. Under the RCP4.5, the flood distributions in the different regions for the different flood generating process are similar between 1975-2005 and 2070-2100 (figure S5), except for some of the most extreme flood events as mentioned above. For the RCP8.5, there are more marked changes, notably with a shift of the flood distribution towards lower values for the cluster 4 corresponding to the -{mountainous regions}, and to a lesser extent for the cluster 2 (Mediterranean regions). On the opposite, there are distribution shifts towards increased flood magnitudes for most processes related to rainfall and soil water excess in the Atlantic and Continental clusters, consistent with the overall increase in flood magnitude in these regions shown in Figure 1, 3 and 4.

Another important question that arises when analyzing the impact of climate change on floods is to understand which factors have the greatest influence on flood trends in different climate projections. Given that 11 climate projections are available here for RCP4.5 and RCP8.5, we can calculate for each model the correlations between flood trends and the trends in the three factors most commonly recognized as flood triggers (Wasko and Nathan, 2019, Zhang et al., 2022) at the event scale (the antecedent soil moisture conditions, the snowmelt contribution,



and the maximal daily rainfall during the event) and the trends for floods with 2-year (Q50) and 20-year (Q95) return periods. The results of this correlation analysis are presented in Figure 8 for each of the four regions identified by the clustering of Figure 7. For the first cluster, i.e. the mountain areas, there are strong positive correlations between snowmelt trends and Q95 flood trends, but much more contrasting correlations with Q50. For the other clusters 2 to 4, the patterns are quite similar. A striking result is that the correlations between trends in floods and trends in SWI are at least equivalent, if not superior, to the correlations with trends in intense rainfall for the different climate projections. This result is not region-specific, and it should also be noted that correlations with SWI are almost systematically stronger for frequent floods (2-year floods) than for 20-year floods, as already found in past observations by Wasko et al., 2021. To a lesser extent, correlations are also weaker between trends in rainfall intensity and the flood trends for the rarest 20-year floods, than for trends in 2-year floods the most frequent floods. This result should however be interpreted with caution given the uncertainties inherent in estimating intense rainfall in climate models. These results imply that the trends projected for the various processes can, to a large extent, explain the trends in flooding in France, and that the evolution of soil saturation plays a role equivalent to that of the evolution of intense rainfall in the various regions to explain the trends in flood hazard in the different climate projections.




Figure 8: Correlations between the trends in event antecedent soil moisture (SWI), snowmelt contribution (Snow) and maximum daily rainfall during the event (MaxRain), and the trends in 2-year (Q50) and 20-year (Q95) floods, for each of the clusters presented in figure 8. The boxplots represent the ranges of the correlation coefficients obtained for each climate projection under the RCP4.5 (top row) and RCP8.5 (bottom row).



#### 4. Discussion

These projections are in line with the expected climate change in France (Strohmenger et al., 2024): a radical change in mountainous regions with a transition to a more temperate climate, an increase in intense rainfall all over the country as in other regions of Europe (Coppola et al., 2021), and an overall North/South contrast for future rainfall changes; with an increasing trends in the northern regions, leading to higher soil moisture content, and conversely a decline in southern regions, yet with a little agreement between models (Marson et al., 2024). It is important to note that this study is one of the few (notably with Zhang et al., 2022, at the global scale) to analyze the evolution of different flood processes in future climates, rather than solely in the past as most published studies do (Tramblay et al., 2022; Tarasova et al., 2023; Kemter et al., 2023). It is interesting to note is that the conclusions of this article largely confirm that the trends and inter-variable relationships observed in the past appear to be consistent with what is happening in future simulations.

As shown in Ivancic and Shaw (2015), Tarasova et al. (2023) or Wasko and Nathan (2019), changes in extreme precipitation alone are not sufficient to explain changes in floods, and it is clearly shown in the present work that a combination of changes in flood types with changes in the frequency and magnitude of the different flood drivers best explain the temporal evolution of flood hazard. In particular, the importance of antecedent soil moisture in modulating flood hazard is expected to remain in future scenarios (as shown in Figure 8), while several observations-based studies have already shown the important role of soil moisture conditions in the genesis of floods (Penna et al., 2011; Tramblay et al., 2019, 2023), a role that is more significant for lower-magnitude floods than for more extreme floods (Wasko and Nathan, 2019; Wasko et al., 2021; Brunner et al., 2021). Given the future projections towards an increase of soil moisture droughts in large parts of Europe (Samaniego et al., 2018; Grillakis, 2019), it is expected that changes in soil moisture are key to understand changes in flood hazard in future scenarios.

For some regions, the future increase of antecedent conditions as in the central-eastern regions of France is combined with a decrease of snowmelt influence, and this was also projected for the greater Rhine basin (Rottler et al., 2021). Overall, the magnitude of snowmelt-related events is declining globally (Zhang et al., 2022) and notably in the Alps (Sikorska-Senoner and Seibert, 2020), and the present study show that the same trend is expected to continue for the future. On the contrary, given the global increasing trends in rainfall extremes, notably at short durations (Fowler et al., 2021), our projections are consistent with previous studies towards an increase of the magnitude of the rarest floods due to intense and short rain episodes in several regions and notably in the Mediterranean (Zhang et al., 2022; Tarasova et al., 2023; Poncet et al., 2025). Yet, it is worth to remind that the uncertainties on flood projections remains high notably due to the spread in climate model simulations (Evin et al., 2025).

#### 4.5. Conclusions

a mis en forme : Police :Gras

a mis en forme : Justifié

621  
622 Flood trends show a contrasting signal in the different regions, with a fairly good consistency  
623 towards an increase in floods in northern France and an absence of signal in southern regions.  
624 In line with the uncertainty in future precipitation changes, it should be emphasized that multi-  
625 model agreement remains fairly weak in most basins for increasing or decreasing trends in  
626 flood hazard. In terms of magnitude, trends are more marked for rare floods (20-year return  
627 period) than for frequent floods (2-year return period). Trends in initial moisture conditions  
628 show a spatial distribution quite similar to that of floods, with an increase mainly in northern  
629 regions and a decrease in antecedent soil moisture in the south. These changes tend to mimic  
630 the seasonal precipitation changes, with an increase in winter precipitation in the north and a  
631 decrease in summer precipitation in the south. The magnitude of rainfall events increases  
632 everywhere, most markedly in the northern half of France, while the contribution of snowmelt  
633 decreases everywhere in the mountain ranges, consistently with the projected decrease in  
634 snowfall as indicated by climate projections. A classification of floods by process reveals that  
635 floods linked to soil saturation account for more than half of all floods in France, and that this  
636 proportion is increasing in the north-east of the country while decreasing in the south. The  
637 relative changes in the importance of the different flood generating processes ~~is~~ are not  
638 spatially homogeneous and varies by region. The analysis highlights that flooding trends can  
639 be decomposed into several distinct signals linked to different flooding processes, and that  
640 the multi-model uncertainty concerning these trends varies according to the process  
641 considered. Trends in soil saturation play as important a role as trends in intense rainfall in  
642 different regions to explain changes in flood hazard in the different climate simulations.

643 Overall, in regions where the proportion of floods linked to antecedent saturated soils (i.e Soil  
644 Water Excess floods) is decreasing, the proportion of floods linked to short or long rains is  
645 increasing, particularly in the south. Both the frequency and magnitude of floods linked to  
646 snowmelt processes are declining in areas where these processes operate, namely  
647 mountainous areas and the north-eastern regions. ~~On the contrary~~In contrast, the flood  
648 magnitudes associated with rainfall on dry soils (short rain and long rain floods) tend to  
649 increase, in line with the increase of maximum event rainfall in the different scenarios. The  
650 trends in the RCP8.5 scenario are both more pronounced and allow better discrimination  
651 between regions. The changes seem fairly consistent with the distribution of French climatic  
652 zones: floods tend to increase in the continental climate zone, with both an increase in rainfall  
653 intensity and soil water content, leading to more extreme floods and a increase in the number  
654 of flood events linked to soil saturation. In the temperate oceanic zone, flooding is also on the  
655 rise, but this time in connection with the increased intensity of ~~intense~~ rainfall leading to higher  
656 direct runoff but no drastic changes in the flood event types. In the Mediterranean regions,  
657 there is no marked trend in flooding, with declining soil moisture and more uncertain trends in  
658 rainfall intensity. Lastly, in mountainous regions, there are drastic changes in flood-generating  
659 conditions, with a marked drop in snowmelt and an absence of trends, or even a decline, in  
660 flooding.

661 In terms of perspectives for this study, there are a number of points to note. The first is that  
662 the analysis carried out here is based on a daily time step, which is unsuitable for the analysis  
663 of flash floods, which occur, for example, in Mediterranean regions with high hourly rainfall  
664 intensities. For small basins, where only until recently high-resolution convection-permitting  
665 models (Lucas-Picher et al., 2021) are able to tackle this problem by improving the modeling  
666 of small-scale convective rainfall events (Kay, 2022; Poncet et al., 2024), there is not yet a

large ensemble of model runs available to assess the uncertainties in the projections. It should be stressed out that the results obtained herein are unlikely to be transposable to flash floods, and large convection-permitting model ensembles would be required to replicate such an analysis for flash floods. Another important aspect is that this study is based on natural hydrology processes in relation to climate change. However, changes in land use (Rogger et al., 2017; Yang et al., 2021) or the construction of infrastructures, such as dams (Zahar et al., 2008; Grill et al., 2019; Blöschl, 2022b), can greatly modulate the risk of flooding. However, this type of non-climate-related change is very difficult, if not impossible, to take into account in such a large scale and multi-basin study. Nevertheless, at more local scales, and in close interaction with land-use planning and water management stakeholders, such multi-criteria analyses could be carried out to better distinguish between climatic and non-climatic influences on the evolution of flood risk.

715  
716 **Data availability**  
717  
718 All the data of this study is available freely in open access. The bias-corrected climate model  
719 simulations are available at: <https://www.drias-climat.fr/>. The hydrological model simulations  
720 are available at: <https://www.drias-eau.fr/>. The discharge measurements over France are  
721 available at: <https://www.hydro.eaufrance.fr/>. The SAFRAN database is available at:  
722 <https://meteo.data.gouv.fr/datasets/donnees-changement-climatique-sim-quotidienne/>

Code de champ modifié

Code de champ modifié

Code de champ modifié

Code de champ modifié

723  
724 All the technical reports of the Explore2 project are available in the repository:  
725 <https://entrepot.recherche.data.gouv.fr/dataverse/explore2>

Code de champ modifié

726  
727 **Code availability**

728  
729 The bias-correction method for climate simulations is available at:  
730 <https://github.com/yrobink/SBCK-python>

Code de champ modifié

731  
732 The hydrological model codes are available at: <https://inrae.github.io/airGRiwrn/> and  
733 <https://hydrogr.github.io/airGR/>

Code de champ modifié

Code de champ modifié

734  
735 **Author contributions**

736  
737 YT designed the study and performed the experiments. GT, LS, ran the GRSD model  
738 simulations, LH managed the database, LC generated the climate scenarios, GE and ES  
739 contributed to the methodology and YT wrote the paper with contributions from all the  
740 authors.

741  
742 **Competing interest declaration**

743  
744 The authors declare that they have no conflict of interest.

745  
746 **Acknowledgements**

747  
748 This research was financed by the Explore2 project with support from the French  
749 Biodiversity Agency (OFB) and the French Ministry of Ecological Transition (MTECT). The  
750 authors would like to thank all the members of the Explore2 consortium who contributed to  
751 the creation of this unique dataset for France.

752  
753 **Financial support**

754  
755 This research has been supported by the Office Français de la Biodiversité (Explore2  
756 project), the Ministère de la Transition écologique et Solidaire (Explore2 project)

757  
758  
759  
760

## References

- Abbenante, S. E., Althoff, I., and Valdes-Abellan, J.: Comparative analysis of lumped and semi-distributed hydrological models in humid Mediterranean environments, *Hydrological Sciences Journal*, 69, 2481–2496, <https://doi.org/10.1080/02626667.2024.2413422>, 2024.
- Alfieri, L., Burek, P., Feyen, L., and Forzieri, G.: Global warming increases the frequency of river floods in Europe, *Hydrol. Earth Syst. Sci.*, 19, 2247–2260, <https://doi.org/10.5194/hess-19-2247-2015>, 2015.
- Bathelemy, R., Brigode, P., Andréassian, V., Perrin, C., Moron, V., Gauchere, C., Tric, E., and Boisson, D.: Simbi: historical hydro-meteorological time series and signatures for 24 catchments in Haiti, *Earth Syst. Sci. Data*, 16, 2073–2098, <https://doi.org/10.5194/essd-16-2073-2024>, 2024.
- Berghuijs, W. R., Woods, R. A., Hutton, C. J., and Sivapalan, M.: Dominant flood generating mechanisms across the United States: Flood Mechanisms Across the U.S., *Geophys. Res. Lett.*, 43, 4382–4390, <https://doi.org/10.1002/2016GL068070>, 2016.
- Berghuijs, W. R., Harrigan, S., Molnar, P., Slater, L. J., and Kirchner, J. W.: The Relative Importance of Different Flood-Generating Mechanisms Across Europe, *Water Resour. Res.*, 2019WR024841, <https://doi.org/10.1029/2019WR024841>, 2019.
- Blöschl, G.: Flood generation: process patterns from the raindrop to the ocean, *Hydrol. Earth Syst. Sci.*, 26, 2469–2480, <https://doi.org/10.5194/hess-26-2469-2022>, 2022a.
- Blöschl, G.: Three hypotheses on changing river flood hazards, *Hydrol. Earth Syst. Sci.*, 26, 5015–5033, <https://doi.org/10.5194/hess-26-5015-2022>, 2022b.
- Blöschl, G., Hall, J., Viglione, A., Perdigão, R. A. P., Parajka, J., Merz, B., Lun, D., Arheimer, B., Aronica, G. T., Bilbashi, A., Boháč, M., Bonacci, O., Borga, M., Čanjevac, I., Castellarin, A., Chirico, G. B., Claps, P., Frolova, N., Ganora, D., Gorbachova, L., Gül, A., Hannaford, J., Harrigan, S., Kireeva, M., Kiss, A., Kjeldsen, T. R., Kohnová, S., Koskela, J. J., Ledvinka, O., Macdonald, N., Mavrova-Guirguinova, M., Mediero, L., Merz, R., Molnar, P., Montanari, A., Murphy, C., Osuch, M., Ovcharuk, V., Radevski, I., Salinas, J. L., Sauquet, E., Šraj, M., Szolgay, J., Volpi, E., Wilson, D., Zaiml, K., and Živković, N.: Changing climate both increases and decreases European river floods, *Nature*, 573, 108–111, <https://doi.org/10.1038/s41586-019-1495-6>, 2019.
- Blöschl, G., Kiss, A., Viglione, A., Barriendos, M., Böhm, O., Brázdil, R., Coeur, D., Demarée, G., Llasat, M. C., Macdonald, N., Retsö, D., Roald, L., Schmocker-Fackel, P., Amorim, I., Bělínová, M., Benito, G., Bertolin, C., Camuffo, D., Cornel, D., Doktor, R., Elleder, L., Enzi, S., Garcia, J. C., Glaser, R., Hall, J., Haslinger, K., Hofstätter, M., Komma, J., Limanówka, D., Lun, D., Panin, A., Parajka, J., Petrić, H., Rodrigo, F. S., Rohr, C., Schönbein, J., Schulte, L., Silva, L. P., Toonen, W. H. J., Valent, P., Waser, J., and Wetter, O.: Current European flood-rich period exceptional compared with past 500 years, *Nature*, 583, 560–566, <https://doi.org/10.1038/s41586-020-2478-3>, 2020.
- Brigode, P., Oudin, L., and Perrin, C.: Hydrological model parameter instability: A source of additional uncertainty in estimating the hydrological impacts of climate change?, *Journal of Hydrology*, 476, 410–425, <https://doi.org/10.1016/j.jhydrol.2012.11.012>, 2013.
- Brunner, M. I., Swain, D. L., Wood, R. R., Willkofer, F., Done, J. M., Gilleland, E., and Ludwig, R.: An extremeness threshold determines the regional response of floods to

806 changes in rainfall extremes, *Commun Earth Environ*, 2, 173,  
807 <https://doi.org/10.1038/s43247-021-00248-x>, 2021.

808 Chauveau, M., Chazot, S., Perrin, C., Bourgin, P.-Y., Sauquet, E., Vidal, J.-P., Rouchy, N.,  
809 Martin, E., David, J., Norotte, T., Maugis, P., and De Lacaze, X.: Quels impacts des  
810 changements climatiques sur les eaux de surface en France à l'horizon 2070 ?, *La Houille*  
811 *Blanche*, 99, 5–15, <https://doi.org/10.1051/lhb/2013027>, 2013.

812 [Coppola, E., Nogherotto, R., Ciarlo', J. M., Giorgi, F., Van Meijgaard, E., Kadygrov, N., Iles,](#)  
813 [C., Corre, L., Sandstad, M., Somot, S., Nabat, P., Vautard, R., Levavasseur, G.,](#)  
814 [Schwingshackl, C., Sillmann, J., Kjellström, E., Nikulin, G., Aalbers, E., Lenderink, G.,](#)  
815 [Christensen, O. B., Boberg, F., Sørland, S. L., Demory, M., Bülow, K., Teichmann, C.,](#)  
816 [Warrach-Sagi, K., and Wulfmeyer, V.: Assessment of the European Climate Projections as](#)  
817 [Simulated by the Large EURO-CORDEX Regional and Global Climate Model Ensemble,](#)  
818 [JGR Atmospheres](#), 126, e2019JD032356, <https://doi.org/10.1029/2019JD032356>, 2021.

819 Clark, M. P., Wilby, R. L., Gutmann, E. D., Vano, J. A., Gangopadhyay, S., Wood, A. W.,  
820 Fowler, H. J., Prudhomme, C., Arnold, J. R., and Brekke, L. D.: Characterizing Uncertainty of  
821 the Hydrologic Impacts of Climate Change, *Curr Clim Change Rep*, 2, 55–64,  
822 <https://doi.org/10.1007/s40641-016-0034-x>, 2016.

823 Coppola, E., Nogherotto, R., Ciarlo', J. M., Giorgi, F., Van Meijgaard, E., Kadygrov, N., Iles,  
824 C., Corre, L., Sandstad, M., Somot, S., Nabat, P., Vautard, R., Levavasseur, G.,  
825 Schwingshackl, C., Sillmann, J., Kjellström, E., Nikulin, G., Aalbers, E., Lenderink, G.,  
826 Christensen, O. B., Boberg, F., Sørland, S. L., Demory, M., Bülow, K., Teichmann, C.,  
827 Warrach-Sagi, K., and Wulfmeyer, V.: Assessment of the European Climate Projections as  
828 Simulated by the Large EURO-CORDEX Regional and Global Climate Model Ensemble,  
829 *JGR Atmospheres*, 126, e2019JD032356, <https://doi.org/10.1029/2019JD032356>, 2021.

830 Cornelissen, T., Diekkrüger, B., and Giertz, S.: A comparison of hydrological models for  
831 assessing the impact of land use and climate change on discharge in a tropical catchment,  
832 *Journal of Hydrology*, 498, 221–236, <https://doi.org/10.1016/j.jhydrol.2013.06.016>, 2013.

833 Coron, L., Thirel, G., Delaigue, O., Perrin, C., and Andréassian, V.: The suite of lumped GR  
834 hydrological models in an R package, *Environmental Modelling & Software*, 94, 166–171,  
835 <https://doi.org/10.1016/j.envsoft.2017.05.002>, 2017.

836 Corre, L., Ribes, A., Bernus, S., Drouin, A., Morin, S., and Soubeyroux, J.-M.: Using regional  
837 warming levels to describe future climate change for services and adaptation: Application to  
838 the French reference trajectory for adaptation, *Climate Services*, 38, 100553,  
839 <https://doi.org/10.1016/j.cliser.2025.100553>, 2025.

840 Dakhlaoui, H., Ruelland, D., Trambay, Y., and Bargaoui, Z.: Evaluating the robustness of  
841 conceptual rainfall-runoff models under climate variability in northern Tunisia, *Journal of*  
842 *Hydrology*, 550, 201–217, <https://doi.org/10.1016/j.jhydrol.2017.04.032>, 2017.

843 De Lavenne, A., Andréassian, V., Thirel, G., Ramos, M. -H., and Perrin, C.: A Regularization  
844 Approach to Improve the Sequential Calibration of a Semidistributed Hydrological Model,  
845 *Water Resources Research*, 55, 8821–8839, <https://doi.org/10.1029/2018WR024266>, 2019.

846 Di Sante, F., Coppola, E., and Giorgi, F.: Projections of river floods in Europe using EURO-  
847 CORDEX, CMIP5 and CMIP6 simulations, *Int J Climatol*, 41, 3203–3221,  
848 <https://doi.org/10.1002/joc.7014>, 2021.

849 Do, H. X., Zhao, F., Westra, S., Leonard, M., Gudmundsson, L., Boulange, J. E. S., Chang,

850 J., Ciais, P., Gerten, D., Gosling, S. N., Müller Schmied, H., Stacke, T., Telteu, C.-E., and  
851 Wada, Y.: Historical and future changes in global flood magnitude – evidence from a model–  
852 observation investigation, *Hydrol. Earth Syst. Sci.*, 24, 1543–1564,  
853 <https://doi.org/10.5194/hess-24-1543-2020>, 2020.

854 Dorchie, D.: hubeau: an R package for the Hub'Eau APIs,  
855 <https://doi.org/10.57745/XKN6NC>, 2022.

856 Efron, B.: Bootstrap Methods: Another Look at the Jackknife, *Ann. Statist.*, 7,  
857 <https://doi.org/10.1214/aos/1176344552>, 1979.

858 [Evin, G., Hingray, B., Thirel, G., Ducharme, A., Strohmenger, L., Corre, L., Trambly, Y.,](#)  
859 [Vidal, J.-P., Bonneau, J., Colleoni, F., Gailhard, J., Habets, F., Hendrickx, F., Hérault, L.,](#)  
860 [Huang, P., Le Lay, M., Magand, C., Marson, P., Monteil, C., Munier, S., Reverdy, A.,](#)  
861 [Soubeyrou, J.-M., Robin, Y., Vergnes, J.-P., Vrac, M., and Sauquet, E.: Uncertainty](#)  
862 [sources in a large ensemble of hydrological projections: Regional Climate Models and](#)  
863 [Internal Variability matter, \*EGUsphere\* \[preprint\], \[https://doi.org/10.5194/egusphere-2025-\]\(https://doi.org/10.5194/egusphere-2025-2727\)](#)  
864 [2727](#), 2025.

865 Fabre, J., Ruelland, D., Dezetter, A., and Grouillet, B.: Sustainability of water uses in  
866 managed hydrosystems: human-and climate-induced changes for the mid-21st century,  
867 *Hydrol. Earth Syst. Sci.*, 20, 3129–3147, <https://doi.org/10.5194/hess-20-3129-2016>, 2016.

868 [Froidevaux, P., Schwanbeck, J., Weingartner, R., Chevalier, C., and Martius, O.: Flood](#)  
869 [triggering in Switzerland: the role of daily to monthly preceding precipitation, \*Hydrol. Earth\*](#)  
870 [Syst. Sci., 19, 3903–3924, <https://doi.org/10.5194/hess-19-3903-2015>.](#)

871 [Fowler, H. J., Ali, H., Allan, R. P., Ban, N., Barbero, R., Berg, P., Blenkinsop, S., Cabi, N. S.,](#)  
872 [Chan, S., Dale, M., Dunn, R. J. H., Ekström, M., Evans, J. P., Fosser, G., Golding, B.,](#)  
873 [Guerreiro, S. B., Hegerl, G. C., Kahraman, A., Kendon, E. J., Lenderink, G., Lewis, E., Li, X.,](#)  
874 [O’Gorman, P. A., Orr, H. G., Peat, K. L., Prein, A. F., Pritchard, D., Schär, C., Sharma, A.,](#)  
875 [Stott, P. A., Villalobos-Herrera, R., Villarini, G., Wasko, C., Wehner, M. F., Westra, S., and](#)  
876 [Whitford, A.: Towards advancing scientific knowledge of climate change impacts on short-](#)  
877 [duration rainfall extremes, \*Phil. Trans. R. Soc. A.\*, 379, 20190542,](#)  
878 <https://doi.org/10.1098/rsta.2019.0542>, 2021.

879

880 Givati, A., Thirel, G., Rosenfeld, D., and Paz, D.: Climate change impacts on streamflow at  
881 the upper Jordan River based on an ensemble of regional climate models, *Journal of*  
882 *Hydrology: Regional Studies*, 21, 92–109, <https://doi.org/10.1016/j.ejrh.2018.12.004>, 2019.

883 Grill, G., Lehner, B., Thieme, M., Geenen, B., Tickner, D., Antonelli, F., Babu, S., Borrelli, P.,  
884 Cheng, L., Crochetiere, H., Ehalt Macedo, H., Filgueiras, R., Goichot, M., Higgins, J., Hogan,  
885 Z., Lip, B., McClain, M. E., Meng, J., Mulligan, M., Nilsson, C., Olden, J. D., Opperman, J. J.,  
886 Petry, P., Reidy Liermann, C., Sáenz, L., Salinas-Rodríguez, S., Schelle, P., Schmitt, R. J.  
887 P., Snider, J., Tan, F., Tockner, K., Valdujo, P. H., van Soesbergen, A., and Zarfl, C.:  
888 Mapping the world’s free-flowing rivers, *Nature*, 569, 215–221,  
889 <https://doi.org/10.1038/s41586-019-1111-9>, 2019.

890 [Grillakis, M. G.: Increase in severe and extreme soil moisture droughts for Europe under](#)  
891 [climate change, \*Science of The Total Environment\*, 660, 1245–1255,](#)  
892 <https://doi.org/10.1016/j.scitotenv.2019.01.001>, 2019.

893 Gupta, H. V., Kling, H., Yilmaz, K. K., and Martinez, G. F.: Decomposition of the mean



894 squared error and NSE performance criteria: Implications for improving hydrological  
895 modelling, *Journal of Hydrology*, 377, 80–91, <https://doi.org/10.1016/j.jhydrol.2009.08.003>,  
896 2009.

897 Hamed, K. H. and Ramachandra Rao, A.: A modified Mann-Kendall trend test for  
898 autocorrelated data, *Journal of Hydrology*, 204, 182–196, [https://doi.org/10.1016/S0022-](https://doi.org/10.1016/S0022-1694(97)00125-X)  
899 1694(97)00125-X, 1998.

900 Héraud L., Sauquet E. : "Fiches de diagnostic régional des modèles hydrologiques de  
901 surface du projet Explore2", <https://doi.org/10.57745/LNTOKL>, Recherche Data Gouv, 2024.

902 Ho, M., Nathan, R., Wasko, C., Vogel, E., and Sharma, A.: Projecting changes in flood event  
903 runoff coefficients under climate change, *Journal of Hydrology*, 615, 128689,  
904 <https://doi.org/10.1016/j.jhydrol.2022.128689>, 2022.

905 Hodgkins, G. A., Whitfield, P. H., Burn, D. H., Hannaford, J., Renard, B., Stahl, K., Fleig, A.  
906 K., Madsen, H., Mediero, L., Korhonen, J., Murphy, C., and Wilson, D.: Climate-driven  
907 variability in the occurrence of major floods across North America and Europe, *Journal of*  
908 *Hydrology*, 552, 704–717, <https://doi.org/10.1016/j.jhydrol.2017.07.027>, 2017.

909 Hrachowitz, M. and Clark, M. P.: HESS Opinions: The complementary merits of competing  
910 modelling philosophies in hydrology, *Hydrol. Earth Syst. Sci.*, 21, 3953–3973,  
911 <https://doi.org/10.5194/hess-21-3953-2017>, 2017.

912 Intergovernmental Panel On Climate Change (Ippc): Climate Change 2021 – The Physical  
913 Science Basis: Working Group I Contribution to the Sixth Assessment Report of the  
914 Intergovernmental Panel on Climate Change, 1st ed., Cambridge University Press,  
915 <https://doi.org/10.1017/9781009157896>, 2023.

916 Ivancic, T. J. and Shaw, S. B.: Examining why trends in very heavy precipitation should not  
917 be mistaken for trends in very high river discharge, *Climatic Change*, 133, 681–693,  
918 <https://doi.org/10.1007/s10584-015-1476-1>, 2015.

919 Jacob, D., Kotova, L., Teichmann, C., Sobolowski, S. P., Vautard, R., Donnelly, C.,  
920 Koutroulis, A. G., Grillakis, M. G., Tsanis, I. K., Damm, A., Sakalli, A., and van Vliet, M. T. H.:  
921 Climate Impacts in Europe Under +1.5°C Global Warming, *Earth's Future*, 6, 264–285,  
922 <https://doi.org/10.1002/2017EF000710>, 2018.

923 Jahanshahi, A., Booij, M. J., and Patil, S. D.: Dependence of rainfall–runoff model  
924 performance on calibration conditions under changing climatic conditions, *Hydrological*  
925 *Sciences Journal*, 1–16, <https://doi.org/10.1080/02626667.2024.2441325>, 2025.

926 Jiang, S., Bevacqua, E., and Zscheischler, J.: River flooding mechanisms and their changes  
927 in Europe revealed by explainable machine learning, *Hydrol. Earth Syst. Sci.*, 26, 6339–  
928 6359, <https://doi.org/10.5194/hess-26-6339-2022>, 2022.

929 Kay, A.: Differences in hydrological impacts using regional climate model and nested  
930 convection-permitting model data, *Climatic Change*, 173, 11, [https://doi.org/10.1007/s10584-](https://doi.org/10.1007/s10584-022-03405-z)  
931 022-03405-z, 2022.

932 Kemter, M., Merz, B., Marwan, N., Vorogushyn, S., and Blöschl, G.: Joint Trends in Flood  
933 Magnitudes and Spatial Extents Across Europe, *Geophys. Res. Lett.*, 47,  
934 <https://doi.org/10.1029/2020GL087464>, 2020.

935 Kemter, M., Marwan, N., Villarini, G., and Merz, B.: Controls on Flood Trends Across the

a mis en forme : Français (France)

a mis en forme : Français (France)

a mis en forme : Français (France)

a mis en forme : Français (France)

936 United States, Water Resources Research, 59, e2021WR031673,  
937 <https://doi.org/10.1029/2021WR031673>, 2023.

938 Killick, R., Fearnhead, P., and Eckley, I. A.: Optimal Detection of Change-points With a Linear  
939 Computational Cost, *Journal of the American Statistical Association*, 107, 1590–1598,  
940 <https://doi.org/10.1080/01621459.2012.737745>, 2012.

941 ~~Kim, J., Johnson, L., Cifelli, R., Thorstensen, A., and Chandrasekar, V.: Assessment of~~  
942 ~~antecedent moisture condition on flood frequency: An experimental study in Napa River~~  
943 ~~Basin, CA, *Journal of Hydrology: Regional Studies*, 26, 100620,~~  
944 ~~<https://doi.org/10.1016/j.ejrh.2019.100620>, 2019.~~

945 Klemeš, V.: Operational testing of hydrological simulation models, *Hydrological Sciences*  
946 *Journal*, 31, 13–24, <https://doi.org/10.1080/02626668609491024>, 1986.

947 Knutti, R. and Sedláček, J.: Robustness and uncertainties in the new CMIP5 climate model  
948 projections, *Nature Clim Change*, 3, 369–373, <https://doi.org/10.1038/nclimate1716>, 2013.

949 Koenker, R. and Bassett, G.: Regression Quantiles, *Econometrica*, 46, 33,  
950 <https://doi.org/10.2307/1913643>, 1978.

951 Kuana, L. A., Almeida, A. S., Mercuri, E. G. F., and Noe, S. M.: Regionalization of GR4J  
952 model parameters for river flow prediction in Paraná, Brazil, *Hydrol. Earth Syst. Sci.*, 28,  
953 3367–3390, <https://doi.org/10.5194/hess-28-3367-2024>, 2024.

954 Lemaitre-Basset, T., Collet, L., Thirel, G., Parajka, J., Evin, G., and Hingray, B.: Climate  
955 change impact and uncertainty analysis on hydrological extremes in a French Mediterranean  
956 catchment, *Hydrological Sciences Journal*, 66, 888–903,  
957 <https://doi.org/10.1080/02626667.2021.1895437>, 2021.

958 Lemaitre-Basset, T., Thirel, G., Oudin, L., and Dorchies, D.: Water use scenarios versus  
959 climate change: Investigating future water management of the French part of the Moselle,  
960 *Journal of Hydrology: Regional Studies*, 54, 101855,  
961 <https://doi.org/10.1016/j.ejrh.2024.101855>, 2024.

962 Lucas-Picher, P., Argüeso, D., Brisson, E., Tramblay, Y., Berg, P., Lemonsu, A., Kotlarski,  
963 S., and Caillaud, C.: CONVECTION -permitting modeling with regional climate models: Latest  
964 developments and next steps, *WIREs Climate Change*, 12, e731,  
965 <https://doi.org/10.1002/wcc.731>, 2021.

966 Lun, D., Fischer, S., Viglione, A., and Blöschl, G.: Detecting Flood-Rich and Flood-Poor  
967 Periods in Annual Peak Discharges Across Europe, *Water Resources Research*, 56,  
968 e2019WR026575, <https://doi.org/10.1029/2019WR026575>, 2020.

969 Lyne, V. D. and Hollick, M.: Stochastic time-variable rainfall runoff modelling, *Hydrology and*  
970 *Water Resources Symposium*, Institution of Engineers, Australia, Perth (1979), Barton,  
971 Australia, 1979.

972 Maraun, D., Shepherd, T. G., Widmann, M., Zappa, G., Walton, D., Gutiérrez, J. M.,  
973 Hagemann, S., Richter, I., Soares, P. M. M., Hall, A., and Mearns, L. O.: Towards process-  
974 informed bias correction of climate change simulations, *Nature Clim Change*, 7, 764–773,  
975 <https://doi.org/10.1038/nclimate3418>, 2017.

976 Marson, P., Corre, L., Soubeyroux, J.-M., Sauquet, É., and Explore2: Rapport de synthèse  
977 sur les projections climatiques régionalisées, <https://doi.org/10.57745/PUR7ML>, 2024.

978 McMillan, H., Araki, R., Bolotin, L., Kim, D.-H., Coxon, G., Clark, M., and Seibert, J.: Global  
979 patterns in observed hydrologic processes, *Nat Water*, [https://doi.org/10.1038/s44221-025-](https://doi.org/10.1038/s44221-025-00407-w)  
980 00407-w, 2025.

981 Meinshausen, M., Nicholls, Z. R. J., Lewis, J., Gidden, M. J., Vogel, E., Freund, M., Beyerle,  
982 U., Gessner, C., Nauels, A., Bauer, N., Canadell, J. G., Daniel, J. S., John, A., Krummel, P.  
983 B., Luderer, G., Meinshausen, N., Montzka, S. A., Rayner, P. J., Reimann, S., Smith, S. J.,  
984 Van Den Berg, M., Velders, G. J. M., Vollmer, M. K., and Wang, R. H. J.: The shared socio-  
985 economic pathway (SSP) greenhouse gas concentrations and their extensions to 2500,  
986 *Geosci. Model Dev.*, 13, 3571–3605, <https://doi.org/10.5194/gmd-13-3571-2020>, 2020.

987 Nepal, S., Chen, J., Penton, D. J., Neumann, L. E., Zheng, H., and Wahid, S.: Spatial GR4J  
988 conceptualization of the Tamor glaciated alpine catchment in Eastern Nepal: evaluation of  
989 GR4JSG against streamflow and MODIS snow extent, *Hydrological Processes*, 31, 51–68,  
990 <https://doi.org/10.1002/hyp.10962>, 2017.

991 [Penna, D., Tromp-van Meerveld, H. J., Gobbi, A., Borga, M., and Dalla Fontana, G.: The](#)  
992 [influence of soil moisture on threshold runoff generation processes in an alpine headwater](#)  
993 [catchment, \*Hydrol. Earth Syst. Sci.\*, 15, 689–702, <https://doi.org/10.5194/hess-15-689-2011>,](#)  
994 [2011.](#)

995 Perrin, C., Michel, C., and Andréassian, V.: Improvement of a parsimonious model for  
996 streamflow simulation, *Journal of Hydrology*, 279, 275–289, [https://doi.org/10.1016/S0022-](https://doi.org/10.1016/S0022-1694(03)00225-7)  
997 1694(03)00225-7, 2003.

998 Poncet, N., Lucas-Picher, P., Trambly, Y., Thirel, G., Vergara, H., Gourley, J., and Alias, A.:  
999 Does a convection-permitting regional climate model bring new perspectives on the  
1000 projection of Mediterranean floods?, *Nat. Hazards Earth Syst. Sci.*, 24, 1163–1183,  
1001 <https://doi.org/10.5194/nhess-24-1163-2024>, 2024.

1002 [Poncet, N., Trambly, Y., Lucas-Picher, P., Thirel, G., and Caillaud, C.: Projections of](#)  
1003 [extreme rainfall and floods in Mediterranean basins from an ensemble of convection-](#)  
1004 [permitting models, \*Climatic Change\*, 178, 141, <https://doi.org/10.1007/s10584-025-03983-8>,](#)  
1005 [2025.](#)

1006 Rogger, M., Agnoletti, M., Alaoui, A., Bathurst, J. C., Bodner, G., Borga, M., Chaplot, V.,  
1007 Gallart, F., Glatzel, G., Hall, J., Holden, J., Holko, L., Horn, R., Kiss, A., Kohnová, S.,  
1008 Leitingner, G., Lennartz, B., Parajka, J., Perdigão, R., Peth, S., Plavcová, L., Quinton, J. N.,  
1009 Robinson, M., Salinas, J. L., Santoro, A., Szolgay, J., Tron, S., van den Akker, J. J. H.,  
1010 Viglione, A., and Blöschl, G.: Land use change impacts on floods at the catchment scale:  
1011 Challenges and opportunities for future research: LAND USE CHANGE IMPACTS ON  
1012 FLOODS, *Water Resour. Res.*, 53, 5209–5219, <https://doi.org/10.1002/2017WR020723>,  
1013 2017.

1014 [Rottler, E., Bronstert, A., Bürger, G., and Rakovec, O.: Projected changes in Rhine River](#)  
1015 [flood seasonality under global warming, \*Hydrol. Earth Syst. Sci.\*, 25, 2353–2371,](#)  
1016 [https://doi.org/10.5194/hess-25-2353-2021, 2021.](#)

1017 Roudier, P., Andersson, J. C. M., Donnelly, C., Feyen, L., Greuell, W., and Ludwig, F.:  
1018 Projections of future floods and hydrological droughts in Europe under a +2°C global  
1019 warming, *Climatic Change*, 135, 341–355, <https://doi.org/10.1007/s10584-015-1570-4>, 2016.

1020 [Samaniego, L., Thober, S., Kumar, R., Wanders, N., Rakovec, O., Pan, M., Zink, M.,](#)  
1021 [Sheffield, J., Wood, E. F., and Marx, A.: Anthropogenic warming exacerbates European soil](#)

1022 [moisture droughts, \*Nature Clim Change\*, 8, 421–426, \[https://doi.org/10.1038/s41558-018-\]\(https://doi.org/10.1038/s41558-018-0138-5\)](https://doi.org/10.1038/s41558-018-0138-5)  
1023 [0138-5](https://doi.org/10.1038/s41558-018-0138-5), 2018.

1024 Santos, L., Andréassian, V., Sonnenborg, T. O., Lindström, G., De Lavenne, A., Perrin, C.,  
1025 Collet, L., and Thirel, G.: Lack of robustness of hydrological models: a large-sample  
1026 diagnosis and an attempt to identify hydrological and climatic drivers, *Hydrol. Earth Syst.*  
1027 *Sci.*, 29, 683–700, <https://doi.org/10.5194/hess-29-683-2025>, 2025.

1028 Sauquet, É., Hérault, L., Bonneau, J., Reverdy, A., Strohmenger, L., Vidal, J.-P., and  
1029 Explore2: Diagnostic des modèles hydrologiques : Des données aux résultats,  
1030 <https://doi.org/10.57745/S6PQXD>, 2024.

1031 Sauquet E., Evin G., Siauue S., Aissat R., Arnaud P., Berel M., Bonneau J., Branger F.,  
1032 Caballero Y., Colleoni F., Ducharme A., Gailhard J., Habets F., Hendrickx F., Hérault L.,  
1033 Hingray B., Huang P., Jaouen T., Jeantet A., Lanini S., Le Lay M., Magand C., Mimeau L.,  
1034 Monteil C., Munier S., Perrin C., Robelin O., Rousset F., Soubeyroux J.-M., Strohmenger L.,  
1035 Thirel G., Tocquer F., Trambly Y., Vergnes J.-P., Vidal J.-P., A large transient multi-  
1036 scenario multi-model ensemble of future streamflows and groundwater projections in France,  
1037 *Hydrology and Earth System Sciences*, [https://doi.org/10.5194/egusphere-2025-](https://doi.org/10.5194/egusphere-2025-1788)  
1038 [1788](https://doi.org/10.5194/egusphere-2025-1788)submitted, 2025.

1039 Scussolini, P., Luu, L. N., Philip, S., Berghuijs, W. R., Eilander, D., Aerts, J. C. J. H., Kew, S.  
1040 F., Van Oldenborgh, G. J., Toonen, W. H. J., Volkholz, J., and Coumou, D.: Challenges in  
1041 the attribution of river flood events, *WIREs Climate Change*, e874,  
1042 <https://doi.org/10.1002/wcc.874>, 2023.

1043 Sen, P. K.: Estimates of the Regression Coefficient Based on Kendall's Tau, *Journal of the*  
1044 *American Statistical Association*, 63, 1379–1389,  
1045 <https://doi.org/10.1080/01621459.1968.10480934>, 1968.

1046 Sharma, A., Wasko, C., and Lettenmaier, D. P.: If Precipitation Extremes Are Increasing,  
1047 Why Aren't Floods?, *Water Resour. Res.*, 54, 8545–8551,  
1048 <https://doi.org/10.1029/2018WR023749>, 2018.

1049 [Sikorska-Senoner, A. E. and Seibert, J.: Flood-type trend analysis for alpine catchments,](https://doi.org/10.1080/02626667.2020.1749761)  
1050 [Hydrological Sciences Journal. 65, 1281–1299,](https://doi.org/10.1080/02626667.2020.1749761)  
1051 <https://doi.org/10.1080/02626667.2020.1749761>, 2020

1052 Stein, L., Pianosi, F., and Woods, R.: Event-based classification for global study of river  
1053 flood generating processes, *Hydrological Processes*, 34, 1514–1529,  
1054 <https://doi.org/10.1002/hyp.13678>, 2020.

1055 Stein, L., Clark, M. P., Knoben, W. J. M., Pianosi, F., and Woods, R. A.: How Do Climate  
1056 and Catchment Attributes Influence Flood Generating Processes? A Large-Sample Study for  
1057 671 Catchments Across the Contiguous USA, *Water Res*, 57,  
1058 <https://doi.org/10.1029/2020WR028300>, 2021.

1059 Stephens, C. M., Johnson, F. M., and Marshall, L. A.: Implications of future climate change  
1060 for event-based hydrologic models, *Advances in Water Resources*, 119, 95–110,  
1061 <https://doi.org/10.1016/j.advwatres.2018.07.004>, 2018.

1062 Stephens, C. M., Marshall, L. A., and Johnson, F. M.: Investigating strategies to improve  
1063 hydrologic model performance in a changing climate, *Journal of Hydrology*, 579, 124219,  
1064 <https://doi.org/10.1016/j.jhydrol.2019.124219>, 2019.

1065 Strohmeier, L., Sauquet, E., Bernard, C., Bonneau, J., Branger, F., Bresson, A., Brigode,  
1066 P., Buzier, R., Delaigue, O., Devers, A., Evin, G., Fournier, M., Hsu, S.-C., Lanini, S., De  
1067 Lavenne, A., Lemaitre-Basset, T., Magand, C., Mendoza Guimarães, G., Mentha, M.,  
1068 Munier, S., Perrin, C., Pouchard, T., Rouchy, L., Sadki, M., Soutif-Bellenger, M., Tilmant,  
1069 F., Trambay, Y., Véron, A.-L., Vidal, J.-P., and Thirel, G.: On the visual detection of non-  
1070 natural records in streamflow time series: challenges and impacts, *Hydrol. Earth Syst. Sci.*,  
1071 27, 3375–3391, <https://doi.org/10.5194/hess-27-3375-2023>, 2023.

1072 Strohmeier, L., Collet, L., Andréassian, V., Corre, L., Rousset, F., and Thirel, G.: Köppen–  
1073 Geiger climate classification across France based on an ensemble of high-resolution climate  
1074 projections, *Comptes Rendus. Géoscience*, 356, 67–82, <https://doi.org/10.5802/crgeos.263>,  
1075 2024.

1076 Tarasova, L., Basso, S., Zink, M., and Merz, R.: Exploring Controls on Rainfall-Runoff  
1077 Events: 1. Time Series-Based Event Separation and Temporal Dynamics of Event Runoff  
1078 Response in Germany, *Water Resour. Res.*, 54, 7711–7732,  
1079 <https://doi.org/10.1029/2018WR022587>, 2018.

1080 Tarasova, L., Merz, R., Kiss, A., Basso, S., Blöschl, G., Merz, B., Viglione, A., Plötner, S.,  
1081 Guse, B., Schumann, A., Fischer, S., Ahrens, B., Anwar, F., Bárdossy, A., Bühler, P.,  
1082 Haberlandt, U., Kreibich, H., Krug, A., Lun, D., Müller-Thomy, H., Pidoto, R., Primo, C.,  
1083 Seidel, J., Vorogushyn, S., and Wietzke, L.: Causative classification of river flood events,  
1084 *WIREs Water*, 6, <https://doi.org/10.1002/wat2.1353>, 2019.

1085 Tarasova, L., Basso, S., Wendi, D., Viglione, A., Kumar, R., and Merz, R.: A Process-Based  
1086 Framework to Characterize and Classify Runoff Events: The Event Typology of Germany,  
1087 *Water Resour. Res.*, 56, <https://doi.org/10.1029/2019WR026951>, 2020.

1088 Tarasova, L., Lun, D., Merz, R., Blöschl, G., Basso, S., Bertola, M., Miniussi, A., Rakovec,  
1089 O., Samaniego, L., Thober, S., and Kumar, R.: Shifts in flood generation processes  
1090 exacerbate regional flood anomalies in Europe, *Commun Earth Environ*, 4, 49,  
1091 <https://doi.org/10.1038/s43247-023-00714-8>, 2023.

1092 Tarek, M., Brissette, F., and Arsenault, R.: Uncertainty of gridded precipitation and  
1093 temperature reference datasets in climate change impact studies, *Hydrol. Earth Syst. Sci.*,  
1094 25, 3331–3350, <https://doi.org/10.5194/hess-25-3331-2021>, 2021.

1095 Taylor, K. E., Stouffer, R. J., and Meehl, G. A.: An Overview of CMIP5 and the Experiment  
1096 Design, *Bulletin of the American Meteorological Society*, 93, 485–498,  
1097 <https://doi.org/10.1175/BAMS-D-11-00094.1>, 2012.

1098 Terray, L. and Boé, J.: Quantifying 21st-century France climate change and related  
1099 uncertainties, *Comptes Rendus. Géoscience*, 345, 136–149,  
1100 <https://doi.org/10.1016/j.crte.2013.02.003>, 2013.

1101 Teutschbein, C. and Seibert, J.: Bias correction of regional climate model simulations for  
1102 hydrological climate-change impact studies: Review and evaluation of different methods,  
1103 *Journal of Hydrology*, 456–457, 12–29, <https://doi.org/10.1016/j.jhydrol.2012.05.052>, 2012.

1104 Thirel, G., Andréassian, V., Perrin, C., Audouy, J.-N., Berthet, L., Edwards, P., Folton, N.,  
1105 Furusho, C., Kuentz, A., Lerat, J., Lindström, G., Martin, E., Mathevet, T., Merz, R., Parajka,  
1106 J., Ruelland, D., and Vaze, J.: Hydrology under change: an evaluation protocol to investigate  
1107 how hydrological models deal with changing catchments, *Hydrological Sciences Journal*, 60,  
1108 1184–1199, <https://doi.org/10.1080/02626667.2014.967248>, 2015a.

- 1109 Thirel, G., Andréassian, V., and Perrin, C.: On the need to test hydrological models under  
1110 changing conditions, *Hydrological Sciences Journal*, 60, 1165–1173,  
1111 <https://doi.org/10.1080/02626667.2015.1050027>, 2015b.
- 1112 Thirel, G., Gerlinger, K., Perrin, C., Drogue, G., Renard, B., and Wagner, J.-P.: Quels futurs  
1113 possibles pour les débits des affluents français du Rhin (Moselle, Sarre, Ill) ?, *La Houille*  
1114 *Blanche*, 105, 140–149, <https://doi.org/10.1051/lhb/2019039>, 2019.
- 1115 Tian, Y., Xu, Y.-P., and Zhang, X.-J.: Assessment of Climate Change Impacts on River High  
1116 Flows through Comparative Use of GR4J, HBV and Xinanjiang Models, *Water Resour*  
1117 *Manage*, 27, 2871–2888, <https://doi.org/10.1007/s11269-013-0321-4>, 2013.
- 1118 Tramblay, Y. and Somot, S.: Future evolution of extreme precipitation in the Mediterranean,  
1119 *Climatic Change*, 151, 289–302, <https://doi.org/10.1007/s10584-018-2300-5>, 2018.
- 1120 Tramblay, Y., Mimeau, L., Neppel, L., Vinet, F., and Sauquet, E.: Detection and attribution of  
1121 flood trends in Mediterranean basins, *Hydrol. Earth Syst. Sci.*, 23, 4419–4431,  
1122 <https://doi.org/10.5194/hess-23-4419-2019>, 2019.
- 1123 Tramblay, Y., Villarini, G., Khalki, E. M., Gründemann, G., and Hughes, D.: Evaluation of the  
1124 Drivers Responsible for Flooding in Africa, *Water Resources Research*, 57,  
1125 <https://doi.org/10.1029/2021WR029595>, 2021.
- 1126 Tramblay, Y., Villarini, G., Saidi, M. E., Massari, C., and Stein, L.: Classification of flood-  
1127 generating processes in Africa, *Sci Rep*, 12, 18920, [https://doi.org/10.1038/s41598-022-](https://doi.org/10.1038/s41598-022-23725-5)  
1128 [23725-5](https://doi.org/10.1038/s41598-022-23725-5), 2022.
- 1129 Tramblay, Y., Arnaud, P., Artigue, G., Lang, M., Paquet, E., Neppel, L., and Sauquet, E.:  
1130 Changes in Mediterranean flood processes and seasonality, *Hydrol. Earth Syst. Sci.*, 27,  
1131 2973–2987, <https://doi.org/10.5194/hess-27-2973-2023>, 2023.
- 1132 Tramblay, Y., Sauquet, É., Arnaud, P., Rousset, F., Soubeyroux, J.-M., Hingray, B., Jaouen,  
1133 T., Jeantet, A., Munier, S., Vergnes, J.-P., and Explore2: Scénarios d'extrêmes  
1134 hydrologiques, <https://doi.org/10.57745/2XDJ5H>, 2024.
- 1135 Valéry, A., Andréassian, V., and Perrin, C.: 'As simple as possible but not simpler': What is  
1136 useful in a temperature-based snow-accounting routine? Part 2 – Sensitivity analysis of the  
1137 Cemaneige snow accounting routine on 380 catchments, *Journal of Hydrology*, 517, 1176–  
1138 1187, <https://doi.org/10.1016/j.jhydrol.2014.04.058>, 2014.
- 1139 Vautard, R., Kadyrov, N., Iles, C., Boberg, F., Buonomo, E., Bülow, K., Coppola, E., Corre,  
1140 L., Van Meijgaard, E., Nogherotto, R., Sandstad, M., Schwingshackl, C., Somot, S., Aalbers,  
1141 E., Christensen, O. B., Ciarlo, J. M., Demory, M., Giorgi, F., Jacob, D., Jones, R. G., Keuler,  
1142 K., Kjellström, E., Lenderink, G., Levavasseur, G., Nikulin, G., Sillmann, J., Solidoro, C.,  
1143 Sørland, S. L., Steger, C., Teichmann, C., Warrach-Sagi, K., and Wulfmeyer, V.: Evaluation  
1144 of the Large EURO-CORDEX Regional Climate Model Ensemble, *JGR Atmospheres*, 126,  
1145 e2019JD032344, <https://doi.org/10.1029/2019JD032344>, 2021.
- 1146 Verfaillie, D., Déqué, M., Morin, S., and Lafaysse, M.: The method ADAMONT v1.0 for  
1147 statistical adjustment of climate projections applicable to energy balance land surface  
1148 models, *Geosci. Model Dev.*, 10, 4257–4283, <https://doi.org/10.5194/gmd-10-4257-2017>,  
1149 2017.
- 1150 Vidal, J., Martin, E., Franchistéguy, L., Baillon, M., and Soubeyroux, J.: A 50-year high-  
1151 resolution atmospheric reanalysis over France with the Safran system, *Intl Journal of*

1152 Climatology, 30, 1627–1644, <https://doi.org/10.1002/joc.2003>, 2010.

1153 Vidal, J.-P., Hingray, B., Magand, C., Sauquet, E., and Ducharne, A.: Hierarchy of climate  
 1154 and hydrological uncertainties in transient low-flow projections, Hydrol. Earth Syst. Sci., 20,  
 1155 3651–3672, <https://doi.org/10.5194/hess-20-3651-2016>, 2016.

1156 Wasko, C. and Nathan, R.: Influence of changes in rainfall and soil moisture on trends in  
 1157 flooding, Journal of Hydrology, 575, 432–441, <https://doi.org/10.1016/j.jhydrol.2019.05.054>,  
 1158 2019.

1159 [Wasko, C., Nathan, R., Stein, L., and O'Shea, D. Evidence of shorter more extreme rainfalls  
 1160 and increased flood variability under climate change. Journal of Hydrology, 603, 126994,  
 1161 <https://doi.org/10.1016/j.jhydrol.2021.126994>, 2021.](https://doi.org/10.1016/j.jhydrol.2021.126994)

1162 Wasko, C., Guo, D., Ho, M., Nathan, R., and Vogel, E.: Diverging projections for flood and  
 1163 rainfall frequency curves, Journal of Hydrology, 620, 129403,  
 1164 <https://doi.org/10.1016/j.jhydrol.2023.129403>, 2023.

1165 Yang, L., Yang, Y., Villarini, G., Li, X., Hu, H., Wang, L., Blöschl, G., and Tian, F.: Climate  
 1166 More Important for Chinese Flood Changes Than Reservoirs and Land Use, Geophys Res  
 1167 Lett, 48, <https://doi.org/10.1029/2021GL093061>, 2021.

1168 Zahar, Y., Ghorbel, A., and Albergel, J.: Impacts of large dams on downstream flow  
 1169 conditions of rivers: Aggradation and reduction of the Medjerda channel capacity  
 1170 downstream of the Sidi Salem dam (Tunisia), Journal of Hydrology, 351, 318–330,  
 1171 <https://doi.org/10.1016/j.jhydrol.2007.12.019>, 2008.

1172 Zhang, S., Zhou, L., Zhang, L., Yang, Y., Wei, Z., Zhou, S., Yang, D., Yang, X., Wu, X.,  
 1173 Zhang, Y., Li, X., and Dai, Y.: Reconciling disagreement on global river flood changes in a  
 1174 warming climate, Nat. Clim. Chang., 12, 1160–1167, <https://doi.org/10.1038/s41558-022-01539-7>, 2022.

1175

1176