

Acid-catalyzed hydrolysis kinetics of organic hydroperoxides: Computational strategy and structure-activity relationship

Qiaojing Zhao¹, Fangfang Ma^{1,2}, Hui Zhao¹, Qian Xu¹, Rujing Yin^{1*}, Hong-Bin Xie^{1*}, Xin Wang^{1,3}, Jingwen Chen¹

¹ Key Laboratory of Industrial Ecology and Environmental Engineering (Ministry of Education), School of Environmental Science and Technology, Dalian University of Technology, Dalian 116024, China

² College of Resources and Environmental Engineering, Guizhou University, Guiyang 550025, China

³ Key Laboratory for Semi-Arid Climate Change of the Ministry of Education, College of Atmospheric Sciences, Lanzhou University, Lanzhou 730000, China

Correspondence to: Rujing Yin (yinrj@dlut.edu.cn) and Hong-Bin Xie (hbxie@dlut.edu.cn)

Abstract. Organic hydroperoxides (ROOHs) are key components of atmospheric aerosols. Determining the acid-catalyzed hydrolysis rate constants (k_A) of ROOHs is crucial for assessing their atmospheric fate and environmental impacts. However, available k_A values are limited due to the difficulty in obtaining authentic ROOH standards. Herein, we addressed this limitation by developing a computational strategy and probing the structure-activity relationship of k_A values. We screened the protonated water cluster ($H^+(H_2O)_n$) model, a critical prerequisite for density functional theory (DFT) calculations of k_A , by comparing experimental k_A values of four ROOHs with DFT-calculated values using $H^+(H_2O)_n$ ($n = 1, 2, 3, 4$) models. Results show that the $H^+(H_2O)_2$ model reliably predicts k_A values with DFT method. Further investigation of 53 additional ROOHs including 45 model compounds and 8 atmospherically relevant species reveals that substituents at the C_α (the carbon atom directly bonded to the -OOH group) site, including -NH₂, -N(CH₃)₂, -OH, -OCH₃, -CH=CH₂, -SH, and -PH₂, can facilitate acid-catalyzed hydrolysis. Notably, the -NH₂ and -N(CH₃)₂ substituents exhibit stronger facilitating effect than the well-documented -OH and -OCH₃ substituents. Additionally, we clarified that not all nitrogen- or oxygen-containing substituents equally enhance k_A , as their efficacy depends on the substituents attached to the O or N atoms. This study provides a reliable computational strategy and essential guidelines for predicting k_A values of ROOHs, enabling more accurate simulations in atmospheric chemistry models.

1 Introduction

Aerosol liquid water, a crucial constituent of atmospheric aerosols, acts as a reactive medium that enables aqueous-phase chemical transformations (Jin et al., 2020; Su et al., 2022; Shi et al., 2024; Wu et al., 2018). The aqueous-phase transformations of organic compounds substantially modify the physicochemical properties of aerosols, such as chemical composition, optical characteristics, and hygroscopicity, ultimately altering the health and climate effects of aerosols (McNeill, 2015; Herrmann et al., 2015; Zheng et al., 2021; Liu et al., 2023; Lei et al., 2022; Zhang et al., 2024). Recently, aqueous-phase chemistry has gained increasing research interest due to its unique reaction mechanism compared to gas-phase chemistry. However, the

complexity and diversity of these reaction pathways have limited our understanding of aqueous-phase chemistry. The lack of kinetic data for the aqueous-phase chemistry of organic compounds, such as acid-catalyzed hydrolysis or esterification, oxidation, photosensitization, and oligomer processes, further hinders its application in precise simulations of three-dimensional (3-D) atmospheric chemistry models (Ervens et al., 2024; Wieser et al., 2024; Abbatt and Ravishankara, 2023).

Organic hydroperoxides (ROOHs) are ubiquitous in aerosols and predominantly derived from atmospheric oxidation processes involving organic peroxy radicals ($\text{RO}_2\cdot$) or Criegee intermediates (CIs) (Wang et al., 2023). Characterized by one or more hydrophilic hydroperoxide groups (-OOH), ROOHs are highly reactive and usually experience rapid aqueous-phase transformations, forming low-volatile multi-functional species or reactive oxygen species (Enami, 2021; Krapf et al., 2016; Dovrou et al., 2019; Wei et al., 2021; Wei et al., 2022a; Wei et al., 2022b; Wang et al., 2019). Laboratory evidence demonstrates that ROOHs can account for 20%-60% isoprene- or monoterpene-derived secondary organic aerosol (SOA) (Wang et al., 2023; Enami, 2021; Epstein et al., 2014). Therefore, investigating the transformation kinetics of ROOHs is essential for improving our understanding of aqueous-phase chemistry.

Hydrolysis has been identified as an important transformation pathway for ROOHs, yielding hydrogen peroxide (H_2O_2) and thereby affect atmospheric oxidation capacity (Enami, 2021; Qiu et al., 2019; Qiu et al., 2020a; Hu et al., 2021a; Dai et al., 2024). Recent studies have found that two types of α -substituted ROOHs, α -hydroxyalkyl-hydroperoxides (α -HHs) and α -alkoxyalkyl-hydroperoxides (α -AHs), exhibit rapid hydrolysis under acid-catalyzed conditions. Their lifetimes are short from seconds to minutes at $\text{pH} < 4$, with first-order rate coefficients increasing significantly as pH decreases (Qiu et al., 2020b; Hu et al., 2021b; Enami, 2022; Hu et al., 2022; Endo et al., 2022). Two distinct mechanisms have been proposed for the acid-catalyzed hydrolysis of α -HHs and α -AHs. For α -HHs, both the -OOH and -OH groups attached to the same carbon atom participate in the reaction simultaneously, yielding the aldehyde and H_2O_2 (Qiu et al., 2020b); while α -AHs decompose directly to release H_2O_2 and generate the carbocation intermediates, which are rapidly hydrated to form the corresponding alcohols (Hu et al., 2021a). Moreover, a latest study further demonstrated that some monoterpene-derived α -acyloxyalkyl-hydroperoxides also decay significantly at low pH (Chang et al., 2025). This highlights the significance of acid-catalyzed hydrolysis in transforming ROOHs and modifying the oxidation capacity of the surrounding atmosphere. However, the currently available data on acid-catalyzed hydrolysis rate constants (k_A) are still very limited, impeding a comprehensive understanding of this aqueous-phase transformation process. The k_A data lack manifests in two aspects: i) current research on α -substituted ROOHs is confined to those with hydroxyl, alkoxy, and acyloxy substituents, while the k_A values of α -substituted ROOHs with other substituents remains unclear; ii) despite recent studies have indicated that non- α -substituted ROOHs, which are more abundant in the atmosphere, have longer lifetimes up to days in aqueous phase, detailed investigations for their k_A values are lacking (Dai et al., 2024; Zhao et al., 2022). Therefore, a comprehensive investigation of k_A values for structurally diverse ROOHs in atmosphere is urgently needed to advance our understanding of this aqueous-phase transformation process.

Acquiring k_A values for diverse ROOHs through laboratory experiments is significantly challenging due to the lack of commercial standards and the prerequisite synthesis of target compounds. Given these experimental limitations and the structural diversity of ROOHs, a viable approach is to use the quantum chemical method to conduct structure-activity

relationship investigation. However, previous DFT calculation studies using the simplest protonated water cluster H_3O^+ (i.e., $\text{H}^+(\text{H}_2\text{O})_1$) model have overestimated hydrolysis rates (Hu et al., 2022), since H_3O^+ would cluster with water molecules in solution and reduce the proton activity (Agmon et al., 2016). Hence, it is urgent to screen a more suitable protonated water cluster model within the DFT calculation strategy to accurately and efficiently predict the k_A values of ROOHs.

In this study, we initially selected four types of ROOHs with experimental k_A values as tested compounds to search for a suitable protonated water cluster model. Utilizing the screened protonated water cluster model and DFT calculations, the acid-catalyzed hydrolysis of additional 45 ROOH model compounds was systematically investigated to elucidate the structure-activity relationship of k_A values and the corresponding quantitative structure-activity relationship (QSAR) models were developed. These model compounds were selected mainly considering the diverse structures at the C_α (the carbon atom directly bonded to the -OOH group) and C_β (the carbon atom attached to C_α) sites. The investigation was further extended to other eight atmospheric ROOHs to validate the structure-activity relationship revealed by the model compounds. The screened protonated water cluster model and the revealed structure-activity relationship can be instrumental in predicting the k_A values of atmospheric ROOHs. The established kinetic database for ROOHs would further improve the predictions of 3-D chemical models and deepen our understanding of atmospheric aqueous chemistry.

2 Computational details

2.1 Electronic structure calculations

All structure optimizations and energy calculations were performed using the Gaussian 09 program (Frisch et al., 2009). Geometry optimization and frequency calculation for the reactants (R), products (P), transition states (TS), pre-reactive complexes (RC), intermediates (IM), and post-reactive complexes (PC) were conducted at the M06-2X/6-31+G(d,p) level of theory (Zhao and Truhlar, 2008). Single-point energy (SPE) calculations were performed at the M06-2X/6-311++G(3df,2pd) level. The employed quantum chemistry calculation method can be succinctly represented as M06-2X/6-311++G(3df,2pd)//M06-2X/6-31+G(d,p). Previous studies have demonstrated that the M06-2X method is suitable for predicting energies and kinetics for the aqueous-phase reactions of organic compounds with a good balance between accuracy and computational efficiency (Ji et al., 2020; Shi et al., 2024; Zhang et al., 2022; Piletic et al., 2013). The solvation model based on density (SMD) was applied to account for the effects of solvent water molecules in the aqueous phase (Marenich et al., 2009). Intrinsic reaction coordinate calculations were used to confirm that the well-defined TSs connect with the corresponding reactants and products (Fukui, 1981). For reaction pathways where TSs could not be successfully located, relaxed scan methods were employed to obtain free-energy surfaces (Zhao et al., 2023; Ryu et al., 2018). Gibbs free energy (G) values for each structure at 298.15 K were calculated by combining the SPE with the Gibbs correction energy calculated at the theoretical level of geometry optimization. A correction factor of $1.89 \text{ kcal mol}^{-1}$ was applied to the activation free energy (ΔG^\ddagger) and reaction free energy ($\Delta_r G$) calculations for reactions where the number of molecules decreases or increases by one from R to TS or from R to P to explain the free energy change from the gas phase standard state of 1 atm to the aqueous phase

standard state of 1 mol L⁻¹ (Sadlej-Sosnowska, 2007; Zhang et al., 2015). The combined use of Molclus 1.9.9.9 and Gaussian 09 programs was applied to search for the global minimum of the reactants (Lu., 2022).

100 2.2 H⁺(H₂O)_n model selection

A previous study found that the simplest protonated water cluster H⁺(H₂O)₁ model overestimated acid-catalyzed hydrolysis rates of ROOHs by combining with feasible DFT method (Hu et al., 2022). This is probably because H⁺(H₂O)₁ clusters with water molecules in the aqueous solution and the proton activity decreases (Agmon et al., 2016). The protonated water cluster model has been found to predict the acid-catalyzed hydrolysis kinetics of other organic compounds such as epoxydiols with high accuracy (Piletic et al., 2013). Here, we screened an appropriate protonated water cluster model by
105 comparing calculated k_A values for reactions between H⁺(H₂O)_n ($n = 1, 2, 3, 4$) and the selected ROOHs with experimental values. Specially, C₁₃ α -AH, C₁₂ α -AH₍₁₎, C₁₂ α -AH₍₂₎, and C₁₀ α -HH were chosen due to the available experimental kinetic data under different pH values (Qiu et al., 2020b; Enami, 2022; Hu et al., 2022). The structures of these four ROOHs and H⁺(H₂O)_n ($n = 1, 2, 3, 4$) are presented in Figs. S1 and S2.

110 2.3 Reaction rate constants calculation

The second-order reaction rate constants k_A of the elementary reactions in acid-catalyzed hydrolysis pathways were calculated using transition state theory (Eq. (1)) (Zhao et al., 2023; Xu et al., 2019). The pseudo-first-order rate constants (k'_A) were calculated by combining k_A with the concentration of protonated water clusters, which are determined by pH. Corresponding acid-catalyzed hydrolysis lifetimes ($\tau_{1/e}$) can be subsequently derived by $\tau_{1/e} = 1/k'_A$.

$$115 \quad k_A = \sigma \frac{k_B T}{h} \exp\left(-\frac{\Delta G^\ddagger}{RT}\right), \quad (1)$$

where σ is the reaction path degeneracy, T is the temperature (298.15 K), k_B is the Boltzmann constant (J K⁻¹), h is the Planck constant (J s), R is the gas constant (8.314 J mol⁻¹ K⁻¹), ΔG^\ddagger is the activation free energy.

2.4 QSAR modeling

The QSAR models were developed for k_A of α - and β -substituted compounds, respectively, among the selected 45 ROOH
120 model compounds. The Taft (σ^*) constants, which are widely used to quantify the inductive effects of substituents in aliphatic systems (Lee and von Gunten, 2012; Kim and Huang, 2021; Ra et al., 2025), were employed to correlate with the logarithms of k_A ($\log k_A$) values in this study. The σ^* values for different substituents, as shown in Table S1, were obtained from the literature (Perrin et al., 1981). Herein, the total effects were evaluated by summing the σ^* values ($\sum \sigma^*$) of each substituent (except the -OOH group) at the reaction center, i.e., the C _{α} site (Table S2).

3.1 Selection of suitable $\text{H}^+(\text{H}_2\text{O})_n$ model

We investigated the acid-catalyzed hydrolysis of four ROOHs, i.e., C_{13} α -AH, C_{12} α -AH₍₁₎, C_{12} α -AH₍₂₎, and C_{10} α -HH, using $\text{H}^+(\text{H}_2\text{O})_n$ ($n = 1, 2, 3, 4$) models with the DFT method. The calculated free-energy profiles for these reactions are presented in Figs. S3-S6. Concluded from the reaction profiles, a two-step acid-catalyzed hydrolysis pathway is revealed for these four ROOHs, with the first step being the rate-limiting step. As shown in Fig. S7, the H^+ of $\text{H}^+(\text{H}_2\text{O})_n$ attacks the -OOH group of the selected four compounds, leading to $\text{C}_\alpha\text{-O}$ bond rupture and simultaneous formation of a carbocation intermediate and H_2O_2 in the first step. Subsequently, the carbocation reacts with water molecules and yields the corresponding protonated alcohols.

Using the activation free energies of the rate-limiting step, the pseudo-first-order rate constants k'_A were calculated over a pH range of 0-14 using the M06-2X/6-311++G(3df,2pd)//M06-2X/6-31+G(d,p) method. The calculated k'_A values for the four selected ROOHs with $\text{H}^+(\text{H}_2\text{O})_n$ models are presented in Fig. 1, along with the experimentally determined hydrolysis rates at different pH values (Qiu et al., 2020b; Enami, 2022; Hu et al., 2022; Endo et al., 2022). The $\text{H}^+(\text{H}_2\text{O})_2$ model was found to best reproduce the experimental data among $\text{H}^+(\text{H}_2\text{O})_n$ ($n = 1, 2, 3, 4$) models. In most cases, the discrepancies between the calculated and the experimental k'_A values are within a factor of 0.5 to 2.2. Two exceptions are C_{10} α -HH at pH 3.3 and C_{12} α -AH₍₂₎ at pH 4.5, but their uncertainties remain within one order of magnitude, an acceptable error range. In contrast, the $\text{H}^+(\text{H}_2\text{O})_4$, $\text{H}^+(\text{H}_2\text{O})_3$, and $\text{H}^+(\text{H}_2\text{O})_1$ models result in much larger discrepancies, with errors reaching 1.0×10^{-5} , 1.1×10^{-3} , and 2.1×10^8 times, respectively. Therefore, the computational strategy combining the $\text{H}^+(\text{H}_2\text{O})_2$ model with the M06-2X/6-311++G(3df,2pd)//M06-2X/6-31+G(d,p) method is well-suited for predicting k_A values of ROOHs.




Figure 1. Variation of calculated pseudo-first-order acid-catalyzed hydrolysis rate constants (k'_A) for four ROOHs along with the available experimental values (Qiu et al., 2020b; Enami, 2022; Hu et al., 2022; Endo et al., 2022). (a) C₁₃ α -AH, (b) C₁₂ α -AH₍₁₎, (c) C₁₂ α -AH₍₂₎, and (d) C₁₀ α -HH.

3.2 Structure-activity relationship of acid-catalyzed hydrolysis rate constants

Employing the $H^+(H_2O)_2$ model and the M06-2X/6-311++G(3df,2pd)//M06-2X/6-31+G(d,p) method, we conducted a systematic investigation for structure-activity relationships of k_A values across 45 ROOH model compounds (Fig. 2a). We selected the simplest unsubstituted primary, secondary, and tertiary ROOHs, i.e., CH_3CH_2OOH , $CH(CH_3)_2OOH$, and $C(CH_3)_3OOH$, as the reference compounds to assess the influence of substituents. Specifically, the influence of substituents was examined by comparing the k_A values of α -substituted primary, secondary, and tertiary ROOHs, as well as β -substituted tertiary ROOHs, represented as $CH_2(X)OOH$, $CH(CH_3)(X)OOH$, $C(CH_3)_2(X)OOH$, and $C(CH_3)_2(CH_2(X))OOH$, with the corresponding unsubstituted ones, respectively. Here, X denotes substituents including $-NH_2$, $-OH$, $-OCH_3$, $-CH=CH_2$, $-SH$, $-PH_2$, $-F$, $-Cl$, and $-CHO$, covering the major functional group types found in atmospheric ROOHs. Therefore, the effects of

different carbon skeletons at the C_α site (primary, secondary, and tertiary) and different substituents at the C_α and C_β sites of the -OOH group on the *k_A* values of ROOHs were revealed.





Figure 2. Molecular structure and acid-catalyzed hydrolysis pathways of ROOH model compounds. (a) Molecular structure of 45 ROOH model compounds; (b) One-step and two-step acid-catalyzed hydrolysis pathways of ROOHs.

For the selected 45 ROOH model compounds, their reaction free-energy profiles were calculated and shown in Figs. S8-S12. They follow two distinct reaction pathways, which are simplified and illustrated in Fig. 2b. CH₃CH₂OOH, CH₂(X)OOH, and CH(CH₃)(X)OOH (X = F, Cl, CHO) undergo a one-step reaction pathway, where H⁺(H₂O)₂ and an H₂O molecule simultaneously attack the -OOH group and the C_α, respectively, leading to the formation of protonated alcohols and H₂O₂. Meanwhile, the other ROOH model compounds follow the two-step reaction pathway via the formation of a carbocation intermediate, similar to that of C₁₃ α-AH, C₁₂ α-AH₍₁₎, C₁₂ α-AH₍₂₎, and C₁₀ α-HH. It is found that these model compounds share nearly identical reaction sites (i.e., the -OOH group and its adjacent C_α atom) and follow analogous reaction pathways to C₁₃ α-AH, C₁₂ α-AH₍₁₎, C₁₂ α-AH₍₂₎, and C₁₀ α-HH. This mechanistic similarity implies that despite variations in their carbon skeletons and substituents, protonation and subsequent reactions are primarily localized at the C_α-OOH site, which is a common structural feature among ROOHs. Therefore, even though experimental data for these model compounds are currently lacking to validate the computational results, we believe the screened protonated water cluster model can be reasonably extended to investigate the acid-catalyzed hydrolysis of diverse ROOHs with acceptable uncertainty.

Based on reaction activation free energies, the *k_A* values for 45 ROOH model compounds were calculated, as well as *k'_A* values under different pH conditions (Table S3 and Fig. 3). As shown in Fig. 3a, the *k'_A* of tertiary ROOH is the highest among the three unsubstituted ROOHs at the same pH, followed by secondary and primary ROOHs, aligning with the previous research findings (Hu et al., 2022). Furthermore, to clarify the effects of substituent types, we calculated the enhancement factors (EF) for *k'_A* of substituted ROOHs relative to their unsubstituted reference compounds. Because the predicted *k'_A* values

180 for ROOHs exhibit a linear dependence on pH, the resulting EF values remain constant across different pH values. These EF values for α -substituted ROOH model compounds, $\text{CH}_2(\text{X})\text{OOH}$, $\text{CH}(\text{CH}_3)(\text{X})\text{OOH}$, and $\text{C}(\text{CH}_3)_2(\text{X})\text{OOH}$ are shown in Fig. 3b-d, with $\text{CH}_3\text{CH}_2\text{OOH}$, $\text{CH}(\text{CH}_3)_2\text{OOH}$, and $\text{C}(\text{CH}_3)_3\text{OOH}$ as reference compounds, respectively. Substituents such as $-\text{NH}_2$, $-\text{OH}$, $-\text{OCH}_3$, $-\text{SH}$, and $-\text{CH}=\text{CH}_2$ increase k'_A , with EF values ranging from 5.1×10^3 - 6.4×10^{23} for primary, 1.3×10^5 - 5.0×10^{18} for secondary, and 1.4 - 5.7×10^{13} for tertiary ROOHs. The $-\text{PH}_2$ substituent shows a divergent effect, increasing k'_A of
 185 primary and secondary ROOHs with the EF values of 1.3×10^5 and 2.5 , respectively, but decreasing that of tertiary ones. While $-\text{OH}$ and $-\text{OCH}_3$ are known to facilitate the acid-catalyzed hydrolysis of ROOHs (Hu et al., 2021a; Qiu et al., 2020b; Hu et al., 2021b; Enami, 2022; Hu et al., 2022; Endo et al., 2022), our study is the first to show that $-\text{NH}_2$, $-\text{SH}$, $-\text{CH}=\text{CH}_2$, and $-\text{PH}_2$ can also enhance it. Among them, $-\text{NH}_2$ has the highest EF, followed by $-\text{OH}$ and $-\text{OCH}_3$, $-\text{SH}$, $-\text{CH}=\text{CH}_2$, and $-\text{PH}_2$. In contrast, $-\text{F}$, $-\text{Cl}$, and $-\text{CHO}$ significantly decrease the k'_A .



190
 195 **Figure 3. Calculated pH-dependent pseudo-first-order rate constants (k'_A) of ROOH model compounds and substituent enhancement effects on k'_A . (a) Calculated k'_A -pH profiles for reference ROOHs: $\text{CH}_3\text{CH}_2\text{OOH}$, $\text{CH}(\text{CH}_3)_2\text{OOH}$, $\text{C}(\text{CH}_3)_3\text{OOH}$, and $\text{C}(\text{CH}_3)_2\text{OCH}_3\text{OOH}$. (b-f) Enhancement factors ($\text{EF} = k'_\text{A, ROOHs}/k'_\text{A, reference}$) for (b) $\text{CH}_2(\text{X})\text{OOH}$ vs. $\text{CH}_3\text{CH}_2\text{OOH}$, (c) $\text{CH}(\text{CH}_3)(\text{X})\text{OOH}$ vs. $\text{CH}(\text{CH}_3)_2\text{OOH}$, (d) $\text{C}(\text{CH}_3)_2(\text{X})\text{OOH}$ vs. $\text{C}(\text{CH}_3)_3\text{OOH}$, (e) $\text{C}(\text{CH}_3)_2(\text{CH}_2(\text{X}))\text{OOH}$ vs. $\text{C}(\text{CH}_3)_3\text{OOH}$, and (f) $\text{C}(\text{CH}_3)_2(\text{OY})\text{OOH}$ vs. $\text{C}(\text{CH}_3)_2\text{OCH}_3\text{OOH}$.**

To further evaluate the effectiveness of acid-catalyzed hydrolysis for these α -substituted model compounds in the atmosphere, we calculated their $\tau_{1/e}$ values at pH 3.8 and 0.9, representing typical aerosol conditions in inland China and the Southeastern United States, respectively (Wang et al., 2022; Jia et al., 2018; Ding et al., 2019; Shi et al., 2019; Xu et al., 2020; Wang et al., 2020; Guo et al., 2015). Compared to the typical atmospheric retention time of ambient aerosols with

approximately 1-2 weeks (Hodzic et al., 2016; Kristiansen et al., 2016), Table 1 shows the calculated k_A and $\tau_{1/e}$ values of the α -substituted ROOH model compounds that undergo effective acid-catalyzed hydrolysis on this time scale at pH 3.8 and 0.9. The $\tau_{1/e}$ values for -NH₂-substituted primary ROOH, -NH₂- and -OH-substituted secondary ROOHs, as well as -NH₂-, -OH-, and -OCH₃-substituted tertiary ROOHs range from less than 1 second to 5.5 hours at pH 3.8. As pH decreases to 0.9, effective hydrolysis occurs for -OH-substituted primary ROOH, -OCH₃-substituted secondary ROOH, and -SH-substituted tertiary ROOH, with $\tau_{1/e}$ ranging from 40.9 min to 6.8 d. The $\tau_{1/e}$ values for other compounds exceeding 2 weeks are shown in Table S3, indicating the limited effectiveness of acid-catalyzed hydrolysis under typical atmospheric conditions. In summary, substituents such as -NH₂, -OH, and -OCH₃ can facilitate effective acid-catalyzed hydrolysis of ROOHs under typical ambient conditions, while the -SH substituent is only effective under exceptionally low pH. Although -OH and -OCH₃ substituents are well-documented, we show that the -NH₂ substituent leads to significantly shorter $\tau_{1/e}$ for the corresponding ROOHs. Despite the -NH₂ group prefers to exist in its protonated form (-NH₃⁺) in the aqueous phase, the rapid reaction of ROOHs with unprotonated -NH₂ group may shift the equilibrium -NH₃⁺ + H₂O \leftrightarrow -NH₂ + H₃O⁺ forward. Recent studies (Ji et al., 2020; Shi et al., 2024) have reported fast reactions between small α -dicarbonyls and amines or ammonia under acidic aqueous conditions, with corresponding products detected experimentally. This indicates that the ROOHs with unprotonated -NH₂ group can still participate in the reaction. In addition, Enami et al. (2010) found that trimethylamine remains largely unprotonated at the air-water interface even at pH 4.0. These findings inspire us that both aqueous-phase and interfacial transformations of α -NH₂-substituted ROOHs are plausible. Previous studies demonstrate that α -NH₂-substituted ROOHs and their analogues α -N(CH₃)₂-substituted ones are primarily formed via the reactions of CIs with ammonia or the oxidation of tertiary amines (Li et al., 2024; Kjaergaard et al., 2023; Ma et al., 2021). Therefore, investigating the transformation of these nitrogen-containing ROOHs should be important, particularly in regions with high amine and ammonia concentrations.

Table 1. Calculated acid-catalyzed hydrolysis second-order reaction rate constants (k_A) and corresponding lifetimes ($\tau_{1/e}$) of 14 ROOH model compounds under two selected scenarios, Inland China (pH 3.8) and Southeastern United States (pH 0.9).

Compounds	Formulas	k_A (L mol ⁻¹ s ⁻¹)	$\tau_{1/e}$, pH 3.8	$\tau_{1/e}$, pH 0.9
CH ₂ (X)OOH	CH ₂ (NH ₂)OOH	1.01×10^3	6.3 s	< 1 s
	CH ₂ (OH)OOH	1.94×10^{-5}	—	4.7 d
CH ₃ CH(X)OOH	CH ₃ CH(NH ₂)OOH	5.60×10^4	< 1 s	< 1 s
	CH ₃ CH(OH)OOH	3.20×10^{-1}	5.5 h	24.8 s
	CH ₃ CH(OCH ₃)OOH	3.24×10^{-3}	—	40.9 min
C(CH ₃) ₂ (X)OOH	C(CH ₃) ₂ (NH ₂)OOH	3.10×10^7	< 1 s	< 1 s
	C(CH ₃) ₂ (OH)OOH	4.87×10^2	13.0 s	< 1 s
	C(CH ₃) ₂ (OCH ₃)OOH	3.93×10^1	2.7 min	< 1 s
	C(CH ₃) ₂ (SH)OOH	1.36×10^{-5}	—	6.8 d
C(CH ₃) ₂ (OY)OOH	C(CH ₃) ₂ (OCH ₂ CH ₃)OOH	2.17×10^1	4.9 min	< 1 s
	C(CH ₃) ₂ (OCH ₂ CH ₂ CH ₃)OOH	1.76×10^1	6.0 min	< 1 s
	C(CH ₃) ₂ (OCH(CH ₃) ₂)OOH	4.16×10^2	15.2 s	< 1 s
	C(CH ₃) ₂ (OC(CH ₃) ₃)OOH	1.81×10^3	3.5 s	< 1 s
	C(CH ₃) ₂ (OCH=CH ₂)OOH	1.62×10^{-3}	—	1.4 h

It is intriguing to discuss why substituents such as -NH₂, -OH, -OCH₃, -CH=CH₂, -SH, and -PH₂ can enhance the acid-catalyzed hydrolysis of ROOHs. As discussed above, a common feature for the acid-catalyzed hydrolysis of these substituted ROOHs is the formation of carbocation intermediates during their two-step reaction pathway. As illustrated in Fig. 4, the greater the enhancing potential of a substituent, the lower the reaction free energy required for carbocation intermediate formation, resulting in higher intermediate stability. This trend remains consistent when using the other computational method, despite differences in the absolute free energies (Fig. S13). Furthermore, carbocation hydration shows minimal impact on reaction energetics, as demonstrated by comparing results from the monohydrated system to those from the implicit solvation model alone, using the reaction of C(CH₃)₂(OCH₃)OOH as a test case (Fig. S14). These findings suggest that the stabilizing effect of these substituents on carbocation intermediates is the driving force behind the enhanced acid-catalyzed hydrolysis. Exceptions for the -OH- and -OCH₃-substituted ROOHs should be caused by the slight difference in their two-step reaction pathways (Fig. S7). According to electronic effect theory (Naredla and Klumpp, 2013; Olah, 2001), substituents -NH₂, -OH, -OCH₃, -SH, -CH=CH₂, and -PH₂ stabilize carbocation intermediates through conjugated electron donation into the unoccupied orbital of the carbocation. This stabilizing effect follows the order: -NH₂ > -OH ≈ -OCH₃ > -SH ≈ -CH=CH₂ ≈ -PH₂. This order aligns with the observed trend in *k_A* values, where α-NH₂-substituted ROOHs exhibit higher *k_A* values than α-OH- and -OCH₃-substituted ones, which in turn are higher than α-SH-, -CH=CH₂-, and -PH₂-substituted ones (Table S3). Similarly, the stabilizing effect of these substituents on carbocation intermediates can also explain the order of *k_A* values for tertiary ROOHs > secondary ROOHs > primary ROOHs. Tertiary ROOHs exhibit higher *k_A* values due to stronger hyperconjugation interactions between the unoccupied p orbital of the carbocation and the additional C-H σ-bonds from the methyl group, which stabilize the carbocation intermediates (Alamiddine and Humbel, 2013).




Figure 4. Calculated schematic free-energy surfaces for carbocation formation during acid-catalyzed hydrolysis of C(CH₃)₂(X)OOH. Substituents *X* denotes NH₂, OH, OCH₃, CH=CH₂, SH, and PH₂. Free energies are calculated at the M06-2X/6-311++G(3df,2pd)//M06-2X/6-31+G(d,p) level with the SMD solvation model, with ROOH and H⁺(H₂O)₂ set as the reference state (R) at 0 kcal mol⁻¹. Transition states (TS) and intermediates (IM) are labeled.

Based on the above analysis, we deduced that the presence of N and O atoms is the main reason for the strongest facilitating effect of -NH₂, -OH, and -OCH₃ in the acid-catalyzed hydrolysis of ROOHs. This raises an interesting question of how do substituents attached to N or O atoms affects the k'_A values. To explore this, we examined the effect of substituents attached to O atom as a test case and calculated the EF for k'_A of C(CH₃)₂(OY)OOH (Y being substituents) compared to the reference compound C(CH₃)₂OCH₃OOH. As shown in Fig. 3f, substituents -CH₂CH₃ and -CH₂CH₂CH₃ lead to only a minor reduction for k'_A . This is validated by the similar experimental k'_A of C₁₂ α-AH₍₁₎ and C₁₃ α-AH, which possess -OCH₂CH₃ and -OCH₂CH₂CH₃, respectively (Hu et al., 2022). Substituents -CH(CH₃)₂ and -C(CH₃)₃ increase the k'_A , with EF values of 10.6 and 45.9, respectively. However, -CH=CH₂ and -CHO substituents dramatically decrease k'_A by approximately 5 and 10 orders of magnitude, respectively. As shown in Table 1, the C(CH₃)₂(OY)OOH with Y being four alkyl substituents undergo rapid acid-catalyzed hydrolysis with $\tau_{1/e}$ values ranging from 3.5 s to 6.0 min even at pH 3.8, while C(CH₃)₂(OCH=CH₂)OOH only undergoes effective reaction at pH 0.9 ($\tau_{1/e}$ = 1.4 h). In contrast, the acid-catalyzed hydrolysis of C(CH₃)₂(OCHO)OOH is difficult to occur. These findings underscore that not all oxygen-containing substituents at the C_α site are equally effective in promoting acid-catalyzed hydrolysis of ROOHs, highlighting the importance of specific structural features. The substituent effects on the O atom could be reasonably extrapolated to the N atom. It should be noted that although the direct acid-catalyzed hydrolysis of C(CH₃)₂(OCHO)OOH is impossible, the indirect reaction initiated by the hydrolysis of the -OCHO group has been proposed (Chang et al., 2025; Zhao et al., 2018), which is beyond the range of this study.

For the β-substituted ROOHs, tertiary C(CH₃)₂(CH₂(X))OOH (X = -NH₂, -OH, -OCH₃, -CH=CH₂, -SH, -PH₂, -F, -Cl and -CHO) were chosen as model compounds, as they potentially exhibit higher k'_A values compared to primary and secondary ones. By calculating the EF for k'_A values of C(CH₃)₂(CH₂(X))OOH relative to the reference compound C(CH₃)₃OOH, we found that all nine substituents reduce the k'_A (Fig. 3e). Thus, we conclude that substitutes at C_β site could hinder the acid-catalyzed hydrolysis. The unfeasible acid-catalyzed hydrolysis of them aligns with previous experimental results that non-α-substituted monoterpene-derived organic peroxides exhibit greater persistence in aqueous environments (Zhao et al., 2022).

Based on the calculated log k_A values and the $\sum\sigma^*$ constants (Table S2), QSAR models were developed for α- and β-substituted ROOH model compounds. As shown in Fig. S15, a strong linear correlation was established for α-substituted compounds, including CH₂(X)OOH, CH(CH₃)(X)OOH, C(CH₃)₂(X)OOH, and C(CH₃)₂(OY)OOH, with the equation as $\log k_A = -11.555 \sum\sigma^* + 19.251$ ($n = 21$, $R^2 = 0.9083$) after excluding compounds with unknown $\sum\sigma^*$ and those with substituents -CH=CH₂ and -CHO. The poor performance in describing the effects of substituents -CH=CH₂ and -CHO is likely because $\sum\sigma^*$ primarily characterizes the inductive effects of substituents (Perrin et al., 1981), while its description of conjugation effects is inadequate. A relatively weaker fit to the QSAR plot ($\log k_A = -3.499 \sum\sigma^* - 6.437$ ($n = 9$, $R^2 = 0.7446$)) was found for compounds including C(CH₃)₃OOH and β-substituted compounds C(CH₃)₂(CH₂(X))OOH (Fig. S16). Nonetheless, a negative correlation was found in both QSAR models. This indicates that the weaker the inductive effects of substituents (i.e., the smaller the $\sum\sigma^*$), the higher the k_A , which is consistent with the trend explained by the stability order of the formed carbocation intermediates discussed earlier.

3.3 Acid-catalyzed hydrolysis of atmospheric ROOHs

280 To evaluate the applicability of the structure-activity relationship derived from model compounds to more complicated atmospheric ROOHs, we investigated the acid-catalyzed hydrolysis of eight ROOHs derived from isoprene (Wennberg et al., 2018), α -pinene (Zhang et al., 2017; Claflin et al., 2018), trimethylamine (Kjaergaard et al., 2023; Ma et al., 2021), and dimethyl sulphide (Berndt et al., 2019) using the screened $H^+(H_2O)_2$ model via the DFT calculations (Figs. S17-S24). These ROOHs feature distinct substituents at the C_α site, including $-N(CH_3)_2$, $-N(CH_3)(CHO)$, $-OH$, $-CH=CH_2$, $-SCH_3$, $-SCHO$, and
285 $-CHO$, and unsubstituted. As shown in Fig. 5, the order of k'_A values for these ROOHs at the same pH (3.8 or 0.9) according to their substituents is as follows: $-N(CH_3)_2 > -OH > -N(CH_3)(CHO) > \text{unsubstituted} > -CH=CH_2 > -SCH_3 > -CHO > -SCHO$, similar to the trends observed in model compounds in three aspects. First, a $-N(CH_3)_2$ or $-OH$ substituent at the C_α site of trimethylamine- $OOH_{(1)}$ or α -pinene- $OOH_{(1)}$ results in their high k'_A values, with $\tau_{1/e}$ values less than 2 d (pH 3.8) and 3.6 min (pH 0.9), respectively. Second, the introduction of a $-CHO$ group to the N atom in the $-N(CH_3)(CHO)$ substituent of
290 trimethylamine- $OOH_{(2)}$ leads to a remarkable reduction in k'_A values, with $\tau_{1/e}$ values exceeding 45 d, even at pH 0.9 (Table S4). The possible reaction of trimethylamine- $OOH_{(2)}$ initiated by the hydrolysis of $-N(CH_3)(CHO)$ group is not considered in this work (Zhang et al., 2015). Moreover, the attachment of a $-CHO$ group to the S atom in the $-SCHO$ substituent leads the corresponding dimethyl sulfide- $OOH_{(2)}$ to exhibit the lowest k'_A values, even lower than that of α -pinene- $OOH_{(3)}$ featuring a $-CHO$ substituent. Finally, the $-CH=CH_2$ and $-SCH_3$ substituents contribute to the relatively low k'_A values of isoprene- OOH
295 and dimethyl sulfide- OOH , respectively. Although the existence of enhancing substituents, their k'_A values is lower than the unsubstituted α -pinene- $OOH_{(2)}$, which is attributed to the reducing effect of an additional $-OH$ substituent at the C_β site for isoprene- OOH , and the primary C_α of sulfide- OOH compared to tertiary C_α of α -pinene- $OOH_{(2)}$. The reproduction of the substituents effect trend on k'_A values in atmospheric ROOHs demonstrates that the structure-activity relationship derived from model ROOHs can be effectively extended to predict the k_A values of structurally diverse ROOHs.




Figure 5. Calculated pseudo-first-order rate constants (k'_A) of eight atmospheric ROOHs under two selected pH values. pH 3.8 and pH 0.9 represent typical aerosol conditions in inland China and the Southeastern United States, respectively.

In the aerosol aqueous phase, anions such as nitrate (NO_3^-) and sulfate ions (SO_4^{2-}) are important components in addition to H_2O . These anions can either directly nucleophilically attack the C_α of ROOHs or react with the formed carbocation intermediates to produce organic nitrates and organic sulfates (Figs. S3-S6 and S17-S24). Understanding the competition between acid-catalyzed hydrolysis and esterification reactions of atmospheric ROOHs is crucial for elucidating their transformation products in ambient aerosols. Herein, we use C_{13} α -AH, C_{12} α -AH₍₁₎, C_{12} α -AH₍₂₎, and C_{10} α -HH as examples to elucidate the competition between their acid-catalyzed hydrolysis and esterification reactions. As shown in Figs. S3-S6, reactions of their derived carbocations with H_2O and NO_3^- are endothermic, while those with SO_4^{2-} are barrierless and exothermic. Although the formation of hemiacetals/geminal diols through reactions with H_2O is thermodynamically unfavorable, these species can subsequently be transformed into lactols (Qiu et al., 2020b; Enami, 2022; Hu et al., 2022). Therefore, competitive reactions with H_2O and SO_4^{2-} must be considered. If the concentrations of H_2O and SO_4^{2-} are equal, carbocations preferentially react with SO_4^{2-} at near diffusion-controlled rate constants ($\sim 10^{10} \text{ L mol}^{-1} \text{ s}^{-1}$) to form organic sulfates. In fact, the distribution of products, including hemiacetals/geminal diols, transformed lactols, and organic sulfates in aerosols, is determined by the second-order rate constants for carbocation reactions with H_2O and the variable concentrations of H_2O and SO_4^{2-} . The formation mechanisms of organic sulfates have been a hot research topic in atmospheric chemistry. To date, a variety of formation mechanisms have been proposed; among them, the acid-catalyzed ring-opening of epoxides has been widely adopted to explain organic sulfates formation in acidic sulfate aerosol (Surratt et al., 2010; Lin et al., 2012; Cooke et al., 2024). Nevertheless, existing mechanisms still cannot fully explain the observed ambient abundance of organic sulfates. Here we present a plausible carbocation-mediated organic sulfates formation mechanism that occurs during the aqueous-phase transformation of ROOHs in the presence of H^+ and SO_4^{2-} . However, our structure-activity relationship analysis indicates that,

among our studied ROOHs, only those with C_α substituents such as -OH, -OCH₃, -NH₂, and -N(CH₃)₂, can effectively generate the corresponding carbocation intermediates on the lifetime scale of aerosols under the considered pH conditions (pH 3.8 and 0.9). In the atmosphere, such ROOHs primarily derive from the reactions of water, alcohols, or ammonia with CIs produced via ozonolysis of unsaturated hydrocarbons or from ·OH oxidation of tertiary amines (Wang et al., 2023; Li et al., 2024; Kjaergaard et al., 2023). Therefore, we speculate that the proposed carbocation-mediated mechanism would be helpful to explain organic sulfates formation in SOA derived from the aforementioned reactions, whereas its applicability to other sources such as ·OH/NO₃· oxidation of volatile organic compounds (Surratt et al., 2008) could be limited unless the precursor compounds contain activating substituents that support the formation of the α-substituted ROOHs mentioned above.

4 Conclusions

This study demonstrates that combining the protonated water cluster H⁺(H₂O)₂ model with the M06-2X/6-311++G(3df,2pd)//M06-2X/6-31+G(d,p) method can accurately predict the acid-catalyzed hydrolysis kinetics of ROOHs. This approach provides a reliable framework for predicting the reaction kinetics of other atmospheric ROOHs. Furthermore, the study identifies new functional groups, including -NH₂, -N(CH₃)₂, -OH, -OCH₃, -CH=CH₂, -SH, and -PH₂, substituted at the C_α site of the -OOH group, which can enhance the acid-catalyzed hydrolysis kinetics of ROOHs. Notably, the newly identified -NH₂ and -N(CH₃)₂ substituents exhibit a greater enhancing effect than the well-documented -OH and -OCH₃ substituents. Contrary to the assumptions based on chemical intuition, it is clarified that not all nitrogen- or oxygen-containing substituents at the C_α site of the -OOH group can equally enhance acid-catalyzed hydrolysis of ROOHs, and the effects depend on the substituents attached to the O or N atoms. Importantly, the structure-activity relationship for commonly encountered substituents was elucidated for the acid-catalyzed hydrolysis kinetics of ROOHs, providing guidelines for qualitatively assessing the feasibility of acid-catalyzed hydrolysis of atmospheric ROOHs in aerosol water. The predicted kinetic data can be incorporated into 3-D chemical models to better simulate the atmospheric aqueous-phase chemistry of ROOHs. Meanwhile, we also emphasize that future work focusing on the standards synthesis of structurally diverse ROOHs would be very helpful for the laboratory validation of our findings.

This study also reveals that carbocation intermediates are formed during the acid-catalyzed hydrolysis of some ROOHs. The formed carbocation could further react with anions and organics, which are abundant in ambient aerosols besides water. Our preliminary studies indicate that the reaction of the formed carbocation with abundant SO₄²⁻ is barrierless and exothermic, leading to the formation of less-volatile organic sulfates. Meanwhile, these carbocations could also mediate the formation of oligomers and N-heterocycles according to the aqueous-phase carbocations chemistry of the α-dicarbonyls (Ji et al., 2020; Shi et al., 2024; Zhang et al., 2022). These findings emphasize the importance of investigating the impact of acid-catalyzed transformations of ROOHs on aerosol composition and properties. Additionally, the aqueous-phase transformations of atmospheric ROOHs with multiple hydrolysable functional groups should be further investigated.

Data availability. All data were available in the main text or supplementary materials. The other relevant data are available upon request from the corresponding authors.

355 **Author contributions.** HX and RY contributed to conceiving the idea, editing, and revision. QZ contributed to conceiving the idea, performing DFT calculations, analyzing results and interpreting data, and writing the original draft. FM, HZ, QX, XW, and JC contributed to editing and revision. All authors read and approved the final manuscript.

Competing interest. The authors declare no conflict of interest.

Financial support. This work was supported by the National Natural Science Foundation of China (22176022, 22406017,
360 22206020, 22236004).

References

- Abbatt, J. P. D., and Ravishankara, A. R.: Opinion: atmospheric multiphase chemistry - past, present, and future, *Atmos. Chem. Phys.*, 23, 9765-9785, 10.5194/acp-23-9765-2023, 2023.
- Agmon, N., Bakker, H. J., Campen, R. K., Henchman, R. H., Pohl, P., Roke, S., Thaemer, M., and Hassanali, A.: Protons and
365 hydroxide ions in aqueous systems, *Chem. Rev.*, 116, 7642-7672, 10.1021/acs.chemrev.5b00736, 2016.
- Alamiddine, Z., and Humbel, S.: Hyperconjugation in carbocations, a BLW Study with DFT approximation, *Front. Chem.*, 1,
37, 10.3389/fchem.2013.00037, 2013.
- Berndt, T., Scholz, W., Mentler, B., Fischer, L., Hoffmann, E. H., Tilgner, A., Hyttinen, N., Prisle, N. L., Hansel, A., and
Herrmann, H.: Fast peroxy radical isomerization and OH recycling in the reaction of OH radicals with dimethyl sulfide, *J.*
370 *Phys. Chem. Lett.*, 10, 6478-6483, 10.1021/acs.jpcllett.9b02567, 2019.
- Chang, C., Zang, H., Yao, M., Li, C., Li, Z., Wang, S., Huang, R., and Zhao, Y.: Rapid iron-mediated aqueous-phase reactions
of organic peroxides from monoterpene-derived Criegee intermediates and implications for aerosol and cloud chemistry,
Environ. Sci. Technol., 59, 2157-2168, 10.1021/acs.est.4c08340, 2025.
- Clafilin, M. S., Krechmer, J. E., Hu, W., Jimenez, J. L., and Ziemann, P. J.: Functional group composition of secondary organic
375 aerosol formed from ozonolysis of α -pinene under high VOC and autoxidation conditions, *ACS Earth Space Chem.*, 2,
1196-1210, 10.1021/acsearthspacechem.8b00117, 2018.
- Cooke, M. E., Armstrong, N. C., Fankhauser, A. M., Chen, Y., Lei, Z., Zhang, Y., Ledsy, I. R., Turpin, B. J., Zhang, Z., Gold,
A., McNeill, V. F., Surratt, J. D., and Ault, A. P.: Decreases in epoxide-driven secondary organic aerosol production under
highly acidic conditions: the importance of acid-base equilibria, *Environ. Sci. Technol.*, 58, 10675-10684,
380 10.1021/acs.est.3c10851, 2024.
- Dai, Y., Chen, Z., Qin, X., Dong, P., Xu, J., Hu, J., Gu, L., and Chen, S.: Hydrolysis reactivity reveals significant seasonal
variation in the composition of organic peroxides in ambient PM_{2.5}, *Sci. Total Environ.*, 927, 172143,
10.1016/j.scitotenv.2024.172143, 2024.
- Ding, J., Zhao, P., Su, J., Dong, Q., Du, X., and Zhang, Y.: Aerosol pH and its driving factors in Beijing, *Atmos. Chem. Phys.*,
385 19, 7939-7954, 10.5194/acp-19-7939-2019, 2019.
- Dovrou, E., Rivera-Rios, J. C., Bates, K. H., and Keutsch, F. N.: Sulfate formation via cloud processing from isoprene hydroxyl
hydroperoxides (ISOPOOH), *Environ. Sci. Technol.*, 53, 12476-12484, 10.1021/acs.est.9b04645, 2019.
- Enami, S.: Fates of organic hydroperoxides in atmospheric condensed phases, *J. Phys. Chem. A*, 125, 4513-4523,
10.1021/acs.jpca.1c01513, 2021.
- 390 Enami, S.: Proton-catalyzed decomposition of multifunctionalized organic hydroperoxides derived from the reactions of
criegee intermediates with ethylene glycol in aqueous organic media, *ACS Earth Space Chem.*, 6, 1937-1947,

- 10.1021/acsearthspacechem.2c00142, 2022.
- Enami, S., Hoffmann, M. R., and Colussi, A. J.: Proton availability at the air/water interface, *J. Phys. Chem. Lett.*, 1, 1599-1604, 10.1021/jz100322w, 2010.
- 395 Endo, Y., Sakamoto, Y., Kajii, Y., and Enami, S.: Decomposition of multifunctionalized α -alkoxyalkyl-hydroperoxides derived from the reactions of Criegee intermediates with diols in liquid phases, *Phys. Chem. Chem. Phys.*, 24, 11562-11572, 10.1039/d2cp00915c, 2022.
- Epstein, S. A., Blair, S. L., and Nizkorodov, S. A.: Direct photolysis of α -pinene ozonolysis secondary organic aerosol: effect on particle mass and peroxide content, *Environ. Sci. Technol.*, 48, 11251-11258, 10.1021/es502350u, 2014.
- 400 Ervens, B., Rickard, A., Aumont, B., Carter, W. P. L., McGillen, M., Mellouki, A., Orlando, J., Picquet-Varraut, B., Seakins, P., Stockwell, W. R., Vereecken, L., and Wallington, T. J.: Opinion: challenges and needs of tropospheric chemical mechanism development, *Atmos. Chem. Phys.*, 24, 13317-13339, 10.5194/acp-24-13317-2024, 2024.
- Frisch, M. J. T. G., Schlegel, H. B., Scuseria, G. E., Robb, M. A., Cheeseman, J. R., Scalmani, G., Barone, V., Mennucci, B., Petersson, G. A., Nakatsuji, H., Caricato, M., Li, X., Hratchian, H. P., Izmaylov, A. F., Bloino, J., Zheng, G., Sonnenberg, J. L., Hada, M., Ehara, M., Toyota, K., Fukuda, R., Hasegawa, J., Ishida, M., Nakajima, T., Honda, Y., Kitao, O., Nakai, H., Vreven, T., Montgomery, J. A. J., Peralta, J. E., Ogliaro, F., Bearpark, M., Heyd, J. J., Brothers, E., Kudin, K. N., Staroverov, V. N., Kobayashi, R., Normand, J., Raghavachari, K., Rendell, A., Burant, J. C., Iyengar, S. S., Tomasi, J., Cossi, M., Rega, N., Millam, J. M., Klene, M., Knox, J. E., Cross, J. B., Bakken, V., Adamo, C., Jaramillo, J., Gomperts, R., Stratmann, R. E., Yazyev, O., Austin, A. J., Cammi, R., Pomelli, C., Ochterski, J. W., Martin, R. L., Morokuma, K., Zakrzewski, V. G., Voth, G. A., Salvador, P., Dannenberg, J. J., Dapprich, S., Daniels, A. D., Farkas, O., Foresman, J. B., Ortiz, J. V., Cioslowski, J., and Fox, D. J.: *Gaussian 09*, Gaussian, Inc.: Wallingford, CT., 2009.
- Fukui, K.: The path of chemical reactions - the IRC approach, *Acc. Chem. Res.*, 14, 363-368, DOI 10.1021/ar00072a001, 1981.
- Guo, H., Xu, L., Bougiatioti, A., Cerully, K. M., Capps, S. L., Hite, J. R., Carlton, A. G., Lee, S. H., Bergin, M. H., Ng, N. L., Nenes, A., and Weber, R. J.: Fine-particle water and pH in the southeastern United States, *Atmos. Chem. Phys.*, 15, 5211-5228, 10.5194/acp-15-5211-2015, 2015.
- 415 Herrmann, H., Schaefer, T., Tilgner, A., Styler, S. A., Weller, C., Teich, M., and Otto, T.: Tropospheric aqueous-phase chemistry: kinetics, mechanisms, and its coupling to a changing gas phase, *Chem. Rev.*, 115, 4259-4334, 10.1021/cr500447k, 2015.
- Hodzic, A., Kasibhatla, P. S., Jo, D. S., Cappa, C. D., Jimenez, J. L., Madronich, S., and Park, R. J.: Rethinking the global secondary organic aerosol (SOA) budget: stronger production, faster removal, shorter lifetime, *Atmos. Chem. Phys.*, 16, 7917-7941, 10.5194/acp-16-7917-2016, 2016.
- Hu, M., Qiu, J., Tonokura, K., and Enami, S.: Aqueous-phase fates of α -alkoxyalkyl-hydroperoxides derived from the reactions of Criegee intermediates with alcohols, *Phys. Chem. Chem. Phys.*, 23, 4605-4614, 10.1039/d0cp06308h, 2021a.
- Hu, M., Tonokura, K., Morino, Y., Sato, K., and Enami, S.: Effects of metal ions on aqueous-phase decomposition of α -hydroxyalkyl-hydroperoxides derived from terpene alcohols, *Environ. Sci. Technol.*, 55, 12893-12901, 10.1021/acs.est.1c04635, 2021b.
- 425 Hu, M., Chen, K., Qiu, J., Lin, Y., Tonokura, K., and Enami, S.: Decomposition mechanism of α -alkoxyalkyl-hydroperoxides in the liquid phase: temperature dependent kinetics and theoretical calculations, *Environ Sci Atmos*, 2, 241-251, 10.1039/d1ea00076d, 2022.
- 430 Ji, Y., Shi, Q., Li, Y., An, T., Zheng, J., Peng, J., Gao, Y., Chen, J., Li, G., Wang, Y., Zhang, F., Zhang, A. L., Zhao, J., Molina, M. J., and Zhang, R.: Carbenium ion-mediated oligomerization of methylglyoxal for secondary organic aerosol formation, *Proc. Natl. Acad. Sci. U. S. A.*, 117, 13294-13299, 10.1073/pnas.1912235117, 2020.
- Jia, S., Wang, X., Zhang, Q., Sarkar, S., Wu, L., Huang, M., Zhang, J., and Yang, L.: Technical note: comparison and interconversion of pH based on different standard states for aerosol acidity characterization, *Atmos. Chem. Phys.*, 18, 11125-11133, 10.5194/acp-18-11125-2018, 2018.
- 435 Jin, X., Wang, Y., Li, Z., Zhang, F., Xu, W., Sun, Y., Fan, X., Chen, G., Wu, H., Ren, J., Wang, Q., and Cribb, M.: Significant contribution of organics to aerosol liquid water content in winter in Beijing, China, *Atmos. Chem. Phys.*, 20, 901-914, 10.5194/acp-20-901-2020, 2020.
- Kim, J., and Huang, C. H.: Reactivity of peracetic acid with organic compounds: a critical review, *ACS ES&T Water*, 1, 15-33, 10.1021/acsestwater.0c00029, 2021.
- 440 Kjaergaard, E. R., Moller, K. H., Berndt, T., and Kjaergaard, H. G.: Highly efficient autoxidation of triethylamine, *J. Phys.*

- Chem. A, 127, 8623-8632, 10.1021/acs.jpca.3c04341, 2023.
- Krapf, M., Haddad, I. E., Bruns, E. A., Molteni, U., Daellenbach, K. R., Prevot, A. S. H., Baltensperger, U., and Dommen, J.: Labile peroxides in secondary organic aerosol, *Chem*, 1, 603-616, 10.1016/j.chempr.2016.09.007, 2016.
- 445 Kristiansen, N. I., Stohl, A., Olivie, D. J. L., Croft, B., Sovde, O. A., Klein, H., Christoudias, T., Kunkel, D., Leadbetter, S. J., Lee, Y. H., Zhang, K., Tsigaridis, K., Bergman, T., Evangeliou, N., Wang, H., Ma, P. L., Easter, R. C., Rasch, P. J., Liu, X., Pitari, G., Di Genova, G., Zhao, S. Y., Balkanski, Y., Bauer, S. E., Faluvegi, G. S., Kokkola, H., Martin, R. V., Pierce, J. R., Schulz, M., Shindell, D., Tost, H., and Zhang, H.: Evaluation of observed and modelled aerosol lifetimes using radioactive tracers of opportunity and an ensemble of 19 global models, *Atmos. Chem. Phys.*, 16, 3525-3561, 10.5194/acp-16-3525-2016, 2016.
- 450 Lee, Y., and von Gunten, U.: Quantitative structure-activity relationships (QSARs) for the transformation of organic micropollutants during oxidative water treatment, *Water Res.*, 46, 6177-6195, 10.1016/j.watres.2012.06.006, 2012.
- Lei, Z., Chen, Y., Zhang, Y., Cooke, M. E., Ledsy, I. R., Armstrong, N. C., Olson, N. E., Zhang, Z., Gold, A., Surratt, J. D., and Ault, A. P.: Initial pH governs secondary organic aerosol phase state and morphology after uptake of isoprene epoxydiols (IEPOX), *Environ. Sci. Technol.*, 56, 10596-10607, 10.1021/acs.est.2c01579, 2022.
- 455 Li, X., Jia, L., Xu, Y., and Pan, Y.: A novel reaction between ammonia and Criegee intermediates can form amines and suppress oligomers from isoprene, *Sci. Total Environ.*, 956, 177389, 10.1016/j.scitotenv.2024.177389, 2024.
- Lin, P., Yu, J. Z., Engling, G., and Kalberer, M.: Organosulfates in humic-like substance fraction isolated from aerosols at seven locations in East Asia: a study by ultra-high-resolution mass spectrometry, *Environ. Sci. Technol.*, 46, 13118-13127, 10.1021/es303570v, 2012.
- 460 Liu, X., Wang, H., Wang, F., Lv, S., Wu, C., Zhao, Y., Zhang, S., Liu, S., Xu, X., Lei, Y., and Wang, G.: Secondary formation of atmospheric brown carbon in China haze: implication for an enhancing role of ammonia, *Environ. Sci. Technol.*, 57, 11163-11172, 10.1021/acs.est.3c03948, 2023.
- Lu, T.: Molclus program, version 1.9.9.9, <http://www.keinsci.com/research/molclus.html>, 2022.
- 465 Ma, F., Xie, H., Li, M., Wang, S., Zhang, R., and Chen, J.: Autoxidation mechanism for atmospheric oxidation of tertiary amines: implications for secondary organic aerosol formation, *Chemosphere*, 273, 129207, 10.1016/j.chemosphere.2020.129207, 2021.
- Marenich, A. V., Cramer, C. J., and Truhlar, D. G.: Universal solvation model based on solute electron density and on a continuum model of the solvent defined by the bulk dielectric constant and atomic surface tensions, *J. Phys. Chem. B*, 113, 6378-6396, 10.1021/jp810292n, 2009.
- 470 McNeill, V. F.: Aqueous organic chemistry in the atmosphere: sources and chemical processing of organic aerosols, *Environ. Sci. Technol.*, 49, 1237-1244, 10.1021/es5043707, 2015.
- Naredla, R. R., and Klumpp, D. A.: Contemporary carbocation chemistry: applications in organic synthesis, *Chem. Rev.*, 113, 6905-6948, 10.1021/cr4001385, 2013.
- 475 Olah, G. A.: 100 years of carbocations and their significance in chemistry, *J. Org. Chem.*, 66, 5943-5957, 10.1021/jo010438x, 2001.
- Perrin, D. D., Dempsey, B., and Serjeant, E. P.: pK_a prediction for organic acids and bases, Chapman and Hall: New York, 1981.
- Piletic, I. R., Edney, E. O., and Bartolotti, L. J.: A computational study of acid catalyzed aerosol reactions of atmospherically relevant epoxides, *Phys. Chem. Chem. Phys.*, 15, 18065-18076, 10.1039/c3cp52851k, 2013.
- 480 Qiu, J., Ishizuka, S., Tonokura, K., Colussi, A. J., and Enami, S.: Water dramatically accelerates the decomposition of α -hydroxyalkyl-hydroperoxides in aerosol particles, *J. Phys. Chem. Lett.*, 10, 5748-5755, 10.1021/acs.jpcllett.9b01953, 2019.
- Qiu, J., Liang, Z., Tonokura, K., Colussi, A. J., and Enami, S.: Stability of Monoterpene-derived α -hydroxyalkyl-hydroperoxides in aqueous organic media: relevance to the fate of hydroperoxides in aerosol particle phases, *Environ. Sci. Technol.*, 54, 3890-3899, 10.1021/acs.est.9b07497, 2020a.
- 485 Qiu, J., Tonokura, K., and Enami, S.: Proton-catalyzed decomposition of α -hydroxyalkyl-hydroperoxides in water, *Environ. Sci. Technol.*, 54, 10561-10569, 10.1021/acs.est.0c03438, 2020b.
- Ra, J., Tolu, J., Rentsch, D., Manasfi, T., and von Gunten, U.: Unveiling the reaction chemistry of sulfoxides during water chlorination, *Water Res.*, 270, 122806, 10.1016/j.watres.2024.122806, 2025.
- 490 Ryu, H., Park, J., Kim, H. K., Park, J. Y., Kim, S., and Baik, M.: Pitfalls in computational modeling of chemical reactions and how to avoid them, *Organometallics*, 37, 3228-3239, 10.1021/acs.organomet.8b00456, 2018.

- Sadlej-Sosnowska, N.: Calculation of acidic dissociation constants in water: solvation free energy terms. Their accuracy and impact, *Theor. Chem. Acc.*, 118, 281-293, 10.1007/s00214-006-0232-z, 2007.
- 495 Shi, Q., Gao, L., Li, W., Wang, J., Shi, Z., Li, Y., Chen, J., Ji, Y., and An, T.: Oligomerization mechanism of methylglyoxal regulated by the methyl groups in reduced nitrogen species: Implications for brown carbon formation, *Environ. Sci. Technol.*, 58, 1563-1576, 10.1021/acs.est.3c05983, 2024.
- Shi, R., Zhang, F., Shen, Y., Shen, J., Xu, B., Kuang, B., Xu, Z., Jin, L., Tang, Q., Tian, X., and Wang, Z.: Aerosol liquid water in PM_{2.5} and its roles in secondary aerosol formation at a regional site of Yangtze River Delta, *J. Environ. Sci.*, 138, 684-696, 2024.
- 500 Shi, X., Nenes, A., Xiao, Z., Song, S., Yu, H., Shi, G., Zhao, Q., Chen, K., Feng, Y., and Russell, A. G.: High-resolution data sets unravel the effects of sources and meteorological conditions on nitrate and its gas-particle partitioning, *Environ. Sci. Technol.*, 53, 3048-3057, 10.1021/acs.est.8b06524, 2019.
- Su, J., Zhao, P., Ge, S., and Ding, J.: Aerosol liquid water content of PM_{2.5} and its influencing factors in Beijing, China, *Sci. Total Environ.*, 839, 10, 10.1016/j.scitotenv.2022.156342, 2022.
- 505 Surratt, J. D., Gómez-González, Y., Chan, A. W. H., Vermeylen, R., Shahgholi, M., Kleindienst, T. E., Edney, E. O., Offenberg, J. H., Lewandowski, M., Jaoui, M., Maenhaut, W., Claeys, M., Flagan, R. C., and Seinfeld, J. H.: Organosulfate formation in biogenic secondary organic aerosol, *J. Phys. Chem. A*, 112, 8345-8378, 10.1021/jp802310p, 2008.
- Surratt, J. D., Chan, A. W. H., Eddingsaas, N. C., Chan, M., Loza, C. L., Kwan, A. J., Hersey, S. P., Flagan, R. C., Wennberg, P. O., and Seinfeld, J. H.: Reactive intermediates revealed in secondary organic aerosol formation from isoprene, *Proc. Natl. Acad. Sci. U. S. A.*, 107, 6640-6645, 10.1073/pnas.0911114107, 2010.
- 510 Wang, G., Tao, Y., Chen, J., Liu, C., Qin, X., Li, H., Yun, L., Zhang, M., Zheng, H., Gui, H., Liu, J., Huo, J., Fu, Q., Deng, C., and Huang, K.: Quantitative decomposition of influencing factors to aerosol pH variation over the coasts of the South China Sea, East China Sea, and Bohai Sea, *Environ. Sci. Technol. Lett.*, 9, 815-821, 10.1021/acs.estlett.2c00527, 2022.
- Wang, S., Wang, L., Li, Y., Wang, C., Wang, W., Yin, S., and Zhang, R.: Effect of ammonia on fine-particle pH in agricultural regions of China: comparison between urban and rural sites, *Atmos. Chem. Phys.*, 20, 2719-2734, 10.5194/acp-20-2719-2020, 2020.
- 515 Wang, S., Zhao, Y., Chan, A. W. H., Yao, M., Chen, Z., and Abbatt, J. P. D.: Organic peroxides in aerosol: key reactive intermediates for multiphase processes in the atmosphere, *Chem. Rev.*, 123, 1635-1679, 10.1021/acs.chemrev.2c00430, 2023.
- 520 Wang, S., Zhou, S., Tao, Y., Tsui, W. G., Ye, J., Yu, J. Z., Murphy, J. G., McNeill, V. F., Abbatt, J. P. D., and Chan, A. W. H.: Organic peroxides and sulfur dioxide in aerosol: source of particulate sulfate, *Environ. Sci. Technol.*, 53, 10695-10704, 10.1021/acs.est.9b02591, 2019.
- Wei, J., Fang, T., Wong, C., Lakey, P. S. J., Nizkorodov, S. A., and Shiraiwa, M.: Superoxide formation from aqueous reactions of biogenic secondary organic aerosols, *Environ. Sci. Technol.*, 55, 260-270, 10.1021/acs.est.0c07789, 2021.
- 525 Wei, J., Fang, T., Lakey, P. S. J., and Shiraiwa, M.: Iron-facilitated organic radical formation from secondary organic aerosols in surrogate lung fluid, *Environ. Sci. Technol.*, 56, 7234-7243, 10.1021/acs.est.1c04334, 2022a.
- Wei, J., Fang, T., and Shiraiwa, M.: Effects of acidity on reactive oxygen species formation from secondary organic aerosols, *ACS Environ. Au*, 2, 336-345, 10.1021/acsenvironau.2c00018, 2022b.
- 530 Wennberg, P. O., Bates, K. H., Crounse, J. D., Dodson, L. G., McVay, R. C., Mertens, L. A., Nguyen, T. B., Praske, E., Schwantes, R. H., Smarte, M. D., St Clair, J. M., Teng, A. P., Zhang, X., and Seinfeld, J. H.: Gas-phase reactions of isoprene and its major oxidation products, *Chem. Rev.*, 118, 3337-3390, 10.1021/acs.chemrev.7b00439, 2018.
- Wieser, F., Sander, R., Cho, C., Fuchs, H., Hohaus, T., Novelli, A., Tillmann, R., and Taraborrelli, D.: Development of a multiphase chemical mechanism to improve secondary organic aerosol formation in CAABA/MECCA (version 4.7.0), *Geosci. Model Dev.*, 17, 4311-4330, 10.5194/gmd-17-4311-2024, 2024.
- 535 Wu, Z., Wang, Y., Tan, T., Zhu, Y., Li, M., Shang, D., Wang, H., Lu, K., Guo, S., Zeng, L., and Zhang, Y.: Aerosol liquid water driven by anthropogenic inorganic salts: implying its key role in haze formation over the North China Plain, *Environ. Sci. Technol. Lett.*, 5, 160-166, 10.1021/acs.estlett.8b00021, 2018.
- Xu, J., Chen, J., Zhao, N., Wang, G., Yu, G., Li, H., Huo, J., Lin, Y., Fu, Q., Guo, H., Deng, C., Lee, S., Chen, J., and Huang, K.: Importance of gas-particle partitioning of ammonia in haze formation in the rural agricultural environment, *Atmos. Chem. Phys.*, 20, 7259-7269, 10.5194/acp-20-7259-2020, 2020.
- 540 Xu, T., Chen, J., Wang, Z., Tang, W., Xia, D., Fu, Z., and Xie, H.: Development of prediction models on base-catalyzed

- hydrolysis kinetics of phthalate esters with density functional theory calculation, *Environ. Sci. Technol.*, 53, 5828-5837, 10.1021/acs.est.9b00574, 2019.
- 545 Zhang, H., Xie, H., Chen, J., and Zhang, S.: Prediction of hydrolysis pathways and kinetics for antibiotics under environmental pH conditions: a quantum chemical study on cephradine, *Environ. Sci. Technol.*, 49, 1552-1558, 10.1021/es505383b, 2015.
- Zhang, J., Shrivastava, M., Ma, L., Jiang, W., Anastasio, C., Zhang, Q., and Zelenyuk, A.: Modeling novel aqueous particle and cloud chemistry processes of biomass burning phenols and their potential to form secondary organic aerosols, *Environ. Sci. Technol.*, 58, 3776-3786, 10.1021/acs.est.3c07762, 2024.
- 550 Zhang, X., Lambe, A. T., Upshur, M. A., Brooks, W. A., Gray Be, A., Thomson, R. J., Geiger, F. M., Surratt, J. D., Zhang, Z., Gold, A., Graf, S., Cubison, M. J., Groessl, M., Jayne, J. T., Worsnop, D. R., and Canagaratna, M. R.: Highly oxygenated multifunctional compounds in α -pinene secondary organic aerosol, *Environ. Sci. Technol.*, 51, 5932-5940, 10.1021/acs.est.6b06588, 2017.
- 555 Zhang, Y., He, L., Sun, X., Ventura, O. N., and Herrmann, H.: Theoretical investigation on the oligomerization of methylglyoxal and glyoxal in aqueous atmospheric aerosol particles, *ACS Earth Space Chem.*, 6, 1031-1043, 10.1021/acsearthspacechem.1c00422, 2022.
- Zhao, Q., Xie, H., Ma, F., Nie, W., Yan, C., Huang, D., Elm, J., and Chen, J.: Mechanism-based structure-activity relationship investigation on hydrolysis kinetics of atmospheric organic nitrates, *npj Clim. Atmos. Sci.*, 6, 10, 10.1038/s41612-023-00517-w, 2023.
- 560 Zhao, R., Kenseth, C. M., Huang, Y., Dalleska, N. F., Kuang, X. M., Chen, J., Paulson, S. E., and Seinfeld, J. H.: Rapid aqueous-phase hydrolysis of ester hydroperoxides arising from Criegee intermediates and organic acids, *J. Phys. Chem. A*, 122, 5190-5201, 10.1021/acs.jpca.8b02195, 2018.
- Zhao, Y., Liu, Z., Yao, M., Wang, S., Li, Z., Li, C., and Xiao, H.: Persistence of monoterpene-derived organic peroxides in the atmospheric aqueous phase, *ACS Earth Space Chem.*, 6, 2226-2235, 10.1021/acsearthspacechem.2c00145, 2022.
- 565 Zhao, Y., and Truhlar, D. G.: The M06 suite of density functionals for main group thermochemistry, thermochemical kinetics, noncovalent interactions, excited states, and transition elements: two new functionals and systematic testing of four M06-class functionals and 12 other functionals, *Theor. Chem. Acc.*, 120, 215-241, 10.1007/s00214-007-0310-x, 2008.
- Zheng, Y., Chen, Q., Cheng, X., Mohr, C., Cai, J., Huang, W., Shrivastava, M., Ye, P., Fu, P., Shi, X., Ge, Y., Liao, K., Miao, R., Qiu, X., Koenig, T. K., and Chen, S.: Precursors and pathways leading to enhanced secondary organic aerosol formation during severe haze episodes, *Environ. Sci. Technol.*, 55, 15680-15693, 10.1021/acs.est.1c04255, 2021.