

Improved management increases soil mineral-protected organic carbon storage via plant-microbial-nutrient mediation in semi-arid grasslands

Alejandro Carrascosa^a, Gerardo Moreno^a, M. Francesca Cotrufo^b, Cristina Frade^c, Sara Rodrigo^a,
V́ctor Rolo^a

^a Forest Research Group, INDEHESA, University of Extremadura, 10600, Plasencia, Cáceres, Spain

^b Department of Soil and Crop Sciences, Colorado State University, Fort Collins, CO, USA

^c Institute of Natural Resources and Agrobiology of Salamanca (IRNASA-CSIC), Salamanca, Spain

Correspondence to: V́ctor Rolo (rolo@unex.es)

Abstract

Soil organic carbon (SOC) storage in semi-arid grasslands is threatened by both climate change and land degradation, impacting food production and climate regulation. Improved management has been proposed to increase SOC stocks and overcome these challenges. However, the benefits of improved management practices in semi-arid regions are in question. Little is known about the effects of management on the functional components of SOC, particulate (POC) and mineral-associated organic carbon (MAOC), which are expected to respond differently, and about the pathways that mediate these responses, such as changes in vegetation and soil microbial communities.

This work analyses the effect of rotational grazing, legume sowing and grazing exclusion on topsoil (0-8 cm) SOC, POC and MAOC stocks in Mediterranean wooded grasslands compared to continuous conventional grazing. Changes in plant diversity and morpho-biochemical traits, soil fertility and microbial composition were also evaluated. A total of 188 plots were sampled in 9 farms across a wide environmental gradient.

More resource-acquisitive, nitrogen-rich and less lignified plant community, higher soil microbial biomass with lower Gram+/Gram- ratio, and higher soil fertility were associated with higher SOC storage, with similar impacts on POC and MAOC. Rotational grazing increased POC, MAOC and total SOC stocks by 15%, 15% and 14% respectively, compared to continuous grazing. This effect was mediated by an increase in soil fertility in the rotationally grazed paddocks. On the other hand, grazing exclusion reduced POC stocks by 12% compared to continuous grazing. This depletion was mainly due to a reduction in microbial biomass and litter quality of vegetation in non-grazed paddocks. Both POC and MAOC stocks were lower at the warmer sites.

We conclude that rotational grazing can enhance long-term SOC storage in semi-arid grasslands, thereby increasing their resilience and climate mitigation capacity, whereas abandoning grazing could lead to SOC losses.

Keywords

Soil organic matter fractions; Soil microbial community; Plant traits; Rotational grazing; Legume ~~sowing-enrichment~~; Grazing exclusion; Functional diversity; Mediterranean grassland

1. Introduction

Grasslands cover 40.5% of the world's ice-free land area and store one third of the terrestrial carbon (C) stocks, mainly (over 80%) in the form of soil organic matter (SOM) (White, 2000). This C pool in grassland soils surpasses the global aboveground vegetation C pool (IPCC, 2023a), highlighting the importance of grasslands SOM in global climate regulation (Bai and Cotrufo, 2022). In addition, SOM is a cornerstone of grassland productivity and functioning (Cotrufo and Lavalley, 2022; Tiessen et al., 1994). However, C storage in grassland soils is threatened by increasing C outputs (i.e., soil respiration and erosion) and decreasing C inputs (i.e., primary productivity) due to anthropogenic impacts such as land degradation and climate change (Crowther et al., 2016; Gang et al., 2014; Godde et al., 2020; Lei et al., 2021). Improved grassland management offers a key opportunity to counter these threats by increasing soil organic C (SOC) stocks and supporting both climate change mitigation and adaptation (Bai and Cotrufo, 2022; Conant, 2012; Dondini et al., 2023; Stanley et al., 2024). This is particularly valuable in the case of semi-arid grasslands, which represent the majority of the global grassland area (White, 2000) but are also more vulnerable to climate change impacts than wetter grasslands (Smith et al., 2024).

Practices such as rotational grazing, legumes sowing and grazing exclusion have been proposed to increase and conserve SOC stocks (Conant et al., 2017; Yu et al., 2021). Rotational grazing encompasses a variety of practices where small paddocks are grazed at high intensity for short periods of time, allowing for longer pasture rest than continuous grazing (Teague et al., 2013). This management favors vegetation recovery after defoliation and reduces grazing patchiness and livestock selectivity (Jacobo et al., 2006; Teague et al., 2013). The effects of rotational grazing on grassland productivity and animal performance are under debate (Briske et al., 2008; Teague et al., 2013; di Virgilio et al., 2019) but most studies have found positive effects on SOC stocks associated with this practice (Byrnes et al., 2018; Conant et al., 2017; Phukubye et al., 2022; Teague et al., 2011). Sowing legumes in natural pastures is practiced worldwide, having clear positive impacts on SOC stocks (Carranca et al., 2022; Conant et al., 2017; Moreno et al., 2021) and grassland productivity (Bartholomew and Williams, 2010; Carrascosa et al., 2024; Jaurena et al., 2016; Khatiwada et al., 2020; Rama et al., 2022). Grazing exclusion is widely advocated as a tool for ecosystem restoration (Cheng et al., 2016; Novelly and Watson, 2007) and is also a global trend driven by land abandonment, particularly in high-income countries (Li and Li, 2017). The effects of grazing exclusion on SOC in grasslands are mixed, showing both positive (Cheng et al., 2016; Yu et al., 2021) and negative outcomes (Wilson et al., 2018). Moreover, the net effect of management practices has been shown to depend on the environmental context (Maestre et al., 2022; McSherry and Ritchie, 2013; Niu et al., 2025), and global meta-analyses remain inconclusive. While some global studies have reported greater benefits on SOC stocks from grazing exclusion and rotational grazing in arid climates (Zhou et al., 2017), others found these management practices to be more beneficial in wetter climates (Byrnes et al., 2018; Hawkins, 2017; Zhou et al., 2019). To the best of our knowledge, no similar studies have addressed the interaction between environmental conditions and the effect of legume sowings on SOC stocks. Thus, some questions remain open and further research is needed to clarify the net effects of improved management practices on SOC storage under different environmental conditions and particularly in semi-arid grasslands.

To better understand the controls and vulnerability of C in soils, SOM can be conceptualized into particulate organic matter (POM) and mineral-associated organic matter (MAOM) fractions (Cotrufo and Lavalley, 2022; Lavalley et al., 2020). POM originates from fragmented structural plant inputs and, to a lesser extent, microbial recalcitrant compounds (Angst et al., 2021; Six et al., 2001). In contrast, MAOM forms through the sorption of microbial necromass and soluble

76 plant inputs onto soil mineral surfaces (Angst et al., 2021; Cotrufo et al., 2022). The mineral bonds partially protect
 77 MAOM from decomposition (Baldock and Skjemstad, 2000), while POM is readily accessible to microbial degradation,
 78 although occlusion within soil aggregates can reduce its accessibility (Angst et al., 2017). Hence, POM accumulation is
 79 mainly controlled by environmental constraints on microbial activity, e.g. low temperatures and highly acidic soils
 80 (Hansen et al., 2024; Yu et al., 2022; Zhou et al., 2024). For the same reasons, POM is more vulnerable to climatic
 81 warming (Benbi et al., 2014; Georgiou et al., 2024; Rocci et al., 2021) and the mean residence time of C in POM (years
 82 to decades) is on average shorter than in MAOM (decades to centuries; Zhou et al., 2024). However, MAOM storage in
 83 soil is theoretically limited by the availability of free mineral surface area (i.e. clay and silt content; Six et al., 2002), and
 84 a saturation point can be observed, where no more MAOM accumulate despite increases in total SOM contents (Cotrufo
 85 et al., 2019a; Georgiou et al., 2022). Management changes have been shown to influence SOM fractions in grassland soils
 86 (Khatri-Chhetri et al., 2024; Mosier et al., 2021; Oliveira Filho et al., 2019), and a conceptual framework for relating
 87 grazing management to SOM distribution has recently been proposed (Stanley et al., 2024). Yet, the underlying processes
 88 mediating these effects, such as alterations in vegetation or soil microbial communities (Laliberté and Tylianakis, 2012;
 89 Peco et al., 2017; Wilson et al., 2018), remain poorly understood.

90 Primary productivity is the point of entry of C into soil, and consequently the amount of plant inputs regulates SOM
 91 accrual (King et al., 2023; Zhou et al., 2024), but microbial processing largely determines the fate of that C (Cotrufo and
 92 Lavallee, 2022; Crowther et al., 2019). In this sense, the chemical composition of plant inputs, and the soil microbiota
 93 carbon use efficiency (CUE), i.e. the amount of C used for microbial growth and products relative to total C uptake, play
 94 a crucial role in the SOM formation process (Cotrufo and Lavallee, 2022; Tao et al., 2023). Recalcitrant plant inputs [i.e.,
 95 high carbon-nitrogen ratio (C/N) and lignin content] tend to promote short-term SOM accumulation, primarily as POM,
 96 due to their chemical resistance to decomposition (Cheng et al., 2023; Cotrufo and Lavallee, 2022). However, as outlined
 97 in the Microbial Efficiency-Matrix Stabilization (MEMS) framework (Cotrufo et al., 2013), recalcitrant inputs are less
 98 efficiently decomposed by microbes, leading to greater C losses in the long term, compared to labile (i.e., water-soluble,
 99 low C/N and lignin content) plant inputs (Cotrufo and Lavallee, 2022; Ridgeway et al., 2022). Thus, labile plant inputs
 100 are expected to enhance MAOM formation and SOM stocks in the long term, due to their faster and more efficient
 101 decomposition (Cheng et al., 2023; Haddix et al., 2016). Elias et al. (2024) added complexity to these assumptions,
 102 showing that plant input characteristics may favor certain microbial groups over others, altering the overall CUE of the
 103 microbial community. For example, fungi, which are often assumed to have a higher CUE than bacteria (Kallenbach et
 104 al., 2016; Strickland and Rousk, 2010), are favored by the addition of recalcitrant inputs (Bai et al., 2024; Strickland and
 105 Rousk, 2010). Substrate preferences have been also identify for Gram-positive (Gram+) and Gram-negative (Gram-)
 106 bacteria (Fanin et al., 2019; Kramer and Gleixner, 2008), with consequences for SOC accrual (Klump et al., 2009).
 107 Importantly, much of the research on the influence of plant input characteristics and microbial communities on SOM
 108 formation dynamics has relied on incubation experiments (Cheng et al., 2023; Haddix et al., 2016; Ridgeway et al., 2022)
 109 and there is limited information on how these findings translate to natural field conditions. Other vegetation characteristics
 110 such as species richness have been shown to positively influence SOC stocks (Lange et al., 2015; Steinbeiss et al., 2008),
 111 but its effects on SOC fractions have been poorly evaluated, with inconclusive results in grasslands (Mortensen et al.,
 112 2025). In addition, the relationships between SOM stocks and fractions and plant functional traits have rarely been studied
 113 (Manning et al., 2015; Mortensen et al., 2025; Xu et al., 2021) despite the latter being widely used to predict ecosystem
 114 functioning and responses (Funk et al., 2017). Plant functional traits are highly correlated with processes such as litter
 115 decomposition (Cornwell et al., 2008; Fortunel et al., 2009; Kazakou et al., 2009) or root exudates production (Guyonnet
 116 et al., 2018) and may therefore be a promising tool to study the relationships between vegetation, soil microbiota and
 117 SOM formation dynamics (Faucon et al., 2017).

118 The aim of this work is to evaluate the impact of rotational grazing, legume sowing, and grazing exclusion on topsoil (0–
119 8 cm) SOC stocks, and their distribution between POM and MAOM fractions, in semi-arid grasslands, compared to
120 conventional continuous grazing. We also evaluate changes in vegetation characteristics (identity and diversity of
121 biochemical and morphological traits), soil nutrients, and soil microbial communities as possible pathways through which
122 management might indirectly affect SOC stocks and fractions. This study focuses on the Iberian *dehesas* Mediterranean
123 woody grassland, the main example of semi-arid grasslands in Europe (Porqueddu et al., 2016). The most widespread
124 livestock management in this ecosystem is continuous grazing, but in recent decades rotational grazing and legume sowing
125 have gained importance (Frangia et al., 2023; Pulina et al., 2023). At the same time, the amount of ungrazed pastures has
126 increased due to land abandonment (Palomo-Campesino et al., 2018). Iberian *dehesas* occupy 3.1 million hectares,
127 spanning a wide environmental gradient, and have been subject to extensive grazing for centuries (Moreno and Pulido,
128 2009), making them an ideal model system for assessing the effects of improved management on SOC stocks in semi-
129 arid grasslands. In particular, we designed our study to answer the following questions: 1) What are the effects of the
130 different management practices on bulk SOC and fractions stocks? 2) Are these effects mediated by changes in vegetation
131 or soil microbial communities? 3) Is SOC storage in these grasslands, and its enhancement, modulated or limited by
132 environmental factors such as climate or soil properties? And 4) are these mechanisms and controls the same or different
133 for C in POM or MAOM? Understanding the potential and limitations of improved management on SOC storage in POM
134 and MAOM in semi-arid grasslands can guide policymakers in enhancing the climate change adaptation and mitigation
135 capacity of these ecosystems, while supporting productivity and soil fertility.

136

137 2. Methods

138 2.1. Study area and experimental design

139 The study was carried out at nine commercial *dehesa* farms located along a north-south gradient in the western part of
140 the Iberian Peninsula (Fig. 1a). The region has a continental Mediterranean climate, but on a local scale, in relative terms,
141 farms can be grouped into three main climatic regions (Fig. 1b, c) according to the average climate for the period 1980–
142 2018 (García Bravo et al., 2023). A cold-dry region [12.9 °C mean annual temperature (MAT) and 445 mm mean annual
143 precipitation (MAP)] in the north; a warm-wet region (17.3 °C MAT and 603 mm MAP) in the middle of the latitudinal
144 gradient; and a warm-dry region (17.0 °C MAT and 510 mm MAP) in the south. The soils of the farms share a common
145 development from granites, shales and sandy tertiary sediments, are acidic and poor in organic matter, but cover a wide
146 texture gradient (Fig. 1d). In these farms, native pastures are often combined with scattered trees, as is common in
147 Mediterranean and semi-arid rangelands (den Herder et al., 2017; Soliveres et al., 2014). The tree layer is dominated by
148 holm oaks (*Quercus ilex* L.) with scattered cork oaks (*Quercus suber* L.) or gall oaks (*Quercus faginea* Lam.). The
149 herbaceous layer is composed of species typical of Mediterranean pastures and presents a high diversity and proportion
150 of annual C3 plants (Table S1). The growing season of the pasture is very limited by the Mediterranean summer drought
151 with the annual species germinating in autumn (early-to mid-October), reaching their peak productivity in mid-spring
152 (late April of the following year) and senescing in June.

153 *Dehesa* farms are typically managed by extensive continuous grazing, where livestock (mainly cattle) graze freely on
154 large areas following a loosely defined grazing plan. This management has been traditionally practiced in all the farms
155 studied, until, in recent decades, some paddocks were converted to other management practices. As a result, on each farm
156 we selected five paddocks, each with one of the following management regimes:

157 - Abandoned (Ab): Paddocks devoid of grazing for at least the last 10-20 years.

158 - Continuous grazing (Ct): Paddocks where livestock stand most of the year, fed by grazing and supplemented. This is the
159 control treatment in our study as it is the most widespread management in Iberian *dehesas*.

160 - Rotational grazing (Ro): Paddocks intensively grazed in short periods and with resting periods (i.e. without livestock)
161 lasting for more than six months. Rotational grazing has been applied in these paddocks for the last 10-15 years.

162 - Recent legume sowing (Lr): Paddocks where pastures have recently (≤ 5 years) been sown with legume mixtures. In
163 *dehesa* farms, legume sowing consists of sowing a mixture of seeds (at a rate of 20 kg ha⁻¹) from various species of annual
164 legumes (pre-inoculated with Rhizobium) such as *Trifolium subterraneum* L., *T. incarnatum* L., *T. michelianum* Savi., *T.*
165 *resupinatum* L., *T. mutabile* Port. and *Ornithopus sativus* Brot., along with some highly productive annual grass species
166 such as *Lolium multiflorum* Lam. and *Lolium rigidum* Gaud., [all of which have relatively shallow root systems](#) (Teixeira
167 et al., 2015). These sowings are preceded by surface tillage and phosphorus application to meet the needs of the legumes
168 and stimulate N-fixation (Jongen et al., 2019). Farmers ~~sow~~ legumes mixture only once, because its effect persists over
169 the years thanks to the natural seeds production and the resulting soil seed bank of the sown species. None of the sown
170 plots studied was resown.

171 - Old legume sowing (Lo): Paddocks sown with legume mixtures more than 10 years ago.

172 The distinction between recent and old legumes sown paddocks allows us to compare short and long-term effects after
173 sowing. In two out of the nine farms, we added an additional control (continuous grazing) paddock close to the legume
174 sowing paddocks because they were at a considerable distance from the other control paddock, limiting their
175 comparability. Additional information on paddocks characteristics is provided in Table S2.

176 In each paddock, we installed four permanent monitoring plots, in open grassland and away from the direct influence of
177 the trees (at least 10 m away from the trees). This setup allowed us to have a total of 188 sampling plots where we
178 measured soil properties, SOC fractions, and soil microbial communities in 2021 and vegetation traits for 3 years (2021
179 to 2023).

180




Figure 1. Iberian “dehesa” cover according to the CORINE 2018 land cover survey and geographical location of the studied farms (a). Mean annual temperature (b) and precipitation (c) during the period 1980–2018 in the study area. Soil texture in all sampled plots (d).

2.3. Vegetation sampling

We assessed above-ground (ABG) pasture productivity over three growing periods at all plots using 50 x 50 x 50 cm grazing exclusion cages. In the 2020–2021 period (hereafter referred to as 2021) we installed one cage on each monitoring plot at the beginning of spring (late March 2021). In the 2021–2022 and 2022–2023 periods (hereafter referred to as 2022 and 2023), we installed the cages at the beginning of the green-up in early November. Standing biomass at the time of

192 cage establishment (t0) was determined by hand clipping within two 25 x 25 cm squares, randomly placed in the proximity
193 of the cage. In the three years we harvested ABG biomass inside the cages at the beginning of summer in early June (t1),
194 when most of the vegetation was already senescent. Biomass collected both in t0 and t1 was oven-dried at 60°C for 48h
195 and weighed. We calculated ABG productivity as the difference between t0 and t1 biomass. Therefore, only spring
196 productivity was measured in 2021, which represents the largest proportion of annual productivity in this system, while
197 a closer estimate of annual productivity was measured in 2022 and 2023.

198 Plant traits were collected in early May 2021, at the peak of the growing season. We considered a circular area of three
199 meters in diameter around each exclusion cage and within this area, we identified the species present at 25 regularly
200 spaced points using the line intercept method (Godínez-Alvarez et al., 2009), with three concentric circular transects going
201 around the cage. After completing the species inventory for all the sampling plots in a paddock, we collected at least three
202 to ten full individuals, in that same paddock, of each of the identified species, as proposed in the abundance-weighted
203 trait sampling scheme (Carmona et al., 2015). Thus, at least ten individuals of the most abundant species were collected
204 in the same paddock, five individuals for the medium abundant species and three individuals for the rare species. The leaf
205 area (LA), specific leaf area (SLA), specific root length (SLR), leaf dry matter content (LDMC), root dry matter content
206 (RDMC), leaf nitrogen content (LNC) and plant maximum height (see Fig. 2 for more detail) of all the collected
207 individuals was measured following the standard protocols (Pérez-Harguindeguy et al., 2016). In total, more than 10000
208 individuals were measured across all the plots. This extensive sampling allowed us to account not only for interspecific
209 differences in trait values, but also for intraspecific variability in trait values across managements, farms and regions.
210 Intraspecific variability has proven to be very important for accurately defining the functional composition of plant
211 communities and their relationship to ecosystem processes (Siefert et al., 2015; Westerland et al., 2021). We also
212 conducted species inventories in the spring of 2022 and 2023, but on these later samplings we only collected and measured
213 the traits of individuals from the species not found in 2021. We used the trait values measured in 2021 for each species in
214 each paddock to impute trait values in 2022 and 2023. The proportion of legumes in each plot was quantified from the
215 species inventories as a measure of the number of plants with N fixation capacity in the communities.

216 Pasture chemical composition was assessed in the three years, after the species inventories and traits samplings. We
217 collected standing biomass by hand-clipping in three 25 x 25 cm quadrats randomly placed in the monitoring plot, outside
218 the exclusion cages. These samples were dried at 60°C for 48h, then grounded and analyzed with Dumas method in
219 DUMATHERM® N Pro analyzer (C. Gerhardt GmbH & Co. Germany) to obtain the nitrogen content of each sample.
220 Acid detergent lignin (ADL) and fiber (ADF) and neutral detergent fiber (NDF) content were also measured. NDF, ADF
221 and ADL were determined using a fiber analyzer (ANKOM A2000, ANKOM Technology, USA), following the official
222 procedures (Latimer, 2023).

Trait	Description	
Height 	Height to top photosynthetic tissue (cm). - Positively related to competitive ability for light and seed dispersal ^{1,2}	Traits associated with the plant size spectrum ³
Leaf area (LA) 	One-sided area of a fresh leaf (mm²) - Positively related to light interception . It also has implications for the water and energy balance of the leaf ³	
Specific leaf area (SLA) 	Leaf area/ Leaf dry mass (mm² mg⁻¹) - Positively related to plant relative growth rate and litter decomposability . Negatively related to stress tolerance ⁴	
Leaf dry matter content (LDMC) 	Leaf dry mass/ Leaf fresh mass (mg g⁻¹) - Negatively related to relative growth rate and litter decomposability . Positively related to stress tolerance ^{5,6}	Traits related with the Leaf/Root Economic Spectrum (conservative – acquisitive resources strategy gradient) ^{8,11}
Leaf nitrogen content (LNC) 	Nitrogen content in dry leaf mass (mg g⁻¹) - Positively related to photosynthetic rate , leaf respiration and litter decomposability ^{7,8}	
Root dry matter content (RDMC) 	Roots dry mass/ Roots fresh mass (m g⁻¹) - Correlated positively with root tissue density . Negatively correlated with root respiration and exudate production ^{9,10}	
Specific root length (SRL) 	Roots length / Roots dry mass (mm mg⁻¹) -Negatively correlated with root diameter and mycorrhizal association . Positively correlated with soil exploration and nutrient acquisition ^{11,12}	Traits related with the fungal collaboration gradient ¹¹



Trait	Description	
Height 	Height to top photosynthetic tissue (cm). - Positively related to competitive ability for light and seed dispersal ^{1,2}	Traits associated with the plant size spectrum ³
Leaf area (LA) 	One-sided area of a fresh leaf (mm²) - Positively related to light interception . It also has implications for the water and energy balance of the leaf ³	
Specific leaf area (SLA) 	Leaf area/ Leaf dry mass (mm² mg⁻¹) - Positively related to plant relative growth rate and litter decomposability . Negatively related to stress tolerance ⁴	
Leaf dry matter content (LDMC) 	Leaf dry mass/ Leaf fresh mass (mg g⁻¹) - Negatively related to relative growth rate and litter decomposability . Positively related to stress tolerance ^{5,6}	Traits related with the Leaf/Root Economic Spectrum (conservative – acquisitive resources strategy gradient) ^{8,11}
Leaf nitrogen content (LNC) 	Nitrogen content in dry leaf mass (mg g⁻¹) - Positively related to photosynthetic rate , leaf respiration and litter decomposability ^{7,8}	
Root dry matter content (RDMC) 	Roots dry mass/ Roots fresh mass (m g⁻¹) - Correlated positively with root tissue density . Negatively correlated with root respiration and exudate production ^{9,10}	
Specific root length (SRL) 	Roots length / Roots dry mass (mm mg⁻¹) -Negatively correlated with root diameter and mycorrhizal association . Positively correlated with soil exploration and nutrient acquisition ^{11,12}	Traits related with the fungal collaboration gradient ¹¹

Figure 2. Graphical representation and description of plant functional traits measured. References embedded in the figure are provided below: ¹Gaudet and Keddy (1988), ²Thomson et al. (2011), ³Díaz et al. (2016), ⁴Poorter et al. (2009), ⁵Kazakou et al. (2009), ⁶Wilson et al. (1999), ⁷Freschet et al. (2012), ⁸Wright et al. (2004), ⁹Guyonnet et al. (2018), ¹⁰Roumet et al. (2016), ¹¹Bergmann et al. (2020), ¹²Kramer-Walter et al. (2016).

2.4. Soil sampling and soil characteristics measurement

In spring 2021, four soil cores were collected with a push sampler (5 cm diameter) at a depth of 8 cm around each exclusion cage, after removal of above-ground vegetation and litter from the sampled surfaces. The four soil cores were combined to obtain a composite sample from each plot, and an aliquot of 40 g of this sample were sieved (<2 mm),

reserved and stored at 4°C for microbial community analysis in the following days. The remainder of the composite soil samples were air-dried and sieved (<2 mm). Coarse mineral material greater than 2 mm (hereinafter referred to as gravel) was weighed and used for bulk density correction (Eq.1).

Soil texture (sand, clay and silt content) was determined using a laser diffraction particle size analyzer (Mastersizer 2000, Malvern Instruments Ltd. UK) after dispersion with sodium hexametaphosphate. Soil pH was measured with a glass electrode (soil:water 1:2.5) pH meter (CRISON Basic20, Alella, Spain). Soils were extracted with 1M KCL and determined colorimetrically in a Bran+Luebbe Autoanalyzer 3 (Norderstedt, Germany), following the manufacturer's protocol, to measure nitrate (NO₃⁻) and ammonium (NH₄⁺) concentrations. Available P (Olsen P) was determined colorimetrically in an Agilent Cary 60 UV-Vis spectrophotometer (Agilent Technologies, USA), after extraction with 0.5 M NaHCO₃ at pH 8.5. Available K, Ca and Mg were determined by ICP-OES (Varian Mod. 720, Palo Alto, California, USA), after extraction with 1M ammonium acetate (pH 7).

245

2.5. Soil organic matter fractionation


Representative subsamples of each 2mm sieved air-dry composite soil sample were used to measure both bulk SOC and total nitrogen (TN) as well as their distribution across distinct physical fractions. First, 10 g aliquots of all samples were ground in a ball mill and the Bernard's calcimeter method (Sherrod et al., 2002) was used to test for the presence of inorganic C. Only four samples (0.02 % of the database) contained traces of carbonates, with CaCO₃ contents between 0.2 % and 0.8 %. These samples were excluded from subsequent analyses. For bulk soil analyses, aliquots of 10g of soil were ground in a ball mill and then analyzed in the DUMATHERM® N Pro analyzer (C. Gerhardt GmbH & Co. Germany) to determine %C (SOC) and %N (TN). For soil fractionation, we followed a combined size and density procedure as described in Leuthold et al. (2024). Briefly, an aliquot of 6 g of soil was mixed with sodium polytungstate (1.85 g cm⁻³) and shaken reciprocally for 18 h to disperse the aggregates. After dispersion samples were centrifuged for density fractionation and the light particulate organic matter "POM" (<1.85 g cm⁻³) was aspirated from the rest of the soil. We then thoroughly rinsed the residual heavy fraction and separated it by wet sieving into coarse heavy mineral-associated organic matter "chaOM" (>53 µm) and fine mineral-associated organic matter "MAOM" (<53 µm). All fractions were analyzed for %C and %N on an elemental analyzer as described above. Fractionation was accepted when mass recovery was within ±5%, and C recovery was within ±15%. For samples that did not satisfy one of these conditions, fractionation was repeated.

Since the chaOM and the MAOM shared similar C/N ratios, which were lower than POM C/N ratios (Fig. 3a), we merged these two mineral associated OM fractions and present them together as MAOM (Santos et al., 2024; Zhang et al., 2021). Thus, in this work, POM and MAOM represented the light (<1.85 g cm⁻³) and heavy (>1.85 g cm⁻³) fractions respectively. This contrasts with other studies based on size fractionation, in which light POM and chaOM are pooled together (Cotrufo et al., 2019b; Dai et al., 2025; Díaz-Martínez et al., 2024). However, our decision is supported by recent findings of Leuthold et al. (2024) who observed greater chemical similarity between chaOM and MAOM than between chaOM and POM. The pooling of the two mineral associated OM fractions is not expected to modify the results or interpretation of this work, as the C content of chaOM only accounts for an average of 8% of the total SOC, and its relative importance is similar across all managements (Fig. S1). Carbon data are presented as SOC, POC and MAOC stocks (Mg ha⁻¹), calculated following Poeplau et al. (2017):

$$\text{OC}_{\text{stock}} = \text{OC}_{\text{content}} \times \frac{\text{mass}_{\text{fine soil (<2mm)}}}{\text{Volume}_{\text{sample}}} \times \text{depth}_{\text{sample}} \quad (1)$$

273 where $OC_{content}$ is the organic C content (as a proportion) of the soil fraction, $mass_{fine\ soil(<2mm)}$ is the dry weight of the soil
 274 excluding gravel and large roots, and considering the dry mass of the aliquot reserved for microbial community analysis.
 275 $Volume_{sample}$ and $depth_{sample}$ are respectively the volume and depth of the soil sampled with the core.
 276 Additionally, we calculated the proportion of MAOC in total SOC (MAOC/SOC) following Cotrufo et al. (2019).

277



278

279 **Figure 3.** Violin plot and boxplot (with median and quartiles) and mean values (expressed with labels and red dots) for the ratio
 280 of carbon to nitrogen content (C/N ratio) in the soil organic matter fractions (a), and relation between carbon content in the
 281 soil organic matter fractions (in g of C per kg of soil) and the total soil organic carbon (SOC) content (b). “chaOM” refers to
 282 the coarse heavy mineral-associated organic matter, “IPOM” to the light particulate organic matter and “MAOM” to the fine
 283 mineral-associated organic matter. In panel “b”, a linear regression line depicts the relationship between IPOM and total SOC,
 284 while logistic curves illustrate the relationships of chaOM and MAOM with total SOC.

285

286 2.6. Soil microbial communities:

287 Soil microbial communities were characterized by phospholipid fatty acids analysis (PLFA; Willers et al., 2015). A ~~2~~^{two}
 288 g fine soil aliquot of lyophilized soil was used for lipid extraction with a one-phase chloroform–methanol–phosphate

289 buffer solvent. Phospholipids were separated from non-polar lipids and converted to fatty acid methyl esters (FAMES),
 290 which were then separated by gas chromatography as described in details in Moreno et al. (2021). The sum of all
 291 individual PLFAs was used as a proxy for microbial biomass (in nmol PLFAs g⁻¹ of soil). Further, we estimated microbial
 292 biomass stocks (in mol ha⁻¹) by substituting the OC_{content} by the total PLFAs concentration in equation 1. Specific PLFAs
 293 were used as biomarkers to estimate the relative abundance of broad taxonomic microbial groups, according to their
 294 characteristic fatty acids: eukaryote, Gram- and Gram+ bacteria, saprophytic fungi and arbuscular mycorrhizal fungi
 295 (Frostegård and Bååth, 1996). The ratios among Gram+ and Gram- bacteria (Gram+/Gram-) and fungi and bacteria
 296 (Fungi/Bacteria) were also calculated to describe the composition of the microbial community.

297

298 2.7. Mineral-associated carbon capacity and saturation:

299 To evaluate the degree of MAOC saturation, we used the boundary line approach reported by Georgiou et al. (2022)
 300 adjusted with global soil samples as a reference of the maximum observed mineralogical capacity (sensu Georgiou et al.,
 301 2025) of our samples (Fig. 4). Therefore, the saturation deficit in each sample was calculated as one minus the ratio of
 302 the current C content and the observed maximum C content according to the mineralogical capacity (% clay + silt).

303


304 2.6. Data analysis:

305 All analyses were carried out in R 4.3.3 (R Core Team, 2024). Species abundance data and species trait values were used
 306 to compute, for each year and plot, the community weighted means (CWM) of all traits as well as the RaoQ multi-trait
 307 functional diversity index (Ricotta and Moretti, 2011). In addition, we calculated the species richness as a measure of the
 308 taxonomic diversity in each plot. These analyses were performed using the package “FD” (Laliberté and Legendre, 2010).
 309 Regarding pasture chemical composition, the hemicellulose content was calculated as NDF minus ADF and the cellulose
 310 content as ADF minus ADL, with ADL being considered a good proxy for the lignin content (Van Soest et al., 1991).

311 For the vegetation analysis, measurements from the three years were combined to obtain mean estimates of each
 312 vegetation attribute. For each year, the ABG production and chemical attributes values, as well as the CWM of all the
 313 functional traits, were centered and scaled between -1 and 1, so that values closer to -1 represent plots with traits values
 314 lower than the mean, while values closer to 1 represent plots above the mean. The three years centered and scaled values
 315 were then averaged to obtain estimates of each vegetation characteristic across the study period, as shown in this equation
 316 2:

$$317 \text{Vegetation attribute}_{\text{mean}} = \frac{\sum_{\text{year } i=2021}^{2023} \text{Vegetation attribute (centered and scaled)}_{\text{year } i}}{3} \quad (2)$$

318 All mean values of vegetation characteristics obtained with equation 2 were used to build a Principal Component Analysis
 319 (PCA). From this PCA, we extracted 4 main axes of variation of the vegetation characteristics with eigenvalues greater
 320 than one (Fig .4).






Figure 4. Representation of the 4 main axis of variation in the principal component analysis (PCA) summarizing the vegetation characteristics variables. Panels a, b, c and d show the correlation between the different variables included in the PCA and the new axis. Panels e, f, g and h represent the plant communities characteristics at the extreme of each axis. Representative species of each axis are represented in Fig. S2.

Nutrients stocks in the soil samples were calculated by substituting the $OC_{content}$, in equation 1, for the concentration of each nutrient. PCA was used to summarize these nutrient stocks into a single composite index, as all nutrients were positively correlated. The first component of this PCA, which absorbed 46.9% of the variance, was extracted as a new variable, henceforth called “soil fertility” (Fig. S3). This soil fertility index increases as ammonium, nitrate, P, Ca, Mg, and K stocks in soil increase.

332 To analyze the direct and indirect effects of management, vegetation characteristics, microbial communities and
 333 environmental variables on POC, MAOC and SOC stocks and MAOC/SOC ratio we built a structural equation model
 334 (SEM) using the “piecewiseSEM” package (Lefcheck, 2016). These models assess the extent to which a defined structure
 335 of causal relationships between variables fits the actual correlations between-in the data. In this causal structure, direct
 336 effects describe the influence of one variable on another without intermediaries, whereas indirect effects occur when a
 337 variable affects another through its impact on a third variable. We started from the assumption that management, climate
 338 and soil properties could affect SOC, POC and MAOC stocks and the MAOC/SOC directly or indirectly, by modifying
 339 microbial composition and vegetation characteristics. This provided us with a theoretical basis for the construction of the
 340 SEMs. The linear sub-models within these SEMs were fitted with the “lme4” package (Bates et al., 2015) using the farms
 341 as random factors to avoid spatial pseudo-replication. Model fitting was performed according to Zuur et al. (2009)
 342 selecting the model with the lowest corrected Akaike Information Criterion (AICc) value after adjusting for random and
 343 fixed factors. All predictor variables used in the model selection are summarised in Table 1. While gravel content is
 344 correlated to BD, its inclusion as predictor in the models helped to control for inter-site variability in gravel content.
 345 Interactions between management and climate or soil texture were also included during model selection. All model
 346 assumptions were checked and satisfactorily met. Predictor variables were scaled and centered prior to inclusion in the
 347 models.

348 To calculate the total effect of each explanatory factor (i.e. the sum of direct and indirect effects) on SOC, POC and
 349 MAOC stocks, and MAOC/SOC ratio in the SEMs, we used the “semEff” package (Murphy, 2022). This package
 350 generates estimates and confidence intervals of the total effects of the explanatory factors across multiple permutations
 351 of the data (9999 bootstraps in our case).

352 Given that average bulk density values partly differed among managements (Figure S4), and that this variation may affect
 353 carbon stocks, the same analysis procedure used for POC, MAOC and SOC stocks was used to analyze the change in
 354 POC, MAOC and SOC contents.

355 **Table 1. List of the variables and their units, which were included as fixed factors during the structural equation model**
 356 **(SEM) sub-models selection.**

Variables	Units
Management	Categorical (5 levels)
Sand content	%
Gravel content	%
Soil fertility	unitless
Mean annual precipitation (MAP)	mm
Mean annal temperature (MAT)	°C
Vegetation PCA- Axis 1 (N-C axis)	unitless
Vegetation PCA- Axis 2 (PES axis)	unitless
Vegetation PCA- Axis 3 (P-L axis)	unitless
Vegetation PCA- Axis 4 (Size axis)	unitless
Functional diversity (Rao multi-trait index)	unitless
Taxonomic diversity (species richness)	count
Microbial biomass (bulk)	nmol ha ⁻¹
Gram+/Gram-	ratio

Fungi/Bacteria	ratio
----------------	-------

357

358 3. Results:

359 The mean values of POC, MAOC and SOC stocks were 9.8 ± 4.3 , 7.1 ± 3.1 and 16.5 ± 6.3 Mg C ha⁻¹ respectively. On
360 average, MAOC represented 44% of the total SOC. The MAOC content of all the soils analyzed was well below the
361 theoretical saturation level established by Georgiou et al. (2022) in relation to the clay + silt content (Fig. 5). Mean MAOC
362 saturation deficit was 80%. Soil organic carbon fractions and stocks were affected directly and indirectly by mean plant
363 traits, microbial communities, soil properties and management, and indirectly by climatic conditions (Fig. 6). Plant
364 diversity indices and interactions between management and climate or soil texture were not retained in the model selection
365 due to lack of significance.





Figure 5. Relation between the mineral-associated organic carbon (MAOC) and the percent Clay+Silt (CS) in the soil samples and the boundary line (in blue) with confidence intervals adjusted by Georgiou et al. (2022) for high-activity mineral soils, indicating the maximum observed mineralogical capacity for each CS content. The equation for the boundary line is provided in the panel. Points are colored based on their MAOC deficit (1- MAOC content / MAOC maximum observed capacity).

Total microbial biomass, estimated as total PLFA stocks, averaged 66.3 ± 25.9 mol ha⁻¹, with mean values for Fungi/Bacteria and Gram+/Gram ratios of 0.08 ± 0.03 and 1.1 ± 0.2 , respectively, although there were some differences between treatments (Fig. S5). Microbial biomass was positively correlated with POC, MAOC and SOC stocks (Fig. 6f;


Fig. 7a, b & c). The Fungi/Bacteria ratio was negatively correlated with MAOC stocks (Fig. 6f; Fig. 7b) meaning that these stocks were higher in soils with a greater bacterial predominance. The Gram+/Gram- ratio had a significant direct negative effect on POC, MAOC and SOC ~~SOC, MAOC and PC~~ stocks (Fig. 6f; Fig. 7a, b & c).

The first axis of the PCA of the vegetation characteristics, which reflects the N-C stoichiometry of the vegetation, had a negative direct effect on POC, MAOC and SOC stocks (Fig. 6h; Fig. 7a, b & c). The second axis, related to the plant economic spectrum (PES), had a positive direct effect on SOC stocks (Fig. 6h; Fig. 7c). In addition, the PES axis was positively correlated with the microbial biomass, thus having a positive indirect effect, and a significant total effect on POC, MAOC and SOC stocks (Fig. 6e; Fig. 7a, b & c). The third axis of the PCA (P-L axis), positively correlated with the lignin content and negatively related to vegetation productivity, was negatively correlated with the POC, MAOC and SOC stocks and the microbial biomass (Fig. 6e & h; Fig. 7a, b & c). The fourth axis, reporting the plant size, was positively correlated with the MAOC stocks, the MAOC/SOC ratio and the Fungi/Bacteria ratio (Fig. 6e & h; Fig. 7a & d). No index of plant functional and taxonomic diversity was retained during model selection. On the other hand, soil fertility had a direct and indirect positive effect on POC, MAOC and SOC stocks (Fig. 6g). Soil fertility was negatively correlated with the Gram+/Gram- ratio, the P-L axis and the N-C axis, and positively correlated with the PES axis of the vegetation PCA (Fig. 6i & j).

Plot under continuous grazing had a mean POC, MAOC and SOC stocks of 9.1, 6.8 and 16.2 Mg C ha⁻¹ respectively, and a MAOC/SOC ratio of 0.44 (Fig. 8). Rotational grazing significantly increased soil fertility compared to continuous grazing (Fig. 6k). This led to a significant indirect effect of rotational grazing on POC, MAOC and SOC stocks (Fig. 7b & c). POC, MAOC and SOC stocks under rotational grazing had a mean value of 10.5, 7.8 and 18.5 Mg C/ha, respectively, which were 15% (in the case of POC and MAOC) and 14% ~~higher~~ (for SOC stocks) higher than mean values in continuous grazing (Fig. 8b & c). Recent legume sowing had a negative direct effect on POC, MAOC and SOC (Fig. 6b). However, Lr also increased significantly the plant size axis, the microbial biomass and the soil fertility and decreased the plant N-C axis compared to Ct (Fig. 6a, c & k). These changes resulted in a positive indirect effect of Lr over POC, MAOC and SOC stocks and a null total effect (Fig. 7a, b, c & d). Grazing abandonment increased the vegetation N-C axis and reduced the microbial biomass compared to Ct (Fig. 6a & c), resulting in a negative indirect effect on POC, MAOC and SOC stocks and a significant negative total effect on POC stocks (Fig. 7a, b & c). Thus, the mean POC stock on abandoned plots was 8.1 Mg C/ha, 11% lower than in ~~Ct continuous grazing~~ plots (Fig. 8a).

Sand content in soil was negatively correlated with MAOC stocks and MAOC/SOC ratio (Fig. 6j). Sand content was also negatively correlated with soil fertility and the PES axis and positively correlated with the P-L and N-C axis (Fig. 6i), thus having a negative indirect effect on POC, MAOC and SOC stocks (Fig. 7a, b & c). As expected, due to its negative correlation with BD, gravel content was also negatively correlated with microbial biomass (Fig. 6j), leading to a significant negative indirect effect on POC, MAOC and SOC stocks (Fig. 7a, b & c). MAT was positively correlated with the N-C axis and negatively correlated with the soil fertility index (Fig. 6d & l). This implied a negative indirect effect of MAT over POC, MAOC and SOC stocks (Fig. 7a, b & c). On the other hand, MAP was negatively correlated with the N-C and the size axis (Fig. 6d). Thus, MAP had a positive indirect effect on POC, MAOC and SOC stocks (Fig. 7a, b, c & e).

Fisher C: 127.842
P-value: 0.905








Figure 6. Structural equation model representation. Factors included in the model are grouped by factor type (management, climate, vegetation traits, soil properties, microbial communities). Arrows between groups of factors indicate significant relationships between any of the factors included in both groups. The width of these arrows is proportional to the mean absolute size of the estimates between the factors in the groups. The plots embedded into these arrows show the standardized estimates of the significant relationships between the factors connected by the arrow. Negative standardized estimates are represented in red, and positive ones in blue. The size of the estimate circles represents the absolute value of the standardized estimate. Causal relationships between factors in the same group are represented by individual arrows.


POC stock




MAOC stock



SOC stock



MAOC/SOC






Figure 7. Direct, indirect and total standardized effects of all studied variables included in the structural equation model (Figure6) over the (a) particulate organic carbon (POC), (b) mineral-associated organic carbon (MAOC), and (c) soil organic carbon (SOC) stocks and (d) the relative MAOC abundance (MAOC/SOC). Bars indicate direct (dark colors) and indirect (light colors) effects, and the black points-ranges indicate the total (i.e. direct + indirect) effect (with its 95% confidence interval). Stars over the total effect values indicate significant effects at a level of 0.05.




Figure 8. Violin plot and boxplot (with median and quartiles) for the (a) particulate organic carbon (POC), (b) mineral-associated organic carbon (MAOC), and (c) soil organic carbon (SOC) stocks and (d) the relative MAOC abundance (MAOC/SOC) in each management (Ct = Continuous grazing; Ab = grazing abandonment; Lr = Recent legume sowing; Lo = old legume sowing; Ro = Rotational grazing). The red point and the label indicate mean values predicted by the structure equation model (Fig.6). Lower case letters indicate significant differences ($p < 0.05$) between managements.

4. Discussion

Our results highlight the potential of management to control carbon storage in semi-arid grasslands. In this sense, rotational grazing arises as a promising tool for enhancing long-term carbon storage in the topsoil. We also identified

several pathways by which management influences SOC, POC and MAOC stocks, showing the importance of changes in vegetation and microbial composition. Furthermore, management effects were consistent across the wide soil-climate gradient examined, enabling generalization of our results to a broad set of semi-arid grasslands in various environmental contexts. It should be noted that the predicted capacity of POC, MAOC and SOC stocks models was high (62%, 69% 73% of the variance explained by the fixed factors, respectively), indicating the robustness of the results presented and their importance for expanding the understanding of soil carbon dynamics in grasslands. Our results are limited to the upper topsoil (the first 8 cm), and it would be necessary to analyze deeper layers to fully understand the processes of SOC formation and stabilization and the effects of management changes. However this top layer is the most important for mediterranean grassland functioning as it contains the majority of roots and therefore most of the microbial, nutrients and water dynamics (Acosta-Gallo et al., 2011; Moreno et al., 2005). Furthermore, changes in management primarily affect the SOC of the topsoil layer, especially in the short term (Ward et al., 2016).

453

4.1. Soil organic carbon fractions, stocks, and saturation levels

SOC stocks and contents (Fig.8 & S4) in the paddocks studied ($16.5 \text{ Mg C ha}^{-1}$ and 19.5 g C kg^{-1} on average, respectively) were consistent with values found in the first 7-10 cm of soil in other grasslands under similar climatic conditions (Díaz-Martínez et al., 2024; Oggioni et al., 2020; Parras-Alcántara et al., 2014). On the other hand, the MAOC/SOC ratio of our soils (0.44 on average) was lower than the values generally reported in both global and semi-arid grasslands (Hansen et al., 2024; Rocci et al., 2022; Shi et al., 2024), which are around 0.6-0.7. The shallow sampling depth in this work (8 cm) may influence the observed MAOC/SOC ratios, as POM decrease more sharply than MAOM with soil depth (Galluzzi et al., 2025; Sanderman et al., 2021; Tang et al., 2024). For instance, Plaza et al. (2022) reported MAOC/SOC ratios between 0.3 and 0.4 in the first 10 cm of soil, while Cappai et al. (2017) found MAOC/SOC ratios around 0.7 in the first 20 cm, both in Mediterranean grasslands.

Carbon concentrations in the fine soil fraction (clay + silt) were well below the saturation point observed in previous studies (Cotrufo et al., 2019a; Georgiou et al., 2022). However, we observed a certain limit to MAOC accumulation in our system, as its content remained stable above SOC contents of 30 g kg^{-1} and stayed below 20 g kg^{-1} even when SOC reached values above 60 g kg^{-1} , following a saturation curve (Fig. 3b). These results support the observations that MAOC accrual is more limited by the amount of C inputs rather than the mineralogical capacity of the soil (Poeplau et al., 2024). In this sense, the maximum C in the clay+silt fraction observed in this work (around $19 \text{ g C/kg clay+silt}$) would represent the ~~effective~~ ~~maximum~~ capacity of the ecosystem to ~~stabilise~~ ~~stabilize~~ C at current levels of productivity. Therefore, MAOC saturation should be a minor concern when planning management strategies to improve C storage in Mediterranean grasslands.

473

4.2. Soil microbial communities regulate SOC storage

Total microbial biomass had a similar and substantial positive effect on POC, MAOC and total SOC stocks, such that the MAOC/SOC ratio was unaffected. Typically, microbial biomass C only represents about 2% of SOC (Xu et al., 2013; Yao et al., 2000), but it is closely correlated with the accumulation of microbial necromass C (Hou et al., 2024). The latter can account for more than 50% of SOC (Liang et al., 2019), with a similar contribution to the POC and MAOC fractions (Zhang et al., 2024). The positive effect of microbial biomass on both POC and MAOC, as well as the relatively low C:N ratio of these fractions found in this study (Fig. 3a) compared to other works (Chang et al., 2024; Yu et al., 2022), indicate

a prominent role of microbial transformation of plant inputs in SOC formation in semi-arid grasslands, which would also explain the relatively low SOC stocks of these ecosystems.

Fungi/bacteria ratio was negatively correlated with MAOC fraction, as fungi residues tend to contribute more to POC (Griepentrog et al., 2014; Lavallee et al., 2020; Tang et al., 2023), but no significant effect of fungi/bacteria ratio over POC stocks or MAOC/SOC ratio was observed. Gram+/Gram- ratio was negatively correlated with POC, MAOC and SOC stock, as observed in previous studies (Khatri-Chhetri et al., 2024). Gram- bacteria are more dependent on plant C inputs, whereas Gram+ bacteria tend to use more of the organic C already present in the soil (Fanin et al., 2019; Kramer and Gleixner, 2008; Waldrop and Firestone, 2004). Thus, the proliferation of Gram+ bacteria may promote the decomposition of pre-existing SOM (Klump et al., 2009).

490

4.3. Plant-soil interactions and their effect on soil carbon stocks

The analysis of vegetation functional traits and chemical composition revealed four main axes of variation. Two of these axes (axes 1 and 4) correspond to the spectrum of plant form and function, found in several global vegetation analyses (Díaz et al., 2016; Weigelt et al., 2021). This two-dimensional spectrum is defined by the leaf and root economic gradient (Kramer-Walter et al., 2016; Wright et al., 2004), that moves from slow-growing and resource-conserving to fast-growing and resource-acquisitive species (Wright et al., 2004), and the size gradient, which increases with increasing plant height and leaf area and reflects the competitive ability of plants for the light (Díaz et al., 2016). The other two axes (axes 2 and 3) are more related to the aboveground chemical composition of the vegetation. Axis 1 reflects the N-C stoichiometry of plant tissues, driven mainly by the ABG C/N ratio, both of which are obviously negatively correlated with certain traits such as LNC and the proportion of legumes. The N-fixing capacity of legumes has a fertilising effect on companion plants, increasing its tissues nitrogen content (Pino et al., 2016). In addition, legumes themselves tend to have a high concentration of nitrogen in leaves and stems (Carranca et al., 2015). Axis 3 is positively related to the lignin content of the vegetation. This independence between lignin and nitrogen content in plant litter has already been observed (Cornwell et al., 2008). ABG productivity of vegetation was correlated with most of these axes, especially with axis 3, being higher in more acquisitive, less lignified and bigger vegetation, as found in other works (Laliberté and Tylianakis, 2012; Zhang et al., 2017).

POC, MAOC and SOC stocks were higher in communities dominated by resource-acquisitive, highly productive plants with low lignin content and low C/N ratio. These communities are expected to provide a higher level of plant inputs, thereby increasing the incorporation of organic matter into the soil (King et al., 2023; Mortensen et al., 2025; Zhou et al., 2024). Increased inputs may come from higher plant litter production but also from increased root exudates, which tend to be higher in acquisitive plants (Guyonnet et al., 2018). Part of the effects of plant traits on SOC was mediated by increases in microbial biomass, which would also benefit from higher levels of plant inputs and increased root exudation (Eisenhauer et al., 2017). In addition, inputs with low lignin and C/N ratios are degraded more efficiently, reducing C losses and promoting long-term SOC storage (Cotrufo and Lavallee, 2022; Ridgeway et al., 2022). However, based on previous work, we would have expected a direct contribution of plant structural input to POC (Cotrufo et al., 2022), and thus a higher proportion of POC in communities that produce more recalcitrant litter (Cheng et al., 2023; Haddix et al., 2016; Mortensen et al., 2025). This was not the case in this work, where the MAOC/SOC ratio remained unaffected by the chemical composition of the ABG vegetation tissues. This unexpected result could be explained by the fact that in these systems, POC also appears to be the product of microbial transformation of plant inputs, as suggested by its positive relationship with microbial biomass and its relatively low C/N ratio. In addition, photodegradation, an important driver of litter degradation in semi-arid ecosystems (Austin and Vivanco, 2006), can promote litter lignin biodegradability and

the production of litter soluble compounds that are readily accessible to soil microbes (Wang et al., 2015), thus reducing the influence of vegetation chemical properties on POC and MAOC formation. Grazing also decouples the quality of plant tissues and the final quality of inputs to the soil, as livestock excreta have a lower C/N ratio than the plant material consumed (Soussana and Lemaire, 2014). We also observed that taller and larger plants (high values on the size axis) were associated with higher MAOC stocks and MAOC/SOC ratios, although the mechanism driving this correlation was unclear. Generally, plant height is positively correlated with shoot:root ratio (Li et al., 2008), and several studies have found a higher contribution of shoots, rather than roots, to the MAOC fraction due to the higher recalcitrance of root tissues (Huang et al., 2021; Lavalley et al., 2018; Ridgeway et al., 2022). However, rhizodeposition, which is closely linked to root-~~growth and turnover~~~~biomass~~, has been shown to promote MAOC formation over POC (Berenstecher et al., 2023; Engedal et al., 2023; Villarino et al., 2021). On the other hand, a greater accumulation of standing litter, rather than surface litter, might be expected in communities with bigger plants, and some studies in semi-arid grasslands have observed higher rates of microbial degradation and release of soluble compounds (thus contributing more to MAOC) in standing litter, compared to surface litter, due to greater retention of night-time moisture in the former (Gliksman et al., 2018; Wang et al., 2017). In any case, our results and their interpretation are limited by the correlation in our vegetation data between input quantity (ABG productivity) and input quality, which could mask the effect of the latter on SOC formation.

No significant effects of taxonomic or functional diversity on POC, MAOC or SOC stocks were found, which could be explained by the negative correlation of these biodiversity indicators with the ABG productivity and soil fertility in our plots (Fig. S7). This negative correlation has been observed in other natural grassland research and may be due to the dominance of some highly productive species in nutrient-rich environments (Feßel et al., 2016; Helsen et al., 2014; Luo et al., 2019; Rolo et al., 2016). In this sense, our results would be consistent with previous research that has found a greater importance of functional identity (i.e. mean traits values) relative to functional diversity in predicting ecosystem processes (Mokany et al., 2008; Zhang et al., 2017).

Soil fertility plays a central role in our models, increasing POC, MAOC and SOC stocks. Previous studies in grasslands have already observed a positive correlation between POC and MAOC and soil nutrients content (Wang et al., 2025). Global N addition experiments have also showed to increase topsoil C storage in grasslands (Hu et al., 2024; Liu et al., 2023). In this work, we observe a direct effect of soil fertility on C storage, but we also identify some indirect effects mediated by changes in vegetation and microbial communities. More resource-acquisitive and productive plant communities, with lignin content and lower C/N ratio, were found at higher soil fertility. Resource-acquisitive plants are expected to be more competitive in nutrient-rich environments (Daou et al., 2021; Ordoñez et al., 2009). The absence of nutrient limitation also promotes plant productivity (Fay et al., 2015). In addition, soil fertility was negatively correlated with the Gram+/Gram- ratio. Gram- bacteria have a more copiotroph strategy than Gram+ bacteria and therefore benefit more from nutrient-rich environments (De Deyn et al., 2010; Luo et al., 2020; Zhong et al., 2010). The direct positive effect of soil fertility on C storage in soils may be explained by changes in other factors that we did not measure. For instance, nutrient addition have been shown to promote below-ground productivity in mediterranean woody pastures (Nair et al., 2019) and in global grasslands (Liu et al., 2023; Lu et al., 2011). Moreover, nutrient availability in soils tends to increase CUE (Poeplau et al., 2019; Spohn et al., 2016) and to alleviate the need for microbial N-mining, reducing C losses and old SOM decomposition (Blagodatskaya and Kuzyakov, 2008).

4.5. Management effects

562 Rotational grazing increased both POC, MAOC and SOC stocks by 15%, 15% and 14%, respectively, compared to
 563 conventional continuous grazing. Previous studies found the same positive effect of rotational grazing on MAOC (Khatri-
 564 Chhetri et al., 2024; Mosier et al., 2021; Stanley et al., 2025) and total SOC content (Byrnes et al., 2018; Conant et al.,
 565 2017; Liu et al., 2024; Phukubye et al., 2022; Teague et al., 2011). According to our results, this effect is mainly explained
 566 by the higher soil fertility observed in rotational grazing plots, cascading in the above explained effects on plant traits and
 567 microbial transformation leading to higher POC, MAOC and SOC stocks. Soil fertility may be higher under rotational
 568 grazing than under continuous grazing because fecal and urine excretions tend to be less spatially clustered in rotationally
 569 grazed paddocks, resulting in more homogeneous fertilization of the entire paddock (Augustine et al., 2023; Dubeux Jr.
 570 et al., 2014). In contrast, in continuous grazing, excreta tend to accumulate in areas of highest animal use, such as near
 571 feeders or water points (Tate et al., 2003), which are areas that have been avoided in our sampling. In turn, the
 572 homogeneous grazing and fertilization maintains a greater amount of ground covered by vegetation (Stanley et al., 2025;
 573 Teague et al., 2004), which limits topsoil losses through erosion (Sanjari et al., 2009).

574 Surprisingly legume sowings had a negative direct effect on soil C stocks, but a positive indirect effect, resulting in a non-
 575 significant total effect. The negative impact was more evident in recent sowings, especially on the MAOC fraction, and
 576 could be attributed to the impact of the pre-sowing tillage. Previous works have observed a reduction in SOC stocks after
 577 tilling in mediterranean grasslands (Parras-Alcántara et al., 2015; Uribe et al., 2015). This would explain why the direct
 578 effect of legume sowing in C storage is more negative in recent compared to old-sowed paddocks. The effect of tillage
 579 on SOC stocks may arise from changes in soil bulk density rather than C content, resulting in misleading conclusions (Du
 580 et al., 2017; Rovira et al., 2022). However, in this work the analysis of POC, MAOC and SOC contents (in g/kg bulk soil;
 581 Fig. S6 & S8) shows similar effects of legume sowing as those observed for C stocks (Fig. 7a, b & c). In the recently
 582 sown plots, the negative impact in C stocks was countered by an increase in soil fertility, plant productivity and plant
 583 nitrogen content, possibly due to increased legume cover (Gómez-Rey et al., 2012; Gou et al., 2023; Hernández-Esteban
 584 et al., 2019). In addition, recent legume sowing also implied an increase in microbial biomass and a decrease in the
 585 Gram+/Gram- ratio, as observed in previous studies (Moreno et al., 2021).

586 Grazing abandonment led to a 12% reduction in the POC stocks, compared to continuous grazing. This result contradicts
 587 the observations of previous studies in grasslands, which reported positive effects of grazing exclusion on SOC storage,
 588 both globally (Eze et al., 2018) and particularly in semi-arid climates (Cheng et al., 2016; Gao et al., 2025; Yu et al.,
 589 2021). However McSherry and Ritchie (2013) meta-analysis found that light grazing could promote SOC stocks in C3
 590 dominated grasslands (such as our study grassland). Other global and regional studies also observed null or even positive
 591 effects of light grazing, compared to grazing exclusion, in SOC stocks in grasslands (Liu et al., 2024; Zhou et al., 2017).
 592 Our results ~~would~~ are in line with previous studies in mediterranean woody pastures that found a decrease in SOC storage
 593 with grazing abandonment (Oggioni et al., 2020; Peco et al., 2017). The fact that grazing abandonment particularly
 594 affected the POC fraction is surprising, since in abandoned paddocks there were a greater accumulation of plant structural
 595 biomass, that should promote POC accrual (Cotrufo et al., 2022). According to our results, the negative effect of grazing
 596 exclusion on POC stocks was mainly mediated by a reduction in microbial biomass, and a proliferation of plants with
 597 higher C/N ratios. Once again pointing to the microbially transformed nature of POC in these soils. Grazing has been
 598 proven to increase microbial activity and biomass (Bardgett et al., 2001; Zhou et al., 2017), in part by increasing root
 599 exudation (Hamilton et al., 2008; Wilson et al., 2018). Previous studies have also observed an increase in vegetation C/N
 600 ratio with cessation of grazing (He et al., 2020; Wang et al., 2016), which could be explained by the accumulation of older
 601 plant tissues and senescent standing biomass, with higher C/N than green biomass and new leaves (Sanaullah et al., 2010).

602 Importantly, we observed no interactions between management effects and climate or soil texture variables, showing that
 603 the observed management effects were consistent across the entire environmental gradient. Although such interactions

604 have been observed for wider climatic gradients (Byrnes et al., 2018; Phukubye et al., 2022), our results show that
605 management practices such as rotational grazing can have net positive impacts on SOC storage in grasslands with varying
606 conditions within semi-arid regions.

607

608 4.6. Climate effects and implications for global warming

609 Even within the small climatic gradient covered in this study, we found lower stocks of POC, MAOC and SOC in the
610 warmer areas, as observed in previous research (Díaz-Martínez et al., 2024; Georgiou et al., 2024; Hansen et al., 2024;
611 Shi et al., 2024). According to our results, these negative effects were entirely mediated by a decrease in soil fertility and
612 vegetation input quality (increased N-C axis) with increasing temperature. Increased aridity has been shown to be
613 associated with the alteration of several ecosystem processes that regulate nutrient cycling and availability (Berdugo et
614 al., 2020; Moreno-Jiménez et al., 2019). Increased fiber for tissue protection, typical from plants growing at higher
615 temperatures, may explain the increase in vegetation C/N ratios (Arroyo et al., 2024). On the other hand, annual
616 precipitation exerted a small positive effect on POC and SOC stocks, mediated by the reduction of C/N ratios in vegetation
617 in wetter farms. Based on future projections of increasing temperatures and decreasing annual precipitation in our study
618 area (IPCC, 2023b) we could expect significant reductions in SOC stocks in Mediterranean grasslands in the next decades,
619 affecting both the POC and MAOC fractions.

620

621 **5. Conclusions:**

622 Implementing rotational grazing can improve both POC and MAOC stocks in Mediterranean grasslands. On the other
623 hand, the abandonment of grazing, ~~far from functioning as a tool for ecological restoration,~~ can lead to a loss of soil
624 carbon storage capacity in these ecosystems.

625 Changes in microbial communities and vegetation attributes were the main drivers of changes in SOC stocks and fractions.
626 In this sense, this work has proven the usefulness of plant functional traits as tools for the study of plant-soil interactions
627 and SOC formation dynamics.

628 We can expect a loss of soil carbon in the studied grasslands over the next few decades due to grazing abandonment and
629 climate warming, so selecting management approaches that mitigate or counteract these losses is vital to maintaining the
630 fertility, productivity and functioning of semi-arid grasslands.

631

632 **Data and code availability**

633 The data and code that support the findings of this study are available from the corresponding author upon reasonable
634 request.

635

636 **CRedit authorship contribution statement**

637 ~~AC-Carraseosa~~: Data curation, Formal analysis, Investigation, Software, Visualization, Writing – original draft. ~~GM-~~
638 ~~Moreno~~: Conceptualization, Funding acquisition, Project administration, Supervision, Writing – review and editing.
639 ~~MFC-F. Cotrufo~~: Conceptualization, Methodology, Supervision, Writing – review and editing. ~~CF-Frade~~: Data
640 curation, Methodology, Investigation, Resources, Writing – review and editing. ~~SR-Rodrigo~~: Data curation,

641 Methodology, Investigation, Resources, Writing – review and editing. **VR-Role:** Conceptualization, Data curation,
642 Formal analysis, Funding acquisition, Investigation, Project administration, Supervision, Writing – review and editing.

643

644 **Competing interests**

645 The authors declare that they have no conflict of interest.

646

647 **Financial support**

648 This work was part of the projects “PID2019-108313RB-C33”, “PID2019-108313RB-C31” and “PCIN2021-122100-2”
649 funded by MICIU/AEI/10.13039/501100011033/ and “European Union NextGenerationEU/PRTR; the project “CLU-
650 2019-05 – IRNASA/CSIC Unit of Excellence” funded by the “Junta de Castilla y León” and co-financed by the European
651 Union (ERDF “Europe drives our growth”); and European Union’s Horizon Europe research and innovation program.
652 Grant agreement: 101059794. Alejandro Carrascosa was supported by the Spanish Ministry of Science and Innovation
653 through an FPU grant (FPU21/02668). Cristina Frade was supported by a JCyL fellowship (E-37-2023-0024090) co-
654 financed by the European Social Fund Plus (FSE+).

655

656 **References:**

- 657 Acosta-Gallo, B., Casado, M. A., Montalvo, J., and Pineda, F. D.: Allometric patterns of below-ground biomass in
658 Mediterranean grasslands, *Plant Biosyst. - Int. J. Deal. Asp. Plant Biol.*, 145, 584–595,
659 <https://doi.org/10.1080/11263504.2011.578836>, 2011.
- 660 Angst, G., Mueller, K. E., Kögel-Knabner, I., Freeman, K. H., and Mueller, C. W.: Aggregation controls the stability of
661 lignin and lipids in clay-sized particulate and mineral associated organic matter, *Biogeochemistry*, 132, 307–324,
662 <https://doi.org/10.1007/s10533-017-0304-2>, 2017.
- 663 Angst, G., Mueller, K. E., Nierop, K. G. J., and Simpson, M. J.: Plant- or microbial-derived? A review on the molecular
664 composition of stabilized soil organic matter, *Soil Biol. Biochem.*, 156, 108189,
665 <https://doi.org/10.1016/j.soilbio.2021.108189>, 2021.
- 666 Arroyo, A. I., Pueyo, Y., Barrantes, O., and Alados, C. L.: Interplay between Livestock Grazing and Aridity on the
667 Ecological and Nutritional Value of Forage in Semi-arid Mediterranean Rangelands (NE Spain), *Environ. Manage.*,
668 <https://doi.org/10.1007/s00267-024-01939-9>, 2024.
- 669 Augustine, D. J., Kearney, S. P., Raynor, E. J., Porensky, L. M., and Derner, J. D.: Adaptive, multi-paddock, rotational
670 grazing management alters foraging behavior and spatial grazing distribution of free-ranging cattle, *Agric. Ecosyst.*
671 *Environ.*, 352, 108521, <https://doi.org/10.1016/j.agee.2023.108521>, 2023.
- 672 Austin, A. T. and Vivanco, L.: Plant litter decomposition in a semi-arid ecosystem controlled by photodegradation, *Nature*,
673 442, 555–558, <https://doi.org/10.1038/nature05038>, 2006.
- 674 Bai, X., Zhai, G., Wang, B., An, S., Liu, J., Xue, Z., and Dippold, M. A.: Litter quality controls the contribution of
675 microbial carbon to main microbial groups and soil organic carbon during its decomposition, *Biol. Fertil. Soils*, 60, 167–
676 181, <https://doi.org/10.1007/s00374-023-01792-8>, 2024.
- 677 Bai, Y. and Cotrufo, M. F.: Grassland soil carbon sequestration: Current understanding, challenges, and solutions, *Science*,
678 377, 603–608, <https://doi.org/10.1126/science.abo2380>, 2022.
- 679 Baldock, J. A. and Skjemstad, J. O.: Role of the soil matrix and minerals in protecting natural organic materials against
680 biological attack, *Org. Geochem.*, 31, 697–710, [https://doi.org/10.1016/S0146-6380\(00\)00049-8](https://doi.org/10.1016/S0146-6380(00)00049-8), 2000.

681 Bardgett, R. D., Jones, A. C., Jones, D. L., Kemmitt, S. J., Cook, R., and Hobbs, P. J.: Soil microbial community patterns
682 related to the history and intensity of grazing in sub-montane ecosystems, *Soil Biol. Biochem.*, 33, 1653–1664,
683 [https://doi.org/10.1016/S0038-0717\(01\)00086-4](https://doi.org/10.1016/S0038-0717(01)00086-4), 2001.

684 Bartholomew, P. W. and Williams, R. D.: Overseeding Unimproved Warm-Season Pasture with Cool- and Warm-Season
685 Legumes to Enhance Forage Productivity, *J. Sustain. Agric.*, 34, 125–140, <https://doi.org/10.1080/10440040903482407>,
686 2010.

687 Bates, D., Mächler, M., Bolker, B., and Walker, S.: Fitting Linear Mixed-Effects Models Using lme4, *J. Stat. Softw.*, 67,
688 1–48, <https://doi.org/10.18637/jss.v067.i01>, 2015.

689 Benbi, D. K., Boparai, A. K., and Brar, K.: Decomposition of particulate organic matter is more sensitive to temperature
690 than the mineral associated organic matter, *Soil Biol. Biochem.*, 70, 183–192,
691 <https://doi.org/10.1016/j.soilbio.2013.12.032>, 2014.

692 Berdugo, M., Delgado-Baquerizo, M., Soliveres, S., Hernández-Clemente, R., Zhao, Y., Gaitán, J. J., Gross, N., Saiz, H.,
693 Maire, V., Lehmann, A., Rillig, M. C., Solé, R. V., and Maestre, F. T.: Global ecosystem thresholds driven by aridity,
694 *Science*, 367, 787–790, <https://doi.org/10.1126/science.aay5958>, 2020.

695 Berenstecher, P., Conti, G., Faigón, A., and Piñeiro, G.: Tracing service crops' net carbon and nitrogen rhizodeposition
696 into soil organic matter fractions using dual isotopic brush-labeling, *Soil Biol. Biochem.*, 184, 109096,
697 <https://doi.org/10.1016/j.soilbio.2023.109096>, 2023.

698 Bergmann, J., Weigelt, A., van der Plas, F., Laughlin, D. C., Kuyper, T. W., Guerrero-Ramirez, N., Valverde-Barrantes,
699 O. J., Bruehlheide, H., Freschet, G. T., Iversen, C. M., Kattge, J., McCormack, M. L., Meier, I. C., Rillig, M. C., Roumet,
700 C., Semchenko, M., Sweeney, C. J., van Ruijven, J., York, L. M., and Mommer, L.: The fungal collaboration gradient
701 dominates the root economics space in plants, *Sci. Adv.*, 6, eaba3756, <https://doi.org/10.1126/sciadv.aba3756>, 2020.

702 Blagodatskaya, E. and Kuzyakov, Y.: Mechanisms of real and apparent priming effects and their dependence on soil
703 microbial biomass and community structure: critical review, *Biol. Fertil. Soils*, 45, 115–131,
704 <https://doi.org/10.1007/s00374-008-0334-y>, 2008.

705 Briske, D. D., Derner, J. D., Brown, J. R., Fuhlendorf, S. D., Teague, W. R., Havstad, K. M., Gillen, R. L., Ash, A. J., and
706 Willms, W. D.: Rotational Grazing on Rangelands: Reconciliation of Perception and Experimental Evidence, *Rangel.*
707 *Ecol. Manag.*, 61, 3–17, <https://doi.org/10.2111/06-159R.1>, 2008.

708 Byrnes, R. C., Eastburn, D. J., Tate, K. W., and Roche, L. M.: A Global Meta-Analysis of Grazing Impacts on Soil Health
709 Indicators, *J. Environ. Qual.*, 47, 758–765, <https://doi.org/10.2134/jeq2017.08.0313>, 2018.

710 Cappai, C., Kemanian, A. R., Lagomarsino, A., Roggero, P. P., Lai, R., Agnelli, A. E., and Seddaiu, G.: Small-scale spatial
711 variation of soil organic matter pools generated by cork oak trees in Mediterranean agro-silvo-pastoral systems,
712 *Geoderma*, 304, 59–67, <https://doi.org/10.1016/j.geoderma.2016.07.021>, 2017.

713 Carmona, C. P., Rota, C., Azcárate, F. M., and Peco, B.: More for less: sampling strategies of plant functional traits across
714 local environmental gradients, *Funct. Ecol.*, 29, 579–588, <https://doi.org/10.1111/1365-2435.12366>, 2015.

715 Carranca, C., Castro, I. V., Figueiredo, N., Redondo, R., Rodrigues, A. R. F., Saraiva, I., Maricato, R., and Madeira, M.
716 A. V.: Influence of tree canopy on N₂ fixation by pasture legumes and soil rhizobial abundance in Mediterranean oak
717 woodlands, *Sci. Total Environ.*, 506–507, 86–94, <https://doi.org/10.1016/j.scitotenv.2014.10.111>, 2015.

718 Carranca, C., Pedra, F., and Madeira, M.: Enhancing Carbon Sequestration in Mediterranean Agroforestry Systems: A
719 Review, *Agriculture*, 12, 1598, <https://doi.org/10.3390/agriculture12101598>, 2022.

720 Carrascosa, A., Moreno, G., Rodrigo, S., and Rolo, V.: Unravelling the contribution of soil, climate and management to
721 the productivity of ecologically intensified Mediterranean wood pastures, *Sci. Total Environ.*, 957, 177575,
722 <https://doi.org/10.1016/j.scitotenv.2024.177575>, 2024.

723 Chang, Y., Sokol, N. W., van Groenigen, K. J., Bradford, M. A., Ji, D., Crowther, T. W., Liang, C., Luo, Y., Kuzyakov, Y.,
724 Wang, J., and Ding, F.: A stoichiometric approach to estimate sources of mineral-associated soil organic matter, *Glob.*
725 *Change Biol.*, 30, e17092, <https://doi.org/10.1111/gcb.17092>, 2024.

726 Cheng, J., Jing, G., Wei, L., and Jing, Z.: Long-term grazing exclusion effects on vegetation characteristics, soil properties
727 and bacterial communities in the semi-arid grasslands of China, *Ecol. Eng.*, 97, 170–178,
728 <https://doi.org/10.1016/j.ecoleng.2016.09.003>, 2016.

729 Cheng, X., Xing, W., and Liu, J.: Litter chemical traits, microbial and soil stoichiometry regulate organic carbon accrual
730 of particulate and mineral-associated organic matter, *Biol. Fertil. Soils*, 59, 777–790, [https://doi.org/10.1007/s00374-023-](https://doi.org/10.1007/s00374-023-01746-0)
731 01746-0, 2023.

732 Conant, R. T.: Grassland Soil Organic Carbon Stocks: Status, Opportunities, Vulnerability, in: *Recarbonization of the*
733 *Biosphere: Ecosystems and the Global Carbon Cycle*, edited by: Lal, R., Lorenz, K., Hüttel, R. F., Schneider, B. U., and
734 von Braun, J., Springer Netherlands, Dordrecht, 275–302, https://doi.org/10.1007/978-94-007-4159-1_13, 2012.

735 Conant, R. T., Cerri, C. E. P., Osborne, B. B., and Paustian, K.: Grassland management impacts on soil carbon stocks: a
736 new synthesis, *Ecol. Appl.*, 27, 662–668, <https://doi.org/10.1002/eap.1473>, 2017.

737 Cornwell, W. K., Cornelissen, J. H. C., Amatangelo, K., Dorrepaal, E., Eviner, V. T., Godoy, O., Hobbie, S. E., Hoorens,
738 B., Kurokawa, H., Pérez-Harguindeguy, N., Quested, H. M., Santiago, L. S., Wardle, D. A., Wright, I. J., Aerts, R., Allison,
739 S. D., Van Bodegom, P., Brovkin, V., Chatain, A., Callaghan, T. V., Díaz, S., Garnier, E., Gurvich, D. E., Kazakou, E.,
740 Klein, J. A., Read, J., Reich, P. B., Soudzilovskaia, N. A., Vaieretti, M. V., and Westoby, M.: Plant species traits are the
741 predominant control on litter decomposition rates within biomes worldwide, *Ecol. Lett.*, 11, 1065–1071,
742 <https://doi.org/10.1111/j.1461-0248.2008.01219.x>, 2008.

743 Cotrufo, M. F. and Lavelle, J. M.: Chapter One - Soil organic matter formation, persistence, and functioning: A synthesis
744 of current understanding to inform its conservation and regeneration, in: *Advances in Agronomy*, vol. 172, edited by:
745 Sparks, D. L., Academic Press, 1–66, <https://doi.org/10.1016/bs.agron.2021.11.002>, 2022.

746 Cotrufo, M. F., Wallenstein, M. D., Boot, C. M., Denef, K., and Paul, E.: The Microbial Efficiency-Matrix Stabilization
747 (MEMS) framework integrates plant litter decomposition with soil organic matter stabilization: do labile plant inputs form
748 stable soil organic matter?, *Glob. Change Biol.*, 19, 988–995, <https://doi.org/10.1111/gcb.12113>, 2013.

749 Cotrufo, M. F., Ranalli, M. G., Haddix, M. L., Six, J., and Lugato, E.: Soil carbon storage informed by particulate and
750 mineral-associated organic matter, *Nat. Geosci.*, 12, 989–994, <https://doi.org/10.1038/s41561-019-0484-6>, 2019a.

751 Cotrufo, M. F., Ranalli, M. G., Haddix, M. L., Six, J., and Lugato, E.: Soil carbon storage informed by particulate and
752 mineral-associated organic matter, *Nat. Geosci.*, 12, 989–994, <https://doi.org/10.1038/s41561-019-0484-6>, 2019b.

753 Cotrufo, M. F., Haddix, M. L., Kroeger, M. E., and Stewart, C. E.: The role of plant input physical-chemical properties,
754 and microbial and soil chemical diversity on the formation of particulate and mineral-associated organic matter, *Soil Biol.*
755 *Biochem.*, 168, 108648, <https://doi.org/10.1016/j.soilbio.2022.108648>, 2022.

756 Crowther, T. W., Todd-Brown, K. E. O., Rowe, C. W., Wieder, W. R., Carey, J. C., Machmuller, M. B., Snoek, B. L., Fang,
757 S., Zhou, G., Allison, S. D., Blair, J. M., Bridgham, S. D., Burton, A. J., Carrillo, Y., Reich, P. B., Clark, J. S., Classen, A.
758 T., Dijkstra, F. A., Elberling, B., Emmett, B. A., Estiarte, M., Frey, S. D., Guo, J., Harte, J., Jiang, L., Johnson, B. R.,
759 Kröel-Dulay, G., Larsen, K. S., Laudon, H., Lavelle, J. M., Luo, Y., Lupascu, M., Ma, L. N., Marhan, S., Michelsen, A.,
760 Mohan, J., Niu, S., Pendall, E., Peñuelas, J., Pfeifer-Meister, L., Poll, C., Reinsch, S., Reynolds, L. L., Schmidt, I. K.,
761 Sistla, S., Sokol, N. W., Templer, P. H., Treseder, K. K., Welker, J. M., and Bradford, M. A.: Quantifying global soil carbon
762 losses in response to warming, *Nature*, 540, 104–108, <https://doi.org/10.1038/nature20150>, 2016.

763 Crowther, T. W., van den Hoogen, J., Wan, J., Mayes, M. A., Keiser, A. D., Mo, L., Averill, C., and Maynard, D. S.: The
764 global soil community and its influence on biogeochemistry, *Science*, 365, eaav0550,
765 <https://doi.org/10.1126/science.aav0550>, 2019.

766 Dai, W., Xiao, R., Wei, C., and Yang, F.: Plant litter traits control the accumulation of mineral-associated organic carbon
767 by influencing its molecular composition and diversity, *Soil Tillage Res.*, 253, 106667,
768 <https://doi.org/10.1016/j.still.2025.106667>, 2025.

769 Daou, L., Garnier, É., and Shipley, B.: Quantifying the relationship linking the community-weighted means of plant traits
770 and soil fertility, *Ecology*, 102, e03454, <https://doi.org/10.1002/ecy.3454>, 2021.

771 De Deyn, G. B., Quirk, H., and Bardgett, R. D.: Plant species richness, identity and productivity differentially influence
772 key groups of microbes in grassland soils of contrasting fertility, *Biol. Lett.*, 7, 75–78,
773 <https://doi.org/10.1098/rsbl.2010.0575>, 2010.

774 Díaz, S., Kattge, J., Cornelissen, J. H. C., Wright, I. J., Lavorel, S., Dray, S., Reu, B., Kleyer, M., Wirth, C., Colin Prentice,
775 I., Garnier, E., Bönsch, G., Westoby, M., Poorter, H., Reich, P. B., Moles, A. T., Dickie, J., Gillison, A. N., Zanne, A. E.,
776 Chave, J., Joseph Wright, S., Sheremet'ev, S. N., Jactel, H., Baraloto, C., Cerabolini, B., Pierce, S., Shipley, B., Kirkup,
777 D., Casanoves, F., Joswig, J. S., Günther, A., Falczuk, V., Rüger, N., Mahecha, M. D., and Gorné, L. D.: The global
778 spectrum of plant form and function, *Nature*, 529, 167–171, <https://doi.org/10.1038/nature16489>, 2016.

779 Díaz-Martínez, P., Maestre, F. T., Moreno-Jiménez, E., Delgado-Baquerizo, M., Eldridge, D. J., Saiz, H., Gross, N., Le
780 Bagousse-Pinguet, Y., Gozalo, B., Ochoa, V., Guirado, E., García-Gómez, M., Valencia, E., Asensio, S., Berdugo, M.,
781 Martínez-Valderrama, J., Mendoza, B. J., García-Gil, J. C., Zaccone, C., Panettieri, M., García-Palacios, P., Fan, W.,
782 Benavente-Ferraces, I., Rey, A., Eisenhauer, N., Cesarz, S., Abedi, M., Ahumada, R. J., Alcántara, J. M., Amghar, F.,
783 Aramayo, V., Arroyo, A. I., Bahalkeh, K., Ben Salem, F., Blaum, N., Boldgiv, B., Bowker, M. A., Bran, D., Branquinho,
784 C., Bu, C., Cáceres, Y., Canessa, R., Castillo-Monroy, A. P., Castro, I., Castro-Quezada, P., Chibani, R., Conceição, A. A.,
785 Currier, C. M., Darrouzet-Nardi, A., Deák, B., Dickman, C. R., Donoso, D. A., Dougill, A. J., Durán, J., Ejtehadi, H.,
786 Espinosa, C., Fajardo, A., Farzam, M., Ferrante, D., Fraser, L. H., Gaitán, J. J., Gusman Montalván, E., Hernández-
787 Hernández, R. M., von Hessberg, A., Hölzel, N., Huber-Sannwald, E., Hughes, F. M., Jadán-Maza, O., Geissler, K.,
788 Jentsch, A., Ju, M., Kaseke, K. F., Kindermann, L., Koopman, J. E., Le Roux, P. C., Liancourt, P., Linstädter, A., Liu, J.,
789 Louw, M. A., Maggs-Kölling, G., Makhalanyane, T. P., Issa, O. M., Marais, E., Margerie, P., Mazaneda, A. J., McClaran,
790 M. P., Messeder, J. V. S., Mora, J. P., Moreno, G., Munson, S. M., Nunes, A., Oliva, G., Oñatibia, G. R., Osborne, B.,
791 Peter, G., Pueyo, Y., Quiroga, R. E., Reed, S. C., Reyes, V. M., et al.: Vulnerability of mineral-associated soil organic
792 carbon to climate across global drylands, *Nat. Clim. Change*, 14, 976–982, <https://doi.org/10.1038/s41558-024-02087-y>,
793 2024.

794 Dondini, M., Martin, M., De Camillis, C., Uwizeye, A., Soussana, J.-F., Robinson, T., and Steinfeld, H.: Global assessment
795 of soil carbon in grasslands – From current stock estimates to sequestration potential., *FAO*, Rome, 2023.

796 Du, Z., Angers, D. A., Ren, T., Zhang, Q., and Li, G.: The effect of no-till on organic C storage in Chinese soils should
797 not be overemphasized: A meta-analysis, *Agric. Ecosyst. Environ.*, 236, 1–11, <https://doi.org/10.1016/j.agee.2016.11.007>,
798 2017.

799 Dubeux Jr., J. c. b., Sollenberger, L. e., Vendramini, J. m. b., Interrante, S. m., and Lira Jr., M. a.: Stocking Method,
800 Animal Behavior, and Soil Nutrient Redistribution: How are They Linked?, *Crop Sci.*, 54, 2341–2350,
801 <https://doi.org/10.2135/cropsci2014.01.0076>, 2014.

802 Eisenhauer, N., Lanoue, A., Strecker, T., Scheu, S., Steinauer, K., Thakur, M. P., and Mommer, L.: Root biomass and
803 exudates link plant diversity with soil bacterial and fungal biomass, *Sci. Rep.*, 7, 44641,
804 <https://doi.org/10.1038/srep44641>, 2017.

805 Elias, D. M. O., Mason, K. E., Goodall, T., Taylor, A., Zhao, P., Otero-Fariña, A., Chen, H., Peacock, C. L., Ostle, N. J.,
806 Griffiths, R., Chapman, P. J., Holden, J., Banwart, S., McNamara, N. P., and Whitaker, J.: Microbial and mineral
807 interactions decouple litter quality from soil organic matter formation, *Nat. Commun.*, 15, 10063,
808 <https://doi.org/10.1038/s41467-024-54446-0>, 2024.

809 Engedal, T., Magid, J., Hansen, V., Rasmussen, J., Sørensen, H., and Stoumann Jensen, L.: Cover crop root morphology
810 rather than quality controls the fate of root and rhizodeposition C into distinct soil C pools, *Glob. Change Biol.*, 29, 5677–
811 5690, <https://doi.org/10.1111/gcb.16870>, 2023.

812 Eze, S., Palmer, S. M., and Chapman, P. J.: Soil organic carbon stock in grasslands: Effects of inorganic fertilizers, liming
813 and grazing in different climate settings, *J. Environ. Manage.*, 223, 74–84, <https://doi.org/10.1016/j.jenvman.2018.06.013>,
814 2018.

815 Fanin, N., Kardol, P., Farrell, M., Nilsson, M.-C., Gundale, M. J., and Wardle, D. A.: The ratio of Gram-positive to Gram-
816 negative bacterial PLFA markers as an indicator of carbon availability in organic soils, *Soil Biol. Biochem.*, 128, 111–
817 114, <https://doi.org/10.1016/j.soilbio.2018.10.010>, 2019.

818 Faucon, M.-P., Houben, D., and Lambers, H.: Plant Functional Traits: Soil and Ecosystem Services, *Trends Plant Sci.*, 22,
819 385–394, <https://doi.org/10.1016/j.tplants.2017.01.005>, 2017.

820 Fay, P. A., Prober, S. M., Harpole, W. S., Knops, J. M. H., Bakker, J. D., Borer, E. T., Lind, E. M., MacDougall, A. S.,
821 Seabloom, E. W., Wragg, P. D., Adler, P. B., Blumenthal, D. M., Buckley, Y. M., Chu, C., Cleland, E. E., Collins, S. L.,
822 Davies, K. F., Du, G., Feng, X., Firn, J., Gruner, D. S., Hagenah, N., Hautier, Y., Heckman, R. W., Jin, V. L., Kirkman, K.
823 P., Klein, J., Ladwig, L. M., Li, Q., McCulley, R. L., Melbourne, B. A., Mitchell, C. E., Moore, J. L., Morgan, J. W., Risch,
824 A. C., Schütz, M., Stevens, C. J., Wedin, D. A., and Yang, L. H.: Grassland productivity limited by multiple nutrients,
825 *Nat. Plants*, 1, 1–5, <https://doi.org/10.1038/nplants.2015.80>, 2015.

826 Feßel, C., Meier, I. C., and Leuschner, C.: Relationship between species diversity, biomass and light transmittance in
827 temperate semi-natural grasslands: is productivity enhanced by complementary light capture?, *J. Veg. Sci.*, 27, 144–155,
828 <https://doi.org/10.1111/jvs.12326>, 2016.

829 Fortunel, C., Garnier, E., Joffre, R., Kazakou, E., Quested, H., Grigulis, K., Lavorel, S., Ansquer, P., Castro, H., Cruz, P.,
830 Doležal, J., Eriksson, O., Freitas, H., Golodets, C., Jouany, C., Kigel, J., Kleyer, M., Lehsten, V., Lepš, J., Meier, T.,

831 Pakeman, R., Papadimitriou, M., Papanastasis, V. P., Quétier, F., Robson, M., Sternberg, M., Theau, J.-P., Thébault, A.,
832 and Zarovali, M.: Leaf traits capture the effects of land use changes and climate on litter decomposability of grasslands
833 across Europe, *Ecology*, 90, 598–611, <https://doi.org/10.1890/08-0418.1>, 2009.

834 Freschet, G. T., Aerts, R., and Cornelissen, J. H. C.: A plant economics spectrum of litter decomposability, *Funct. Ecol.*,
835 26, 56–65, <https://doi.org/10.1111/j.1365-2435.2011.01913.x>, 2012.

836 Frongia, A., Pulina, A., Tanda, A., Seddaiu, G., Roggero, P. P., and Moreno, G.: Assessing the effect of rotational grazing
837 adoption in Iberian silvopastoral systems with Normalized Difference Vegetation Index time series, *Ital. J. Agron.*,
838 <https://doi.org/10.4081/ija.2023.2185>, 2023.

839 Frostegård, A. and Bååth, E.: The use of phospholipid fatty acid analysis to estimate bacterial and fungal biomass in soil,
840 *Biol. Fertil. Soils*, 22, 59–65, <https://doi.org/10.1007/BF00384433>, 1996.

841 Funk, J. L., Larson, J. E., Ames, G. M., Butterfield, B. J., Cavender-Bares, J., Firn, J., Laughlin, D. C., Sutton-Grier, A.
842 E., Williams, L., and Wright, J.: Revisiting the Holy Grail: using plant functional traits to understand ecological processes,
843 *Biol. Rev.*, 92, 1156–1173, <https://doi.org/10.1111/brev.12275>, 2017.

844 Galluzzi, G., Plaza, C., Giannetta, B., Priori, S., and Zaccone, C.: Time and climate roles in driving soil carbon distribution
845 and stability in particulate and mineral-associated organic matter pools, *Sci. Total Environ.*, 963, 178511,
846 <https://doi.org/10.1016/j.scitotenv.2025.178511>, 2025.

847 Gang, C., Zhou, W., Chen, Y., Wang, Z., Sun, Z., Li, J., Qi, J., and Odeh, I.: Quantitative assessment of the contributions
848 of climate change and human activities on global grassland degradation, *Environ. Earth Sci.*, 72, 4273–4282,
849 <https://doi.org/10.1007/s12665-014-3322-6>, 2014.

850 Gao, Y., Liu, J., Wang, D., An, Y., Ma, H., and Tong, S.: Synergy and trade-off between plant functional traits enhance
851 grassland multifunctionality under grazing exclusion in a semi-arid region, *J. Environ. Manage.*, 373, 123877,
852 <https://doi.org/10.1016/j.jenvman.2024.123877>, 2025.

853 García Bravo, N., Dimas, M., Murillo Díaz, J. M., Barranco Sanz, L. M., and Ruiz Hernández, J. M.: Ejemplo de
854 aplicación del nuevo modelo hidrogeológico en SIMPA para mejorar la evaluación de los recursos hídricos a escala
855 nacional en España, Centro Nacional de Información Geográfica (España), 2023.

856 Gaudet, C. L. and Keddy, P. A.: A comparative approach to predicting competitive ability from plant traits, *Nature*, 334,
857 242–243, <https://doi.org/10.1038/334242a0>, 1988.

858 Georgiou, K., Jackson, R. B., Vindušková, O., Abramoff, R. Z., Ahlström, A., Feng, W., Harden, J. W., Pellegrini, A. F.
859 A., Polley, H. W., Soong, J. L., Riley, W. J., and Torn, M. S.: Global stocks and capacity of mineral-associated soil organic
860 carbon, *Nat. Commun.*, 13, 3797, <https://doi.org/10.1038/s41467-022-31540-9>, 2022.

861 Georgiou, K., Koven, C. D., Wieder, W. R., Hartman, M. D., Riley, W. J., Pett-Ridge, J., Bouskill, N. J., Abramoff, R. Z.,
862 Slessarev, E. W., Ahlström, A., Parton, W. J., Pellegrini, A. F. A., Pierson, D., Sulman, B. N., Zhu, Q., and Jackson, R. B.:
863 Emergent temperature sensitivity of soil organic carbon driven by mineral associations, *Nat. Geosci.*, 17, 205–212,
864 <https://doi.org/10.1038/s41561-024-01384-7>, 2024.

865 Georgiou, K., Angers, D., Champigny, R. E., Cotrufo, M. F., Craig, M. E., Doetterl, S., Grandy, A. S., Lavalley, J. M., Lin,
866 Y., Lugato, E., Poeplau, C., Rocci, K. S., Schweizer, S. A., Six, J., and Wieder, W. R.: Soil Carbon Saturation: What Do
867 We Really Know?, *Glob. Change Biol.*, 31, e70197, <https://doi.org/10.1111/gcb.70197>, 2025.

868 Gliksman, D., Navon, Y., Dumbur, R., Haenel, S., and Grünzweig, J. M.: Higher rates of decomposition in standing vs.
869 surface litter in a Mediterranean ecosystem during the dry and the wet seasons, *Plant Soil*, 428, 427–439,
870 <https://doi.org/10.1007/s11104-018-3696-4>, 2018.

871 Godde, C. M., Boone, R. B., Ash, A. J., Waha, K., Sloat, L. L., Thornton, P. K., and Herrero, M.: Global rangeland
872 production systems and livelihoods at threat under climate change and variability, *Environ. Res. Lett.*, 15, 044021,
873 <https://doi.org/10.1088/1748-9326/ab7395>, 2020.

874 Godínez-Alvarez, H., Herrick, J. E., Mattocks, M., Toledo, D., and Van Zee, J.: Comparison of three vegetation monitoring
875 methods: Their relative utility for ecological assessment and monitoring, *Ecol. Indic.*, 9, 1001–1008,
876 <https://doi.org/10.1016/j.ecolind.2008.11.011>, 2009.

877 Gómez-Rey, M. X., Garcês, A., and Madeira, M.: Soil organic-C accumulation and N availability under improved pastures
878 established in Mediterranean oak woodlands, *Soil Use Manag.*, 28, 497–507, [https://doi.org/10.1111/j.1475-](https://doi.org/10.1111/j.1475-2743.2012.00428.x)
879 2743.2012.00428.x, 2012.

880 Gou, X., Reich, P. B., Qiu, L., Shao, M., Wei, G., Wang, J., and Wei, X.: Leguminous plants significantly increase soil
881 nitrogen cycling across global climates and ecosystem types, *Glob. Change Biol.*, 29, 4028–4043,
882 <https://doi.org/10.1111/gcb.16742>, 2023.

883 Griepentrog, M., Bodé, S., Boeckx, P., Hagedorn, F., Heim, A., and Schmidt, M. W. I.: Nitrogen deposition promotes the
884 production of new fungal residues but retards the decomposition of old residues in forest soil fractions, *Glob. Change*
885 *Biol.*, 20, 327–340, <https://doi.org/10.1111/gcb.12374>, 2014.

886 Guyonnet, J. P., Cantarel, A. A. M., Simon, L., and Haichar, F. el Z.: Root exudation rate as functional trait involved in
887 plant nutrient-use strategy classification, *Ecol. Evol.*, 8, 8573–8581, <https://doi.org/10.1002/ece3.4383>, 2018.

888 Haddix, M. L., Paul, E. A., and Cotrufo, M. F.: Dual, differential isotope labeling shows the preferential movement of
889 labile plant constituents into mineral-bonded soil organic matter, *Glob. Change Biol.*, 22, 2301–2312,
890 <https://doi.org/10.1111/gcb.13237>, 2016.

891 Hamilton, E. W., Frank, D. A., Hinchey, P. M., and Murray, T. R.: Defoliation induces root exudation and triggers positive
892 rhizospheric feedbacks in a temperate grassland, *Soil Biol. Biochem.*, 40, 2865–2873,
893 <https://doi.org/10.1016/j.soilbio.2008.08.007>, 2008.

894 Hansen, P. M., Even, R., King, A. E., Lavalée, J., Schipanski, M., and Cotrufo, M. F.: Distinct, direct and climate-
895 mediated environmental controls on global particulate and mineral-associated organic carbon storage, *Glob. Change Biol.*,
896 30, e17080, <https://doi.org/10.1111/gcb.17080>, 2024.

897 Hawkins, H.-J.: A global assessment of Holistic Planned Grazing™ compared with season-long, continuous grazing:
898 meta-analysis findings, *Afr. J. Range Forage Sci.*, 34, 65–75, <https://doi.org/10.2989/10220119.2017.1358213>, 2017.

899 He, M., Zhou, G., Yuan, T., van Groenigen, K. J., Shao, J., and Zhou, X.: Grazing intensity significantly changes the C :
900 N : P stoichiometry in grassland ecosystems, *Glob. Ecol. Biogeogr.*, 29, 355–369, <https://doi.org/10.1111/geb.13028>,
901 2020.

902 Helsen, K., Ceulemans, T., Stevens, C. J., and Honnay, O.: Increasing Soil Nutrient Loads of European Semi-natural
903 Grasslands Strongly Alter Plant Functional Diversity Independently of Species Loss, *Ecosystems*, 17, 169–181,
904 <https://doi.org/10.1007/s10021-013-9714-8>, 2014.

905 den Herder, M., Moreno, G., Mosquera-Losada, R. M., Palma, J. H. N., Sidiropoulou, A., Santiago Freijanes, J. J., Crous-
906 Duran, J., Paulo, J. A., Tomé, M., Pantera, A., Papanastasis, V. P., Mantzanas, K., Pachana, P., Papadopoulos, A.,
907 Plieninger, T., and Burgess, P. J.: Current extent and stratification of agroforestry in the European Union, *Agric. Ecosyst.*
908 *Environ.*, 241, 121–132, <https://doi.org/10.1016/j.agee.2017.03.005>, 2017.

909 Hernández-Esteban, A., López-Díaz, M. L., Cáceres, Y., and Moreno, G.: Are sown legume-rich pastures effective allies
910 for the profitability and sustainability of Mediterranean dehesas?, *Agrofor. Syst.*, 93, 2047–2065,
911 <https://doi.org/10.1007/s10457-018-0307-6>, 2019.

912 Hou, Z., Wang, R., Chang, S., Zheng, Y., Ma, T., Xu, S., Zhang, X., Shi, X., Lu, J., Luo, D., Wang, B., Du, Z., and Wei,
913 Y.: The contribution of microbial necromass to soil organic carbon and influencing factors along a variation of habitats in
914 alpine ecosystems, *Sci. Total Environ.*, 921, 171126, <https://doi.org/10.1016/j.scitotenv.2024.171126>, 2024.

915 Hu, Y., Deng, Q., Kätterer, T., Olesen, J. E., Ying, S. C., Ochoa-Hueso, R., Mueller, C. W., Weintraub, M. N., and Chen,
916 J.: Depth-dependent responses of soil organic carbon under nitrogen deposition, *Glob. Change Biol.*, 30, e17247,
917 <https://doi.org/10.1111/gcb.17247>, 2024.

918 Huang, J., Liu, W., Pan, S., Wang, Z., Yang, S., Jia, Z., Wang, Z., Deng, M., Yang, L., Liu, C., Chang, P., and Liu, L.:
919 Divergent contributions of living roots to turnover of different soil organic carbon pools and their links to plant traits,
920 *Funct. Ecol.*, 35, 2821–2830, <https://doi.org/10.1111/1365-2435.13934>, 2021.

921 Intergovernmental Panel on Climate Change (IPCC) (Ed.): Climate Change Information for Regional Impact and for Risk
922 Assessment, in: Climate Change 2021 – The Physical Science Basis: Working Group I Contribution to the Sixth
923 Assessment Report of the Intergovernmental Panel on Climate Change, Cambridge University Press, Cambridge, 1767–
924 1926, <https://doi.org/10.1017/9781009157896.014>, 2023a.

925 Intergovernmental Panel on Climate Change (IPCC) (Ed.): Global Carbon and Other Biogeochemical Cycles and
926 Feedbacks, in: Climate Change 2021 – The Physical Science Basis: Working Group I Contribution to the Sixth Assessment
927 Report of the Intergovernmental Panel on Climate Change, Cambridge University Press, Cambridge, 673–816,
928 <https://doi.org/10.1017/9781009157896.007>, 2023b.

929 Jacobo, E. J., Rodríguez, A. M., Bartoloni, N., and Deregibus, V. A.: Rotational Grazing Effects on Rangeland Vegetation
930 at a Farm Scale, *Rangel. Ecol. Manag.*, 59, 249–257, <https://doi.org/10.2111/05-129R1.1>, 2006.

931 Jaurena, M., Lezama, F., Salvo, L., Cardozo, G., Ayala, W., Terra, J., and Nabinger, C.: The Dilemma of Improving Native
932 Grasslands by Overseeding Legumes: Production Intensification or Diversity Conservation, *Rangel. Ecol. Manag.*, 69,
933 35–42, <https://doi.org/10.1016/j.rama.2015.10.006>, 2016.

934 Jongen, M., Förster, A. C., and Unger, S.: Overwhelming effects of autumn-time drought during seedling establishment
935 impair recovery potential in sown and semi-natural pastures in Portugal, *Plant Ecol.*, 220, 183–197,
936 <https://doi.org/10.1007/s11258-018-0869-4>, 2019.

937 Kallenbach, C. M., Frey, S. D., and Grandy, A. S.: Direct evidence for microbial-derived soil organic matter formation
938 and its ecophysiological controls, *Nat. Commun.*, 7, 13630, <https://doi.org/10.1038/ncomms13630>, 2016.

939 Kazakou, E., Violle, C., Roumet, C., Pintor, C., Gimenez, O., and Garnier, E.: Litter quality and decomposability of
940 species from a Mediterranean succession depend on leaf traits but not on nitrogen supply, *Ann. Bot.*, 104, 1151–1161,
941 <https://doi.org/10.1093/aob/mcp202>, 2009.

942 Khatiwada, B., Acharya, S. N., Larney, F. J., Lupwayi, N. Z., Smith, E. G., Islam, M. A., and Thomas, J. E.: Benefits of
943 mixed grass–legume pastures and pasture rejuvenation using bloat-free legumes in western Canada: a review, *Can. J.*
944 *Plant Sci.*, 100, 463–476, <https://doi.org/10.1139/cjps-2019-0212>, 2020.

945 Khatri-Chhetri, U., Thompson, K. A., Quideau, S. A., Boyce, M. S., Chang, S. X., Bork, E. W., and Carlyle, C. N.:
946 Adaptive multi-paddock grazing increases mineral associated soil carbon in Northern grasslands, *Agric. Ecosyst.*
947 *Environ.*, 369, 109000, <https://doi.org/10.1016/j.agee.2024.109000>, 2024.

948 King, A. E., Amsili, J. P., Córdova, S. C., Culman, S., Fonte, S. J., Kotcon, J., Liebig, M., Masters, M. D., McVay, K.,
949 Olk, D. C., Schipanski, M., Schneider, S. K., Stewart, C. E., and Cotrufo, M. F.: A soil matrix capacity index to predict
950 mineral-associated but not particulate organic carbon across a range of climate and soil pH, *Biogeochemistry*, 165, 1–14,
951 <https://doi.org/10.1007/s10533-023-01066-3>, 2023.

952 Klumpp, K., Fontaine, S., Attard, E., Le Roux, X., Gleixner, G., and Soussana, J.-F.: Grazing triggers soil carbon loss by
953 altering plant roots and their control on soil microbial community, *J. Ecol.*, 97, 876–885, <https://doi.org/10.1111/j.1365-2745.2009.01549.x>, 2009.

955 Kramer, C. and Gleixner, G.: Soil organic matter in soil depth profiles: Distinct carbon preferences of microbial groups
956 during carbon transformation, *Soil Biol. Biochem.*, 40, 425–433, <https://doi.org/10.1016/j.soilbio.2007.09.016>, 2008.

957 Kramer-Walter, K. R., Bellingham, P. J., Millar, T. R., Smissen, R. D., Richardson, S. J., and Laughlin, D. C.: Root traits
958 are multidimensional: specific root length is independent from root tissue density and the plant economic spectrum, *J.*
959 *Ecol.*, 104, 1299–1310, <https://doi.org/10.1111/1365-2745.12562>, 2016.

960 Laliberté, E. and Legendre, P.: A distance-based framework for measuring functional diversity from multiple traits,
961 *Ecology*, 91, 299–305, <https://doi.org/10.1890/08-2244.1>, 2010.

962 Laliberté, E. and Tylianakis, J. M.: Cascading effects of long-term land-use changes on plant traits and ecosystem
963 functioning, *Ecology*, 93, 145–155, <https://doi.org/10.1890/11-0338.1>, 2012.

964 Lange, M., Eisenhauer, N., Sierra, C. A., Bessler, H., Engels, C., Griffiths, R. I., Mellado-Vázquez, P. G., Malik, A. A.,
965 Roy, J., Scheu, S., Steinbeiss, S., Thomson, B. C., Trumbore, S. E., and Gleixner, G.: Plant diversity increases soil
966 microbial activity and soil carbon storage, *Nat. Commun.*, 6, 6707, <https://doi.org/10.1038/ncomms7707>, 2015.

967 Latimer, G. W., Jr. (Ed.): Official Methods of Analysis of AOAC INTERNATIONAL, Oxford University Press,
968 <https://doi.org/10.1093/9780197610145.001.0001>, 2023.

969 Lavalley, J. M., Conant, R. T., Paul, E. A., and Cotrufo, M. F.: Incorporation of shoot versus root-derived ¹³C and ¹⁵N
970 into mineral-associated organic matter fractions: results of a soil slurry incubation with dual-labelled plant material,
971 *Biogeochemistry*, 137, 379–393, <https://doi.org/10.1007/s10533-018-0428-z>, 2018.

972 Lavalley, J. M., Soong, J. L., and Cotrufo, M. F.: Conceptualizing soil organic matter into particulate and mineral-
973 associated forms to address global change in the 21st century, *Glob. Change Biol.*, 26, 261–273,
974 <https://doi.org/10.1111/gcb.14859>, 2020.

975 Lefcheck, J. S.: piecewiseSEM: Piecewise structural equation modelling in r for ecology, evolution, and systematics,
976 *Methods Ecol. Evol.*, 7, 573–579, <https://doi.org/10.1111/2041-210X.12512>, 2016.

977 Lei, J., Guo, X., Zeng, Y., Zhou, J., Gao, Q., and Yang, Y.: Temporal changes in global soil respiration since 1987, *Nat.*
978 *Commun.*, 12, 403, <https://doi.org/10.1038/s41467-020-20616-z>, 2021.

979 Leuthold, S., Lavalley, J. M., Haddix, M. L., and Cotrufo, M. F.: Contrasting properties of soil organic matter fractions
980 isolated by different physical separation methodologies, *Geoderma*, 445, 116870,
981 <https://doi.org/10.1016/j.geoderma.2024.116870>, 2024.

982 Li, S. and Li, X.: Global understanding of farmland abandonment: A review and prospects, *J. Geogr. Sci.*, 27, 1123–1150,
983 <https://doi.org/10.1007/s11442-017-1426-0>, 2017.

984 Li, Y., Luo, T., and Lu, Q.: Plant height as a simple predictor of the root to shoot ratio: Evidence from alpine grasslands
985 on the Tibetan Plateau, *J. Veg. Sci.*, 19, 245–252, <https://doi.org/10.3170/2007-8-18365>, 2008.

986 Liang, C., Amelung, W., Lehmann, J., and Kästner, M.: Quantitative assessment of microbial necromass contribution to
987 soil organic matter, *Glob. Change Biol.*, 25, 3578–3590, <https://doi.org/10.1111/gcb.14781>, 2019.

988 Liu, H. Y., Huang, N., Zhao, C. M., and Li, J. H.: Responses of carbon cycling and soil organic carbon content to nitrogen
989 addition in grasslands globally, *Soil Biol. Biochem.*, 186, 109164, <https://doi.org/10.1016/j.soilbio.2023.109164>, 2023.

990 Liu, Y., Zhang, M., Wang, X., and Wang, C.: The impact of different grazing intensity and management measures on soil
991 organic carbon density in Zhangye grassland, *Sci. Rep.*, 14, 17556, <https://doi.org/10.1038/s41598-024-68277-y>, 2024.

992 Lu, M., Zhou, X., Luo, Y., Yang, Y., Fang, C., Chen, J., and Li, B.: Minor stimulation of soil carbon storage by nitrogen
993 addition: A meta-analysis, *Agric. Ecosyst. Environ.*, 140, 234–244, <https://doi.org/10.1016/j.agee.2010.12.010>, 2011.

994 Luo, R., Fan, J., Wang, W., Luo, J., Kuzyakov, Y., He, J.-S., Chu, H., and Ding, W.: Nitrogen and phosphorus enrichment
995 accelerates soil organic carbon loss in alpine grassland on the Qinghai-Tibetan Plateau, *Sci. Total Environ.*, 650, 303–
996 312, <https://doi.org/10.1016/j.scitotenv.2018.09.038>, 2019.

997 Luo, R., Kuzyakov, Y., Liu, D., Fan, J., Luo, J., Lindsey, S., He, J.-S., and Ding, W.: Nutrient addition reduces carbon
998 sequestration in a Tibetan grassland soil: Disentangling microbial and physical controls, *Soil Biol. Biochem.*, 144,
999 107764, <https://doi.org/10.1016/j.soilbio.2020.107764>, 2020.

1000 Maestre, F. T., Le Bagousse-Pinguet, Y., Delgado-Baquerizo, M., Eldridge, D. J., Saiz, H., Berdugo, M., Gozalo, B.,
1001 Ochoa, V., Guirado, E., García-Gómez, M., Valencia, E., Gaitán, J. J., Asensio, S., Mendoza, B. J., Plaza, C., Díaz-
1002 Martínez, P., Rey, A., Hu, H.-W., He, J.-Z., Wang, J.-T., Lehmann, A., Rillig, M. C., Cesarz, S., Eisenhauer, N., Martínez-
1003 Valderrama, J., Moreno-Jiménez, E., Sala, O., Abedi, M., Ahmadian, N., Alados, C. L., Aramayo, V., Amghar, F.,
1004 Arredondo, T., Ahumada, R. J., Bahalkeh, K., Ben Salem, F., Blaum, N., Boldgiv, B., Bowker, M. A., Bran, D., Bu, C.,
1005 Canessa, R., Castillo-Monroy, A. P., Castro, H., Castro, I., Castro-Quezada, P., Chibani, R., Conceição, A. A., Currier, C.
1006 M., Darrouzet-Nardi, A., Deák, B., Donoso, D. A., Dougill, A. J., Durán, J., Erdenetsetseg, B., Espinosa, C. I., Fajardo,
1007 A., Farzam, M., Ferrante, D., Frank, A. S. K., Fraser, L. H., Gherardi, L. A., Greenville, A. C., Guerra, C. A., Gusmán-
1008 Montalvan, E., Hernández-Hernández, R. M., Hölzel, N., Huber-Sannwald, E., Hughes, F. M., Jadán-Maza, O., Jeltsch,
1009 F., Jentsch, A., Kaseke, K. F., Köbel, M., Koopman, J. E., Leder, C. V., Linstädter, A., le Roux, P. C., Li, X., Liancourt,
1010 P., Liu, J., Louw, M. A., Maggs-Kölling, G., Makhilanyane, T. P., Issa, O. M., Manzaneda, A. J., Marais, E., Mora, J. P.,
1011 Moreno, G., Munson, S. M., Nunes, A., Oliva, G., Oñatibia, G. R., Peter, G., Pivari, M. O. D., Pueyo, Y., Quiroga, R. E.,
1012 Rahmanian, S., Reed, S. C., et al.: Grazing and ecosystem service delivery in global drylands, *Science*, 378, 915–920,
1013 <https://doi.org/10.1126/science.abq4062>, 2022.

1014 Manning, P., de Vries, F. T., Tallowin, J. R. B., Smith, R., Mortimer, S. R., Pilgrim, E. S., Harrison, K. A., Wright, D. G.,
1015 Quirk, H., Benson, J., Shipley, B., Cornelissen, J. H. C., Kattge, J., Bönisch, G., Wirth, C., and Bardgett, R. D.: Simple
1016 measures of climate, soil properties and plant traits predict national-scale grassland soil carbon stocks, *J. Appl. Ecol.*, 52,
1017 1188–1196, <https://doi.org/10.1111/1365-2664.12478>, 2015.

1018 McSherry, M. E. and Ritchie, M. E.: Effects of grazing on grassland soil carbon: a global review, *Glob. Change Biol.*, 19,
1019 1347–1357, <https://doi.org/10.1111/gcb.12144>, 2013.

- 1020 Mokany, K., Ash, J., and Roxburgh, S.: Functional identity is more important than diversity in influencing ecosystem
1021 processes in a temperate native grassland, *J. Ecol.*, 96, 884–893, <https://doi.org/10.1111/j.1365-2745.2008.01395.x>, 2008.
- 1022 Moreno, G. and Pulido, F. J.: The Functioning, Management and Persistence of Dehesas, in: *Agroforestry in Europe:*
1023 *Current Status and Future Prospects*, edited by: Rigueiro-Rodríguez, A., McAdam, J., and Mosquera-Losada, M. R.,
1024 Springer Netherlands, Dordrecht, 127–160, https://doi.org/10.1007/978-1-4020-8272-6_7, 2009.
- 1025 Moreno, G., Obrador, J. J., Cubera, E., and Dupraz, C.: Fine Root Distribution in Dehesas of Central-Western Spain, *Plant*
1026 *Soil*, 277, 153–162, <https://doi.org/10.1007/s11104-005-6805-0>, 2005.
- 1027 Moreno, G., Hernández-Esteban, A., Rolo, V., and Igual, J. M.: The enduring effects of sowing legume-rich mixtures on
1028 the soil microbial community and soil carbon in semi-arid wood pastures, *Plant Soil*, 465, 563–582,
1029 <https://doi.org/10.1007/s11104-021-05023-7>, 2021.
- 1030 Moreno-Jiménez, E., Plaza, C., Saiz, H., Manzano, R., Flagmeier, M., and Maestre, F. T.: Aridity and reduced soil
1031 micronutrient availability in global drylands, *Nat. Sustain.*, 2, 371–377, <https://doi.org/10.1038/s41893-019-0262-x>,
1032 2019.
- 1033 Mortensen, E. Ø., Abalos, D., Engedal, T., Lægsgaard, A. K., Enggrob, K., Mueller, C. W., and Rasmussen, J.: Smart
1034 Mixture Design Can Steer the Fate of Root-Derived Carbon Into Mineral-Associated and Particulate Organic Matter in
1035 Intensively Managed Grasslands, *Glob. Change Biol.*, 31, e70117, <https://doi.org/10.1111/gcb.70117>, 2025.
- 1036 Mosier, S., Apfelbaum, S., Byck, P., Calderon, F., Teague, R., Thompson, R., and Cotrufo, M. F.: Adaptive multi-paddock
1037 grazing enhances soil carbon and nitrogen stocks and stabilization through mineral association in southeastern U.S.
1038 grazing lands, *J. Environ. Manage.*, 288, 112409, <https://doi.org/10.1016/j.jenvman.2021.112409>, 2021.
- 1039 Murphy, M. V.: *semEff: Automatic Calculation of Effects for Piecewise Structural Equation Models*, 2022.
- 1040 Nair, R. K. F., Morris, K. A., Hertel, M., Luo, Y., Moreno, G., Reichstein, M., Schrumpf, M., and Migliavacca, M.:
1041 N  P stoichiometry and habitat effects on Mediterranean savanna seasonal root dynamics,
1042 *Biogeosciences*, 16, 1883–1901, <https://doi.org/10.5194/bg-16-1883-2019>, 2019.
- 1043 Niu, W., Ding, J., Fu, B., Zhao, W., and Eldridge, D.: Global effects of livestock grazing on ecosystem functions vary
1044 with grazing management and environment, *Agric. Ecosyst. Environ.*, 378, 109296,
1045 <https://doi.org/10.1016/j.agee.2024.109296>, 2025.
- 1046 Novelty, P. E. and Watson, I. W.: Successful Grassland Regeneration in a Severely Degraded Catchment: a Whole of
1047 Government Approach in North West Australia, in: *Climate and Land Degradation*, edited by: Sivakumar, M. V. K. and
1048 Ndiang’ui, N., Springer, Berlin, Heidelberg, 469–485, https://doi.org/10.1007/978-3-540-72438-4_26, 2007.
- 1049 Oggioni, S. D., Ochoa-Hueso, R., and Peco, B.: Livestock grazing abandonment reduces soil microbial activity and carbon
1050 storage in a Mediterranean Dehesa, *Appl. Soil Ecol.*, 153, 103588, <https://doi.org/10.1016/j.apsoil.2020.103588>, 2020.
- 1051 Oliveira Filho, J. de S., Vieira, J. N., Ribeiro da Silva, E. M., Beserra de Oliveira, J. G., Pereira, M. G., and Brasileiro, F.
1052 G.: Assessing the effects of 17 years of grazing exclusion in degraded semi-arid soils: Evaluation of soil fertility, nutrients
1053 pools and stoichiometry, *J. Arid Environ.*, 166, 1–10, <https://doi.org/10.1016/j.jaridenv.2019.03.006>, 2019.
- 1054 Ordoñez, J. C., Van Bodegom, P. M., Witte, J.-P. M., Wright, I. J., Reich, P. B., and Aerts, R.: A global study of
1055 relationships between leaf traits, climate and soil measures of nutrient fertility, *Glob. Ecol. Biogeogr.*, 18, 137–149,
1056 <https://doi.org/10.1111/j.1466-8238.2008.00441.x>, 2009.
- 1057 Palomo-Campesino, S., Ravera, F., González, J. A., and García-Llorente, M.: Exploring Current and Future Situation of
1058 Mediterranean Silvopastoral Systems: Case Study in Southern Spain, *Rangel. Ecol. Manag.*, 71, 578–591,
1059 <https://doi.org/10.1016/j.rama.2017.12.013>, 2018.
- 1060 Parras-Alcántara, L., Díaz-Jaimes, L., Lozano-García, B., Rebollo, P. F., Elcure, F. M., and Muñoz, M. D. C.: Organic
1061 farming has little effect on carbon stock in a Mediterranean dehesa (southern Spain), *CATENA*, 113, 9–17,
1062 <https://doi.org/10.1016/j.catena.2013.09.002>, 2014.
- 1063 Parras-Alcántara, L., Díaz-Jaimes, L., and Lozano-García, B.: Management Effects on Soil Organic Carbon Stock in
1064 Mediterranean Open Rangelands—Treeless Grasslands, *Land Degrad. Dev.*, 26, 22–34, <https://doi.org/10.1002/ldr.2269>,
1065 2015.

1066 Peco, B., Navarro, E., Carmona, C. P., Medina, N. G., and Marques, M. J.: Effects of grazing abandonment on soil
1067 multifunctionality: The role of plant functional traits, *Agric. Ecosyst. Environ.*, 249, 215–225,
1068 <https://doi.org/10.1016/j.agee.2017.08.013>, 2017.

1069 Pérez-Harguindeguy, N., Díaz, S., Garnier, E., Lavorel, S., Poorter, H., Jaureguiberry, P., Bret-Harte, M. S., Cornwell, W.
1070 K., Craine, J. M., Gurvich, D. E., Urcelay, C., Veneklaas, E. J., Reich, P. B., Poorter, L., Wright, I. J., Ray, P., Enrico, L.,
1071 Pausas, J. G., Vos, A. C. de, Buchmann, N., Funes, G., Quétier, F., Hodgson, J. G., Thompson, K., Morgan, H. D., Steege,
1072 H. ter, Sack, L., Blonder, B., Poschlod, P., Vaieretti, M. V., Conti, G., Staver, A. C., Aquino, S., and Cornelissen, J. H. C.:
1073 Corrigendum to: New handbook for standardised measurement of plant functional traits worldwide, *Aust. J. Bot.*, 64,
1074 715–716, https://doi.org/10.1071/bt12225_co, 2016.

1075 Phukubye, K., Mutema, M., Buthelezi, N., Muchaonyerwa, P., Cerri, C., and Chaplot, V.: On the impact of grassland
1076 management on soil carbon stocks: a worldwide meta-analysis, *Geoderma Reg.*, 28, e00479,
1077 <https://doi.org/10.1016/j.geodrs.2021.e00479>, 2022.

1078 Pino, A. D., Rodríguez, T., and Andión, J.: Production improvement through phosphorus fertilization and legume
1079 introduction in grazed native pastures of Uruguay, *J. Agric. Sci.*, 154, 347–358,
1080 <https://doi.org/10.1017/S002185961500101X>, 2016.

1081 Plaza, C., García-Palacios, P., Berhe, A. A., Barquero, J., Bastida, F., Png, G. K., Rey, A., Bardgett, R. D., and Delgado-
1082 Baquerizo, M.: Ecosystem productivity has a stronger influence than soil age on surface soil carbon storage across global
1083 biomes, *Commun. Earth Environ.*, 3, 1–8, <https://doi.org/10.1038/s43247-022-00567-7>, 2022.

1084 Poeplau, C., Vos, C., and Don, A.: Soil organic carbon stocks are systematically overestimated by misuse of the parameters
1085 bulk density and rock fragment content, *SOIL*, 3, 61–66, <https://doi.org/10.5194/soil-3-61-2017>, 2017.

1086 Poeplau, C., Helfrich, M., Dechow, R., Szoboszlay, M., Tebbe, C. C., Don, A., Greiner, B., Zopf, D., Thumm, U.,
1087 Korevaar, H., and Geerts, R.: Increased microbial anabolism contributes to soil carbon sequestration by mineral
1088 fertilization in temperate grasslands, *Soil Biol. Biochem.*, 130, 167–176, <https://doi.org/10.1016/j.soilbio.2018.12.019>,
1089 2019.

1090 Poeplau, C., Dechow, R., Begill, N., and Don, A.: Towards an ecosystem capacity to stabilise organic carbon in soils,
1091 *Glob. Change Biol.*, 30, e17453, <https://doi.org/10.1111/gcb.17453>, 2024.

1092 Poorter, H., Niinemets, Ü., Poorter, L., Wright, I. J., and Villar, R.: Causes and consequences of variation in leaf mass per
1093 area (LMA): a meta-analysis, *New Phytol.*, 182, 565–588, <https://doi.org/10.1111/j.1469-8137.2009.02830.x>, 2009.

1094 Porqueddu, C., Ates, S., Louhaichi, M., Kyriazopoulos, A. P., Moreno, G., del Pozo, A., Ovalle, C., Ewing, M. A., and
1095 Nichols, P. G. H.: Grasslands in ‘Old World’ and ‘New World’ Mediterranean-climate zones: past trends, current status
1096 and future research priorities, *Grass Forage Sci.*, 71, 1–35, <https://doi.org/10.1111/gfs.12212>, 2016.

1097 Pulina, A., Rolo, V., Hernández-Esteban, A., Seddaiu, G., Roggero, P. P., and Moreno, G.: Long-term legacy of sowing
1098 legume-rich mixtures in Mediterranean wooded grasslands, *Agric. Ecosyst. Environ.*, 348, 108397,
1099 <https://doi.org/10.1016/j.agee.2023.108397>, 2023.

1100 R Core Team: R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna,
1101 Austria. URL <https://www.R-project.org/>, 2024.

1102 Rama, G., Oyarzabal, M., Cardozo, G., Lezama, F., and Baeza, S.: Legume Overseeding along with P Fertilization
1103 Increase Forage Production of Temperate Natural Grasslands, *Agronomy*, 12, 2507,
1104 <https://doi.org/10.3390/agronomy12102507>, 2022.

1105 Ricotta, C. and Moretti, M.: CWM and Rao’s quadratic diversity: a unified framework for functional ecology, *Oecologia*,
1106 167, 181–188, <https://doi.org/10.1007/s00442-011-1965-5>, 2011.

1107 Ridgeway, J. R., Morrissey, E. M., and Brzostek, E. R.: Plant litter traits control microbial decomposition and drive soil
1108 carbon stabilization, *Soil Biol. Biochem.*, 175, 108857, <https://doi.org/10.1016/j.soilbio.2022.108857>, 2022.

1109 Rocci, K. S., Lavalley, J. M., Stewart, C. E., and Cotrufo, M. F.: Soil organic carbon response to global environmental
1110 change depends on its distribution between mineral-associated and particulate organic matter: A meta-analysis, *Sci. Total*
1111 *Environ.*, 793, 148569, <https://doi.org/10.1016/j.scitotenv.2021.148569>, 2021.

1112 Rocci, K. S., Barker, K. S., Seabloom, E. W., Borer, E. T., Hobbie, S. E., Bakker, J. D., MacDougall, A. S., McCulley, R.
1113 L., Moore, J. L., Raynaud, X., Stevens, C. J., and Cotrufo, M. F.: Impacts of nutrient addition on soil carbon and nitrogen

stoichiometry and stability in globally-distributed grasslands, *Biogeochemistry*, 159, 353–370, <https://doi.org/10.1007/s10533-022-00932-w>, 2022.

Rolo, V., Rivest, D., Lorente, M., Kattge, J., and Moreno, G.: Taxonomic and functional diversity in Mediterranean pastures: insights on the biodiversity–productivity trade-off, *J. Appl. Ecol.*, 53, 1575–1584, <https://doi.org/10.1111/1365-2664.12685>, 2016.

Roumet, C., Birouste, M., Picon-Cochard, C., Ghestem, M., Osman, N., Vrignon-Brenas, S., Cao, K., and Stokes, A.: Root structure–function relationships in 74 species: evidence of a root economics spectrum related to carbon economy, *New Phytol.*, 210, 815–826, <https://doi.org/10.1111/nph.13828>, 2016.

Rovira, P., Sauras-Yera, T., and Romanyà, J.: Equivalent-mass versus fixed-depth as criteria for quantifying soil carbon sequestration: How relevant is the difference?, *CATENA*, 214, 106283, <https://doi.org/10.1016/j.catena.2022.106283>, 2022.

Sanaullah, M., Chabbi, A., Lemaire, G., Charrier, X., and Rumpel, C.: How does plant leaf senescence of grassland species influence decomposition kinetics and litter compounds dynamics?, *Nutr. Cycl. Agroecosystems*, 88, 159–171, <https://doi.org/10.1007/s10705-009-9323-2>, 2010.

Sanderman, J., Baldock, J. A., Dangal, S. R. S., Ludwig, S., Potter, S., Rivard, C., and Savage, K.: Soil organic carbon fractions in the Great Plains of the United States: an application of mid-infrared spectroscopy, *Biogeochemistry*, 156, 97–114, <https://doi.org/10.1007/s10533-021-00755-1>, 2021.

Sanjari, G., Yu, B., Ghadiri, H., Ciesiolka, C. A. A., and Rose, C. W.: Effects of time-controlled grazing on runoff and sediment loss, *Soil Res.*, 47, 796–808, <https://doi.org/10.1071/SR09032>, 2009.

Santos, R. S., Hamilton, E. K., Stanley, P. L., Paustian, K., Cotrufo, M. F., and Zhang, Y.: Simulating adaptive grazing management on soil organic carbon in the Southeast U.S.A. using MEMS 2, *J. Environ. Manage.*, 365, 121657, <https://doi.org/10.1016/j.jenvman.2024.121657>, 2024.

Sherrod, L. A., Dunn, G., Peterson, G. A., and Kolberg, R. L.: Inorganic Carbon Analysis by Modified Pressure-*Calcimeter Method*, *Soil Sci. Soc. Am. J.*, 66, 299–305, <https://doi.org/10.2136/sssaj2002.2990>, 2002.

Shi, B., Delgado-Baquerizo, M., Knapp, A. K., Smith, M. D., Reed, S., Osborne, B., Carrillo, Y., Maestre, F. T., Zhu, Y., Chen, A., Wilkins, K., Holdrege, M. C., Kulmatiski, A., Picon-Cochard, C., Roscher, C., Power, S., Byrne, K. M., Churchill, A. C., Jentsch, A., Henry, H. A. L., Beard, K. H., Schuchardt, M. A., Eisenhauer, N., Otfinowski, R., Hautier, Y., Shen, H., Wang, Y., Wang, Z., Wang, C., Cusack, D. F., Petraglia, A., Carbognani, M., Forte, T. G. W., Flory, S., Hou, P., Zhang, T., Gao, W., and Sun, W.: Aridity drives the response of soil total and particulate organic carbon to drought in temperate grasslands and shrublands, *Sci. Adv.*, 10, eadq2654, <https://doi.org/10.1126/sciadv.adq2654>, 2024.

Siefert, A., Violle, C., Chalmandrier, L., Albert, C. H., Taudiere, A., Fajardo, A., Aarssen, L. W., Baraloto, C., Carlucci, M. B., Cianciaruso, M. V., de L. Dantas, V., de Bello, F., Duarte, L. D. S., Fonseca, C. R., Freschet, G. T., Gaucherand, S., Gross, N., Hikosaka, K., Jackson, B., Jung, V., Kamiyama, C., Katabuchi, M., Kembel, S. W., Kichenin, E., Kraft, N. J. B., Lagerström, A., Bagousse-Pinguet, Y. L., Li, Y., Mason, N., Messier, J., Nakashizuka, T., Overton, J. McC., Peltzer, D. A., Pérez-Ramos, I. M., Pillar, V. D., Prentice, H. C., Richardson, S., Sasaki, T., Schamp, B. S., Schöb, C., Shipley, B., Sundqvist, M., Sykes, M. T., Vandewalle, M., and Wardle, D. A.: A global meta-analysis of the relative extent of intraspecific trait variation in plant communities, *Ecol. Lett.*, 18, 1406–1419, <https://doi.org/10.1111/ele.12508>, 2015.

Six, J., Guggenberger, G., Paustian, K., Haumaier, L., Elliott, E. T., and Zech, W.: Sources and composition of soil organic matter fractions between and within soil aggregates, *Eur. J. Soil Sci.*, 52, 607–618, <https://doi.org/10.1046/j.1365-2389.2001.00406.x>, 2001.

Six, J., Conant, R. T., Paul, E. A., and Paustian, K.: Stabilization mechanisms of soil organic matter: Implications for C-saturation of soils, *Plant Soil*, 241, 155–176, <https://doi.org/10.1023/A:1016125726789>, 2002.

Smith, M. D., Wilkins, K. D., Holdrege, M. C., Wilfahrt, P., Collins, S. L., Knapp, A. K., Sala, O. E., Dukes, J. S., Phillips, R. P., Yahdjian, L., Gherardi, L. A., Ohlert, T., Beier, C., Fraser, L. H., Jentsch, A., Loik, M. E., Maestre, F. T., Power, S. A., Yu, Q., Felton, A. J., Munson, S. M., Luo, Y., Abdoli, H., Abedi, M., Alados, C. L., Alberti, J., Alon, M., An, H., Anacker, B., Anderson, M., Auge, H., Bachle, S., Bahalkheh, K., Bahn, M., Batbaatar, A., Bauerle, T., Beard, K. H., Behn, K., Beil, I., Biancarini, L., Blindow, I., Bondaruk, V. F., Borer, E. T., Bork, E. W., Bruschetti, C. M., Byrne, S. X., Cahill Jr., J. F., Calvo, D. A., Carbognani, M., Cardoni, A., Carlyle, C. N., Castillo-Garcia, M., Chang, S. X., Chieppa, J., Cianciaruso, M. V., Cohen, O., Cordeiro, A. L., Cusack, D. F., Dahlke, S., Daleo, P., D’Antonio, C. M., Dietterich, L. H., S. Doherty, T., Dubbert, M., Ebeling, A., Eisenhauer, N., Fischer, F. M., Forte, T. G. W., Gebauer, T., Gozalo, B., Greenville, A. C., Guidoni-Martins, K. G., Hannusch, H. J., Vatsø Haugum, S., Hautier, Y., Hefting, M., Henry, H. A. L.,

- Hoss, D., Ingrisch, J., Iribarne, O., Isbell, F., Johnson, Y., Jordan, S., Kelly, E. F., Kimmel, K., Kreyling, J., Kröel-Dulay, G., Kröpfl, A., Kübert, A., Kulmatiski, A., Lamb, E. G., Larsen, K. S., Larson, J., Lawson, J., Leder, C. V., Linstädter, A., Liu, J., Liu, S., Lodge, A. G., et al.: Extreme drought impacts have been underestimated in grasslands and shrublands globally, *Proc. Natl. Acad. Sci.*, 121, e2309881120, <https://doi.org/10.1073/pnas.2309881120>, 2024.
- Soliveres, S., Maestre, F. T., Eldridge, D. J., Delgado-Baquerizo, M., Quero, J. L., Bowker, M. A., and Gallardo, A.: Plant diversity and ecosystem multifunctionality peak at intermediate levels of woody cover in global drylands, *Glob. Ecol. Biogeogr.*, 23, 1408–1416, <https://doi.org/10.1111/geb.12215>, 2014.
- Soussana, J.-F. and Lemaire, G.: Coupling carbon and nitrogen cycles for environmentally sustainable intensification of grasslands and crop-livestock systems, *Agric. Ecosyst. Environ.*, 190, 9–17, <https://doi.org/10.1016/j.agee.2013.10.012>, 2014.
- Spohn, M., Pötsch, E. M., Eichorst, S. A., Wobken, D., Wanek, W., and Richter, A.: Soil microbial carbon use efficiency and biomass turnover in a long-term fertilization experiment in a temperate grassland, *Soil Biol. Biochem.*, 97, 168–175, <https://doi.org/10.1016/j.soilbio.2016.03.008>, 2016.
- Stanley, P., Roche, L., and Bowles, T.: Amping up soil carbon: soil carbon stocks in California rangelands under adaptive multi-paddock and conventional grazing management, *Int. J. Agric. Sustain.*, 2025.
- Stanley, P. L., Wilson, C., Patterson, E., Machmuller, M. B., and Cotrufo, M. F.: Ruminating on soil carbon: Applying current understanding to inform grazing management, *Glob. Change Biol.*, 30, e17223, <https://doi.org/10.1111/gcb.17223>, 2024.
- Steinbeiss, S., BEßLER, H., Engels, C., Temperton, V. M., Buchmann, N., Roscher, C., Kreutziger, Y., Baade, J., Habekost, M., and Gleixner, G.: Plant diversity positively affects short-term soil carbon storage in experimental grasslands, *Glob. Change Biol.*, 14, 2937–2949, <https://doi.org/10.1111/j.1365-2486.2008.01697.x>, 2008.
- Strickland, M. S. and Rousk, J.: Considering fungal:bacterial dominance in soils – Methods, controls, and ecosystem implications, *Soil Biol. Biochem.*, 42, 1385–1395, <https://doi.org/10.1016/j.soilbio.2010.05.007>, 2010.
- Tang, K., Wu, C., Wang, S., Liao, W., Yin, L., Zhou, W., and Cui, H.-J.: Distribution characteristics of soil organic carbon fractions in paddy profiles with 40 years of fertilization under two groundwater levels, *J. Soils Sediments*, 24, 681–691, <https://doi.org/10.1007/s11368-023-03664-y>, 2024.
- Tang, Z., Feng, J., Chen, L., Chen, Z., Shao, X., and Xia, T.: Coupling amendment of microbial and compound fertilizers increases fungal necromass carbon and soil organic carbon by regulating microbial activity in flue-cured tobacco-planted field, *Eur. J. Soil Biol.*, 117, 103518, <https://doi.org/10.1016/j.ejsobi.2023.103518>, 2023.
- Tao, F., Huang, Y., Hungate, B. A., Manzoni, S., Frey, S. D., Schmidt, M. W. I., Reichstein, M., Carvalhais, N., Ciais, P., Jiang, L., Lehmann, J., Wang, Y.-P., Houlton, B. Z., Ahrens, B., Mishra, U., Hugelius, G., Hocking, T. D., Lu, X., Shi, Z., Viatkin, K., Vargas, R., Yigini, Y., Omuto, C., Malik, A. A., Peralta, G., Cuevas-Corona, R., Di Paolo, L. E., Luotto, I., Liao, C., Liang, Y.-S., Saynes, V. S., Huang, X., and Luo, Y.: Microbial carbon use efficiency promotes global soil carbon storage, *Nature*, 618, 981–985, <https://doi.org/10.1038/s41586-023-06042-3>, 2023.
- Tate, K. W., Atwill, E. R., McDougald, N. K., and George, M. R.: Spatial and Temporal Patterns of Cattle Feces Deposition on Rangeland, *J. Range Manag.*, 56, 432–438, <https://doi.org/10.2307/4003833>, 2003.
- Teague, R., Provenza, F., Kreuter, U., Steffens, T., and Barnes, M.: Multi-paddock grazing on rangelands: Why the perceptual dichotomy between research results and rancher experience?, *J. Environ. Manage.*, 128, 699–717, <https://doi.org/10.1016/j.jenvman.2013.05.064>, 2013.
- Teague, W. R., Dowhower, S. L., and Waggoner, J. A.: Drought and grazing patch dynamics under different grazing management, *J. Arid Environ.*, 58, 97–117, [https://doi.org/10.1016/S0140-1963\(03\)00122-8](https://doi.org/10.1016/S0140-1963(03)00122-8), 2004.
- Teague, W. R., Dowhower, S. L., Baker, S. A., Haile, N., DeLaune, P. B., and Conover, D. M.: Grazing management impacts on vegetation, soil biota and soil chemical, physical and hydrological properties in tall grass prairie, *Agric. Ecosyst. Environ.*, 141, 310–322, <https://doi.org/10.1016/j.agee.2011.03.009>, 2011.
- Teixeira, R. F. M., Proença, V., Crespo, D., Valada, T., and Domingos, T.: A conceptual framework for the analysis of engineered biodiverse pastures, *Ecol. Eng.*, 77, 85–97, <https://doi.org/10.1016/j.ecoleng.2015.01.002>, 2015.
- Thomson, F. J., Moles, A. T., Auld, T. D., and Kingsford, R. T.: Seed dispersal distance is more strongly correlated with plant height than with seed mass, *J. Ecol.*, 99, 1299–1307, <https://doi.org/10.1111/j.1365-2745.2011.01867.x>, 2011.

- 1213 Tiessen, H., Cuevas, E., and Chacon, P.: The role of soil organic matter in sustaining soil fertility, *Nature*, 371, 783–785,
1214 <https://doi.org/10.1038/371783a0>, 1994.
- 1215 Uribe, C., Inclán, R., Hernando, L., Román, M., Clavero, M. A., Roig, S., and Van Miegroet, H.: Grazing, tilling and
1216 canopy effects on carbon dioxide fluxes in a Spanish dehesa, *Agrofor. Syst.*, 89, 305–318, [https://doi.org/10.1007/s10457-](https://doi.org/10.1007/s10457-014-9767-5)
1217 014-9767-5, 2015.
- 1218 Van Soest, P. J., Robertson, J. B., and Lewis, B. A.: Methods for Dietary Fiber, Neutral Detergent Fiber, and Nonstarch
1219 Polysaccharides in Relation to Animal Nutrition, *J. Dairy Sci.*, 74, 3583–3597, [https://doi.org/10.3168/jds.S0022-](https://doi.org/10.3168/jds.S0022-0302(91)78551-2)
1220 0302(91)78551-2, 1991.
- 1221 Villarino, S. H., Pinto, P., Jackson, R. B., and Piñeiro, G.: Plant rhizodeposition: A key factor for soil organic matter
1222 formation in stable fractions, *Sci. Adv.*, 7, eabd3176, <https://doi.org/10.1126/sciadv.abd3176>, 2021.
- 1223 di Virgilio, A., Lambertucci, S. A., and Morales, J. M.: Sustainable grazing management in rangelands: Over a century
1224 searching for a silver bullet, *Agric. Ecosyst. Environ.*, 283, 106561, <https://doi.org/10.1016/j.agee.2019.05.020>, 2019.
- 1225 Waldrop, M. P. and Firestone, M. K.: Microbial community utilization of recalcitrant and simple carbon compounds:
1226 impact of oak-woodland plant communities, *Oecologia*, 138, 275–284, <https://doi.org/10.1007/s00442-003-1419-9>, 2004.
- 1227 Wang, J., Liu, L., Wang, X., and Chen, Y.: The interaction between abiotic photodegradation and microbial decomposition
1228 under ultraviolet radiation, *Glob. Change Biol.*, 21, 2095–2104, <https://doi.org/10.1111/gcb.12812>, 2015.
- 1229 Wang, J., Liu, L., Wang, X., Yang, S., Zhang, B., Li, P., Qiao, C., Deng, M., and Liu, W.: High night-time humidity and
1230 dissolved organic carbon content support rapid decomposition of standing litter in a semi-arid landscape, *Funct. Ecol.*,
1231 31, 1659–1668, <https://doi.org/10.1111/1365-2435.12854>, 2017.
- 1232 Wang, K., Ma, Z., Qin, W., Li, X., Shi, H., Hasi, B., and Liu, X.: Soil nutrients and pH modulate carbon dynamics in
1233 particulate and mineral-associated organic matter during restoration of a Tibetan alpine grassland, *Ecol. Eng.*, 212,
1234 107522, <https://doi.org/10.1016/j.ecoleng.2025.107522>, 2025.
- 1235 Wang, X., McConkey, B. G., VandenBygaart, A. J., Fan, J., Iwaasa, A., and Schellenberg, M.: Grazing improves C and N
1236 cycling in the Northern Great Plains: a meta-analysis, *Sci. Rep.*, 6, 33190, <https://doi.org/10.1038/srep33190>, 2016.
- 1237 Ward, S. E., Smart, S. M., Quirk, H., Tallowin, J. R. B., Mortimer, S. R., Shiel, R. S., Wilby, A., and Bardgett, R. D.:
1238 Legacy effects of grassland management on soil carbon to depth, *Glob. Change Biol.*, 22, 2929–2938,
1239 <https://doi.org/10.1111/gcb.13246>, 2016.
- 1240 Weigelt, A., Mommer, L., Andrzejek, K., Iversen, C. M., Bergmann, J., Bruelheide, H., Fan, Y., Freschet, G. T., Guerrero-
1241 Ramírez, N. R., Kattge, J., Kuiper, T. W., Laughlin, D. C., Meier, I. C., van der Plas, F., Poorter, H., Roumet, C., van
1242 Ruijven, J., Sabatini, F. M., Semchenko, M., Sweeney, C. J., Valverde-Barrantes, O. J., York, L. M., and McCormack, M.
1243 L.: An integrated framework of plant form and function: the belowground perspective, *New Phytol.*, 232, 42–59,
1244 <https://doi.org/10.1111/nph.17590>, 2021.
- 1245 Westerband, A. C., Funk, J. L., and Barton, K. E.: Intraspecific trait variation in plants: a renewed focus on its role in
1246 ecological processes, *Ann. Bot.*, 127, 397–410, <https://doi.org/10.1093/aob/mcab011>, 2021.
- 1247 White, R.: Pilot analysis of global ecosystems: Grassland ecosystems, 2000.
- 1248 Willers, C., Jansen van Rensburg, P. J., and Claassens, S.: Phospholipid fatty acid profiling of microbial communities—a
1249 review of interpretations and recent applications, *J. Appl. Microbiol.*, 119, 1207–1218, <https://doi.org/10.1111/jam.12902>,
1250 2015.
- 1251 Wilson, C. H., Strickland, M. S., Hutchings, J. A., Bianchi, T. S., and Flory, S. L.: Grazing enhances belowground carbon
1252 allocation, microbial biomass, and soil carbon in a subtropical grassland, *Glob. Change Biol.*, 24, 2997–3009,
1253 <https://doi.org/10.1111/gcb.14070>, 2018.
- 1254 Wilson, P. J., Thompson, K., and Hodgson, J. G.: Specific leaf area and leaf dry matter content as alternative predictors
1255 of plant strategies, *New Phytol.*, 143, 155–162, <https://doi.org/10.1046/j.1469-8137.1999.00427.x>, 1999.
- 1256 Wright, I. J., Reich, P. B., Westoby, M., Ackerly, D. D., Baruch, Z., Bongers, F., Cavender-Bares, J., Chapin, T.,
1257 Cornelissen, J. H. C., Diemer, M., Flexas, J., Garnier, E., Groom, P. K., Gulias, J., Hikosaka, K., Lamont, B. B., Lee, T.,
1258 Lee, W., Lusk, C., Midgley, J. J., Navas, M.-L., Niinemets, Ü., Oleksyn, J., Osada, N., Poorter, H., Poot, P., Prior, L.,

- 1259 Pyankov, V. I., Roumet, C., Thomas, S. C., Tjoelker, M. G., Veneklaas, E. J., and Villar, R.: The worldwide leaf economics
1260 spectrum, *Nature*, 428, 821–827, <https://doi.org/10.1038/nature02403>, 2004.
- 1261 Xu, H., Zhu, B., Wei, X., Yu, M., and Cheng, X.: Root functional traits mediate rhizosphere soil carbon stability in a
1262 subtropical forest, *Soil Biol. Biochem.*, 162, 108431, <https://doi.org/10.1016/j.soilbio.2021.108431>, 2021.
- 1263 Xu, X., Thornton, P. E., and Post, W. M.: A global analysis of soil microbial biomass carbon, nitrogen and phosphorus in
1264 terrestrial ecosystems, *Glob. Ecol. Biogeogr.*, 22, 737–749, <https://doi.org/10.1111/geb.12029>, 2013.
- 1265 Yao, H., He, Z., Wilson, M. J., and Campbell, C. D.: Microbial Biomass and Community Structure in a Sequence of Soils
1266 with Increasing Fertility and Changing Land Use, *Microb. Ecol.*, 40, 223–237, <https://doi.org/10.1007/s002480000053>,
1267 2000.
- 1268 Yu, L., Sun, W., and Huang, Y.: Grazing exclusion enhances plant and topsoil carbon stocks in arid and semiarid
1269 grasslands, *Agric. Ecosyst. Environ.*, 320, 107605, <https://doi.org/10.1016/j.agee.2021.107605>, 2021.
- 1270 Yu, W., Huang, W., Weintraub-Leff, S. R., and Hall, S. J.: Where and why do particulate organic matter (POM) and
1271 mineral-associated organic matter (MAOM) differ among diverse soils?, *Soil Biol. Biochem.*, 172, 108756,
1272 <https://doi.org/10.1016/j.soilbio.2022.108756>, 2022.
- 1273 Zhang, Q., Buyantuev, A., Li, F. Y., Jiang, L., Niu, J., Ding, Y., Kang, S., and Ma, W.: Functional dominance rather than
1274 taxonomic diversity and functional diversity mainly affects community aboveground biomass in the Inner Mongolia
1275 grassland, *Ecol. Evol.*, 7, 1605–1615, <https://doi.org/10.1002/ece3.2778>, 2017.
- 1276 Zhang, Y., Lavalley, J. M., Robertson, A. D., Even, R., Ogle, S. M., Paustian, K., and Cotrufo, M. F.: Simulating
1277 measurable ecosystem carbon and nitrogen dynamics with the mechanistically defined MEMS 2.0 model, *Biogeosciences*,
1278 18, 3147–3171, <https://doi.org/10.5194/bg-18-3147-2021>, 2021.
- 1279 Zhang, Y., Wang, T., Yan, C., Li, Y., Mo, F., and Han, J.: Microbial life-history strategies and particulate organic carbon
1280 mediate formation of microbial necromass carbon and stabilization in response to biochar addition, *Sci. Total Environ.*,
1281 950, 175041, <https://doi.org/10.1016/j.scitotenv.2024.175041>, 2024.
- 1282 Zhong, W., Gu, T., Wang, W., Zhang, B., Lin, X., Huang, Q., and Shen, W.: The effects of mineral fertilizer and organic
1283 manure on soil microbial community and diversity, *Plant Soil*, 326, 511–522, <https://doi.org/10.1007/s11104-009-9988->
1284 y, 2010.
- 1285 Zhou, G., Zhou, X., He, Y., Shao, J., Hu, Z., Liu, R., Zhou, H., and Hosseinibai, S.: Grazing intensity significantly affects
1286 belowground carbon and nitrogen cycling in grassland ecosystems: a meta-analysis, *Glob. Change Biol.*, 23, 1167–1179,
1287 <https://doi.org/10.1111/gcb.13431>, 2017.
- 1288 Zhou, G., Luo, Q., Chen, Y., He, M., Zhou, L., Frank, D., He, Y., Fu, Y., Zhang, B., and Zhou, X.: Effects of livestock
1289 grazing on grassland carbon storage and release override impacts associated with global climate change, *Glob. Change*
1290 *Biol.*, 25, 1119–1132, <https://doi.org/10.1111/gcb.14533>, 2019.
- 1291 Zhou, Z., Ren, C., Wang, C., Delgado-Baquerizo, M., Luo, Y., Luo, Z., Du, Z., Zhu, B., Yang, Y., Jiao, S., Zhao, F., Cai,
1292 A., Yang, G., and Wei, G.: Global turnover of soil mineral-associated and particulate organic carbon, *Nat. Commun.*, 15,
1293 5329, <https://doi.org/10.1038/s41467-024-49743-7>, 2024.
- 1294 Zuur, A. F., Ieno, E. N., Walker, N., Saveliev, A. A., and Smith, G. M.: Mixed effects models and extensions in ecology
1295 with R, Springer, New York, NY, <https://doi.org/10.1007/978-0-387-87458-6>, 2009.

1296