

Improved management increases soil mineral-protected organic carbon storage via plant-microbial-nutrient mediation in semi-arid grasslands

Alejandro Carrascosa^a, Gerardo Moreno^a, M. Francesca Cotrufo^b, Cristina Frade^c, Sara Rodrigo^a,
V́ctor Rolo^a

^a Forest Research Group, INDEHESA, University of Extremadura, 10600, Plasencia, Cáceres, Spain

^b Department of Soil and Crop Sciences, Colorado State University, Fort Collins, CO, USA

^c Institute of Natural Resources and Agrobiology of Salamanca (IRNASA-CSIC), Salamanca, Spain

Correspondence to: V́ctor Rolo (rolo@unex.es)

Abstract

Soil organic carbon (SOC) storage in semi-arid grasslands is threatened by both climate change and land degradation, impacting food production and climate regulation. Improved management has been proposed to increase SOC stocks and overcome these challenges. However, the benefits of improved management practices in semi-arid regions are in question. Little is known about the effects of management on the functional components of SOC, particulate (POC) and mineral-associated organic carbon (MAOC), which are expected to respond differently, and about the pathways that mediate these responses, such as changes in vegetation and soil microbial communities.

This work analyses the effect of rotational grazing, legumes sowing and grazing exclusion on topsoil (0-8 cm) SOC, POC and MAOC stocks in Mediterranean wooded grasslands compared to continuous conventional grazing. Changes in plant diversity and morpho-biochemical traits, soil fertility and microbial composition were also evaluated. A total of 188 plots were sampled in 9 farms across a wide environmental gradient.

More resource-acquisitive, nitrogen-rich and less lignified plant community, higher soil microbial biomass with lower Gram+/Gram- ratio, and higher soil fertility were associated with higher SOC storage, with similar impacts on POC and MAOC. Rotational grazing increased POC, MAOC and total SOC stocks by 15%, 15% and 14% respectively, compared to continuous grazing. This effect was mediated by an increase in soil fertility in the rotationally grazed paddocks. On the other hand, grazing exclusion reduced POC stocks by 12% compared to continuous grazing. This depletion was mainly due to a reduction in microbial biomass and litter quality of vegetation in non-grazed paddocks. Both POC and MAOC stocks were lower at the warmer sites.

We conclude that rotational grazing can enhance long-term SOC storage in semi-arid grasslands, thereby increasing their resilience and climate mitigation capacity, whereas abandoning grazing could lead to SOC losses.

Keywords

35 Soil organic matter fractions; Soil microbial community; Plant traits; Rotational grazing; Legume enrichment; Grazing
36 exclusion; Functional diversity; Mediterranean grassland

37

38 1. Introduction

39 Grasslands cover 40.5% of the world's ice-free land area and store one third of the terrestrial carbon (C) stocks, mainly
40 (over 80%) in the form of soil organic matter (SOM; White, 2000). This C pool in grassland soils surpasses the global
41 aboveground vegetation C pool (IPCC, 2023a), highlighting the importance of grasslands SOM in global climate
42 regulation (Bai and Cotrufo, 2022). In addition, SOM is a cornerstone of grassland productivity and functioning (Cotrufo
43 and Lavallee, 2022; Tiessen et al., 1994). However, C storage in grassland soils is threatened by increasing C outputs (i.e.,
44 soil respiration and erosion) and decreasing C inputs (i.e., primary productivity) due to anthropogenic impacts such as
45 land degradation and climate change (Crowther et al., 2016; Gang et al., 2014; Godde et al., 2020; Lei et al., 2021).
46 Improved grassland management offers a key opportunity to counter these threats by increasing soil organic C (SOC)
47 stocks and supporting both climate change mitigation and adaptation (Bai and Cotrufo, 2022; Conant, 2012; Dondini et
48 al., 2023; Stanley et al., 2024). This is particularly valuable in the case of semi-arid grasslands, which represent the
49 majority of the global grassland area (White, 2000) but are also more vulnerable to climate change impacts than wetter
50 grasslands (Smith et al., 2024).

51 Practices such as rotational grazing, legumes sowing and grazing exclusion have been proposed to increase and conserve
52 SOC stocks (Conant et al., 2017; Yu et al., 2021). Rotational grazing encompasses a variety of practices where small
53 paddocks are grazed at high intensity for short periods of time, allowing for longer pasture rest than continuous grazing
54 (Teague et al., 2013). This management favors vegetation recovery after defoliation and reduces grazing patchiness and
55 livestock selectivity (Jacobo et al., 2006; Teague et al., 2013). The effects of rotational grazing on grassland productivity
56 and animal performance are under debate (Briske et al., 2008; Teague et al., 2013; di Virgilio et al., 2019) but most studies
57 have found positive effects on SOC stocks associated with this practice (Byrnes et al., 2018; Conant et al., 2017; Phukubye
58 et al., 2022; Teague et al., 2011). Sowing legumes in natural pastures is practiced worldwide, having clear positive impacts
59 on SOC stocks (Carranca et al., 2022; Conant et al., 2017; Moreno et al., 2021) and grassland productivity (Bartholomew
60 and Williams, 2010; Carrascosa et al., 2024; Jaurena et al., 2016; Khatiwada et al., 2020; Rama et al., 2022). Grazing
61 exclusion is widely advocated as a tool for ecosystem restoration (Cheng et al., 2016; Novelly and Watson, 2007) and is
62 also a global trend driven by land abandonment, particularly in high-income countries (Li and Li, 2017). The effects of
63 grazing exclusion on SOC in grasslands are mixed, showing both positive (Cheng et al., 2016; Yu et al., 2021) and negative
64 outcomes (Wilson et al., 2018). Moreover, the net effect of management practices has been shown to depend on the
65 environmental context (Maestre et al., 2022; McSherry and Ritchie, 2013; Niu et al., 2025), and global meta-analyses
66 remain inconclusive. While some global studies have reported greater benefits on SOC stocks from grazing exclusion and
67 rotational grazing in arid climates (Zhou et al., 2017), others found these management practices to be more beneficial in
68 wetter climates (Byrnes et al., 2018; Hawkins, 2017; Zhou et al., 2019). To the best of our knowledge, no similar studies
69 have addressed the interaction between environmental conditions and the effect of legume sowings on SOC stocks. Thus,
70 some questions remain open and further research is needed to clarify the net effects of improved management practices
71 on SOC storage under different environmental conditions and particularly in semi-arid grasslands.

72 To better understand the controls and vulnerability of C in soils, SOM can be conceptualized into particulate organic
73 matter (POM) and mineral-associated organic matter (MAOM) fractions (Cotrufo and Lavallee, 2022; Lavallee et al.,
74 2020). POM originates from fragmented structural plant inputs and, to a lesser extent, microbial recalcitrant compounds
75 (Angst et al., 2021; Six et al., 2001). In contrast, MAOM forms through the sorption of microbial necromass and soluble

76 plant inputs onto soil mineral surfaces (Angst et al., 2021; Cotrufo et al., 2022). The mineral bonds partially protect
 77 MAOM from decomposition (Baldock and Skjemstad, 2000), while POM is readily accessible to microbial degradation,
 78 although occlusion within soil aggregates can reduce its accessibility (Angst et al., 2017). Hence, POM accumulation is
 79 mainly controlled by environmental constraints on microbial activity, e.g. low temperatures and highly acidic soils
 80 (Hansen et al., 2024; Yu et al., 2022; Zhou et al., 2024). For the same reasons, POM is more vulnerable to climatic
 81 warming (Benbi et al., 2014; Georgiou et al., 2024; Rocci et al., 2021) and the mean residence time of C in POM (years
 82 to decades) is on average shorter than in MAOM (decades to centuries; Zhou et al., 2024). However, MAOM storage in
 83 soil is theoretically limited by the availability of free mineral surface area (i.e. clay and silt content; Six et al., 2002), and
 84 a saturation point can be observed, where no more MAOM accumulate despite increases in total SOM contents (Cotrufo
 85 et al., 2019a; Georgiou et al., 2022). Management changes have been shown to influence SOM fractions in grassland soils
 86 (Khatri-Chhetri et al., 2024; Mosier et al., 2021; Oliveira Filho et al., 2019), and a conceptual framework for relating
 87 grazing management to SOM distribution has recently been proposed (Stanley et al., 2024). Yet, the underlying processes
 88 mediating these effects, such as alterations in vegetation or soil microbial communities (Laliberté and Tylianakis, 2012;
 89 Peco et al., 2017; Wilson et al., 2018), remain poorly understood.

90 Primary productivity is the point of entry of C into soil, and consequently the amount of plant inputs regulates SOM
 91 accrual (King et al., 2023; Zhou et al., 2024), but microbial processing largely determines the fate of that C (Cotrufo and
 92 Lavallee, 2022; Crowther et al., 2019). In this sense, the chemical composition of plant inputs, and the soil microbiota
 93 carbon use efficiency (CUE), i.e. the amount of C used for microbial growth and products relative to total C uptake, play
 94 a crucial role in the SOM formation process (Cotrufo and Lavallee, 2022; Tao et al., 2023). Recalcitrant plant inputs [i.e.,
 95 high carbon-nitrogen ratio (C/N) and lignin content] tend to promote short-term SOM accumulation, primarily as POM,
 96 due to their chemical resistance to decomposition (Cheng et al., 2023; Cotrufo and Lavallee, 2022). However, as outlined
 97 in the Microbial Efficiency-Matrix Stabilization (MEMS) framework (Cotrufo et al., 2013), recalcitrant inputs are less
 98 efficiently decomposed by microbes, leading to greater C losses in the long term, compared to labile (i.e., water-soluble,
 99 low C/N and lignin content) plant inputs (Cotrufo and Lavallee, 2022; Ridgeway et al., 2022). Thus, labile plant inputs
 100 are expected to enhance MAOM formation and SOM stocks in the long term, due to their faster and more efficient
 101 decomposition (Cheng et al., 2023; Haddix et al., 2016). Elias et al. (2024) added complexity to these assumptions,
 102 showing that plant input characteristics may favor certain microbial groups over others, altering the overall CUE of the
 103 microbial community. For example, fungi, which are often assumed to have a higher CUE than bacteria (Kallenbach et
 104 al., 2016; Strickland and Rousk, 2010), are favored by the addition of recalcitrant inputs (Bai et al., 2024; Strickland and
 105 Rousk, 2010). Substrate preferences have been also identify for Gram-positive (Gram+) and Gram-negative (Gram-)
 106 bacteria (Fanin et al., 2019; Kramer and Gleixner, 2008), with consequences for SOC accrual (Klumpp et al., 2009).
 107 Importantly, much of the research on the influence of plant input characteristics and microbial communities on SOM
 108 formation dynamics has relied on incubation experiments (Cheng et al., 2023; Haddix et al., 2016; Ridgeway et al., 2022)
 109 and there is limited information on how these findings translate to natural field conditions. Other vegetation characteristics
 110 such as species richness has been shown to positively influence SOC stocks (Lange et al., 2015; Steinbeiss et al., 2008),
 111 but its effects on SOC fractions have been poorly evaluated, with inconclusive results in grasslands (Mortensen et al.,
 112 2025). In addition, the relationships between SOM stocks and fractions and plant functional traits have rarely been studied
 113 (Manning et al., 2015; Mortensen et al., 2025; Xu et al., 2021) despite the latter being widely used to predict ecosystem
 114 functioning and responses (Funk et al., 2017). Plant functional traits are highly correlated with processes such as litter
 115 decomposition (Cornwell et al., 2008; Fortunel et al., 2009; Kazakou et al., 2009) or root exudates production (Guyonnet
 116 et al., 2018) and may therefore be a promising tool to study the relationships between vegetation, soil microbiota and
 117 SOM formation dynamics (Faucon et al., 2017).

118 The aim of this work is to evaluate the impact of rotational grazing, legume sowing, and grazing exclusion on topsoil (0-
 119 8 cm) SOC stocks, and their distribution between POM and MAOM fractions, in semi-arid grasslands, compared to
 120 conventional continuous grazing. We also evaluate changes in vegetation characteristics (identity and diversity of
 121 biochemical and morphological traits), soil nutrients, and soil microbial communities as possible pathways through which
 122 management might indirectly affect SOC stocks and fractions. This study focuses on the Iberian *dehesas* Mediterranean
 123 woody grassland, the main example of semi-arid grasslands in Europe (Porqueddu et al., 2016). The most widespread
 124 livestock management in this ecosystem is continuous grazing, but in recent decades rotational grazing and legume sowing
 125 have gained importance (Frangia et al., 2023; Pulina et al., 2023). At the same time, the amount of ungrazed pastures has
 126 increased due to land abandonment (Palomo-Campesino et al., 2018). Iberian *dehesas* occupy 3.1 million hectares,
 127 spanning a wide environmental gradient, and have been subject to extensive grazing for centuries (Moreno and Pulido,
 128 2009), making them an ideal model system for assessing the effects of improved management on SOC stocks in semi-
 129 arid grasslands. In particular, we designed our study to answer the following questions: 1) What are the effects of the
 130 different management practices on bulk SOC and fractions stocks? 2) Are these effects mediated by changes in vegetation
 131 or soil microbial communities? 3) Is SOC storage in these grasslands, and its enhancement, modulated or limited by
 132 environmental factors such as climate or soil properties? And 4) are these mechanisms and controls the same or different
 133 for C in POM or MAOM? Understanding the potential and limitations of improved management on SOC storage in POM
 134 and MAOM in semi-arid grasslands can guide policymakers in enhancing the climate change adaptation and mitigation
 135 capacity of these ecosystems, while supporting productivity and soil fertility.

136

137 2. Methods

138 2.1. Study area and experimental design

139 The study was carried out at nine commercial *dehesa* farms located along a north-south gradient in the western part of
 140 the Iberian Peninsula (Fig. 1a). The region has a continental Mediterranean climate, but on a local scale, in relative terms,
 141 farms can be grouped into three main climatic regions (Fig. 1b, c) according to the average climate for the period 1980–
 142 2018 (García Bravo et al., 2023). A cold-dry region [12.9 °C mean annual temperature (MAT) and 445 mm mean annual
 143 precipitation (MAP)] in the north; a warm-wet region (17.3 °C MAT and 603 mm MAP) in the middle of the latitudinal
 144 gradient; and a warm-dry region (17.0 °C MAT and 510 mm MAP) in the south. The soils of the farms share a common
 145 development from granites, shales and sandy tertiary sediments, are acidic and poor in organic matter, but cover a wide
 146 texture gradient (Fig. 1d). In these farms, native pastures are often combined with scattered trees, as is common in
 147 Mediterranean and semi-arid rangelands (den Herder et al., 2017; Soliveres et al., 2014). The tree layer is dominated by
 148 holm oaks (*Quercus ilex* L.) with scattered cork oaks (*Quercus suber* L.) or gall oaks (*Quercus faginea* Lam.). The
 149 herbaceous layer is composed of species typical of Mediterranean pastures and presents a high diversity and proportion
 150 of annual C3 plants (Table S1). The growing season of the pasture is very limited by the Mediterranean summer drought
 151 with the annual species germinating in autumn (early-to mid-October), reaching their peak productivity in mid-spring
 152 (late April of the following year) and senescing in June.

153 *Dehesa* farms are typically managed by extensive continuous grazing, where livestock (mainly cattle) graze freely on
 154 large areas following a loosely defined grazing plan. This management has been traditionally practiced in all the farms
 155 studied, until, in recent decades, some paddocks were converted to other management practices. As a result, on each farm
 156 we selected five paddocks, each with one of the following management regimes:

157 - Abandoned (Ab): Paddocks devoid of grazing for at least the last 10-20 years.

158 - Continuous grazing (Ct): Paddocks where livestock stand most of the year, fed by grazing and supplemented. This is the
159 control treatment in our study as it is the most widespread management in Iberian *dehesas*.

160 - Rotational grazing (Ro): Paddocks intensively grazed in short periods and with resting periods (i.e. without livestock)
161 lasting for more than six months. Rotational grazing has been applied in these paddocks for the last 10-15 years.

162 - Recent legume sowing (Lr): Paddocks where pastures have recently (≤ 5 years) been sown with legume mixtures. In
163 *dehesa* farms, legume sowing consists of sowing a mixture of seeds (at a rate of 20 kg ha⁻¹) from various species of annual
164 legumes (pre-inoculated with Rhizobium) such as *Trifolium subterraneum* L., *T. incarnatum* L., *T. michelianum* Savi., *T.*
165 *resupinatum* L., *T. mutabile* Port. and *Ornithopus sativus* Brot., along with some highly productive annual grass species
166 such as *Lolium multiflorum* Lam. and *Lolium rigidum* Gaud. (Teixeira et al., 2015). These sowings are preceded by surface
167 tillage and phosphorus application to meet the needs of the legumes and stimulate N-fixation (Jongen et al., 2019).
168 Farmers saw legumes mixture only once, because its effect persists over the years thanks to the natural seeds production
169 and the resulting soil seed bank of the sown species. None of the sown plots studied was resown.

170 - Old legume sowing (Lo): Paddocks sown with legume mixtures more than 10 years ago.

171 The distinction between recent and old legumes sown paddocks allows us to compare short and long-term effects after
172 sowing. In two out of the nine farms, we added an additional control (continuous grazing) paddock close to the legume
173 sowing paddocks because they were at a considerable distance from the other control paddock, limiting their
174 comparability. Additional information on paddocks characteristics is provided in Table S2.

175 In each paddock, we installed four permanent monitoring plots, in open grassland and away from the direct influence of
176 the trees (at least 10 m away from the trees). This setup allowed us to have a total of 188 sampling plots where we
177 measured soil properties, SOC fractions, and soil microbial communities in 2021 and vegetation traits for 3 years (2021
178 to 2023).

179




Figure 1. Iberian “dehesa” cover according to the CORINE 2018 land cover survey and geographical location of the studied farms (a). Mean annual temperature (b) and precipitation (c) during the period 1980-2018 in the study area. Soil texture in all sampled plots (d).

2.3. Vegetation sampling

We assessed above-ground (ABG) pasture productivity over three growing periods at all plots using 50 x 50 x 50 cm grazing exclusion cages. In the 2020-2021 period (hereafter referred to as 2021) we installed one cage on each monitoring plot at the beginning of spring (late March 2021). In the 2021-2022 and 2022-2023 periods (hereafter referred to as 2022 and 2023), we installed the cages at the beginning of the green-up in early November. Standing biomass at the time of cage establishment (t_0) was determined by hand clipping within two 25 x 25 cm squares, randomly placed in the proximity of the cage. In the three years we harvested ABG biomass inside the cages at the beginning of summer in early June (t_1), when most of the vegetation was already senescent. Biomass collected both in t_0 and t_1 was oven-dried at 60°C for 48h and weighed. We calculated ABG productivity as the difference between t_0 and t_1 biomass. Therefore, only spring productivity was measured in 2021, which represents the largest proportion of annual productivity in this system, while a closer estimate of annual productivity was measured in 2022 and 2023.

Plant traits were collected in early May 2021, at the peak of the growing season. We considered a circular area of three meters in diameter around each exclusion cage and within this area, we identified the species present at 25 regularly spaced points using the line intercept method (Godínez-Alvarez et al., 2009), with three concentric circular transects going around the cage. After completing the species inventory for all the sampling plots in a paddock, we collected at least three to ten full individuals, in that same paddock, of each of the identified species, as proposed in the abundance-weighted trait sampling scheme (Carmona et al., 2015). Thus, at least ten individuals of the most abundant species were collected in the same paddock, five individuals for the medium abundant species and three individuals for the rare species. The leaf area (LA), specific leaf area (SLA), specific root length (SLR), leaf dry matter content (LDMC), root dry matter content (RDMC), leaf nitrogen content (LNC) and plant maximum height (see Fig. 2 for more detail) of all the collected individuals was measured following the standard protocols (Pérez-Harguindeguy et al., 2016). In total, more than 10000

206 individuals were measured across all the plots. This extensive sampling allowed us to account not only for interspecific
207 differences in trait values, but also for intraspecific variability in trait values across managements, farms and regions.
208 Intraspecific variability has proven to be very important for accurately defining the functional composition of plant
209 communities and their relationship to ecosystem processes (Siefert et al., 2015; Westerband et al., 2021). We also
210 conducted species inventories in the spring of 2022 and 2023, but on these later samplings we only collected and measured
211 the traits of individuals from the species not found in 2021. We used the trait values measured in 2021 for each species in
212 each paddock to impute trait values in 2022 and 2023. The proportion of legumes in each plot was quantified from the
213 species inventories as a measure of the number of plants with N fixation capacity in the communities.

214 Pasture chemical composition was assessed in the three years, after the species inventories and traits samplings. We
215 collected standing biomass by hand-clipping in three 25 x 25 cm quadrats randomly placed in the monitoring plot, outside
216 the exclusion cages. These samples were dried at 60°C for 48h, then grounded and analyzed with Dumas method in
217 DUMATHERM® N Pro analyzer (C. Gerhardt GmbH & Co. Germany) to obtain the nitrogen content of each sample.
218 Acid detergent lignin (ADL) and fiber (ADF) and neutral detergent fiber (NDF) content were also measured. NDF, ADF
219 and ADL were determined using a fiber analyzer (ANKOM A2000, ANKOM Technology, USA), following the official
220 procedures (Latimer, 2023).




Trait	Description	
Height 	Height to top photosynthetic tissue (cm). - Positively related to competitive ability for light and seed dispersal ^{1,2}	Traits associated with the plant size spectrum ³
Leaf area (LA) 	One-sided area of a fresh leaf (mm²) - Positively related to light interception . It also has implications for the water and energy balance of the leaf ³	
Specific leaf area (SLA) 	Leaf area/ Leaf dry mass (mm² mg⁻¹) - Positively related to plant relative growth rate and litter decomposability . Negatively related to stress tolerance ⁴	
Leaf dry matter content (LDMC) 	Leaf dry mass/ Leaf fresh mass (mg g⁻¹) - Negatively related to relative growth rate and litter decomposability . Positively related to stress tolerance ^{5,6}	Traits related with the Leaf/Root Economic Spectrum (conservative – acquisitive resources strategy gradient) ^{8,11}
Leaf nitrogen content (LNC) 	Nitrogen content in dry leaf mass (mg g⁻¹) - Positively related to photosynthetic rate , leaf respiration and litter decomposability ^{7,8}	
Root dry matter content (RDMC) 	Roots dry mass/ Roots fresh mass (m g⁻¹) - Correlated positively with root tissue density . Negatively correlated with root respiration and exudate production ^{9,10}	
Specific root length (SRL) 	Roots lenght / Roots dry mass (mm mg⁻¹) -Negatively correlated with root diameter and mycorrhizal association . Positively correlated with soil exploration and nutrient adquisition ^{11,12}	Traits related with the fungal collaboration gradient ¹¹

Figure 2. Graphical representation and description of plant functional traits measured. References embedded in the figure are provided below: ¹Gaudet and Keddy (1988), ²Thomson et al. (2011), ³Diaz et al. (2016), ⁴Poorter et al. (2009), ⁵Kazakou et al. (2009), ⁶Wilson et al. (1999), ⁷Freschet et al. (2012), ⁸Wright et al. (2004), ⁹Guyonnet et al. (2018), ¹⁰Roumet et al. (2016), ¹¹Bergmann et al. (2020), ¹²Kramer-Walter et al. (2016).

2.4. Soil sampling and soil characteristics measurement

In spring 2021, four soil cores were collected with a push sampler (5 cm diameter) at a depth of 8 cm around each exclusion cage, after removal of above-ground vegetation and litter from the sampled surfaces. The four soil cores were combined to obtain a composite sample from each plot, and an aliquot of 40 g of this sample were sieved (<2 mm), reserved and stored at 4°C for microbial community analysis in the following days. The remainder of the composite soil samples were air-dried and sieved (<2 mm). Coarse mineral material greater than 2 mm (hereinafter referred to as gravel) was weighed and used for bulk density correction (Eq.1).

Soil texture (sand, clay and silt content) was determined using a laser diffraction particle size analyzer (Mastersizer 2000, Malvern Instruments Ltd. UK) after dispersion with sodium hexametaphosphate. Soil pH was measured with a glass electrode (soil:water 1:2.5) pH meter (CRISON Basic20, Alella, Spain). Soils were extracted with 1M KCL and determined colorimetrically in a Bran+Luebbe Autoanalyzer 3 (Norderstedt, Germany), following the manufacturer's protocol, to measure nitrate (NO₃⁻) and ammonium (NH₄⁺) concentrations. Available P (Olsen P) was determined colorimetrically in an Agilent Cary 60 UV-Vis spectrophotometer (Agilent Technologies, USA), after extraction with 0.5 M NaHCO₃ at pH 8.5. Available K, Ca and Mg were determined by ICP-OES (Varian Mod. 720, Palo Alto, California, USA), after extraction with 1M ammonium acetate (pH 7).

242

2.5. Soil organic matter fractionation


Representative subsamples of each 2mm sieved air-dry composite soil sample were used to measure both bulk SOC and total nitrogen (TN) as well as their distribution across distinct physical fractions. First, 10 g aliquots of all samples were ground in a ball mill and the Bernard's calcimeter method (Sherrod et al., 2002) was used to test for the presence of inorganic C. Only four samples (0.02 % of the database) contained traces of carbonates, with CaCO₃ contents between 0.2 % and 0.8 %. These samples were excluded from subsequent analyses. For bulk soil analyses, aliquots of 10g of soil were ground in a ball mill and then analyzed in the DUMATHERM® N Pro analyzer (C. Gerhardt GmbH & Co. Germany) to determine %C (SOC) and %N (TN). For soil fractionation, we followed a combined size and density procedure as described in Leuthold et al. (2024). Briefly, an aliquot of 6 g of soil was mixed with sodium polytungstate (1.85 g cm⁻³) and shaken reciprocally for 18 h to disperse the aggregates. After dispersion samples were centrifuged for density fractionation and the light particulate organic matter "POM" (<1.85 g cm⁻³) was aspirated from the rest of the soil. We then thoroughly rinsed the residual heavy fraction and separated it by wet sieving into coarse heavy mineral-associated organic matter "chaOM" (>53 µm) and fine mineral-associated organic matter "MAOM" (<53 µm). All fractions were analyzed for %C and %N on an elemental analyzer as described above. Fractionation was accepted when mass recovery was within ±5%, and C recovery was within ±15%. For samples that did not satisfy one of these conditions, fractionation was repeated.

Since the chaOM and the MAOM shared similar C/N ratios, which were lower than POM C/N ratios (Fig. 3a), we merged these two mineral associated OM fractions and present them together as MAOM (Santos et al., 2024; Zhang et al., 2021). Thus, in this work, POM and MAOM represented the light (<1.85 g cm⁻³) and heavy (>1.85 g cm⁻³) fractions respectively. This contrasts with other studies based on size fractionation, in which light POM and chaOM are pooled together (Cotrufo et al., 2019b; Dai et al., 2025; Díaz-Martínez et al., 2024). However, our decision is supported by recent findings of Leuthold et al. (2024) who observed greater chemical similarity between chaOM and MAOM than between chaOM and POM. The pooling of the two mineral associated OM fractions is not expected to modify the results or interpretation of this work, as the C content of chaOM only accounts for an average of 8% of the total SOC, and its relative importance is similar across all managements (Fig. S1). Carbon data are presented as SOC, POC and MAOC stocks (Mg ha⁻¹), calculated following Poeplau et al. (2017):

$$\text{OC}_{\text{stock}} = \text{OC}_{\text{content}} \times \frac{\text{mass}_{\text{fine soil (<2mm)}}}{\text{Volume}_{\text{sample}}} \times \text{depth}_{\text{sample}} \quad (1)$$

where OC_{content} is the organic C content (as a proportion) of the soil fraction, mass_{fine soil(<2mm)} is the dry weight of the soil excluding gravel and large roots, and considering the dry mass of the aliquot reserved for microbial community analysis. Volume_{sample} and depth_{sample} are respectively the volume and depth of the soil sampled with the core.

273 Additionally, we calculated the proportion of MAOC in total SOC (MAOC/SOC) following Cotrufo et al. (2019).



274
275 **Figure 3.** Violin plot and boxplot (with median and quartiles) and mean values (expressed with labels and red dots) for the ratio
276 of carbon to nitrogen content (C/N ratio) in the soil organic matter fractions (a), and relation between carbon content in the
277 soil organic matter fractions (in g of C per kg of soil) and the total soil organic carbon (SOC) content (b). “chaOM” refers to
278 the coarse heavy mineral-associated organic matter, “IPOM” to the light particulate organic matter and “MAOM” to the fine
279 mineral-associated organic matter. In panel “b”, a linear regression line depicts the relationship between IPOM and total SOC,
280 while logistic curves illustrate the relationships of chaOM and MAOM with total SOC.

281

282 2.6. Soil microbial communities:

283 Soil microbial communities were characterized by phospholipid fatty acids analysis (PLFA; Willers et al., 2015). A two
284 g fine soil aliquot of lyophilized soil was used for lipid extraction with a one-phase chloroform–methanol-phosphate
285 buffer solvent. Phospholipids were separated from non-polar lipids and converted to fatty acid methyl esters (FAMES),
286 which were then separated by gas chromatography as described in details in Moreno et al. (2021). The sum of all
287 individual PLFAs was used as a proxy for microbial biomass (in nmol PLFAs g⁻¹ of soil). Further, we estimated microbial
288 biomass stocks (in mol ha⁻¹) by substituting the OC_{content} by the total PLFAs concentration in equation 1. Specific PLFAs
289 were used as biomarkers to estimate the relative abundance of broad taxonomic microbial groups, according to their
290 characteristic fatty acids: eukaryote, Gram- and Gram+ bacteria, saprophytic fungi and arbuscular mycorrhizal fungi
291 (Frostegård and Bååth, 1996). The ratios among Gram+ and Gram- bacteria (Gram+/Gram-) and fungi and bacteria
292 (Fungi/Bacteria) were also calculated to describe the composition of the microbial community.

293

294 2.7. Mineral-associated carbon capacity and saturation:

295 To evaluate the degree of MAOC saturation, we used the boundary line approach reported by Georgiou et al. (2022)
296 adjusted with global soil samples as a reference of the maximum observed mineralogical capacity (sensu Georgiou et al.,
297 2025) of our samples (Fig. 4). Therefore, the saturation deficit in each sample was calculated as one minus the ratio of
298 the current C content and the observed maximum C content according to the mineralogical capacity (% clay + silt).

299

300 2.6. Data analysis:

301 All analyses were carried out in R 4.3.3 (R Core Team, 2024). Species abundance data and species trait values were used
302 to compute, for each year and plot, the community weighted means (CWM) of all traits as well as the RaoQ multi-trait
303 functional diversity index (Ricotta and Moretti, 2011). In addition, we calculated the species richness as a measure of the
304 taxonomic diversity in each plot. These analyses were performed using the package “FD” (Laliberté and Legendre, 2010).
305 Regarding pasture chemical composition, the hemicellulose content was calculated as NDF minus ADF and the cellulose
306 content as ADF minus ADL, with ADL being considered a good proxy for the lignin content (Van Soest et al., 1991).

307 For the vegetation analysis, measurements from the three years were combined to obtain mean estimates of each
308 vegetation attribute. For each year, the ABG production and chemical attributes values, as well as the CWM of all the
309 functional traits, were centered and scaled between -1 and 1, so that values closer to -1 represent plots with traits values
310 lower than the mean, while values closer to 1 represent plots above the mean. The three years centered and scaled values
311 were then averaged to obtain estimates of each vegetation characteristic across the study period, as shown in this equation
312 2:

313
$$\text{Vegetation attribute}_{\text{mean}} = \frac{\sum_{\text{year } i=2021}^{2023} \text{Vegetation attribute (centered and scaled)}_{\text{year } i}}{3} \quad (2)$$

314 All mean values of vegetation characteristics obtained with equation 2 were used to build a Principal Component Analysis
315 (PCA). From this PCA, we extracted 4 main axes of variation of the vegetation characteristics with eigenvalues greater
316 than one (Fig .4).




Figure 4. Representation of the 4 main axes of variation in the principal component analysis (PCA) summarizing the vegetation characteristics variables. Panels a, b, c and d show the correlation between the different variables included in the PCA and the new axis. Panels e, f, g and h represent the plant communities characteristics at the extreme of each axis. Representative species of each axis are represented in Fig. S2.

Nutrients stocks in the soil samples were calculated by substituting the $OC_{content}$, in equation 1, for the concentration of each nutrient. PCA was used to summarize these nutrient stocks into a single composite index, as all nutrients were positively correlated. The first component of this PCA, which absorbed 46.9% of the variance, was extracted as a new variable, henceforth called “soil fertility” (Fig. S3). This soil fertility index increases as ammonium, nitrate, P, Ca, Mg, and K stocks in soil increase.

327 To analyze the direct and indirect effects of management, vegetation characteristics, microbial communities and
 328 environmental variables on POC, MAOC and SOC stocks and MAOC/SOC ratio we built a structural equation model
 329 (SEM) using the “piecewiseSEM” package (Lefcheck, 2016). These models assess the extent to which a defined structure
 330 of causal relationships fits the actual correlations between the data. We started from the assumption that management,
 331 climate and soil properties could affect SOC, POC and MAOC stocks and the MAOC/SOC directly or indirectly, by
 332 modifying microbial composition and vegetation characteristics. This provided us with a theoretical basis for the
 333 construction of the SEMs. The linear sub-models within these SEMs were fitted with the “lme4” package (Bates et al.,
 334 2015) using the farms as random factors to avoid spatial pseudo-replication. Model fitting was performed according to
 335 Zuur et al. (2009) selecting the model with the lowest corrected Akaike Information Criterion (AICc) value after adjusting
 336 for random and fixed factors. All predictor variables used in the model selection are summarised in Table 1. While gravel
 337 content is correlated to BD, its inclusion as predictor in the models helped to control for inter-site variability in gravel
 338 content. Interactions between management and climate or soil texture were also included during model selection. All
 339 model assumptions were checked and satisfactorily met. Predictor variables were scaled and centered prior to inclusion
 340 in the models.

341 To calculate the total effect of each explanatory factor (i.e. the sum of direct and indirect effects) on SOC, POC and
 342 MAOC stocks, and MAOC/SOC ratio in the SEMs, we used the “semEff” package (Murphy, 2022). This package
 343 generates estimates and confidence intervals of the total effects of the explanatory factors across multiple permutations
 344 of the data (9999 bootstraps in our case).

345 Given that average bulk density values partly differed among managements (Figure S4), and that this variation may affect
 346 carbon stocks, the same analysis procedure used for POC, MAOC and SOC stocks was used to analyze the change in
 347 POC, MAOC and SOC contents.

348 **Table 1. List of the variables and their units, which were included as fixed factors during the structural equation model**
 349 **(SEM) sub-models selection.**

Variables	Units
Management	Categorical (5 levels)
Sand content	%
Gravel content	%
Soil fertility	unitless
Mean annual precipitation (MAP)	mm
Mean annal temperature (MAT)	°C
Vegetation PCA- Axis 1 (N-C axis)	unitless
Vegetation PCA- Axis 2 (PES axis)	unitless
Vegetation PCA- Axis 3 (P-L axis)	unitless
Vegetation PCA- Axis 4 (Size axis)	unitless
Functional diversity (Rao multi-trait index)	unitless
Taxonomic diversity (species richness)	count
Microbial biomass (bulk)	nmol ha ⁻¹
Gram+/Gram-	ratio
Fungi/Bacteria	ratio

350

3. Results:

The mean values of POC, MAOC and SOC stocks were 9.8 ± 4.3 , 7.1 ± 3.1 and 16.5 ± 6.3 Mg C ha⁻¹ respectively. On average, MAOC represented 44% of the total SOC. The MAOC content of all the soils analyzed was well below the theoretical saturation level established by Georgiou et al. (2022) in relation to the clay + silt content (Fig. 5). Mean MAOC saturation deficit was 80%. Soil organic carbon fractions and stocks were affected directly and indirectly by mean plant traits, microbial communities, soil properties and management, and indirectly by climatic conditions (Fig. 6). Plant diversity indices and interactions between management and climate or soil texture were not retained in the model selection due to lack of significance.




Figure 5. Relation between the mineral-associated organic carbon (MAOC) and the percent Clay+Silt (CS) in the soil samples and the boundary line (in blue) with confidence intervals adjusted by Georgiou et al. (2022) for high-activity mineral soils, indicating the maximum observed mineralogical capacity for each CS content. The equation for the boundary line is provided in the panel. Points are colored based on their MAOC deficit (1- MAOC content / MAOC maximum observed capacity).

Total microbial biomass, estimated as total PLFA stocks, averaged 66.3 ± 25.9 mol ha⁻¹, with mean values for Fungi/Bacteria and Gram+/Gram ratios of 0.08 ± 0.03 and 1.1 ± 0.2 , respectively, although there were some differences between treatments (Fig. S5). Microbial biomass was positively correlated with POC, MAOC and SOC stocks (Fig. 6f; Fig. 7a, b & c). The Fungi/Bacteria ratio was negatively correlated with MAOC stocks (Fig. 6f; Fig. 7b) meaning that these stocks were higher in soils with a greater bacterial predominance. The Gram+/Gram- ratio had a significant direct negative effect on SOC, MAOC and PC stocks (Fig. 6f; Fig. 7a, b & c).

The first axis of the PCA of the vegetation characteristics, which reflects the N-C stoichiometry of the vegetation, had a negative direct effect on POC, MAOC and SOC stocks (Fig. 6h; Fig. 7a, b & c). The second axis, related to the plant economic spectrum (PES), had a positive direct effect on SOC stocks (Fig. 6h; Fig. 7c). In addition, the PES axis was positively correlated with the microbial biomass, thus having a positive indirect effect, and a significant total effect on POC, MAOC and SOC stocks (Fig. 6e; Fig. 7a, b & c). The third axis of the PCA (P-L axis), positively correlated with the lignin content and negatively related to vegetation productivity, was negatively correlated with the POC, MAOC and SOC stocks and the microbial biomass (Fig. 6e & h; Fig. 7a, b & c). The fourth axis, reporting the plant size, was positively correlated with the MAOC stocks, the MAOC/SOC ratio and the Fungi/Bacteria ratio (Fig. 6e & h; Fig. 7a & d). No index

379 of plant functional and taxonomic diversity was retained during model selection. On the other hand, soil fertility had a
 380 direct and indirect positive effect on POC, MAOC and SOC stocks (Fig. 6g). Soil fertility was negatively correlated with
 381 the Gram+/Gram- ratio, the P-L axis and the N-C axis, and positively correlated with the PES axis of the vegetation PCA
 382 (Fig. 6i & j).

383 Plot under continuous grazing had a mean POC, MAOC and SOC stocks of 9.1, 6.8 and 16.2 Mg C ha⁻¹ respectively, and
 384 a MAOC/SOC ratio of 0.44 (Fig. 8). Rotational grazing significantly increased soil fertility compared to continuous
 385 grazing (Fig. 6k). This led to a significant indirect effect of rotational grazing on POC, MAOC and SOC stocks (Fig. 7b
 386 & c). POC, MAOC and SOC stocks under rotational grazing had a mean value of 10.5, 7.8 and 18.5 Mg C/ha, respectively,
 387 which were 15% (in the case of POC and MAOC) and 14% higher (for SOC stocks) than mean values in continuous
 388 grazing (Fig. 8b & c). Recent legume sowing had a negative direct effect on POC, MAOC and SOC (Fig. 6b). However,
 389 Lr also increased significantly the plant size axis, the microbial biomass and the soil fertility and decreased the plant N-
 390 C axis compared to Ct (Fig. 6a, c & k). These changes resulted in a positive indirect effect of Lr over POC, MAOC and
 391 SOC stocks and a null total effect (Fig. 7a, b, c & d). Grazing abandonment increased the vegetation N-C axis and reduced
 392 the microbial biomass compared to Ct (Fig. 6a & c), resulting in a negative indirect effect on POC, MAOC and SOC
 393 stocks and a significant negative total effect on POC stocks (Fig. 7a, b & c). Thus, the mean POC stock on abandoned
 394 plots was 8.1 Mg C/ha, 11% lower than in continuous grazing plots (Fig. 8a).

395 Sand content in soil was negatively correlated with MAOC stocks and MAOC/SOC ratio (Fig. 6j). Sand content was also
 396 negatively correlated with soil fertility and the PES axis and positively correlated with the P-L and N-C axis (Fig. 6i),
 397 thus having a negative indirect effect on POC, MAOC and SOC stocks (Fig. 7a, b & c). As expected, due to its negative
 398 correlation with BD, gravel content was also negatively correlated with microbial biomass (Fig. 6j), leading to a
 399 significant negative indirect effect on POC, MAOC and SOC stocks (Fig. 7a, b & c). MAT was positively correlated with
 400 the N-C axis and negatively correlated with the soil fertility index (Fig. 6d & l). This implied a negative indirect effect of
 401 MAT over POC, MAOC and SOC stocks (Fig. 7a, b & c). On the other hand, MAP was negatively correlated with the N-
 402 C and the size axis (Fig. 6d). Thus, MAP had a positive indirect effect on POC, MAOC and SOC stocks (Fig. 7a, b, c &
 403 e).

404




Figure 6. Structural equation model representation. Factors included in the model are grouped by factor type (management, climate, vegetation traits, soil properties, microbial communities). Arrows between groups of factors indicate significant relationships between any of the factors included in both groups. The width of these arrows is proportional to the mean absolute size of the estimates between the factors in the groups. The plots embedded into these arrows show the standardized estimates of the significant relationships between the factors connected by the arrow. Negative standardized estimates are represented in red, and positive ones in blue. The size of the estimate circles represents the absolute value of the standardized estimate. Causal relationships between factors in the same group are represented by individual arrows.




Figure 7. Direct, indirect and total standardized effects of all studied variables included in the structural equation model (Figure6) over the (a) particulate organic carbon (POC), (b) mineral-associated organic carbon (MAOC), and (c) soil organic carbon (SOC) stocks and (d) the relative MAOC abundance (MAOC/SOC). Bars indicate direct (dark colors) and indirect (light colors) effects, and the black points-ranges indicate the total (i.e. direct + indirect) effect (with its 95% confidence interval). Stars over the total effect values indicate significant effects at a level of 0.05.




Figure 8. Violin plot and boxplot (with median and quartiles) for the (a) particulate organic carbon (POC), (b) mineral-associated organic carbon (MAOC), and (c) soil organic carbon (SOC) stocks and (d) the relative MAOC abundance (MAOC/SOC) in each management (Ct = Continuous grazing; Ab = grazing abandonment; Lr = Recent legume sowing; Lo = old legume sowing; Ro = Rotational grazing). The red point and the label indicate mean values predicted by the structure equation model (Fig.6). Lower case letters indicate significant differences ($p < 0.05$) between managements.

4. Discussion

Our results highlight the potential of management to control carbon storage in semi-arid grasslands. In this sense, rotational grazing arises as a promising tool for enhancing long-term carbon storage in the topsoil. We also identified several pathways by which management influences SOC, POC and MAOC stocks, showing the importance of changes in vegetation and microbial composition. Furthermore, management effects were consistent across the wide soil-climate gradient examined, enabling generalization of our results to a broad set of semi-arid grasslands in various environmental contexts. It should be noted that the predicted capacity of POC, MAOC and SOC stocks models was high (62%, 69% 73% of the variance explained by the fixed factors, respectively), indicating the robustness of the results presented and their importance for expanding the understanding of soil carbon dynamics in grasslands. Our results are limited to the upper topsoil (the first 8 cm), and it would be necessary to analyze deeper layers to fully understand the processes of SOC formation and stabilization and the effects of management changes. However this top layer is the most important for mediterranean grassland functioning as it contains the majority of roots and therefore most of the microbial, nutrients and water dynamics (Acosta-Gallo et al., 2011; Moreno et al., 2005). Furthermore, changes in management primarily affect the SOC of the topsoil layer, especially in the short term (Ward et al., 2016).

4.1. Soil organic carbon fractions, stocks, and saturation levels

SOC stocks and contents (Fig.8 & S4) in the paddocks studied ($16.5 \text{ Mg C ha}^{-1}$ and 19.5 g C kg^{-1} on average, respectively) were consistent with values found in the first 7-10 cm of soil in other grasslands under similar climatic conditions (Díaz-

446 Martínez et al., 2024; Oggioni et al., 2020; Parras-Alcántara et al., 2014). On the other hand, the MAOC/SOC ratio of our
447 soils (0.44 on average) was lower than the values generally reported in both global and semi-arid grasslands (Hansen et
448 al., 2024; Rocci et al., 2022; Shi et al., 2024), which are around 0.6-0.7. The shallow sampling depth in this work (8 cm)
449 may influence the observed MAOC/SOC ratios, as POM decrease more sharply than MAOM with soil depth (Galluzzi et
450 al., 2025; Sanderman et al., 2021; Tang et al., 2024). For instance, Plaza et al. (2022) reported MAOC/SOC ratios between
451 0.3 and 0.4 in the first 10 cm of soil, while Cappai et al. (2017) found MAOC/SOC ratios around 0.7 in the first 20 cm,
452 both in Mediterranean grasslands.

453 Carbon concentrations in the fine soil fraction (clay + silt) were well below the saturation point observed in previous
454 studies (Cotrufo et al., 2019a; Georgiou et al., 2022). However, we observed a certain limit to MAOC accumulation in
455 our system, as its content remained stable above SOC contents of 30 g/kg⁻¹ and stayed below 20 g/kg⁻¹ even when SOC
456 reached values above 60 g/kg⁻¹, following a saturation curve (Fig. 3b). These results support the observations that MAOC
457 accrual is more limited by the amount of C inputs rather than the mineralogical capacity of the soil (Poeplau et al., 2024).
458 In this sense, the maximum C in the clay+silt fraction observed in this work (around 19 g C/kg clay+silt) would represent
459 the maximum capacity of the ecosystem to stabilise C at current levels of productivity. Therefore, MAOC saturation
460 should be a minor concern when planning management strategies to improve C storage in Mediterranean grasslands.

461

462 4.2. Soil microbial communities regulate SOC storage

463 Total microbial biomass had a similar and substantial positive effect on POC, MAOC and total SOC stocks, such that the
464 MAOC/SOC ratio was unaffected. Typically, microbial biomass C only represents about 2% of SOC (Xu et al., 2013; Yao
465 et al., 2000), but it is closely correlated with the accumulation of microbial necromass C (Hou et al., 2024). The latter can
466 account for more than 50% of SOC (Liang et al., 2019), with a similar contribution to the POC and MAOC fractions
467 (Zhang et al., 2024). The positive effect of microbial biomass on both POC and MAOC, as well as the relatively low C:N
468 ratio of these fractions found in this study (Fig. 3a) compared to other works (Chang et al., 2024; Yu et al., 2022), indicate
469 a prominent role of microbial transformation of plant inputs in SOC formation in semi-arid grasslands, which would also
470 explain the relatively low SOC stocks of these ecosystems.

471 Fungi/bacteria ratio was negatively correlated with MAOC fraction, as fungi residues tend to contribute more to POC
472 (Griepentrog et al., 2014; Lavalley et al., 2020; Tang et al., 2023), but no significant effect of fungi/bacteria ratio over
473 POC stocks or MAOC/SOC ratio was observed. Gram⁺/Gram⁻ ratio was negatively correlated with POC, MAOC and
474 SOC stock, as observed in previous studies (Khatri-Chhetri et al., 2024). Gram⁻ bacteria are more dependent on plant C
475 inputs, whereas Gram⁺ bacteria tend to use more of the organic C already present in the soil (Fanin et al., 2019; Kramer
476 and Gleixner, 2008; Waldrop and Firestone, 2004). Thus, the proliferation of Gram⁺ bacteria may promote the
477 decomposition of pre-existing SOM (Klumpp et al., 2009).

478

479 4.3. Plant-soil interactions and their effect on soil carbon stocks

480 The analysis of vegetation functional traits and chemical composition revealed four main axes of variation. Two of these
481 axes (axes 1 and 4) correspond to the spectrum of plant form and function, found in several global vegetation analyses
482 (Díaz et al., 2016; Weigelt et al., 2021). This two-dimensional spectrum is defined by the leaf and root economic gradient
483 (Kramer-Walter et al., 2016; Wright et al., 2004), that moves from slow-growing and resource-conserving to fast-growing
484 and resource-acquisitive species (Wright et al., 2004), and the size gradient, which increases with increasing plant height

485 and leaf area and reflects the competitive ability of plants for the light (Díaz et al., 2016). The other two axes (axes 2 and
486 3) are more related to the aboveground chemical composition of the vegetation. Axis 1 reflects the N-C stoichiometry of
487 plant tissues, driven mainly by the ABG C/N ratio, both of which are obviously negatively correlated with certain traits
488 such as LNC and the proportion of legumes. The N-fixing capacity of legumes has a fertilising effect on companion
489 plants, increasing its tissues nitrogen content (Pino et al., 2016). In addition, legumes themselves tend to have a high
490 concentration of nitrogen in leaves and stems (Carranca et al., 2015). Axis 3 is positively related to the lignin content of
491 the vegetation. This independence between lignin and nitrogen content in plant litter has already been observed (Cornwell
492 et al., 2008). ABG productivity of vegetation was correlated with most of these axes, especially with axis 3, being higher
493 in more acquisitive, less lignified and bigger vegetation, as found in other works (Laliberté and Tylianakis, 2012; Zhang
494 et al., 2017).

495 POC, MAOC and SOC stocks were higher in communities dominated by resource-acquisitive, highly productive plants
496 with low lignin content and low C/N ratio. These communities are expected to provide a higher level of plant inputs,
497 thereby increasing the incorporation of organic matter into the soil (King et al., 2023; Mortensen et al., 2025; Zhou et al.,
498 2024). Increased inputs may come from higher plant litter production but also from increased root exudates, which tend
499 to be higher in acquisitive plants (Guyonnet et al., 2018). Part of the effects of plant traits in SOC was mediated by
500 increases in microbial biomass, which would also benefit from higher levels of plant inputs and increased root exudation
501 (Eisenhauer et al., 2017). In addition, inputs with low lignin and C/N ratios are degraded more efficiently, reducing C
502 losses and promoting long-term SOC storage (Cotrufo and Lavelle, 2022; Ridgeway et al., 2022). However, based on
503 previous work, we would have expected a direct contribution of plant structural input to POC (Cotrufo et al., 2022), and
504 thus a higher proportion of POC in communities that produce more recalcitrant litter (Cheng et al., 2023; Haddix et al.,
505 2016; Mortensen et al., 2025). This was not the case in this work, where the MAOC/SOC ratio remained unaffected by
506 the chemical composition of the ABG vegetation tissues. This unexpected result could be explained by the fact that in
507 these systems, POC also appears to be the product of microbial transformation of plant inputs, as suggested by its positive
508 relationship with microbial biomass and its relatively low C/N ratio. In addition, photodegradation, an important driver
509 of litter degradation in semi-arid ecosystems (Austin and Vivanco, 2006), can promote litter lignin biodegradability and
510 the production of litter soluble compounds that are readily accessible to soil microbes (Wang et al., 2015), thus reducing
511 the influence of vegetation chemical properties on POC and MAOC formation. Grazing also decouples the quality of
512 plant tissues and the final quality of inputs to the soil, as livestock excreta have a lower C/N ratio than the plant material
513 consumed (Soussana and Lemaire, 2014). We also observed that taller and larger plants (high values on the size axis)
514 were associated with higher MAOC stocks and MAOC/SOC ratios, although the mechanism driving this correlation was
515 unclear. Generally, plant height is positively correlated with shoot:root ratio (Li et al., 2008), and several studies have
516 found a higher contribution of shoots, rather than roots, to the MAOC fraction due to the higher recalcitrance of root
517 tissues (Huang et al., 2021; Lavelle et al., 2018; Ridgeway et al., 2022). However, rhizodeposition, which is closely
518 linked to root biomass, has been shown to promote MAOC formation over POC (Berenstecher et al., 2023; Villarino et
519 al., 2021). On the other hand, a greater accumulation of standing litter, rather than surface litter, might be expected in
520 communities with bigger plants, and some studies in semi-arid grasslands have observed higher rates of microbial
521 degradation and release of soluble compounds (thus contributing more to MAOC) in standing litter, compared to surface
522 litter, due to greater retention of night-time moisture in the former (Gliksman et al., 2018; Wang et al., 2017). In any case,
523 our results and their interpretation are limited by the correlation in our vegetation data between input quantity (ABG
524 productivity) and input quality, which could mask the effect of the latter on SOC formation.

525 No significant effects of taxonomic or functional diversity on POC, MAOC or SOC stocks were found, which could be
526 explained by the negative correlation of these biodiversity indicators with the ABG productivity and soil fertility in our

plots (Fig. S7). This negative correlation has been observed in other natural grassland research and may be due to the dominance of some highly productive species in nutrient-rich environments (Feßel et al., 2016; Helsen et al., 2014; Luo et al., 2019; Rolo et al., 2016). In this sense, our results would be consistent with previous research that has found a greater importance of functional identity (i.e. mean traits values) relative to functional diversity in predicting ecosystem processes (Mokany et al., 2008; Zhang et al., 2017).

Soil fertility plays a central role in our models, increasing POC, MAOC and SOC stocks. Previous studies in grasslands have already observed a positive correlation between POC and MAOC and soil nutrients content (Wang et al., 2025). Global N addition experiments have also showed to increase topsoil C storage in grasslands (Hu et al., 2024; Liu et al., 2023). In this work, we observe a direct effect of soil fertility on C storage, but we also identify some indirect effects mediated by changes in vegetation and microbial communities. More resource-acquisitive and productive plant communities, with lignin content and lower C/N ratio, were found at higher soil fertility. Resource-acquisitive plants are expected to be more competitive in nutrient-rich environments (Daou et al., 2021; Ordoñez et al., 2009). The absence of nutrient limitation also promotes plant productivity (Fay et al., 2015). In addition, soil fertility was negatively correlated with the Gram+/Gram- ratio. Gram- bacteria have a more copiotroph strategy than Gram+ bacteria and therefore benefit more from nutrient-rich environments (De Deyn et al., 2010; Luo et al., 2020; Zhong et al., 2010). The direct positive effect of soil fertility on C storage in soils may be explained by changes in other factors that we did not measure. For instance, nutrient addition have been shown to promote below-ground productivity in mediterranean woody pastures (Nair et al., 2019) and in global grasslands (Liu et al., 2023; Lu et al., 2011). Moreover, nutrient availability in soils tends to increase CUE (Poeplau et al., 2019; Spohn et al., 2016) and to alleviate the need for microbial N-mining, reducing C losses and old SOM decomposition (Blagodatskaya and Kuzyakov, 2008).

4.5. Management effects

Rotational grazing increased both POC, MAOC and SOC stocks by 15%, 15% and 14%, respectively, compared to conventional continuous grazing. Previous studies found the same positive effect of rotational grazing on MAOC (Khatri-Chhetri et al., 2024; Mosier et al., 2021; Stanley et al., 2025) and total SOC content (Byrnes et al., 2018; Conant et al., 2017; Liu et al., 2024; Phukubye et al., 2022; Teague et al., 2011). According to our results, this effect is mainly explained by the higher soil fertility observed in rotational grazing plots, cascading in the above explained effects on plant traits and microbial transformation leading to higher POC, MAOC and SOC stocks. Soil fertility may be higher under rotational grazing than under continuous grazing because fecal and urine excretions tend to be less spatially clustered in rotationally grazed paddocks, resulting in more homogeneous fertilization of the entire paddock (Augustine et al., 2023; Dubeux Jr. et al., 2014). In contrast, in continuous grazing, excreta tend to accumulate in areas of highest animal use, such as near feeders or water points (Tate et al., 2003), which are areas that have been avoided in our sampling. In turn, the homogeneous grazing and fertilization maintains a greater amount of ground covered by vegetation (Stanley et al., 2025; Teague et al., 2004), which limits topsoil losses through erosion (Sanjari et al., 2009).

Surprisingly legume sowings had a negative direct effect on soil C stocks, but a positive indirect effect, resulting in a non-significant total effect. The negative impact was more evident in recent sowings, especially on the MAOC fraction, and could be attributed to the impact of the pre-sowing tillage. Previous works have observed a reduction in SOC stocks after tilling in mediterranean grasslands (Parras-Alcántara et al., 2015; Uribe et al., 2015). This would explain why the direct effect of legume sowing in C storage is more negative in recent compared to old-sowed paddocks. The effect of tillage on SOC stocks may arise from changes in soil bulk density rather than C content, resulting in misleading conclusions (Du et al., 2017; Rovira et al., 2022). However, in this work the analysis of POC, MAOC and SOC contents (in g/kg bulk soil;

Fig. S6 & S8) shows similar effects of legume sowing as those observed for C stocks (Fig. 7a, b & c). In the recently sown plots, the negative impact in C stocks was countered by an increase in soil fertility, plant productivity and plant nitrogen content, possibly due to increased legume cover (Gómez-Rey et al., 2012; Gou et al., 2023; Hernández-Esteban et al., 2019). In addition, recent legume sowing also implied an increase in microbial biomass and a decrease in the Gram+/Gram- ratio, as observed in previous studies (Moreno et al., 2021).

Grazing abandonment led to a 12% reduction in the POC stocks, compared to continuous grazing. This result contradicts the observations of previous studies in grasslands, which reported positive effects of grazing exclusion on SOC storage, both globally (Eze et al., 2018) and particularly in semi-arid climates (Cheng et al., 2016; Gao et al., 2025; Yu et al., 2021). However McSherry and Ritchie (2013) meta-analysis found that light grazing could promote SOC stocks in C3 dominated grasslands (such as our study grassland). Other global and regional studies also observed null or even positive effects of light grazing, compared to grazing exclusion, in SOC stocks in grasslands (Liu et al., 2024; Zhou et al., 2017). Our results would be in line with previous studies in mediterranean woody pastures that found a decrease in SOC storage with grazing abandonment (Oggioni et al., 2020; Peco et al., 2017). The fact that grazing abandonment particularly affected the POC fraction is surprising, since in abandoned paddocks there was a greater accumulation of plant structural biomass, that should promote POC accrual (Cotrufo et al., 2022). According to our results, the negative effect of grazing exclusion on POC stocks was mainly mediated by a reduction in microbial biomass, and a proliferation of plants with higher C/N ratios. Once again pointing to the microbially transformed nature of POC in these soils. Grazing has been proven to increase microbial activity and biomass (Bardgett et al., 2001; Zhou et al., 2017), in part by increasing root exudation (Hamilton et al., 2008; Wilson et al., 2018). Previous studies have also observed an increase in vegetation C/N ratio with cessation of grazing (He et al., 2020; Wang et al., 2016), which could be explained by the accumulation of older plant tissues and senescent standing biomass, with higher C/N than green biomass and new leaves (Sanaullah et al., 2010). Importantly, we observed no interactions between management effects and climate or soil texture variables, showing that the observed management effects were consistent across the entire environmental gradient. Although such interactions have been observed for wider climatic gradients (Byrnes et al., 2018; Phukubye et al., 2022), our results show that management practices such as rotational grazing can have net positive impacts on SOC storage in grasslands with varying conditions within semi-arid regions.

594

595 Climate effects and implications for global warming

Even within the small climatic gradient covered in this study, we found lower stocks of POC, MAOC and SOC in the warmer areas, as observed in previous research (Díaz-Martínez et al., 2024; Georgiou et al., 2024; Hansen et al., 2024; Shi et al., 2024). According to our results, these negative effects were entirely mediated by a decrease in soil fertility and vegetation input quality (increased N-C axis) with increasing temperature. Increased aridity has been shown to be associated with the alteration of several ecosystem processes that regulate nutrient cycling and availability (Berdugo et al., 2020; Moreno-Jiménez et al., 2019). Increased fiber for tissue protection, typical from plants growing at higher temperatures, may explain the increase in vegetation C/N ratios (Arroyo et al., 2024). On the other hand, annual precipitation exerted a small positive effect on POC and SOC stocks, mediated by the reduction of C/N ratios in vegetation in wetter farms. Based on future projections of increasing temperatures and decreasing annual precipitation in our study area (IPCC, 2023b) we could expect significant reductions in SOC stocks in Mediterranean grasslands in the next decades, affecting both the POC and MAOC fractions.

607

608 **5. Conclusions:**

609 Implementing rotational grazing can improve both POC and MAOC stocks in Mediterranean grasslands. On the other
610 hand, the abandonment of grazing, far from functioning as a tool for ecological restoration, can lead to a loss of soil
611 carbon storage capacity in these ecosystems.

612 Changes in microbial communities and vegetation attributes were the main drivers of changes in SOC stocks and fractions.
613 In this sense, this work has proven the usefulness of plant functional traits as tools for the study of plant-soil interactions
614 and SOC formation dynamics.

615 We can expect a loss of soil carbon in the studied grasslands over the next few decades due to grazing abandonment and
616 climate warming, so selecting management approaches that mitigate or counteract these losses is vital to maintaining the
617 fertility, productivity and functioning of semi-arid grasslands.

618

619 **Data and code availability**

620 The data and code that support the findings of this study are available from the corresponding author upon reasonable
621 request.

622

623 **CRedit authorship contribution statement**

624 **A. Carrascosa:** Data curation, Formal analysis, Investigation, Software, Visualization, Writing – original draft. **G.**
625 **Moreno:** Conceptualization, Funding acquisition, Project administration, Supervision, Writing – review and editing. **M.**
626 **F. Cotrufo:** Conceptualization, Methodology, Supervision, Writing – review and editing. **C. Frade:** Data curation,
627 Methodology, Investigation, Resources, Writing – review and editing. **S. Rodrigo:** Data curation, Methodology,
628 Investigation, Resources, Writing – review and editing. **V. Rolo:** Conceptualization, Data curation, Formal analysis,
629 Funding acquisition, Investigation, Project administration, Supervision, Writing – review and editing.

630

631 **Competing interests**

632 The authors declare that they have no conflict of interest.

633

634 **Financial support**

635 This work was part of the projects “PID2019-108313RB-C33”, “PID2019-108313RB-C31” and “PCIN2021-122100-2”
636 funded by MICIU/AEI/10.13039/501100011033/ and “European Union NextGenerationEU/PRTR; the project “CLU-
637 2019-05 – IRNASA/CSIC Unit of Excellence” funded by the “Junta de Castilla y León” and co-financed by the European
638 Union (ERDF “Europe drives our growth”); and European Union’s Horizon Europe research and innovation program.
639 Grant agreement: 101059794. Alejandro Carrascosa was supported by the Spanish Ministry of Science and Innovation
640 through an FPU grant (FPU21/02668). Cristina Frade was supported by a JCyL fellowship (E-37-2023-0024090) co-
641 financed by the European Social Fund Plus (FSE+).

642

643 **References:**

- 644 Acosta-Gallo, B., Casado, M. A., Montalvo, J., and Pineda, F. D.: Allometric patterns of below-ground biomass in
645 Mediterranean grasslands, *Plant Biosyst.* - *Int. J. Deal. Asp. Plant Biol.*, 145, 584–595,
646 <https://doi.org/10.1080/11263504.2011.578836>, 2011.
- 647 Angst, G., Mueller, K. E., Kögel-Knabner, I., Freeman, K. H., and Mueller, C. W.: Aggregation controls the stability of
648 lignin and lipids in clay-sized particulate and mineral associated organic matter, *Biogeochemistry*, 132, 307–324,
649 <https://doi.org/10.1007/s10533-017-0304-2>, 2017.
- 650 Angst, G., Mueller, K. E., Nierop, K. G. J., and Simpson, M. J.: Plant- or microbial-derived? A review on the molecular
651 composition of stabilized soil organic matter, *Soil Biol. Biochem.*, 156, 108189,
652 <https://doi.org/10.1016/j.soilbio.2021.108189>, 2021.
- 653 Arroyo, A. I., Pueyo, Y., Barrantes, O., and Alados, C. L.: Interplay between Livestock Grazing and Aridity on the
654 Ecological and Nutritional Value of Forage in Semi-arid Mediterranean Rangelands (NE Spain), *Environ.*
655 *Manage.*, <https://doi.org/10.1007/s00267-024-01939-9>, 2024.
- 656 Augustine, D. J., Kearney, S. P., Raynor, E. J., Porensky, L. M., and Derner, J. D.: Adaptive, multi-paddock, rotational
657 grazing management alters foraging behavior and spatial grazing distribution of free-ranging cattle, *Agric.*
658 *Ecosyst. Environ.*, 352, 108521, <https://doi.org/10.1016/j.agee.2023.108521>, 2023.
- 659 Austin, A. T. and Vivanco, L.: Plant litter decomposition in a semi-arid ecosystem controlled by photodegradation, *Nature*,
660 442, 555–558, <https://doi.org/10.1038/nature05038>, 2006.
- 661 Bai, X., Zhai, G., Wang, B., An, S., Liu, J., Xue, Z., and Dippold, M. A.: Litter quality controls the contribution of
662 microbial carbon to main microbial groups and soil organic carbon during its decomposition, *Biol. Fertil. Soils*,
663 60, 167–181, <https://doi.org/10.1007/s00374-023-01792-8>, 2024.
- 664 Bai, Y. and Cotrufo, M. F.: Grassland soil carbon sequestration: Current understanding, challenges, and solutions, *Science*,
665 377, 603–608, <https://doi.org/10.1126/science.abo2380>, 2022.
- 666 Baldock, J. A. and Skjemstad, J. O.: Role of the soil matrix and minerals in protecting natural organic materials against
667 biological attack, *Org. Geochem.*, 31, 697–710, [https://doi.org/10.1016/S0146-6380\(00\)00049-8](https://doi.org/10.1016/S0146-6380(00)00049-8), 2000.
- 668 Bardgett, R. D., Jones, A. C., Jones, D. L., Kemmitt, S. J., Cook, R., and Hobbs, P. J.: Soil microbial community patterns
669 related to the history and intensity of grazing in sub-montane ecosystems, *Soil Biol. Biochem.*, 33, 1653–1664,
670 [https://doi.org/10.1016/S0038-0717\(01\)00086-4](https://doi.org/10.1016/S0038-0717(01)00086-4), 2001.
- 671 Bartholomew, P. W. and Williams, R. D.: Overseeding Unimproved Warm-Season Pasture with Cool- and Warm-Season
672 Legumes to Enhance Forage Productivity, *J. Sustain. Agric.*, 34, 125–140,
673 <https://doi.org/10.1080/10440040903482407>, 2010.
- 674 Bates, D., Mächler, M., Bolker, B., and Walker, S.: Fitting Linear Mixed-Effects Models Using lme4, *J. Stat. Softw.*, 67,
675 1–48, <https://doi.org/10.18637/jss.v067.i01>, 2015.
- 676 Benbi, D. K., Boparai, A. K., and Brar, K.: Decomposition of particulate organic matter is more sensitive to temperature
677 than the mineral associated organic matter, *Soil Biol. Biochem.*, 70, 183–192,
678 <https://doi.org/10.1016/j.soilbio.2013.12.032>, 2014.
- 679 Berdugo, M., Delgado-Baquerizo, M., Soliveres, S., Hernández-Clemente, R., Zhao, Y., Gaitán, J. J., Gross, N., Saiz, H.,
680 Maire, V., Lehmann, A., Rillig, M. C., Solé, R. V., and Maestre, F. T.: Global ecosystem thresholds driven by
681 aridity, *Science*, 367, 787–790, <https://doi.org/10.1126/science.aay5958>, 2020.
- 682 Berenstecher, P., Conti, G., Faigón, A., and Piñeiro, G.: Tracing service crops’ net carbon and nitrogen rhizodeposition
683 into soil organic matter fractions using dual isotopic brush-labeling, *Soil Biol. Biochem.*, 184, 109096,
684 <https://doi.org/10.1016/j.soilbio.2023.109096>, 2023.
- 685 Bergmann, J., Weigelt, A., van der Plas, F., Laughlin, D. C., Kuyper, T. W., Guerrero-Ramirez, N., Valverde-Barrantes,
686 O. J., Bruehlheide, H., Freschet, G. T., Iversen, C. M., Kattge, J., McCormack, M. L., Meier, I. C., Rillig, M. C.,
687 Roumet, C., Semchenko, M., Sweeney, C. J., van Ruijven, J., York, L. M., and Mommer, L.: The fungal
688 collaboration gradient dominates the root economics space in plants, *Sci. Adv.*, 6, eaba3756,
689 <https://doi.org/10.1126/sciadv.aba3756>, 2020.

- Blagodatskaya, E. and Kuzyakov, Y.: Mechanisms of real and apparent priming effects and their dependence on soil microbial biomass and community structure: critical review, *Biol. Fertil. Soils*, 45, 115–131, <https://doi.org/10.1007/s00374-008-0334-y>, 2008.
- Briske, D. D., Derner, J. D., Brown, J. R., Fuhlendorf, S. D., Teague, W. R., Havstad, K. M., Gillen, R. L., Ash, A. J., and Willms, W. D.: Rotational Grazing on Rangelands: Reconciliation of Perception and Experimental Evidence, *Rangel. Ecol. Manag.*, 61, 3–17, <https://doi.org/10.2111/06-159R.1>, 2008.
- Byrnes, R. C., Eastburn, D. J., Tate, K. W., and Roche, L. M.: A Global Meta-Analysis of Grazing Impacts on Soil Health Indicators, *J. Environ. Qual.*, 47, 758–765, <https://doi.org/10.2134/jeq2017.08.0313>, 2018.
- Cappai, C., Kemanian, A. R., Lagomarsino, A., Roggero, P. P., Lai, R., Agnelli, A. E., and Seddaiu, G.: Small-scale spatial variation of soil organic matter pools generated by cork oak trees in Mediterranean agro-silvo-pastoral systems, *Geoderma*, 304, 59–67, <https://doi.org/10.1016/j.geoderma.2016.07.021>, 2017.
- Carmona, C. P., Rota, C., Azcárate, F. M., and Peco, B.: More for less: sampling strategies of plant functional traits across local environmental gradients, *Funct. Ecol.*, 29, 579–588, <https://doi.org/10.1111/1365-2435.12366>, 2015.
- Carranca, C., Castro, I. V., Figueiredo, N., Redondo, R., Rodrigues, A. R. F., Saraiva, I., Maricato, R., and Madeira, M. A. V.: Influence of tree canopy on N₂ fixation by pasture legumes and soil rhizobial abundance in Mediterranean oak woodlands, *Sci. Total Environ.*, 506–507, 86–94, <https://doi.org/10.1016/j.scitotenv.2014.10.111>, 2015.
- Carranca, C., Pedra, F., and Madeira, M.: Enhancing Carbon Sequestration in Mediterranean Agroforestry Systems: A Review, *Agriculture*, 12, 1598, <https://doi.org/10.3390/agriculture12101598>, 2022.
- Carrascosa, A., Moreno, G., Rodrigo, S., and Rolo, V.: Unravelling the contribution of soil, climate and management to the productivity of ecologically intensified Mediterranean wood pastures, *Sci. Total Environ.*, 957, 177575, <https://doi.org/10.1016/j.scitotenv.2024.177575>, 2024.
- Chang, Y., Sokol, N. W., van Groenigen, K. J., Bradford, M. A., Ji, D., Crowther, T. W., Liang, C., Luo, Y., Kuzyakov, Y., Wang, J., and Ding, F.: A stoichiometric approach to estimate sources of mineral-associated soil organic matter, *Glob. Change Biol.*, 30, e17092, <https://doi.org/10.1111/gcb.17092>, 2024.
- Cheng, J., Jing, G., Wei, L., and Jing, Z.: Long-term grazing exclusion effects on vegetation characteristics, soil properties and bacterial communities in the semi-arid grasslands of China, *Ecol. Eng.*, 97, 170–178, <https://doi.org/10.1016/j.ecoleng.2016.09.003>, 2016.
- Cheng, X., Xing, W., and Liu, J.: Litter chemical traits, microbial and soil stoichiometry regulate organic carbon accrual of particulate and mineral-associated organic matter, *Biol. Fertil. Soils*, 59, 777–790, <https://doi.org/10.1007/s00374-023-01746-0>, 2023.
- Conant, R. T.: Grassland Soil Organic Carbon Stocks: Status, Opportunities, Vulnerability, in: *Recarbonization of the Biosphere: Ecosystems and the Global Carbon Cycle*, edited by: Lal, R., Lorenz, K., Hüttl, R. F., Schneider, B. U., and von Braun, J., Springer Netherlands, Dordrecht, 275–302, https://doi.org/10.1007/978-94-007-4159-1_13, 2012.
- Conant, R. T., Cerri, C. E. P., Osborne, B. B., and Paustian, K.: Grassland management impacts on soil carbon stocks: a new synthesis, *Ecol. Appl.*, 27, 662–668, <https://doi.org/10.1002/eap.1473>, 2017.
- Cornwell, W. K., Cornelissen, J. H. C., Amatangelo, K., Dorrepaal, E., Eviner, V. T., Godoy, O., Hobbie, S. E., Hoorens, B., Kurokawa, H., Pérez-Harguindeguy, N., Quested, H. M., Santiago, L. S., Wardle, D. A., Wright, I. J., Aerts, R., Allison, S. D., Van Bodegom, P., Brovkin, V., Chatain, A., Callaghan, T. V., Díaz, S., Garnier, E., Gurvich, D. E., Kazakou, E., Klein, J. A., Read, J., Reich, P. B., Soudzilovskaia, N. A., Vaieretti, M. V., and Westoby, M.: Plant species traits are the predominant control on litter decomposition rates within biomes worldwide, *Ecol. Lett.*, 11, 1065–1071, <https://doi.org/10.1111/j.1461-0248.2008.01219.x>, 2008.
- Cotrufo, M. F. and Lavelle, J. M.: Chapter One - Soil organic matter formation, persistence, and functioning: A synthesis of current understanding to inform its conservation and regeneration, in: *Advances in Agronomy*, vol. 172, edited by: Sparks, D. L., Academic Press, 1–66, <https://doi.org/10.1016/bs.agron.2021.11.002>, 2022.
- Cotrufo, M. F., Wallenstein, M. D., Boot, C. M., Denef, K., and Paul, E.: The Microbial Efficiency-Matrix Stabilization (MEMS) framework integrates plant litter decomposition with soil organic matter stabilization: do labile plant inputs form stable soil organic matter?, *Glob. Change Biol.*, 19, 988–995, <https://doi.org/10.1111/gcb.12113>, 2013.

- 739 Cotrufo, M. F., Ranalli, M. G., Haddix, M. L., Six, J., and Lugato, E.: Soil carbon storage informed by particulate and
740 mineral-associated organic matter, *Nat. Geosci.*, 12, 989–994, <https://doi.org/10.1038/s41561-019-0484-6>,
741 2019a.
- 742 Cotrufo, M. F., Ranalli, M. G., Haddix, M. L., Six, J., and Lugato, E.: Soil carbon storage informed by particulate and
743 mineral-associated organic matter, *Nat. Geosci.*, 12, 989–994, <https://doi.org/10.1038/s41561-019-0484-6>,
744 2019b.
- 745 Cotrufo, M. F., Haddix, M. L., Kroeger, M. E., and Stewart, C. E.: The role of plant input physical-chemical properties,
746 and microbial and soil chemical diversity on the formation of particulate and mineral-associated organic matter,
747 *Soil Biol. Biochem.*, 168, 108648, <https://doi.org/10.1016/j.soilbio.2022.108648>, 2022.
- 748 Crowther, T. W., Todd-Brown, K. E. O., Rowe, C. W., Wieder, W. R., Carey, J. C., Machmuller, M. B., Snoek, B. L., Fang,
749 S., Zhou, G., Allison, S. D., Blair, J. M., Bridgham, S. D., Burton, A. J., Carrillo, Y., Reich, P. B., Clark, J. S.,
750 Classen, A. T., Dijkstra, F. A., Elberling, B., Emmett, B. A., Estiarte, M., Frey, S. D., Guo, J., Harte, J., Jiang, L.,
751 Johnson, B. R., Kröel-Dulay, G., Larsen, K. S., Laudon, H., Lavallee, J. M., Luo, Y., Lupascu, M., Ma, L. N.,
752 Marhan, S., Michelsen, A., Mohan, J., Niu, S., Pendall, E., Peñuelas, J., Pfeifer-Meister, L., Poll, C., Reinsch, S.,
753 Reynolds, L. L., Schmidt, I. K., Sistla, S., Sokol, N. W., Templer, P. H., Treseder, K. K., Welker, J. M., and
754 Bradford, M. A.: Quantifying global soil carbon losses in response to warming, *Nature*, 540, 104–108,
755 <https://doi.org/10.1038/nature20150>, 2016.
- 756 Crowther, T. W., van den Hoogen, J., Wan, J., Mayes, M. A., Keiser, A. D., Mo, L., Averill, C., and Maynard, D. S.: The
757 global soil community and its influence on biogeochemistry, *Science*, 365, eaav0550,
758 <https://doi.org/10.1126/science.aav0550>, 2019.
- 759 Dai, W., Xiao, R., Wei, C., and Yang, F.: Plant litter traits control the accumulation of mineral-associated organic carbon
760 by influencing its molecular composition and diversity, *Soil Tillage Res.*, 253, 106667,
761 <https://doi.org/10.1016/j.still.2025.106667>, 2025.
- 762 Daou, L., Garnier, É., and Shipley, B.: Quantifying the relationship linking the community-weighted means of plant traits
763 and soil fertility, *Ecology*, 102, e03454, <https://doi.org/10.1002/ecy.3454>, 2021.
- 764 De Deyn, G. B., Quirk, H., and Bardgett, R. D.: Plant species richness, identity and productivity differentially influence
765 key groups of microbes in grassland soils of contrasting fertility, *Biol. Lett.*, 7, 75–78,
766 <https://doi.org/10.1098/rsbl.2010.0575>, 2010.
- 767 Díaz, S., Kattge, J., Cornelissen, J. H. C., Wright, I. J., Lavorel, S., Dray, S., Reu, B., Kleyer, M., Wirth, C., Colin Prentice,
768 I., Garnier, E., Bönsch, G., Westoby, M., Poorter, H., Reich, P. B., Moles, A. T., Dickie, J., Gillison, A. N., Zanne,
769 A. E., Chave, J., Joseph Wright, S., Sheremet'ev, S. N., Jactel, H., Baraloto, C., Cerabolini, B., Pierce, S., Shipley,
770 B., Kirkup, D., Casanoves, F., Joswig, J. S., Günther, A., Falczuk, V., Rüger, N., Mahecha, M. D., and Gorné, L.
771 D.: The global spectrum of plant form and function, *Nature*, 529, 167–171, <https://doi.org/10.1038/nature16489>,
772 2016.
- 773 Díaz-Martínez, P., Maestre, F. T., Moreno-Jiménez, E., Delgado-Baquerizo, M., Eldridge, D. J., Saiz, H., Gross, N., Le
774 Bagousse-Pinguet, Y., Gozalo, B., Ochoa, V., Guirado, E., García-Gómez, M., Valencia, E., Asensio, S., Berdugo,
775 M., Martínez-Valderrama, J., Mendoza, B. J., García-Gil, J. C., Zaccane, C., Panettieri, M., García-Palacios, P.,
776 Fan, W., Benavente-Ferraces, I., Rey, A., Eisenhauer, N., Cesarz, S., Abedi, M., Ahumada, R. J., Alcántara, J.
777 M., Amghar, F., Aramayo, V., Arroyo, A. I., Bahalkeh, K., Ben Salem, F., Blaum, N., Boldgiv, B., Bowker, M.
778 A., Bran, D., Branquinho, C., Bu, C., Cáceres, Y., Canessa, R., Castillo-Monroy, A. P., Castro, I., Castro-Quezada,
779 P., Chibani, R., Conceição, A. A., Currier, C. M., Darrouzet-Nardi, A., Deák, B., Dickman, C. R., Donoso, D. A.,
780 Dougill, A. J., Durán, J., Ejtehadi, H., Espinosa, C., Fajardo, A., Farzam, M., Ferrante, D., Fraser, L. H., Gaitán,
781 J. J., Gusman Montalván, E., Hernández-Hernández, R. M., von Hessberg, A., Hölzel, N., Huber-Sannwald, E.,
782 Hughes, F. M., Jadán-Maza, O., Geissler, K., Jentsch, A., Ju, M., Kaseke, K. F., Kindermann, L., Koopman, J.
783 E., Le Roux, P. C., Liancourt, P., Linstädter, A., Liu, J., Louw, M. A., Maggs-Köling, G., Makhallanyane, T. P.,
784 Issa, O. M., Marais, E., Margerie, P., Mazaneda, A. J., McClaran, M. P., Messeder, J. V. S., Mora, J. P., Moreno,
785 G., Munson, S. M., Nunes, A., Oliva, G., Oñativia, G. R., Osborne, B., Peter, G., Pueyo, Y., Quiroga, R. E., Reed,
786 S. C., Reyes, V. M., et al.: Vulnerability of mineral-associated soil organic carbon to climate across global
787 drylands, *Nat. Clim. Change*, 14, 976–982, <https://doi.org/10.1038/s41558-024-02087-y>, 2024.
- 788 Dondini, M., Martin, M., De Camillis, C., Uwizeye, A., Soussana, J.-F., Robinson, T., and Steinfeld, H.: Global assessment
789 of soil carbon in grasslands – From current stock estimates to sequestration potential., *FAO*, Rome, 2023.

790 Du, Z., Angers, D. A., Ren, T., Zhang, Q., and Li, G.: The effect of no-till on organic C storage in Chinese soils should
791 not be overemphasized: A meta-analysis, *Agric. Ecosyst. Environ.*, 236, 1–11,
792 <https://doi.org/10.1016/j.agee.2016.11.007>, 2017.

793 Dubeux Jr., J. c. b., Sollenberger, L. e., Vendramini, J. m. b., Interrante, S. m., and Lira Jr., M. a.: Stocking Method,
794 Animal Behavior, and Soil Nutrient Redistribution: How are They Linked?, *Crop Sci.*, 54, 2341–2350,
795 <https://doi.org/10.2135/cropsci2014.01.0076>, 2014.

796 Eisenhauer, N., Lanoue, A., Strecker, T., Scheu, S., Steinauer, K., Thakur, M. P., and Mommer, L.: Root biomass and
797 exudates link plant diversity with soil bacterial and fungal biomass, *Sci. Rep.*, 7, 44641,
798 <https://doi.org/10.1038/srep44641>, 2017.

799 Elias, D. M. O., Mason, K. E., Goodall, T., Taylor, A., Zhao, P., Otero-Fariña, A., Chen, H., Peacock, C. L., Ostle, N. J.,
800 Griffiths, R., Chapman, P. J., Holden, J., Banwart, S., McNamara, N. P., and Whitaker, J.: Microbial and mineral
801 interactions decouple litter quality from soil organic matter formation, *Nat. Commun.*, 15, 10063,
802 <https://doi.org/10.1038/s41467-024-54446-0>, 2024.

803 Eze, S., Palmer, S. M., and Chapman, P. J.: Soil organic carbon stock in grasslands: Effects of inorganic fertilizers, liming
804 and grazing in different climate settings, *J. Environ. Manage.*, 223, 74–84,
805 <https://doi.org/10.1016/j.jenvman.2018.06.013>, 2018.

806 Fanin, N., Kardol, P., Farrell, M., Nilsson, M.-C., Gundale, M. J., and Wardle, D. A.: The ratio of Gram-positive to Gram-
807 negative bacterial PLFA markers as an indicator of carbon availability in organic soils, *Soil Biol. Biochem.*, 128,
808 111–114, <https://doi.org/10.1016/j.soilbio.2018.10.010>, 2019.

809 Faucon, M.-P., Houben, D., and Lambers, H.: Plant Functional Traits: Soil and Ecosystem Services, *Trends Plant Sci.*, 22,
810 385–394, <https://doi.org/10.1016/j.tplants.2017.01.005>, 2017.

811 Fay, P. A., Prober, S. M., Harpole, W. S., Knops, J. M. H., Bakker, J. D., Borer, E. T., Lind, E. M., MacDougall, A. S.,
812 Seabloom, E. W., Wragg, P. D., Adler, P. B., Blumenthal, D. M., Buckley, Y. M., Chu, C., Cleland, E. E., Collins,
813 S. L., Davies, K. F., Du, G., Feng, X., Firn, J., Gruner, D. S., Hagenah, N., Hautier, Y., Heckman, R. W., Jin, V.
814 L., Kirkman, K. P., Klein, J., Ladwig, L. M., Li, Q., McCulley, R. L., Melbourne, B. A., Mitchell, C. E., Moore,
815 J. L., Morgan, J. W., Risch, A. C., Schütz, M., Stevens, C. J., Wedin, D. A., and Yang, L. H.: Grassland
816 productivity limited by multiple nutrients, *Nat. Plants*, 1, 1–5, <https://doi.org/10.1038/nplants.2015.80>, 2015.

817 Feßel, C., Meier, I. C., and Leuschner, C.: Relationship between species diversity, biomass and light transmittance in
818 temperate semi-natural grasslands: is productivity enhanced by complementary light capture?, *J. Veg. Sci.*, 27,
819 144–155, <https://doi.org/10.1111/jvs.12326>, 2016.

820 Fortunel, C., Garnier, E., Joffre, R., Kazakou, E., Quested, H., Grigulis, K., Lavorel, S., Ansquer, P., Castro, H., Cruz, P.,
821 Doležal, J., Eriksson, O., Freitas, H., Golodets, C., Jouany, C., Kigel, J., Kleyer, M., Lehsten, V., Lepš, J., Meier,
822 T., Pakeman, R., Papadimitriou, M., Papanastasis, V. P., Quétier, F., Robson, M., Sternberg, M., Theau, J.-P.,
823 Thébaud, A., and Zarovali, M.: Leaf traits capture the effects of land use changes and climate on litter
824 decomposability of grasslands across Europe, *Ecology*, 90, 598–611, <https://doi.org/10.1890/08-0418.1>, 2009.

825 Freschet, G. T., Aerts, R., and Cornelissen, J. H. C.: A plant economics spectrum of litter decomposability, *Funct. Ecol.*,
826 26, 56–65, <https://doi.org/10.1111/j.1365-2435.2011.01913.x>, 2012.

827 Frongia, A., Pulina, A., Tanda, A., Seddaiu, G., Roggero, P. P., and Moreno, G.: Assessing the effect of rotational grazing
828 adoption in Iberian silvopastoral systems with Normalized Difference Vegetation Index time series, *Ital. J.*
829 *Agron.*, <https://doi.org/10.4081/ija.2023.2185>, 2023.

830 Frostegård, A. and Bååth, E.: The use of phospholipid fatty acid analysis to estimate bacterial and fungal biomass in soil,
831 *Biol. Fertil. Soils*, 22, 59–65, <https://doi.org/10.1007/BF00384433>, 1996.

832 Funk, J. L., Larson, J. E., Ames, G. M., Butterfield, B. J., Cavender-Bares, J., Firn, J., Laughlin, D. C., Sutton-Grier, A.
833 E., Williams, L., and Wright, J.: Revisiting the Holy Grail: using plant functional traits to understand ecological
834 processes, *Biol. Rev.*, 92, 1156–1173, <https://doi.org/10.1111/brv.12275>, 2017.

835 Galluzzi, G., Plaza, C., Giannetta, B., Priori, S., and Zaccone, C.: Time and climate roles in driving soil carbon distribution
836 and stability in particulate and mineral-associated organic matter pools, *Sci. Total Environ.*, 963, 178511,
837 <https://doi.org/10.1016/j.scitotenv.2025.178511>, 2025.

838 Gang, C., Zhou, W., Chen, Y., Wang, Z., Sun, Z., Li, J., Qi, J., and Odeh, I.: Quantitative assessment of the contributions
839 of climate change and human activities on global grassland degradation, *Environ. Earth Sci.*, 72, 4273–4282,
840 <https://doi.org/10.1007/s12665-014-3322-6>, 2014.

841 Gao, Y., Liu, J., Wang, D., An, Y., Ma, H., and Tong, S.: Synergy and trade-off between plant functional traits enhance
842 grassland multifunctionality under grazing exclusion in a semi-arid region, *J. Environ. Manage.*, 373, 123877,
843 <https://doi.org/10.1016/j.jenvman.2024.123877>, 2025.

844 García Bravo, N., Dimas, M., Murillo Díaz, J. M., Barranco Sanz, L. M., and Ruiz Hernández, J. M.: Ejemplo de
845 aplicación del nuevo modelo hidrogeológico en SIMPA para mejorar la evaluación de los recursos hídricos a
846 escala nacional en España, Centro Nacional de Información Geográfica (España), 2023.

847 Gaudet, C. L. and Keddy, P. A.: A comparative approach to predicting competitive ability from plant traits, *Nature*, 334,
848 242–243, <https://doi.org/10.1038/334242a0>, 1988.

849 Georgiou, K., Jackson, R. B., Vindušková, O., Abramoff, R. Z., Ahlström, A., Feng, W., Harden, J. W., Pellegrini, A. F.
850 A., Polley, H. W., Soong, J. L., Riley, W. J., and Torn, M. S.: Global stocks and capacity of mineral-associated
851 soil organic carbon, *Nat. Commun.*, 13, 3797, <https://doi.org/10.1038/s41467-022-31540-9>, 2022.

852 Georgiou, K., Koven, C. D., Wieder, W. R., Hartman, M. D., Riley, W. J., Pett-Ridge, J., Bouskill, N. J., Abramoff, R. Z.,
853 Slessarev, E. W., Ahlström, A., Parton, W. J., Pellegrini, A. F. A., Pierson, D., Sulman, B. N., Zhu, Q., and
854 Jackson, R. B.: Emergent temperature sensitivity of soil organic carbon driven by mineral associations, *Nat.*
855 *Geosci.*, 17, 205–212, <https://doi.org/10.1038/s41561-024-01384-7>, 2024.

856 Georgiou, K., Angers, D., Champiny, R. E., Cotrufo, M. F., Craig, M. E., Doetterl, S., Grandy, A. S., Lavallee, J. M., Lin,
857 Y., Lugato, E., Poeplau, C., Rocci, K. S., Schweizer, S. A., Six, J., and Wieder, W. R.: Soil Carbon Saturation:
858 What Do We Really Know?, *Glob. Change Biol.*, 31, e70197, <https://doi.org/10.1111/gcb.70197>, 2025.

859 Gliksman, D., Navon, Y., Dumbur, R., Haenel, S., and Grünzweig, J. M.: Higher rates of decomposition in standing vs.
860 surface litter in a Mediterranean ecosystem during the dry and the wet seasons, *Plant Soil*, 428, 427–439,
861 <https://doi.org/10.1007/s11104-018-3696-4>, 2018.

862 Godde, C. M., Boone, R. B., Ash, A. J., Waha, K., Sloat, L. L., Thornton, P. K., and Herrero, M.: Global rangeland
863 production systems and livelihoods at threat under climate change and variability, *Environ. Res. Lett.*, 15,
864 044021, <https://doi.org/10.1088/1748-9326/ab7395>, 2020.

865 Godínez-Alvarez, H., Herrick, J. E., Mattocks, M., Toledo, D., and Van Zee, J.: Comparison of three vegetation monitoring
866 methods: Their relative utility for ecological assessment and monitoring, *Ecol. Indic.*, 9, 1001–1008,
867 <https://doi.org/10.1016/j.ecolind.2008.11.011>, 2009.

868 Gómez-Rey, M. X., Garcês, A., and Madeira, M.: Soil organic-C accumulation and N availability under improved pastures
869 established in Mediterranean oak woodlands, *Soil Use Manag.*, 28, 497–507, <https://doi.org/10.1111/j.1475-2743.2012.00428.x>, 2012.

871 Gou, X., Reich, P. B., Qiu, L., Shao, M., Wei, G., Wang, J., and Wei, X.: Leguminous plants significantly increase soil
872 nitrogen cycling across global climates and ecosystem types, *Glob. Change Biol.*, 29, 4028–4043,
873 <https://doi.org/10.1111/gcb.16742>, 2023.

874 Griepentrog, M., Bodé, S., Boeckx, P., Hagedorn, F., Heim, A., and Schmidt, M. W. I.: Nitrogen deposition promotes the
875 production of new fungal residues but retards the decomposition of old residues in forest soil fractions, *Glob.*
876 *Change Biol.*, 20, 327–340, <https://doi.org/10.1111/gcb.12374>, 2014.

877 Guyonnet, J. P., Cantarel, A. A. M., Simon, L., and Haichar, F. el Z.: Root exudation rate as functional trait involved in
878 plant nutrient-use strategy classification, *Ecol. Evol.*, 8, 8573–8581, <https://doi.org/10.1002/ece3.4383>, 2018.

879 Haddix, M. L., Paul, E. A., and Cotrufo, M. F.: Dual, differential isotope labeling shows the preferential movement of
880 labile plant constituents into mineral-bonded soil organic matter, *Glob. Change Biol.*, 22, 2301–2312,
881 <https://doi.org/10.1111/gcb.13237>, 2016.

882 Hamilton, E. W., Frank, D. A., Hinchey, P. M., and Murray, T. R.: Defoliation induces root exudation and triggers positive
883 rhizospheric feedbacks in a temperate grassland, *Soil Biol. Biochem.*, 40, 2865–2873,
884 <https://doi.org/10.1016/j.soilbio.2008.08.007>, 2008.

885 Hansen, P. M., Even, R., King, A. E., Lavallee, J., Schipanski, M., and Cotrufo, M. F.: Distinct, direct and climate-
886 mediated environmental controls on global particulate and mineral-associated organic carbon storage, *Glob.*
887 *Change Biol.*, 30, e17080, <https://doi.org/10.1111/gcb.17080>, 2024.

888 Hawkins, H.-J.: A global assessment of Holistic Planned GrazingTM compared with season-long, continuous grazing:
889 meta-analysis findings, *Afr. J. Range Forage Sci.*, 34, 65–75, <https://doi.org/10.2989/10220119.2017.1358213>,
890 2017.

891 He, M., Zhou, G., Yuan, T., van Groenigen, K. J., Shao, J., and Zhou, X.: Grazing intensity significantly changes the C :
892 N : P stoichiometry in grassland ecosystems, *Glob. Ecol. Biogeogr.*, 29, 355–369,
893 <https://doi.org/10.1111/geb.13028>, 2020.

894 Helsen, K., Ceulemans, T., Stevens, C. J., and Honnay, O.: Increasing Soil Nutrient Loads of European Semi-natural
895 Grasslands Strongly Alter Plant Functional Diversity Independently of Species Loss, *Ecosystems*, 17, 169–181,
896 <https://doi.org/10.1007/s10021-013-9714-8>, 2014.

897 den Herder, M., Moreno, G., Mosquera-Losada, R. M., Palma, J. H. N., Sidiropoulou, A., Santiago Freijanes, J. J., Crous-
898 Duran, J., Paulo, J. A., Tomé, M., Pantera, A., Papanastasis, V. P., Mantzanas, K., Pachana, P., Papadopoulos, A.,
899 Plieninger, T., and Burgess, P. J.: Current extent and stratification of agroforestry in the European Union, *Agric.*
900 *Ecosyst. Environ.*, 241, 121–132, <https://doi.org/10.1016/j.agee.2017.03.005>, 2017.

901 Hernández-Esteban, A., López-Díaz, M. L., Cáceres, Y., and Moreno, G.: Are sown legume-rich pastures effective allies
902 for the profitability and sustainability of Mediterranean dehesas?, *Agrofor. Syst.*, 93, 2047–2065,
903 <https://doi.org/10.1007/s10457-018-0307-6>, 2019.

904 Hou, Z., Wang, R., Chang, S., Zheng, Y., Ma, T., Xu, S., Zhang, X., Shi, X., Lu, J., Luo, D., Wang, B., Du, Z., and Wei,
905 Y.: The contribution of microbial necromass to soil organic carbon and influencing factors along a variation of
906 habitats in alpine ecosystems, *Sci. Total Environ.*, 921, 171126, <https://doi.org/10.1016/j.scitotenv.2024.171126>,
907 2024.

908 Hu, Y., Deng, Q., Kätterer, T., Olesen, J. E., Ying, S. C., Ochoa-Hueso, R., Mueller, C. W., Weintraub, M. N., and Chen,
909 J.: Depth-dependent responses of soil organic carbon under nitrogen deposition, *Glob. Change Biol.*, 30, e17247,
910 <https://doi.org/10.1111/gcb.17247>, 2024.

911 Huang, J., Liu, W., Pan, S., Wang, Z., Yang, S., Jia, Z., Wang, Z., Deng, M., Yang, L., Liu, C., Chang, P., and Liu, L.:
912 Divergent contributions of living roots to turnover of different soil organic carbon pools and their links to plant
913 traits, *Funct. Ecol.*, 35, 2821–2830, <https://doi.org/10.1111/1365-2435.13934>, 2021.

914 Intergovernmental Panel on Climate Change (IPCC) (Ed.): Climate Change Information for Regional Impact and for Risk
915 Assessment, in: *Climate Change 2021 – The Physical Science Basis: Working Group I Contribution to the Sixth*
916 *Assessment Report of the Intergovernmental Panel on Climate Change*, Cambridge University Press, Cambridge,
917 1767–1926, <https://doi.org/10.1017/9781009157896.014>, 2023a.

918 Intergovernmental Panel on Climate Change (IPCC) (Ed.): Global Carbon and Other Biogeochemical Cycles and
919 Feedbacks, in: *Climate Change 2021 – The Physical Science Basis: Working Group I Contribution to the Sixth*
920 *Assessment Report of the Intergovernmental Panel on Climate Change*, Cambridge University Press, Cambridge,
921 673–816, <https://doi.org/10.1017/9781009157896.007>, 2023b.

922 Jacobo, E. J., Rodríguez, A. M., Bartoloni, N., and Deregibus, V. A.: Rotational Grazing Effects on Rangeland Vegetation
923 at a Farm Scale, *Rangel. Ecol. Manag.*, 59, 249–257, <https://doi.org/10.2111/05-129R1.1>, 2006.

924 Jaurena, M., Lezama, F., Salvo, L., Cardozo, G., Ayala, W., Terra, J., and Nabinger, C.: The Dilemma of Improving Native
925 Grasslands by Overseeding Legumes: Production Intensification or Diversity Conservation, *Rangel. Ecol.*
926 *Manag.*, 69, 35–42, <https://doi.org/10.1016/j.rama.2015.10.006>, 2016.

927 Jongen, M., Förster, A. C., and Unger, S.: Overwhelming effects of autumn-time drought during seedling establishment
928 impair recovery potential in sown and semi-natural pastures in Portugal, *Plant Ecol.*, 220, 183–197,
929 <https://doi.org/10.1007/s11258-018-0869-4>, 2019.

930 Kallenbach, C. M., Frey, S. D., and Grandy, A. S.: Direct evidence for microbial-derived soil organic matter formation
931 and its ecophysiological controls, *Nat. Commun.*, 7, 13630, <https://doi.org/10.1038/ncomms13630>, 2016.

- 932 Kazakou, E., Violle, C., Roumet, C., Pintor, C., Gimenez, O., and Garnier, E.: Litter quality and decomposability of
933 species from a Mediterranean succession depend on leaf traits but not on nitrogen supply, *Ann. Bot.*, 104, 1151–
934 1161, <https://doi.org/10.1093/aob/mcp202>, 2009.
- 935 Khatiwada, B., Acharya, S. N., Larney, F. J., Lupwayi, N. Z., Smith, E. G., Islam, M. A., and Thomas, J. E.: Benefits of
936 mixed grass–legume pastures and pasture rejuvenation using bloat-free legumes in western Canada: a review,
937 *Can. J. Plant Sci.*, 100, 463–476, <https://doi.org/10.1139/cjps-2019-0212>, 2020.
- 938 Khatri-Chhetri, U., Thompson, K. A., Quideau, S. A., Boyce, M. S., Chang, S. X., Bork, E. W., and Carlyle, C. N.:
939 Adaptive multi-paddock grazing increases mineral associated soil carbon in Northern grasslands, *Agric. Ecosyst.*
940 *Environ.*, 369, 109000, <https://doi.org/10.1016/j.agee.2024.109000>, 2024.
- 941 King, A. E., Amsili, J. P., Córdova, S. C., Culman, S., Fonte, S. J., Kotcon, J., Liebig, M., Masters, M. D., McVay, K.,
942 Olk, D. C., Schipanski, M., Schneider, S. K., Stewart, C. E., and Cotrufo, M. F.: A soil matrix capacity index to
943 predict mineral-associated but not particulate organic carbon across a range of climate and soil pH,
944 *Biogeochemistry*, 165, 1–14, <https://doi.org/10.1007/s10533-023-01066-3>, 2023.
- 945 Klumpp, K., Fontaine, S., Attard, E., Le Roux, X., Gleixner, G., and Soussana, J.-F.: Grazing triggers soil carbon loss by
946 altering plant roots and their control on soil microbial community, *J. Ecol.*, 97, 876–885,
947 <https://doi.org/10.1111/j.1365-2745.2009.01549.x>, 2009.
- 948 Kramer, C. and Gleixner, G.: Soil organic matter in soil depth profiles: Distinct carbon preferences of microbial groups
949 during carbon transformation, *Soil Biol. Biochem.*, 40, 425–433, <https://doi.org/10.1016/j.soilbio.2007.09.016>,
950 2008.
- 951 Kramer-Walter, K. R., Bellingham, P. J., Millar, T. R., Smissen, R. D., Richardson, S. J., and Laughlin, D. C.: Root traits
952 are multidimensional: specific root length is independent from root tissue density and the plant economic
953 spectrum, *J. Ecol.*, 104, 1299–1310, <https://doi.org/10.1111/1365-2745.12562>, 2016.
- 954 Laliberté, E. and Legendre, P.: A distance-based framework for measuring functional diversity from multiple traits,
955 *Ecology*, 91, 299–305, <https://doi.org/10.1890/08-2244.1>, 2010.
- 956 Laliberté, E. and Tylianakis, J. M.: Cascading effects of long-term land-use changes on plant traits and ecosystem
957 functioning, *Ecology*, 93, 145–155, <https://doi.org/10.1890/11-0338.1>, 2012.
- 958 Lange, M., Eisenhauer, N., Sierra, C. A., Bessler, H., Engels, C., Griffiths, R. I., Mellado-Vázquez, P. G., Malik, A. A.,
959 Roy, J., Scheu, S., Steinbeiss, S., Thomson, B. C., Trumbore, S. E., and Gleixner, G.: Plant diversity increases
960 soil microbial activity and soil carbon storage, *Nat. Commun.*, 6, 6707, <https://doi.org/10.1038/ncomms7707>,
961 2015.
- 962 Latimer, G. W., Jr. (Ed.): *Official Methods of Analysis of AOAC INTERNATIONAL*, Oxford University Press,
963 <https://doi.org/10.1093/9780197610145.001.0001>, 2023.
- 964 Lavalley, J. M., Conant, R. T., Paul, E. A., and Cotrufo, M. F.: Incorporation of shoot versus root-derived ¹³C and ¹⁵N
965 into mineral-associated organic matter fractions: results of a soil slurry incubation with dual-labelled plant
966 material, *Biogeochemistry*, 137, 379–393, <https://doi.org/10.1007/s10533-018-0428-z>, 2018.
- 967 Lavalley, J. M., Soong, J. L., and Cotrufo, M. F.: Conceptualizing soil organic matter into particulate and mineral-
968 associated forms to address global change in the 21st century, *Glob. Change Biol.*, 26, 261–273,
969 <https://doi.org/10.1111/gcb.14859>, 2020.
- 970 Lefcheck, J. S.: piecewiseSEM: Piecewise structural equation modelling in r for ecology, evolution, and systematics,
971 *Methods Ecol. Evol.*, 7, 573–579, <https://doi.org/10.1111/2041-210X.12512>, 2016.
- 972 Lei, J., Guo, X., Zeng, Y., Zhou, J., Gao, Q., and Yang, Y.: Temporal changes in global soil respiration since 1987, *Nat.*
973 *Commun.*, 12, 403, <https://doi.org/10.1038/s41467-020-20616-z>, 2021.
- 974 Leuthold, S., Lavalley, J. M., Haddix, M. L., and Cotrufo, M. F.: Contrasting properties of soil organic matter fractions
975 isolated by different physical separation methodologies, *Geoderma*, 445, 116870,
976 <https://doi.org/10.1016/j.geoderma.2024.116870>, 2024.
- 977 Li, S. and Li, X.: Global understanding of farmland abandonment: A review and prospects, *J. Geogr. Sci.*, 27, 1123–1150,
978 <https://doi.org/10.1007/s11442-017-1426-0>, 2017.

- 979 Li, Y., Luo, T., and Lu, Q.: Plant height as a simple predictor of the root to shoot ratio: Evidence from alpine grasslands
980 on the Tibetan Plateau, *J. Veg. Sci.*, 19, 245–252, <https://doi.org/10.3170/2007-8-18365>, 2008.
- 981 Liang, C., Amelung, W., Lehmann, J., and Kästner, M.: Quantitative assessment of microbial necromass contribution to
982 soil organic matter, *Glob. Change Biol.*, 25, 3578–3590, <https://doi.org/10.1111/gcb.14781>, 2019.
- 983 Liu, H. Y., Huang, N., Zhao, C. M., and Li, J. H.: Responses of carbon cycling and soil organic carbon content to nitrogen
984 addition in grasslands globally, *Soil Biol. Biochem.*, 186, 109164, <https://doi.org/10.1016/j.soilbio.2023.109164>,
985 2023.
- 986 Liu, Y., Zhang, M., Wang, X., and Wang, C.: The impact of different grazing intensity and management measures on soil
987 organic carbon density in Zhangye grassland, *Sci. Rep.*, 14, 17556, [https://doi.org/10.1038/s41598-024-68277-](https://doi.org/10.1038/s41598-024-68277-y)
988 y, 2024.
- 989 Lu, M., Zhou, X., Luo, Y., Yang, Y., Fang, C., Chen, J., and Li, B.: Minor stimulation of soil carbon storage by nitrogen
990 addition: A meta-analysis, *Agric. Ecosyst. Environ.*, 140, 234–244, <https://doi.org/10.1016/j.agee.2010.12.010>,
991 2011.
- 992 Luo, R., Fan, J., Wang, W., Luo, J., Kuzyakov, Y., He, J.-S., Chu, H., and Ding, W.: Nitrogen and phosphorus enrichment
993 accelerates soil organic carbon loss in alpine grassland on the Qinghai-Tibetan Plateau, *Sci. Total Environ.*, 650,
994 303–312, <https://doi.org/10.1016/j.scitotenv.2018.09.038>, 2019.
- 995 Luo, R., Kuzyakov, Y., Liu, D., Fan, J., Luo, J., Lindsey, S., He, J.-S., and Ding, W.: Nutrient addition reduces carbon
996 sequestration in a Tibetan grassland soil: Disentangling microbial and physical controls, *Soil Biol. Biochem.*,
997 144, 107764, <https://doi.org/10.1016/j.soilbio.2020.107764>, 2020.
- 998 Maestre, F. T., Le Bagousse-Pinguet, Y., Delgado-Baquerizo, M., Eldridge, D. J., Saiz, H., Berdugo, M., Gozalo, B.,
999 Ochoa, V., Guirado, E., García-Gómez, M., Valencia, E., Gaitán, J. J., Asensio, S., Mendoza, B. J., Plaza, C.,
1000 Díaz-Martínez, P., Rey, A., Hu, H.-W., He, J.-Z., Wang, J.-T., Lehmann, A., Rillig, M. C., Cesarz, S., Eisenhauer,
1001 N., Martínez-Valderrama, J., Moreno-Jiménez, E., Sala, O., Abedi, M., Ahmadian, N., Alados, C. L., Aramayo,
1002 V., Amghar, F., Arredondo, T., Ahumada, R. J., Bahalkeh, K., Ben Salem, F., Blaum, N., Boldgiv, B., Bowker,
1003 M. A., Bran, D., Bu, C., Canessa, R., Castillo-Monroy, A. P., Castro, H., Castro, I., Castro-Quezada, P., Chibani,
1004 R., Conceição, A. A., Currier, C. M., Darrouzet-Nardi, A., Deák, B., Donoso, D. A., Dougill, A. J., Durán, J.,
1005 Erdenetsetseg, B., Espinosa, C. I., Fajardo, A., Farzam, M., Ferrante, D., Frank, A. S. K., Fraser, L. H., Gherardi,
1006 L. A., Greenville, A. C., Guerra, C. A., Gusmán-Montalvan, E., Hernández-Hernández, R. M., Hölzel, N., Huber-
1007 Sannwald, E., Hughes, F. M., Jadán-Maza, O., Jeltsch, F., Jentsch, A., Kaseke, K. F., Köbel, M., Koopman, J. E.,
1008 Leder, C. V., Linstädter, A., le Roux, P. C., Li, X., Liancourt, P., Liu, J., Louw, M. A., Maggs-Kölling, G.,
1009 Makhalanyane, T. P., Issa, O. M., Manzaneda, A. J., Marais, E., Mora, J. P., Moreno, G., Munson, S. M., Nunes,
1010 A., Oliva, G., Oñatibia, G. R., Peter, G., Pivari, M. O. D., Pueyo, Y., Quiroga, R. E., Rahmanian, S., Reed, S. C.,
1011 et al.: Grazing and ecosystem service delivery in global drylands, *Science*, 378, 915–920,
1012 <https://doi.org/10.1126/science.abq4062>, 2022.
- 1013 Manning, P., de Vries, F. T., Tallowin, J. R. B., Smith, R., Mortimer, S. R., Pilgrim, E. S., Harrison, K. A., Wright, D. G.,
1014 Quirk, H., Benson, J., Shipley, B., Cornelissen, J. H. C., Kattge, J., Bönisch, G., Wirth, C., and Bardgett, R. D.:
1015 Simple measures of climate, soil properties and plant traits predict national-scale grassland soil carbon stocks, *J.*
1016 *Appl. Ecol.*, 52, 1188–1196, <https://doi.org/10.1111/1365-2664.12478>, 2015.
- 1017 McSherry, M. E. and Ritchie, M. E.: Effects of grazing on grassland soil carbon: a global review, *Glob. Change Biol.*, 19,
1018 1347–1357, <https://doi.org/10.1111/gcb.12144>, 2013.
- 1019 Mokany, K., Ash, J., and Roxburgh, S.: Functional identity is more important than diversity in influencing ecosystem
1020 processes in a temperate native grassland, *J. Ecol.*, 96, 884–893, [https://doi.org/10.1111/j.1365-](https://doi.org/10.1111/j.1365-2745.2008.01395.x)
1021 2745.2008.01395.x, 2008.
- 1022 Moreno, G. and Pulido, F. J.: The Functioning, Management and Persistence of Dehesas, in: *Agroforestry in Europe:*
1023 *Current Status and Future Prospects*, edited by: Rigueiro-Rodríguez, A., McAdam, J., and Mosquera-Losada,
1024 M. R., Springer Netherlands, Dordrecht, 127–160, https://doi.org/10.1007/978-1-4020-8272-6_7, 2009.
- 1025 Moreno, G., Obrador, J. J., Cubera, E., and Dupraz, C.: Fine Root Distribution in Dehesas of Central-Western Spain, *Plant*
1026 *Soil*, 277, 153–162, <https://doi.org/10.1007/s11104-005-6805-0>, 2005.
- 1027 Moreno, G., Hernández-Esteban, A., Rolo, V., and Igual, J. M.: The enduring effects of sowing legume-rich mixtures on
1028 the soil microbial community and soil carbon in semi-arid wood pastures, *Plant Soil*, 465, 563–582,
1029 <https://doi.org/10.1007/s11104-021-05023-7>, 2021.

- 1030 Moreno-Jiménez, E., Plaza, C., Saiz, H., Manzano, R., Flagmeier, M., and Maestre, F. T.: Aridity and reduced soil
1031 micronutrient availability in global drylands, *Nat. Sustain.*, 2, 371–377, [https://doi.org/10.1038/s41893-019-](https://doi.org/10.1038/s41893-019-0262-x)
1032 0262-x, 2019.
- 1033 Mortensen, E. Ø., Abalos, D., Engedal, T., Lægsgaard, A. K., Enggrob, K., Mueller, C. W., and Rasmussen, J.: Smart
1034 Mixture Design Can Steer the Fate of Root-Derived Carbon Into Mineral-Associated and Particulate Organic
1035 Matter in Intensively Managed Grasslands, *Glob. Change Biol.*, 31, e70117, <https://doi.org/10.1111/gcb.70117>,
1036 2025.
- 1037 Mosier, S., Apfelbaum, S., Byck, P., Calderon, F., Teague, R., Thompson, R., and Cotrufo, M. F.: Adaptive multi-paddock
1038 grazing enhances soil carbon and nitrogen stocks and stabilization through mineral association in southeastern
1039 U.S. grazing lands, *J. Environ. Manage.*, 288, 112409, <https://doi.org/10.1016/j.jenvman.2021.112409>, 2021.
- 1040 Murphy, M. V.: *semEff: Automatic Calculation of Effects for Piecewise Structural Equation Models*, 2022.
- 1041 Nair, R. K. F., Morris, K. A., Hertel, M., Luo, Y., Moreno, G., Reichstein, M., Schrumpf, M., and Migliavacca, M.:
1042 N  P stoichiometry and habitat effects on Mediterranean savanna seasonal root dynamics,
1043 *Biogeosciences*, 16, 1883–1901, <https://doi.org/10.5194/bg-16-1883-2019>, 2019.
- 1044 Niu, W., Ding, J., Fu, B., Zhao, W., and Eldridge, D.: Global effects of livestock grazing on ecosystem functions vary
1045 with grazing management and environment, *Agric. Ecosyst. Environ.*, 378, 109296,
1046 <https://doi.org/10.1016/j.agee.2024.109296>, 2025.
- 1047 Novelty, P. E. and Watson, I. W.: Successful Grassland Regeneration in a Severely Degraded Catchment: a Whole of
1048 Government Approach in North West Australia, in: *Climate and Land Degradation*, edited by: Sivakumar, M. V.
1049 K. and Ndiang'ui, N., Springer, Berlin, Heidelberg, 469–485, https://doi.org/10.1007/978-3-540-72438-4_26,
1050 2007.
- 1051 Oggioni, S. D., Ochoa-Hueso, R., and Peco, B.: Livestock grazing abandonment reduces soil microbial activity and carbon
1052 storage in a Mediterranean Dehesa, *Appl. Soil Ecol.*, 153, 103588, <https://doi.org/10.1016/j.apsoil.2020.103588>,
1053 2020.
- 1054 Oliveira Filho, J. de S., Vieira, J. N., Ribeiro da Silva, E. M., Beserra de Oliveira, J. G., Pereira, M. G., and Brasileiro, F.
1055 G.: Assessing the effects of 17 years of grazing exclusion in degraded semi-arid soils: Evaluation of soil fertility,
1056 nutrients pools and stoichiometry, *J. Arid Environ.*, 166, 1–10, <https://doi.org/10.1016/j.jaridenv.2019.03.006>,
1057 2019.
- 1058 Ordoñez, J. C., Van Bodegom, P. M., Witte, J.-P. M., Wright, I. J., Reich, P. B., and Aerts, R.: A global study of
1059 relationships between leaf traits, climate and soil measures of nutrient fertility, *Glob. Ecol. Biogeogr.*, 18, 137–
1060 149, <https://doi.org/10.1111/j.1466-8238.2008.00441.x>, 2009.
- 1061 Palomo-Campesino, S., Ravera, F., González, J. A., and García-Llorente, M.: Exploring Current and Future Situation of
1062 Mediterranean Silvopastoral Systems: Case Study in Southern Spain, *Rangel. Ecol. Manag.*, 71, 578–591,
1063 <https://doi.org/10.1016/j.rama.2017.12.013>, 2018.
- 1064 Parras-Alcántara, L., Díaz-Jaimes, L., Lozano-García, B., Rebollo, P. F., Elcure, F. M., and Muñoz, M. D. C.: Organic
1065 farming has little effect on carbon stock in a Mediterranean dehesa (southern Spain), *CATENA*, 113, 9–17,
1066 <https://doi.org/10.1016/j.catena.2013.09.002>, 2014.
- 1067 Parras-Alcántara, L., Díaz-Jaimes, L., and Lozano-García, B.: Management Effects on Soil Organic Carbon Stock in
1068 Mediterranean Open Rangelands—Treeless Grasslands, *Land Degrad. Dev.*, 26, 22–34,
1069 <https://doi.org/10.1002/ldr.2269>, 2015.
- 1070 Peco, B., Navarro, E., Carmona, C. P., Medina, N. G., and Marques, M. J.: Effects of grazing abandonment on soil
1071 multifunctionality: The role of plant functional traits, *Agric. Ecosyst. Environ.*, 249, 215–225,
1072 <https://doi.org/10.1016/j.agee.2017.08.013>, 2017.
- 1073 Pérez-Harguindeguy, N., Díaz, S., Garnier, E., Lavorel, S., Poorter, H., Jaureguiberry, P., Bret-Harte, M. S., Cornwell, W.
1074 K., Craine, J. M., Gurvich, D. E., Urcelay, C., Veneklaas, E. J., Reich, P. B., Poorter, L., Wright, I. J., Ray, P.,
1075 Enrico, L., Pausas, J. G., Vos, A. C. de, Buchmann, N., Funes, G., Quétier, F., Hodgson, J. G., Thompson, K.,
1076 Morgan, H. D., Steege, H. ter, Sack, L., Blonder, B., Poschlod, P., Vaieretti, M. V., Conti, G., Staver, A. C.,
1077 Aquino, S., and Cornelissen, J. H. C.: Corrigendum to: New handbook for standardised measurement of plant
1078 functional traits worldwide, *Aust. J. Bot.*, 64, 715–716, https://doi.org/10.1071/bt12225_co, 2016.

- 1079 Phukubye, K., Mutema, M., Buthelezi, N., Muchaonyerwa, P., Cerri, C., and Chaplot, V.: On the impact of grassland
1080 management on soil carbon stocks: a worldwide meta-analysis, *Geoderma Reg.*, 28, e00479,
1081 <https://doi.org/10.1016/j.geodrs.2021.e00479>, 2022.
- 1082 Pino, A. D., Rodríguez, T., and Andión, J.: Production improvement through phosphorus fertilization and legume
1083 introduction in grazed native pastures of Uruguay, *J. Agric. Sci.*, 154, 347–358,
1084 <https://doi.org/10.1017/S002185961500101X>, 2016.
- 1085 Plaza, C., García-Palacios, P., Berhe, A. A., Barquero, J., Bastida, F., Png, G. K., Rey, A., Bardgett, R. D., and Delgado-
1086 Baquerizo, M.: Ecosystem productivity has a stronger influence than soil age on surface soil carbon storage
1087 across global biomes, *Commun. Earth Environ.*, 3, 1–8, <https://doi.org/10.1038/s43247-022-00567-7>, 2022.
- 1088 Poeplau, C., Vos, C., and Don, A.: Soil organic carbon stocks are systematically overestimated by misuse of the parameters
1089 bulk density and rock fragment content, *SOIL*, 3, 61–66, <https://doi.org/10.5194/soil-3-61-2017>, 2017.
- 1090 Poeplau, C., Helfrich, M., Dechow, R., Szoboszlay, M., Tebbe, C. C., Don, A., Greiner, B., Zopf, D., Thumm, U.,
1091 Korevaar, H., and Geerts, R.: Increased microbial anabolism contributes to soil carbon sequestration by mineral
1092 fertilization in temperate grasslands, *Soil Biol. Biochem.*, 130, 167–176,
1093 <https://doi.org/10.1016/j.soilbio.2018.12.019>, 2019.
- 1094 Poeplau, C., Dechow, R., Begill, N., and Don, A.: Towards an ecosystem capacity to stabilise organic carbon in soils,
1095 *Glob. Change Biol.*, 30, e17453, <https://doi.org/10.1111/gcb.17453>, 2024.
- 1096 Poorter, H., Niinemets, Ü., Poorter, L., Wright, I. J., and Villar, R.: Causes and consequences of variation in leaf mass per
1097 area (LMA): a meta-analysis, *New Phytol.*, 182, 565–588, <https://doi.org/10.1111/j.1469-8137.2009.02830.x>,
1098 2009.
- 1099 Porqueddu, C., Ates, S., Louhaichi, M., Kyriazopoulos, A. P., Moreno, G., del Pozo, A., Ovalle, C., Ewing, M. A., and
1100 Nichols, P. G. H.: Grasslands in ‘Old World’ and ‘New World’ Mediterranean-climate zones: past trends, current
1101 status and future research priorities, *Grass Forage Sci.*, 71, 1–35, <https://doi.org/10.1111/gfs.12212>, 2016.
- 1102 Pulina, A., Rolo, V., Hernández-Esteban, A., Seddaiu, G., Roggero, P. P., and Moreno, G.: Long-term legacy of sowing
1103 legume-rich mixtures in Mediterranean wooded grasslands, *Agric. Ecosyst. Environ.*, 348, 108397,
1104 <https://doi.org/10.1016/j.agee.2023.108397>, 2023.
- 1105 R Core Team: R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna,
1106 Austria. URL <https://www.R-project.org/>, 2024.
- 1107 Rama, G., Oyarzabal, M., Cardozo, G., Lezama, F., and Baeza, S.: Legume Overseeding along with P Fertilization
1108 Increase Forage Production of Temperate Natural Grasslands, *Agronomy*, 12, 2507,
1109 <https://doi.org/10.3390/agronomy12102507>, 2022.
- 1110 Ricotta, C. and Moretti, M.: CWM and Rao’s quadratic diversity: a unified framework for functional ecology, *Oecologia*,
1111 167, 181–188, <https://doi.org/10.1007/s00442-011-1965-5>, 2011.
- 1112 Ridgeway, J. R., Morrissey, E. M., and Brzostek, E. R.: Plant litter traits control microbial decomposition and drive soil
1113 carbon stabilization, *Soil Biol. Biochem.*, 175, 108857, <https://doi.org/10.1016/j.soilbio.2022.108857>, 2022.
- 1114 Rocci, K. S., Lavalley, J. M., Stewart, C. E., and Cotrufo, M. F.: Soil organic carbon response to global environmental
1115 change depends on its distribution between mineral-associated and particulate organic matter: A meta-analysis,
1116 *Sci. Total Environ.*, 793, 148569, <https://doi.org/10.1016/j.scitotenv.2021.148569>, 2021.
- 1117 Rocci, K. S., Barker, K. S., Seabloom, E. W., Borer, E. T., Hobbie, S. E., Bakker, J. D., MacDougall, A. S., McCulley, R.
1118 L., Moore, J. L., Raynaud, X., Stevens, C. J., and Cotrufo, M. F.: Impacts of nutrient addition on soil carbon and
1119 nitrogen stoichiometry and stability in globally-distributed grasslands, *Biogeochemistry*, 159, 353–370,
1120 <https://doi.org/10.1007/s10533-022-00932-w>, 2022.
- 1121 Rolo, V., Rivest, D., Lorente, M., Kattge, J., and Moreno, G.: Taxonomic and functional diversity in Mediterranean
1122 pastures: insights on the biodiversity–productivity trade-off, *J. Appl. Ecol.*, 53, 1575–1584,
1123 <https://doi.org/10.1111/1365-2664.12685>, 2016.
- 1124 Roumet, C., Birouste, M., Picon-Cochard, C., Ghestem, M., Osman, N., Vrignon-Brenas, S., Cao, K., and Stokes, A.:
1125 Root structure–function relationships in 74 species: evidence of a root economics spectrum related to carbon
1126 economy, *New Phytol.*, 210, 815–826, <https://doi.org/10.1111/nph.13828>, 2016.

- 1127 Rovira, P., Sauras-Yera, T., and Romanyà, J.: Equivalent-mass versus fixed-depth as criteria for quantifying soil carbon
1128 sequestration: How relevant is the difference?, CATENA, 214, 106283,
1129 <https://doi.org/10.1016/j.catena.2022.106283>, 2022.
- 1130 Sanaullah, M., Chabbi, A., Lemaire, G., Charrier, X., and Rumpel, C.: How does plant leaf senescence of grassland species
1131 influence decomposition kinetics and litter compounds dynamics?, Nutr. Cycl. Agroecosystems, 88, 159–171,
1132 <https://doi.org/10.1007/s10705-009-9323-2>, 2010.
- 1133 Sanderman, J., Baldock, J. A., Dangal, S. R. S., Ludwig, S., Potter, S., Rivard, C., and Savage, K.: Soil organic carbon
1134 fractions in the Great Plains of the United States: an application of mid-infrared spectroscopy, Biogeochemistry,
1135 156, 97–114, <https://doi.org/10.1007/s10533-021-00755-1>, 2021.
- 1136 Sanjari, G., Yu, B., Ghadiri, H., Ciesiolka, C. A. A., and Rose, C. W.: Effects of time-controlled grazing on runoff and
1137 sediment loss, Soil Res., 47, 796–808, <https://doi.org/10.1071/SR09032>, 2009.
- 1138 Santos, R. S., Hamilton, E. K., Stanley, P. L., Paustian, K., Cotrufo, M. F., and Zhang, Y.: Simulating adaptive grazing
1139 management on soil organic carbon in the Southeast U.S.A. using MEMS 2, J. Environ. Manage., 365, 121657,
1140 <https://doi.org/10.1016/j.jenvman.2024.121657>, 2024.
- 1141 Sherrod, L. A., Dunn, G., Peterson, G. A., and Kolberg, R. L.: Inorganic Carbon Analysis by Modified Pressure-
1142 Calcimeter Method, Soil Sci. Soc. Am. J., 66, 299–305, <https://doi.org/10.2136/sssaj2002.2990>, 2002.
- 1143 Shi, B., Delgado-Baquerizo, M., Knapp, A. K., Smith, M. D., Reed, S., Osborne, B., Carrillo, Y., Maestre, F. T., Zhu, Y.,
1144 Chen, A., Wilkins, K., Holdrege, M. C., Kulmatiski, A., Picon-Cochard, C., Roscher, C., Power, S., Byrne, K.
1145 M., Churchill, A. C., Jentsch, A., Henry, H. A. L., Beard, K. H., Schuchardt, M. A., Eisenhauer, N., Otfinowski,
1146 R., Hautier, Y., Shen, H., Wang, Y., Wang, Z., Wang, C., Cusack, D. F., Petraglia, A., Carbognani, M., Forte, T.
1147 G. W., Flory, S., Hou, P., Zhang, T., Gao, W., and Sun, W.: Aridity drives the response of soil total and particulate
1148 organic carbon to drought in temperate grasslands and shrublands, Sci. Adv., 10, eadq2654,
1149 <https://doi.org/10.1126/sciadv.adq2654>, 2024.
- 1150 Siefert, A., Violle, C., Chalmandrier, L., Albert, C. H., Taudiere, A., Fajardo, A., Aarssen, L. W., Baraloto, C., Carlucci,
1151 M. B., Cianciaruso, M. V., de L. Dantas, V., de Bello, F., Duarte, L. D. S., Fonseca, C. R., Freschet, G. T.,
1152 Gaucherand, S., Gross, N., Hikosaka, K., Jackson, B., Jung, V., Kamiyama, C., Katabuchi, M., Kembel, S. W.,
1153 Kichenin, E., Kraft, N. J. B., Lagerström, A., Bagousse-Pinguet, Y. L., Li, Y., Mason, N., Messier, J.,
1154 Nakashizuka, T., Overton, J. McC., Peltzer, D. A., Pérez-Ramos, I. M., Pillar, V. D., Prentice, H. C., Richardson,
1155 S., Sasaki, T., Schamp, B. S., Schöb, C., Shipley, B., Sundqvist, M., Sykes, M. T., Vandewalle, M., and Wardle,
1156 D. A.: A global meta-analysis of the relative extent of intraspecific trait variation in plant communities, Ecol.
1157 Lett., 18, 1406–1419, <https://doi.org/10.1111/ele.12508>, 2015.
- 1158 Six, J., Guggenberger, G., Paustian, K., Haumaier, L., Elliott, E. T., and Zech, W.: Sources and composition of soil organic
1159 matter fractions between and within soil aggregates, Eur. J. Soil Sci., 52, 607–618,
1160 <https://doi.org/10.1046/j.1365-2389.2001.00406.x>, 2001.
- 1161 Six, J., Conant, R. T., Paul, E. A., and Paustian, K.: Stabilization mechanisms of soil organic matter: Implications for C-
1162 saturation of soils, Plant Soil, 241, 155–176, <https://doi.org/10.1023/A:1016125726789>, 2002.
- 1163 Smith, M. D., Wilkins, K. D., Holdrege, M. C., Wilfahrt, P., Collins, S. L., Knapp, A. K., Sala, O. E., Dukes, J. S., Phillips,
1164 R. P., Yahdjian, L., Gherardi, L. A., Ohlert, T., Beier, C., Fraser, L. H., Jentsch, A., Loik, M. E., Maestre, F. T.,
1165 Power, S. A., Yu, Q., Felton, A. J., Munson, S. M., Luo, Y., Abdoli, H., Abedi, M., Alados, C. L., Alberti, J., Alon,
1166 M., An, H., Anacker, B., Anderson, M., Auge, H., Bachle, S., Bahalkeh, K., Bahn, M., Batbaatar, A., Bauerle, T.,
1167 Beard, K. H., Behn, K., Beil, I., Biancari, L., Blindow, I., Bondaruk, V. F., Borer, E. T., Bork, E. W., Bruschetti,
1168 C. M., Byrne, K. M., Cahill Jr., J. F., Calvo, D. A., Carbognani, M., Cardoni, A., Carlyle, C. N., Castillo-Garcia,
1169 M., Chang, S. X., Chieppa, J., Cianciaruso, M. V., Cohen, O., Cordeiro, A. L., Cusack, D. F., Dahlke, S., Daleo,
1170 P., D'Antonio, C. M., Dietterich, L. H., S. Doherty, T., Dubbert, M., Ebeling, A., Eisenhauer, N., Fischer, F. M.,
1171 Forte, T. G. W., Gebauer, T., Gozalo, B., Greenville, A. C., Guidoni-Martins, K. G., Hannusch, H. J., Vatsø
1172 Haugum, S., Hautier, Y., Hefting, M., Henry, H. A. L., Hoss, D., Ingrisch, J., Iribarne, O., Isbell, F., Johnson, Y.,
1173 Jordan, S., Kelly, E. F., Kimmel, K., Kreyling, J., Kröel-Dulay, G., Kröpf, A., Kübert, A., Kulmatiski, A., Lamb,
1174 E. G., Larsen, K. S., Larson, J., Lawson, J., Leder, C. V., Linstädter, A., Liu, J., Liu, S., Lodge, A. G., et al.:
1175 Extreme drought impacts have been underestimated in grasslands and shrublands globally, Proc. Natl. Acad. Sci.,
1176 121, e2309881120, <https://doi.org/10.1073/pnas.2309881120>, 2024.
- 1177 Soliveres, S., Maestre, F. T., Eldridge, D. J., Delgado-Baquerizo, M., Quero, J. L., Bowker, M. A., and Gallardo, A.: Plant
1178 diversity and ecosystem multifunctionality peak at intermediate levels of woody cover in global drylands, Glob.
1179 Ecol. Biogeogr., 23, 1408–1416, <https://doi.org/10.1111/geb.12215>, 2014.

1180 Soussana, J.-F. and Lemaire, G.: Coupling carbon and nitrogen cycles for environmentally sustainable intensification of
 1181 grasslands and crop-livestock systems, *Agric. Ecosyst. Environ.*, 190, 9–17,
 1182 <https://doi.org/10.1016/j.agee.2013.10.012>, 2014.

1183 Spohn, M., Pötsch, E. M., Eichorst, S. A., Woebken, D., Wanek, W., and Richter, A.: Soil microbial carbon use efficiency
 1184 and biomass turnover in a long-term fertilization experiment in a temperate grassland, *Soil Biol. Biochem.*, 97,
 1185 168–175, <https://doi.org/10.1016/j.soilbio.2016.03.008>, 2016.

1186 Stanley, P., Roche, L., and Bowles, T.: Amping up soil carbon: soil carbon stocks in California rangelands under adaptive
 1187 multi-paddock and conventional grazing management, *Int. J. Agric. Sustain.*, 2025.

1188 Stanley, P. L., Wilson, C., Patterson, E., Machmuller, M. B., and Cotrufo, M. F.: Ruminating on soil carbon: Applying
 1189 current understanding to inform grazing management, *Glob. Change Biol.*, 30, e17223,
 1190 <https://doi.org/10.1111/gcb.17223>, 2024.

1191 Steinbeiss, S., BEßLER, H., Engels, C., Temperton, V. M., Buchmann, N., Roscher, C., Kreuziger, Y., Baade, J.,
 1192 Habekost, M., and Gleixner, G.: Plant diversity positively affects short-term soil carbon storage in experimental
 1193 grasslands, *Glob. Change Biol.*, 14, 2937–2949, <https://doi.org/10.1111/j.1365-2486.2008.01697.x>, 2008.

1194 Strickland, M. S. and Rousk, J.: Considering fungal:bacterial dominance in soils – Methods, controls, and ecosystem
 1195 implications, *Soil Biol. Biochem.*, 42, 1385–1395, <https://doi.org/10.1016/j.soilbio.2010.05.007>, 2010.

1196 Tang, K., Wu, C., Wang, S., Liao, W., Yin, L., Zhou, W., and Cui, H.-J.: Distribution characteristics of soil organic carbon
 1197 fractions in paddy profiles with 40 years of fertilization under two groundwater levels, *J. Soils Sediments*, 24,
 1198 681–691, <https://doi.org/10.1007/s11368-023-03664-y>, 2024.

1199 Tang, Z., Feng, J., Chen, L., Chen, Z., Shao, X., and Xia, T.: Coupling amendment of microbial and compound fertilizers
 1200 increases fungal necromass carbon and soil organic carbon by regulating microbial activity in flue-cured tobacco-
 1201 planted field, *Eur. J. Soil Biol.*, 117, 103518, <https://doi.org/10.1016/j.ejsobi.2023.103518>, 2023.

1202 Tao, F., Huang, Y., Hungate, B. A., Manzoni, S., Frey, S. D., Schmidt, M. W. I., Reichstein, M., Carvalhais, N., Ciais, P.,
 1203 Jiang, L., Lehmann, J., Wang, Y.-P., Houlton, B. Z., Ahrens, B., Mishra, U., Hugelius, G., Hocking, T. D., Lu,
 1204 X., Shi, Z., Viatkin, K., Vargas, R., Yigini, Y., Omuto, C., Malik, A. A., Peralta, G., Cuevas-Corona, R., Di Paolo,
 1205 L. E., Luotto, I., Liao, C., Liang, Y.-S., Saynes, V. S., Huang, X., and Luo, Y.: Microbial carbon use efficiency
 1206 promotes global soil carbon storage, *Nature*, 618, 981–985, <https://doi.org/10.1038/s41586-023-06042-3>, 2023.

1207 Tate, K. W., Atwill, E. R., McDougald, N. K., and George, M. R.: Spatial and Temporal Patterns of Cattle Feces Deposition
 1208 on Rangeland, *J. Range Manag.*, 56, 432–438, <https://doi.org/10.2307/4003833>, 2003.

1209 Teague, R., Provenza, F., Kreuter, U., Steffens, T., and Barnes, M.: Multi-paddock grazing on rangelands: Why the
 1210 perceptual dichotomy between research results and rancher experience?, *J. Environ. Manage.*, 128, 699–717,
 1211 <https://doi.org/10.1016/j.jenvman.2013.05.064>, 2013.

1212 Teague, W. R., Dowhower, S. L., and Waggoner, J. A.: Drought and grazing patch dynamics under different grazing
 1213 management, *J. Arid Environ.*, 58, 97–117, [https://doi.org/10.1016/S0140-1963\(03\)00122-8](https://doi.org/10.1016/S0140-1963(03)00122-8), 2004.

1214 Teague, W. R., Dowhower, S. L., Baker, S. A., Haile, N., DeLaune, P. B., and Conover, D. M.: Grazing management
 1215 impacts on vegetation, soil biota and soil chemical, physical and hydrological properties in tall grass prairie,
 1216 *Agric. Ecosyst. Environ.*, 141, 310–322, <https://doi.org/10.1016/j.agee.2011.03.009>, 2011.

1217 Teixeira, R. F. M., Proença, V., Crespo, D., Valada, T., and Domingos, T.: A conceptual framework for the analysis of
 1218 engineered biodiverse pastures, *Ecol. Eng.*, 77, 85–97, <https://doi.org/10.1016/j.ecoleng.2015.01.002>, 2015.

1219 Thomson, F. J., Moles, A. T., Auld, T. D., and Kingsford, R. T.: Seed dispersal distance is more strongly correlated with
 1220 plant height than with seed mass, *J. Ecol.*, 99, 1299–1307, <https://doi.org/10.1111/j.1365-2745.2011.01867.x>,
 1221 2011.

1222 Tiessen, H., Cuevas, E., and Chacon, P.: The role of soil organic matter in sustaining soil fertility, *Nature*, 371, 783–785,
 1223 <https://doi.org/10.1038/371783a0>, 1994.

1224 Uribe, C., Inclán, R., Hernando, L., Román, M., Clavero, M. A., Roig, S., and Van Miegroet, H.: Grazing, tilling and
 1225 canopy effects on carbon dioxide fluxes in a Spanish dehesa, *Agrofor. Syst.*, 89, 305–318,
 1226 <https://doi.org/10.1007/s10457-014-9767-5>, 2015.

- 1227 Van Soest, P. J., Robertson, J. B., and Lewis, B. A.: Methods for Dietary Fiber, Neutral Detergent Fiber, and Nonstarch
1228 Polysaccharides in Relation to Animal Nutrition, *J. Dairy Sci.*, 74, 3583–3597,
1229 [https://doi.org/10.3168/jds.S0022-0302\(91\)78551-2](https://doi.org/10.3168/jds.S0022-0302(91)78551-2), 1991.
- 1230 Villarino, S. H., Pinto, P., Jackson, R. B., and Piñeiro, G.: Plant rhizodeposition: A key factor for soil organic matter
1231 formation in stable fractions, *Sci. Adv.*, 7, eabd3176, <https://doi.org/10.1126/sciadv.abd3176>, 2021.
- 1232 di Virgilio, A., Lambertucci, S. A., and Morales, J. M.: Sustainable grazing management in rangelands: Over a century
1233 searching for a silver bullet, *Agric. Ecosyst. Environ.*, 283, 106561, <https://doi.org/10.1016/j.agee.2019.05.020>,
1234 2019.
- 1235 Waldrop, M. P. and Firestone, M. K.: Microbial community utilization of recalcitrant and simple carbon compounds:
1236 impact of oak-woodland plant communities, *Oecologia*, 138, 275–284, [https://doi.org/10.1007/s00442-003-](https://doi.org/10.1007/s00442-003-1419-9)
1237 1419-9, 2004.
- 1238 Wang, J., Liu, L., Wang, X., and Chen, Y.: The interaction between abiotic photodegradation and microbial decomposition
1239 under ultraviolet radiation, *Glob. Change Biol.*, 21, 2095–2104, <https://doi.org/10.1111/gcb.12812>, 2015.
- 1240 Wang, J., Liu, L., Wang, X., Yang, S., Zhang, B., Li, P., Qiao, C., Deng, M., and Liu, W.: High night-time humidity and
1241 dissolved organic carbon content support rapid decomposition of standing litter in a semi-arid landscape, *Funct.*
1242 *Ecol.*, 31, 1659–1668, <https://doi.org/10.1111/1365-2435.12854>, 2017.
- 1243 Wang, K., Ma, Z., Qin, W., Li, X., Shi, H., Hasi, B., and Liu, X.: Soil nutrients and pH modulate carbon dynamics in
1244 particulate and mineral-associated organic matter during restoration of a Tibetan alpine grassland, *Ecol. Eng.*,
1245 212, 107522, <https://doi.org/10.1016/j.ecoleng.2025.107522>, 2025.
- 1246 Wang, X., McConkey, B. G., VandenBygaart, A. J., Fan, J., Iwaasa, A., and Schellenberg, M.: Grazing improves C and N
1247 cycling in the Northern Great Plains: a meta-analysis, *Sci. Rep.*, 6, 33190, <https://doi.org/10.1038/srep33190>,
1248 2016.
- 1249 Ward, S. E., Smart, S. M., Quirk, H., Tallowin, J. R. B., Mortimer, S. R., Shiel, R. S., Wilby, A., and Bardgett, R. D.:
1250 Legacy effects of grassland management on soil carbon to depth, *Glob. Change Biol.*, 22, 2929–2938,
1251 <https://doi.org/10.1111/gcb.13246>, 2016.
- 1252 Weigelt, A., Mommer, L., Andrzejak, K., Iversen, C. M., Bergmann, J., Bruehlheide, H., Fan, Y., Freschet, G. T., Guerrero-
1253 Ramirez, N. R., Kattge, J., Kuyper, T. W., Laughlin, D. C., Meier, I. C., van der Plas, F., Poorter, H., Roumet, C.,
1254 van Ruijven, J., Sabatini, F. M., Semchenko, M., Sweeney, C. J., Valverde-Barrantes, O. J., York, L. M., and
1255 McCormack, M. L.: An integrated framework of plant form and function: the belowground perspective, *New*
1256 *Phytol.*, 232, 42–59, <https://doi.org/10.1111/nph.17590>, 2021.
- 1257 Westerband, A. C., Funk, J. L., and Barton, K. E.: Intraspecific trait variation in plants: a renewed focus on its role in
1258 ecological processes, *Ann. Bot.*, 127, 397–410, <https://doi.org/10.1093/aob/mcab011>, 2021.
- 1259 White, R.: Pilot analysis of global ecosystems: Grassland ecosystems, 2000.
- 1260 Willers, C., Jansen van Rensburg, P. J., and Claassens, S.: Phospholipid fatty acid profiling of microbial communities—a
1261 review of interpretations and recent applications, *J. Appl. Microbiol.*, 119, 1207–1218,
1262 <https://doi.org/10.1111/jam.12902>, 2015.
- 1263 Wilson, C. H., Strickland, M. S., Hutchings, J. A., Bianchi, T. S., and Flory, S. L.: Grazing enhances belowground carbon
1264 allocation, microbial biomass, and soil carbon in a subtropical grassland, *Glob. Change Biol.*, 24, 2997–3009,
1265 <https://doi.org/10.1111/gcb.14070>, 2018.
- 1266 Wilson, P. J., Thompson, K., and Hodgson, J. G.: Specific leaf area and leaf dry matter content as alternative predictors
1267 of plant strategies, *New Phytol.*, 143, 155–162, <https://doi.org/10.1046/j.1469-8137.1999.00427.x>, 1999.
- 1268 Wright, I. J., Reich, P. B., Westoby, M., Ackerly, D. D., Baruch, Z., Bongers, F., Cavender-Bares, J., Chapin, T.,
1269 Cornelissen, J. H. C., Diemer, M., Flexas, J., Garnier, E., Groom, P. K., Gulias, J., Hikosaka, K., Lamont, B. B.,
1270 Lee, T., Lee, W., Lusk, C., Midgley, J. J., Navas, M.-L., Niinemets, Ü., Oleksyn, J., Osada, N., Poorter, H., Poot,
1271 P., Prior, L., Pyankov, V. I., Roumet, C., Thomas, S. C., Tjoelker, M. G., Veneklaas, E. J., and Villar, R.: The
1272 worldwide leaf economics spectrum, *Nature*, 428, 821–827, <https://doi.org/10.1038/nature02403>, 2004.
- 1273 Xu, H., Zhu, B., Wei, X., Yu, M., and Cheng, X.: Root functional traits mediate rhizosphere soil carbon stability in a
1274 subtropical forest, *Soil Biol. Biochem.*, 162, 108431, <https://doi.org/10.1016/j.soilbio.2021.108431>, 2021.

1275 Xu, X., Thornton, P. E., and Post, W. M.: A global analysis of soil microbial biomass carbon, nitrogen and phosphorus in
1276 terrestrial ecosystems, *Glob. Ecol. Biogeogr.*, 22, 737–749, <https://doi.org/10.1111/geb.12029>, 2013.

1277 Yao, H., He, Z., Wilson, M. J., and Campbell, C. D.: Microbial Biomass and Community Structure in a Sequence of Soils
1278 with Increasing Fertility and Changing Land Use, *Microb. Ecol.*, 40, 223–237,
1279 <https://doi.org/10.1007/s002480000053>, 2000.

1280 Yu, L., Sun, W., and Huang, Y.: Grazing exclusion enhances plant and topsoil carbon stocks in arid and semiarid
1281 grasslands, *Agric. Ecosyst. Environ.*, 320, 107605, <https://doi.org/10.1016/j.agee.2021.107605>, 2021.

1282 Yu, W., Huang, W., Weintraub-Leff, S. R., and Hall, S. J.: Where and why do particulate organic matter (POM) and
1283 mineral-associated organic matter (MAOM) differ among diverse soils?, *Soil Biol. Biochem.*, 172, 108756,
1284 <https://doi.org/10.1016/j.soilbio.2022.108756>, 2022.

1285 Zhang, Q., Buyantuev, A., Li, F. Y., Jiang, L., Niu, J., Ding, Y., Kang, S., and Ma, W.: Functional dominance rather than
1286 taxonomic diversity and functional diversity mainly affects community aboveground biomass in the Inner
1287 Mongolia grassland, *Ecol. Evol.*, 7, 1605–1615, <https://doi.org/10.1002/ece3.2778>, 2017.

1288 Zhang, Y., Lavallee, J. M., Robertson, A. D., Even, R., Ogle, S. M., Paustian, K., and Cotrufo, M. F.: Simulating
1289 measurable ecosystem carbon and nitrogen dynamics with the mechanistically defined MEMS 2.0 model,
1290 *Biogeosciences*, 18, 3147–3171, <https://doi.org/10.5194/bg-18-3147-2021>, 2021.

1291 Zhang, Y., Wang, T., Yan, C., Li, Y., Mo, F., and Han, J.: Microbial life-history strategies and particulate organic carbon
1292 mediate formation of microbial necromass carbon and stabilization in response to biochar addition, *Sci. Total*
1293 *Environ.*, 950, 175041, <https://doi.org/10.1016/j.scitotenv.2024.175041>, 2024.

1294 Zhong, W., Gu, T., Wang, W., Zhang, B., Lin, X., Huang, Q., and Shen, W.: The effects of mineral fertilizer and organic
1295 manure on soil microbial community and diversity, *Plant Soil*, 326, 511–522, [https://doi.org/10.1007/s11104-](https://doi.org/10.1007/s11104-009-9988-y)
1296 [009-9988-y](https://doi.org/10.1007/s11104-009-9988-y), 2010.

1297 Zhou, G., Zhou, X., He, Y., Shao, J., Hu, Z., Liu, R., Zhou, H., and Hosseinibai, S.: Grazing intensity significantly affects
1298 belowground carbon and nitrogen cycling in grassland ecosystems: a meta-analysis, *Glob. Change Biol.*, 23,
1299 1167–1179, <https://doi.org/10.1111/gcb.13431>, 2017.

1300 Zhou, G., Luo, Q., Chen, Y., He, M., Zhou, L., Frank, D., He, Y., Fu, Y., Zhang, B., and Zhou, X.: Effects of livestock
1301 grazing on grassland carbon storage and release override impacts associated with global climate change, *Glob.*
1302 *Change Biol.*, 25, 1119–1132, <https://doi.org/10.1111/gcb.14533>, 2019.

1303 Zhou, Z., Ren, C., Wang, C., Delgado-Baquerizo, M., Luo, Y., Luo, Z., Du, Z., Zhu, B., Yang, Y., Jiao, S., Zhao, F., Cai,
1304 A., Yang, G., and Wei, G.: Global turnover of soil mineral-associated and particulate organic carbon, *Nat.*
1305 *Commun.*, 15, 5329, <https://doi.org/10.1038/s41467-024-49743-7>, 2024.

1306 Zuur, A. F., Ieno, E. N., Walker, N., Saveliev, A. A., and Smith, G. M.: Mixed effects models and extensions in ecology
1307 with R, Springer, New York, NY, <https://doi.org/10.1007/978-0-387-87458-6>, 2009.

1308