

Improved management increases soil mineral-protected organic carbon storage via plant-microbial-nutrient mediation in semi-arid grasslands

Alejandro Carrascosa^a, Gerardo Moreno^a, M. Francesca Cotrufo^b, Cristina Frade^c, Sara Rodrigo^a, Víctor Rolo^a

^a Forest Research Group, INDEHESA, University of Extremadura, 10600, Plasencia, Cáceres, Spain

^b Department of Soil and Crop Sciences, Colorado State University, Fort Collins, CO, USA

^c Institute of Natural Resources and Agrobiology of Salamanca (IRNASA-CSIC), Salamanca, Spain

Correspondence to: Víctor Rolo (rolo@unex.es)

Abstract

Soil organic carbon (SOC) storage in semi-arid grasslands is threatened by both climate change and land degradation, impacting food production and climate regulation. Improved management has been proposed to increase SOC stocks and overcome these challenges. However, the benefits of improved management practices in semi-arid regions are in question. Little is known about the effects of management on the functional components of SOC, particulate (POC) and mineral-associated organic carbon (MAOC), which are expected to respond differently, and about the pathways that mediate these responses, such as changes in vegetation and soil microbial communities.

This work analyses the effect of rotational grazing, legume sowing and grazing exclusion on topsoil (0-8 cm) SOC, POC and MAOC stocks in Mediterranean wooded grasslands compared to continuous conventional grazing. Changes in plant diversity and morpho-biochemical traits, soil fertility and microbial composition were also evaluated. A total of 188 plots were sampled in 9 farms across a wide environmental gradient.

More resource-acquisitive, nitrogen-rich and less lignified plant community, higher soil microbial biomass with lower Gram+/Gram- ratio, and higher soil fertility were associated with higher SOC storage, with similar impacts on POC and MAOC. Rotational grazing increased POC, MAOC and total SOC stocks by 15%, 15% and 14% respectively, compared to continuous grazing. This effect was mediated by an increase in soil fertility in the rotationally grazed paddocks. On the other hand, grazing exclusion reduced POC stocks by 12% compared to continuous grazing. This depletion was mainly due to a reduction in microbial biomass and litter quality of vegetation in non-grazed paddocks. Both POC and MAOC stocks were lower at the warmer sites.

We conclude that rotational grazing can enhance long-term SOC storage in semi-arid grasslands, thereby increasing their resilience and climate mitigation capacity, whereas abandoning grazing could lead to SOC losses.

Keywords

35 Soil organic matter fractions; Soil microbial community; Plant traits; Rotational grazing; Legume sowing; Grazing
36 exclusion; Functional diversity; Mediterranean grassland

37

38 1. Introduction

39 Grasslands cover 40.5% of the world's ice-free land area and store one third of the terrestrial carbon (C) stocks, mainly
40 (over 80%) in the form of soil organic matter (SOM) (White, 2000). This C pool in grassland soils surpasses the global
41 aboveground vegetation C pool (IPCC, 2023a), highlighting the importance of grasslands SOM in global climate
42 regulation (Bai and Cotrufo, 2022). In addition, SOM is a cornerstone of grassland productivity and functioning (Cotrufo
43 and Lavallee, 2022; Tiessen et al., 1994). However, C storage in grassland soils is threatened by increasing C outputs (i.e.,
44 soil respiration and erosion) and decreasing C inputs (i.e., primary productivity) due to anthropogenic impacts such as
45 land degradation and climate change (Crowther et al., 2016; Gang et al., 2014; Godde et al., 2020; Lei et al., 2021).
46 Improved grassland management offers a key opportunity to counter these threats by increasing soil organic C (SOC)
47 stocks and supporting both climate change mitigation and adaptation (Bai and Cotrufo, 2022; Conant, 2012; Dondini et
48 al., 2023; Stanley et al., 2024). This is particularly valuable in the case of semi-arid grasslands, which represent the
49 majority of the global grassland area (White, 2000) but are also more vulnerable to climate change impacts than wetter
50 grasslands (Smith et al., 2024).

51 Practices such as rotational grazing, legume sowing and grazing exclusion have been proposed to increase and conserve
52 SOC stocks (Conant et al., 2017; Yu et al., 2021). Rotational grazing encompasses a variety of practices where small
53 paddocks are grazed at high intensity for short periods of time, allowing for longer pasture rest than continuous grazing
54 (Teague et al., 2013). This management favors vegetation recovery after defoliation and reduces grazing patchiness and
55 livestock selectivity (Jacobo et al., 2006; Teague et al., 2013). The effects of rotational grazing on grassland productivity
56 and animal performance are under debate (Briske et al., 2008; Teague et al., 2013; di Virgilio et al., 2019) but most studies
57 have found positive effects on SOC stocks associated with this practice (Byrnes et al., 2018; Conant et al., 2017; Phukubye
58 et al., 2022; Teague et al., 2011). Sowing legumes in natural pastures is practiced worldwide, having clear positive impacts
59 on SOC stocks (Carranca et al., 2022; Conant et al., 2017; Moreno et al., 2021) and grassland productivity (Bartholomew
60 and Williams, 2010; Carrascosa et al., 2024; Jaurena et al., 2016; Khatiwada et al., 2020; Rama et al., 2022). Grazing
61 exclusion is widely advocated as a tool for ecosystem restoration (Cheng et al., 2016; Novelly and Watson, 2007) and is
62 also a global trend driven by land abandonment, particularly in high-income countries (Li and Li, 2017). The effects of
63 grazing exclusion on SOC in grasslands are mixed, showing both positive (Cheng et al., 2016; Yu et al., 2021) and negative
64 outcomes (Wilson et al., 2018). Moreover, the net effect of management practices has been shown to depend on the
65 environmental context (Maestre et al., 2022; McSherry and Ritchie, 2013; Niu et al., 2025), and global meta-analyses
66 remain inconclusive. While some global studies have reported greater benefits on SOC stocks from grazing exclusion and
67 rotational grazing in arid climates (Zhou et al., 2017), others found these management practices to be more beneficial in
68 wetter climates (Byrnes et al., 2018; Hawkins, 2017; Zhou et al., 2019). To the best of our knowledge, no similar studies
69 have addressed the interaction between environmental conditions and the effect of legume sowings on SOC stocks. Thus,
70 some questions remain open and further research is needed to clarify the net effects of improved management practices
71 on SOC storage under different environmental conditions and particularly in semi-arid grasslands.

72 To better understand the controls and vulnerability of C in soils, SOM can be conceptualized into particulate organic
73 matter (POM) and mineral-associated organic matter (MAOM) fractions (Cotrufo and Lavallee, 2022; Lavallee et al.,
74 2020). POM originates from fragmented structural plant inputs and, to a lesser extent, microbial recalcitrant compounds
75 (Angst et al., 2021; Six et al., 2001). In contrast, MAOM forms through the sorption of microbial necromass and soluble

76 plant inputs onto soil mineral surfaces (Angst et al., 2021; Cotrufo et al., 2022). The mineral bonds partially protect
 77 MAOM from decomposition (Baldock and Skjemstad, 2000), while POM is readily accessible to microbial degradation,
 78 although occlusion within soil aggregates can reduce its accessibility (Angst et al., 2017). Hence, POM accumulation is
 79 mainly controlled by environmental constraints on microbial activity, e.g. low temperatures and highly acidic soils
 80 (Hansen et al., 2024; Yu et al., 2022; Zhou et al., 2024). For the same reasons, POM is more vulnerable to climatic
 81 warming (Benbi et al., 2014; Georgiou et al., 2024; Rocci et al., 2021) and the mean residence time of C in POM (years
 82 to decades) is on average shorter than in MAOM (decades to centuries; Zhou et al., 2024). However, MAOM storage in
 83 soil is theoretically limited by the availability of free mineral surface area (i.e. clay and silt content; Six et al., 2002), and
 84 a saturation point can be observed, where no more MAOM accumulate despite increases in total SOM contents (Cotrufo
 85 et al., 2019a; Georgiou et al., 2022). Management changes have been shown to influence SOM fractions in grassland soils
 86 (Khatri-Chhetri et al., 2024; Mosier et al., 2021; Oliveira Filho et al., 2019), and a conceptual framework for relating
 87 grazing management to SOM distribution has recently been proposed (Stanley et al., 2024). Yet, the underlying processes
 88 mediating these effects, such as alterations in vegetation or soil microbial communities (Laliberté and Tylianakis, 2012;
 89 Peco et al., 2017; Wilson et al., 2018), remain poorly understood.

90 Primary productivity is the point of entry of C into soil, and consequently the amount of plant inputs regulates SOM
 91 accrual (King et al., 2023; Zhou et al., 2024), but microbial processing largely determines the fate of that C (Cotrufo and
 92 Lavallee, 2022; Crowther et al., 2019). In this sense, the chemical composition of plant inputs, and the soil microbiota
 93 carbon use efficiency (CUE), i.e. the amount of C used for microbial growth and products relative to total C uptake, play
 94 a crucial role in the SOM formation process (Cotrufo and Lavallee, 2022; Tao et al., 2023). Recalcitrant plant inputs [i.e.,
 95 high carbon-nitrogen ratio (C/N) and lignin content] tend to promote short-term SOM accumulation, primarily as POM,
 96 due to their chemical resistance to decomposition (Cheng et al., 2023; Cotrufo and Lavallee, 2022). However, as outlined
 97 in the Microbial Efficiency-Matrix Stabilization (MEMS) framework (Cotrufo et al., 2013), recalcitrant inputs are less
 98 efficiently decomposed by microbes, leading to greater C losses in the long term, compared to labile (i.e., water-soluble,
 99 low C/N and lignin content) plant inputs (Cotrufo and Lavallee, 2022; Ridgeway et al., 2022). Thus, labile plant inputs
 100 are expected to enhance MAOM formation and SOM stocks in the long term, due to their faster and more efficient
 101 decomposition (Cheng et al., 2023; Haddix et al., 2016). Elias et al. (2024) added complexity to these assumptions,
 102 showing that plant input characteristics may favor certain microbial groups over others, altering the overall CUE of the
 103 microbial community. For example, fungi, which are often assumed to have a higher CUE than bacteria (Kallenbach et
 104 al., 2016; Strickland and Rousk, 2010), are favored by the addition of recalcitrant inputs (Bai et al., 2024; Strickland and
 105 Rousk, 2010). Substrate preferences have been also identify for Gram-positive (Gram+) and Gram-negative (Gram-)
 106 bacteria (Fanin et al., 2019; Kramer and Gleixner, 2008), with consequences for SOC accrual (Klump et al., 2009).
 107 Importantly, much of the research on the influence of plant input characteristics and microbial communities on SOM
 108 formation dynamics has relied on incubation experiments (Cheng et al., 2023; Haddix et al., 2016; Ridgeway et al., 2022)
 109 and there is limited information on how these findings translate to natural field conditions. Other vegetation characteristics
 110 such as species richness have been shown to positively influence SOC stocks (Lange et al., 2015; Steinbeiss et al., 2008),
 111 but its effects on SOC fractions have been poorly evaluated, with inconclusive results in grasslands (Mortensen et al.,
 112 2025). In addition, the relationships between SOM stocks and fractions and plant functional traits have rarely been studied
 113 (Manning et al., 2015; Mortensen et al., 2025; Xu et al., 2021) despite the latter being widely used to predict ecosystem
 114 functioning and responses (Funk et al., 2017). Plant functional traits are highly correlated with processes such as litter
 115 decomposition (Cornwell et al., 2008; Fortunel et al., 2009; Kazakou et al., 2009) or root exudates production (Guyonnet
 116 et al., 2018) and may therefore be a promising tool to study the relationships between vegetation, soil microbiota and
 117 SOM formation dynamics (Faucon et al., 2017).

118 The aim of this work is to evaluate the impact of rotational grazing, legume sowing, and grazing exclusion on topsoil (0–
119 8 cm) SOC stocks, and their distribution between POM and MAOM fractions, in semi-arid grasslands, compared to
120 conventional continuous grazing. We also evaluate changes in vegetation characteristics (identity and diversity of
121 biochemical and morphological traits), soil nutrients, and soil microbial communities as possible pathways through which
122 management might indirectly affect SOC stocks and fractions. This study focuses on the Iberian *dehesas* Mediterranean
123 woody grassland, the main example of semi-arid grasslands in Europe (Porqueddu et al., 2016). The most widespread
124 livestock management in this ecosystem is continuous grazing, but in recent decades rotational grazing and legume sowing
125 have gained importance (Frangia et al., 2023; Pulina et al., 2023). At the same time, the amount of ungrazed pastures has
126 increased due to land abandonment (Palomo-Campesino et al., 2018). Iberian *dehesas* occupy 3.1 million hectares,
127 spanning a wide environmental gradient, and have been subject to extensive grazing for centuries (Moreno and Pulido,
128 2009), making them an ideal model system for assessing the effects of improved management on SOC stocks in semi-
129 arid grasslands. In particular, we designed our study to answer the following questions: 1) What are the effects of the
130 different management practices on bulk SOC and fractions stocks? 2) Are these effects mediated by changes in vegetation
131 or soil microbial communities? 3) Is SOC storage in these grasslands, and its enhancement, modulated or limited by
132 environmental factors such as climate or soil properties? And 4) are these mechanisms and controls the same or different
133 for C in POM or MAOM? Understanding the potential and limitations of improved management on SOC storage in POM
134 and MAOM in semi-arid grasslands can guide policymakers in enhancing the climate change adaptation and mitigation
135 capacity of these ecosystems, while supporting productivity and soil fertility.

136

137 2. Methods

138 2.1. Study area and experimental design

139 The study was carried out at nine commercial *dehesa* farms located along a north-south gradient in the western part of
140 the Iberian Peninsula (Fig. 1a). The region has a continental Mediterranean climate, but on a local scale, in relative terms,
141 farms can be grouped into three main climatic regions (Fig. 1b, c) according to the average climate for the period 1980–
142 2018 (García Bravo et al., 2023). A cold-dry region [12.9 °C mean annual temperature (MAT) and 445 mm mean annual
143 precipitation (MAP)] in the north; a warm-wet region (17.3 °C MAT and 603 mm MAP) in the middle of the latitudinal
144 gradient; and a warm-dry region (17.0 °C MAT and 510 mm MAP) in the south. The soils of the farms share a common
145 development from granites, shales and sandy tertiary sediments, are acidic and poor in organic matter, but cover a wide
146 texture gradient (Fig. 1d). In these farms, native pastures are often combined with scattered trees, as is common in
147 Mediterranean and semi-arid rangelands (den Herder et al., 2017; Soliveres et al., 2014). The tree layer is dominated by
148 holm oaks (*Quercus ilex* L.) with scattered cork oaks (*Quercus suber* L.) or gall oaks (*Quercus faginea* Lam.). The
149 herbaceous layer is composed of species typical of Mediterranean pastures and presents a high diversity and proportion
150 of annual C3 plants (Table S1). The growing season of the pasture is very limited by the Mediterranean summer drought
151 with the annual species germinating in autumn (early-to mid-October), reaching their peak productivity in mid-spring
152 (late April of the following year) and senescing in June.

153 *Dehesa* farms are typically managed by extensive continuous grazing, where livestock (mainly cattle) graze freely on
154 large areas following a loosely defined grazing plan. This management has been traditionally practiced in all the farms
155 studied, until, in recent decades, some paddocks were converted to other management practices. As a result, on each farm
156 we selected five paddocks, each with one of the following management regimes:

157 - Abandoned (Ab): Paddocks devoid of grazing for at least the last 10-20 years.

158 - Continuous grazing (Ct): Paddocks where livestock stand most of the year, fed by grazing and supplemented. This is the
159 control treatment in our study as it is the most widespread management in Iberian *dehesas*.

160 - Rotational grazing (Ro): Paddocks intensively grazed in short periods and with resting periods (i.e. without livestock)
161 lasting for more than six months. Rotational grazing has been applied in these paddocks for the last 10-15 years.

162 - Recent legume sowing (Lr): Paddocks where pastures have recently (≤ 5 years) been sown with legume mixtures. In
163 *dehesa* farms, legume sowing consists of sowing a mixture of seeds (at a rate of 20 kg ha⁻¹) from various species of annual
164 legumes (pre-inoculated with Rhizobium) such as *Trifolium subterraneum* L., *T. incarnatum* L., *T. michelianum* Savi., *T.*
165 *resupinatum* L., *T. mutabile* Port. and *Ornithopus sativus* Brot., along with some highly productive annual grass species
166 such as *Lolium multiflorum* Lam. and *Lolium rigidum* Gaud., all of which have relatively shallow root systems (Teixeira
167 et al., 2015). These sowings are preceded by surface tillage and phosphorus application to meet the needs of the legumes
168 and stimulate N-fixation (Jongen et al., 2019). Farmers sow legumes mixture only once, because its effect persists over
169 the years thanks to the natural seeds production and the resulting soil seed bank of the sown species. None of the sown
170 plots studied was resown.

171 - Old legume sowing (Lo): Paddocks sown with legume mixtures more than 10 years ago.

172 The distinction between recent and old legumes sown paddocks allows us to compare short and long-term effects after
173 sowing. In two out of the nine farms, we added an additional control (continuous grazing) paddock close to the legume
174 sowing paddocks because they were at a considerable distance from the other control paddock, limiting their
175 comparability. Additional information on paddocks characteristics is provided in Table S2.

176 In each paddock, we installed four permanent monitoring plots, in open grassland and away from the direct influence of
177 the trees (at least 10 m away from the trees). This setup allowed us to have a total of 188 sampling plots where we
178 measured soil properties, SOC fractions, and soil microbial communities in 2021 and vegetation traits for 3 years (2021
179 to 2023).

180




Figure 1. Iberian “dehesa” cover according to the CORINE 2018 land cover survey and geographical location of the studied farms (a). Mean annual temperature (b) and precipitation (c) during the period 1980-2018 in the study area. Soil texture in all sampled plots (d).

2.3. Vegetation sampling

We assessed above-ground (ABG) pasture productivity over three growing periods at all plots using 50 x 50 x 50 cm grazing exclusion cages. In the 2020-2021 period (hereafter referred to as 2021) we installed one cage on each monitoring plot at the beginning of spring (late March 2021). In the 2021-2022 and 2022-2023 periods (hereafter referred to as 2022 and 2023), we installed the cages at the beginning of the green-up in early November. Standing biomass at the time of cage establishment (t_0) was determined by hand clipping within two 25 x 25 cm squares, randomly placed in the proximity of the cage. In the three years we harvested ABG biomass inside the cages at the beginning of summer in early June (t_1), when most of the vegetation was already senescent. Biomass collected both in t_0 and t_1 was oven-dried at 60°C for 48h and weighed. We calculated ABG productivity as the difference between t_0 and t_1 biomass. Therefore, only spring productivity was measured in 2021, which represents the largest proportion of annual productivity in this system, while a closer estimate of annual productivity was measured in 2022 and 2023.

Plant traits were collected in early May 2021, at the peak of the growing season. We considered a circular area of three meters in diameter around each exclusion cage and within this area, we identified the species present at 25 regularly spaced points using the line intercept method (Godínez-Alvarez et al., 2009), with three concentric circular transects going around the cage. After completing the species inventory for all the sampling plots in a paddock, we collected at least three to ten full individuals, in that same paddock, of each of the identified species, as proposed in the abundance-weighted trait sampling scheme (Carmona et al., 2015). Thus, at least ten individuals of the most abundant species were collected in the same paddock, five individuals for the medium abundant species and three individuals for the rare species. The leaf area (LA), specific leaf area (SLA), specific root length (SLR), leaf dry matter content (LDMC), root dry matter content (RDMC), leaf nitrogen content (LNC) and plant maximum height (see Fig. 2 for more detail) of all the collected

206 individuals was measured following the standard protocols (Pérez-Harguindeguy et al., 2016). In total, more than 10000
207 individuals were measured across all the plots. This extensive sampling allowed us to account not only for interspecific
208 differences in trait values, but also for intraspecific variability in trait values across managements, farms and regions.
209 Intraspecific variability has proven to be very important for accurately defining the functional composition of plant
210 communities and their relationship to ecosystem processes (Siefert et al., 2015; Westerland et al., 2021). We also
211 conducted species inventories in the spring of 2022 and 2023, but on these later samplings we only collected and measured
212 the traits of individuals from the species not found in 2021. We used the trait values measured in 2021 for each species in
213 each paddock to impute trait values in 2022 and 2023. The proportion of legumes in each plot was quantified from the
214 species inventories as a measure of the number of plants with N fixation capacity in the communities.

215 Pasture chemical composition was assessed in the three years, after the species inventories and traits samplings. We
216 collected standing biomass by hand-clipping in three 25 x 25 cm quadrats randomly placed in the monitoring plot, outside
217 the exclusion cages. These samples were dried at 60°C for 48h, then grounded and analyzed with Dumas method in
218 DUMATHERM® N Pro analyzer (C. Gerhardt GmbH & Co. Germany) to obtain the nitrogen content of each sample.
219 Acid detergent lignin (ADL) and fiber (ADF) and neutral detergent fiber (NDF) content were also measured. NDF, ADF
220 and ADL were determined using a fiber analyzer (ANKOM A2000, ANKOM Technology, USA), following the official
221 procedures (Latimer, 2023).


Trait	Description	
Height 	Height to top photosynthetic tissue (cm). - Positively related to competitive ability for light and seed dispersal ^{1,2}	Traits associated with the plant size spectrum ³
Leaf area (LA) 	One-sided area of a fresh leaf (mm²) - Positively related to light interception . It also has implications for the water and energy balance of the leaf ³	
Specific leaf area (SLA) 	Leaf area/ Leaf dry mass (mm² mg⁻¹) - Positively related to plant relative growth rate and litter decomposability . Negatively related to stress tolerance ⁴	
Leaf dry matter content (LDMC) 	Leaf dry mass/ Leaf fresh mass (mg g⁻¹) - Negatively related to relative growth rate and litter decomposability . Positively related to stress tolerance ^{5,6}	Traits related with the Leaf/Root Economic Spectrum (conservative – acquisitive resources strategy gradient) ^{8,11}
Leaf nitrogen content (LNC) 	Nitrogen content in dry leaf mass (mg g⁻¹) - Positively related to photosynthetic rate , leaf respiration and litter decomposability ^{7,8}	
Root dry matter content (RDMC) 	Roots dry mass/ Roots fresh mass (m g⁻¹) - Correlated positively with root tissue density . Negatively correlated with root respiration and exudate production ^{9,10}	
Specific root length (SRL) 	Roots length / Roots dry mass (mm mg⁻¹) -Negatively correlated with root diameter and mycorrhizal association . Positively correlated with soil exploration and nutrient acquisition ^{11,12}	Traits related with the fungal collaboration gradient ¹¹

Figure 2. Graphical representation and description of plant functional traits measured. References embedded in the figure are provided below: ¹Gaudet and Keddy (1988), ²Thomson et al. (2011), ³Díaz et al. (2016), ⁴Poorter et al. (2009), ⁵Kazakou et al. (2009), ⁶Wilson et al. (1999), ⁷Freschet et al. (2012), ⁸Wright et al. (2004), ⁹Guyonnet et al. (2018), ¹⁰Roumet et al. (2016), ¹¹Bergmann et al. (2020), ¹²Kramer-Walter et al. (2016).

2.4. Soil sampling and soil characteristics measurement

In spring 2021, four soil cores were collected with a push sampler (5 cm diameter) at a depth of 8 cm around each exclusion cage, after removal of above-ground vegetation and litter from the sampled surfaces. The four soil cores were combined to obtain a composite sample from each plot, and an aliquot of 40 g of this sample were sieved (<2 mm),

reserved and stored at 4°C for microbial community analysis in the following days. The remainder of the composite soil samples were air-dried and sieved (<2 mm). Coarse mineral material greater than 2 mm (hereinafter referred to as gravel) was weighed and used for bulk density correction (Eq.1).

Soil texture (sand, clay and silt content) was determined using a laser diffraction particle size analyzer (Mastersizer 2000, Malvern Instruments Ltd. UK) after dispersion with sodium hexametaphosphate. Soil pH was measured with a glass electrode (soil:water 1:2.5) pH meter (CRISON Basic20, Alella, Spain). Soils were extracted with 1M KCL and determined colorimetrically in a Bran+Luebbe Autoanalyzer 3 (Norderstedt, Germany), following the manufacturer's protocol, to measure nitrate (NO₃⁻) and ammonium (NH₄⁺) concentrations. Available P (Olsen P) was determined colorimetrically in an Agilent Cary 60 UV-Vis spectrophotometer (Agilent Technologies, USA), after extraction with 0.5 M NaHCO₃ at pH 8.5. Available K, Ca and Mg were determined by ICP-OES (Varian Mod. 720, Palo Alto, California, USA), after extraction with 1M ammonium acetate (pH 7).


2.5. Soil organic matter fractionation

Representative subsamples of each 2mm sieved air-dry composite soil sample were used to measure both bulk SOC and total nitrogen (TN) as well as their distribution across distinct physical fractions. First, 10 g aliquots of all samples were ground in a ball mill and the Bernard's calcimeter method (Sherrod et al., 2002) was used to test for the presence of inorganic C. Only four samples (0.02 % of the database) contained traces of carbonates, with CaCO₃ contents between 0.2 % and 0.8 %. These samples were excluded from subsequent analyses. For bulk soil analyses, aliquots of 10g of soil were ground in a ball mill and then analyzed in the DUMATHERM® N Pro analyzer (C. Gerhardt GmbH & Co. Germany) to determine %C (SOC) and %N (TN). For soil fractionation, we followed a combined size and density procedure as described in Leuthold et al. (2024). Briefly, an aliquot of 6 g of soil was mixed with sodium polytungstate (1.85 g cm⁻³) and shaken reciprocally for 18 h to disperse the aggregates. After dispersion samples were centrifuged for density fractionation and the light particulate organic matter "POM" (<1.85 g cm⁻³) was aspirated from the rest of the soil. We then thoroughly rinsed the residual heavy fraction and separated it by wet sieving into coarse heavy mineral-associated organic matter "chaOM" (>53 µm) and fine mineral-associated organic matter "MAOM" (<53 µm). All fractions were analyzed for %C and %N on an elemental analyzer as described above. Fractionation was accepted when mass recovery was within ±5%, and C recovery was within ±15%. For samples that did not satisfy one of these conditions, fractionation was repeated.

Since the chaOM and the MAOM shared similar C/N ratios, which were lower than POM C/N ratios (Fig. 3a), we merged these two mineral associated OM fractions and present them together as MAOM (Santos et al., 2024; Zhang et al., 2021). Thus, in this work, POM and MAOM represented the light (<1.85 g cm⁻³) and heavy (>1.85 g cm⁻³) fractions respectively. This contrasts with other studies based on size fractionation, in which light POM and chaOM are pooled together (Cotrufo et al., 2019b; Dai et al., 2025; Díaz-Martínez et al., 2024). However, our decision is supported by recent findings of Leuthold et al. (2024) who observed greater chemical similarity between chaOM and MAOM than between chaOM and POM. The pooling of the two mineral associated OM fractions is not expected to modify the results or interpretation of this work, as the C content of chaOM only accounts for an average of 8% of the total SOC, and its relative importance is similar across all managements (Fig. S1). Carbon data are presented as SOC, POC and MAOC stocks (Mg ha⁻¹), calculated following Poeplau et al. (2017):

$$OC_{stock} = OC_{content} \times \frac{mass_{fine\ soil\ (<2mm)}}{Volume_{sample}} \times depth_{sample} \quad (1)$$

271 where $OC_{content}$ is the organic C content (as a proportion) of the soil fraction, $mass_{fine\ soil(<2mm)}$ is the dry weight of the soil
 272 excluding gravel and large roots, and considering the dry mass of the aliquot reserved for microbial community analysis.
 273 $Volume_{sample}$ and $depth_{sample}$ are respectively the volume and depth of the soil sampled with the core.
 274 Additionally, we calculated the proportion of MAOC in total SOC (MAOC/SOC) following Cotrufo et al. (2019).



275
 276 **Figure 3.** Violin plot and boxplot (with median and quartiles) and mean values (expressed with labels and red dots) for the ratio
 277 of carbon to nitrogen content (C/N ratio) in the soil organic matter fractions (a), and relation between carbon content in the
 278 soil organic matter fractions (in g of C per kg of soil) and the total soil organic carbon (SOC) content (b). “chaOM” refers to
 279 the coarse heavy mineral-associated organic matter, “IPOM” to the light particulate organic matter and “MAOM” to the fine
 280 mineral-associated organic matter. In panel “b”, a linear regression line depicts the relationship between IPOM and total SOC,
 281 while logistic curves illustrate the relationships of chaOM and MAOM with total SOC.

282

283 2.6. Soil microbial communities:

284 Soil microbial communities were characterized by phospholipid fatty acids analysis (PLFA; Willers et al., 2015). A 2 g
 285 fine soil aliquot of lyophilized soil was used for lipid extraction with a one-phase chloroform–methanol-phosphate buffer
 286 solvent. Phospholipids were separated from non-polar lipids and converted to fatty acid methyl esters (FAMES), which
 287 were then separated by gas chromatography as described in details in Moreno et al. (2021). The sum of all individual
 288 PLFAs was used as a proxy for microbial biomass (in nmol PLFAs g⁻¹ of soil). Further, we estimated microbial biomass
 289 stocks (in mol ha⁻¹) by substituting the $OC_{content}$ by the total PLFAs concentration in equation 1. Specific PLFAs were
 290 used as biomarkers to estimate the relative abundance of broad taxonomic microbial groups, according to their
 291 characteristic fatty acids: eukaryote, Gram- and Gram+ bacteria, saprophytic fungi and arbuscular mycorrhizal fungi
 292 (Frostegård and Bååth, 1996). The ratios among Gram+ and Gram- bacteria (Gram+/Gram-) and fungi and bacteria
 293 (Fungi/Bacteria) were also calculated to describe the composition of the microbial community.

294

295 2.7. Mineral-associated carbon capacity and saturation:

296 To evaluate the degree of MAOC saturation, we used the boundary line approach reported by Georgiou et al. (2022)
 297 adjusted with global soil samples as a reference of the maximum observed mineralogical capacity (sensu Georgiou et al.,

298 2025) of our samples (Fig. 4). Therefore, the saturation deficit in each sample was calculated as one minus the ratio of
 299 the current C content and the observed maximum C content according to the mineralogical capacity (% clay + silt).

300

301 2.6. Data analysis:

302 All analyses were carried out in R 4.3.3 (R Core Team, 2024). Species abundance data and species trait values were used
 303 to compute, for each year and plot, the community weighted means (CWM) of all traits as well as the RaoQ multi-trait
 304 functional diversity index (Ricotta and Moretti, 2011). In addition, we calculated the species richness as a measure of the
 305 taxonomic diversity in each plot. These analyses were performed using the package “FD” (Laliberté and Legendre, 2010).
 306 Regarding pasture chemical composition, the hemicellulose content was calculated as NDF minus ADF and the cellulose
 307 content as ADF minus ADL, with ADL being considered a good proxy for the lignin content (Van Soest et al., 1991).

308 For the vegetation analysis, measurements from the three years were combined to obtain mean estimates of each
 309 vegetation attribute. For each year, the ABG production and chemical attributes values, as well as the CWM of all the
 310 functional traits, were centered and scaled between -1 and 1, so that values closer to -1 represent plots with traits values
 311 lower than the mean, while values closer to 1 represent plots above the mean. The three years centered and scaled values
 312 were then averaged to obtain estimates of each vegetation characteristic across the study period, as shown in this equation
 313 2:

$$314 \text{Vegetation attribute}_{\text{mean}} = \frac{\sum_{year\ i=2021}^{2023} \text{Vegetation attribute (centered and scaled)}_{year\ i}}{3} \quad (2)$$

315 All mean values of vegetation characteristics obtained with equation 2 were used to build a Principal Component Analysis
 316 (PCA). From this PCA, we extracted 4 main axes of variation of the vegetation characteristics with eigenvalues greater
 317 than one (Fig .4).




Figure 4. Representation of the 4 main axis of variation in the principal component analysis (PCA) summarizing the vegetation characteristics variables. Panels a, b, c and d show the correlation between the different variables included in the PCA and the new axis. Panels e, f, g and h represent the plant communities characteristics at the extreme of each axis. Representative species of each axis are represented in Fig. S2.

Nutrients stocks in the soil samples were calculated by substituting the OC_{contents} in equation 1, for the concentration of each nutrient. PCA was used to summarize these nutrient stocks into a single composite index, as all nutrients were positively correlated. The first component of this PCA, which absorbed 46.9% of the variance, was extracted as a new variable, henceforth called “soil fertility” (Fig. S3). This soil fertility index increases as ammonium, nitrate, P, Ca, Mg, and K stocks in soil increase.

328 To analyze the direct and indirect effects of management, vegetation characteristics, microbial communities and
 329 environmental variables on POC, MAOC and SOC stocks and MAOC/SOC ratio we built a structural equation model
 330 (SEM) using the “piecewiseSEM” package (Lefcheck, 2016). These models assess the extent to which a defined structure
 331 of causal relationships between variables fits the actual correlations in the data. In this causal structure, direct effects
 332 describe the influence of one variable on another without intermediaries, whereas indirect effects occur when a variable
 333 affects another through its impact on a third variable. We started from the assumption that management, climate and soil
 334 properties could affect SOC, POC and MAOC stocks and the MAOC/SOC directly or indirectly, by modifying microbial
 335 composition and vegetation characteristics. This provided us with a theoretical basis for the construction of the SEMs.
 336 The linear sub-models within these SEMs were fitted with the “lme4” package (Bates et al., 2015) using the farms as
 337 random factors to avoid spatial pseudo-replication. Model fitting was performed according to Zuur et al. (2009) selecting
 338 the model with the lowest corrected Akaike Information Criterion (AICc) value after adjusting for random and fixed
 339 factors. All predictor variables used in the model selection are summarised in Table 1. While gravel content is correlated
 340 to BD, its inclusion as predictor in the models helped to control for inter-site variability in gravel content. Interactions
 341 between management and climate or soil texture were also included during model selection. All model assumptions were
 342 checked and satisfactorily met. Predictor variables were scaled and centered prior to inclusion in the models.

343 To calculate the total effect of each explanatory factor (i.e. the sum of direct and indirect effects) on SOC, POC and
 344 MAOC stocks, and MAOC/SOC ratio in the SEMs, we used the “semEff” package (Murphy, 2022). This package
 345 generates estimates and confidence intervals of the total effects of the explanatory factors across multiple permutations
 346 of the data (9999 bootstraps in our case).

347 Given that average bulk density values partly differed among managements (Figure S4), and that this variation may affect
 348 carbon stocks, the same analysis procedure used for POC, MAOC and SOC stocks was used to analyze the change in
 349 POC, MAOC and SOC contents.


350 **Table 1. List of the variables and their units, which were included as fixed factors during the structural equation model**
 351 **(SEM) sub-models selection.**

Variables	Units
Management	Categorical (5 levels)
Sand content	%
Gravel content	%
Soil fertility	unitless
Mean annual precipitation (MAP)	mm
Mean annual temperature (MAT)	°C
Vegetation PCA- Axis 1 (N-C axis)	unitless
Vegetation PCA- Axis 2 (PES axis)	unitless
Vegetation PCA- Axis 3 (P-L axis)	unitless
Vegetation PCA- Axis 4 (Size axis)	unitless
Functional diversity (Rao multi-trait index)	unitless
Taxonomic diversity (species richness)	count
Microbial biomass (bulk)	nmol ha ⁻¹
Gram+/Gram-	ratio
Fungi/Bacteria	ratio

352

353 **3. Results:**

354 The mean values of POC, MAOC and SOC stocks were 9.8 ± 4.3 , 7.1 ± 3.1 and 16.5 ± 6.3 Mg C ha⁻¹ respectively. On
355 average, MAOC represented 44% of the total SOC. The MAOC content of all the soils analyzed was well below the
356 theoretical saturation level established by Georgiou et al. (2022) in relation to the clay + silt content (Fig. 5). Mean MAOC
357 saturation deficit was 80%. Soil organic carbon fractions and stocks were affected directly and indirectly by mean plant
358 traits, microbial communities, soil properties and management, and indirectly by climatic conditions (Fig. 6). Plant
359 diversity indices and interactions between management and climate or soil texture were not retained in the model selection
360 due to lack of significance.



361

362 **Figure 5. Relation between the mineral-associated organic carbon (MAOC) and the percent Clay+Silt (CS) in the soil samples**
363 **and the boundary line (in blue) with confidence intervals adjusted by Georgiou et al. (2022) for high-activity mineral soils,**
364 **indicating the maximum observed mineralogical capacity for each CS content. The equation for the boundary line is provided**
365 **in the panel. Points are colored based on their MAOC deficit (1- MAOC content / MAOC maximum observed capacity).**

366


367 Total microbial biomass, estimated as total PLFA stocks, averaged 66.3 ± 25.9 mol ha⁻¹, with mean values for
368 Fungi/Bacteria and Gram+/Gram ratios of 0.08 ± 0.03 and 1.1 ± 0.2 , respectively, although there were some differences
369 between treatments (Fig. S5). Microbial biomass was positively correlated with POC, MAOC and SOC stocks (Fig. 6f;
370 Fig. 7a, b & c). The Fungi/Bacteria ratio was negatively correlated with MAOC stocks (Fig. 6f; Fig. 7b) meaning that
371 these stocks were higher in soils with a greater bacterial predominance. The Gram+/Gram- ratio had a significant direct
372 negative effect on POC, MAOC and SOC stocks (Fig. 6f; Fig. 7a, b & c).

373 The first axis of the PCA of the vegetation characteristics, which reflects the N-C stoichiometry of the vegetation, had a
 374 negative direct effect on POC, MAOC and SOC stocks (Fig. 6h; Fig. 7a, b & c). The second axis, related to the plant
 375 economic spectrum (PES), had a positive direct effect on SOC stocks (Fig. 6h; Fig. 7c). In addition, the PES axis was
 376 positively correlated with the microbial biomass, thus having a positive indirect effect, and a significant total effect on
 377 POC, MAOC and SOC stocks (Fig. 6e; Fig. 7a, b & c). The third axis of the PCA (P-L axis), positively correlated with
 378 the lignin content and negatively related to vegetation productivity, was negatively correlated with the POC, MAOC and
 379 SOC stocks and the microbial biomass (Fig. 6e & h; Fig. 7a, b & c). The fourth axis, reporting the plant size, was positively
 380 correlated with the MAOC stocks, the MAOC/SOC ratio and the Fungi/Bacteria ratio (Fig. 6e & h; Fig. 7a & d). No index
 381 of plant functional and taxonomic diversity was retained during model selection. On the other hand, soil fertility had a
 382 direct and indirect positive effect on POC, MAOC and SOC stocks (Fig. 6g). Soil fertility was negatively correlated with
 383 the Gram+/Gram- ratio, the P-L axis and the N-C axis, and positively correlated with the PES axis of the vegetation PCA
 384 (Fig. 6i & j).

385 Plot under continuous grazing had a mean POC, MAOC and SOC stocks of 9.1, 6.8 and 16.2 Mg C ha⁻¹ respectively, and
 386 a MAOC/SOC ratio of 0.44 (Fig. 8). Rotational grazing significantly increased soil fertility compared to continuous
 387 grazing (Fig. 6k). This led to a significant indirect effect of rotational grazing on POC, MAOC and SOC stocks (Fig. 7b
 388 & c). POC, MAOC and SOC stocks under rotational grazing had a mean value of 10.5, 7.8 and 18.5 Mg C/ha, respectively,
 389 which were 15% (in the case of POC and MAOC) and 14% (for SOC stocks) higher than mean values in continuous
 390 grazing (Fig. 8b & c). Recent legume sowing had a negative direct effect on POC, MAOC and SOC (Fig. 6b). However,
 391 Lr also increased significantly the plant size axis, the microbial biomass and the soil fertility and decreased the plant N-
 392 C axis compared to Ct (Fig. 6a, c & k). These changes resulted in a positive indirect effect of Lr over POC, MAOC and
 393 SOC stocks and a null total effect (Fig. 7a, b, c & d). Grazing abandonment increased the vegetation N-C axis and reduced
 394 the microbial biomass compared to Ct (Fig. 6a & c), resulting in a negative indirect effect on POC, MAOC and SOC
 395 stocks and a significant negative total effect on POC stocks (Fig. 7a, b & c). Thus, the mean POC stock on abandoned
 396 plots was 8.1 Mg C/ha, 11% lower than in Ct plots (Fig. 8a).

397 Sand content in soil was negatively correlated with MAOC stocks and MAOC/SOC ratio (Fig. 6j). Sand content was also
 398 negatively correlated with soil fertility and the PES axis and positively correlated with the P-L and N-C axis (Fig. 6i),
 399 thus having a negative indirect effect on POC, MAOC and SOC stocks (Fig. 7a, b & c). As expected, due to its negative
 400 correlation with BD, gravel content was also negatively correlated with microbial biomass (Fig. 6j), leading to a
 401 significant negative indirect effect on POC, MAOC and SOC stocks (Fig. 7a, b & c). MAT was positively correlated with
 402 the N-C axis and negatively correlated with the soil fertility index (Fig. 6d & l). This implied a negative indirect effect of
 403 MAT over POC, MAOC and SOC stocks (Fig. 7a, b & c). On the other hand, MAP was negatively correlated with the N-
 404 C and the size axis (Fig. 6d). Thus, MAP had a positive indirect effect on POC, MAOC and SOC stocks (Fig. 7a, b, c &
 405 e).

406



417

418 **Figure 7. Direct, indirect and total standardized effects of all studied variables included in the structural equation model**
 419 **(Figure6) over the (a) particulate organic carbon (POC), (b) mineral-associated organic carbon (MAOC), and (c) soil organic**
 420 **carbon (SOC) stocks and (d) the relative MAOC abundance (MAOC/SOC). Bars indicate direct (dark colors) and indirect**
 421 **(light colors) effects, and the black points-ranges indicate the total (i.e. direct + indirect) effect (with its 95% confidence**
 422 **interval). Stars over the total effect values indicate significant effects at a level of 0.05.**




Figure 8. Violin plot and boxplot (with median and quartiles) for the (a) particulate organic carbon (POC), (b) mineral-associated organic carbon (MAOC), and (c) soil organic carbon (SOC) stocks and (d) the relative MAOC abundance (MAOC/SOC) in each management (Ct = Continuous grazing; Ab = grazing abandonment; Lr = Recent legume sowing; Lo = old legume sowing; Ro = Rotational grazing). The red point and the label indicate mean values predicted by the structure equation model (Fig.6). Lower case letters indicate significant differences ($p < 0.05$) between managements.

4. Discussion

Our results highlight the potential of management to control carbon storage in semi-arid grasslands. In this sense, rotational grazing arises as a promising tool for enhancing long-term carbon storage in the topsoil. We also identified several pathways by which management influences SOC, POC and MAOC stocks, showing the importance of changes in vegetation and microbial composition. Furthermore, management effects were consistent across the wide soil-climate gradient examined, enabling generalization of our results to a broad set of semi-arid grasslands in various environmental contexts. It should be noted that the predicted capacity of POC, MAOC and SOC stocks models was high (62%, 69% 73% of the variance explained by the fixed factors, respectively), indicating the robustness of the results presented and their importance for expanding the understanding of soil carbon dynamics in grasslands. Our results are limited to the upper topsoil (the first 8 cm), and it would be necessary to analyze deeper layers to fully understand the processes of SOC formation and stabilization and the effects of management changes. However this top layer is the most important for mediterranean grassland functioning as it contains the majority of roots and therefore most of the microbial, nutrients and water dynamics (Acosta-Gallo et al., 2011; Moreno et al., 2005). Furthermore, changes in management primarily affect the SOC of the topsoil layer, especially in the short term (Ward et al., 2016).

4.1. Soil organic carbon fractions, stocks, and saturation levels

SOC stocks and contents (Fig.8 & S4) in the paddocks studied ($16.5 \text{ Mg C ha}^{-1}$ and 19.5 g C kg^{-1} on average, respectively) were consistent with values found in the first 7-10 cm of soil in other grasslands under similar climatic conditions (Díaz-

448 Martínez et al., 2024; Oggioni et al., 2020; Parras-Alcántara et al., 2014). On the other hand, the MAOC/SOC ratio of our
449 soils (0.44 on average) was lower than the values generally reported in both global and semi-arid grasslands (Hansen et
450 al., 2024; Rocci et al., 2022; Shi et al., 2024), which are around 0.6-0.7. The shallow sampling depth in this work (8 cm)
451 may influence the observed MAOC/SOC ratios, as POM decrease more sharply than MAOM with soil depth (Galluzzi et
452 al., 2025; Sanderman et al., 2021; Tang et al., 2024). For instance, Plaza et al. (2022) reported MAOC/SOC ratios between
453 0.3 and 0.4 in the first 10 cm of soil, while Cappai et al. (2017) found MAOC/SOC ratios around 0.7 in the first 20 cm,
454 both in Mediterranean grasslands.

455 Carbon concentrations in the fine soil fraction (clay + silt) were well below the saturation point observed in previous
456 studies (Cotrufo et al., 2019a; Georgiou et al., 2022). However, we observed a certain limit to MAOC accumulation in
457 our system, as its content remained stable above SOC contents of 30 g/kg⁻¹ and stayed below 20 g kg⁻¹ even when SOC
458 reached values above 60 g kg⁻¹, following a saturation curve (Fig. 3b). These results support the observations that MAOC
459 accrual is more limited by the amount of C inputs rather than the mineralogical capacity of the soil (Poeplau et al., 2024).
460 In this sense, the maximum C in the clay+silt fraction observed in this work (around 19 g C/kg clay+silt) would represent
461 the effective capacity of the ecosystem to stabilize C at current levels of productivity. Therefore, MAOC saturation should
462 be a minor concern when planning management strategies to improve C storage in Mediterranean grasslands.

463

464 4.2. Soil microbial communities regulate SOC storage

465 Total microbial biomass had a similar and substantial positive effect on POC, MAOC and total SOC stocks, such that the
466 MAOC/SOC ratio was unaffected. Typically, microbial biomass C only represents about 2% of SOC (Xu et al., 2013; Yao
467 et al., 2000), but it is closely correlated with the accumulation of microbial necromass C (Hou et al., 2024). The latter can
468 account for more than 50% of SOC (Liang et al., 2019), with a similar contribution to the POC and MAOC fractions
469 (Zhang et al., 2024). The positive effect of microbial biomass on both POC and MAOC, as well as the relatively low C:N
470 ratio of these fractions found in this study (Fig. 3a) compared to other works (Chang et al., 2024; Yu et al., 2022), indicate
471 a prominent role of microbial transformation of plant inputs in SOC formation in semi-arid grasslands, which would also
472 explain the relatively low SOC stocks of these ecosystems.

473 Fungi/bacteria ratio was negatively correlated with MAOC fraction, as fungi residues tend to contribute more to POC
474 (Griepentrog et al., 2014; Lavalley et al., 2020; Tang et al., 2023), but no significant effect of fungi/bacteria ratio over
475 POC stocks or MAOC/SOC ratio was observed. Gram⁺/Gram⁻ ratio was negatively correlated with POC, MAOC and
476 SOC stock, as observed in previous studies (Khatri-Chhetri et al., 2024). Gram⁻ bacteria are more dependent on plant C
477 inputs, whereas Gram⁺ bacteria tend to use more of the organic C already present in the soil (Fanin et al., 2019; Kramer
478 and Gleixner, 2008; Waldrop and Firestone, 2004). Thus, the proliferation of Gram⁺ bacteria may promote the
479 decomposition of pre-existing SOM (Klumpp et al., 2009).

480

481 4.3. Plant-soil interactions and their effect on soil carbon stocks

482 The analysis of vegetation functional traits and chemical composition revealed four main axes of variation. Two of these
483 axes (axes 1 and 4) correspond to the spectrum of plant form and function, found in several global vegetation analyses
484 (Díaz et al., 2016; Weigelt et al., 2021). This two-dimensional spectrum is defined by the leaf and root economic gradient
485 (Kramer-Walter et al., 2016; Wright et al., 2004), that moves from slow-growing and resource-conserving to fast-growing
486 and resource-acquisitive species (Wright et al., 2004), and the size gradient, which increases with increasing plant height

487 and leaf area and reflects the competitive ability of plants for the light (Díaz et al., 2016). The other two axes (axes 2 and
488 3) are more related to the aboveground chemical composition of the vegetation. Axis 1 reflects the N-C stoichiometry of
489 plant tissues, driven mainly by the ABG C/N ratio, both of which are obviously negatively correlated with certain traits
490 such as LNC and the proportion of legumes. The N-fixing capacity of legumes has a fertilising effect on companion
491 plants, increasing its tissues nitrogen content (Pino et al., 2016). In addition, legumes themselves tend to have a high
492 concentration of nitrogen in leaves and stems (Carranca et al., 2015). Axis 3 is positively related to the lignin content of
493 the vegetation. This independence between lignin and nitrogen content in plant litter has already been observed (Cornwell
494 et al., 2008). ABG productivity of vegetation was correlated with most of these axes, especially with axis 3, being higher
495 in more acquisitive, less lignified and bigger vegetation, as found in other works (Laliberté and Tylianakis, 2012; Zhang
496 et al., 2017).

497 POC, MAOC and SOC stocks were higher in communities dominated by resource-acquisitive, highly productive plants
498 with low lignin content and low C/N ratio. These communities are expected to provide a higher level of plant inputs,
499 thereby increasing the incorporation of organic matter into the soil (King et al., 2023; Mortensen et al., 2025; Zhou et al.,
500 2024). Increased inputs may come from higher plant litter production but also from increased root exudates, which tend
501 to be higher in acquisitive plants (Guyonnet et al., 2018). Part of the effects of plant traits on SOC was mediated by
502 increases in microbial biomass, which would also benefit from higher levels of plant inputs and increased root exudation
503 (Eisenhauer et al., 2017). In addition, inputs with low lignin and C/N ratios are degraded more efficiently, reducing C
504 losses and promoting long-term SOC storage (Cotrufo and Lavelle, 2022; Ridgeway et al., 2022). However, based on
505 previous work, we would have expected a direct contribution of plant structural input to POC (Cotrufo et al., 2022), and
506 thus a higher proportion of POC in communities that produce more recalcitrant litter (Cheng et al., 2023; Haddix et al.,
507 2016; Mortensen et al., 2025). This was not the case in this work, where the MAOC/SOC ratio remained unaffected by
508 the chemical composition of the ABG vegetation tissues. This unexpected result could be explained by the fact that in
509 these systems, POC also appears to be the product of microbial transformation of plant inputs, as suggested by its positive
510 relationship with microbial biomass and its relatively low C/N ratio. In addition, photodegradation, an important driver
511 of litter degradation in semi-arid ecosystems (Austin and Vivanco, 2006), can promote litter lignin biodegradability and
512 the production of litter soluble compounds that are readily accessible to soil microbes (Wang et al., 2015), thus reducing
513 the influence of vegetation chemical properties on POC and MAOC formation. Grazing also decouples the quality of
514 plant tissues and the final quality of inputs to the soil, as livestock excreta have a lower C/N ratio than the plant material
515 consumed (Soussana and Lemaire, 2014). We also observed that taller and larger plants (high values on the size axis)
516 were associated with higher MAOC stocks and MAOC/SOC ratios, although the mechanism driving this correlation was
517 unclear. Generally, plant height is positively correlated with shoot:root ratio (Li et al., 2008), and several studies have
518 found a higher contribution of shoots, rather than roots, to the MAOC fraction due to the higher recalcitrance of root
519 tissues (Huang et al., 2021; Lavelle et al., 2018; Ridgeway et al., 2022). However, rhizodeposition, which is closely
520 linked to root growth and turnover, has been shown to promote MAOC formation over POC (Berenstecher et al., 2023;
521 Engedal et al., 2023; Villarino et al., 2021). On the other hand, a greater accumulation of standing litter, rather than surface
522 litter, might be expected in communities with bigger plants, and some studies in semi-arid grasslands have observed
523 higher rates of microbial degradation and release of soluble compounds (thus contributing more to MAOC) in standing
524 litter, compared to surface litter, due to greater retention of night-time moisture in the former (Gliksman et al., 2018; Wang
525 et al., 2017). In any case, our results and their interpretation are limited by the correlation in our vegetation data between
526 input quantity (ABG productivity) and input quality, which could mask the effect of the latter on SOC formation.

527 No significant effects of taxonomic or functional diversity on POC, MAOC or SOC stocks were found, which could be
528 explained by the negative correlation of these biodiversity indicators with the ABG productivity and soil fertility in our

plots (Fig. S7). This negative correlation has been observed in other natural grassland research and may be due to the dominance of some highly productive species in nutrient-rich environments (Feßel et al., 2016; Helsen et al., 2014; Luo et al., 2019; Rolo et al., 2016). In this sense, our results would be consistent with previous research that has found a greater importance of functional identity (i.e. mean traits values) relative to functional diversity in predicting ecosystem processes (Mokany et al., 2008; Zhang et al., 2017).

Soil fertility plays a central role in our models, increasing POC, MAOC and SOC stocks. Previous studies in grasslands have already observed a positive correlation between POC and MAOC and soil nutrients content (Wang et al., 2025). Global N addition experiments have also showed to increase topsoil C storage in grasslands (Hu et al., 2024; Liu et al., 2023). In this work, we observe a direct effect of soil fertility on C storage, but we also identify some indirect effects mediated by changes in vegetation and microbial communities. More resource-acquisitive and productive plant communities, with lignin content and lower C/N ratio, were found at higher soil fertility. Resource-acquisitive plants are expected to be more competitive in nutrient-rich environments (Daou et al., 2021; Ordoñez et al., 2009). The absence of nutrient limitation also promotes plant productivity (Fay et al., 2015). In addition, soil fertility was negatively correlated with the Gram+/Gram- ratio. Gram- bacteria have a more copiotroph strategy than Gram+ bacteria and therefore benefit more from nutrient-rich environments (De Deyn et al., 2010; Luo et al., 2020; Zhong et al., 2010). The direct positive effect of soil fertility on C storage in soils may be explained by changes in other factors that we did not measure. For instance, nutrient addition have been shown to promote below-ground productivity in mediterranean woody pastures (Nair et al., 2019) and in global grasslands (Liu et al., 2023; Lu et al., 2011). Moreover, nutrient availability in soils tends to increase CUE (Poeplau et al., 2019; Spohn et al., 2016) and to alleviate the need for microbial N-mining, reducing C losses and old SOM decomposition (Blagodatskaya and Kuzyakov, 2008).

4.5. Management effects

Rotational grazing increased both POC, MAOC and SOC stocks by 15%, 15% and 14%, respectively, compared to conventional continuous grazing. Previous studies found the same positive effect of rotational grazing on MAOC (Khatri-Chhetri et al., 2024; Mosier et al., 2021; Stanley et al., 2025) and total SOC content (Byrnes et al., 2018; Conant et al., 2017; Liu et al., 2024; Phukubye et al., 2022; Teague et al., 2011). According to our results, this effect is mainly explained by the higher soil fertility observed in rotational grazing plots, cascading in the above explained effects on plant traits and microbial transformation leading to higher POC, MAOC and SOC stocks. Soil fertility may be higher under rotational grazing than under continuous grazing because fecal and urine excretions tend to be less spatially clustered in rotationally grazed paddocks, resulting in more homogeneous fertilization of the entire paddock (Augustine et al., 2023; Dubeux Jr. et al., 2014). In contrast, in continuous grazing, excreta tend to accumulate in areas of highest animal use, such as near feeders or water points (Tate et al., 2003), which are areas that have been avoided in our sampling. In turn, the homogeneous grazing and fertilization maintains a greater amount of ground covered by vegetation (Stanley et al., 2025; Teague et al., 2004), which limits topsoil losses through erosion (Sanjari et al., 2009).

Surprisingly legume sowings had a negative direct effect on soil C stocks, but a positive indirect effect, resulting in a non-significant total effect. The negative impact was more evident in recent sowings, especially on the MAOC fraction, and could be attributed to the impact of the pre-sowing tillage. Previous works have observed a reduction in SOC stocks after tilling in mediterranean grasslands (Parras-Alcántara et al., 2015; Uribe et al., 2015). This would explain why the direct effect of legume sowing in C storage is more negative in recent compared to old-sowed paddocks. The effect of tillage on SOC stocks may arise from changes in soil bulk density rather than C content, resulting in misleading conclusions (Du et al., 2017; Rovira et al., 2022). However, in this work the analysis of POC, MAOC and SOC contents (in g/kg bulk soil;

Fig. S6 & S8) shows similar effects of legume sowing as those observed for C stocks (Fig. 7a, b & c). In the recently sown plots, the negative impact in C stocks was countered by an increase in soil fertility, plant productivity and plant nitrogen content, possibly due to increased legume cover (Gómez-Rey et al., 2012; Gou et al., 2023; Hernández-Esteban et al., 2019). In addition, recent legume sowing also implied an increase in microbial biomass and a decrease in the Gram+/Gram- ratio, as observed in previous studies (Moreno et al., 2021).

Grazing abandonment led to a 12% reduction in the POC stocks, compared to continuous grazing. This result contradicts the observations of previous studies in grasslands, which reported positive effects of grazing exclusion on SOC storage, both globally (Eze et al., 2018) and particularly in semi-arid climates (Cheng et al., 2016; Gao et al., 2025; Yu et al., 2021). However McSherry and Ritchie (2013) meta-analysis found that light grazing could promote SOC stocks in C3 dominated grasslands (such as our study grassland). Other global and regional studies also observed null or even positive effects of light grazing, compared to grazing exclusion, in SOC stocks in grasslands (Liu et al., 2024; Zhou et al., 2017). Our results are in line with previous studies in mediterranean woody pastures that found a decrease in SOC storage with grazing abandonment (Oggioni et al., 2020; Peco et al., 2017). The fact that grazing abandonment particularly affected the POC fraction is surprising, since in abandoned paddocks there were a greater accumulation of plant structural biomass, that should promote POC accrual (Cotrufo et al., 2022). According to our results, the negative effect of grazing exclusion on POC stocks was mainly mediated by a reduction in microbial biomass, and a proliferation of plants with higher C/N ratios. Once again pointing to the microbially transformed nature of POC in these soils. Grazing has been proven to increase microbial activity and biomass (Bardgett et al., 2001; Zhou et al., 2017), in part by increasing root exudation (Hamilton et al., 2008; Wilson et al., 2018). Previous studies have also observed an increase in vegetation C/N ratio with cessation of grazing (He et al., 2020; Wang et al., 2016), which could be explained by the accumulation of older plant tissues and senescent standing biomass, with higher C/N than green biomass and new leaves (Sanaullah et al., 2010).

Importantly, we observed no interactions between management effects and climate or soil texture variables, showing that the observed management effects were consistent across the entire environmental gradient. Although such interactions have been observed for wider climatic gradients (Byrnes et al., 2018; Phukubye et al., 2022), our results show that management practices such as rotational grazing can have net positive impacts on SOC storage in grasslands with varying conditions within semi-arid regions.

596

597 4.6. Climate effects and implications for global warming

Even within the small climatic gradient covered in this study, we found lower stocks of POC, MAOC and SOC in the warmer areas, as observed in previous research (Díaz-Martínez et al., 2024; Georgiou et al., 2024; Hansen et al., 2024; Shi et al., 2024). According to our results, these negative effects were entirely mediated by a decrease in soil fertility and vegetation input quality (increased N-C axis) with increasing temperature. Increased aridity has been shown to be associated with the alteration of several ecosystem processes that regulate nutrient cycling and availability (Berdugo et al., 2020; Moreno-Jiménez et al., 2019). Increased fiber for tissue protection, typical from plants growing at higher temperatures, may explain the increase in vegetation C/N ratios (Arroyo et al., 2024). On the other hand, annual precipitation exerted a small positive effect on POC and SOC stocks, mediated by the reduction of C/N ratios in vegetation in wetter farms. Based on future projections of increasing temperatures and decreasing annual precipitation in our study area (IPCC, 2023b) we could expect significant reductions in SOC stocks in Mediterranean grasslands in the next decades, affecting both the POC and MAOC fractions.

609

610 **5. Conclusions:**

611 Implementing rotational grazing can improve both POC and MAOC stocks in Mediterranean grasslands. On the other
612 hand, the abandonment of grazing can lead to a loss of soil carbon storage capacity in these ecosystems.

613 Changes in microbial communities and vegetation attributes were the main drivers of changes in SOC stocks and fractions.
614 In this sense, this work has proven the usefulness of plant functional traits as tools for the study of plant-soil interactions
615 and SOC formation dynamics.

616 We can expect a loss of soil carbon in the studied grasslands over the next few decades due to grazing abandonment and
617 climate warming, so selecting management approaches that mitigate or counteract these losses is vital to maintaining the
618 fertility, productivity and functioning of semi-arid grasslands.

619

620 **Data and code availability**

621 The data and code that support the findings of this study are available from the corresponding author upon reasonable
622 request.

623

624 **CRedit authorship contribution statement**

625 **AC:** Data curation, Formal analysis, Investigation, Software, Visualization, Writing – original draft. **GM:**
626 Conceptualization, Funding acquisition, Project administration, Supervision, Writing – review and editing. **MFC:**
627 Conceptualization, Methodology, Supervision, Writing – review and editing. **CF:** Data curation, Methodology,
628 Investigation, Resources, Writing – review and editing. **SR:** Data curation, Methodology, Investigation, Resources,
629 Writing – review and editing. **VR:** Conceptualization, Data curation, Formal analysis, Funding acquisition, Investigation,
630 Project administration, Supervision, Writing – review and editing.

631

632 **Competing interests**

633 The authors declare that they have no conflict of interest.

634

635 **Financial support**

636 This work was part of the projects “PID2019-108313RB-C33”, “PID2019-108313RB-C31” and “PCIN2021-122100-2”
637 funded by MICIU/AEI/10.13039/501100011033/ and “European Union NextGenerationEU/PRTR; the project “CLU-
638 2019-05 – IRNASA/CSIC Unit of Excellence” funded by the “Junta de Castilla y León” and co-financed by the European
639 Union (ERDF “Europe drives our growth”); and European Union’s Horizon Europe research and innovation program.
640 Grant agreement: 101059794. Alejandro Carrascosa was supported by the Spanish Ministry of Science and Innovation
641 through an FPU grant (FPU21/02668). Cristina Frade was supported by a JCyL fellowship (E-37-2023-0024090) co-
642 financed by the European Social Fund Plus (FSE+).

643

644 **References:**

645 Acosta-Gallo, B., Casado, M. A., Montalvo, J., and Pineda, F. D.: Allometric patterns of below-ground biomass in
646 Mediterranean grasslands, *Plant Biosyst. - Int. J. Deal. Asp. Plant Biol.*, 145, 584–595,
647 <https://doi.org/10.1080/11263504.2011.578836>, 2011.

648 Angst, G., Mueller, K. E., Kögel-Knabner, I., Freeman, K. H., and Mueller, C. W.: Aggregation controls the stability of
649 lignin and lipids in clay-sized particulate and mineral associated organic matter, *Biogeochemistry*, 132, 307–324,
650 <https://doi.org/10.1007/s10533-017-0304-2>, 2017.

651 Angst, G., Mueller, K. E., Nierop, K. G. J., and Simpson, M. J.: Plant- or microbial-derived? A review on the molecular
652 composition of stabilized soil organic matter, *Soil Biol. Biochem.*, 156, 108189,
653 <https://doi.org/10.1016/j.soilbio.2021.108189>, 2021.

654 Arroyo, A. I., Pueyo, Y., Barrantes, O., and Alados, C. L.: Interplay between Livestock Grazing and Aridity on the
655 Ecological and Nutritional Value of Forage in Semi-arid Mediterranean Rangelands (NE Spain), *Environ. Manage.*,
656 <https://doi.org/10.1007/s00267-024-01939-9>, 2024.

657 Augustine, D. J., Kearney, S. P., Raynor, E. J., Porensky, L. M., and Derner, J. D.: Adaptive, multi-paddock, rotational
658 grazing management alters foraging behavior and spatial grazing distribution of free-ranging cattle, *Agric. Ecosyst.*
659 *Environ.*, 352, 108521, <https://doi.org/10.1016/j.agee.2023.108521>, 2023.

660 Austin, A. T. and Vivanco, L.: Plant litter decomposition in a semi-arid ecosystem controlled by photodegradation, *Nature*,
661 442, 555–558, <https://doi.org/10.1038/nature05038>, 2006.

662 Bai, X., Zhai, G., Wang, B., An, S., Liu, J., Xue, Z., and Dippold, M. A.: Litter quality controls the contribution of
663 microbial carbon to main microbial groups and soil organic carbon during its decomposition, *Biol. Fertil. Soils*, 60, 167–
664 181, <https://doi.org/10.1007/s00374-023-01792-8>, 2024.

665 Bai, Y. and Cotrufo, M. F.: Grassland soil carbon sequestration: Current understanding, challenges, and solutions, *Science*,
666 377, 603–608, <https://doi.org/10.1126/science.abo2380>, 2022.

667 Baldock, J. A. and Skjemstad, J. O.: Role of the soil matrix and minerals in protecting natural organic materials against
668 biological attack, *Org. Geochem.*, 31, 697–710, [https://doi.org/10.1016/S0146-6380\(00\)00049-8](https://doi.org/10.1016/S0146-6380(00)00049-8), 2000.

669 Bardgett, R. D., Jones, A. C., Jones, D. L., Kemmitt, S. J., Cook, R., and Hobbs, P. J.: Soil microbial community patterns
670 related to the history and intensity of grazing in sub-montane ecosystems, *Soil Biol. Biochem.*, 33, 1653–1664,
671 [https://doi.org/10.1016/S0038-0717\(01\)00086-4](https://doi.org/10.1016/S0038-0717(01)00086-4), 2001.

672 Bartholomew, P. W. and Williams, R. D.: Overseeding Unimproved Warm-Season Pasture with Cool- and Warm-Season
673 Legumes to Enhance Forage Productivity, *J. Sustain. Agric.*, 34, 125–140, <https://doi.org/10.1080/10440040903482407>,
674 2010.

675 Bates, D., Mächler, M., Bolker, B., and Walker, S.: Fitting Linear Mixed-Effects Models Using lme4, *J. Stat. Softw.*, 67,
676 1–48, <https://doi.org/10.18637/jss.v067.i01>, 2015.

677 Benbi, D. K., Boparai, A. K., and Brar, K.: Decomposition of particulate organic matter is more sensitive to temperature
678 than the mineral associated organic matter, *Soil Biol. Biochem.*, 70, 183–192,
679 <https://doi.org/10.1016/j.soilbio.2013.12.032>, 2014.

680 Berdugo, M., Delgado-Baquerizo, M., Soliveres, S., Hernández-Clemente, R., Zhao, Y., Gaitán, J. J., Gross, N., Saiz, H.,
681 Maire, V., Lehmann, A., Rillig, M. C., Solé, R. V., and Maestre, F. T.: Global ecosystem thresholds driven by aridity,
682 *Science*, 367, 787–790, <https://doi.org/10.1126/science.aay5958>, 2020.

683 Berenstecher, P., Conti, G., Faigón, A., and Piñeiro, G.: Tracing service crops' net carbon and nitrogen rhizodeposition
684 into soil organic matter fractions using dual isotopic brush-labeling, *Soil Biol. Biochem.*, 184, 109096,
685 <https://doi.org/10.1016/j.soilbio.2023.109096>, 2023.

686 Bergmann, J., Weigelt, A., van der Plas, F., Laughlin, D. C., Kuyper, T. W., Guerrero-Ramirez, N., Valverde-Barrantes,
687 O. J., Bruelheide, H., Freschet, G. T., Iversen, C. M., Kattge, J., McCormack, M. L., Meier, I. C., Rillig, M. C., Roumet,
688 C., Semchenko, M., Sweeney, C. J., van Ruijven, J., York, L. M., and Mommer, L.: The fungal collaboration gradient
689 dominates the root economics space in plants, *Sci. Adv.*, 6, eaba3756, <https://doi.org/10.1126/sciadv.aba3756>, 2020.

690 Blagodatskaya, E. and Kuzyakov, Y.: Mechanisms of real and apparent priming effects and their dependence on soil
691 microbial biomass and community structure: critical review, *Biol. Fertil. Soils*, 45, 115–131,
692 <https://doi.org/10.1007/s00374-008-0334-y>, 2008.

- 693 Briske, D. D., Derner, J. D., Brown, J. R., Fuhlendorf, S. D., Teague, W. R., Havstad, K. M., Gillen, R. L., Ash, A. J., and
694 Willms, W. D.: Rotational Grazing on Rangelands: Reconciliation of Perception and Experimental Evidence, *Rangel.*
695 *Ecol. Manag.*, 61, 3–17, <https://doi.org/10.2111/06-159R.1>, 2008.
- 696 Byrnes, R. C., Eastburn, D. J., Tate, K. W., and Roche, L. M.: A Global Meta-Analysis of Grazing Impacts on Soil Health
697 Indicators, *J. Environ. Qual.*, 47, 758–765, <https://doi.org/10.2134/jeq2017.08.0313>, 2018.
- 698 Cappai, C., Kemanian, A. R., Lagomarsino, A., Roggero, P. P., Lai, R., Agnelli, A. E., and Seddaiu, G.: Small-scale spatial
699 variation of soil organic matter pools generated by cork oak trees in Mediterranean agro-silvo-pastoral systems,
700 *Geoderma*, 304, 59–67, <https://doi.org/10.1016/j.geoderma.2016.07.021>, 2017.
- 701 Carmona, C. P., Rota, C., Azcárate, F. M., and Peco, B.: More for less: sampling strategies of plant functional traits across
702 local environmental gradients, *Funct. Ecol.*, 29, 579–588, <https://doi.org/10.1111/1365-2435.12366>, 2015.
- 703 Carranca, C., Castro, I. V., Figueiredo, N., Redondo, R., Rodrigues, A. R. F., Saraiva, I., Maricato, R., and Madeira, M.
704 A. V.: Influence of tree canopy on N₂ fixation by pasture legumes and soil rhizobial abundance in Mediterranean oak
705 woodlands, *Sci. Total Environ.*, 506–507, 86–94, <https://doi.org/10.1016/j.scitotenv.2014.10.111>, 2015.
- 706 Carranca, C., Pedra, F., and Madeira, M.: Enhancing Carbon Sequestration in Mediterranean Agroforestry Systems: A
707 Review, *Agriculture*, 12, 1598, <https://doi.org/10.3390/agriculture12101598>, 2022.
- 708 Carrascosa, A., Moreno, G., Rodrigo, S., and Rolo, V.: Unravelling the contribution of soil, climate and management to
709 the productivity of ecologically intensified Mediterranean wood pastures, *Sci. Total Environ.*, 957, 177575,
710 <https://doi.org/10.1016/j.scitotenv.2024.177575>, 2024.
- 711 Chang, Y., Sokol, N. W., van Groenigen, K. J., Bradford, M. A., Ji, D., Crowther, T. W., Liang, C., Luo, Y., Kuzyakov, Y.,
712 Wang, J., and Ding, F.: A stoichiometric approach to estimate sources of mineral-associated soil organic matter, *Glob.*
713 *Change Biol.*, 30, e17092, <https://doi.org/10.1111/gcb.17092>, 2024.
- 714 Cheng, J., Jing, G., Wei, L., and Jing, Z.: Long-term grazing exclusion effects on vegetation characteristics, soil properties
715 and bacterial communities in the semi-arid grasslands of China, *Ecol. Eng.*, 97, 170–178,
716 <https://doi.org/10.1016/j.ecoleng.2016.09.003>, 2016.
- 717 Cheng, X., Xing, W., and Liu, J.: Litter chemical traits, microbial and soil stoichiometry regulate organic carbon accrual
718 of particulate and mineral-associated organic matter, *Biol. Fertil. Soils*, 59, 777–790, [https://doi.org/10.1007/s00374-023-](https://doi.org/10.1007/s00374-023-01746-0)
719 [01746-0](https://doi.org/10.1007/s00374-023-01746-0), 2023.
- 720 Conant, R. T.: Grassland Soil Organic Carbon Stocks: Status, Opportunities, Vulnerability, in: *Recarbonization of the*
721 *Biosphere: Ecosystems and the Global Carbon Cycle*, edited by: Lal, R., Lorenz, K., Hüttl, R. F., Schneider, B. U., and
722 von Braun, J., Springer Netherlands, Dordrecht, 275–302, https://doi.org/10.1007/978-94-007-4159-1_13, 2012.
- 723 Conant, R. T., Cerri, C. E. P., Osborne, B. B., and Paustian, K.: Grassland management impacts on soil carbon stocks: a
724 new synthesis, *Ecol. Appl.*, 27, 662–668, <https://doi.org/10.1002/eap.1473>, 2017.
- 725 Cornwell, W. K., Cornelissen, J. H. C., Amatangelo, K., Dorrepaal, E., Eviner, V. T., Godoy, O., Hobbie, S. E., Hoorens,
726 B., Kurokawa, H., Pérez-Harguindeguy, N., Quested, H. M., Santiago, L. S., Wardle, D. A., Wright, I. J., Aerts, R., Allison,
727 S. D., Van Bodegom, P., Brovkin, V., Chatain, A., Callaghan, T. V., Díaz, S., Garnier, E., Gurvich, D. E., Kazakou, E.,
728 Klein, J. A., Read, J., Reich, P. B., Soudzilovskaia, N. A., Vaieretti, M. V., and Westoby, M.: Plant species traits are the
729 predominant control on litter decomposition rates within biomes worldwide, *Ecol. Lett.*, 11, 1065–1071,
730 <https://doi.org/10.1111/j.1461-0248.2008.01219.x>, 2008.
- 731 Cotrufo, M. F. and Lavelle, J. M.: Chapter One - Soil organic matter formation, persistence, and functioning: A synthesis
732 of current understanding to inform its conservation and regeneration, in: *Advances in Agronomy*, vol. 172, edited by:
733 Sparks, D. L., Academic Press, 1–66, <https://doi.org/10.1016/bs.agron.2021.11.002>, 2022.
- 734 Cotrufo, M. F., Wallenstein, M. D., Boot, C. M., Denef, K., and Paul, E.: The Microbial Efficiency-Matrix Stabilization
735 (MEMS) framework integrates plant litter decomposition with soil organic matter stabilization: do labile plant inputs form
736 stable soil organic matter?, *Glob. Change Biol.*, 19, 988–995, <https://doi.org/10.1111/gcb.12113>, 2013.
- 737 Cotrufo, M. F., Ranalli, M. G., Haddix, M. L., Six, J., and Lugato, E.: Soil carbon storage informed by particulate and
738 mineral-associated organic matter, *Nat. Geosci.*, 12, 989–994, <https://doi.org/10.1038/s41561-019-0484-6>, 2019a.
- 739 Cotrufo, M. F., Ranalli, M. G., Haddix, M. L., Six, J., and Lugato, E.: Soil carbon storage informed by particulate and
740 mineral-associated organic matter, *Nat. Geosci.*, 12, 989–994, <https://doi.org/10.1038/s41561-019-0484-6>, 2019b.

741 Cotrufo, M. F., Haddix, M. L., Kroeger, M. E., and Stewart, C. E.: The role of plant input physical-chemical properties,
742 and microbial and soil chemical diversity on the formation of particulate and mineral-associated organic matter, *Soil Biol.*
743 *Biochem.*, 168, 108648, <https://doi.org/10.1016/j.soilbio.2022.108648>, 2022.

744 Crowther, T. W., Todd-Brown, K. E. O., Rowe, C. W., Wieder, W. R., Carey, J. C., Machmuller, M. B., Snoek, B. L., Fang,
745 S., Zhou, G., Allison, S. D., Blair, J. M., Bridgham, S. D., Burton, A. J., Carrillo, Y., Reich, P. B., Clark, J. S., Classen, A.
746 T., Dijkstra, F. A., Elberling, B., Emmett, B. A., Estiarte, M., Frey, S. D., Guo, J., Harte, J., Jiang, L., Johnson, B. R.,
747 Kröel-Dulay, G., Larsen, K. S., Laudon, H., Lavallee, J. M., Luo, Y., Lupascu, M., Ma, L. N., Marhan, S., Michelsen, A.,
748 Mohan, J., Niu, S., Pendall, E., Peñuelas, J., Pfeifer-Meister, L., Poll, C., Reinsch, S., Reynolds, L. L., Schmidt, I. K.,
749 Sistla, S., Sokol, N. W., Templer, P. H., Treseder, K. K., Welker, J. M., and Bradford, M. A.: Quantifying global soil carbon
750 losses in response to warming, *Nature*, 540, 104–108, <https://doi.org/10.1038/nature20150>, 2016.

751 Crowther, T. W., van den Hoogen, J., Wan, J., Mayes, M. A., Keiser, A. D., Mo, L., Averill, C., and Maynard, D. S.: The
752 global soil community and its influence on biogeochemistry, *Science*, 365, eaav0550,
753 <https://doi.org/10.1126/science.aav0550>, 2019.

754 Dai, W., Xiao, R., Wei, C., and Yang, F.: Plant litter traits control the accumulation of mineral-associated organic carbon
755 by influencing its molecular composition and diversity, *Soil Tillage Res.*, 253, 106667,
756 <https://doi.org/10.1016/j.still.2025.106667>, 2025.

757 Daou, L., Garnier, É., and Shipley, B.: Quantifying the relationship linking the community-weighted means of plant traits
758 and soil fertility, *Ecology*, 102, e03454, <https://doi.org/10.1002/ecy.3454>, 2021.

759 De Deyn, G. B., Quirk, H., and Bardgett, R. D.: Plant species richness, identity and productivity differentially influence
760 key groups of microbes in grassland soils of contrasting fertility, *Biol. Lett.*, 7, 75–78,
761 <https://doi.org/10.1098/rsbl.2010.0575>, 2010.

762 Díaz, S., Kattge, J., Cornelissen, J. H. C., Wright, I. J., Lavorel, S., Dray, S., Reu, B., Kleyer, M., Wirth, C., Colin Prentice,
763 I., Garnier, E., Bönsch, G., Westoby, M., Poorter, H., Reich, P. B., Moles, A. T., Dickie, J., Gillison, A. N., Zanne, A. E.,
764 Chave, J., Joseph Wright, S., Sheremet'ev, S. N., Jactel, H., Baraloto, C., Cerabolini, B., Pierce, S., Shipley, B., Kirkup,
765 D., Casanoves, F., Joswig, J. S., Günther, A., Falczuk, V., Rüger, N., Mahecha, M. D., and Gorné, L. D.: The global
766 spectrum of plant form and function, *Nature*, 529, 167–171, <https://doi.org/10.1038/nature16489>, 2016.

767 Díaz-Martínez, P., Maestre, F. T., Moreno-Jiménez, E., Delgado-Baquerizo, M., Eldridge, D. J., Saiz, H., Gross, N., Le
768 Bagousse-Pinguet, Y., Gozalo, B., Ochoa, V., Guirado, E., García-Gómez, M., Valencia, E., Asensio, S., Berdugo, M.,
769 Martínez-Valderrama, J., Mendoza, B. J., García-Gil, J. C., Zacccone, C., Panettieri, M., García-Palacios, P., Fan, W.,
770 Benavente-Ferraces, I., Rey, A., Eisenhauer, N., Cesarz, S., Abedi, M., Ahumada, R. J., Alcántara, J. M., Amghar, F.,
771 Aramayo, V., Arroyo, A. I., Bahalkeh, K., Ben Salem, F., Blaum, N., Boldgiv, B., Bowker, M. A., Bran, D., Branquinho,
772 C., Bu, C., Cáceres, Y., Canessa, R., Castillo-Monroy, A. P., Castro, I., Castro-Quezada, P., Chibani, R., Conceição, A. A.,
773 Currier, C. M., Darrouzet-Nardi, A., Deák, B., Dickman, C. R., Donoso, D. A., Dougill, A. J., Durán, J., Ejtehadi, H.,
774 Espinosa, C., Fajardo, A., Farzam, M., Ferrante, D., Fraser, L. H., Gaitán, J. J., Gusman Montalván, E., Hernández-
775 Hernández, R. M., von Hessberg, A., Hölzel, N., Huber-Sannwald, E., Hughes, F. M., Jadán-Maza, O., Geissler, K.,
776 Jentsch, A., Ju, M., Kaseke, K. F., Kindermann, L., Koopman, J. E., Le Roux, P. C., Liancourt, P., Linstädter, A., Liu, J.,
777 Louw, M. A., Maggs-Kölling, G., Makhallanyane, T. P., Issa, O. M., Marais, E., Margerie, P., Mazaneda, A. J., McClaran,
778 M. P., Messeder, J. V. S., Mora, J. P., Moreno, G., Munson, S. M., Nunes, A., Oliva, G., Oñatibia, G. R., Osborne, B.,
779 Peter, G., Pueyo, Y., Quiroga, R. E., Reed, S. C., Reyes, V. M., et al.: Vulnerability of mineral-associated soil organic
780 carbon to climate across global drylands, *Nat. Clim. Change*, 14, 976–982, <https://doi.org/10.1038/s41558-024-02087-y>,
781 2024.

782 Dondini, M., Martin, M., De Camillis, C., Uwizeye, A., Soussana, J.-F., Robinson, T., and Steinfeld, H.: Global assessment
783 of soil carbon in grasslands – From current stock estimates to sequestration potential., *FAO*, Rome, 2023.

784 Du, Z., Angers, D. A., Ren, T., Zhang, Q., and Li, G.: The effect of no-till on organic C storage in Chinese soils should
785 not be overemphasized: A meta-analysis, *Agric. Ecosyst. Environ.*, 236, 1–11, <https://doi.org/10.1016/j.agee.2016.11.007>,
786 2017.

787 Dubeux Jr., J. c. b., Sollenberger, L. e., Vendramini, J. m. b., Interrante, S. m., and Lira Jr., M. a.: Stocking Method,
788 Animal Behavior, and Soil Nutrient Redistribution: How are They Linked?, *Crop Sci.*, 54, 2341–2350,
789 <https://doi.org/10.2135/cropsci2014.01.0076>, 2014.

790 Eisenhauer, N., Lanoue, A., Strecker, T., Scheu, S., Steinauer, K., Thakur, M. P., and Mommer, L.: Root biomass and
791 exudates link plant diversity with soil bacterial and fungal biomass, *Sci. Rep.*, 7, 44641,
792 <https://doi.org/10.1038/srep44641>, 2017.

793 Elias, D. M. O., Mason, K. E., Goodall, T., Taylor, A., Zhao, P., Otero-Fariña, A., Chen, H., Peacock, C. L., Ostle, N. J.,
794 Griffiths, R., Chapman, P. J., Holden, J., Banwart, S., McNamara, N. P., and Whitaker, J.: Microbial and mineral
795 interactions decouple litter quality from soil organic matter formation, *Nat. Commun.*, 15, 10063,
796 <https://doi.org/10.1038/s41467-024-54446-0>, 2024.

797 Engedal, T., Magid, J., Hansen, V., Rasmussen, J., Sørensen, H., and Stoumann Jensen, L.: Cover crop root morphology
798 rather than quality controls the fate of root and rhizodeposition C into distinct soil C pools, *Glob. Change Biol.*, 29, 5677–
799 5690, <https://doi.org/10.1111/gcb.16870>, 2023.

800 Eze, S., Palmer, S. M., and Chapman, P. J.: Soil organic carbon stock in grasslands: Effects of inorganic fertilizers, liming
801 and grazing in different climate settings, *J. Environ. Manage.*, 223, 74–84, <https://doi.org/10.1016/j.jenvman.2018.06.013>,
802 2018.

803 Fanin, N., Kardol, P., Farrell, M., Nilsson, M.-C., Gundale, M. J., and Wardle, D. A.: The ratio of Gram-positive to Gram-
804 negative bacterial PLFA markers as an indicator of carbon availability in organic soils, *Soil Biol. Biochem.*, 128, 111–
805 114, <https://doi.org/10.1016/j.soilbio.2018.10.010>, 2019.

806 Faucon, M.-P., Houben, D., and Lambers, H.: Plant Functional Traits: Soil and Ecosystem Services, *Trends Plant Sci.*, 22,
807 385–394, <https://doi.org/10.1016/j.tplants.2017.01.005>, 2017.

808 Fay, P. A., Prober, S. M., Harpole, W. S., Knops, J. M. H., Bakker, J. D., Borer, E. T., Lind, E. M., MacDougall, A. S.,
809 Seabloom, E. W., Wragg, P. D., Adler, P. B., Blumenthal, D. M., Buckley, Y. M., Chu, C., Cleland, E. E., Collins, S. L.,
810 Davies, K. F., Du, G., Feng, X., Firn, J., Gruner, D. S., Hagenah, N., Hautier, Y., Heckman, R. W., Jin, V. L., Kirkman, K.
811 P., Klein, J., Ladwig, L. M., Li, Q., McCulley, R. L., Melbourne, B. A., Mitchell, C. E., Moore, J. L., Morgan, J. W., Risch,
812 A. C., Schütz, M., Stevens, C. J., Wedin, D. A., and Yang, L. H.: Grassland productivity limited by multiple nutrients,
813 *Nat. Plants*, 1, 1–5, <https://doi.org/10.1038/nplants.2015.80>, 2015.

814 Feßel, C., Meier, I. C., and Leuschner, C.: Relationship between species diversity, biomass and light transmittance in
815 temperate semi-natural grasslands: is productivity enhanced by complementary light capture?, *J. Veg. Sci.*, 27, 144–155,
816 <https://doi.org/10.1111/jvs.12326>, 2016.

817 Fortunel, C., Garnier, E., Joffre, R., Kazakou, E., Quested, H., Grigulis, K., Lavorel, S., Ansquer, P., Castro, H., Cruz, P.,
818 Doležal, J., Eriksson, O., Freitas, H., Golodets, C., Jouany, C., Kigel, J., Kleyer, M., Lehsten, V., Lepš, J., Meier, T.,
819 Pakeman, R., Papadimitriou, M., Papanastasis, V. P., Quétier, F., Robson, M., Sternberg, M., Theau, J.-P., Thébault, A.,
820 and Zarovali, M.: Leaf traits capture the effects of land use changes and climate on litter decomposability of grasslands
821 across Europe, *Ecology*, 90, 598–611, <https://doi.org/10.1890/08-0418.1>, 2009.

822 Freschet, G. T., Aerts, R., and Cornelissen, J. H. C.: A plant economics spectrum of litter decomposability, *Funct. Ecol.*,
823 26, 56–65, <https://doi.org/10.1111/j.1365-2435.2011.01913.x>, 2012.

824 Frongia, A., Pulina, A., Tanda, A., Seddaiu, G., Roggero, P. P., and Moreno, G.: Assessing the effect of rotational grazing
825 adoption in Iberian silvopastoral systems with Normalized Difference Vegetation Index time series, *Ital. J. Agron.*,
826 <https://doi.org/10.4081/ija.2023.2185>, 2023.

827 Frostegård, A. and Bååth, E.: The use of phospholipid fatty acid analysis to estimate bacterial and fungal biomass in soil,
828 *Biol. Fertil. Soils*, 22, 59–65, <https://doi.org/10.1007/BF00384433>, 1996.

829 Funk, J. L., Larson, J. E., Ames, G. M., Butterfield, B. J., Cavender-Bares, J., Firn, J., Laughlin, D. C., Sutton-Grier, A.
830 E., Williams, L., and Wright, J.: Revisiting the Holy Grail: using plant functional traits to understand ecological processes,
831 *Biol. Rev.*, 92, 1156–1173, <https://doi.org/10.1111/brv.12275>, 2017.

832 Galluzzi, G., Plaza, C., Giannetta, B., Priori, S., and Zaccone, C.: Time and climate roles in driving soil carbon distribution
833 and stability in particulate and mineral-associated organic matter pools, *Sci. Total Environ.*, 963, 178511,
834 <https://doi.org/10.1016/j.scitotenv.2025.178511>, 2025.

835 Gang, C., Zhou, W., Chen, Y., Wang, Z., Sun, Z., Li, J., Qi, J., and Odeh, I.: Quantitative assessment of the contributions
836 of climate change and human activities on global grassland degradation, *Environ. Earth Sci.*, 72, 4273–4282,
837 <https://doi.org/10.1007/s12665-014-3322-6>, 2014.

838 Gao, Y., Liu, J., Wang, D., An, Y., Ma, H., and Tong, S.: Synergy and trade-off between plant functional traits enhance
839 grassland multifunctionality under grazing exclusion in a semi-arid region, *J. Environ. Manage.*, 373, 123877,
840 <https://doi.org/10.1016/j.jenvman.2024.123877>, 2025.

841 García Bravo, N., Dimas, M., Murillo Díaz, J. M., Barranco Sanz, L. M., and Ruiz Hernández, J. M.: Ejemplo de
842 aplicación del nuevo modelo hidrogeológico en SIMPA para mejorar la evaluación de los recursos hídricos a escala
843 nacional en España, Centro Nacional de Información Geográfica (España), 2023.

844 Gaudet, C. L. and Keddy, P. A.: A comparative approach to predicting competitive ability from plant traits, *Nature*, 334,
845 242–243, <https://doi.org/10.1038/334242a0>, 1988.

846 Georgiou, K., Jackson, R. B., Vindušková, O., Abramoff, R. Z., Ahlström, A., Feng, W., Harden, J. W., Pellegrini, A. F.
847 A., Polley, H. W., Soong, J. L., Riley, W. J., and Torn, M. S.: Global stocks and capacity of mineral-associated soil organic
848 carbon, *Nat. Commun.*, 13, 3797, <https://doi.org/10.1038/s41467-022-31540-9>, 2022.

849 Georgiou, K., Koven, C. D., Wieder, W. R., Hartman, M. D., Riley, W. J., Pett-Ridge, J., Bouskill, N. J., Abramoff, R. Z.,
850 Slessarev, E. W., Ahlström, A., Parton, W. J., Pellegrini, A. F. A., Pierson, D., Sulman, B. N., Zhu, Q., and Jackson, R. B.:
851 Emergent temperature sensitivity of soil organic carbon driven by mineral associations, *Nat. Geosci.*, 17, 205–212,
852 <https://doi.org/10.1038/s41561-024-01384-7>, 2024.

853 Georgiou, K., Angers, D., Champigny, R. E., Cotrufo, M. F., Craig, M. E., Doetterl, S., Grandy, A. S., Lavalley, J. M., Lin,
854 Y., Lugato, E., Poeplau, C., Rocci, K. S., Schweizer, S. A., Six, J., and Wieder, W. R.: Soil Carbon Saturation: What Do
855 We Really Know?, *Glob. Change Biol.*, 31, e70197, <https://doi.org/10.1111/gcb.70197>, 2025.

856 Gliksman, D., Navon, Y., Dumbur, R., Haenel, S., and Grünzweig, J. M.: Higher rates of decomposition in standing vs.
857 surface litter in a Mediterranean ecosystem during the dry and the wet seasons, *Plant Soil*, 428, 427–439,
858 <https://doi.org/10.1007/s11104-018-3696-4>, 2018.

859 Godde, C. M., Boone, R. B., Ash, A. J., Waha, K., Sloat, L. L., Thornton, P. K., and Herrero, M.: Global rangeland
860 production systems and livelihoods at threat under climate change and variability, *Environ. Res. Lett.*, 15, 044021,
861 <https://doi.org/10.1088/1748-9326/ab7395>, 2020.

862 Godínez-Alvarez, H., Herrick, J. E., Mattocks, M., Toledo, D., and Van Zee, J.: Comparison of three vegetation monitoring
863 methods: Their relative utility for ecological assessment and monitoring, *Ecol. Indic.*, 9, 1001–1008,
864 <https://doi.org/10.1016/j.ecolind.2008.11.011>, 2009.

865 Gómez-Rey, M. X., Garcês, A., and Madeira, M.: Soil organic-C accumulation and N availability under improved pastures
866 established in Mediterranean oak woodlands, *Soil Use Manag.*, 28, 497–507, <https://doi.org/10.1111/j.1475-2743.2012.00428.x>, 2012.

868 Gou, X., Reich, P. B., Qiu, L., Shao, M., Wei, G., Wang, J., and Wei, X.: Leguminous plants significantly increase soil
869 nitrogen cycling across global climates and ecosystem types, *Glob. Change Biol.*, 29, 4028–4043,
870 <https://doi.org/10.1111/gcb.16742>, 2023.

871 Griepentrog, M., Bodé, S., Boeckx, P., Hagedorn, F., Heim, A., and Schmidt, M. W. I.: Nitrogen deposition promotes the
872 production of new fungal residues but retards the decomposition of old residues in forest soil fractions, *Glob. Change*
873 *Biol.*, 20, 327–340, <https://doi.org/10.1111/gcb.12374>, 2014.

874 Guyonnet, J. P., Cantarel, A. A. M., Simon, L., and Haichar, F. el Z.: Root exudation rate as functional trait involved in
875 plant nutrient-use strategy classification, *Ecol. Evol.*, 8, 8573–8581, <https://doi.org/10.1002/ece3.4383>, 2018.

876 Haddix, M. L., Paul, E. A., and Cotrufo, M. F.: Dual, differential isotope labeling shows the preferential movement of
877 labile plant constituents into mineral-bonded soil organic matter, *Glob. Change Biol.*, 22, 2301–2312,
878 <https://doi.org/10.1111/gcb.13237>, 2016.

879 Hamilton, E. W., Frank, D. A., Hinchey, P. M., and Murray, T. R.: Defoliation induces root exudation and triggers positive
880 rhizospheric feedbacks in a temperate grassland, *Soil Biol. Biochem.*, 40, 2865–2873,
881 <https://doi.org/10.1016/j.soilbio.2008.08.007>, 2008.

882 Hansen, P. M., Even, R., King, A. E., Lavalley, J., Schipanski, M., and Cotrufo, M. F.: Distinct, direct and climate-
883 mediated environmental controls on global particulate and mineral-associated organic carbon storage, *Glob. Change Biol.*,
884 30, e17080, <https://doi.org/10.1111/gcb.17080>, 2024.

885 Hawkins, H.-J.: A global assessment of Holistic Planned Grazing™ compared with season-long, continuous grazing:
886 meta-analysis findings, *Afr. J. Range Forage Sci.*, 34, 65–75, <https://doi.org/10.2989/10220119.2017.1358213>, 2017.

887 He, M., Zhou, G., Yuan, T., van Groenigen, K. J., Shao, J., and Zhou, X.: Grazing intensity significantly changes the C :
888 N : P stoichiometry in grassland ecosystems, *Glob. Ecol. Biogeogr.*, 29, 355–369, <https://doi.org/10.1111/geb.13028>,
889 2020.

890 Helsen, K., Ceulemans, T., Stevens, C. J., and Honnay, O.: Increasing Soil Nutrient Loads of European Semi-natural
891 Grasslands Strongly Alter Plant Functional Diversity Independently of Species Loss, *Ecosystems*, 17, 169–181,
892 <https://doi.org/10.1007/s10021-013-9714-8>, 2014.

893 den Herder, M., Moreno, G., Mosquera-Losada, R. M., Palma, J. H. N., Sidiropoulou, A., Santiago Freijanes, J. J., Crous-
894 Duran, J., Paulo, J. A., Tomé, M., Pantera, A., Papanastasis, V. P., Mantzanas, K., Pachana, P., Papadopoulos, A.,
895 Plieninger, T., and Burgess, P. J.: Current extent and stratification of agroforestry in the European Union, *Agric. Ecosyst.*
896 *Environ.*, 241, 121–132, <https://doi.org/10.1016/j.agee.2017.03.005>, 2017.

897 Hernández-Esteban, A., López-Díaz, M. L., Cáceres, Y., and Moreno, G.: Are sown legume-rich pastures effective allies
898 for the profitability and sustainability of Mediterranean dehesas?, *Agrofor. Syst.*, 93, 2047–2065,
899 <https://doi.org/10.1007/s10457-018-0307-6>, 2019.

900 Hou, Z., Wang, R., Chang, S., Zheng, Y., Ma, T., Xu, S., Zhang, X., Shi, X., Lu, J., Luo, D., Wang, B., Du, Z., and Wei,
901 Y.: The contribution of microbial necromass to soil organic carbon and influencing factors along a variation of habitats in
902 alpine ecosystems, *Sci. Total Environ.*, 921, 171126, <https://doi.org/10.1016/j.scitotenv.2024.171126>, 2024.

903 Hu, Y., Deng, Q., Kätterer, T., Olesen, J. E., Ying, S. C., Ochoa-Hueso, R., Mueller, C. W., Weintraub, M. N., and Chen,
904 J.: Depth-dependent responses of soil organic carbon under nitrogen deposition, *Glob. Change Biol.*, 30, e17247,
905 <https://doi.org/10.1111/gcb.17247>, 2024.

906 Huang, J., Liu, W., Pan, S., Wang, Z., Yang, S., Jia, Z., Wang, Z., Deng, M., Yang, L., Liu, C., Chang, P., and Liu, L.:
907 Divergent contributions of living roots to turnover of different soil organic carbon pools and their links to plant traits,
908 *Funct. Ecol.*, 35, 2821–2830, <https://doi.org/10.1111/1365-2435.13934>, 2021.

909 Intergovernmental Panel on Climate Change (IPCC) (Ed.): Climate Change Information for Regional Impact and for Risk
910 Assessment, in: Climate Change 2021 – The Physical Science Basis: Working Group I Contribution to the Sixth
911 Assessment Report of the Intergovernmental Panel on Climate Change, Cambridge University Press, Cambridge, 1767–
912 1926, <https://doi.org/10.1017/9781009157896.014>, 2023a.

913 Intergovernmental Panel on Climate Change (IPCC) (Ed.): Global Carbon and Other Biogeochemical Cycles and
914 Feedbacks, in: Climate Change 2021 – The Physical Science Basis: Working Group I Contribution to the Sixth Assessment
915 Report of the Intergovernmental Panel on Climate Change, Cambridge University Press, Cambridge, 673–816,
916 <https://doi.org/10.1017/9781009157896.007>, 2023b.

917 Jacobo, E. J., Rodríguez, A. M., Bartoloni, N., and Deregibus, V. A.: Rotational Grazing Effects on Rangeland Vegetation
918 at a Farm Scale, *Rangel. Ecol. Manag.*, 59, 249–257, <https://doi.org/10.2111/05-129R1.1>, 2006.

919 Jaurena, M., Lezama, F., Salvo, L., Cardozo, G., Ayala, W., Terra, J., and Nabinger, C.: The Dilemma of Improving Native
920 Grasslands by Overseeding Legumes: Production Intensification or Diversity Conservation, *Rangel. Ecol. Manag.*, 69,
921 35–42, <https://doi.org/10.1016/j.rama.2015.10.006>, 2016.

922 Jongen, M., Förster, A. C., and Unger, S.: Overwhelming effects of autumn-time drought during seedling establishment
923 impair recovery potential in sown and semi-natural pastures in Portugal, *Plant Ecol.*, 220, 183–197,
924 <https://doi.org/10.1007/s11258-018-0869-4>, 2019.

925 Kallenbach, C. M., Frey, S. D., and Grandy, A. S.: Direct evidence for microbial-derived soil organic matter formation
926 and its ecophysiological controls, *Nat. Commun.*, 7, 13630, <https://doi.org/10.1038/ncomms13630>, 2016.

927 Kazakou, E., Violle, C., Roumet, C., Pintor, C., Gimenez, O., and Garnier, E.: Litter quality and decomposability of
928 species from a Mediterranean succession depend on leaf traits but not on nitrogen supply, *Ann. Bot.*, 104, 1151–1161,
929 <https://doi.org/10.1093/aob/mcp202>, 2009.

930 Khatiwada, B., Acharya, S. N., Larney, F. J., Lupwayi, N. Z., Smith, E. G., Islam, M. A., and Thomas, J. E.: Benefits of
931 mixed grass–legume pastures and pasture rejuvenation using bloat-free legumes in western Canada: a review, *Can. J.*
932 *Plant Sci.*, 100, 463–476, <https://doi.org/10.1139/cjps-2019-0212>, 2020.

933 Khatri-Chhetri, U., Thompson, K. A., Quideau, S. A., Boyce, M. S., Chang, S. X., Bork, E. W., and Carlyle, C. N.:
934 Adaptive multi-paddock grazing increases mineral associated soil carbon in Northern grasslands, *Agric. Ecosyst.*
935 *Environ.*, 369, 109000, <https://doi.org/10.1016/j.agee.2024.109000>, 2024.

King, A. E., Amsili, J. P., Córdova, S. C., Culman, S., Fonte, S. J., Kotcon, J., Liebig, M., Masters, M. D., McVay, K., Olk, D. C., Schipanski, M., Schneider, S. K., Stewart, C. E., and Cotrufo, M. F.: A soil matrix capacity index to predict mineral-associated but not particulate organic carbon across a range of climate and soil pH, *Biogeochemistry*, 165, 1–14, <https://doi.org/10.1007/s10533-023-01066-3>, 2023.

Klump, K., Fontaine, S., Attard, E., Le Roux, X., Gleixner, G., and Soussana, J.-F.: Grazing triggers soil carbon loss by altering plant roots and their control on soil microbial community, *J. Ecol.*, 97, 876–885, <https://doi.org/10.1111/j.1365-2745.2009.01549.x>, 2009.

Kramer, C. and Gleixner, G.: Soil organic matter in soil depth profiles: Distinct carbon preferences of microbial groups during carbon transformation, *Soil Biol. Biochem.*, 40, 425–433, <https://doi.org/10.1016/j.soilbio.2007.09.016>, 2008.

Kramer-Walter, K. R., Bellingham, P. J., Millar, T. R., Smissen, R. D., Richardson, S. J., and Laughlin, D. C.: Root traits are multidimensional: specific root length is independent from root tissue density and the plant economic spectrum, *J. Ecol.*, 104, 1299–1310, <https://doi.org/10.1111/1365-2745.12562>, 2016.

Laliberté, E. and Legendre, P.: A distance-based framework for measuring functional diversity from multiple traits, *Ecology*, 91, 299–305, <https://doi.org/10.1890/08-2244.1>, 2010.

Laliberté, E. and Tylianakis, J. M.: Cascading effects of long-term land-use changes on plant traits and ecosystem functioning, *Ecology*, 93, 145–155, <https://doi.org/10.1890/11-0338.1>, 2012.

Lange, M., Eisenhauer, N., Sierra, C. A., Bessler, H., Engels, C., Griffiths, R. I., Mellado-Vázquez, P. G., Malik, A. A., Roy, J., Scheu, S., Steinbeiss, S., Thomson, B. C., Trumbore, S. E., and Gleixner, G.: Plant diversity increases soil microbial activity and soil carbon storage, *Nat. Commun.*, 6, 6707, <https://doi.org/10.1038/ncomms7707>, 2015.

Latimer, G. W., Jr. (Ed.): *Official Methods of Analysis of AOAC INTERNATIONAL*, Oxford University Press, <https://doi.org/10.1093/9780197610145.001.0001>, 2023.

Lavallee, J. M., Conant, R. T., Paul, E. A., and Cotrufo, M. F.: Incorporation of shoot versus root-derived ¹³C and ¹⁵N into mineral-associated organic matter fractions: results of a soil slurry incubation with dual-labelled plant material, *Biogeochemistry*, 137, 379–393, <https://doi.org/10.1007/s10533-018-0428-z>, 2018.

Lavallee, J. M., Soong, J. L., and Cotrufo, M. F.: Conceptualizing soil organic matter into particulate and mineral-associated forms to address global change in the 21st century, *Glob. Change Biol.*, 26, 261–273, <https://doi.org/10.1111/gcb.14859>, 2020.

Lefcheck, J. S.: piecewiseSEM: Piecewise structural equation modelling in R for ecology, evolution, and systematics, *Methods Ecol. Evol.*, 7, 573–579, <https://doi.org/10.1111/2041-210X.12512>, 2016.

Lei, J., Guo, X., Zeng, Y., Zhou, J., Gao, Q., and Yang, Y.: Temporal changes in global soil respiration since 1987, *Nat. Commun.*, 12, 403, <https://doi.org/10.1038/s41467-020-20616-z>, 2021.

Leuthold, S., Lavallee, J. M., Haddix, M. L., and Cotrufo, M. F.: Contrasting properties of soil organic matter fractions isolated by different physical separation methodologies, *Geoderma*, 445, 116870, <https://doi.org/10.1016/j.geoderma.2024.116870>, 2024.

Li, S. and Li, X.: Global understanding of farmland abandonment: A review and prospects, *J. Geogr. Sci.*, 27, 1123–1150, <https://doi.org/10.1007/s11442-017-1426-0>, 2017.

Li, Y., Luo, T., and Lu, Q.: Plant height as a simple predictor of the root to shoot ratio: Evidence from alpine grasslands on the Tibetan Plateau, *J. Veg. Sci.*, 19, 245–252, <https://doi.org/10.3170/2007-8-18365>, 2008.

Liang, C., Amelung, W., Lehmann, J., and Kästner, M.: Quantitative assessment of microbial necromass contribution to soil organic matter, *Glob. Change Biol.*, 25, 3578–3590, <https://doi.org/10.1111/gcb.14781>, 2019.

Liu, H. Y., Huang, N., Zhao, C. M., and Li, J. H.: Responses of carbon cycling and soil organic carbon content to nitrogen addition in grasslands globally, *Soil Biol. Biochem.*, 186, 109164, <https://doi.org/10.1016/j.soilbio.2023.109164>, 2023.

Liu, Y., Zhang, M., Wang, X., and Wang, C.: The impact of different grazing intensity and management measures on soil organic carbon density in Zhangye grassland, *Sci. Rep.*, 14, 17556, <https://doi.org/10.1038/s41598-024-68277-y>, 2024.

Lu, M., Zhou, X., Luo, Y., Yang, Y., Fang, C., Chen, J., and Li, B.: Minor stimulation of soil carbon storage by nitrogen addition: A meta-analysis, *Agric. Ecosyst. Environ.*, 140, 234–244, <https://doi.org/10.1016/j.agee.2010.12.010>, 2011.

- 982 Luo, R., Fan, J., Wang, W., Luo, J., Kuzyakov, Y., He, J.-S., Chu, H., and Ding, W.: Nitrogen and phosphorus enrichment
983 accelerates soil organic carbon loss in alpine grassland on the Qinghai-Tibetan Plateau, *Sci. Total Environ.*, 650, 303–
984 312, <https://doi.org/10.1016/j.scitotenv.2018.09.038>, 2019.
- 985 Luo, R., Kuzyakov, Y., Liu, D., Fan, J., Luo, J., Lindsey, S., He, J.-S., and Ding, W.: Nutrient addition reduces carbon
986 sequestration in a Tibetan grassland soil: Disentangling microbial and physical controls, *Soil Biol. Biochem.*, 144,
987 107764, <https://doi.org/10.1016/j.soilbio.2020.107764>, 2020.
- 988 Maestre, F. T., Le Bagousse-Pinguet, Y., Delgado-Baquerizo, M., Eldridge, D. J., Saiz, H., Berdugo, M., Gozalo, B.,
989 Ochoa, V., Guirado, E., García-Gómez, M., Valencia, E., Gaitán, J. J., Asensio, S., Mendoza, B. J., Plaza, C., Díaz-
990 Martínez, P., Rey, A., Hu, H.-W., He, J.-Z., Wang, J.-T., Lehmann, A., Rillig, M. C., Cesarz, S., Eisenhauer, N., Martínez-
991 Valderrama, J., Moreno-Jiménez, E., Sala, O., Abedi, M., Ahmadian, N., Alados, C. L., Aramayo, V., Amghar, F.,
992 Arredondo, T., Ahumada, R. J., Bahalkeh, K., Ben Salem, F., Blaum, N., Boldgiv, B., Bowker, M. A., Bran, D., Bu, C.,
993 Canessa, R., Castillo-Monroy, A. P., Castro, H., Castro, I., Castro-Quezada, P., Chibani, R., Conceição, A. A., Currier, C.
994 M., Darrouzet-Nardi, A., Deák, B., Donoso, D. A., Dougill, A. J., Durán, J., Erdenetsetseg, B., Espinosa, C. I., Fajardo,
995 A., Farzam, M., Ferrante, D., Frank, A. S. K., Fraser, L. H., Gherardi, L. A., Greenville, A. C., Guerra, C. A., Gusmán-
996 Montalvan, E., Hernández-Hernández, R. M., Hölzel, N., Huber-Sannwald, E., Hughes, F. M., Jadán-Maza, O., Jeltsch,
997 F., Jentsch, A., Kaseke, K. F., Köbel, M., Koopman, J. E., Leder, C. V., Linstädter, A., le Roux, P. C., Li, X., Liancourt,
998 P., Liu, J., Louw, M. A., Maggs-Köling, G., Makhalanyane, T. P., Issa, O. M., Manzaneda, A. J., Marais, E., Mora, J. P.,
999 Moreno, G., Munson, S. M., Nunes, A., Oliva, G., Oñatibia, G. R., Peter, G., Pivari, M. O. D., Pueyo, Y., Quiroga, R. E.,
1000 Rahmanian, S., Reed, S. C., et al.: Grazing and ecosystem service delivery in global drylands, *Science*, 378, 915–920,
1001 <https://doi.org/10.1126/science.abq4062>, 2022.
- 1002 Manning, P., de Vries, F. T., Tallowin, J. R. B., Smith, R., Mortimer, S. R., Pilgrim, E. S., Harrison, K. A., Wright, D. G.,
1003 Quirk, H., Benson, J., Shipley, B., Cornelissen, J. H. C., Kattge, J., Bönisch, G., Wirth, C., and Bardgett, R. D.: Simple
1004 measures of climate, soil properties and plant traits predict national-scale grassland soil carbon stocks, *J. Appl. Ecol.*, 52,
1005 1188–1196, <https://doi.org/10.1111/1365-2664.12478>, 2015.
- 1006 McSherry, M. E. and Ritchie, M. E.: Effects of grazing on grassland soil carbon: a global review, *Glob. Change Biol.*, 19,
1007 1347–1357, <https://doi.org/10.1111/gcb.12144>, 2013.
- 1008 Mokany, K., Ash, J., and Roxburgh, S.: Functional identity is more important than diversity in influencing ecosystem
1009 processes in a temperate native grassland, *J. Ecol.*, 96, 884–893, <https://doi.org/10.1111/j.1365-2745.2008.01395.x>, 2008.
- 1010 Moreno, G. and Pulido, F. J.: The Functioning, Management and Persistence of Dehesas, in: *Agroforestry in Europe:*
1011 *Current Status and Future Prospects*, edited by: Rigueiro-Rodríguez, A., McAdam, J., and Mosquera-Losada, M. R.,
1012 Springer Netherlands, Dordrecht, 127–160, https://doi.org/10.1007/978-1-4020-8272-6_7, 2009.
- 1013 Moreno, G., Obrador, J. J., Cubera, E., and Dupraz, C.: Fine Root Distribution in Dehesas of Central-Western Spain, *Plant*
1014 *Soil*, 277, 153–162, <https://doi.org/10.1007/s11104-005-6805-0>, 2005.
- 1015 Moreno, G., Hernández-Esteban, A., Rolo, V., and Igual, J. M.: The enduring effects of sowing legume-rich mixtures on
1016 the soil microbial community and soil carbon in semi-arid wood pastures, *Plant Soil*, 465, 563–582,
1017 <https://doi.org/10.1007/s11104-021-05023-7>, 2021.
- 1018 Moreno-Jiménez, E., Plaza, C., Saiz, H., Manzano, R., Flagmeier, M., and Maestre, F. T.: Aridity and reduced soil
1019 micronutrient availability in global drylands, *Nat. Sustain.*, 2, 371–377, <https://doi.org/10.1038/s41893-019-0262-x>,
1020 2019.
- 1021 Mortensen, E. Ø., Abalos, D., Engedal, T., Lægsgaard, A. K., Enggrob, K., Mueller, C. W., and Rasmussen, J.: Smart
1022 Mixture Design Can Steer the Fate of Root-Derived Carbon Into Mineral-Associated and Particulate Organic Matter in
1023 Intensively Managed Grasslands, *Glob. Change Biol.*, 31, e70117, <https://doi.org/10.1111/gcb.70117>, 2025.
- 1024 Mosier, S., Apfelbaum, S., Byck, P., Calderon, F., Teague, R., Thompson, R., and Cotrufo, M. F.: Adaptive multi-paddock
1025 grazing enhances soil carbon and nitrogen stocks and stabilization through mineral association in southeastern U.S.
1026 grazing lands, *J. Environ. Manage.*, 288, 112409, <https://doi.org/10.1016/j.jenvman.2021.112409>, 2021.
- 1027 Murphy, M. V.: *semEff: Automatic Calculation of Effects for Piecewise Structural Equation Models*, 2022.
- 1028 Nair, R. K. F., Morris, K. A., Hertel, M., Luo, Y., Moreno, G., Reichstein, M., Schrumpf, M., and Migliavacca, M.:
1029 N:P stoichiometry and habitat effects on Mediterranean savanna seasonal root dynamics,
1030 *Biogeosciences*, 16, 1883–1901, <https://doi.org/10.5194/bg-16-1883-2019>, 2019.

- 1031 Niu, W., Ding, J., Fu, B., Zhao, W., and Eldridge, D.: Global effects of livestock grazing on ecosystem functions vary
1032 with grazing management and environment, *Agric. Ecosyst. Environ.*, 378, 109296,
1033 <https://doi.org/10.1016/j.agee.2024.109296>, 2025.
- 1034 Novelty, P. E. and Watson, I. W.: Successful Grassland Regeneration in a Severely Degraded Catchment: a Whole of
1035 Government Approach in North West Australia, in: *Climate and Land Degradation*, edited by: Sivakumar, M. V. K. and
1036 Ndiang’ui, N., Springer, Berlin, Heidelberg, 469–485, https://doi.org/10.1007/978-3-540-72438-4_26, 2007.
- 1037 Oggioni, S. D., Ochoa-Hueso, R., and Peco, B.: Livestock grazing abandonment reduces soil microbial activity and carbon
1038 storage in a Mediterranean Dehesa, *Appl. Soil Ecol.*, 153, 103588, <https://doi.org/10.1016/j.apsoil.2020.103588>, 2020.
- 1039 Oliveira Filho, J. de S., Vieira, J. N., Ribeiro da Silva, E. M., Beserra de Oliveira, J. G., Pereira, M. G., and Brasileiro, F.
1040 G.: Assessing the effects of 17 years of grazing exclusion in degraded semi-arid soils: Evaluation of soil fertility, nutrients
1041 pools and stoichiometry, *J. Arid Environ.*, 166, 1–10, <https://doi.org/10.1016/j.jaridenv.2019.03.006>, 2019.
- 1042 Ordoñez, J. C., Van Bodegom, P. M., Witte, J.-P. M., Wright, I. J., Reich, P. B., and Aerts, R.: A global study of
1043 relationships between leaf traits, climate and soil measures of nutrient fertility, *Glob. Ecol. Biogeogr.*, 18, 137–149,
1044 <https://doi.org/10.1111/j.1466-8238.2008.00441.x>, 2009.
- 1045 Palomo-Campesino, S., Ravera, F., González, J. A., and García-Llorente, M.: Exploring Current and Future Situation of
1046 Mediterranean Silvopastoral Systems: Case Study in Southern Spain, *Rangel. Ecol. Manag.*, 71, 578–591,
1047 <https://doi.org/10.1016/j.rama.2017.12.013>, 2018.
- 1048 Parras-Alcántara, L., Díaz-Jaimes, L., Lozano-García, B., Rebollo, P. F., Elcure, F. M., and Muñoz, M. D. C.: Organic
1049 farming has little effect on carbon stock in a Mediterranean dehesa (southern Spain), *CATENA*, 113, 9–17,
1050 <https://doi.org/10.1016/j.catena.2013.09.002>, 2014.
- 1051 Parras-Alcántara, L., Díaz-Jaimes, L., and Lozano-García, B.: Management Effects on Soil Organic Carbon Stock in
1052 Mediterranean Open Rangelands—Treeless Grasslands, *Land Degrad. Dev.*, 26, 22–34, <https://doi.org/10.1002/ldr.2269>,
1053 2015.
- 1054 Peco, B., Navarro, E., Carmona, C. P., Medina, N. G., and Marques, M. J.: Effects of grazing abandonment on soil
1055 multifunctionality: The role of plant functional traits, *Agric. Ecosyst. Environ.*, 249, 215–225,
1056 <https://doi.org/10.1016/j.agee.2017.08.013>, 2017.
- 1057 Pérez-Harguindeguy, N., Díaz, S., Garnier, E., Lavorel, S., Poorter, H., Jaureguiberry, P., Bret-Harte, M. S., Cornwell, W.
1058 K., Craine, J. M., Gurvich, D. E., Urcelay, C., Veneklaas, E. J., Reich, P. B., Poorter, L., Wright, I. J., Ray, P., Enrico, L.,
1059 Pausas, J. G., Vos, A. C. de, Buchmann, N., Funes, G., Quétier, F., Hodgson, J. G., Thompson, K., Morgan, H. D., Steege,
1060 H. ter, Sack, L., Blonder, B., Poschlod, P., Vaieretti, M. V., Conti, G., Staver, A. C., Aquino, S., and Cornelissen, J. H. C.:
1061 Corrigendum to: New handbook for standardised measurement of plant functional traits worldwide, *Aust. J. Bot.*, 64,
1062 715–716, https://doi.org/10.1071/bt12225_co, 2016.
- 1063 Phukubye, K., Mutema, M., Buthelezi, N., Muchaonyerwa, P., Cerri, C., and Chaplot, V.: On the impact of grassland
1064 management on soil carbon stocks: a worldwide meta-analysis, *Geoderma Reg.*, 28, e00479,
1065 <https://doi.org/10.1016/j.geodrs.2021.e00479>, 2022.
- 1066 Pino, A. D., Rodríguez, T., and Andión, J.: Production improvement through phosphorus fertilization and legume
1067 introduction in grazed native pastures of Uruguay, *J. Agric. Sci.*, 154, 347–358,
1068 <https://doi.org/10.1017/S002185961500101X>, 2016.
- 1069 Plaza, C., García-Palacios, P., Berhe, A. A., Barquero, J., Bastida, F., Png, G. K., Rey, A., Bardgett, R. D., and Delgado-
1070 Baquerizo, M.: Ecosystem productivity has a stronger influence than soil age on surface soil carbon storage across global
1071 biomes, *Commun. Earth Environ.*, 3, 1–8, <https://doi.org/10.1038/s43247-022-00567-7>, 2022.
- 1072 Poeplau, C., Vos, C., and Don, A.: Soil organic carbon stocks are systematically overestimated by misuse of the parameters
1073 bulk density and rock fragment content, *SOIL*, 3, 61–66, <https://doi.org/10.5194/soil-3-61-2017>, 2017.
- 1074 Poeplau, C., Helfrich, M., Dechow, R., Szoboszlay, M., Tebbe, C. C., Don, A., Greiner, B., Zopf, D., Thumm, U.,
1075 Korevaar, H., and Geerts, R.: Increased microbial anabolism contributes to soil carbon sequestration by mineral
1076 fertilization in temperate grasslands, *Soil Biol. Biochem.*, 130, 167–176, <https://doi.org/10.1016/j.soilbio.2018.12.019>,
1077 2019.
- 1078 Poeplau, C., Dechow, R., Begill, N., and Don, A.: Towards an ecosystem capacity to stabilise organic carbon in soils,
1079 *Glob. Change Biol.*, 30, e17453, <https://doi.org/10.1111/gcb.17453>, 2024.

1080 Poorter, H., Niinemets, Ü., Poorter, L., Wright, I. J., and Villar, R.: Causes and consequences of variation in leaf mass per
1081 area (LMA): a meta-analysis, *New Phytol.*, 182, 565–588, <https://doi.org/10.1111/j.1469-8137.2009.02830.x>, 2009.

1082 Porqueddu, C., Ates, S., Louhaichi, M., Kyriazopoulos, A. P., Moreno, G., del Pozo, A., Ovalle, C., Ewing, M. A., and
1083 Nichols, P. G. H.: Grasslands in ‘Old World’ and ‘New World’ Mediterranean-climate zones: past trends, current status
1084 and future research priorities, *Grass Forage Sci.*, 71, 1–35, <https://doi.org/10.1111/gfs.12212>, 2016.

1085 Pulina, A., Rolo, V., Hernández-Esteban, A., Seddaiu, G., Roggero, P. P., and Moreno, G.: Long-term legacy of sowing
1086 legume-rich mixtures in Mediterranean wooded grasslands, *Agric. Ecosyst. Environ.*, 348, 108397,
1087 <https://doi.org/10.1016/j.agee.2023.108397>, 2023.

1088 R Core Team: R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna,
1089 Austria. URL <https://www.R-project.org/>, 2024.

1090 Rama, G., Oyarzabal, M., Cardozo, G., Lezama, F., and Baeza, S.: Legume Overseeding along with P Fertilization
1091 Increase Forage Production of Temperate Natural Grasslands, *Agronomy*, 12, 2507,
1092 <https://doi.org/10.3390/agronomy12102507>, 2022.

1093 Ricotta, C. and Moretti, M.: CWM and Rao’s quadratic diversity: a unified framework for functional ecology, *Oecologia*,
1094 167, 181–188, <https://doi.org/10.1007/s00442-011-1965-5>, 2011.

1095 Ridgeway, J. R., Morrissey, E. M., and Brzostek, E. R.: Plant litter traits control microbial decomposition and drive soil
1096 carbon stabilization, *Soil Biol. Biochem.*, 175, 108857, <https://doi.org/10.1016/j.soilbio.2022.108857>, 2022.

1097 Rocci, K. S., Lavalley, J. M., Stewart, C. E., and Cotrufo, M. F.: Soil organic carbon response to global environmental
1098 change depends on its distribution between mineral-associated and particulate organic matter: A meta-analysis, *Sci. Total*
1099 *Environ.*, 793, 148569, <https://doi.org/10.1016/j.scitotenv.2021.148569>, 2021.

1100 Rocci, K. S., Barker, K. S., Seabloom, E. W., Borer, E. T., Hobbie, S. E., Bakker, J. D., MacDougall, A. S., McCulley, R.
1101 L., Moore, J. L., Raynaud, X., Stevens, C. J., and Cotrufo, M. F.: Impacts of nutrient addition on soil carbon and nitrogen
1102 stoichiometry and stability in globally-distributed grasslands, *Biogeochemistry*, 159, 353–370,
1103 <https://doi.org/10.1007/s10533-022-00932-w>, 2022.

1104 Rolo, V., Rivest, D., Lorente, M., Kattge, J., and Moreno, G.: Taxonomic and functional diversity in Mediterranean
1105 pastures: insights on the biodiversity–productivity trade-off, *J. Appl. Ecol.*, 53, 1575–1584, <https://doi.org/10.1111/1365-2664.12685>, 2016.

1107 Roumet, C., Birouste, M., Picon-Cochard, C., Ghestem, M., Osman, N., Vrignon-Brenas, S., Cao, K., and Stokes, A.:
1108 Root structure–function relationships in 74 species: evidence of a root economics spectrum related to carbon economy,
1109 *New Phytol.*, 210, 815–826, <https://doi.org/10.1111/nph.13828>, 2016.

1110 Rovira, P., Sauras-Yera, T., and Romanyà, J.: Equivalent-mass versus fixed-depth as criteria for quantifying soil carbon
1111 sequestration: How relevant is the difference?, *CATENA*, 214, 106283, <https://doi.org/10.1016/j.catena.2022.106283>,
1112 2022.

1113 Sanaullah, M., Chabbi, A., Lemaire, G., Charrier, X., and Rumpel, C.: How does plant leaf senescence of grassland species
1114 influence decomposition kinetics and litter compounds dynamics?, *Nutr. Cycl. Agroecosystems*, 88, 159–171,
1115 <https://doi.org/10.1007/s10705-009-9323-2>, 2010.

1116 Sanderman, J., Baldock, J. A., Dangal, S. R. S., Ludwig, S., Potter, S., Rivard, C., and Savage, K.: Soil organic carbon
1117 fractions in the Great Plains of the United States: an application of mid-infrared spectroscopy, *Biogeochemistry*, 156, 97–
1118 114, <https://doi.org/10.1007/s10533-021-00755-1>, 2021.

1119 Sanjari, G., Yu, B., Ghadiri, H., Ciesiolka, C. A. A., and Rose, C. W.: Effects of time-controlled grazing on runoff and
1120 sediment loss, *Soil Res.*, 47, 796–808, <https://doi.org/10.1071/SR09032>, 2009.

1121 Santos, R. S., Hamilton, E. K., Stanley, P. L., Paustian, K., Cotrufo, M. F., and Zhang, Y.: Simulating adaptive grazing
1122 management on soil organic carbon in the Southeast U.S.A. using MEMS 2, *J. Environ. Manage.*, 365, 121657,
1123 <https://doi.org/10.1016/j.jenvman.2024.121657>, 2024.

1124 Sherrod, L. A., Dunn, G., Peterson, G. A., and Kolberg, R. L.: Inorganic Carbon Analysis by Modified Pressure-
1125 Calcimeter Method, *Soil Sci. Soc. Am. J.*, 66, 299–305, <https://doi.org/10.2136/sssaj2002.2990>, 2002.

- 1126 Shi, B., Delgado-Baquerizo, M., Knapp, A. K., Smith, M. D., Reed, S., Osborne, B., Carrillo, Y., Maestre, F. T., Zhu, Y.,
1127 Chen, A., Wilkins, K., Holdrege, M. C., Kulmatiski, A., Picon-Cochard, C., Roscher, C., Power, S., Byrne, K. M.,
1128 Churchill, A. C., Jentsch, A., Henry, H. A. L., Beard, K. H., Schuchardt, M. A., Eisenhauer, N., Otfinowski, R., Hautier,
1129 Y., Shen, H., Wang, Y., Wang, Z., Wang, C., Cusack, D. F., Petraglia, A., Carbognani, M., Forte, T. G. W., Flory, S., Hou,
1130 P., Zhang, T., Gao, W., and Sun, W.: Aridity drives the response of soil total and particulate organic carbon to drought in
1131 temperate grasslands and shrublands, *Sci. Adv.*, 10, eadq2654, <https://doi.org/10.1126/sciadv.adq2654>, 2024.
- 1132 Siefert, A., Violle, C., Chalmandrier, L., Albert, C. H., Taudiere, A., Fajardo, A., Aarssen, L. W., Baraloto, C., Carlucci,
1133 M. B., Cianciaruso, M. V., de L. Dantas, V., de Bello, F., Duarte, L. D. S., Fonseca, C. R., Freschet, G. T., Gaucherand,
1134 S., Gross, N., Hikosaka, K., Jackson, B., Jung, V., Kamiyama, C., Katabuchi, M., Kembel, S. W., Kichenin, E., Kraft, N.
1135 J. B., Lagerström, A., Bagousse-Pinguet, Y. L., Li, Y., Mason, N., Messier, J., Nakashizuka, T., Overton, J. McC., Peltzer,
1136 D. A., Pérez-Ramos, I. M., Pillar, V. D., Prentice, H. C., Richardson, S., Sasaki, T., Schamp, B. S., Schöb, C., Shipley, B.,
1137 Sundqvist, M., Sykes, M. T., Vandewalle, M., and Wardle, D. A.: A global meta-analysis of the relative extent of
1138 intraspecific trait variation in plant communities, *Ecol. Lett.*, 18, 1406–1419, <https://doi.org/10.1111/ele.12508>, 2015.
- 1139 Six, J., Guggenberger, G., Paustian, K., Haumaier, L., Elliott, E. T., and Zech, W.: Sources and composition of soil organic
1140 matter fractions between and within soil aggregates, *Eur. J. Soil Sci.*, 52, 607–618, <https://doi.org/10.1046/j.1365-2389.2001.00406.x>, 2001.
- 1142 Six, J., Conant, R. T., Paul, E. A., and Paustian, K.: Stabilization mechanisms of soil organic matter: Implications for C-
1143 saturation of soils, *Plant Soil*, 241, 155–176, <https://doi.org/10.1023/A:1016125726789>, 2002.
- 1144 Smith, M. D., Wilkins, K. D., Holdrege, M. C., Wilfahrt, P., Collins, S. L., Knapp, A. K., Sala, O. E., Dukes, J. S., Phillips,
1145 R. P., Yahdjian, L., Gherardi, L. A., Ohlert, T., Beier, C., Fraser, L. H., Jentsch, A., Loik, M. E., Maestre, F. T., Power, S.
1146 A., Yu, Q., Felton, A. J., Munson, S. M., Luo, Y., Abdoli, H., Abedi, M., Alados, C. L., Alberti, J., Alon, M., An, H.,
1147 Anacker, B., Anderson, M., Auge, H., Bachle, S., Bahalkeh, K., Bahn, M., Batbaatar, A., Bauerle, T., Beard, K. H., Behn,
1148 K., Beil, I., Biancari, L., Blindow, I., Bondaruk, V. F., Borer, E. T., Bork, E. W., Bruschetti, C. M., Byrne, K. M., Cahill
1149 Jr., J. F., Calvo, D. A., Carbognani, M., Cardoni, A., Carlyle, C. N., Castillo-Garcia, M., Chang, S. X., Chieppa, J.,
1150 Cianciaruso, M. V., Cohen, O., Cordeiro, A. L., Cusack, D. F., Dahlke, S., Daleo, P., D’Antonio, C. M., Dietterich, L. H.,
1151 S. Doherty, T., Dubbert, M., Ebeling, A., Eisenhauer, N., Fischer, F. M., Forte, T. G. W., Gebauer, T., Gozalo, B.,
1152 Greenville, A. C., Guidoni-Martins, K. G., Hannusch, H. J., Vatsø Haugum, S., Hautier, Y., Hefting, M., Henry, H. A. L.,
1153 Hoss, D., Ingrisch, J., Iribarne, O., Isbell, F., Johnson, Y., Jordan, S., Kelly, E. F., Kimmel, K., Kreyling, J., Kröel-Dulay,
1154 G., Kröpfl, A., Kübert, A., Kulmatiski, A., Lamb, E. G., Larsen, K. S., Larson, J., Lawson, J., Leder, C. V., Linstädter, A.,
1155 Liu, J., Liu, S., Lodge, A. G., et al.: Extreme drought impacts have been underestimated in grasslands and shrublands
1156 globally, *Proc. Natl. Acad. Sci.*, 121, e2309881120, <https://doi.org/10.1073/pnas.2309881120>, 2024.
- 1157 Soliveres, S., Maestre, F. T., Eldridge, D. J., Delgado-Baquerizo, M., Quero, J. L., Bowker, M. A., and Gallardo, A.: Plant
1158 diversity and ecosystem multifunctionality peak at intermediate levels of woody cover in global drylands, *Glob. Ecol.*
1159 *Biogeogr.*, 23, 1408–1416, <https://doi.org/10.1111/geb.12215>, 2014.
- 1160 Soussana, J.-F. and Lemaire, G.: Coupling carbon and nitrogen cycles for environmentally sustainable intensification of
1161 grasslands and crop-livestock systems, *Agric. Ecosyst. Environ.*, 190, 9–17, <https://doi.org/10.1016/j.agee.2013.10.012>,
1162 2014.
- 1163 Spohn, M., Pötsch, E. M., Eichorst, S. A., Wobken, D., Wanek, W., and Richter, A.: Soil microbial carbon use efficiency
1164 and biomass turnover in a long-term fertilization experiment in a temperate grassland, *Soil Biol. Biochem.*, 97, 168–175,
1165 <https://doi.org/10.1016/j.soilbio.2016.03.008>, 2016.
- 1166 Stanley, P., Roche, L., and Bowles, T.: Amping up soil carbon: soil carbon stocks in California rangelands under adaptive
1167 multi-paddock and conventional grazing management, *Int. J. Agric. Sustain.*, 2025.
- 1168 Stanley, P. L., Wilson, C., Patterson, E., Machmuller, M. B., and Cotrufo, M. F.: Ruminating on soil carbon: Applying
1169 current understanding to inform grazing management, *Glob. Change Biol.*, 30, e17223, <https://doi.org/10.1111/gcb.17223>,
1170 2024.
- 1171 Steinbeiss, S., BEßLER, H., Engels, C., Temperton, V. M., Buchmann, N., Roscher, C., Kreutziger, Y., Baade, J.,
1172 Habekost, M., and Gleixner, G.: Plant diversity positively affects short-term soil carbon storage in experimental
1173 grasslands, *Glob. Change Biol.*, 14, 2937–2949, <https://doi.org/10.1111/j.1365-2486.2008.01697.x>, 2008.
- 1174 Strickland, M. S. and Rousk, J.: Considering fungal:bacterial dominance in soils – Methods, controls, and ecosystem
1175 implications, *Soil Biol. Biochem.*, 42, 1385–1395, <https://doi.org/10.1016/j.soilbio.2010.05.007>, 2010.

- 1176 Tang, K., Wu, C., Wang, S., Liao, W., Yin, L., Zhou, W., and Cui, H.-J.: Distribution characteristics of soil organic carbon
1177 fractions in paddy profiles with 40 years of fertilization under two groundwater levels, *J. Soils Sediments*, 24, 681–691,
1178 <https://doi.org/10.1007/s11368-023-03664-y>, 2024.
- 1179 Tang, Z., Feng, J., Chen, L., Chen, Z., Shao, X., and Xia, T.: Coupling amendment of microbial and compound fertilizers
1180 increases fungal necromass carbon and soil organic carbon by regulating microbial activity in flue-cured tobacco-planted
1181 field, *Eur. J. Soil Biol.*, 117, 103518, <https://doi.org/10.1016/j.ejsobi.2023.103518>, 2023.
- 1182 Tao, F., Huang, Y., Hungate, B. A., Manzoni, S., Frey, S. D., Schmidt, M. W. I., Reichstein, M., Carvalhais, N., Ciais, P.,
1183 Jiang, L., Lehmann, J., Wang, Y.-P., Houlton, B. Z., Ahrens, B., Mishra, U., Hugelius, G., Hocking, T. D., Lu, X., Shi, Z.,
1184 Viatkin, K., Vargas, R., Yigini, Y., Omuto, C., Malik, A. A., Peralta, G., Cuevas-Corona, R., Di Paolo, L. E., Luotto, I.,
1185 Liao, C., Liang, Y.-S., Saynes, V. S., Huang, X., and Luo, Y.: Microbial carbon use efficiency promotes global soil carbon
1186 storage, *Nature*, 618, 981–985, <https://doi.org/10.1038/s41586-023-06042-3>, 2023.
- 1187 Tate, K. W., Atwill, E. R., McDougald, N. K., and George, M. R.: Spatial and Temporal Patterns of Cattle Feces Deposition
1188 on Rangeland, *J. Range Manag.*, 56, 432–438, <https://doi.org/10.2307/4003833>, 2003.
- 1189 Teague, R., Provenza, F., Kreuter, U., Steffens, T., and Barnes, M.: Multi-paddock grazing on rangelands: Why the
1190 perceptual dichotomy between research results and rancher experience?, *J. Environ. Manage.*, 128, 699–717,
1191 <https://doi.org/10.1016/j.jenvman.2013.05.064>, 2013.
- 1192 Teague, W. R., Dowhower, S. L., and Waggoner, J. A.: Drought and grazing patch dynamics under different grazing
1193 management, *J. Arid Environ.*, 58, 97–117, [https://doi.org/10.1016/S0140-1963\(03\)00122-8](https://doi.org/10.1016/S0140-1963(03)00122-8), 2004.
- 1194 Teague, W. R., Dowhower, S. L., Baker, S. A., Haile, N., DeLaune, P. B., and Conover, D. M.: Grazing management
1195 impacts on vegetation, soil biota and soil chemical, physical and hydrological properties in tall grass prairie, *Agric.*
1196 *Ecosyst. Environ.*, 141, 310–322, <https://doi.org/10.1016/j.agee.2011.03.009>, 2011.
- 1197 Teixeira, R. F. M., Proença, V., Crespo, D., Valada, T., and Domingos, T.: A conceptual framework for the analysis of
1198 engineered biodiverse pastures, *Ecol. Eng.*, 77, 85–97, <https://doi.org/10.1016/j.ecoleng.2015.01.002>, 2015.
- 1199 Thomson, F. J., Moles, A. T., Auld, T. D., and Kingsford, R. T.: Seed dispersal distance is more strongly correlated with
1200 plant height than with seed mass, *J. Ecol.*, 99, 1299–1307, <https://doi.org/10.1111/j.1365-2745.2011.01867.x>, 2011.
- 1201 Tiessen, H., Cuevas, E., and Chacon, P.: The role of soil organic matter in sustaining soil fertility, *Nature*, 371, 783–785,
1202 <https://doi.org/10.1038/371783a0>, 1994.
- 1203 Uribe, C., Inclán, R., Hernando, L., Román, M., Clavero, M. A., Roig, S., and Van Miegroet, H.: Grazing, tilling and
1204 canopy effects on carbon dioxide fluxes in a Spanish dehesa, *Agrofor. Syst.*, 89, 305–318, [https://doi.org/10.1007/s10457-](https://doi.org/10.1007/s10457-014-9767-5)
1205 [014-9767-5](https://doi.org/10.1007/s10457-014-9767-5), 2015.
- 1206 Van Soest, P. J., Robertson, J. B., and Lewis, B. A.: Methods for Dietary Fiber, Neutral Detergent Fiber, and Nonstarch
1207 Polysaccharides in Relation to Animal Nutrition, *J. Dairy Sci.*, 74, 3583–3597, [https://doi.org/10.3168/jds.S0022-](https://doi.org/10.3168/jds.S0022-0302(91)78551-2)
1208 [0302\(91\)78551-2](https://doi.org/10.3168/jds.S0022-0302(91)78551-2), 1991.
- 1209 Villarino, S. H., Pinto, P., Jackson, R. B., and Piñeiro, G.: Plant rhizodeposition: A key factor for soil organic matter
1210 formation in stable fractions, *Sci. Adv.*, 7, eabd3176, <https://doi.org/10.1126/sciadv.abd3176>, 2021.
- 1211 di Virgilio, A., Lambertucci, S. A., and Morales, J. M.: Sustainable grazing management in rangelands: Over a century
1212 searching for a silver bullet, *Agric. Ecosyst. Environ.*, 283, 106561, <https://doi.org/10.1016/j.agee.2019.05.020>, 2019.
- 1213 Waldrop, M. P. and Firestone, M. K.: Microbial community utilization of recalcitrant and simple carbon compounds:
1214 impact of oak-woodland plant communities, *Oecologia*, 138, 275–284, <https://doi.org/10.1007/s00442-003-1419-9>, 2004.
- 1215 Wang, J., Liu, L., Wang, X., and Chen, Y.: The interaction between abiotic photodegradation and microbial decomposition
1216 under ultraviolet radiation, *Glob. Change Biol.*, 21, 2095–2104, <https://doi.org/10.1111/gcb.12812>, 2015.
- 1217 Wang, J., Liu, L., Wang, X., Yang, S., Zhang, B., Li, P., Qiao, C., Deng, M., and Liu, W.: High night-time humidity and
1218 dissolved organic carbon content support rapid decomposition of standing litter in a semi-arid landscape, *Funct. Ecol.*,
1219 31, 1659–1668, <https://doi.org/10.1111/1365-2435.12854>, 2017.
- 1220 Wang, K., Ma, Z., Qin, W., Li, X., Shi, H., Hasi, B., and Liu, X.: Soil nutrients and pH modulate carbon dynamics in
1221 particulate and mineral-associated organic matter during restoration of a Tibetan alpine grassland, *Ecol. Eng.*, 212,
1222 107522, <https://doi.org/10.1016/j.ecoleng.2025.107522>, 2025.

- 1223 Wang, X., McConkey, B. G., VandenBygaart, A. J., Fan, J., Iwaasa, A., and Schellenberg, M.: Grazing improves C and N
1224 cycling in the Northern Great Plains: a meta-analysis, *Sci. Rep.*, 6, 33190, <https://doi.org/10.1038/srep33190>, 2016.
- 1225 Ward, S. E., Smart, S. M., Quirk, H., Tallowin, J. R. B., Mortimer, S. R., Shiel, R. S., Wilby, A., and Bardgett, R. D.:
1226 Legacy effects of grassland management on soil carbon to depth, *Glob. Change Biol.*, 22, 2929–2938,
1227 <https://doi.org/10.1111/gcb.13246>, 2016.
- 1228 Weigelt, A., Mommer, L., Andrzejak, K., Iversen, C. M., Bergmann, J., Bruehlheide, H., Fan, Y., Freschet, G. T., Guerrero-
1229 Ramírez, N. R., Kattge, J., Kuiper, T. W., Laughlin, D. C., Meier, I. C., van der Plas, F., Poorter, H., Roumet, C., van
1230 Ruijven, J., Sabatini, F. M., Semchenko, M., Sweeney, C. J., Valverde-Barrantes, O. J., York, L. M., and McCormack, M.
1231 L.: An integrated framework of plant form and function: the belowground perspective, *New Phytol.*, 232, 42–59,
1232 <https://doi.org/10.1111/nph.17590>, 2021.
- 1233 Westerband, A. C., Funk, J. L., and Barton, K. E.: Intraspecific trait variation in plants: a renewed focus on its role in
1234 ecological processes, *Ann. Bot.*, 127, 397–410, <https://doi.org/10.1093/aob/mcab011>, 2021.
- 1235 White, R.: Pilot analysis of global ecosystems: Grassland ecosystems, 2000.
- 1236 Willers, C., Jansen van Rensburg, P. J., and Claassens, S.: Phospholipid fatty acid profiling of microbial communities—a
1237 review of interpretations and recent applications, *J. Appl. Microbiol.*, 119, 1207–1218, <https://doi.org/10.1111/jam.12902>,
1238 2015.
- 1239 Wilson, C. H., Strickland, M. S., Hutchings, J. A., Bianchi, T. S., and Flory, S. L.: Grazing enhances belowground carbon
1240 allocation, microbial biomass, and soil carbon in a subtropical grassland, *Glob. Change Biol.*, 24, 2997–3009,
1241 <https://doi.org/10.1111/gcb.14070>, 2018.
- 1242 Wilson, P. J., Thompson, K., and Hodgson, J. G.: Specific leaf area and leaf dry matter content as alternative predictors
1243 of plant strategies, *New Phytol.*, 143, 155–162, <https://doi.org/10.1046/j.1469-8137.1999.00427.x>, 1999.
- 1244 Wright, I. J., Reich, P. B., Westoby, M., Ackerly, D. D., Baruch, Z., Bongers, F., Cavender-Bares, J., Chapin, T.,
1245 Cornelissen, J. H. C., Diemer, M., Flexas, J., Garnier, E., Groom, P. K., Gulias, J., Hikosaka, K., Lamont, B. B., Lee, T.,
1246 Lee, W., Lusk, C., Midgley, J. J., Navas, M.-L., Niinemets, Ü., Oleksyn, J., Osada, N., Poorter, H., Poot, P., Prior, L.,
1247 Pyankov, V. I., Roumet, C., Thomas, S. C., Tjoelker, M. G., Veneklaas, E. J., and Villar, R.: The worldwide leaf economics
1248 spectrum, *Nature*, 428, 821–827, <https://doi.org/10.1038/nature02403>, 2004.
- 1249 Xu, H., Zhu, B., Wei, X., Yu, M., and Cheng, X.: Root functional traits mediate rhizosphere soil carbon stability in a
1250 subtropical forest, *Soil Biol. Biochem.*, 162, 108431, <https://doi.org/10.1016/j.soilbio.2021.108431>, 2021.
- 1251 Xu, X., Thornton, P. E., and Post, W. M.: A global analysis of soil microbial biomass carbon, nitrogen and phosphorus in
1252 terrestrial ecosystems, *Glob. Ecol. Biogeogr.*, 22, 737–749, <https://doi.org/10.1111/geb.12029>, 2013.
- 1253 Yao, H., He, Z., Wilson, M. J., and Campbell, C. D.: Microbial Biomass and Community Structure in a Sequence of Soils
1254 with Increasing Fertility and Changing Land Use, *Microb. Ecol.*, 40, 223–237, <https://doi.org/10.1007/s002480000053>,
1255 2000.
- 1256 Yu, L., Sun, W., and Huang, Y.: Grazing exclusion enhances plant and topsoil carbon stocks in arid and semiarid
1257 grasslands, *Agric. Ecosyst. Environ.*, 320, 107605, <https://doi.org/10.1016/j.agee.2021.107605>, 2021.
- 1258 Yu, W., Huang, W., Weintraub-Leff, S. R., and Hall, S. J.: Where and why do particulate organic matter (POM) and
1259 mineral-associated organic matter (MAOM) differ among diverse soils?, *Soil Biol. Biochem.*, 172, 108756,
1260 <https://doi.org/10.1016/j.soilbio.2022.108756>, 2022.
- 1261 Zhang, Q., Buyantuev, A., Li, F. Y., Jiang, L., Niu, J., Ding, Y., Kang, S., and Ma, W.: Functional dominance rather than
1262 taxonomic diversity and functional diversity mainly affects community aboveground biomass in the Inner Mongolia
1263 grassland, *Ecol. Evol.*, 7, 1605–1615, <https://doi.org/10.1002/ece3.2778>, 2017.
- 1264 Zhang, Y., Lavalley, J. M., Robertson, A. D., Even, R., Ogle, S. M., Paustian, K., and Cotrufo, M. F.: Simulating
1265 measurable ecosystem carbon and nitrogen dynamics with the mechanistically defined MEMS 2.0 model, *Biogeosciences*,
1266 18, 3147–3171, <https://doi.org/10.5194/bg-18-3147-2021>, 2021.
- 1267 Zhang, Y., Wang, T., Yan, C., Li, Y., Mo, F., and Han, J.: Microbial life-history strategies and particulate organic carbon
1268 mediate formation of microbial necromass carbon and stabilization in response to biochar addition, *Sci. Total Environ.*,
1269 950, 175041, <https://doi.org/10.1016/j.scitotenv.2024.175041>, 2024.

1270

1271

1272

1273

1274

1275

1276

1277

1278

1279

1280

1281

1282

1283

1284

Zhong, W., Gu, T., Wang, W., Zhang, B., Lin, X., Huang, Q., and Shen, W.: The effects of mineral fertilizer and organic manure on soil microbial community and diversity, *Plant Soil*, 326, 511–522, <https://doi.org/10.1007/s11104-009-9988-y>, 2010.

Zhou, G., Zhou, X., He, Y., Shao, J., Hu, Z., Liu, R., Zhou, H., and Hosseinibai, S.: Grazing intensity significantly affects belowground carbon and nitrogen cycling in grassland ecosystems: a meta-analysis, *Glob. Change Biol.*, 23, 1167–1179, <https://doi.org/10.1111/gcb.13431>, 2017.

Zhou, G., Luo, Q., Chen, Y., He, M., Zhou, L., Frank, D., He, Y., Fu, Y., Zhang, B., and Zhou, X.: Effects of livestock grazing on grassland carbon storage and release override impacts associated with global climate change, *Glob. Change Biol.*, 25, 1119–1132, <https://doi.org/10.1111/gcb.14533>, 2019.

Zhou, Z., Ren, C., Wang, C., Delgado-Baquerizo, M., Luo, Y., Luo, Z., Du, Z., Zhu, B., Yang, Y., Jiao, S., Zhao, F., Cai, A., Yang, G., and Wei, G.: Global turnover of soil mineral-associated and particulate organic carbon, *Nat. Commun.*, 15, 5329, <https://doi.org/10.1038/s41467-024-49743-7>, 2024.

Zuur, A. F., Ieno, E. N., Walker, N., Saveliev, A. A., and Smith, G. M.: *Mixed effects models and extensions in ecology with R*, Springer, New York, NY, <https://doi.org/10.1007/978-0-387-87458-6>, 2009.