

# Detailed responses to reviews

## Saharan dust impacts on the surface mass balance of Argentière Glacier (French Alps)

Léon Roussel, Marie Dumont, Marion Réveillet, Delphine Six,  
Marin Kneib, Pierre Nabat, Kévin Fourteau, Diego Monteiro,  
Simon Gascoin, Emmanuel Thibert, Antoine Rabatel, Jean-Emmanuel Sicart,  
Mylène Bonnefoy, Luc Piard, Olivier Laarman, Bruno Jourdain,  
Mathieu Fructus, Matthieu Vernay, and Matthieu Lafaysse

September 22, 2025

The authors thank the editor and the two reviewers for the time dedicated to reading the manuscript and for the constructive feedbacks, which we believe contributed to improving its quality. Answers to the two reviewers are listed in the coloured boxes and our responses in black below the boxes. The quotation marks are used when citing the manuscript in blue, and the modifications are highlighted in bold. We also provide a revised version of the manuscript, as well as a version with tracked changes.

## 1 Summary of the corrections

The main adjustments carried out in this revised version of the manuscript are the following:

- broader details on the ALADIN model (R1.2.)
- justifying our choice of attribution concerning the dust impact (R2.6.)
- addition of uncertainty tests regarding dust scavenging (R1.3.), and mass absorption efficiency of dust (R2.7.)
- adjustment of the mass absorption efficiency (MAE) of LAPs and its justification in a detailed analysis in section 3 of this response

## 2 Answers to Reviewers

### 2.1 Reviewer #1

**R1.1.** This is an interesting and rigorous study investigating the impacts of Saharan dust on a French Alpine glacier. The study nicely combines remote sensing data of albedo, in-situ measurements of glacier surface mass balance, snowpack radiative transfer modeling, and glacier surface mass balance modeling. Uncertainty is explored via numerous perturbed simulations. Overall, I find the conclusions to be compelling and robustly supported through analysis and discussion. The authors find a large impact on glacier SMB from the dust deposition in 2022, supporting previous studies, and adding context to earlier studies (e.g., Gabbi et al, 2015) that explored impacts of dust and black carbon on other glaciers in the Alps. Moreover, the manuscript is well-written and the figures are excellent. I recommend publication after relatively minor issues are addressed.

We thank reviewer #1 for his review, his positive feedback, and for his interesting and constructive comments that help improving the manuscript. We hope that our answers and modifications help to clarify our choices.

### 2.1.1 More important issues

**R1.2.** Section 2.3: Please include more detail about how the dust and black carbon emissions and deposition fluxes are simulated in ALADIN, as the deposition fluxes critically impact the results of this study. In particular, are the dust emissions simulated prognostically, or are they prescribed? Which inventory of BC emissions is used? Do the BC emissions vary between years? Does the regional domain of the model include the Sahara desert?

The ALADIN model, providing the deposition fluxes of dust and BC for this study, includes seven aerosol species, namely dust, sea salt, sulfate, black carbon, organic matter, nitrate and ammonium.

- *Does the regional domain of the model include the Sahara desert? Are the dust emissions simulated prognostically, or are they prescribed?* The domain of the model indeed includes the Saharan desert with a southern limit roughly corresponding to +15°N (see Fig. 1 of [Nabat et al. \(2020\)](#)). The emissions of mineral dust are "dynamically computed by the model as a function of surface wind as well as soil characteristics" ([Nabat et al., 2020](#)). Mineral dust is a prognostic variable of the model subject to atmospheric processes (emission, transport and deposition).
- *Which inventory of BC emissions is used? Do the BC emissions vary between years?* Anthropogenic and biomass burning emissions (including BC) are based on "monthly CMIP6 historical inventories provided, respectively, by [Hoesly et al. \(2018\)](#) and [Marle van et al. \(2017\)](#)" ([Nabat et al., 2020](#)).

Anthropogenic and biomass burning emissions (including BC) are based on monthly inventories commonly used in climate models. The CEDS v2021-04-21 dataset accounts for historical anthropogenic emissions ([O'Rourke et al., 2021](#)) up to 2019, extended up to 2022 following the methodology of [Lamboll et al. \(2021\)](#) to account for the decrease in anthropogenic activities due to lockdown periods of the year 2020. Biomass burning emissions come from the GFED4.1s dataset ([Randerson et al., 2017](#)). All these emissions vary from year to year.

To clarify the manuscript, we added the following summary in section 2.3: "Mineral dust (dust) and black carbon (BC) hourly deposition fluxes were obtained from the CNRM-ALADIN63 regional climate model driven by the ERA5 reanalysis ([Nabat et al., 2020](#)). **The prognostic aerosol scheme of ALADIN is called TACTIC (Tropospheric Aerosols for Climate In CNRM, coming from [Morcrette et al. \(2009\)](#) and used in [Michou et al. \(2015, 2020\)](#)**). The domain of ALADIN included the Saharan desert with a southern limit roughly corresponding to +15°N, as shown in Fig. 1 of [Nabat et al. \(2020\)](#). The emissions of dust were computed by the model as a function of surface wind and soil characteristics ([Nabat et al., 2020](#)). Then, mineral dust was a prognostic variable of the model subject to atmospheric processes (transport and deposition). Anthropogenic and biomass burning emissions (including BC) were based on monthly inventories varying from year to year, namely the CEDS v2021-04-21 dataset for historical anthropogenic emissions ([O'Rourke et al., 2021](#)) up to 2019, extended up to 2022 following the study of [Lamboll et al. \(2021\)](#), and the GFED4.1s dataset ([Randerson et al., 2017](#)) for biomass burning emissions. The horizontal grid resolution of the CNRM-ALADIN63 outputs is **12.5 km**. Only the nearest grid point output was used for all simulation points."

**R1.3.** Figure 8 and associated discussion: The sensitivity of dust impacts to the BC melt scavenging factor is certainly an interesting result. The authors may disagree, but I suspect that uncertainty in melt scavenging efficiency of dust itself is similarly large, and if so, the simulated SMB impacts would also be sensitive to uncertainty in this parameter. One way of exploring and presenting this would be to add additional markers to Figure 8 that depict the sensitivity to dust scavenging. Regardless, I think it would be helpful to include a bit more acknowledgment and discussion of this.

We thank the reviewer for pointing out this point, and we agree with its comment. In the reference version of the simulations, we chose not to allow dust percolation, in agreement with the literature, as mentioned in section 2.4.1 of the initial version of the manuscript: "For large LAPs such as mineral

dust, with a typical radius of 5 microns, Conway et al. (1996) explained that these particles stay near the snow surface, almost not percolating with liquid water, hence we set dust scavenging coefficient  $k_{\text{dust}} = 0$ ." It applies here since the size of the dust particles of this study are likely  $\geq 1 \mu\text{m}$  (e.g., Fig. 2 and Fig 8 of Dumont et al. (2023)). In addition, we did a sensitivity test with  $k_{\text{dust}} = 0.04$  in the initial version of the paper (Fig. 7) to account for a light scavenging of dust. We chose this value  $k_{\text{dust}} = 0.04 < k_{\text{BC}} = 0.2$ , because it is known that the scavenging efficiency of dust is lower than the scavenging efficiency of BC (e.g., Doherty et al. (2013)).

However, although this is poorly documented in the existing literature, we also suspect that dust scavenging efficiency at the snow surface could be non-zero and higher than 0.04 (e.g., during intense rain on snow events, or when liquid water flows on an inclined snow surface).

Overall, these uncertainties regarding LAP scavenging efficiencies highlights the need to better understand the motion of the different LAP inside of the snowpack at the micro-scale.

Therefore, following the advice of reviewer #1, we included an additional simulation and the corresponding marker in Fig. 8 for a dust scavenging efficiency of  $k_{\text{dust}} = 0.2$ . Regarding the previous remarks on the available literature, this is likely too high compared to  $k_{\text{BC}} = 0.2$ , so this sensitivity test has to be taken with cautious. We added the following paragraph in section 4.1: **"The dust scavenging efficiency  $k_{\text{dust}}$  that was set to 0 in the reference version, i.e., no scavenging of dust (see Method Section 2.4.1). A higher  $k_{\text{dust}} = 0.04$  led to a lower impact of dust (Fig. 7). Although this was poorly documented in the existing literature, we suspected that dust scavenging efficiency at the snow surface could be significantly higher than the previous sensitivity test. So we added a test of uncertainty using  $k_{\text{dust}} = 0.2 = k_{\text{BC}}$  (Fig. 8), which is likely too high since BC particles scavenge more easily than large particles such as dust (e.g., Conway et al. (1996); Doherty et al. (2013)). The dust impacts using those two dust scavenging efficiencies were still inside the uncertainty induced by the meteorological ensemble."** We also modified the last paragraph of section 4.2: Overall, the uncertainties on LAP scavenging efficiencies in snow result in large uncertainties on the respective absolute impacts of dust and BC. **This highlights the need to better understand the motion of the different LAPs inside of the snowpack at the micro-scale.** Here, BC scavenging in snow, [...].

### 2.1.2 Minor issues

**R1.4.** line 30-31: "LAPs have advanced the snow melt-out date on average by 18 days in the French Alps in the past 40 years..." - As written, there is potential for ambiguity in whether this a trend towards earlier melt-out date over the last 40 years, or a mean impact over the last 40 years. Please clarify.

We agree that there is a potential confusion. This is here a mean impact over 40 years, not a trend. To insist on the mean, we propose the following formula: **"For instance, on average over the past 40 years, the presence of LAPs in snow has advanced the snow melt-out date by 18 days in the French Alps, and even more at high elevations (more than 20 days at 3,000 m a.s.l., Réveillet et al., 2022)."**

**R1.5.** Section 2.2.3: Please list the wavelengths of the spectral bands used for the NDSI and RGND calculations.

We added in Section 2.2.3 **"The central wavelengths of bands B3, B4 and B11 are respectively 560, 665 and 1610 nm."**

**R1.6.** Section 2.2.5: Here, BC is varied within SNICAR to match the Sentinel-2 albedos. Dust also could have been varied to match the satellite albedos. Are the results at all sensitive to the choice of LAP used for this purpose?

We understand the comment and the concerns. We agree that using an more advanced method could increase the accuracy of the retrieved ice albedo.

However, (i) since only the broadband albedo was prescribed for ice albedo, and (ii) since we quantified here the impact of LAP in snow, and not in ice, we chose this simple method varying only one parameter. Varying other parameters, such as dust concentration, air bubble size or density, could indeed have slightly modified the broadband ice albedo and probably increase the accuracy. But retrieving a very accurate ice albedo is not the aim of this study. We argue that the error between our method and a more accurate method is small compared to the spatial variability of ice albedo on the glacier (that we do not represent in the model), and for which we performed the sensitivity tests of  $\pm 20\%$  on Figure 7.

To reflect this limitation, we added the following sentence in section 2.2.5: **"This method is rather simple since we here aimed to quantify the impact of dust in snow, not in ice. Nevertheless, the sensitivity to this parameter was investigated in Section 3.3."**

**R1.7.** line 138: The use of "(-)" is unfamiliar to me, but is perhaps used to indicate that albedo is a dimensionless quantity. I don't think this is needed, but if there is precedence for using such notation, it is fine.

"(-)" was indeed used for unit-less variables. We removed this notation from the text and from the figures.

**R1.8.** line 171: "right-hand side moraine" seems to be perspective-dependent, but perhaps there is precedence for using this terminology. Regardless, please clarify.

From an hydrological point-of-view, the right and left-hand sides are determined using the flow direction. Therefore, here the moraine AWS is on the right-hand side of the glacier considering the glacier flow direction. To clarify this point, we added this information: **"Two automatic weather stations (AWS) were located at the surface and on the right-hand side moraine with respect to the glacier flow of Argentière Glacier (Fig. 1d)."**

**R1.9.** line 203: Please clarify this sentence. It may be as simple as changing "then" to "so".

Thanks for spotting this. We switched to **"so"**.

**R1.10.** Equation 1: Is there an upper bound of temperature to which this scaling is applied?

There is no upper bound in Equation 1. However, for example, at both the moraine or glacier AWS, located on the part of the glacier experiencing the highest air temperatures, the 2 m air temperature rarely exceeded 20 °C during the summer 2021, as shown in the example of Appendix Fig. C1. We added this information in the manuscript, section 2.3.4: **"Therefore, we adjusted the SAFRAN temperature  $T^*$ , above  $T_0 = 3.5$  °C without any upper bound, ..."**

**R1.11.** Section 2.3.6: Although this is clarified in the appendix, please briefly communicate here whether the perturbations are random within the confines of individual variances, as in a Monte Carlo approach, or whether variables are perturbed in combination by their full variance. (Or in general, please provide a little more detail on the perturbation approach here).

We are not completely sure to understand correctly the comment. It seems that the reviewer is asking: "Are the perturbations univariate or multivariate?". If we understood correctly the request, in this study, the approach chosen is univariate. This means that each variable was perturbed independently from other variables, *i.e.*, here each variable is perturbed uniquely using its own standard deviation  $\sigma_X$ . This implies that the different perturbations do not account for initial correlations between the meteorological variables, *e.g.*, an concurrent increase in air temperature and a decrease in long-shortwave radiation; are just as likely as; an concurrent increase in air temperature and a increase in long-shortwave radiation.

To account for your comment, we clarified this point in the revised manuscript, in the section 2.3.6, where you can now read: "... For a given member, the same perturbations were spatially applied to all simulation points. **Each variable was perturbed independently from other variables. For each variable, the perturbation term  $X_t$  depends on a decorrelation time  $\tau$  and an amplitude  $\sigma_X^2$ . The exact relationship is explained in detail in Appendix C1. Given an additive variable  $V_t$  (T or LW) at time-step  $t$ , the perturbed variable  $V_t^*$  is defined as  $V_t^* = V_t + X_t$ , and in case the variable is multiplicative (SW or W),  $V_t^*$  is defined as  $V_t^* = V_t \cdot (1 + X_t)$ .** We applied the perturbations on a daily scale with a decorrelation time ..." The method is detailed in Appendix C1 that remains unchanged.

**R1.12.** Figure 5: The month abbreviations on the x-axis appear to be in French.

Thanks. We switched to English.

**R1.13.** line 457: "be" → "been"

We modified accordingly: "the dust impact in the ablation area would have likely **been** higher."

**R1.14.** line 489: "We could not directly compare the MAC of dust..." - Despite this, please include the MAC values of dust that were assumed in your study.

As explained and justified in section 3 of this review, the MAE (mass absorption efficiency) of dust and BC were modified in this revised version. We added the revised MAE of dust and BC in the revised manuscript: "To compute the albedo of snow, the refractive indices of ice were set as Réveillet et al. (2022). **The mass absorption efficiency (MAE) of dust  $\text{MAE}_{\text{dust}}(\lambda)$  used for this study was the MAE of a PM10 Lybian dust from Caponi et al. (2017), i.e.,  $\text{MAE}_{\text{dust}}(\lambda) = \text{MAE}_{\text{dust}}(\lambda_0) \times (\frac{\lambda}{\lambda_0})^{-\text{AAE}_{\text{dust}}}$  where  $\lambda_0 = 400$  nm,  $\text{MAE}_{\text{dust}}(\lambda_0) = 71 \times 10^{-3} \text{ m}^2 \text{ g}^{-1}$ , and the Ångström absorption exponent  $\text{AAE}_{\text{dust}} = 3.2$ . The MAE of BC  $\text{MAE}_{\text{BC}}$  at 550 nm used for this study was  $7.5 \text{ m}^2 \text{ g}^{-1}$  (Bond, Bergstrom, 2006).** To quantify the impact... To harmonize we also switched the occurrences of "MAC" (mass absorption cross-section) to "**MAE**" (mass absorption efficiency) in the whole manuscript.

## 2.2 Reviewer #2

**R2.1.** This study examines the impact of mineral dust and BC on the surface mass balance of a glacier in the French Alps. The study uses a variety of assimilated and reanalysis data to inform a multilayer snow model, CROCUS, which was adapted to account for LAP layer deposition and exposure within the snowpack structure. Results show that dust contributed significantly to water equivalent loss from the glacier over all years investigated, but especially in 2022 when compounding effects of dust deposition during the previous season resulted in exception melt loss. I think this study is very well structured and organized. I appreciate the wide array of data used and the thorough discussion of methods and limitations. The Figure and results are mostly well designed and can be clearly understood.

We thank reviewer #2 for his general comment and for the specific comments that helps clarifying the limitations of our study. We hope that our answers help to better apprehend our choices and results.

### 2.2.1 Primary concerns

Each of those primary concerns were detailed below by reviewer #2. So we gave an answer to each of those primary concerns directly under the linked comment below.

**R2.2.** My primary concern lies within the discussion associated with BC scavenging and increased dust impact, detailed in the comments below.

We addressed this concern in the response to specific comment R2.6.

**R2.3.** I would also request some clarity regarding the choice of both the LAP representation within the snow representation, [...]

We addressed this concern in the response to specific comment R2.7.

**R2.4.** I would also request some clarity regarding [...] the choice of CROCUS itself.

We addressed this concern in the response to specific comment R2.9.

**R2.5.** Overall, I think the approach in this study is great for assessing specific, well-observed glaciers. However, I question some broader claims of scalability and transferability to other regions.

We addressed this concern in the response to specific comment R2.13.

### 2.2.2 Specific comments

**R2.6.** Line 530, Section 4.2, Figure 8, and elsewhere: There is mention of higher dust impacts when BC is low or highly scavenged within the snowpack. I agree that this increases the relative impact of dust on snowmelt (compared to BC), but as I understand it, this shouldn't directly affect the impact that dust has in generating melt water. In other words, 1g/m<sup>2</sup> of exposed dust would generate X m w.e. of melt, and 1 g/m<sup>2</sup> of dust with 0.1 g/m<sup>2</sup> black carbon would generate X m w.e. of melt (from dust) and Y m w.e. of melt (from BC). The point being that X m w.e. is the same in both scenarios. If this is not the case, then it is currently unclear what the mechanism is that would drive the same amount of dust to have a greater impact on melt in the absence of BC (because of higher scavenging). I would argue that having more BC retained in the snowpack with dust would enhance the impact of dust indirectly, by resulting in more rapid exposure of buried deeper dust layers from X+Y vs. just X melt. Please explain and/or clarify.

This is indeed an interesting remark and it is a question of attribution. To clarify the situation, we introduce Figure 1 of this document (not added in the revised manuscript) that illustrates the difference between the two possible approaches to quantify the impact of dust: the method suggested by reviewer #2 and the one adopted in this study.

In this study, the impact of dust was computed by comparing the simulation with [BC only] and the simulation with [BC+dust]. In other words, only the dust is removed in order to quantify its impact, by comparison with 'contaminated' snow, which we consider to be closer to reality. However, as mentioned in the manuscript in section 4.2, when using this reference, the impact of one LAP depends on the concentration of other LAP, because albedo is a non-linear function of LAP concentration (*e.g.*, Fig. 12 of Skiles, Painter (2017)). For instance, at a given timestep, less BC at the snow surface increase the dust radiative forcing.

In the second method suggested by reviewer #2, our understanding is that the impact of dust would be based on comparing the simulation [without LAP] and the simulation with [dust only]. However, this method, (i) is a comparison between 2 virtual scenarios ([no LAP] and [only dust]) and (ii) probably lead to overestimate the impact of dust (see Fig. 1).

Both methods have been used in the literature to quantify this impact (*e.g.*, Flanner et al. (2007, 2012) used the first approach, and Réveillet et al. (2022) the second approach). As mentioned above, each has advantages and limitations. Nevertheless, based on the work of Mark Flanner and others




Figure 1: Choice on the attribution of the dust impact  $\Delta_{\text{dust}}$  from different point-of-view. Note that we have intentionally (i)  $\Delta_{\text{dust}} + \Delta_{\text{BC}} \leq \Delta_{\text{LAP}}$  and (ii)  $\Delta_{\text{dust}} \leq \Delta'_{\text{dust}}$ , this is due (among other factors) to the sub-linearity of the albedo decrease with respect to the LAP concentration.

(cited below) and considering that the potential overestimation of the impact inherent to the second approach represents a limitation, we prefer to retain the approach described in this study.

Citations from Flanner’s work: Flanner et al. (2007): ”We calculate instantaneous radiative forcing of carbon aerosols in both snow and the atmosphere at each radiative transfer time step as the difference in absorbed radiation with all aerosols and all aerosols except carbon aerosols.” Flanner et al. (2012): ”We calculate radiative forcing [of BC] each timestep as the instantaneous difference in absorbed surface energy with and without BC.” In this case, adding other LAPs to the simulation decreases the RF of the initial LAPs: ”These forcings are smaller than previous estimates (Flanner et al., 2007, 2009), demonstrating the importance of other factors. Reasons why forcing is different in this study are [...] (4) light absorption by dust in snow, which decreases BC forcing, was not considered by Flanner et al. (2007)” (Flanner et al., 2012).

To clarify and better justify our approach, we added the following clarifications in section 2.5: ”The forcing ensemble was used as input to the simulation model to get the simulations considering the impact of mineral dust (simulations named ”dust”), and the simulations without the impact of mineral dust (simulations named ”no-dust”). Simulations without the impact of dust were obtained without the radiative effect of **dust LAP** as explained in Section 2.4.2. As a result, we obtained 40 paired simulations (dust and no-dust). **An important point is that BC remained present in all simulations. An alternative could have been to use the difference between (i) the simulation with dust and without BC, and (ii) the simulation without LAP at all as in Réveillet et al. (2022). This would have likely lead to larger quantified impacts (Flanner et al., 2012). However, we chose the simulation with both BC and dust as the reference as it is closer to the observed state of the snowpack, hence avoiding an overestimation of the dust impact and staying in line with other studies, e.g., Flanner et al. (2007, 2012).**”

**R2.7.** Section 2.4.1: As I understand it, the LAP implementation and resulting modeled albedo depends on optical properties derived from dust during previous years at the same glacier. Is there ample evidence that the optical properties do not vary that much? Explain.

We thank the reviewer to point out this limitation of the study.

Indeed, depending on the region of origin or its size distribution, the mineral dust deposited over the European Alps can have different optical properties (*e.g.*, Dumont et al. (2023); Caponi et al. (2017)). After deposition the optical properties of dust can also evolve inside the snowpack with (i) the location of the LAP with respect to the snow grains, *i.e.*, inside or outside of the snow grains, *e.g.*, He et al. (2018), or with (ii) the aggregation of the dust particles, *e.g.*, Bond, Bergstrom (2006).

To assess the impact of different absorption properties of dust, we did 2 additional sensitivity tests (added in Fig. 8 of the manuscript) using a low and a high value of mass absorption efficiency (MAE) of dust, respectively [ $\text{MAE}_{\text{dust}}(\lambda_0) = 27 \times 10^{-3} \text{ m}^2 \text{ g}^{-1}$ ,  $\text{AAE}_{\text{dust}} = 3.3$ ] and [ $\text{MAE}_{\text{dust}}(\lambda_0) = 630 \times 10^{-3} \text{ m}^2 \text{ g}^{-1}$ ,  $\text{AAE}_{\text{dust}} = 3.4$ ]. Those values are the extrema of MAE listed in Caponi et al. (2017) when considering the regions of Sahel and Sahara, the main sources of mineral dust for European Alps (Collaud Coen et al. (2004), or more recently Collaud Coen et al. (2025)).

The sensitivity tests show that increasing the dust MAE results in a higher dust impact, leading to enhanced melt and consequently a decrease in glacier surface mass balance. We noted that there was a significant difference in dust impact in 2022 between these extreme values of MAE ( $\Delta_{\text{dust}} < 0.4 \text{ m w.e.}$  for the low MAE and  $\Delta_{\text{dust}} > 1.4 \text{ m w.e.}$  for a high MAE). However, these extreme dust impacts are likely unrealistic since the simulations using extreme dust MAE (low and high) had lower evaluation rates than the reference scenario used throughout the study (see Table. 3).

To address your comment and the additional sensitivity tests mentioned below, the Methods and Results sections, as well as Figure 8, have been updated in the revised version as follows:

Method section of section 2.4.1: **For the two uncertainty tests on the dust MAE (Fig. 2), we used the highest and the lowest values of dust MAE considering the regions of Sahel and Sahara of Caponi et al. (2017), respectively [ $\text{MAE}_{\text{dust}}(\lambda_0) = 27 \times 10^{-3} \text{ m}^2 \text{ g}^{-1}$ ,  $\text{AAE}_{\text{dust}} = 3.3$ ] and [ $\text{MAE}_{\text{dust}}(\lambda_0) = 630 \times 10^{-3} \text{ m}^2 \text{ g}^{-1}$ ,  $\text{AAE}_{\text{dust}} = 3.4$ ].**

We added the following sentences in a new paragraph of section 4.1: **In addition, the temporal variability of the absorption efficiency of dust was not represented in the model: the MAE used in this study was constant in time. However, the absorption efficiency of mineral dust can vary for each deposition event depending on the region of origin or the size distribution (*e.g.*, Dumont et al. (2023); Caponi et al. (2017)), as well as by its location within the snowpack either inside or outside the snow grains (He et al., 2018) and by the aggregation state of the dust particles (Bond, Bergstrom, 2006). As shown in Fig. 2, a low and a high value of dust MAE (Section 2.4.1) strongly affects the resulting dust impact ( $\Delta_{\text{dust}} < 0.4 \text{ m w.e.}$  for the low MAE and  $\Delta_{\text{dust}} > 1.4 \text{ m w.e.}$  for a high MAE). However, these extreme dust impacts are unlikely since the simulations using extreme dust MAE (low and high) results in degraded accuracy metrics with respect to in situ measurements compared to the reference scenario used throughout the study (not shown).** Since this MAE remains poorly constrained and represent an interesting research perspective, we also propose to mention this limitation in the conclusion section: **Furthermore, large uncertainties linked to the percolation of BC within the snowpack might lead to significantly higher impacts of dust. Using different mass absorption efficiency for the dust particles in snow also significantly modulated the results.** Further research is needed to understand these processes and to quantify their impacts."

**R2.8.** The LAP representation could be assessed indirectly by comparison to remotely sensed snow albedo. Why was a snow albedo comparison from Sentinel or another sensor omitted? Remotely sensed snow albedo could also inform modeling directly and may be more scalable. Explain why this approach was not used in this paper. If Sentinel has a pixel saturation issue that prevents good snow albedo detection, mention this.

We thank reviewer #2 to mention this idea. Indeed, broadband albedo derived from remote measured reflectances has been increasingly used for glaciological studies. For instance, Naegeli et al. (2019) derived ice albedo trends using remote-sensing data using the method of Liang et al. (2003). However, we should remain cautious when comparing values of simulated broadband albedo and broadband albedo retrieved from remote-sensed reflectances, here Sentinel-2. First, Sentinel-2 measures the hemispherical-conical reflectance at given wavelengths. Since the reflection of snow is not isotropic (*e.g.*, Dumont et al. (2010)) (*i.e.*, depends on the observation angle, and also varies with the solar




Figure 2: Figure 8: Impact of uncertainties relative to LAP scavenging efficiencies and deposition fluxes on the glacier-wide annual SMB (y-axis) and dust impact  $\Delta_{\text{dust}}$  (x-axis) for hydrological year 2022. Reference simulation is represented by the central black dot. The 40 members of the ensemble are represented with the orange dots.

zenith angle), a reflectance of snow on a given surface may differ from the albedo (bi-hemispherical reflectance) of this surface. Exploiting correctly the relationship between snow reflectance observed from S2 and albedo would require a more complex processing, *e.g.*, Lamare et al. (2020). The method of Liang et al. (2003), initially developed for other land surfaces than snow, does not account for the anisotropy of snow. Second, the uncertainties of the reflectance retrievals in complex terrain may be too high to estimate the accuracy of the simulated albedo, as shown in Cluzet et al. (2020) for Sentinel-2 (S2) images. For instance, the strong dependency to the slope is illustrated for S2 derived albedo in Fig. 3a. That is why we initially based our evaluation on surface type (snow, ice), instead of broadband albedo values derived from S2 images.

Considering your comment, and the limitations mentioned above, we added a comparison between the simulated albedo and the albedo derived from the observed S2 reflectances in the Appendix of the revised version of the manuscript (Figure 3 of this review). The method is explained in the new section 2.2.6: **”Broadband snow albedo derived from Sentinel-2 reflectances”** of the revised manuscript: **”Using all images of Table B2, we compared the simulated broadband albedo at noon, and the observed broadband albedo derived from Liang et al. (2003), *i.e.*,  $S2 \text{ albedo} = 0.356 * R_{B2} + 0.130 * R_{B4} + 0.373 * R_{B8} + 0.085 * R_{B11} + 0.072 * R_{B12} - 0.018$  where  $R_{Bi}$  is the reflectance of Sentinel-2 band  $i$ . Only the points where the surface type (Section 2.2.3) was snow for both the model and S2 were used. Appendix Fig. 3 shows this comparison.”** The precautions to be taken regarding this comparison are mentioned in the caption of Appendix Fig.3.

It should also be noted that these results highlighted a number of limitations, particularly concerning the choice of MAE, and led to additional modifications in the revised manuscript, detailed in section 3 of this document.




Figure 3: **Figure B3. Comparison between simulated broadband albedo and broadband albedo derived from S2.** See section 2.2.6 for methods and limitations. (a) Influence of the slope. (b) Comparison for slopes  $< 20^\circ$ . The mean of all dates, except 2, shows less than 0.1 of difference between the modelled albedo and the albedo derived from S2. This comparison of broadband albedo has to be taken with cautious. Indeed, there are several limitations. First, the broadband albedo derived from S2 (Liang et al., 2003) was not developed for snow, an anisotropic surface (e.g., Dumont et al. (2010)), i.e., the reflectance depends on the observation angle. Second, the uncertainties of the reflectance retrievals in complex terrain may be too high to estimate the accuracy of the simulated albedo, e.g., Cluzet et al. (2020), or panel (a) of this figure where the broadband albedo derived from S2 strongly depends on the slope.

**R2.9.** Section 4.3: I appreciate the thorough discussion of uncertainties regarding LAPs, but think there also needs to be more discussion of broader model uncertainties. What are the limits of CROCUS? How would using different [...] physically based snow models vary the results? If word count is an issue, I would recommend condensing some lines from the detailed discussion of SMB and LAPs.

*This comment was initially combined with the next comment (R2.10.). Please refer to the next comment for the uncertainties concerning the choice of the model providing the meteorological forcing.*

In the initial version of the manuscript, we did not discuss the snow model choice. We used here the SURFEX-ISBA/Crocus model because (i) it has already been widely used in the European Alps to model glacier SMB (e.g., Gerbaux et al. (2005); Dumont et al. (2012); Réveillet et al. (2018)), (ii) it has a detailed representation of the physical processes governing the snowpack evolution (which is important for our study assessing impacts in snow), and (iii) this model is one of the only models that combined a detailed layered description of snow microstructure with an explicit representation of the radiative effects of LAPs. Indeed, Crocus is coupled with the radiative transfer model TARTES (G. Picard, Q. Libois, 2024) and this coupling has already been used in multiple studies (e.g., Tuzet et al. (2020); Dumont et al. (2020); Réveillet et al. (2022)). For such reasons, we think that Crocus is our best option to quantify the impact of LAP in the Alps.

Nevertheless, Crocus, as every model, is subject to modelling uncertainties related to the physical processes driving snow evolution, or from the radiative transfer model. This is now discussed in section 4.3 of the revised version of the manuscript: **"The uncertainties in the results can also be related to the choice of the model and the parameters used in the model (e.g., Lafaysse et al. (2017)). In this study, this primarily concerns the physical processes driving snow**

evolution and the radiative transfer model. Concerning the radiative transfer model, here TARTES coupled with the snow model, comparisons between radiative models (SNICAR and TARTES in G. Picard, Q. Libois (2024)) indicate that the choice of the radiative transfer model has a significantly smaller impact on the results than variations in the optical properties of the LAP, here represented using the MAE of each LAP. Therefore, the uncertainty related to the radiative model is accounted for in this study by varying the dust MAE factor (Section 4.1). Concerning the snow model, an ensemble approach can provide estimates of snow model uncertainties. Réveillet et al. (2022) applied this method to Crocus and reported results on the impact of the LAPs comparable to deterministic simulations (Supplementary Fig. S10 of Réveillet et al. (2022)).”

**R2.10.** Section 4.3: [...] How would using different NWP [...] vary the results?

For the meteorological forcing, we used the SAFRAN reanalysis. It is indeed NWP-based, however SAFRAN also ”combines information from numerical weather prediction models (ERA-40 reanalysis from 1958 to 2002, ARPEGE from 2002 to 2021) and the best possible set of available in situ meteorological observations” (Vernay et al., 2022), *i.e.*, SAFRAN is a reanalysis and not just a numerical weather prediction. In addition, it was the most performant meteorological reanalysis over the French Alps given our usage. Thus, using a different NWP without any assimilated observation would very likely lead to higher uncertainties.

Nevertheless, this dataset has identified errors, *e.g.*, incoming longwave bias in Réveillet et al. (2018). So, we used in-situ data using 2 AWS to improve the accuracy of this forcing (section 2.3.2). And to account for the remaining errors and to avoid using meteorological forcings from different sources, we applied stochastic perturbations on the SAFRAN data (section 2.3.6), taking into account the errors of SAFRAN w.r.t. the AWS moraine. This way, we propagated the forcing uncertainties and we quantified their impact on the results. The median-Q10-Q90 of the results using the meteorological ensemble are often given in the manuscript in the form of XXX [XXX, XXX]. At each point, this led to meteorological uncertainties of around  $\pm 1$  m w.e. on the annual simulated SMB, with meteorological uncertainties on the fraction of melt due to dust varying between a few percent for the lowest impact (*e.g.*, at point P1) to  $\pm 10\%$  for the highest impacts (*e.g.*, at point P2).

Finally, our study is mostly based on relative differences between simulations, thus lowering the impact of uncertainties related to the meteorological variables (except the LAP deposition fluxes) and to the surface model. That is actually also a reason why the different result in term of dust impact do not diverge a lot (often small standard deviation compared to the median).

Continuing the paragraph of the previous comment R2.9. on the model uncertainties, we propose to add the following sentences: **”Furthermore, uncertainties related to the snow model choice are generally smaller than those associated with the forcing (*e.g.*, Etchevers et al. (2004); Günther et al. (2019)). To explore the uncertainties related to the meteorological variables, we did the choice to use an ensemble of forcing produced using stochastic perturbations on the adjusted SAFRAN data (section 2.3.6). Ultimately, the dust impact presented in this study is expressed as relative differences between simulations rather than as absolute values, which limits the influence of uncertainties related to the forcing and to the snow model choice.”**

**R2.11.** What if there is no information about the optical properties of snow? Is CROCUS best suited for very large glaciers? This also relates to my comment on line 623.

Concerning the optical properties of snow itself in Crocus, they are calculated from Crocus prognostic variables (*e.g.*, snow grain size, density). Thus there is no need for any information other than the meteorological forcing to compute them. We added the following sentence in section 2.4 to clarify this point: **”Crocus simulates the evolution of the snow cover with a 15 minutes time-step representing different processes including thermal diffusion, phase change, water flow, snow metamorphism and compaction. Snow optical properties (single scattering albedo and absorption cross section for each snow layer) are calculated from the prognostic variables of the model (grain size, density, layer thickness).** In Crocus, the snowpack is discretized as a one dimensional column with up to 50 layers, [...]”




Figure 4: Revised Figure 6.

For "very large glaciers", if it refers to large glaciers in the European Alps, Crocus has already been used widely and demonstrated its ability to perform well regardless of the glacier size (*e.g.*, Gerbaux et al. (2005); Dumont et al. (2012); Réveillet et al. (2018)). We propose the following modification in section 2.4: "which was already used to simulate alpine glacier SMB regardless of the glacier size, *e.g.*, Gerbaux et al. (2005); Dumont et al. (2012); Réveillet et al. (2018)". If it refers to glaciers outside of the Alps, we answered below under the comment R2.13.

**R2.12.** Figure 6: Basic cartography issues. Please add a scale bar and a north arrow to the maps. Some indication of elevations (maybe very faint topo lines) would help the reader understand the distribution of these changes over terrain. I would recommend including the same basemap as Figure 1 (d) in these panels. This is the take-home figure of this paper, and it would be helpful to orient readers, especially those just "skimming" and those not familiar with the glacier.

These element were indeed missing in Figure 6. We thank the reviewer for pointing this out. A scale bar and a north arrow have need added to the first plot. Grey isolines have been added to every plot of this figure. We did not added the glacier outlines, to keep a light and readable figure. The resulting revised figure is Fig. 4 of this document.

**R2.13.** Line 623: Is this truly “easily” scalable if this modeling work relies heavily on bias correction and adjustments from automated in situ measurements (as discussed in section 2.3.3)? Are these relationships scalable to other glaciers at other latitudes? Most glaciers do not have automated measurements, let alone a glacier observatory, and modeled forcings have much higher uncertainties in more remote regions (Himalayas, Andes, Rockies). Would this approach to such glaciers truly be reasonable? Please clarify how such an approach would be transferable with ease. *Combined with the general comment:* Overall, I think the approach in this study is great for assessing specific, well-observed glaciers. However, I question some broader claims of scalability and transferability to other regions.

We agree with this comment, there are some limits. It should be noted that the Crocus model has already been used in other regions than the European Alps to model the snow cover, *e.g.*, Gaillard et al. (2025); Luijting et al. (2018); Fréville et al. (2014); Lejeune et al. (2007). When claiming the scalability and transferability, we were indeed implicitly referring to monitored areas (*e.g.*, the European Alps) where the meteorological forcing (including solid precipitations) have less uncertainty than in remote areas.

We modified the manuscript accordingly: “The simulation setup of this study is easily transferable in time and space for glaciers where measurements of winter solid precipitation and summer melt exist. An automatic weather station is also useful to ensure the accuracy of the meteorological forcings used in the model. As a result, this work paves the way towards a comprehensive regional assessment of the impacts of LAPs on glaciers in monitored regions.”

**R2.14.** Line 447: Do you mean the sensible heat flux? Latent heat fluxes are generally negative during melt.

Thank you for this remark on the following sentence of the initial version: “Hence, the impact of dust alone at the glacier-wide scale in summer is likely higher than the impact of the latent heat flux, which represents a few percents of the total melt.” We wanted here to focus on the mass loss due to the latent heat flux, *i.e.*, negative latent heat flux, *i.e.*, sublimation. However, this sentence is poorly link with the scope of the study. We removed that sentence in the revised version of the manuscript.

### 2.2.3 Technical corrections

**R2.15.** Line 1: color → turn (Just a recommendation for slightly smoother wording. Also consider “... deposits frequently darken alpine glaciers”)

We changed from ‘color’ to ‘turn’, and mentioned the darkening afterwards: “Saharan dust deposits frequently turn alpine glaciers orange and darken their surface.”

**R2.16.** Line 15: to account → accounting

We modified accordingly “Hence, we recommend accounting for impurities to simulate the distributed surface mass balance of glaciers.”

**R2.17.** Line 588: depositions → deposition, radiations → radiation

We modified accordingly “... because of: (i) higher dust deposition; and (ii) higher shortwave radiation.”

**R2.18.** Figure 5c: Change month abbreviations to English (Jan., Jul.) or spell out the month. Also, decrease the size of x-axis labels (or rotate slightly) on red/purple plots on the right. Currently, it looks like “19202122” instead of “19 20 21 22”. Please specify that this is year.

Thanks. We switched to English. An indeed the years were not readable, we tuned the x-axis labels and added this information in the legend: **"Double digits refers to the years."**

**R2.19.** Line 620: their  $\rightarrow$  its

We modified accordingly "... for the period 2019-2022, **its** impact can not be totally neglected...".

### 3 Authors additional corrections

During the review process, we identified some inaccuracies that have been corrected, and some perspectives of evolution that have been investigated in this revised version. They are listed and detailed below, together with the corresponding modifications implemented in the revised version of the manuscript. While these corrections (especially the point 1) slightly change some of the numbers related to the quantification of dust impact, they do not alter the conclusions of the paper.

1. In the initial version of the paper, we did not redistribute the ALADIN wet and dry deposition fluxes with respect to the SAFRAN precipitations, hence deposited LAP were lost (*e.g.*, when ALADIN wet depositions without SAFRAN precipitations). We added this redistribution in this revised version following the method of Réveillet et al. (2022). Such a modification increased the simulated concentration of dust and BC at the snow surface, hence lowering albedo and advancing snow melt-out date compared to the initial version. So we adjusted and lowered (i) the mass absorption efficiency (MAE) of LAP of dust to  $[\text{MAE}_{\text{dust}}(400 \text{ nm}) = 71 \cdot 10^{-3} \text{ m}^2/\text{g}; \text{AAE}_{\text{dust}} = 3.2]$  and the MAE of BC at 550 nm to  $7.5 \text{ m}^2/\text{g}$ , (ii) the roughness length of snow  $z_{0,\text{snow}}$  from  $5 \cdot 10^{-4} \text{ m}$  to  $10^{-4} \text{ m}$ , now better reproducing the ratio  $z_{0,\text{snow}}/z_{0,\text{ice}}$  of 10 in Réveillet et al. (2018), and of  $\approx 7$  in Gerbaux et al. (2005).

To assess the accuracy of this revised parametrization, we tested different values of MAE (presented in Fig. 5), and we evaluated them using the following evaluation metrics:

- comparison to the summer melt measured at the stakes (*in the initial version*)
- comparison to the surface type evaluated from S2 (*in the initial version*)
- comparison to the broadband albedo derived from S2 (Fig. 6 b) recommended by reviewer #2 (*new in this review*)
- comparison to the broadband albedo measured at the AWS (Fig. 7) (*new in this review*)

Table 3 summarizes the evaluation results. The new parametrization used in this revised version showed accurate results for all the evaluation metrics. The summer melts of the initial parametrization of MAE were too large compared to the stake measurements, and the simulated albedo are lower than the albedo retrieved from S2. Using low values of dust MAE (low parametrization), melt metrics were the best, however the comparison against S2 albedo and the AWS SMOD showed worse results than for the new parametrization. So, the new parametrization appeared as the best trade-off. The new MAE values were therefore used as reference in this revised version.

The MAE values of this revised version were lower than that of the initial version (Fig. 5). However, the initial MAE values (from Tuzet et al. (2020); Réveillet et al. (2022)) were used in in the framework of a seasonal snowpack at intermediate elevations (1'500 to 3'000 m a.s.l.). On glaciers, the LAP surface concentration may reach higher values due to longer melt seasons. This could likely lead to more aggregation of LAPs in the snowpack, which decrease their MAE (*e.g.*, Bond, Bergstrom (2006); Schwarz et al. (2013)). Therefore, it is reasonable to find lower optimal MAE on glaciers.

As a result, those revised MAE values did not affect the conclusions of this article.

We propose the following modifications. For the method section 2.3.1: **"ALADIN wet and dry deposition were redistributed with respect to the the SAFRAN precipitation as in Réveillet et al. (2022), *i.e.*, wet depositions when SAFRAN precipitations are positive, dry depositions otherwise."** The revised values of the dust MAE have been integrated in the manuscript as indicated under comment R1.14., and the values of low and high dust MAE were also integrated in the manuscript as indicated under comment R2.7. We removed words from section 2.4.1: ~~To compute the albedo of snow, the refractive indices of ice, mineral dust and BC were was set as in Réveillet et al. (2022), which quantified the impact of LAPs in the same region as in this study (the~~

European Alps). Particularly, the mass absorption efficiency of dust [...]” and to discuss the choice of MAE, we added the following discussion in section 4.1: **”The MAE of dust used in this study was slightly lower than initial values used in previous studies in the framework of a seasonal alpine snowpack at intermediate elevations (Tuzet et al., 2020; Réveillet et al., 2022). The reliability of this parametrization was assessed by the evaluations of section 3.1 and the evaluation of the modelled snow albedo in Appendix Fig. 3. Such lower MAE implicitly accounts for the probable aggregation of LAP (likely due their high concentrations during long melt seasons) that lowers the MAE (e.g., Bond, Bergstrom (2006); Schwarz et al. (2013)).”**

The numeric values, concerning the results, of Sections 2.3.5, 3.1, 3.2, 3.3, 4, 4.1, 4.2, 4.3 have been modified (highlighted in the track change version attached to this response). Figures 3, 4, 5, 6, 7, 8 and Appendix Fig. D1, E1, E2, E3 have also been modified accordingly. In section 4.2, the word ”almost” was removed. In section 4.3, CPU costs were adjusted to account for the addition of the sensitivity tests.

2. We modified the unit of the third row of Figure E1, from ”m w.e.” to **”%**”.

3. The total melt was computed as the difference between the minimum and the maximum of SWE during the year, it is now computed accordingly to the formulation stated in section 2.5 of the manuscript, *i.e.*, as the sum of every snow or ice melt of the year.

4. We cited a recent article in the introduction (Menounos et al., 2025) dealing with glacier and LAP: ”By lowering the albedo, LAPs contribute to significantly accelerate snow and glacier melt around the world, *e.g.*, [...] **Menounos et al. (2025).**”

5. All  $k_{BC}$  were adjusted to  $k_{BC}$ .

6. In the Conclusion section, **”This year,”** was added to link to the previous sentence.

7. In the Acknowledgements section, **”Ghislain Picard”** was added.

8. Code and data availability have been revised, now providing access to the code repository, and forcing files in a Zenodo repository.




Figure 5: Numerical values used for mass absorption efficiencies (MAE) for both dust and BC. New and old values are respectively the ones of the revised and the initial version. The MAE of dust are from Caponi et al. (2017) with couples  $[\text{MAE}_{\text{dust}}(\lambda_0); \text{AAE}_{\text{dust}}]$  being respectively equal to  $[110 \cdot 10^{-3} \text{ m}^2/\text{g}; 4.1]$ ,  $[71 \cdot 10^{-3} \text{ m}^2/\text{g}; 3.2]$ ,  $[27 \cdot 10^{-3} \text{ m}^2/\text{g}; 3.3]$  and  $[630 \cdot 10^{-3} \text{ m}^2/\text{g}; 3.4]$  for old, new, low and high parametrization. The old BC MAE was an enhanced MAE of  $11.25 \text{ m}^2/\text{g}$  at 550 nm previously used in Tuzet et al. (2020) to ”implicitly account for the potential absorption enhancement due to internal particle mixing or particle coating”. The new BC MAE of  $7.5 \text{ m}^2/\text{g}$  at 550 nm for BC is a non-enhanced MAE close to the external-mixing of BC used in SNICAR (Flanner et al., 2021).




Figure 6: Comparison between modelled broadband albedo and observed broadband S2 albedo using the method of Liang et al. (2003) ( $S2 \text{ albedo} = 0.356 * R_2 + 0.130 * R_4 + 0.373 * R_5 + 0.085 * R_{11} + 0.072 * R_{12} - 0.018$  where  $R_i$  is the reflectance of Sentinel-2 band  $i$ ). All S2 dates mentioned in Table B2 were used. Only the points where the surface type was snow for both the model and S2 were used. a) Using the new MAE parametrization of the model, blue point are for slopes  $\leq 20^\circ$ , and red points for slopes  $> 20^\circ$ . b) Using only slopes  $\leq 20^\circ$  for the different model MAE parametrizations found plotted in Fig. 7. Large opaque dots are the means for each date.




Figure 7: Albedo at the AWS glacier location. Albedo AWS (green crosses) is the measured albedo at the AWS computed as the ratio of reflected and incoming shortwave radiation between 9:00AM and 15:00PM. The simulated albedo (coloured contour envelopes) are the albedo values at noon plotted for the 4 simulation points near the AWS glacier.

$z_{0,\text{snow}}$ (mm)	MAE dust, MAE BC	stakes summer mean error				stakes summer RMSE				vs. Sentinel-2		vs. AWS	
		2019	2020	2021	2022	2019	2020	2021	2022	snow albedo	surface type	snow albedo	SMOD
0.1	high, new	1.12	0.94	0.97	1.08	1.30	1.08	1.09	1.38	too low	too much ice	too low	5–6 d early
0.1	old, old	0.67	0.28	0.40	0.60	0.89	0.52	0.56	0.97	too low	OK	too low	2 d late
0.1	new, new	0.49	0.10	0.25	0.46	0.74	0.43	0.45	0.86	OK	more snow	OK	4 d late
0.2	new, new	0.57	0.19	0.31	0.57	0.78	0.44	0.48	0.90	OK	OK	?	?
0.05	new, new	0.33	-0.08	0.12	0.24	0.69	0.47	0.38	0.84	OK	more snow	?	?
0.1	low, new	0.33	-0.16	0.02	0.28	0.63	0.44	0.31	0.76	too high	too much snow	OK	6–9 d late

Table 1: Summary of the evaluations for different parametrizations. See Fig. 5 for the numeric MAE values. Melt evaluation against the stakes measurements (in m w.e.) and surface type evaluation against S2 are the metrics initially used in Figure 3 of the manuscript. SMOD stands for snow melt-out date. Evaluations, except evaluation with stakes, are qualitative. Question marks stand for uncalculated metrics.

## References

- Bond Tami C., Bergstrom Robert W.* Light Absorption by Carbonaceous Particles: An Investigative Review // *Aerosol Science and Technology*. I 2006. 40, 1. 27–67. Publisher: Taylor & Francis .eprint: <https://doi.org/10.1080/02786820500421521>.
- Caponi Lorenzo, Formenti Paola, Massabó Dario, Di Biagio Claudia, Cazaunau Mathieu, Pangui Edouard, Chevaillier Servanne, Landrot Gautier, Andreae Meinrat O., Kandler Konrad, Piketh Stuart, Saeed Thuraya, Seibert Dave, Williams Earle, Balkanski Yves, Prati Paolo, Doussin Jean-François.* Spectral- and size-resolved mass absorption efficiency of mineral dust aerosols in the shortwave spectrum: a simulation chamber study // *Atmospheric Chemistry and Physics*. VI 2017. 17, 11. 7175–7191. Publisher: Copernicus GmbH.
- Cluzet Bertrand, Revuelto Jesus, Lafaysse Matthieu, Tuzet François, Cosme Emmanuel, Picard Ghislain, Arnaud Laurent, Dumont Marie.* Towards the assimilation of satellite reflectance into semi-distributed ensemble snowpack simulations // *Cold Regions Science and Technology*. II 2020. 170. 102918.
- Collaud Coen M., Weingartner E., Schaub D., Hueglin C., Corrigan C., Henning S., Schwikowski M., Baltensperger U.* Saharan dust events at the Jungfraujoch: detection by wavelength dependence of the single scattering albedo and first climatology analysis // *Atmospheric Chemistry and Physics*. XII 2004. 4, 11/12. 2465–2480. Publisher: Copernicus GmbH.
- Collaud Coen Martine, Brem Benjamin Tobias, Gysel-Beer Martin, Modini Robin, Henne Stephan, Steinbacher Martin, Putero Davide, Gini Maria I., Eleftheriadis Kostantinos.* Detection and climatology of Saharan dust frequency and mass at the Jungfraujoch (3580 m asl, Switzerland) // *EGUsphere*. IX 2025. 1–35. Publisher: Copernicus GmbH.
- Conway H., Gades A., Raymond C. F.* Albedo of dirty snow during conditions of melt // *Water Resources Research*. 1996. 32, 6. 1713–1718. .eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/96WR00712>.
- Doherty Sarah J., Grenfell Thomas C., Forsström Sanja, Hegg Dean L., Brandt Richard E., Warren Stephen G.* Observed vertical redistribution of black carbon and other insoluble light-absorbing particles in melting snow // *Journal of Geophysical Research: Atmospheres*. 2013. 118, 11. 5553–5569. .eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1002/jgrd.50235>.
- Dumont M., Brissaud O., Picard G., Schmitt B., Gallet J.-C., Arnaud Y.* High-accuracy measurements of snow Bidirectional Reflectance Distribution Function at visible and NIR wavelengths – comparison with modelling results // *Atmospheric Chemistry and Physics*. III 2010. 10, 5. 2507–2520. Publisher: Copernicus GmbH.
- Dumont M., Tuzet F., Gascoïn S., Picard G., Kutuzov S., Lafaysse M., Cluzet B., Nheili R., Painter T. H.* Accelerated Snow Melt in the Russian Caucasus Mountains After the Saharan Dust Outbreak in March 2018 // *Journal of Geophysical Research: Earth Surface*. 2020. 125, 9. e2020JF005641. .eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2020JF005641>.
- Dumont Marie, Durand Yves, Arnaud Yves, Six Delphine.* Variational assimilation of albedo in a snowpack model and reconstruction of the spatial mass-balance distribution of an alpine glacier // *Journal of Glaciology*. I 2012. 58, 207. 151–164. Publisher: Cambridge University Press.
- Dumont Marie, Gascoïn Simon, Réveillet Marion, Voisin Didier, Tuzet François, Arnaud Laurent, Bonnefoy Mylène, Bacardit Peñarroya Montse, Carmagnola Carlo, Deguine Alexandre, Diacre Aurélie, Dürr Lukas, Evrard Olivier, Fontaine Firmin, Frankl Amaury, Fructus Mathieu, Gandois Laure, Gouttevin Isabelle, Gherab Abdelfateh, Hagenmuller Pascal, Hansson Sophia, Herbin Hervé, Josse Béatrice, Jourdain Bruno, Lefevre Irene, Le Roux Gaël, Libois Quentin, Liger Lucie, Morin Samuel, Petitprez Denis, Robledano Alvaro, Schneebeli Martin, Salze Pascal, Six Delphine, Thibert Emmanuel, Trachsel Jürg, Vernay Matthieu, Viallon-Galinier Léo, Voiron Céline.* Spatial variability of Saharan dust deposition revealed through a citizen science campaign // *Earth System Science Data*. VII 2023. 15, 7. 3075–3094. Publisher: Copernicus GmbH.

- Etchevers Pierre, Martin Eric, Brown Ross, Fierz Charles, Lejeune Yves, Bazile Eric, Boone Aaron, Dai Yong-Jiu, Essery Richard, Fernandez Alberto, Gusev Yeugeniy, Jordan Rachel, Koren Victor, Kowalczyk Eva, Nasonova N. Olga, Pyles R. David, Schlosser Adam, Shmakin Andrey B., Smirnova Tatiana G., Strasser Ulrich, Verseghy Diana, Yamazaki Takeshi, Yang Zong-Liang.* Validation of the energy budget of an alpine snowpack simulated by several snow models (Snow MIP project) // *Annals of Glaciology*. I 2004. 38. 150–158.
- Flanner M. G., Liu X., Zhou C., Penner J. E., Jiao C.* Enhanced solar energy absorption by internally-mixed black carbon in snow grains // *Atmospheric Chemistry and Physics*. V 2012. 12, 10. 4699–4721. Publisher: Copernicus GmbH.
- Flanner Mark G., Arnheim Julian B., Cook Joseph M., Dang Cheng, He Cenlin, Huang Xianglei, Singh Deepak, Skiles S. McKenzie, Whicker Chloe A., Zender Charles S.* SNICAR-ADv3: a community tool for modeling spectral snow albedo // *Geoscientific Model Development*. XII 2021. 14, 12. 7673–7704. Publisher: Copernicus GmbH.
- Flanner Mark G., Zender Charles S., Randerson James T., Rasch Philip J.* Present-day climate forcing and response from black carbon in snow // *Journal of Geophysical Research: Atmospheres*. 2007. 112, D11. \_eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2006JD008003>.
- Fréville H., Brun E., Picard G., Tatarinova N., Arnaud L., Lanconelli C., Reijmer C., Broeke M. van den.* Using MODIS land surface temperatures and the Crocus snow model to understand the warm bias of ERA-Interim reanalyses at the surface in Antarctica // *The Cryosphere*. VII 2014. 8, 4. 1361–1373. Publisher: Copernicus GmbH.
- Gaillard Manon, Vionnet Vincent, Lafaysse Matthieu, Dumont Marie, Ginoux Paul.* Improving large-scale snow albedo modeling using a climatology of light-absorbing particle deposition // *The Cryosphere*. II 2025. 19, 2. 769–792. Publisher: Copernicus GmbH.
- Gerbaux M., Genthon C., Etchevers P., Vincent C., Dedieu J. P.* Surface mass balance of glaciers in the French Alps: distributed modeling and sensitivity to climate change // *Journal of Glaciology*. I 2005. 51, 175. 561–572. Publisher: Cambridge University Press.
- Günther Daniel, Marke Thomas, Essery Richard, Strasser Ulrich.* Uncertainties in Snowpack Simulations—Assessing the Impact of Model Structure, Parameter Choice, and Forcing Data Error on Point-Scale Energy Balance Snow Model Performance // *Water Resources Research*. 2019. 55, 4. 2779–2800. \_eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2018WR023403>.
- He Cenlin, Flanner Mark G., Chen Fei, Barlage Michael, Liou Kuo-Nan, Kang Shichang, Ming Jing, Qian Yun.* Black carbon-induced snow albedo reduction over the Tibetan Plateau: uncertainties from snow grain shape and aerosol–snow mixing state based on an updated SNICAR model // *Atmospheric Chemistry and Physics*. VIII 2018. 18, 15. 11507–11527. Publisher: Copernicus GmbH.
- Hoesly Rachel M., Smith Steven J., Feng Leyang, Klimont Zbigniew, Janssens-Maenhout Greet, Pitkanen Tyler, Seibert Jonathan J., Vu Linh, Andres Robert J., Bolt Ryan M., Bond Tamì C., Dawidowski Laura, Kholod Nazar, Kurokawa June-ichi, Li Meng, Liu Liang, Lu Zifeng, Moura Maria Cecilia P., O'Rourke Patrick R., Zhang Qiang.* Historical (1750–2014) anthropogenic emissions of reactive gases and aerosols from the Community Emissions Data System (CEDS) // *Geoscientific Model Development*. I 2018. 11, 1. 369–408. Publisher: Copernicus GmbH.
- Lafaysse Matthieu, Cluzet Bertrand, Dumont Marie, Lejeune Yves, Vionnet Vincent, Morin Samuel.* A multiphysical ensemble system of numerical snow modelling // *The Cryosphere*. V 2017. 11, 3. 1173–1198. Publisher: Copernicus GmbH.
- Lamare Maxim, Dumont Marie, Picard Ghislain, Larue Fanny, Tuzet François, Delcourt Clément, Arnaud Laurent.* Simulating optical top-of-atmosphere radiance satellite images over snow-covered rugged terrain // *The Cryosphere*. XI 2020. 14, 11. 3995–4020. Publisher: Copernicus GmbH.
- Lamboll Robin D., Jones Chris D., Skeie Ragnhild B., Fiedler Stephanie, Samset Bjørn H., Gillett Nathan P., Rogelj Joeri, Forster Piers M.* Modifying emissions scenario projections to account for the effects of COVID-19: protocol for CovidMIP // *Geoscientific Model Development*. VI 2021. 14, 6. 3683–3695. Publisher: Copernicus GmbH.

- Lejeune Yves, Bouilloud Ludovic, Etchevers Pierre, Wagon Patrick, Chevallier Pierre, Sicart Jean-Emmanuel, Martin Eric, Habets Florence.* Melting of Snow Cover in a Tropical Mountain Environment in Bolivia: Processes and Modeling // *Journal of Hydrometeorology*. VIII 2007. 8, 4. 922–937. Publisher: American Meteorological Society Section: *Journal of Hydrometeorology*.
- Liang Shunlin, Shuey Chad J, Russ Andrew L, Fang Hongliang, Chen Mingzhen, Walthall Charles L, Daughtry Craig S. T, Hunt Raymond.* Narrowband to broadband conversions of land surface albedo: II. Validation // *Remote Sensing of Environment*. I 2003. 84, 1. 25–41.
- Luijting Hanneke, Vikhamar-Schuler Dagrun, Aspelien Trygve, Bakketun Åsmund, Homleid Mariken.* Forcing the SURFEX/Crocus snow model with combined hourly meteorological forecasts and gridded observations in southern Norway // *The Cryosphere*. VI 2018. 12, 6. 2123–2145. Publisher: Copernicus GmbH.
- Marle Margreet J. E. van, Kloster Silvia, Magi Brian I., Marlon Jennifer R., Daniau Anne-Laure, Field Robert D., Arneth Almut, Forrest Matthew, Hantson Stijn, Kehrwald Natalie M., Knorr Wolfgang, Lasslop Gitta, Li Fang, Mangeon Stéphane, Yue Chao, Kaiser Johannes W., Werf Guido R. van der.* Historic global biomass burning emissions for CMIP6 (BB4CMIP) based on merging satellite observations with proxies and fire models (1750–2015) // *Geoscientific Model Development*. IX 2017. 10, 9. 3329–3357. Publisher: Copernicus GmbH.
- Menounos Brian, Huss Matthias, Marshall Shawn, Ednie Mark, Florentine Caitlyn, Hartl Lea.* Glaciers in Western Canada-Conterminous US and Switzerland Experience Unprecedented Mass Loss Over the Last Four Years (2021–2024) // *Geophysical Research Letters*. 2025. 52, 12. e2025GL115235. [\\_eprint: https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2025GL115235](https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2025GL115235).
- Michou M., Nabat P., Saint-Martin D.* Development and basic evaluation of a prognostic aerosol scheme (v1) in the CNRM Climate Model CNRM-CM6 // *Geoscientific Model Development*. III 2015. 8, 3. 501–531. Publisher: Copernicus GmbH.
- Michou M., Nabat P., Saint-Martin D., Bock J., Decharme B., Mallet M., Roehrig R., Séférian R., Sénési S., Voldoire A.* Present-Day and Historical Aerosol and Ozone Characteristics in CNRM CMIP6 Simulations // *Journal of Advances in Modeling Earth Systems*. 2020. 12, 1. e2019MS001816. [\\_eprint: https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2019MS001816](https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2019MS001816).
- Morcrette J.-J., Boucher O., Jones L., Salmond D., Bechtold P., Beljaars A., Benedetti A., Bonet A., Kaiser J. W., Razingar M., Schulz M., Serrar S., Simmons A. J., Sofiev M., Suttie M., Tompkins A. M., Untch A.* Aerosol analysis and forecast in the European Centre for Medium-Range Weather Forecasts Integrated Forecast System: Forward modeling // *Journal of Geophysical Research: Atmospheres*. 2009. 114, D6. [\\_eprint: https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2008JD011235](https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2008JD011235).
- Nabat Pierre, Somot Samuel, Cassou Christophe, Mallet Marc, Michou Martine, Bouniol Dominique, Decharme Bertrand, Drugé Thomas, Roehrig Romain, Saint-Martin David.* Modulation of radiative aerosols effects by atmospheric circulation over the Euro-Mediterranean region // *Atmospheric Chemistry and Physics*. VII 2020. 20, 14. 8315–8349. Publisher: Copernicus GmbH.
- Naegeli Kathrin, Huss Matthias, Hoelzle Martin.* Change detection of bare-ice albedo in the Swiss Alps // *The Cryosphere*. II 2019. 13, 1. 397–412. Publisher: Copernicus GmbH.
- O'Rourke Patrick, Smith Steven, Mott Andrea, Ahsan Hamza, Mcduffie Erin, Crippa Monica, Klimont Zbigniew, Mcdonald Brian, Wang Shuxiao, Nicholson Matthew, Hoesly Rachel, Feng Leyang.* CEDS v.2021.04.21 Gridded emissions data. 2021.
- Picard Ghislain, Libois Quentin.* Simulation of snow albedo and solar irradiance profile with the Two-streAm Radiative TransfEr in Snow (TARTES) v2.0 model // *Geoscientific Model Development*. XII 2024. 17, 24. 8927–8953. Publisher: Copernicus GmbH.
- Randerson J. T., Werf G. R. van der, Giglio L., Collatz G. J., Kasibhatla P. S.* Global Fire Emissions Database, Version 4.1 (GFEDv4) // ORNL Distributed Active Archive Center (DAAC) dataset 10.3334/ORNLDAAC/1293 (2017. I 2017. 1293. ADS Bibcode: 2017ornl.data.1293R.

- Réveillet Marion, Dumont Marie, Gascoïn Simon, Lafaysse Matthieu, Nabat Pierre, Ribes Aurélien, Nheili Rafife, Tuzet François, Ménégoz Martin, Morin Samuel, Picard Ghislain, Ginoux Paul.* Black carbon and dust alter the response of mountain snow cover under climate change // *Nature Communications*. IX 2022. 13, 1. 5279. Number: 1 Publisher: Nature Publishing Group.
- Réveillet Marion, Six Delphine, Vincent Christian, Rabatel Antoine, Dumont Marie, Lafaysse Matthieu, Morin Samuel, Vionnet Vincent, Litt Maxime.* Relative performance of empirical and physical models in assessing the seasonal and annual glacier surface mass balance of Saint-Sorlin Glacier (French Alps) // *The Cryosphere*. IV 2018. 12, 4. 1367–1386. Publisher: Copernicus GmbH.
- Schwarz J. P., Gao R. S., Perring A. E., Spackman J. R., Fahey D. W.* Black carbon aerosol size in snow // *Scientific Reports*. III 2013. 3, 1. 1356. Publisher: Nature Publishing Group.
- Skiles S. McKENZIE, Painter Thomas.* Daily evolution in dust and black carbon content, snow grain size, and snow albedo during snowmelt, Rocky Mountains, Colorado // *Journal of Glaciology*. II 2017. 63, 237. 118–132.
- Tuzet François, Dumont Marie, Picard Ghislain, Lamare Maxim, Voisin Didier, Nabat Pierre, Lafaysse Mathieu, Larue Fanny, Revuelto Jesus, Arnaud Laurent.* Quantification of the radiative impact of light-absorbing particles during two contrasted snow seasons at Col du Lautaret (2058&thinsp;m&thinsp;a.s.l., French Alps) // *The Cryosphere*. XII 2020. 14, 12. 4553–4579. Publisher: Copernicus GmbH.
- Vernay Matthieu, Lafaysse Matthieu, Monteiro Diego, Hagenmuller Pascal, Nheili Rafife, Samacoïts Raphaëlle, Verfaillie Deborah, Morin Samuel.* The S2M meteorological and snow cover reanalysis over the French mountainous areas: description and evaluation (1958–2021) // *Earth System Science Data*. IV 2022. 14, 4. 1707–1733. Publisher: Copernicus GmbH.