

Bacteriohopanepolyols track past environmental transitions in the Black Sea

Anna Cutmore^{1,2*}, Nora Richter^{1*}, Nicole Bale¹, Stefan Schouten^{1,4}, and Darci Rush¹

*Joint first authors

¹NIOZ Royal Netherlands Institute for Sea Research, Texel, the Netherlands

²Durham University, Department of Geography, Durham, United Kingdom

³EAWAG Swiss Federal Institute of Aquatic Science and Technology, Dübendorf, Switzerland

⁴Utrecht University, Utrecht, the Netherlands

Correspondence to: Anna Cutmore, anna.v.cutmore@durham.ac.uk and Nora Richter, nora.richter@eawag.ch

Abstract

Bacteriohopanepolyols (BHPs) are structurally diverse compounds produced by a wide range of bacteria making them ideal candidates as chemotaxonomic biomarkers and indicators of bacterially-driven biogeochemical processes in the geological record. In this study, we characterize changes in the BHP distribution in the Black Sea over the past 20 thousand years (ka), as the basin underwent three distinct environmental phases: (i) an oxic lacustrine phase where the Black Sea was disconnected from the global ocean; (ii) a transition period marked by the initial influx of marine water into the basin; and (iii) a marine phase where the basin was permanently euxinic. During the lacustrine phase we observe a high abundance and diversity of nucleoside BHPs (Nu-BHPs) that are likely derived from elevated terrigenous inputs as well as production of Nu-BHPs in the brackish-to-fresh water column. The transition phase is marked by a decrease in the abundance of most Nu-BHPs and an increase in the abundance of methoxylated-BHPs as well as BHPs such as aminobacteriohopanetriol which are ubiquitous across a wide range of environments including soils as well as marine and freshwater settings. The euxinic marine phase (7.2 ka-present) can be divided into two stages based on changes in BHP composition. The early stage is characterised by a high abundance of aminobacteriohopanetetrol and aminobacteriohopanepentol, which were likely produced by methanotrophs at the oxycline. A shallow oxycline likely allowed for increased transport of these BHPs to the sediment. The later marine phase is characterised by a decline in these BHPs, likely due to a deepening of the oxycline and reduced transport of BHPs from the oxycline to the sediment. The changes in BHP distributions throughout the record may either be attributed to shifts in the bacterial communities or physiological adaptations of bacteria to the changing environment. Throughout the record, diagenetic products of BHPs (e.g., anhydro-bacteriohopanetetrol) were detected. These degradation products, however, remain a small proportion of the overall BHP composition, indicating excellent preservation conditions throughout the record. This study offers new insights into changes in microbial communities and biogeochemical processes that occurred in the Black Sea during the Last Deglaciation and Holocene in response to substantial shifts in the hydrology and oxygen conditions of the basin.

Keywords: Bacteriohopanepolyols, Black Sea, Holocene, Last Deglaciation, Anoxia, Euxinia

Deleted: us

1. Introduction

Bacteriohopanepolyols (BHPs) are pentacyclic triterpenoids produced by a wide-range of bacteria that are considered to be the precursors to one of the most abundant lipids in the geological record, hopanoids (Rohmer et al., 1984; Ourisson & Albrecht, 1992). BHPs themselves are also abundant in the geological record and are well-preserved in sedimentary archives for >1 million years (Handley et al., 2010). In some cases, BHPs are linked to certain source organisms (Cvejic et al., 2000; Kool et al., 2014; Rush et al., 2014; van Winden et al., 2012) or biogeochemical processes (Talbot et al., 2003; 2007; Coolen et al., 2008; Blumenberg et al., 2013). Consequently, shifts in BHP compositions in the geological record have become a useful tool for palaeoclimate reconstructions to assess changes in redox conditions (Blumenberg et al., 2013; Matys et al., 2017; Rush et al., 2019; Zindorf et al., 2020), salinity (Coolen et al., 2008), methanotrophy (Talbot et al., 2014), and soil inputs (Blumenberg et al., 2013). For example, 35-aminobacteriohopane-31,32,33,34-tetrol (aminotetrol from herein), 35-aminobacteriohopane-30,31,32,33,34-pentol (aminopentol from herein), and methylcarbamate-amino BHPs are primarily associated with aerobic methane-oxidizing bacteria (MOB; Rohmer et al., 1984; Jahnke et al., 1999; Cvejic et al., 2000; Talbot et al., 2001; Zhu et al., 2011; van Winden et al., 2012; Rush et al., 2016) and have been used to reconstruct past changes in methane-oxidation in the Congo River Delta (Talbot et al., 2014; Spencer-Jones et al., 2017), the Baltic Sea (Blumenberg et al., 2013), and in an Antarctic lake (Coolen et al., 2008). Recent studies show that bacteriohopane-32,33,34,35-tetrol-x (BHT-x) is synthesized by anaerobic ammonium oxidation (anammox) bacteria (Rush et al., 2014; Schwartz-Narbonne et al., 2020) and thus has been used to reconstruct changes in N-cycling under low-oxygen conditions in Mediterranean sapropels (Rush et al., 2019; van Kemenade et al., 2023) and the Black Sea (Cutmore et al., 2025). Nucleoside BHPs (Nu-BHPs) occur in high abundance in soils (Seemann et al., 1999; Bravo et al., 2001; Cooke et al., 2008a; Xu et al., 2009; Rethemeyer et al., 2010), and, therefore, have been applied to track the transport of soil organic matter to aquatic sediments (Talbot & Farrimond, 2007; Cooke et al., 2008b). Recent studies, however, have identified potential production of Nu-BHPs along redoxclines in marine oxygen minimum zones (Kusch et al., 2021) and near the oxycline in lakes (Richter et al., 2023). While this highlights a small number of the known BHPs and their potential application to paleo-records, there are still many unknowns surrounding the origin and sources of other BHPs in the environment. Recent analytical advancements in BHP analysis have led to the identification of additional novel BHPs in both cultures and environmental samples (Talbot et al., 2016a; Hopmans et al., 2021; Richter et al., 2023). The large structural diversity of BHPs along with their preservation potential in the recent geological record, points to a useful new tool in biomarker research that can provide insights into past changes in microbial communities and/or biogeochemical processes.

The Black Sea provides a compelling location to study the impact of bacterial community composition and environmental changes on BHP composition due to its unique hydrological history and present-day characteristics. Today, it is the largest permanently stratified anoxic basin in the world with limited connection to the global ocean through the Bosphorus Strait. Over the past 20 thousand years (ka), however, the Black Sea underwent significant hydrological changes (Fig. 1). During the Last Glacial Maximum (LGM), the basin was an oxygenated brackish-to-

Deleted: has

Deleted:

76 freshwater environment (Schrader, 1979), then, during the subsequent deglaciation, the basin experienced many
77 environmental shifts, including changes in temperature (Bahr et al., 2005; 2008; Ion et al., 2022), water-level
78 (Ivanova et al., 2007; Nicholas et al., 2011; Piper and Calvert, 2011), and freshwater input. This freshwater influx was
79 influenced by changes in regional precipitation and melting of Eurasian ice sheets and alpine glaciers (Bahr et al.,
80 2005; 2006; 2008; Badertscher et al., 2011; Shumilovskikh et al., 2012). The basin reconnected with the global ocean
81 around 9.6 ka when post-glacial sea-level rise led to an initial marine inflow (IMI) over the Bosphorus sill (Aksu et al.,
82 2002; Major et al., 2006; Bahr et al., 2008; Ankindinova et al., 2019), leading to ~~higher~~ salinity of the water column
83 (Marret et al., 2009; Verleye et al., 2009; Filipova-Marinova et al., 2013) and permanent euxinia developing in the
84 water column after ~7.2 ka (Arthur and Dean, 1998; Eckert et al., 2013).

Deleted: enhanced

85 Several studies have demonstrated the usefulness of lipid biomarkers as records of biogeochemical
86 processes and as indicators of ecological niches within the modern-day water column and surface sediments of the
87 Black Sea (Schubert et al., 2006; Coolen et al., 2007; Wakeham et al., 2007; Schubotz et al., 2009; Sollai et al., 2019;
88 Bale et al., 2021; Kusch et al., 2022). BHPs produced in the lower suboxic zone and upper sulfidic zone, for instance,
89 were shown to record activities of aerobic methanotrophs and anammox bacteria, respectively, demonstrating their
90 reliability as potential biomarkers for microbial communities (Kusch et al., 2022). Furthermore, previous work has
91 explored BHP changes in the Black Sea over the Holocene, however, this applies to a limited number of ~~BHPs~~
92 (Blumenberg et al., 2009a). With significant advances in analytical techniques, there is now the opportunity to
93 explore the BHP geolipidome of the Black Sea in unprecedented detail. Indeed, we recently used the ratio of
94 bacteriohopanetetrol (BHT)-34S and the later eluting stereoisomer BHT-x to trace past anammox activity in the Black
95 Sea over the last 20 ka, complementing other nitrogen cycling biomarker tools (Cutmore et al., 2025). In this study,
96 we now fully characterize the distribution of BHPs in a Black Sea sediment core over the Last Deglaciation and
97 Holocene to shed light on changing environmental conditions and bacterial communities.

Deleted: compounds

Deleted:

98 99 2. Methods

100 Piston core 64PE418 (235 cm length) was recovered from the Black Sea (42°56' N, 30°02' E; 1970 ~~metres~~
101 ~~below sea level (mbsl)~~) during the 2017 Pelagia cruise (Fig. 1). The core was subsampled at the Royal Netherlands
102 Institute for Sea Research, and all samples were stored frozen until further analysis. The contemporary Black Sea
103 water column is characterized by pronounced redox gradients. At the location of the core site in the western gyre of
104 the basin, the oxic zone, found between 0 – 75 m depth range, has an oxygen concentration of ~121 µmol/kg at 50
105 m depth, while sulphide concentrations are not detected (Sollai et al., 2019). The suboxic zone lies below, situated
106 between 75 – 115 m depth range, with traces of sulphide found at the bottom of this layer between 105 – 115 m
107 (Sollai et al., 2019). Beneath this is the euxinic zone, located between 115 – 2000 m depth range, with sulphide
108 concentrations increasing significantly with depth, reaching ~400 µmol/L at 2,000 m water depth (Sollai et al., 2019).

Deleted: mbsl depth

109 110 2.1 Lipid extraction and analysis

Sediment samples (n=44) were taken at 5 cm intervals throughout the core's depth before freeze drying. Using a modified Bligh and Dyer extraction method (as described previously by Bale et al., 2021), lipids were extracted from the dry sediment samples, as described in detail in Cutmore et al. (2025). Extraction blanks were carried out alongside the sediment extractions, using the same glassware, solvents and extraction methodology.

Extract analysis was performed on an ultra-high performance liquid chromatography-high resolution mass spectrometer (UHPLC-HRMS) in reverse phase as described in Hopmans et al. (2021). An Agilent 1290 Infinity I UHPLC, featuring a thermostatted auto-injector and column oven, was utilised, coupled to a Q Exactive Orbitrap MS with Ion Max source with heated electrospray ionization (HESI) probe (Thermo Fisher Scientific, Waltham, MA). Separation was carried out with an Acquity BEH C18 column (Waters, 2.1 × 150 mm, 1.7 µm) maintained at 30°C, using an eluent composition of (1) MeOH/H₂O/formic acid/14.8 M NH₃aq [85:15:0.12:0.04 (v:v)] and (2) IPA/MeOH/formic acid/14.8 M NH₃aq [50:50:0.12:0.04 (v:v)]. The flow rate was 0.2 mL min⁻¹ with an elution program of: 95% A for 3 minutes, followed by a 40% linear gradient at 12 minutes, then 0% A at 50 minutes which was maintained until 80 minutes. The positive ion HESI parameters were as follows: capillary temperature at 300°C; sheath gas (N₂) pressure at 40 arbitrary units (AU); spray voltage at 4.5 kV; auxiliary gas (N₂) pressure at 10 AU; S-lens at 70 V; probe heater temperature at 50°C. Lipid analysis was conducted across a mass range of m/z 350–2000 (resolving power 70,000 ppm at m/z 200), followed by data-dependent tandem MS/MS (resolving power 17,500 ppm), in which the 10 most abundant masses in the mass spectrum were fragmented successively. Optimal fragmentation was achieved with a stepped normalized collision energy of 22.5 and 40 (isolation width 1.0 m/z) for BHP analysis. The Q Exactive was calibrated to a mass accuracy of ±1 ppm using the Thermo Scientific Pierce LTQ Velos ESI Positive Ion Calibration Solution. During analysis, dynamic exclusion was applied to temporarily exclude masses for 6 s, enabling selection of less abundant ions for MS/MS.

BHPs were tentatively identified based on their retention time, exact mass, and fragmentation spectra as described in Hopmans et al. (2021) and Richter et al. (2023). Note, the BHPs discussed in this study are tentative structures and we rely on comparisons with previous studies for structural elucidation, in particular for determining the stereochemistry of the bacteriohopanetetrol isomers (Schwartz-Narbonne et al., 2020). Integrations were performed on (summed) mass chromatograms of relevant molecular ions ([M+H]⁺, [M+NH₄]⁺, and [M+Na]⁺). The BHP absolute abundances are all presented as peak area per gram of total organic carbon (TOC) since we lack appropriate standards to correct for differences in ionization efficiency between BHPs (Cutmore et al., 2025).

2.2. Age Model

The age model for core 64PE418 was previously published in Cutmore et al. (2025). Seven bulk organic matter ¹⁴C dates were used in its production. Six of these were from core 64PE418, combined with an additional bulk organic carbon ¹⁴C date from core KNR 134-08 BC17 (Jones and Gagnon, 1994), which is sourced from the same location and water depth as 64PE418. This additional ¹⁴C date is from the widely observed Unit I/II boundary and was employed to refine the age model for the upper section of the core. Variable reservoir-ages, calculated by

Deleted: BHPs were tentatively identified based on their retention time, exact mass, and fragmentation spectra.

Deleted: &

153 Kwiecien et al. (2008) for intermediate water depths in the Black Sea over the last ~20 ka, were added to the
154 calibration. The ¹⁴C dates were calibrated using the Marine20 calibration curve (Heaton et al., 2020) for the upper
155 three ¹⁴C dates (reflecting the period after the IMI), while the lower four ¹⁴C dates were calibrated using the IntCal20
156 calibration curve (Reimer et al., 2020), (reflecting the period when then Black Sea was a lacustrine environment).
157 The R-code, CLAM (Blaauw, 2010), was used to create the age–depth model. The BHP record from core 64PE418
158 spans the last 19.5 ka, with an average resolution of ~450 years. The key transitions (identified by colour and
159 elemental changes in the core [Cutmore et al., 2025]) are as follows: the onset of the IMI is dated at 9.6 ka ± 237 yrs;
160 Unit II/III occurs at 7.2 ka ± 202 yrs; and the Unit I/II boundary is dated at 2.6 ka ± 402 yrs.

161
162 *2.3. Statistical analyses*

163 A principal component analysis (PCA) was used to assess the variability in BHP distributions downcore after
164 applying a Hellinger transformation and scaling the BHP dataset. All analyses were performed and visualized in R
165 (version 4.2.2; R Core Team, 2023) with the vegan package (version 2.6-4; Oksanen et al., 2020), factoextra (version
166 1.0.7; Kassambara and Mundt, 2020), FactoMineR (version 2.7; Lê et al., 2008), and ggplot2 (version 3.4.0; Wickham
167 and Chang, 2016).

168
169 **3. Results & Discussion**

170 Significant changes in total organic carbon (TOC) content, colour and elemental signatures of the core are
171 observed (Cutmore et al., 2025), corresponding to changing environmental conditions in the Black Sea basin over
172 the last 20 ka (Arthur & Dean, 1998; Bahr et al., 2005; Jones & Gagnon, 1994; Ankindinova et al., 2019). Based on
173 this, the core is divided into three widely acknowledged key periods: the oxic lacustrine phase (19.5-9.6 ka) where
174 the basin was disconnected from the global ocean; the transition phase (9.6 - 7.2 ka) after the IMI over the Bosphorus
175 sill at ~9.6 ka, whereby the basin moved towards a marine environment; and the marine phase (7.2 ka to the present)
176 where the Black Sea became a euxinic brackish-to-marine environment (Jones & Gagnon, 1994; Arthur & Dean, 1998;
177 Aksu et al., 2002; Major et al., 2006; Bahr et al., 2008; Ankindinova et al., 2019; Cutmore et al., 2025).

178
179 *3.1. BHP distribution changes over time*

180 In total, 63 BHPs were detected throughout the Black Sea record. **BHP structures and other compounds**
181 **discussed in this study are shown in Fig. 2. A PCA of the relative distribution of BHPs reveals four distinct clusters**
182 **(Fig. 3), which correspond with the above defined phases of the Black Sea: an oxic lacustrine phase (19.5 - 9.6 ka), a**
183 **transition phase (9.6 - 7.2 ka), and a marine phase that is further divided into an early marine phase (7.2 - 5 ka) and**
184 **a late marine phase (5 ka to present). PC1 explains 40.8% of the variance and is positively correlated with Nu-BHPs,**
185 **while PC2 explains 18.2% of the variance and is negatively correlated with BHPs such as 35-aminobacteriohopane-**
186 **32,33,34-triol (aminotriol from herein), N-formylated-aminotriol, and ethenolamine-bacteriohopane-32,33,34,35-**
187 **tetrol (ethenolamine-BHT from herein). BHT-34S, aminotriol and ethenolamine-BHT are present throughout the**

Deleted: A PCA of the relative distribution of BHPs reveals four distinct clusters (Fig. 2),
Deleted: which correspond with the above defined phases of the Black Sea: ...

Deleted: 22

record (Fig. 3), which is in line with the ubiquitous nature of these BHPs that are found in contemporary marine, freshwater and terrestrial environments (Talbot and Farrimond 2007; Hopmans et al., 2021; Richter et al., 2023). Note, due to a lack of standards, absolute concentrations cannot be determined and BHP relative abundances must be interpreted with caution. However, despite this limitation, this semi-quantitative approach still provides a valuable overview of major changes in BHP distributions throughout the record.

Each phase of the Black Sea is defined by several notable changes in BHP distributions. Throughout the oxic lacustrine phase, during the Last Deglaciation and early Holocene (19.5 – 9.6 ka), a wider range of BHPs were detected compared to other phases (Fig. 4). This coincides with low TOC levels and low concentrations of elements that accumulate in sediments under anoxic conditions (i.e., uranium [U], vanadium [V], and molybdenum [Mo]) (Cutmore et al., 2025), pointing to a well-oxygenated environment. Nu-BHPs, for instance, are present in higher absolute abundance and demonstrate a more diverse distribution during the lacustrine phase relative to the other periods (Fig. 3). Bacteriohopane-32,33,34-triol (BHtriol), bacteriohopane-31,32,33,34,35-pentol (BHpentol), and bacteriohopane-30,31,32,33,34,35-hexol (BHhexol) are also present in higher absolute abundance during the oxic lacustrine phase, relative to the rest of the record where they are largely absent. The transition phase (9.6 - 7.2 ka) occurs after the IMI at ~9.6 ka and is characterized by a decrease in almost all Nu-BHPs with the exception of an increase in aminotriol, ethenolamine-BHT, and N-formylated-aminotriol.

At 7.2 ka, there is a shift to a permanent euxinic marine system which led to increased preservation of organic matter resulting from permanent water column anoxia. Despite the enhanced preservation, previous studies have reported a decrease in both homohopanoid and BHP concentrations in the Black Sea during this period (Blumenberg et al., 2009a) indicating either a decline in BHP production or a decrease in BHP transport to the sediment, as observed in the modern-day Black Sea water column (Wakeham et al., 2007; Blumenberg et al., 2009a; Kusch et al., 2022). In our record, the early marine phase (7.2 – 5 ka) is marked by an increase in methoxylated-BHT (methoxy-BHT from herein), methoxylated-ethenolamine-BHT (methoxy-ethenolamine-BHT from herein), and propenolamine-BHT. There is a temporary increase in aminotetrol, aminopentol, N-acylated-aminotriols, ethenolamine-BHpentol, and ethenolamine-BHhexol that spans the transition of the early and late marine phases. The absolute abundances of individual BHPs are considerably lower after ~3.9 ka, with the exception of methoxy-BHT and methoxy-ethenolamine-BHT, which increase from 0.9 ka to present (Fig. S2-S7). A decrease in BHP and homohopanoid concentrations during the late marine phase has also been reported in a previous reconstruction of the Black Sea (Blumenberg et al., 2009a). This highlights how distinct BHP distributions characterize the different environmental phases of the Black Sea record.

3.2. Diagenetic products of BHPs

BHP preservation was assessed throughout the core by tracking known early diagenetic products of BHP degradation: 32,35-anhydrobacteriohopanetetrol (anhydro-BHT) and anhydrobacteriohopanepentol (anhydro-BHpentol) which are formed from the dehydration and cyclization of BHT and BHpentol, respectively (Bednarczyk et

Deleted: 3

Deleted: 2

Deleted: enhanced

Deleted: The absolute abundances of BHPs are considerably lower after ~3.9 ka, with the exception of methoxy-BHT and methoxy-ethenolamine-BHT, which increase from 0.9 ka to present. ...

al., 2005; Talbot et al., 2005; Schaeffer et al., 2008; 2010). In addition, it has been shown that anhydro-BHT is also a degradation product of composite BHPs, such as BHT-cyclitol ether and adenosylhopane (Schaeffer et al., 2010; Eickhoff et al., 2014). Note, the diagenetic BHPs discussed here are only the early degradation products of BHPs, and we do not discuss BHPs preserved as e.g. organic sulfur compounds (a novel C₃₅ hopanoid in sediments [Valisolalao et al., 1984]) or the fate of C₂₇-C₃₅ hopanes. However, previous studies have shown that the concentrations of S-bound hopanoids only make up to 7.5% of the sum of free BHPs and S-bound hopanoids, with the lowest abundance of S-bound hopanoids occurring during the lacustrine phase (Blumenberg et al. 2009a). Non-extended hopanoids were found to be more abundant during the marine phase and decreased with depth, further non-extended hopanoids were found to be less abundant than BHPs during the transition and lacustrine phases of the Black Sea record (Blumeberg et al., 2009a).

Throughout the record, the absolute abundance of anhydro-BHT and -BHpentol follow opposite trends to each other, with a higher absolute abundance of anhydro-BHpentol and anhydro-BHT during the lacustrine oxic phase and marine phase, respectively (Fig. S2 and Fig. S8). In a previous study of Black Sea surface sediments, anhydro-BHT increased below the top 2 cm of the core, suggesting rapid early diagenetic modifications of BHPs (Kusch et al., 2022). Previous studies point to the possibility of selective degradation of BHPs (Kusch et al., 2022), in particular the degradation of Nu-BHPs relative to co-occurring composite BHPs, such as aminotriol (Talbot et al., 2016b). However, Nu-BHPs occur in high absolute abundance throughout the record, particularly during the lacustrine oxic phase where we would anticipate increased degradation due to an oxygenated water column relative to the more recent marine phase whereby the water column became anoxic (Schrader, 1979; Arthur and Dean, 1998; Eckert et al., 2013).

BHtriol and unsaturated BHtriol, found previously in lacustrine environments (Rodier et al., 1999; Watson and Farrimond, 2000)., follow a similar trend to anhydro-BHpentol, with all three BHPs present in higher absolute abundance during the lacustrine phase relative to the late transition period and the marine phase, suggesting that BHtriol and unsaturated BHtriol could also be diagenetic products (Fig. S2 and Fig. S8). Indeed, BHtriol only contains 7 carbons on the side-chain instead of the typical 8 carbons found in other BHPs and in unsaturated BHtriol the unsaturation occurs on the side-chain, indicating that they may have formed from decarboxylation and dehydration of functionalized BHPs. It is possible that other early diagenetic products of BHPs (i.e. diols and triols identified in Watson & Farrimond, 2000) were present during the lake phase, however, the mass spectra were not clear enough to identify these compounds.

The decline in BHtriol and anhydro-BHpentol throughout the transition period (between 9.6 - 7.2 ka) may indicate reduced degradation due to intermittent anoxia in the water column. It may, however, also indicate a decrease in the abundance of composite BHPs that are precursors to these diagenetic BHPs, with both BHHexol and BHpentol also declining in absolute abundance during the transition period compared to the lacustrine phase. Anhydro-BHT, in contrast, does not decline during the transition period, instead increasing during the transition and marine periods. This could suggest that there is an increase in the source of the diagenetic precursors for anhydro-

Formatted: Subscript

Formatted: Subscript

Formatted: Subscript

Deleted: Fig. S4

Deleted: decreased

Deleted: diagenesis

Deleted: ; Kusch et al., 2022

Deleted:

Deleted: Fig. S4

276 BHT (i.e., BHT and adenosylhopane; Eickhoff et al., 2014), while the precursors to the other diagenetic BHPs
277 decreased. The changes in the absolute abundance of these BHPs are likely associated with the changes in precursor
278 BHP concentrations. Overall, there are BHP diagenetic products present throughout the record, averaging 11%
279 during the lacustrine period, 6% during the transition, and 11% and 16% during the early and late marine period,
280 respectively. These results taken together with previous analyses of S-bound hopanes and non-extended hopanoids
281 (Blumenberg et al., 2009a), suggests generally good preservation conditions throughout the record.

Deleted: Overall, there are BHP diagenetic products present throughout the record, and the changes in the absolute abundance of these BHPs are likely associated with the changes in precursor BHP concentrations.

282
283 3.3. Sources of Nu-BHPs

284 Nu-BHPs were particularly abundant and diverse during the lacustrine phase (Fig. 5), with the highest Nu-
285 BHP abundance occurring between 19.5 - 13.8 ka, during the last deglaciation, dominated by adenosylhopane_{HG-diMe}
286 followed by adenosylhopane and adenosylhopane_{HG-Me}. Many of the Nu-BHPs are present in higher abundance early
287 in the record and decline during the later part of the lacustrine phase. This pattern corresponds with changes in
288 terrigenous input, as indicated by a decline in the elemental records of titanium to calcium (Ti/Ca) and of potassium
289 (K) (Cutmore et al., 2025). Furthermore, it corresponds with a decrease in the branched and isoprenoid tetraether
290 (BIT) index over the lacustrine period (Bingjie Yang, personal communications, 2024), which indicates a decline in
291 the delivery of terrestrial organic matter to our site (Hopmans et al., 2004), likely primarily supplied by three major
292 ivers: the Danube, the Dnieper, and the Don. Between 18 and 14.8 ka, there were multiple significant meltwater
293 pulses in the Black Sea (Major et al., 2006; Bahr et al., 2008; Badertscher et al., 2011; Yanchilina et al., 2019) followed
294 by higher precipitation during the Bølling Allrød (BA) (14.5-13 ka [Shumilovskikh et al., 2012]). Consequently, the
295 increased terrestrial input is likely the result of this increased freshwater influx. Many of the Nu-BHPs may therefore
296 be sourced from this increased contribution of terrestrial organic matter, with Nu-BHPs primarily associated with
297 soils (Talbot and Farrimond, 2007; Cooke et al., 2008a; Xu et al., 2009; Rethemeyer et al., 2010).

Deleted: 4

Deleted: t

Deleted: as indicated by the elemental records of titanium to calcium (Ti/Ca) and of potassium (K), as well as the BIT index based on branched glycerol dialkyl glycerol tetraether (br-GDGT) and crenarchaeol (Hopmans et al., 2004), in this core

Deleted: pers. comms.

Deleted: .

Deleted: ;

Deleted: .

Deleted: This indicates a high contribution of terrigenous material early in the record and a decline thereafter.

Deleted: enhanced

Deleted: enhanced

Deleted: enhanced

298 The absolute abundance of Nu-BHPs declines after 13 ka, corresponding with low Ti/Ca and K values
299 (Cutmore et al., 2025), which indicate reduced input of terrestrial organic matter. This corresponds with drier
300 regional conditions during the Younger Dryas (YD) and Early Holocene (Shumilovskikh et al., 2012) and reduced influx
301 of meltwater into the Black Sea due to the retreat of ice sheets across Northern Europe (Major et al., 2006). During
302 this time, the composition of the Nu-BHPs also changes with a decline in the absolute abundance of
303 adenosylhopane_{HG-diMe}. After 13 ka, 2Me-adenosylhopane_{HG-diMe} increases, peaking at 10.6 ka and declining abruptly
304 at the IMI at 9.6 ka. While the majority of Nu-BHPs identified in this record are likely linked to soil input and closely
305 follow the terrigenous elements, some, such as 2Me-adenosylhopane_{HG-diMe}, demonstrate a different pattern at 11
306 ka. 2Me-adenosylhopane_{HG-diMe}, for instance, peaks at the start of the Holocene (11-9 ka) followed by an abrupt
307 decline at the IMI when anoxic conditions started to prevail in the lower part of the water column (Fig. S7). 2Me-
308 adenosylhopane_{HG-diMe} was previously detected in higher abundance near the chemocline of a lake (Richter et al.,
309 2023), suggesting potential in situ production rather than a terrestrial source for this Nu-BHP between 11-9 ka.

Deleted: While the majority of Nu-BHPs identified in this record are likely linked to soil input and closely follow the terrigenous elements, some, such as 2Me-adenosylhopane_{HG-diMe}, demonstrate a different pattern throughout the core and are likely being produced within the lacustrine water column (Fig. 5). 2Me-adenosylhopane_{HG-diMe}, for example, peaks at the start of the Holocene (11-9 ka), and may therefore be linked to higher temperatures or other factors associated with warming in the Black Sea. Its abundance declines abruptly at the IMI, indicating that it is not produced under anoxic conditions. ¶

341 During the transition period, most Nu-BHPs are still detected in the record (e.g., adenosylhopane_{HG-Me} and
342 2Me-adenosylhopane_{HG-diMe}), but in much lower absolute abundance than during the oxic lacustrine period, likely
343 due to reduced input of terrestrial organic matter, corresponding with low Ti/Ca and K values (Cutmore et al., 2025).
344 Me-adenosylhopane_{HG-diMe}, however, demonstrates a higher abundance during the transition period compared to
345 the end of the oxic lacustrine period (Fig. S6). As this compound has previously been found near the chemocline in
346 lakes (Richter et al., 2023), it is possible that it was being produced near the oxic-anoxic transition zone during
347 periods of water column anoxia.

348 The diversity of Nu-BHPs is low throughout the marine phase with many Nu-BHPs below detection limit
349 throughout this period (e.g., 2Me-Adenosylhopane, 3Me-Adenosylhopane, diMe-adenosylhopane_{HG-Me}, 3Me-
350 Adenosylhopane_{HG-diMe}, 2Me-N1-methyl-inosylhopane and Me-N1-methyl-inosylhopane). The Nu-BHPs that are
351 detected are present in low absolute abundance compared to other phases. This is likely due to the low contribution
352 of terrestrial organic matter to the site, demonstrated by the low Ti/Ca values (Cutmore et al., 2025), which reduced
353 the delivery of soil-sourced Nu-BHPs to this site. Notably, there is an increase in several Nu-BHPs after 1 ka, including
354 adenosylhopane, 2Me-adenosylhopane_{HG-Me}, Me-adenosylhopane_{HG-diMe}, inosylhopane, N1-methylinosylhopane and
355 Me-N1-methyl-inosylhopane. If these BHPs are not derived from soil inputs, then they are likely being produced in
356 the water column of the Black Sea. Indeed, adenosylhopane, 2Me-adenosylhopane_{HG-Me}, and inosylhopane are
357 detected in the modern-day Black Sea water column (Kusch et al., 2022).

358 In summary, most Nu-BHPs detected in the Black Sea core are likely sourced from soils, linked to freshwater
359 influx into the basin. Several Nu-BHPs, such as adenosylhopane, 2Me-adenosylhopane_{HG-Me}, Me-adenosylhopane_{HG-}
360 diMe, 2Me-adenosylhopane_{HG-diMe}, inosylhopane, N1-methylinosylhopane and Me-N1-methyl-inosylhopane,
361 however, exhibit patterns that suggest these Nu-BHPs were produced in the water column of the Black Sea. As
362 ongoing research continues to explore the sources of Nu-BHPs both in soils and in the water column of lakes and
363 marine environments, it may be possible to expand the interpretation of these records.

364 365 3.4. BHPs associated with the N-cycle

366 BHT-x, produced by anaerobic ammonium oxidising (anammox) bacteria, was previously reported,
367 revealing increased anammox activity after 7.2 ka leading to the loss of bioavailable nitrogen during the marine
368 euxinic period (Cutmore et al., 2025). To further advance our understanding of the nitrogen cycle over the last 20
369 kyr in the Black Sea, we highlight other BHPs as potential biomarkers for investigating this biogeochemical cycle.

370 Oxazinone-aminotriol, for instance, was detected during the lacustrine and transition phases, but absent
371 during the marine euxinic period. As a potential marker for nitrite-oxidizing bacteria (NOB) (Elling et al., 2022; Richter
372 et al., 2023), its presence during the lacustrine and transition phases suggests that nitrite oxidation, performed by
373 NOB as the second step of nitrification (Stein & Nicol, 2018), was an important process when the upper water column
374 was oxygenated. Its absence during the marine phase indicates a decline under euxinic conditions. This pattern
375 mirrors the crenarchaeol record from the same core (Cutmore et al., 2025), a biomarker exclusively produced by the

Deleted: 8

Deleted: surface waters of the

Deleted: future studies discover

Deleted: more about the

abundant and widespread archaea Thaumarchaeota (Nitrososphaerota) (Sinninghe Damsté et al., 2002), which performs the crucial first step in nitrification in the Black Sea (Lam et al., 2007) by aerobically oxidizing ammonia to nitrite (Könneke et al., 2005; Wuchter et al., 2006). This reinforces the interpretation that nitrification was enhanced when the water column was well-oxygenated.

Other BHPs identified in this study may also be associated with the nitrogen-cycle. Notably, BHT-CE abundance closely tracks that of crenarchaeol (Fig. 6; Cutmore et al., 2025), indicating a potential association between the dominant bacterial producer of BHT-CE and the ammonia oxidizing archaea, Nitrososphaerota. This pattern may reflect a functional or ecological coupling between the dominant BHT-CE-producing bacteria and the nitrification process in the Black Sea. Similar associations between archaea and bacteria have been observed elsewhere, such as between *ca. Methanoperedens nitroreducens* and *ca. Methyloirabilis oxyfera* (NC10 phylum) during methane oxidation (Smit et al., 2019), and with anammox bacteria utilising nitrite produced by Nitrososphaerota and ammonia-oxidising bacteria (AOB) in the modern-day Black Sea (Kuypers et al., 2003; Lam et al., 2007). Although BHT-CE has been detected in an enrichment culture from a peat sample for the NC10 bacteria, its exact microbial source remains uncertain (Kool et al., 2014). Consequently, a functional coupling between Nitrososphaerota and an unknown BHT-CE-producing bacteria may have existed in the Black Sea water column. Alternatively, both microbes may have inhabited similar ecological niches and depth habitats, leading to parallel responses to changes in water column conditions.

We identified BHPs with a methylation in the second carbon position of the A ring (2Me-BHPs) in our record. 2Me-BHT was largely absent during the lacustrine phase until the early Holocene (10.3 ka), where it increased before peaking at the start of the transition phase (9.6 ka) and again increasing during the marine phase (6–4.2 ka) (Fig. S2). 2Me-BHPs, precursors for 2Me-hopanes, were initially linked to cyanobacteria (Summons et al., 1999; Kuypers et al., 2004; Talbot et al., 2008), but are now known to also be produced in low amounts by alpha (α)-proteobacteria (Rashby et al., 2007; Welander et al., 2010; Naafs et al., 2022), such as NOB *Nitrobacter vulgaris* (Elling et al., 2020), as well as Actinobacteria and one strain of Acidobacteria (Welander et al., 2010; Sinninghe Damsté et al., 2017; Naafs et al., 2022). In our record, the increase in 2Me-BHT coincides with increasing absolute abundance of hexose HGs, known indicators of increased N₂-fixation by freshwater or brackish heterocystous cyanobacteria (Cutmore et al., 2025). It is therefore possible that 2Me-BHT is related to N₂-fixing cyanobacteria in the Black Sea record, with 2Me-BHT previously identified in various heterocystous cyanobacteria, including *Nostoc* sp., *Calothrix* sp. and *Chlorogloeopsis fritschii* (Talbot et al., 2008). While α -proteobacteria may also contribute to the presence of 2Me-BHT in this record, the lack of alignment with oxazinone-aminotriol (a potential marker for NOB [Elling et al., 2022]) supports the theory that the primary origin of 2Me-BHT in this record is from N₂-fixing cyanobacteria. While further research is needed to refine their use, the BHPs in this record show potential as both novel indicators for microbial nitrogen cycling and as supporting evidence for established biomarkers like those of heterocystous cyanobacteria.

Deleted: BHT-x, produced by anammox bacteria, has already successfully been applied to this record in a previous study (Cutmore et al., 2025) to detect anammox activity, which is occurring in the present day Black Sea (Kuypers et al., 2003; Dalsgaard et al., 2012). At 7.2 ka, there is an increase in concentration suggesting expansion of anammox activity, due to the permanently anoxic water column which enabled anammox bacteria to expand their habitat from the anoxic sediments; consequently, during the marine euxinic period, loss of bioavailable nitrogen by anammox activity was enhanced (Cutmore et al., 2025). In contrast, oxazinone-aminotriol was detected throughout the lacustrine and transition phases, and absent during the marine euxinic period. Oxazinone-aminotriol was first identified in cultures of nitrite-oxidizing bacteria and later found in the surface sediments of lakes (Elling et al., 2022; Richter et al., 2023). Its presence in the Black Sea record may therefore indicate that nitrite-oxidation was an important process in the water column when the upper water column was oxygenated during the lacustrine and transition phase, and less so when the water column was euxinic, during the later marine phase.

Other BHPs might also be associated with the nitrogen-cycle. For instance, there appears to be a close coupling between the absolute abundance of BHT-CE and crenarchaeol throughout this record (Fig. 5; Cutmore et al., 2025). As crenarchaeol is exclusively produced by the abundant and widespread archaea Thaumarchaeota (Nitrososphaerota) (Sinninghe Damsté et al., 2002), the close coupling of these records indicates an association between the dominant bacterial producer of BHT-CE in the Black Sea over this period and Nitrososphaerota. As Nitrososphaerota plays a crucial role in nitrification in the Black Sea (Lam et al., 2007) by aerobically oxidizing ammonia to nitrite (Könneke et al., 2005; Wuchter et al., 2006), it may be that the dominant bacterial producer of BHT-CE was coupled to this nitrification process. Previous studies have indeed shown associations between archaea and bacteria, with a coupling between *ca. Methanoperedens nitroreducens* and *ca. Methyloirabilis oxyfera* (NC10 phylum) in order to perform methane oxidation (Smit et al., 2019), and with anammox bacteria utilising the nitrite produced by both Nitrososphaerota and ammonia-oxidising bacteria (AOB) in the modern-day Black Sea (Kuypers et al., 2003; Lam et al., 2007). For instance, BHT-CE was detected in an enrichment culture from a peat sample for the NC10 bacteria; however, it is unclear whether BHT-CE was being produced by NC10 bacteria or other microbial communities in the enrichment (Kool et al., 2014). Consequently, a coupling may have existed in the Black Sea water column between Nitrososphaerota and unknown bacteria with regards to nitrification. Alternatively, both microbes may have inhabited the same ecological niche and depth habitat, meaning their abundance was similarly affected by changes in the water column conditions.

—BHPs with a methylation in the second carbon position of the A ring (2Me-BHPs) are precursors for 2Me-hopanes, which were initially proposed as biomarkers for N₂-fixing ... [1]

3.5. BHPs associated with methane-oxidizing bacteria

Aminotetrol and aminopentol are present throughout the lacustrine phase of the Black Sea, albeit in low absolute abundance compared to the other phases (Fig. S3). Aminotetrol and aminopentol are associated with MOB (Rohmer et al., 1984; Neunlist and Rohmer, 1985a,b; Jahnke et al., 1999; Cvejic et al., 2000; Talbot et al., 2001; Zhu et al., 2011; van Winden et al., 2012; Rush et al., 2016), and are produced in small proportions by sulfate-reducing bacteria (SRB) (Blumenberg et al., 2006; 2009b; 2012). Given the low sulfate availability and aerobic conditions during the lacustrine phase, (Cutmore et al., 2025), aminotetrol and aminopentol were more likely being produced by methane-oxidizing bacteria than SRB, which are considered obligate anaerobes (Hao et al., 1996). Although NOB have also been proposed as a source of aminopentol and aminotetrol (Elling et al., 2022), the lack of co-variation with other biomarkers associated with the nitrogen-cycle (i.e. crenarchaeol), suggest that NOB are not the main contributors of these BHPs. Other independent biomarkers, such as high concentrations of 17 β (H)-moret-22(29)-ene with depleted $\delta^{13}\text{C}$ values, support a possible aerobic methanotrophic source (Uemura and Ishiwatari, 1995; Blumenberg et al., 2009a). The continuous presence of dioxanone-methylaminotriol, which was previously identified in the surface sediments and bottom waters of lakes that experience seasonal hypoxia or anoxia (Richter et al., 2023), throughout this period suggests that the lake phase of Black Sea may have experienced seasonal stratification and the formation of an oxycline. In modern lakes, methane-oxidizing bacteria are known to thrive at the oxycline of the water column (Hanson & Hanson, 1996), but are also active at the sediment-water interface in well-mixed lakes (e.g., Kuivila et al., 1988; Frenzel et al., 1990), in anoxic parts of the water column (Schubert et al., 2010; Brees et al., 2014; Oswald et al., 2016), and in anoxic sediments (Martinez-Cruz et al., 2017; He et al., 2021; Su et al., 2022). Although it is unclear how much of these different pools are contributing to the sedimentary aminotetrol and aminopentol pool identified in the Black Sea record, we can still conclude that aminotetrol and aminopentol are likely sourced from methane-oxidizing bacteria during the lacustrine phase.

Several BHPs increased during the transition period relative to the preceding lacustrine phase (i.e., methoxy-BHT, aminotriol, aminotetrol, EC-aminotriol, ethenolamine-BHT, unsaturated ethenolamine-BHT, ethenolamine-BHpentol, methoxy-ethenolamine, propenolamine-BHT and unsaturated propenolamine-BHT), with several BHPs gradually increasing in absolute abundance over the transition phase (i.e., aminotriol and N-acyl-aminotriols C_{12:0}, C_{14:0}, C_{15:0}, C_{16:0} and C_{18:0}) (Fig. S5). The continuous presence of aminotetrol during this period is likely associated with methane-oxidizing bacteria as the pelagic redoxcline was established. In the Black Sea, evidence for aerobic methanotrophs during the transition phase is supported by the presence of ethenolamine-BHpentol, which has so far only been detected near a terrestrial methane seep and in lakes with low oxygen conditions in the bottom waters (Hopmans et al., 2021; Richter et al., 2023). Similarly, in the Baltic Sea, the transition from the Ancyclus Lake phase to the Littorina Sea was marked with a peak in aminotriol and aminotetrol, attributed to methane-oxidizing bacteria (Blumenberg et al., 2013). The presence of MOB-associated BHPs would suggest that aerobic methane-oxidation occurred during this phase.

Deleted: 5

Deleted: Aminotetrol and aminopentol are associated with MOB (Rohmer et al., 1984; Jahnke et al., 1999; Cvejic et al., 2000; Talbot et al., 2001; Zhu et al., 2011; van Winden et al., 2012; Rush et al., 2016), but are also produced in small proportions by sulfate-reducing bacteria (Blumenberg et al., 2006; 2009b; 2012). In addition, aminopentol was detected in cultured nitrite-oxidizing bacteria, *Nitrospira defluvii* and *Nitrobacter vulgaris* (Elling et al., 2022). Indicators of a well-oxygenated water column during the lacustrine phase (Cutmore et al., 2025), would suggest that methane-oxidizing bacteria are a more likely source of aminotetrol and aminopentol within the lacustrine water column than sulfate-reducing bacteria, which are typically considered to be obligate anaerobes (Hao et al., 1996). Although we cannot rule out possible sulfate-reduction in the sediment, other independent biomarkers suggest that aerobic methanotrophs were active in the basin and are a more likely source for these biomarkers. For instance, high concentrations of 17 β (H)-moret-22(29)-ene with depleted $\delta^{13}\text{C}$ values were previously detected in sediments from the lacustrine phase from the Black Sea, and are thought to be sourced from aerobic methanotrophs in either the water column or the surface sediments (Uemura and Ishiwatari, 1995; Blumenberg et al., 2009a). Nitrite-oxidizing bacteria are also a potential source for aminopentol (Elling et al., 2022), but as both aminopentol and aminotetrol vary independently of BHPs and other biomarkers associated with the nitrogen-cycle (such as crenarchaeol), it is likely that nitrite-oxidizing bacteria are not a major source of these BHPs. Thus, aminotetrol and aminopentol were likely produced by methane-oxidizing bacteria near the oxycline of the basin, or, if the basin was fully oxygenated, in the surface sediments (Neunlist and Rohmer, 1985a,b; Cvejic et al., 2000; Talbot et al., 2001). The occurrence of temporary anoxia in the bottom waters or surface sediments during the lacustrine phase is supported by

Deleted: t

Deleted: throughout this period

Deleted: with the highest absolute abundance occurring between 19.5 and 16.4 ka. Dioxanone-methylaminotriol

Deleted: basins

Deleted: .

Formatted: Font colour: Red

Deleted: 4.2

Deleted: and high absolute abundance of aminotriol

Deleted: Similarly, in the Baltic Sea, the transition from the Ancyclus Lake phase to the Littorina Sea was marked with a peak in aminotriol and aminotetrol, attributed to methane-oxidizing bacteria (Blumenberg et al., 2013). This interpretation is also

During the early marine phase, ethenolamine-BHpentol, ethenolamine-BHhexol, propenolamine-BHT, aminotetrol and aminopentol are all present in higher relative abundance. While propenolamine-BHT has been observed near a terrestrial methane seep, as well as in lakes and in the sediments of coastal lagoons, its microbial source remains uncertain (Hopmans et al., 2021; Richter et al., 2023). Methanotrophic bacteria, known to thrive in the oxic-anoxic transition zone in the modern Black Sea water column (Kuypers et al., 2003; Durisch-Kaiser et al., 2005; Schubert et al. 2006), are known sources of aminotetrol and aminopentol (Blumenberg et al., 2007; Wakeham et al., 2007), however, the distributions and concentrations of BHPs detected in the modern-day Black Sea water column are not reflected in the geological record as there is no effective transport mode from the oxic-anoxic transition zone to the sediment (Wakeham et al., 2007; Blumenberg et al., 2009a; Kusch et al., 2022). However, the high absolute abundance of isorenieratene throughout the early marine phase indicates that euxinic conditions reached the photic zone during this period (Cutmore et al., 2025), pointing to a shallower oxic-anoxic interface compared to the later marine phase. The large accumulation of particles just below the photic zone at the oxycline likely led to a heightened transport of particulate matter, including BHPs, from the oxic-anoxic interface to the sediments (e.g., Repeta, 1993; Sinninghe Damsté et al., 1993; Coolen et al., 2008; Blumenberg et al., 2009a). Between 5 and 3.9 ka, SRB likely occurred in the deeper waters and sediments of the Black Sea and are another potential source for aminotetrol and aminopentol.. This signal is unlikely to have been reflected in the sedimentary record as there is no effective transport of lipids from the deeper waters to the sediments (Schouten et al., 2001; Wakeham et al., 2003), and BHP-producing species of SRB were not detected in genetic analyses of a Black Sea sediment core (Blumenberg et al., 2006; 2009b; 2012; Leloup et al., 2006). Thus, unless there are previously unknown BHP-producing SRBs in the Black Sea sediments, SRBs are likely only a minor source of BHPs during the marine phase (Blumenberg et al., 2009a). N-acyl-aminotriols (i.e., C_{12:0}, C_{14:0}, C_{15:0}, C_{16:0}, C_{17:0}, and C_{18:0}) are also present in high absolute abundance during the early marine phase compared to the rest of the core and peak at 6.1 ka. There is no known distinct source for N-acyl-aminotriols; however, they have been detected in a methanotroph culture as well as various environmental settings (e.g., Hopmans et al., 2021; Kusch et al., 2022; Richter et al., 2023). The high abundance of these long-chain compounds might be associated with heightened transport from the oxycline to the sediments and increased preservation during this time (Blumenberg et al., 2009a; Cutmore et al., 2025). Thus, methane-oxidizing bacteria were likely active in the early marine phase near the oxycline of the Black Sea, as reflected in the sediment record.

High-rates of methane-oxidation in the modern-day Black Sea are mediated by anaerobic methanotrophs in the euxinic water column, with lower rates of anaerobic methane oxidation occurring in the sediments (Reeburgh et al., 1991; Lin et al., 2006; Schubert et al., 2006). MOB are present at the oxic-anoxic transition zone of the modern-day Black Sea water column; however, their associated rates of aerobic methane-oxidation are several orders of magnitude lower than the anaerobic oxidation of methane (AOM) occurring in the anoxic water column (Reeburgh et al., 1991; Schubert et al., 2006). Consequently, the limited transport of suspended particulate matter from the chemocline and euxinic bottom waters to the sediments, leads to an underrepresentation of these lipids in the

Deleted: Similarly, during the transition phase in the Black Sea there is an increase in archaeal lipids derived from anaerobic methanotrophs that might indicate heightened activity by anaerobic methane-oxidizing archaeal communities; alternatively, these lipids could be derived from overprinting in the sediments (Zhu et al., 2024). The presence of MOB-associated BHPs would suggest that aerobic methane-oxidation occurred during this phase, but it is unclear whether this coincided with an increase in anaerobic methane-oxidation. ¶

Deleted:

Deleted: there is a distinct increase in aminotetrol, aminopentol,

Deleted: and

Deleted: If the oxic-anoxic transition zone occurs near the photic zone, however, and there is a large accumulation of particles, it is possible for particulate matter to be transported from this zone to the sediments (e.g., Repeta, 1993; Sinninghe Damsté et al., 1993; Coolen et al., 2008). The

Deleted: indicating

Deleted: This likely led to a heightened transport of BHPs from the oxic-anoxic interface to the sediments (Blumenberg et al., 2009a).

Deleted: Between 5 and 3.9 ka, aminotetrol and aminopentol may have also been produced by sulfate-reducing bacteria (SRB) in the water column or sediments. There is, however, also no effective transport of lipids from the deeper waters to the sediments (Schouten et al., 2001; Wakeham et al., 2003). Further, genetic analyses of a Black Sea sediment core shows that although sulfate-reducing bacteria are abundant in the sulfate-methane transition zone, the *Desulfovibrio* genus (which is known to produce BHPs [Blumenberg et al., 2006; 2009b; 2012]) plays a minor role in the Black Sea sediments compared to *Desulfobacter* (Leloup et al., 2006). *Desulfobacter*, in contrast, does not contain the squalene-hopene cyclase gene (accession no. WP_038942977.1) required for BHP synthesis as confirmed by a protein blast search (NCBI, National Center for Biotechnology Information). Aside from aminotetrol and aminopentol, the only other known BHPs produced by SRB are BHT-22S, BHT-CE, and aminotriol (Blumenberg et al. 2006; 2009b; 2012). Aminotriol is present in low abundance throughout the marine phase. Both BHT-22S and BHT-CE increase during the late marine phase, however, these BHPs could also be derived from other bacterial sources as th... [2]

Formatted: Indent: First line: 1.27 cm

Deleted: occur

Deleted: ,

Deleted: rates

Deleted: compared to the anoxic water column (Reeburgh et al., 1991; Schubert et al., 2006).

geological record and restricts any lipid-based interpretations of methane-oxidizing microbial communities or associated **SRB** in the Black Sea water column during the late marine phase (Schouten et al. 2001; Wakeham et al., 2003; Schubert et al., 2006). Studies of lipid biomarkers associated with anaerobic methanotrophs (e.g., compound-specific carbon isotope compositions of archaeal lipids) confirm that there was minimal overprinting from active methane-oxidizing archaea in the sediments and fossil lipids derived from anaerobic methanotrophs were only a minor portion of the archaeal lipid assemblage during the marine phase (Zhu et al., 2024). In summary, MOB-associated BHPs in our record indicate that aerobic methane-oxidation occurred throughout the Black Sea record. However, archaeal lipids and modern studies demonstrate that anaerobic methane **oxidation** became a more significant “methane sink” in the Black Sea water column during the marine phase relative to the lacustrine phase. Additional studies are needed to confirm this hypothesis by quantifying the contributions of aerobic versus anaerobic methanotrophs and identifying **SRB** during the lacustrine and transition phase.

Deleted: sulfate-reducing bacteria

Deleted: -

Deleted: sulfate-reducing bacteria

3.6. BHPs as indicators for surface salinity

Methoxy-BHT has previously been detected in marine environments and at present, has not been detected in lake samples (Richter et al., 2023). In the Black Sea record, we detect methoxy-BHT intermittently during the oxic lacustrine phase, possibly due to the slightly brackish conditions that existed during this period (Huang et al., 2021). After the IMI, the absolute abundance of this BHP increases, possibly due to the influx of saline water. The highest abundance of methoxy-BHT occurs between 6.1 and 4.2 ka, when surface water salinities reach a peak (Filipova-Marinova et al., 2013; Huang et al., 2021). The reduction in methoxy-BHT between 3.8 and 1.5 ka coincides with the non-detection of isorenieratene (Cutmore et al., 2025) which likely resulted from the erosion of the chemocline (Sinninghe Damsté et al., 1993) due to higher freshwater input, with a short reoccurrence of freshwater/brackish species occurring at this time (Filipova-Marinova et al., 2013; Huang et al., 2021). Consequently, the lower salinity of the surface waters may have also led to the decline in methoxy-BHT. Methoxy-BHT and methoxy-ethanolamine-BHT show similar patterns throughout this period (Fig. 6), both peaking between 6.1 and 4.2 ka, declining between 3.8 and 1.5 ka and increasing towards the end of the record. To our knowledge, methoxy-ethanolamine-BHT has not previously been linked to salinity, but its close correspondence with methoxy-BHT during this period indicates that both BHPs may be a specific adaptation of BHP compositions to saline environments or are linked to specific microbes adapted to brackish to marine systems.

Deleted: The reduction in methoxy-BHT between 3.8 and 1.5 ka coincides with the absence of isorenieratene which likely resulted from the erosion of the chemocline (Sinninghe Damsté et al., 1993) due to enhanced freshwater input, with a short reoccurrence of freshwater/brackish species occurring at this time (Filipova-Marinova et al., 2013; Huang et al., 2021).

Deleted: 5

Deleted: .

Deleted: This indicates that both BHPs may be a specific adaptation of BHP compositions to saline environments or are linked to specific microbes adapted to brackish to marine systems.¶

Deleted:

Deleted: enhanced

4. Conclusions

The Black Sea record shows substantial shifts in BHP absolute abundance and composition over the last 20 ka, which correspond to major environmental transitions. During the lacustrine phase (19.5 – 9.6 ka), high Nu-BHP absolute abundance and diversity likely reflects **increased** terrigenous input during the Last Deglaciation and/or in situ production in the water column. The transition phase (9.6 – 7.2 ka) is marked by a decrease in many BHPs, such as Nu-BHPs, and an increase in methoxylated-BHPs and more ubiquitous BHPs (e.g., aminotriol), likely linked to

807 changing oxygenation and salinity of the water column. The early euxinic marine phase (7.2 ka – 5 ka) is characterized
808 by higher aminotetrol and aminopentol relative to the late marine phase, likely transported with sinking particulate
809 matter to the sediment from a shallow oxycline. The late marine phase (5 ka – present) is marked by a decline in
810 these BHPs, likely due to a deepening of the oxycline. BHP distribution changes throughout the record are attributed
811 either to microbial adaptations to shifts in the oxygen levels or hydrology of the basin, or are associated with specific
812 groups of bacteria suited to these conditions. For instance, BHT-CE was closely associated with crenarchaeol
813 throughout the record, indicating that these compounds are either produced in a similar ecological niche or by
814 related microbes. Aminotetrol, aminopentol, ethenolamine-BHPentol and -BHhexol were detected in the Black Sea
815 record as likely biomarkers for methane-oxidizing bacteria, while methoxylated-BHPs, detected only during the
816 transition and marine phases, may be associated with higher salinities or marine environments. The changes in the
817 Nu-BHP distribution likely reflects changing terrestrial input over this period, with the exception of some Nu-BHPs
818 (e.g., 2Me-adenosylhopane_{HG-diMe} and Me-adenosylhopane_{HG-diMe}) that are likely produced in the water column. BHP
819 degradation products were detected as a minor component of the overall BHP distribution, suggesting good BHP
820 preservation throughout the record. This new BHP record provides valuable insights into ecological changes as well
821 as changes in the nitrogen and methane cycle in the Black Sea over the last 20 ka.

Deleted: a high aminotetrol and aminopentol absolute abundance,

824 **Data Availability**

825 All data generated for this study are archived and publicly available via the Mendeley Data repository online at
826 <http://doi.org/10.17632/m6v5mj5gtp.2> (Cutmore and Richter et al., 2025).

Deleted: *** (***).

828 **Author Contribution**

829 AC- Conceptualization, Formal analysis, Investigation, Data Curation, Visualization, Writing - Original Draft, Writing -
830 Review & Editing. NR- Conceptualization, Formal analysis, Investigation, Data Curation, Visualization, Writing -
831 Original Draft, Writing - Review & Editing. DR- Supervision, Writing - Review & Editing. NB- Supervision, Writing -
832 Review & Editing. SS- Supervision, Funding acquisition, Writing - Review & Editing.

834 **Competing Interests**

835 The authors declare that they have no conflict of interest.

837 **Acknowledgements**

838 We thank the Chief Scientist Prof. Laura Villanueva and Dr. Rick Hennekam, as well as the captain and crew of the
839 R/V Pelagia for the collection of core 64PE418. For laboratory support we thank Anhelique Mets, Denise Dorhout
840 and Monique Verweij. Thank you to Diana Sahonero Canavesi for helpful discussions and Bingjie Yang for providing

helpful insight into the BIT index of this core. This study was funded by the Netherlands Earth System Science Centre (024.002.001) from the Dutch Ministry of Education, Culture and Science (OCW) and the Swiss National Science Foundation (SNSF) Ambizione Fellowship (PZ00P2_216050).

References:

- Aksu, A., Hiscott, R.N., Kaminski, M.A., Mudie, P.J., Gillespie, H., Abrajano, T and Yasar, D.: Last glacial–Holocene paleoceanography of the Black Sea and Marmara Sea: stable isotopic, foraminiferal and coccolith evidence, *Mar. Geol.*, **190**, 119–149, doi.org/10.1016/S0025-3227(02)00345-6, 2002.
-
- Ankindinova, O., Hiscott, R.N., Aksu, A.E and Grimes, V.: High-resolution Sr-isotopic evolution of Black Sea water during the Holocene: Implications for reconnection with the global ocean, *Mar. Geol.*, **407**, 213–228, doi.org/10.1016/j.margeo.2018.11.004, 2019.
-
- Arthur, M.A and Dean, W.E.: Organic-matter production and preservation and evolution of anoxia in the Holocene Black Sea, *Paleoceanogr. Paleoclimatol.*, **13**, 395–411, doi.org/10.1029/98PA01161, 1998.
-
- Badertscher, S., Fleitmann, D., Cheng, H., Edwards, R.L., Göktürk, O.M., Zumbühl, A., Leuenberger, M and Tüysüz, O.: Pleistocene water intrusions from the Mediterranean and Caspian seas into the Black Sea, *Nat. Geosci.*, **4**, 236–239, doi.org/10.1038/ngeo1106, 2011.
-
- Bahr, A., Lamy, F., Arz, H., Kuhlmann, H and Wefer, G.: Late glacial to Holocene climate and sedimentation history in the NW Black Sea, *Mar. Geol.*, **214**, 309–322, doi.org/10.1016/j.margeo.2004.11.013, 2005.
-
- Bahr, A., Arz, H., Lamy, F and Wefer, G.: Late glacial to Holocene paleoenvironmental evolution of the Black Sea, reconstructed with stable oxygen isotope records obtained on ostracod shells, *Earth Planet. Sc. Lett.*, **241**, 863–875, doi.org/10.1016/j.epsl.2005.10.036, 2006.
-
- Bahr, A., Lamy, F., Arz, H., Major, C., Kwiecien, O and Wefer, G.: Abrupt changes of temperature and water chemistry in the late Pleistocene and early Holocene Black Sea, *Geochem., Geophys., Geosy.*, **9**, 1–16, doi.org/10.1029/2007GC001683, 2008.
-
- Bale, N., Ding, S., Hopmans, E.C., Arts, M.G.I., Villanueva, L., Boschman, C., Haas, A.F., Schouten, S and Sinninghe Damsté, J.S.: Lipidomics of Environmental Microbial Communities. I: Visualization of Component Distributions Using

Untargeted Analysis of High-Resolution Mass Spectrometry Data, *Front. Microbiol.*, **12**, 1-15, doi.org/10.3389/fmicb.2021.659302, 2021.

Bednarczyk, A., Hernandez, T. C., Schaeffer, P., Adam, P., Talbot, H. M., Farrimond, P., Riboulleau, A., Largeau, C., Derenne, S., Rohmer, M., and Albrecht, P.: 32,35-Anhydrobacteriohopanetetrol: an unusual bacteriohopanepolyol widespread in recent and past environments, *Org. Geochem.*, **36**, 673–677, doi.org/10.1016/j.orggeochem.2004.10.014, 2005.

Bisseret, P., Zundel, M., and Rohmer, M.: Prokaryotic triterpenoids, *Eur. J. Biochem.*, **150**, 29–34, doi.org/10.1111/j.1432-1033.1985.tb08982.x, 1985.

Blaaw, M.: Methods and code for ‘classical’ age-modelling of radiocarbon sequences, *Quat. Geochron.*, **5**, 512-518, doi.org/10.1016/j.quageo.2010.01.002, 2010.

Blees, J., Niemann, H., Wenk, C. B., Zopfi, J., Schubert, C. J., Kirf, M. K., Veronesi, M. L., Hitz, C., and Lehmann, M. F.: Micro-aerobic bacterial methane oxidation in the chemocline and anoxic water column of deep south-Alpine Lake Lugano (Switzerland), *Limnology and Oceanography*, **59**, 311–324, https://doi.org/10.4319/lo.2014.59.2.0311, 2014.

Blumenberg, M., Krüger, M., Nauhaus, K., Talbot, H. M., Oppermann, B. I., Seifert, R., Pape, T., and Michaelis, W.: Biosynthesis of hopanoids by sulfate-reducing bacteria (genus *Desulfovibrio*), *Environ. Microbiol.*, **8**, 1220–1227, doi.org/10.1111/j.1462-2920.2006.01014.x, 2006.

Blumenberg, M., Seifert, R., and Michaelis, W.: Aerobic methanotrophy in the oxic–anoxic transition zone of the Black Sea water column, *Org. Geochem.*, **38**, 84–91, doi.org/10.1016/j.orggeochem.2006.08.011, 2007.

Blumenberg, M., Hoppert, M., Krüger, M., Dreier, A., and Thiel, V.: Novel findings on hopanoid occurrences among sulfate reducing bacteria: Is there a direct link to nitrogen fixation?, *Org. Geochem.*, **49**, 1–5, doi.org/10.1016/j.orggeochem.2012.05.003, 2012.

Blumenberg, M., Seifert, R., Kasten, S., Bahlmann, E and Michaelis, W.: Euphotic zone bacterioplankton sources major sedimentary bacteriohopanepolyols in the Holocene Black Sea, *Geochim. Cosmochim. Ac.*, **73**, 750-766, doi.org/10.1016/j.gca.2008.11.005, 2009a.

911 [Blumenberg, M., Oppermann, B. I., Guyoneaud, R., and Michaelis, W.: Hopanoid production by *Desulfovibrio bastinii*](#)
912 [isolated from oilfield formation water, *FEMS Microbiol. Lett.*, **293**, 73–78, doi.org/10.1111/j.1574-](#)
913 [6968.2009.01520.x, 2009b.](#)

914 -

915 [Blumenberg, M., Berndmeyer, C., Moros, M., Muschalla, M., Schmale, O and Thiel, V.: Bacteriohopanepolyols record](#)
916 [stratification, nitrogen fixation and other biogeochemical perturbations in Holocene sediments of the central Baltic](#)
917 [Sea, *Biogeosciences*, **10**, 2725–2735, doi:10.5194/bg-10-2725-2013, 2013.](#)

918 -

919 [Bravo, J.-M., Perzl, M., Härtner, T., Kannenberg, E. L and Rohmer, M.: Novel methylated triterpenoids of the](#)
920 [gammacerane series from the nitrogen-fixing bacterium *Bradyrhizobium japonicum* USDA 110, *Eur. J. Biochem.*, **268**,](#)
921 [1323–1331, doi.org/10.1046/j.1432-1327.2001.01998.x, 2001.](#)

922 -

923 [Coolen, M.J.L., Abbas, B., Van Bleijswijk, J., Hopmans, E.C., Kuypers, M.M.M., Wakeham, S.G and Sinninghe Damsté,](#)
924 [J.S.: Putative ammonia-oxidizing Crenarchaeota in suboxic waters of the Black Sea: a basin-wide ecological study](#)
925 [using 16S ribosomal and functional genes and membrane lipids, *Environ. Microbiol.*, **9**, 1001-1016,](#)
926 [doi.org/10.1111/j.1462-2920.2006.01227.x, 2007.](#)

927 -

928 [Coolen M.J.L., Talbot, H.M., Abbas, B., Ward, C., Schouten, S., Volkman, J.K and Sinninghe Damsté, J.S.: Sources for](#)
929 [sedimentary bacteriohopanepolyols as revealed by 16S rDNA stratigraphy, *Environ. Microbiol.*, **10**, 1783-1803,](#)
930 [doi.org/10.1111/j.1462-2920.2008.01601.x, 2008.](#)

931 -

932 [Cooke, M. P., Talbot, H. M and Farrimond, P.: Bacterial populations recorded in bacteriohopanepolyol distributions](#)
933 [in soils from Northern England, *Org. Geochem.*, **39**, 1347–1358, doi.org/10.1016/j.orggeochem.2008.05.003, 2008a.](#)

934 -

935 [Cooke, M.P., Talbot, H.M and Wagner, T.: Tracking soil organic carbon transport to continental margin sediments](#)
936 [using soil-specific hopanoid biomarkers: a case study from the Congo fan \(ODP site 1075\), *Org. Geochem.*, **39**, 965-](#)
937 [971, https://doi.org/10.1016/j.orggeochem.2008.03.009, 2008b.](#)

938 -

939 [Cutmore, A., Bale, N., Hennekman, R., Yang, B., Rush, D., Schouten, Reichart, G.-J., Hopmans, E and S, Schouten.: Impact of deoxygenation and hydrological changes on the Black Sea nitrogen cycle during the Last Deglaciation and](#)
940 [Holocene, *Clim. Past*, doi.org/10.5194/cp-2024-59, 2025.](#)

941 -

942 -

943 [Cvejic, J.H., Bodrossy, L., Kovács, K.L and Rohmer, M.: Bacterial triterpenoids of the hopane series from the](#)
944 [methanotrophic bacteria *Methylocaldum* spp.: phylogenetic implications and first evidence for an unsaturated](#)

aminobacteriohopanepolyol, *FEMS Microbiol. Lett.*, **182**, 361–365, doi.org/10.1111/j.1574-6968.2000.tb08922.x, 2000.

Dalsgaard, T., Thamdrup, B., Fariás, L and Revsbech, N.P.: Anammox and denitrification in the oxygen minimum zone of the eastern South Pacific, *Limnol. Oceanogr.*, **57**, 1331-1346, doi.org/10.4319/lo.2012.57.5.1331, 2012.

Durisch-Kaiser, E., Klauser, L., Wehrli, B., and Schubert, C.: Evidence of Intense Archaeal and Bacterial Methanotrophic Activity in the Black Sea Water Column, *Appl. Environ. Microbiol.*, **71**, 8099–8106, doi.org/10.1128/AEM.71.12.8099-8106.2005, 2005.

Deutzmann, J.S., Stief, P., Brandes, J and Schink, B.: Anaerobic methane oxidation coupled to denitrification is the dominant methane sink in a deep lake, *Proc. Nat. Acad. Sci. USA.*, **111**, 18273-18278, doi.org/10.1073/pnas.1411617111, 2014.

Eckert, S., Brumsack, H.-J., Severmann, S., Schnetger, B., März, C and Fröllje, H.: Establishment of euxinic conditions in the Holocene Black, *Geology*, **41**, 431–434, doi.org/10.1130/G33826.1, 2013.

Eickhoff, M., Birgel, D., Talbot, H. M., Peckmann, J., and Kappler, A.: Diagenetic degradation products of bacteriohopanepolyols produced by *Rhodopseudomonas palustris* strain TIE-1, *Org. Geochem.*, **68**, 31–38, doi.org/10.1016/j.orggeochem.2014.01.002, 2014.

Elling, F. J., Hemingway, J. D., Evans, T. W., Kharbush, J. J., Spieck, E., Summons, R. E and Pearson, A.: Vitamin B12 - dependent biosynthesis ties amplified 2-methylhopanoid production during oceanic anoxic events to nitrification, *Proc. Nat. Acad. Sci. USA.*, **117**, 32996–33004, doi.org/10.1073/pnas.2012357117, 2020.

Elling, F. J., Hemingway, J. D., Kharbush, J. J., Becker, K.W., Polik, C.A and Pearson, A.: Linking diatom-diazotroph symbioses to nitrogen cycle perturbations and deep-water anoxia: Insights from Mediterranean sapropel events, *Earth Planet. Sc. Lett.*, **571**, 1-11, doi.org/10.1016/j.epsl.2021.117110, 2021.

Elling, F. J., Evans, T. W., Nathan, V., Hemingway, J. D., Kharbush, J. J., Bayer, B., Spieck, E., Husain, F., Summons, R. E and Pearson, A.: Marine and terrestrial nitrifying bacteria are sources of diverse bacteriohopanepolyols, *Geobiology*, **20**, 399-420, doi.org/10.1111/gbi.12484, 2022.

Ettwig, K. F., Butler, M. K., Le Paslier, D., Pelletier, E., Mangenot, S., Kuypers, M. M. M., Schreiber, F., Dutilh, B. E., Zedelius, J., De Beer, D., Gloerich, J., Wessels, H. J. C. T., Van Alen, T., Luesken, F., Wu, M. L., Van De Pas-Schoonen,

980 [K. T., Op Den Camp, H. J. M., Janssen-Megens, E. M., Francoijs, K.-J., Stunnenberg, H., Weissenbach, J., Jetten, M. S.](#)
981 [M., and Strous, M.: Nitrite-driven anaerobic methane oxidation by oxygenic bacteria, *Nature*, **464**, 543–548,](#)
982 [doi.org/10.1038/nature08883, 2010.](#)

983

984 [Filipova-Marinova, M., Pavlov, D., Coolen, M and Giosan, L.: First high-resolution marinopolynological stratigraphy](#)
985 [of Late Quaternary sediments from the central part of the Bulgarian Black Sea area, *Quatern. Int.*, **293**, 170-183,](#)
986 [doi.org/10.1016/j.quaint.2012.05.002, 2013.](#)

987 -

988 [Frenzel, P., Thebrath, B., and Conrad, R.: Oxidation of methane in the oxic surface layer of a deep lake sediment](#)
989 [\(Lake Constance\), *FEMS Microbiology Letters*, **73**, 149–158, <https://doi.org/10.1111/j.1574-6968.1990.tb03935.x>,](#)
990 [1990.](#)

991

992 [Handley, L., Talbot, H.M., Cooke, M.P., Anderson, K.E and Wagner, T.: Bacteriohopanepolyols as tracers for](#)
993 [continental and marine organic matter supply and phases of enhanced nitrogen cycling on the late Quaternary Congo](#)
994 [deep sea fan, *Org. Geochem.*, **41**, 910-914, doi.org/10.1016/j.orggeochem.2010.04.016, 2010.](#)

995

996 [Hanson, R. S. and Hanson, T. E.: Methanotrophic bacteria, *Microbiol. Rev.*, **60**, 439–471,](#)
997 [https://doi.org/10.1128/mr.60.2.439-471.1996, 1996.](#)

998

999 [Hao, O.J., Chen, J.M., Huang, L and Buglass, R.L.: Sulfate- reducing bacteria, *Crit. Rev. Env. Sci. Tec.*, **26**, 155-187,](#)
1000 [doi.org/10.1080/10643389609388489, 1996.](#)

1001

1002 [He, R., Wang, J., Pohlman, J. W., Jia, Z., Chu, Y.-X., Wooller, M. J., and Leigh, M. B.: Metabolic flexibility of aerobic](#)
1003 [methanotrophs under anoxic conditions in Arctic lake sediments, *The ISME Journal*, **16**, 78–90,](#)
1004 [https://doi.org/10.1038/s41396-021-01049-y, 2022.](#)

1005

1006 [Heaton, T.J., Köhler, P., Butzin, M., Bard, E., Reimer, R.W., Austin, W.E.N., Bronk Ramsey, C., Grootes, P.M., Hughen,](#)
1007 [K.A., Kromer, B., Reimer, P.J., Adkins, J., Burke, A., Cook, M.S., Olsen, J and Skinner, L.C.: Marine20—The Marine](#)
1008 [Radiocarbon Age Calibration Curve \(0–55,000 cal BP\), *Radiocarbon*, **62**, 779–820, doi.org/10.1017/RDC.2020.68,](#)
1009 [2020.](#)

1010

1011 [Hiscott, R.N., Aksu, A.E., Mudie, P.J., Marret, F., Abrajano, T., Kaminski, M.A., Evans, J., Çakiroğlu, A.İ., Yaşar, D.: A](#)
1012 [gradual drowning of the southwestern Black Sea shelf: Evidence for a progressive rather than abrupt Holocene](#)

1013 [reconnection with the eastern Mediterranean Sea through the Marmara Sea Gateway, *Quatern. Int.*, **167–168**, 19–](#)
1014 [34, doi.org/10.1016/j.quaint.2006.11.007, 2007.](#)

1015

1016 [Hopmans, E.C., Weijers, J.W.H., Schefuß, E., Herfort, L., Sinninghe Damsté, J.S and Schouten, S.: A novel proxy for](#)
1017 [terrestrial organic matter in sediments based on branched and isoprenoid tetraether lipids, *Earth Planet. Sc. Lett.*,](#)
1018 [224, 107–116, doi.org/10.1016/j.epsl.2004.05.012, 2004.](#)

1019

1020 [Hopmans, E.C., Smit, N.T., Schwartz-Narbonne, R., Sinninghe Damsté, J.S and Rush, D.: Analysis of non-derivatized](#)
1021 [bacteriohopanepolyols using UHPLC-HRMS reveals great structural diversity in environmental lipid assemblages,](#)
1022 [Org. Geochem., **160**, 1–17, doi.org/10.1016/j.orggeochem.2021.104285, 2021.](#)

1023

1024 [Huang, Y., Zheng, Y., Heng, P., Giosan, L and Coolen, M.J.L.: Black Sea paleosalinity evolution since the last](#)
1025 [deglaciation reconstructed from alkenone-inferred Isochrysidales diversity, *Earth Planet. Sc. Lett.*, **564**, 1–9,](#)
1026 [doi.org/10.1016/j.epsl.2021.116881, 2021.](#)

1027

1028 [Ion, G., Briceag, A., Vasiliu, D., Lupaşcu, N and Melinte-Dobrinescu, M.: A multiproxy reconstruction of the Late](#)
1029 [Pleistocene-Holocene paleoenvironment: New insights from the NW Black Sea, *Mar. Geol.*, **443**, 1–19,](#)
1030 [doi.org/10.1016/j.margeo.2021.106648, 2022.](#)

1031 -

1032 [Ivanova, E.V., Murdmaa, I.O., Chepalyga, A.L., Cronin, T.M., Pasechnik, I.V., Levchenko, O.V., Howe, S.S., Manushkina,](#)
1033 [A.V and Platonova, E.A.: Holocene sea-level oscillations and environmental changes on the Eastern Black Sea shelf,](#)
1034 [Palaeogeogr. Palaeoclimatol. Palaeoecol., **246**, 228–259, doi.org/10.1016/j.palaeo.2006.09.014, 2007.](#)

1035 -

1036 [Jahnke, L.L., Summons, R., Hope, J.M and Des Marais, D.J.: Carbon isotopic fractionation in lipids from](#)
1037 [methanotrophic bacteria II: the effects of physiology and environmental parameters on the biosynthesis and isotopic](#)
1038 [signatures of biomarkers, *Geochim. Cosmochim. Ac.*, **63**, 79–93, doi.org/10.1016/S0016-7037\(98\)00270-1, 1999.](#)

1039 -

1040 [Jones, G.A and Gagnon, A.R.: Radiocarbon chronology of Black Sea sediments, *Deep-Sea Res. Pt. 1.*, **41**, 531–557,](#)
1041 [doi.org/10.1016/0967-0637\(94\)90094-9, 1994.](#)

1042 -

1043 [Kassambara, A and Mundt, F.: Factoextra: Extract and Visualize the Results of Multivariate Data Analyses, R Package](#)
1044 [Version 1.0.7, <http://CRAN.R-project.org/package=factoextra>, 2020.](#)

1045

1046 [Könneke, M., Bernhard, A.E., de la Torre, J.R., Walker, C.B., Waterbury, J.B and Stahl, D.A.: Isolation of an autotrophic](#)
1047 [ammonia-oxidizing marine archaeon, *Nature*, **437**, 543–546, doi.org/10.1038/nature03911, 2005.](#)

1048 -
1049 [Kool, D.M., Talbot, H.M., Rush, D., Ettwig, K and Sinninghe Damsté, J.S.: Rare bacteriohopanepolyols as markers for](#)
1050 [an autotrophic, intra-aerobic methanotroph, *Geochim. Cosmochim. Ac.*, **136**, 114-125,](#)
1051 [doi.org/10.1016/j.gca.2014.04.002, 2014.](#)
1052
1053 [Kuivila, K. M., Murray, J. W., Devol, A. H., Lidstrom, M. E., and Reimers, C. E.: Methane cycling in the sediments of](#)
1054 [Lake Washington, *Limnology and Oceanography*, **33**, 571–581, <https://doi.org/10.4319/lo.1988.33.4.0571>, 1988.](#)
1055
1056 [Kulkarni, G., Busset, N., Molinaro, A., Gargani, D., Chaintreuil, C., Silipo, A., Giraud, E and Newman, D. K.: Specific](#)
1057 [Hopanoid Classes Differentially Affect Free-Living and Symbiotic States of *Bradyrhizobium diazoefficiens*. *mBio*, **6**,](#)
1058 [e01251-15, doi.org/10.1128/mBio.01251-15, 2015.](#)
1059
1060 [Kusch, S., Sepúlveda, J and Wakeham, S.G.: Origin of Sedimentary BHPs Along a Mississippi River–Gulf of Mexico](#)
1061 [Export Transect: Insights From Spatial and Density Distributions, *Front. Mar. Sci.*, **6**, 1-14,](#)
1062 [doi.org/10.3389/fmars.2019.00729, 2019.](#)
1063 -
1064 [Kusch, S., Wakeham, S. G and Sepúlveda, J.: Diverse origins of “soil marker” bacteriohopanepolyols in marine oxygen](#)
1065 [deficient zones, *Org. Geochem.*, **151**, 104150, doi.org/10.1016/j.orggeochem.2020.104150, 2021.](#)
1066
1067 [Kusch, S., Wakeham, S.G and Sepúlveda, J.: Bacteriohopanepolyols across the Black Sea redoxcline trace diverse](#)
1068 [bacterial metabolisms, *Org. Geochem.*, **172**, 1-18, doi.org/10.1016/j.orggeochem.2022.104462, 2022.](#)
1069
1070 [Kuypers, M.M.M., Sliekers, A.O., Lavik, G., Schmid, M., Barker Jørgensen, B., Kuenen, J.G., Sinninghe Damsté, J.S.,](#)
1071 [Strous, M and Jetten, M.S.M.: Anaerobic ammonium oxidation by anammox bacteria in the Black Sea. *Nature*, **422**,](#)
1072 [608-611, doi.org/10.1038/nature01472, 2003.](#)
1073 -
1074 [Kuypers, M. M. M., Van Breugel, Y., Schouten, S., Erba, E and Sinninghe Damsté, J. S.: N₂-fixing cyanobacteria](#)
1075 [supplied nutrient N for Cretaceous oceanic anoxic events, *Geology*, **32**, 853, doi.org/10.1130/G20458.1, 2004.](#)
1076
1077 [Kwiecien, O., Arz, H.W., Lamy, F., Wulf, S., Bahr, A., Röhl, U and Haug, G.H.: Estimated Reservoir Ages of the Black](#)
1078 [Sea Since the Last Glacial, *Radiocarbon*, **50**, 99-118, doi.org/10.1017/S0033822200043393, 2008.](#)
1079
1080 [Lam, P., Jensen, M.M., Lavik, G., McGinnis, D.F., Müller, B., Schubert, C.J., Amann, R., Thamdrup, B., and Kuypers,](#)
1081 [M.M.M.: Linking crenarchaeal and bacterial nitrification to anammox in the Black Sea, *Proc. Nat. Acad. Sci. USA.*,](#)
1082 [104, 7104-7109, doi.org/10.1073/pnas.061108110, 2007.](#)

1083
1084 [Lê, S., Josse, J and Husson, F.: FactoMineR: An R Package for Multivariate Analysis, *J. Stat. Softw.*, 25\(1\), 1–18,](#)
1085 [doi.org/10.18637/jss.v025.i01, 2008.](#)
1086 -
1087 [Leloup, J., Loy, A., Knab, N.J., Borowski, C., Wagner, M. and Jørgensen, B.B.: Diversity and abundance of sulfate-](#)
1088 [reducing microorganisms in the sulfate and methane zones of a marine sediment, Black Sea, *Environ. Microbiol.*, 9,](#)
1089 [131-142, doi.org/10.1111/j.1462-2920.2006.01122.x, 2007.](#)
1090
1091 [Lin, X., Wakeham, S. G., Putnam, I. F., Astor, Y. M., Scranton, M. I., Chistoserdov, A. Y and Taylor, G. T.: Comparison](#)
1092 [of Vertical Distributions of Prokaryotic Assemblages in the Anoxic Cariaco Basin and Black Sea by Use of Fluorescence](#)
1093 [In Situ Hybridization, *Appl. Environ. Microbiol.*, 72, 2679–2690, doi.org/10.1128/AEM.72.4.2679-2690.2006, 2006.](#)
1094
1095 [Major, C., Goldstein, S., Ryan, W., Lericolais, G., Piotrowski, A.M and Hajdas, I.: The co-evolution of Black Sea level](#)
1096 [and composition through the last deglaciation and its paleoclimatic significance, *Quaternary Sci. Rev.*, 25, 2031-2047,](#)
1097 [doi.org/10.1016/j.quascirev.2006.01.032, 2006.](#)
1098 -
1099 [Marret, F., Mudie, P., Aksu, A and Hiscott, R.N.: A Holocene dinocyst record of a two-step transformation of the](#)
1100 [Neoeuxinian brackish water lake into the Black Sea, *Quatern. Int.*, 197, 72-86, doi.org/10.1016/j.quaint.2007.01.010,](#)
1101 [2009.](#)
1102 -
1103 [Martinez-Cruz, K., Leewis, M.-C., Herriott, I. C., Sepúlveda-Jauregui, A., Anthony, K. W., Thalasso, F., and Leigh, M.](#)
1104 [B.: Anaerobic oxidation of methane by aerobic methanotrophs in sub-Arctic lake sediments, *Science of The Total*](#)
1105 [Environment, 607–608, 23–31, https://doi.org/10.1016/j.scitotenv.2017.06.187, 2017.](#)
1106
1107 [Matys, E.D., Sepúlveda, J., Pantoja, S., Lange, C.B., Caniupán, M., Lamy, F and Summons, R. E.: Bacteriohopanepolyols](#)
1108 [along redox gradients in the Humboldt Current System off northern Chile, *Geobiology*, 15, 844-857,](#)
1109 [doi.org/10.1111/gbi.12250, 2017.](#)
1110
1111 [Naafs, B. D. A., Bianchini, G., Monteiro, F. M and Sánchez-Baracaldo, P.: The occurrence of 2-methylhopanoids in](#)
1112 [modern bacteria and the geological record, *Geobiology*, 20, 41–59, doi.org/10.1111/gbi.12465, 2022.](#)
1113
1114 [Neunlist, S. and Rohmer, M.: Novel hopanoids from the methylotrophic bacteria *Methylococcus capsulatus* and](#)
1115 [Methylomonas methanica. \(22S\)-35-aminobacteriohopane-30,31,32,33,34-pentol and \(22S\)-35-amino-3 beta-](#)
1116 [methylbacteriohopane-30,31,32,33,34-pentol, *Biochemical Journal*, 231, 635–639, doi:10.1042/bj2310635, 1985a.](#)
1117

1118 [Neunlist, S. and Rohmer, M.: The Hopanoids of 'Methylosinus trichosporium': Aminobacteriohopanetriol and](#)
1119 [Aminobacteriohopanetetrol, *Microbiology*, **131**, 1363–1367, doi.org/10.1099/00221287-131-6-1363, 1985b.](#)

1120

1121 [Nicholas, W.A., Chivas, A.R., Murray-Wallace, C.V and Fink, D.: Prompt transgression and gradual salinisation of the](#)
1122 [Black Sea during the early Holocene constrained by amino acid racemization and radiocarbon dating, *Quaternary*](#)
1123 [Sci. Rev.](#), **30**, 3769-3790, doi.org/10.1016/j.quascirev.2011.09.018, 2011.

1124 -

1125 [Oksanen, J., Simpson, G. L., Blanchet, F. G., Kindt, R., Legendre, P., Minchin, P. R., O'Hara, R. B., Solymos, P., Stevens,](#)
1126 [M. H. H., Szoecs, E., Wagner, H., Barbour, M., Bedward, M., Bolker, B., Borcard, D., Carvalho, G., Chirico, M., Caceres,](#)
1127 [M. D., Durand, S., Evangelista, H. B. A., FitzJohn, R., Friendly, M., Furneaux, B., Hannigan, G., Hill, M. O., Lahti, L.,](#)
1128 [McGlinn, D., Ouellette, M.-H., Cunha, E. R., Smith, T., Stier, A., Braak, C. J. F. T., and Weedon, J.: vegan: Community](#)
1129 [Ecology Package, 2022.](#)

1130 -

1131 [Oswald, K., Milucka, J., Brand, A., Hach, P., Littmann, S., Wehrli, B., Kuypers, M. M. M., and Schubert, C. J.: Aerobic](#)
1132 [gammaproteobacterial methanotrophs mitigate methane emissions from oxic and anoxic lake waters, *Limnology*](#)
1133 [and *Oceanography*, **61**, S101–S118, https://doi.org/10.1002/lno.10312, 2016.](#)

1134

1135 [Ourisson, G and Albrecht, P.: Hopanoids. 1. Geohopanoids: the most abundant natural products on Earth?, *Accounts*](#)
1136 [Chem. Res.](#), **25**, 398-402, doi.org/10.1021/ar00021a003, 1992.

1137 -

1138 [Piper, D.Z and Calvert, S.E.: Holocene and late glacial palaeoceanography and palaeolimnology of the Black Sea:](#)
1139 [Changing sediment provenance and basin hydrography over the past 20,000 years, *Geochim. Cosmochim. Ac.*, **75**,](#)
1140 [5597-5624, doi.org/10.1016/j.gca.2011.07.016, 2011.](#)

1141

1142 [R Core Team.: R: A Language and Environment for Statistical Computing, Vienna, Austria, https://www.R-](#)
1143 [project.org/, 2023.](#)

1144

1145 [Rashby, S. E., Sessions, A. L., Summons, R. E and Newman, D. K.: Biosynthesis of 2-methylbacteriohopanepolyols by](#)
1146 [an anoxygenic phototroph, *Proc. Nat. Acad. Sci. USA.*, **104**, 15099–15104, doi.org/10.1073/pnas.0704912104, 2007.](#)

1147

1148 [Reeburgh, W. S., Ward, B. B., Whalen, S. C., Sandbeck, K. A., Kilpatrick, K. A and Kerkhof, L. J.: Black Sea methane](#)
1149 [geochemistry, *Deep-Sea Res. Pt. 1.*, **38**, S1189–S1210, doi.org/10.1016/S0198-0149\(10\)80030-5, 1991.](#)

1150

1151 [Reimer, P.J., Austin, W.E.N., Bard, E., Bayliss, A., Blackwell, P.G., Bronk Ramsey, C., Butzin, M., Cheng, H., Edwards,](#)
1152 [R.L., Friedrich, N., Grootes, P.M., Guilderson, T.P., Haidas, L., Heaton, T.J., Hogg, A.G., Hughen, K.A., Kromer, B.,](#)

1153 [Manning, S.W., Muscheler, R., Palmer, J.G., Pearson, C., van der Plicht, J., Reimer, R.W., Richards, D.A., Scott, E.M.,](#)
1154 [Southon, J.R., Turney, C.S.M., Wacker, L., Adolphi, F., Büntgen, U., Capano, M., Fahrni, S.M., Fogtmann-Schulz, A.,](#)
1155 [Friedrich, R., Köhler, P., Kudsk, S., Miyake, F., Olsen, J., Reinig, F., Sakamoto, M., Sookdeo, A and Talamo, S.: The](#)
1156 [IntCal20 Northern Hemisphere Radiocarbon Age Calibration Curve \(0–55 cal kBP\), *Radiocarbon*, **62**, 725–757,](#)
1157 [doi.org/10.1017/RDC.2020.41, 2020.](#)
1158 -
1159 [Rethemeyer, J., Schubotz, F., Talbot, H.M., Cooke, M.P., Hinrichs, K.-U., Mollenhauer, G.: Distribution of polar](#)
1160 [membrane lipids in permafrost soils and sediments of a small high Arctic catchment, *Org. Geochem.*, **41**, 1130–1145,](#)
1161 [doi.org/10.1016/j.orggeochem.2010.06.004, 2010.](#)
1162
1163 [Repeta, D.J.: A high resolution historical record of Holocene anoxygenic primary production in the Black Sea,](#)
1164 [*Geochim. Cosmochim. Ac.*, **57**, 4337–4342, doi.org/10.1016/0016-7037\(93\)90334-S, 1993.](#)
1165
1166 [Řezanka, T., Siristova, L., Melzoch, K and Sigler, K.: N-Acylated Bacteriohopanehexol-Mannosamides from the](#)
1167 [Thermophilic Bacterium *Alicyclobacillus acidoterrestris*, *Lipids*, **46**, 249–261, doi.org/10.1007/s11745-010-3482-4,](#)
1168 [2011.](#)
1169
1170 [Ricci, J. N., Coleman, M. L., Welander, P. V., Sessions, A. L., Summons, R. E., Spear, J. R., and Newman, D. K.: Diverse](#)
1171 [capacity for 2-methylhopanoid production correlates with a specific ecological niche, *ISME J.*, **8**, 675–684,](#)
1172 [doi.org/10.1038/ismej.2013.191, 2014.](#)
1173
1174 [Richter, N., Hopmans, E.C., Mitrovic, D., Raposeiro, O.M., Gonçalves, P., Costa, A.C., Amaral-Zettler, L.A., Villanueva,](#)
1175 [L and Rush, D.: Distributions of bacteriohopanepolyols in lakes and coastal lagoons of the Azores Archipelago,](#)
1176 [*Biogeosciences*, **20**, 2065–2098, doi.org/10.5194/bg-20-2065-2023, 2023.](#)
1177
1178 [Rodier, C., Llopiz, P and Neunlist, S.: C32 and C34 hopanoids in recent sediments of European lakes: novel](#)
1179 [intermediates in the early diagenesis of biohopanoids. *Org. Geochem.*, **30**, 713–716, doi.org/10.1016/S0146-](#)
1180 [6380\(99\)00045-5, 1999.](#)
1181
1182 [Rohmer, M., Bouvier-Nave, P and Ourisson, G.: Distribution of Hopanoid Triterpenes in Prokaryotes, *Microbiology*,](#)
1183 [130, 1137–1150, doi.org/10.1099/00221287-130-5-1137, 1984.](#)
1184 -
1185 [Rush, D., Sinninghe Damsté, J.S., Poulton, S.W., Thamdrup, B., Garside, A.L., Acuña González, J., Schouten, S., Jetten,](#)
1186 [M.S.M and Talbot, H.M.: Anaerobic ammonium-oxidising bacteria: A biological source of the bacteriohopanetetrol](#)
1187 [stereoisomer in marine sediments, *Geochim. Cosmochim. Ac.*, **140**, 50–64, doi.org/10.1016/j.gca.2014.05.014, 2014.](#)

1188 -
1189 [Rush, D., Osborne, K.A., Birgel, D., Kappler, A., Hirayama, H., Peckmann, J., Poulton, S.W., Nickel, J.C., Mangelsdorf,](#)
1190 [K., Kalyuzhnaya, M., Sidgwick, F.R and Talbot, H.M.: The Bacteriohopanepolyol Inventory of Novel Aerobic Methane](#)
1191 [Oxidising Bacteria Reveals New Biomarker Signatures of Aerobic Methanotrophy in Marine Systems, *PLOS One*, **11**,](#)
1192 [1-27, doi.org/10.1371/journal.pone.0165635, 2016.](#)
1193
1194 [Rush, D., Talbot, H.M., van der Meer, M.T.J., Hopmans, E.C., Douglas, B and Sinninghe Damsté, J.S.: Biomarker](#)
1195 [evidence for the occurrence of anaerobic ammonium oxidation in the eastern Mediterranean Sea during Quaternary](#)
1196 [and Pliocene sapropel formation, *Biogeochemistry*, **16**, 2467-2479, doi.org/10.5194/bg-16-2467-2019, 2019.](#)
1197
1198 [Schaeffer, P., Schmitt, G., Adam, P., and Rohmer, M.: Acid-catalyzed formation of 32,35-](#)
1199 [anhydrobacteriohopanetetrol from bacteriohopanetetrol, *Org. Geochem.*, **39**, 1479–1482,](#)
1200 [doi.org/10.1016/j.orggeochem.2008.07.007, 2008.](#)
1201
1202 [Schaeffer, P., Schmitt, G., Adam, P., and Rohmer, M.: Abiotic formation of 32,35-anhydrobacteriohopanetetrol: A](#)
1203 [geomimetic approach, *Org. Geochem.*, **41**, 1005–1008, doi.org/10.1016/j.orggeochem.2010.04.013.](#)
1204
1205 [Schrader, H.-J. 1979. Quaternary Paleoclimatology of the Black Sea basin, *Sedimentary Geol.*, **23**, 165-180,](#)
1206 [doi.org/10.1016/0037-0738\(79\)90013-7, 2010.](#)
1207
1208 [Schouten, S., Wakeham, S. G and Damsté, J. S. S.: Evidence for anaerobic methane oxidation by archaea in euxinic](#)
1209 [waters of the Black Sea, *Org. Geochem.*, **32**, 1277–1281, doi.org/10.1016/S0146-6380\(01\)00110-3, 2001.](#)
1210
1211 [Schubert, C. J., Coolen, M. J. L., Neretin, L. N., Schippers, A., Abbas, B., Durisch-Kaiser, E., Wehrli, B., Hopmans, E. C.,](#)
1212 [Damsté, J. S. S., Wakeham, S., and Kuypers, M. M. M.: Aerobic and anaerobic methanotrophs in the Black Sea water](#)
1213 [column, *Environ. Microbiol.*, **8**, 1844–1856, doi.org/10.1111/j.1462-2920.2006.01079.x, 2006.](#)
1214
1215 [Schubert, C. J., Lucas, F. S., Durisch-Kaiser, E., Stierli, R., Diem, T., Scheidegger, O., Vazquez, F., and Müller, B.:](#)
1216 [Oxidation and emission of methane in a monomictic lake \(Rotsee, Switzerland\), *Aquat. Sci.*, **72**, 455–466,](#)
1217 [https://doi.org/10.1007/s00027-010-0148-5, 2010.](#)
1218
1219 [Schubotz, F., Wakeham, S. G., Lipp, J. S., Fredricks, H. F., and Hinrichs, K.: Detection of microbial biomass by intact](#)
1220 [polar membrane lipid analysis in the water column and surface sediments of the Black Sea, *Environ. Microbiol.*, **11**,](#)
1221 [2720–2734, doi.org/10.1111/j.1462-2920.2009.01999.x, 2009.](#)
1222 -

1223 [Schwartz-Narbonne, N., Schaeffer, P., Hopmans, E.C., Schenese, M., Charlton, E.A., Jones, D.M., Sinninghe Damsté,](#)
1224 [J.S., Farhan, M., Haque, U., Jetten, M.S.M., Lenggler, S.K., Murrell, J.C., Normand, P., Nuijten, G.H.L., Talbot, H.M and](#)
1225 [Rush, D.: A unique bacteriohopanetetrol stereoisomer of marine anammox, *Org. Geochem.*, **143**, 1-10,](#)
1226 [doi.org/10.1016/j.orggeochem.2020.103994, 2020.](#)
1227
1228 [Seemann, M., Bisset, P., Tritz, J.-P., Hooper, A. B and Rohmer, M.: Novel bacterial triterpenoids of the hopane](#)
1229 [series from *Nitrosomonas europaea* and their significance for the formation of the C35 bacteriohopane skeleton,](#)
1230 [Tetrahedron Lett., **40**, 1681–1684, doi.org/10.1016/S0040-4039\(99\)00064-7, 1999.](#)
1231
1232 [Shatz, M., Yosief, T and Kashman, Y.: Bacteriohopanehexol, a New Triterpene from the Marine Sponge *Petrosia*](#)
1233 [Species, *J. Nat. Prod.*, **63**, 1554–1556, doi.org/10.1021/np000190r, 2000.](#)
1234
1235 [Shumilovskikh, L.S., Tarasov, P., Arz, H.W., Fleitmann, D., Marret, F., Nowaczyk, N., Plessen, B., Schlütz, F and Behling,](#)
1236 [H.: Vegetation and environmental dynamics in the southern Black Sea region since 18 kyr BP derived from the marine](#)
1237 [core 22-GC3, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, **337–338**, 177-193, doi.org/10.1016/j.palaeo.2012.04.015,](#)
1238 [2012.](#)
1239
1240 [Sinninghe Damsté, J.S., Wakeham, S.G., Kohnen, M.E.L., Hayes, J.M., de Leeuw, J.W.: A 6,000-year sedimentary](#)
1241 [molecular record of chemocline excursions in the Black Sea, *Nature*, **362**, 827–829, doi.org/10.1038/362827a0,](#)
1242 [1993.](#)
1243
1244 [Sinninghe Damsté, J. S., Rijpstra, W. I. C., Dedysh, S. N., Foesel, B. U and Villanueva, L.: Pheno- and Genotyping of](#)
1245 [Hopanoid Production in Acidobacteria, *Front. Microbiol.*, **8**, 1-20, doi.org/10.3389/fmicb.2017.00968, 2017.](#)
1246
1247 [Sinninghe Damsté, J.S., Schouten, S., Hopmans, E.C., van Duin, A.C.T and Geenevasen, A.J.A.: Crenarchaeol, *J. Lipid*](#)
1248 [Res., **43**, 1641-1651, doi:10.1194/jlr.M200148-JLR200, 2002.](#)
1249
1250 [Sollai, M. Villanueva, L., Hopmans, E.C., Reichart, G.-J and Sinninghe Damsté, J.S.: A combined lipidomic and 16S](#)
1251 [rRNA gene amplicon sequencing approach reveals archaeal sources of intact polar lipids in the stratified Black Sea](#)
1252 [water column, *Geobiology*, **17**, 91-109, doi.org/10.1111/gbi.12316, 2019.](#)
1253
1254 [Spencer-Jones, C.L., Wagner, T and Talbot, H.M.: A record of aerobic methane oxidation in tropical Africa over the](#)
1255 [last 2.5 Ma, *Org. Geochem.*, **89-90**, 1-13, doi.org/10.1016/j.gca.2017.08.042, 2017.](#)
1256

1257 Stein, L.Y. and Nicol, G.W.: Nitrification, eLS., John Wiley & Sons: Chichester, UK, 1-9,
1258 doi.org/10.1002/9780470015902.a0021154.pub2, 2018.

1259

1260 Su, G., Zopfi, J., Niemann, H., and Lehmann, M. F.: Multiple Groups of Methanotrophic Bacteria Mediate Methane
1261 Oxidation in Anoxic Lake Sediments, *Front. Microbiol.*, **13**, 864630, <https://doi.org/10.3389/fmicb.2022.864630>,
1262 2022.

1263

1264 Summons, R. E., Jahnke, L. L., Hope, J. M and Logan, G. A.: 2-Methylhopanoids as biomarkers for cyanobacterial
1265 oxygenic photosynthesis, *Nature*, **400**, 554–557, doi.org/10.1038/23005, 1999.

1266

1267 Talbot, H. M., Watson, D. F., Murrell, J. C., Carter, J. F and Farrimond, P.: Analysis of intact bacteriohopanepolyols
1268 from methanotrophic bacteria by reversed-phase high-performance liquid chromatography–atmospheric pressure
1269 chemical ionisation mass spectrometry, *J. Chromatogr. A.*, **921**, 175–185, [doi.org/10.1016/S0021-9673\(01\)00871-8](https://doi.org/10.1016/S0021-9673(01)00871-8),
1270 2001.

1271

1272 Talbot, H. M., Watson, D.F., Pearson, E.J and Farrimond, P.: Diverse biohopanoid compositions of non-marine
1273 sediments, *Org. Geochem.*, **34**, 1353–1371, [doi.org/10.1016/S0146-6380\(03\)00159-1](https://doi.org/10.1016/S0146-6380(03)00159-1), 2003.

1274

1275 Talbot, H. M., Farrimond, P., Schaeffer, P., and Pancost, R. D.: Bacteriohopanepolyols in hydrothermal vent biogenic
1276 silicates, *Org. Geochem.*, **36**, 663–672, doi.org/10.1016/j.orggeochem.2004.10.015, 2005.

1277

1278 Talbot, H.M and Farrimond, P.: Bacterial populations recorded in diverse sedimentary biohopanoid distributions,
1279 *Org. Geochem.*, **38**, 1212–1225, <https://doi.org/10.1016/j.orggeochem.2007.04.006>, 2007.

1280

1281 Talbot, H. M., Summons, R. E., Jahnke, L. L., Cockell, C. S., Rohmer, M., and Farrimond, P.: Cyanobacterial
1282 bacteriohopanepolyol signatures from cultures and natural environmental settings, *Org. Geochem.*, **39**, 232–263,
1283 doi.org/10.1016/j.orggeochem.2007.08.006, 2008.

1284

1285 Talbot, H.M., Handley, L., Spencer-Jones, C.L., Dinga, B.J., Schefuß, E., Mann, P.J., Poulsen, J.R., Spencer, R.G.M.,
1286 Wabakanghanzi, J.N and Wagner, T.: Variability in aerobic methane oxidation over the past 1.2 Myrs recorded in
1287 microbial biomarker signatures from Congo fan sediments, *Geochim. Cosmochim. Ac.*, **133**, 387–401,
1288 doi.org/10.1016/j.gca.2014.02.035, 2014.

1289 -

1290 [Talbot, H.M., Sidgwick, F.R., Bischoff, J., Osborne, K.A., Rush, D., Sherry, A and Spencer-Jones, C.L.: Analysis of non-](#)
 1291 [derivatised bacteriohopanepolyols by ultrahigh-performance liquid chromatography/tandem mass spectrometry.](#)
 1292 [Rapid Commun. Mass Sp., **30**, 2087-2098, doi.org/10.1002/rcm.7696, 2016a.](#)
 1293
 1294 [Talbot, H.M., McClymont, E.L., Inglis, G.N., Evershed, R.P and Pancost, R.D.: Origin and preservation of](#)
 1295 [bacteriohopanepolyol signatures in Sphagnum peat from Bissendorfer Moor \(Germany\), *Org. Geochem.*, **97**, 95-110,](#)
 1296 [doi.org/10.1016/j.orggeochem.2016.04.011, 2016b.](#)
 1297
 1298 [Uemura, H. and Ishiwatari, R.: Identification of unusual 17 \$\beta\$ \(H\)-moret-22\(29\)-ene in lake sediments, *Org. Geochem.*,](#)
 1299 [23, 675-680, doi.org/10.1016/0146-6380\(95\)00036-E, 1995.](#)
 1300
 1301 [van Kemenade, Z.R., Cutmore, A., Hennekam, R., Hopmans, E.C., van der Meer, M.T.J., Mojtahid, M., Jorissen, F.J.,](#)
 1302 [Bale, N.J., Reichart, G.-J., Sinninghe Damsté, J.S and Rush, D.: Marine nitrogen cycling dynamics under altering redox](#)
 1303 [conditions: insights from deposition of sapropels S1 and the ambiguous S2 in the Eastern Mediterranean Sea,](#)
 1304 [Geochim. Cosmochim. Ac., **354**, doi.org/10.1016/j.gca.2023.06.018, 2023.](#)
 1305
 1306 [van Winden, J.F., Talbot, H.M., Kip, N., Reichart, J.-G., Pol, A., McNamara, N.P., Jetten, M.S.M., Op den Camp, H.J.M](#)
 1307 [and Sinninghe Damsté, J.S.: Bacteriohopanepolyol signatures as markers for methanotrophic bacteria in peat moss,](#)
 1308 [Geochim. Cosmochim. Ac., **77**, 52-61, doi.org/10.1016/j.gca.2011.10.026, 2012.](#)
 1309
 1310 [Valisolalao, J., Perakis, N., Chappe, B., Albrecht, P.: A novel sulfur containing C35 hopanoid in sediments, *Tetrahedron*](#)
 1311 [Lett., **25**, 1183-1186, doi.org/10.1016/S0040-4039\(01\)91555-2, 1984.](#)
 1312
 1313 [Verleye, T.J., Mertens, K.N., Louwye, S and Arz, H.W.: Holocene salinity changes in the southwestern black sea: A](#)
 1314 [reconstruction based on dinoflagellate cysts, *Palynology*, **33**, 77-100, doi:10.1080/01916122.2009.9989666, 2009.](#)
 1315
 1316 [Wakeham, S. G. and Beier, J. A.: Fatty acid and sterol biomarkers as indicators of particulate matter source and](#)
 1317 [alteration processes in the Black Sea, *Deep-Sea Res. Pt. 1.*, **38**, S943–S968, doi.org/10.1016/S0198-0149\(10\)80018-](#)
 1318 [4, 1991.](#)
 1319
 1319 [Wakeham, S. G., Beier, J. A., and Clifford, C. H.: Organic matter sources in the Black Sea as inferred from hydrocarbon](#)
 1320 [distributions, in: Black Sea Oceanography, edited by: İzdar, E., and Murray, J.W., Dordrecht: Springer Netherlands,](#)
 1321 [319-341, https://doi.org/10.1007/978-94-011-2608-3_20, 1991.](#)

1322 [Wakeham, S. G., Lewis, C. M., Hopmans, E. C., Schouten, S., and Damsté, J. S. S.: Archaea mediate anaerobic oxidation](#)
1323 [of methane in deep euxinic waters of the Black Sea, *Geochim. Cosmochim. Ac.*, **67**, 1359-1374,](#)
1324 [doi.org/10.1016/S0016-7037\(02\)01220-6, 2003.](#)

1325

1326 [Wakeham, S.G., Amann, R., Freeman, K.H., Hopmans, E.C., Barker Jørgensen, B., Putnam, I.F., Schouten, S., Sinninghe](#)
1327 [Damsté, J.S., Talbot, H.M and Woebken, D.: Microbial ecology of the stratified water column of the Black Sea as](#)
1328 [revealed by a comprehensive biomarker study, *Org. Geochem.*, **38**, 2070-2097,](#)
1329 [doi.org/10.1016/j.orggeochem.2007.08.003, 2007.](#)

1330

1331 [Watson, D.F and Farrimond, P.: Novel polyfunctionalised geohopanooids in a recent lacustrine sediment \(Priest Pot,](#)
1332 [UK\). *Org. Geochem.*, **31**, 1247-1252, doi.org/10.1016/S0146-6380\(00\)00148-0, 2000.](#)

1333

1334 [Welander, P. V., Coleman, M. L., Sessions, A. L., Summons, R. E., and Newman, D. K.: Identification of a methylase](#)
1335 [required for 2-methylhopanoid production and implications for the interpretation of sedimentary hopanes, *Proc.*](#)
1336 [Nat. Acad. Sci. USA., **107**, 8537–8542, doi.org/10.1073/pnas.0912949107, 2010.](#)

1337

1338 [Welte, C.U., Rasigraf, O., Vaksmaa, A., Versantvoort, W., Arshad, A., Op den Camp, H.J.M., Jetten, M.S.M., Lüke, C.](#)
1339 [and Reimann, J.: Nitrate- and nitrite-dependent anaerobic oxidation of methane, *Environ. Microbiol. Rep.*, **8**, 941-](#)
1340 [955, doi.org/10.1111/1758-2229.12487, 2016.](#)

1341 -

1342 [Wickham, H and Chang, W.: Package 'ggplot2': Create elegant data visualisations using the grammar of graphics. R-](#)
1343 [package Version 3.4.0, <https://ggplot2.tidyverse.org/reference/ggplot2-package.html>, 2016.](#)

1344

1345 [Wuchter, C., Abbas, B., Coolen, M.J.L., Herfort, L., van Bleijswijk, J., Timmers, P., Strous, M., Teira, E., Herndl, G.J.,](#)
1346 [Middelburg, J.J., Schouten, S and Sinninghe Damsté, J.S.: Archaeal nitrification in the ocean, *Proc. Nat. Acad. Sci.*](#)
1347 [USA, **103**, 12317-12322, doi.org/10.1073/pnas.0600756103, 2006.](#)

1348

1349 [Xu, Y., Cooke, M.P., Talbot, H.M and Simpson, M.J.: Bacteriohopanepolyol signatures of bacterial populations in](#)
1350 [Western Canadian soils, *Org. Geochem.*, **40**, 79-86, doi.org/10.1016/j.orggeochem.2008.09.003, 2009.](#)

1351

1352 [Yanchilina, A. G., Ryan, W. B. F., Kenna, T. C., and McManus, J. F.: Meltwater floods into the Black and Caspian Seas](#)
1353 [during Heinrich Stadial 1, *Earth-Sci. Rev.*, **198**, 102931, doi.org/10.1016/j.earscirev.2019.102931, 2019.](#)

1354 -

1355 [Zindorf, M., Rush, D., Jaeger, J., Mix, A., Penkrot, M.L., Schnetgere, B., Sidgwick, F.R., Talbot, H.M., van der Land, C.,](#)
1356 [Wagner T., Walczak, M and März, C.: Reconstructing oxygen deficiency in the glacial Gulf of Alaska: Combining](#)

1357 biomarkers and trace metals as paleo-redox proxies, *Chem. Geol.*, **558**, 1-11,
1358 doi.org/10.1016/j.chemgeo.2020.119864, 2020.

1359

1360 Zhao, N., Berova, N., Nakanishi, K., Rohmer, M., Mougnot, P., and Jürgens, U. J.: Structures of two
1361 bacteriohopanoids with acyclic pentol side-chains from the cyanobacterium *Nostoc* PCC 6720, *Tetrahedron*, **52**,
1362 2777–2788, [doi.org/10.1016/0040-4020\(96\)00013-0](https://doi.org/10.1016/0040-4020(96)00013-0), 1996.

1363

1364 Zhu, Q.-Z., Elvert, M., Meador, T. B., Schröder, J. M., Doeana, K. D., Becker, K. W., Elling, F. J., Lipp, J. S., Heuer, V. B.,
1365 Zabel, M and Hinrichs, K.-U.: Comprehensive molecular-isotopic characterization of archaeal lipids in the Black Sea
1366 water column and underlying sediments, *Geobiology*, **22**, e12589, doi.org/10.1111/gbi.12589, 2024.

1367

1368 Zhu, C., Talbot, H. M., Wagner, T., Pan, J.-M., and Pancost, R. D.: Distribution of hopanoids along a land to sea
1369 transect: Implications for microbial ecology and the use of hopanoids in environmental studies, *Limnol. Oceanog.*,
1370 **56**, 1850–1865, doi.org/10.4319/lo.2011.56.5.1850, 2011.

1371

1372

1373

1374

1375

1376

1377

1378

1379

1380

1381

1382

1383

1384

1385

1386

1387

1388

Deleted: ¶

Deleted: Aksu, A., Hiscott, R.N., Kaminski, M.A., Mudie, P.J., Gillespie, H., Abrajano, T and Yasar, D. 2002. Last glacial–Holocene paleoceanography of the Black Sea and Marmara Sea: stable isotopic, foraminiferal and coccolith evidence. *Marine Geology*, **190**, 119–149. [doi.org/10.1016/S0025-3227\(02\)00345-6](https://doi.org/10.1016/S0025-3227(02)00345-6) ¶

Ankindinova, O., Hiscott, R.N., Aksu, A.E and Grimes, V. 2019. High-resolution Sr-isotopic evolution of Black Sea water during the Holocene: Implications for reconnection with the global ocean. *Marine Geology*, **407**, 213–228. doi.org/10.1016/j.margeo.2018.11.004 ¶

Arthur & Dean, 1998. Organic-matter production and preservation and evolution of anoxia in the Holocene Black Sea. *Palaeoceanography & Palaeoclimatology*, **13**, 395–411. doi.org/10.1029/98PA01161 ¶

Badertscher, S., Fleitmann, D., Cheng, H., Edwards, R.L., Göktürk, O.M., Zumbühl, A., Leuenberger, M and Tüysüz, O. 2011. Pleistocene water intrusions from the Mediterranean and Caspian seas into the Black Sea. *Nature Geoscience*, **4**, 236–239. doi.org/10.1038/ngeo1106 ¶

Bahr, A., Lamy, F., Arz, H., Kuhlmann, H and Wefer, G. 2005. Late glacial to Holocene climate and sedimentation history in the NW Black Sea. *Marine Geology*, **214**, 309–322. doi.org/10.1016/j.margeo.2004.11.013 ¶

Bahr, A., Arz, H., Lamy, F and Wefer, G. 2006. Late glacial to Holocene paleoenvironmental evolution of the Black Sea, reconstructed with stable oxygen isotope records obtained on ostracod shells. *Earth and Planetary Science Letters Volume*, **241**, 863–875. doi.org/10.1016/j.epsl.2005.10.036 ¶


Bahr, A., Lamy, F., Arz, H., Major, C., Kwiecien, O and Wefer, G. 2008. Abrupt changes of temperature and water chemistry in the late Pleistocene and early Holocene Black Sea. *Geochemistry, Geophysics, Geosystems*, **9**, 1–16. doi.org/10.1029/2007GC001683 ¶

Bale, N., Ding, S., Hopmans, E.C., Arts, M.G.I., Villanueva, L., Boschman, C., Haas, A.F., Schouten, S and Sinninghe Damsté, J.S. 2021. Lipidomics of Environmental Microbial Communities. I: Visualization of Component Distributions Using Untargeted Analysis of High-Resolution Mass Spectrometry Data. *Frontiers in Microbiology*, **12**, 1–15. doi.org/10.3389/fmicb.2021.659302 ¶

Bednarczyk, A., Hernandez, T. C., Schaeffer, P., Adam, P., Talbot, H. M., Farrimond, P., Riboulleau, A., Largeau, C., Derenne, S., Rohmer, M., and Albrecht, P. 2005. 32,35-Anhydrobacteriohopanetetrol: an unusual bacteriohopanepolyol widespread in recent and past environments, *Organic Geochemistry*, **36**, 673–677. doi.org/10.1016/j.orggeochem.2004.10.014 ¶


1518
1519
1520
1521

Figures




1522
1523
1524
1525
1526


Figure 1: a. Site of Black Sea core 64PE418 taken at a water depth of 1970 mbsl (Adapted from: Giorgi Balakhadze, English Wikipedia, 2016); and a schematic diagram of the Black Sea basin during: b. the marine euxinic phase; c. the transition phase; d. the oxic lacustrine phase.



a. General bacteriohopanepolyol (BHP) structure




b. BHP core structure modifications:




c. BHP side structure modifications:


Nucleoside bacteriohopanepolyols




Hydroxy-bacteriohopanepolyols



Ethenolamine- and propenolamine-bacteriohopanepolyols



N-formylated-35-aminobacteriohopanetriol



d. Other compounds:





Figure 2: (A) A general bacteriohopanepolyol (BHP) structure with the core hopane and extended side chain (example of BHT). (B) Common modifications to the core BHP structure, including additional methyl groups and unsaturations on the core structure. (C) Side chain configurations discussed in this study.

1532 Note: for methoxy-BHT and methoxy-ethanolamine-BHT, the placement of the methoxy group is arbitrary.
1533 (D) Structures of other compounds discussed in this study: crenarchaeol and isorenieratene.




1534 **Figure 3:** Results from a principal component analysis of the bacteriohopanepolyol (BHP) abundance
1535 with depth. The individuals plot shows the sample depths with the ellipses highlighting major groupings.
1536 The biplot shows the sample depth (colored circles) and the contributions of each BHP (where BHT =
1537 bacteriohopanetetrol, unsat. = unsaturated, Me = methyl, and HG = headgroup). Note: BHPs with low
1538 contributions (< 1%) are not shown.

Deleted: 2

Deleted: anhydro = anhydrous,

1542



1543


1544

1545

1546

1547


Figure 4: Relative abundance of peak areas (%) of all BHPs identified in core 64PE418 over the last 19.5 ka. Changes in abundance of each BHP (peak area per g TOC) over time is shown in the supplementary material.



Deleted:

Deleted: 3

1550




1551

1552

1553 **Figure 5:** Changes in the peak area per g TOC of Nu-BHPs in Black Sea core 64PE418: **a)** over the past
1554 19.5 ka; highlighting changes during: **b)** the oxic lacustrine phase; **c)** the transition phase; and **d)** the
1555 euxinic marine phase. Note the different scales on the y-axes. 'Early' and 'Late' Me-adenosylhopane_{HG}-
1556 diMe represents two different isomers.

1557



Deleted: 4

Deleted: during: a) the oxic lacustrine phase; b) the transition phase; and c) the euxinic marine phase. Note the different scales on the y-axes.




Figure 6: Changes in the peak area per g TOC of a number of potential diagnostic BHPs.

Deleted: 5

Page 10: [1] Deleted	CUTMORE, ANNA V.	21/08/2025 13:03:00
----------------------	------------------	---------------------

▼

▲

Page 12: [2] Deleted	CUTMORE, ANNA V.	21/08/2025 13:22:00
----------------------	------------------	---------------------

✖

Page 30: [3] Deleted	CUTMORE, ANNA V.	21/08/2025 11:04:00
----------------------	------------------	---------------------

✖