

Evaluation of Extreme Sea-Levels and Flood Return Period using Tidal Day Maxima at Coastal Locations in the United Kingdom

Stephen E. Taylor¹

¹Geomatix Ltd. UK

Correspondence to: [Dr. Stephen E. Taylor \(set@geomatix.net\)](mailto:set@geomatix.net)

Abstract. Tidal storm surges can result in significant inundation and damage if sea defences are insufficiently robust. Coastal planners need to know the risk of flooding so that sea defences and coastal developments can be specified and located appropriately. Since the original work on extreme value statistics by Gumbel & Lieblein (1954), several alternatives have been proposed for evaluating the risk of tidal inundation, with the Skew Surge Joint Probability Method (SSJPM) gaining popularity. However, SSJPM is complex and cannot always be applied generally. Guided by the search for a general method having wide application and amenable to automation, this paper re-examines the original approach of Gumbel & Lieblein and proposes a simple modification for combined peak selection and declustering: it is termed here TMAX, since it selects one maximum per tide. In comparison with the Gumbel & Lieblein (1954) method using annual maxima (later termed AMAX), the TMAX method offers more efficient use of extreme data events, and in addition simpler handling of missing data. The results of the TMAX method are compared with those of a recent UK study using the SSJPM method at the same 39 United Kingdom coastal locations. The broadly applicable TMAX method has the potential to offer more widespread calculation of flood return period, thereby improving strategies regarding coastal management and resilience.

1. Introduction

Coastal planners and developers require accurate estimates of coastal flood risk to design and site sea defences, coastal buildings, harbours, nuclear power stations, and other critical infrastructure. Coastal floods generally occur when three factors converge; a significant storm surge, a spring tide and the time of high tide occurs at or near the peak of the storm surge. It is predicted the frequency of such events will double by 2050 due to rising sea levels (Vitousek et al., 2017). This challenge is especially critical for low-lying coastal nations, including parts of the United Kingdom (UK), (Williams et al., 2016).

The required height of sea defences can vary quite rapidly along the coastline. This is, because the coastal seabed topography can locally magnify or diminish the effects of tidal and surge effects. Tide gauge data provides a statistical snapshot relevant to its location; whether it is sited in an exposed coastal location or in a sheltered harbour, the recorded information can serve as a vital local source for estimating flood probability and return period. Tide gauge data is traditionally used for tidal harmonic analysis and for the establishment of tidal levels such as the Highest Astronomical Tide

Deleted: and inundation...sited Gumbel's ...modifications and new methods...recently ..., often requiring manual intervention, and is difficult to automate. specifically applicable to tides that is ...t... proposes several modifications to...Gumbel's ... The novel technique...s initial time unit... is one tidal day, ...rather than the usual ... (...). Compared to AMAX..., ...more ...provides reduced variance in design height, more efficient...ly handles ... are d... ..35 ... [1]

Formatted: Font color: Auto

Deleted: identical ..., showing broad agreement...is... new approach provides a robust mechanism for extreme tide analysis and ...better informs strategies ...for ... [2]

Deleted:

Deleted: , as...Coastal tide...gauges...data s...probabilities ...periods..., providing the local records necessary for deriving flood risk parameters.... often ..., ... [3]

30 (HAT), see Doodson & Warburg 1941); more recently it finds use in the determination of boundary conditions for tidal
modelling and in their numerical validation (Zhang et al., 2003). However, extreme sea-level analysis can leverage the same
tide-gauge data to predict the flood return probability. It is important here to distinguish between HAT and extreme flood
height. HAT assumes average conditions for the meteorological component, whereas the conditions leading to extreme
flooding include the effects of the meteorological component. Throughout this paper, the term tide refers to the total
35 fluctuating sea level, while the astronomical tide refers to the deterministic predicted component caused by tidal forces
generated by the earth-solar and lunar-earth orbits. The difference between the two is termed the residual. The residual,
especially when large, may be caused by a storm surge but this is not necessarily the case. Similarly, although the residual
may often result from meteorological effects and may appear as noise it may contain a deterministic component, and the
terms noise and storm surge are generally best avoided in this context. The term 'ESLs' is used throughout to represent the
40 expression 'estimates of the exceedance probabilities of Extreme Sea-Levels'. Several statistical methods have been
developed to assess Extreme Sea Levels (ESL) from measurements of sea level, these are now briefly reviewed.

The original work by Gumbel & Lieblein in 1954 describes a method for the conversion of selected extreme
maxima into ESLs, with the majority of cases discussed being based upon the analysis of annual maxima; this later became
known as the AMAX method. Since AMAX uses annual maxima, significant events that rank second or lower each year are
45 not utilised. However, in his original paper, Gumbel & Lieblein also examined some extreme events without using
annualized data, such as the breaking point of yarn, and the breakdown voltage of electrical capacitors. Subsequently an
approach based upon threshold rather than grouping has become known as the "Peaks over Threshold" (POT) method, see
Coles (2001), while an approach based upon selection of the largest, r, peaks within each time block has become known as
the "r-largest" method (Smith 1986; Tawn 1988).

50 A different approach by Pugh and Vassie (1978), splits the total sea level into two components which sum to the
total tide: these are the deterministic astronomical tide component and the residual component. Each component is converted
into separate probability distributions which are then combined by convolution to produce a joint probability distribution
function (PDF). The method is known as the joint probability method (JPM). Its main advantage lies in its efficient use of
source data; all values of the residual contribute to the final probability distribution function (PDF), even if they are not at or
55 near high tide. However, there are two notable drawbacks to the JPM. First, the conversion of the PDF into design risk is
challenging, as it can depend on sampling period (Middleton & Thompson 1986; Tawn & Vassie, 1989). Second, the timing
of actual high tide is often shifted in relation to the underlying predicted astronomical tide; this introduces correlation
between two components, undermining the convolution (Tawn 1992). Nevertheless the JPM is widely used (McInnes et al.,
2013), despite criticism on the above grounds, (e.g. from Batstone et al., 2013).

60 The skew surge joint probability method, SSJPM, aims to overcome these shortcomings of the JPM, by replacing
the residual (i.e. the instantaneous difference of measured and predicted water level) in JPM with the height difference
between the maxima of the measured and predicted water level for each tidal cycle. This new height difference, referred to as
the "skew surge height," is claimed to have minimal correlation with tidal height at most locations and is therefore

Deleted: , (...can ... accurately
variation of ... period with
height... inherently ... that tidal
height noise... averages out...
flood defence heights, take ... into
account the effect of noise...
make this ... ment... which ... [4]

Deleted: The Gumbel Type I
AMAX ... uses only the largest
annual extreme events for its
extreme value analysis; with...
significant events that rank second
or lower not being utilised. To
address this apparent inefficient
use of available data, several
alternative techniques have been
developed. One such technique is
the "r-largest" method (Smith
1986; Tawn 1988), which involves
using a fixed number, r, of the
largest annual maxima for
analysis. Another approach is the
"Peaks over Threshold" (POT)
method described by Coles (2001),
which utilizes maxima that exceed
a certain threshold value to create a
new probability distribution ... [5]

Deleted: (...)... a ... andom
surge... converting ... JPM
output ... requires an understanding
of flood duration... and perhaps
more critically, research has
indicated that the residual and
astronomical components are not
statistically independent and are
not sufficiently uncorrelated
(Tawn and Vassie, 1989; Tawn,
1992), making the convolution of
these components questionable.
The correlation becomes
particularly evident during storm
surges, as ..., which ... [6]

Deleted: undermines ... u ... [7]

Formatted: Not Highlight

Deleted: though... it has also
faced ... [8]

Deleted: (Batstone et al., 2013)

Formatted: Not Highlight

Formatted: Not Highlight

Deleted: ... this ... by
using... the ... maxima... rather
than the difference between
instantaneous height (Batstone et
al. 2013). ... [9]

65 considered an ideal parameter for characterizing surge statistics (Williams et al., 2016). To accurately extrapolate the tail of the probability curve, an extreme value probability distribution, such as the generalized Pareto distribution (GPD), is used. The method was used in the UK Environment Agency Study, 2011, EA (2011) as described by Batstone et al. (2013), and referred to onwards as EA2011, which used source data from the UK National Tide Gauge Network (UKNTGN) locations. However, Batstone et al. reported: "*The ESL's values derived from the SSJPM were compared with the AMAX time series of sea-level*". "At approximately one-quarter of the UKNTGN sites, it became clear that the GPD fitted on the skew surge distribution was leading to a seemingly implausible representation of the most extreme sea levels." "*Adjustments to the GPD shape parameter were performed by averaging the value with four immediate neighbours weighted by the length of data at each site*". The problem was attributed to correlation between skew surge and predicted tide. Williams et al., 2016, reported that when "seasonal relationships between tides and the storm season were removed, then skew surge and associated HW are completely independent at 68 of our 77 study sites." Thus, even after removing seasonal effects, skew surge and associated HW were not completely independent at a minority, but significant number, of locations. Here is the dilemma of the SSJPM method. It works well when the skew surge is independent of the tide but how can we know in advance when this is the case? As we have seen, in EA2011, approximately one-quarter in that study had to be examined and re-calculated using other adjustments.

80 The above situation indicates that a generally applicable means of delivering quality ESLs from tide gauge data would be a most useful tool if, even if the results were somewhat less accurate than the very best results of the SSJPM, i.e. where skew tides and the tidal heights are substantially independent. This paper pursues such a goal by re-examining the work of Gumbel & Lieblein (1954) and incorporating a relatively simple modification. The results of this approach, called here TMAX, are compared with the EA2011 study which used the SSJPM method. Unlike in EA2011, the TMAX method did not require manual intervention, and used an identical algorithm for each location. Note that the primary purpose of this study is to examine the suitability of the TMAX technique, by comparing its results with those of the AMAX and SSJPM methods using the results of EA2011. Note that, although it would be useful to make comparisons at as many locations world-wide as possible, the author has not been able to locate the necessary suitable data sources for carrying out such a study. Therefore this study is limited to the UK locations in the EA2011. Its purpose is to examine the proof of concept for the described TMAX method, rather than being an attempt to redefine UK ESL flood defence heights.

90 2. Background

W.E. Fuller (1914) claimed that, on a purely empirical basis, the size of floods increases proportionately to the logarithm of time. Some 40 years later Gumbel & Lieblein, in their classic 1954 paper "Statistical Theory of Extreme Values and Some Practical Applications", theoretically justified Fuller's claim, provided the extreme values are derived from a stationary series, i.e. one whose mean did not drift uniformly with time, and have a probability distribution of an exponential type. This latter condition, later known as a Gumbel/Fisher-Tippett Type I distribution, applies but is not limited to

Deleted: suitable candidate... However, there are only a limited number of skew extreme high tides, resulting in fewer output points compared to the Joint Probability Method (JPM). ... needed... For a complete set of equations, the reader should refer to Batstone (2013). SSJPM was applied to tidal locations around the UK... but as ... [10]

Formatted: Font: Italic

Deleted: The values

Formatted: [11]

Deleted: were corrected by manual intervention using the AMAX method and also data values from nearby tidal s [12]

Formatted: [13]

Deleted: ¶

Deleted: review highlights ... the need for... method ... analysing that can consistently... deliver design risk as a function of height, without requiring manual intervention. ... discusses the development of such a method, building on the AMAX approach but utilising more extremes each year, when applicable. The method initially identifies the highest sea level values recorded each tidal day rather than each year, leading to the acronym TMAX. These maxima are sorted and truncated, leaving a set of the most extreme, n values as inputs to the analysis. The paper discusses how a judicious choice of n reduces uncertainty in the resulting output; the flood return period as [14]

Formatted: Highlight

Deleted: he results of t... is... TMAX ... demonstrated ... good agreement with findings fr [15]

Formatted: Highlight

Deleted: the UK Environment Agency... study..., running ... the same ... very tidal... The f [16]

Formatted: Highlight

Deleted: Throughout this text, the term tide refers to the total sea level, while the astronomi [17]

Deleted: his

exponential, normal, chi-squared, logistical and log-normal distributions. (See also Leadbetter 1983, Tawn 1988). Gumbel gave many practical examples ranging from floods, radioactive decay, human life expectation, electrical capacitor breakdown, the strength of yarn, and the stock market share value. In Gumbel's original description (see his Eq.(2.17), a number, N, of observed peak values (generally annual), were ranked in ascending order with each ranked value i, being converted into the cumulative probability of a value not exceeding the ranked value, F_i by the formula.

$$F_i = i / (N+1) \quad (1)$$

The probability F_i is related to the return period T (generally in years) by Gumbel's Eq.(2.8), i.e.

$$T_i = 1 / (1 - F_i) \quad (2)$$

Gumbel showed that for a Type I distribution, the probability of the value being below a given value can be written as

$$F = \exp(- \exp(- y)) \quad (3)$$

The reduced variate, y exhibits a linear relationship with observed extreme value x, via a scale factor α and location factor, μ being given by

$$y = \alpha (x - \mu) \quad (4)$$

Therefore, in a plot of observed extreme height x, versus reduced variate y, each extreme value falls approximately in a straight line. In Gumbel's time, each ranked value was physically plotted on probability paper, where the horizontal scale had been marked out according to Eq.(3), in terms of either return period or reduced variate or both, depending upon the manufacturer of the paper. A straight line was fitted to the points, and extrapolation of the line gave the probability of a value not exceeding a given value x, without explicitly requiring the calculation of the scale factor, α or location factor, μ . Returning now to Eq.(1), Gringorten (1963) proposed a widely accepted correction to improve the plotting accuracy as

$$F_i = (i - 0.44) / (N + 0.12) \quad (5)$$

3. TMAX Method

The method described in this study named TMAX, differs from the method described by Gumbel in three main ways. Firstly, the ascending order of maxima used by Gumbel is replaced by a descending rank of maxima. Secondly, only one maximum

Deleted:

Deleted:

Formatted: Subscript

Formatted: Subscript

Formatted: Subscript

Deleted: equation

Deleted:

Deleted:

Deleted: equation

Deleted:

Deleted: which

Deleted: d

Deleted: and has been widely adopted

Deleted:

Deleted: It has already been mentioned that there is a requirement for an accurate, consistent and reliable method of establishing the relationship between flood design risk and sea defence height. The proposed technique described here, named TMAX, is designed to meet this need and to be suitable for automation without needing manual intervention, and which uses the data efficiently. TMAX

Deleted: (and subsequently called AMAX)

Deleted: a tidal day replaces the annual grouping used in AMAX (hence the name TMAX), s

is detected within each tide. Finally, only a subset of the total number of peak values are selected and used to fit the straight line in the probability plot. These differences are now considered in more detail.

3.1 Descending Rank

There are advantages in reversing the rank order as compared with Gumbel's original scheme, see Harris (1996), and this reversal is used here. Since the largest most extreme value is known it can readily be indexed as 1 while successively smaller values are indexed with rising integers (2,3, etc). We consider here $F(x)$ to be the probability of the tide exceeding a height value x (i.e. to be the flood probability), rather than the opposite as in Gumbel's original formulation. Adding a prime to those equations of Gumbel i.e. F' in Eq.s (2 & 3) and noting that $F' = 1 - F$, Eq.(2) & Eq.(3) now become,

$$T = 1 / (1 - F') = 1 / F \quad (6)$$

$$F = 1 - \exp(-\exp(-y)) \quad (7)$$

where y is the reduced variate. Making y the subject of Eq.(7) gives,

$$y = -\log_e(-\log_e(1 - F)) \quad (8)$$

We note that F is very much less than 1, hence the right hand logarithm can be represented by $-F$, and since F and T are inversely related we obtain $y = \log(T)$, confirming Fuller's original claim that maximum flood height varies with the logarithm of time. Writing Gumbel's descending rank, as i' , we write the ascending rank, i , as $i = N + 1 - i'$. Substituting this into Eq.(1) we obtain

$$F_i = 1 - F_{i'} = 1 - i' / (N+1) = i / (N+1) \quad (9)$$

and applying Gringorten's Correction we obtain

$$F_i = 1 - F_{i'} = 1 - (i' - 0.44) / (N + 0.12) = (i - 0.44) / (N + 0.12) \quad (10)$$

Since (10) is identical in form to Eq.(5), the reversal of rank order does not affect Gumbel's original plotting formulae, nor Gringorten's correction to it. The Gringorten formula, rather than Eq.(1), was used in all of the relevant calculations from hereon. The design risk $D(x)$, i.e. the probability that a given value of x will be exceeded during a design life consisting of n durations (usually years), is given by.

Deleted: , and finally a descending rank replaces the ascending order used by Gumbel. Further

Deleted: etails

Deleted:

Deleted: described

Deleted: ¶

3.1 Tidal Day¶

Most but not all of the examples given by Gumbel use the annual maxima, which later became known as the AMAX method. However, in his original paper, Gumbel also examines some extreme events without using annualized data, such as the breaking point of yarn, and the breakdown voltage of electrical capacitors. He states in his ... [18]

Deleted: 2

Deleted: Harris (1996) i ... [19]

Deleted: s

Deleted: s are

Deleted: but the least ar ... [20]

Formatted ... [21]

Deleted: refer to the abc ... [22]

Deleted: from Gumbel's ... [23]

Deleted: ,

Deleted: hence equation

Deleted: s

Deleted:

Deleted: now become

Deleted: ¶

Deleted: . .

Deleted: ¶

Deleted: w

Deleted: Conversion fro ... [24]

Deleted: namely.

Formatted ... [25]

Formatted ... [26]

Formatted: Subscript

Formatted: Subscript

Deleted: Remembering ... [27]

160
165
170
175
180
185
190

$$D(x) = 1 - (1 - F(x))^n$$

(11)

3.2 Tide Peak Detection Algorithm

Most but not all of the examples given by Gumbel use the annual maxima. Gumbel stated in his conclusion, "If the number of observed extremes N is not excessive, do not group the observations." Therefore, although extrema are generally grouped into annual time blocks, it is not necessary to do so. In the TMAX method described, maxima are not explicitly grouped. Only one peak is detected per tide (semi-diurnal or diurnal) with the largest peaks in the tide gauge record being identified as follows. The data is initially scanned and the mean height value, msl, is approximately found. The data is again re-scanned, and when the height value transitions above msl a search flag is set and the date and time of the upwards transition is stored. Once the search flag is set, the highest value is determined until the sequence transitions downwards below the msl. At this point the search flag is unset and the date and height of the highest value found during the period when the tide is above msl. Provided that the date-time of the downwards transition minus the date-time of the upwards transition is less than half of a tidal day the event is stored in a list as a maximum. This latter condition avoids counting transitions when the data has large gaps. Furthermore, assuming the storm surge duration is less than that of the tide, it also prevents the selection of multiple peaks from within the same storm event. Therefore, in addition to providing a determination of relevant peak value, it is also a form of declustering (see below). This list produced may include smaller peaks associated with noise and smaller semi-diurnal tides, but their existence in the list is irrelevant as only a small number n of the most significant are ultimately used, as is now described.

3.3 Subset Selection

Clearly, for a duration of tidal data of many decades, there will be many thousands of peak values. However, using all of the peak values on a probability plot is not advisable because the plot of height against the logarithm of return period sometimes deviates from a straight line at lower values of height; Fig. 1 shows this curvature and how it adversely affects the straight line fit. The curvature is probably related to shallow water effects during the neap of a spring-neap cycle. We use a fit of only the n largest values, since these fall, more or less, in a straight line as shown in Fig. 1, where n is calculated from an average number per year, n_A multiplied by the tide gauge record duration in years n_y , i.e. $n = n_A \times n_y$. The optimization of n_A is discussed in Sect. 4.1 and a value of 5 per year seems appropriate. The reader may consider here some similarities with r-largest, AMAX and Peaks Over Threshold (POT) methods. In r-largest, a fixed number of the greatest extreme events are selected for each time unit, whereas in TMAX the selection process applies to all of the data. In AMAX, one annual maxima is used, so TMAX could be viewed as being similar to AMAX but with a time unit of one or half a tidal day, depending upon whether the tide is diurnal or semidiurnal. However, TMAX incorporates further selection, using only the n largest values, whereas AMAX does not. The greatest similarity is with POT, where peaks are selected over the entire dataset, but in POT selection uses a certain threshold value, as opposed to in TMAX, where a given number of the largest values are selected.

- Deleted: 9
- Deleted: ¶
¶
- Deleted: 3
- Deleted: Peak
- Deleted: Selection
- Deleted: <sp>
- Deleted: In practice, it is relatively easy to determine t
- Deleted: a
- Deleted: As t
- Formatted: Font color: Brown
- Deleted: sequentially scanned, a transition of sea-level above mean-sea-level (MSL) is identified, and the largest value is found and stored before the sea-level scan returns below MSL. Continuing this scan generates a list of maxima.
- Formatted: Font:
- Deleted: <sp>
- Deleted: <sp>
- Formatted: Font: Bold

This is more much convenient than employing a threshold, since the number selected corresponds directly to the rank number which is already known; whereas using a threshold definition involves tidal range and datum, which are different for each location. Furthermore, in TMAX, the algorithm for peak detection (as described above), provides a level of isolation between successively selected peaks to select only one significant maximum per tidal day time block. This declustering avoids selection of multiple maxima from within the same extreme event "cluster", since such maxima would not be statistically "Independent and Identically Distributed" (IID). The key assumption here is that surges are of a sufficiently short duration that none will persist for more than one tidal day. Indeed on the list of maxima produced, none were found to occur on adjacent dates. One other key difference is that POT uses the generalised Pareto distribution (GPD) for fitting, whereas in TMAX, a simple least square regression linear fit is used.




Figure 1. Probability extreme tide plot showing its curvature in a 54-year record. Setting a threshold of 50 greatest extremes tides per year (2439 total), the LSQ intercept is 11.95m \pm 0.02m $R^2=0.9776$ (black). At 5 greatest extreme tides per year (242 points) the straighter red section gives an improved LSQ intercept of 11.6m (red) \pm 0.009m $R^2=0.9939$. Location Heysham UK See Sect. 4.1 & Figs. 2&3.

But what should be the value of N be in TMAX? If TMAX is considered as a modified form of AMAX, but with a tidal day rather than the annual time window, then N becomes the data duration D_D divided by tidal day length, T_D (in the same units), and the number of tidal days in a year is given by $1/T_D$, hence the usual ranking Eq.(9) becomes

$$T_i = N / i \text{ in tidal days} \quad (12)$$

Formatted: Font: Bold

Formatted: Keep with next

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Caption

Formatted: Font: Not Bold

Formatted: Indent: First line: 1.27 cm

Deleted: This list may include smaller peaks associated with noise and smaller semi-diurnal tides, but as only the most significant are ultimately used, their existence in the list is ... [28]

Formatted: Font color: Auto

Formatted: Font color: Auto

Formatted: Font color: Auto

Formatted: Font color: Auto

Formatted: ... [29]

Formatted: Font color: Auto

Formatted: Font color: Auto

Formatted: Font color: Auto

Formatted: ... [30]

Formatted: Font color: Auto

Formatted: Font color: Auto

Formatted: Font color: Auto

Formatted: ... [31]

Formatted: Font color: Auto

Formatted: ... [32]

Formatted: Font color: Auto

Formatted: Font color: Auto

Formatted: Font color: Auto

215 where $N=D_D/T_D$. Multiplying by T_D (in years) to convert to years, eliminates N becoming

$$T_i = D_D / i \text{ in years}$$

where 'i' is 1 based and corresponds to the rank having a maximum value of 'n' as defined earlier in this section.

220 Incorporating Gringorten's Correction Eq.(10) in Eq.(12) leads to

$$T_i = (N + 0.12) / (i - 0.44) \text{ in tidal days} \quad (13)$$

However, Eq.(14) can again be converted to years by multiplying by T_D giving

$$T_i = (D_D + 0.12 T_D) / (i - 0.44) \text{ years} \quad (14)$$

225 In practice, on TMAX, eliminating Gringorten's correction and using Eq.(12) rather than Eq.(13) was found on average to increase the ESL values by a few centimetres, and increased the 95% confidence limits slightly by a few millimetres; on AMAX the results were similar, except the 95% confidence limits were also increases by centimetres. Accordingly, this study utilised Eq.(13) with N specified in tidal days, the results being converted to years for display.

3.4 Straight Line Fit

235 As discussed above, the ordinary least squares (LSQ) fit, also known as linear regression, was used to determine the line fit (Morrison 2021). This is despite the existence of many other more complex methods including the method of maximum likelihood estimation (MME), method of moments (MOM), method of L-moments (MLM), method of probability-weighted moments (PWM), and the generalized least-squares methods GLSM/V. See Harris (1996), Coles (2000), Hong et al (2013), van Zyl & Schall (2012). Consideration of these other fitting methods in conjunction with TMAX is viewed as a potential refinement for future research. With both AMAX and TMAX methods, the expected value of a predicted new point (i.e. a flood) is the mean y value of the intersection of the extrapolated regression line with the ordinate corresponding to the required return period. Morrison gives expressions for the mean slope m_m , the mean intercept c_m , the variance of the expected mean σ_m and the variance of the new predicted mean σ_p , where n represents the number of data points, x_m , and y_m are the mean x and y values respectively, as

$$SS_{xy} = \sum (x_i - x_m)(y_i - y_m)$$

$$SS_{xx} = \sum (x_i - x_m)^2$$

$$m_m = SS_{xy} / SS_{xx}$$

$$c_m = ((\sum x_i)^2 \sum y_i - \sum x_i y_i \sum x_i) / (n \sum x_i^2 - (\sum x_i)^2)$$

$$\sigma_m = s^2_{y,x} (1/n + (x_p - x_m)^2 / SS_{xx}) \quad (15)$$

$$\sigma_p = s^2_{y,x} (1 + 1/n + (x_p - x_m)^2 / SS_{xx}) \quad (16)$$

$$s^2_{y,x} = (1/(n-2)) \sum (y_i - y_m)^2$$

$$sd = \sigma_p^{1/2} \quad (17)$$

Given the standard deviation, sd, the mean value of a predicted value y at a given value xp is given by

$$y = m_m x_p + c_m \pm t_{\alpha/2, n-2} \cdot sd$$

where 95% confidence interval, $t_{\alpha/2, n-2}$ signifies the t distribution and is commonly taken as approximately 2. The variance of the expected mean σ_m , Eq.(15), is always less than the variance of the new predicted value σ_p , Eq.(16), since the variance in the expected mean value is subject only to the distribution of data points, and tends to zero as the number of points, n increases, whereas the variance in a new estimated point is subjected also to variation in the process under examination, and does not tend to zero as n increases.

3.5 Missing Data

Tide gauge records often contain temporal data gaps due to faulty instrumentation, or damage to the gauge installation.

Using the TMAX method, files containing missing tidal data can be accommodated by simply reducing the value of N Eq.(13) to correspond to the number of tidal days of valid data. By contrast, it is not so straightforward in the AMAX or r-largest methods to accommodate missing data, since either synthetic data must be provided to fill the gaps, as was discussed by Gumbel, or whole years of data must be rejected if a significant portion of the annual data content is missing.

4. Comparative Study

The AMAX and TMAX methods were applied to calculate the ESLs for 39 of the 41 UK stations used in the UK study EA2011; the difference in number of ports arising because two, Hilbre Island and Exmouth were not included in the available BODC download. After downloading the tide gauge data, it was first concatenated for each port to form a single data file for each port; these were then imported and manually inspected. Most of the data before 1993 had been recorded at hourly intervals whereas subsequently, data was at 15-minute intervals. Although a spline fit had been considered as a strategy to artificially fill-in hourly results at 15 minute intervals it was considered advantageous to use the data directly, especially bearing in mind that the spline fit could result in new high tide values which had not actually been recorded. Bearing in mind that the TMAX method is dependant upon the height of the recorded peaks rather than their exact timing, the advantage of using much longer records would easily outweigh the potential small inaccuracies induced due to the change in sample rate. In addition a down-sampling experiment was carried out, by degrading the 15-min periods to hourly sampling. It changed the 200 year ESL by <1cm and the 95% confidence level by 1~2mm. Therefore, to obtain the longest data duration possible, both 15-minute and one-hourly data were used within a single file if necessary. Each file was

Formatted ... [46]

Deleted: 10

Deleted: 11

Deleted: where

Formatted: Font: (Default) Times New Roman

Deleted: ... SSxx = $\sum (x_i - x_m)^2$

Formatted ... [47]

Formatted ... [48]

Formatted ... [49]

Deleted: The ... variance of the new predicted mean σ_p is always greater than the variance of the mean σ_m ; practically this is because ..., whereas the variance in the intersection is subject to variation in the regression line parameters, the latter tending towards zero as the number of data points increases. ... [50]

Deleted: 5

Deleted: Frequently tide

Formatted: Normal

Deleted: data ... data ... taken into account ... tidal ... handle ... the below a given threshold... This is a considerable advantage of the TMAX over the AMAX ... [51]

Deleted: method. [51]

Formatted: Heading 2

Deleted: A comparison study was carried out of the AMAX, TMAX and SSJPM methods by repeating the study of Batstone 2013

Formatted: Normal

Deleted: at exactly the same 35 UK stations

Formatted ... [52]

Deleted: .

examined for tide gauge malfunctions and faulty data, and any data suspected of being faulty was deleted; this included isolated extreme maxima of non-contiguous data, gauge drifts and gauge slips. The maximum 5 values in each file were re-checked to verify the peaks represented a surge and not intermittent tide gauge faults

4.1 Optimising the average number of peaks per year.

As discussed above, selecting only the highest peaks avoids the graph curvature of Fig 1, improving its accuracy. Therefore, although larger sample values are generally associated with a decrease in variance, in this case the opposite may be true. This section investigates how, using the TMAX method, the variance of the mean, σ_m , and variance of a predicted new point, σ_p , as determined using Eqs.(10) & (11), varies with n_a , the average number of extremes selected per year. Figure 2 shows how σ_p attains a minimum at a value of n_a of around five extremes per year. This concurs with the value of five per year found by Robson and Reed (1999) for extreme river flow studies and used by Smith (1986), although this may be coincidental. Nevertheless, since we obtain a minimum variance at the value of $n_a = 5$, this was adopted for all subsequent results.




Figure 2. The mean-variance σ_m and of the predicted new point variance σ_p plotted against the number of extrema per year n_a for the TMAX method as determined from the tidal data.

Formatted: Centered

Formatted: Font: Times New Roman, Not Bold

Field Code Changed

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Font: Times New Roman, Not Bold

Formatted: Font: Not Bold

295

4.2 Analysis

For compatibility with EA2011, all data was sea-level rise de-trended, using the constant value and date origin adopted in that study of 2mm per year, with zero being applied on 1 January 2008. For further comparison purposes, the records were truncated to have a maximum date of 1 Jan 2009, which corresponded to the data used in EA2011, see Table A1.3. The analysis for dates to 1 Jan 2009 was run using both AMAX and TMAX methods, which are referred to here as AMAX2009 and TMAX 2009. During the analysis the peak selection algorithm as described above was used. The number of valid days of data was determined and only those in the top rank corresponding to a total number, n , corresponding to n_A of 5 per year were used. The peaks were ranked and plotted using Gringorten's formula, where the total number of samples was calculated on the basis of one per tidal day for each tidal day of valid data. The LSQ regression line analysis was then used to give an intercept for the required return periods. The entire analysis was then repeated using the complete data set to 1 May 2018, being references as AMAX2018 and TMAX2018. The periods from tidal days to years were converted as required. The results were derived solely from the tide gauge data and have not been re-processed in any other way. These are now described. Comparisons with the results from EA2011 are indicated where appropriate.

300

305

4.3 Results

Table 1 compares the results of TMAX2009 with EA2011 with ESLs in metres above ODN relative to MSL2008 for all 39 ports. The results are shown for each of the return periods of 20, 100, 200, and 1000 years. The column headed SD shows standard deviation for the TMAX2009 ESLs as obtained from the Least Squares Fit using Eq.(17). The column headed "Difference" shows the values returned from TMAX2009 minus those from EA2011. The lower two rows show the mean values and their standard deviations. The mean differences were of the order of centimetres while the standard deviations are of the order of 10 to 20 centimetres, with both rising for the longer return periods. Figure 3 shows the estimated ESL values for a 100-year return period at 39 UK sites on a map as derived from TMAX2018 rather than from TMAX2009, utilising all of the tidal data examined. The results are discussed in the error analysis in the following section.

310

315

Deleted: The tide gauge data was downloaded and manually inspected, and isolated non-contiguous data points and points showing clear evidence of gauge slips were removed. Most data before 1993 had been recorded at hourly intervals whereas subsequently, data was at 15-minute intervals. To obtain the longest data duration possible, both 15-minute and one-hourly data were used within a single file for many of the locations. Although there is a risk that the use of hourly data may result in missing or under-recording of the true peak value, most significant surges are at least one hour in duration, reducing this likelihood. The advantage of using much longer records should therefore easily outweigh the potential small inaccuracy due to the change in sample rate within the file.

Deleted: In the first phase of analysis, the optimum value of n_A was determined using the data processed above.

Deleted: In the second phase, the extreme sea levels corresponding to the return period were determined using the optimum value of n_A .

Deleted: and c

Deleted: Batstone

Deleted: ¶
4.1 Optimising the average number of peaks per year. ¶
 As discussed above, larger values of n_A cause a higher proportion of extreme peaks under investigation to originate from smaller astronomical tides (neap tides) whose probability distribution may differ. Therefore, although larger sample values are generally associated with a decrease in variance, in this case the opposite may be true. ¶

... [53]

Formatted: Heading 2

Formatted: Normal, Left

Table 1. ESLs (m) above ODN relative to MSL2008.
Results of TMAX2009, Standard Deviation of LSQ fit, Values of EA2011, and Difference TMAX2009-EA2011

Return Period (Yrs)	TMAX2009				SD				EA2011				Difference			
	20	100	200	1000	20	100	200	1000	20	100	200	1000	20	100	200	1000
Aberdeen	3.0	3.1	3.2	3.3	0.1	0.1	0.1	0.1	3.0	3.1	3.2	3.3	0.0	0.0	0.0	0.1
Avonmouth	8.7	8.9	9.0	9.3	0.2	0.2	0.2	0.2	8.7	9.0	9.1	9.4	0.0	-0.1	-0.1	-0.1
Barmouth	3.9	4.1	4.2	4.3	0.1	0.1	0.1	0.2	3.9	4.1	4.2	4.4	0.0	-0.1	-0.1	-0.1
Bournemouth	1.6	1.8	1.8	2.0	0.1	0.1	0.1	0.1	1.7	1.8	1.9	2.0	-0.1	0.0	0.0	0.0
Cromer	3.5	3.8	3.9	4.1	0.2	0.2	0.2	0.2	3.7	4.1	4.3	4.7	-0.2	-0.3	-0.4	-0.6
Devonport	3.2	3.3	3.4	3.5	0.1	0.1	0.1	0.1	3.2	3.4	3.5	3.6	-0.1	-0.1	-0.1	-0.1
Dover	4.3	4.5	4.6	4.8	0.1	0.2	0.2	0.2	4.2	4.5	4.6	4.8	0.0	0.0	0.0	0.0
Felixstowe Pier	3.0	3.3	3.3	3.6	0.1	0.2	0.2	0.2	3.3	3.7	3.9	4.4	-0.3	-0.5	-0.6	-0.8
Fishguard	3.3	3.5	3.5	3.7	0.1	0.1	0.1	0.1	3.4	3.5	3.6	3.7	0.0	-0.1	-0.1	-0.1
Hevsham	6.4	6.7	6.8	7.1	0.2	0.2	0.2	0.2	6.4	6.7	6.8	7.1	0.0	0.0	0.0	0.0
Hinkley Point	7.4	7.5	7.6	7.7	0.1	0.1	0.1	0.1	7.5	7.7	7.8	8.1	-0.1	-0.2	-0.3	-0.3
Holyhead	3.7	3.8	3.9	4.1	0.1	0.1	0.1	0.1	3.7	3.8	3.9	4.0	0.0	0.0	0.0	0.0
Jlfracombe	5.7	5.9	5.9	6.1	0.1	0.1	0.1	0.1	5.7	5.9	5.9	6.1	0.0	0.0	0.0	0.0
Jmningham	4.6	4.8	4.9	5.1	0.1	0.1	0.1	0.2	4.6	4.9	5.0	5.3	-0.1	-0.1	-0.1	-0.1
Kinlochbervie	3.8	4.0	4.1	4.4	0.2	0.2	0.2	0.2	3.6	3.8	3.9	4.2	0.2	0.2	0.2	0.2
Leith	3.7	3.8	3.9	4.0	0.1	0.1	0.1	0.1	3.7	3.9	4.0	4.2	0.0	-0.1	-0.1	-0.2
Lerwick	1.8	1.9	2.0	2.1	0.1	0.1	0.1	0.1	1.7	1.8	1.9	2.0	0.1	0.1	0.1	0.2
Llandudno	5.1	5.3	5.4	5.6	0.1	0.2	0.2	0.2	5.1	5.3	5.4	5.6	0.0	0.0	0.0	0.0
Lowestoft	2.7	3.0	3.2	3.6	0.2	0.2	0.2	0.3	2.7	3.1	3.3	3.8	0.0	0.0	-0.1	-0.2
Milford Haven	4.6	4.8	4.9	5.0	0.1	0.1	0.1	0.1	4.5	4.7	4.8	5.0	0.1	0.1	0.1	0.1
Millport	3.3	3.6	3.7	4.0	0.2	0.2	0.2	0.2	3.2	3.5	3.7	4.0	0.1	0.0	0.0	0.0
Moray Firth	3.2	3.4	3.4	3.5	0.1	0.1	0.1	0.1	3.1	3.3	3.4	3.5	0.1	0.1	0.1	0.0
Mumbles	5.8	6.0	6.1	6.3	0.1	0.1	0.1	0.2	5.8	6.1	6.2	6.4	0.0	0.0	0.0	-0.1
Newhaven	4.1	4.3	4.3	4.5	0.1	0.1	0.1	0.1	4.2	4.4	4.5	4.6	-0.1	-0.1	-0.1	-0.1
Newlyn	3.4	3.5	3.5	3.7	0.1	0.1	0.1	0.1	3.3	3.5	3.5	3.6	0.0	0.0	0.0	0.0
Newport	8.0	8.2	8.3	8.5	0.2	0.2	0.2	0.2	8.0	8.3	8.4	8.7	0.0	-0.1	-0.1	-0.2
North Shields	3.5	3.7	3.8	4.0	0.1	0.1	0.1	0.1	3.6	3.8	3.9	4.1	0.0	0.0	-0.1	-0.1
Portpatrick	3.2	3.4	3.5	3.7	0.1	0.1	0.1	0.1	3.2	3.4	3.5	3.6	0.0	0.1	0.1	0.1
Portsmouth	2.9	3.0	3.1	3.2	0.1	0.1	0.1	0.1	2.9	3.1	3.1	3.3	0.0	0.0	0.0	-0.1
Port Ellen	1.7	2.0	2.1	2.3	0.2	0.2	0.2	0.2	1.9	2.1	2.2	2.4	-0.2	-0.2	-0.2	-0.1
Port Erin	3.7	3.9	4.0	4.2	0.1	0.2	0.2	0.2	3.7	3.8	3.9	4.0	0.0	0.1	0.1	0.2
Sheerness	4.0	4.2	4.3	4.5	0.1	0.1	0.1	0.2	4.1	4.5	4.6	5.1	-0.1	-0.2	-0.3	-0.5
Stomoway	3.2	3.4	3.5	3.6	0.1	0.1	0.1	0.1	3.2	3.3	3.4	3.5	0.0	0.1	0.1	0.1
Tobermory	3.4	3.6	3.7	3.9	0.1	0.2	0.2	0.2	3.5	3.7	3.9	4.2	0.0	-0.1	-0.1	-0.2
Ullapool	3.7	3.9	3.9	4.1	0.1	0.1	0.1	0.1	3.6	3.8	3.8	4.0	0.1	0.1	0.1	0.2
Weymouth	2.1	2.2	2.3	2.4	0.1	0.1	0.1	0.1	2.1	2.2	2.3	2.4	0.1	0.0	0.0	0.0
Whitby	3.6	3.7	3.8	3.9	0.1	0.1	0.1	0.1	3.8	4.0	4.1	4.4	-0.2	-0.3	-0.3	-0.5
Wick	2.8	2.9	3.0	3.1	0.1	0.1	0.1	0.1	2.7	2.8	2.9	3.0	0.1	0.1	0.1	0.1
Workington	5.7	6.0	6.1	6.3	0.2	0.2	0.2	0.2	5.6	5.8	5.9	6.2	0.1	0.1	0.2	0.2
mean (m)					0.1	0.1	0.1	0.1					0.0	0.0	0.0	-0.1
stdev (m)													0.1	0.1	0.2	0.2

- Formatted ... [54]
- Formatted ... [55]
- Formatted Table ... [56]
- Formatted ... [57]
- Formatted ... [58]
- Formatted ... [59]
- Formatted ... [60]
- Formatted ... [61]
- Formatted ... [62]
- Formatted ... [63]
- Formatted ... [64]
- Formatted ... [65]
- Formatted ... [66]
- Formatted ... [67]
- Formatted ... [68]
- Formatted ... [69]
- Formatted ... [70]
- Formatted ... [71]
- Formatted ... [72]
- Formatted ... [73]
- Formatted ... [74]
- Formatted ... [75]
- Formatted ... [76]
- Formatted ... [77]
- Formatted ... [78]
- Formatted ... [79]
- Formatted ... [80]
- Formatted ... [81]
- Formatted ... [82]
- Formatted ... [83]
- Formatted ... [84]
- Formatted ... [85]
- Formatted ... [86]
- Formatted ... [87]
- Formatted ... [88]
- Formatted ... [89]
- Formatted ... [90]
- Formatted ... [91]
- Formatted ... [92]
- Formatted ... [93]
- Formatted ... [94]
- Formatted ... [95]
- Formatted ... [96]
- Formatted ... [97]
- Formatted ... [98]
- Formatted ... [99]
- Formatted ... [100]
- Formatted ... [101]
- Formatted ... [102]
- Formatted ... [103]
- Formatted ... [104]
- Formatted ... [105]
- Formatted ... [106]
- Formatted ... [107]



Figure 3. 100 Year ESL ODN (m) at 39 UK locations From TMAX2018 relative to MSL 2008

Deleted: <sp><sp><sp>Table 1: AMAX and TMAX methods. Quality of Fit of Regres(... [109]

Formatted: Caption, Centered, Keep with next

Formatted: Font: (Default) Arial, 8 pt

Formatted Table

Formatted: Highlight

Formatted: Keep with next

Formatted: Left

4.4 Error Analysis

Table 2 shows the standard deviation for each given return period and were calculated from the best fit to the source "n" maxima, using Eq.(10) and those of Sect 3.4. The standard deviation is calculated from the expected variance in a new forecast point σ_p Eq.(16) rather than the smaller variance in the mean σ_m Eq.(15), as indicated in Sect. 3.5. It can be seen that there is a significant difference between the standard deviation for the two methods, AMAX and TMAX, with TMAX significantly outperforming AMAX, reflecting the larger number of points involved in the TMAX method. On average the standard deviation for TMAX were approximately 60% of those of AMAX which suggest a useful margin of improvement in accuracy. The level of agreement between AMAX and TMAX with the results of EA2011 are now described.

Table 2: Mean of Standard Deviation for AMAX and TMAX based upon variance in predicted level, $\sigma_p^{1/2}$ of the LSO fit.

Return Period (years)	20	100	200	1000
AMAX2009	0.19	0.22	0.23	0.27
AMAX2018	0.20	0.22	0.23	0.26
TMAX2009	0.12	0.13	0.14	0.15
TMAX2018	0.12	0.13	0.14	0.14

* means of confidence bounds in Table A6.1 EA2011

Figure 4 shows the differences between the ESLs from TMAX and AMAX with EA2011 as histograms. Table 3 shows the means differences, and standard deviations of the differences. With AMAX the mean of the differences were generally slightly positive, indicating marginally higher ESLs as compared to EA2011, whereas with TMAX the ESLs were generally lower, indicating a slight reduction in sea-defence requirements. For AMAX the mean differences were slightly smaller than TMAX, indicating marginally better agreement, perhaps reflecting the use of AMAX in EA2011 at those locations where SSJPM was not used. However, the standard deviation of the difference were much greater than the mean difference, for all return periods listed. The standard deviation of the difference ranged from 0.09m for the 20 year return period with TMAX2009/2018 to 0.25 m for the 1000 years return period with AMAX. The Students "t" test was used establish the significance of the mean difference values, ($\alpha=0.05$, 39 pairs, $df=38$ tails=2), giving a critical value at a 95% confidence level of $t_{crit} = 2.02$. Out of sixteen comparisons made, fifteen determined "t" to be considerably smaller than this critical value, the exception being AMAX09 at a 1000 year return period where only a marginal significance was found. Thus, according to the paired t-test the values of the mean difference is not significant. Across all studies for TMAX2009, TMAX2018, AMAX2009 and AMAX2018 and their comparison with EA2011 there were no consistent positive outliers

- Formatted ... [110]
- Formatted ... [111]
- Formatted ... [112]
- Formatted ... [113]
- Formatted ... [114]
- Formatted ... [115]
- Formatted ... [116]
- Formatted ... [117]
- Formatted ... [118]
- Formatted ... [119]
- Formatted ... [120]
- Formatted ... [121]
- Formatted ... [122]
- Formatted ... [123]
- Formatted ... [124]
- Formatted ... [125]
- Formatted ... [126]
- Formatted ... [127]
- Formatted ... [128]
- Formatted ... [129]
- Formatted ... [130]
- Formatted ... [131]
- Formatted ... [132]
- Formatted ... [133]
- Formatted ... [134]
- Formatted ... [135]
- Formatted ... [136]
- Formatted ... [137]
- Formatted ... [138]
- Formatted ... [139]
- Formatted ... [140]
- Formatted ... [141]
- Formatted ... [142]
- Formatted ... [143]
- Formatted ... [144]
- Formatted ... [145]
- Formatted ... [146]
- Formatted ... [147]
- Formatted ... [148]
- Formatted ... [149]

360

exceeding two standard deviations above the mean. However there were consistently significant negative outliers, with over two standard deviations below the mean; these were Cromer, Felixstowe and Sheerness. Significantly, these ports had been singled out for discussion of levels in the EA2011 study. The port of Felixstowe consistently had return levels which are showed the largest significant difference from those of EA2011. Further study using TMAX of tidal records for the port of Harwich, at a distance of only 3 nautical miles from Felixstowe, gave ESL's showing a considerably improved level of agreement with the results of EA2011 for Felixstowe, although the implication of this are not clear.



365

Table 3. Standard Error in Mean and Mean Difference between AMAX, TMAX and EA2011.

	Return Period (years)	20	100	200	1000
AMAX2009	Mean Difference (m)	0.01	0.01	0.01	-0.01
	Std Dev(m)	0.10	0.15	0.18	0.25
AMAX2018	Mean Difference (m)	0.02	0.04	0.04	0.05
	Std Dev(m)	0.11	0.16	0.18	0.25
TMAX2009	Mean Difference (m)	-0.01	-0.04	-0.05	-0.08
	Std Dev	0.09	0.13	0.15	0.22
TMAX2018	Mean Difference (m)	-0.01	-0.03	-0.04	-0.07
	Std Dev(m)	0.09	0.13	0.15	0.22

- Formatted:** Caption, Centered, Keep with next
- Formatted:** Font: (Default) Times New Roman
- Formatted:** Font: (Default) Times New Roman, Bold
- Formatted:** Right, Line spacing: single
- Formatted Table**
- Formatted** (... [150])
- Formatted:** Font color: Brown
- Formatted** (... [151])
- Formatted:** Right, Line spacing: single
- Formatted** (... [152])
- Formatted:** Font: (Default) Times New Roman
- Formatted:** Line spacing: single
- Formatted** (... [153])
- Formatted** (... [154])
- Formatted:** Right, Line spacing: single
- Formatted** (... [155])
- Formatted:** Font: (Default) Times New Roman
- Formatted:** Line spacing: single
- Formatted** (... [156])
- Formatted** (... [157])
- Formatted:** Right, Line spacing: single
- Formatted** (... [158])
- Formatted:** Font: (Default) Times New Roman
- Formatted:** Line spacing: single
- Formatted** (... [159])
- Formatted** (... [160])
- Formatted** (... [161])

Formatted: Keep with next



370





Figure 4. Differences between TMAX, AMAX and EA2011, using data to 2009 and 2018.

Formatted: Caption

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Field Code Changed

Formatted: Font: Not Bold

375
380
385
390
395
400

5. Conclusion

The paper describes a method named here as "TMAX", which is designed to derive extreme sea level statistics (ESLs) from tide gauge data, based upon a modification of Gumbel and Lieblein's method of 1954. When compared with the AMAX method, TMAX confidence bounds are approximately 40% smaller than those for AMAX, suggesting a significant margin of improvement in accuracy. Further, it is shown that TMAX generates estimates of ESLs which are in broad agreement when compared with those published by UK Environment Agency 2011 (EA 2011) at all 39 locations, and good agreement at 36 locations, with agreement at 3 location in the southern North Sea, in particular at the port of Felixstowe being outliers. Overall mean differences between the two methods were of the order of centimetres and standard deviation of the order of decimetres. However, the TMAX method is much simpler than the SSJPM used in EA2011. Unlike in the EA2011 study, where one quarter of sites were singled out for special attention, the same algorithm was used at each location without "manual intervention". Also, unlike the SSJPM, the TMAX assumes a Gumbel Type 1 statistic and does not split the tide into two components, i.e. the predicted tide and skew tide. Hence it does require a) harmonic tidal analysis, b) the use of a probability density function, or c) the fit of a generalized Pareto distribution (GPD). TMAX therefore does not suffer from any errors induced in this process, which may, in a minority of cases, significantly influence the accuracy of the result. A further advantage of the TMAX method is that missing data, quite common in tide gauge records, is handled in a simple, efficient and elegant way. The study indicates that the TMAX method shows promise as a general screening method for extreme sea level analysis. It is hoped the approach may facilitate the generation of ESLs from tide gauge data, thereby informing and improving strategies for coastal management and resilience.

Acknowledgments

The author is grateful to the British Oceanographic Data Centre (BODC), who provided the tidal data used in this research via their "Sea Level Portal" at <https://www.bodc.ac.uk>. BODC do not accept any liability for the correctness and/or appropriate interpretation of the data or their suitability for any use.

Competing Interests

The author declares that he has no conflict of interest.




Figure 4. Number of ports showing a given rise in the level of the return period between data to 2009 and 2018.

Deleted: ¶ [162]

Formatted: Centered, Indent: Left: 0 cm, Right: 0 cm

Deleted: Colour indicates the change between 2009 and 2018, red: increase, green: decrease.

Deleted: ¶ Table 5: Extreme Sea-level (ESL) at UK coastal locations calculated from data to 2018 using TMAX method. The differences listed are re [164]

Formatted: Indent: Left: 1 cm, Right: 0.7 cm, Tabs: 1 cm, Left + Not at 2.5 cm

Formatted Table

Deleted: Acknowledgments¶ The author is grateful to the British Oceanographic Data Centre (BODC), who provided the tidal data used in this research via their "Sea Level Portal" at [166]

405

References

Formatted: Font color: Auto

1. Batstone, C., Lawless, M., Tawn, J., Horsburgh, K., Blackman, D., McMillan, A., Worth, D., Laeger, S., Hunt, T., 2013. A UK best-practice approach for extreme sea-level analysis along complex topographic coastlines. Ocean Engineering 71, 28–39. <https://doi.org/10.1016/j.oceaneng.2013.02.003>

410

2. Coles, S.G., Dixon, M.J., Likelihood-Based Inference for Extreme Value Models. Extremes 2, 5–23. <https://doi.org/10.1023/A:1009905222644>, 2000.

Deleted: <#>Church, J.A., White, N.J., 2011. Sea-Level Rise from the Late 19th to the Early 21st Century. *Surv Geophys* 32, 585–602. <https://doi.org/10.1007/s10712-011-9119-1>

3. Coles, S., 2001. An Introduction to Statistical Modeling of Extreme Values. Springer Series in Statistics. Springer London, London. <https://doi.org/10.1007/978-1-4471-3675-0>, 2001

Deleted: 2000.

4. Doodson A.T. and Warburg H.D. 1941. Admiralty Manual of Tides. Her Majesty's Stationary Office UK 1941 Reprinted 1980.

Formatted: Bullets and Numbering

415

5. Fuller, W.E., 1914 Flood flows, Trans. A. S. C. E. Paper 1293, LXXVII, 564 (1914)

Formatted: Font color: Auto

6. Gringorten, I.J., 1963. A Plotting Rule for Extreme Probability Paper. Journal of Geophysical Research Vol68 No 3 February 1, 1963

Deleted:

7. Gumbel, E.J., Lieblein, J., 1954 "Statistical Theory of Extreme Values and Some Practical Applications: A Series of Lectures (Vol. 33)". 1954. National Bureau of Standards, US Government Printing Office, Washington

Deleted: ¶

420

8. Harris, R. I. (1996). Gumbel re-visited - a new look at extreme value statistics applied to wind speeds. Journal of Wind Engineering and Industrial Aerodynamics, 59(1), 1–22. [https://doi.org/10.1016/0167-6105\(95\)00029-1](https://doi.org/10.1016/0167-6105(95)00029-1)

Formatted: Font color: Auto

Formatted: Bullets and Numbering

9. Hong, H.P., Li, S.H., Mara, T.G., 2013. Performance of the generalized least-squares method for the Gumbel distribution and its application to annual maximum wind speeds. Journal of Wind Engineering and Industrial Aerodynamics 119, 121–132. <https://doi.org/10.1016/j.jweia.2013.05.012>

425

10. Leadbetter M.R. 1983. Extremes and Local Dependence in Stationary Sequences. Z. Wahrscheinlichkeitstheorie verw. Gebiete 65, 291-306 (1983)

Deleted: <#>IPCC 2021: Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University Press, UK & NY, USA. doi:10.1017/9781009157896¶

Deleted:

11. Middleton J. F. and Thompson K.R. 1986. Return Periods of Extreme Sea Levels from Short Records. Journal of Geophysical Research Vol 91, NO C10, pages 11,707-11,716 October 15, 1986.

Deleted: <#>Kirezci, E., Young, I.R., Ranasinghe, R. et al. Projections of global-scale extreme sea levels and resulting episodic coastal flooding over the 21st Century. *Sci Rep* 10, 11629 (2020). <https://doi.org/10.1038/s41598-020-67736-6>¶

430

12. McInnes, K.L., Macadam, I., Hubbert, G., O'Grady, J., 2013. An assessment of current and future vulnerability to coastal inundation due to sea-level extremes in Victoria, southeast Australia: Current And Future Coastal Inundation In Victoria, Australia. Int. J. Climatol. 33, 33–47. <https://doi.org/10.1002/joc.3405>

Formatted: Font color: Auto

Formatted: Font color: Auto

13. Morrison, F.A., Uncertainty Analysis for Engineers and Scientists: A Practical Guide, 1st ed. Cambridge University Press. <https://doi.org/10.1017/9781108777513>, 2020.

Formatted: Font color: Auto

435

14. Pugh, D.T., Vassie, J.M., Extreme Sea Levels from Tide and Surge Probability. 1978. Coastal Engineering 1978: 911-930

Deleted: 2020.

15. Robson, A., Reed, D., 1999. Flood Estimation Handbook Volume 3. Statistical procedures for flood frequency estimation. ISBN: 978-1-906698-03-4. Institute of Hydrology 1999

- 440 16. Smith, R.L., 1986. Extreme value theory based on the r largest annual events. *Journal of Hydrology* 86, 27–43.
[https://doi.org/10.1016/0022-1694\(86\)90004-1](https://doi.org/10.1016/0022-1694(86)90004-1)
17. Tawn, J.A., 1988. An extreme-value theory model for dependent observations. *Journal of Hydrology* 101, 227-250.
18. Tawn, J.A., Vassie, J.M., 1989. Extreme Sea Levels: The Joint Probabilities Method Revisited and Revised. *Proc. Inst. Civ. Engrs. Part 2*, 87, Sept., 429-442. <https://doi.org/10.1680/iicep.1989.2975>
19. Tawn, J.A., 1992. Estimating Probabilities of Extreme Sea-Levels. *Applied Statistics* 41, 77.
<https://doi.org/10.2307/2347619>
- 445 20. UK Environment Agency 2011. Coastal flood boundary conditions for UK mainland and islands, Project: SC060064/TR2: Design sea levels, Environment Agency UK. ISBN 978-1-84911-212-3, © Environment Agency, February 2011.
21. van Zyl, J. M., & Schall, R. (2012). Parameter Estimation Through Weighted Least-Squares Rank Regression with Specific Reference to the Weibull and Gumbel Distributions. *Communications in Statistics - Simulation and Computation*, 41(9), 1654–1666. <https://doi.org/10.1080/03610918.2011.611315>
- 450 22. Vitousek, S., Barnard, P., Fletcher, C. et al. Doubling of coastal flooding frequency within decades due to sea-level rise. *Sci Rep* 7, 1399 (2017). <https://doi.org/10.1038/s41598-017-01362-7>
23. Williams, J., Horsburgh, K.J., Williams, J.A., Proctor, R.N.F., 2016. Tide and skew surge independence: New insights for flood risk: Skew Surge-Tide Independence. *Geophys. Res. Lett.* 43, 6410–6417.
<https://doi.org/10.1002/2016GL069522>
- 455 24. Zhang A., Wei E., Parker B, Optimal estimation of tidal open boundary conditions using predicted tides and adjoint data assimilation technique. *Continental Shelf Research*, Volume 23, Issues 11&13, 2003, Pages 1055-1070, ISSN 0278-4343. [https://doi.org/10.1016/S0278-4343\(03\)00105-5](https://doi.org/10.1016/S0278-4343(03)00105-5)

Deleted:

Deleted:

Deleted: <#>Tebaldi, C., Ranasinghe, R., Vousdoukas, M., Rasmussen, D.J., Vega-Westhoff, B., Kirezci, E., Kopp, R.E., Sriver, R., Mentaschi, L., 2021. Extreme sea levels at different global warming levels. *Nat. Clim. Chang.* 11, 746–751.
<https://doi.org/10.1038/s41558-021-01127-1>

Deleted: <#>Wahl, T., Haigh, I.D., Woodworth, P.L., Albrecht, F., Dillingh, D., Jensen, J., Nicholls, R.J., Weisse, R., Wöppelmann, G., 2013. Observed mean sea level changes around the North Sea coastline from 1800 to present. *Earth-Science Reviews* 124, 51–67.
<https://doi.org/10.1016/j.earscirev.2013.05.003>

<#>Wahl, T., Haigh, I.D., Nicholls, R.J., Arns, A., Dangendorf, S., Hinkel, J., Slangen, A.B.A., 2017. Understanding extreme sea levels for broad-scale coastal impact and adaptation analysis. *Nat Commun* 8, 16075.
<https://doi.org/10.1038/ncomms16075>

Formatted: Font color: Auto

Page 1: [1] Deleted and inundation	User	11/11/2025 5:44 PM
Page 1: [1] Deleted sited	User	19/12/2025 1:35 AM
Page 1: [1] Deleted Gumbel's	User	11/11/2025 5:50 PM
Page 1: [1] Deleted modifications and new	User	11/11/2025 5:21 PM
Page 1: [1] Deleted methods	User	18/11/2025 6:15 PM
Page 1: [1] Deleted recently	User	16/11/2025 5:28 PM
Page 1: [1] Deleted , often requiring manual intervent	User	20/11/2025 9:02 PM
Page 1: [1] Deleted ion, and is difficult to automate.	User	20/11/2025 9:02 PM
Page 1: [1] Deleted specifically applicable to tides that is	User	11/11/2025 5:42 PM
Page 1: [1] Deleted t	User	11/11/2025 5:23 PM
Page 1: [1] Deleted proposes several modifications to	User	20/11/2025 9:03 PM
Page 1: [1] Deleted Gumbel's	User	11/11/2025 5:52 PM
Page 1: [1] Deleted . The novel technique	User	11/11/2025 5:25 PM
Page 1: [1] Deleted s initial time unit	User	16/11/2025 5:31 PM
Page 1: [1] Deleted is one tidal day,	User	16/11/2025 5:31 PM
Page 1: [1] Deleted rather than the usual	User	11/11/2025 5:28 PM
Page 1: [1] Deleted (User	11/11/2025 5:29 PM
Page 1: [1] Deleted	User	11/11/2025 5:29 PM

). Compared to AMAX

Page 1: [1] Deleted	User	11/11/2025 5:30 PM
---------------------	------	--------------------

,

Page 1: [1] Deleted	User	18/11/2025 6:16 PM
---------------------	------	--------------------

more

Page 1: [1] Deleted	User	11/11/2025 5:30 PM
---------------------	------	--------------------

provides reduced variance in design height,

Page 1: [1] Deleted	User	18/11/2025 6:16 PM
---------------------	------	--------------------

more efficient

Page 1: [1] Deleted	User	12/11/2025 11:51 AM
---------------------	------	---------------------

ly handles

Page 1: [1] Deleted	User	15/12/2025 6:46 PM
---------------------	------	--------------------

are

Page 1: [1] Deleted	User	11/11/2025 5:35 PM
---------------------	------	--------------------

d

Page 1: [1] Deleted	User	16/11/2025 8:29 PM
---------------------	------	--------------------

Page 1: [1] Deleted	User	11/11/2025 5:41 PM
---------------------	------	--------------------

Page 1: [1] Deleted	User	03/12/2025 2:20 PM
---------------------	------	--------------------

35

Page 1: [2] Deleted	User	03/12/2025 2:21 PM
---------------------	------	--------------------

identical

Page 1: [2] Deleted	User	11/11/2025 5:35 PM
---------------------	------	--------------------

, showing broad agreement.

Page 1: [2] Deleted	User	11/11/2025 5:32 PM
---------------------	------	--------------------

is

Page 1: [2] Deleted	User	11/11/2025 5:32 PM
---------------------	------	--------------------

new approach provides a robust mechanism for extreme tide analysis and

Page 1: [2] Deleted	User	20/11/2025 9:05 PM
---------------------	------	--------------------

better informs strategies

Page 1: [2] Deleted	User	06/12/2025 11:56 PM
---------------------	------	---------------------

for

Page 1: [3] Deleted	User	13/11/2025 2:59 PM
---------------------	------	--------------------

, as

Page 1: [3] Deleted	User	13/11/2025 3:21 PM
Coastal tide		
Page 1: [3] Deleted	User	18/11/2025 6:23 PM
gauges		
Page 1: [3] Deleted	User	18/11/2025 6:24 PM
data		
Page 1: [3] Deleted	User	18/11/2025 6:24 PM
s		
Page 1: [3] Deleted	User	18/11/2025 6:24 PM
probabilities		
Page 1: [3] Deleted	User	18/11/2025 6:24 PM
periods		
Page 1: [3] Deleted	User	20/11/2025 9:08 PM
, providing the local records necessary for deriving flood risk parameters		
Page 1: [3] Deleted	User	11/11/2025 6:14 PM
.		
Page 1: [3] Deleted	User	13/11/2025 3:12 PM
often		
Page 1: [3] Deleted	User	17/12/2025 1:35 PM
, and		
Page 2: [4] Deleted	User	17/12/2025 1:36 PM
, (
Page 2: [4] Deleted	User	16/11/2025 5:45 PM
can		
Page 2: [4] Deleted	User	22/12/2025 11:35 AM
accurately		
Page 2: [4] Deleted	User	13/11/2025 3:13 PM
variation of		
Page 2: [4] Deleted	User	13/11/2025 3:13 PM
period with height		
Page 2: [4] Deleted	User	11/11/2025 7:21 PM
inherently		
Page 2: [4] Deleted	User	11/11/2025 9:13 PM
that tidal height noise		
Page 2: [4] Deleted	User	11/11/2025 9:14 PM

Page 2: [4] Deleted **User** **11/11/2025 9:14 PM**
averages out

Page 2: [4] Deleted **User** **13/12/2025 11:23 PM**
flood defence heights, take

Page 2: [4] Deleted **User** **13/12/2025 11:23 PM**
into account the effect of noise

Page 2: [4] Deleted **User** **11/11/2025 7:23 PM**
.

Page 2: [4] Deleted **User** **16/11/2025 8:32 PM**
make this

Page 2: [4] Deleted **User** **16/11/2025 8:32 PM**
ment

Page 2: [4] Deleted **User** **16/11/2025 8:33 PM**
which

Page 2: [5] Deleted **User** **10/12/2025 4:08 PM**
The Gumbel Type I AMAX

Page 2: [5] Deleted **User** **10/12/2025 4:08 PM**
uses only the largest annual extreme events for its extreme value analysis; with

Page 2: [5] Deleted **User** **12/11/2025 4:17 PM**
significant events that rank second or lower not being utilised. To address this apparent inefficient use of available data, several alternative techniques have been developed. One such technique is the "r-largest" method (Smith 1986; Tawn 1988), which involves using a fixed number, r , of the largest annual maxima for analysis. Another approach is the "Peaks over Threshold" (POT) method described by Coles (2001), which utilizes maxima that exceed a certain threshold value to create a new probability distribution.

Page 2: [6] Deleted **User** **18/12/2025 10:41 PM**
(

Page 2: [6] Deleted **User** **17/12/2025 12:53 AM**
)

Page 2: [6] Deleted **User** **12/11/2025 4:19 PM**

Page 2: [6] Deleted **User** **12/11/2025 4:19 PM**
a

Page 2: [6] Deleted andom surge	User	11/11/2025 9:21 PM
Page 2: [6] Deleted	User	17/12/2025 12:20 AM
Page 2: [6] Deleted converting	User	16/12/2025 10:43 PM
Page 2: [6] Deleted JPM output	User	16/12/2025 10:44 PM
Page 2: [6] Deleted requires an understanding of flood duration	User	11/11/2025 8:08 PM
Page 2: [6] Deleted	User	11/11/2025 7:41 PM
Page 2: [6] Deleted and perhaps more critically, research has indicated that the residual and astronomical components are not statistically independent and are not sufficiently uncorrelated (Tawn and Vassie, 1989; Tawn, 1992), making the convolution of these components questionable. The correlation becomes particularly evident during storm surges, as	User	11/11/2025 7:42 PM
Page 2: [6] Deleted , which	User	12/11/2025 4:21 PM
Page 2: [7] Deleted undermines	User	11/11/2025 9:22 PM
Page 2: [7] Deleted utilized	User	12/11/2025 4:21 PM
Page 2: [8] Deleted though	User	11/11/2025 8:14 PM
Page 2: [8] Deleted it has also faced	User	11/11/2025 8:14 PM
Page 2: [9] Deleted	User	20/11/2025 9:10 PM
Page 2: [9] Deleted this	User	16/11/2025 8:34 PM
Page 2: [9] Deleted by using	User	18/11/2025 6:30 PM
Page 2: [9] Deleted	User	18/11/2025 6:31 PM

the

Page 2: [9] Deleted	User	18/11/2025 6:32 PM
maxima		
Page 2: [9] Deleted	User	18/11/2025 6:33 PM
,		
Page 2: [9] Deleted	User	18/11/2025 6:31 PM
rather than the difference between instantaneous height (Batstone et al. 2013).		
Page 3: [10] Deleted	User	12/11/2025 9:27 PM
suitable candidate		
Page 3: [10] Deleted	User	11/11/2025 8:26 PM
However, there are only a limited number of skew extreme high tides, resulting in fewer output points compared to the Joint Probability Method (JPM).		
Page 3: [10] Deleted	User	11/11/2025 8:26 PM
needed		
Page 3: [10] Deleted	User	11/11/2025 8:57 PM
For a complete set of equations, the reader should refer to Batstone (2013). SSJPM was applied to tidal locations around the UK		
Page 3: [10] Deleted	User	11/11/2025 8:27 PM
but as		
Page 3: [10] Deleted	User	11/11/2025 8:58 PM
reported		
Page 3: [11] Formatted	User	11/11/2025 8:58 PM
Font: Italic		
Page 3: [11] Formatted	User	11/11/2025 8:58 PM
Font: Italic		
Page 3: [12] Deleted	User	11/11/2025 8:53 PM
were corrected by manual intervention using the AMAX method and also data values from nearby tidal stations		
Page 3: [12] Deleted	User	14/11/2025 3:28 PM
Page 3: [13] Formatted	User	13/12/2025 11:27 PM
Font: Times New Roman, 10 pt, Font color: Auto		
Page 3: [13] Formatted	User	13/12/2025 11:27 PM
Font: Times New Roman, 10 pt, Font color: Auto		
Page 3: [13] Formatted	User	18/11/2025 10:21 PM
Not Highlight		
Page 3: [13] Formatted	User	18/11/2025 10:21 PM

Not Highlight

Page 3: [13] Formatted	User	18/11/2025 10:21 PM
------------------------	------	---------------------

Not Highlight

Page 3: [13] Formatted	User	18/11/2025 10:21 PM
------------------------	------	---------------------

Not Highlight

Page 3: [14] Deleted	User	12/11/2025 4:25 PM
----------------------	------	--------------------

review highlights

Page 3: [14] Deleted	User	12/11/2025 4:24 PM
----------------------	------	--------------------

the need for

Page 3: [14] Deleted	User	15/11/2025 9:54 PM
----------------------	------	--------------------

method

Page 3: [14] Deleted	User	18/11/2025 10:09 PM
----------------------	------	---------------------

analysing

Page 3: [14] Deleted	User	18/11/2025 10:09 PM
----------------------	------	---------------------

that can consistently

Page 3: [14] Deleted	User	18/11/2025 10:09 PM
----------------------	------	---------------------

deliver design risk as a function of height, without requiring manual intervention.

Page 3: [14] Deleted	User	15/11/2025 10:04 PM
----------------------	------	---------------------

discusses the development of such a method, building on the AMAX approach but utilising more extremes each year, when applicable. The method initially identifies the highest sea level values recorded each tidal day rather than each year, leading to the acronym TMAX. These maxima are sorted and truncated, leaving a set of the most extreme, n values as inputs to the analysis. The paper discusses how a judicious choice of n reduces uncertainty in the resulting output; the flood return period as a function of height.

Page 3: [15] Deleted	User	21/12/2025 11:42 PM
----------------------	------	---------------------

he results of t

Page 3: [15] Deleted	User	22/12/2025 11:46 AM
----------------------	------	---------------------

is

Page 3: [15] Deleted	User	10/12/2025 4:20 PM
----------------------	------	--------------------

TMAX

Page 3: [15] Deleted	User	10/12/2025 4:21 PM
----------------------	------	--------------------

Page 3: [15] Deleted	User	14/11/2025 5:19 PM
----------------------	------	--------------------

demonstrated

Page 3: [15] Deleted	User	21/12/2025 11:42 PM
good agreement with findings from		
Page 3: [15] Deleted	User	07/12/2025 11:43 PM
a		
Page 3: [15] Deleted	User	07/12/2025 11:43 PM
commissioned by the UK Environment Agency 2011		
Page 3: [15] Deleted	User	11/11/2025 9:30 PM
at 35 identical coastal locations. However, u		
Page 3: [16] Deleted	User	07/12/2025 11:44 PM
the UK Environment Agency		
Page 3: [16] Deleted	User	07/12/2025 11:44 PM
study		
Page 3: [16] Deleted	User	15/11/2025 10:10 PM
, running		
Page 3: [16] Deleted	User	11/11/2025 9:24 PM
the same		
Page 3: [16] Deleted	User	11/11/2025 9:27 PM
very tidal		
Page 3: [16] Deleted	User	15/11/2025 1:18 PM
<p>The primary goal of this study is to evaluate the application of the TMAX technique, rather than to provide a conclusive re-evaluation of the UK flood defence heights.</p>		
Page 3: [17] Deleted	User	11/11/2025 9:01 PM
<p>Throughout this text, the term tide refers to the total sea level, while the astronomical tide refers to the deterministic component caused by tidal forces generated by the earth-solar and lunar-earth orbits. The difference between the two is termed the residual, the storm surge, or the random noise component.</p>		
Page 5: [18] Deleted	User	16/11/2025 6:15 PM

3.1 Tidal Day

Most but not all of the examples given by Gumbel use the annual maxima, which later became known as the AMAX method. However, in his original paper, Gumbel also examines some extreme events without using annualized data, such as the breaking point of yarn, and the breakdown voltage of electrical capacitors. He states in his conclusion, "If the number of observed extremes N is not excessive, do not group the

observations." Therefore, although extrema are generally grouped annually before applying Gumbel's method it is not necessarily the case. Annual grouping is thus replaced here with a time unit of one tidal day which, also known as a lunar day, is the time it takes for a specific site on Earth to rotate to the same point under the moon and has a value of 1.034989 solar days. Changing the time unit from annual to tidal days considerably simplifies the identification and selection of tidal maxima which naturally occur in phase with the passing of each tidal day.

Page 5: [19] Deleted	User	12/11/2025 4:48 PM
-----------------------------	-------------	---------------------------

Harris (1996) indicated there were some mathematical advantages in reversing the rank order as compared with Gumbel's original scheme, and this reversal is used here. This is particularly appropriate for this implementation

Page 5: [20] Deleted	User	12/11/2025 6:33 PM
-----------------------------	-------------	---------------------------

but the least are not and would be much more difficult to reliably determine. However, the total number of observations, N , is precisely known, **since it is the source duration in units of tidal days**.

Page 5: [21] Formatted	User	16/11/2025 5:40 PM
-------------------------------	-------------	---------------------------

Font color: Rose, Highlight

Page 5: [22] Deleted	User	12/11/2025 4:57 PM
-----------------------------	-------------	---------------------------

refer to the above equation

Page 5: [23] Deleted	User	12/11/2025 4:57 PM
-----------------------------	-------------	---------------------------

from Gumbel's original paper and substituting for Gumbel's descending rank, i' with an ascending rank, i , where $i = N + 1 - i'$

Page 5: [24] Deleted	User	12/11/2025 5:00 PM
-----------------------------	-------------	---------------------------

Conversion from flood probability F to the reduced variate, y is carried out using the inverse of equation (7),

Page 5: [25] Formatted	User	12/11/2025 6:00 PM
-------------------------------	-------------	---------------------------

Indent: First line: 1.27 cm

Page 5: [26] Formatted	User	12/11/2025 5:53 PM
-------------------------------	-------------	---------------------------

Indent: First line: 1.27 cm

Page 5: [27] Deleted	User	12/11/2025 5:12 PM
-----------------------------	-------------	---------------------------

Remembering that the reduced variate, y relates to flood height x via equation (4), this formula approximates to $y = \log(T)$ since F is very small, confirming Fuller's original claim that maximum flood height varies with the logarithm of time. Since the tidal day has been adopted as the basic time unit, the flood return period is returned in tidal days; it can be converted to solar days by dividing by the length of a tidal day which is 1.034989 days. The reversal of rank order does not affect equation (1) nor

Gringorten's correction equation (5), since it can be shown by making the above substitutions that the formulae remain the same and hence apply to both ascending and descending ranks. The Gringorten formula, rather than equation (1), was used in all of the relevant calculations from hereon.

Page 7: [28] Deleted	User	17/11/2025 11:23 PM
This list may include smaller peaks associated with noise and smaller semi-diurnal tides, but as only the most significant are ultimately used, their existence in the list is entirely irrelevant. Clearly, for a duration of tidal data of many decades, there will be many thousands of valid tidal-daily peak values. However, using all of the tidal-daily peak values on a probability plot is not advisable because of curvature in the distribution at less extreme values. Figure 1 shows this curvature and how it influences a straight line fit.		
Page 7: [29] Formatted	User	16/12/2025 10:55 PM
Font color: Auto, Subscript		
Page 7: [30] Formatted	User	16/12/2025 10:55 PM
Font color: Auto, Subscript		
Page 7: [31] Formatted	User	16/12/2025 10:56 PM
Font color: Auto, Subscript		
Page 7: [32] Formatted	User	13/12/2025 11:17 PM
Font color: Auto, Subscript		
Page 8: [33] Formatted	User	13/12/2025 11:17 PM
Font color: Auto		
Page 8: [33] Formatted	User	13/12/2025 11:17 PM
Font color: Auto, Subscript		
Page 8: [33] Formatted	User	13/12/2025 11:17 PM
Font color: Auto		
Page 8: [33] Formatted	User	13/12/2025 11:17 PM
Font color: Auto		
Page 8: [33] Formatted	User	13/12/2025 11:17 PM
Font color: Auto, Subscript		
Page 8: [34] Formatted	User	13/12/2025 11:17 PM
Font color: Auto		
Page 8: [34] Formatted	User	13/12/2025 11:17 PM
Font color: Auto		
Page 8: [35] Formatted	User	13/12/2025 11:17 PM

Font color: Auto

Page 8: [35] Formatted	User	13/12/2025 11:17 PM
-------------------------------	-------------	----------------------------

Font color: Auto

Page 8: [35] Formatted	User	13/12/2025 11:17 PM
-------------------------------	-------------	----------------------------

Font color: Auto

Page 8: [36] Formatted	User	18/12/2025 10:27 PM
-------------------------------	-------------	----------------------------

Font color: Auto, Subscript

Page 8: [36] Formatted	User	13/12/2025 11:17 PM
-------------------------------	-------------	----------------------------

Font color: Auto

Page 8: [36] Formatted	User	13/12/2025 11:17 PM
-------------------------------	-------------	----------------------------

Font color: Auto

Page 8: [36] Formatted	User	13/12/2025 11:17 PM
-------------------------------	-------------	----------------------------

Font color: Auto

Page 8: [37] Formatted	User	13/12/2025 11:17 PM
-------------------------------	-------------	----------------------------

Font color: Auto

Page 8: [37] Formatted	User	13/12/2025 11:17 PM
-------------------------------	-------------	----------------------------

Font color: Auto

Page 8: [37] Formatted	User	13/12/2025 11:17 PM
-------------------------------	-------------	----------------------------

Font color: Auto

Page 8: [37] Formatted	User	17/12/2025 12:21 AM
-------------------------------	-------------	----------------------------

Font color: Auto, Subscript

Page 8: [37] Formatted	User	13/12/2025 11:17 PM
-------------------------------	-------------	----------------------------

Font color: Auto

Page 8: [38] Formatted	User	18/12/2025 10:27 PM
-------------------------------	-------------	----------------------------

Font color: Auto, Subscript

Page 8: [38] Formatted	User	13/12/2025 11:17 PM
-------------------------------	-------------	----------------------------

Font color: Auto

Page 8: [38] Formatted	User	13/12/2025 11:17 PM
-------------------------------	-------------	----------------------------

Font color: Auto

Page 8: [39] Formatted	User	14/12/2025 1:51 AM
-------------------------------	-------------	---------------------------

Font color: Auto

Page 8: [39] Formatted	User	14/12/2025 1:51 AM
-------------------------------	-------------	---------------------------

Font color: Auto

Page 8: [39] Formatted	User	14/12/2025 1:51 AM
-------------------------------	-------------	---------------------------

Font color: Auto

Page 8: [39] Formatted	User	14/12/2025 1:51 AM
-------------------------------	-------------	---------------------------

Font color: Auto

Page 8: [39] Formatted	User	14/12/2025 1:51 AM
-------------------------------	-------------	---------------------------

Font color: Auto

Page 8: [39] Formatted	User	14/12/2025 1:51 AM
-------------------------------	-------------	---------------------------

Font color: Auto

Page 8: [39] Formatted	User	14/12/2025 1:51 AM
-------------------------------	-------------	---------------------------

Font color: Auto

Page 8: [39] Formatted	User	14/12/2025 1:51 AM
-------------------------------	-------------	---------------------------

Font color: Auto

Page 8: [39] Formatted	User	14/12/2025 1:51 AM
Font color: Auto		
Page 8: [39] Formatted	User	14/12/2025 1:51 AM
Font color: Auto		
Page 8: [39] Formatted	User	14/12/2025 1:51 AM
Font color: Auto		
Page 8: [39] Formatted	User	14/12/2025 1:51 AM
Font color: Auto		
Page 8: [39] Formatted	User	14/12/2025 1:51 AM
Font color: Auto		
Page 8: [40] Deleted	User	12/11/2025 8:46 PM

In harmonic tidal analysis, shallow water coefficients are necessary to take account of non-linearity in the ocean response occurring at lower tides (Doodson and Warburg 1941).




Figure 1. Probability extreme tide plot showing its curvature in a 54-year record. Setting a threshold of 50 greatest extremes tides per year (2439 total), the LSQ intercept is 11.95m ± 0.02m R2= 0.9776 (black). At 5 greatest extreme tides per year (242 points) the straighter red section gives an improved LSQ intercept of 11.6m (red) ± 0.009m R2=0.9939. Location Heysham UK See 3.3 & 4.1 & Fig 3.

Similarly, it is suggested that the curvature of the probability plot is probably related to shallow water effects during the neap of a spring-neap cycle. Rather than attempting to fit the statistics of all high tides, we use a fit of only the n largest

values, since these fall, more or less, in a straight line as shown in Fig. 1, where n is calculated from an average number per year, n_A multiplied by the tide gauge record duration in years n_y , i.e. $n = n_A \times n_y$. The number of points used for the straight-line fit is much smaller than the number of high tides, N . Nevertheless, N must be used in equations (1) and (5) since it determines the scaling of the points plotted. The optimization of n_A is discussed in Section 4.1. Note that this proposed method is quite different from the Points Over Threshold (POT) method (Coles 2001), or R-Largest (Smith 1986). Here in the TMAX method like in the original Gumble AMAX method, ranking alone determines return period.

Page 8: [41] Formatted Formatted	User	11/12/2025 12:07 AM
Page 8: [41] Formatted Highlight	User	11/12/2025 12:07 AM
Page 8: [42] Deleted T	User	04/12/2025 5:35 PM
Page 8: [42] Deleted method	User	11/12/2025 12:07 AM
Page 8: [42] Deleted . Other	User	14/11/2025 5:25 PM
Page 8: [42] Deleted exist	User	11/12/2025 12:12 AM
Page 8: [42] Deleted	User	17/12/2025 12:21 AM
Page 8: [42] Deleted Although it is stated that LSQ is less accurate than these other methods, LSQ uses simple closed-loop expressions to provide slope, intercept, and mean variance, providing a reasonable estimate.	User	14/11/2025 5:27 PM
Page 8: [42] Deleted using	User	14/11/2025 5:27 PM
Page 8: [42] Deleted of the method	User	16/11/2025 9:01 PM
Page 8: [42] Deleted	User	17/12/2025 12:21 AM
Page 8: [43] Formatted Subscript	User	09/12/2025 10:30 PM

Page 8: [43] Formatted	User	09/12/2025 10:30 PM
Subscript		
Page 8: [44] Formatted	User	09/12/2025 10:28 PM
Subscript		
Page 8: [44] Formatted	User	09/12/2025 10:28 PM
Font color: Auto		
Page 8: [45] Formatted	User	09/12/2025 10:28 PM
Subscript		
Page 8: [45] Formatted	User	09/12/2025 10:28 PM
Subscript		
Page 9: [46] Formatted	User	09/12/2025 10:28 PM
Subscript		
Page 9: [46] Formatted	User	09/12/2025 10:28 PM
Not Superscript/ Subscript		
Page 9: [46] Formatted	User	09/12/2025 10:28 PM
Font color: Auto		
Page 9: [46] Formatted	User	09/12/2025 10:28 PM
Font color: Auto		
Page 9: [47] Formatted	User	22/12/2025 11:59 AM
Font: (Default) Times New Roman, Not Superscript/ Subscript		
Page 9: [47] Formatted	User	22/12/2025 11:59 AM
Font: (Default) Times New Roman		
Page 9: [47] Formatted	User	22/12/2025 11:59 AM
Font: (Default) Times New Roman		
Page 9: [47] Formatted	User	22/12/2025 11:59 AM
Font: (Default) Times New Roman		
Page 9: [47] Formatted	User	22/12/2025 11:59 AM
Font: (Default) Times New Roman, Not Superscript/ Subscript		
Page 9: [47] Formatted	User	22/12/2025 11:59 AM
Font: (Default) Times New Roman		
Page 9: [47] Formatted	User	22/12/2025 11:59 AM
Font: (Default) Times New Roman, Not Superscript/ Subscript		
Page 9: [47] Formatted	User	22/12/2025 11:59 AM
Font: (Default) Times New Roman		
Page 9: [47] Formatted	User	22/12/2025 11:59 AM
Font: (Default) Times New Roman, Not Superscript/ Subscript		
Page 9: [47] Formatted	User	22/12/2025 11:59 AM
Font: (Default) Times New Roman		
Page 9: [48] Formatted	User	09/12/2025 10:28 PM
Subscript		
Page 9: [48] Formatted	User	09/12/2025 10:28 PM

Subscript

Page 9: [48] Formatted	User	09/12/2025 10:28 PM
-------------------------------	-------------	----------------------------

Font: Symbol, Subscript

Page 9: [48] Formatted	User	09/12/2025 10:28 PM
-------------------------------	-------------	----------------------------

Subscript

Page 9: [48] Formatted	User	22/12/2025 12:42 AM
-------------------------------	-------------	----------------------------

Font: (Default) Times New Roman

Page 9: [48] Formatted	User	22/12/2025 12:43 AM
-------------------------------	-------------	----------------------------

Font: (Default) Calibri

Page 9: [48] Formatted	User	18/12/2025 9:16 PM
-------------------------------	-------------	---------------------------

Not Superscript/ Subscript

Page 9: [49] Formatted	User	22/12/2025 11:58 AM
-------------------------------	-------------	----------------------------

Font: (Default) Times New Roman

Page 9: [49] Formatted	User	22/12/2025 11:58 AM
-------------------------------	-------------	----------------------------

Not Superscript/ Subscript

Page 9: [49] Formatted	User	22/12/2025 11:58 AM
-------------------------------	-------------	----------------------------

Font: (Default) Times New Roman, Not Superscript/ Subscript

Page 9: [49] Formatted	User	22/12/2025 11:58 AM
-------------------------------	-------------	----------------------------

Font: (Default) Times New Roman

Page 9: [50] Deleted	User	09/12/2025 10:40 PM
-----------------------------	-------------	----------------------------

The

Page 9: [50] Deleted	User	09/12/2025 10:40 PM
-----------------------------	-------------	----------------------------

variance of the new predicted mean σ_p is always greater than the variance of the mean

σ_m ; practically this is

Page 9: [50] Deleted	User	09/12/2025 10:24 PM
-----------------------------	-------------	----------------------------

because

Page 9: [50] Deleted	User	09/12/2025 10:27 PM
-----------------------------	-------------	----------------------------

, whereas the variance in the intersection is subject to variation in the regression line

parameters, the latter tending towards zero as the number of data points increases.

Page 9: [50] Deleted	User	18/11/2025 10:45 PM
-----------------------------	-------------	----------------------------

Page 9: [51] Deleted	User	16/11/2025 9:07 PM
-----------------------------	-------------	---------------------------

data

Page 9: [51] Deleted	User	19/11/2025 11:46 AM
-----------------------------	-------------	----------------------------

s

Page 9: [51] Deleted	User	03/12/2025 2:31 PM
-----------------------------	-------------	---------------------------

data

Page 9: [51] Deleted	User	15/11/2025 1:07 PM
-----------------------------	-------------	---------------------------

taken into account

Page 9: [51] Deleted	User	16/11/2025 9:01 PM
-----------------------------	-------------	---------------------------

tidal

Page 9: [51] Deleted	User	18/11/2025 10:42 PM
handle		
Page 9: [51] Deleted	User	18/11/2025 10:43 PM
the		
Page 9: [51] Deleted	User	18/11/2025 10:43 PM
below a given threshold		
Page 9: [51] Deleted	User	19/12/2025 10:50 PM
This is a considerable advantage of the TMAX over the AMAX		
Page 9: [52] Formatted	User	12/11/2025 11:04 PM
Highlight		
Page 9: [52] Formatted	User	28/11/2025 12:40 AM
Not Highlight		
Page 11: [53] Deleted	User	16/12/2025 11:59 PM

4.1 Optimising the average number of peaks per year.

As discussed above, larger values of n_A cause a higher proportion of extreme peaks under investigation to originate from smaller astronomical tides (neap tides) whose probability distribution may differ. Therefore, although larger sample values are generally associated with a decrease in variance, in this case the opposite may be true.




Figure 2. The mean-variance σ_m and of the predicted new point variance σ_p plotted against the number of extrema per year n_A for the TMAX method as practically determined from the tidal data.

Figure 2 shows how in practice the variance of the mean, σ_m , and variance of a predicted new point, σ_p varies with n_A , the average number of extremes selected per

year, using the TMAX method. The values were determined from the tidal data using equations (10) & (11). The variance of the mean, σ_m decreases as n_A increases; however, σ_p shows a minimum with a value of n_A of around five extremes per year. This concurs with the value of five per year found by Robson and Reed 1999 for extreme river flow studies and used by Smith 1986, although this may be coincidental. Nevertheless, since we obtain a minimum variance at the value of $n_A = 5$ this was adopted for all subsequent results.

Page 12: [54] Formatted	User	18/12/2025 9:04 PM
Caption, Centered, Keep with next		
Page 12: [55] Formatted	User	22/12/2025 2:26 PM
Font: 8 pt		
Page 12: [56] Change	User	22/12/2025 9:55 PM
Formatted Table		
Page 12: [57] Formatted	User	22/12/2025 2:26 PM
Font: 8 pt		
Page 12: [58] Formatted	User	22/12/2025 2:31 PM
Justified		
Page 12: [59] Formatted	User	22/12/2025 9:55 PM
Centered		
Page 12: [60] Formatted	User	22/12/2025 2:26 PM
Font: 8 pt		
Page 12: [61] Formatted	User	22/12/2025 2:31 PM
Right		
Page 12: [62] Formatted	User	22/12/2025 2:31 PM
Justified		
Page 12: [62] Formatted	User	22/12/2025 2:31 PM
Right		
Page 12: [62] Formatted	User	22/12/2025 2:55 PM
Centered		
Page 12: [63] Formatted	User	22/12/2025 2:26 PM
Font: 8 pt		
Page 12: [64] Formatted	User	22/12/2025 2:31 PM
Justified		
Page 12: [65] Formatted	User	22/12/2025 2:26 PM
Font: 8 pt		
Page 12: [66] Formatted	User	22/12/2025 2:31 PM
Justified		
Page 12: [67] Formatted	User	22/12/2025 2:26 PM
Font: 8 pt		
Page 12: [68] Formatted	User	22/12/2025 2:31 PM
Justified		
Page 12: [69] Formatted	User	22/12/2025 2:26 PM

Font: 8 pt

Page 12: [70] Formatted	User	22/12/2025 2:31 PM
--------------------------------	-------------	---------------------------

Justified

Page 12: [71] Formatted	User	22/12/2025 2:26 PM
--------------------------------	-------------	---------------------------

Font: 8 pt

Page 12: [72] Formatted	User	22/12/2025 2:31 PM
--------------------------------	-------------	---------------------------

Justified

Page 12: [73] Formatted	User	22/12/2025 2:26 PM
--------------------------------	-------------	---------------------------

Font: 8 pt

Page 12: [74] Formatted	User	22/12/2025 2:31 PM
--------------------------------	-------------	---------------------------

Justified

Page 12: [75] Formatted	User	22/12/2025 2:26 PM
--------------------------------	-------------	---------------------------

Font: 8 pt

Page 12: [76] Formatted	User	22/12/2025 2:31 PM
--------------------------------	-------------	---------------------------

Justified

Page 12: [77] Formatted	User	22/12/2025 2:26 PM
--------------------------------	-------------	---------------------------

Font: 8 pt

Page 12: [78] Formatted	User	22/12/2025 2:31 PM
--------------------------------	-------------	---------------------------

Justified

Page 12: [79] Formatted	User	22/12/2025 2:26 PM
--------------------------------	-------------	---------------------------

Font: 8 pt

Page 12: [80] Formatted	User	22/12/2025 2:31 PM
--------------------------------	-------------	---------------------------

Justified

Page 12: [81] Formatted	User	22/12/2025 2:26 PM
--------------------------------	-------------	---------------------------

Font: 8 pt

Page 12: [82] Formatted	User	22/12/2025 2:31 PM
--------------------------------	-------------	---------------------------

Justified

Page 12: [83] Formatted	User	22/12/2025 2:26 PM
--------------------------------	-------------	---------------------------

Font: 8 pt

Page 12: [84] Formatted	User	22/12/2025 2:31 PM
--------------------------------	-------------	---------------------------

Justified

Page 12: [85] Formatted	User	22/12/2025 2:26 PM
--------------------------------	-------------	---------------------------

Font: 8 pt

Page 12: [86] Formatted	User	22/12/2025 2:31 PM
--------------------------------	-------------	---------------------------

Justified

Page 12: [87] Formatted	User	22/12/2025 2:26 PM
--------------------------------	-------------	---------------------------

Font: 8 pt

Page 12: [88] Formatted	User	22/12/2025 2:31 PM
--------------------------------	-------------	---------------------------

Justified

Page 12: [89] Formatted	User	22/12/2025 2:26 PM
--------------------------------	-------------	---------------------------

Font: 8 pt

Page 12: [90] Formatted	User	22/12/2025 2:31 PM
--------------------------------	-------------	---------------------------


Justified

Page 12: [91] Formatted	User	22/12/2025 2:26 PM
--------------------------------	-------------	---------------------------

Font: 8 pt

Page 12: [92] Formatted Justified	User	22/12/2025 2:31 PM
Page 12: [93] Formatted Font: 8 pt	User	22/12/2025 2:26 PM
Page 12: [94] Formatted Justified	User	22/12/2025 2:31 PM
Page 12: [95] Formatted Font: 8 pt	User	22/12/2025 2:26 PM
Page 12: [96] Formatted Justified	User	22/12/2025 2:31 PM
Page 12: [97] Formatted Font: 8 pt	User	22/12/2025 2:26 PM
Page 12: [98] Formatted Justified	User	22/12/2025 2:31 PM
Page 12: [99] Formatted Font: 8 pt	User	22/12/2025 2:26 PM
Page 12: [100] Formatted Justified	User	22/12/2025 2:31 PM
Page 12: [101] Formatted Font: 8 pt	User	22/12/2025 2:26 PM
Page 12: [102] Formatted Justified	User	22/12/2025 2:31 PM
Page 12: [103] Formatted Font: 8 pt	User	22/12/2025 2:26 PM
Page 12: [104] Formatted Justified	User	22/12/2025 2:31 PM
Page 12: [105] Formatted Font: 8 pt	User	22/12/2025 2:26 PM
Page 12: [106] Formatted Justified	User	22/12/2025 2:31 PM
Page 12: [107] Formatted Font: 8 pt	User	22/12/2025 2:26 PM
Page 12: [108] Formatted Justified	User	22/12/2025 2:31 PM
Page 13: [109] Deleted	User	04/12/2025 11:10 PM

Table 1: AMAX and TMAX methods. Quality of Fit of Regression Line



Return Period (yr)	Variance of Mean σ_M (m)				Variance of Predicted σ_P (m)			
	20	100	200	1000	20	100	200	1000
AMAX to 2009	0.09	0.10	0.11	0.13	0.39	0.39	0.39	0.40
TMAX to 2009	0.02	0.03	0.03	0.03	0.24	0.24	0.24	0.24
AMAX to 2018	0.08	0.09	0.09	0.11	0.41	0.41	0.41	0.41
TMAX to 2018	0.02	0.02	0.02	0.03	0.26	0.26	0.26	0.26

4.2 Determination of Extreme Sea Levels

Figure 3 shows a typical extreme tide probability plot for a single location, Heysham. The corresponding top four plotted points/heights are also listed in Table 2.

Table 2. Tabulated data for the TMAX method at Heysham, UK. Input duration span 19844 days, with valid data 17834 days i.e. 17232 tidal days.

Rank	Date	Height(H_0) (m)	$F(H_t > H_0)$ * per tidal day	Return Period [†] years
1	01/02/2002 13:45:00	11.36482	3.2E-05	87.20
2	10/02/1997 13:15:00	11.30877	9.1E-05	31.30
3	30/03/2006 11:45:00	11.25951	1.5E-04	19.07
4	02/01/2018 23:30:00	11.18499	2.1E-04	13.72

* Calculated using Eq. 5 with N=17232 samples.

† Calculated using Eq. 6 with one tidal day=1.034989 solar days.

All of the results were derived directly from the tide gauge data and have not been re-processed in any way to enforce the agreement with any other data set or with neighbouring local values. The analysis was carried out for the entire data set consisting of 35 ports, once with the tidal records truncated to 1 Jan 2009 for comparison with the results of Batstone's analysis, and once using the complete data set to 1 May 2018. Each data set was analysed using the AMAX and TMAX methods. For compatibility with Batstone, and in order to comply with the requirement for stationarity, all data was sea-level rise de-trended, using the constant value adopted by Batstone of 2mm per year. The de-trending was arranged to apply a zero correction on 1 January 2008, to facilitate comparison with the results from Batstone who used the same date origin. The analysis used Gringorten's correction throughout for compatibility with Batstone's research.




Figure 3. Extreme value probability plot of 242 highest tides at Heysham over a 54-year duration, showing regression line. Height is in meters relative to Admiralty Chart Datum (CD). A 200-year flood return period gives 11.6m CD, 6.7m OD. (OD-CD Heysham = 4.9m). See Table 5.

Table 3 compares the AMAX and TMAX results with those of Batstone. The mean differences for data to 2009 using either method were generally in the centimetre region. In either case, the standard deviation was around 0.1m rising to approximately 0.24m for the 1000-year return level and was somewhat higher for the AMAX than for TMAX. Figure 4 shows the frequency distribution of the differences between TMAX results derived from data to 2009 and data to 2018 across the 35 UK ports, showing a rise in level for each return period T.

Table 3. AMAX and TMAX methods: Comparison with Batstone 2013.

Return Period (yr)	Mean Difference (m)				Standard Deviation (m)			
	20	100	200	1000	20	100	200	1000
AMAX to 2009	0.00	0.03	0.04	0.06	0.11	0.16	0.19	0.27
TMAX to 2009	-0.01	-0.03	-0.04	-0.07	0.09	0.13	0.15	0.22
AMAX- to 2018	0.02	0.06	0.07	0.09	0.10	0.14	0.16	0.22
TMAX to 2018	0.02	0.01	0.00	-0.02	0.09	0.12	0.14	0.20

Page 14: [110] Formatted **User** **17/12/2025 1:13 AM**
 Font: Bold

Page 14: [111] Formatted **User** **17/12/2025 1:13 AM**
 Left

Page 14: [112] Formatted **User** **22/12/2025 10:54 PM**
 Font: (Default) Times New Roman, Font color: Black

Page 14: [113] Formatted	User	22/12/2025 10:54 PM
Font color: Black		
Page 14: [114] Formatted	User	12/12/2025 12:31 PM
Font color: Auto		
Page 14: [115] Formatted	User	17/12/2025 12:10 AM
Font: (Default) Times New Roman		
Page 14: [116] Formatted	User	17/12/2025 12:10 AM
Font: (Default) Times New Roman		
Page 14: [117] Formatted	User	17/12/2025 12:10 AM
Font: (Default) Times New Roman		
Page 14: [118] Formatted	User	17/12/2025 12:10 AM
Font: (Default) Times New Roman		
Page 14: [119] Formatted	User	17/12/2025 12:10 AM
Font: (Default) Times New Roman		
Page 14: [120] Formatted	User	17/12/2025 12:09 AM
Font: (Default) Times New Roman, Bold		
Page 14: [121] Formatted	User	17/12/2025 12:17 PM
Left		
Page 14: [122] Formatted	User	17/12/2025 12:17 PM
Left		
Page 14: [123] Formatted	User	17/12/2025 12:18 PM
Font: (Default) Times New Roman		
Page 14: [124] Formatted	User	17/12/2025 12:09 AM
Font: (Default) Times New Roman		
Page 14: [125] Formatted	User	17/12/2025 12:09 AM
Font: (Default) Times New Roman		
Page 14: [126] Formatted	User	17/12/2025 12:09 AM
Font: (Default) Times New Roman		
Page 14: [127] Formatted	User	17/12/2025 12:09 AM
Font: (Default) Times New Roman		
Page 14: [128] Formatted	User	17/12/2025 12:17 PM
Left		
Page 14: [129] Formatted	User	17/12/2025 12:18 PM
Font: (Default) Times New Roman		
Page 14: [130] Formatted	User	17/12/2025 12:09 AM
Font: (Default) Times New Roman		
Page 14: [131] Formatted	User	17/12/2025 12:09 AM
Font: (Default) Times New Roman		
Page 14: [132] Formatted	User	17/12/2025 12:09 AM
Font: (Default) Times New Roman		
Page 14: [133] Formatted	User	17/12/2025 12:09 AM
Font: (Default) Times New Roman		
Page 14: [134] Formatted	User	17/12/2025 12:17 PM
Left		
Page 14: [135] Formatted	User	17/12/2025 12:18 PM

Font: (Default) Times New Roman

Page 14: [136] Formatted	User	17/12/2025 12:09 AM
---------------------------------	-------------	----------------------------

Font: (Default) Times New Roman

Page 14: [137] Formatted	User	17/12/2025 12:09 AM
---------------------------------	-------------	----------------------------

Font: (Default) Times New Roman

Page 14: [138] Formatted	User	17/12/2025 12:09 AM
---------------------------------	-------------	----------------------------

Font: (Default) Times New Roman

Page 14: [139] Formatted	User	17/12/2025 12:09 AM
---------------------------------	-------------	----------------------------

Font: (Default) Times New Roman

Page 14: [140] Formatted	User	17/12/2025 12:18 PM
---------------------------------	-------------	----------------------------

Font: (Default) Times New Roman

Page 14: [141] Formatted	User	17/12/2025 12:09 AM
---------------------------------	-------------	----------------------------

Font: (Default) Times New Roman

Page 14: [142] Formatted	User	17/12/2025 12:09 AM
---------------------------------	-------------	----------------------------

Font: (Default) Times New Roman

Page 14: [143] Formatted	User	17/12/2025 12:09 AM
---------------------------------	-------------	----------------------------

Font: (Default) Times New Roman

Page 14: [144] Formatted	User	17/12/2025 12:09 AM
---------------------------------	-------------	----------------------------

Font: (Default) Times New Roman

Page 14: [145] Formatted	User	17/12/2025 12:17 PM
---------------------------------	-------------	----------------------------

Left

Page 14: [146] Formatted	User	15/12/2025 6:40 PM
---------------------------------	-------------	---------------------------

Font color: Auto

Page 14: [147] Formatted	User	15/12/2025 6:40 PM
---------------------------------	-------------	---------------------------

Font color: Auto

Page 14: [148] Formatted	User	22/12/2025 12:06 PM
---------------------------------	-------------	----------------------------

Font: Symbol

Page 14: [149] Formatted	User	22/12/2025 12:46 AM
---------------------------------	-------------	----------------------------

Font color: Red

Page 15: [150] Formatted	User	17/12/2025 12:09 AM
---------------------------------	-------------	----------------------------

Font: 9 pt, Bold

Page 15: [150] Formatted	User	17/12/2025 12:09 AM
---------------------------------	-------------	----------------------------

Font: (Default) Times New Roman, Bold

Page 15: [150] Formatted	User	17/12/2025 12:09 AM
---------------------------------	-------------	----------------------------

Font: 9 pt, Bold

Page 15: [150] Formatted	User	17/12/2025 12:09 AM
---------------------------------	-------------	----------------------------

Font: (Default) Times New Roman, Bold

Page 15: [150] Formatted	User	17/12/2025 12:09 AM
---------------------------------	-------------	----------------------------

Font: 9 pt, Bold

Page 15: [150] Formatted	User	17/12/2025 12:09 AM
---------------------------------	-------------	----------------------------

Font: (Default) Times New Roman, Bold

Page 15: [150] Formatted	User	17/12/2025 12:09 AM
---------------------------------	-------------	----------------------------

Font: 9 pt, Bold

Page 15: [150] Formatted	User	17/12/2025 12:09 AM
---------------------------------	-------------	----------------------------

Font: (Default) Times New Roman, Bold

Page 15: [150] Formatted	User	17/12/2025 12:09 AM
Font: 9 pt, Bold		
Page 15: [151] Formatted	User	17/12/2025 12:09 AM
Font: (Default) Times New Roman		
Page 15: [151] Formatted	User	17/12/2025 12:09 AM
Font: (Default) Times New Roman, Not Bold		
Page 15: [152] Formatted	User	17/12/2025 12:09 AM
Font: 9 pt, Font color: Auto		
Page 15: [152] Formatted	User	17/12/2025 12:09 AM
Font: (Default) Times New Roman, 9 pt		
Page 15: [152] Formatted	User	17/12/2025 12:09 AM
Font: 9 pt, Font color: Auto		
Page 15: [152] Formatted	User	17/12/2025 12:09 AM
Font: (Default) Times New Roman, 9 pt		
Page 15: [152] Formatted	User	17/12/2025 12:09 AM
Font: 9 pt, Font color: Auto		
Page 15: [152] Formatted	User	17/12/2025 12:09 AM
Font: (Default) Times New Roman, 9 pt		
Page 15: [152] Formatted	User	17/12/2025 12:09 AM
Font: 9 pt, Font color: Auto		
Page 15: [152] Formatted	User	17/12/2025 12:09 AM
Font: (Default) Times New Roman, 9 pt		
Page 15: [152] Formatted	User	17/12/2025 12:09 AM
Font: 9 pt, Font color: Auto		
Page 15: [153] Formatted	User	17/12/2025 12:09 AM
Font: (Default) Times New Roman, Not Bold		
Page 15: [153] Formatted	User	17/12/2025 12:09 AM
Font: (Default) Times New Roman		
Page 15: [153] Formatted	User	17/12/2025 12:09 AM
Font: (Default) Times New Roman, 9 pt		
Page 15: [154] Formatted	User	17/12/2025 12:09 AM
Font: (Default) Times New Roman		
Page 15: [154] Formatted	User	17/12/2025 12:09 AM
Font: (Default) Times New Roman, Not Bold		
Page 15: [155] Formatted	User	17/12/2025 12:09 AM
Font: 9 pt		
Page 15: [155] Formatted	User	17/12/2025 12:09 AM
Font: (Default) Times New Roman, 9 pt		
Page 15: [155] Formatted	User	17/12/2025 12:09 AM
Font: 9 pt		
Page 15: [155] Formatted	User	17/12/2025 12:09 AM
Font: (Default) Times New Roman, 9 pt		
Page 15: [155] Formatted	User	17/12/2025 12:09 AM
Font: 9 pt		
Page 15: [155] Formatted	User	17/12/2025 12:09 AM

Font: (Default) Times New Roman, 9 pt

Page 15: [155] Formatted	User	17/12/2025 12:09 AM
---------------------------------	-------------	----------------------------

Font: 9 pt

Page 15: [155] Formatted	User	17/12/2025 12:09 AM
---------------------------------	-------------	----------------------------

Font: (Default) Times New Roman, 9 pt

Page 15: [155] Formatted	User	17/12/2025 12:09 AM
---------------------------------	-------------	----------------------------

Font: 9 pt

Page 15: [156] Formatted	User	17/12/2025 12:09 AM
---------------------------------	-------------	----------------------------

Font: (Default) Times New Roman

Page 15: [156] Formatted	User	17/12/2025 12:09 AM
---------------------------------	-------------	----------------------------

Font: (Default) Times New Roman

Page 15: [156] Formatted	User	17/12/2025 12:09 AM
---------------------------------	-------------	----------------------------

Font: (Default) Times New Roman, 9 pt

Page 15: [157] Formatted	User	17/12/2025 12:09 AM
---------------------------------	-------------	----------------------------

Font: (Default) Times New Roman

Page 15: [157] Formatted	User	17/12/2025 12:09 AM
---------------------------------	-------------	----------------------------

Font: (Default) Times New Roman, Not Bold

Page 15: [158] Formatted	User	17/12/2025 12:09 AM
---------------------------------	-------------	----------------------------

Font: 9 pt, Font color: Auto

Page 15: [158] Formatted	User	17/12/2025 12:09 AM
---------------------------------	-------------	----------------------------

Font: (Default) Times New Roman, 9 pt

Page 15: [158] Formatted	User	17/12/2025 12:09 AM
---------------------------------	-------------	----------------------------

Font: 9 pt, Font color: Auto

Page 15: [158] Formatted	User	17/12/2025 12:09 AM
---------------------------------	-------------	----------------------------

Font: (Default) Times New Roman, 9 pt

Page 15: [158] Formatted	User	17/12/2025 12:09 AM
---------------------------------	-------------	----------------------------

Font: 9 pt, Font color: Auto

Page 15: [158] Formatted	User	17/12/2025 12:09 AM
---------------------------------	-------------	----------------------------

Font: (Default) Times New Roman, 9 pt

Page 15: [158] Formatted	User	17/12/2025 12:09 AM
---------------------------------	-------------	----------------------------

Font: 9 pt, Font color: Auto

Page 15: [158] Formatted	User	17/12/2025 12:09 AM
---------------------------------	-------------	----------------------------

Font: (Default) Times New Roman, 9 pt

Page 15: [158] Formatted	User	17/12/2025 12:09 AM
---------------------------------	-------------	----------------------------

Font: 9 pt, Font color: Auto

Page 15: [159] Formatted	User	17/12/2025 12:09 AM
---------------------------------	-------------	----------------------------

Font: (Default) Times New Roman

Page 15: [159] Formatted	User	17/12/2025 12:09 AM
---------------------------------	-------------	----------------------------

Font: (Default) Times New Roman, 9 pt

Page 15: [160] Formatted	User	17/12/2025 12:09 AM
---------------------------------	-------------	----------------------------

Font: (Default) Times New Roman

Page 15: [160] Formatted	User	17/12/2025 12:09 AM
---------------------------------	-------------	----------------------------

Font: (Default) Times New Roman, Not Bold

Page 15: [160] Formatted	User	17/12/2025 12:09 AM
---------------------------------	-------------	----------------------------

Font: (Default) Times New Roman, 9 pt

Page 15: [161] Formatted	User	17/12/2025 12:09 AM
Font: (Default) Times New Roman		
Page 15: [161] Formatted	User	17/12/2025 12:09 AM
Font: (Default) Times New Roman, Not Bold		
Page 15: [161] Formatted	User	17/12/2025 12:09 AM
Font: (Default) Times New Roman, Not Bold		
Page 15: [161] Formatted	User	17/12/2025 12:09 AM
Font: (Default) Times New Roman, Not Bold		
Page 15: [161] Formatted	User	17/12/2025 12:09 AM
Font: (Default) Times New Roman, 9 pt		
Page 17: [162] Deleted	User	28/11/2025 11:39 PM




Figure 4. Number of ports showing a given rise in the level of the return period between data to 2009 and data to 2018 as analysed across 35 UK ports. Table 4 indicates an average rate of rise over the 9 years of the extreme 20-year sea level of $3.4 \times 10^{-3} \text{ yr}^{-1}$ and for the 1000-year level of $6.1 \times 10^{-3} \text{ yr}^{-1}$. This compares with a rate of rise calculated for the whole dataset of $1.85 \times 10^{-3} \text{ yr}^{-1}$. When the increase is analysed by the paired-sample left-tailed t-test it gives P values of 0.005 to 0.006 indicating the increase in level is significant.

Table 4. Rise in level averaged across 35 UK ports between TMAX (2009) and TMAX (2018).

Return Period (yr)	20	100	200	1000
Mean Difference (m)	0.031	0.041	0.044	0.055
SD (m)	0.066	0.088	0.096	0.121
t score= $M\sqrt{N}/SD$	2.706	2.678	2.653	2.623
P (Significance=0.01)	0.005	0.006	0.006	0.006




Figure 5. Extreme Sea Level ODN (m) for a 100-year return period at 35 selected locations in the UK.

Table 5: Extreme Sea-level (ESL) at UK coastal locations calculated from data to 2018 using TMAX method. The differences listed are relative to Batstone 2013 using data to 2009. Heights are in meters above Ordnance Datum Newlyn (ODN), relative to the mean sea level on 1 January 2008.

Location	Return Period (yrs) Difference from Batstone (m)							
	100	200	1000	D20	D100	D200	D1000	
Aberdeen	3.14	3.20	3.35	0.02	0.03	0.03	0.06	
Avonmouth	8.75	8.85	9.08	-0.14	-0.23	-0.26	-0.35	
Barmouth	4.08	4.16	4.36	-0.04	-0.05	-0.06	-0.05	
Bournemouth	1.84	1.91	2.08	0.00	0.03	0.04	0.09	
Dover	4.62	4.74	5.01	0.11	0.14	0.17	0.21	
Felixstowe	3.59	3.72	4.03	-0.06	-0.13	-0.18	-0.34	
Fishguard	3.52	3.58	3.73	0.01	0.00	0.00	0.00	
Heysham	6.59	6.70	6.95	-0.08	-0.11	-0.12	-0.14	
Hinkley Point	7.57	7.64	7.81	-0.10	-0.17	-0.20	-0.28	
Holyhead	4.15	4.26	4.52	0.22	0.32	0.37	0.48	
Immingham	5.11	5.24	5.55	0.16	0.22	0.24	0.30	
Kinlochberv	3.94	4.04	4.28	0.09	0.10	0.10	0.11	

ie								
Leith	3.81	3.87	4.02	-0.02	-0.07	-0.10		-0.18
Lerwick	1.88	1.93	2.06	0.02	0.05	0.06		0.11
Llandudno	5.30	5.38	5.58	0.01	0.01	0.00		0.00
Lowestoft	3.07	3.23	3.60	0.04	0.00	-0.04		-0.18
Milford Haven	4.75	4.83	5.02	0.09	0.08	0.08		0.07
Millport	3.55	3.67	3.96	0.06	0.03	0.00		-0.07
Moray Firth	3.36	3.41	3.54	0.09	0.07	0.06		0.03
Mumbles	6.03	6.11	6.30	0.02	-0.02	-0.04		-0.09
Newhaven	4.26	4.32	4.47	-0.07	-0.11	-0.13		-0.17
Newlyn	3.59	3.66	3.81	0.10	0.13	0.15		0.18
Newport	8.20	8.28	8.47	0.00	-0.08	-0.13		-0.25
North Shields	3.93	4.03	4.27	0.14	0.17	0.17		0.16
Portpatrick	3.47	3.57	3.79	0.07	0.10	0.12		0.18
Portsmouth	3.03	3.10	3.25	0.00	-0.02	-0.02		-0.03
Port Ellen	1.95	2.05	2.27	-0.20	-0.19	-0.17		-0.14
Sheerness	4.45	4.56	4.80	0.08	-0.02	-0.08		-0.25
Stornoway	3.43	3.50	3.67	0.04	0.09	0.10		0.15
Tobermory	3.50	3.57	3.75	-0.15	-0.24	-0.30		-0.43
Ullapool	3.84	3.92	4.11	0.06	0.08	0.10		0.15
Weymouth	2.27	2.32	2.45	0.08	0.07	0.06		0.05
Whitby	3.89	3.97	4.16	-0.08	-0.13	-0.17		-0.25
Wick	2.90	2.97	3.12	0.06	0.07	0.08		0.09
Workington	5.91	6.02	6.28	0.10	0.10	0.11		0.13
Mean				0.02	0.01	0.00	0.02	
Stdev				0.09	0.12	0.14	0.20	

Page Break

The values for sea level rate of rise do not include glacial isostatic adjustment (GIA) or geoid adjustments and are reasonably in line with reported values, (Wahl et al. 2013; Church and White 2011, IPCC 2021). Table 5 shows the heights using the TMAX method corresponding to the return periods of 20, 100, 200, and 1000 years for the 35 locations; it also lists the differences from Batstone's results. Figure 5 shows the ESL values for a 100-year return period deduced by the TMAX using the data to 2018. The colours indicate whether the values for 2018 are higher (red) or lower (green) compared with the TMAX analysis to 2009. Investigation of the individual contributing tidal locations reveals four main outliers when comparing this 2018 TMAX study to Batstone's results for the 200-year return period (with heights shown in brackets) are; Tobermory (-0.3), Avonmouth (-0.26), Holyhead (0.37), Immingham (0.24). Attention is drawn to these locations as their ESLs may require revision.

5. Conclusion

When applied to UK tide gauge data, the described TMAX method generates reasonable estimates of extreme sea level return period and height, broadly in agreement with published values (UK Environment Agency 2011), with mean differences of the order of centimetres and standard deviation of the order of decimetres. However, the TMAX method is much simpler than the SSJPM used in EA2011. Unlike the SSJPM, TMAX does not use or require a) harmonic tidal analysis, b) the use of a probability density function, or c) the fit of a generalized

Pareto distribution (GPD), nor does it require manual intervention so is also amenable to automation. Compared to the well-known AMAX method, the TMAX method gives a significantly better internal fit and reduced variance. Missing data, quite common in tide gauge records, is also handled by TMAX in a much simpler, more efficient and elegant way, rather than with AMAX. The study indicates that the TMAX method is eminently suitable for coastal extreme sea level analysis and is at least as accurate as the AMAX and SSJPM methods. This innovative approach can more effectively inform strategies for coastal management and resilience.

Acknowledgments

The author is grateful to the British Oceanographic Data Centre (BODC), who provided the tidal data used in this research via their "Sea Level Portal" at <https://www.bodc.ac.uk>. BODC do not accept any liability for the correctness and/or appropriate interpretation of the data or their suitability for any use.

Competing Interests

The author declares that he has no conflict of interest.