



Prescribing the aerosol effective radiative forcing in the Simple Cloud-Resolving E3SM Atmosphere Model v1

Naser Mahfouz¹, Hassan Beydoun², Johannes Mülmenstädt¹, Noel Keen³, Adam C. Varble¹, Luca Bertagna⁴, Peter Bogenschütz², Andrew Bradley⁴, Matthew W. Christensen¹, T. Conrad Clevenger⁴, Aaron Donahue², Jerome Fast¹, James Foucar⁴, Jean-Christophe Golaz², Oksana Guba⁴, Walter Hannah², Benjamin Hillman⁴, Robert Jacob⁵, Wuyin Lin⁶, Po-Lun Ma¹, Yun Qian¹, Balwinder Singh¹, Christopher Terai², Hailong Wang¹, Mingxuan Wu¹, Kai Zhang¹, Andrew Gettelman¹, Mark Taylor⁴, L. Ruby Leung¹, Peter Caldwell², and Susannah Burrows¹

¹Pacific Northwest National Laboratory, Richland, WA, USA

²Lawrence Livermore National Laboratory, Livermore, CA, USA

³Lawrence Berkeley National Laboratory, Berkeley, CA, USA

⁴Sandia National Laboratories, Albuquerque, NM, USA

⁵Argonne National Laboratory, Lemont, IL, USA

⁶Brookhaven National Laboratory, Upton, NY, USA

Correspondence: Naser Mahfouz (naser.mahfouz@pnnl.gov) and Susannah Burrows (susannah.burrows@pnnl.gov)

Abstract. Aerosol effective radiative forcing critically influences climate projections but remains poorly constrained. Using the Energy Exascale Earth System Model (E3SM) Simple Cloud-Resolving E3SM Atmosphere Model (SCREAM) v1 configuration, we quantify the forcing due to anthropogenic aerosol changes using a simplified prescribed aerosol scheme (SPA) derived from E3SM v3. Nudged simulations at 3 km and 12 km horizontal grid spacings reveal a more negative aerosol forcing than the reference 100-km E3SM v3 whence the SPA properties are derived. The resulting globally averaged aerosol forcing signal is largely due to aerosol–cloud interactions and exhibits little overall resolution sensitivity. While the default SPA scheme overestimates cloud droplet dependence on aerosols, parameterization adjustments in the activation process reconcile forcing estimates with the reference model. Our results demonstrate the ability to use a prescribed aerosol scheme to hold aerosol forcing to a desired strength across resolutions.

1 Introduction

Atmospheric aerosol particles are an essential component of climate models (Intergovernmental Panel On Climate Change, 2014; Carslaw, 2022). Aerosols are known to influence the atmosphere state directly by altering its radiation budget through aerosol–radiation interactions and indirectly by altering cloud properties through aerosol–cloud interactions (Bellouin et al., 2020, and references therein). Through their role as cloud condensation nuclei and ice nuclei, aerosol particles modify liquid and ice clouds (e.g., Mülmenstädt et al., 2019; Burrows et al., 2022). Despite significant progress in aerosol-related studies over recent decades, many questions remain about the role of aerosols in climate projections — particularly those involving aerosol–cloud interactions (Kreidenweis et al., 2019).



Accurately modeling aerosols and their effects on the atmosphere is complicated by their variability in source, composition, and size (e.g., Seinfeld and Pandis, 2016; Topping and Bane, 2022). Additional complexity arises in processes they undergo and processes they influence (Carslaw, 2022). Aerosols can be emitted into the atmosphere as fully formed particles from natural and anthropogenic sources (Seinfeld and Pandis, 2016), or they can be formed in the atmosphere from precursor gases via gas-to-particle conversion and new-particle formation that have their origin in natural or anthropogenic sources (Dunne et al., 2016). They can undergo a variety of processes in the atmosphere, such as growing or shrinking via condensation or evaporation as well as coagulation growth or loss via collisions with other particles (Fuchs, 1964; Fuchs and Sutugin, 1970; Seinfeld and Pandis, 2016). They can additionally be scavenged by clouds and precipitation, sometimes reemerging by resuspension, and they can be removed from the atmosphere by wet and dry deposition (Adams and Seinfeld, 2002; Wang et al., 2020). Whether direct or indirect, aerosol effects on the atmosphere and thus climate depend closely on the aforementioned processing as well as their concentration, composition, and size (Adams et al., 2013; Carslaw, 2022).

Due to the complexity associated with aerosols, they are generally represented through simplified schemes in atmosphere models, varying from prescribed (e.g., Stevens et al., 2017) to modal (e.g., Stier et al., 2005; Liu et al., 2012) and interactive sectional (e.g., Tilmes et al., 2023) schemes. In all, the multi-scale complexity of aerosol representation is usually reduced to balance process realism with computational performance depending on the need. Notwithstanding, fully constraining aerosol effects on the atmosphere remains elusive. Notably, definitively constraining the effective aerosol forcing to anthropogenic aerosol perturbation remains mired in uncertainty (Intergovernmental Panel On Climate Change, 2023). One of the main challenges is understanding the adjustments (Quaas et al., 2024) to aerosol perturbations. While the instantaneous radiative response to an aerosol change is relatively well understood (Twomey effect; Twomey, 1977; Quaas et al., 2020), subsequent adjustment processes like precipitation suppression (Albrecht, 1989) and entrainment feedback (Ackerman et al., 2004; Bretherton et al., 2007) lead to changes in macroscopic cloud properties (like cloud water amount and cloud fraction) that are far less constrained (Gryspeerdt et al., 2019; Bellouin et al., 2020).

With the dawn of new high-resolution global atmosphere models, it is hoped that some of the uncertainty in aerosol–cloud interactions can be reduced (e.g., Sato et al., 2018; Terai et al., 2020; Mülmenstädt and Wilcox, 2021; Herbert et al., 2024; Weiss et al., 2024). Current state-of-the-art climate models, which typically have an effective resolution of about 100 km, tend to have interactive aerosol schemes. On the other hand, next-generation high-resolution models running with an effective resolution below 10 km are computationally expensive, and so they often use the simplest and cheapest representation for aerosols. Most high-resolution models use prescribed aerosol schemes but some are being developed with interactive aerosol schemes (Weiss et al., 2024, and references therein). Even with prescribed aerosols, high-resolution models can still provide valuable insights into aerosol–cloud interactions (e.g., Herbert et al., 2024).

In this study, we examine the sensitivity of the Energy Exascale Earth System Model (E3SM) Simple Cloud-Resolving E3SM Atmosphere Model (SCREAM) v1 configuration to aerosol changes during the industrial era. Our focus is the global quantification of the aerosol effective radiative forcing (ERFaer), and especially its dominant component: the aerosol–cloud interactions. Goto et al. (2020) report three-year-long global simulations at a 14-km horizontal grid spacing with an interactive aerosol scheme, including sensitivity studies to anthropogenic aerosols. They show that most features related to aerosols



(especially primary aerosols) improve slightly with finer grid spacing, but the ERFaer sensitivity to resolution is small (Goto et al., 2020). Additionally, Weiss et al. (2024) report a one-year-long global simulation at a 5-km horizontal grid spacing with interactive modal aerosols at present-day conditions using a one-moment scheme that predicts number concentration with other aerosol properties prescribed. They highlight the emergence of mesoscale features of natural aerosols like dust aerosol patterns in desert storms and the interplay of sea salt aerosol with tropical cyclones (Weiss et al., 2024). Both studies utilized free-running simulations that are not nudged to reanalysis data. Our study differs from both in its exclusive focus on the global ERFaer signal obtained from a prescribed aerosol scheme, using pairs of one-year-long simulations at 3-km, 12-km, and 100-km effective resolutions, all nudged to reanalysis data in order to deduce a robust ERFaer signal. While an interactive treatment of aerosols offers more realism overall, it is also more computationally expensive and less flexible for disentangling process pathways due to the two-way nature of the coupling between aerosols and the model state. In the following manuscript, we focus on the aerosol activation process and highlight its importance in determining the ERFaer signal. We note that an interactive aerosol scheme is in development for the SCREAM configuration and should be available in the near term.

2 Methods

2.1 Modeling framework

2.1.1 SCREAM v1

We use the Energy Exascale Earth System Model (E3SM) Simple Cloud-Resolving E3SM Atmosphere Model (SCREAM) v1 configuration in this study. The SCREAM v1 configuration is described in detail by Donahue et al. (2024), and we adopt changes outlined in their Section 5.5. We briefly outline parts of the configuration most pertinent to this study. The simplified higher-order closure scheme (SHOC; Bogenschutz and Krueger, 2013) is used to parameterize cloud macrophysics and turbulent vertical mixing. A modified version of the predicted particle properties scheme (P3; Morrison and Milbrandt, 2015) is used for cloud microphysics parameterization. In the modified P3 version, there exists only one ice category and the liquid supersaturation is removed for consistency with SHOC's liquid saturation adjustment assumption (Caldwell et al., 2021). Additionally, the ice cloud fraction is set based on a grid-mean ice mass mixing ratio threshold — all or nothing based on the condition: $q_i > 10^{-14} \text{ kg kg}^{-1}$.

Aerosol effects on the model physics are parameterized by the simple prescribed aerosol (SPA) scheme. The SPA scheme prescribes the optical properties of aerosols, so that they are accounted for in the radiation scheme (RRTMGP; Pincus et al., 2019). The SPA aerosol optical properties are the three-dimensionally resolved and band-resolved single scattering albedo, asymmetry parameter, as well as the shortwave and longwave optical depths. Furthermore, SPA is linked to the model microphysics via its impact on the prognostic liquid cloud droplet number concentration, N_c . SPA prescribes a three-dimensional grid-mean cloud condensation nuclei concentration, N_{ccn} , whose value scaled by the three-dimensional liquid cloud fraction (C_f) is used to update grid-mean N_c according to Equation 1.

$$N_c = \max(N_c, \alpha(N_{ccn}C_f)) \quad (1)$$



85 The function $\alpha(N_{\text{ccn}}C_f)$ represents the activation process of cloud condensation nuclei into liquid cloud droplets. For simplicity, the default SCREAM v1 configuration assumes a linear function of activation α , defined as $\alpha(N_{\text{ccn}}C_f) = N_{\text{ccn}}C_f$. This represents the upper limit of theoretically possible values (e.g., Chen et al., 2011, Equation 5), though some global models show values exceeding that (e.g., Gryspeerdt et al., 2023, Section 2.2). By default, SPA uses N_{ccn} at 0.10 % supersaturation as a reasonable proxy for cloud condensation nuclei concentration and for its availability as an output and as mean representative

90 for all cloud conditions. For the rest of the manuscript, unless explicitly stated otherwise, we use N_{ccn} as a shorthand for cloud condensation nuclei concentration at 0.10 % supersaturation. Both the optical properties and N_{ccn} are prescribed as climatological monthly in-grid means at a coarser resolution than the model's (see Section 2.1.2); during the simulation, the values are interpolated to the model's resolution and time step. The P3 ice nucleation process is not aerosol-aware, and so it is not directly affected by the SPA scheme (Donahue et al., 2024).

95 Although generally similar to the MACv2-SP scheme (Stevens et al., 2017) and variations thereof (Fiedler et al., 2017; Herbert et al., 2024), the SPA scheme in SCREAM v1 has some key differences. First, even though MACv2-SP is derived from an observational dataset, it still requires a reference N_c (i.e., a simulated preindustrial background N_c) obtained from a model run to estimate anthropogenic aerosol changes (Stevens et al., 2017, Section 3). On the other hand, SPA uses a more consistent approach by prescribing all aerosol optical and cloud properties from a previous model run. Second, while the MACv2-SP

100 scheme prescribes only shortwave aerosol optical properties (Fiedler et al., 2017), the SPA scheme prescribes both shortwave and longwave aerosol optical depths. Third and most substantially, the MACv2-SP and SPA schemes differ significantly in how they are coupled to cloud microphysics. The original MACv2-SP accounts for the Twomey effect and its associated radiative adjustments but is not coupled to cloud microphysics (Fiedler et al., 2017, Section 2.2.1). A modified version of MACv2-SP (Herbert et al., 2024) includes further coupling to cloud microphysics but only through changing the autoconversion rate of

105 cloud droplets to raindrops (Herbert et al., 2024, Section 2.1). In SCREAM v1, SPA is coupled to the cloud microphysics scheme directly by setting a climatological minimum cloud droplet number concentration, N_c . The SPA-derived minimum replenishes the prognostic N_c at each time step right before the cloud microphysics calculations. This ensures that aerosol effects are accounted for in the prognostic tracer N_c , while allowing for cloud processes in P3, vertical mixing in SHOC, and large-scale transport to modify it (Donahue et al., 2024, Section 2.1). The P3 cloud processes include autoconversion, accretion,

110 sedimentation, and collisions with ice crystals (Morrison and Milbrandt, 2015, Appendices B and C).

In summary, SPA replaces the “aerosol state” (e.g., Karset et al., 2020, “aerosol physics” in Figure 2) with a climatological mean state derived from long model runs with interactive aerosols. In doing so, it prescribes aerosol–radiation interactions through aerosol optical properties and sets a climatological minimum N_c to account for aerosol–cloud interactions. This represents a one-way coupling between the aerosol state and the rest of the model, wherein the aerosol state influences the model's

115 state but not vice versa. Of special note, this one-way coupling allows for the study of the instantaneous effect of aerosol perturbation as well as adjustments to it, without the complications of feedbacks from the new cloud state to the aerosol state.



2.1.2 E3SM v3

The Energy Exascale Earth System Model (E3SM) v3 is a milestone release (E3SM Project, 2024), with significant improvements in the model's ability to reproduce the historical temperature record (Golaz et al. 2025, Xie et al. 2025, both in preparation). We use the state-of-the-art E3SM v3 coarse-resolution atmosphere model, which is run on an unstructured grid whose effective horizontal resolution is 100 km with output at 150 km near the equator and 80 vertical levels. We briefly describe features of the E3SM v3 atmosphere model most pertinent to the study at hand. Unlike SCREAM v1, it uses the Cloud Layers Unified By Binomials scheme (CLUBB; Golaz et al., 2002) for the macrophysics parameterization of clouds and turbulent vertical mixing. Similar to SCREAM v1, it uses the P3 scheme for stratiform cloud microphysics parameterization (Teraï et al., 2024), but with slightly different tuning parameters (Shan et al., 2024). While SCREAM v1 has no parameterized deep convection, E3SM v3 uses a Zhang and McFarlane (1995) type convection scheme that also includes cloud microphysics effects (Teraï et al., 2024). E3SM v3 also uses the RRTMG radiation scheme (Clough et al., 2005). Most importantly, it has an interactive modal aerosol model (MAM) scheme, based on MAM4 in previous E3SM versions (Wang et al., 2020), with new features that are being characterized and evaluated in detail for upcoming publications (e.g., Xie et al, 2025; Wang et al., 2025, Burrows et al., 2025, all in preparation).

We use the default E3SM v3 configuration in this study to produce the reference climatological mean state for SPA as described in Section 2.2.1. We also use the default E3SM v3 configuration to conduct additional sensitivity studies detailed in Section 2.2.2. Furthermore, we reconfigure E3SM v3 to prescribe aerosols and chemistry using SPA as described in Section 2.1.1. We call the default E3SM v3 configuration E3SM-MAM and we call the modified E3SM v3 configuration with prescribed aerosols E3SM-SPA. The only differences between E3SM-MAM and E3SM-SPA are the nature of aerosols and chemistry (interactive vs. prescribed, respectively) and the radiation scheme (RRTMG vs. RRTMGP, respectively), with minor differences in coupling interfaces between components.

2.2 Simulation protocol

2.2.1 Reference simulation

To reduce potential sampling biases in the SPA climatological means, we run the E3SM-MAM v3 model for 31 years with only the atmosphere and land components with prescribed sea ice extent and sea surface temperature. For the present-day simulation, all initial conditions and prescribed settings correspond to a climatological state representative of around 2010. For the pre-industrial run, we keep everything the same as the present-day run, but change the settings for aerosols and their precursors to a pre-industrial state in 1850. The simulations are free-running, that is, without the use of nudging to control the meteorological state. We discard the first year and obtain climatological monthly means for the last 30 years of the SPA fields of interest described in Section 2.1.



2.2.2 Sensitivity simulations

The main goal of this study is to examine the sensitivity of the SCREAM v1 configuration to aerosol perturbation, namely from pre-industrial to present-day conditions. To do so, we conduct two SCREAM v1 simulations: one with SPA pre-industrial aerosol conditions and another with SPA present-day aerosol conditions. With the SPA input files obtained from the reference simulation (Section 2.2.1), we run SCREAM v1 at two horizontal grid spacings (3 km and 12 km) for 13 months starting on August 1, 2019 and ending on September 1, 2020. This time period was selected based on the availability of boundary conditions as part of a series for sensitivity studies of the SCREAM v1 configuration. Specifically, a recent 13-month period was chosen with a weak El Niño Southern Oscillation starting from a northern hemisphere summer for a parallel radiative feedback study by Terai et al. (2025). The sea surface temperature is obtained from the Operational Sea Surface Temperature and Ice Analysis (Donlon et al., 2012; UKMO, 2012) and the sea ice coverage is obtained from the The European Organization for the Exploitation of Meteorological Satellite Ocean and Sea Ice Satellite Application Facility (UKMO, 2012). The atmosphere initial condition is obtained from the fifth-generation European Centre for Medium-Range Weather Forecasts atmospheric reanalyses (ERA5) data at a horizontal resolution of $0.25^\circ \times 0.25^\circ$ and native vertical level (Hersbach et al., 2020). The atmosphere initial condition is then interpolated to SCREAM v1's horizontal and vertical levels. The land initial condition is obtained from running the land model forced by atmospheric reanalysis from CRUNCEP data (Viovy, 2018) until the SCREAM v1 simulation start date (for 20 years for the 3-km runs and for 28 years for the 12-km runs).

To enhance the signal-to-noise ratio, we nudge the model horizontal winds towards the Modern-Era Retrospective Analysis for Research and Applications, version 2 (MERRA-2) reanalysis data (Gelaro et al., 2017; Global Modeling And Assimilation Office, 2015). The MERRA-2 data is obtained at a temporal resolution of six hours, horizontal resolution of $0.5^\circ \times 0.625^\circ$, and native vertical levels (Global Modeling And Assimilation Office, 2015). The MERRA-2 data is then interpolated offline to an unstructured grid whose horizontal resolution is 100 km at the equator. During SCREAM v1 runs, the nudging data is interpolated online to the model's temporal, horizontal, and vertical levels and applied to the model's horizontal winds. The application of nudging is consistent with best practices established in the E3SM Project to diagnose aerosol forcing while controlling for variability due to large-scale circulation (e.g., Zhang et al., 2022a, b; Mahfouz et al., 2024). Specifically, the nudging data is read at a six-hourly cadence; it is temporally interpolated linearly in between analysis time-steps. Additionally, the nudging is applied with a relaxation timescale of six hours to the horizontal winds at all vertical levels.

We repeat the above procedure for the E3SM-MAM v3 and E3SM-SPA v3 configurations at 100 km, with the following difference: the model is run for 13 months starting on December 1, 2009, and nudged with MERRA-2 data corresponding to that period. We note that the source of the nudging data (MERRA-2 vs. ERA5) and the period of interest (2019–2020 like in SCREAM v1 runs vs. 2009–2010 in E3SM v3 runs) have a negligible effect on the ERFaer signal, and the choice of nudging data and period was made purely for practical reasons of using readily available, existing data. The use of nudging is solely to control for the large-scale circulation and to enhance the signal-to-noise ratio in the ERFaer calculations (e.g., Kooperman et al., 2012; Zhang et al., 2014; Forster et al., 2016; Zhang et al., 2022b); as long as the same nudging data is used for the present-day and pre-industrial runs, the ERFaer signal will be consistent regardless of the year or the reanalysis data used. In



Table 1. Summary of simulations conducted in this study, labeled as they appear in the figures. Because the simulations are used to calculate ERFaer, the simulations appear in pairs. For each pair, the only difference between the two simulations is the aerosol state, either pre-industrial or present-day. The simulations are described in detail in Section 2.2.

	Simulation pair	Resolution	Length	Nudging	Notes
1	Reference 100 km	100 km	31 years	No	Create SPA files
2	SCREAM v1 12 km	12 km	13 months	Yes	vs. 3: resolution sensitivity
3	SCREAM v1 3 km	3 km	13 months	Yes	SCREAM v1 ERFaer
4	E3SM-MAM v3; $N_c \geq 20 \text{ cm}^{-3}$	100 km	13 months	Yes	vs. 1: nudging sensitivity
5	E3SM-MAM v3	100 km	13 months	Yes	vs. 4: N_c limiter sensitivity
6	E3SM-SPA v3; $\alpha = 2000 (N_{\text{ccn}} C_f)^{0.55}$	100 km	13 months	Yes	vs. 7: activation sensitivity
7	E3SM-SPA v3; $\alpha = N_{\text{ccn}} C_f$	100 km	13 months	Yes	vs. 3: SPA-model sensitivity

Note that the N_c limiter is applied in-cloud, that is, in-cloud $N_c \geq 20 \text{ cm}^{-3}$. All other quantities throughout the manuscript are grid-mean values, unless explicitly noted otherwise.

E3SM v3 testing we conducted, but not shown here, we found that systematic uncertainties in ERFaer due to choice of nudging data and period are approximately 0.1 W m^{-2} .

3 Results

3.1 SCREAM v1 ERFaer

185 The central question of this study is: Can the high-resolution SCREAM v1 configuration reproduce an aerosol effective radiative forcing (ERFaer) from pre-industrial to present-day conditions based on the SPA prescription from a low-resolution model that is similar to that produced by low-resolution models with fully interactive aerosols? In Figures 1 and 2, we show the time series and spatial distribution of the SCREAM v1 ERFaer, which is defined as the difference in the top-of-model radiative imbalance between the present-day and pre-industrial simulations (Forster et al., 2016). It is immediately clear that

190 SCREAM v1 does not reproduce the ERFaer from the reference E3SM v3 simulations; instead, it produces a significantly more negative ERFaer value, even though aerosols in SCREAM are prescribed based on the E3SM v3 simulations. Both horizontal resolutions of SCREAM v1 produce similar seasonal cycles and total numbers (Figure 1), but with slightly differing spatial distributions (Figure 2). Compared to E3SM v3, the SCREAM v1 configuration produces a more accentuated ERFaer signal spatially (Figure 2), resulting in a more negative global ERFaer.

195 While the ERFaer resolution sensitivity of the SCREAM v1 configuration appears to be minimal, the 12-km configuration produces a slightly more negative ERFaer signal than the 3-km configuration. At first glance, this may be surprising because of the significant differences in the base cloud and climate states between the two resolutions. The summary in Table 2 shows that while the base states at present-day settings are different, the perturbation from pre-industrial to present-day settings are broadly similar. The cloud droplet concentration at cloud top is similar, showing that SPA is effectively translating the cloud




Figure 1. Seasonal cycle of area-weighted globally averaged aerosol effective radiative forcing (ERFaer) from the SCREAM v1 configuration and the reference from which the SPA files were obtained (spatiotemporal average in the legend). The climatological reference SPA values are obtained from a simulation run at a 100-km horizontal resolution with E3SM v3 for 31 years with only the atmosphere and land components interactive without nudging (the first year is discarded). The SCREAM v1 configuration is run at two horizontal grid spacings (3 km and 12 km) forced with the same SPA files with other boundary conditions, while nudged to MERRA-2 data from August 1, 2019 to September 1, 2020; the first month (August 2019) is discarded.

condensation nuclei prescription into cloud droplets consistently across resolutions, as designed in Equation 1. Additionally, nudging is employed in these studies to control for confounding factors that may distort the ERFaer signal. Specifically, the nudging is strong enough to keep large-scale features roughly constant across the perturbation (e.g., Gettelman et al., 2020).

Yet, we note that in the 12-km simulation shown in Figure 2, the more negative ERFaer signal appears in convectively active land regions (Amazon, East Africa, South Asia, Maritime Continent) where aerosol perturbations exist and/or orographic precipitation is common (Maritime Continent, Himalayas, Western Ghats, Ethiopian Highlands, tropical Andes). This potentially indicates that explicit convection representation and topographic resolution modulate regional ERFaer through their effects on clouds and precipitation. On the other hand, the reverse appears to be true in the marine stratocumulus regions west of continental masses that are so often the focus of aerosol–cloud interactions studies. In those regions, the ERFaer signal appears to be broadly more negative in the 3-km run than the 12-km run. Together, the opposing trends in the two regions suggest a globally insensitive ERFaer signal, but with a regional dependence on the resolution. Such details can help guide future research efforts to probe the regional aspect of aerosol–cloud interactions in high-resolution models.




Figure 2. Same as Figure 1, but showing a spatial distribution of the temporally averaged ERFaer (spatiotemporal average at top left). Note that here and elsewhere, all data are interpolated onto the coarsest unstructured output grid (150 km) for visualization.

We assess that the ERFaer signal is mostly due to aerosol–cloud interactions (indirect effects; approximately 90 % of the total ERFaer signal). Aerosol–radiation interactions (direct effects) are relatively small in the SCREAM v1 configuration, as they are relatively small in the reference simulations, compared to the indirect effects. The most significant contributions to the ERFaer signal are from the instantaneous radiative forcing due to changes in cloud droplet concentration (Twomey effect) and the adjustment of cloud content to the cloud droplet changes (cloud content adjustments). Through partial radiative perturbation studies (Mülmenstädt et al., 2019), we find that the liquid and ice content adjustment terms account only for about 15 % (-0.23 and -0.05 W m^{-2} for liquid and ice water path adjustments, respectively) of the total ERFaer signal in the 12-km simulations, broadly consistent with previous studies (e.g., Zelinka et al., 2023, Figure 9, “cloud amount” component). Taken together, these



Table 2. Summary of area-weighted, globally averaged quantities in present-day (PD) conditions and the PD–PI differences in parentheses (Δ).

SCREAM v1 quantity	12 km PD (Δ)	3 km PD (Δ)
TOA ΔF (W m^{-2})	−2.06 (−1.95)	3.23 (−1.89)
SW CRE (W m^{-2})	−51.18 (−1.20)	−46.29 (−0.98)
LW CRE (W m^{-2})	26.00 (−0.22)	25.30 (−0.22)
Cloud-top liquid C_f	0.14 (0.00)	0.13 (0.00)
Cloud-top ice C_f	0.78 (−0.00)	0.79 (−0.01)
In-cloud LWP (g m^{-2})	31.89 (2.48)	26.51 (1.45)
In-cloud IWP (g m^{-2})	7.30 (0.17)	10.70 (−0.09)
In-cloud $N_{c,\text{top}}$ (cm^{-3})	64.49 (20.41)	65.10 (19.63)
Cloud-top R_c (μm)	1.11 (−0.08)	0.98 (−0.04)
Cloud-top R_i (μm)	10.90 (−0.03)	12.38 (−0.07)

TOA ΔF is the top-of-atmosphere net radiation imbalance. CRE is cloud radiative effects; values for shortwave (SW) and longwave (LW) are shown. Cloud-top quantities were calculated using the cloud-top statistical definition (Tiedtke et al., 1979, and references therein). Liquid water path (LWP) and ice water path (IWP) are calculated as the vertically integrated mass of liquid and ice water, respectively. The effective cloud droplet radius R_c and effective ice crystal radius R_i are shown at cloud top.

220 results indicate that the Twomey effect is the most important component of the ERFaer signal in the SCREAM v1 configuration, which depends strongly on the aerosol activation process.

The default SPA translation of aerosol activation (Equation 1, $\alpha = N_{\text{ccn}} C_f$) shows a significant overestimation in the indirect effect signal going from the reference E3SM v3 simulation to SCREAM v1 simulations. There are several potential explanations for the discrepancy. First, the models use different macrophysics schemes; second, they use different tunings of the same cloud microphysics scheme; third, one model has explicit convection while the other parameterized; fourth, one model uses a fully interactive aerosol scheme while the other completely bypasses that with a simple prescribed one. In the following sections, we show that the two most important factors are the tuning of the cloud droplet number concentration limiter employed in the E3SM v3 cloud microphysics scheme and the nature of the aerosol activation from cloud condensation nuclei to cloud droplets in SCREAM v1. To do so, we use the relatively inexpensive E3SM v3 (Section 2.1.2) to conduct additional sensitivity studies in Section 3.3.

3.2 Droplet activation

Because pure-water cloud droplets require an atmospherically infeasible supersaturation to form, cloud droplets must form on cloud condensation nuclei (CCN) that activate at a given supersaturation (Twomey, 1991, Section 5). Representation of the so-called CCN spectrum varies from simplified power-law functions (e.g., Twomey, 1959) to more complex mappings from




Figure 3. Relationship between grid-mean $N_{\text{ccn}} C_f$ and grid-mean N_c in E3SM-MAM v3 model without the N_c limiter. The N_{ccn} shown here is the cloud condensation nuclei concentration at 0.10 % supersaturation. Because E3SM v3 tends to produce a high frequency of low- N_c values, we filter the data by two different cutoffs in N_c : 10^{-5} and 10^{-6} kg^{-1} . The circles represent 150 logarithmic bins of the data obtained from 75 randomly selected temporal samples fully resolved in space and height. The count frequency in the legend is the number of data points in each bin, with the total bins being 129600000 (75 samples, 21600 horizontal columns, and 80 vertical levels). The solid lines are denoted by the equations in the legend. The $d \ln N_c / d \ln N_{\text{ccn}}$ slope of 0.55 is inspired by observational studies cited in the main text, while the prefactor (2000) is chosen to best fit the data with the higher cutoff.

235 aerosol size space to the supersaturation space (e.g., Nenes and Seinfeld, 2003). In the supersaturation space, it is possible to calculate N_{ccn} from aerosol information at any supersaturation (Ghan et al., 2011, Equation 20). The activation of N_{ccn} into N_c depends on the underlying aerosol size distribution and its composition as well as atmospheric dynamics, especially updraft velocity, thus resulting in distinct regimes of activation (Reutter et al., 2009). While simplified activation schemes (e.g., Seifert and Beheng, 2006, Equation 17) and more complex activation schemes (e.g., Abdul-Razzak and Ghan, 2000; Ming et al., 2006; Morales Betancourt and Nenes, 2014) are routinely used, the relationship between N_{ccn} and N_c predicted by activation schemes remains variable and uncertain from a globally aggregated perspective (Gryspeerdt et al., 2023), despite the models' general agreement with more accurate but computationally expensive parcel models (e.g., Ghan et al., 2011; Rothenberg et al., 2018, and references therein). In summary, while theoretical studies along with laboratory and field experiments have improved our understanding of the aerosol activation process, uncertainties persist in its their remotely sensed and globally modeled impact

245 on cloud droplet concentrations (McCoy et al., 2017; Hasekamp et al., 2019; Gryspeerdt et al., 2023).




Figure 4. Like Figure 1, but for the E3SM v3 configurations tested.

Despite the uncertainty, it is thought that the relationship between N_{ccn} (at constant supersaturation) and N_c exhibits a log-log slope that is less than 1, specifically, $0.3 \leq d \ln N_c / d \ln N_{\text{ccn}} \leq 0.8$ (McCoy et al., 2017; Bellouin et al., 2020). This resulting sublinear relationship is often understood in the context of supersaturation variability and the transition from aerosol-limited to updraft-limited activation regimes. In order to better constrain $\alpha (N_{\text{ccn}} C_f)$ in Equation 1, we examine the correlations between these two variables in an E3SM v3 run in Figure 3. We choose to construct a correlation between $N_{\text{ccn}} C_f$ and N_c as they appear in Equation 1 using the present-day simulation of experiment 5 in Table 2, which does not include the N_c limiter. We note that E3SM v2 and v3 runs tend to have a frequent occurrence of low- N_c values (Shan et al., 2024; Wan et al., 2025), which may distort the correlation; we thus filter by two different cutoffs in Figure 3. The $d \ln N_c / d \ln N_{\text{ccn}}$ slope of 0.55 is inspired by observational studies (Hasekamp et al., 2019; Gryspeerdt et al., 2023) and lies in the middle of the aforementioned assessed range (McCoy et al., 2017; Bellouin et al., 2020), but it generally varies depending on atmospheric conditions and underlying aerosol information. Based on the results in Figure 3, we estimate that $\alpha = 2000 (N_{\text{ccn}} C_f)^{0.55}$ likely summarizes the relationship in E3SM v3 without the potentially too-frequent low- N_c values; as such, we use the new formulation in the sensitivity test in Section 3.3. We acknowledge the crude nature of this approach due to the confounding factors across space and time when it comes to the relationship between N_c and N_{ccn} (e.g., Andreae, 2009; Chen et al., 2011; Moore et al., 2013; Gryspeerdt et al., 2023; Varble et al., 2023; Ghosh et al., 2024).




Figure 5. Like Figure 2, but for the E3SM v3 configurations tested.

3.3 E3SM-SPA v3 ERFaer

In Figures 4 and 5, we show the results from the sensitivity tests conducted using the E3SM v3 configurations described in Section 2.1.2. As a confirmation, we performed a one-year nudged version (Figure 4) of the reference 31-year simulation



(Figure 1), and we find a good agreement between the ERFaer signal from the reference and the one-year simulation. We find that the N_c limiter (in-cloud $N_c \geq 20 \text{ cm}^{-3}$) has an effect of about a global average of 0.40 W m^{-2} on the ERFaer signal compared to the case where it is not applied. The remaining difference between the reference simulation and the E3SM-SPA v3 simulation (about 1 W m^{-2}) is due to the activation process in the SPA scheme. We find that we are able to roughly reproduce the SCREAM v1 ERFaer signal using the E3SM-SPA v3 configuration, with a global average of -1.89 W m^{-2} in the SCREAM v1 3-km configuration, -1.95 W m^{-2} in the SCREAM v1 12-km configuration, and -2.12 W m^{-2} in the E3SM-SPA v3 100-km configuration. In Table 3, we show the results of further sensitivity tests to confirm the robustness of the ERFaer signal to the log–log slope of the relationship between N_{ccn} and N_c in the E3SM-SPA v3 configuration, showing a consistent trend of increasing ERFaer magnitude with increasing slope. Most importantly, we are able to almost exactly reproduce the global ERFaer signal of E3SM-MAM v3 with the modified activation relationship in the E3SM-SPA v3 configuration using the fit from Figure 3.

In Figure 5, we note the modified activation parameterization ($\alpha = 2000 (N_{\text{ccn}} C_f)^{0.55}$) in the E3SM-SPA v3 configuration produces a broadly similar spatial distribution of the ERFaer signal compared to the interactive and more complex activation scheme in the E3SM-MAM v3 configuration. Nonetheless, there appear to be subtle differences between the two signals in several regions. The E3SM-MAM configuration appears to produce a more accentuated signal (positive and negative) than the E3SM-SPA configuration, though the overall global average is similar. This could be due to the deviations from the fit line in Figure 3 which effectively balances deviations above and below such that the net global result is similar. To investigate further, we examine the cloud droplet number concentration at cloud top in Figure 6. We find that the activation formulation in the E3SM-SPA v3 configuration has a significant effect on the cloud droplet number concentration at cloud top. Going from $\alpha = N_{\text{ccn}} C_f$ to $\alpha = 2000 (N_{\text{ccn}} C_f)^{0.55}$ we see an approximate decrease of 40 % in the present-day simulations (Figure 6 left) and an approximate decrease of 66 % in the PD–PI differences (Figure 6 right). Moreover, the revised activation formulation with the slope of 0.55 does not exactly match the more complex interactive MAM implementation, with a slight underestimation of cloud-top N_c , likely explainable by the fact that the data averages lie above the line corresponding to the slope of 0.55 in Figure 3. The fact that the globally averaged ERFaer signal calculated from experiments 5 and 6 in Table 1 are quite similar, but the cloud-top N_c values are different, suggests that the end climatological state may be achieved through different means. For example, it is plausible that the nature of the Twomey effect and adjustments to it may be fundamentally changed between the two configurations. We leave a more thorough investigation of these process pathways to future work.

4 Conclusions

Our main conclusion is that the global aerosol effective radiative forcing (ERFaer) can be constrained across different resolutions and configurations using a simple prescribed aerosol scheme. We find that SCREAM v1 ERFaer is dominated by indirect cloud effects, principally through the Twomey effect from aerosol-induced cloud droplet number perturbations. This radiative forcing term (Twomey effect) is sensitive to aerosol activation assumptions, while global-scale cloud adjustments to the




Figure 6. Like Figure 5, but showing the present-day values (left) and PD-PI differences (right) in grid-mean cloud-top N_c . We emphasize that the values represented are grid-mean values, which are often smaller than their in-cloud counterparts.

Twomey effect remain secondary. Consequently, we hypothesize that ERFaer resolution dependencies may emerge primarily from how models resolve aerosol lifecycle processes and activation physics in aerosol schemes.



Table 3. The sensitivity of ERFaer to the log–log slope of the relationship between N_{ccn} and N_c in the E3SM-SPA v3 configuration while controlling for the prefactor. The slope is chosen first, and then the prefactor is modified to ensure all log–log fit lines approximately pass through the intersection of the two lines in Figure 3. These experiments are not reported in Table 1; they follow experiment 6 exactly but with different values of slope and prefactor in the activation equation.

SPA activation	ERFaer (W m^{-2})
$\alpha = (N_{\text{ccn}} C_f)^{0.50} \times 4654$	–1.03
$\alpha = (N_{\text{ccn}} C_f)^{0.55} \times 2000$	–1.16
$\alpha = (N_{\text{ccn}} C_f)^{0.60} \times 860$	–1.25
$\alpha = (N_{\text{ccn}} C_f)^{0.80} \times 30$	–1.70
$\alpha = (N_{\text{ccn}} C_f)^{1.00} \times 1$	–2.12

The default SPA linear activation formulation in our study using E3SM v3 N_{ccn} overestimates cloud droplet sensitivity to aerosol perturbations. Using a simplified activation formulation that inaccurately translates aerosol effects is not unprecedented. For example, the MACv2-SP scheme as originally formulated (Stevens et al., 2017, Figure 8 top) underestimated the cloud droplet sensitivity to aerosol perturbation (Herbert et al., 2021, Figure 3 c). As a result, subsequent studies adopted a stronger aerosol sensitivity formulation (Herbert et al., 2024). We modified the prescribed relationship between N_c and N_{ccn} , which resulted in agreement of global ERFaer between E3SM v3 and SCREAM v1 across resolutions. Thus, this modified SPA formulation will be used as the default going forward. This adaptability constitutes a fundamental strength of prescribed schemes, enabling targeted hypothesis testing through controlled decoupling of aerosol–cloud interactions (Fiedler et al., 2017).

By holding the aerosol state (and thus forcing) to a preferred reference state, our framework enables different types of applications and studies. First, it enables studies of extreme weather and climate feedbacks independent from aerosol forcing uncertainty by imposing a targeted ERFaer. Second, it allows for the high-resolution investigation of cloud responses to aerosol perturbations without cloud-state feedback complications. The scheme’s one-way coupling eliminates interactive feedback challenges like wet removal parameterization uncertainties (McCoy et al., 2020; Shan et al., 2024), allowing modular analysis of aerosol–cloud interactions. This proves particularly valuable for disentangling explicitly parameterized aerosol–cloud interactions mechanisms (e.g., precipitation suppression) from implicitly represented aerosol–cloud interactions mechanisms that arise from the interplay of multiple processes like entrainment feedbacks — the interplay in these process pathways remains critical for understanding aerosol–cloud interactions (e.g., Mahfouz et al., 2024; Mülmenstädt et al., 2024b, a). Finally, our framework opens the door for systematic identification of ERFaer deviation hot spots between simple (SPA) and complex (MAM) schemes for process-level scrutiny. Ultimately, the “true” ERFaer signal is not well constrained, and uncertainties persist in how close global circulation models are to representing it faithfully. The iterative approach of using a simple scheme to constrain the signal and then using that signal to better understand and improve complex schemes is a promising avenue for future research.



320 While SPA currently utilizes E3SM v3 data, future implementations could integrate observationally constrained datasets like
MACv2-SP (Stevens et al., 2017) while preserving aerosol spatial heterogeneity (Hassan et al., 2024). Further, the documented
challenges in constraining the relationship between N_{ccn} and N_c (e.g., Gryspeerdt et al., 2023) indicate the need for a more
advanced droplet activation treatment that takes into account aerosol information as well as atmospheric factors and cloud
dynamics (e.g., Ghan et al., 2011). In the future, the SPA implementation could include a dynamic set of N_{ccn} tracers with
325 climatological relaxation to better capture the effects of large-scale circulation, macrophysics sub-grid variability, and cloud
microphysics on aerosols. Such enhancements may bring SPA closer to the MAM implementation, while still retaining the
advantages of efficiency and interpretability.

While SPA has advantages in efficiency, interpretability, and simplicity, the soon-to-be-available MAM implementation
into SCREAM will explicitly predict the aerosol lifecycle, and thus likely offer a far superior representation aerosol–cloud
330 interactions. The one-way coupling of SPA with clouds allows for easier access to certain process pathways, but the two-way
coupling nature of MAM allows for a more realistic representation of aerosol effects. Moreover, the promise of high-resolution
models like SCREAM to improve predictions relative to traditional, coarser-resolution models lies to a substantial extent in
their ability to represent convection explicitly, and it is of keen interest to know how explicitly represented convection interacts
with aerosols. In its current form, SPA hardly captures any of the links between convection and aerosols, such as the effects
335 of variable updraft velocity on activation, efficient wet removal of aerosols, and vertical transport of aerosols as well as their
precursors.

In summary, while constraining the global ERFaer signal is an important first step, it is only the beginning of fully under-
standing global aerosol–cloud interactions at higher resolution in SCREAM v1. The ability to constrain global states allows re-
searchers to proceed with confidence in adjacent studies, such as those examining extreme weather events or climate feedbacks.
340 However, focusing solely on global states may obscure important process-level insights, which remain crucial for comprehen-
sive understanding. Having reproduced known ERFaer results with a global convection-permitting model opens opportunities
to leverage these models for rigorous process-level studies through carefully designed experiments targeting specific aerosol–
cloud interaction pathways. It also sets the stage for a detailed comparison against the soon-to-be-available interactive aerosol
scheme for the SCREAM configuration, as well as an iterative process to improve both the prescribed and interactive schemes.

345 *Code and data availability.* The default E3SM v3 code is publicly available at <https://github.com/E3SM-Project/E3SM> (specifically commit
d9553ab is used in this study; last access: 30 December 2024). The SCREAM v1 code is available at <https://github.com/E3SM-Project/scream>
(specifically commit 29bdb81 is used in this study; last access: 30 December 2024). We additionally provide a Zenodo archival repository
with the following: (1) the SCREAM v1 code; (2) the default E3SM v3 code for E3SM-MAM runs; (3) the modified E3SM v3 code
for E3SM-SPA runs; (4) the code to reproduce the plots in the manuscripts; (5) instructions to reproduce the runs in the manuscript; (6)
350 instructions to retrieve the data files produced by the simulations in this study, which have been publicly long-term archived using NERSC
HPSS. The archival repository is available at <https://doi.org/10.5281/zenodo.15224730> (Mahfouz, 2025).



Author contributions. All authors contributed to the study design, model development, model runs, data analysis, and/or manuscript writing.

Competing interests. At least one of the (co-)authors is a member of the editorial board of *Atmospheric Chemistry and Physics*.

Acknowledgements. This research was supported as part of the Energy Exascale Earth System Model (E3SM) project, funded by the U.S. Department of Energy (DOE), Office of Science, Office of Biological and Environmental Research. The Pacific Northwest National Laboratory (PNNL) is operated for DOE by the Battelle Memorial Institute under contract no. DE-AC06-76RLO1830. The data (E3SM v3 simulations) were produced using a high-performance computing cluster provided by the DOE Office of Science and Office of Biological and Environmental Research Earth System Model Development program area of the Earth and Environmental System Modeling program and were operated by the Laboratory Computing Resource Center at Argonne National Laboratory. This research (SCREAM v1 simulations) used resources of the National Energy Research Scientific Computing Center (NERSC), a US Department of Energy Office of Science user facility located at Lawrence Berkeley National Laboratory, operated under contract no. DE-AC02-05CH11231 using NERSC award BER-ERCAP-0027116. Sandia National Laboratories is a multi-mission laboratory managed and operated by National Technology & Engineering Solutions of Sandia, LLC (NTESS), a wholly owned subsidiary of Honeywell International Inc., for the U.S. Department of Energy's National Nuclear Security Administration (DOE/NNSA) under contract DE-NA0003525. This written work is authored by an employee of NTESS. The employee, not NTESS, owns the right, title and interest in and to the written work and is responsible for its contents. Any subjective views or opinions that might be expressed in the written work do not necessarily represent the views of the U.S. Government. The publisher acknowledges that the U.S. Government retains a non-exclusive, paid-up, irrevocable, world-wide license to publish or reproduce the published form of this written work or allow others to do so, for U.S. Government purposes. The DOE will provide public access to results of federally sponsored research in accordance with the DOE Public Access Plan. This paper describes objective technical results and analysis. Any subjective views or opinions that might be expressed in the paper do not necessarily represent the views of the U.S. Department of Energy or the United States Government. This paper was performed under the auspices of the U.S. Department of Energy by Lawrence Livermore National Laboratory under Contract DE-AC52-07NA27344.



References

- Abdul-Razzak, H. and Ghan, S. J.: A parameterization of aerosol activation: 2. Multiple aerosol types, *Journal of Geophysical Research: Atmospheres*, 105, 6837–6844, <https://doi.org/10.1029/1999JD901161>, 2000.
- Ackerman, A. S., Kirkpatrick, M. P., Stevens, D. E., and Toon, O. B.: The impact of humidity above stratiform clouds on indirect aerosol climate forcing, *Nature*, 432, 1014–1017, <https://doi.org/10.1038/nature03174>, 2004.
- Adams, P. J. and Seinfeld, J. H.: Predicting global aerosol size distributions in general circulation models, *Journal of Geophysical Research: Atmospheres*, 107, <https://doi.org/10.1029/2001JD001010>, 2002.
- Adams, P. J., Donahue, N. M., and Pandis, S. N.: Atmospheric nanoparticles and climate change, *AIChE Journal*, 59, 4006–4019, <https://doi.org/10.1002/aic.14242>, 2013.
- Albrecht, B. A.: Aerosols, Cloud Microphysics, and Fractional Cloudiness, *Science*, 245, 1227–1230, <https://doi.org/10.1126/science.245.4923.1227>, 1989.
- Andreae, M. O.: Correlation between cloud condensation nuclei concentration and aerosol optical thickness in remote and polluted regions, *Atmospheric Chemistry and Physics*, 9, 543–556, <https://doi.org/10.5194/acp-9-543-2009>, 2009.
- Bellouin, N., Quaas, J., Gryspeerdt, E., Kinne, S., Stier, P., Watson-Parris, D., Boucher, O., Carslaw, K. S., Christensen, M., Daniau, A., Dufresne, J., Feingold, G., Fiedler, S., Forster, P., Gettelman, A., Haywood, J. M., Lohmann, U., Malavelle, F., Mauritsen, T., McCoy, D. T., Myhre, G., Mülmenstädt, J., Neubauer, D., Possner, A., Rugenstein, M., Sato, Y., Schulz, M., Schwartz, S. E., Sourdeval, O., Storelvmo, T., Toll, V., Winker, D., and Stevens, B.: Bounding Global Aerosol Radiative Forcing of Climate Change, *Reviews of Geophysics*, 58, e2019RG000 660, <https://doi.org/10.1029/2019RG000660>, 2020.
- Bogenschutz, P. A. and Krueger, S. K.: A simplified PDF parameterization of subgrid-scale clouds and turbulence for cloud-resolving models, *Journal of Advances in Modeling Earth Systems*, 5, 195–211, <https://doi.org/10.1002/jame.20018>, 2013.
- Bretherton, C. S., Blossey, P. N., and Uchida, J.: Cloud droplet sedimentation, entrainment efficiency, and subtropical stratocumulus albedo, *Geophysical Research Letters*, 34, 2006GL027 648, <https://doi.org/10.1029/2006GL027648>, 2007.
- Burrows, S. M., McCluskey, C. S., Cornwell, G., Steinke, I., Zhang, K., Zhao, B., Zawadowicz, M., Raman, A., Kulkarni, G., China, S., Zelenyuk, A., and DeMott, P. J.: Ice-Nucleating Particles That Impact Clouds and Climate: Observational and Modeling Research Needs, *Reviews of Geophysics*, 60, e2021RG000 745, <https://doi.org/10.1029/2021RG000745>, 2022.
- Caldwell, P. M., Terai, C. R., Hillman, B., Keen, N. D., Bogenschutz, P., Lin, W., Beydoun, H., Taylor, M., Bertagna, L., Bradley, A. M., Clevenger, T. C., Donahue, A. S., Eldred, C., Foucar, J., Golaz, J., Guba, O., Jacob, R., Johnson, J., Krishna, J., Liu, W., Pressel, K., Salinger, A. G., Singh, B., Steyer, A., Ullrich, P., Wu, D., Yuan, X., Shpund, J., Ma, H., and Zender, C. S.: Convection-Permitting Simulations With the E3SM Global Atmosphere Model, *Journal of Advances in Modeling Earth Systems*, 13, e2021MS002 544, <https://doi.org/10.1029/2021MS002544>, 2021.
- Carslaw, K., ed.: *Aerosols and Climate*, Elsevier, Amsterdam, Netherlands, 1 edn., ISBN 978-0-12-819766-0, oCLC: on1285708858, 2022.
- Chen, Y.-C., Xue, L., Lebo, Z. J., Wang, H., Rasmussen, R. M., and Seinfeld, J. H.: A comprehensive numerical study of aerosol-cloud-precipitation interactions in marine stratocumulus, *Atmospheric Chemistry and Physics*, 11, 9749–9769, <https://doi.org/10.5194/acp-11-9749-2011>, 2011.
- Clough, S., Shephard, M., Mlawer, E., Delamere, J., Iacono, M., Cady-Pereira, K., Boukabara, S., and Brown, P.: Atmospheric radiative transfer modeling: a summary of the AER codes, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 91, 233–244, <https://doi.org/10.1016/j.jqsrt.2004.05.058>, 2005.



- 410 Donahue, A. S., Caldwell, P. M., Bertagna, L., Beydoun, H., Bogenschutz, P. A., Bradley, A. M., Clevenger, T. C., Foucar, J., Golaz, C., Guba, O., Hannah, W., Hillman, B. R., Johnson, J. N., Keen, N., Lin, W., Singh, B., Sreepathi, S., Taylor, M. A., Tian, J., Terai, C. R., Ullrich, P. A., Yuan, X., and Zhang, Y.: To Exascale and Beyond—the Simple Cloud-Resolving E3sm Atmosphere Model (scream), a Performance Portable Global Atmosphere Model for Cloud-Resolving Scales, *Journal of Advances in Modeling Earth Systems*, 16, e2024MS004314, <https://doi.org/10.1029/2024MS004314>, 2024.
- 415 Donlon, C. J., Martin, M., Stark, J., Roberts-Jones, J., Fiedler, E., and Wimmer, W.: The Operational Sea Surface Temperature and Sea Ice Analysis (OSTIA) system, *Remote Sensing of Environment*, 116, 140–158, <https://doi.org/10.1016/j.rse.2010.10.017>, 2012.
- Dunne, E. M., Gordon, H., Kürten, A., Almeida, J., Duplissy, J., Williamson, C., Ortega, I. K., Pringle, K. J., Adamov, A., Baltensperger, U., Barmet, P., Benduhn, F., Bianchi, F., Breitenlechner, M., Clarke, A., Curtius, J., Dommen, J., Donahue, N. M., Ehrhart, S., Flagan, R. C., Franchin, A., Guida, R., Hakala, J., Hansel, A., Heinritzi, M., Jokinen, T., Kangasluoma, J., Kirkby, J., Kulmala, M., Kupc, A., Lawler, M. J., Lehtipalo, K., Makhmutov, V., Mann, G., Mathot, S., Merikanto, J., Miettinen, P., Nenes, A., Onnela, A., Rap, A., Reddington, C.
- 420 L. S., Riccobono, F., Richards, N. A. D., Rissanen, M. P., Rondo, L., Sarnela, N., Schobesberger, S., Sengupta, K., Simon, M., Sipilä, M., Smith, J. N., Stozkhov, Y., Tomé, A., Tröstl, J., Wagner, P. E., Wimmer, D., Winkler, P. M., Worsnop, D. R., and Carslaw, K. S.: Global atmospheric particle formation from CERN CLOUD measurements, *Science*, 354, 1119–1124, <https://doi.org/10.1126/science.aaf2649>, 2016.
- 425 E3SM Project, D.: Energy Exascale Earth System Model v3.0.0, <https://doi.org/10.11578/E3SM/DC.20240301.3>, language: en, 2024.
- Fiedler, S., Stevens, B., and Mauritsen, T.: On the sensitivity of anthropogenic aerosol forcing to model-internal variability and parameterizing a T womey effect, *Journal of Advances in Modeling Earth Systems*, 9, 1325–1341, <https://doi.org/10.1002/2017MS000932>, 2017.
- Forster, P. M., Richardson, T., Maycock, A. C., Smith, C. J., Samset, B. H., Myhre, G., Andrews, T., Pincus, R., and Schulz, M.: Recommendations for diagnosing effective radiative forcing from climate models for CMIP6, *Journal of Geophysical Research: Atmospheres*, 121,
- 430 <https://doi.org/10.1002/2016JD025320>, 2016.
- Fuchs, N. A.: *The mechanics of aerosols*, Pergamon Press; [distributed in the Western Hemisphere by Macmillan, New York], Oxford, rev. and enl. ed. edn., 1964.
- Fuchs, N. A. and Sutugin, A. G.: *Highly dispersed aerosols*, Ann Arbor Science Publishers, Ann Arbor, ISBN 978-0-250-39996-3, 1970.
- 435 Gelaro, R., McCarty, W., Suárez, M. J., Todling, R., Molod, A., Takacs, L., Randles, C. A., Darmenov, A., Bosilovich, M. G., Reichle, R., Wargan, K., Coy, L., Cullather, R., Draper, C., Akella, S., Buchard, V., Conaty, A., Da Silva, A. M., Gu, W., Kim, G.-K., Koster, R., Lucchesi, R., Merkova, D., Nielsen, J. E., Partyka, G., Pawson, S., Putman, W., Rienecker, M., Schubert, S. D., Sienkiewicz, M., and Zhao, B.: The Modern-Era Retrospective Analysis for Research and Applications, Version 2 (MERRA-2), *Journal of Climate*, 30, 5419–5454, <https://doi.org/10.1175/JCLI-D-16-0758.1>, 2017.
- 440 Gettelman, A., Bardeen, C. G., McCluskey, C. S., Järvinen, E., Stith, J., Bretherton, C., McFarquhar, G., Twohy, C., D’Alessandro, J., and Wu, W.: Simulating Observations of Southern Ocean Clouds and Implications for Climate, *Journal of Geophysical Research: Atmospheres*, 125, e2020JD032619, <https://doi.org/10.1029/2020JD032619>, 2020.
- Ghan, S. J., Abdul-Razzak, H., Nenes, A., Ming, Y., Liu, X., Ovchinnikov, M., Shipway, B., Meskhidze, N., Xu, J., and Shi, X.: Droplet nucleation: Physically-based parameterizations and comparative evaluation: DROPLET NUCLEATION, *Journal of Advances in Modeling Earth Systems*, 3, <https://doi.org/10.1029/2011MS000074>, 2011.
- 445



- Ghosh, P., Evans, K. J., Grosvenor, D. P., Kang, H.-G., Mahajan, S., Xu, M., Zhang, W., and Gordon, H.: Modifying the Abdul-Razzak & Ghan aerosol activation parameterization (version ARG2000) impacts simulated cloud radiative effects (shown in the UK Met Office Unified Model, version 13.0), <https://doi.org/10.5194/egusphere-2024-2423>, 2024.
- Global Modeling And Assimilation Office: MERRA-2 inst3_3d_asm_Nv: 3d,3-Hourly,Instantaneous,Model-Level,Assimilation,Assimilated Meteorological Fields V5.12.4, <https://doi.org/10.5067/WWQSQ8IVFW8>, 2015.
- Golaz, J.-C., Larson, V. E., and Cotton, W. R.: A PDF-Based Model for Boundary Layer Clouds. Part I: Method and Model Description, *Journal of the Atmospheric Sciences*, 59, 3540–3551, [https://doi.org/10.1175/1520-0469\(2002\)059<3540:APBMFB>2.0.CO;2](https://doi.org/10.1175/1520-0469(2002)059<3540:APBMFB>2.0.CO;2), 2002.
- Goto, D., Sato, Y., Yashiro, H., Suzuki, K., Oikawa, E., Kudo, R., Nagao, T. M., and Nakajima, T.: Global aerosol simulations using NICAM.16 on a 14 km grid spacing for a climate study: improved and remaining issues relative to a lower-resolution model, *Geoscientific Model Development*, 13, 3731–3768, <https://doi.org/10.5194/gmd-13-3731-2020>, 2020.
- Gryspeerdt, E., Goren, T., Sourdeval, O., Quaas, J., Mülmenstädt, J., Dipu, S., Unglaub, C., Gettelman, A., and Christensen, M.: Constraining the aerosol influence on cloud liquid water path, *Atmospheric Chemistry and Physics*, 19, 5331–5347, <https://doi.org/10.5194/acp-19-5331-2019>, 2019.
- Gryspeerdt, E., Povey, A. C., Grainger, R. G., Hasekamp, O., Hsu, N. C., Mulcahy, J. P., Sayer, A. M., and Sorooshian, A.: Uncertainty in aerosol–cloud radiative forcing is driven by clean conditions, *Atmospheric Chemistry and Physics*, 23, 4115–4122, <https://doi.org/10.5194/acp-23-4115-2023>, 2023.
- Hasekamp, O. P., Gryspeerdt, E., and Quaas, J.: Analysis of polarimetric satellite measurements suggests stronger cooling due to aerosol–cloud interactions, *Nature Communications*, 10, 5405, <https://doi.org/10.1038/s41467-019-13372-2>, 2019.
- Hassan, T., Zhang, K., Li, J., Singh, B., Zhang, S., Wang, H., and Ma, P.-L.: Impacts of spatial heterogeneity of anthropogenic aerosol emissions in a regionally refined global aerosol–climate model, *Geoscientific Model Development*, 17, 3507–3532, <https://doi.org/10.5194/gmd-17-3507-2024>, 2024.
- Herbert, R., Stier, P., and Dagan, G.: Isolating Large-Scale Smoke Impacts on Cloud and Precipitation Processes Over the Amazon With Convection Permitting Resolution, *Journal of Geophysical Research: Atmospheres*, 126, e2021JD034615, <https://doi.org/10.1029/2021JD034615>, 2021.
- Herbert, R. J., Williams, A. I. L., Weiss, P., Watson-Parris, D., Dingley, E., Klocke, D., and Stier, P.: Isolating aerosol–climate interactions in global kilometre-scale simulations, <https://doi.org/10.5194/egusphere-2024-1689>, 2024.
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., De Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J.: The ERA5 global reanalysis, *Quarterly Journal of the Royal Meteorological Society*, 146, 1999–2049, <https://doi.org/10.1002/qj.3803>, 2020.
- Intergovernmental Panel On Climate Change: Clouds and Aerosols, in: *Climate Change 2013 - The Physical Science Basis*, pp. 571–658, Cambridge University Press, Cambridge, ISBN 978-1-107-41532-4, <https://doi.org/10.1017/CBO9781107415324.016>, 2014.
- Intergovernmental Panel On Climate Change: Climate Change 2021 – The Physical Science Basis: Working Group I Contribution to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change, Cambridge University Press, 1 edn., ISBN 978-1-109-15789-6, <https://doi.org/10.1017/97811009157896>, 2023.



- Karset, I. H. H., Gettelman, A., Storelvmo, T., Alterskjær, K., and Berntsen, T. K.: Exploring Impacts of Size-Dependent Evaporation and Entrainment in a Global Model, *Journal of Geophysical Research: Atmospheres*, 125, e2019JD031817, <https://doi.org/10.1029/2019JD031817>, 2020.
- Kooperman, G. J., Pritchard, M. S., Ghan, S. J., Wang, M., Somerville, R. C. J., and Russell, L. M.: Constraining the influence of natural variability to improve estimates of global aerosol indirect effects in a nudged version of the Community Atmosphere Model 5, *Journal of Geophysical Research: Atmospheres*, 117, 2012JD018588, <https://doi.org/10.1029/2012JD018588>, 2012.
- Kreidenweis, S. M., Petters, M., and Lohmann, U.: 100 Years of Progress in Cloud Physics, Aerosols, and Aerosol Chemistry Research, *Meteorological Monographs*, 59, 11.1–11.72, <https://doi.org/10.1175/AMSMONOGRAPHIS-D-18-0024.1>, 2019.
- Liu, X., Easter, R. C., Ghan, S. J., Zaveri, R., Rasch, P., Shi, X., Lamarque, J.-F., Gettelman, A., Morrison, H., Vitt, F., Conley, A., Park, S., Neale, R., Hannay, C., Ekman, A. M. L., Hess, P., Mahowald, N., Collins, W., Iacono, M. J., Bretherton, C. S., Flanner, M. G., and Mitchell, D.: Toward a minimal representation of aerosols in climate models: description and evaluation in the Community Atmosphere Model CAM5, *Geoscientific Model Development*, 5, 709–739, <https://doi.org/10.5194/gmd-5-709-2012>, 2012.
- Mahfouz, N.: mahf708/repros: tmp-eamxx-acp-erfaer-rel, <https://doi.org/10.5281/zenodo.15224730>, 2025.
- Mahfouz, N., Mülmenstädt, J., and Burrows, S.: Present-day correlations are insufficient to predict cloud albedo change by anthropogenic aerosols in E3SM v2, *Atmospheric Chemistry and Physics*, 24, 7253–7260, <https://doi.org/10.5194/acp-24-7253-2024>, 2024.
- McCoy, D. T., Bender, F. A., Mohrmann, J. K. C., Hartmann, D. L., Wood, R., and Grosvenor, D. P.: The global aerosol-cloud first indirect effect estimated using MODIS, MERRA, and AeroCom, *Journal of Geophysical Research: Atmospheres*, 122, 1779–1796, <https://doi.org/10.1002/2016JD026141>, 2017.
- McCoy, D. T., Field, P., Gordon, H., Elsaesser, G. S., and Grosvenor, D. P.: Untangling causality in midlatitude aerosol–cloud adjustments, *Atmospheric Chemistry and Physics*, 20, 4085–4103, <https://doi.org/10.5194/acp-20-4085-2020>, 2020.
- Ming, Y., Ramaswamy, V., Donner, L. J., and Phillips, V. T. J.: A New Parameterization of Cloud Droplet Activation Applicable to General Circulation Models, *Journal of the Atmospheric Sciences*, 63, 1348–1356, <https://doi.org/10.1175/JAS3686.1>, 2006.
- Moore, R. H., Karydis, V. A., Capps, S. L., Latham, T. L., and Nenes, A.: Droplet number uncertainties associated with CCN: an assessment using observations and a global model adjoint, *Atmospheric Chemistry and Physics*, 13, 4235–4251, <https://doi.org/10.5194/acp-13-4235-2013>, 2013.
- Morales Betancourt, R. and Nenes, A.: Droplet activation parameterization: the population-splitting concept revisited, *Geoscientific Model Development*, 7, 2345–2357, <https://doi.org/10.5194/gmd-7-2345-2014>, 2014.
- Morrison, H. and Milbrandt, J. A.: Parameterization of Cloud Microphysics Based on the Prediction of Bulk Ice Particle Properties. Part I: Scheme Description and Idealized Tests, *Journal of the Atmospheric Sciences*, 72, 287–311, <https://doi.org/10.1175/JAS-D-14-0065.1>, 2015.
- Mülmenstädt, J. and Wilcox, L. J.: The Fall and Rise of the Global Climate Model, *Journal of Advances in Modeling Earth Systems*, 13, e2021MS002781, <https://doi.org/10.1029/2021MS002781>, 2021.
- Mülmenstädt, J., Gryspeerdt, E., Salzmann, M., Ma, P.-L., Dipu, S., and Quaas, J.: Separating radiative forcing by aerosol–cloud interactions and rapid cloud adjustments in the ECHAM–HAMMOZ aerosol–climate model using the method of partial radiative perturbations, *Atmospheric Chemistry and Physics*, 19, 15415–15429, <https://doi.org/10.5194/acp-19-15415-2019>, 2019.
- Mülmenstädt, J., Ackerman, A. S., Fridlind, A. M., Huang, M., Ma, P.-L., Mahfouz, N., Bauer, S. E., Burrows, S. M., Christensen, M. W., Dipu, S., Gettelman, A., Leung, L. R., Tornow, F., Quaas, J., Varble, A. C., Wang, H., Zhang, K., and Zheng, Y.: Can general circulation



- 520 models (GCMs) represent cloud liquid water path adjustments to aerosol–cloud interactions?, *Atmospheric Chemistry and Physics*, 24, 13 633–13 652, <https://doi.org/10.5194/acp-24-13633-2024>, 2024a.
- Mülmenstädt, J., Gryspeerdt, E., Dipu, S., Quaas, J., Ackerman, A. S., Fridlind, A. M., Tornow, F., Bauer, S. E., Gettelman, A., Ming, Y., Zheng, Y., Ma, P.-L., Wang, H., Zhang, K., Christensen, M. W., Varble, A. C., Leung, L. R., Liu, X., Neubauer, D., Partridge, D. G., Stier, P., and Takemura, T.: General circulation models simulate negative liquid water path–droplet number correlations, but anthropogenic
525 aerosols still increase simulated liquid water path, *Atmospheric Chemistry and Physics*, 24, 7331–7345, <https://doi.org/10.5194/acp-24-7331-2024>, 2024b.
- Nenes, A. and Seinfeld, J. H.: Parameterization of cloud droplet formation in global climate models, *Journal of Geophysical Research: Atmospheres*, 108, 2002JD002 911, <https://doi.org/10.1029/2002JD002911>, 2003.
- Pincus, R., Mlawer, E. J., and Delamere, J. S.: Balancing Accuracy, Efficiency, and Flexibility in Radiation Calculations for Dynamical
530 Models, *Journal of Advances in Modeling Earth Systems*, 11, 3074–3089, <https://doi.org/10.1029/2019MS001621>, 2019.
- Quaas, J., Arola, A., Cairns, B., Christensen, M., Deneke, H., Ekman, A. M. L., Feingold, G., Fridlind, A., Gryspeerdt, E., Hasekamp, O., Li, Z., Lipponen, A., Ma, P.-L., Mülmenstädt, J., Nenes, A., Penner, J. E., Rosenfeld, D., Schrödner, R., Sinclair, K., Sourdeval, O., Stier, P., Tesche, M., Van Diedenhoven, B., and Wendisch, M.: Constraining the Twomey effect from satellite observations: issues and perspectives, *Atmospheric Chemistry and Physics*, 20, 15 079–15 099, <https://doi.org/10.5194/acp-20-15079-2020>, 2020.
- 535 Quaas, J., Andrews, T., Bellouin, N., Block, K., Boucher, O., Ceppi, P., Dagan, G., Doktorowski, S., Eichholz, H. M., Forster, P., Goren, T., Gryspeerdt, E., Hodnebrog, O., Jia, H., Kramer, R., Lange, C., Maycock, A. C., Mülmenstädt, J., Myhre, G., O'Connor, F. M., Pincus, R., Samset, B. H., Senf, F., Shine, K. P., Smith, C., Stjern, C. W., Takemura, T., Toll, V., and Wall, C. J.: Adjustments to Climate Perturbations—Mechanisms, Implications, Observational Constraints, *AGU Advances*, 5, e2023AV001 144, <https://doi.org/10.1029/2023AV001144>, 2024.
- 540 Reutter, P., Su, H., Trentmann, J., Simmel, M., Rose, D., Gunthe, S. S., Wernli, H., Andreae, M. O., and Pöschl, U.: Aerosol- and updraft-limited regimes of cloud droplet formation: influence of particle number, size and hygroscopicity on the activation of cloud condensation nuclei (CCN), *Atmospheric Chemistry and Physics*, 9, 7067–7080, <https://doi.org/10.5194/acp-9-7067-2009>, 2009.
- Rothenberg, D., Avramov, A., and Wang, C.: On the representation of aerosol activation and its influence on model-derived estimates of the aerosol indirect effect, *Atmospheric Chemistry and Physics*, 18, 7961–7983, <https://doi.org/10.5194/acp-18-7961-2018>, 2018.
- 545 Sato, Y., Goto, D., Michibata, T., Suzuki, K., Takemura, T., Tomita, H., and Nakajima, T.: Aerosol effects on cloud water amounts were successfully simulated by a global cloud-system resolving model, *Nature Communications*, 9, 985, <https://doi.org/10.1038/s41467-018-03379-6>, 2018.
- Seifert, A. and Beheng, K. D.: A two-moment cloud microphysics parameterization for mixed-phase clouds. Part 1: Model description, *Meteorology and Atmospheric Physics*, 92, 45–66, <https://doi.org/10.1007/s00703-005-0112-4>, 2006.
- 550 Seinfeld, J. H. and Pandis, S. N.: *Atmospheric chemistry and physics: from air pollution to climate change*, John Wiley & Sons, Hoboken, New Jersey, third edition edn., ISBN 978-1-119-22116-6 978-1-119-22117-3, 2016.
- Shan, Y., Fan, J., Zhang, K., Shpund, J., Terai, C., Zhang, G. J., Song, X., Chen, C., Lin, W., Liu, X., Shrivastava, M., Wang, H., and Xie, S.: Improving Aerosol Radiative Forcing and Climate in E3SM: Impacts of New Cloud Microphysics and Improved Wet Removal Treatments, *Journal of Advances in Modeling Earth Systems*, 16, e2023MS004 059, <https://doi.org/10.1029/2023MS004059>, 2024.
- 555 Stevens, B., Fiedler, S., Kinne, S., Peters, K., Rast, S., Müsse, J., Smith, S. J., and Mauritsen, T.: MACv2-SP: a parameterization of anthropogenic aerosol optical properties and an associated Twomey effect for use in CMIP6, *Geoscientific Model Development*, 10, 433–452, <https://doi.org/10.5194/gmd-10-433-2017>, 2017.



- Stier, P., Feichter, J., Kinne, S., Kloster, S., Vignati, E., Wilson, J., Ganzeveld, L., Tegen, I., Werner, M., Balkanski, Y., Schulz, M., Boucher, O., Minikin, A., and Petzold, A.: The aerosol-climate model ECHAM5-HAM, *Atmospheric Chemistry and Physics*, 5, 1125–1156, <https://doi.org/10.5194/acp-5-1125-2005>, 2005.
- Terai, C. R., Pritchard, M. S., Blosssey, P., and Bretherton, C. S.: The Impact of Resolving Subkilometer Processes on Aerosol-Cloud Interactions of Low-Level Clouds in Global Model Simulations, *Journal of Advances in Modeling Earth Systems*, 12, e2020MS002274, <https://doi.org/10.1029/2020MS002274>, 2020.
- Terai, C. R., Xie, S., Song, X., Chen, C.-C., Fan, J., Zhang, G. J., Richter, J. H., Shpund, K., Lin, W., Golaz, J.-C., Larson, V. E., Moncrieff, M. W., Shan, Y., Zhang, C., Zhang, K., and Zhang, Y.: Impact of microphysics and convection schemes on the mean-state and variability of clouds and precipitation in the E3SM Atmosphere Model, <https://doi.org/10.22541/au.172252420.09883031/v1>, 2024.
- Terai, C. R., Keen, N. D., Caldwell, P. M., Beydoun, H., Bogenschutz, P. A., Chao, L.-W., Hillman, B. R., Ma, H.-Y., Zelinka, M. D., Bertagna, L., Bradley, A., Clevenger, T. C., Donahue, A. S., Foucar, J. G., Golaz, J.-C., Guba, O., Hannah, W. M., Lee, J., Lin, W., Mahfouz, N. G. A., Mülmenstädt, J., Salinger, A. G., Singh, B., Sreepathi, S., Qin, Y., Taylor, M. A., Ullrich, P. A., Wu, W.-Y., Yuan, X., Zender, C. S., and Zhang, Y.: Climate Response to Warming in Cess-Potter Simulations Using the Global 3-km SCREAM, <https://doi.org/10.22541/essoar.173655643.33295443/v1>, 2025.
- Tiedtke, M., Geleyn, J.-F., Hollingsworth, A., and Louis, J.-F.: ECMWF model parameterisation of sub-grid scale processes, Technical Report, ECMWF, Shinfield Park, Reading, <https://www.ecmwf.int/node/12743>, 10, 1979.
- Tilmes, S., Mills, M. J., Zhu, Y., Bardeen, C. G., Vitt, F., Yu, P., Fillmore, D., Liu, X., Toon, B., and Deshler, T.: Description and performance of a sectional aerosol microphysical model in the Community Earth System Model (CESM2), *Geoscientific Model Development*, 16, 6087–6125, <https://doi.org/10.5194/gmd-16-6087-2023>, 2023.
- Topping, D. and Bane, M., eds.: Introduction to aerosol modelling: from theory to code, John Wiley & Sons, Hoboken, NJ, ISBN 978-1-119-62565-0, 2022.
- Twomey, S.: The nuclei of natural cloud formation part II: The supersaturation in natural clouds and the variation of cloud droplet concentration, *Geofisica pura e applicata*, 43, 243–249, <https://doi.org/10.1007/BF01993560>, 1959.
- Twomey, S.: The Influence of Pollution on the Shortwave Albedo of Clouds, *Journal of the Atmospheric Sciences*, 34, 1149–1152, [https://doi.org/10.1175/1520-0469\(1977\)034<1149:TIOPOT>2.0.CO;2](https://doi.org/10.1175/1520-0469(1977)034<1149:TIOPOT>2.0.CO;2), 1977.
- Twomey, S.: Aerosols, clouds and radiation, *Atmospheric Environment. Part A. General Topics*, 25, 2435–2442, [https://doi.org/10.1016/0960-1686\(91\)90159-5](https://doi.org/10.1016/0960-1686(91)90159-5), 1991.
- UKMO: GHRSSST Level 4 OSTIA Global Foundation Sea Surface Temperature Analysis (GDS version 2), <https://doi.org/10.5067/GHOST-4FK02>, 2012.
- Varble, A. C., Ma, P.-L., Christensen, M. W., Mülmenstädt, J., Tang, S., and Fast, J.: Evaluation of liquid cloud albedo susceptibility in E3SM using coupled eastern North Atlantic surface and satellite retrievals, *Atmospheric Chemistry and Physics*, 23, 13 523–13 553, <https://doi.org/10.5194/acp-23-13523-2023>, 2023.
- Viovy, N.: CRUNCEP Version 7 - Atmospheric Forcing Data for the Community Land Model, <https://doi.org/10.5065/PZ8F-F017>, artwork Size: 1.238 Tbytes Pages: 1.238 Tbytes, 2018.
- Wan, H., Yenpure, A., Geveci, B., Easter, R. C., Rasch, P. J., Zhang, K., and Zeng, X.: Features of mid- and high-latitude low-level clouds and their relation to strong aerosol effects in the Energy Exascale Earth System Model version 2 (E3SMv2), <https://doi.org/10.5194/egusphere-2024-4020>, 2025.



- 595 Wang, H., Easter, R. C., Zhang, R., Ma, P., Singh, B., Zhang, K., Ganguly, D., Rasch, P. J., Burrows, S. M., Ghan, S. J., Lou, S., Qian, Y., Yang, Y., Feng, Y., Flanner, M., Leung, L. R., Liu, X., Shrivastava, M., Sun, J., Tang, Q., Xie, S., and Yoon, J.: Aerosols in the E3SM Version 1: New Developments and Their Impacts on Radiative Forcing, *Journal of Advances in Modeling Earth Systems*, 12, e2019MS001851, <https://doi.org/10.1029/2019MS001851>, 2020.
- Weiss, P., Herbert, R., and Stier, P.: ICON-HAM-lite: simulating the Earth system with interactive aerosols at kilometer scales, <https://doi.org/10.5194/egusphere-2024-3325>, 2024.
- 600 Zelinka, M. D., Smith, C. J., Qin, Y., and Taylor, K. E.: Comparison of methods to estimate aerosol effective radiative forcings in climate models, *Atmospheric Chemistry and Physics*, 23, 8879–8898, <https://doi.org/10.5194/acp-23-8879-2023>, 2023.
- Zhang, G. and McFarlane, N. A.: Sensitivity of climate simulations to the parameterization of cumulus convection in the Canadian climate centre general circulation model, *Atmosphere-Ocean*, 33, 407–446, <https://doi.org/10.1080/07055900.1995.9649539>, 1995.
- 605 Zhang, K., Wan, H., Liu, X., Ghan, S. J., Kooperman, G. J., Ma, P.-L., Rasch, P. J., Neubauer, D., and Lohmann, U.: Technical Note: On the use of nudging for aerosol–climate model intercomparison studies, *Atmospheric Chemistry and Physics*, 14, 8631–8645, <https://doi.org/10.5194/acp-14-8631-2014>, 2014.
- Zhang, K., Zhang, W., Wan, H., Rasch, P. J., Ghan, S. J., Easter, R. C., Shi, X., Wang, Y., Wang, H., Ma, P.-L., Zhang, S., Sun, J., Burrows, S. M., Shrivastava, M., Singh, B., Qian, Y., Liu, X., Golaz, J.-C., Tang, Q., Zheng, X., Xie, S., Lin, W., Feng, Y., Wang, M., Yoon, J.-H., and
- 610 Leung, L. R.: Effective radiative forcing of anthropogenic aerosols in E3SM version 1: historical changes, causality, decomposition, and parameterization sensitivities, *Atmospheric Chemistry and Physics*, 22, 9129–9160, <https://doi.org/10.5194/acp-22-9129-2022>, 2022a.
- Zhang, S., Zhang, K., Wan, H., and Sun, J.: Further improvement and evaluation of nudging in the E3SM Atmosphere Model version 1 (EAMv1): simulations of the mean climate, weather events, and anthropogenic aerosol effects, *Geoscientific Model Development*, 15, 6787–6816, <https://doi.org/10.5194/gmd-15-6787-2022>, 2022b.