

Development of the global maize yield model MATCRO-Maize version 1.0

Marin Nagata¹, Astrid Yusara², Tomomichi Kato^{3*}, Yuji Masutomi⁴

¹Graduate School of Global Food Resources, Hokkaido University, Sapporo, Hokkaido 060-0809, Japan

²Graduate School of Agriculture, Hokkaido University, Sapporo, Hokkaido 060-0808, Japan

³Research Faculty of Agriculture, Hokkaido University, Sapporo, Hokkaido 060-8589, Japan

⁴Center for Climate Change Adaptation, National Institute for Environmental Studies, Tsukuba, Ibaraki 305-8506, Japan

Correspondence to: Tomomichi Kato (tkato@agr.hokudai.ac.jp)

Abstract. Process-based crop models combined with land surface models are useful tools for accurately quantifying the impacts of climate change on crops while considering the interactions between agricultural land and climate. We developed a new process-based crop model for maize, named MATCRO-Maize, by incorporating leaf-level photosynthesis of C4 plants and adjusting crop-specific parameters into the original MATCRO model, which is a process-based crop model initially developed for paddy rice combined with a land surface model. The model was evaluated at both a point scale and a global scale through comparisons with observational values. The evaluation at the point scale was conducted at four globally distributed sites based on global parameters. It showed statistically significant correlation for final yield with correlation coefficient (COR) of 0.34 with a p value < 0.01. For the global scale evaluation, the simulated yield was statistically compared with the reference data at the country level and total global level. Although the absolute value of the simulated yield tended to be overestimated, MATCRO-Maize could capture spatial variability, as indicated by a COR of 0.58 (p value < 0.01) for the 30-year average yield comparison of the top 20 maize-producing countries. In addition, the comparisons of the interannual variability derived from detrended deviation were statistically significant for the total global yield (COR of 0.54 with p value < 0.01) and for half of the top 20 countries (COR of 0.64-0.90 with p value < 0.001 for 6 countries; COR of 0.50-0.51 with p value < 0.01 for 2 countries; COR of 0.48-0.55 with p value < 0.05 for 2 countries), which are comparable with those of other global crop models. One of the reasons for this overestimation could be related to the strong nitrogen fertilization effect observed in MATCRO-Maize. With experimental field data under more comprehensive conditions, improvements in the functions of nitrogen fertilizer in the model would be needed to simulate the maize yield more accurately.

1 Introduction

Maize (*Zea mays* L.) is one of the most important cereals not only because of its large production (FAO, 2022) but also because of its various roles in human food, feed, and industrial uses. Maize has high photosynthetic efficiency as a C4 plant. It contains

Style Definition: Strong

Deleted: production

Deleted: valida

Deleted: validation

Deleted: three variables (leaf area index: correlation coefficient (COR) of 0.76 with a p value < 0.01; total aboveground biomass: COR of 0.89 with a p value < 0.001;

Deleted: : COR

Deleted:)

Deleted: validation

Deleted: FAOSTAT

phosphoenolpyruvate (PEP) carboxylase in mesophyll cells, which concentrates CO₂ in bundle sheath cells. The concentrated CO₂ increases the relative amount of carboxylation versus oxygenation performed by ribulose-1,5-bisphosphate carboxylase/oxygenase (Rubisco) (Kanai and Edwards, 1999), allowing C4 plants to operate at lower stomatal conductance rates than C3 plants (Sage, 1999). This mechanism results in high efficiencies of light, water, and nitrogen use (Knapp and Medina, 1999; Long, 1999). These features, such as multipurpose crops and high photosynthetic efficiency, enable the cultivated area to range over wide environments from wet to dry and from low to midlatitude. However, climate change impacts and climate-related extremes negatively affect the productivity of the agricultural sector, which leads to negative consequences for food security (Intergovernmental Panel on Climate Change (IPCC), 2023). Therefore, it is important to accurately quantify the impact of climate change on crop growth and yield and to identify effective adaptation strategies to mitigate climate risk.

Process-based crop models are useful tools for climate change studies because they consider the response of the physiological processes of crop growth and development to the environment and management (Tubiello and Ewert, 2002). The ensemble of process-based crop model simulations has shown good agreement with observed maize yields both at the site scale and at the global scale (Bassu et al., 2014; Jägermeyr et al., 2021), showing its potential to quantify the uncertainty in studies on the impacts of climate change on crop yields (Asseng et al., 2013). Crop models combined with land surface models (LSMs) or earth system models (ESMs) (as classified by Peng et al., 2017) have the ability to consider the effects of agricultural land on the climate globally through the exchange of fluxes of heat, water, and gases, as well as the effects of climate on crops. Some studies have revealed that agricultural land affects the climate through fluxes (Bondeau et al., 2007; Levis et al., 2012; Maruyama and Kuwagata, 2010; Tsvetsinskaya et al., 2001) and subsequently affects crop production (Osborne et al., 2009). This indicates the importance of considering the interaction between agricultural land and climate to accurately quantify the impacts of climate change on crops. Despite this importance, few LSM/ESM-based crop models exist (Lin et al., 2021; Lombardozzi et al., 2020; Osborne et al., 2015; Wu et al., 2016).

MATCRO is a process-based crop growth model developed for C3 plants (Masutomi et al., 2016a, b; Yusara et al., in prep). It was initially combined with a land surface model of Minimal Advanced Treatments of Surface Interaction and Runoff, called MATSIRO (Takata et al., 2003). MATSIRO is embedded in an earth system model, which is the Model for Interdisciplinary Research on Climate, Earth System version 2 for Long-term simulations called MIROC-ES2L (Hajima et al., 2020). MATCRO simulates crop growth based on leaf-level photosynthesis and parameterized crop-specific parameters determined from experimental data, and can run simulations both at a point scale and at a global scale. The model was applied to assess the impact of climate change at the country and local levels (Kinose et al., 2020; Kinose and Masutomi, 2019) and was used in the study investigating factors to improve the simulation performance of global gridded crop models (GGCMs) (Iizumi et al., 2021). MATCRO is applicable to other crops, including maize as a C4 plant, with adjusted parameters from experimental datasets and the literature.

We extended MATCRO for global maize yield simulation, called MATCRO-Maize, by adjusting crop-specific parameters for maize and incorporating the C4 photosynthetic mechanism. MATCRO-Rice can simulate latent heat flux, sensible heat flux, net carbon uptake by crops, and rice yield, indicating its application in studies on climate change impacts

75 as an LSM-based model (Masutomi et al. 2016b). However, this study focused only on crop growth and yields, omitting water and heat fluxes to increase computational efficiency. This paper aims to describe MATCRO-Maize in detail (Section 2) and model [evaluation](#) on simulated yields both at a point scale and at a global scale (Section 3), with a discussion of the [evaluation](#) and model limitations (Section 4).

Deleted: validation

Deleted: alidatio

Deleted: n

2. Model description

80 MATCRO consists of four modules: radiation, net carbon assimilation, crop growth, and soil water balance. It requires the following input data: (i) phenological data (i.e., crop calendar), (ii) water management data (i.e., the land is rainfed or irrigated), (iii) nitrogen fertilizer application data (N_{fert}) [kg N ha^{-1}], (iv) soil classification data (i.e., soil texture classification), (v) annual CO_2 data [ppm], and (vi) 6 types of daily meteorological data: air pressure (P_s) [Pa], precipitation (P_{rc}) [$\text{kg m}^{-2} \text{s}^{-1}$], specific humidity [S_h] [kg kg^{-1}], downwards shortwave radiation (R_s) [W m^{-2}], maximum, minimum, and mean air temperature (T_{max} , T_{min} , T_a) [K], and wind speed (U) [m s^{-1}]. Based on input data, MATCRO simulates crop growth during a growing period. It is controlled by the crop developmental stage (D_{vs}) based on (Bouman et al., 2001), which is the index used to quantify crop development. The final crop yield is determined by the dry weight of the storage organ with a parameter (K_{yld}) when $D_{vs} = 1$. To adapt MATCRO for maize, crop-specific parameters and equations were improved, as shown in Table 1 and Eq. (1)–(35). The details are described in the following sections.

90 2.1 Photosynthetic mechanism

MATCRO-Maize calculates net carbon assimilation for the entire canopy (A_n) via the big-leaf model, where [C4](#) leaf-level photosynthesis is separately calculated for sunlit and shaded leaves from the coupled photosynthesis–stomatal conductance model (Dai et al., 2004).

Deleted: (Dai et al., 2004)

Deleted: C3

Deleted: model (Dai(Collatz et al., 1991).2004).

A_n for the entire canopy is given by:

$$95 \quad A_n = \bar{A}_{n,sn} L_{sn} + \bar{A}_{n,sh} L_{sh}, \quad (1)$$

where $\bar{A}_{n,sn}$ and $\bar{A}_{n,sh}$ represent the net carbon assimilation per unit leaf area [$\mu \text{mol m}^{-2} \text{s}^{-1}$] and where L_{sn} and L_{sh} represent the leaf area index (LAI) [$\text{m}^2 (\text{leaf})^{-2}$]. sn and sh indicate sunlit and shaded leaves, respectively. $\bar{A}_{n,sn}$ and $\bar{A}_{n,sh}$ are defined in the following equations:

$$100 \quad \bar{A}_{n,x} = \bar{A}_{g,x} - \bar{R}_{d,x}, \quad (2)$$

where $\bar{A}_{g,x}$ and $\bar{R}_{d,x}$ represent gross carbon assimilation and dark respiration per unit leaf area [$\mu \text{mol m}^{-2} \text{s}^{-1}$], respectively.

Suffix x means sn or sh . L_{sn} and L_{sh} are determined following the approach of Masutomi et al., (2016a). $\bar{R}_{d,x}$ is calculated via the following equation (Bonan et al., 2011):

$$110 \quad \bar{R}_{d,x} = 0.025 V_{cmax,x} \left(\frac{2^{(T_v-298.15)/10}}{1+\exp(1.3(T_v-328.15))} \right), \quad (3)$$

where $V_{cmax,x}$ [μ mol m^{-2} s^{-1}] is the maximum rate of carboxylation and where T_v is the leaf temperature [K] (assumed to be the same as the air temperature: T_a). $\bar{A}_{g,x}$ is determined by the smaller root of the following equations:

$$\beta_{cj} \bar{A}_{i,x}^2 - (\bar{A}_{c,x} + \bar{A}_{j,x}) \bar{A}_{i,x} + \bar{A}_{c,x} \bar{A}_{j,x} = 0, \quad (4)$$

$$\beta_{ip} \bar{A}_{g,x}^2 - (\bar{A}_{i,x} + \bar{A}_{p,x}) \bar{A}_{g,x} + \bar{A}_{i,x} \bar{A}_{p,x} = 0, \quad (5)$$

115 where β_{cj} and β_{ip} are the transition factors (Table 1) and where $\bar{A}_{i,x}$ [μ mol m^{-2} s^{-1}] is the **carbon fixation rate**. Here, we introduced the C4 leaf-level photosynthesis model based on Collatz et al., (1992) into MATCRO, in which some parameters were taken from Oleson et al., (2013) and Lawrence et al., (2020) (Table 1). In C4 photosynthesis, $\bar{A}_{c,x}$, $\bar{A}_{j,x}$, and $\bar{A}_{p,x}$ [μ mol m^{-2} s^{-1}] represent Rubisco-limited, RUBP-limited, and PEP-limited photosynthesis, respectively, and are given by the following equations:

$$120 \quad \bar{A}_{c,x} = V_{cmax,x}, \quad (6)$$

$$\bar{A}_{j,x} = \alpha(4.6Q_{ab,x}), \quad (7)$$

$$\bar{A}_{p,x} = k_{p,x}C_{i,x}, \quad (8)$$

where $C_{i,x}$ [ppm] is the internal leaf CO_2 concentration, $Q_{ab,x}$ [$W m^{-2}$] is the absorbed photosynthetically active radiation (PAR), α [$mol mol^{-1}$] is the quantum efficiency, and $k_{p,x}$ [$mol m^{-2} s^{-1}$] is the initial slope of the CO_2 response curve for the C4 CO_2 response curve. $Q_{ab,x}$ is calculated from R_s via the same methods as in Masutomi et al. (2016a) and is converted to photosynthetic photon flux by multiplying by 4.6 [μ mol (photons) J^{-1}]. $V_{cmax,x}$ and $k_{p,x}$ are functions of T_v and are based on Lawrence et al. (2020),

$$V_{cmax,x} = f_v V_{cmax25,x} \left[\frac{Q_{10}^{(T_v-298.15)/10}}{f_H(T_v)f_L(T_v)} \right], \quad (9)$$

$$f_H(T_v) = 1 + \exp[S_1(T_v - S_2)], \quad (10)$$

$$130 \quad f_L(T_v) = 1 + \exp[S_3(S_4 - T_v)], \quad (11)$$

$$k_{p,x} = \begin{cases} k_{p25,x} Q_{10}^{(T_v-298.15)/10}, & V_{cmax25,x} > 0, \\ 0.7, & V_{cmax25,x} = 0, \end{cases} \quad (12)$$

$$k_{p25,x} = 20000 V_{cmax25,x}, \quad (13)$$

with $Q_{10} = 2$, $S_1 = 0.3 K^{-1}$, $S_2 = 313.15 K$, $S_3 = 0.2 K^{-1}$, and $S_4 = 288.15 K$ (Table 1). Notably, $k_{p,x}$ is adjusted to be **Formatted: Justified** **Deleted: 7μ** $0.7 mol m^{-2} s^{-1}$ (Collatz et al., 1992) when $V_{cmax25,x} = 0$ because of the process of the photosynthesis calculation (Eq. (20)).

135 $V_{cmax25,x}$ is the maximum Rubisco carboxylation rate per unit leaf area at $25^\circ C$ (the details are described in Section 2.2.2). f_v is the water stress factor calculated in the soil water balance module, which indirectly affects A_n through $V_{cmax,x}$ (Sellers et al., 1996). f_v is derived from the following equations:

Deleted: colimited photosynthesis.

Deleted: μ

Deleted: $\begin{cases} k_{p25,x} Q_{10}^{(T_v-298.15)/10}, & V_{cmax25,x} > 0, \\ 0.7\mu, & V_{cmax25,x} = 0, \end{cases}$

Formatted: Justified

Deleted: 7μ

$$f_v = \sum_{i=1}^{NSL} \begin{cases} 1 * ETF(i), & FAW(i) > 0.45, \\ \frac{FAW(i)}{0.45} * ETF(i), & \text{otherwise}, \end{cases} \quad (14)$$

$$FAW(i) = \min\left(\frac{\max((WSL(i) - WILT), 0)}{FC - WILT}, 1\right), \quad (15)$$

$$ETF(i) = \frac{3(z_{rt}^2 - z^2)}{2z_{rt}^3}, \quad (16)$$

145 where NSL represents the number of soil layers, ETF represents the fraction of transpiration from root distribution, FAW represents the fraction of available water, WSL represents the soil water content [$\text{m}^3 \text{m}^{-3}$], $WILT$ represents the wilting point, FC represents the field capacity, and z_{rt} and z represent the root depth and the soil depth, respectively, for each layer. MATCRO assumes $NSL = 5$, where each of the soil layers has thicknesses of 0.05, 0.2, 0.75, 1, and 2 [m], respectively. MATCRO uses the soil texture data as input data, where the soil is classified into 13 types, leading to differences in $WILT$ and FC based on Campbell and Norman (1998). WSL is calculated considering transpiration from the canopy, evaporation from the soil, and water flux (those calculations are the same as those of the original MATCRO). The ETF calculation assumes that the root has no spatial orientation and is equally distributed in the soil (Masutomi et al., 2016a). z_{rt} is determined by the same calculation as the original MATCRO, where the crop-specific parameter ($z_{rt, mx}$) was changed to maize (Table 1). The conditional branch ($FAW(i) > 0.45$) is based on the FAO 56 guidelines (Allen et al., 1998).

155 Stomatal conductance influences CO_2 uptake during photosynthesis. MATCRO-Maize represents stomatal conductance for CO_2 , $G_{sc,x}$ [$\mu \text{mol m}^{-2} \text{s}^{-1}$], based on Ball et al. (1987) as follows:

$$G_{sc,x} = \begin{cases} G_{0c} + G_{1c} R_h \frac{\bar{A}_{n,x}}{C_{s,x}}, & \bar{A}_{n,x} \geq 0, \\ G_{0c}, & \text{otherwise}, \end{cases} \quad (17)$$

160 where $C_{s,x}$ [ppm] is the CO_2 concentration at the leaf surface and where R_h [-] is the relative humidity at the leaf surface. G_{0c} and G_{1c} are derived from parameters b and m (shown in Table 1), respectively, by adjusting their ratio of 1:1.6, which is the ratio of diffusivity of H_2O to CO_2 . Here, the leaf-level net carbon assimilation rate ($\bar{A}_{n,x}$), stomatal conductance for CO_2 ($G_{sc,x}$), and boundary layer conductance for CO_2 (G_{bc}) were calculated to satisfy the following physical flux equations.

$$\bar{A}_{n,x} = G_{sc,x}(C_{s,x} - C_{i,x}), \quad (18)$$

$$\bar{A}_{n,x} = G_{bc}(C_a - C_{s,x}), \quad (19)$$

165 where C_a [ppm] is the atmospheric CO_2 concentration. G_{bc} is a function of air pressure (P_s [Pa]) and the wind speed in the canopy (U [m s^{-1}]) (Masutomi, 2023).

Here, T_v , $Q_{ab,x}$, R_h , U , and C_a are environmental variables derived from input meteorological climate data. There are four relationships (Eqs. (2), (17)-(19)) in terms of internal variables ($\bar{A}_{n,x}$, $G_{sc,x}$, $C_{s,x}$, $C_{i,x}$). MATCRO for C3 photosynthesis obtains analytical solutions from relationships via the method shown in Masutomi (2023). For C4 photosynthesis, it is also possible to solve these equations analytically. In the case of Rubisco-limited and RuBP-limited photosynthesis, exact

Deleted: meet

Deleted: :

expressions for $\bar{A}_{c,x}$ and $\bar{A}_{j,x}$ are obtained. Under $\bar{A}_{n,x} \geq 0$, PEP-limited photosynthesis ($\bar{A}_{p,x}$) can be represented by quadratic equations by the algebraic procedures as follows:

$$0 = \{G_{bc}^2 G_{1c} R_h - G_{bc} G_{0c} - k_{p,x}(G_{0c} - G_{bc} G_{1c} R_h + G_{bc})\} \bar{A}_{p,x}^2 + \{C_a G_{bc}^2 G_{0c} - G_{bc} G_{0c} R_d + G_{bc}^2 G_{1c} R_h R_d - k_{p,x} C_a (G_{bc}^2 G_{1c} R_h - 2 G_{bc} G_{0c} - G_{bc}^2)\} \bar{A}_{p,x} + C_a G_{bc}^2 G_{0c} (R_d - k_{p,x} C_a). \quad (20)$$

Under $\bar{A}_{n,x} < 0$, the PEP-limited photosynthesis rate can be expressed as

$$\bar{A}_{p,x} = \frac{k_{p,x} C_a - R_d}{1 + k_{p,x} \left(\frac{1}{G_{bc}} + \frac{1}{G_{0c}} \right)}. \quad (21)$$

According to these equations, in the case of PEP-limited photosynthesis, there are three possible solutions. Following the criteria described by Masutomi (2023), only one analytical solution can be selected when the following requirements are satisfied: (i) under $\bar{A}_{n,x} \geq 0$, the solution must be a positive or zero real solution, and under $\bar{A}_{n,x} < 0$, it must be a negative real solution; (ii) $G_{sc,x} > 0$; and (iii) $C_l > 0$.

2.2 Crop-specific parameterization

2.2.1 Phenology

The crop growing period in MATCRO is controlled by D_{vs} based on Bouman et al. (2001). Here, $D_{vs} = 0$ means sowing, and $D_{vs} = 1$ means maturity (harvesting). It is calculated from the following equations:

$$D_{vs,t} = G_{dd,t} / G_{ddm,t}, \quad (22)$$

$$G_{dd} = \int_0^t D_{vr} dt', \quad (23)$$

$$D_{vr} = \begin{cases} 0, & T_t < T_b \mid T_h \leq T_t, \\ T_t - T_b, & T_b \leq T_t < T_o, \\ \frac{(T_b - T_o)(T_t - T_h)}{(T_h - T_o)}, & T_o \leq T_t < T_h, \end{cases} \quad (24)$$

where $G_{dd,t}$ is the growing degree days at t (time) for specific grid cell number i , $G_{ddm,i}$ is the growing degree day at maturity, D_{vr} is the developmental rate at time t , and T_t is the temperature at time t . T_b , T_h , and T_o are the crop-specific cardinal temperatures (minimum, maximum, and optimal temperatures for development, respectively, as shown in Table 1). $G_{dd,m}$ were calibrated for each point scale simulation and global scale simulation (Section 2.3). In addition, one parameter that represents the timing of flowering (known as silking; $D_{vs,flw}$) was calibrated based on observational data for the point scale simulation (Table 1).

Deleted: $D_{vs} = G_{dd} / G_{dd,m}$

Deleted: G_{dd}

Deleted:),

Deleted: $G_{dd,m}$

Deleted: parameterized

Deleted: $D_{vs,flw}$ was parameterized

2.2.2 Leaf nitrogen and Rubisco capacity

Maximum Rubisco carboxylation rate

$V_{cmax25,x}$ used in the photosynthesis module (Section 2.1) is obtained by dividing the maximum Rubisco carboxylation rate at a LAI depth of l ($V_{cmax25,x}(l)$) by L_x separately for sunlit and shaded leaves based on Bonan et al. (2011). The vertical

Deleted: the canopy level

distribution of $V_{cmax25}(l)$, which is the sum of $V_{cmax25,sn}(l)$ and $V_{cmax25,sh}(l)$, follows the exponential profile:

$$V_{cmax25}(l) = V_{cmax25}(0) \exp(-K_n l), \quad (25)$$

where $V_{cmax25}(0)$ is the maximum Rubisco carboxylation rate at the canopy top, K_n is a parameter for the vertical distribution of nitrogen (Table 1), and l represents the LAI depth from the top. The maximum Rubisco carboxylation rate in sunlit leaves ($V_{cmax25,sn}(l)$) is also calculated by the same relationship considering the light distribution:

$$V_{cmax25,sn}(l) = V_{cmax25}(0) [1 - \exp(-l(K_n + K))]^{\frac{1}{K_n + K}}, \quad (26)$$

where K is the direct beam extinction coefficient (the calculation is the same as that for Masutomi et al., 2016a). $V_{cmax25,sh}(l)$ is given by the subtraction of Eq. (25) and Eq. (26).

Here, while Bonan et al. (2011) uses the fixed value of $V_{cmax25}(0)$ value over time, $V_{cmax25}(0)$ in MATCRO is calculated dynamically as a function of specific leaf nitrogen (S_{ln} [g N m⁻²]). The function is established based on the experimental literature data. Notably, we applied the relationship between S_{ln} and light-saturated CO₂ assimilation (A_{max}) from the literature, although MATCRO-Rice and MATCRO-Soybean utilize the direct relationship between S_{ln} and $V_{cmax25}(0)$ based on the experimental literature data. The reasons are that we assume that A_{max} could be used as Rubisco-limited photosynthesis in C4 photosynthesis and that Rubisco-limited photosynthesis could be equal to the maximum Rubisco carboxylation rate from Eq. (6). Several studies have shown that A_{max} has a close relationship with S_{ln} , as shown by the logistic equation for maize (Drouet and Bonhomme, 2004; Muchow and Sinclair, 1994; Paponov and Engels, 2003; Paponov et al., 2005; Sinclair and Horie, 1989; Vos et al., 2005). We used two functions from them for different D_{vs} as follows:

$$V_{cmax25}(0) = \begin{cases} 45.1 * \left\{ \frac{2}{1 + \exp[-2.9 * (S_{ln} - 0.25)]} - 1 \right\}, & D_{vs} < D_{vs,flw}, \\ 40.2 * \left\{ \frac{2}{1 + \exp[-1.41 * (S_{ln} - 0.43)]} - 1 \right\}, & D_{vs} \geq D_{vs,flw}, \end{cases} \quad (27)$$

where $D_{vs} < D_{vs,flw}$ represents the vegetative stage at which the equation was based on Vos et al. (2005); then, for the reproductive stage, the equation was from Drouet and Bonhomme (2004). Stage-specific parameterizations were applied to reflect the lower photosynthetic activity observed during the reproductive phase compared to the vegetative phase since no single dataset adequately represents both growth phase.

Deleted: in each plant functional type

Specific leaf nitrogen

S_{ln} , which is used in the calculation of $V_{cmax25}(0)$, is obtained from the function of D_{vs} in MATCRO. The function is established based on the observational data. We utilized the study by Muchow (1988), in which S_{ln} was measured under various levels of N_{fert} (0, 60, 120, 240, 420 [kg ha⁻¹]), as follows: (i) we traced S_{ln} data using digitizer software (<https://apps.automeris.io/wpd4/>) and obtained the measurement and phenological data from the paper; and (ii) we conducted

the fitting based on the assumption that S_{ln} linearly increased until flowering and then decreased towards maturity. The parameterization given by Eqs. (28)-(30) is shown in Figure 1.

$$S_{ln} = \begin{cases} \frac{S_{ln,mx} - S_{ln,plt}}{D_{vs,flw}} D_{vs} + S_{ln,plt}, & D_{vs} < D_{vs,flw}, \\ \frac{S_{ln,matu} - S_{ln,mx}}{1 - D_{vs,flw}} (D_{vs} - 1) + S_{ln,matu}, & D_{vs} \geq D_{vs,flw}, \end{cases} \quad (28)$$

where $S_{ln,matu}$, $S_{ln,mx}$, and $S_{ln,plt}$ are S_{ln} at maturity, maximum S_{ln} , and S_{ln} at planting, respectively (Table 1). $S_{ln,mx}$ and $S_{ln,matu}$ are empirically parameterized as functions of N_{fert} as follows:

$$S_{ln,mx} = \begin{cases} -0.00001 N_{fert}^2 + 0.0064 N_{fert} + 0.6891, & N_{fert} \leq 240, \\ 1.75, & N_{fert} > 240. \end{cases} \quad (29)$$

$$S_{ln,matu} = \begin{cases} 0.001 N_{fert} + 0.57, & N_{fert} \leq 240, \\ 1, & N_{fert} > 240. \end{cases} \quad (30)$$

We set fixed values of 1.75 for $S_{ln,mx}$ and 1.0 for $S_{ln,matu}$ when N_{fert} exceeds 240 [kg ha⁻¹], as $S_{ln,mx}$ and $S_{ln,matu}$ exhibit minimal increases beyond this threshold.




Figure 1. Relationship between developmental stage (D_{vs}) and specific leaf nitrogen (S_{ln}) in MATCRO-Maize. Shapes show observational data from Muchow (1988) with the 5 types of N_{fert} : 0 kg ha⁻¹ (square), 60 kg ha⁻¹ (circle), 120 kg ha⁻¹ (triangle), 240 kg ha⁻¹ (diamond), and 420 kg ha⁻¹ (inverted triangle). The red lines represent the fitted line parameters used in MATCRO-Maize, while the dashed line represents D_{vs} at flowering ($D_{vs,flw}$).

2.2.3 Crop growth

Glucose partitioning

MATCRO calculates crop growth by partitioning net carbon assimilation (A_n) in the form of glucose, which is calculated in the photosynthesis module (Section 2.1). Partitioned glucose is supplied through photosynthesis in leaves and remobilization from the stem. The ratio of glucose partition to each organ (leaf, stem, root, and storage organ; ear) depends on D_{vs} . The term

Formatted: Font color: Text 1, English (US)

“ear” in maize represents the organ that supports the development and storage of grain. The grain developed later than the ear with approximately 83% of ear at maturity in this study (see Section 2.2.5). The dry matter for each organ is obtained from the partitioned glucose considering the carbon fraction for each organ ($C_{glu,ear}$, $C_{glu,leaf}$, $C_{glu,rot}$, $C_{glu,stm}$ in Table 1). We calibrated the partitioning ratio to leaf and ear based on the observational biomass data from Ciampitti et al. (2013a, b), whereas the ratio to shoots/roots was derived from the value from Penning de Vries et al. (1989). The stem partitioning was determined by reducing the shoot ratio with respect to the leaf and ear. Figure 2 shows the partition ratio to the leaf ($P_{r,leaf}$) and ear ($P_{r,ear}$) established via the following equations:

$$P_{r,leaf} = \begin{cases} P_{leaf}, & D_{vs} < D_{vs,leaf1}, \\ \frac{P_{leaf}(D_{vs,leaf2} - D_{vs})}{D_{vs,leaf2} - D_{vs,leaf1}}, & D_{vs} < D_{vs,leaf2}, \\ 0, & \text{otherwise}, \\ 0, & D_{vs} < D_{vs,ear1}, \end{cases} \quad (31)$$

$$P_{r,ear} = \begin{cases} \frac{D_{vs} - D_{vs,ear1}}{D_{vs,ear2} - D_{vs,ear1}}, & D_{vs} < D_{vs,leaf2}, \\ 1, & \text{otherwise}, \end{cases} \quad (32)$$

where $D_{vs,leaf1}$, $D_{vs,leaf2}$, $D_{vs,ear1}$ and $D_{vs,ear2}$ represent the D_{vs} at which the corresponding partition changes, as determined in Table 1 based on Figure 2, and where P_{leaf} is the ratio of glucose partitioned to the leaf from glucose partitioned to the shoot.




Figure 2. The ratio of glucose partitioning to leaves (a) and ears (b). Points show the ratio of glucose partition with different N_{fert} : 0 kg ha⁻¹ (square), 112 kg ha⁻¹ (cycle), and 224 kg ha⁻¹ (triangle) measured in Ciampitti et al. (2013a, b). The red lines in Figure 2 show the fitted line parameters used in MATCRO-Maize, while the dashed line represents D_{vs} at flowering ($D_{vs,flw}$).

Specific leaf weight


The specific leaf weight (S_{lw}) is used to calculate the total leaf area index (L) in MATCRO. It is a function of D_{vs} and is given by:

- Deleted:** reproductive
- Deleted:** responsible for grain
- Deleted:** .
- Formatted:** Font color: Text 1, English (US)
- Formatted:** Font color: Text 1, English (US)
- Deleted:** additional details in
- Formatted:** Font: Italic, Font color: Text 1, English (US)
- Deleted:** parameterized
- Deleted:** , stem,

280

$$S_{lw} = S_{lw,mx} + (S_{lw,mn} - S_{lw,mx}) \exp(-k_{slw} D_{vs}) \tag{33}$$

where $S_{lw,mn}$, $S_{lw,mx}$, and k_{slw} are crop-specific parameters derived from the observational data expressed in Table 1. We conducted curve fitting to S_{lw} calculated from the dry weight of the leaf biomass and the leaf area index based on Ciampitti et al. (2013a, b) and established a relationship (Figure 3).



285

Figure 3. Relationships between specific leaf weights and developmental stages. Similar to Fig. 2.

2.2.4 Crop height

Crop height (H_{gt}) is related to the calculation of evaporation in MATCRO. It assumes that the dependence of the crop height on D_{vs} is based on Penning de Vries et al. (1989) and is given by

$$H_{gt} = \begin{cases} h_{aa} D_{vs} / D_{vs,flw}, & D_{vs} < D_{vs,flw} \\ h_{aa}, & D_{vs} \geq D_{vs,flw} \end{cases} \tag{34}$$

290

where h_{aa} is the crop height at flowering, as shown in Table 1.

2.2.5 Crop yield

MATCRO calculates the final crop yield, Y_{ld} , from the dry weight of the storage organ at maturity ($W_{ear,mt}$) as follows:

$$Y_{ld} = k_{yld} W_{ear,mt}. \tag{35}$$

Here, k_{yld} is the crop-specific parameter (Table 1), which represents the ratio of Y_{ld} to $W_{ear,mt}$. The dry weight of the ear is a consistent predictor of the plant's potential yield at maturity. We parameterized K_{yld} based on Ciampitti et al. (2013b).

Table 1. Parameters in MATCRO-Maize

Variable	Value	Units	Description	Source
Crop-specific (maize)				
b	0.04	$\text{mol (H}_2\text{O) m}^{-2} \text{ s}^{-1}$	intercept of the Ball-Berry model	Sellers et al., (1996)

Formatted Table

Variable	Value	Units	Description	Source
$C_{glu,ear}$	0.815	ratio	conversion factor of dry weight from glucose to ear	Penning de Vries et al., (1989)
$C_{glu,leaf}$	0.871	ratio	conversion factor of dry weight from glucose to leaf	Penning de Vries et al., (1989)
$C_{glu,rot}$	0.857	ratio	conversion factor of dry weight from glucose to root	Penning de Vries et al., (1989)
$C_{glu,stm}$	0.810	ratio	conversion factor of dry weight from glucose to stem	Penning de Vries et al., (1989)
$D_{vs,rot1}$	0.35	ratio	1st point of D_{vs} at which the partition pattern to root changes	Penning de Vries et al., (1989)
Crop-specific (maize)				
$D_{vs,rot2}$	0.72	ratio	2nd point of D_{vs} at which the partition pattern to root changes	Penning de Vries et al., (1989)
$D_{vs,ear1}$	0.37	ratio	1st point of D_{vs} at which the partition pattern to ear changes	Parameterized in this study
$D_{vs,ear2}$	0.6	ratio	2nd point of D_{vs} at which the partition pattern to ear changes	Parameterized in this study
$D_{vs,flw}$	0.52	ratio	D_{vs} at flowering	Parameterized in this study
$D_{vs,lef1}$	0.25	ratio	1st point of D_{vs} at which the partition pattern to leaf changes	Parameterized in this study
$D_{vs,lef2}$	0.48	ratio	2nd point of D_{vs} at which the partition pattern to leaf changes	Parameterized in this study
f_{stc}	0.35	ratio	fraction of glucose allocated to starch reserves	Penning de Vries et al., (1989)
h_{aa}	2	m	crop height at flowering	Penning de Vries et al., (1989)
k_{yld}	0.83	ratio	ratio of crop yield to dry weight of ear at maturity	Parameterized in this study
k_{slw}	3	ratio	parameter that represents the relationship between S_{lw} and D_{vs}	Parameterized in this study
m	4	ratio	the slope of the Ball-Berry model	Sellers et al., (1996)
$G_{dd,m}$	—	K day	growing degree day at maturity	Parameterized in this study
P_{lef}	0.49	ratio	partition ratio of glucose to leaf from glucose partitioned to the shoot	Parameterized in this study
P_{rot}	0.25	ratio	partition ratio of glucose to root	Penning de Vries et al., (1989)
$r_{dl,lef}$	3.0×10^{-7}	s^{-1}	ratio of dead leaf at harvest	Masutomi et al., (2016)
r_{rt}	0.06	$m s^{-1}$	growth ratio of root	Penning de Vries et al., (1989)
$S_{ln,plt}$	0.825	$g m^{-2}$	specific leaf nitrogen at planting	Parameterized in this study
$S_{ln,mx}$	See Eq. (29)	$g m^{-2}$	maximum specific leaf nitrogen	Parameterized in this study
$S_{ln,matu}$	See Eq. (30)	$g m^{-2}$	specific leaf nitrogen at maturity	Parameterized in this study
$S_{lw,mn}$	400	$kg ha^{-1}$	minimum specific leaf weight	Parameterized in this study
$S_{lw,mx}$	700	$kg ha^{-1}$	maximum specific leaf weight	Parameterized in this study
T_b	8.6	$^{\circ}C$	minimum temperature for development	Osborne et al., (2015)
T_h	42.0	$^{\circ}C$	maximum temperature for development	Osborne et al., (2015)
T_o	10.0	$^{\circ}C$	optimal temperature for development	Osborne et al., (2015)
$z_{rt,mx}$	1.5	m	maximum root depth	Penning de Vries et al., (1989)
α	0.05	$mol mol^{-1}$	quantum efficiency	Sellers et al., (1996)
β_{ej}	0.8	ratio	GPP transition factor	Lawrence et al., (2020)
Others				
k_n	0.3	ratio	vertical distribution of nitrogen	Oleson et al., (2013)
S_1	0.3	K^{-1}	temperature dependence of $V_{cmax,x}$	Lawrence et al., (2020)
S_2	313.15	K	temperature dependence of $V_{cmax,x}$	Lawrence et al., (2020)
S_3	0.2	K^{-1}	temperature dependence of $V_{cmax,x}$	Lawrence et al., (2020)
S_4	288.15	K	temperature dependence of $V_{cmax,x}$	Lawrence et al., (2020)
β_{ip}	0.95	ratio	GPP transition factor	Lawrence et al., (2020)

Formatted Table

Deleted: 281.75

Deleted: K

Deleted: 315.15

Deleted: K

Deleted: 303.15

Deleted: K

Formatted Table

2.3 Model evaluation

MATCRO can run the simulation both at a point scale and at a global scale. The developed model was evaluated both at a point scale and at a global scale. For point scale levels, the two model output datasets, LAI and total aboveground were compared with the observation data from the four sites. Meanwhile, we use yield data for evaluation. After confirming the ability of the model to simulate maize growth, two types of evaluations were conducted at the global scale. First, the simulated yields at the grid cell were compared with the gridded yield data of the Global Dataset of Historical Yields (GDHY) (Iizumi, and Sakai, 2020). Second, the simulated yields at the country and total global levels were compared with the country yield report and global data from the Food and Agriculture Organization (FAOSTAT, 2024). To quantify the model performance, four statistical values were used in this study: the Pearson correlation coefficient (COR), root mean square error (RMSE), relative root mean square error (RRMSE) and normalized mean absolute error (NMAE). RRMSE and NMAE were calculated as follows:

$$RMSE = \sqrt{\frac{1}{n} \sum_{i=1}^n (y_i - \hat{y}_i)^2}, \quad (36)$$

$$RRMSE = \frac{RMSE}{\bar{y}}, \quad (37)$$

$$NMAE = \frac{1}{n} \sum_{i=1}^n \frac{|y_i - \hat{y}_i|}{y_i}, \quad (38)$$

where y_i is the actual value, $\hat{y}_i$ is the predicted value, and $\bar{y}$ is the mean of the actual value.

2.3.1 Model evaluation at a point scale

To evaluate the model performance at a field scale, we used observational data from four sites (Brazil, France, Tanzania, and the USA; Table 2) used in the Agricultural Model Intercomparison and Improvement Project (AgMIP) study (Bassu et al., 2014). We used local daily climate data of precipitation, downwards shortwave radiation, air temperature, wind speed (P_{rc} , R_s , T_a , U respectively), management data (N_{fert} and irrigation regime) and phenological data (planting, flowering, and maturity dates) for model input data at each site. We identified the soil texture from the gridded soil texture dataset of ISIMIP (Volkholz and Müller, 2020). Annual CO₂ data were also taken from the same data used for the global simulation. Climatic data were estimated from the NASA Modern Era Retrospective-Analysis for Research and Applications (AgMERRA; Ruane, et al., 2015) when measured data were unavailable (Bassu et al., 2014).

Table 2. Evaluation site information in the point-scale simulation

Country	Site	Latitude	Longitude	Soil type	Sowing date	Hybrid	Total N fertilizer [kg N ha ⁻¹]	Irrigation
Brazil	Rio Verde	17.52°S	51.43°W	Geri-Gibbsic Ferralsol	Oct. 22 nd 2003	Pioneer 30K75	0	No
France	Lusignan	46.25°N	00.07°E	Cambisol	Apr. 26 th 1996	Furio	255	Yes
Tanzania	Morogoro	06.50°S	37.39°E	Haplic Arenosol	Oct. 26 th 2009	TMV1	61	Yes

Deleted: validation

Deleted: validated

Deleted: validation

Deleted: three

Deleted: ,

Deleted: biomass, and yield,

Deleted: validation

Deleted: validations

Deleted: , 2019)

Deleted: .

Deleted: validation

Deleted: validate

Deleted: accuracy

Deleted: There was no information on

Deleted: classification of

Deleted: , so we extracted one grid

Deleted: from the same data used for the global simulation (Section 2.3.2)....

Deleted: MERRA; Rienecker

Deleted: 2011

Deleted: Validation

Formatted Table

350

355

360

365

370

USA	Iowa	42.01°N	93.45°W	Gleysols	May 4 th 2010	Golden Harvest GH-9014	167	No
-----	------	---------	---------	----------	--------------------------	------------------------	-----	----

Notably, air pressure (P_s) and specific humidity (S_h) were not provided. We used the same data as the global simulation for the soil classification and P_s . S_h was converted from R_h using T_a and the vapour pressure. We parameterized $G_{dd,m}$ and $D_{vs,flw}$ based on T_a and phenological data (sowing, flowering, and maturity dates). $G_{dd,m}$ calibrated for each site is used for the simulations, while the average $D_{vs,flw}$ over the 4 sites is used (0.52 in Table 1). As a result, the mean average errors were estimated as 4.25 and 7 days for flowering and maturity, respectively (Figure 4). MATCRO was run with these parameters, and then the model output was evaluated with the observations for the following 3 variables: seasonal change in the LAI, total aboveground biomass, and final yield.

Model calibration was conducted based on phenological data (Table 2, Bassu et al., 2014) and biomass data for carbon partitioning of leaf and ear derived from Ciampitti et al. (2013a, b). In this study, a global parameter from the literature was applied uniformly across all regions at the grid-cell level instead of using site-specific calibrated parameters in the simulations. The model was then assessed at the point scale to check the calibration for phenology (flowering and maturity) and was evaluated against time-series data of LAI, aboveground biomass, and harvested yield (see Section 3.1) that were not included in the model calibration.




Figure 4. Model-fit comparison of the flowering and maturity date simulations (SIM on the y-axis) and observations (OBS on the x-axis). DOY represents the number of days from January 1st. Shapes show each site: Brazil (square), France (circle), Tanzania (triangle), and the USA (diamond). The colours indicate the phenological stages: flowering (red) and maturity (blue).

2.3.2 Model evaluation at a global scale

Simulation settings

Deleted: parameterized

Deleted: valida

Deleted: ion

Deleted: Comparison

Deleted:

Deleted: validation

For the global-scale simulation, the model was run at a spatial resolution of $0.5^\circ \times 0.5^\circ$ from 1980–2010 under both rainfed and irrigated conditions. The required input data were as follows. (i) Crop calendar data were from the Global Gridded Crop Model Intercomparison (GGCMI) phase 3 protocol (Jägermeyr et al., 2021). It provides planting and maturity dates for 18 different crops, including maize, separated by rainfed and irrigated systems. We parameterized the average $G_{dd,m}$ at each grid over the period 1980–2010 for the growing season from the planting to maturity dates for each of the rainfed and irrigated conditions. Both the planting date and the simulated $G_{dd,m}$ were used as the input data for the global-scale simulations. (ii) Water management data (i.e., irrigation regime) from the MIRCA2000 dataset (Portmann et al., 2010). In the case of irrigated conditions, the soil moisture was set to field capacity during the growing season. (iii) N_{fert} from the Inter-Sectoral Impact Model Intercomparison Project (ISIMIP; Volkholz and Ostberg, 2022). It provides the annual nitrogen fertilizer inputs for five canonical crop types, including C4 annual crops for maize. (iv) Soil texture classification from ISIMIP3a protocol soil input data (Volkholz and Müller, 2020). (v) Annual atmospheric CO₂ data from the ISIMIP3a (Büchner and Reyer, 2022). (vi) Six types of daily meteorological for model inputs (P_s , P_{rc} , S_h , T_{max} , T_{min} , T_a , U) from the GSWP3-W5E5 dataset for the ISIMIP3a dataset (Lange et al., 2022). We set the data from (i), (ii), and (iv) as constants across the simulation period, whereas the data from (iii), (v), and (vi) are variables.

Analysis

MATCRO-Maize was assessed for the phenological simulation of harvest time against the phenological dataset GGCMI (Jägermeyr et al., 2021) and global datasets of crop phenological events for agricultural and earth system modeling which was derived from various field experiments and a phenology model (GCPE; Mori et al., 2023). These datasets were compared under both rainfed and irrigated conditions in $0.5^\circ \times 0.5^\circ$ resolution to check the model performance. The simulated final yields in each grid cell under irrigated and rainfed conditions then were aggregated by grid cell, country and global level with the harvested area from MIRCA2000 data (Portmann et al., 2010) via the following equation for each year from 1981–2010:

$$Yield_{aggregated} = \frac{\sum_{i=1}^n (Yield_{i,rf} \times Area_{i,rf}) + \sum_{i=1}^n (Yield_{i,irr} \times Area_{i,irr})}{\sum_{i=1}^n (Area_{i,rf} + Area_{i,irr})} \quad (39)$$

where $Yield_{aggregated}$ is the aggregated yield with the total grid cells (n) in grid cell i . $Yield_{rf}$ and $Yield_{irr}$ are the simulated yields under rainfed and irrigated conditions, respectively, and $Area_{rf}$ and $Area_{irr}$ are the harvested areas from MIRCA2000 for rainfed and irrigated conditions, respectively.

The model performance was evaluated by comparing its output with the historical yield dataset. The grid cell-level yield was averaged across a 30-year period and compared with the Global Dataset of Historical Yields (GDHY) (Iizumi and Sakai, 2020), 290-year period of GlobalCropYield (GCY, Cao et al., 2025), and the Spatial Production Allocation Model by (SPAM; IFPRI, 2019) at year 2010. The country- and global-level yields were compared with FAOSTAT data (FAOSTAT, 2024) for the average and annual variabilities over the 30 years. In the comparison at the country level, we focus on the top 20 maize-producing countries that account for more than 85% of total maize production.

Deleted: , 2019)

Deleted: .

We focused on two perspectives for [evaluation](#): (i) the ability of the model to capture the spatial distribution of yield in both low- and high-producing countries and (ii) the ability of the model to reproduce the climatic effect reflected in the interannual variability at the country and global scales. The first perspective was analysed using NMAE to quantify model error for both the global yield and the yield of the top 20 producing countries. The 30-year average yields were also compared on the basis of the statistics of COR, RMSE, and RRMSE to confirm the accuracy. The second perspective was analysed via the COR of the detrended deviation between the simulated and FAOSTAT yields to assess the interannual variability.

Deleted: validation

3 Results

3.1 Point-scale simulations

A comparison of the time series changes in the LAI at each experimental site is shown in Figure 5. In general, MATCRO-Maize captured the increasing trend towards flowering time and then decreasing trend towards the end of maturity. Especially during the vegetative stage ($D_{vs} < D_{vs,flw}:0.52$), the simulated LAI showed relatively good agreement. However, the simulated LAI was notably underestimated in Brazil and France immediately before the reproductive stage (near the dashed black line in Fig. 5). Figure 6 compares the time series of total aboveground biomass between the simulated and experimental data. Except for Tanzania, MATCRO-Maize accurately estimated the increasing trend of total aboveground biomass towards maturity, although the simulated biomass in Brazil was underestimated at maturity. The simulated total aboveground biomass in Tanzania increased until maturity, while the observations gradually decreased towards maturity time (Fig. 6 (c)).

Moved down [1]: The effects of photosynthesis and N fertilizer
In addition to the yield comparison, we analysed the effect of nitrogen fertilizer (N_{fert}) on maize yield, as it is a key determinant of crop yield. This analysis compared both FAOSTAT data and

Deleted: This analysis compared both FAOSTAT data and simulated data from N_{fert} for a 30-year average with simple linear regression. We also conducted two tests to quantify the effects of the N_{fert} -related function and parameters as follows: (i) Eq. (27) during the vegetative stage is derived from Drouet and Bonhomme (2004), defined as “test $S_{in}-V_{emax}$ ”, was changed to:
$$V_{emax}(0) = 36.8 * \frac{2}{1 + \exp[-2.45 * (S_{in} - 0.27)]} - 1, D_{vs} < D_{vs,flw}$$

(40)
and (ii) $S_{in,plt}$ from 0.825 (Table 1) to 0.5 (defined as “test $S_{in,plt}$ ”).

Formatted: Indent: First line: 0 ch

Deleted:

Figure 7 compares the 1:1 line between the simulated and experimental data for the seasonal LAI (Fig. 7 (a)), seasonal total aboveground biomass (Fig. 7 (b)), and harvested yield (Fig. 7 (c)). The LAI underestimation in France and Brazil (Fig. 5) could also be seen with a large RMSE, which is approximately 50% of the average LAI across all observational values at 3 sites except for Tanzania, although overall, the comparison was statistically significant (p value < 0.01), with a COR of 0.762. The comparison of total aboveground biomass was statistically significant (p value < 0.001), with a COR of 0.895, although the RMSE was 3,628.3 [kg ha⁻¹], which corresponds to approximately 35% of the average of all observed total aboveground biomass. While the comparison of the final crop yield was statistically significant (p value < 0.01), there was a relatively low COR compared with the LAI and total aboveground biomass due to the small sample size (N=4) and the overestimation for Tanzania. The RMSE was 2,575.0 [kg ha⁻¹], which is approximately 30% of the average observational yield at all the sites. It is noted that Figures 5 – 7 present the model evaluation using independent data. Evaluation was performed using a global parameter from the literature to simulate the plant organs in the global-scale simulation, which may have resulted in some deviations.

Deleted: lead

Deleted: to




Figure 5. Temporal evaluation of leaf area index (LAI) simulated by MATCRO-Maize (red line) at each site: (a) Brazil, (b) France, (c) Tanzania and (d) the USA across the developmental stage (D_{vs}). The observation data in each site is shown by black point. Notably, there were no observational data in Tanzania. The error bars were provided only for Brazil. The dashed black line shows the flowering time.




Figure 6. Temporal evaluation of total aboveground biomass (AGB) simulated by MATCRO-Maize (red line) at each site: (a) Brazil, (b) France, (c) Tanzania and (d) the USA across the developmental stage (D_{VS}). The observation data in each site is shown by black point. The error bars were only provided for Brazil and Tanzania. The dashed black line shows the flowering time.





Figure 7. Statistical comparison (COR, RMSE, and RRMSE) of maize yield. The x-axis (OBS) represents the observational data, and the y-axis (SIM.) is the simulated data. Shapes show each site: Brazil (square), France (circle), Tanzania (triangle), and the USA (diamond). Notably, there was no observed LAI in Tanzania. The symbols ***, **, indicate p values < 0.001 and 0.01, respectively.

3.2 Global-scale simulations

3.2.1 Phenology

The timing of seasonal biological events (i.e. harvest time) has a significant impact on crop growth and yield outcomes. Global yield is affected by global phenology. We assessed agreement to check the model performance by comparing the difference between simulated global harvest time (1981–2010 mean) with gridded global dataset of phenological datasets of GGCM (Jägermeyr et al., 2021; Figs. 8(a and b)), and GCPE (Mori et al., 2023; Figs. 8(c and d)). The maps show consistent spatial patterns for later harvest time between the simulation and the reference datasets, in parts of Brazil, USA, southern and central Africa. The discrepancies between dataset are likely produced due to the difference in phenology parameterization and management assumptions where GGCM and GCPE used different methodology and data sources. Moreover, the use of the growing degree day method in the simulations led to year-to-year differences in harvest time compared with the reference crop calendar used for the input data (Figs. 8(a and b)). The mean absolute differences in harvest time (Figs. 8(e and f)) indicated that the largest biases occur mostly in tropical regions.



Deleted:

Deleted: (a) LAI, (b) Total aboveground biomass, and (c) Yield

Deleted: A comparison of the global distributions is shown in Figure 8 (simulation: Fig. 8(a); GDHY dataset: Fig. 8(b)).




Figure 8. The difference between simulated harvest time (days) in MATCRO-Maize simulations with (a) GGCMI in the rainfed, and (b) irrigated conditions, (c) GCPE in the irrigated, and (d) rainfed conditions. Blue indicates underestimation, while red indicates overestimation between simulation and references. Panels (e) and (f) show the mean of absolute differences (days) under the rainfed (a, c) and irrigated (b, d) comparisons, respectively.


3.2.2 Yield

A comparison of the global distributions is shown in Figure 9 (simulation: Fig. 9(a); observation dataset: Figs. 9(b, c, and d)). All datasets were harmonized to a $0.5^\circ \times 0.5^\circ$ resolution, including simulated yield from MATCRO-Maize (Fig. 9(a)), the Global Dataset of Historical Yield (GDHY; Iizumi and Sakai, 2020; Fig. 9(b)), GlobalCropYield (GCY; Cao et al., 2025; Fig.

Deleted: €

9(c)), and the Spatial Production Allocation Model by (SPAM; IFPRI, 2019; Fig. 9(d)). The data were averaged over 1981–2014 for GDHY, 1982–2014 for GlobalCropYield, and for the year 2010 for SPAM. While the overestimation could be seen mainly in tropical regions, the simulated yield could capture high-yielding regions, including the Corn Belt in the United States and the northern part of China, in agreement with the reference datasets.

Temporal changes in the global yield across 30 years indicated that although the global yield had an NMAE of 0.67, indicating a simulation error of 67% with respect to the average FAO yield, the comparison of the interannual variability between the simulations and observations was statistically significant (p value < 0.01), with a COR of 0.549 (Figure 10). For the top 20 producing countries, MATCRO-Maize also tended to overestimate the yield in terms of the annual yield (Figure 11) and the average yield over a 30-year period (Figure 12). The overestimation was strong in Egypt, where the simulated yield was approximately four times greater across 30 years. In terms of interannual variability, half of the 20 countries were statistically significant, with p values < 0.001 for 6 countries, < 0.01 for 2 countries, and < 0.05 for 2 countries (Fig. 11). The 30-year average comparison was also statistically significant (p value < 0.01), with a COR of 0.58, although the RMSE was 4,007.7 [kg ha⁻¹], which is almost the same as the average yield of the top 20 maize-producing countries (Fig. 12).



Deleted: .

Formatted: Indent: First line: 0.5 cm


Deleted: 9

Deleted: 10

Deleted: 11


Deleted: 10

Deleted: 11



Deleted:

520 **Figure 9.** Global distribution of the 30-year average (1981-2010) maize yield by (a) simulations from the MATCRO-Maize and (b) the GDHY dataset. For comparison, yield estimates from shorter periods are also shown from (c) GlobalCropYield for 29-year average (1982-2014) and (d) SPAM2010 for year 2010. The yield is aggregated based on the harvested area from MIRCA2000.



525 **Figure 10.** Interannual variability in global maize yield from 1981 to 2010 for our simulation (red circles) and FAOSTAT (black) yields. COR represents the correlation coefficient of interannual variability. NMAE means normalized mean absolute error. Asterisks ** indicate p value < 0.01.

Deleted: 8


Deleted: <object>

Deleted: 9

Formatted: Font: 9 pt

Formatted: Font: 9 pt


Formatted: Normal



535 Figure 11. Comparison of interannual variability for the top 20 maize-producing countries. Similar to Fig. 9. Notably, the simulated yield in Egypt is not shown as it extends beyond the range of the y-axis. The symbols ***, **, and * indicate p values < 0.001, 0.01, and 0.05, respectively.

Deleted: 10

Formatted: Normal



540 **Figure 12.** Accuracy of the 30-year average of the simulated yield (SIM) to the observed yield (OBS from FAOSTAT data) for the top 20 countries. Notably, the Egypt data points are not shown as exceeding the range of the y-axis. Asterisks ** indicate a p value < 0.01.

3.3 The effects of photosynthesis and N fertilizer

In addition to the yield comparison, we analysed the effect of nitrogen fertilizer (N_{fert}) on maize yield, as it is a key determinant of crop yield. It compared both FAOSTAT data and simulated data from N_{fert} for a 30-year average using a fitted polynomial curve (quadratic polynomial regression). We also conducted two tests to quantify the effects of the N_{fert} -related function and parameters as follows: (i) Eq. (27) during the vegetative stage is derived from Drouet and Bonhomme (2004), defined as “test $S_{ln-V_{cmax}}$ ”, was changed to:

$$V_{cmax}(0) = 36.8 * \left\{ \frac{2}{1 + \exp[-2.45 * (S_{ln} - 0.27)]} - 1 \right\}, D_{vs} < D_{vs,flw} \quad (40)$$

and (ii) $S_{ln,plt}$ from 0.825 (Table 1) to 0.5 (defined as “test $S_{ln,plt}$ ”).

550 Figure 13 illustrates the comparison of country-level yield data with nitrogen fertilizer levels: (a) FAOSTAT data, (b) simulated yield by MATCRO-Maize, (c) the impact of reduced Rubisco activity on photosynthetic rates based on experimental data from Drouet and Bonhomme (2004) in the “test Sln-Vcmax” scenario, and (d) the effect of reduced photosynthetic rates due to lower initial specific leaf nitrogen at planting time in the “test Sln,plt” scenario. The nitrogen fertilizer values were derived from gridded dataset (ISIMIP; Volkholz and Ostberg, 2022).

555 Figures 13 (a) and (b) show the comparisons based on N_{fert} for each FAOSTAT and simulated yield, respectively. MATCRO has a strong N_{fert} effect on the yield reflected in the steep upward trend of the fitted curves. This effect was scarcely alleviated by the intentionally reduced effect of photosynthesis (Figs. 13(c and d)), mainly because of the effect of Egypt as an outlier, with higher values. Without Egypt, as an outlier, the curves for FAOSTAT and MATCRO-Maize were more comparable. The maize yield in Egypt shows high value compared to other countries where significant overestimation was observed.

560

Deleted: 11

Moved (insertion) [1]

Formatted: Font: Not Bold

Formatted: Heading 2

Deleted: 12

Deleted: slope

Deleted: regression line with relatively similar intercepts (slope: 27.6 and 47.3; intercept: 1,701.0 and 2,187.4 for FAOSTAT and MATCRO-Maize, respectively).

Deleted: changed setting (Fig. 12 (c), (

Deleted: . Figure 13 shows the same comparison as Fig. 12 without

Deleted: .

Deleted: slope

Deleted: similar to that of FAOSTAT (33.8 for FAOSTAT and 30.35 for MATCRO-Maize), whereas the intercept was approximately 2.5 times greater (1,310.6 for FAOSTAT and 3,247.0 for MATCRO-Maize) (Fig. 13 (a), (b)). The changed equation and parameter partly alleviated the large intercept, although the slope became slightly smaller than that of FAOSTAT (Fig. 13 (c), (d)).





Figure 13. N_{fert} impact on yield of (a) FAOSTAT, (b) simulated yield with the original setting (Default), (c) simulated yield with the changed $S_{In-Vcmax}$ relationship (test Sln-Vcmax), (d) simulated yield with the changed parameter related to the $D_{vs}-S_{In}$ function (test Sln, plt). N_{fert} (N fertilizer) and country yield were averaged across 30 years for each country. The legends for symbols are the same as those in Fig. 11. The solid lines are fitted curve for the data, while the dashed lines in (b), (c), and (d) indicate fitted curve based on the data in (a). All lines were fitted using a quadratic polynomial regression.

4 Discussion

4.1 Point-scale simulations

The point-scale simulations were evaluated using global parameters to assess their ability to capture broad yield patterns across different regions. The simulated harvested yield showed statistically significant correlations at the point scale (Fig. 7), indicating that the MATCRO-Maize model could simulate maize growth and yield, but its performance was limited at the point-scale. However, there were some discrepancies between the simulations and observations remain due to the limitations of using global parameters, such as the underestimation of the LAI in Brazil and France, the underestimation of the total aboveground biomass in Brazil, and the different growth trends of the total aboveground biomass in Tanzania. The underestimation of LAI is primarily due to the use of global morphological parameters at the site scale. Further investigation will improve site-specific performance by coupling LAI to key soil properties (soil organic carbon, total nitrogen, and water-holding capacity) and by incorporating canopy cover fraction following Hasegawa et al. (2008). Global parameters at the point scale enable testing the model's applicability across various regions, although local variations in soil, climate, or crop




Deleted:

Deleted: 12

Formatted: Tab stops: 16.5 cm, Left

Deleted: The...he solid lines are fitted curve for the data, while the dashed lines in (b), (c), and (d) indicate the ...itted curve based on the data in (a). All lines were fitted using a quadratic polynomial regression lines in (a).

(... [1])



620 management may not be fully captured. The MATCRO-Maize model could simulate maize growth and yield, but its performance was limited at the point scale.

625 One potential factor contributing to the underestimation of the LAI in France might be related to the effect of plant density, which is not currently considered in MATCRO. The actual plant density [plants m⁻²] at each site was 9.5 (France), 7.5 (USA), 6.6 (Brazil), and 9.5 (Tanzania) (Bassu et al., 2014). Some studies have shown that LAI trends are affected primarily by the plant density factor relative to N_{fert} and hybrids (Boomsma et al., 2009; Ciampitti et al., 2013a; Ciampitti and Vyn, 2011). This may be the reason for the underestimation that MATCRO could not reproduce the trends driven by plant density, although other important factors (e.g., management practices, climatic conditions), which are quite different from each site in the literature, would also affect crop growth variables, including the LAI.

630 Both the underestimation of the LAI and total aboveground biomass in Brazil were probably caused by the field experimental conditions of $N_{fert} = 0$, given its effect on crop growth in MATCRO. The reason for the lack of fertilization in the field experiment was that sufficient N was released by organic matter mineralization (Bassu et al., 2014), which was not considered in the model. Moreover, N_{fert} directly affects S_{ln} in MATCRO, with an increasing trend towards flowering and then a decreasing trend towards maturity (Fig. 1). S_{ln} is related to $V_{cmax25}(0)$, which in turn affects the photosynthesis calculation (Section 2.1 and Section 2.2.2). In particular, during the reproductive stage, we used Eq. (27), which results in a low $V_{cmax25}(0)$ under low S_{ln} due to the more gradual slope of the curve compared with the vegetative stage (1.41 for the reproductive stage, and 2.9 for the vegetative stage, in Eq. (27)), indirectly leading to low biomass accumulation through photosynthesis. This could be attributed to the underestimation of total aboveground biomass at maturity (Fig. 6 (a)). For underestimation of the LAI, low leaf biomass accumulation, which is derived from the same mechanism, would be the reason considering the calculation process of the LAI in MATCRO. The LAI is determined by the division of the leaf biomass weight by S_{lw} , which depends on D_{vs} . Because S_{lw} is calculated from the same parameter at all sites (Eq. (33) and Fig. 3), leaf weight is the factor that causes differences between sites, leading to the underestimation of the LAI in Brazil. Therefore, the condition of $N_{fert} = 0$ might be the reason for both underestimations.

645 One possible reason for the difference in the growth trend of biomass in Tanzania might be related to growing season length. The cultivar used in Tanzania was a short season type with 99 days of observed growing season length, whereas the cultivars at other sites were medium or long season type with lengths ranging from 122 to 173 days (Bassu et al., 2014). Capristo et al. (2007) reported that, compared with medium- and long-season cultivars, short-season cultivars presented the lowest biomass accumulation from flowering to maturity, which was reflected in the observed biomass (Fig. 6 (c)). This might indicate that the trend of biomass accumulation differs across growing season types, although other factors, such as climatic conditions or biotic stresses, could also affect accumulation. While MATCRO considers the growing season length as $G_{dd,m}$ to judge the harvesting time, this does not mean that MATCRO could capture the difference in trends due to growing season types, possibly leading to the gap between the simulations and observations shown in Tanzania.

Deleted: The reason for

4.2 Global-scale simulations

A comparison of the global distribution of maize yield revealed that MATCRO-Maize could capture the distribution of high-yield regions but could not capture the yield in tropical regions (Figures 8 and 9). Similar overestimations in tropical regions have also been reported in other global models, possibly because of the lack of representation of extreme weather or crop pests (Lombardozzi et al., 2020; Osborne et al., 2015). Moreover, soil fertility also an important source of model error and contributes to spatial variation.

Notably, MATCRO-Maize tended to overestimate the absolute values for both the total global yield and the yields of the top 20 countries, as reflected in the NMAE and RMSE values (Figures 10, 11, and 12). The simulated total global yield is determined mainly by the yield of the top 3 maize-producing countries, the United States, China, and Brazil, which have large cultivated areas (Table 3). All three countries' yields were overestimated, where the simulated yields were approximately 1.2, 1.7, and 1.8 times greater for the 30-year averages in the United States, China, and Brazil, respectively, leading to overestimation of the total global yield. Such overestimations in the main producing countries, especially in China and Brazil, are also observed in other global crop models (Von Bloh et al., 2018; Osborne et al., 2015; Schaphoff et al., 2018). This might indicate that there are factors that are important for determining yields but are not considered in most crop models.

For the top 20 producing countries, the overestimation was strong in Egypt, with an approximately fourfold greater simulated yield than that of FAOSTAT. This overestimation might be caused by the irrigated conditions in all grids in Egypt. Under manually changed rainfed conditions, crop growth in Egypt in the model was almost not simulated because of the inhibited photosynthesis rate caused by strong water stress. Under irrigated conditions, this strong water stress was alleviated. In addition, the radiation in Egypt was consistently strong throughout the growing period, and N_{fert} was highest among the top 20 countries across the 30 years simulated, increasing from approximately 180 kg ha⁻¹ in 1980 to 360 kg ha⁻¹ in 2010. This caused the colimited photosynthesis rate to be high (Eq. (4)) across the growing seasons, leading to marked overestimation.

The current version of MATCRO-Maize can reproduce yield responses to nitrogen fertilization across a range of fertilizer levels, but it tends to overestimate yields under certain conditions (e.g., Egypt) likely because the model assumes higher nitrogen use efficiency and idealized irrigation conditions where actual yields are constrained by soil quality, management, and local cultivar traits that are not explicitly represented. This suggests that the representation of nitrogen effects in the model remains simplified, and further refinement is needed for region-specific scale simulation.

Although the simulated yield has the large error in terms of the absolute value, the comparison of the 30-year average yield was statistically significant, with a COR of 0.58 (p value < 0.01) and an RMSE of 4,008 kg ha⁻¹ (Fig. 12), showing the ability to capture the spatial distribution of the yield both in low- and high-producing countries from the first perspective of the comparison (Section 2.3.2). This result was comparable with the similar result of another model: LPJ-GUESS (Olin et al., 2015), with a COR of 0.46 and an RMSE of 4,300 kg ha⁻¹ (Table 4), although the targeted countries were different (top 20 producing countries for MATCRO-Maize, whole countries for LPJ-GUESS).

Deleted: Figure

Deleted: 9,

Deleted: 11

Deleted: 11

690 In terms of interannual variability from the second perspective, the total global yield and approximately one-third of the
top 20 producing countries were statistically significant, with p values < 0.01 (Figs. 10 and 11), indicating that MATCRO-
Maize could reproduce the climatic effect globally to some extent. This might also be supported by the similar comparisons of
other global crop models in terms of statistics (Table 4), although it is difficult to simply compare the statistical values between
the models owing to the differences in periods, input data, and methods for detrending and aggregating the yield. The COR of
interannual variability for total global yield in MATCRO-Maize was in the range of those of the other models (0.55; 0.42–0.89,
695 respectively). For the top 20 countries, almost all the COR values also ranged between those of the other models. Therefore,
these comparisons from two perspectives might indicate that MATCRO-Maize could yield reasonable results. The moderate
correlations observed reflect the typical influence of yield data variability and uncertainty in management practices across
regions.

Deleted: 9

Deleted: 10

700 **Table 3.** Maize cultivated land area for 20 major producer countries from MIRCA2000 (Portmann et al., 2010).

Country	Total area [ha]	Rainfed area [ha]	Irrigated area [ha]
Argentina	3,248,715.9	3,147,580.7	101,135.3
Brazil	11,223,262.5	11,120,154.9	103,107.6
Canada	1,364,585.3	1,328,206.2	36,379.1
China	24,376,805.2	11,615,190.0	12,761,615.2
Egypt	827,766.1	0.0	827,766.1
Ethiopia	1,172,231.1	1,084,795.6	87,435.5
France	3,128,401.0	2,257,380.0	871,021.0
Hungary	1,057,610.7	1,052,622.6	4,988.1
India	6,294,770.9	4,833,685.9	1,461,085.0
Indonesia	3,479,825.7	3,135,443.9	344,381.8
Italy	1,322,692.9	534,281.4	788,411.5
Mexico	7,459,039.5	5,852,617.4	1,606,422.1
Nigeria	3,686,757.3	3,667,564.5	19,192.8
Philippines	2,590,081.0	2,590,081.0	0.0
Romania	3,139,981.1	3,016,990.5	122,990.6
Russia	4,206,747.0	3,594,403.2	612,343.9
Serbia	1,074,614.2	1,062,985.8	11,628.4
South Africa	3,060,053.5	2,930,208.2	129,845.4
Ukraine	3,382,783.5	3,194,146.2	188,637.3
United States	31,307,667.3	26,508,600.7	4,799,066.7

Formatted Table

Table 4. Statics of model simulation accuracy of the MATCRO-Maize and other crop models. Notably, the asterisks for GGCMi phase I indicate the p values: *** for p values < 0.001, ** for p values < 0.05, * for p values < 0.1, whereas those of LPJmL4 and MATCRO-Maize indicate the p values: *** for p values < 0.001, ** for p values < 0.01, * for p values < 0.05.

			COR of interannual variability						
References	Period	Global	USA	China	Brazil	Mexico	France	Argentina	
MATCRO-Maize	-	1981-2010	0.549**	0.692***	0.518**	0.349	0.015	0.654***	0.694***
JULES-crop ¹	Osborne et al., 2015	1961-2008	0.48	0.43	0.12	0.12	0.061	0.52	0.57
LPJmL4 ²	Schaphoff et al., 2018	1981-2010	–	0.675***	0.676***	0.169	-0.124	-0.331	0.717***
LPJmL5 ³	Bloh et al., 2018	1981-2010	–	0.686***	0.641***	0.0591	0.0618	0.461*	0.650***
GGCMI phase 3 ⁴	Jägermeyr et al., 2021	1981-2015	–	0.817	0.245	0.029	–	0.649	0.727
GGCMI phase 1 ⁵	Müller et al., 2017	1982-2006	0.42**~0.89***	0.89	0.75	0.66	0.85	0.87	0.85
			COR of interannual variability						
References	Period	Romania	South Africa	India	Italy	Hungary	Indonesia	Ukraine	
MATCRO-Maize	-	1981-2010	0.719***	0.646***	0.046	0.276	0.900***	0.252	0.339
JULES-crop ¹	Osborne et al., 2015	1961-2008	0.32	0.41	0.34	0.34	0.33	0.065	–
LPJmL4 ²	Schaphoff et al., 2018	1981-2010	–	0.711***	-0.22	–	–	0.124	-0.046
LPJmL5 ³	Bloh et al., 2018	1981-2010	–	0.667***	0.496**	–	–	-0.163	0.152
GGCMI phase 3 ⁴	Jägermeyr et al., 2021	1981-2015	–	–	–	–	–	–	–
GGCMI phase 1 ⁵	Müller et al., 2017	1982-2006	0.90	0.91	0.76	0.76	0.90	0.42	0.61
			30-year averaged yield						
References	Period	COR	RMSE [kg ha ⁻¹]						
MATCRO-Maize	-	1981-2010	0.580**	4,008					
LPJ-GUESS ⁶	Olin et al., 2015	1996-2005	0.46	4,300					

¹ Countries-level comparison was conducted for 12 countries, which were detrended only for observation. p values are not shown.

^{2,3} Countries-level comparison was conducted for the top 10 producing countries, which were detrended via a 5-year moving average.

⁴ Twelve global gridded crop models were used. The COR shown here is the ensembled mean value for the 5 largest producing countries after detrending. p values are not shown.

⁵ Fourteen global gridded crop models were used. The COR of the global yield shown here is the minimum and maximum value, except for one nonsignificant correlation with the default setting. The COR of each country is the best correlation among the 14 models, including 3 different settings with statistical significance (p values are not shown). For both the global and country-level comparisons, a 5-year moving average was used to remove trends.

⁶ The 10-year average comparisons included all countries. p values are not shown.

Formatted Table

735 4.3 Model limitations

740 ~~MATCRO-Maize currently lacks explicit simulation of soil organic carbon and soil nitrogen mineralization. Instead, the effects of nitrogen supply are represented by describing the relationship between a broad range of nitrogen fertilization levels (Muchow, 1988) and specific leaf nitrogen (SLN), which subsequently affects photosynthetic capacity (V_{cmax}). While this simplification allows for global-scale application, it limits the model ability to represent nitrogen balance in maize yield at specific sites. Yield variations can be influenced by soil organic carbon and nitrogen, which are affected by farming practices and contribute to soil fertility (Ma et al., 2023). Future development could involve coupling MATCRO with a mechanistic soil nitrogen and carbon module to dynamic plant nitrogen balance. This would enhance the model ability to capture nitrogen dynamics under varying soil types and management practices.~~

745 The strong N_{fert} effect shown in the ~~evaluation~~ (the site in Brazil for the point scale) and comparison based on the N_{fert} (Figure 13). In the model, N_{fert} has the direct relationship with S_{ln} (Eq. (28)) and consequently affects V_{cmax25} (0) through the function $S_{ln}-V_{cmax25}$ (0) (Eq. (27)). Therefore, the strong N_{fert} effect is caused by either the former, the latter, or both processes. Few studies have explicitly shown time series changes in S_{ln} and $S_{ln}-V_{cmax}$ relationships from experiments. We used some of them to establish the functions shown in Eqs. (27) and (28) (Section 2.2.2) at this stage, resulting in a strong N_{fert} effect in the model. However, the intentional experiment indicated that the changed relationships could partly reproduce the adequate effect, which was observed in the FAOSTAT yield. This might mean that the established functions include a degree of uncertainty, and if we establish robust relationships based on other experimental data under more comprehensive conditions, it might be possible to improve the model in terms of the N_{fert} effect, leading to a more accurate simulation of maize yield. ~~Nitrogen effects are represented indirectly via SLN as a function of fertilizer rate and developmental stage, which constrains the model ability to capture nitrogen cycling in soils and plants.~~

755 In this study, we applied identical parameters to simulate the global yield across all grid cells and throughout the years without considering cultivar differences. As mentioned in Section 3.1.2, the trend of biomass accumulation would differ across growing season types. ~~A limitation of the current study is the use of global parameters at the site scale leads to discrepancies between site-level and country-level simulations. It partly arises from applying global parameters across different environments. Although MATCRO-Maize shows relatively weak correlations at the site scale due to the use of generalized parameters that do not account for local varieties and management, the model demonstrates consistent and statistically significant performance at country and global levels. This indicates that MATCRO-Maize is well suited for capturing large-scale yield patterns and for application in global gridded crop modeling, while recognizing its limited capacity for precise site-specific prediction. However, global-scale simulation results tend to overestimate yield due to LAI being directly driven by carbon balance, which can create feedbacks that produce excessively high LAI. Future improvements should incorporate constraints on LAI expansion and adjust leaf partitioning when LAI exceeds realistic levels.~~

765 Moreover, in major producing countries, such as the United States and China, some studies have shown that there is genetic gain in terms of maize yield (Cooper et al., 2014; Duvick et al., 2003; Liu et al., 2021). Such cultivar differences and long-

Deleted: The limitation of the current MATCRO-Maize is the

Deleted: validation

Formatted: Indent: First line: 0.5 cm

Deleted: Figures 12 and

term genetic improvements are not included in the current MATCRO-Maize. This finding indicates that the generic parameterization used in the model ~~are~~ simple in accounting for the diversity of crop cultivars (Lombardozi et al., 2020), partly leading to a gap between the simulations and observations, which is recognized as a limitation of the global model (Osborne et al., 2015). In addition, other important factors that are not considered in the current MATCRO also affect crop growth and final yield. These factors include biotic stresses (e.g., diseases, pests) and detailed management practices (e.g., plant density, as mentioned in Section 4.1). Further improvement to incorporate such factors with reliable N_{fert} -related functions could be needed to contribute to more accurate simulations and contribute to studies on the interaction between climate and agriculture.

Deleted: would be

Deleted: ¶

5 Conclusions

We developed a process-based crop model for maize yield estimation, called MATCRO-Maize, by incorporating C4 leaf-level photosynthesis and some crop-specific parameters into MATCRO. The model was first ~~evaluated~~ at the point scale, showing a somewhat reasonable accuracy considered with insufficient field-based information for parameterization. The calibrated parameters were set from point-scale experimental data and used uniformly in the global-scale simulation. MATCRO-Maize could represent the spatial distribution well and showed reasonable responses to climatic variability, where the results were comparable with those of other studies in terms of statistics. The strong nitrogen fertilizer effect was one of the MATCRO limitations, while the established functions related to nitrogen fertilizer in the model have a degree of uncertainty. Further experimental data under more comprehensive conditions might improve the model. Overall, MATCRO-Maize could contribute to climate effect studies through its ability to be integrated with the LSM for crop growth and the interactions between climate and agriculture.

Deleted: validated

790 Code and data availability.

The source code used in this study is archived at <https://doi.org/10.5281/zenodo.14869445> (Nagata et al., 2025).

Author contributions.

TK supervised this study and acquired the funding. YM developed the model source code and supervised this study. MN adjusted the source code for this study, performed the model simulations, and analysed the model output. MN prepared the original draft, which was supervised by TK, YM, and AY. The final manuscript was written by MN and approved by all the authors.

800 *Acknowledgements.*

This research was funded by JSPS KAKENHI grant number JP 22k18497. This study was partly supported by the Environment Research and Technology Development Fund (JPMEERF20202005) of the Environmental Restoration and Conservation Agency (ERCA) and JSPS KAKENHI Grant Number JP23H00351.

We thank all the contributors. Dr. Jean-Louis Durand at the French National Research Institute for Agriculture, Food, and Environment (INRAE) provided the observational data at the four sites, which we used to [evaluate](#) our model at the point scale.

Deleted: validate

References

- Allen R.G., Pereira L.S., Raes D., and Smith M.: Crop Evapotranspiration: Guidelines for Computing Crop Water Requirements - FAO Irrigation and drainage paper 56, FAO - Food and Agriculture Organization of the United Nations, Rome, 1998.
- Asseng, S., Ewert, F., Rosenzweig, C., Jones, J. W., Hatfield, J. L., Ruane, A. C., Boote, K. J., Thorburn, P. J., Rötter, R. P., Cammarano, D., Brisson, N., Basso, B., Martre, P., Aggarwal, P. K., Angulo, C., Bertuzzi, P., Biernath, C., Challinor, A. J., Doltra, J., Gayler, S., Goldberg, R., Grant, R., Heng, L., Hooker, J., Hunt, L. A., Ingwersen, J., Izaurralde, R. C., Kersebaum, K. C., Müller, C., Naresh Kumar, S., Nendel, C., O'Leary, G., Olesen, J. E., Osborne, T. M., Palosuo, T., Priesack, E., Ripoche, D., Semenov, M. A., Shcherbak, I., Steduto, P., Stöckle, C., Stratonovitch, P., Streck, T., Supit, I., Tao, F., Travasso, M., Waha, K., Wallach, D., White, J. W., Williams, J. R., and Wolf, J.: Uncertainty in simulating wheat yields under climate change, *Nat Clim Chang*, 3, 827–832, <https://doi.org/10.1038/NCLIMATE1916>, 2013.
- Ball J.T.: An analysis of Stomatal Conductance, Stanford University, CA, USA, 1988.
- Bassu, S., Brisson, N., Durand, J. L., Boote, K., Lizaso, J., Jones, J. W., Rosenzweig, C., Ruane, A. C., Adam, M., Baron, C., Basso, B., Biernath, C., Boogaard, H., Conijn, S., Corbeels, M., Deryng, D., De Sanctis, G., Gayler, S., Grassini, P., Hatfield, J., Hoek, S., Izaurralde, C., Jongschaap, R., Kemanian, A. R., Kersebaum, K. C., Kim, S. H., Kumar, N. S., Makowski, D., Müller, C., Nendel, C., Priesack, E., Pravia, M. V., Sau, F., Shcherbak, I., Tao, F., Teixeira, E., Timlin, D., and Waha, K.: How do various maize crop models vary in their responses to climate change factors?, *Glob Chang Biol*, 20, 2301–2320, <https://doi.org/10.1111/gcb.12520>, 2014.
- Von Bloh, W., Schaphoff, S., Müller, C., Rolinski, S., Waha, K., and Zachle, S.: Implementing the nitrogen cycle into the dynamic global vegetation, hydrology, and crop growth model LPJmL (version 5.0), *Geosci Model Dev*, 11, 2789–2812, <https://doi.org/10.5194/gmd-11-2789-2018>, 2018.
- Bonan, G. B., Lawrence, P. J., Oleson, K. W., Levis, S., Jung, M., Reichstein, M., Lawrence, D. M., and Swenson, S. C.: Improving canopy processes in the Community Land Model version 4 (CLM4) using global flux fields empirically inferred from FLUXNET data, *J Geophys Res*, 116, 1–22, <https://doi.org/10.1029/2010JG001593>, 2011.

- Bondeau, A., Smith, P. C., Zaehle, S., Schaphoff, S., Lucht, W., Cramer, W., Gerten, D., Lotze-campen, H., Müller, C., Reichstein, M., and Smith, B.: Modelling the role of agriculture for the 20th century global terrestrial carbon balance, *Glob Chang Biol*, 13, 679–706, <https://doi.org/10.1111/J.1365-2486.2006.01305.X>, 2007.
- 835 Boomsma, C. R., Santini, J. B., Tollenaar, M., and Vyn, T. J.: Maize Morphophysiological Responses to Intense Crowding and Low Nitrogen Availability: An Analysis and Review, *Agron J*, 101, 1426–1452, <https://doi.org/10.2134/AGRONJ2009.0082>, 2009.
- Bouman, B. A. M., Kropff, M., Tuong, T., Wopereis, M., ten Berge, H., and van Laar, H.: *Oryza2000: modeling lowland rice*, International Rice Research Institute, and Wageningen: Wageningen University and Research Centre, Philippines and
840 Wageningen, the Netherlands, 235 pp., 2001.
- Büchner, M. and Reyer, P. O. , C.: ISIMIP3a atmospheric composition input data (v1.2). ISIMIP Repository., <https://doi.org/https://doi.org/10.48364/ISIMIP.664235.2>, 2022.
- Campbell, G. S. and Norman, J. M.: *An Introduction to Environmental Biophysics*, Springer New York, NY, New York, USA, <https://doi.org/10.1007/978-1-4612-1626-1>, 1998.
- 845 [Cao, J., Zhao Z., Xiangzhong L., Yuchuan L., Jialu X., Jun X., Jichong H., and Fulu T. Mapping global yields of four major crops at 5-minute resolution from 1982 to 2015 using multi-source data and machine learning. *Scientific Data* 12, 1, 357, 2025.](#)
- Capristo, P. R., Rizzalli, R. H., and Andrade, F. H.: Ecophysiological Yield Components of Maize Hybrids with Contrasting Maturity, *Agronomy Jomal*, 99, 1111–1118, <https://doi.org/10.2134/agronj2006.0360>, 2007.
- 850 Ciampitti, I. A. and Vyn, T. J.: A comprehensive study of plant density consequences on nitrogen uptake dynamics of maize plants from vegetative to reproductive stages, *Field Crops Res*, 121, 2–18, <https://doi.org/10.1016/J.FCR.2010.10.009>, 2011.
- Ciampitti, I. A., Murrell, S. T., Camberato, J. J., Tuinstra, M., Xia, Y., Friedemann, P., and Vyn, T. J.: Physiological dynamics of maize nitrogen uptake and partitioning in response to plant density and N stress factors: I. Vegetative phase, *Crop Sci*, 53, 2105–2119, <https://doi.org/10.2135/cropsci2013.01.0040>, 2013a.
- 855 Ciampitti, I. A., Murrell, S. T., Camberato, J. J., Tuinstra, M., Xia, Y., Friedemann, P., and Vyn, T. J.: Physiological dynamics of maize nitrogen uptake and partitioning in response to plant density and nitrogen stress factors: II. reproductive phase, *Crop Sci*, 53, 2588–2602, <https://doi.org/10.2135/cropsci2013.01.0041>, 2013b.
- Collatz, G., Ribas-Carbo, M., and Berry, J.: Coupled Photosynthesis-Stomatal Conductance Model for Leaves of C4 Plants, *Functional Plant Biology*, 19, 519, <https://doi.org/10.1071/PP9920519>, 1992.
- 860 Collatz, G. J., Ball, J. T., Grivet, C., and Berry, J. A.: Physiological and environmental regulation of stomatal conductance, photosynthesis and transpiration: a model that includes a laminar boundary layer, *Agric For Meteorol*, 54, 107–136, [https://doi.org/10.1016/0168-1923\(91\)90002-8](https://doi.org/10.1016/0168-1923(91)90002-8), 1991.
- Cooper, M., Gho, C., Leafgren, R., Tang, T., and Messina, C.: Breeding drought-tolerant maize hybrids for the US corn-belt: discovery to product, *J Exp Bot*, 65, 6191–6204, <https://doi.org/10.1093/jxb/eru064>, 2014.
- 865

Dai, Y., Dickinson, R. E., and Wang, Y. P.: A two-big-leaf model for canopy temperature, photosynthesis, and stomatal conductance, *J Clim*, 17, 2281–2299, [https://doi.org/10.1175/1520-0442\(2004\)017<2281:ATMFCT>2.0.CO;2](https://doi.org/10.1175/1520-0442(2004)017<2281:ATMFCT>2.0.CO;2), 2004.

Drouet, J. L. and Bonhomme, R.: Effect of 3D Nitrogen, Dry Mass per Area and Local Irradiance on Canopy Photosynthesis Within Leaves of Contrasted Heterogeneous Maize Crops, *Ann Bot*, 93, 699–710, <https://doi.org/10.1093/aob/mch099>, 2004.

Duvick, D. N., Smith, J. S. C., and Cooper, M.: Long-Term Selection in a Commercial Hybrid Maize Breeding Program, in: *Plant Breeding Reviews*, Wiley, 109–151, <https://doi.org/10.1002/9780470650288.ch4>, 2003.

FAO: Agricultural production statistics 2000–2021, FAO, Rome, Italy, <https://doi.org/10.4060/cc3751en>, 2022.

FAOSTAT Production / Crops and livestock products - Metadata: <https://www.fao.org/faostat/en/#data/QCL/metadata>, last access: 31 December 2024.

Hajima, T., Watanabe, M., Yamamoto, A., Tatebe, H., Noguchi, M. A., Abe, M., Ohgaito, R., Ito, A., Yamazaki, D., Okajima, H., Ito, A., Takata, K., Ogochi, K., Watanabe, S., and Kawamiya, M.: Development of the MIROC-ES2L Earth system model and the evaluation of biogeochemical processes and feedbacks, *Geosci Model Dev*, 13, 2197–2244, <https://doi.org/10.5194/GMD-13-2197-2020>, 2020.

Hasegawa, T., Sawano, S., Goto, S. et al. A model driven by crop water use and nitrogen supply for simulating changes in the regional yield of rain-fed lowland rice in Northeast Thailand. *Paddy Water Environ* 6, 73–82, <https://doi.org/10.1007/s10333-007-0099-1>, 2008.

International Food Policy Research Institute (IFPRI), Global Spatially-Disaggregated Crop Production Statistics Data for 2010 Version 2.0, <https://doi.org/10.7910/DVN/PRFF8V>, Harvard Dataverse, V4, 2019.

Iizumi, T., Sakai, T. The global dataset of historical yields for major crops 1981-2016. *Scientific Data* 7, 97, <https://doi.org/10.1038/s41597-020-0433-7>, 2020.

Iizumi, T., Masaki, Y., Takimoto, T., and Masutomi, Y.: Aligning the harvesting year in global gridded crop model simulations with that in census reports is pivotal to national-level model performance evaluations for rice, *European Journal of Agronomy*, 130, 126367, <https://doi.org/10.1016/J.EJA.2021.126367>, 2021.

Intergovernmental Panel on Climate Change (IPCC): Technical Summary, in: *Climate Change 2022 – Impacts, Adaptation and Vulnerability*, Cambridge University Press, 37–118, <https://doi.org/10.1017/9781009325844.002>, 2023.

Jägermeyr, J., Müller, C., Ruane, A. C., Elliott, J., Balkovic, J., Castillo, O., Faye, B., Foster, I., Folberth, C., Franke, J. A., Fuchs, K., Guarin, J. R., Heinke, J., Hoogenboom, G., Iizumi, T., Jain, A. K., Kelly, D., Khabarov, N., Lange, S., Lin, T. S., Liu, W., Mialyk, O., Minoli, S., Moyer, E. J., Okada, M., Phillips, M., Porter, C., Rabin, S. S., Scheer, C., Schneider, J. M., Schyns, J. F., Skalsky, R., Smerald, A., Stella, T., Stephens, H., Webber, H., Zabel, F., and Rosenzweig, C.: Climate impacts on global agriculture emerge earlier in new generation of climate and crop models, *Nat Food*, 2, 873–885, <https://doi.org/10.1038/s43016-021-00400-y>, 2021.

Kanai, R. and Edwards, G. E.: The Biochemistry of C 4 Photosynthesis, in: *C4 plant biology*, Elsevier, 49–87, 1999.

Deleted: ∴ Global

Deleted: v1.2 and v1.3 aligned version [dataset], PANGAEA

Deleted: <https://doi.org/>

Deleted: 1594/PANGAEA.909132, 2019.

Kinose, Y. and Masutomi, Y.: Impact Assessment of Climate Change on Rice Yield Using a Crop Growth Model and Activities Toward Adaptation: Targeting Three Provinces in Indonesia, in: *Adaptation to Climate Change in Agriculture: Research and Practices*, Springer, Singapore, 67–80, https://doi.org/10.1007/978-981-13-9235-1_5, 2019.

Kinose, Y., Masutomi, Y., Shiotsu, F., Hayashi, K., Ogawada, D., Gomez-Garcia, M., Matsumura, A., Takahashi, K., and Fukushima, K.: Impact assessment of climate change on the major rice cultivar ciherang in Indonesia, *Journal of Agricultural Meteorology*, 76, 19–28, <https://doi.org/10.2480/agrmet.D-19-00045>, 2020.

Knapp, A. K. and Medina, E.: Success of C4 photosynthesis in the Field; Lessons from Communities Dominated by C4 plants, in: *C4 plant biology*, Elsevier, 251–277, 1999.

Lange, S., Mengel, M., Treu, S., and Büchner, M.: ISIMIP3a atmospheric climate input data (v1.0). ISIMIP Repository., <https://doi.org/10.48364/ISIMIP.982724>, 2022.

Lawrence D., Fisher R., Koven C., Oleson K., Swenson S., and Vertenstein M: Technical Description of Version 5.0 of the Community Land Model (CLM), NCAR, 2020.

Levis, S., Gordon, B. B., Erik Kluzek, Thornton, P. E., Jones, A., Sacks, W. J., and Kucharik, C. J.: Interactive Crop Management in the Community Earth System Model (CESM1): Seasonal influences on land-atmosphere fluxes, *J Clim*, 25, 4839–4859, <https://doi.org/10.1175/JCLI-D-11-00446.1>, 2012.

Lin, T., Song, Y., Lawrence, P., Khesghi, H. S., and Jain, A. K.: Worldwide Maize and Soybean Yield Response to Environmental and Management Factors Over the 20th and 21st Centuries, *J Geophys Res Biogeosci*, 126, 1–21, <https://doi.org/10.1029/2021JG006304>, 2021.

Liu, G., Yang, H., Xie, R., Yang, Y., Liu, W., Guo, X., Xue, J., Ming, B., Wang, K., Hou, P., and Li, S.: Genetic gains in maize yield and related traits for high-yielding cultivars released during 1980s to 2010s in China, *Field Crops Res*, 270, 108223, <https://doi.org/10.1016/j.fcr.2021.108223>, 2021.

Lombardozzi, D. L., Lu, Y., Lawrence, P. J., Lawrence, D. M., Swenson, S., Oleson, K. W., Wieder, W. R., and Ainsworth, E. A.: Simulating Agriculture in the Community Land Model Version 5, *J Geophys Res Biogeosci*, 125, 1–19, <https://doi.org/10.1029/2019JG005529>, 2020.

Long, S. P.: Environmental Responses, in: *C4 plant biology*, Elsevier, 12–21, 1999.

[Ma, Y., Dominic W., Mingsheng F., Lei Q., Rong L., and Johannes L. Global crop production increase by soil organic carbon. Nature Geoscience 16 \(12\): 1159-1165, 2023.](#)

Maruyama, A. and Kuwagata, T.: Coupling land surface and crop growth models to estimate the effects of changes in the growing season on energy balance and water use of rice paddies, *Agric For Meteorol*, 150, 919–930, <https://doi.org/10.1016/J.AGRFORMET.2010.02.011>, 2010.

Masutomi, Y.: The appropriate analytical solution for coupled leaf photosynthesis and stomatal conductance models for C3 plants, *Ecol Modell*, 481, 110306, <https://doi.org/10.1016/j.ecolmodel.2023.110306>, 2023.

- 935 Masutomi, Y., Ono, K., Mano, M., Maruyama, A., and Miyata, A.: A land surface model combined with a crop growth model
for paddy rice (MATCRO-Rice v. 1) - Part 1: Model description, *Geosci Model Dev*, 9, 4133–4154,
<https://doi.org/10.5194/gmd-9-4133-2016>, 2016a.
- Masutomi, Y., Ono, K., Takimoto, T., Mano, M., Maruyama, A., and Miyata, A.: A land surface model combined with a crop
growth model for paddy rice (MATCRO-Rice v. 1) - Part 2: Model validation, *Geosci Model Dev*, 9, 4155–4167,
940 <https://doi.org/10.5194/gmd-9-4155-2016>, 2016b.
- Muchow, R. C.: Effect of nitrogen supply on the comparative productivity of maize and sorghum in a semi-arid tropical
environment I. Leaf growth and leaf nitrogen, *Field Crops Res*, 18, 1–16, [https://doi.org/10.1016/0378-4290\(88\)90055-X](https://doi.org/10.1016/0378-4290(88)90055-X),
1988.
- Muchow, R. C. and Sinclair, T. R.: Nitrogen response of leaf photosynthesis and canopy radiation use efficiency in field-
945 grown maize and sorghum, *Crop Sci*, 34, 721–727, <https://doi.org/10.2135/CROPSCI1994.0011183X003400030022X>,
1994.
- Nagata, M., Yusara, A., Kato, T., and Masutomi, Y.: Development of global maize production model MATCRO-Maize version
1.0 (MATCRO-Maize ver1.0) , Zenodo [code and dataset], <https://doi.org/10.5281/zenodo.14869444>, 2025.
- Oleson, K. W., Lead, D. M. L., Bonan, G. B., Drewniak, B., Huang, M., Koven, C. D., Levis, S., Li, F., Riley, W. J., Subin,
950 Z. M., Swenson, S. C., Thornton, P. E., Bozbiyik, A., Fisher, R., Heald, C. L., Kluzek, E., Lamarque, J.-F., Lawrence, P.
J., Leung, L. R., Lipscomb, W., Muszala, S., Ricciuto, D. M., Sacks, W., Sun, Y., Tang, J., and Yang, Z.-L.: Technical
Description of version 4.5 of the Community Land Model (CLM) Coordinating Lead Authors, NCAR, 2013.
- Olin, S., Lindeskog, M., Pugh, T. A. M., Schurgers, G., Wärlind, D., Mishurov, M., Zaehle, S., Stocker, B. D., Smith, B., and
Armeth, A.: Soil carbon management in large-scale Earth system modelling: Implications for crop yields and nitrogen
955 leaching, *Earth System Dynamics*, 6, 745–768, <https://doi.org/10.5194/ESD-6-745-2015>, 2015.
- Osborne, T., Slingo, J., Lawrence, D., and Wheeler, T.: Examining the interaction of growing crops with local climate using a
coupled crop-climate model, *J Clim*, 22, 1393–1411, <https://doi.org/10.1175/2008JCLI2494.1>, 2009.
- Osborne, T., Gornall, J., Hooker, J., Williams, K., Wiltshire, A., Betts, R., and Wheeler, T.: JULES-crop: a parametrisation of
crops in the Joint UK Land Environment Simulator, *Geosci. Model Dev*, 8, 1139–1155, [https://doi.org/10.5194/gmd-8-](https://doi.org/10.5194/gmd-8-1139-2015)
960 [1139-2015](https://doi.org/10.5194/gmd-8-1139-2015), 2015.
- Paponov, I. A. and Engels, C.: Effect of nitrogen supply on leaf traits related to photosynthesis during grain filling in two
maize genotypes with different N efficiency, *Journal of Plant Nutrition and Soil Science*, 166, 756–763,
<https://doi.org/10.1002/jpln.200320339>, 2003.
- Paponov, I. A., Sambo, P., Erley, G. S. auf'm., Presterl, T., Geiger, H. H., and Engels, C.: Grain yield and kernel weight of
965 two maize genotypes differing in nitrogen use efficiency at various levels of nitrogen and carbohydrate availability during
flowering and grain filling, *Plant Soil*, 272, 111–123, <https://doi.org/10.1007/s11104-004-4211-7>, 2005.

- Peng, B., Guan, K., Chen, M., Lawrence, D. M., Pokhrel, Y., Suyker, A., Arkebauer, T., and Lu, Y.: Improving maize growth processes in the community land model: Implementation and evaluation, *Agric For Meteorol*, 250–251, 64–89, <https://doi.org/10.1016/j.agrformet.2017.11.012>, 2018.
- 970 Penning de Vries, F. W. T., Jansen, D. M., ten Berge, H. F. M., and Bakema, A.: Simulation of ecophysiological process of growth in several annual crops, Centre for Agricultural Publishing and Documentation (Pudoc), Wageningen, the Netherlands, 1989.
- Portmann, F. T., Siebert, S., and Döll, P.: MIRCA2000—Global monthly irrigated and rainfed crop areas around the year 2000: A new high-resolution data set for agricultural and hydrological modeling, *Global Biogeochem Cycles*, 24, GB1011, <https://doi.org/10.1029/2008GB003435>, 2010.
- 975 [Ruane, A.C., Goldberg, R., and Chrysanthacopoulos, J. AgMIP climate forcing datasets for agricultural modeling: Merged products for gap-filling and historical climate series estimation, *Agr. Forest Meteorol.*, 200, 233–248, doi:10.1016/j.agrformet.2014.09.016, 2015.](https://doi.org/10.1016/j.agrformet.2014.09.016)
- Sage, R. F.: Why C4 photosynthesis?, in: *C4 plant biology*, Elsevier, 3–16, 1999.
- 980 Schaphoff, S., Forkel, M., Müller, C., Knauer, J., Von Bloh, W., Gerten, D., Jägermeyr, J., Lucht, W., Rammig, A., Thonicke, K., and Waha, K.: LPJmL4 - A dynamic global vegetation model with managed land - Part 2: Model evaluation, *Geosci Model Dev*, 11, 1377–1403, <https://doi.org/10.5194/gmd-11-1377-2018>, 2018.
- Sellers, P. J., Randall, D. A., Collatz, G. J., Berry, J. A., Field, C. B., Dazlich, D. A., Zhang, C., Collelo, G. D., and Bounoua, L.: A Revised Land Surface Parameterization (SiB2) for Atmospheric GCMS. Part I: Model Formulation, *J Clim*, 9, 676–705, [https://doi.org/10.1175/1520-0442\(1996\)009<0676:ARLSPF>2.0.CO;2](https://doi.org/10.1175/1520-0442(1996)009<0676:ARLSPF>2.0.CO;2), 1996.
- 985 Sinclair, T. R. and Horie, T.: Leaf Nitrogen, Photosynthesis, and Crop Radiation Use Efficiency: A Review, *Crop Sci*, 29, 90–98, <https://doi.org/10.2135/cropsci1989.0011183X002900010023x>, 1989.
- Takata, K., Emori, S., and Watanabe, T.: Development of the minimal advanced treatments of surface interaction and runoff, *Glob Planet Change*, 38, 209–222, [https://doi.org/10.1016/S0921-8181\(03\)00030-4](https://doi.org/10.1016/S0921-8181(03)00030-4), 2003.
- 990 Tsvetinskaya, E. A., Mearns, L. O., and Easterling, W. E.: Investigating the Effect of Seasonal Plant Growth and Development in Three-Dimensional Atmospheric Simulations. Part II: Atmospheric Response to Crop Growth and Development, *J Clim*, 14, 711–729, [https://doi.org/10.1175/1520-0442\(2001\)014<0711:ITEOSP>2.0.CO;2](https://doi.org/10.1175/1520-0442(2001)014<0711:ITEOSP>2.0.CO;2), 2001.
- Tubiello, F. N. and Ewert, F.: Simulating the effects of elevated CO2 on crops: approaches and applications for climate change, *European Journal of Agronomy*, 18, 57–74, [https://doi.org/10.1016/S1161-0301\(02\)00097-7](https://doi.org/10.1016/S1161-0301(02)00097-7), 2002.
- 995 Volkholz, J. and Müller, C.: ISIMIP3 soil input data (v1.0). ISIMIP Repository., <https://doi.org/10.48364/ISIMIP.942125>, 2020.
- Volkholz, J. and Ostberg, S.: ISIMIP3a N-fertilizer input data (v1.2). ISIMIP Repository., <https://doi.org/10.48364/ISIMIP.311496.2>, 2022.

Deleted: Rienecker, M. M., Suarez, M. J., Gelaro, R., Todling, R., Bacmeister, J., Liu, E., Bosilovich, M. G., Schubert, S. D., Takacs, L., Kim, G. K., Bloom, S., Chen, J., Collins, D., Conaty, A., Da Silva, A., Gu, W., Joiner, J., Koster, R. D., Lucchesi, R., Molod, A., Owens, T., Pawson, S., Pegion, P., Redder, C. R., Reichle, R., Robertson, F. R., Ruddick, A. G., Sienkiewicz, M., and Woollen, J.: MERRA: NASA's modern-era retrospective analysis for research and applications, *J Clim*, 24, 3624–3648, <https://doi.org/10.1175/JCLI-D-11-00015.1>, 2011.

Vos, J., Van Der Putten, P. E. L., and Birch, C. J.: Effect of nitrogen supply on leaf appearance, leaf growth, leaf nitrogen economy and photosynthetic capacity in maize (*Zea mays* L.), *Field Crops Res*, 93, 64–73, <https://doi.org/10.1016/J.FCR.2004.09.013>, 2005.

1010 Wu, X., Vuichard, N., Ciais, P., Viovy, N., De Noblet-Ducoudré, N., Wang, X., Magliulo, V., Wattenbach, M., Vitale, L., Di Tommasi, P., Moors, E. J., Jans, W., Elbers, J., Ceschia, E., Tallec, T., Bernhofer, C., Grünwald, T., Moureaux, C., Manise, T., Ligne, A., Cellier, P., Loubet, B., Larmanou, E., and Ripoche, D.: ORCHIDEE-CROP (v0), a new process-based agro-land surface model: Model description and evaluation over Europe, *Geosci Model Dev*, 9, 857–873, <https://doi.org/10.5194/gmd-9-857-2016>, 2016.

1015 Yusara, A., Kato, T., Ainsworth, E. A., Battisti, R., Kumagai, E., Nakano, S., Wu, Y., Tsusumi-Morita, Y., Kobayashi, K., and Masutomi, Y.: Developing an eco-physiological process-based model of soybean growth and yield (MATCRO-Soy v.1): Model calibration and evaluation, *EGUsphere*[preprint], <https://doi.org/10.5194/egusphere-2025-453>, 2025.

Page 24: [1] Deleted	Astrid Yusara	9/23/25 6:07:00 PM
----------------------	---------------	--------------------

▼
▲

Page 24: [1] Deleted	Astrid Yusara	9/23/25 6:07:00 PM
----------------------	---------------	--------------------

▼
▲

Page 24: [1] Deleted	Astrid Yusara	9/23/25 6:07:00 PM
----------------------	---------------	--------------------

▼
▲

Page 24: [2] Deleted	Astrid Yusara	9/23/25 6:07:00 PM
----------------------	---------------	--------------------

✕

Page 24: [3] Deleted	Astrid Yusara	9/23/25 6:07:00 PM
----------------------	---------------	--------------------

▼

Page 24: [3] Deleted	Astrid Yusara	9/23/25 6:07:00 PM
----------------------	---------------	--------------------

▼
▲

Page 24: [3] Deleted	Astrid Yusara	9/23/25 6:07:00 PM
----------------------	---------------	--------------------

▼
▲