

Metal Layer Depletion during the Super Substorm on 4 November 2021

Gang Chen^{1*}, Yimeng Xu¹, Guotao Yang², Shaodong Zhang^{1,3}, Zhipeng Ren⁴, Pengfei Hu¹, Tingting Yu⁴, Fujun Wu⁵, Lifang Du², Haoran Zheng², Xuewu Cheng⁶, Faquan Li⁶, Min Zhang¹

¹School of Earth and Space Science and Technology, Wuhan University, Wuhan 430072, China

²National Space Science Center, Chinese Academy of Sciences, Beijing 100190, China

³School of Physics and Electronic Science, Guizhou Normal University, Guiyang 550025, China

⁴Key Laboratory of Earth and Planetary Physics, Institute of Geology and Geophysics, Chinese Academy of Sciences, Beijing 100029, China

⁵School of Physics, Henan Normal University, Xinxiang 453007, China

⁶Innovation Academy for Precision Measurement Science and Technology, Chinese Academy of Sciences, Wuhan 430071, China

Corresponding Author: Gang Chen (g.chen@whu.edu.cn)

This expression is not rigorous. The sentence is revised to avoid misunderstanding.

Abstract. Metal layer forms as a result of meteoric ablation and exist as a layer of metal elements between approximately 80 and 105 km altitude, and it provides information about the physics and chemistry of the boundary between the atmosphere and space. There are some studies about the wind field disturbances in Mesosphere and Low Thermosphere (MLT) region and the plasma variations in ionospheric E-region during magnetic storms, but no study on the impact of storms on the metal atom layers in mesosphere. During the super substorm on 4 Nov. 2021~~Due to the viscous force of air, the wind and electric field disturbances of a magnetic storm is hard to penetrate deep into the Earth's dense atmospheric region. It is generally believed that the influence of storms cannot reach the metal layers.~~ However, during the super substorm on 4 Nov. 2021, the atmospheric metal layers were observed to ~~decreasedeplete~~ by observations from three lidars at the mid-latitudes of China. The Na-, Ca and Ni densities on the storm day were significantly lower than ~~those on~~ in other days in October and November. The O/N₂ column density ratio observed by the Global Ultraviolet Imager (GUVI) on the storm day was much higher than ~~that on the~~ quiet days, and the numerical simulation results

demonstrate a substantial increase in atomic oxygen density at the heights of the metal layer. The increase in oxygen density may lead to the formation of more metal compounds, thus more metal atoms are consumed. This is an interesting phenomenon that magnetic storm can perturb the atmospheric metal layer through chemical reactions.

1 Introduction

Earth's metal layer is located at the height of the Kármán line, namely the boundary separating Earth's atmosphere and outer space. It is composed by the metallic elements, such as Fe, Mg, Na, K, Ca, Ni, etc, which are due to meteoroid ablation through collision with atmospheric molecules (Plane et al., 2015). These metallic elements undergo complex chemical reactions with atmospheric composition as well as dynamic processes with background wind field, eventually they are highly concentrated between 80-105 km altitude and form a metal layer that envelops the earth (Gardner et al., 2005, 2011; Hunten, 1967; Kopp, 1997). Components of the metal layer turn into compounds that deposit on the Earth's surface in about four years, and become the nourishment of the biosphere (Plane et al., 2015). Our knowledge of metal layers is still very limited, especially the mechanism for its distribution features in the Earth's atmosphere.

We have revised this sentence and provided the correct and appropriate expression.

Temporal variation of the metal layer is of primary concern, therein, Na layer is the most studied due to its high concentration of atomic particles and large scattering cross section. The density of Na peaks in winter, and reaches its lowest level in summer at all latitudes (Gong et al., 2013; Yi et al., 2009). ~~It is widely accepted that the Na layer exists at an altitude of 80-105 kilometers, and at lower altitudes, Na atoms are mostly converted into compounds due to chemical reactions. It is widely accepted that chemical reactions below 85 km altitude have an important impact on the density variation of Na atoms, but the chemical reaction above 85 km has little influence.~~ The background temperature and wind field at these heights are suggested to be responsible for the Na layer variation (Hickey and Plane, 1995). The density variation of Ca layer has two peaks in winter and summer, and the density peak in winter is higher (Gerding et al., 2000; Xun et al., 2024). It suggests to be relevant to the meteor input function (Plane et al., 2018). The density of Ni layer is maximum in winter and minimum in summer, similar to the seasonal cycle of Na layer (Jiao et al., 2022). At middle and high latitudes, the density of Fe layer peaks in winter and is lowest in summer. Conversely, at low latitudes,

Fe layer density reaches its maximum in spring and becomes lowest in the fall. The global distribution features of the mesospheric ozone and O₂ is attributed to result in these latitudinal differences (Yi et al., 2009; Raizada et al., 2003). K layer is different to other layers, its density is maximum in winter and summer, and minimum in spring and autumn (Eskola et al., 1998; Yue et al., 2017). There is a correlation between the Na and Ni layers, but not with other metal layers (Wu et al., 2022).

Solar flare and coronal mass ejection can induce geomagnetic storm on the earth, including intense currents in magnetosphere, changes in radiation belts, variations in the ionosphere, and heating of the thermosphere. Although the wind field and electric field disturbances in storms are hard to penetrate deep into the dense atmospheric region due to the viscous force of air, the wind field disturbances in Mesosphere and Low Thermosphere (MLT) region as well as the plasma variations in ionospheric E-region have been recorded and analyzed (Chen et al., 2023; Li et al., 2019; Li et al., 2024; Resende et al., 2020). However, due to the viscous force of air, the wind field and electric field disturbances are hard to penetrate deep into the dense atmospheric region. Whether a storm can affect the atmospheric layers as low as the Mesosphere and Low Thermosphere (MLT) region is a very interesting topic. In this height range, a number of studies on the wind field disturbances and variations in ionospheric E-region have been conducted (Chen et al., 2023; Li et al., 2024). The metal layer is also located in the MLT region, and the variations of the metal atom density in the metal layer they can be observed by lidar in the night with good weather condition. During the intense magnetic storm on 4 Nov. 2021, we have recorded the density decrease of different compositions in the metal layers at two locations, and ~~this~~ is the first time to introduce the influence of a storm on Earth's the metal atoms in the MLT region layer is recorded.

2 Data and Analysis Methods

In Oct. and Nov. 2021, two lidars in Yangqing, Beijing, China (40.42°N, 116.02°E) conducted the observations of Na, Ni and Ca layers, and one lidar in Pingquan, Hebei, China (41°N, 118.7°E) conducted the observations of Na layer. The Pingquan station is 237 km east of the Yangqing station. On the storm day of 4 Nov. 2021, all the three lidars were well operated. The Na and Ni observations were implemented by the dual-wavelength simultaneous detection system (Du et al., 2020; Jiao et al., 2015; Wu et al., 2021). The all-solid-state narrowband lidar is used to observe the Ca layer (Du et al., 2023).

We completely agree with the reviewer. The expression is somewhat oversimplified. We have revised this sentence and made the expression more rigorous.

This sentence is improved.

The dual-wavelength simultaneous detection system is used for Na and Ni observation and the all-solid-state narrowband lidar is used to observe the Ca layer (Du et al., 2020, 2023; Jiao et al., 2015; Wu et al., 2021). The time and height resolutions of these lidars are 33s and 96m, respectively. After long time integration to improve the Signal-to-Noise Ratio (SNR), the time resolutions of the output data are one hour. Due to the very low received signal power, the range integration is executed for the Ni layer observation and the range resolution of the output data becomes one kilometer.

The detection threshold of the lidars is defined as the background noise plus two times the noise standard deviation (Gao et al., 2015). The error of the lidar photon counting is inversely proportional to the square root of the return photon number, and the error of the measured metal atom density determined by the error of the photon counting is no more than 5%.

The absolute density can be estimated by the following standard lidar equation (Megie et al., 1978; Gerding et al., 2018; Chu and Papen, 2005).

$$n_{\text{atom}}(z) \equiv n_R(z_R) \frac{N_S(\lambda, z) - \frac{N_B \Delta t}{N_R(\lambda, z_R)} \frac{z^2}{z_R^2} \frac{\sigma_R(\lambda)}{\sigma_{\text{atom}}(\lambda)}}{1 - \frac{N_B \Delta t}{N_R(\lambda, z_R)} \frac{z^2}{z_R^2} \frac{\sigma_R(\lambda)}{\sigma_{\text{atom}}(\lambda)}}$$

Detection limits, uncertainty, and calibration methods of the lidars are added in the manuscript.

The reference altitude z_R is usually selected between 30-45 km, and z represents the altitude of the resonant fluorescence scattering region, and $n_R(z_R)$ is the atmospheric total number density at the reference altitude, which can be obtained from the NRLMSISE-00 model. $N_S(\lambda, z)$ and $N_R(\lambda, z_R)$ are the return photon numbers from the altitude z and the reference altitude z_R , respectively. N_B is the return photon number from the noise altitude generally at 150-190 km. λ represents the wavelength of Na (589.158 nm), Ca (422.6728 nm), and Ni (341.5744 nm). At 30 km altitude, the Rayleigh backscattering cross section $\sigma_R(589.158\text{nm})$ for Na is $4.14 \times 10^{-32} \text{ m}^2\text{sr}^{-1}$ and the effective backscattering cross section of resonance $\sigma_{\text{atom}}(589.158\text{nm})$ for Na is $5.22 \times 10^{-16} \text{ m}^2\text{sr}^{-1}$, the $\sigma_R(422.6728\text{nm})$ for Ca is $1.28 \times 10^{-31} \text{ m}^2\text{sr}^{-1}$ and the $\sigma_{\text{atom}}(422.6728\text{nm})$ for Ca is $2.23 \times 10^{-16} \text{ m}^2\text{sr}^{-1}$. At 40 km altitude, the $\sigma_R(341.5744\text{nm})$ for Ni is $3.55 \times 10^{-31} \text{ m}^2\text{sr}^{-1}$ and the $\sigma_{\text{atom}}(341.5744\text{nm})$ is $1.58 \times 10^{-17} \text{ m}^2\text{sr}^{-1}$. When calculating the absolute density of the Ni layer, it is also necessary to consider the branching fraction of the transition lines of the received return photons (Wu et al., 2021).

The number of return photons should be proportional to the flux of the transmitted laser photons. However, for large laser intensities, the returned fluorescence photons are no longer proportional to the laser power, but less than expectations. Thus, the saturation effect occurs. The content of Ni atoms in

metal layer is relatively low, so its saturation effect is generally not considered. The energy of Na lidar's laser is not high enough to produce saturation effect. The laser energy emitted by the Ca lidar is relatively large and the divergence angle of the laser beam is very small, thus the saturation effect of the Ca lidar must be considered. Refer to the method of Megie et al., (1978), Welsh and Gardner (1989), and Chu and Papen (2005) to estimate saturations, the ratio $\frac{N^{\text{Sat}}(z)}{N(z)} = 91.42\%$ is obtained (Wu et al., 2020), where $N^{\text{Sat}}(z)$ is the number of the photons received at height z , when the saturation effect occurs, and $N(z)$ is the number of photons received at height z , when there is no saturation effect. Thus, we have taken this ratio into account in the effective scattering cross sections to correct the Ca densities.


All the measured metal densities, including the density values of different metal atoms at different places, are quantitative and can be compared directly.–

GUUVI is a spectrograph imager on the Thermosphere Ionosphere Mesosphere Energy and Dynamics (TIMED) spacecraft. It can provide cross-track scanned images of the Earth's airglow and auroral emission in the Far UltraViolet (FUV) at wavelengths between 110.0 and 185.0 nm (Christensen et al., 2003). The OI 135.6 nm and N₂ Lyman-Birge-Hopfield (140-150 nm) data were extracted from the FUV spectral data and can be used to calculate the global thermospheric column density O/N₂ ratio, $\Sigma\text{O}/\text{N}_2$ (Zhang et al., 2004). The global O/N₂ variations can be used to describe the response of thermosphere-neutral components to geomagnetic activities.

Thermosphere Ionosphere Electrodynamics General Circulation Model (TIEGCM) is a self-consistent general circulation model of the coupled ionosphere-thermosphere system. It can solve the three-dimensional energy, continuity, momentum and electrodynamic equations of ions and neutral species (Richmond et al., 1992; Roble et al., 1988). The altitude range of the model is between 97 and 600 km and the spatial resolution is $1.25^\circ \times 1.25^\circ \times 0.25$ scale heights. The bottom boundary of TIEGCM is ~97 km, solving a series of equations in the thermosphere and ionosphere system self-consistently.

Solar extreme ultraviolet irradiance and the auroral electron precipitation are input in this model (Solomon and Qian, 2005; Roble and Ridley, 1987). The Heelis empirical model (Heelis et al., 1982) and the monthly tidal climatology (Hagan and Forbes, 2002, 2003) are used as the high-latitude and the lower boundary inputs, respectively.

141 3.1 Geomagnetic index



142

A red up arrow is added in the bottom plot to indicate the beginning of the simulated O density enhancement in MLT region.




Figure 1. Geomagnetic indexes. (a) Bz, (b) SYM/H, (c) AE, (d) PC and (e) kp indexes on 3-5 November 2021.


The green and red dashed boxes indicate the observation periods of the lidars in the nights of 3 and 4 Nov..

The red up arrow in the bottom plot is used to indicate the beginning of the simulated O density enhancement in MLT region.

Figure 1 shows the variations of Bz, SYM-H (can be seen as a high-resolution Dst), AE, PC and Kp indices on the three days of 3-5 Nov. 2021 from top to bottom. These geomagnetic indexes can be freely accessed in the OMNIWeb (1963). Before 17:00 UT on 3 Nov., there was no geomagnetic activity. The Interplanetary Magnetic Field (IMF) Bz component in Fig. 1a began to turn southward firstly at ~20:00 UT indicating the beginning of the storm and then it turned southward and northward for many times until ~12:38 UT on the next day. The SYM-H index in Fig. 1b shows that the Storm Sudden Commencement (SSC) began at ~20:00 UT on 3 Nov. and the storm continued to 5 Nov. The AE index in Fig. 1c presented several peaks. The first peak emerged at 21:45 UT on 3 Nov. with the maximum value of 2379 nT, and many AE peaks followed. There are two peaks exceeded 3000 nT at

9:13 UT and 11:25 UT on 34 Nov., indicating the occurrence of super substorm. The variation of the Polar Cap (PC) index of the Northern Hemisphere in Fig. 1d can be used as an indicator of the level of Joule heating. The PC index has exceeded 10 mV/m for many times, and the maximum PC index of 18.43 mV/m emerged at ~22:00 UT on 3 Nov., indicating the polar region Joule heating events have occurred many times, including one intense heating event at least. The Kp index used to characterize the magnitude of geomagnetic storms is shown in Fig. 1e. The Kp index exceeded 5 at 21:00 UT on 3 Nov. and reached its maximum of 7.7 at 9:00 UT on 4 Nov., indicating the strong geomagnetic storm happened. The green and red dashed boxes in Figure 1 are used to indicate the observation periods of the lidars in the nights of 3 and 4 Nov.. The geomagnetic indexes show that the influence of the storm began in the daytime of 4 Nov., at that time, the lidars didn't work.

3.2 Observation results of the lidars



This figure is rotated to make it consist of 4×5 subplots, and we enlarge this figure to make it occupy an entire page.





Figure 2. Lidar observations of the metal layers. The lidar observations of the Na density in Pingquan, as well as the Na, Ca and Ni density in Yanqing are listed from top to bottom row. The observations on the substorm day of 4 Nov. 2021 are displayed in the third column and highlighted by the red dashed box. The observations on the reference days are shown in the second and forth columns. The monthly average of November is shown in the fifth column. The first column shows the altitude profiles of the average metal density for a day, and the error bar indicates the uncertainty in density measurements. LT = UT + 8 h.

Morphological structures of the metal layers between 75-120 km height in the night are displayed in Fig. 2. The metal atoms dataset [can-beis](#) accessed [inat](#) the Zenodo website (Xu and Chen, 2025). The observations on the storm day of 4 Nov. 2021 are shown in the third column as highlighted by the red dashed box. The observations on the reference days are shown in the first and third columns. The monthly average of the observations in Nov. 2021 is displayed in the fifth column. The observed Na layer in Pingquan, the Na, Ca, and Ni layers in Yanqing are shown from top to bottom rows in sequence. The observation time of the lidars is set between 10 and 21 UT (18 and 5 LT of the next day), usually, and the weather conditions will determine the exact time of observation. Compared to the conference days and the monthly average, the density of the Na layer density in Pingquan, the Na, Ca and Ni layer density in Yanqing were obviously lower on 4 Nov. The maximum Na density in Figs. 2b and 2f was no more than 3600 ~~atomparticles~~/ cm^{-3} , and the Na layers were distributed in a narrow altitude range of 90-96 km, but the Na layer on other days emerged in wider altitude range with much higher peak metal density. The Ca layer in Fig. 2j on 4 Nov. almost disappeared and the maximum Ca density was as lower as 16.6 ~~atoms~~ $\cdot\text{cm}^{-3}$ ~~partieles~~/ cm^3 . The Ca layer presented very high density on the reference days, and the peak density had exceeded 80 ~~atoms~~ $\cdot\text{cm}^{-3}$ ~~partieles~~/ cm^3 . For a long time in Fig. 2i, the monthly average of Ca layer density exceeded 40 ~~atoms~~ $\cdot\text{cm}^{-3}$ ~~partieles~~/ cm^3 . The maximum density of the Ni layer in Fig. 2n was lower than 110 ~~atoms~~ $\cdot\text{cm}^{-3}$ ~~partieles~~/ cm^3 , also lower than ~~those~~ on the reference days in Figs. 2m and 2o, and much lower than ~~the~~ monthly averages in Fig. 2p.

The average altitude profiles of the measured metal layer density in a day are displayed in the first column of Figure 2. The average profiles recorded on the storm day are shown as red curves, those recorded on the reference days before and after are shown as blue and green dashed curves, respectively, and the average profiles of the monthly average are shown as the gray dashed curves. There is an error in the measurement of photon counts by a lidar, thus there is also uncertainty in metal density

measurements. The error bars on the profiles are used to indicate the uncertainty of the estimated metal density at different altitudes. The maximum Na, Ca and Ni densities on the storm day in Figure 2q, 2s, 2t are lower than those on the reference days as well as those of month averages, and their height distributions are also narrower. The maximum density of the red Na density profile in Figure 2r is lower than that of the green and gray profiles, but it is almost equal to that of the blue profile. The height distribution of red profile is narrower than that of the blue one, thus the Na layer density on the storm day was generally lower than that on 3 Nov.






Figure 3. Average column abundances of the metal layers. The observations of the (a) Na lidar in Pingquan, as well as the (b) Na lidar, (c) Ca lidar and (d) Ni lidar in Yanqing are listed from top to bottom row. The red bars indicate the observations on the substorm day. Each vertical bar shows the average column abundance between 75-120 km altitude in one night and all the valid observation results in October and November 2021 are displayed.— The uncertainty of the column abundance is exhibited on the top of each bar. The uncertainty of the column abundance is present as the error bar on the top of each bar.

The uncertainty of the column abundance is exhibited on the top of each bar.

Figure 3 presents the average column abundance of the metal layers on each observation day of Oct. and Nov. 2021 and the observations of the Na column abundance in Pingquan, the Na, Ca and Ni column abundance in Yanqing are listed from top to bottom in sequence. The average column abundance of a day can be obtained by integrating the density profile in the first column of Figure 2 along the height between 75-120 km. The error bar on the top of each bar shows the uncertainty of the column abundance. The uncertainty is about one thousandth of the column abundance, thus error bar is

compressed into almost a horizontal line.

In Oct. and Nov. 2021, except for the storm on 4 Nov., there was only a moderate storm occurring on 12 Oct. with the Kp index of 6.3 and all the lidars were not operated on that day. There is no obvious regularity in the day-to-day variation of the metal layer in the two months, but the lowest column abundances in all ~~the~~ four plots of Fig. 3 occurred on 4 Nov. 2021 as shown by the red bars. We have estimated the correlations between the different metal layer data in Fig. 3, and the correlation coefficients of Pingquan Na-Yanqing Na, Pingquan Na-Yanqing Ca, Pingquan Na-Yanqing Ni, Yangqing Na-Ca, Yangqing Na-Ni, and Yangqing Ca-Ni are 0.397, 3.578×10^{-4} , 0.507, 0.346, 0.603 and 0.169, respectively. There is little correlation between them. However, this does not mean that they are not correlated over a longer time scale. If there is no magnetic storm, the probability of all the four observations recording the lowest column abundances on the same day is 1.334×10^{-7} . Therefore, the metal layer depletion on 4 Nov. 2021 should be related to the storm.

We also can find that the abundance variation of the Ca has presented less correlation with other atoms, and the correlation coefficient of Pingquan Na-Yanqing Ca is much lower. Observations indicate that the Ni and Na layers show close correlations on the scale of hours (Wu et al., 2022). And the calcium layer is somewhat unique. Although Ca abundance has a similar elemental abundance to Na in meteorites, the Ca atom abundance in metal layer is roughly 2 orders of magnitude smaller than Na. Plane et al., (2018) suggested that CaOH and CaCO₃ are stable reservoirs for Ca in metal layer, as a result, more Ca atoms are converted to compounds and the Ca atom abundance in metal layer is much less than the Na atom abundance. The Ca abundance variation is more affected by chemical reactions, so it has less correlation with other metal atoms. The relevance between the Ca and Na in different places further decreases, thus the estimated correlation coefficient becomes very low.

We discuss the correlation coefficient in this paragraph.




Figure 4. GUVI O/N_2 distribution in the Eastern Hemisphere. The observations on (a) 3 November, (b) 4 November, (c) 5 November, 2021 and (d) the quiet day for reference are displayed. The black and gray dots represent the locations of Yanqing and Pingquan, respectively.

Joule heating in polar region during magnetic storm can produce equatorward wind surges and have influence on the MLT region wind field at mid-latitude (Li et al., 2019). However, it is difficult to explain the depletion of metal layers only by the variation of wind field. And the metal layer is electrically neutral and free from the electric field disturbance. During this storm, the atmospheric composition has changed. The data of the thermospheric $\Sigma O/N_2$ can be accessed from the NASA GUVI Team (2002). Figure 4d shows the average $\Sigma O/N_2$ in the quiet days of Oct. and Nov. 2021, and there was no obvious $\Sigma O/N_2$ enhancement region around the world. But in Figs. 4a-4c, the $\Sigma O/N_2$ enhancement region had extended to mid-latitudes on 3-5 Nov., and on the storm day of 4 Nov., the Yanqing and Pingquan observation stations had entered the maximum enhancement region as indicated


by the yellow, indicating that the thermospheric $\Sigma O/N_2$ over the two locations increased greatly.

4 Discussion

4.1 Mechanism for storm time O/N_2 variation

It is difficult to explain the density variations of the neutral metal atoms in the metal layer by only dynamic process, and there was no ~~significant~~ changes in the MLT region wind field as the observation of the meteor radar located in Beijing. The storm time O/N_2 variation as observed by the GUVI is very likely to induce metal layer depletion. Polar temperature increase due to the Joule heating leads to strong upwelling, and thermosphere O and N_2 move upwards with the upwelling (Prölss, 1980, 2011). Vertical advection during the upwelling will reduce O and enhance N_2 , thus lead to the depletion of the O/N_2 . When the downwelling occurs due to the decreasing temperature gradient, the O density increases and N_2 density decreases to enhance the O/N_2 (Yu et al., 2023). Kil et al. (2011) studied the O and N_2 disturbances in F layer during the storm on 20 Nov. 2003. They proposed that the different behaviors of O and N_2 perturbations can't be simply explained by the upwelling and downwelling in the upper atmosphere, and may be related to the global wind circulation, which may transfer the O perturbations from mid-high latitudes to middle-low latitudes, and this transferring process is more remarkable in the low F-region than the high F-region (Mayr et al., 1978; Meier et al., 1995). More importantly, it is necessary to distinguish the changes in oxygen and nitrogen atom density.

4.2 Simulation results of the TIEGCM






Figure 5. Simulated global distribution of the column O/N_2 ($\Sigma O/N_2$). Latitude and longitude/ local time maps show the $\Sigma O/N_2$ variations from 8:20 to 18:20 UT on 4 Nov. 2021 with 2 hour step. The gray curves in each panel are the zero values. The horizontal white dashed lines are the latitude of $\pm 60^\circ$. The gray circle is used to indicate the lidar stations [40°N, 116°E].

We apply the TIEGCM to investigate the O and N_2 variation in thermosphere and mesosphere. The simulated longitude-latitude distribution of the percentage changes of the $\Sigma O/N_2$ from 8:20 to 18:20 UT on 4 Nov. compared to the quiet days is shown in Figure 5. The O/N_2 dataset can be accessed in the Zenodo website (Yu, 2024). In this simulation, the O/N_2 variations are mainly attributed to the equatorward wind disturbances driven by the Joule heating and the influence of Coriolis force is in consideration, the details of the simulated results can refer to Yu et al., (2021; 2023). The horizontal white dashed lines show the latitudes of $\pm 60^\circ$. The simulated $\Sigma O/N_2$ over Yanqing and Pingquan is 1.0109, close to and the GUVI measurement of 1.0181 on the storm day, and the simulation results are quite accurate. Between 8:20-18:20 UT in Figure 5, the $\Sigma O/N_2$ over the lidar stations continued increasing. At 8:20 UT in Figure 5a, the increase of the $\Sigma O/N_2$ near the gray circle didn't exceed 20%,

and then it increased gradually until reaching its maximum at 16:20 UT in Figure 5e. At 18:20 UT in Figure 5f, the $\Sigma O/N_2$ began to decrease, but it is still higher than ~~that on~~ the quiet days.




Figure 6. Simulated oxygen density variations at different altitudes. (a) Storm-quiet time ratio of O and N₂ density between 95-200 km altitude from 5:40 to 20:00 UT on 4 November 2021. Oxygen density variation at (b) 150, 160 and 170 km altitudes, as well as at (c) 100, 102 and 104 km altitudes on the substorm day and reference day.

Both increase of O density and decrease of N₂ density can enhance the O/N₂. Thus, we investigate the O and N₂ density variation at different heights as shown in Figure 6. Each curve in Fig. 6a shows the height profile of the storm-quiet time percentage of O density (solid curve) and N₂ density (dashed curve) between 97-200 km. The quiet time values come from the simulated data on 3 Nov.. The profiles of different time are displayed by different colors. Between 5:40-20:00 UT on 4 Nov., the AE index in Figure 1c reached its maximum and the lidars recorded the lowest metal layer density in this night. As indicated by the height profiles in the same duration in Figure 6a, the enhancement of the O/N₂ ratio between 100-200 km altitude was mainly due to the increase of O as well as ~~the both~~ ~~great~~ decrease of N₂ and increase of O density. N₂ is inert gas and its density variation is unlikely to have a great influence on the density of the metal atoms, thus we mainly focus on the O density variation. Figures 6b and 6c show the O density variations on the storm day compared with ~~those~~ on the reference day ~~of 4 Nov.~~ at ~~difference~~ lower altitudes. ~~I~~in Figure 6b, the O density enhancements at 150-170 km altitudes were not obvious, and only appeared in a short duration between ~10:00-13:00 UT. However, at the lower heights in Figure 6c, the O density increased even higher and earlier. The enhancement occurred at ~2:30 UT on 4 Nov. and continued to the next day. The beginning of the O enhancement is indicated as a red up arrow in Figure 1e, and the metal layer dissipation should start in the daytime, when the lidars didn't work. The largest increase was 1.9×10^{10} ~~atoms~~ cm^{-3} at 100 km altitude at 14:00 UT. This increment is much higher than the density of the metal layer. As suggested by previous studies, the storm-time O density enhancement is more in the low F-layer due to the global wind circulation, and the increased oxygen in this region can extend to the E region due to the effect of downwelling (Yu et al., 2023; Kil et al., 2011). Thus, the O increases in lower thermosphere were bound to influence the O variations in the metal layer at an altitude of 80-100 km by molecular diffusion and downwelling, resulting in the O density increase there.

TIEGCM bottom boundary is ~97 km, But this does not affect our conclusion.

4.3 Mechanism for metal layer depletion

Metallic elements are not stable in the Earth's atmosphere, and they will constantly shift between ions, atoms, and compounds. The released Na atoms from meteoroids will be oxidized by O₃ to form NaO (R1) in MLT region, and further react with H₂O or H₂ to form NaOH (R2 and R3). NaHCO₃ is the recombination of NaOH with CO₂ (R4) and it is the major reservoir for Na on the bottom side of the metal layer (Gómez-Martín et al., 2017; Plane et al., 2015; Yuan et al., 2019). NaHCO₃ is converted

The detailed chemical pathways for Na and Ni compounds affecting Na and Ni atom abundances are also presented.

back to Na either by photolysis ($h\nu$) or by reaction with atom H (R5 & R6). While the O density increases, more Na atoms will become NaHCO_3 to reduce the Na density.







Ni atoms in MLT region is oxidized by O_3 and O_2 to form NiO and NiO_2 , respectively (R7 and R8). These two Ni compounds further react with O_3 , O_2 , CO_2 , and H_2O to form higher oxides, carbonates, and hydroxides (e.g. R9-R14), therein, NiOH , $\text{Ni}(\text{OH})_2$ and NiCO_3 is the major reservoir for Ni. These higher compounds will be converted to NiOH and NiO , which are finally converted back to Ni by reaction with atom H and O, as well as CO (R15-R17) (Daly et al., 2020). The increased O will convert more Ni atoms into the compound reservoir to dissipate the Ni layer.



The increase in O density during storm time can intervene in this cycle and lead to the depletion of the metal layer. Take the example of the Ca density, its dissipation is most severe compared to the monthly average. The Ca atoms in atmosphere will be converted into the compounds, such as CaO , CaOH , O_2CaCO_3 , OCaOH , CaCO_3 and so on (e.g. R18-R26). When the O density at the metal layer heights continues to increase, it is likely to promote the formation of O_2CaOH (R24), which slows the

conversion of CaOH to atomic Ca by reacting with atomic H (R27) (Gomez Martin and Plane, 2017). And then O₂CaOH will convert to the stable compound CaOH (R25 and R28) as indicated by the Equations (1) and (2). CaOH will further convert to the stable compound CaCO₃ and enter into a stable chemical cycle as indicated by R24, R29-R31 the Equations (3)–(6). CaOH and CaCO₃ are stable reservoirs for Ca (Plane et al., 2018). As a result, the increased O density converts more Ca atoms to the stable compound reservoirs, thus the Ca layer has suffered much severer dissipation during the storm as shown in Figure 2. As a result, the increased O density converts more Ca atoms to compounds, and thus the Ca density in the metal layer decreases significantly. Similarly, the increased O density also converts other metal atoms to the stable compounds to decrease their density.







Figure 7. Schematic of the metal layer depletion during a storm. The red circular arrows indicate that high-energy particles have entered the earth. The white transparent area indicates the atmospheric

expansion in polar region due to auroral heating, and the yellow arrows show the directions of the heating driven winds with the influence of Coriolis force. The color on the earth indicates the magnitude of the O/N₂, decreasing in order from red, yellow, green to blue. The translucent dots represent oxygen atoms. Layers of earth's atmosphere are displayed. The red circle around the earth is used to represent the metal layer, and lighter colors indicate lower density.

Figure 7 is drawn to demonstrate the possible mechanism for the metal layer depletion during the super substorm. The high-energy particles enter the earth to enhance Joule heating and increase upper atmosphere temperature (Burns et al., 1995). The temperature enhancements change the horizontal pressure gradient, in conjunction with the ion drag, drive horizontal wind perturbations to modify the global thermospheric circulation. It should be noted that the storm-time horizontal winds are usually influenced by the Coriolis force due to the Earth corotation during the storm evolution (Yu et al., 2022). The storm-induced divergent and convergent winds induce the atmospheric upwelling and downwelling, thus composition changes at high latitudes, which are transported to lower latitudes by the horizontal winds. O increases mainly occur at lower ~~latitudes~~altitudes (Yu et al., 2021), like that around the height of the metal layer as shown in Figure 6. Metal elements in metal layer cycle between atomic, ionic, and compound states, reaching a conversion equilibrium. The enhancement of O density in the storm has broken this equilibrium, more metal atoms will react with the increased O to produce more metal compounds, thus the density of metal atoms in the metal layer decreasessd as the lidars observed.

5 Conclusions

Though the density of atmospheric metal layers is varying day to day, the Na, Ca and Ni densities recorded by the Yanqing lidar on the storm day were significantly lower than ~~those~~ on other days, and the Na density recorded by the Pingquan lidar was also lowest in the two months. It is quite certain that the super substorm has induced the depletion of the metal layers. The increase of the O density in the MLT region is considered to consume the metal atoms.

Magnetic storms affect the ionosphere and thermosphere by disturbing the electric field and wind field usually. Metal layers are located at a lower altitude, where the ambient atmospheric density is much greater. It is generally believed that the influence of magnetic storms is difficult to reach the altitudes of metal layer. Our observations present two very interesting phenomena, one is that the

effects of magnetic storm have penetrated through the thermosphere and reached the MLT region, and the other is that the pathway of storm influence is no longer the usual electrodynamic process, but a chemical process. The storm has provided us a very nice opportunity to record the responses of the metal layer to breaking of the chemical equilibrium in MLT region. It implies that magnetic storms have a broader impact on Earth's atmosphere, and a new horizon is opened up for us to study of the impact of magnetic storms on the Earth's space environment.

This observation was carried out only at mid-latitudes of China. However, what happens at other latitudes is unknown, especially more significant impact may occur at higher latitudes as Figure 4 indicated. Thus, it is necessary to record the storm influence on mesospheric metal layers at different latitudes for more latitude coverage. The new observation station in Mohe, Heilongjiang province, China (122.37°E, 53.5°N) is now in operation. We will report more observation results at higher latitude in the coming years.

~~More observations at different latitudes for the storm influence on metal layers are needed, especially more significant impact may occur at higher latitudes as Figure 4 indicated. Furthermore, it is also necessary to pay attention to the influence of magnetic storm on lower atmosphere. These observations will deepen our understanding of the impacts of magnetic storms on the atmospheric chemical balance and environmental quality, and provide us more understanding of the physics and chemistry of the boundary between the atmosphere and space.~~

The conclusion section is revised according to the reviewer's suggestions.

Data availability. The data of metal atoms can be publicly obtained in the Zenodo website (<https://zenodo.org/records/15003984>) (Xu and Chen, 2025). The geomagnetic indices can be freely accessed in the OMNIWeb (1963) (<https://omniweb.gsfc.nasa.gov/>). TIMED GUVI observations of O/N₂ column density ratios are available from <https://guvitimed.jhuapl.edu/> (NASA GUVI Team, 2002). The TIE-GCM simulation datasets supporting this study are available in the Zenodo website (<https://zenodo.org/records/14232871>) (Yu, 2024).

Author contributions. Y.X. wrote the manuscript and G. C. rewrote it. Y. X. conducted comprehensive data analysis. G. C. investigated the physicochemical mechanisms. G. Y., P. H., and M. Z. provided helpful suggestions and revised the manuscript. Z. R. and T. Y. provided TIEGCM simulation results. F. W. assisted in nickel atoms data processing. G. Y., L. D., H. Z., X. C., and F. L. organized the datasets.

450 All the authors discussed the results and commented on the manuscript.

451 **Competing interests.** The contact author has declared that none of the authors has any competing
452 interests.

453 **Acknowledgements.** This project was supported by the National Natural Science Foundation of China
454 (42274197 and 41722404). We acknowledge the use of the lidar data from the Chinese Meridian
455 Project.

456 Reference

457 Burns, A. G., Killeen, T. L., Deng, W., Carignan, G. R., & Roble, R. G. Geomagnetic storm effects in
458 the low- to middle-latitude upper thermosphere. *Journal of Geophysical Research*, 100(A8),
459 14673. <https://doi.org/10.1029/94ja03232>, 1995

460 Chen, G. et al.: Anomalous sporadic-E enhancements and field-aligned irregularities at low-latitudes
461 during the intense geomagnetic activities on 17–18 March 2015. *Journal of Geophysical Research:*
462 *Space Physics*, 128(10), e2023JA031856. <https://doi.org/10.1029/2023JA031856>, 2023.

463 Christensen, A. B. et al.: Initial observations with the Global Ultraviolet Imager (GUVI) in the NASA
464 TIMED satellite mission. *Journal of Geophysical Research: Space Physics*, 108(A12), 1,451.
465 <https://doi.org/10.1029/2003JA009918>, 2003.

466 Chu, X. and Papen, G. C.: Resonance fluorescence lidar for measurements of the middle and upper
467 atmosphere, in *Laser Remote Sensing*, Fujii, T. and Fukuchi, T., eds. (CRC Press), 179-432, 2005.

468 Daly, S. M., Feng, W., Mangan, T. P., Gerding, M., & Plane, J. M. C.: The Meteoric Ni Layer in the
469 Upper Atmosphere. *Journal of Geophysical Research: Space Physics*, 125, e2020JA028083.
470 <https://doi.org/10.1029/2020JA028083>, 2020.

471 Du, L. F. et al.: Continuous detection of diurnal sodium fluorescent lidar over Beijing in China.
472 *Atmosphere*, 11(1), 118. doi:10.3390/atmos11010118, 2020.

473 Du, L. F. et al.: The all-solid-state narrowband lidar developed by optical parametric
474 oscillator/amplifier (OPO/OPA) technology for simultaneous detection of the Ca and Ca⁺ layers.
475 *Remote Sensing*, 15(18), 4,566. doi:10.3390/rs15184566, 2023.

Eska, V., Höffner, J. and von Zahn, U.: Upper atmosphere potassium layer and its seasonal variations at 54°N. *Journal of Geophysical Research: Space Physics*, 103(A12), 29,207–29,214. <https://doi.org/10.1029/98JA02481>, 1998.

Gao, Q., Chu, X., Xue, X., Dou, X., Chen, T., and Chen, J., Lidar observations of thermospheric Na layers up to 170 km with a descending tidal phase at Lijiang (26.7°N, 100.0°E), China, *J. Geophys. Res. Space Physics*, 120, 9213–9220, doi:10.1002/2015JA021808, 2015.

Gardner, C. S. et al.: Seasonal variations of the Na and Fe layers at the south pole and their implications for the chemistry and general circulation of the polar mesosphere. *Journal of Geophysical Research: Atmospheres*, 110(D10), D10302. <https://doi.org/10.1029/2004JD005670>, 2005.

Gardner, C. S. et al.: Seasonal variations of the mesospheric Fe layer at Rothera, Antarctica (67.5°S, 68.0°W). *Journal of Geophysical Research: Atmospheres*, 116(D2), D02304. <https://doi.org/10.1029/2010JD014655>, 2011.

Gerding, M., Alpers, M., von Zahn, U., Rollason, R. J. and Plane, J. M. C.: Atmospheric Ca and Ca⁺ layers: Midlatitude observations and modeling. *Journal of Geophysical Research: Space Physics*, 105(A12), 27,131–27,146. <https://doi.org/10.1029/2000JA900088>, 2000.

Gerding, M., Daly, S., and Plane, J.M.C.. Lidar Soundings of the Mesospheric Nickel Layer Using Ni(3F) and Ni(3D) Transitions. *Geophys. Res. Lett.* 46. doi:10.1029/2018GL080701, 2018.

Gómez Martin, J. C. and Plane, J. M. C.: Reaction kinetics of CaOH with H and O₂ and O₂CaOH with O: Implications for the atmospheric chemistry of meteoric calcium. *ACS Earth and Space Chemistry*, 1(7), 431–441. <https://doi.org/10.1021/acsearthspacechem.7b00072>, 2017.

Gong, S. H. et al.: Lidar studies on the nighttime and seasonal variations of background sodium layer at different latitudes in China. *Chinese Journal of Geophysics*, 56(8), 2511–2521. doi:10.6038/cjg20130802, 2013.

Hagan, M. E., and Forbes, J. M.: Migrating and nonmigrating diurnal tides in the middle and upper atmosphere excited by tropospheric latent heat release. *Journal of Geophysical Research: Atmospheres*, 107(D24), 4,754. <https://doi.org/10.1029/2001JD001236>, 2002.

Hagan, M. E., and Forbes, J. M.: Migrating and nonmigrating semidiurnal tides in the upper atmosphere excited by tropospheric latent heat release. *Journal of Geophysical Research: Space Physics*, 108(A2), 1,062. <https://doi.org/10.1029/2002JA009466>, 2003.

Heelis, R. A., Lowell, J. K. and Spiro, R. W.: A model of the high-latitude ionospheric convection

pattern. Journal of Geophysical Research: Space Physics, 87(A8), 6,339–6,345.
<https://doi.org/10.1029/JA087iA08p06339>, 1982.

Hickey, M. P. and Plane, J. M. C.: A chemical-dynamical model of wave-driven sodium fluctuations. Geophysical Research Letters, 22(20), 2861–2864. <https://doi.org/10.1029/95GL02784>, 1995.

Hunten, D.M.: Spectroscopic studies of the twilight airglow. Space Science Reviews 6, 493–573. <https://doi.org/10.1007/BF00173704>, 1967.

Jiao, J. et al.: First report of sporadic K layers and comparison with sporadic Na layers at Beijing, China (40.6°N, 116.2°E). Journal of Geophysical Research: Space Physics, 120(6), 5,214–5,225. <https://doi.org/10.1002/2014JA020955>, 2015.

Jiao, J. et al.: A Comparison of the midlatitude nickel and sodium layers in the mesosphere: Observations and modeling. Journal of Geophysical Research: Space Physics, 127(2), e2021JA030170. <https://doi.org/10.1029/2021JA030170>, 2022.

Kil, H., Kwak, Y.-S., Paxton, L. J., Meier, R. R. and Zhang Y.: O and N₂ disturbances in the F region during the 20 November 2003 storm seen from TIMED/GUVI. Journal of Geophysical Research: Space Physics, 116(A2), A02314. <https://doi.org/10.1029/2010JA016227>, 2011.

Kopp, E.: On the abundance of metal ions in the lower ionosphere. Journal of Geophysical Research: Space Physics, 102(A5), 9667–9674. <https://doi.org/10.1029/97JA00384>, 1997.

Li, J. Y. et al.: A modeling study of the responses of mesosphere and lower thermosphere winds to geomagnetic storms at middle latitudes. Journal of Geophysical Research: Space Physics, 124(5), 3,666–3,680. <https://doi.org/10.1029/2019JA026533>, 2019.

Li, Y. X. et al.: Observational evidence for the neutral wind responses in the mid-latitude lower thermosphere to the strong geomagnetic activity. Space Weather, 22(9), e2023SW003830. <https://doi.org/10.1029/2023SW003830>, 2024.

Mayr, H. G., Harris, I. and Spencer, N. W.: Some properties of upper atmosphere dynamics. Reviews of Geophysics, 16(4), 539–565. <https://doi.org/10.1029/RG016i004p00539>, 1978.

Megie, G., Bos, F., Blamont, J. and Chanin, M.: Simultaneous nighttime lidar measurements of atmospheric sodium and potassium, Planet. Space Sci. 26, 27-35, [https://doi.org/10.1016/0032-0633\(78\)90034-X](https://doi.org/10.1016/0032-0633(78)90034-X), 1978.

Meier, R. R., Cox, R., Strickland, D. J., Craven, J. D. and Frank L. A.: Interpretation of dynamics explorer far UV images of the quiet time thermosphere. Journal of Geophysical Research: Space

Physics, 100(A4), 5,777–5,794. <https://doi.org/10.1029/94JA02679>, 1995.

NASA GUVI Team.: GUVI level 3 data products [Dataset]. Thermospheric O/N₂.
<https://guvitimed.jhuapl.edu/>, 2002.

OMNIWeb.: Geomagnetic indices [Dataset]. OMNIWeb. Retrieved from
<https://omniweb.gsfc.nasa.gov/>, 1963.

Picone, J. M., Hedin, A. E., Drob, D. P., and Aikin, A. C.: NRLMSISE-00 empirical model of the
atmosphere: Statistical comparison and scientific issues. J. Geophys. Res. 107(A12): 1468.
doi:10.1029/2002JA009430, 2002.

Plane, J. M. C., Feng, W. and Dawkins, E. C. M.: The mesosphere and metals: Chemistry and changes.
Chemical Reviews, 115(10), 4497–4541. <https://doi.org/10.1021/cr500501m>, 2015.

Plane, J. M. C., Feng, W. H., Gómez Martín, J. C., Gerding, M., and Raizada, S.: A new model of
meteoric calcium in the mesosphere and lower thermosphere. Atmospheric Chemistry and Physics,
18(20), 14,799–14,811. <https://doi.org/10.5194/acp-18-14799-2018>, 2018.

Prölss, G. W.: Magnetic storm associated perturbations of the upper atmosphere: Recent results
obtained by satellite-borne gas analyzers. Reviews of Geophysics, 18(1), 183–202.
<https://doi.org/10.1029/RG018i001p00183>, 1980.

Prölss, G. W.: Density perturbations in the upper atmosphere caused by the dissipation of solar wind
energy. Surveys in Geophysics, 32, 101–195. <https://doi.org/10.1007/s10712-010-9104-0>, 2011.

Raizada, S., and Tepley, C. A.: Seasonal variation of mesospheric iron layers at Arecibo: First results
from low-latitudes. Geophysical Research Letters, 30(2), 1082.
<https://doi.org/10.1029/2002GL016537>, 2003.

Resende, L. C. A., Shi, J. K., Denardini, C. M., Batista, I. S., Nogueira, P. A. B., Arras, C., et al.: The
influence of disturbance dynamo electric field in the formation of strong sporadic E layers over
Boa Vista, a low-latitude station in the American sector. Journal of Geophysical Research: Space
Physics, 125(7), e2019JA027519. <https://doi.org/10.1029/2019JA027519>, 2020.

Richmond, A. D., Ridley, E. C., and Roble, R. G.: A thermosphere/ionosphere general circulation
model with coupled electrodynamics. Geophysical Research Letters, 19(6), 601–604.
<https://doi.org/10.1029/92GL00401>, 1992.

Roble, R. G., and Ridley, E. C.: An auroral model for the NCAR thermosphere general circulation
model (TGCM). Annales Geophysicae, 5, 369–382., 1987.

566 Roble, R. G., Ridley, E. C., Richmond, A. D., and Dickinson, R. E.: A coupled
 567 thermosphere/ionosphere general circulation model. *Geophysical Research Letters*, 15(12),
 568 1,325-1,328. <https://doi.org/10.1029/GL015i012p01325>, 1988.

569 Solomon, S. C., and Qian, L.: Solar extreme-ultraviolet irradiance for general circulation models.
 570 *Journal of Geophysical Research: Space Physics*, 110(A10), A10306.
 571 <https://doi.org/10.1029/2005JA011160>, 2005.

572 Welsh, B. and Gardner C.: Nonlinear resonant absorption effects on the design of resonance
 573 fluorescence lidars and laser guide stars, *Appl. Opt.*, 28, 4141-4153,
 574 <https://doi.org/10.1364/AO.28.004141>, 1989.

575 Wu, F., Zheng, H., Cheng, X., Yang, Y., Li, F., Gong, S., Du, L., Wang, J., Yang, G., Simultaneous
 576 detection of the Ca and Ca⁺ layers by a dual-wavelength tunable lidar system, *Appl Opt.*, 59(13),
 577 4122-4130, doi: 10.1364/AO.381699, 2020.

578 Wu, F. J. et al.: Lidar observations of the upper atmospheric nickel layer at Beijing (40°N,116°E).
 579 *Journal of Quantitative Spectroscopy and Radiative Transfer*, 260, 107, 468.
 580 <https://doi.org/10.1016/j.jqsrt.2020.107468>, 2021.

581 Wu, F., Chu, X., Du, L., Jiao, J., Zheng, H., Xun, Y., et al.: First simultaneous lidar observations of
 582 thermosphere-ionosphere sporadic Ni and Na (TISNi and TISNa) layers (~105-120 km) over
 583 Beijing (40.42°N, 116.02°E). *Geophysical Research Letters*, 49, e2022GL100397.
 584 <https://doi.org/10.1029/2022GL100397>, 2022.

585 Xu, Y. M., and Chen, G.: Metal atoms dataset [Dataset]. Zenodo. <https://zenodo.org/records/15003984>,
 586 2025.

587 Xun, Y. et al.: Seasonal variation in the mesospheric Ca layer and Ca⁺ layer simultaneously observed
 588 over Beijing (40.41°N, 116.01°E). *Remote Sensing*, 16(3), 596. doi:10.3390/rs16030596, 2024.

589 Yi, F. et al.: Seasonal variations of the nocturnal mesospheric Na and Fe layers at 30°N. *Journal of*
 590 *Geophysical Research: Atmospheres*, 114(D1), D01301. <https://doi.org/10.1029/2008JD010344>,
 591 2009.

592 Yu, T., Wang, W., Ren, Z., Cai, X., Yue, X., & He, M.. The response of middle thermosphere (~ 160
 593 km) composition to the 20-21 November 2003 superstorm. *Journal of Geophysical Research:*
 594 *Space Physics*, 126, e2021JA029449, <https://doi.org/10.1029/2021ja029449>, 2021.

595 Yu, T., Cai, X., Ren, Z., Li, S., Pedatella, N., & He, M.. Investigation of the ΣO/N₂ depletion with

latitudinally tilted equatorward boundary observed by GOLD during the geomagnetic storm on April 20, 2020. *Journal of Geophysical Research: Space Physics*, 127, e2022JA030889. <https://doi.org/10.1029/2022JA030889>, 2022.

Yu, T., Cai, X., Ren, Z., Li, S., Pedatella, N., & He, M.. Investigation of the $\Sigma O/N_2$ depletion with latitudinally tilted equatorward boundary observed by GOLD during the geomagnetic storm on April 20, 2020. *Journal of Geophysical Research: Space Physics*, 127, e2022JA030889. <https://doi.org/10.1029/2022JA030889>, 2022. Yu, T. T. et al.: Investigation of interhemispheric asymmetry of the thermospheric composition observed by GOLD during the first strong geomagnetic storm in solar-cycle 25, 1: IMF B_y effects. *Journal of Geophysical Research: Space Physics*, 128(10), e2023JA031429. <https://doi.org/10.1029/2023JA031429>, 2023.

Yu, T. T.: TIEGCM 2019_11 storm [Dataset]. Zenodo. <https://zenodo.org/records/14232871>, 2024.

Yuan, T., Feng, W., Plane, J. M. C., and Marsh, D. R.: Photochemistry on the bottom side of the mesospheric Na layer, *Atmos. Chem. Phys.*, 19, 3769-3777, <https://doi.org/10.5194/acp-19-3769-2019>, 2019.

Yue, X. C., Friedman, J. S., Wu, X. B. and Zhou Q. H.: Structure and seasonal variations of the nocturnal mesospheric K layer at Arecibo. *Journal of Geophysical Research: Atmospheres*, 122(14), 7,260–7,275. <https://doi.org/10.1002/2017JD026541>, 2017.

Zhang, Y. et al.: O/N₂ changes during 1–4 October 2002 storms: IMAGE SI-13 and TIMED/GUVI observations. *Journal of Geophysical Research: Space Physics*, 109(A10), A10308. <https://doi.org/10.1029/2004JA010441>, 2004.