



Trends in South Pole Particle Concentrations Imply Holocene Westerly Wind Strengthening

Aaron Chesler^{1,2,*}, Dominic Winski^{1,2}, Karl Kreutz^{1,2}, Bess Koffman³, Erich Osterberg⁴, David Ferris⁴, Zayta Thundercloud⁴, Jihong Cole-Dai⁵, Mark Wells⁶, Aaron Putnam², and Katherine Anderson⁴

5 ¹Climate Change Institute, University of Maine, Orono, Maine, 04469, USA

² School of Earth and Climate Sciences, University of Maine, Orono, Maine, 04469, USA

³Department of Geology, Colby College, Waterville, Maine, 04903, USA

⁴Department of Earth Science, Dartmouth College, Hanover, New Hampshire, 03755, USA

⁵Department of Chemistry and Biochemistry, South Dakota State University, Brookings, South Dakota, 57007

10 ⁶School of Marine Sciences, University of Maine, Orono, Maine, 04469, USA

*now at: Environmental Studies and Science Program, Colorado College, Colorado Springs, 80903, USA

Correspondence to: Aaron Chesler (aaron.chesler@maine.edu)

15

20

25

30



35 **Abstract.** The Southern Hemisphere Westerly Winds (SHWW) play an important role in global climate and Antarctic ice
sheet dynamics; however, high-resolution proxy reconstructions are sparse. Dust microparticles preserved in Antarctic ice
cores provide valuable paleo-perspectives on SHWW behavior. We present South Pole Ice Core (SPC14) dust records spanning
the Holocene. Dust concentrations decrease through the Holocene by ~ 20 particles $\text{mL}^{-1} \text{kyr}^{-1}$, while the coarse particle
percentage (CPP) increases by $\sim 0.10\% \text{kyr}^{-1}$. Dust trends, CMIP6-PMIP4 model results, and SH proxy records are consistent
40 with an increase in wind speed south of the SHWW core ($> \sim 51^\circ\text{S}$) across the mid-Holocene. The decrease in dust concentration
may reflect weakening winds coupled with precipitation and/or vegetation changes over mid-latitude dust source regions,
while the CPP increase may indicate strengthening of the SHWW south of the core and activation of Antarctic dust sources.
Our findings suggest that following a stable early Holocene, the SHWW began contracting towards Antarctica $\sim 7 - 6$ ka.

1 Introduction

45 The Southern Hemisphere Westerly Winds (SHWW) represent the strongest persistent surface wind system on Earth.
Variations in the SHWW influence global carbon cycling and heat distribution, and are believed to play a critical role in Earth's
climate and ice sheet stability across wide-ranging timescales (Hillenbrand et al., 2017; Russell et al., 2006). The strength and
latitudinal position of the SHWW during the past several centuries is related to changes in the equator-to-pole temperature
gradient (i.e., Perren et al., 2020). During warm periods, SHWW migrate southward and increase in strength and during cooler
50 periods migrate northward and weaken. Increases in SHWW strength intensify the ocean-to-atmosphere transfer of CO_2 via
upwelling and impact ice sheet stability and mid-latitude vegetation/precipitation dynamics (Hillenbrand et al., 2017; Anderson
et al., 2009; Toggweiler and Russell, 2008; Moreno et al., 2010). However, while the SHWW are an important climatic feature,
Holocene SHWW dynamics are poorly constrained, with previous work disagreeing between zonally symmetric or asymmetric
change in the strength and/or intensity as well as the mean latitudinal position of the winds (Saunders et al., 2018; Perren et
55 al., 2020; Moreno et al., 2010; Hillenbrand et al., 2017; Van Der Bilt et al., 2022; Zwier et al., 2022; Varma et al., 2012).

Microparticle records of concentration and relative particle size (i.e., coarse particle percentage; CPP) from Antarctic
ice cores offer a unique perspective of SHWW strength and/or position dynamics because they have been interpreted to reflect
source-to-sink conditions and atmospheric variability, specifically distillation (Markle et al., 2018), geomorphology of the
source regions (Sugden et al., 2009), wind strength (Koffman et al., 2014) vegetation cover, precipitation (Tegen and
60 Schepanski, 2018), and atmospheric transport patterns (Albani et al., 2018; Delmonte et al., 2020; Delmonte et al., 2013;
Delmonte et al., 2002; Aarons et al., 2019; Aarons et al., 2017; Aarons et al., 2016; Koffman et al., 2023). However, there are



few complete Holocene dust records from Antarctica, limiting a full interpretation of past SHWW behavior. Here we measure and develop records of dust concentration and CPP in the South Pole Ice Core (SPC14) to better constrain Holocene SHWW dynamics. The South Pole is well-situated because of its geographic position, which facilitates hemispheric-scale
65 interpretations, the relatively high snow accumulation rate that adds precision to the time record, and the 60 years of aerosol and meteorological studies in the region (Lazzara et al., 2012; Sheridan et al., 2016; Clem et al., 2020; Casey et al., 2014).

2 Methods

We measure microparticle concentration and size distributions using a subset of data collected from a continuous flow analysis (CFA) Klotz Abakus laser dust sensor at Dartmouth College (Text S1; Breton et al., 2012; Winski et al., 2021; Osterberg et al., 2006). To facilitate a comparison with other ice core records, we use a density of 2.6 g cm^{-3} (following
70 Koffman et al., 2014) and a spherical shape to estimate flux, rather than the measured prolate particle shapes (Chesler et al., 2023). Following Chesler et al. (2023), we do not use metrics that rely on the particle shape assumption (i.e., particle flux and mass concentration), which can increase particle metric concentrations by 25 – 44%, because the South Pole microparticle shape does not cover the Holocene (50 – 16 ka; Text S1; Chesler et al., 2023). Therefore, we use particle concentration
75 measured between $1.1 - 5.1 \text{ }\mu\text{m}$ and CPP (defined as number of particles mL^{-1} $[3.2 - <5.1] / [1.1 - 5.1] \text{ }\mu\text{m diameter} \times 100$) following other East Antarctic studies (i.e., Delmonte et al., 2020).

2.1 HYSPLIT and CMIP6 models

We ran HYSPLIT NCEP trajectories for December, January, February (austral summer; DJF) and June, July, August (austral winter; JJA) from 1980 – 2022 (inclusive) hourly for 10 days over a 90-day period. We then calculate air mass origin
80 percentage using a $2 \times 2^\circ$ grid to identify where air masses originate 10 days prior to reaching the South Pole.

We also explore changes in atmospheric temperature, wind speed, and precipitation using CMIP6 – PIMP4 model outputs. We use 11 different models (Table 1) to assess model agreement between mid-Holocene runs (6,000 years ago) relative to pre-industrial (1850 – 2014 AD). We calculate model difference by subtracting the mid-Holocene from pre-industrial to assess change in each climate variable from the mid-Holocene to the pre-Industrial. We replace all positive values,
85 which signifying an increase with time, and replace all negative values with a 0 and sum all model runs. Our method allows



us to assess if the models agree to an increase/decrease in each climate variable rather than assessing the magnitude of change between each model.

Table 1: List of CMIP6-PMIP4 Models used with name, institution, resolution using the rli1p1 (r = realization, i = initialization, p = physics, respectively) ensemble, with variables for surface temperature (tas), precipitation, (pr), and surface wind speed (sfc Wind).

Model	Institute	Grid Size		Variables
		Latitude (°)	Longitude (°)	
ACCESS-ESM 1-5	Commonwealth Scientific and Industrial Research Organization, Australia	1.25	1.88	pr, tas, sfcWind
AWI-ESM-1-1-LR	Alfred Wegner Institute, Germany	1.85	1.88	pr, tas, sfcWind
FGOALS-f3-L	Chinese Academy of Sciences, China	1.00	1.25	pr, tas, sfcWind
GISS-E2-1-G	Goddard Institute for Space Studies, USA	2.00	2.50	pr, tas, sfcWind
INM-CM4-8	Institute for Numerical Mathematics, Russian Academy of Science, Russia	1.50	2.00	pr, tas, sfcWind
IPSL-CM6A-LR	L'Institut Pierre-Simon Laplace, France	1.27	2.50	pr, tas, sfcWind
MPI-ESM1-2-LR	Max Planck Institute for Meteorology, Germany	1.85	1.88	sfcWind
MRI-ESM2-0	Meteorological Research Institute, Japan	1.11	1.13	pr, tas, sfcWind
NCAR - CESM3	National Center for Atmospheric Research, USA	0.94	1.25	pr, tas, sfcWind



NUIST - NESM3	Nanjing University of Information Science and Technology, China	1.875	1.875	pr, tas
NorESM2-LM	Norwegian Climate Centre	1.875	2.5	pr, tas

3 Results

95 Fine dust concentrations (i.e., $1.1 - 5.1 \mu\text{m}$ particles mL^{-1}) make up on average $\sim 85 - 96\%$ of all particles measured (median $\pm 1\sigma$; Figure 1). The fine dust concentration and CPP abundance have opposing trends throughout the Holocene, while the timing of relative change within each metric is relatively similar (Figure 1). This inverse relationship is most apparent during the mid-Holocene ($\sim 7000 - 3000$ years BP).

3.1 Holocene dust flux

100 South Pole Holocene dust flux has a similar magnitude and trend to other Antarctic plateau ice core sites (50-year resampled), with a median flux of $0.45 \pm 0.23 \text{ mg m}^{-2} \text{ yr}^{-1}$ (2σ ; Figure S1). The SPC14 Holocene median dust flux is within 2σ of the EPICA Dome C (0.33 ± 0.23) and Vostok dust flux records (0.63 ± 0.21), while non-plateau sites (i.e., TALDICE and WAIS Divide) are significantly higher ($\sim 1.31 \text{ mg m}^{-2} \text{ yr}^{-1}$ and $\sim 1.36 \text{ mg m}^{-2} \text{ yr}^{-1}$, $[< 5.1 \mu\text{m}]$, respectively; Koffman et al., 2014; Albani et al., 2012; Lambert et al., 2012; Petit et al., 1990). NOAA HYSPLIT analysis from austral summer (DJF) and winter (JJA; 1980 – 2022) highlights an Atlantic sector signal (Figure 2; Rolph et al., 2017; Stein et al., 2015). Previous South Pole dust analyses identified a primary austral summer peak (Lazzara et al., 2012; Carrasco et al., 2003; Parungo et al., 1981; Hogan et al., 1984; Bodhaine et al., 1986). Our HYSPLIT analysis identifies 10-day air mass origin locations during the summer with an average coordinate of $70.9 \pm 9.4^\circ \text{ S}$ $-1.4 \pm 80.03^\circ \text{ W}$, indicating a primarily Weddell Sea/Atlantic Sector modern signal. Following Albani et al. (2012), we compare our Holocene dust flux record against $\delta^{18}\text{O}$ to assess a primarily

105



110 remote or mixed dust source. Albani et al. (2012) compared the TALDICE dust- $\delta^{18}\text{O}$ profile against EPICA Dome C as EPICA
Dome C is considered to represent remote dust sources. We define remote dust sources similarly, as those from the mid-
latitudes (i.e., southern South America and Australia; Delmonte et al., 2020), local sources from Antarctica (i.e., Aarons et al.,
2019), and mixed being a combination of the remote and local. The Holocene South Pole dust record and $\delta^{18}\text{O}$ have a weak
significant relationship ($r = -0.17$, $p = <0.01$), which suggests a primarily remote dust source signal. Albani et al. (2012)
115 interpreted the close relationship between dust and $\delta^{18}\text{O}$ trends at Talos Dome as evidence of a local/regional climate influence.
Conversely at Dome C during the Holocene, the dust and $\delta^{18}\text{O}$ did not vary proportionally, which is interpreted to reflect mid-
to-high latitudinal processes (Lambert et al., 2008; Albani et al., 2012). The similarity to other East Antarctic sites (i.e., Dome
C and Vostok; Figure S1), modern HYSPLIT back-trajectory modelling, and internal dust-water isotopes relationships suggests
that the South Pole represents a site typical of Antarctic plateau locations, with dust likely sourced from remote region,
120 facilitating comparisons (Figure S1 and Figure 2).






Figure 1: 50-year median resampled SPC14 dust concentration (black line), fine dust concentration (light grey shading; $1.1 - <3.2\mu\text{m}$), coarse particles (dark grey shading; $3.2 - <5.1\mu\text{m}$), and 50-year median resampled CPP (light red) and CPP 11-point rolling Gaussian mean (2σ ; red dashed line). The figure is limited to 11,300 – 250 years BP to highlight Holocene variability recorded in the South Pole annually resolved record. Inset shows histograms of dust concentration from each of the three time periods highlighted in the record. Note the bimodal distribution during the mid-Holocene highlighting the transition from early to late Holocene. Average median absolute deviation (MAD) for the particle concentration is ~ 84 (Figure S2).

3.2 Holocene trends in dust deposited at the South Pole

The SPC14 dust concentration decreases relatively linearly during the Holocene by an average of ~ 20 particles mL^{-1} kyr^{-1} (Figure 1; Table S1). To describe the SPC14 dust record, we divide the Holocene into early (11,300-7,000 years BP), middle (7,000-3,000 years BP) and late (3,000 - 250 years BP) periods. During the early Holocene, dust concentration remained relatively stable, followed by a nearly two-fold decrease to the modern era (Table S1). Dust concentration distributions in all three time periods are statistically different (Figure 1 inset; $p\text{-value} < 0.01$; Table S2a), further supporting a reduction in Holocene dust deposition.

In contrast, CPP increases over the Holocene from 8.6% to 9.4%. The CPP during the early and late Holocene are distinct, with significantly different distributions (two-sided t-test; $p\text{-value} < 0.01$; Table S2b). The increase in CPP is driven primarily by a steeper decrease in fine particles ($1.1 - <3.2 \mu\text{m}$) rather than an increase in coarse particles. Although the coarse dust concentration also decreases slightly throughout the Holocene (~ -0.09 particles $\text{mL}^{-1} \text{kyr}^{-1}$), the decrease is much less than that of the total dust concentration slope such that the proportion of coarse particles rises, especially during the middle Holocene. CPP has a weak statistical relationship with dust concentration during the Holocene (Spearman $r = -0.28$; $p < 0.01$), suggesting independent factors driving most of their variability. During the late Holocene, both the dust concentration and CPP have the lowest variability compared to the middle or early Holocene (Figure 1, Table S1).

3.3 Modern air mass sources

Our HYSPLIT results (Figure 2) identify a high percentage of 10-day air mass origins in the Weddell, Dronning Maud Land, Enderby and Kemp Land (grid north through east northeast) during DJF. During JJA, air masses at their 10-day origin are more latitudinally and longitudinally diverse, with a greater percentage of air mass origins over the Atlantic and Pacific sectors of the Southern Ocean (Figure 2c and d).



4 Discussion

4.1 Dust source(s)

The South Pole receives air masses from multiple different regions, including Western Antarctica (Nicolas and Bromwich, 2011), the Weddell Sea (Winski et al., 2021; Lazzara et al., 2012; Arimoto et al., 2008), and downslope from the East Antarctic Plateau (Lazzara et al., 2012). The dominant advection of remote air masses are from grid north (Weddell Sea) and grid southwest (WAIS) with a minor source from grid east (EAIS; Harris, 1992). Our HYSPLIT data analysis then is reflective of past studies and highlights modern seasonal latitudinal changes in wind strength and position of 10-day air mass origins. While we recognize that some air masses and dust are from the East Antarctic Plateau, this does not predominantly reflect the decrease in the fine dust particles during the mid-Holocene since dust sources are predominantly from southern South America and Australia during the Holocene and in the modern period (Wegner et al., 2012; Vecchio et al., 2024; Delmonte et al., 2020). Rather, we suggest that fluctuations in fine particle deposition are advected to the South Pole largely by strengthening of air masses in the Weddell Sea or West Antarctica, consistent with previously published data (Harris, 1992; Lazzara et al., 2012; Clem et al., 2020; Winski et al., 2021; Arimoto et al., 2008). Because the fine particles decrease significantly during the mid-Holocene compared to the relatively small decrease in the total number of coarse particles, we explore the possibility of different sources of fine vs coarse particles. Given the multiple trajectories of air masses to the South Pole, we do not specify a specific dust source contributor but rather limit our interpretation to local or remote dust sources.




Figure 2a - d: NOAA HYSPLIT analysis of 10-day hourly back-trajectory of air masses to the South Pole from 1980 – 2020 (Rolph et al., 2017; Stein et al., 2015). Height elevation of back trajectory is 1000m. 2a and b indicate the percentage of 10-day air masses (240 hour) in a 2x2° grid. 2c and d are histograms of frequency of seasonal data by latitude and longitude, respectively.

170 4.2 Proxy and modelled Holocene wind changes

We suggest that the SPC14 dust concentration and CPP records indicate a mid-to-high latitude ($>51^{\circ}\text{S}$) strengthening and southward shift of the SHWW during the mid-Holocene ($\sim 7000\text{--}3000$ y BP), consistent with other proxy-based reconstructions of SHWW migration and CMIP6-PMIP4 model output (Figure 3a). Simulations from an ensemble of CMIP6-PMIP4 models (Figure 3a), all show a clear pattern of decreased surface wind strength over South American and mid-latitude



175 (40°–50°) oceans concurrent with increasing surface wind strength over the Antarctic periphery between the mid-Holocene (6,000 BP) and historical periods (1850 – 2014 AD; Figure 3a). The weakening of the SHWW over latitudes $>\sim 51^\circ\text{S}$ is consistent with the observed decrease in the SPC14 dust concentration, indicating a reduced entrainment of fine remote dust sources in surface winds. We note broad agreement spatially with Southern Hemisphere temperature and precipitation increase between the mid-Holocene and historical periods over common dust source regions (i.e., South America, Australia, and
180 Antarctica; Figure 3b and c). A southward shift in the core of the SHWW can simultaneously explain the opposing trends in CPP and total particle concentration at the South Pole while being consistent with proxy records reflecting changes in winds, precipitation and/or glacier retreat across the mid-to-high-latitude Southern Hemisphere during the mid-Holocene.

More northern proxy records corroborate a poleward shift in the SHWW during the mid-Holocene. Lacustrine sediments from Emerald Lake, Macquarie Island ($\sim 55^\circ\text{S}$, $\sim 159^\circ\text{E}$) reflect a weakening of the SHWW between 11200 – 9200
185 years BP, followed by a period of increased variability between 9200 – 5300 years BP, and sustained and strengthened winds between 5300 – 200 years BP (Figure 4f; Saunders et al., 2018). In southern South America, Moreno et al. (2010), compared changes in precipitation via plant community changes in two lacustrine records (Lago Guanaco and Lago Condorito; 51° [Figure 4e] and 41°S , respectively, $\sim 72^\circ\text{W}$). Their results show a relatively dry period during the Early Holocene between 10500 – 7800 years BP, followed by an increase in precipitation in both records following 7800 years BP (Figure 4e). These
190 changes were interpreted to reflect a relatively weak SHWW influence during the Early Holocene followed by a strengthening of the SHWW influence during the middle and late Holocene, which is consistent with Saunders et al. (2018; Figure 4e and f).

The timing of inferred changes in the SHWW are not zonally consistent, suggesting regional variability. In the eastern Pacific Ocean in New Zealand (43°S , 171°E), progressive glacier retreat and southward shift in the ITCZ and SHWW occur between ~ 9800 and ~ 6900 years ago, suggesting a warming early Holocene (Putnam et al., 2012). Whereas in South America,
195 the changes located at Lago Condorito and Lago Guanaco suggest southward migration starting at the earliest at 7800 years ago. While southern South America might suggest a relatively linear change in the SHWW (Figure 4e) this does not negate a non-linear shift in the SHWW belt in other regions. The offset between these two records would suggest zonal variability across the Pacific, in agreement with (Kaplan et al., 2016).



CMIP6-PMIP4 Model Agreement






Figure 3a – c: CMIP6-PMIP4 models agreement displaying difference between mid-Holocene and historical surface wind speed (3a), temperature (3b), and precipitation (3c). The color bar indicates the number of models that infer a positive or no/negative change in each variable between 6000 years BP and 1850–2014 AD. Red regions indicate a stronger model consensus towards an increase and white indicates a consensus towards a decrease in the respective variable. Values in the middle represent a lack of consensus. The black dotted line represents the average position of peak surface wind speed for both time periods.

4.3 Changes in SPC14 dust concentration and hydrologic conditions

From 7000 – 3000 years BP, South Pole dust concentration decreased by ~60% (Table S1). While we have shown this decrease is qualitatively consistent with a reduction in Patagonian surface winds, possibly related to SHWW variability (<~51°S; Figure 3a), increased precipitation in the remote dust source regions (Figure 3c and Figure 4d and e; Tonello et al., 2009; Moreno et al., 2010) or during transport may also have contributed to this change. Precipitation in Patagonian source regions is closely related to changes in surface wind speed on the Andes western slope with an opposing pattern between wind speed and precipitation on the eastern slope, due to increasing cyclonic activity and the presence of the mountains causing a rain shadow effect (Berman et al., 2012; Moreno et al., 2010; Garreaud, 2007). Precipitation trends in South America vary across regions north and south during the Holocene with icefields in Patagonia (47 – 48°S, 73.4° W) and Andes experiencing arid conditions during the early (9400 – 6300 cal years BP) and late (2400 – 1600 cal years BP) Holocene, while regions in Southern Patagonia (~50°S; ~72°W) received increased precipitation during the mid-late Holocene (Glasser et al., 2004; Tonello et al., 2009; Figure 4d; Moreno et al., 2010). However, both regions show increased in precipitation during the mid-Holocene (~7000- >3000 cal years BP; Figure 4d; Glasser et al., 2004; Tonello et al., 2009; Moreno et al., 2010; Bamonte et al., 2015). At lower latitudes (~44°S), there was a relative increase in effective moisture during the Holocene (~9500 - 4200 years BP; De Porras et al., 2014). We note that the CMIP6-PIMP4 output models suggest a broad increase in precipitation over most of South America, with disagreement in the models over the southern South America (Figure 3c). Paleoproxy lake records (Figure 4) also support a progressive increase in precipitation throughout the Holocene suggesting southward migration of the SHWW belt.

Modelled and paleoproxy increases in precipitation are also observed with increases in *Nothofagus* compared to grassland vegetation cover (Figure 4e; i.e., Moreno et al., 2010). The increase in vegetation cover and precipitation can reduce soil erosion and dust entrainment, while increasing dust fallout (i.e., Mahowald et al., 2005; Tegen and Schepanski, 2018).



Therefore, the Holocene decrease in South Pole dust may not merely be related to a southward migration of the SHWW away from the dust sources, but also to a rise in precipitation resulting from this shift that also suppressed dust mobilization.

230 4.4 Antarctic ice sheet thinning and CPP increase

Previous studies have linked CPP variability with changes in air mass advection and/or activation of local dust sources (Delmonte et al., 2020; Koffman et al., 2014; Albani et al., 2012; Wegner et al., 2012; Aarons et al., 2017; Gabrielli et al., 2010). We therefore interpret the significant increase in SPC14 CPP as evidence of both the decrease in fine remote dust and transport of local Antarctic dust. The increase in SPC14 CPP begins ~8000 – 6000 yr B.P. (Figure 1 and 4a) coinciding with
235 the timing of glacial retreat and dust source exposure in lower-elevation regions of Antarctica between ~7500 – 6000 y B.P. (Figure 4h; Jones et al., 2020; Jones et al., 2015; Stutz et al., 2021; Stone et al., 2003; Hein et al., 2016; Kawamata et al., 2020). For consistency, we recalibrate ¹⁰Be surface exposure ages using production rates from Putnam et al. (2010) and (Kaplan et al., 2011; Supplementary S1). Newly exposed source areas in the Weddell Sea sector are plausible origins of dust in SPC14 given prevailing low-level wind patterns (Lazzara et al., 2012; Winski et al., 2021; Clem et al., 2020). However, because the
240 South Pole has multiple air sources, either moving into the East Antarctic plateau via in the Weddell Sea (Winski et al., 2021; Clem et al., 2020; Lazzara et al., 2012) or across West Antarctica (Nicolas and Bromwich, 2011), we interpret the changes in coarse particles to reflect broader scale Antarctic changes rather than to indicate activation of a specific local source (Figure 2).

Holocene icesheet thinning, leading to exposure of Antarctic dust sources, is primarily related to 1) ocean warming
245 (Jones et al., 2020; Stutz et al., 2021), 2) change in glacial flow patterns (Hein et al., 2016), and/or 3) reduced buttressing (Stutz et al., 2021). Strengthening and/or southward latitudinal shifts in the SHWW have been shown to increase surface upwelling of relatively warmer circumpolar deep water (CDW) and increase air mass incursions driven by cyclogenesis into the Antarctic plateau (Hillenbrand et al., 2017; Carrasco et al., 2003). It seems likely that contracting, southward shifting, and accelerating SHWW, leading to warming ocean and atmosphere, is linked to the mid-Holocene glacial retreat (i.e., Albani et al., 2012; Koffman et al., 2014) and would increase Antarctic dust sources and the increase in CPP observed in SPC14,
250 however, the coarse dust concentration has a slight decrease, which could be related to deflation of local Antarctic dust sources, similar to Holocene dynamics at Talos Dome (i.e., Albani et al., 2012). As further evidence, the timing of decreasing surface



elevation around Antarctica corresponds to increases in SPC14 CPP as a southward shift in the Atlantic ITCZ, representing warming of the Southern Hemisphere relative to the Northern Hemisphere and the Lago Guanaco which was interpreted to reflect latitudinal migrations in the SHWW (Figure 4a, c, and e; Haug et al., 2001; Moreno et al., 2010; respectively). We suggest that the CPP increase is primarily related to changes in remote dust contributions.






Figure 4 Southern Hemisphere perspective of climate variability (11300 – 250 years BP). Figure a – j are stacks of different proxies spanning from the ITCZ to the South Pole. 3a) is the SPC14 dust concentration (blue; # mL⁻¹) and CPP (light red shading) and CPP 11-point rolling mean (dashed red line; %). b) summer (December) insolation at 45°S (Laskar et al., 2004), c) Cariaco Basin Ti% (Haug et al., 2001), d) Cerro Frias southern Patagonia precipitation reconstruction (Tonello et al., 2009), e) Lago Guanaco precipitation reconstruction from Torres del Paine (Moreno et al., 2010), f) Macquarie Island (South Pacific) diatom-inferred (D-I) conductivity measurements interpreted to reflect relative wind strength over the island (~55°S; Saunders et al., 2018), g) Pine Island Bay (PIB) δ¹³C reflective of circumpolar deep water (CDW) advection, h) ¹⁰Be surface exposure from the Pine Island Glacier (Johnson et al., 2014), Ross Sea (Jones et al., 2020; Jones et al., 2015), and Dronning Maud Land (Kawamata et al., 2020). Scaling of each distribution kernel density estimate (KDE) is altered to show relative timing rather than magnitude of exposure dates, i) Siple Dome melt layer frequency record (1000-year running mean; Das and Alley, 2008), and j) compiled Antarctic CO₂ (Bereiter et al., 2015).

4.5 Southern Hemisphere Westerly Winds and Holocene climate

There is evidence indicating that the poleward shift in Holocene SHWW we infer at the South Pole is related to broader, hemisphere-scale changes in temperature and orbital changes. Mid-to-high latitude Southern Hemisphere Holocene proxy records provide ample evidence for an orbitally-driven southward shift of Holocene thermal gradients resulting in the strengthening of the SHWW. South Pole dust concentrations and CPP have significant relationships ($r = -0.76$ and 0.52 , respectively, $p < 0.01$) with the composite Antarctic CO₂ record throughout the SPC14 Holocene record (Figure 4j). Moreno et al. (2010) interpreted the close relationship between Antarctic CO₂ and their NPI record at Lago Guanaco (Figure 4e) as a metric of southward migration of the SHWW and strengthening of the wind belt. The close relationship between the SPC14 dust concentration and CPP with Antarctic CO₂ bolsters our interpretations of southward migration of the SHWW during the Holocene. SHWW model and proxy-based latitudinal reconstructions suggest that the latitudinal position and/or strength can reflect changes in regional temperature variability over millennial to centennial timescales (i.e., Jiang and Yan, 2020; Thompson and Solomon, 2002; Toggweiler et al., 2006). High-latitude zonally averaged Holocene temperature reconstructions (60 – 90°S latitude) differ, where Kaufman et al. (2020) identify a relatively stable Holocene temperature of 0.04 ± 0.22 °C (slope = < 0.01 °C kyr⁻¹; relative to pre-industrial temperature) whereas Osman et al. (2021) suggest a more insolation-like increase (slope = 0.11 °C kyr⁻¹; relative to last two millennia). The insolation-like structure in the South Pole dust record seems most consistent with the reconstruction of Osman et al. (2021). Together, the data suggest that high-southern latitude wind dynamics and resulting precipitation changes are driven by high-latitude hemispheric temperature change on orbital timescales. While there is broad agreement among many proxy records, the Pine Island Bay upwelling record shows a different



pattern, with reduced wind-driven upwelling near West Antarctica after ~8 ka (Figure 4g; Hillenbrand et al., 2017). Zonal differences in wind strength may account for these apparently conflicting observations (Kaplan et al., 2016).

290 Opportunities for future research include describing the asymmetry in the timing of SHWW wind shifts between Western and Eastern Antarctica, which could influence the output of ocean derived CO₂ to the atmosphere. Our results further suggest the early Holocene may be a poor analogue for modern climate change. The SHWW southward shift and CO₂ transfer from the ocean to the atmosphere increased throughout the Holocene and has continued to increase today (Menviel et al., 2023). Identifying and examining WAIS stability within the context of Holocene changes therefore becomes critical if the
295 early Holocene can no longer be used as an analogue to modern temperature change.

5 Conclusions

The SPC14 record over the course of the Holocene documents a decrease in dust concentration accompanied by an increase in the proportion of coarse particles (i.e., CPP). We interpret these trends as reflecting; 1) a linear southward shift in SHWW during the mid-Holocene leading to decreasing surface wind speeds and increasing precipitation affecting fine dust
300 mobilization and transport in the mid-latitudes and 2) increased Antarctic wind strength and temperature associated with a southward shift in SHWW, leading to marginal ice retreat and exposure of local dust sources. Together the timing and structure of these changes are consistent with an orbitally driven southward shift of the SHWW and global temperature belts. The interpreted shifts in the SHWW are coeval with changes in the ITCZ, suggesting that over millennial time scales, the SHWW are driven by Southern Hemisphere temperature change modulated by earth's orbital characteristics and rising CO₂.

305 Data Availability

Dust datasets used in this paper are available at the U.S. Antarctic Program Data Center (DOI: 10.15784/601553.).

Author Contributions

AC analyzed the data and prepared the manuscript with contributions from all co-authors. KK, EO, JCD, and MW designed the experiments and guided the overall research effort. DF and ZT melted the core and measured the Abakus data



310 and collected discrete samples. KA wrote the Abakus cleaning code. DW and BK provided data analysis and writing support.
AP provided writing support. MH provided laboratory analysis feedback.

Competing Interests

The authors declare that they have no conflicts of interest.

Acknowledgements

315 This research was funded by the US National Science Foundation grants [1443336 (Osterberg); 1443397 (Kreutz);
1443663 (Cole-Dai); and 1443105, 1141839 (Steig)]. We thank Mark Twickler, Joseph Souney and the South Pole Ice Core
(SPICEcore) Science Coordination Office for administering the project; the U.S. Ice Drilling Program for support activities
through NSF Cooperative Agreement 1836328; the 109th New York Air National Guard for airlift in Antarctica; the field
team who helped collect the core; the members of South Pole station who facilitated the field operations; the National
320 Science Foundation Ice Core Facility for ice core processing; and the many student researchers involved in producing the
data used in this research. We thank two anonymous reviewers, whose thoughtful suggestions and questions served to clarify
and improve the paper.

References

- 325 Aarons, S. M., Aciego, S. M., McConnell, J. R., Delmonte, B., and Baccolo, G.: Dust Transport to the Taylor Glacier,
Antarctica, During the Last Interglacial, *Geophysical Research Letters*, 46, 2261-2270, 10.1029/2018GL081887, 2019.
- Aarons, S. M., Aciego, S. M., Gabrielli, P., Delmonte, B., Koornneef, J. M., Wegner, A., and Blakowski, M. A.: The impact
of glacier retreat from the Ross Sea on local climate: Characterization of mineral dust in the Taylor Dome ice core, East
Antarctica, *Earth and Planetary Science Letters*, 444, 34-44, 10.1016/j.epsl.2016.03.035, 2016.
- 330 Aarons, S. M., Aciego, S. M., Arendt, C. A., Blakowski, M. A., Steigmeyer, A., Gabrielli, P., Sierra-Hernández, M. R.,
Beaudon, E., Delmonte, B., Baccolo, G., May, N. W., and Pratt, K. A.: Dust composition changes from Taylor Glacier (East
Antarctica) during the last glacial-interglacial transition: A multi-proxy approach, *Quaternary Science Reviews*, 162, 60-71,
10.1016/j.quascirev.2017.03.011, 2017.
- Albani, S., Balkanski, Y., Mahowald, N., Winckler, G., Maggi, V., and Delmonte, B.: Aerosol-Climate Interactions During
the Last Glacial Maximum, *Current Climate Change Reports*, 4, 99-114, 10.1007/s40641-018-0100-7, 2018.
- 335 Albani, S., Delmonte, B., Maggi, V., Baroni, C., Petit, J. R., Stenni, B., Mazzola, C., and Frezzotti, M.: Interpreting last glacial
to Holocene dust changes at Talos Dome (East Antarctica): implications for atmospheric variations from regional to
hemispheric scales, *Climate of the Past*, 8, 741-750, 10.5194/cp-8-741-2012, 2012.
- Anderson, R. F., Ali, S., Bradtmiller, L. I., Nielsen, S. H. H., Fleisher, M. Q., Anderson, B. E., and Burckle, L. H.: Wind-
Driven Upwelling in the Southern Ocean and the Deglacial Rise in Atmospheric CO₂, *Science*, 323, 1443-1448,
340 10.1126/science.1167441, 2009.
- Arimoto, R., Zeng, T., Davis, D., Wang, Y., Khaing, H., Nesbit, C., and Huey, G.: Concentrations and sources of aerosol ions
and trace elements during ANTICI-2003, *Atmospheric Environment*, 42, 2864-2876, 10.1016/j.atmosenv.2007.05.054, 2008.



- Bamonte, F. P., Mancini, M. V., Sottile, G. D., Marcos, M. A., and Gogorza, C.: Vegetation dynamics from Lago San Martín area (Southwest Patagonia, Argentina) during the last 6,500 years, *Vegetation History and Archaeobotany*, 24, 267-277, 10.1007/s00334-014-0473-z, 2015.
- Bereiter, B., Eggleston, S., Schmitt, J., Nehrbass-Ahles, C., Stocker, T. F., Fischer, H., Kipfstuhl, S., and Chappellaz, J.: Revision of the EPICA Dome C CO₂ record from 800 to 600 kyr before present, *Geophysical Research Letters*, 42, 542-549, 10.1002/2014GL061957, 2015.
- Berman, A. L., Silvestri, G., and Compagnucci, R.: Eastern Patagonia Seasonal Precipitation: Influence of Southern Hemisphere Circulation and Links with Subtropical South American Precipitation, *Journal of Climate*, 25, 6781-6795, <https://doi.org/10.1175/JCLI-D-11-00514.1>, 2012.
- Bodhaine, B. A., Deluisi, J. J., Harris, J. M., Houmère, P., and Bauman, S.: Aerosol measurements at the South Pole, *Tellus B: Chemical and Physical Meteorology*, 38, 223-235, 10.3402/tellusb.v38i3-4.15131, 1986.
- Breton, D. J., Koffman, B. G., Kurbatov, A. V., Kreutz, K. J., and Hamilton, G. S.: Quantifying Signal Dispersion in a Hybrid Ice Core Melting System, *Environmental Science & Technology*, 46, 11922-11928, 10.1021/es302041k, 2012.
- Carrasco, J. F., Bromwich, D. H., and Monaghan, A. J.: Distribution and Characteristics of Mesoscale Cyclones in the Antarctic: Ross Sea Eastward to the Weddell Sea, *Monthly Weather Review*, 131, 289-301, 10.1175/1520-0493(2003)131<0289:dacomc>2.0.co;2, 2003.
- Casey, K. A., Fudge, T. J., Neumann, T. A., Steig, E. J., and Cavitte, M. G. P.: The 1500 m South Pole ice core: recovering a 40 ka environmental record, *Annals of glaciology*, 55, 137-146, 10.3189/2014AoG68A016, 2014.
- Chesler, A., Winski, D., Kreutz, K., Koffman, B., Osterberg, E., Ferris, D., Thundercloud, Z., Mohan, J., Cole-Dai, J., Wells, M., Handley, M., Putnam, A., Anderson, K., and Harmon, N.: Non-spherical microparticle shape in Antarctica during the last glacial period affects dust volume-related metrics, *Clim. Past*, 19, 477-492, 10.5194/cp-19-477-2023, 2023.
- Clem, K. R., Fogt, R. L., Turner, J., Lintner, B. R., Marshall, G. J., Miller, J. R., and Renwick, J. A.: Record warming at the South Pole during the past three decades, *Nature Climate Change*, 10, 762-770, 10.1038/s41558-020-0815-z, 2020.
- Das, S. B. and Alley, R. B.: Rise in frequency of surface melting at Siple Dome through the Holocene: Evidence for increasing marine influence on the climate of West Antarctica, *Journal of Geophysical Research: Atmospheres*, 113, 10.1029/2007JD008790, 2008.
- de Porras, M. E., Maldonado, A., Quintana, F. A., Martel-Cea, A., Reyes, O., and Méndez, C.: Environmental and climatic changes in central Chilean Patagonia since the Late Glacial (Mallín El Embudo, 44° S), *Clim. Past*, 10, 1063-1078, 10.5194/cp-10-1063-2014, 2014.
- Delmonte, B., Petit, J., and Maggi, V.: Glacial to Holocene implications of the new 27000-year dust record from the EPICA Dome C (East Antarctica) ice core, *Climate Dynamics*, 18, 647-660, 10.1007/s00382-001-0193-9, 2002.
- Delmonte, B., Baroni, C., Andersson, P. S., Narcisi, B., Salvatore, M. C., Petit, J. R., Scarchilli, C., Frezzotti, M., Albani, S., and Maggi, V.: Modern and Holocene aeolian dust variability from Talos Dome (Northern Victoria Land) to the interior of the Antarctic ice sheet, *Quaternary Science Reviews*, 64, 76-89, 10.1016/j.quascirev.2012.11.033, 2013.
- Delmonte, B., Winton, H., Baroni, M., Baccolo, G., Hansson, M., Andersson, P., Baroni, C., Salvatore, M. C., Lanci, L., and Maggi, V.: Holocene dust in East Antarctica: Provenance and variability in time and space, *The Holocene*, 30, 546-558, 10.1177/0959683619875188, 2020.
- Gabrielli, P., Wegner, A., Petit, J. R., Delmonte, B., and De Deckker, P.: A major glacial-interglacial change in aeolian dust composition inferred from Rare Earth Elements in Antarctic ice, *Quaternary science reviews*, 29, 265-273, 10.1016/j.quascirev.2009.09.002, 2010.
- Garreaud, R.: Precipitation and Circulation Covariability in the Extratropics, *Journal of Climate*, 20, 4789-4797, <https://doi.org/10.1175/JCLI4257.1>, 2007.
- Glasser, N. F., Harrison, S., Winchester, V., and Aniya, M.: Late Pleistocene and Holocene palaeoclimate and glacier fluctuations in Patagonia, *Global and Planetary Change*, 43, 79-101, 10.1016/j.gloplacha.2004.03.002, 2004.
- Harris, J. M.: An analysis of 5-day midtropospheric flow patterns for the South Pole: 1985-1989, *Tellus B: Chemical and Physical Meteorology*, 10.3402/tellusb.v44i4.15466, 1992.
- Haug, G. H., Hughen, K. A., Sigman, D. M., Peterson, L. C., and Röhl, U.: Southward Migration of the Intertropical Convergence Zone Through the Holocene, *Science*, 293, 1304-1308, 10.1126/science.1059725, 2001.



- Hein, A. S., Marrero, S. M., Woodward, J., Dunning, S. A., Winter, K., Westoby, M. J., Freeman, S. P. H. T., Shanks, R. P., and Sugden, D. E.: Mid-Holocene pulse of thinning in the Weddell Sea sector of the West Antarctic ice sheet, *Nature Communications*, 7, 12511, 10.1038/ncomms12511, 2016.
- 395 Hillenbrand, C.-D., Smith, J. A., Hodell, D. A., Greaves, M., Poole, C. R., Kender, S., Williams, M., Andersen, T. J., Jernas, P. E., Elderfield, H., Klages, J. P., Roberts, S. J., Gohl, K., Larter, R. D., and Kuhn, G.: West Antarctic Ice Sheet retreat driven by Holocene warm water incursions, *Nature*, 547, 43–48, 10.1038/nature22995, 2017.
- Hogan, A., Kebschull, K., Townsend, R., Murphey, B., Samson, J., and Barnard, S.: Particle concentrations at the South Pole, on meteorological and climatological time scales; Is the difference important?, *Geophysical Research Letters*, 11, 850–853, 10.1029/GL011i009p00850, 1984.
- 400 Jiang, N. X. and Yan, Q.: Evolution of the meridional shift of the subtropical and subpolar westerly jet over the Southern Hemisphere during the past 21,000 years, *Quaternary Science Reviews*, 246, 10.1016/j.quascirev.2020.106544, 2020.
- Johnson, J. S., Bentley, M. J., Smith, J. A., Finkel, R. C., Rood, D. H., Gohl, K., Balco, G., Larter, R. D., and Schaefer, J. M.: Rapid Thinning of Pine Island Glacier in the Early Holocene, *Science*, 343, 999–1001, doi:10.1126/science.1247385, 2014.
- Jones, R. S., Whitmore, R. J., Mackintosh, A. N., Norton, K. P., Eaves, S. R., Stutz, J., and Christl, M.: Regional-scale abrupt
405 Mid-Holocene ice sheet thinning in the western Ross Sea, *Antarctica, Geology*, 49, 278–282, 10.1130/g48347.1, 2020.
- Jones, R. S., Mackintosh, A. N., Norton, K. P., Gollledge, N. R., Fogwill, C. J., Kubik, P. W., Christl, M., and Greenwood, S. L.: Rapid Holocene thinning of an East Antarctic outlet glacier driven by marine ice sheet instability, *Nature Communications*, 6, 8910, 10.1038/ncomms9910, 2015.
- Kaplan, M. R., Schaefer, J. M., Strelin, J. A., Denton, G. H., and Anderson, R. F.: Patagonian and southern South Atlantic
410 view of Holocene climate, *Quaternary science reviews*, 141, 112–125, 10.1016/j.quascirev.2016.03.014, 2016.
- Kaplan, M. R., Strelin, J. A., Schaefer, J. M., Denton, G. H., Finkel, R. C., Schwartz, R., Putnam, A. E., Vandergoes, M. J., Goehring, B. M., and Travis, S. G.: In-situ cosmogenic ¹⁰Be production rate at Lago Argentino, Patagonia: Implications for late-glacial climate chronology, *Earth and Planetary Science Letters*, 309, 21–32, 10.1016/j.epsl.2011.06.018, 2011.
- Kaufman, D., McKay, N., Routson, C., Erb, M., Dätwyler, C., Sommer, P. S., Heiri, O., and Davis, B.: Holocene global mean
415 surface temperature, a multi-method reconstruction approach, *Scientific Data*, 7, 201, 10.1038/s41597-020-0530-7, 2020.
- Kawamata, M., Suganuma, Y., Doi, K., Misawa, K., Hirabayashi, M., Hattori, A., and Sawagaki, T.: Abrupt Holocene ice-sheet thinning along the southern Soya Coast, Lützow-Holm Bay, East Antarctica, revealed by glacial geomorphology and surface exposure dating, *Quaternary Science Reviews*, 247, 106540, 10.1016/j.quascirev.2020.106540, 2020.
- Koffman, B. G., Kreutz, K. J., Breton, D. J., Kane, E. J., and Winski, D. A.: Centennial-scale variability of the Southern
420 Hemisphere westerly wind belt in the eastern Pacific over the past two millennia, *Climate of the past*, 10, 1125–1144, 10.5194/cp-10-1125-2014, 2014.
- Koffman, B. G., Goldstein, S. L., Winckler, G., Kaplan, M. R., Bolge, L., and Biscaye, P.: Abrupt Changes in Atmospheric Circulation During the Medieval Climate Anomaly and Little Ice Age Recorded by Sr-Nd Isotopes in the Siple Dome Ice Core, Antarctica, *Paleoceanography and Paleoclimatology*, 38, e2022PA004543, <https://doi.org/10.1029/2022PA004543>, 2023.
- 425 Lambert, F., Bigler, M., Steffensen, J. P., Hutterli, M., and Fischer, H.: Centennial mineral dust variability in high-resolution ice core data from Dome C, Antarctica, *Clim. Past*, 8, 609–623, 10.5194/cp-8-609-2012, 2012.
- Lambert, F., Delmonte, B., Petit, J. R., Bigler, M., and Kaufmann, P. R.: Dust-climate couplings over the past 800,000 years from the EPICA Dome C ice core, *Nature (London)*, 452, 616–619, 10.1038/nature06763, 2008.
- Laskar, J., Robutel, P., Joutel, F., Gastineau, M., Correia, A. C. M., and Levrard, B.: A long-term numerical solution for the insolation quantities of the Earth, *A&A*, 428, 261–285, doi.org/10.1051/0004-6361:20041335, 2004.
- 430 Lazzara, M. A., Keller, L. M., Markle, T., and Gallagher, J.: Fifty-year Amundsen–Scott South Pole station surface climatology, *Atmospheric Research*, 118, 240–259, doi.org/10.1016/j.atmosres.2012.06.027, 2012.
- Mahowald, N. M., Baker, A. R., Bergametti, G., Brooks, N., Duce, R. A., Jickells, T. D., Kubilay, N., Prospero, J. M., and Tegen, I.: Atmospheric global dust cycle and iron inputs to the ocean, *Global Biogeochemical Cycles*, 19, doi.org/10.1029/2004GB002402, 2005.
- 435 Markle, B. R., Steig, E. J., Roe, G. H., Winckler, G., and McConnell, J. R.: Concomitant variability in high-latitude aerosols, water isotopes and the hydrologic cycle, *Nature Geoscience*, 11, 853–859, 10.1038/s41561-018-0210-9, 2018.
- Menviel, L. C., Spence, P., Kiss, A. E., Chamberlain, M. A., Hayashida, H., England, M. H., and Waugh, D.: Enhanced Southern Ocean CO₂ outgassing as a result of stronger and poleward shifted southern hemispheric westerlies, *Biogeosciences*,
440 20, 4413–4431, 10.5194/bg-20-4413-2023, 2023.



- Moreno, P. I., Francois, J. P., Moy, C. M., and Villa-Martínez, R.: Covariability of the Southern Westerlies and atmospheric CO₂ during the Holocene, *Geology*, 38, 727-730, 10.1130/g30962.1, 2010.
- Nicolas, J. P. and Bromwich, D. H.: Climate of West Antarctica and Influence of Marine Air Intrusions, *Journal of Climate*, 24, 49-67, 10.1175/2010jcli3522.1, 2011.
- 445 Osman, M. B., Tierney, J. E., Zhu, J., Tardif, R., Hakim, G. J., King, J., and Poulsen, C. J.: Globally resolved surface temperatures since the Last Glacial Maximum, *Nature*, 599, 239-244, 10.1038/s41586-021-03984-4, 2021.
- Osterberg, E. C., Handley, M. J., Sneed, S. B., Mayewski, P. A., and Kreutz, K. J.: Continuous ice core melter system with discrete sampling for major ion, trace element, and stable isotope analyses, *Environmental Science & Technology*, 40, 3355-3361, 10.1021/es052536w, 2006.
- 450 Parungo, F., Bodhaine, B., and Bortniak, J.: Seasonal variation in antarctic aerosol, *Journal of Aerosol Science*, 12, 491-504, 10.1016/0021-8502(81)90052-5, 1981.
- Perren, B. B., Hodgson, D. A., Roberts, S. J., Sime, L., Van Nieuwenhuyze, W., Verleyen, E., and Vyverman, W.: Southward migration of the Southern Hemisphere westerly winds corresponds with warming climate over centennial timescales, *Communications Earth & Environment*, 1, 58, 10.1038/s43247-020-00059-6, 2020.
- 455 Petit, J. R., Mournier, L., Jouzel, J., Korotkevich, Y. S., Kotlyakov, V. I., and Lorius, C.: Palaeoclimatological and chronological implications of the Vostok core dust record, *Nature*, 343, 56-58, 1990.
- Putnam, A. E., Schaefer, J. M., Denton, G. H., Barrell, D. J. A., Finkel, R. C., Andersen, B. G., Schwartz, R., Chinn, T. J. H., and Doughty, A. M.: Regional climate control of glaciers in New Zealand and Europe during the pre-industrial Holocene, *Nature Geoscience*, 5, 627-630, 10.1038/ngeo1548, 2012.
- 460 Putnam, A. E., Schaefer, J. M., Barrell, D. J. A., Vandergoes, M., Denton, G. H., Kaplan, M. R., Finkel, R. C., Schwartz, R., Goehring, B. M., and Kelley, S. E.: In situ cosmogenic ¹⁰Be production-rate calibration from the Southern Alps, New Zealand, *Quaternary Geochronology*, 5, 392-409, 10.1016/j.quageo.2009.12.001, 2010.
- Rolph, G., Stein, A., and Stunder, B.: Real-time Environmental Applications and Display sYstem: READY, *Environmental Modelling & Software*, 95, 210-228, <https://doi.org/10.1016/j.envsoft.2017.06.025>, 2017.
- 465 Russell, J. L., Dixon, K. W., Gnanadesikan, A., Stouffer, R. J., and Toggweiler, J. R.: The Southern Hemisphere Westerlies in a Warming World: Propping Open the Door to the Deep Ocean, *Journal of Climate*, 19, 6382-6390, 10.1175/jcli3984.1, 2006.
- Saunders, K. M., Roberts, S. J., Perren, B., Butz, C., Sime, L., Davies, S., Van Nieuwenhuyze, W., Grosjean, M., and Hodgson, D. A.: Holocene dynamics of the Southern Hemisphere westerly winds and possible links to CO₂ outgassing, *Nature Geoscience*, 11, 650-655, 10.1038/s41561-018-0186-5, 2018.
- 470 Sheridan, P., Andrews, E., Schmeisser, L., Vassel, B., and Ogren, J.: Aerosol Measurements at South Pole: Climatology and Impact of Local Contamination, *Aerosol and Air Quality Research*, 16, 855-872, 10.4209/aaqr.2015.05.0358, 2016.
- Stein, A. F., Draxler, R. R., Rolph, G. D., Stunder, B. J. B., Cohen, M. D., and Ngan, F.: NOAA's HYSPLIT Atmospheric Transport and Dispersion Modeling System, *Bulletin of the American Meteorological Society*, 96, 2059-2077, <https://doi.org/10.1175/BAMS-D-14-00110.1>, 2015.
- 475 Stone, J. O., Balco, G. A., Sugden, D. E., Caffee, M. W., Sass, L. C., Cowdery, S. G., and Siddoway, C.: Holocene Deglaciation of Marie Byrd Land, West Antarctica, *Science*, 299, 99-102, 2003.
- Stutz, J., Mackintosh, A., Norton, K., Whitmore, R., Baroni, C., Jamieson, S. S. R., Jones, R. S., Balco, G., Salvatore, M. C., Casale, S., Lee, J. I., Seong, Y. B., McKay, R., Vargo, L. J., Lowry, D., Spector, P., Christl, M., Ivy Ochs, S., Di Nicola, L., Iarossi, M., Stuart, F., and Woodruff, T.: Mid-Holocene thinning of David Glacier, Antarctica: chronology and controls, *The Cryosphere*, 15, 5447-5471, 10.5194/tc-15-5447-2021, 2021.
- 480 Sugden, D. E., McCulloch, R. D., Bory, A. J. M., and Hein, A. S.: Influence of Patagonian glaciers on Antarctic dust deposition during the last glacial period, *Nature geoscience*, 2, 281-285, 10.1038/ngeo474, 2009.
- Tegen, I. and Schepanski, K.: Climate Feedback on Aerosol Emission and Atmospheric Concentrations, *Current Climate Change Reports*, 4, 1-10, 10.1007/s40641-018-0086-1, 2018.
- 485 Thompson, D. W. J. and Solomon, S.: Interpretation of Recent Southern Hemisphere Climate Change, *Science*, 296, 895-899, doi:10.1126/science.1069270, 2002.
- Toggweiler, J. R. and Russell, J.: Ocean circulation in a warming climate, *Nature*, 451, 286-288, 2008.
- Toggweiler, J. R., Russell, J. L., and Carson, S.: Midlatitude westerlies, atmospheric CO₂, and climate change during the ice ages, *Paleoceanography*, 21, n-a, 2006.



- 490 Tonello, M. S., Mancini, M. V., and Seppä, H.: Quantitative reconstruction of Holocene precipitation changes in southern Patagonia, *Quaternary Research*, 72, 410-420, [10.1016/j.yqres.2009.06.011](https://doi.org/10.1016/j.yqres.2009.06.011), 2009.
- van der Bilt, W. G. M., D'Andrea, W. J., Oppedal, L. T., Bakke, J., Bjune, A. E., and Zwier, M.: Stable Southern Hemisphere westerly winds throughout the Holocene until intensification in the last two millennia, *Communications Earth & Environment*, 3, 186, [10.1038/s43247-022-00512-8](https://doi.org/10.1038/s43247-022-00512-8), 2022.
- 495 Varma, V., Prange, M., Merkel, U., Kleinen, T., Lohmann, G., Pfeiffer, M., Renssen, H., Wagner, A., Wagner, S., and Schulz, M.: Holocene evolution of the Southern Hemisphere westerly winds in transient simulations with global climate models, *Clim. Past*, 8, 391-402, [10.5194/cp-8-391-2012](https://doi.org/10.5194/cp-8-391-2012), 2012.
- Vecchio, M. A., Costas-Rodríguez, M., Caiazzo, L., Bruschi, F., Hobin, K., Vanhaecke, F., and Grotti, M.: Provenance of mineral dust deposited on Antarctica over the last sixty years by strontium isotopic analysis of snow from Dome C, *Atmospheric Environment*, 338, 120850, <https://doi.org/10.1016/j.atmosenv.2024.120850>, 2024.
- 500 Wegner, A., Gabrielli, P., Wilhelms-Dick, D., Ruth, U., Kriews, M., De Deckker, P., Barbante, C., Cozzi, G., Delmonte, B., and Fischer, H.: Change in dust variability in the Atlantic sector of Antarctica at the end of the last deglaciation, *Clim. Past*, 8, 135-147, [10.5194/cp-8-135-2012](https://doi.org/10.5194/cp-8-135-2012), 2012.
- Winski, D. A., Osterberg, E. C., Kreutz, K. J., Ferris, D. G., Cole-Dai, J., Thundercloud, Z., Huang, J., Alexander, B., Jaeglé, L., Kennedy, J. A., Larrick, C., Kahle, E. C., Steig, E. J., and Jones, T. R.: Seasonally Resolved Holocene Sea Ice Variability Inferred From South Pole Ice Core Chemistry, *Geophysical Research Letters*, 48, e2020GL091602, [10.1029/2020GL091602](https://doi.org/10.1029/2020GL091602), 2021.
- 505 Zwier, M., van der Bilt, W. G., de Stigter, H., and Bjune, A. E.: Pollen evidence of variations in Holocene climate and Southern Hemisphere Westerly Wind strength on sub-Antarctic South Georgia, *The Holocene*, 32, 147-158, [10.1177/09596836211060495](https://doi.org/10.1177/09596836211060495), 2022.
- 510