

1 **Composition and Formation Mechanism of Brown Carbon: Identification and**
2 **Quantification of Phenolic Precursors**

3 Md. Al-amin Hossen¹, Mehedi Hasan Shakil¹, Md. Fahim Ehasan², Abu Rayhan Mohammad Tareq²,
4 Abdus Salam^{1*}

5 ¹Department of Chemistry, Faculty of Science, University of Dhaka, Dhaka-1000, Bangladesh

6 ²Environmental and Organic Chemistry Laboratory, Chemistry Division, Atomic Energy Centre Dhaka,
7 Bangladesh

8 *Corresponding author: asalam@gmail.com; asalam@du.ac.bd

Abstract

Light-absorbing organic carbon, collectively known as brown carbon (BrC), significantly influence climate and air quality, particularly in urban environments like Dhaka, Bangladesh. Despite their significance, the contributions and transformation pathways of phenolic compounds—major precursors of brown carbon (BrC)—are still insufficiently understood in the South Asian megacities. This study addresses this gap by investigating the surface morphology of PM_{2.5}, quantifying seven phenolic BrC precursors, and exploring the aqueous-phase formation pathway of nitrophenols at two urban sites (Dhaka South and Dhaka North) from July 2023 to January 2024. Phenolic compounds, including phenol, methylphenols, methoxyphenol, hydroxyphenol, and nitrophenol were identified and quantified using gas chromatography–flame ionization detection (GC-FID). PM_{2.5} surface morphology and elemental composition were analyzed via Field Emission Scanning Electron Microscopy – Energy Dispersive X-ray Spectroscopy (FESEM-EDX), and functional groups were characterized using Attenuated Total Reflectance - Fourier Transform Infrared Spectroscopy (ATR-FTIR). Results revealed that PM_{2.5} particles were predominantly spherical or chain-like with carbonaceous elements (C, O, N, S), mineral dust, and trace metals. The dominant functional groups included aromatic conjugate double bond, carbonyl, and nitro group. Aqueous-phase nitration of 2-hydroxyphenol under acidic conditions, analyzed via UV-Vis spectroscopy, demonstrated an alternative nitrophenol formation pathway. Among the detected compounds, 2-hydroxyphenol and 4-nitrophenol showed the highest average concentrations (2.31 ± 1.39 and $2.20 \pm 1.21 \mu\text{g m}^{-3}$, respectively). Seasonal variations showed elevated nitrophenol levels during winter, especially in Dhaka South ($4.54 \pm 2.94 \mu\text{g m}^{-3}$). These findings highlight the quantification of phenolic precursors and the role of aqueous-phase reactions in BrC formation, providing valuable insights for future atmospheric modeling and air quality management strategies in South Asia.

Keywords: Phenolic compounds, functional groups, surface morphology, nitrophenol

1. Introduction

Brown carbon (BrC) is a type of organic carbon that absorbs light at shorter wavelengths, such as 300-400 nm (Laskin et al., 2015). In contrast to black carbon, brown carbon exhibits a wavelength-dependent absorptivity that increases significantly towards the higher energy end of the spectrum (Wang et al., 2023). Similar to black carbon, brown carbon (BrC) also has a positive radiative effect, which lowers the total cooling effect of atmospheric aerosols caused by light scattering (Feng et al., 2013). Because of the complexity and variety of its origins, composition, and atmospheric aging mechanisms, the direct radiative effect of BrC is still unknown despite intensive research (Lee et al., 2014). The two main sources of BrC were automobile emissions and biomass burning. The composition of primary emissions can differ significantly depending on the type of fuel and the conditions under which it burns (Laskin et al., 2015c). There are probably thousands of organic molecules in atmospheric BrC, many of which are unidentified. However, phenolic compounds are a class of components that are commonly discovered (Wang et al., 2018). Phenolic and nitrated phenols are organic substances that contain at least one hydroxyl group and a nitro group in the benzene ring. Phenolic derivatives are the major component of BrC. They originate from diverse processes of production and different emission sources. Large volumes of nitrated phenolic compounds are released into the atmosphere by primary sources such as vehicle exhaust, burning of biomass, and combustion of coal (Li et al., 2020; Wang et al., 2020). The secondary synthesis of these substances from precursors like aromatic hydrocarbons and phenolic compounds via nitration in both gaseous or aqueous phases (Harrison et al., 2005b).

According to earlier research, one of the main contributors of brown carbon is nitrated phenolic compounds and R-NO₂ nitroaromatic compounds (Hossen et al., 2023; Li et al., 2022). The main reasons for the increased focus on phenolic compounds are their human toxicity (Rahman et al., 2022) and their role in producing secondary organic aerosols (SOAs) (Nakao et al., 2011). According to previous research,

75 phenol and cresols (such as m-cresol, p-cresol, and o-cresol) can oxidize to produce nitrophenol and its
76 derivatives (Harrison et al., 2005). Cresols released during the burning of biomass can undergo
77 photooxidation, producing nitro-catechol and its derivatives (Finewax et al., 2018; Iinuma et al., 2010).
78 Furthermore, studies conducted by Andreozzi et al., (2006) and Hummel et al., (2010) have demonstrated
79 that the primary pathway for the creation of 3-nitrosalicylic acid and 5-nitrosalicylic acid in the aqueous
80 phase is the nitration of salicylic acid. Laboratory investigations have established the importance of
81 aqueous processes as a production mechanism for nitrated phenolic compounds. In particular, when
82 phenol was subjected to aqueous-phase interactions with nitronium ions (NO_2^+), 4-nitrophenol was
83 produced (Heal et al., 2007). The most prevalent nitro-aromatic species were nitrophenols and nitro
84 catechols, which accounted for 31% and 32% of all nitro-aromatic compounds, respectively (Wang et al.,
85 2021). To fully understand the precursor of BrC in the atmosphere, it is crucial to analyze the
86 concentrations, chemical compositions, and sources of their phenolic precursors, along with the
87 derivatives of nitrated and methyl derivatives of phenols. Aerosol surface morphology plays an important
88 role in the light-absorbing properties and reactivity of BrC. Irregular or porous particle surfaces can
89 enhance the adsorption of gaseous precursors, such as phenolic compounds, thereby influencing their
90 heterogeneous oxidation and nitration pathways in the atmosphere. Moreover, morphological features can
91 affect how BrC and its phenolic components interact with cloud or fog droplets, altering aqueous-phase
92 reaction rates and the formation of light-absorbing secondary products.

93 In Bangladesh, most pollutants originate from vehicle emissions, solid waste burning, biomass burning,
94 and brick kiln emissions (Hossain et al., 2024; Kumar et al., 2021). All these are major sources of BrC.
95 Therefore, studies on BrC composition, such as phenolic compounds and their derivatives, in Bangladesh
96 are essential. In previous studies, Ankhy et al., (2024); Hossen et al., (2023); and Runa et al., (2022)
97 determined the optical properties of BrC emissions from biomass burning as well as ambient air in Dhaka,

Bangladesh, but did not investigate the individual composition of BrC. Previous studies have primarily emphasized gas-phase reactions between phenolic VOCs and NO_x, followed by photochemical oxidation and subsequent partitioning into the particle phase (Hems et al., 2021; Laskin et al., 2015). Yang et al., (2021) reported that photooxidation of phenolic compounds forms BrC, and phenolic compounds are a major class of aromatic compounds known as BrC precursors. Li et al., (2020) show the formation pathway of nitro aromatic compounds via gas-phase reactions, occurring through $^{\bullet}\text{OH} + \text{NO}_2$ in daytime and $^{\bullet}\text{NO}_3 + \text{NO}_2$ at nighttime. Our study reports an alternative pathway for the formation of nitroaromatic compounds in aqueous-phase reactions through electrophilic substitution (mediated by NO_2^+ or HONO) under acidic or alkaline conditions in the absence of sunlight, with demonstrated applicability to biomass samples. The main focus of this work, however, is the quantification of phenolic precursors of BrC. Specifically, we investigate the surface composition of PM_{2.5} and phenolic precursors in the atmosphere of Dhaka, Bangladesh, from July 2023 to January 2024, along with an experiment on the aqueous nitration of 2-hydroxyphenol, a common BrC precursor.

2. Methodology

2.1 Sampling sites

PM_{2.5} samples were collected from ambient air at two distinct locations in Dhaka city: Dhaka North (Shah Alibag, Mirpur-1) and Dhaka South (Department of Chemistry, University of Dhaka) (Fig. S1). Both sampling sites were situated approximately 21 meters above the ground level. The Dhaka South site, while located within an educational institution, was adjacent to a busy highway with frequent traffic congestion. Additionally, metro rail construction was ongoing nearby, further contributing to the area's pollution levels. The Dhaka North site, a mixed residential and commercial zone, was surrounded by garment factories, pharmaceutical plants, shopping centers, and restaurants. It was situated about 100 meters from the Mirpur bus station and approximately 4 kilometers from the Taltola solid waste dumping yard.

2.2 Sample Collection

PM_{2.5} samples were collected using a High-Volume Air Sampler equipped with a PM_{2.5} impactor, operating at a flow rate of 16.7 L/min, and PM_{2.5} particulates were captured on Quartz filter paper (Gelman, Membrane Filters, type TISSU Quartz 2500QAT-UP, 47 mm diameter). Before sampling, each Quartz filter was heated at 800°C for 4 hours in a furnace to remove any organic impurities. Sampling was conducted simultaneously at both locations (Dhaka South and Dhaka North sites) from July 2023 to January 2024. Each month, four samples were collected from both sites. The sampling period was 24 hours. In total, 26 samples, including blanks, were collected from each site. The PM_{2.5} mass was determined by calculating the difference between the weight of the blank filter and the PM_{2.5}-loaded filter. The total volume of air passing through the filter was calculated using the difference between the initial and final readings of the gas meter. After sampling, each sample was wrapped in aluminum foil, stored in a desiccator to regulate humidity, and finally preserved in a refrigerator at 4°C until further analysis. Flow diagrams are given below Fig.1.




Fig. 1: Schematic diagram of the overall sampling and analysis procedure from July 2023 to January 2024 at the Dhaka North site and the Dhaka South site

Fig. 1 illustrates the sampling and analysis workflow. In short, PM_{2.5} samples were collected for 24 hours from two locations in Dhaka city. Each PM_{2.5}-loaded filter was divided into four equal parts. One-quarter was extracted and analyzed for phenolic precursors of brown carbon using gas chromatography–flame ionization detection (GC-FID). Another quarter was used to identify functional groups via attenuated total reflectance–Fourier transform infrared spectroscopy (ATR-FTIR). A further quarter was examined for surface morphology and elemental composition using field emission scanning electron microscopy coupled with energy-dispersive X-ray spectroscopy (FESEM-EDX).

2.3 SEM-EDX Analysis

Morphological and chemical analyses of PM_{2.5} were performed by field emission scanning electron microscopy (FE-SEM, Zeiss Sigma 300 VP with 2 Bruker Quantax Xflash 60 mm² SDD EDS Detectors). To obtain a high-resolution SEM image, each sample was kept in a desiccator for 1 hour and then freeze-dried for 6 hours to completely remove the moisture. Next, a small section of the filter was precisely severed and arranged on a sample holder using carbon nanotape. The sample surface was subjected to gold sputtering to enhance its conductivity. Due to the presence of non-conducting particles on the surface, the electron beam will collide with the sample, resulting in the accumulation of charge and causing a charging effect that leads to fuzzy pictures, which were minimized by gold sputtering. The microscope is controlled entirely by the SmartSEM software suite developed by Zeiss. After gold sputtering under reduced pressure, the sample holder was inserted into the chamber of the SEM-EDX instrument. Air from the chamber was evacuated. When the pressure was reduced below ~ 0.009 Pa, extra high tension (EHT) was enabled. SEM image was obtained with an EHT value of 2.0 kV, and EDX was performed with an EHT value ranging from 10kV -15kV. The SDD detector was utilized for EDX analysis. For obtaining the EDX report ESPRIT Spectrum software package was used. The images obtained from SEM and EDX

were analyzed by the ImageJ software package. For the morphology study, samples collected on filters may introduce positive artifacts and uncertainties due to the fiber structure of the filters. However, SEM-EDX analysis of these filter samples has been widely employed for the morphological and elemental characterization of individual airborne particles (Park et al., 2023).

2.4 ATR-FTIR analysis

An attenuated total reflection-Fourier transform infrared (ATR-FTIR) spectrometer (Shimadzu, IRPrestige-21, Japan) was utilized to examine the chemical functional group of brown carbon according to Hossen et al., (2023). IR spectra of the samples were obtained within the range of wave number 700–4000 cm^{-1} . Since an ATR probe was utilized in ATR-FTIR, sample preparation was not needed. It is a non-destructive process. During ATR-FTIR analysis, the sample was kept on the crystal (ATR probe), which is composed of high refractive index materials such as AgCl and Si. In this process, infrared (IR) radiation incident on the crystal interacts with the sample on its surface.

2.5 Sample Extraction and Chemical Analysis

To identify and quantify of phenolic precursors of BrC, half of each sample filter paper was cut into small pieces and taken into a conical flask. Then added 30.0 mL of solvent 1:1 (dichloromethane: ethylacetate) and kept it to soak overnight for better extraction. After 24 hours, each sample was sonicated for 40 min at 30°C, followed by shaking for 40 min in an orbital shaker. During the extraction process mouth of the conical flask is completely closed by a cap. The extracted solution was filtered by a syringe filter (PTFE, 0.45 μm). The volume was reduced to approximately 1.0 mL by the rotary evaporator. The final extracted samples were kept in the refrigerator at 4°C until analysis in a gas chromatography-flame ionization detector (GC-FID). The phenolic precursors of BrC in the extracted sample solutions were identified and

184 quantified by gas chromatography coupled with flame ionization detector (GC-FID) (**model:** Varian 450-
185 GC, **column:** VF-5 ms (30m x 0.25mm, ID 0.25 μ m), **carrier gas:** N₂, Air, H₂, **detector temperature:**
186 280°C, **Split ratio:** 20, **injection volume:** 1.0 μ L, **column flow:** 1.2 mL/min, **column temperature:** Initial
187 60°C (2min hold), 180°C (8°C ram/min), 180°C (10°C ram/min,) 4 min hold, **total run time:** 27 min).

188 Standard solution of seven phenolic compounds, such as Phenol (MERCK-Schuchardt), 2-methylphenol
189 (MERCK-Schuchardt), 3-methylphenol (MERCK-Schuchardt), 2-methoxyphenol (MERCK-
190 Schuchardt), 2,4-dimethylphenol (MERCK-Schuchardt), 2-hydroxyphenol (MERCK-Schuchardt), 4-
191 nitrophenol (MERCK-Schuchardt) was prepared using the solvent 1:1 dichloromethane and ethylacetate.

192 Also prepared a mixed standard including these seven standard phenolic compounds. The concentrations
193 of the prepared standard solution were 0.5, 2.5, 5.0, 10.0, 20.0, 40.0, 60.0, 80.0, and 100.0 ppm. At first
194 run the individual standard solution to identify the retention time of target compounds. Then the mixed
195 standard solution was injected into the gas chromatography to separate the individual phenolic
196 compounds. The presence of phenolic compounds in the ambient sample was identified by the presence
197 of a peak at a characteristic retention time in the sample chromatogram compared to that of the
198 chromatogram of the standard solution. On the other hand, the quantitative analysis was conducted by
199 comparing the peak areas of the compounds to those of a standard solution, achieved through the creation
200 of a calibration curve (Fig.S2). Method validation, including the limit of detection (LOD), limit of
201 quantification (LOQ), and recovery rate, was also carried out for this method of quantification of phenolic
202 precursor. The details are described in the supplementary file in section 2. In a brief, the recovery
203 experiment was carried out in three samples by spiking a known amount of standard solution. A blank
204 sample and the two samples that gave very little signal/peak in the chromatogram were used as a sample
205 matrix for the recovery test. A known amount of standard solutions was spiked to the filter sample and
206 allowed the sample to stand for an hour to let the phenolic compound be absorbed into the filter matrix.

207 Then extract this spiked sample by the same process of sample extraction. The average recovery rate of
208 the samples was 85% which is in the acceptable range (Table S1).

209

210 **2.6. Correlation Analysis**

211 The degree and direction of a linear relationship between two variables (in this case, compounds) are
212 measured by Pearson's correlation coefficient or r . The r -value is between -1 and 1: $r=1$ (perfectly positive
213 correlation), $r=-1$ (perfectly negative correlation), $r=0$ no correlation. $r > 0.7$ means strongly correlation,
214 $0.3 > r > 0.7$ means moderately correlation, and $0.3 > r$ means weak correlation (Hauke and Kossowski,
215 2011).

216

217 **2.7 Nitration Experiment of Catechol to Formation of Light-Absorbing Product**

218 Nitrophenols are the major component of light-absorbing atmospheric organic aerosol, commonly referred
219 as brown carbon. The primary constituent of BrC is the nitroaromatic compounds, which account for
220 around 40-60% of the total BrC composition. Most of the nitrophenol formation pathways involve the
221 reactions of phenolic compounds with NO_3 and NO_2 in the gas phase in the ambient atmosphere (Wang et
222 al., 2023). Aqueous phase nitration of 1,2-dihydroxy phenol (catechol) forms a major component of brown
223 carbon, such as 4-nitrocatechol. Catechol is produced from biomass burning and vehicle emissions. This
224 catechol reacts with nitrous acid in the aqueous phase to form secondary aerosol nitrophenol, which is the
225 major light-absorbing component of organic aerosol. Multiple investigations have demonstrated the
226 significant involvement of NO_x in the creation of nitroaromatic compounds (Chow et al., 2016). However,
227 this study presents a method for the production of nitro-aromatic compounds by simple aqueous phase
228 nitration of 2-hydroxyphenol (catechol).


229 Table-1: List of required reagents for the laboratory experiments to study the formation of nitro-aromatic
 230 compounds.

S/N	Reagents	Required Concentration
1.	Catechol (1,2-dihydroxybenzene), C ₆ H ₄ (OH) ₂	1 mM
2.	Sodium Nitrite, NaNO ₂	1 mM
3.	Sulfuric Acid, H ₂ SO ₄	0.1 M
4.	Ammonia Buffer (pH~8) & 6% H ₂ O ₂	Used for alkaline conditions
5.	Hydrogen Peroxide, H ₂ O ₂	6%

231

232 For the nitration reaction, a solution of 1 mM 1,2-dihydroxybenzene and 1 mM NaNO₂ is prepared by
 233 deionized water in a 50 mL volumetric flask. The 1:1 ratio of catechol and sodium nitrite undergoes a
 234 chemical reaction both in acidic (pH=3-4, pH adjusted by 0.1 M H₂SO₄ and NaOH) and alkaline medium
 235 (used ammonia buffer to maintain pH~9, and H₂O₂, which facilitates more NO₂⁺) to form nitrated phenol.
 236 The pH of the solution was measured by a pH meter (model L-510, voltage DC 5V, S/N
 237 PK51028202310104). During the reaction continuously measured the absorbance at different times and
 238 observed the change of absorbance with time.

239 **Reaction:**




240

241


3. Results and Discussion

3.1 Morphological Study of PM_{2.5}


In this research investigated the surface elements that are most prevalent in the Earth's atmosphere. The chemical components of PM_{2.5} were categorized into distinct categories according to their morphology, as indicated by the SEM images (Fig. 2a–c). It shows that PM_{2.5} particles exhibit predominantly irregular, aggregate, and chain-like morphologies. Most individual particles are $\leq 2.5 \mu\text{m}$, while aggregates and chain-like structures range from $1.88 \mu\text{m}$ to $4.99 \mu\text{m}$ (Fig. 2c). Such coarse aggregates indicate particle growth through coagulation/agglomeration of smaller particles and the condensation of vapours onto pre-existing seed particles, rather than formation by fresh homogeneous nucleation (Lee et al., 2019). Lots of pre-existing particles in the atmosphere are released from the combustion event of biomass in our daily life, fossil fuels, emissions from vehicles, and other sources. As a result, the gaseous pollutants in the atmosphere (NO_x , SO_x , NH_3 , etc) undergo heterogeneous condensation on the surface of the pre-existing particle to form aggregates, chain-like aerosols of sulfate and nitrate. The morphology of the particles in PM_{2.5}, including diameter and shape, is similar in Dhaka North and Dhaka South sites, but the surface chemical compositions are varied with season and sampling locations. The observed surface morphologies of aerosol particles, including irregular, aggregate, and chain-like structures, play a significant role in the formation and optical properties of BrC. These morphologies provide a large reactive surface area that facilitates the heterogeneous adsorption and condensation of gaseous phenolic precursors and other organic compounds onto existing particles. Such interactions promote the chemical aging processes like nitration and oxidation, which enhance the light-absorbing capacity characteristic of BrC (Hossen et al., 2025).




(a)




(b)



(c)



(d)



(e)

Elements	Atomic No.	Weight %
C	12	29.84
O	16	44.61
Al	13	0.62
Na	11	2.96
K	19	0.63
N	7	0.02
S	16	0.61
Ca	20	0.55
Cr	24	0.01
Pb	82	0.04
As	33	0.06
Hg	80	0.09

(f)

270 **Fig. 2:** Scanning Electron Microscope images of PM_{2.5} loaded filter in (a) Dhaka North site, (b) Dhaka,
271 South site, and (c) size distribution of particles in PM_{2.5} loaded sample; (d) Energy Dispersive X-ray
272 Spectroscopy mapping, (e) Spectra of PM_{2.5}, and Table (f) relative abundance of elements in PM_{2.5}
273

274 3.2 Surface Elements Analysis

275 According to the EDX data, surface chemical compositions were categorized into carbonaceous, mineral
276 dust, and trace metal. Fig.2f shows that the quantity of carbon (C) and oxygen (O) was very high in PM_{2.5},
277 along with varying amounts of minerals (Ca, Fe, K, Al), sulfur, nitrogen, and trace metals (As, Cr, Cd,
278 Pb, Hg). The sequence of carbonaceous elements (C, O, N, and S) in Dhaka North & South sites was O >
279 C > N > S. Their possible emission sources are the combustion of fossil fuels and biomass burning. Sulfur
280 dioxide (containing S) is primarily emitted from coal combustion, while nitrogen dioxide (containing N)
281 mainly originates from agricultural land and fertilizers. The sequence of abundance for minerals is Al >
282 Ca > K > Fe. These minerals originate from mineral dust, biomass burning, and industrial waste. The trace
283 heavy metals were present in meager amounts. Among the five harmful trace metals (Hg, Cr, Pb, Cd, As),
284 lead has the highest abundance of surface elements, while the abundance of Hg was the lowest. The
285 composition suggests that particle growth was mainly due to condensation and coagulation, with mixing
286 of combustion-related and mineral dust aerosols at the sampling sites.

287

288 3.3 Characterization of Functional Group

289 The IR spectra (Fig.S3) demonstrate that the majority of the material comprises the aromatic ring system
290 because sharp peaks are found at $\sim 1630\text{-}1660\text{ cm}^{-1}$, and $\sim 1440\text{-}1460\text{ cm}^{-1}$. At this range, conjugated
291 double bonds display the IR absorption. Peak obtained at $1600\text{-}1690\text{ cm}^{-1}$, which is most probably the
292 characteristic peak of stretching of the C-O-C bond. The peak at $1500\text{-}1600\text{ cm}^{-1}$ suggested that the
293 stretching of the aromatic ring, which is a common peak in all the PM_{2.5} samples (Table 2). So, it's giving

an obvious understanding that the particulate matter contains the aromatic ring with a conjugated double bond. Few samples exhibit a peak at 2400-2500 cm^{-1} in this range $\text{-C}\equiv\text{N}$ and $\text{C}\equiv\text{C}$ undergo stretching vibration. The peak at the wave number 1400-1450 cm^{-1} is the bending of methyl groups and the methylene group. Examining the fingerprint region reveals intriguing characteristics. This investigation identified a substantial number of peaks in the fingerprint region, specifically at 1325 and 1340 cm^{-1} , which indicate the existence of organic nitrate aerosol in $\text{PM}_{2.5}$.

Table-2: Primary band assignments for ATR-FTIR spectra of $\text{PM}_{2.5}$ particulate matter

Range of Wavenumber	Vibrational Assignment
2500 – 2000 cm^{-1}	$\text{C} \equiv \text{N}$ and $\text{C} \equiv \text{C}$ bond stretching
1800 – 1650 cm^{-1}	Carbonyl ($\text{C} = \text{O}$) bond stretching
1600 – 1475 cm^{-1}	Aromatic ring stretching
1640-1550 cm^{-1}	Bending of aromatic N-H bond
1445 cm^{-1}	Bending of -CH_3 , -CH_2 and stretching of aromatic ring
1355 – 1315 cm^{-1}	$\text{R} - \text{ONO}_2$ organic nitrate stretching
1440 – 1220 cm^{-1}	Bending of C-O-H
1200-1000 cm^{-1}	Stretching vibration of $\text{C} - \text{O}$
900 – 690 cm^{-1}	Out plane bending of aromatic ring

3.4 Identification and Quantification of Phenolic Compounds (precursor of BrC)

3.4.1 Chromatogram of GC-FID Analysis

The chromatogram generated by GC-FID displayed the retention time, peak area, and identified compound. The identity of the unknown compound is ascertained through a comparison of its chromatogram with that of the standard sample. A calibration curve represents quantitative information regarding the identified compound. The chromatogram was given in the supplementary file in section 5 (Fig. S4 to S8).

3.4.2 Phenolic Precursor of BrC

The phenolic components serving as primary precursors of brown carbon include phenol, 4-nitrophenol, methoxy phenol, and cresols (m-cresol, p-cresol, and o-cresol). The precursors are initially emitted from the source as primary aerosols. Fig.3 illustrates that the mean concentration of nitrophenol and catechol in PM_{2.5} was comparatively greater than that of other phenolic compounds. The average concentrations of nitrophenol in Dhaka South (DU) during the monsoon, post-monsoon, and winter seasons were 1.52 ± 0.64 , 2.50 ± 0.43 , and $4.54 \pm 2.94 \mu\text{g m}^{-3}$, respectively. In contrast, at Mirpur, Dhaka North, the concentrations were 1.34 ± 0.09 , 1.65 ± 0.44 , and $1.62 \pm 0.52 \mu\text{g m}^{-3}$ during the same seasons. These concentrations were approximately 40-70% lower than the levels observed at the Dhaka South site. In Dhaka South, the second-highest concentration was found to be 2-hydroxyphenol (catechol) with values of 1.34 ± 0.31 , 2.44 ± 0.16 , and $5.04 \pm 3.41 \mu\text{g m}^{-3}$ during the monsoon, post-monsoon, and winter seasons, respectively, while the values recorded at Mirpur were 1.43 ± 0.07 , 1.78 ± 0.53 , and $1.82 \pm 0.51 \mu\text{g m}^{-3}$, respectively (Table S9). The levels of phenol and methyl derivatives of phenol were significantly lower compared to nitrophenol. In Mirpur, Dhaka North, the concentration of 2-hydroxyphenol is higher compared to nitrophenol. This is due to emissions from cooking, burning coal in restaurants, and biomass burning, which directly release 2-hydroxyphenol or catechol. During the winter, the increased combustion of coal and biomass for heating

purposes is responsible for the significant rises in phenolic compound concentrations (Wang et al., 2018). After the release of 2-hydroxyphenol, it undergoes a secondary chemical reaction with NO_x or aqueous nitrous acid in the atmosphere to form nitrophenol.

The average concentration of phenol in Dhaka city was 0.15 ± 0.08 , 0.17 ± 0.04 , and $0.30 \pm 0.25 \mu\text{g m}^{-3}$ in monsoon, post-monsoon, and winter seasons, respectively. The concentration of phenol in the winter season was comparatively higher than in other seasons because the major emission source of phenol is biomass burning. And most of the biomass burning occurred in the winter seasons. Li (2020) reported that the average concentration of phenol in urban Jinan, China was $(1.67 \pm 0.37) \times 10^{-2}$, $(7.1 \pm 2.6) \times 10^{-3}$, and $(2.6 \pm 0.4) \times 10^{-3} \mu\text{g m}^{-3}$ in winter, spring, and summer, respectively. Among the methyl derivatives measured in Dhaka, Bangladesh, from July 2023 to January 2024, 2-methoxyphenol exhibited the highest concentration at $0.42 \pm 0.29 \mu\text{g m}^{-3}$, while 2,4-dimethylphenol had the lowest concentration, recorded at $0.05 \pm 0.04 \mu\text{g m}^{-3}$. The high concentration of 2-methoxyphenol, which is directly emitted from biomass burning, likely indicates frequent biomass burning activities occurring near the sampling site in Dhaka, Bangladesh.




Fig. 3: Variation of concentration of phenolic compounds in (a) Dhaka, South and (b) Dhaka North site during monsoon, post-monsoon and winter

344 In contrast to findings from other research, the mean concentration of nitrated phenols observed in this
345 study is significantly greater than the concentration of phenol. Li (2020) reported that the average
346 concentration of phenol in urban Jinan, China was $(1.67 \pm 0.37) \times 10^{-2}$, $(7.1 \pm 2.6) \times 10^{-3}$, and (2.6 ± 0.4)
347 $\times 10^{-3} \mu\text{g m}^{-3}$ in winter, spring, and summer, respectively. Delhomme (2010) quantified the phenolic and
348 nitrated compounds in the ambient air and obtained the concentrations of phenol, m-cresol (3-
349 methylphenol), and o-cresol (2-methylphenol) were 1.04×10^{-2} , 2.20×10^{-3} , $1.20 \times 10^{-3} \mu\text{g m}^{-3}$,
350 respectively, in an urban site. Fig. 3 also shows that season and sampling location had an impact on the
351 fractional compositions of phenolic compounds. The most prevalent phenolic compounds in PM_{2.5} during
352 the monsoon, post-monsoon, and winter seasons were 4-nitrophenol and 2-hydroxyphenol, which
353 accounted for 37-40% and 39-41% of the quantified phenolic precursors, respectively, in Dhaka city from
354 July 2023 to January 2024. 2,4-dimethylphenol lies in the lowest portion among the identified phenolic
355 compounds, accounting for 0.8-1% of the total identified phenolic precursors. The contribution of 3-
356 methylphenol increased by 0.2% from the monsoon to the post-monsoon season and saw a further increase
357 of 0.5% during winter (Fig. 3).

358 Previous research found that catechols and 4-nitrophenol were the most abundant phenolic compounds
359 detected in Jinan and Beijing, with the highest observed concentrations (Wang et al., 2018; Wang et al.,
360 2019). The prevalence of 4-nitrophenol and 2-hydroxyphenol in both Dhaka South and Dhaka North may
361 be attributed to primary emissions, such as combustion activities, as well as the subsequent formation of
362 precursors of BrC. The high value of nitrophenol also indicated the higher concentration of NO_x in the
363 atmosphere. Liu et al. (2016) reported an enhancement in the light absorption of secondary organic
364 aerosols (SOA) when exposed to elevated NO_x levels and moderate humidity, attributing this primarily to
365 the presence of nitro-aromatic compounds as key chromophores. According to Laskin et al. (2025), the
366 optical characteristics of brown carbon (BrC) can be altered through atmospheric aging, including

processes such as photolysis and multiphase reactions that either produce or degrade light-absorbing species.

3.4.3 Correlation analysis of phenolic compounds

Table-3: Correlation analysis of phenolic compounds

	4-NPh	2-HPh	Ph	2-MPh	3-MPh	2-MOPh	2,4-DMPH
4-NPh	1						
2-HPh	0.974	1					
Ph	0.900	0.889	1				
2-MPh	0.907	0.9199	0.9879	1			
3-MPh	0.957	0.9558	0.9813	0.9898	1		
2-MOPh	0.915	0.9442	0.966	0.9701	0.9723	1	
2,4-DMPH	0.662	0.6809	0.8341	0.8659	0.8077	0.7708	1

Note: 4-NPh= 4-nitrophenol, 2-HPh= 2-hydroxyphenol, Ph= phenol, 2-MPh= 2-methylphenol, 3-MPh= 3-methylphenol, 2-MOPh= 2-methoxyphenol, 2,4-DMPH= 2,4-dimethylphenol

The correlation analysis (Table-3) shows that most of the compounds exhibit strong correlations with each other (close to 1), indicating that they behave similarly, possibly due to shared structural or chemical properties. This analysis also indicated that most of the phenolic compounds emitted from the same sources were most probably it was biomass burning and vehicle emissions. $r=0.974$ for 4-NPh and 2-HPh, and $r=0.9879$ for Ph and 2-MPh, suggesting that an extremely strong positive correlation exists between each pair, and they behave very similarly. There is a moderate to significant association between 2,4-dimethylphenol (2,4-DMPH) and 2-methoxyphenol (2-MOPh) ($r=0.7708$). Their relationship is not as strongly connected as some other pairs in Table-3. 4-NPh and 2,4-DMPH show a moderate positive

383 correlation. They may exhibit different behaviors. 2,4-DMPH shows less correlation with other phenolic
384 compounds.

385

386 **3.5 Formation of light-absorbing nitro-aromatic compounds**

387 In our study, we quantified several common brown carbon (BrC) precursors in the ambient atmosphere.
388 Among them, 2-hydroxyphenol and 4-nitrophenol were found in the highest concentrations in our
389 sampling area, both of which are well-known products of biomass burning. To gain a better understanding
390 of the possible formation processes of nitroaromatic compounds under atmospheric conditions, we
391 conducted controlled aqueous-phase nitration experiments in the laboratory.

392 In these experiments, catechol (2-hydroxyphenol), selected because it was the most abundant precursor
393 detected in our samples, was reacted with sodium nitrite under both acidic conditions (pH adjusted using
394 H₂SO₄ and NaOH) and basic conditions (pH adjusted using ammonia buffer, with H₂O₂ added to enhance
395 NO₂⁻ production) to form light absorbing nitro aromatic compound.

396

397 **3.5.1. UV-visible light absorption properties of the catechol-nitrite reaction system**

398

399 The catechol (2-hydroxyphenol) absorbs the UV-visible light at 280 nm (peak at 280 nm) (Scalzone et al.,
400 2020). When it reacted with the sodium nitrite in the presence of H₂SO₄, the peak (λ_{max}) was shifted to a
401 longer wavelength. This shift of λ_{max} indicated that there was an interaction between 2-hydroxyphenol and
402 the nitrite ions in the reaction system (Fig. 4).







Fig. 4: Absorbance of (a) single catechol, $\lambda_{\text{max}}=275$ nm, (b) NaNO₂ + H₂SO₄, and (c) reaction system (catechol-nitrite).

Fig. 5 shows the UV–Vis spectrum of the NaNO₂ and 2-hydroxyphenol reaction system in acidic medium shows a clear time-dependent evolution in absorbance over the 1–30 min period, indicating progressive formation of new light-absorbing species. At pH=3.5, the absorbance gradually increases with time in the wavelength range from 300 – 700 nm, and the λ_{max} shifts to a shorter wavelength with time increase. Initially, at 1 min the λ_{max} obtained at 355 nm then as time increased the absorbance increased gradually and the λ_{max} shifted to a shorter wavelength such as 338 nm after half an hour and no change further increase in time (Fig.5a). It is clear from this change that the nitro group has been added to the aromatic ring, leading to an increase in absorption and a shift in the position of the peak.



Time (min)	$\lambda_{\max}$ (nm)	Change of absorbance
1	365.5	0.236
3	355	0.899
6	351	1.246
9	348.5	1.598
12	344	2.216
15	340.5	2.466
20	338.5	2.628
30	338	3.094



419

Fig. 5: (a) Change of the absorbance with times of the reaction system containing 1 mM 2-hydroxyphenol and 1 mM NaNO₂ at pH=3.5. A distinct color denotes the duration of the sample run in the ultraviolet-visible spectrophotometer. The initial reactant contribution was subtracted from each spectrum. Table (b) Data for the shift of peak and change of absorbance, (c) Shift of $\lambda_{\max}$ with time, and (d) Change of absorbance with time

At the early stage (1 min), as shown in Fig. 5a, the spectrum is dominated by the characteristic $\pi \rightarrow \pi^*$ transition of the aromatic ring, centered near 270–280 nm, with relatively low intensity in the higher wavelength region. As the reaction proceeds, both the main aromatic band and a second band in the 300–350 nm region gradually increase in intensity, with the latter showing a slight bathochromic shift. This hyperchromic effect and modest red shift are consistent with the formation of nitro-substituted aromatic compounds such as o- or p-nitrophenol. The strong electron-withdrawing nature of the nitro group extends

conjugation and enhances $n \rightarrow \pi^*$ and charge-transfer transitions, leading to stronger and slightly longer wavelength absorption than the parent phenol. The continuous increase in absorbance with time, without any later decline, confirms the accumulation of these nitro-aromatic products in solution. This spectral change therefore provides strong evidence that the reaction proceeds through electrophilic aromatic substitution of 2-hydroxyphenol by nitronium ions generated from NaNO_2 in the acidic medium, yielding stable nitrophenol derivatives with enhanced UV absorption.

Reaction in alkaline medium

At high pH, nitrite remains primarily in the form of NO_2^- , and the formation of electrophiles such as HONO , NO_2^+ , or N_2O_3 is strongly suppressed. Although phenolates are highly activating toward substitution, the lack of these electrophilic species makes aromatic nitration largely ineffective. However, the presence of peroxide facilitates the formation of NO_2^+ .




Fig. 6: UV-visible absorption spectra of the nitrite-catechol reaction system in alkaline medium, pH~7-8 to form a light-absorbing nitro aromatic compound.

Fig. 6, the UV–Vis spectra of the NaNO_2 –2-hydroxyphenol system in alkaline medium with H_2O_2 show a time-dependent evolution of absorption features, indicating the formation of light-absorbing products. Initially, the spectrum is dominated by the characteristic UV absorption of phenolate anions (~ 280 – 300 nm), but with reaction time (0–15 min), the peak intensity increases and a broad shoulder emerges and intensifies in the 400–450 nm region. This indicated the completed protonated form of nitrophenol (Syed-León et al., 2020). This bathochromic shift and visible-range growth reflect the formation of conjugated nitroaromatic and quinone-like species with extended π -systems, likely via oxidation of phenolate by HO_2^- /oxidants and subsequent nitration by reactive nitrogen species from nitrite. The progressive increase in visible absorption is consistent with the generation of chromophoric, brown carbon-type products with potential atmospheric light-absorbing properties. These spectral changes strongly suggest nitro-substituted aromatic and quinone-type chromophores are forming in the system.

Under acidic conditions, nitrite generates the potent electrophile NO_2^+ (via $\text{HNO}_2/\text{HNO}_2^+$) and $\text{N}_2\text{O}_3/\text{NO}^+$ equilibria. The activated aromatic ring of phenol or catechol then undergoes rate-determining electrophilic attack by NO_2^+ to form the Wheland (σ) intermediate, directed ortho/para by the OH group(s). Subsequent deprotonation restores aromaticity, yielding the nitroaromatic product. The reaction mechanism of the nitration of aromatic compounds takes place through the reaction with an aqueous nitrite solution. Esteves et al., (2003) have also suggested that the exceedingly potent electrophile NO_2^+ facilitated the reaction at a low pH. The NO_2^+ electrophile attacks the benzene ring and forms a C–N bond. According to Wang et al., (2023) HONO acts as an electrophile in the aqueous nitration of catechol (2-hydroxybenzene). This explains the accelerated nitration rate in acidic environments, as HONO, being a stronger electrophile than NO_2^- , is more readily formed at lower pH levels.

Under alkaline conditions, nitration is generally unfavorable. At high pH, nitrite remains primarily in the form of NO_2^- , and the formation of electrophiles such as HONO, NO_2^+ , or N_2O_3 is strongly suppressed.

468 Although phenolates are highly activating toward substitution, the lack of these electrophilic species
469 makes aromatic nitration largely ineffective. However, the presence of peroxide facilitates the formation
470 of NO_2^+ .

471 In the atmosphere, a similar pathway occurs inside aqueous aerosol droplets or cloud/fog water, where
472 phenolic compounds (e.g., catechol) are taken up. These droplets adsorb NO_2 and interact with OH radicals
473 under sunlight, generating reactive nitrogen species (HONO , NO_2^+ , peroxyxynitrite) that nitrate and oxidize
474 the phenolic precursors. This process yields nitro-aromatic and quinone-type chromophores, which extend
475 absorption into the visible range (Yang, X., et al. 2023). Thus, the laboratory results directly mimic the
476 aqueous-phase reactions in atmospheric droplets that lead to the formation of light-absorbing brown
477 carbon.

478

479 **4. Conclusions**

480 This study provides comprehensive insights into the compositions, surface elements, and formation
481 mechanism of BrC in the atmosphere of Dhaka, Bangladesh, through the identification and quantification
482 of seven key phenolic precursors and an aqueous-phase nitration experiment. We observed that $\text{PM}_{2.5}$
483 particles predominantly exhibited spherical, irregular, and chain-like morphologies, with carbonaceous
484 (C, O, N, S), mineral (Ca, Fe, K, Al), and trace elements (Pb, Cr, Hg, Cd, As) identified on their surfaces.
485 ATR-FTIR analysis revealed prominent functional groups, including carbonyls, nitro groups ($-\text{NO}_2$), and
486 aromatic conjugated systems, suggesting the presence of complex organic structures. Among the phenolic
487 precursors, 4-nitrophenol ($2.20 \pm 1.21 \mu\text{g m}^{-3}$) and 2-hydroxyphenol ($2.31 \pm 1.39 \mu\text{g m}^{-3}$) showed the
488 highest concentrations, especially during winter. A strong seasonal and spatial variation was evident, with
489 higher levels in Dhaka South, indicating the influence of urban emissions and combustion activities.

490 Correlation analysis revealed strong positive relationships among most compounds ($r > 0.9$), suggesting
491 shared emission sources such as biomass and fossil fuel combustion.
492 These findings suggest that phenolic compounds are not only abundant in the urban air of Dhaka but also
493 highly reactive, forming nitrated derivatives such as 4-nitrophenol that contribute substantially to BrC.
494 The detection of nitrated and hydroxylated phenols further highlights their transformation via atmospheric
495 oxidation processes. The similarity in chemical behavior and high co-occurrence among these phenolic
496 compounds confirms their common origin and potential for secondary aerosol formation.

497 Compared to other urban regions like Jinan, China, and Strasbourg, France, the concentrations of phenolic
498 and nitrated phenolic compounds in Dhaka are substantially higher, likely due to more intense and
499 unregulated combustion activities (Li et al. 2020, Delhomme et al. 2010). Our findings align with prior
500 studies (e.g., Wang et al., 2018; Li et al., 2020) identifying catechols and nitrophenols as dominant BrC
501 components but extend the understanding by quantifying these compounds across seasons and locations
502 within a South Asian megacity.

503 While this study presents robust data on PM_{2.5} composition and BrC precursors, limitations include the
504 relatively short sampling period and limited spatial coverage. Additionally, although aqueous-phase
505 nitration experiments were conducted, the ambient contribution of such processes remains to be
506 quantified. Future studies should explore broader geographic regions, perform year-round monitoring, and
507 couple laboratory findings with atmospheric modeling for deeper mechanistic insights.
508 The results underscore the significant contribution of phenolic and nitrated phenolic compounds to
509 atmospheric BrC in Dhaka, with implications for both regional air quality and climate forcing. The
510 demonstrated aqueous-phase nitration pathway of catechol at acidic pH suggests a potentially
511 underappreciated route for BrC formation under humid and polluted conditions. This highlights the

512 importance of regulating combustion-related emissions and considering aqueous-phase chemistry in
513 climate models to accurately assess the radiative effects of organic aerosols.

514 **Acknowledgment**

515 The authors express their gratitude to the Department of Chemistry, University of Dhaka, and the
516 Environmental and Organic Chemistry Laboratory, Chemistry Division, Atomic Energy Centre Dhaka,
517 Bangladesh

518

519 **Funding:** Internal funding of the Department of Chemistry, University of Dhaka, Bangladesh

520

521 **Author's contribution**

522 MAH: Conceptualization, Sampling, Methodology, Chemical analysis, Writing-original draft, Review &
523 editing

524 MHS: Chemical Analysis

525 ARMT: Chemical Analysis, Methodology, and Supervision

526 MFE: Chemical Analysis

527 AS: Conceptualization, Review & editing, Supervision

528

529 **Conflict of interest statement:** The authors declared that they had no known competing financial
530 interests or personal relationships that could have appeared to influence the work reported in this paper.

531

532 **Date access statement:** Data will be available upon request.

533 **Ethics statement:** The research work was done according to the ethical standard of the University of
534 Dhaka, Bangladesh. There is no ethical clearance was required during this research work - because this
535 study did not include any human and animal subjects.

536

537 **5. References**

538 Andreozzi, R., Canterino, M., Caprio, V., Di Somma, I., and Sanchirico, R.: Salicylic acid nitration by
539 means of nitric acid/acetic acid system: Chemical and kinetic characterization, *Org Process Res Dev*, 10,
540 1199–1204, <https://doi.org/10.1021/op060148o>, 2006.

541 Ankhy, R. S., Roy, S., Nahar, A., Akbor, A., Al-amin Hossen, M., Jeba, F., Safiqul Islam, M.,
542 Moniruzzaman, M., and Salam, A.: Optical Characteristics of Brown Carbon in the Atmospheric
543 Particulate Matter of Dhaka, Bangladesh: Analysis of Solvent Effects and Chromophore Identification,
544 *Heliyon*, e36213, <https://doi.org/10.1016/j.heliyon.2024.e36213>, 2024.

545 Chow, K. S., Huang, X. H. H., and Yu, J. Z.: Quantification of nitroaromatic compounds in atmospheric
546 fine particulate matter in Hong Kong over 3 years: Field measurement evidence for secondary formation
547 derived from biomass burning emissions, *Environmental Chemistry*, 13, 665–673,
548 <https://doi.org/10.1071/EN15174>, 2016.

549 Delhomme, O., Morville, S., and Millet, M.: Seasonal and diurnal variations of atmospheric
550 concentrations of phenols and nitrophenols measured in the Strasbourg area, France, *Atmos Pollut Res*, 1,
551 16–22, <https://doi.org/10.5094/APR.2010.003>, 2010.

552 Esteves, P. M., Carneiro, J. W. de M., Cardoso, S. P., Barbosa, A. G. H., Laali, K. K., Rasul, G., Prakash,
553 G. K. S., and Olah, G. A.: Unified mechanistic concept of electrophilic aromatic nitration: Convergence
554 of computational results and experimental data, *J Am Chem Soc*, 125, 4836–4849,
555 <https://doi.org/10.1021/ja021307w>, 2003.

556 Feng, Y., Ramanathan, V., and Kotamarthi, V. R.: Brown carbon: A significant atmospheric absorber of
557 solar radiation, *Atmos Chem Phys*, 13, 8607–8621, <https://doi.org/10.5194/acp-13-8607-2013>, 2013.

558 Finewax, Z., De Gouw, J. A., and Ziemann, P. J.: Identification and Quantification of 4-Nitrocatechol
559 Formed from OH and NO₃ Radical-Initiated Reactions of Catechol in Air in the Presence of NO_x:
560 Implications for Secondary Organic Aerosol Formation from Biomass Burning, *Environ Sci Technol*, 52,
561 1981–1989, <https://doi.org/10.1021/acs.est.7b05864>, 2018.

562 Harrison, M. A. J., Barra, S., Borghesi, D., Vione, D., Arsene, C., and Iulian Olariu, R.: Nitrated phenols
563 in the atmosphere: A review, *Atmos Environ*, 39, 231–248,
564 <https://doi.org/10.1016/j.atmosenv.2004.09.044>, 2005.

565 Hauke, J. and Kossowski, T.: Comparison of values of pearson's and spearman's correlation coefficients
 566 on the same sets of data, *Quaestiones Geographicae*, 30, 87–93, [https://doi.org/10.2478/v10117-011-](https://doi.org/10.2478/v10117-011-0021-1)
 567 0021-1, 2011.

568 Heal, M. R., Harrison, M. A. J., and Neil Cape, J.: Aqueous-phase nitration of phenol by N₂O₅ and
 569 ClNO₂, *Atmos Environ*, 41, 3515–3520, <https://doi.org/10.1016/j.atmosenv.2007.02.003>, 2007.

570 Hems, R. F., Schnitzler, E. G., Liu-Kang, C., Cappa, C. D., and Abbatt, J. P. D.: Aging of Atmospheric
 571 Brown Carbon Aerosol, <https://doi.org/10.1021/acsearthspacechem.0c00346>, 15 April 2021

572 Hossain, M. A., Zaman, S. U., Roy, S., Islam, M. S., Zerín, I., and Salam, A.: Emission of particulate and
 573 gaseous air pollutants from municipal solid waste in Dhaka City, Bangladesh, *J Mater Cycles Waste*
 574 *Manag*, 26, 552–561, <https://doi.org/10.1007/s10163-023-01855-w>, 2024.

575 Hossen, M. A. A., Roy, S., Zaman, S. U., and Salam, A.: Emission of water soluble brown carbon from
 576 different combustion sources: Optical properties and functional group characterisation, *Environ Res*
 577 *Commun*, 5, <https://doi.org/10.1088/2515-7620/aceald>, 2023.

578 Hossen, M. A. amin, Roy, S., Nahian, S., Zaman, S. U., Selim, A., and Salam, A.: Spectral characteristics
 579 of water-soluble Brown carbon and it's radiative impacts on the atmosphere of Dhaka, Bangladesh, *Atmos*
 580 *Environ*, 351, 121185, <https://doi.org/10.1016/J.ATMOSENV.2025.121185>, 2025.

581 Hummel, M., Laus, G., Nerdinger, S., and Schottenberger, H.: Improved synthesis of 3-nitrosalicylic acid,
 582 *Synth Commun*, 40, 3353–3357, <https://doi.org/10.1080/00397910903419852>, 2010.

583 Iinuma, Y., Böge, O., and Herrmann, H.: Methyl-nitrocatechols: Atmospheric tracer compounds for
 584 biomass burning secondary organic aerosols, *Environ Sci Technol*, 44, 8453–8459,
 585 <https://doi.org/10.1021/es102938a>, 2010.

586 Kulmala, M., Dal Maso, M., Mañkela, J. M., Mañkela, M., Mañkela, M., Pirjola, L., Vañkeva, Vañ, M.,
 587 Vañkeva, V., Aalto, V. P., Miikkulainen, P., Hañmeri, K., Hañmeri, H., and O'dowd, C. D.: On the
 588 formation, growth and composition of nucleation mode particles, 53, 479–490, 2001.

589 Kumar, P., Hama, S., Nogueira, T., Abbass, R. A., Brand, V. S., Andrade, M. de F., Asfaw, A., Aziz, K.
 590 H., Cao, S. J., El-Gendy, A., Islam, S., Jeba, F., Khare, M., Mamuya, S. H., Martinez, J., Meng, M. R.,
 591 Morawska, L., Muula, A. S., Shiva Nagendra, S. M., Ngowi, A. V., Omer, K., Olaya, Y., Osano, P., and
 592 Salam, A.: In-car particulate matter exposure across ten global cities, *Science of the Total Environment*,
 593 750, <https://doi.org/10.1016/j.scitotenv.2020.141395>, 2021.

594 Laskin, A., Laskin, J., and Nizkorodov, S. A.: Chemistry of Atmospheric Brown Carbon,
 595 <https://doi.org/10.1021/cr5006167>, 27 May 2015.

596 Laskin, A., West, C. P., and Hettiyadura, A. P. S.: Molecular insights into the composition, sources, and
 597 aging of atmospheric brown carbon, <https://doi.org/10.1039/d3cs00609c>, 2 January 2025.

598 Lee, H. J., Aiona, P. K., Laskin, A., Laskin, J., and Nizkorodov, S. A.: Effect of solar radiation on the
 599 optical properties and molecular composition of laboratory proxies of atmospheric brown carbon, *Environ*
 600 *Sci Technol*, 48, 10217–10226, <https://doi.org/10.1021/es502515r>, 2014.

601 Lee, S. H., Gordon, H., Yu, H., Lehtipalo, K., Haley, R., Li, Y., and Zhang, R.: New Particle Formation
 602 in the Atmosphere: From Molecular Clusters to Global Climate, <https://doi.org/10.1029/2018JD029356>,
 603 16 July 2019.

604 Li, M., Wang, X., Lu, C., Li, R., Zhang, J., Dong, S., Yang, L., Xue, L., Chen, J., and Wang, W.: Nitrated
 605 phenols and the phenolic precursors in the atmosphere in urban Jinan, China, *Science of the Total*
 606 *Environment*, 714, <https://doi.org/10.1016/j.scitotenv.2020.136760>, 2020.

607 Li, M., Wang, X., Zhao, Y., Du, P., Li, H., Li, J., Shen, H., Liu, Z., Jiang, Y., Chen, J., Bi, Y., Zhao, Y.,
 608 Xue, L., Wang, Y., Chen, J., and Wang, W.: Atmospheric Nitrated Phenolic Compounds in Particle,
 609 Gaseous, and Aqueous Phases During Cloud Events at a Mountain Site in North China: Distribution
 610 Characteristics and Aqueous-Phase Formation, *Journal of Geophysical Research: Atmospheres*, 127,
 611 <https://doi.org/10.1029/2022JD037130>, 2022.

612 Nakao, S., Clark, C., Tang, P., Sato, K., and Cocker, D.: Secondary organic aerosol formation from
 613 phenolic compounds in the absence of NO_x, *Atmos Chem Phys*, 11, 10649–10660,
 614 <https://doi.org/10.5194/acp-11-10649-2011>, 2011.

615 Park, J. I., Kim, M. S., Yeo, M., Choi, M., Lee, J. Y., Natsagdorj, A., Kim, C., Song, M., and Jang, K. S.:
 616 Chemical and morphological characterization by SEM–EDS of PM_{2.5} collected during winter in
 617 Ulaanbaatar, Mongolia, *Atmos Environ*, 303, <https://doi.org/10.1016/j.atmosenv.2023.119752>, 2023.

618 Rahman, M. M., Rahaman, M. S., Islam, M. R., Rahman, F., Mithi, F. M., Alqahtani, T., Almikhlafi, M.
 619 A., Alghamdi, S. Q., Alruwaili, A. S., Hossain, M. S., Ahmed, M., Das, R., Emran, T. Bin, and Uddin, M.
 620 S.: Role of phenolic compounds in human disease: Current knowledge and future prospects,
 621 <https://doi.org/10.3390/molecules27010233>, 1 January 2022.

622 Runa, F., Islam, M. S., Jeba, F., and Salam, A.: Light absorption properties of brown carbon from biomass
 623 burning emissions, *Environmental Science and Pollution Research*, 29, 21012–21022,
 624 <https://doi.org/10.1007/s11356-021-17220-z>, 2022.

625 Scalzone, A., Bonifacio, M. A., Cometa, S., Cucinotta, F., De Giglio, E., Ferreira, A. M., and Gentile, P.:
 626 pH-Triggered Adhesiveness and Cohesiveness of Chondroitin Sulfate-Catechol Biopolymer for
 627 Biomedical Applications, *Front Bioeng Biotechnol*, 8, <https://doi.org/10.3389/fbioe.2020.00712>, 2020.

628 Wang, H., Gao, Y., Wang, S., Wu, X., Liu, Y., Li, X., Huang, D., Lou, S., Wu, Z., Guo, S., Jing, S., Li,
 629 Y., Huang, C., Tyndall, G. S., Orlando, J. J., and Zhang, X.: Atmospheric Processing of Nitrophenols and
 630 Nitrocresols From Biomass Burning Emissions, *Journal of Geophysical Research: Atmospheres*, 125,
 631 <https://doi.org/10.1029/2020JD033401>, 2020.

632 Wang, J., Nie, W., Cheng, Y., Shen, Y., Chi, X., Wang, J., Huang, X., Xie, Y., Sun, P., Xu, Z., Qi, X., Su,
 633 H., and Ding, A.: Light absorption of brown carbon in eastern China based on 3-year multi-wavelength
 634 aerosol optical property observations and an improved absorption Ångström exponent segregation
 635 method, *Atmos Chem Phys*, 18, 9061–9074, <https://doi.org/10.5194/acp-18-9061-2018>, 2018a.

636 Wang, L., Wang, X., Gu, R., Wang, H., Yao, L., Wen, L., Zhu, F., Wang, W., Xue, L., Yang, L., Lu, K.,
 637 Chen, J., Wang, T., Zhang, Y., and Wang, W.: Observations of fine particulate nitrated phenols in four

638 sites in northern China: Concentrations, source apportionment, and secondary formation, *Atmos Chem*
639 *Phys*, 18, 4349–4359, <https://doi.org/10.5194/acp-18-4349-2018>, 2018b.

640 Wang, Y., Hu, M., Wang, Y., Zheng, J., Shang, D., Yang, Y., Liu, Y., Li, X., Tang, R., Zhu, W., Du, Z.,
641 Wu, Y., Guo, S., Wu, Z., Lou, S., Hallquist, M., and Yu, J. Z.: The formation of nitro-aromatic compounds
642 under high NO_x and anthropogenic VOC conditions in urban Beijing, China, *Atmos Chem Phys*, 19,
643 7649–7665, <https://doi.org/10.5194/acp-19-7649-2019>, 2019.

644 Wang, Y., Jorga, S., and Abbatt, J.: Nitration of Phenols by Reaction with Aqueous Nitrite: A Pathway
645 for the Formation of Atmospheric Brown Carbon, *ACS Earth Space Chem*, 7, 632–641,
646 <https://doi.org/10.1021/acsearthspacechem.2c00396>, 2023.

647 Wang, Z., Zhang, J., Zhang, L., Liang, Y., and Shi, Q.: Characterization of nitroaromatic compounds in
648 atmospheric particulate matter from Beijing, *Atmos Environ*, 246,
649 <https://doi.org/10.1016/j.atmosenv.2020.118046>, 2021.

650 Yang, J., Au, W. C., Law, H., Lam, C. H., and Nah, T.: Formation and evolution of brown carbon during
651 aqueous-phase nitrate-mediated photooxidation of guaiacol and 5-nitroguaiacol, *Atmos Environ*, 254,
652 <https://doi.org/10.1016/j.atmosenv.2021.118401>, 2021.

653