

Deformation and exhumation in thick continental crusts induced by valley incision of elevated plateaux

Thomas Geffroy¹, Philippe Yamato^{1,2}, Philippe Steer^{1,2}, Benjamin Guillaume¹, Thibault Duretz³

¹Univ.Rennes, CNRS, Géosciences Rennes UMR 6118, 35000 Rennes, France

²Institut universitaire de France (IUF)

³Institut für Geowissenschaften, Goethe-Universität Frankfurt, Frankfurt, Germany

Correspondence to: Philippe Yamato (philippe.yamato@univ-rennes.fr)

Short summary

While erosion's role in mountain building is well known, deformation from valley incision in inactive regions is less understood. Using our numerical models, we show that incision alone can cause significant crustal deformation and drive lower crust exhumation. This is favored in areas with thick crust, weak lower crust, and high plateaux. Our results show surface processes can reshape Earth's surface over time.

Abstract

Surface processes such as erosion and sedimentation play a critical role in crustal deformation, particularly in actively deforming orogenic belts. While these processes have been extensively studied in large-scale erosive and tectonically active regions, the specific effects of valley incision on crustal deformation, especially in tectonically inactive regions, remain poorly understood. In this study, we hypothesize that crustal deformation induced by valley incision is primarily governed by three parameters: incision velocity, crustal thickness, and the elevation difference between the plateau and the valley base level. Using two-dimensional thermo-mechanical models, we investigate the influence of valley incision on crustal deformation and exhumation by varying these parameters. Our results show that valley incision alone can induce significant crustal deformation, associated with lateral viscous flow in the lower crust leading to near-vertical channel flow and extensional brittle deformation in the upper crust below the valley. This deformation leads to lower crust exhumation, within a 10 Myr timeframe, if crustal thickness is greater than 50 km, the initial plateau elevation is greater or equal to 2 km, and the long-term effective erosion rate exceeds 0.5 mm.yr⁻¹. Furthermore, while the onset of lower crust exhumation is primarily controlled by the initial plateau elevation, the total amount of exhumed lower crust after 10 Myr strongly increases with the initial thickness of the lower crust which favors viscous flow. *Moreover, natural systems that exhibit the required crustal thickness, plateau elevation, and erosion rates for lower crustal exhumation, as highlighted in our models, also demonstrate active lower crustal exhumation, as it is the case in regions such as Nanga Parbat and Namche Barwa.* These findings offer new insights into the coupling between surface processes and deep crustal dynamics, highlighting the potential for valley incision to drive substantial crustal deformation and promote lower crustal exhumation.

1 Introduction

Surface processes represent a pivotal element in the evolution of mountain ranges (e.g., Beaumont et al., 1992; Koons, 1990). While tectonic forces increase topographic relief and slope, numerous modelling studies have shown that surface processes and the associated mass redistribution exert a significant influence on tectonic processes (Avouac and Burov, 1996; Braun and Yamato, 2010; Steer et al., 2014; Sternai et al., 2019; Thieulot et al., 2014; Vernant et al., 2013; Willett, 1999; Wolf et al., 2022). Previous studies have demonstrated the critical role of erosion in controlling deformation within active orogens (e.g. Thieulot et al., 2014; Wolf et al., 2022), triggering complex feedback mechanisms, such as isostatic rebound (e.g., Sternai et al., 2019), or the setting up of large lower crustal flow (Beaumont et al., 2001), involving mechanical, thermal and time-dependent processes. These mechanisms enhance strain localization through intensified erosion, which in turn promotes increased rock uplift and accentuates orogen asymmetry (e.g. Braun and Yamato, 2010; Thieulot et al., 2014; Wolf et al., 2022). However, these studies specifically focused on active orogens which limits a comprehensive understanding of the isolated influence of erosion on crustal deformation, particularly in regions without active tectonics. Moreover, these studies do not specifically focus on the impact of valley incision on crustal deformation.

Nevertheless some studies demonstrate that valley incision can significantly influence topography through the response of lithosphere deformation to erosion (Montgomery & Stolar, 2006; Simpson, 2004; Zeitler et al., 2001). Baiadori et al. (2024) show that, even in the absence of regional deformation, the incision of fluvial valleys can lead to lithospheric bending and significant change in landscape dynamics. However, these studies have not explored the role of valley incision on large-amplitude deformation, accounting for visco-elasto-plastic rheologies, over timescales relevant for orogen building. Recently, Yang et al. (2023) proposed that the erosion of wide valleys (50 km), could drive significant crustal deformation, leading to the development of substantial relief (1–2 km) and, in extreme cases, the exhumation of lower crustal material. While this study presents important findings, the inclusion of a horizontal shortening rate of approximately 1.6 mm.yr^{-1} limits our ability to isolate the effects of valley incision on crustal deformation.

If rivers influence crustal deformation, different river systems should exert distinct impacts depending on various parameters, such as incision velocity, crustal thickness, or base level. Specifically, when considering a plateau in a non-convergent setting, we expect the difference of altitude between the plateau and the base level (what we refer to as “erosion potential” here in after) to be a main controlling factor of the induced tectonic response. Indeed, we hypothesize that a river with a low erosion potential would have a limited impact on crustal deformation, whereas a river on a high plateau, implying a high erosion potential, could induce larger crustal deformation (Figure 1). We expect the deformation to be accommodated by a flow of the viscous lower crust, leading to the formation of high relief and, in extreme cases, to the exhumation of lower crustal material at the surface (Figure 1).

The potential for valley incision to drive the exhumation of lower crustal material has been previously debated in specific natural settings, such as the Nanga Parbat–Haramosh Massif and the Namche Barwa Massif (Zeitler et al., 2001). However, the tectonic complexity of these regions makes the distinction between erosionally and tectonically driven exhumation difficult.

In this study, we therefore focus on a scenario in which regional tectonic forces are absent, ensuring that all observed crustal deformation results solely from valley incision. *This does not imply that our models are limited to inactive tectonic settings (thick continental crust indeed generally requires important shortening). However, as our objective is to isolate the specific impact of valley incision on crustal deformation, we voluntarily imposed a fixed horizontal velocity boundary conditions set to nearly 0 (i.e., 0.015 mm.yr^{-1}). Applying convergence would modify our model behavior and would make difficult to quantify what is due solely to the effect of the river incision..*

To systematically investigate the impact of valley incision on crustal deformation, exhumation, and their feedback on valley morphology, we employ a two-dimensional (2D) thermo-mechanical numerical model incorporating a visco-elasto-plastic rheology and surface processes *consisting in a simple erosion law coupling a source term for river incision with diffusion (see Section 2.1)*. Our objective is to gain a deeper understanding of the mechanical response of the lithosphere to valley incision and evaluate the potential for this process to drive the exhumation of lower crustal material, as well as whether such exhumation can occur within a geological timeframe of less than 10 Myr. *This value of 10 Myr was chosen as representative for the upper limit of river persistence at a fixed location over geological timescale. Examples of rivers maintaining the same course for more than 10 Myr are indeed extremely rare in nature (see section 4.3). Moreover, natural examples of valley incision associated with lower crust exhumation indicate that such exhumation typically occurs within less than 10 Myr (e.g., Nanga Parbat and Namche Barwa; Zeitler et al., 2001). Therefore, we assume that if this process does not take place within a 10 Myr timeframe in our model, it is unlikely to be geologically realistic.*




Figure 1: Schematic representation of the impact of a valley incision on crustal deformation, depending on the erosion potential (see text for details).

2 Materials and Methods

2.1 Code description

Numerical simulations are performed using the 2D thermo-mechanical code MDoodz (Duretz et al., 2021; Yamato et al., 2015). This numerical code was applied to the study of grain- (e.g., Luisier et al., 2023; Yamato et al., 2019), to lithospheric-scale (e.g., Auzemery et al., 2022; Candiotti et al., 2022; Poh et al., 2020; Porkoláb et al., 2021)

geological processes. It accounts for visco-elasto-plastic rheologies (Duretz et al., 2020) and a true free surface (Duretz et al., 2016). Based on the marker-in-cell method (Gerya & Yuen, 2003; Harlow & Welch, 1965), MDoodz employs the finite difference method, to solve the governing equations of momentum (1), mass conservation (2), and heat (3) on a staggered-grid. These equations are expressed as follows:

$$\frac{\partial \tau_{ij}}{\partial x_j} - \frac{\partial P}{\partial x_i} = -\rho^{\text{eff}} g_i, \quad (1)$$

$$\frac{\partial v_i}{\partial x_i} = 0, \quad (2)$$

$$\rho^{\text{eff}} C_p^{\text{eff}} \frac{DT}{Dt} = \frac{\partial}{\partial x_i} \left(k \frac{\partial T}{\partial x_i} \right) + Q_r + Q_{\text{SH}}, \quad (3)$$

where τ_{ij} is the deviatoric stress tensor, x_i and x_j the spatial coordinates, P is the pressure (Pa), ρ^{eff} is the effective density (kg.m^{-3}), g_i is the gravity acceleration vector *in the direction i (here $g_x = 0$, and $g_y = 9.81 \text{ m}^2 \cdot \text{s}^{-1}$, v_i the velocity component in the direction i (m.s^{-1}), C_p^{eff} is the effective heat capacity ($\text{J.kg}^{-1} \cdot \text{K}^{-1}$), t is time (s), T is the temperature (K), k is the thermal conductivity ($\text{W.m}^{-1} \cdot \text{K}^{-1}$), Q_r is the radiogenic heat production (W.m^{-3}), ~~Q_L is the latent heat of production or consumption, and Q_A is the adiabatic heat.~~ Q_{SH} describes the production of heat by visco-plastic dissipation (shear heating). In Eq. (1 and 3) the rock's effective density (ρ^{eff}) and heat capacity (C_p^{eff}) can indeed change when partial melting of rocks is involved (see Appendix B).*

Material properties are stored on particles that are advected through time using a fourth-order Runge-Kutta scheme. The evolution of the density field is described by the following equation:

$$\rho = \rho_0 (1 - \alpha(T - T_0))(1 + \beta(P - P_0)), \quad (4)$$

where ρ_0 is the reference density, α is the thermal expansivity, β is compressibility, T_0 and P_0 are the reference temperature and pressure, set to 0°C and 10^5 Pa, respectively. The initial temperature field is obtained by solving the equilibrium heat equation using the reference thermal parameters (Table 1), except for the lithospheric mantle, for which the conductivity has been artificially set to a very high value to obtain an adiabatic asthenosphere. The Courant condition is set at 0.25 for all the simulations.

To prevent numerical instabilities, we impose a minimum viscosity threshold of 10^{18} Pa.s in the model and use a free surface stabilization algorithm (Duretz et al., 2011). This threshold is set to avoid viscosity variations exceeding seven orders of magnitude within the model. It should be noted that 10^{18} Pa.s is still higher than the viscosity of magmas ($\sim 10^2 - 10^{14}$ Pa.s; (e.g., Dingwell, 2006)) or partially molten rocks ($\sim 10^{17} - 10^{18}$ Pa.s; (e.g., Molitor et al., 2024)).

To account for a visco-elasto-plastic rheology, the deviatoric strain rate tensor is additively decomposed such as:

$$\dot{\epsilon}_{ij} = \dot{\epsilon}_{ij}^v + \dot{\epsilon}_{ij}^e + \dot{\epsilon}_{ij}^p, \quad (5)$$

where the exponents v, e, and p correspond to the viscous, elastic, and plastic (frictional) components. We detail in Appendix A each component of the strain rate. Partial melting is also included in the numerical code (see Appendix B).

At a specific location, the vertical erosion of a plateau can occur through river or glacier incision, modulated by climatic conditions, upstream conditions (e.g., water and sediment fluxes) and local slope. In addition, the lateral erosion of the valley wall can occur through multiple processes, including river lateral mobility and hillslope destabilization, mass wasting processes, peri-glacial processes, soil production and creep (e.g., Brocard and van der Beek, 2006; Hancock and Anderson, 2002; Martin et al., 2011). Accounting for all these different and interrelated processes in a single numerical model represents a challenge, clearly out of the scope of this manuscript. Instead, we rely on the use of a simple erosion law that combines a constant incision velocity imposed at the top of the lithosphere, to simulate vertical incision by glaciers or rivers, with a topographic diffusion coefficient to simulate lateral valley erosion and sedimentation. In our models, the incision rate (V_i) remains constant as long as the valley floor remains above the river base level, set at 0 km elevation. Once this base level is reached, V_i is set to 0 and remains so unless the elevation increases above this base level. In the following, we impose a prescribed valley incision velocity V_i at the model center over an initial width W_i . In addition, diffusion is applied to surface, considering a constant diffusivity coefficient (K). The equation controlling the time-evolution of elevation h of the free surface is therefore:

$$\frac{dh}{dt} = \frac{\partial}{\partial x} \left(K \frac{\partial h}{\partial x} \right) + V_i. \quad (6)$$

2.2 Reference model design

To define the reference model set-up, we first compiled global data on crustal thickness (Crust 1.0 model; Laske et al., 2013) and Earth continental surface elevation (ETOPO1; NOAA, 2009). Based on these datasets, we constructed a diagram illustrating the correlation between both parameters (Figure 2a). The reference model was then selected to represent an intermediate position in the observed distribution.

The reference model includes a crust with a thickness (H) of 55 km and an initial plateau elevation (h_P) of 3 km (Figure 2a). The model domain used in this study is 500 km wide and 160 km deep. The reference model therefore includes a continental lithosphere made of a 55 km thick continental crust resting on top of a 65 km thick lithospheric mantle and 30 km of asthenospheric mantle (Figure 2b). We perform simulations at 1 km resolution for the full parametric study, as resolution tests show good convergence for resolutions lower than 1 km (see Figure S1).

The incision width W_i in our study has been set at a constant value of 1 km. In the reference model, we apply an incision velocity (V_i) of 30 mm.yr⁻¹ and a topographic diffusion coefficient (K) of 3.0×10^{-6} m².s⁻¹. The value of K lie in the range of previous numerical modeling studies involving surface processes (e.g., Avouac & Burov, 1996; Munch et al., 2022; Yamato et al., 2008). To avoid material loss, we compensate the volume loss due to the incision of the free surface by the addition of asthenospheric mantle material evenly distributed across the model base. The bottom, right, and left boundaries are free-slip boundaries.

The temperature at the top and lithosphere-asthenosphere boundaries are maintained at 0 and 1330°C, respectively. The asthenosphere is assumed to be adiabatic. Heat flux is set to zero at the right and left boundaries of the model (Neumann boundary conditions). The initial brittle/ductile transition occurs at 6 km of depth for the reference model (Figure 2b). *We define the boundary between the upper and lower crust based on the brittle/ductile transition of the Westerly Granite rheology. In our models, the lower crust corresponds to the portion of the crust*

186 *located beneath this transition zone (represented in green in Figure 2b) and is considered as the ductile portion*
187 *of the crust.*

188 The parametrization of the reference model leads to conditions favorable to crustal convection, with a Rayleigh
189 number of $\sim 1 \times 10^5$.

190 All material parameters are presented in Table 1 for the reference model.

191




Figure 2: a) Continental crustal thickness vs. elevation from natural data. Data for crustal thickness are from the Crust 1.0 model (Laske et al., 2013). Data for global earth altitude are from ETOPO1 (NOAA, 2009). Black points represent the combination of plateau elevation and crustal thickness used in the parametric study, and black star represent the reference model. *The black line represents the linear relationship between crustal thickness and elevation assuming that this relationship is only driven by isostasy and using the material properties of our models (see Table 1).* b) Initial model setup for the reference model. The horizontal limit between the red and green layers in the continental crust corresponds to the depth of the initial peak of strength in the crust that we define as the limit between the upper and the lower crust

(considering a homogeneous strain rate of 10^{-18} s^{-1}). White lines represent each 200°C isotherm. Black arrows indicate the input of material to balance the amount of material removed by the valley incision (see text for details). Initial topography (Plateau elevation) is here set to 3 km and the valley base level is at 0 km. *c) Yield strength profile from the surface to 70-km depth, considering a homogeneous strain rate of 10^{-18} s^{-1} . d) Temperature profile from the surface to 150-km depth.*

	Crust	Lithospheric and astenospheric mantle		
Rheological and materials parameters				
Material	Westerly Granite	Dry Olivine dislocation creep	Dry Olivine diffusion creep	Olivine Peierls creep
n	3.3	3.5	1	-
$A \text{ (Pa}^{-n} \cdot \text{s}^{-1})$	3.1623×10^{-26}	1.1×10^{-16}	1.5×10^{-15}	-
$Q \text{ (J.mol}^{-1})$	1.865×10^5	5.3×10^5	3.75×10^5	-
m	-	-	3	-
$d \text{ (m)}$	-	-	2×10^{-3}	-
$G \text{ (Pa)}$	3.0×10^{10}	3.0×10^{10}	3.0×10^{10}	-
$C \text{ (MPa)}$	50	50	50	-
$\phi \text{ (}^\circ\text{)}$	30	30	30	-
Crustal Thickness (km)	60	-	-	-
$k \text{ (W.m}^{-1} \cdot \text{K}^{-1})$	2	3.2	3.2	-
$\rho_0 \text{ (kg.m}^{-3})$	2800	3260	3260	-
$C_P \text{ (J.kg}^{-1} \cdot \text{K}^{-1})$	1050	1050	1050	-
$Q_r \text{ (W.m}^{-3})$	1.0×10^{-6}	1.0×10^{-10}	1.0×10^{-10}	-
$\alpha \text{ (K}^{-1})$	3.0×10^{-6}	3.0×10^{-6}	3.0×10^{-6}	-
$\beta \text{ (Pa}^{-1})$	1.0×10^{-11}	1.0×10^{-11}	1.0×10^{-11}	-
$\eta_{yp} \text{ (Pa.s)}$	1.0×10^{21}	1.0×10^{21}	1.0×10^{21}	-
$Q^{\text{Peierls}} \text{ (kJ.mol}^{-1})$	-	-	-	540
$\sigma^{\text{Peierls}} \text{ (Pa)}$	-	-	-	8.5×10^9
$E^{\text{Peierls}} \text{ (s}^{-1})$	-	-	-	5.7×10^{11}
q	-	-	-	2
γ	-	-	-	0.1
s	-	-	-	1

Table 1: Rheological and material parameters values used in our study for the reference model. The dislocation creep parameters used for the crust are those for Westerly Granite (Hansen & Carter, 1983). For the lithospheric and asthenospheric mantle, the chosen dislocation and diffusion creep parameters are those for Dry Olivine (Hirth & Kohlstedt, 2003) and the chosen Peierls creep are those for Olivine (Evans & Goetze, 1979; Kameyama et al., 1999). For elasticity, the shear modulus G is 30 GPa for all materials, which is in the range of values proposed for natural rocks at lithospheric depths (e.g., Turcotte & Schubert, 2002). Friction angle (ϕ) and cohesion (C) are set to be representative of Byerlee's law (J. Byerlee, 1978). Values of Q_r are from Jaupart & Mareschal, (2021) and Rudnick & Gao, (2003).

2.3 Parametric analysis

Two different systematic studies were performed from this reference model.

- (1) We systematically vary (i) crustal thickness (H) from 40 to 65 km (5 km increments) and (ii) initial plateau elevation (h_p) from 1 to 5 km (1 km increments), following their observed natural correlation (Figure 2a).

We assume that, by using constant crustal and mantle densities and excluding horizontal velocities at the lateral boundaries of our models, isostasy imposes a linear correlation between crustal thickness and plateau elevation. This relationship implies that a single plateau elevation corresponds to each crustal

thickness (Figure 2a). However, to explore a broader range of geodynamic scenarios and to capture the diversity of natural systems, we deliberately extend our models beyond this constraint. This approach allows us to simulate configurations where thick crust is associated with anomalously low/high plateau elevations, cases that cannot be explained by isostasy alone. Relying strictly on the isostatic relationship would prevent us from independently assessing the roles of crustal thickness and plateau elevation. Nonetheless, most of our models, particularly the reference model, remain close to this linear relationship (Figure 2a).

We use the same incision velocity (V_i) of 30 mm.yr⁻¹ used for the reference model.

- (2) We perform models with two different incision velocities (V_i) of 10 and 50 mm.yr⁻¹ considering the same combination of crustal thickness and initial elevation as for the previous systematic study.

A total of 48 models were performed. All models with their respective parameters are available in Table S1.

2.4 Model metrics

For each model, we design a set of metrics to quantify their evolution in terms of mechanical strength, temperature, exhumation and topography. To characterize valley morphology, we compute the valley maximum relief (Δh), corresponding to the difference between the maximum ($h_{\max}$) and minimum ($h_{\min}$) elevation of the model surface (Figure 3a and 3b). We also compute the effective erosion rate (E_{eff}) in the center of the valley by calculating the *time required to remove successive 2-km thick layers of material eroded by valley incision at the center of the valley*. From this, we deduce the cumulative amount of vertical exhumation at different time step in the center of the valley. Because of lateral topographic diffusion, the incision velocity V_i does not directly correspond to the effective erosion velocity in the center of the valley as E_{eff} depends on both V_i and K . Hence, in the reference model for instance (with $V_i = 30$ mm.yr⁻¹), the mean E_{eff} value in the center of the valley is 0.8 ± 0.25 mm.yr⁻¹ between 1 and 10 Myr of model simulations (Figure 3c).

3 Results

3.1 Reference model

3.1.1 Valley morphology and effective erosion

The valley relief (Δh) evolves over time, reaching a maximum value of 3.5 km after 4.5 Myr (Figure 3b). After this peak, Δh decreases almost linearly until the end of the simulation, reaching 3.2 km after 10 Myr. Notably, half of the total relief increase occurs within the first 1 Myr, emphasizing the transient nature of the topographic evolution under a prescribed incision rate. The primary driver of relief development is the decrease in $h_{\min}$, which decreases by 2.8 km after 4.5 Myr, rather than an increase in $h_{\max}$, which rises by only 700 m over the initial plateau elevation during the same period (Figure 3b).

After a sharp decrease during the first 1 Myr, $h_{\min}$ decreases gradually, particularly between 1.5 and 3 Myr. Between 3 and 5 Myr, $h_{\min}$ stabilizes at approximately 200 m before increasing again, reaching 600 m by the end of the simulation (10 Myr). In contrast, $h_{\max}$ increases almost linearly over the first 3 Myr, reaching 3.6 km, before gradually rising to a stable value of 3.8 km after 6 Myr (Figure 3b).

A bump of 100 to 200 m appears in both $h_{\min}$ and $h_{\max}$ curves after ~ 1.5 Myr, corresponding to the initiation of crustal convection (Fig. 4), which temporarily increases both values (Figure 3b).

The effective erosion rate (E_{eff}) follows a different evolution. It starts at ~ 9.5 mm.yr $^{-1}$ during the first 100 kyr, rapidly decreases to reach ~ 1 mm.yr $^{-1}$ after 1 Myr, and remains nearly constant at ~ 0.8 mm.yr $^{-1}$ afterwards (Figure 3c).




Figure 3: (a) Topographic profile of the reference model after 1 Myr of simulation. Measurement of the relief (Δh) and of the maximum ($h_{\max}$) and minimum ($h_{\min}$) altitude are indicated on the profile. (b) Time evolution of Δh , $h_{\max}$ and $h_{\min}$ for the reference model. The black triangle represents the moment when the lower crust is exhumed to the surface. The vertical dotted line represents when the initiation of the crustal convection starts. (c) Time evolution of the effective erosion rate at the center of the valley (E_{eff}) for the reference model.

3.1.2 Crustal deformation and exhumation

Due to the presence of crustal convection, which obscures a clear visualization of the deformation patterns, our analysis focuses exclusively on the upper crustal layers unaffected by convective motion. Valley incision induces crustal deformation and alters crustal strength. Over the 10 Myr of the simulation, the reference model exhibits significant upward motion of the lower crust, with 3.5 km of exhumation to the surface (Figure 4a). The lower crust reaches the surface after 4.6 Myr (Figure 3b). The upward motion of the lower crust is accompanied by the upward advection of isotherms, at a slower rate due to heat diffusion (Figure 4a and S2a). Throughout the simulation, the Moho depth remains largely stable, with only minor variations associated with the onset of crustal convection.

Crustal motions result in a non-uniform spatial distribution of strain rates (Figure 4b, left). *Over time, strain rates remain largely diffuse within the non-partially melted portion of the crust. However, zones of elevated strain rates ($>10^{-16} \text{ s}^{-1}$) develop within the viscous part of the non-melted crust, particularly beneath the center of the valley. After 5 Myr, this zone of localized high strain rates (up to 10^{-15} s^{-1}) reaches a width of approximately 10 km (Figure 4b). —two distinct high-strain rate zones ($>10^{-16} \text{ s}^{-1}$) develop in the viscous part of the crust (i.e., below 6 km (Figure S3 depth). The first is located beneath the valley center, with a width of 10 km after 5 Myr, where strain rates reach 10^{-15} s^{-1} . The second zone appears 20 km from the valley center, expanding to 30 km after 5 Myr, with strain rates of $7 \times 10^{-16} \text{ s}^{-1}$ (Figure 4b, left).* This high-strain rate zone in the viscous part of the crust is associated with brittle deformation in the upper crust (i.e., above 6 km of depth), exhibiting similar spatial dimensions and strain rate values (Figure 4b).

The analysis of the stress field and of the direction of the maximum principal stress (σ_1) suggest that the brittle deformation in the center of the valley corresponds to extensional deformation, whereas the brittle zones further away from the valley center experiences compressional deformation (Figure 4b, right). In the viscous part of the crust, this is the opposite, with compressional deformation below the valley and extensional deformation 20-30 km away from the valley center (Figure 4b, right). As the simulation progresses, the region of viscous compressional deformation beneath the valley center migrates upward, reducing the extent of extensional deformation near the surface (Figure 4b, right). Elsewhere in the crust, the deformation pattern remains stable, with compressional deformation at the surface and extensional deformation at greater depths.

Partial melting is also important in the crust, as seen with the convection cells, and can reach values up to 60% of rocks melted in this lower part of the crust (Figure S2b). This fraction tends to decrease to 40 % after 10 Myr of simulation.




Figure 4: Time evolution of the reference model after 1, 5 and 10 Myr. (a) material and thermal evolution. White lines represent each 200°C isotherm. Color legend is as for Figure 2b. (b) left: Evolution of the second invariant of the strain rate. right: Evolution of the angle between the maximum principal stress (σ_1) and the vertical axis.

3.2 Parametric study

3.2.1 Individual impact of crustal thickness and plateau elevation

Although crustal thickness (H) and plateau elevation (h_p) are closely linked, a given crustal thickness can correspond to different plateau elevations, and conversely (Figure 2a).


Varying crustal thickness while maintaining a constant plateau elevation of 3 km has only a minor impact on the evolution of relief (Δh) (Figure 5a). The main difference lies in the maximum relief, which reaches 3900 m for the thinner crust model ($H = 50$ km) and 3300 m for the thicker crust model ($H = 60$ km). This difference of 600 m is partly due to differences in the evolution of h_{min} . Indeed, in the thinner crust model, the river reaches its base level after 5.5 Myr, whereas in the thicker crust model, h_{min} only decreases down to 400 m after 3.5 Myr (Figure 5b).

Compared to the reference model, a model involving a thicker crust leads to a lower value of Δh . In such a model, the onset of exhumation to the surface of the lower crust is slightly earlier (4.6 Myr) and the river does not reach its base level (Figure 5a and 5b). In contrast, the presence of a thinner crust has the opposite effect, leading to higher Δh values, a delayed onset of lower crust exhumation to the surface (5.7 Myr), and a river reaching its base level (Figure 5a and 5b).

Instead, varying the initial plateau elevation, and hence the erosion potential, while maintaining a constant crustal thickness of 55 km results in larger differences. First, plateau elevation plays a key role in valley morphology. Models with a lower h_P value enable the river to reach the base level (Figure 5c and 5d). When $h_P = 1$ km, the river reaches its base level in 200 kyr and remains at this altitude throughout the entire simulation (Figure 5d). This has a major impact on the relief, with Δh not exceeding 1.2 km and remaining at this value for the duration of the simulation (Figure 5c). A similar pattern is observed for the model where $h_P = 2$ km, with differences in timing—the river reaches its base level at 1.1 Myr—and in the maximum Δh value, reaching 2.4 km. For $h_P \geq 3$ km (i.e., including the reference model), the models behave identically with the river unable to reach its base level (Figure 5c and 5d). Consequently, increasing the plateau elevation above 3 km, while keeping the same combination of incision velocity and crustal thickness (i.e., 30 mm.yr⁻¹ and 55 km), produces the same outcome (Figure 5c, 5d and S3b).

In addition, the initial plateau elevation also significantly impacts lower crust exhumation. In the model where $h_P = 1$ km, the exhumation of the lower crust to the surface does not occur within the 10 Myr of the simulation (Figure 5c). In the model where $h_P = 2$ km, lower crust exhumation is possible but occurs at 8.5 Myr. This is later than in the reference model ($h_P = 3$ km) or for models with a higher plateau elevation ($h_P > 3$ km), which all show the onset of lower crust exhumation at 4.6 Myr.

Effect of crustal thickness



Effect of plateau elevation




Figure 5: Time evolution of Δh (left), and $h_{\min}$ (right) for models with different values of H (a and b) or different values of h_P (c and d). For all models, the incision velocity (V_i) is fixed at 30 mm.yr⁻¹. Downward triangles represent the time when the lower crust is exhumed to the surface if it happens. Vertical dotted lines represent the time of the convection initiation within the crust.

3.2.2 Combined impact of crustal thickness and plateau elevation

Looking at the entire parametric study conducted for an incision velocity of 30 mm.yr⁻¹, the timing of lower crust exhumation appears to be mostly influenced by the initial plateau elevation rather than the crustal thickness (Figure 6a). Indeed, for models with $h_P \geq 4$ km, exhumation occurs at 4.6-4.7 Myr, while models with $h_P < 1$ km do not exhume lower crust over 10 Myr of simulation time. For intermediate h_P of 2-3 km, we observe an abrupt change in the timing of lower crust exhumation, with a certain sensitivity to crustal thickness as well (i.e., a thinner crust leading to a delayed exhumation time). For instance, at $h_P = 2$ km, exhumation time varies from more than 10 Myr to 6.9 Myr, by varying H from 45 to 60 km, respectively. A similar trend can be depicted at $h_P = 3$ km, with a range of 5.7 to 4.6 Myr when varying H between 50 to 65 km.

Apart from this difference in timing of the lower crust exhumation, the overall dynamics of the models are similar than in the reference model for the tested values of h_P and H (Figure S4 and S5). In terms of strain rate patterns, changes in crustal thickness or plateau elevation do not significantly alter the results either. The primary effect is a variation in strain rate magnitude that increases with an increase of crustal thickness as well as an increase of the initial plateau elevation (Figure S4 and S5).




Figure 6: Time for the exhumation to the surface of lower crust with a) $V_i = 30$ mm.yr⁻¹ and b) $V_i = 50$ mm.yr⁻¹. Values presented inside the circles are in Myr. The models with $V_i = 10$ mm.yr⁻¹ are not presented due to the absence of lower crust exhumation in all models in the 10 Myr timeframe.

3.2.3 Individual impact of the imposed incision velocity

Two different incision velocities (10 mm.yr^{-1} and 50 mm.yr^{-1}) were tested on the same set of model combinations as in the previous section with 30 mm.yr^{-1} . Results show that the observed strain rate pattern as well as the deformation regime of the model remains almost the same as in the reference model (Figure S6).

Incision velocity also plays a crucial role in the timing of lower crust exhumation to the surface. With an incision velocity of 10 mm.yr^{-1} , no exhumation occurs within the 10 Myr timeframe. With an incision velocity of 50 mm.yr^{-1} , the conditions leading to lower crust exhumation to the surface within the 10 Myr timeframe (Figure 6b) remain similar to those obtained for an incision velocity of 30 mm.yr^{-1} , even if exhumation of the lower crust occurs systematically earlier.

Incision velocity also has a significant impact on valley morphology and lower crust exhumation (Figure 7a). As expected, incision velocity directly influences the river ability to reach its base level. Keeping the same combination of crustal thickness and plateau elevation than in the reference model, the river fails to reach its base level in both the low-incision and reference models, whereas it successfully reaches its base level in the high-incision model (Figure 7b). Consequently, the evolution of Δh is closely tied to the capacity for the river to reach its base level. In the high-incision model, *Δh reaches a plateau after around 1 Myr, which roughly corresponds to the time when the valley floor reaches the base level (Figure 7b). Thereafter, Δh remains nearly stable over time, exhibiting only a slight increase accompanied by minor oscillations, which correspond to fluctuations in $h_{\min}$ and reaches a near steady state afterwards.* In contrast, for the other two models, Δh evolution is therefore not limited by the river base level and evolves according to the combination of imposed surface processes and crustal response (Figure 7a).

Incision velocity also plays a crucial role in the timing of lower crust exhumation to the surface. In the low-incision model, no exhumation occurs within the 10 Myr timeframe, whereas in the high-incision model, exhumation begins earlier (at 3.6 Myr) than in the reference model.




Figure 7: Time evolution of Δh (a), and $h_{\min}$ (b) for models with different initial incision velocity. For all models, the values of h_p and H are fixed at 3 and 55 km, respectively. Downward triangles represent the time when the lower crust is exhumed to the surface if it happens. Vertical dotted lines represent the time of the convection initiation within the crust.

4 Discussion

4.1 Effective erosion and comparison to natural river erosion

As explained in Section 2.5, the incision velocity does not directly correspond to the calculated effective erosion. The evolution of effective erosion can be divided into two distinct phases (Figure S7): (1) the initial phase (< 1 Myr), when most models exhibit a rapid decrease in E_{eff} (Figures S7 and 3c) and (2) a stabilization phase (> 1 Myr), when effective erosion values tend to a steady state (Figures S7 and 3c).

During the initial phase, the mean value of E_{eff} , ranging from 2 to 10 mm.yr⁻¹, is highly dependent on the imposed incision velocity, between 10 and 50 mm.yr⁻¹, for $h_p = 3$ km and $H = 55$ km. This phase derives from two different behaviors of our models: (i) For models that do not reach the base level within 1 Myr (see Figure S3), the observed decrease in E_{eff} is mainly driven by the action of topographic diffusion. In addition, this timescale of 1 Myr roughly corresponds to the diffusion timescale $t_K = (W/2)^2/4K$, ranging between 0.6 and 1.6 Myr for a valley width (W) varying between 30 and 50 km, respectively. Importantly, the decrease in effective erosion due to the topographic diffusion does not change the total volume of eroded material over time. (ii) For models in which the valley reaches the valley base level in less than 1 Myr (see Figure S3), the decrease in E_{eff} is primarily controlled by the base level. Once the valley reaches the base level, incision can no longer proceed, leading to a significant reduction in effective erosion. During this initial phase, the tectonic response to incision can be considered negligible with respect to the decrease in effective erosion.

During the stabilization phase, the mean E_{eff} values range from 0.1 to 2 mm.yr⁻¹ (Figure 8). This steady-state phase results from the tectonic response to incision. The development of a vertical crustal flow beneath the valley brings deeper crustal material to the surface, uplifting the river above the base level and enabling its incision to persist. Therefore, while V_i remains a key factor, h_p and H also influence long-term effective erosion values (Figure 8).

Natural valley incision rates can vary significantly due to differences in environmental settings, such as climatic conditions, drainage basin area, or lithology. Another factor contributing to these variations is the timescale over which incision rates are calculated or observed, as shorter- or longer-term measurements can lead to different estimates of incision rates. (e.g., Finnegan et al., 2014; Gallen et al., 2015, 2015; Mills, 2000). Recent studies have explored the relationship between incision rate values and the timescales over which they are measured, using datasets of river incision rates from various locations on Earth (Nativ and Turowski, 2020). This relationship indicates that for timescales shorter than 1 Myr, incision rates range from 0.01 to 5 mm yr⁻¹, with an average of 0.4 mm yr⁻¹. For timescales exceeding 1 Myr, incision rates range from 0.01 to 1 mm yr⁻¹, with an average of 0.08 mm yr⁻¹.

In our models, during the first Myr, effective erosion rates tend to be slightly higher than those observed in natural settings for this timescale. Notably, models with incision velocities of 50 mm yr⁻¹ yield mean effective erosion rates of 10 mm yr⁻¹ (Figure S7). Beyond 1 Myr, our models rarely produce effective erosion rates exceeding a threshold of 1.1 mm yr⁻¹ (Figure 8). Only a limited number of simulations with an incision velocity of 50 mm yr⁻¹ exceed this threshold (Figure 8c).




Figure 8: Mean values of the effective erosion velocity (E_{eff}) between 1 and 10 Myr for each models with, a) $V_i = 10 \text{ mm.yr}^{-1}$, b) $V_i = 30 \text{ mm.yr}^{-1}$, and c) $V_i = 50 \text{ mm.yr}^{-1}$. Values presented inside the circles are in mm.yr^{-1} . Black contours represent models with no exhumation of the lower crust in the 10 Myr timeframe. Red contours represent models presenting lower crust exhumation in the 10 Myr timeframe.

4.2 Crustal deformation and lower crust exhumation

Using visco-elasto-plastic models, Vernant et al. (2013) investigated the role of isostatic rebound induced by erosion in low-convergent mountain ranges. In their study, they performed some models without imposing tectonic boundary conditions and demonstrated that, in this context, erosion can induce regional uplift and contribute to maintaining mountain elevation. More importantly, the deformation regime observed in their models is very close to that obtained in our simulations, with erosion leading to horizontal extension in the uppermost kilometers of the crust and to horizontal compression at greater depths (Figure 4). This superficial extensional deformation is associated with plastic deformation and faulting in both models. Furthermore, the spatial pattern of crustal motion in both studies is similar, with intense vertical flow concentrated in the center of the model where erosion is applied, decreasing progressively away from the erosion zone, and ultimately leading to downward (negative) vertical flow at greater distances (Figure 9b). This vertical crustal flow is associated with the isostatic rebound induced by erosion, as mantle flow patterns in their study also highlight a vertical component. Similarly, our models show that valley incision can generate substantial vertical crustal flow, reaching values of nearly 0.8 mm yr^{-1} at the center of the valley in the reference model (Figure 9a). *An interesting feature observed in our models is that surface uplift can be decoupled from the lithospheric response, and that crust-mantle decoupling explains why the Moho remains stable while the lower crust migrates toward zones of lower pressure. This result can potentially explain the low effective elastic thickness (EET) often inferred from the isostatic response of the lithosphere to surface processes (Champagnac et al., 2007). At the global scale, EET values typically range between 30 and 80 km, reflecting a coupled mantle-crust system (e.g., Tesauro et al., 2012). However, in orogenic settings, EET values used to model uplift induced by erosion are generally significantly lower, often around 10 km (Champagnac et al., 2007). These low EET values likely reflect the mechanical behavior of the crust alone, rather than that of the entire lithosphere. This discrepancy helps to explain why erosion-induced uplift in orogens is best matched by low EET values. In such settings, the lower crust is fully ductile, enabling it to accommodate surface unloading without requiring a whole-lithosphere isostatic adjustment.*

This vertical motion of the crust promotes the rise of low-viscosity material toward the surface (Figure 9b). Despite some differences in the processes responsible for initiating crustal motion, these similarities suggest that whether erosion is applied to an already elevated mountain range (Vernant et al., 2013) or results from localized valley incision (as in our case), the resulting crustal deformation exhibits comparable patterns. In the former case, erosion contributes to maintaining mountain elevation, while in our models, it promotes valley relief development and lower crust exhumation.

The amount of exhumed lower crust at the center of the valley (as we refer as Q_{LC} hereafter), is highly sensitive to the input parameters of our models (Figure 10). Indeed, exhumation of the lower crust, reaching 6 km of depth below the model surface first depends on vertical incision of the valley bottom. Lower crust exhumation is first favored by the incision velocity V_i and the effective erosion rate E_{eff} , but is bounded by plateau elevation h_P , which sets the erosion potential. For instance, models with a 5 km high plateau relative to base level can exhume 5 over 6 km of depth simply by erosion. To reach lower crust material, valley incision is therefore not sufficient and a vertical uplift of the valley bottom, above the base level, is required to continue erosion. This second process, as previously discussed, depends on the upward velocity of viscous flow below the valley, and in turn on the effective viscosity of the lower crust. All other things being equal, increasing crustal thickness H , and in turn the ductile lower crust thickness, decreases this viscosity and increases the velocity of the upward flow (Figure 9b). A more rapid upward flow increases effective erosion rates and shortens the time required to exhume lower crust material.

To illustrate the importance of the ductile lower crust thickness, we performed two additional models. In the first one, we reduced the radiogenic heat production of the crust, resulting in an initially colder thermal structure (i.e., 800 °C at the Moho). In the second one, we applied a different rheology for the lower crust, using the Maryland Diabase flow law (Mackwell et al., 1998) (see Figures S8 and S9 for thermal and rheological parameters). Both configurations result in a stronger/thicker upper crust and a thinner ductile lower crust (Figures S8a and S9a). These models show that lower crustal exhumation is significantly reduced under these conditions, with exhumation occurring only after approximately 50 Myr of simulation time, at best (Figures S8b and S9b). These results emphasize the critical role of the ductile lower crustal thickness in enabling exhumation: a thinner ductile layer substantially delays lower crustal exhumation, or in some cases, inhibits it entirely.

. Therefore, if h_P and V_i mostly control the onset of lower crust exhumation, in particular for the most elevated plateaux, the amount of lower crust exhumation after a few million years is tightly linked to viscous flow of the lower crust in response to incision, and in turn to H *and the ductile lower crust thickness*.




Figure 9: (a) Vertical velocity at $z = -5 \text{ km}$ for different timesteps for the reference model with $H = 55 \text{ km}$ (left) and the model with $H = 65 \text{ km}$ (right). (b) Viscosity distribution for the reference model (left) and the model with $H = 65 \text{ km}$ (right) after 10 Myr. White arrows represent the velocity field. We only show velocity field within the non-partially melted part of the crust to enhance visibility and clarity.




Figure 10: Quantity of exhumed lower crust in the center of the valley (Q_{LC}) for the models with $V_i = 30 \text{ mm.yr}^{-1}$ after (a) 5 Myr and (b) 10 Myr of simulation. Values presented inside the circles are in km.

4.3 Application to natural settings: the conditions required for valley incision to exhume lower crust

Our models allow us to determine the conditions leading to the exhumation of lower crust material when a continental crust is submitted to valley incision only. In this configuration, exhumation of lower crust is only possible:

- (1) for high initial plateau elevation (i.e. for a value of $h_P \geq 2 \text{ km}$ over a 10 Myr timeframe). Over a 5 Myr timeframe, this threshold increases to 3 km.
- (2) for high crustal thickness (i.e. if $H > 50 \text{ km}$ over 10 Myr). Over 5 Myr, the threshold increases to 55 km.
- (3) for an erosion rate above 0.5 mm.yr^{-1} over 10 Myr, with a threshold increasing to 0.8 mm.yr^{-1} over 5 Myr.

Using these criterions, we can compare our findings to natural river systems on Earth associated or not with lower crust exhumation (Figure 11). We emphasize that no natural setting perfectly matches the boundary conditions and setup of our models, which in turn leads to a certain degree of interpretation when comparing models with data.

Rivers such as the Ebro and Potomac do not exhibit lower crust exhumation, which aligns well with our models when considering crustal thickness, elevation, incision rates, and duration of incision (Figure 11).

The Tsangpo and Indus rivers also align with our model predictions, as lower crust exhumation is observed in Namche Barwa and Nanga Parbat (Figure 11). However, given the intense tectonic forcing in these regions, with important convergence, we acknowledge that lower crust exhumation may also result from additional processes beyond valley incision. Nonetheless, our models provide a well-founded hypothesis regarding lower crust exhumation in these areas. A dedicated modeling study including horizontal convergence would be necessary to confirm this hypothesis, but such an investigation lies beyond the scope of this study.

The Yellow River, despite having crustal thickness and elevation within the predicted range for exhumation (Figure 11a), does not exhibit lower crust exhumation. This discrepancy is likely due to the short duration of incision, which began only 1.25 Myr ago (X. Wang et al., 2022) (Figure 11b). This underscores the critical role of incision duration in determining whether a river can induce lower crust exhumation.

Finally, the Grand Canyon is an outlier. It shows lower crust exhumation despite not meeting the required crustal thickness and plateau elevation predicted by our models (Figure 11a). The Grand Canyon presents a complex case and its formation remains debated, with some studies suggesting that regional uplift driven by mantle dynamics — known as epeirogeny — may have contributed to its development (Karlstrom and Timmons, 2012). Epeirogenic uplift corresponds to the vertical rise of large crustal regions without significant tilting, folding, or thrusting. This additional uplift, which is not accounted for in our models, could have contributed to increasing the plateau elevation, thereby enhancing the erosion potential. *Moreover, the presence of a Proterozoic basement beneath the sedimentary cover (Karlstrom and Timmons, 2012; Shufeldt et al., 2010) suggests that the eroded lower crustal material in the Grand Canyon was potentially already located near the surface prior to the onset of incision, which would have facilitated its exhumation.* Indeed, such a regional uplift, and *the presence of old lower crust material close to the surface*, could have enabled the Grand Canyon to exhume lower crustal material without necessarily meeting the specific crustal thickness, plateau elevation, and incision rates required in our models.




Figure 11: Comparison of our models to natural river systems for a) elevation and crustal thickness and b) incision rates and time of incision. The dotted red and orange lines represent the limit between models exhuming or not exhuming lower crust in the 10 Myr and 5 Myr timeframe, respectively. Red points represent river systems with lower crust exhumation. Different river systems and references used are: Eb = Ebro river (Regard et al., 2021), Po = Potomac river (Reusser et al., 2004), GC = Grand Canyon (Darling et al., 2012; Karlstrom et al., 2021; Pederson et al., 2002), Ts = Tsangpo river (near Namcha Barwa) (Finnegan et al., 2014; Koons et al., 2022; Wang et al., 2017), Ye = Yellow river (Tongde) (Harkins et al., 2007; Wang et al., 2022), In = Indus River (near Nanga Parbat) (Ali and De Boer, 2010; Burbank et al., 1996; Zeitler et al., 2001). Data of elevation and crustal thickness are from ETOPO1 (NOAA, 2009) and Crust 1.0 model (Laske et al., 2013), respectively.

Our models demonstrate that valley incision into continental crust can result in recognizable crustal deformation. In scenarios with the highest erosion potential, this deformation manifests as plastic strain within the upper kilometers of the crust. Notably, our models reveal a deformation regime associated with this plastic deformation, primarily extensional in the central part of the valley (Figure 4b). This insight opens the possibility of comparing model-predicted deformation with that observed in natural incised valleys. However, such a comparison is not straightforward. Natural examples chosen in this study exhibiting the largest erosion potential

(Indus, Tsangpo, and Yellow rivers) are located in tectonically active regions. As a result, these valleys do not display extensional deformation at their centers, due to the overriding influence of convergent tectonic settings. Conversely, rivers like the Ebro and Potomac have relatively low erosion potential, likely insufficient to induce observable incision-related surface deformation as observed in our models (Figure S3). Finally, while the Grand Canyon does feature extensional faults in its center (Billingsley et al., 2019), these faults predate the onset of Colorado River incision and therefore cannot be attributed to the valley formation. Consequently, the shallow deformation observed in these natural settings cannot be unequivocally interpreted as a consequence of the incision of their valleys.

5 Conclusions

We performed a series of thermo-mechanical numerical models *including surface processes* to investigate the deformation and exhumation of a thick continental crust subjected to valley incising an elevated plateau. Our results show that the incision of a single localized, but wide, valley is sufficient to generate significant crustal deformation, ultimately leading, for some specific conditions, to the exhumation of lower crustal material. This exhumation is enabled by the development of an upward crustal flow beneath the valley center, which facilitates the rise of low-viscosity material towards the surface and promotes lower crustal exhumation.

Our results indicate that lower crustal exhumation is only possible within a 10 Myr timeframe if favorable conditions are met, such as high elevation plateau (≥ 2 km), thick crust (> 50 km), and a long-term erosion rate (> 0.5 mm.yr⁻¹). Achieving exhumation within 5 Myr requires more restrictive conditions, with elevation plateau over 3 km, crustal thickness over 55 km and a long-term effective erosion over 0.8 mm.yr⁻¹.

We also show that the parameters controlling the onset of lower crustal exhumation differ from those governing the final quantity of exhumed material. While the onset of exhumation is primarily driven by the initial plateau elevation, the final amount of exhumed lower crust is largely controlled by the initial crustal thickness. An increase in crustal thickness leads to a thicker lower crust, which enhances the upward flow of material and ultimately increases the volume of exhumed lower crust.

Last, we show that our models can be informative when compared to natural examples, even though our model set-up does not perfectly replicate natural examples, particularly regarding regional tectonic boundary conditions. These first order results demonstrate that our model outcomes are broadly consistent with observations from natural systems.

Appendices

Appendix A: Rheological model

This appendix detailed the equation of each component of the strain rate solved in the 2D thermo-mechanical code MDoodz.

The viscous strain rate $\dot{\epsilon}_{ij}^v = \dot{\epsilon}_{ij}^{\text{dis}} + \dot{\epsilon}_{ij}^{\text{dif}} + \dot{\epsilon}_{ij}^{\text{Peierls}}$ accounts for the dislocation, *the diffusion and the* Peierls creep mechanism, represented by dis, dif and Peierls exponents, respectively. The dislocation and diffusion creep strain rate ($\dot{\epsilon}_{ij}^{\text{dis,dif}}$) can be expressed as:

$$\dot{\epsilon}_{ij}^{\text{dis,dif}} = \dot{\epsilon}_{\text{II}}^{\text{dis,dif}} \frac{\tau_{ij}}{\tau_{\text{II}}} = A \left(2f_{\text{dis,dif}} \exp\left(\frac{Q}{nRT}\right) \exp\left(\frac{-aM}{n}\right) \right)^{-n} d^{-m} \tau_{\text{II}}^n \frac{\tau_{ij}}{\tau_{\text{II}}}, \quad (\text{A1})$$

where n , A , and Q are the dislocation creep parameters of the material (see Table 1), a is a melt weakening factor set to 0 in the lithospheric *and astenospheric* mantle and 50 in the crust for the partial melting models, M the melt fraction and R is the universal gas constant ($R = 8.314510 \text{ J.K}^{-1}.\text{mol}^{-1}$). The subscript II indicates that this is the second *invariant* of the considered *tensor*. $f_{\text{dis,dif}}$ corresponds to the correction factor (e.g., Schmalholz & Fletcher, 2011) for invariant formulation relative to the type of experiments used for calibration (here uniaxial compression), such as:

$$f_{\text{dis,dif}} = \frac{1}{6} 2^{\frac{1}{n}} 3^{\frac{n-1}{2n}}, \quad (\text{A2})$$

Peierls strain rate is written:

$$\dot{\epsilon}_{ij}^{\text{Peierls}} = \dot{\epsilon}_{\text{II}}^{\text{Peierls}} \frac{\tau_{ij}}{\tau_{\text{II}}}, \quad (\text{A3})$$

with

$$\begin{cases} \dot{\epsilon}_{\text{II}}^{\text{Peierls}} = (2A^{\text{Peierls}})^{-s} \frac{\tau_{ij}}{\tau_{\text{II}}} \\ A^{\text{Peierls}} = f\gamma\sigma^{\text{Peierls}} \left(E^{\text{Peierls}} e^{-\frac{(1-\gamma)^2 Q^{\text{Peierls}}}{RT}} \right)^{-\frac{1}{s}} \\ s = \frac{Q^{\text{Peierls}}}{RT} (1-\gamma)^{(q-1)q\gamma} \end{cases} \quad (\text{A4})$$

s is the effective stress exponent (T -dependent), Q^{Peierls} is the activation energy, σ^{Peierls} is the Peierls stress, E^{Peierls} , q ($=2.0$), and γ ($=0.1$) (Evans & Goetze, 1979) (Table 1). Peierls creep stress is computed using a regularised formulation (Kameyama et al., 1999)

Elastic strain rate is expressed following Hooke's law as:

$$\dot{\epsilon}_{ij}^{\text{e}} = \frac{\dot{\tau}_{ij}}{2G} \frac{\tau_{ij}}{\tau_{\text{II}}}, \quad (\text{A5})$$

where G is the shear modulus (see Table 1) and $\dot{\tau}_{ij}$ is the Jaumann derivative of the deviatoric stress.

The 2D simulations include the effect of frictional rheology. To this end a viscoplastic Drucker-Prager rheological model is employed. The yield function is expressed as:

$$F = \tau_{\text{II}} - P \sin \varphi - C \cos \varphi - 2\eta_{\text{vp}} \dot{\epsilon}_{\text{II}}^{\text{vp}}, \quad (\text{A6})$$

where τ_{II} is the deviatoric second stress invariant, P the pressure, C the cohesion and φ the friction angle. The viscoplastic parameter η_{vp} is set such that the overstress ($2\eta_{\text{vp}} \dot{\epsilon}_{\text{II}}^{\text{vp}}$) is on the order of 1 MPa (i.e. $\eta_{\text{vp}} = 10^{21} \text{ Pa.s}$) in our simulations.

Appendix B: partial melting implementation

This appendix detailed the partial melting equation solved in the 2D thermo-mechanical code MDoodz. In partial melting models, latent heat (Q_L) can be significant during partial melting or crystallization events. During melting, Q_L acts as a heat sink ($Q_L < 0$) whereas crystallization results in heat production ($Q_L > 0$). In models including partial melting, this process is incorporated through: (i) a decrease in density with increasing melt fraction, (ii) the modification of effective viscosity and (iii) the consideration of the thermal impacts associated with melting/crystallization (Stüwe, 1995). The volumetric fraction of melt M is assumed to increase linearly with temperature, according to these relationships (Burg and Gerya, 2005; Gerya and Yuen, 2003):

$$\begin{aligned} M &= 0, \text{ at } T \leq T_{\text{solidus}} \\ M &= \frac{T - T_{\text{solidus}}}{T_{\text{liquidus}} - T_{\text{solidus}}}, \text{ at } T_{\text{solidus}} < T < T_{\text{liquidus}} \\ M &= 1, \text{ at } T \geq T_{\text{liquidus}} \end{aligned} \quad (\text{B1})$$

where T_{solidus} and T_{liquidus} are the wet solidus and dry liquidus temperatures of the rocks under consideration, respectively. The volumetric fraction of melt affects the effective density (ρ^{eff}) of the partially molten rocks as follows:

$$\rho^{\text{eff}} = \rho \left(1 - M + M \frac{\rho_0^{\text{molten}}}{\rho_0^{\text{solid}}} \right), \quad (\text{B2})$$

where $\rho_0^{\text{solid}} = \rho_0$ and $\rho_0^{\text{molten}} = \rho_0 - \Delta\rho$ (see Table 1).

Other processes associated with partial melting, such as solid-melt segregation and melt extraction, are neglected. The effect of latent heating due to equilibrium melting or crystallization is included implicitly by increasing the effective heat capacity (C_p^{eff}) and the thermal expansion (α^{eff}) of the partially molten rocks ($0 < M < 1$) (Burg and Gerya, 2005). Q_L is the latent heat of melting of the considered rocks.

$$C_p^{\text{eff}} = C_p + Q_L \left(\frac{\partial M}{\partial T} \right)_P \quad (\text{B3})$$

$$\alpha^{\text{eff}} = \alpha - \rho \frac{Q_L}{T} \left(\frac{\partial M}{\partial P} \right)_T \quad (\text{B4})$$

Crustal partial melting parameters and equations	Crust
T_{solidus} (K) if $P < 1600\text{MPa}$	$973 - 70400/(P + 354) + 77800000/(P + 354)^2$
T_{solidus} (K) if $P > 1600\text{MPa}$	$935 + 0.0035 \times P + 0.0000062 \times P^2$
T_{liquidus} (K)	$1423 + 0.105 \times P$
$\Delta\rho$ (kg.m ⁻³)	200
a	50
H_L [J.kg ⁻¹]	380,000

Table B1: Equations and parameters used to implement crustal partial melting. Equations for the liquidus and solidus temperatures are from Schmidt and Poli (1998) and Poli and Schmidt (2002), where P corresponds to the pressure in MPa. The value for the latent heat of melting (H_L) is in line with standard values used for crustal material (e.g. Gerya, 2019; Turcotte & Schubert, 2002). The value of the change in density due to partial melting ($\Delta\rho$) is in line with standard values, assuming a drop of density of 5-10% for partially melted granite (e.g., Philpotts & Ague, 2009). The a value used in Eq. A1 is set to produce a drop of viscosity of three orders of magnitude for 10 % of partial melting.

Code availability

The data of this study have been generated by using the code MDoodz7.0. This code is freely accessible and all the information needed to install the code and use it is available through Zenodo Geffroy et al., (2025a), including the updated files of the code specifically written for this study. The Matlab file used to read the Output files and produce the figures of this paper is also given in Geffroy et al., (2025a)

Data availability

All output data of the reference are through Zenodo (Geffroy et al., 2025b)

Authors contribution

TD and PY developed the model code and TG performed the simulations. TG prepared the manuscript with contribution of all co-authors.

Competing interest

The authors declare that they have no conflict of interest

Acknowledgements

This research has been supported by the Université de Rennes, CNRS and the H2020 European Research Council (grant agreement no. 803721) and the Institut Universitaire de France (IUF). P.Y. would particularly like to thank Jean Van Den Driessche for the discussions that initiated this work several years ago. Clément Cholet is also thanked here for having produced the first models of this type using MDoodz during his Master 2 internship. We

also want to sincerely acknowledge the two reviewers, Carole Petit and Guillaume Duclaux for their insightful comments that provided valuable guidance in improving our study.

Financial support

This work was supported by the Institut Universitaire de France (IUF)

References

- Ali, K. F. and De Boer, D. H.: Spatially distributed erosion and sediment yield modeling in the upper Indus River basin, *Water Resources Research*, 46, <https://doi.org/10.1029/2009WR008762>, 2010.
- Auzemery, A., Yamato, P., Duretz, T., Willingshofer, E., Matenco, L., and Porkoláb, K.: Influence of magma-poor versus magma-rich passive margins on subduction initiation, *Gondwana Research*, 103, 172–186, <https://doi.org/10.1016/j.gr.2021.11.012>, 2022.
- Avouac, J. P. and Burov, E. B.: Erosion as a driving mechanism of intracontinental mountain growth, <https://doi.org/10.1029/96JB01344>, 1996.
- Baiadori, F., Sacek, V., Freitas, B., and Almeida, R.: Evidence of crustal flexure induced by fluvial incisions, *Tectonophysics*, 877, 230292, <https://doi.org/10.1016/j.tecto.2024.230292>, 2024.
- Beaumont, C., Fullsack, P., and Hamilton, J.: Erosional control of active compressional orogens, in: *Thrust Tectonics*, edited by: McClay, K. R., Springer Netherlands, Dordrecht, 1–18, https://doi.org/10.1007/978-94-011-3066-0_1, 1992.
- Beaumont, C., Jamieson, R. A., Nguyen, M. H., and Lee, B.: Himalayan tectonics explained by extrusion of a low-viscosity crustal channel coupled to focused surface denudation, *Nature*, 414, 738–742, <https://doi.org/10.1038/414738a>, 2001.
- Braun, J. and Yamato, P.: Structural evolution of a three-dimensional, finite-width crustal wedge, *Tectonophysics*, 484, 181–192, <https://doi.org/10.1016/j.tecto.2009.08.032>, 2010.
- Brocard, G. Y. and van der Beek, P. A.: Influence of incision rate, rock strength, and bedload supply on bedrock river gradients and valley-flat widths: Field-based evidence and calibrations from western Alpine rivers (southeast France), in: *Tectonics, Climate, and Landscape Evolution*, edited by: Willett, S. D., Hovius, N., Brandon, M. T., and Fisher, D. M., Geological Society of America, 0, [https://doi.org/10.1130/2006.2398\(07\)](https://doi.org/10.1130/2006.2398(07)), 2006.
- Burbank, D. W., Leland, J., Fielding, E., Anderson, R. S., Brozovic, N., Reid, M. R., and Duncan, C.: Bedrock incision, rock uplift and threshold hillslopes in the northwestern Himalayas, *Nature*, 379, 505–510, <https://doi.org/10.1038/379505a0>, 1996.
- Burg, J.-P. and Gerya, T. V.: The role of viscous heating in Barrovian metamorphism of collisional orogens: thermomechanical models and application to the Lepontine Dome in the Central Alps, *Journal of Metamorphic Geology*, 23, 75–95, <https://doi.org/10.1111/j.1525-1314.2005.00563.x>, 2005.

738 Candioti, L. G., Duretz, T., and Schmalholz, S. M.: Horizontal Force Required for Subduction
739 Initiation at Passive Margins With Constraints From Slab Detachment, *Front. Earth Sci.*, 10,
740 <https://doi.org/10.3389/feart.2022.785418>, 2022.

741 Champagnac, J. D., Molnar, P., Anderson, R. S., Sue, C., and Delacou, B.: Quaternary
742 erosion-induced isostatic rebound in the western Alps, *Geol*, 35, 195,
743 <https://doi.org/10.1130/G23053A.1>, 2007.

744 Darling, A. L., Karlstrom, K. E., Granger, D. E., Aslan, A., Kirby, E., Ouimet, W. B., Lazear,
745 G. D., Coblenz, D. D., and Cole, R. D.: New incision rates along the Colorado River system
746 based on cosmogenic burial dating of terraces: Implications for regional controls on
747 Quaternary incision, *Geosphere*, 8, 1020–1041, <https://doi.org/10.1130/GES00724.1>, 2012.

748 Dingwell, D. B.: Transport Properties of Magmas: Diffusion and Rheology, *Elements*, 2, 281–
749 286, <https://doi.org/10.2113/gselements.2.5.281>, 2006.

750 Duretz, T., May, D. A., Gerya, T. V., and Tackley, P. J.: Discretization errors and free surface
751 stabilization in the finite difference and marker-in-cell method for applied geodynamics: A
752 numerical study, *Geochemistry, Geophysics, Geosystems*, 12,
753 <https://doi.org/10.1029/2011GC003567>, 2011.

754 Duretz, T., May, D. A., and Yamato, P.: A free surface capturing discretization for the
755 staggered grid finite difference scheme, *Geophys. J. Int.*, 204, 1518–1530,
756 <https://doi.org/10.1093/gji/ggv526>, 2016.

757 Duretz, T., de Borst, R., Yamato, P., and Le Pourhiet, L.: Toward Robust and Predictive
758 Geodynamic Modeling: The Way Forward in Frictional Plasticity, *Geophysical Research*
759 *Letters*, 47, e2019GL086027, <https://doi.org/10.1029/2019GL086027>, 2020.

760 Duretz, T., de Borst, R., and Yamato, P.: Modeling Lithospheric Deformation Using a
761 Compressible Visco-Elasto-Viscoplastic Rheology and the Effective Viscosity Approach,
762 *Geochemistry, Geophysics, Geosystems*, 22, e2021GC009675,
763 <https://doi.org/10.1029/2021GC009675>, 2021.

764 Finnegan, N. J., Schumer, R., and Finnegan, S.: A signature of transience in bedrock river
765 incision rates over timescales of 104–107 years, *Nature*, 505, 391–394,
766 <https://doi.org/10.1038/nature12913>, 2014.

767 Gallen, S. F., Pazzaglia, F. J., Wegmann, K. W., Pederson, J. L., and Gardner, T. W.: The
768 dynamic reference frame of rivers and apparent transience in incision rates, *Geology*, 43, 623–
769 626, <https://doi.org/10.1130/G36692.1>, 2015.

770 Geffroy, T., Yamato, P., Steer, P., Guillaume, B., and Duretz, T.: Files relative to MDoodz7.0
771 code used in Geffroy et al. paper entitled « Deformation and exhumation in thick continental
772 crusts induced by valley incision of elevated plateaus »,
773 <https://doi.org/10.5281/zenodo.15241965>, 2025a.

774 Geffroy, T., Yamato, P., Steer, P., Guillaume, B., and Duretz, T.: Output files of the reference
775 model from Geffroy et al. paper entitled « Deformation and exhumation in thick continental
776 crusts induced by valley incision of elevated plateaus »,
777 <https://doi.org/10.5281/zenodo.15241946>, 2025b.

778 Gerya, T.: Introduction to Numerical Geodynamic Modelling, Cambridge University Press,
779 487 pp., 2019.

780 Gerya, T. V. and Yuen, D. A.: Rayleigh–Taylor instabilities from hydration and melting propel
781 ‘cold plumes’ at subduction zones, *Earth and Planetary Science Letters*, 212, 47–62,
782 [https://doi.org/10.1016/S0012-821X\(03\)00265-6](https://doi.org/10.1016/S0012-821X(03)00265-6), 2003.

783 Hancock, G. S. and Anderson, R. S.: Numerical modeling of fluvial strath-terrace formation in
784 response to oscillating climate, *GSA Bulletin*, 114, 1131–1142, [https://doi.org/10.1130/0016-7606\(2002\)114<1131:NMOFST>2.0.CO;2](https://doi.org/10.1130/0016-7606(2002)114<1131:NMOFST>2.0.CO;2), 2002.

786 Harkins, N., Kirby, E., Heimsath, A., Robinson, R., and Reiser, U.: Transient fluvial incision
787 in the headwaters of the Yellow River, northeastern Tibet, China, *Journal of Geophysical*
788 *Research: Earth Surface*, 112, <https://doi.org/10.1029/2006JF000570>, 2007.

789 Harlow, F. and Welch, J.: Numerical calculation of time-dependent viscous incompressible
790 flow of fluid with free surface, *Physics of fluids*, 1965.

791 Jaupart, C. and Mareschal, J.-C.: Radiogenic Heat Production in the Continental Crust, in:
792 *Encyclopedia of Solid Earth Geophysics*, edited by: Gupta, H. K., Springer International
793 Publishing, Cham, 1298–1303, https://doi.org/10.1007/978-3-030-58631-7_243, 2021.

794 Kameyama, M., Yuen, D. A., and Karato, S.-I.: Thermal-mechanical effects of low-
795 temperature plasticity (the Peierls mechanism) on the deformation of a viscoelastic shear
796 zone, *Earth and Planetary Science Letters*, 168, 159–172, [https://doi.org/10.1016/S0012-821X\(99\)00040-0](https://doi.org/10.1016/S0012-821X(99)00040-0), 1999.

798 Karlstrom, K., Crossey, L., Matthis, A., and Bowman, C.: Telling time at Grand Canyon
799 National Park: 2020 update, National Park Service, <https://doi.org/10.36967/nrr-2285173>,
800 2021.

801 Karlstrom, K. E. and Timmons, J. M.: Faulting and uplift in the Grand Canyon region, in:
802 *Grand Canyon Geology: Two Billion Years of Earth’s History*, Geological Society of America,
803 [https://doi.org/10.1130/2012.2489\(06\)](https://doi.org/10.1130/2012.2489(06)), 2012.

804 Koons, P. O.: Two-sided orogen: Collision and erosion from the sandbox to the Southern Alps,
805 *New Zealand, Geol*, 18, 679, [https://doi.org/10.1130/0091-7613\(1990\)018<0679:TSOCAE>2.3.CO;2](https://doi.org/10.1130/0091-7613(1990)018<0679:TSOCAE>2.3.CO;2), 1990.

807 Koons, P. O., Hallet, B., Zeitler, P. K., and Meltzer, A. S.: Tectonic aneurysm: A culmination
808 of tectonic and geomorphic cooperation in mountain building, 2022.

809 Laske, G., Masters, G., Ma, Z., and Pasyanos, M.: Update on CRUST1.0 - A 1-degree Global
810 Model of Earth’s Crust, 2013.

811 Luisier, C., Tajčmanová, L., Yamato, P., and Duretz, T.: Garnet microstructures suggest ultra-
812 fast decompression of ultrahigh-pressure rocks, *Nat Commun*, 14, 6012,
813 <https://doi.org/10.1038/s41467-023-41310-w>, 2023.

814 Mackwell, S. J., Zimmerman, M. E., and Kohlstedt, D. L.: High-temperature deformation of
815 dry diabase with application to tectonics on Venus, *Journal of Geophysical Research: Solid*
816 *Earth*, 103, 975–984, <https://doi.org/10.1029/97JB02671>, 1998.

817 Martin, J., Cantelli, A., Paola, C., Blum, M., and Wolinsky, M.: Quantitative Modeling of the
818 Evolution and Geometry of Incised Valleys, *Journal of Sedimentary Research*, 81, 64–79,
819 <https://doi.org/10.2110/jsr.2011.5>, 2011.

820 Mills, H. H.: Apparent increasing rates of stream incision in the eastern United States during
821 the late Cenozoic, *Geology*, 28, 955–957, [https://doi.org/10.1130/0091-](https://doi.org/10.1130/0091-7613(2000)28<955:AIROSI>2.0.CO;2)
822 [7613\(2000\)28<955:AIROSI>2.0.CO;2](https://doi.org/10.1130/0091-7613(2000)28<955:AIROSI>2.0.CO;2), 2000.

823 Molitor, Z., Mittal, T., and Jagoutz, O.: The viscosity of a partially molten layer in a paleo-
824 orogenic plateau, *Earth and Planetary Science Letters*, 648, 119060,
825 <https://doi.org/10.1016/j.epsl.2024.119060>, 2024.

826 Montgomery, D. R. and Stolar, D. B.: Reconsidering Himalayan river anticlines,
827 *Geomorphology*, 82, 4–15, <https://doi.org/10.1016/j.geomorph.2005.08.021>, 2006.

828 Munch, J., Ueda, K., Schnydrig, S., May, D. A., and Gerya, T. V.: Contrasting influence of
829 sediments vs surface processes on retreating subduction zones dynamics, *Tectonophysics*,
830 836, 229410, <https://doi.org/10.1016/j.tecto.2022.229410>, 2022.

831 Nativ, R. and Turowski, J. M.: Site Dependence of Fluvial Incision Rate Scaling With
832 Timescale, *Journal of Geophysical Research: Earth Surface*, 125, e2020JF005808,
833 <https://doi.org/10.1029/2020JF005808>, 2020.

834 NOAA: ETOPO1 1 Arc-Minute Global Relief Model, 2009.

835 Pederson, J., Karlstrom, K., Sharp, W., and McIntosh, W.: Differential incision of the Grand
836 Canyon related to Quaternary faulting—Constraints from U-series and Ar/Ar dating, *Geology*,
837 30, 739–742, [https://doi.org/10.1130/0091-7613\(2002\)030<0739:DIOTGC>2.0.CO;2](https://doi.org/10.1130/0091-7613(2002)030<0739:DIOTGC>2.0.CO;2), 2002.

838 Philpotts, A. R. and Ague, J. J.: *Principles of Igneous and Metamorphic Petrology*, Cambridge
839 University Press, 669 pp., 2009.

840 Poh, J., Yamato, P., Duretz, T., Gapais, D., and Ledru, P.: Precambrian deformation belts in
841 compressive tectonic regimes: A numerical perspective, *Tectonophysics*, 777, 228350,
842 <https://doi.org/10.1016/j.tecto.2020.228350>, 2020.

843 Porkoláb, K., Duretz, T., Yamato, P., Auzemery, A., and Willingshofer, E.: Extrusion of
844 subducted crust explains the emplacement of far-travelled ophiolites, *Nat Commun*, 12, 1499,
845 <https://doi.org/10.1038/s41467-021-21866-1>, 2021.

846 Regard, V., Vacherat, A., Bonnet, S., Mouthereau, F., Nørgaard, J., and Knudsen, M. F.: Late
847 Pliocene-Pleistocene incision in the Ebro Basin (North Spain), *BSGF - Earth Sci. Bull.*, 192,
848 30, <https://doi.org/10.1051/bsgf/2021020>, 2021.

849 Reusser, L. J., Bierman, P. R., Pavich, M. J., Zen, E., Larsen, J., and Finkel, R.: Rapid Late
850 Pleistocene Incision of Atlantic Passive-Margin River Gorges, *Science*, 305, 499–502,
851 <https://doi.org/10.1126/science.1097780>, 2004.

852 Rudnick, R. and Gao, S.: Composition of the Continental Crust. *Treatise Geochem* 3:1-64,
853 *Treatise on Geochemistry*, 3, 1–64, <https://doi.org/10.1016/B0-08-043751-6/03016-4>, 2003.

854 Schmalholz, S. M. and Fletcher, R. C.: The exponential flow law applied to necking and
855 folding of a ductile layer: Exponential flow law applied to folding and necking, *Geophysical*
856 *Journal International*, 184, 83–89, <https://doi.org/10.1111/j.1365-246X.2010.04846.x>, 2011.

857 Shufeldt, O. P., Karlstrom, K. E., Gehrels, G. E., and Howard, K. E.: Archean detrital zircons
858 in the Proterozoic Vishnu Schist of the Grand Canyon, Arizona: Implications for crustal
859 architecture and Nuna supercontinent reconstructions, *Geology*, 38, 1099–1102,
860 <https://doi.org/10.1130/G31335.1>, 2010.

861 Simpson, G.: Role of river incision in enhancing deformation, *Geol*, 32, 341,
862 <https://doi.org/10.1130/G20190.2>, 2004.

863 Steer, P., Simoes, M., Cattin, R., and Shyu, J. B. H.: Erosion influences the seismicity of
864 active thrust faults, *Nat Commun*, 5, 5564, <https://doi.org/10.1038/ncomms6564>, 2014.

865 Sternai, P., Sue, C., Husson, L., Serpelloni, E., Becker, T. W., Willett, S. D., Faccenna, C., Di
866 Giulio, A., Spada, G., Jolivet, L., Valla, P., Petit, C., Nocquet, J.-M., Walpersdorf, A., and
867 Castelltort, S.: Present-day uplift of the European Alps: Evaluating mechanisms and models
868 of their relative contributions, *Earth-Science Reviews*, 190, 589–604,
869 <https://doi.org/10.1016/j.earscirev.2019.01.005>, 2019.

870 Stüwe, K.: Thermal buffering effects at the solidus. Implications for the equilibration of
871 partially melted metamorphic rocks, *Tectonophysics*, 248, 39–51,
872 [https://doi.org/10.1016/0040-1951\(94\)00282-E](https://doi.org/10.1016/0040-1951(94)00282-E), 1995.

873 Tesauero, M., Kaban, M. K., and Cloetingh, S. A. P. L.: Global strength and elastic thickness of
874 the lithosphere, *Global and Planetary Change*, 90–91, 51–57,
875 <https://doi.org/10.1016/j.gloplacha.2011.12.003>, 2012.

876 Thieulot, C., Steer, P., and Huismans, R. S.: Three-dimensional numerical simulations of
877 crustal systems undergoing orogeny and subjected to surface processes, *Geochemistry,*
878 *Geophysics, Geosystems*, 15, 4936–4957, <https://doi.org/10.1002/2014GC005490>, 2014.

879 Turcotte, D. L. and Schubert, G.: *Geodynamics*, Cambridge University Press, 484 pp., 2002.

880 Vernant, P., Reilinger, R., and McClusky, S.: Geodetic evidence for low coupling on the
881 Hellenic subduction plate interface, *Earth and Planetary Science Letters*, 385, 122–129,
882 <https://doi.org/10.1016/j.epsl.2013.10.018>, 2014.

883 Wang, X., Hu, G., Saito, Y., Ni, G., Hu, H., Yu, Z., Chen, J., Wang, M., Yuan, X., Wang, L.,
884 Hu, Z., Nie, J., and Pan, B.: Did the modern Yellow River form at the Mid-Pleistocene
885 transition?, *Science Bulletin*, 67, 1603–1610, <https://doi.org/10.1016/j.scib.2022.06.003>,
886 2022.

887 Wang, Y., Zhang, H., Zheng, D., von Dassow, W., Zhang, Z., Yu, J., and Pang, J.: How a
888 stationary knickpoint is sustained: New insights into the formation of the deep Yarlung
889 Tsangpo Gorge, *Geomorphology*, 285, 28–43,
890 <https://doi.org/10.1016/j.geomorph.2017.02.005>, 2017.

891 Willett, S. D.: Orogeny and orography: The effects of erosion on the structure of mountain
892 belts, *Journal of Geophysical Research: Solid Earth*, 104, 28957–28981,
893 <https://doi.org/10.1029/1999JB900248>, 1999.

894 Wolf, S. G., Huismans, R. S., Braun, J., and Yuan, X.: Topography of mountain belts
 895 controlled by rheology and surface processes, *Nature*, 606, 516–521,
 896 <https://doi.org/10.1038/s41586-022-04700-6>, 2022.

897 Yamato, P., Burov, E., Agard, P., Le Pourhiet, L., and Jolivet, L.: HP-UHP exhumation during
 898 slow continental subduction: Self-consistent thermodynamically and thermomechanically
 899 coupled model with application to the Western Alps, *Earth and Planetary Science Letters*, 271,
 900 63–74, <https://doi.org/10.1016/j.epsl.2008.03.049>, 2008.

901 Yamato, P., Duretz, T., May, D. A., and Tartèse, R.: Quantifying magma segregation in dykes,
 902 *Tectonophysics*, 660, 132–147, <https://doi.org/10.1016/j.tecto.2015.08.030>, 2015.

903 Yamato, P., Duretz, T., and Angiboust, S.: Brittle/Ductile Deformation of Eclogites: Insights
 904 From Numerical Models, *Geochemistry, Geophysics, Geosystems*, 20, 3116–3133,
 905 <https://doi.org/10.1029/2019GC008249>, 2019.

906 Zeitler, P. K., Koons, P. O., Bishop, M. P., Chamberlain, C. P., Craw, D., Edwards, M. A.,
 907 Hamidullah, S., Jan, M. Q., Khan, M. A., Khattak, M. U. K., Kidd, W. S. F., Mackie, R. L.,
 908 Meltzer, A. S., Park, S. K., Pecher, A., Poage, M. A., Sarker, G., Schneider, D. A., Seeber, L.,
 909 and Shroder, J. F.: Crustal reworking at Nanga Parbat, Pakistan: Metamorphic consequences
 910 of thermal-mechanical coupling facilitated by erosion, *Tectonics*, 20, 712–728,
 911 <https://doi.org/10.1029/2000TC001243>, 2001.

912
 913
 914