

Investigating fire-induced ozone production from local to global scales

Joseph O. Palmo¹, Colette L. Heald^{1,2}, Donald R. Blake³, Ilann Bourgeois⁴, Matthew Coggon^{5,6}, Jeff Collett⁷, Frank Flocke⁸, Alan Fried⁹, Georgios Gkatzelis¹⁰, Samuel Hall⁸, Lu Hu¹¹, Jose L. Jimenez^{5,12}, Pedro Campuzano-Jost^{5,12}, I-Ting Ku⁷, Benjamin Nault^{13,14}, Brett Palm⁸, Jeff Peischl⁵, Ilana Pollack⁶, Amy Sullivan⁷, Joel Thornton¹⁵, Carsten Warneke⁶, Armin Wisthaler^{16,17}, Lu Xu¹⁸

¹ Department of Civil and Environmental Engineering, Massachusetts Institute of Technology, Cambridge, MA, USA

² Institute for Atmospheric and Climate Science, ETH Zurich, Zurich, Switzerland

³ Department of Chemistry, University of California, Irvine, Irvine CA, USA

10 ⁴ Univ. Savoie Mont Blanc, INRAE, CARRTEL, Thonon-les-Bains, France

⁵ Cooperative Institute for Research in Environmental Sciences, University of Colorado Boulder, Boulder, CO, USA

⁶ NOAA Chemical Sciences Laboratory (CSL), Boulder, CO, USA

⁷ Department of Atmospheric Science, Colorado State University, Fort Collins, CO, USA

⁸ Atmospheric Chemistry Observations & Modeling Laboratory, NCAR, Boulder, CO, USA

15 ⁹ Institute of Arctic and Alpine Research (INSTAAR), University of Colorado, Boulder, CO 80309, USA

¹⁰ Institute of Energy and Climate Research, IEK-8: Troposphere, Forschungszentrum Jülich GmbH, Jülich, Germany

¹¹ Department of Chemistry and Biochemistry, University of Montana, Missoula, MT, USA

¹² Department of Chemistry, Univ. of Colorado-Boulder, Boulder, CO, USA

¹³ Department of Environmental Health and Engineering, Johns Hopkins University, Baltimore, MD, USA

20 ¹⁴ CACC, Aerodyne Research, Inc., Billerica, MA, USA

¹⁵ Department of Atmospheric Sciences, University of Washington, Seattle, WA, USA

¹⁶ Department of Chemistry, University of Oslo, Oslo, Norway

¹⁷ Institut für Ionenphysik und Angewandte Physik, Universität Innsbruck, Innsbruck, Austria

¹⁸ McKelvey School of Engineering, Washington University, St. Louis, MO, USA

25

Correspondence to: Joseph O. Palmo (jpalmo@mit.edu), Colette L. Heald (colette.heald@env.ethz.ch)

Abstract.

Tropospheric ozone (O_3) production from wildfires is highly uncertain; previous studies have identified both production and
30 loss of O_3 in fire-influenced air masses. To capture the total ozone production attributable to a smoke plume, we bridge the gap
between near-field fire plume chemistry and aged smoke in the remote troposphere. Using airborne measurements from several
major campaigns, we find that fire-ozone production increases with age, with a regime transition from NO_x -saturated to NO_x -
limited conditions, showing that O_3 production in well-aged plumes is largely controlled by nitrogen oxides (NO_x).
Observations in fresh smoke demonstrate that suppressed photochemistry reduces O_3 production by ~70% in units of ppb O_x

35 (O_3+NO_2) per ppm CO in the near-field (age<20hr). We demonstrate that anthropogenic NO_x injection into VOC-rich fire plumes drives additional O_3 production, sometimes exceeding 50 ppb above background. Using a box model, we explore the evolving sensitivity of O_3 production to fire emissions and chemical parameters. We demonstrate the importance of aerosol-induced photochemical suppression over heterogeneous HO_2 uptake, validate $HONO$'s importance as an oxidant precursor, and confirm evolving NO_x sensitivity. We evaluate GEOS-Chem's performance against these observations, finding the model captures fire-induced O_3 enhancements at older ages but overestimates near-field enhancements, fails to capture the magnitude and variability of fire emissions, and does not capture the chemical regime transition. These discrepancies drive biases in normalized ozone production ($\Delta O_3/\Delta CO$) across plume lifetime, though the model generally captures observed absolute O_3 enhancements in fire plumes. GEOS-Chem attributes 2.4% of the global tropospheric ozone burden and 3.1% of surface ozone concentrations to fire emissions in 2020, with stronger impacts in regions of frequent burning.

45 1 Introduction

Wildfires have been increasing in frequency and scale in many regions of the world over the past few decades (Abatzoglou et al., 2021; Westerling, 2016). Rising temperatures, drier conditions, and longer fire seasons attributed to climate change are expected to continue to drive this trend (Abatzoglou and Williams, 2016; Williams et al., 2019). Wildfires emit a range of reactive species to the atmosphere (Andreae, 2019; Lindaas et al., 2021; Permar et al., 2021), including particulate matter and the precursors necessary to form ozone (O_3) and particulate matter. Ozone is an important secondary pollutant that degrades air quality and negatively impacts human health while also being the third-most important greenhouse gas behind carbon dioxide and methane (Tarasick et al., 2019). The production of ozone from wildfire emissions has been a persistent source of uncertainty in assessing the global tropospheric ozone burden (Tarasick et al., 2019), which has implications for radiative forcing, global tropospheric oxidizing capacity (Fiore et al., 2024), and associated human health risks. In the US, the increase in wildfires has occurred in parallel with a rise in the number of homes near the wildland-urban interface (Burke et al., 2021), thus increasing human exposure to smoke and making it doubly important that we understand the impact of wildfires on air pollution.

Ozone formation is complex and nonlinear, occurring through a series of reactions between volatile organic compounds ($VOCs$), nitrogen oxides ($NO_x = NO + NO_2$), and oxidizing radical species in the presence of sunlight. Depending on the chemical environment and the relative concentrations of NO_x , $VOCs$, and radicals, the production rate of ozone will have different sensitivities to its precursors. At low NO_x concentrations, ozone production increases linearly with higher NO_x , and we refer to this as the NO_x -sensitive or NO_x -limited regime. As NO_x concentrations increase, ozone production is more sensitive to changes in VOC mixing ratios, and we refer to this regime as NO_x -saturated or VOC -limited (Kleinman, 1994; Sillman et al., 1990). The rate of ozone production reaches its maximum at the transition point between the NO_x -limited and VOC -limited regimes. In fact, the VOC -limited regime might be better characterized as "radical-limited", since ozone production is governed by the availability of peroxy radicals generated by the oxidation of $VOCs$, rather than the $VOCs$ themselves

(Schroeder et al., 2017). This is a distinction that becomes important in wildfire smoke where the availability of *VOCs* does not always correspond to radical availability when limited photolysis slows radical production. While both the photochemical production of ozone and the details of *NO_x-VOC-O₃* chemistry have been widely studied in the urban context and are reasonably well understood, the different and variable chemical environment presented by wildfire smoke plumes has proven challenging to measure and characterize. These uncertainties result from an inadequate understanding of (1) the emissions of *NO_x*, *VOCs*, and oxidant precursors across different fuel types and burning conditions (Gkatzelis et al., 2024; Jin et al., 2023a; Yokelson et al., 2013), (2) the dynamic chemistry that occurs as those emissions interact, age, and mix with other air masses, and (3) local (i.e. plume injection height) and downwind meteorological transport.

Previous studies have leveraged ground observations, and occasionally airborne data, to quantify ozone production in smoke plumes. Figure 1 shows a summary of some of these studies compiled by (Jaffe and Wigder, 2012). They identify both production and loss of *O₃* in fire-influenced air masses, with significant scatter introduced by the aging and mixing of plumes with other sources (Alvarado et al., 2010; DeBell et al., 2004; Pfister et al., 2008; Singh et al., 2010, 2012). The amount of ozone produced in smoke plumes is characterized by the normalized excess mixing ratio (NEMR) between *O₃* and *CO*, which are often well correlated in fire plumes. It is generally observed that this quantity, $\Delta O_3/\Delta CO$, is positive and increases with plume age.




Figure 1. Fire-induced ozone production ($\Delta O_3/\Delta CO$; ppb/ppb) from previous studies, compiled by Jaffe and Wigder 2012. Each data point represents a separate study; in some cases, multiple plumes are aggregated.

Xu et al. 2021 used a single transect analysis method to explore FIREX-AQ measurements of wildfire plumes close to the source, from which they derive a parameterization for ozone formation that agrees reasonably well with observations. Other recent wildfire-ozone studies use observationally constrained box model simulations to assess ozone production rates in fresh

smoke plumes and analyze the sensitivity to certain chemical factors, such as the breakdown of reactive *VOCs* and the changes to the radical budget by species emitted and produced within wildfires (namely nitrous acid (*HONO*), formaldehyde (*CH₂O*), and oxygenated *VOCs* (*OVOCs*)) (Robinson et al., 2021; Wolfe et al., 2022). The comprehensive chemical representation of these box modeling studies provides insight into the ozone chemical regime observed within smoke – they assert that typically a smoke plume begins in a *NO_x*-saturated regime before transitioning to a *NO_x*-limited regime within a few hours. Other studies have used observations together with more sophisticated numerical methods, such as large eddy simulations, to represent the spatial heterogeneity within smoke plumes, and the importance that this plays in determining photolysis rates and radical availability (Decker et al., 2021; Wang et al., 2021a). Near-field studies typically only focus on single plume events close to the source, limiting their utility for evaluating the lifecycle of *O₃* production in an airmass and the global impact of fires on the burden of *O₃*.

Because fire plumes emit large amounts of *NO_x* and *VOCs*, they may potentially produce significant downwind ozone, especially if they mix with anthropogenic emissions, which are typically high in *NO_x* (Jaffe and Wigder, 2012). Ground-based measurements from Mount Bachelor Observatory in Washington suggest that older, more chemically processed smoke plumes have experienced greater ozone production (Baylon et al., 2015), consistent with the results shown in Fig. 1. Brey and Fischer (2016) analyze ground-based observations to show that heavily populated U.S. cities far downwind of the wildfire-prone western U.S. experience increased *O₃* exceedances due to aged smoke. A box modeling study driven by measurements taken during the COVID-19 lockdowns at a site in Boulder, Colorado, similarly demonstrates that transported smoke can bring significant ozone enhancements (up to 30 ppb) while also driving the local urban chemical environment towards a more *NO_x*-sensitive regime (Rickly et al., 2023). These effects have been persistently observed; a study in Western Canada estimated an average surface ozone enhancement of at least 2 ppb across all smoke-influenced periods from 2001 to 2019 (Schneider et al., 2024). Lee and Jaffe (2024) employ a statistical approach to estimate that wildfires contribute ~7.8 ppb of additional surface ozone on smoky days compared to no-smoke days at surface measurement sites across the US. These far-field studies confirm the importance of wildfires in impacting ozone across spatiotemporal scales; however, the chemical insights that can be drawn from their conclusions are limited.

Modeling ozone production in wildfire plumes is challenging and resolution dependent. Global three-dimensional chemical transport models (CTMs) cannot, with current computational constraints, represent wildfire smoke in all its chemical complexity, which raises the question: how well do global models represent smoke plume evolution and what is the impact of that representation on the modeled distribution of tropospheric ozone? Bourgeois et al. (2021) analyzed airborne measurements from the NASA Atmospheric Tomography mission (ATom) and inferred that wildfire-induced ozone is underestimated by global CTMs in most regions of the remote troposphere. In their multi-decadal analysis of tropospheric ozone in GEOS-Chem, H. Wang et al. (2022) observed that years with higher biomass burning emissions experienced an overall 2–3% increase in the tropospheric ozone burden, highlighting the importance of modeling wildfire emissions and chemistry accurately.

Here we present an analysis of wildfire-induced ozone production that spans airborne measurements, zero-dimensional box modeling, and global chemical transport modeling. Our work aims to connect wildfire-induced ozone production across scales

by developing a process-based understanding informed by comparisons between observations and chemical models. Specifically, we use airborne observations to thoroughly define the limitations of our current global models in representing wildfire smoke chemistry and the resulting ozone production.

2 Materials and Methods

2.1 Description of observations

To investigate biomass burning plume chemistry, we focus on airborne measurements that sample fire smoke across a range of chemical environments within the troposphere. We use observations from five different airborne campaigns for this study: ATom (2016–2018), the Fire Influence on Regional to Global Environments and Air Quality (FIREX-AQ; July–September 2019) campaign, the Western wildfire Experiment for Cloud chemistry, Aerosol absorption and Nitrogen (WE-CAN; July–September 2018), the Deep Convective Clouds and Chemistry (DC3; May–June 2012) campaign, and the Arctic Research of the Composition of the Troposphere from Aircraft and Satellites (ARCTAS; March–July 2008) campaign. Within the ATom observations, only spring and summer (N. Hemisphere: April–September; S. Hemisphere: October–March) observations are used to match the other campaigns and avoid seasonal gradients that may impact our analysis. Figure 2a shows the spatial extent of these campaigns. Our observations contain measurements taken at distinct times during the chemical development of smoke, ranging from fresh emissions in the near-field (ARCTAS, FIREX-AQ, WE-CAN), to aged, chemically processed smoke in the far-field (ATom, ARCTAS). ARCTAS and DC3 also provide an intermediate sample with minor influence from aged smoke and some nearby urban sources.




Figure 2. (a) Location of flight tracks for the airborne field campaigns used in this analysis (campaign abbreviations in legend). (b) Flight tracks colored by regime type. (c) Fire-influenced observations colored by photochemical age (section 2.3) in hours.

We re-sample observations to 2-minute resolution to incorporate instruments with longer time resolutions. Measurements (Table S1) used in this study include carbon monoxide (CO), nitrogen oxides (NO_x , NO_y), nitric acid (HNO_3), peroxyacyl

145 nitrates (*PAN*), ozone (O_3), formaldehyde (CH_2O), nitrous acid (*HONO*), photolysis rates, and a collection of *VOCs* including
our fire and anthropogenic *VOC* tracers (see Section 2.2) and those used for photochemical aging (Section 2.3). In some cases,
multiple instruments measured a given compound. We leverage overlapping measurements from multiple instruments if the
instruments showed suitable agreement ($R^2 > 0.75$, normalized mean bias (*NMB*) < 0.2) across a substantial number of
measurements ($N > 50$). Finally, any data point contaminated by marine or stratospheric influence is removed from our analysis
150 following the criteria defined in Bourgeois et al. (2021), keeping only measurements with a $\frac{\Delta O_3}{H_2O}$ ratio between 0.003 and 1
ppb/ppm (Fig. S1). This results in a merged dataset of 19,942 points.

2.2 Regime analysis

We use measurements of tracer species to assess the relative influence of biomass burning and anthropogenic emissions on
each air mass sampled. We first subtract background levels from our observations. We separate our dataset into clean and
155 polluted subsets by splitting at the 40th percentile of carbon monoxide mixing ratios ($[CO] \approx 100$ ppb). The background level
for each measurement is defined as the 25th percentile mixing ratio in each 2km altitude bin in the free troposphere of the clean
subset of points. We define different background levels for the conterminous United States (FIREX-AQ, WE-CAN, DC3), the
remote measurements taken by ATom and the boreal, Arctic measurements taken by ARCTAS (Fig. S2). By subtracting this
background level and putting the observed tracers in enhancement (Δ) space for the regime definition, we avoid the
160 complications that arise from comparing data across a range of latitudes and altitudes in the free troposphere where background
concentrations vary. This is especially important when using longer-lived *VOC* tracers that exhibit structured vertical profiles.
We sort our dataset into different regimes – clean, fire, anthropogenic, or mixed (Fig. 2b) – using the observed enhancements
of measured tracer compounds commonly used to assess air mass influences in the airborne context due to their longer lifetimes
(weeks–months) (Alvarado et al., 2010; Bourgeois et al., 2021; Singh et al., 2010, 2012): acetonitrile (CH_3CN) and hydrogen
165 cyanide (*HCN*) for fire, and tetrachloroethylene (C_2Cl_4) and dichloromethane (CH_2Cl_2) for anthropogenic pollution. Where
possible, we use acetonitrile (CH_3CN) as our fire tracer, and tetrachloroethylene (C_2Cl_4) as our anthropogenic tracer, then opt
for the alternative tracer where the first choice is not available. We discard points where measurement of both a fire tracer and
anthropogenic tracer are not available.

Within the polluted subset of observations, any observation that is greater than the 50th percentile of the measured tracer
170 enhancement is classified as fire or anthropogenically influenced. If the observed air mass tracer concentrations are greater
than the 50th percentile of both the fire and anthropogenic tracer it is considered mixed. Points that lie below the 50th percentile
of both the fire and anthropogenic tracer may be considered lightly mixed or background pollution; these are not considered
in our analysis. The majority of our analysis focuses on the fire regime, which contains 4042 points, or 39% of the polluted
subset (Fig. S4). The anthropogenic regime contains 3242 points (31%), and the mixed regime contains 1552 points (15%).
175 The precise boundaries for our regime definitions (e.g. 40th percentile of CO demarcation of clean/polluted) were chosen based

on an examination of the distribution of measured concentrations and to ensure a sufficient number of data points in each category; the results presented herein are insensitive to modest shifts (i.e. within 10 percentile) in these boundaries.

2.3 Photochemical aging

It is difficult to assess ozone production from smoke from instantaneous measurements. To address this, we derive the age of our fire-influenced points to analyze the time evolution of the smoke plumes in our dataset (Fig. 2c). A common method is to approximate a physical smoke age using back trajectory analysis (Stein et al., 2015) and space-based observations of fires and smoke (Jin et al., 2023b). However, this approach is subject to physical uncertainties, including an assumption of injection height / altitude of emission, that compound over time, and its utility is further limited by the mixing between fresh and aged smoke. Instead, we adopt a chemical aging strategy that employs our in-situ measurements to compute an approximate smoke age. Assuming that (1) the chosen measured *VOCs* are co-emitted from biomass burning (i.e., well-correlated in fire influenced points; see Fig. S5) and (2) the removal of the chosen *VOCs* is dominated by reactions with hydroxyl radicals, one can use the following formula to estimate the photochemical age of smoke measured in sampled air masses (De Gouw, 2005):

$$\Delta t = \frac{1}{[OH] * (k_X - k_Y)} * \left[\ln\left(\frac{X}{Y}\right) \Big|_{t=0} - \ln\left(\frac{X}{Y}\right) \right] \quad (1)$$

190

Where $[OH]$ is the average concentration of hydroxyl radicals, X and Y are chosen VOCs, and k_X and k_Y are the rate constants for the reaction of OH with X and Y, respectively (here following the NASA JPL recommendations (“JPL Data Evaluation,”)). The emission ratio of X to Y is taken from the recent compilation from (Andreae, 2019), using the temperate forest values. The average concentration of OH is specified as $10^6 \frac{molec}{cm^3}$ (following recent work from (Liao et al., 2021)).

195 We strategically choose which X and Y VOC ratios to use based on the measurements available in each campaign and the age of the smoke that they sought to measure (see Fig. S6). For all campaigns, we use benzene as our denominator VOC (Y), as we expect it to remain elevated within the fire air masses that we sample given its longer lifetime ($\tau \approx 9d$). For WE-CAN observations, we calculate photochemical age using furan ($\tau \approx 7h$) as our numerator VOC (X). For FIREX-AQ, we use phenol ($\tau \approx 10h$) vs. benzene, following work by (Xu et al., 2021). For the other three campaigns we use toluene ($\tau \approx 1.9d$) vs. benzene to capture the evolution of aged smoke plumes further downwind.

200 The accuracy of the photochemical age estimate is limited by (1) the assumption of constant $[OH]$ during a plume’s lifetime, (2) the uncertainty in VOC emissions ratios and variation arising from diverse fuel types, different burning conditions, and plume-to-plume variability (3) the measurement uncertainty in our VOC observations, and (4) potential mixing between more and less aged air masses. Given this, we use the derived photochemical ages to qualitatively measure the evolution of a smoke plume, and the results of our study depend on relative rather than absolute ages, as discussed below.

205

2.4 Idealized 0-D box modeling

To explore chemical behavior occurring within wildfire smoke at different ages, we constructed a simplified zero-dimensional box model which represents the key reactions that drive HO_x - NO_x - VOC - O_3 photochemistry ($HO_x = OH + HO_2$; hydrogen oxide radicals). The box model is not used for direct comparison with observations, but for idealized calculations testing the relative importance of different chemical quantities in governing ozone production. Our mechanism is made up of 35 species and 69 reactions: HO_x - NO_x cycling and CO/CH_4 oxidation reactions and their rates are taken from the GEOS-Chem model, a lumped VOC scheme is employed with GEOS-Chem reaction rates for isoprene, and we include a parameter to scale the rates of our photolysis reactions, to maintain a measure of control on HO_x production and cycling. Default photolysis rates are kept constant and calculated using the simplified MCM parameterization assuming a solar zenith angle of 0° , simulating clear-sky photolysis with the sun directly overhead. Temperature is held constant at 288.15 K. Our findings are robust with respect to temperature and solar zenith angle variations, as shown in Fig. S9. The heterogeneous uptake of HO_2 onto aerosol is represented in the box model in the same fashion as GEOS-Chem – using a reaction probability parameterization (Jacob, 2000), with $\gamma = 0.2$. The box model is built using Catalyst.jl, a Julia package for high-performance simulation of chemical reaction networks (Loman et al., 2023). Catalyst.jl represents models symbolically to enable compatibility with other Julia libraries and the large number of numerical solvers Julia offers.

We use a Monte Carlo approach to test the sensitivity of ozone formation to a series of factors, running many steady-state box model simulations under various conditions to assess the resulting spread in ozone production. For each factor, a Monte Carlo experiment ($N=100$) is performed where every other factor is held constant at the default value, while the factor of interest is randomly initialized in each simulation by bootstrapping from the defined distribution. The ozone production rate (P_{O_3}) is computed for each simulation and the spread of the distribution of P_{O_3} reflects the importance of the factor of interest in governing the ozone production rate.

2.5 3D Model description

We use the global CTM GEOS-Chem (<https://geoschem.github.io/>) to simulate concentrations of ozone and its chemical precursors along the flight tracks of the campaigns described in section 2.1. GEOS-Chem Classic v14.3.0 (<https://zenodo.org/records/10640536>) was used in this study, with simulations performed with a horizontal resolution of $2^\circ \times 2.5^\circ$ and 72 vertical hybrid-sigma pressure levels. Additionally, we performed nested-grid simulations for the FIREX-AQ and WE-CAN campaigns at a higher spatial resolution of $0.5^\circ \times 0.625^\circ$ using boundary conditions from the $2^\circ \times 2.5^\circ$ global run. Six-month spin-up simulations were completed prior to the time periods of interest to ensure equilibrated initial conditions. The model is driven by the Modern Era Retrospective analysis for Research and Applications, Version 2 (MERRA-2) assimilated meteorology product from the NASA Global Modeling and Assimilation Office (GMAO). The model includes a HO_x - NO_x - VOC - O_3 -halogen photochemistry scheme (Bates and Jacob, 2019; Wang et al., 2021b) coupled to aerosol thermodynamics (Park et al., 2004).

Global anthropogenic emissions for each year follow the Community Emissions Data System (CEDSv2) (Hoesly et al., 2018) and are overwritten by regional inventories where possible. Biogenic VOC emissions are calculated online based on local meteorological conditions using the Model of Emissions of Gases and Aerosols from Nature version 2.1 (MEGANv2.1) emissions framework (Guenther et al., 2012). Year-specific fire emissions are taken from the satellite-derived global fire emissions database GFED4.1s and simulated at a 3-hour resolution (van der Werf et al., 2010). Pyrogenic VOC emissions are included in GFED using updated emission factors (Carter et al., 2022).

To test GEOS-Chem’s ability to reproduce wildfire-ozone production, we performed a series of simulations from 2008 to 2019, directly sampling the model at the locations and times of our airborne observations. For our analysis, we assume that the observationally defined regimes also apply to the model output, though this is not necessarily true in a coarse-grid global CTM with uncertain emissions inventories. As simulated smoke dilutes across grid boxes, its distribution will not always match the observations, meaning some “fire-influenced” points may include little to no simulated smoke in the model. To mitigate the impact of these “false-negative” points on our analysis, we filter our modeled fire distribution to include only points above the 40th percentile of modeled CO, and the model evaluation considers only this subset of points. To isolate fire-induced emissions and production within the model, as opposed to a more general bias, we calculate model background concentrations using the same method we applied to the observations, taking the modeled 25th percentile of the modeled clean ($[\text{CO}_{\text{model}}] < [\text{CO}_{\text{model}}]_{40\text{th}}$) population in each 2km altitude bin (Fig. S3). We subtract these background concentrations from our model output at each altitude level to obtain modeled enhancements (Δ) of certain compounds.

To estimate the total impact that fires have in governing the tropospheric ozone burden, we perform a perturbation sensitivity study with GEOS-Chem. Generally, source attribution model studies zero out emissions of the source of interest, and compare the results to a base case. In our work we aim to minimize the nonlinearities associated with the ozone chemical system, such as changing the oxidative potential of the global atmosphere. Hence, in our sensitivity simulations we increase and decrease wildfire emissions by 10% (referred to as BB1.1 and BB0.9, respectively) and scale the results up to approximate the total effect, linearized against both a positive and negative perturbation.

2.6 Approximating ozone production sensitivity to precursors

In Figure 3, the data is split using an observationally-derived threshold – the 50th percentile of θ' (within each bin) a reactivity weighted ratio between NO_x and CO (Eq. 3), used to approximate ozone production sensitivity to its precursors. Previous studies have used θ (Eq. 2) to define ozone formation regimes, especially in the urban context (Heald et al., 2020; Kirchner et al., 2001). Calculating θ requires a comprehensive and consistent estimate for total VOC reactivity, which is difficult to achieve, particularly across campaigns, despite improved instrumentation over the past several years. θ' is a more easily observed metric for approximating ozone formation regime, and its validity is confirmed qualitatively in our box modeling experiments (Fig. 7). This metric is especially relevant for analyzing fire-influenced airmasses, as total VOC emissions are extremely well correlated with CO ($R^2 = 0.98$; (Gkatzelis et al., 2024)). Additionally, CO is the largest contributor to OH reactivity (OHR) in the remote troposphere, and its concentration has previously been used as a proxy for assessing total OH reactivity in ATom

measurements (Baublitz et al., 2023). Therefore, we use OHR_{CO} as a qualitative predictor for VOC reactivity across smoke plume ages, in part because of its simplicity and the potential to extend this analysis further (e.g. towards satellite data).

$$\theta = \frac{OHR_{NOx}}{OHR_{ROC}} \quad (2)$$

$$\theta' = \frac{OHR_{NOx}}{OHR_{CO}} \quad (3)$$

where:

$$\begin{aligned} OHR_{CO} &= k_{CO+OH}[CO] \\ OHR_{NOx} &= k_{NO+OH}[NO] + k_{NO_2+OH}[NO_2] \\ OHR_{ROC} &= k_{CO+OH}[CO] + \sum_i k_{VOC_i+OH}[VOC_i]_i \end{aligned}$$

3 Results and Discussion

3.1 Observational analysis

3.1.1 Observed ozone formation regime transition




Figure 3. Observed ozone production ($\Delta O_3/\Delta CO$) plotted across 11 evenly log-spaced bins of photochemical age for all fire-influenced observations, split at the 50th percentile of θ' ($= OHR_{NOx}/OHR_{CO}$). The blue trace represents the median of the top 50th percentile points within each bin, while the shaded area extends from the 16th to the 84th percentile (corresponding to one standard deviation, but in outlier resistant percentile space) within each bin. The black trace represents the same, but for points below the 50th percentile of θ' . The black points represent historical data compiled by Jaffe and Wigder 2012 and shown in Fig. 1.

290 Observed fire ozone production expressed as $\Delta O_3/\Delta CO$ and binned over time is shown in Fig. 3. Because O_3 and CO are often correlated in smoke plumes, the normalized excess mixing ratio (NEMR) $\Delta O_3/\Delta CO$ is used to characterize wildfire-induced ozone production, as in many previous biomass burning studies (Liao et al., 2021; Müller et al., 2016; Robinson et al., 2021; Wolfe et al., 2022). CO is emitted in high concentrations in a wildfire plume and is effectively chemically inert (no production or loss) on timescales considered here; for those reasons, normalizing trace gas concentrations to CO accounts for plume
 295 dilution and entrainment of background air into the plume. When comparing multiple plumes, $\Delta O_3/\Delta CO$ also normalizes measured ozone production across fires of different sizes.

Observed $\Delta O_3/\Delta CO$ matches the results reported by Jaffe and Wigder from Fig. 1. Median $\Delta O_3/\Delta CO$ ranges from close to zero in young plumes, where NO titration of O_3 limits ozone production, to a peak of about 0.8 ppb ppb⁻¹ in aged plumes. In the near-field, observed $\Delta O_3/\Delta CO$ in the top 50th percentile (high NO_x) of θ' experience lower $\Delta O_3/\Delta CO$ than those in the lower
 300 50th percentile (low NO_x). A crossover happens around $t = 5\text{--}10$ hr, and the opposite behavior is observed in the far-field – higher ozone production in the top 50th percentile of θ' . This behavior indicates a chemical regime shift from NO_x -saturated behavior in freshly emitted fire plumes to NO_x -limited behavior in aged plumes. The evolving NO_x sensitivity is also seen when observations are split at the 50th percentile of $[NO_x]$ (Fig. S7) or the formaldehyde to NO_2 ratio (Fig. S8). This confirms the theoretical predictions in previous studies (Robinson et al., 2021; Wang et al., 2021a; Wolfe et al., 2022), which have
 305 estimated the regime shift to occur within the first few hours of a plume's lifetime – the fact that we observe the crossover point at slightly older photochemical ages likely has to do with the limited resolution of our calculated photochemical age related to the assumption of constant $[OH]$ (See Sect. 2.3). In Figure 3, we reproduce the variability seen across previous studies (Jaffe and Wigder, 2012) and offer an explanation of the observed spread in $\Delta O_3/\Delta CO$ based on our suite of measurements – the presence of NO_x in the aged, NO_x -limited environment drives significant ozone production.

310 3.1.2 Near-field photolysis suppression

NO_x -saturated air masses typically produce ozone efficiently in the presence of organic compounds. Photolysis suppression can limit the HO_x source, slowing the rate of VOC oxidation and thus ozone production – an effect that is expected in some freshly emitted smoke where aerosols can limit the radiation in a smoke plume core (Palm et al., 2021). In Figure 4, we explore how this effect manifests itself in the observations. We take the near-field observations with a photochemical age of less than
 315 20 hours and isolate points experiencing smoke-related suppression of photolysis (observed $j_{NO_2} < 0.005\text{ s}^{-1}$, 25th percentile of near-field and organic aerosol $>20\text{ }\mu\text{g}/\text{m}^3$, 40th percentile of near-field). We compare total odd oxygen ($O_x = O_3 + NO_2$) production here to control for the temporary titration of ozone in heavy smoke. Observations with measured photolysis rates higher than the cutoff point experience enhanced O_x production as a function of CO compared to those below: $65.1\text{ }\frac{\text{ppb}}{\text{ppm}}$ vs. $18.0\text{ }\frac{\text{ppb}}{\text{ppm}}$. This analysis demonstrates the importance of photochemical suppression in governing observed ozone production
 320 in young smoke plumes. From our analysis, we cannot conclude how much local ozone production suppression reduces the

overall O_3 production in the lifecycle of a fire plume; future plume-following case study analysis may be able to further explore this question.




Figure 4. Measured ΔO_x ($\Delta O_3 + \Delta NO_2$) vs. ΔCO for young ($t < 20h$) fire-influenced points. Points sampled in heavy-smoke ($j_{NO_2} < 0.005 s^{-1}$ and organic aerosol concentrations $> 20 \mu g m^{-3}$) are shown in grey; others in yellow. Fits are made with reduced major axis regression.

3.1.3 Transport and mixing: ozone production across regimes

In the free troposphere, most air masses are generally NO_x -limited, such that ozone production is primarily driven by the abundance of NO_x (Chameides et al., 1992), as seen in our analysis of aged fire plumes (Fig. S7). It has been suggested that enhanced anthropogenic-fire mixed ozone production is observed when a VOC-rich biomass burning plume entrains additional NO_x from an anthropogenic source, leading to downwind ozone production (Brey and Fischer, 2016; Jaffe and Wigder, 2012; McClure and Jaffe, 2018; Permar et al., 2023; Rickly et al., 2023; Singh et al., 2012). We explore the extent to which this is supported by the airborne data considered here. We find that the mixed regime exhibits larger ΔO_3 values than both the fire and anthropogenic regimes on average ($\mu_{\Delta O_3, fire} = 18.5$ ppb; $\mu_{\Delta O_3, anthro} = 21.1$ ppb; $\mu_{\Delta O_3, mixed} = 27.0$ ppb), as shown in Fig. 5a. We ascribe these disparities to the different mean chemical environments (reactive nitrogen budget, and VOC reactivity) that are observed in each regime.




Figure 5. (a) Probability density distributions of observed ΔO_3 across fire, urban, and mixed regimes. (b) Mean speciated measured NO_y across regimes. Patterns correspond to each of the most abundant gas phase NO_y species. Box plots represent NO_x/NO_y (box plot with star denoting mean) and $VOCR_t$ (box plot with triangle denoting mean) distributions for each regime.

First, we compare the mean reactive nitrogen (NO_y) budget within each regime. NO_y is derived from both anthropogenic and biomass burning sources and contains both fresh nitrogen emissions and reservoir species that are produced during aging. In Figure 5b, we report mean mixing ratios of the five most abundant gas-phase NO_y species measured across all the campaigns considered here: NO_x ($=NO+NO_2$), HNO_3 , PAN , and $HONO$ (which was only measured during FIREX-AQ and WE-CAN; hence the HONO average only reflects these campaigns). In general, NO_y concentrations are largest in fire air masses, followed by mixed, then anthropogenic – but these differences should be interpreted within the airborne in-situ context. Because urban NO_x has a local lifetime on the order of hours (Laughner and Cohen, 2019), and the campaigns used here did not focus on sampling urban air masses, our anthropogenic airborne observations do not directly measure freshly emitted urban NO_x , but instead reflect only the fraction of reactive nitrogen that has escaped the boundary layer and entered the free troposphere. In contrast, the majority of our fire influenced observations are taken in the near-field (67%) and are rich in NO_x , as reflected in Fig. S10. Both the fire and mixed regimes contain a relatively larger amount of PAN as compared to the anthropogenic points. As shown by the NO_x/NO_y box plots with the stars denoting the means, in the mixed population a higher proportion of the observed NO_y remains unprocessed in the form of NO_x compared to the anthropogenic population, which could be due to either the mixing of fresher anthropogenic NO_x emissions, temperature-driven PAN decomposition, or remaining NO_x emitted from fires.




Figure 6. The relationship between NO_x and CH_3CN enhancements for mixed pollution observations in log-log space. Points are colored by C_2Cl_4 enhancements and sized by ozone enhancements. $\Delta\text{NO}_x/\Delta\text{CH}_3\text{CN}$ ratios are overplotted.

To distinguish between NO_x that is leftover in smoke vs. the anthropogenic injection of NO_x in the mixed population, we examine the relationship between ΔNO_x and our anthropogenic and fire tracers. We use the ratio between ΔNO_x and $\Delta\text{CH}_3\text{CN}$ as an indicator for fresh anthropogenic NO_x in mixed smoke plumes (Singh et al. 2012; Juncosa Calahorrano et al., 2020), shown in Fig. 6. Points outlined in yellow, with higher $\Delta\text{NO}_x / \Delta\text{CH}_3\text{CN}$ and higher $\Delta\text{C}_2\text{Cl}_4$ represent the subset of the mixed population that are directly influenced by an injection of anthropogenic NO_x (Fig. S11). It is these measurements that experience the most extreme ozone enhancements – at times greater than 50 ppb – and represent the high tail of the mixed distribution in Fig. 5a.

The additional chemical processing that has taken place in the urban case during its transport from the surface likely means more accumulated ozone production and mixing has occurred before sampling the anthropogenic air masses – which explains the higher ozone production seen in anthropogenic points compared to fires in Fig. 5a. Hence, a fraction of the mixed

population ozone enhancements will likely also be a result of mixing between fire-produced ozone and urban-produced ozone. In Figure 6, the O_3 enhancements observed within the subset of the mixed population that exhibit lower anthropogenic NO_x enhancements (i.e., lower $\Delta NO_x / \Delta CH_3CN$) are primarily due to the mixing of “historically produced” ozone – the combination of both fire-produced ozone that formed within a smoke plume before mixing, and ozone that formed in the anthropogenic context before the smoke arrived. Whereas when VOC-rich aged smoke mixes with fresh anthropogenic NO_x (i.e., higher $\Delta NO_x / \Delta CH_3CN$ and higher ΔC_2Cl_4), ozone is directly produced downwind in addition to that historically produced ozone. This distinction is an important one, where ozone formation in the latter case could be preventable through anthropogenic NO_x reductions.

In Figure 5b we confirm that higher organic emissions from fires lead to higher VOC reactivity in the mixed population compared to the anthropogenic, which enhances ozone production by interaction with NO_x . We use available VOC measurements to compute a partial VOC reactivity ($VOCR_i$) for our observations, a quantity that relates to the production of organic radicals that contribute to ozone production (see S.I.). In the fire and mixed populations, C_2H_4 , *Furans*, and oxygenated *VOCs* such as *MVK*, CH_3OH , and *methacrolein* enhance observed $VOCR_i$. We hypothesize that this enhanced VOC population is the reason we see the largest ozone enhancements for mixed pollution in Fig. S10, spanning the range of observed NO_x/NO_y .

3.2 Idealized box model sensitivity analysis

3.2.1 Exploring the sensitivity of ozone production in fire plumes

We use our box model to test the sensitivity of ozone production against specific factors including (1) the speciation and reactivity of the emitted *VOC* population, (2) the range of wildfire emissions observed across fires that vary in size, fuel, and burning conditions, (3) the emission of radical precursors, (4) the evolving availability of NO_x , and (5) aerosol effects. We use observed conditions to drive the model and examine the spread of the resulting simulated ozone production rate that correspond to Monte Carlo simulations runs for each factor as a metric for assessing importance.

The box model results in Fig. 7 consider three scenarios relating to the physical transport and chemical aging of smoke. The “at emission” population, shown in green, reflects emission ratio estimates from Gkatzelis et al. (2024), in which emissions of various species are scaled against CO . The airborne observations are split into near-field (near field: $t < 20h$; blue) and far-field (far field: $t > 20h$; orange) populations, and for gas phase measurements we also scale concentrations against CO . Using this framework allows CO concentration to act as a metric describing the approximate total magnitude of emissions of the fire / smoke plume, and further isolates the other factors that govern chemical variability within a plume of any given size.

The sensitivity of ozone production rate to certain chemical species changes over time. Notably, HONO is the most important oxidant precursor at the point of emission and, to lesser degree, in the near field (and negligibly in the far-field). As expected

from Fig. 3, ozone production becomes more sensitive to NO_x with age, especially in the far field, reflecting the results from the airborne observations. The sensitivity to VOC shows the opposite trend, peaking in the near-field and decreasing with age. Higher CO concentrations, corresponding to larger fires, also lead to extremely high ozone production rates, manifested by the outlying near-field points in the CO factor that are driven by the tail of the observed CO distribution.

We also explore how smoke aerosol can reduce ozone production, by 1) suppressing photolysis and thus limiting oxidant concentrations (i.e., reducing P_{HO_x}), and 2) acting as a site for heterogeneous uptake and removal of HO_2 (i.e., increasing L_{HO_x}). In general, at the observed aerosol concentrations, our model is much more sensitive to optical effects / photolysis suppression (P_{HO_x}) than to chemical effects (L_{HO_x}), and this behavior is consistent across age. Ozone production rates are negligibly sensitive to HO_2 uptake, given that the enhancements of NO_x and VOC in smoke tend to overwhelm the chemical system, and the increased heterogeneous chemical effect is thus insignificant.





Figure 7. a) Ozone isopleth in log-log space overplotted with ovals representing distributions of NO_x and VOC concentrations for three different populations: time of emission (green), near-field (blue), and far-field (orange). b) Distributions for each factor of interest corresponding to each population. Near-field and far-field distributions are taken directly from the airborne observations, while the time of emission population relies on emission factors from Gkatzelis et al. (2024). c) P_{O_3} distributions corresponding to the Monte Carlo simulations run for each factor, repeated for each population.

3.3.1 Evaluating fire-induced ozone production in GEOS-Chem

We examine how well the GEOS-Chem model captures the airborne observations across the lifecycle of a smoke plume. In what follows, all comparisons are for fire-influenced points only.



425 **Figure 8. Comparison between GEOS-Chem simulated (red) and fire-influenced observations (black), plotted over observationally**
430 **derived photochemical ages, binned in log-space. The lines represent the median, and the shaded area extends from the 16th to the**
435 **84th percentile (corresponding to one standard deviation, in outlier resistant percentile space) within each bin. The dashed line**
represents the approximate transport time for a 2x2.5 grid box at mid-latitudes.

Emissions and resolution

430 GEOS-Chem is biased low in its representation of fresh fire *CO* and *NO_x* (Figures 8a, 8b), as expected given that standard
Eulerian CTMs dilute emissions from localized sources across an entire model grid cell, and thus the model is unable to
reproduce concentrated plumes (such as those observed in FIREX-AQ and WE-CAN) at the default resolution of 2°×2.5°
(Rastigejev et al., 2010). Air mass ages of < 20 hr likely all fall within the grid box of emission. In addition, the model
dramatically underestimates the variability in *CO* and *NO_x*, particularly in the near-field. Current global fire emissions
435 inventories, including GFED4 as used here, do not represent the full range of fuel types, burning conditions, and fire sizes
thereby limiting the ability of GEOS-Chem to represent the variability and range of fire emissions. GEOS-Chem also
underestimates total wildfire *CO* emissions, shown by the consistent ΔCO low bias across all ages. While observed ΔCO is
highest in the near-field and decreases with age, the model remains only moderately elevated across the plume lifetime (Fig.
8a). These comparisons suggest that the *NO_x/CO* emission ratio is overestimated in the model for the fires measured in our

440 dataset. Accurately representing NO_x/CO has implications for ozone production rates, especially after aging has occurred, as shown in Fig. 3, and the smoke chemical environment is NO_x -sensitive.

We find that a higher resolution ($0.5^\circ \times 0.625^\circ$) simulation of the FIREX-AQ and WE-CAN campaigns (which include the majority of the near-field points) does not improve the agreement between near field ΔNO_x and ΔCO (see Fig. S12). This
445 suggests that the 4-fold increase in resolution is insufficient to resolve most sampled fire plumes, and without accurate emissions inventories, it does little to improve the simulation of fire smoke.

We also note that simulated concentrations of radical precursors, $HONO$ and CH_2O (Fig. S14), are biased low in the near-field. These species produce OH when they photolyze and therefore drive ozone production. $HONO$ emissions have been shown to
450 provide the dominant source of OH in near-field wildfire plumes (Peng et al., 2020), and our box model confirms the importance of $HONO$ for ozone production rates in the near-field (Fig. 7). However, wildfire emissions of $HONO$ are currently neglected in GEOS-Chem; this omission contributes to the myriad of challenges for reproducing sub-grid near-field ozone formation.

455 *Fire-induced ozone production*

Estimating the ozone formed from fires is predicated on removing the ozone background. Previous assessments of background ozone in GEOS-Chem have observed a high model bias; these biases are spatially varying (Guo et al., 2018). Bourgeois et al. (2021) found that CTMs typically underestimate wildfire ozone enhancements in the remote troposphere, which they attribute in part to the artificially high background O_3 in their models. After removing the simulated background (see Fig. S3) GEOS-
460 Chem overestimates observed ΔO_3 in fire-influenced air masses in the near-field, but captures observed ΔO_3 in the far field (Fig. 8, near-field: $R^2=0.55$, NMB=54%; far-field $R^2=0.43$, NMB=18%); the latter is in contrast to the results of Bourgeois et al. (2021) in the far-field. We find that the ozone background in GEOS-Chem for the regions sampled by the aircraft campaigns varies sharply from the observed background, emphasizing the importance of carefully accounting for this background when quantifying ozone production from fires.

465

In Figure 8d, we visualize the comparison between simulated and observed ozone in fire-influenced air masses in units of $\Delta O_3/\Delta CO$ (as in Fig. 1 and Fig. 3). This reveals consistent high bias in simulated wildfire ozone production (normalized by a proxy for total emissions, ΔCO) across both the near-field ($R^2=0.44$, NMB=193%) and far-field ($R^2=0.37$, NMB=80%). Given the underestimate in CO emissions (Fig. 8a) this suggests that the model is overly efficient in ozone production across plume
470 ages. Thus, while the model simulates the correct magnitude of ozone enhancements in aged fire plumes in particular (see Fig. 8c), this is the result of offsetting biases in emissions and chemistry.

In the near-field the model is unable to capture the NO_x suppression and photochemical suppression in concentrated smoke plumes because of limitations in resolution and underestimated emissions. This drives more efficient near-field ozone production in the model than observed, similar to previous observations for ship plumes (Rastigejev et al., 2010), producing elevated $\Delta O_3/\Delta CO$.

Figure S12 shows that the GEOS-Chem model generally reproduces the range of the far-field observed ΔNO_x distribution, with the exception of the high tail of values above 1 ppb. While the airborne observations show higher $\Delta O_3/\Delta CO$ for points with higher ΔNO_x in Fig. S12b, this behavior is not reproduced in the model. This suggests that, despite similar NO_x levels, ozone production in the model in these air masses is less NO_x sensitive and tends more towards the transition regime (see Fig. S13). In Figure S14, we compare the observed and modeled age evolution of two common ozone formation regime proxies, HNO_3/H_2O_2 and $HCHO/NO_2$. The respective high and low biases in these ratios confirm that the GEOS-Chem fire-influenced air masses are less sensitive to NO_x in the far-field than the observations. Indeed, the model is biased low across all ages for many of the measured VOCs (such as benzene; Fig. S14), and oxidant precursors (HCHO, HONO; Fig. S14). The ratio between NO and NO_2 in Fig. S14 shows a high bias across all ages, which is consistent with insufficient conversion of NO to NO_2 due to an underestimate in the peroxy radical population, due to the missing VOC reactivity in GEOS-Chem simulated smoke (Carter et al., 2022). Thus, the far-field ozone production high bias in the model likely reflects a combination of higher ozone production in the far-field due to less NO_x -sensitivity in the model, as well as excess ozone production in the near-field transported downwind.

While GEOS-Chem (with GFEDv4.1s) does not capture the magnitude and variability of wildfire emissions, leading to a misrepresentation in the smoke ozone formation regime in both the near-field and far-field, the total amount of ozone produced (ΔO_3) in the far-field is well captured by the model. This suggests that while more work is needed to accurately capture the evolution of fire plume chemistry – such that the changing ozone formation sensitivity is captured – outside of source regions, the model reasonably reproduces the magnitude of ozone produced in wildfires.

3.3.2 Estimating wildfire impact on tropospheric ozone and human health

In Figure 9, we quantify the overall impact that fires have in governing the tropospheric ozone burden in the GEOS-Chem model. For 2020, we estimate that fire emissions contribute 2.4% of the annual mean tropospheric ozone burden, and 3.1% to annual mean surface ozone concentrations. The total burden estimate agrees reasonably well with the estimate made by Jaffe and Wigder 2012 that 3.5% of the global in-situ tropospheric O_3 production is due to fires, based on scaled average observed regional NEMRs ($\Delta O_3/\Delta CO$) against inventory estimates of fire CO emissions.

The contribution of fires to simulated ozone burden is regionally variable and highest near areas of frequent burning. In the Amazon and Central Africa fires account for over 10% of the tropospheric burden (Figure 9a). The transient fires in the Western

505 U.S. and Canada produce a smaller contribution to the annual mean ozone, but seasonally exceed 5% of the tropospheric burden.

In Figure 9b, we show that fire emissions account for up to 30% of annual mean surface ozone in the most affected regions, with a spatial distribution that largely matches the total burden. The mean percentage of population-weighted exposure to ozone due to fires is generally consistent (3–4%) for each continent, but higher contributions in South America (5%) and Africa (10.5%) (Figure 9c). We also highlight three regions of interest (marked on Figure 9b): in the Western U.S., 6.7% of annual mean population-weighted ozone exposure is due to wildfires, and in South and West Africa, that number reaches 21% and 18%, respectively. It is worth noting that fire emissions exhibit interannual variability (Cooper et al., 2024) and that fire emissions inventories struggle with regionally-dependent uncertainties (Wiedinmyer et al., 2023). The reported results describe GEOS-Chem’s representation of fire impacts on global ozone in a single year (2020), subject to the uncertainties in fire emission inventories.




Figure 9. Impact of wildfire emissions on tropospheric and surface ozone in GEOS-Chem for the year 2020. (a) Percent change in the tropospheric ozone column burden due to wildfire emissions, with the largest enhancements over tropical Africa and South America. (b) Percent of surface ozone concentrations attributable to wildfire emissions. (c) Regional, population weighted surface ozone exposure. The regions in (c) correspond to the colored / patterned outlines in (b).

4 Conclusions

This study uses data from several airborne campaigns to provide a comprehensive analysis of wildfire-induced tropospheric ozone production during the lifetime of a smoke plume. We observe the chemical regime transition from NO_x -saturated to NO_x -limited that occurs during smoke aging and identify enhanced ozone production in the far-field when NO_x is available to drive ozone chemistry. We quantify the importance of suppressed photochemistry in limiting ozone production under heavy smoke in the near-field. We show that generally, mixed fire-anthropogenic air masses display higher ozone production than air masses that are solely fire-influenced, and suggest that anthropogenic NO_x mixing with VOC-rich smoke is the primary driver of these enhancements. We expand upon the work of several previous studies (Bourgeois et al., 2021; Brey and Fischer, 2016; Rickly et al., 2023; Singh et al., 2012) and provide a constraint on the amount of ozone that may be attributed to the mixing between fire smoke and anthropogenic airmasses.

We find that the GEOS-Chem model overestimates near-field ozone enhancements, but captures the amount of ozone produced in far-field fire-influenced airmasses. However, spatial resolution and missing emissions contribute to the overestimate of normalized ozone production in both the near-field and far-field. In the near-field, we show that GEOS-Chem cannot capture concentrated fire emissions, and that the model cannot capture the NO_x -suppressed and photochemically-suppressed sub-grid conditions. Other recent work has established that GEOS-Chem struggles to reproduce the impacts of large wildfires in the western US (Carter et al., 2020), in part because chemical transport models with coarse resolution cannot accurately resolve sub-grid processes (Eastham and Jacob, 2017) or the transport of synoptic scale plumes due to numerical diffusion issues (Rastigejev et al., 2010). (Wang et al., 2021a) used a large-eddy simulation coupled to a chemical model to demonstrate that while a resolution of 1km was sufficient for capturing downwind chemistry, a model with 4km resolution failed at representing chemical regime shifts and thus incorrectly estimated O_3 formation. With current computational constraints, GEOS-Chem and other global chemical transport models cannot operate at such fine resolutions. Resolution limitations coupled with uncertain emissions inventories drive persistent biases in the ozone formation regime behavior seen in the model. In the far-field, comparing GEOS-Chem to the observed response of different integrated (HNO_3/H_2O_2 , $HCHO/NO_2$) and instantaneous (NO/NO_2) metrics of ozone chemistry and radical cycling reveals that the model does not capture the observed shift towards NO_x -limited regime, likely due to underestimated VOC reactivity in fire plumes. Hence, the agreement in wildfire-ozone enhancements between GEOS-Chem and the airborne observations in the far-field may be fortuitous – more work is needed to ensure that models capture smoke chemical evolution correctly, so that accurate predictions of ozone air quality can be made, especially in the context of mixing with anthropogenic pollution, which is shown to drive extreme exceedance events.

550 There are additional factors that may influence the formation of ozone from fires that were not explored in this study and could be further investigated in future work. Neither our observational analysis nor our model evaluation focused directly on fire plume injection height, though it contributes uncertainty to the transport process and determines where wildfire-induced ozone production will occur. To define the chemical smoke aging process with more confidence, our photochemical aging method could be validated using other approaches. Having a reliable physical age for smoke could corroborate our photochemical age
555 metric while also providing information about oxidation in wildfire plumes. Limited to our set of observations, this study did not consider the impact that nighttime chemistry has on smoke plumes, though previous work has shown the importance of the diurnal cycle in changing oxidation patterns and plume chemistry (Decker et al., 2021; Robinson et al., 2021; Wang et al., 2021a).

560 Additional observations are needed to connect near-field chemistry to far-field chemistry, both for confirming the phenomena discussed above and to provide additional benchmarks for model development. More observations are needed particularly in regions outside the US – the campaign data that we synthesize does not fully represent the fuel types and meteorological conditions outside North America. Additionally, more observations of smoke mixing with urban air would be valuable. For fire-ozone chemistry studies, observations of the radical population (HO_2 , OHR), radical precursors ($HONO$), and reservoir
565 species (HNO_3 , H_2O_2) would be valuable. Consistent tracers for fires and anthropogenic activity remain essential. The superior coverage of geostationary satellites provides exciting avenues for future analysis, and observations of formaldehyde and nitrogen dioxide may be leveraged to better understand the aging of fire plumes in relation to NO_x - VOC - O_3 photochemistry in near real-time, following work by (Jin et al., 2023b).

570 Improving global simulation of wildfire-induced ozone production will require further investigation into fire dynamics and chemistry, and ongoing refinement of emissions inventories and related parameterizations. Despite resolution limitations, efforts to capture variability in wildfire emission inventories could improve model skill in reproducing the smoke chemical environment. More work is needed to comprehensively characterize leading sources of smoke VOC reactivity and incorporate near-field radical sources such as $HONO$ in models. One potential avenue for representing sub-grid wildfire smoke chemistry
575 within GEOS-Chem (and similar models) would be to use the type of sub-grid parameterization used for ship plumes (Vinken et al., 2011). And finally, improved representation of ozone background concentrations would enable more productive comparisons with observations.

With climate-driven increases in fire activity and frequency in many regions of the world, it is becoming increasingly important to better understand how these fires contribute to ozone formation throughout the troposphere. Additional comprehensive
580 measurements that characterize this evolution over the lifecycle of a fire plume are essential for testing and identifying opportunities to improve models. High fidelity models are needed to facilitate more comprehensive estimates of the tropospheric ozone burden and drive more intelligent policy measures and mitigation strategies in the face of a changing climate.

Code availability

585 The code used to run the analysis and the data used to generate the figures included in this paper are publicly available at:
<https://zenodo.org/records/17127240>.

Data availability

The GEOS-Chem model is publicly available at: <https://zenodo.org/records/10640383>. The ARCTAS campaign data are available at: <https://www-air.larc.nasa.gov/cgi-bin/ArcView/arctas>. The ATom campaign data are available at:
590 <https://espo.nasa.gov/atom>. The DC3 campaign data are available at: https://www.eol.ucar.edu/field_projects/dc3. The FIREX-AQ campaign data are available at: <https://www-air.larc.nasa.gov/cgi-bin/ArcView/firexaq>. The WE-CAN campaign data are available at: https://www.eol.ucar.edu/field_projects/we-can.

Author Contributions

CLH and JOP designed the study. JOP led the analysis and performed the simulations. DRB, IB, MC, JC, FF, AF, GG, SH,
595 LH, JLJ, PCJ, ITK, BN, BP, JP, IP, AS, JT, CW, AW, LX provided measurements used in the analysis (See Table S1). JOP and CLH wrote and edited the paper with input from the co-authors.

Acknowledgements

We also acknowledge the following investigators for providing measurements (see Table S1): Andrew Weinheimer, Bruce Daube, Chelsea Thompson, Colm Sweeney, Delphine Farmer, Dirk Richter, Eric Apel, Glenn Diskin, Glenn Wolfe, Greg
600 Huey, Hannah Allen, James Elkins, Jason St. Clair, Jessica Gilman, John Crounse, Kathryn McKain, Kirk Ullmann, Patrick Veres, Paul Wennberg, Roisin Commane, Steve Wofsy, Teresa Campos, Thomas Hanisco, Thomas Ryerson, Wade Permar, Yong Zhou. The model simulations and analyses presented here were conducted using the “Svante” cluster, a facility located at MIT’s Massachusetts Green High Performance Computing Center and jointly supported by the MIT Joint Program on the Science and Policy of Global Change; the Department of Earth, Atmospheric and Planetary Sciences; the Department of Civil
605 and Environmental Engineering; the Institute for Data, Systems, and Society; and the Center for Global Change Science.

Competing interests

One of the (co-)authors is a member of the editorial board of ACP.

Financial Support

This work was funded by the U.S. National Science Foundation (AGS-2223070). L.H. acknowledges additional support from
610 NSF (EPSCoR-2242802 and AGS-2144896). The CU-Boulder group was supported by NASA 80NSSC23K0828.
S.H. was supported by the NSF National Center for Atmospheric Research, which is a major facility sponsored by the US
National Science Foundation under cooperative agreement no. 1852977. Their research was funded by NASA award nos.
NNX08AD41G, NNX15AG71A and 80NSSC18K0638.

References

- 615 Abatzoglou, J. T. and Williams, A. P.: Impact of anthropogenic climate change on wildfire across western US forests, *Proc. Natl. Acad. Sci. U.S.A.*, 113, 11770–11775, <https://doi.org/10.1073/pnas.1607171113>, 2016.
- Abatzoglou, J. T., Battisti, D. S., Williams, A. P., Hansen, W. D., Harvey, B. J., and Kolden, C. A.: Projected increases in western US forest fire despite growing fuel constraints, *Commun Earth Environ*, 2, 1–8, <https://doi.org/10.1038/s43247-021-00299-0>, 2021.
- 620 Alvarado, M. J., Logan, J. A., Mao, J., Apel, E., Riemer, D., Blake, D., Cohen, R. C., Min, K.-E., Perring, A. E., Browne, E. C., Wooldridge, P. J., Diskin, G. S., Sachse, G. W., Fuelberg, H., Sessions, W. R., Harrigan, D. L., Huey, G., Liao, J., Case-Hanks, A., Jimenez, J. L., Cubison, M. J., Vay, S. A., Weinheimer, A. J., Knapp, D. J., Montzka, D. D., Flocke, F. M., Pollack, I. B., Wennberg, P. O., Kurten, A., Crounse, J., Clair, J. M. S., Wisthaler, A., Mikoviny, T., Yantosca, R. M., Carouge, C. C., and Le Sager, P.: Nitrogen oxides and PAN in plumes from boreal fires during ARCTAS-B and their impact on ozone: an
625 integrated analysis of aircraft and satellite observations, *Atmospheric Chemistry and Physics*, 10, 9739–9760, <https://doi.org/10.5194/acp-10-9739-2010>, 2010.
- Andreae, M. O.: Emission of trace gases and aerosols from biomass burning – an updated assessment, *Atmos. Chem. Phys.*, 19, 8523–8546, <https://doi.org/10.5194/acp-19-8523-2019>, 2019.
- JPL Data Evaluation: <https://jpldataeval.jpl.nasa.gov/>, last access: 25 March 2024.
- 630 Bates, K. H. and Jacob, D. J.: A new model mechanism for atmospheric oxidation of isoprene: global effects on oxidants, nitrogen oxides, organic products, and secondary organic aerosol, *Atmospheric Chemistry and Physics*, 19, 9613–9640, <https://doi.org/10.5194/acp-19-9613-2019>, 2019.
- Baublitz, C. B., Fiore, A. M., Ludwig, S. M., Nicely, J. M., Wolfe, G. M., Murray, L. T., Commane, R., Prather, M. J., Anderson, D. C., Correa, G., Duncan, B. N., Follette-Cook, M., Westervelt, D. M., Bourgeois, I., Brune, W. H., Bui, T. P., DiGangi, J. P., Diskin, G. S., Hall, S. R., McKain, K., Miller, D. O., Peischl, J., Thames, A. B., Thompson, C. R., Ullmann, K., and Wofsy, S. C.: An observation-based, reduced-form model for oxidation in the remote marine troposphere, *Proceedings of the National Academy of Sciences*, 120, e2209735120, <https://doi.org/10.1073/pnas.2209735120>, 2023.
- 635 Baylon, P., Jaffe, D. A., Wigder, N. L., Gao, H., and Hee, J.: Ozone enhancement in western US wildfire plumes at the Mt. Bachelor Observatory: The role of NO_x, *Atmospheric Environment*, 109, 297–304, <https://doi.org/10.1016/j.atmosenv.2014.09.013>, 2015.
- Bourgeois, I., Peischl, J., Neuman, J. A., Brown, S. S., Thompson, C. R., Aikin, K. C., Allen, H. M., Angot, H., Apel, E. C., Baublitz, C. B., Brewer, J. F., Campuzano-Jost, P., Commane, R., Crounse, J. D., Daube, B. C., DiGangi, J. P., Diskin, G. S.,

- Emmons, L. K., Fiore, A. M., Gkatzelis, G. I., Hills, A., Hornbrook, R. S., Huey, L. G., Jimenez, J. L., Kim, M., Lacey, F., McKain, K., Murray, L. T., Nault, B. A., Parrish, D. D., Ray, E., Sweeney, C., Tanner, D., Wofsy, S. C., and Ryerson, T. B.: Large contribution of biomass burning emissions to ozone throughout the global remote troposphere, *Proc. Natl. Acad. Sci. U.S.A.*, 118, e2109628118, <https://doi.org/10.1073/pnas.2109628118>, 2021.
- Brey, S. J. and Fischer, E. V.: Smoke in the City: How Often and Where Does Smoke Impact Summertime Ozone in the United States?, *Environ. Sci. Technol.*, 50, 1288–1294, <https://doi.org/10.1021/acs.est.5b05218>, 2016.
- Burke, M., Driscoll, A., Heft-Neal, S., Xue, J., Burney, J., and Wara, M.: The changing risk and burden of wildfire in the United States, *Proc. Natl. Acad. Sci. U.S.A.*, 118, e2011048118, <https://doi.org/10.1073/pnas.2011048118>, 2021.
- Carter, T. S., Heald, C. L., Jimenez, J. L., Campuzano-Jost, P., Kondo, Y., Moteki, N., Schwarz, J. P., Wiedinmyer, C., Darmenov, A. S., da Silva, A. M., and Kaiser, J. W.: How emissions uncertainty influences the distribution and radiative impacts of smoke from fires in North America, *Atmos. Chem. Phys.*, 20, 2073–2097, <https://doi.org/10.5194/acp-20-2073-2020>, 2020.
- Carter, T. S., Heald, C. L., Kroll, J. H., Apel, E. C., Blake, D., Coggon, M., Edtbauer, A., Gkatzelis, G., Hornbrook, R. S., Peischl, J., Pfannerstill, E. Y., Piel, F., Reijrink, N. G., Ringsdorf, A., Warneke, C., Williams, J., Wisthaler, A., and Xu, L.: An improved representation of fire non-methane organic gases (NMOGs) in models: emissions to reactivity, *Atmospheric Chemistry and Physics*, 22, 12093–12111, <https://doi.org/10.5194/acp-22-12093-2022>, 2022.
- Chameides, W. L., Fehsenfeld, F., Rodgers, M. O., Cardelino, C., Martinez, J., Parrish, D., Lonneman, W., Lawson, D. R., Rasmussen, R. A., Zimmerman, P., Greenberg, J., Middleton, P., and Wang, T.: Ozone precursor relationships in the ambient atmosphere, *Journal of Geophysical Research: Atmospheres*, 97, 6037–6055, <https://doi.org/10.1029/91JD03014>, 1992.
- De Gouw, J. A.: Budget of organic carbon in a polluted atmosphere: Results from the New England Air Quality Study in 2002, *J. Geophys. Res.*, 110, D16305, <https://doi.org/10.1029/2004JD005623>, 2005.
- DeBell, L. J., Talbot, R. W., Dibb, J. E., Munger, J. W., Fischer, E. V., and Frohking, S. E.: A major regional air pollution event in the northeastern United States caused by extensive forest fires in Quebec, Canada, *Journal of Geophysical Research: Atmospheres*, 109, <https://doi.org/10.1029/2004JD004840>, 2004.
- Decker, Z. C. J., Wang, S., Bourgeois, I., Campuzano Jost, P., Coggon, M. M., DiGangi, J. P., Diskin, G. S., Flocke, F. M., Franchin, A., Fredrickson, C. D., Gkatzelis, G. I., Hall, S. R., Halliday, H., Hayden, K., Holmes, C. D., Huey, L. G., Jimenez, J. L., Lee, Y. R., Lindaas, J., Middlebrook, A. M., Montzka, D. D., Neuman, J. A., Nowak, J. B., Pagonis, D., Palm, B. B., Peischl, J., Piel, F., Rickly, P. S., Robinson, M. A., Rollins, A. W., Ryerson, T. B., Sekimoto, K., Thornton, J. A., Tyndall, G. S., Ullmann, K., Veres, P. R., Warneke, C., Washenfelder, R. A., Weinheimer, A. J., Wisthaler, A., Womack, C., and Brown, S. S.: Novel Analysis to Quantify Plume Crosswind Heterogeneity Applied to Biomass Burning Smoke, *Environ. Sci. Technol.*, 55, 15646–15657, <https://doi.org/10.1021/acs.est.1c03803>, 2021.
- Eastham, S. D. and Jacob, D. J.: Limits on the ability of global Eulerian models to resolve intercontinental transport of chemical plumes, *Atmospheric Chemistry and Physics*, 17, 2543–2553, <https://doi.org/10.5194/acp-17-2543-2017>, 2017.
- Fiore, A. M., Mickley, L. J., Zhu, Q., and Baublitz, C. B.: Climate and Tropospheric Oxidizing Capacity, *Annual Review of Earth and Planetary Sciences*, 52, null, <https://doi.org/10.1146/annurev-earth-032320-090307>, 2024.
- Gkatzelis, G. I., Coggon, M. M., Stockwell, C. E., Hornbrook, R. S., Allen, H., Apel, E. C., Bela, M. M., Blake, D. R., Bourgeois, I., Brown, S. S., Campuzano-Jost, P., St. Clair, J. M., Crawford, J. H., Crounse, J. D., Day, D. A., DiGangi, J. P., Diskin, G. S., Fried, A., Gilman, J. B., Guo, H., Hair, J. W., Halliday, H. S., Hanisco, T. F., Hannun, R., Hills, A., Huey, L. G., Jimenez, J. L., Katich, J. M., Lamplugh, A., Lee, Y. R., Liao, J., Lindaas, J., McKeen, S. A., Mikoviny, T., Nault, B. A.,

- Neuman, J. A., Nowak, J. B., Pagonis, D., Peischl, J., Perring, A. E., Piel, F., Rickly, P. S., Robinson, M. A., Rollins, A. W., Ryerson, T. B., Schueneman, M. K., Schwantes, R. H., Schwarz, J. P., Sekimoto, K., Selimovic, V., Shingler, T., Tanner, D. J., Tomsche, L., Vasquez, K. T., Veres, P. R., Washenfelder, R., Weibring, P., Wennberg, P. O., Wisthaler, A., Wolfe, G. M., Womack, C. C., Xu, L., Ball, K., Yokelson, R. J., and Warneke, C.: Parameterizations of US wildfire and prescribed fire emission ratios and emission factors based on FIREX-AQ aircraft measurements, *Atmospheric Chemistry and Physics*, 24, 929–956, <https://doi.org/10.5194/acp-24-929-2024>, 2024.
- Guenther, A. B., Jiang, X., Heald, C. L., Sakulyanontvittaya, T., Duhl, T., Emmons, L. K., and Wang, X.: The Model of Emissions of Gases and Aerosols from Nature version 2.1 (MEGAN2.1): an extended and updated framework for modeling biogenic emissions, *Geoscientific Model Development*, 5, 1471–1492, <https://doi.org/10.5194/gmd-5-1471-2012>, 2012.
- Guo, J. J., Fiore, A. M., Murray, L. T., Jaffe, D. A., Schnell, J. L., Moore, C. T., and Milly, G. P.: Average versus high surface ozone levels over the continental USA: model bias, background influences, and interannual variability, *Atmospheric Chemistry and Physics*, 18, 12123–12140, <https://doi.org/10.5194/acp-18-12123-2018>, 2018.
- Heald, C. L., Gouw, J. de, Goldstein, A. H., Guenther, A. B., Hayes, P. L., Hu, W., Isaacman-VanWertz, G., Jimenez, J. L., Keutsch, F. N., Koss, A. R., Misztal, P. K., Rappenglück, B., Roberts, J. M., Stevens, P. S., Washenfelder, R. A., Warneke, C., and Young, C. J.: Contrasting Reactive Organic Carbon Observations in the Southeast United States (SOAS) and Southern California (CalNex), *Environ. Sci. Technol.*, 54, 14923–14935, <https://doi.org/10.1021/acs.est.0c05027>, 2020.
- Hoesly, R. M., Smith, S. J., Feng, L., Klimont, Z., Janssens-Maenhout, G., Pitkanen, T., Seibert, J. J., Vu, L., Andres, R. J., Bolt, R. M., Bond, T. C., Dawidowski, L., Kholod, N., Kurokawa, J., Li, M., Liu, L., Lu, Z., Moura, M. C. P., O'Rourke, P. R., and Zhang, Q.: Historical (1750–2014) anthropogenic emissions of reactive gases and aerosols from the Community Emissions Data System (CEDS), *Geoscientific Model Development*, 11, 369–408, <https://doi.org/10.5194/gmd-11-369-2018>, 2018.
- Jacob, D. J.: Heterogeneous chemistry and tropospheric ozone, *Atmospheric Environment*, 34, 2131–2159, [https://doi.org/10.1016/S1352-2310\(99\)00462-8](https://doi.org/10.1016/S1352-2310(99)00462-8), 2000.
- Jaffe, D. A. and Wigder, N. L.: Ozone production from wildfires: A critical review, *Atmospheric Environment*, 51, 1–10, <https://doi.org/10.1016/j.atmosenv.2011.11.063>, 2012.
- Jin, L., Permar, W., Selimovic, V., Ketcherside, D., Yokelson, R. J., Hornbrook, R. S., Apel, E. C., Ku, I.-T., Collett Jr., J. L., Sullivan, A. P., Jaffe, D. A., Pierce, J. R., Fried, A., Coggon, M. M., Gkatzelis, G. I., Warneke, C., Fischer, E. V., and Hu, L.: Constraining emissions of volatile organic compounds from western US wildfires with WE-CAN and FIREX-AQ airborne observations, *Atmospheric Chemistry and Physics*, 23, 5969–5991, <https://doi.org/10.5194/acp-23-5969-2023>, 2023a.
- Jin, X., Fiore, A. M., and Cohen, R. C.: Space-Based Observations of Ozone Precursors within California Wildfire Plumes and the Impacts on Ozone-NO_x-VOC Chemistry, *Environ. Sci. Technol.*, 57, 14648–14660, <https://doi.org/10.1021/acs.est.3c04411>, 2023b.
- Juncosa Calahorrano, J. F., Lindaas, J., O'Dell, K., Palm, B. B., Peng, Q., Flocke, F., Pollack, I. B., Garofalo, L. A., Farmer, D. K., Pierce, J. R., Collett, J. L., Weinheimer, A., Campos, T., Hornbrook, R. S., Hall, S. R., Ullmann, K., Pothier, M. A., Apel, E. C., Permar, W., Hu, L., Hills, A. J., Montzka, D., Tyndall, G., Thornton, J. A., and Fischer, E. V.: Daytime Oxidized Reactive Nitrogen Partitioning in Western U.S. Wildfire Smoke Plumes, n.d.
- Kirchner, F., Jeanneret, F., Clappier, A., Krüger, B., van den Bergh, H., and Calpini, B.: Total VOC reactivity in the planetary boundary layer: 2. A new indicator for determining the sensitivity of the ozone production to VOC and NO_x, *Journal of Geophysical Research: Atmospheres*, 106, 3095–3110, <https://doi.org/10.1029/2000JD900603>, 2001.

- Kleinman, L. I.: Low and high NO_x tropospheric photochemistry, *Journal of Geophysical Research: Atmospheres*, 99, 16831–16838, <https://doi.org/10.1029/94JD01028>, 1994.
- Laughner, J. L. and Cohen, R. C.: Direct observation of changing NO_x lifetime in North American cities, *Science*, 366, 723–727, <https://doi.org/10.1126/science.aax6832>, 2019.
- 725 Lee, H. and Jaffe, D. A.: Impact of wildfire smoke on ozone concentrations using a Generalized Additive model in Salt Lake City, Utah, USA, 2006–2022, *Journal of the Air & Waste Management Association*, 74, 116–130, <https://doi.org/10.1080/10962247.2023.2291197>, 2024.
- Liao, J., Wolfe, G. M., Hannun, R. A., St. Clair, J. M., Hanisco, T. F., Gilman, J. B., Lamplugh, A., Selimovic, V., Diskin, G. S., Nowak, J. B., Halliday, H. S., DiGangi, J. P., Hall, S. R., Ullmann, K., Holmes, C. D., Fite, C. H., Agastra, A., Ryerson, T. B., Peischl, J., Bourgeois, I., Warneke, C., Coggon, M. M., Gkatzelis, G. I., Sekimoto, K., Fried, A., Richter, D., Weibring, P.,
730 Apel, E. C., Hornbrook, R. S., Brown, S. S., Womack, C. C., Robinson, M. A., Washenfelder, R. A., Veres, P. R., and Neuman, J. A.: Formaldehyde evolution in US wildfire plumes during the Fire Influence on Regional to Global Environments and Air Quality experiment (FIREX-AQ), *Atmospheric Chemistry and Physics*, 21, 18319–18331, <https://doi.org/10.5194/acp-21-18319-2021>, 2021.
- 735 Lindaas, J., Pollack, I. B., Garofalo, L. A., Pothier, M. A., Farmer, D. K., Kreidenweis, S. M., Campos, T. L., Flocke, F., Weinheimer, A. J., Montzka, D. D., Tyndall, G. S., Palm, B. B., Peng, Q., Thornton, J. A., Permar, W., Wielgasz, C., Hu, L., Ottmar, R. D., Restaino, J. C., Hudak, A. T., Ku, I.-T., Zhou, Y., Sive, B. C., Sullivan, A., Collett Jr, J. L., and Fischer, E. V.: Emissions of Reactive Nitrogen From Western U.S. Wildfires During Summer 2018, *Journal of Geophysical Research: Atmospheres*, 126, e2020JD032657, <https://doi.org/10.1029/2020JD032657>, 2021.
- 740 Loman, T. E., Ma, Y., Ilin, V., Gowda, S., Korsbo, N., Yewale, N., Rackauckas, C., and Isaacson, S. A.: Catalyst: Fast and flexible modeling of reaction networks, *PLOS Computational Biology*, 19, e1011530, <https://doi.org/10.1371/journal.pcbi.1011530>, 2023.
- McClure, C. D. and Jaffe, D. A.: Investigation of high ozone events due to wildfire smoke in an urban area, *Atmospheric Environment*, 194, 146–157, <https://doi.org/10.1016/j.atmosenv.2018.09.021>, 2018.
- 745 Müller, M., Anderson, B. E., Beyersdorf, A. J., Crawford, J. H., Diskin, G. S., Eichler, P., Fried, A., Keutsch, F. N., Mikoviny, T., Thornhill, K. L., Walega, J. G., Weinheimer, A. J., Yang, M., Yokelson, R. J., and Wisthaler, A.: In situ measurements and modeling of reactive trace gases in a small biomass burning plume, *Atmospheric Chemistry and Physics*, 16, 3813–3824, <https://doi.org/10.5194/acp-16-3813-2016>, 2016.
- 750 Park, R. J., Jacob, D. J., Field, B. D., Yantosca, R. M., and Chin, M.: Natural and transboundary pollution influences on sulfate-nitrate-ammonium aerosols in the United States: Implications for policy, *Journal of Geophysical Research: Atmospheres*, 109, <https://doi.org/10.1029/2003JD004473>, 2004.
- Peng, Q., Palm, B. B., Melander, K. E., Lee, B. H., Hall, S. R., Ullmann, K., Campos, T., Weinheimer, A. J., Apel, E. C., Hornbrook, R. S., Hills, A. J., Montzka, D. D., Flocke, F., Hu, L., Permar, W., Wielgasz, C., Lindaas, J., Pollack, I. B., Fischer, E. V., Bertram, T. H., and Thornton, J. A.: HONO Emissions from Western U.S. Wildfires Provide Dominant Radical Source
755 in Fresh Wildfire Smoke, *Environ. Sci. Technol.*, 54, 5954–5963, <https://doi.org/10.1021/acs.est.0c00126>, 2020.
- Permar, W., Wang, Q., Selimovic, V., Wielgasz, C., Yokelson, R. J., Hornbrook, R. S., Hills, A. J., Apel, E. C., Ku, I.-T., Zhou, Y., Sive, B. C., Sullivan, A. P., Collett Jr, J. L., Campos, T. L., Palm, B. B., Peng, Q., Thornton, J. A., Garofalo, L. A., Farmer, D. K., Kreidenweis, S. M., Levin, E. J. T., DeMott, P. J., Flocke, F., Fischer, E. V., and Hu, L.: Emissions of Trace Organic Gases From Western U.S. Wildfires Based on WE-CAN Aircraft Measurements, *Journal of Geophysical Research: Atmospheres*, 126, e2020JD033838, <https://doi.org/10.1029/2020JD033838>, 2021.
- 760

- Permar, W., Jin, L., Peng, Q., O'Dell, K., Lill, E., Selimovic, V., J. Yokelson, R., S. Hornbrook, R., J. Hills, A., C. Apel, E., Ku, I.-T., Zhou, Y., C. Sive, B., P. Sullivan, A., L. Collett, J., B. Palm, B., A. Thornton, J., Flocke, F., V. Fischer, E., and Hu, L.: Atmospheric OH reactivity in the western United States determined from comprehensive gas-phase measurements during WE-CAN, *Environmental Science: Atmospheres*, 3, 97–114, <https://doi.org/10.1039/D2EA00063F>, 2023.
- 765 Pfister, G. G., Wiedinmyer, C., and Emmons, L. K.: Impacts of the fall 2007 California wildfires on surface ozone: Integrating local observations with global model simulations, *Geophysical Research Letters*, 35, <https://doi.org/10.1029/2008GL034747>, 2008.
- Rastigejev, Y., Park, R., Brenner, M. P., and Jacob, D. J.: Resolving intercontinental pollution plumes in global models of atmospheric transport, *Journal of Geophysical Research: Atmospheres*, 115, <https://doi.org/10.1029/2009JD012568>, 2010.
- 770 Rickly, P. S., Coggon, M. M., Aikin, K. C., Alvarez, R. J. I., Baidar, S., Gilman, J. B., Gkatzelis, G. I., Harkins, C., He, J., Lamplugh, A., Langford, A. O., McDonald, B. C., Peischl, J., Robinson, M. A., Rollins, A. W., Schwantes, R. H., Senff, C. J., Warneke, C., and Brown, S. S.: Influence of Wildfire on Urban Ozone: An Observationally Constrained Box Modeling Study at a Site in the Colorado Front Range, *Environ. Sci. Technol.*, 57, 1257–1267, <https://doi.org/10.1021/acs.est.2c06157>, 2023.
- 775 Robinson, M. A., Decker, Z. C. J., Barsanti, K. C., Coggon, M. M., Flocke, F. M., Franchin, A., Fredrickson, C. D., Gilman, J. B., Gkatzelis, G. I., Holmes, C. D., Lamplugh, A., Lavi, A., Middlebrook, A. M., Montzka, D. M., Palm, B. B., Peischl, J., Pierce, B., Schwantes, R. H., Sekimoto, K., Selimovic, V., Tyndall, G. S., Thornton, J. A., Van Rooy, P., Warneke, C., Weinheimer, A. J., and Brown, S. S.: Variability and Time of Day Dependence of Ozone Photochemistry in Western Wildfire Plumes, *Environ. Sci. Technol.*, 55, 10280–10290, <https://doi.org/10.1021/acs.est.1c01963>, 2021.
- 780 Schneider, S. R., Shi, B., and Abbatt, J. P. D.: The Measured Impact of Wildfires on Ozone in Western Canada From 2001 to 2019, *Journal of Geophysical Research: Atmospheres*, 129, e2023JD038866, <https://doi.org/10.1029/2023JD038866>, 2024.
- Schroeder, J. R., Crawford, J. H., Fried, A., Walega, J., Weinheimer, A., Wisthaler, A., Müller, M., Mikoviny, T., Chen, G., Shook, M., Blake, D. R., and Tonnesen, G. S.: New insights into the column CH₂O/NO₂ ratio as an indicator of near-surface ozone sensitivity, *Journal of Geophysical Research: Atmospheres*, 122, 8885–8907, <https://doi.org/10.1002/2017JD026781>, 2017.
- 785 Sillman, S., Logan, J. A., and Wofsy, S. C.: The sensitivity of ozone to nitrogen oxides and hydrocarbons in regional ozone episodes, *Journal of Geophysical Research: Atmospheres*, 95, 1837–1851, <https://doi.org/10.1029/JD095iD02p01837>, 1990.
- 790 Singh, H. B., Anderson, B. E., Brune, W. H., Cai, C., Cohen, R. C., Crawford, J. H., Cubison, M. J., Czech, E. P., Emmons, L., Fuelberg, H. E., Huey, G., Jacob, D. J., Jimenez, J. L., Kaduwela, A., Kondo, Y., Mao, J., Olson, J. R., Sachse, G. W., Vay, S. A., Weinheimer, A., Wennberg, P. O., and Wisthaler, A.: Pollution influences on atmospheric composition and chemistry at high northern latitudes: Boreal and California forest fire emissions, *Atmospheric Environment*, 44, 4553–4564, <https://doi.org/10.1016/j.atmosenv.2010.08.026>, 2010.
- Singh, H. B., Cai, C., Kaduwela, A., Weinheimer, A., and Wisthaler, A.: Interactions of fire emissions and urban pollution over California: Ozone formation and air quality simulations, *Atmospheric Environment*, 56, 45–51, <https://doi.org/10.1016/j.atmosenv.2012.03.046>, 2012.
- 795 Stein, A. F., Draxler, R. R., Rolph, G. D., Stunder, B. J. B., Cohen, M. D., and Ngan, F.: NOAA's HYSPLIT Atmospheric Transport and Dispersion Modeling System, *Bulletin of the American Meteorological Society*, 96, 2059–2077, <https://doi.org/10.1175/BAMS-D-14-00110.1>, 2015.
- Tarasick, D., Galbally, I. E., Cooper, O. R., Schultz, M. G., Ancellet, G., Leblanc, T., Wallington, T. J., Ziemke, J., Liu, X., Steinbacher, M., Stachelin, J., Vigouroux, C., Hannigan, J. W., García, O., Foret, G., Zanis, P., Weatherhead, E.,

- 800 Petropavlovskikh, I., Worden, H., Osman, M., Liu, J., Chang, K.-L., Gaudel, A., Lin, M., Granados-Muñoz, M., Thompson, A. M., Oltmans, S. J., Cuesta, J., Dufour, G., Thouret, V., Hassler, B., Trickl, T., and Neu, J. L.: Tropospheric Ozone Assessment Report: Tropospheric ozone from 1877 to 2016, observed levels, trends and uncertainties, *Elementa: Science of the Anthropocene*, 7, 39, <https://doi.org/10.1525/elementa.376>, 2019.
- Vinken, G. C. M., Boersma, K. F., Jacob, D. J., and Meijer, E. W.: Accounting for non-linear chemistry of ship plumes in the
805 GEOS-Chem global chemistry transport model, *Atmospheric Chemistry and Physics*, 11, 11707–11722, <https://doi.org/10.5194/acp-11-11707-2011>, 2011.
- Wang, H., Lu, X., Jacob, D. J., Cooper, O. R., Chang, K.-L., Li, K., Gao, M., Liu, Y., Sheng, B., Wu, K., Wu, T., Zhang, J., Sauvage, B., Nédélec, P., Blot, R., and Fan, S.: Global tropospheric ozone trends, attributions, and radiative impacts in 1995–
810 2017: an integrated analysis using aircraft (IAGOS) observations, ozonesonde, and multi-decadal chemical model simulations, *Atmospheric Chemistry and Physics*, 22, 13753–13782, <https://doi.org/10.5194/acp-22-13753-2022>, 2022.
- Wang, S., Coggon, M. M., Gkatzelis, G. I., Warneke, C., Bourgeois, I., Ryerson, T., Peischl, J., Veres, P. R., Neuman, J. A., Hair, J., Shingler, T., Fenn, M., Diskin, G., Huey, L. G., Lee, Y. R., Apel, E. C., Hornbrook, R. S., Hills, A. J., Hall, S. R., Ullmann, K., Bela, M. M., Trainer, M. K., Kumar, R., Orlando, J. J., Flocke, F. M., and Emmons, L. K.: Chemical Tomography in a Fresh Wildland Fire Plume: A Large Eddy Simulation (LES) Study, *Journal of Geophysical Research: Atmospheres*, 126, e2021JD035203, <https://doi.org/10.1029/2021JD035203>, 2021a.
815
- Wang, X., Jacob, D. J., Downs, W., Zhai, S., Zhu, L., Shah, V., Holmes, C. D., Sherwen, T., Alexander, B., Evans, M. J., Eastham, S. D., Neuman, J. A., Veres, P. R., Koenig, T. K., Volkamer, R., Huey, L. G., Bannan, T. J., Percival, C. J., Lee, B. H., and Thornton, J. A.: Global tropospheric halogen (Cl, Br, I) chemistry and its impact on oxidants, *Atmospheric Chemistry and Physics*, 21, 13973–13996, <https://doi.org/10.5194/acp-21-13973-2021>, 2021b.
- 820 van der Werf, G. R., Randerson, J. T., Giglio, L., Collatz, G. J., Mu, M., Kasibhatla, P. S., Morton, D. C., DeFries, R. S., Jin, Y., and van Leeuwen, T. T.: Global fire emissions and the contribution of deforestation, savanna, forest, agricultural, and peat fires (1997–2009), *Atmos. Chem. Phys.*, 10, 11707–11735, <https://doi.org/10.5194/acp-10-11707-2010>, 2010.
- Westerling, A. L.: Increasing western US forest wildfire activity: sensitivity to changes in the timing of spring, *Philosophical Transactions of the Royal Society B: Biological Sciences*, 371, 20150178, <https://doi.org/10.1098/rstb.2015.0178>, 2016.
- 825 Williams, A. P., Abatzoglou, J. T., Gershunov, A., Guzman-Morales, J., Bishop, D. A., Balch, J. K., and Lettenmaier, D. P.: Observed Impacts of Anthropogenic Climate Change on Wildfire in California, *Earth's Future*, 7, 892–910, <https://doi.org/10.1029/2019EF001210>, 2019.
- Wolfe, G. M., Hanisco, T. F., Arkinson, H. L., Blake, D. R., Wisthaler, A., Mikoviny, T., Ryerson, T. B., Pollack, I., Peischl, J., Wennberg, P. O., Crounse, J. D., St. Clair, J. M., Teng, A., Huey, L. G., Liu, X., Fried, A., Weibring, P., Richter, D.,
830 Walega, J., Hall, S. R., Ullmann, K., Jimenez, J. L., Campuzano-Jost, P., Bui, T. P., Diskin, G., Podolske, J. R., Sachse, G., and Cohen, R. C.: Photochemical evolution of the 2013 California Rim Fire: synergistic impacts of reactive hydrocarbons and enhanced oxidants, *Atmospheric Chemistry and Physics*, 22, 4253–4275, <https://doi.org/10.5194/acp-22-4253-2022>, 2022.
- Xu, L., Crounse, J. D., Vasquez, K. T., Allen, H., Wennberg, P. O., Bourgeois, I., Brown, S. S., Campuzano-Jost, P., Coggon, M. M., Crawford, J. H., DiGangi, J. P., Diskin, G. S., Fried, A., Gargulinski, E. M., Gilman, J. B., Gkatzelis, G. I., Guo, H., Hair, J. W., Hall, S. R., Halliday, H. A., Hanisco, T. F., Hannun, R. A., Holmes, C. D., Huey, L. G., Jimenez, J. L., Lamplugh, A., Lee, Y. R., Liao, J., Lindaas, J., Neuman, J. A., Nowak, J. B., Peischl, J., Peterson, D. A., Piel, F., Richter, D., Rickly, P. S., Robinson, M. A., Rollins, A. W., Ryerson, T. B., Sekimoto, K., Selimovic, V., Shingler, T., Soja, A. J., St. Clair, J. M., Tanner, D. J., Ullmann, K., Veres, P. R., Walega, J., Warneke, C., Washenfelder, R. A., Weibring, P., Wisthaler, A., Wolfe, G. M., Womack, C. C., and Yokelson, R. J.: Ozone chemistry in western U.S. wildfire plumes, *Science Advances*, 7, eabl3648,
840 <https://doi.org/10.1126/sciadv.abl3648>, 2021.

Yokelson, R. J., Andreae, M. O., and Akagi, S. K.: Pitfalls with the use of enhancement ratios or normalized excess mixing ratios measured in plumes to characterize pollution sources and aging, *Atmospheric Measurement Techniques*, 6, 2155–2158, <https://doi.org/10.5194/amt-6-2155-2013>, 2013.