



Indirect climate impacts of the Hunga eruption

Ewa M Bednarz^{1,2}, Amy H. Butler², Xinyue Wang³, Zhihong Zhuo⁴, Wandi Yu⁵, Georgiy Stenchikov⁶, Matthew Toohey⁷, Yunqian Zhu^{1,2}

5

1. Cooperative Institute for Research in Environmental Sciences (CIRES), University of Colorado Boulder, Boulder, CO, USA.
2. NOAA Chemical Sciences Laboratory (NOAA CSL), Boulder, CO, USA.
3. Department of Atmospheric and Oceanic Sciences, University of Colorado Boulder, Boulder, USA
- 10 4. Department of Earth and Atmospheric Sciences, University of Quebec in Montreal, Montreal (Quebec), Canada.
5. Lawrence Livermore National Laboratory, CA, USA
6. Physical Science and Engineering Division, King Abdullah University of Science and Technology, Jeddah, Saudi Arabia
7. Institute of Space and Atmospheric Studies, University of Saskatchewan, Saskatoon, Canada.

Correspondence to: Ewa M. Bednarz (ewa.bednarz@noaa.gov)

15 **Abstract.** Injection of sulfur and water vapour by the Hunga volcanic eruption significantly altered chemical composition and radiative budget of the stratosphere. Yet, whether the eruption could also affect surface climate, especially via indirect pathways, remains poorly understood. Here we investigate these effects using large ensembles of simulations with the CESM2(WACCM6) Earth system model, incorporating interactive chemistry and aerosols in both coupled ocean and atmosphere-only configurations.

20

We find some statistically significant extratropical regional climate responses to the eruption driven by circulation changes; these are partially linked to the modulation of El Nino Southern Oscillation, and its associated teleconnections, and to perturbations of the stratospheric polar vortex in both hemispheres. The stratospheric anomalies affect surface climate through modulating the North Atlantic Oscillation in the Northern Hemisphere (up to three boreal winters following the 25 eruption) and the Southern Annular Mode in the Southern Hemisphere in late 2023. The latter is partly related to a concurrent reduction in Antarctic ozone, as increased stratospheric aerosols and water vapor reach the polar vortex.

Our study demonstrates that the eruption could have had a non-negligible influence on regional surface climate, and 30 discusses the mechanisms via which such an influence could occur. However, the results also highlight that this forcing is relatively weak compared to interannual variability, and is subject to model uncertainties in the representation of key processes. More research is thus needed before definitive statements on the role of the eruption in contributing to surface climate and weather events in the following years are made.



1. Introduction

The 2022 Hunga eruption erupted on January 15, 2022, injecting large amounts of water vapor and other volcanic materials into the stratosphere and upper atmosphere up to 57 km (Proud et al., 2022). Microwave Limb Sounder remote sensing measurements suggest that the eruption increased global stratospheric water vapor burden by approximately 10-15% (Khaykin et al., 2022), equivalent to ~ 150 Tg of water vapor (Millan et al., 2022), making it the largest water vapor perturbation in the satellite era. At the same time, the eruption injected a moderate amount of SO₂, ~ 0.5 -1.0 Tg of SO₂ (Carn et al., 2023; Sellitto et al., 2024), and likely other species, such as sea salt, chlorine (Zhu et al., 2023), and bromine species (Li et al., 2023). The SO₂ converted to sulfate aerosol and resulted in the largest stratospheric aerosol optical depth (sAOD) since the 1990 Mt. Pinatubo eruption (Taha et al., 2022).

The large volcanic water vapor and sulfur injections by the eruption were shown to significantly impact stratospheric temperatures and chemistry. Upper stratospheric cooling of up to a few degrees K were observed in 2022 and 2023 as the result of the radiative cooling by the Hunga water (Stocker, et al., 2024; Wang et al. 2023; Randel et al., 2024). The enhanced heterogeneous processing on sulfate aerosols in turn contributed to the strong stratospheric ozone depletion observed at Southern Hemisphere mid-latitudes in the austral winter of 2022 (Zhang et al., 2024). The state-of the art climate models participating in the Hunga Tonga-Hunga Ha'apai Volcano Impact Model Observation Comparison (HTHH-MOC) Project reproduce the stratospheric temperature and ozone responses as observed when the Hunga forcings are included (Zhuo et al., 2025).

Unlike the significant impacts in the stratosphere, the direct radiative impact of the eruption on the surface climate is likely small (e.g. Schoerberl et al. 2024; Gupta et al., 2025). However, whether the eruption could lead to regional surface climate changes via other, more indirect pathways remains not well understood. For instance, using WACCM4 climate model simulations, Jucker et al. (2024) reported the existence of robust regional surface climate responses to the stratospheric water injection that emerged 3-8 years following the eruption, albeit without a clear mechanistic understanding of the origin of such responses.

Here we address this using large 30-member ensembles of CESM2(WACCM6) earth system model simulations with interactive chemistry and aerosols forced with and without Hunga SO₂ and H₂O injections; the simulations are also carried out using either the atmosphere-only or coupled-ocean configuration to investigate the role of atmosphere-ocean coupling. Section 2 gives details on the model used, experimental protocol and Section 3 presents the resulting evolution of stratospheric aerosols and water vapour from the eruption. Section 4 discusses the simulated changes in near-surface air temperatures in the simulations. Sections 5 and 6 present changes in the Northern Hemisphere (NH) and Southern



65 Hemisphere (SH) stratospheric polar vortex and their links with the troposphere and the extratropical surface climate. Section 7 summarizes the main results.

2. Methods

2.1. CESM2(WACCM6) model

We use the Community Earth System Model version 2 coupled to the high-top Whole Atmosphere Community Climate Model version 6 (CESM2(WACCM6); Danabasoglu et al., 2020; Gettleman et al. 2019). The configuration includes an interactive Troposphere-Stratosphere-Mesosphere-Lower-Thermosphere (TSMLT) chemistry scheme (Davis et al., 2023), and the interactive Modal Aerosol Microphysics version 4 (MAM4, Lu et al., 2016). The horizontal resolution of the atmospheric model is 0.9° in latitude by 1.25° in longitude, with 70 vertical levels with a hybrid-pressure coordinate up to ~ 140 km. The ocean component is the Parallel Ocean Program version 2 (POP2; Danabasoglu et al., 2012; Smith et al., 75 2010), with 60 vertical levels, and the horizontal resolution is 1.125° in the zonal direction and between about 0.27° and 0.64° in the meridional direction.

2.2. Experimental description.

The simulations follow the ‘Experiment 1’ experimental protocol of the Hunga Tonga-Hunga Ha’apai impacts Model Observation Comparison project (HTHH-MOC) described in detail in Zhu et al. (2024). Briefly, the simulations cover the 10 80 years following the eruption - i.e. January 2022 to December 2031 - and follow the Coupled Model Intercomparison Project phase 6 (CMIP6) middle-of-the-road SSP2-4.5 emission scenario (Meinenhausen et al., 2020). All simulations are initialized from the same initial atmospheric and ocean state using the observed sea-surface temperatures following the procedure described in Richter et al. (2022). The simulations are run with a free-running meteorology, with the exception of the first 1- 2 months where the atmospheric meteorology is nudged to the Goddard Earth Observing System (GEOS) Modern-Era 85 Retrospective Analysis for Research and Applications, Version 2 (MERRA-2) reanalysis (Gelaro, et al., 2017) to ensure the eruption occurs during realistic background conditions.

For each experiment, there are two pairs of simulations: a ‘forced’ simulation with simultaneous injection of 0.5 Tg SO₂ and 90 150 Tg H₂O at 25-30 km on 15 January and a ‘control’ simulation without the Hunga injection. The first pair, denoted here as “HUNGA_fix”, is the core atmosphere-only simulation of the HTHH-MOC that uses prescribed climatological sea-surface temperatures and sea-ice (same for both control and perturbation simulation) derived from the NOAA high-resolution blended analysis (Banzon et al., 2022). The second pair, denoted here as “HUNGA_cpl”, is analogous but runs in a fully coupled mode with interactive ocean and sea-ice. In order to thoroughly account for the role of interannual variability, each simulation consists of 30 ensemble members - a significant improvement upon the original 10 required by the HTHH-



95 MOC - obtained by modifying the end date of the initial nudging in the increment of one day (with resulting end date between 27 January - 23 February 2022 depending on the member).

3 The simulated evolution of sulfate aerosols and water vapour

The resulting evolution of the anomalous (forced minus control) stratospheric Aerosol Optical Depth (sAOD) and stratospheric water vapour during the first 5 years simulated in the two pairs of experiments is shown in Fig. 1. In general, 100 the coupled ocean and atmosphere-only simulations give a similar evolution of aerosols and H₂O. In 2022, most of the anomalous aerosols and water vapour is confined to the tropics and SH mid-latitudes. While some aerosols reach also the SH high latitudes in the second part of 2022, most of the Antarctic sAOD and H₂O enhancement does not occur until 2023, when the largest Antarctic sAOD and H₂O anomalies are found. Similarly, the transport of Hunga water and aerosols to the NH 105 does not occur until early 2023. Unlike the aerosols, the anomalous H₂O is uplifted into the ascending branch of the Brewer Dobson Circulation and reaches the upper stratosphere, with peak values at the stratopause simulated in the second part of 2023, and persisting there until ~2027. The evolution of aerosols and water vapor simulated in CESM2(WACCM) is thus similar to that simulated by other models participating in the HTHH-MOC and inferred from available satellite data (Zhuo et al., 2025).




Figure 1. Evolution of the ensemble mean changes in zonal mean (a-b) sAOD, (c-d) H₂O at 25 hPa and (e-f) H₂O at 1 hPa between the forced simulation and the control. Left panels are for the coupled ocean simulations, and right panels for the atmosphere-only simulations. Stippling denotes regions where the response is statistically significant, here taken as larger than ± 2 standard errors in the difference in means.



115 4. Simulated changes in near-surface air temperatures following the eruption

Figures 2 and 3 show the simulated changes in annual mean surface temperatures for each of the five years following the eruption. The mean over the years 1-3 (2022-2024), i.e. when most of the direct radiative forcing from the eruption is present (e.g. Quaglia et al in prep), is also included. In agreement with the small radiative forcing estimated for the eruption (Schoeberl et al., 2024), the simulations do not show any significant cooling in the SH; this is unlike the small SH cooling 120 (~0.1 K) estimated from a radiative transfer and energy balance models in Gupta et al. (2025). However, the coupled ocean simulations (Fig. 2) show some significant, largely negative, temperature anomalies in the NH extratropics, especially over North America, Europe and central Asia, in years 1-3. These NH anomalies are unlikely to be due to direct radiative forcing 125 from aerosols or water vapor because most of the aerosol perturbation is found in the SH; rather, they are likely indicative of atmospheric circulation changes, as will be discussed in more detail in Section 4. In contrast, the atmosphere-only simulations (Fig. 3) show few significant near surface temperature changes. This occurs partly because the use of atmosphere-only set-up constrains and dampens any surface responses; in addition, the associated NH circulation responses 130 are much weaker, as will be discussed in Section 4.

In addition to the extratropical anomalies, the coupled ocean simulations show a significant cooling in the equatorial Pacific 130 in 2022 and 2023; the anomaly dominates the concurrent tropical temperature changes (Fig. 4a,c) and is evident also in the global mean (Fig. 4c). This Pacific response corresponds to a negative phase of the El Nino-Southern Oscillation (ENSO; Trenberth, 1997), or a La Nina-like response. Figure 4(e-f) shows the changes in the Equatorial Southern Oscillation Index (eqSOI), defined as the difference in 5°S-5°N sea-level pressure between 80°W-130°W and 90°E-140°E. Positive changes in eqSOI correspond to La Nina-like responses, and negative changes in eqSOI correspond to El Nino-like responses. 135 Comparison between panels 4e and 4c shows that changes in the eqSOI track and mirror changes in tropical and global mean near-surface temperatures, illustrating that the modulation of ENSO by the eruption is the dominant contributor to the large-scale temperature changes following the eruption.

The simulated La Nina-like response in 2022-2023 following the eruption is consistent with previous studies that found a 140 similar response to past SH volcanic eruptions (Pausata et al. 2020, 2023; Ward et al. 2021) and SH wildfire aerosols (Fasullo et al. 2023). These studies concluded that changes in the inter-tropical convergence zone (ITCZ) following a hemispherically asymmetrical aerosol forcing are the main driver of the La Nina-like response, and it is thus likely that the same mechanism operates here. Note that the ENSO response (and hence tropical temperature response) changes sign in year 145 3 and 4 (2024-2025), with a significant El Nino-like response in the Pacific, suggesting that the Hunga forcing might have longer-term impacts due to the much longer timescales governing ocean processes.



Shifts in the ENSO phase are associated with shifts in tropical convective heating that drive poleward-propagating planetary waves (Hoskins and Karoly, 1981), forming teleconnections that remotely affect regional climate patterns around the world (Mo and Livezey, 1986). In the observations, La Nina is associated with anomalous high pressure over the North Pacific and 150 southeastern US, low pressure over Canada, and a dipole in pressure over the North Atlantic that arises in part due to ENSO's stratospheric pathway (Butler et al. 2014, Domeisen et al. 2019). This circulation pattern generally means colder conditions over northwestern North America and subtropical Asia, and warmer conditions over the southeastern US and northern Eurasia during boreal winter. La Nina teleconnections to the SH can similarly lead to cooling over the Maritime continent, southern Africa, and northern South America. These patterns are qualitatively present in the coupled simulation 155 responses to the eruption, especially in 2023 (year 2) and to some extent in the year 1-3 average (Figure 2f), suggesting that these regional responses may arise in part from the shift towards a more La Nina-like teleconnection in the first two years. That these responses are largely not significant may be a reflection of either sampling error – previous studies have noted that some regional ENSO teleconnection responses are subject to large internal variability (Deser et al. 2017, 2018) - or an 160 inability of coupled models to capture ENSO teleconnections adequately (Williams et al. 2023, Fang et al. 2024). Such anomalies are not found in the atmosphere-only simulations, since by definition, ENSO (and its related teleconnections) is the same as in the control experiment (Fig. 3, Fig. 4b,d,f).

In addition to the ENSO changes in the coupled ocean simulations, which can drive remote extratropical responses in regional surface climate via teleconnections, the anomalous stratospheric aerosols and water vapor could drive changes in 165 stratospheric circulation that then influence extratropical surface climate. In the next two sections, we consider stratospheric polar vortex response to the Hunga forcing in the Northern (section 4) and Southern (section 5) hemisphere.




Figure 2. Yearly mean changes in near-surface air temperature between the forced simulation and the control in the coupled ocean simulations for each of the 5 years following the eruption (i.e. 2022-2027). The bottom right panel shows the response averaged over the first 3 years (i.e. 2022-2024). Stippling indicates statistical significance (defined as in Fig. 1).




Figure 3. As in Fig. 2 but for the changes in the atmosphere-only simulations.




Figure 4. Time series of changes in (a-b) zonal mean near-surface air temperature, (c-d) global and tropical mean near-surface air temperature, and (e-f) the eqSOI index between the forced simulation and the control in the coupled ocean (left) and atmosphere-only (right) simulations. Dashed lines in (c-f) indicate ensemble mean changes and solid lines their 5-month running means. Error bars in (c-f) denote confidence intervals of the ensemble mean changes (± 2 standard errors).



5. Seasonal changes in the NH polar vortex and impacts on the extratropical surface climate

This section examines the response of the NH stratospheric vortex to the eruption and the resulting impacts on the NH
180 extratropical surface climate. Figure 5 shows the evolution of anomalies in zonal winds at 65°N during the 5 years following the eruption, as well as the associated changes in geopotential heights over the Arctic polar cap (65°N-90°N). We find a statistically significant strengthening of the stratospheric vortex in the coupled ocean simulations in late 2022 (Fig. 5a); the response propagates down to the troposphere and is accompanied by a statistically significant reduction in geopotential heights over the Arctic (so colder, more compact air) at the same time (Fig. 5c). At the surface, the response manifests as the
185 pattern of changes in sea-level pressure projecting onto the positive phase of the Arctic Oscillation (AO) and, for the Atlantic sector, the North Atlantic Oscillation (NAO; Hurrell et al., 2003), i.e. decrease in sea-level pressure over the Arctic and increase over the NH mid-latitudes in early winter (November-December-January, NDJ; Fig. 6a). The response is also accompanied by some statistically significant near-surface air temperature changes, especially a cooling over Greenland (Fig. 6d). The stratospheric and surface responses change sign in late winter (February-March-April, FMA), with a
190 weakening (albeit not statistically significant in the zonal means) of the zonal winds (Fig. 5a) and a pattern of changes in sea-level pressure projection onto the negative phase of NAO (Fig. 7a). The response is also accompanied by statistically significant surface temperature changes, including a cooling over Europe and warming over Eastern Canada (Fig. 7d).

While the analogous early winter zonal wind changes in the second (i.e. 2023/2024) and third winter (2024/2025) are not
195 statistically significant (Fig. 5a), the pattern of changes in sea-level pressure and near-surface air temperatures also resemble those associated with a positive NAO, i.e. increase sea-level pressure in the NH mid-latitudes (Fig. 6b-c), a warming over northern Eurasia and a cooling over southern Europe and continental Asia (Fig. 6e-f). In the late period of the third winter, the coupled ocean simulations show a significant weakening of the stratospheric vortex (Fig. 5a) and increase in geopotential heights over the Arctic (Fig. 5c); the response propagates down to the surface and results in a negative AO/NAO pattern
200 (Fig. 7c) as well as increases in surface temperatures in southern Europe (Fig. 7f).

To summarise, while there is some variability between the individual winters, all three Arctic winters following the eruption simulated in the coupled ocean simulations suggest a strengthening of the stratospheric polar vortex and tropospheric polar jet stream in early winter, with the accompanied positive NAO-like anomalies in sea-level pressure and near-surface
205 temperatures (with the opposite sign stratospheric and surface climate responses found in late winters). The modulation of the polar vortex by the eruption could be driven by changes in meridional temperature gradients and wind shear associated with anomalous stratospheric radiative heating and cooling from Hunga aerosols and water vapour, and the resulting wave-mean flow feedbacks with planetary wave propagation and breaking. The increased stratospheric aerosols act to increase tropical lower stratospheric temperatures (Fig. S1a-b) and thus meridional temperature gradients at these altitudes; by
210 thermal wind relationship this can act to strengthen the polar vortex in the lower stratosphere. Evidence for this mechanism



has been found for previous volcanic eruptions (Polvani et al., 2019; Paik et al., 2023), although some studies point that this polar vortex signal only emerges for eruptions with large aerosol loading (Azoulay et al. 2021; DallaSanta and Polvani, 2022). However, a unique aspect of the Hunga eruption was the exceptional water vapor injection. The increased stratospheric water vapour results in cooling of the mid-to-upper stratosphere (Fig. S1c-f), with the upper stratospheric 215 cooling being amplified by the H₂O-induced ozone reduction at those levels and the resulting reduction in ozone shortwave heating (Randel et al. 2024). This acts to reduce meridional temperature gradients at these altitudes, thereby potentially weakening the upper stratospheric jet but strengthening the lower stratospheric jet. The enhancement of stratospheric H₂O also contributes to the warming in the tropical lower stratosphere below the water plume by trapping and re-radiating the outgoing terrestrial radiation, and thus amplifying the aerosol-induced heating there (e.g. Wang et al., 2023; Yook et al., 220 2025). Nonetheless, full details behind the relative interplay of changes in meridional temperature gradients and wind shear to Hunga water vapour and aerosols, and their coupling with wave feedbacks, on the polar vortex strength remains to be fully understood.

The simulated changes in the polar vortex are also likely to be at least partly related to the modulation of ENSO by the 225 eruption, and the associated changes in tropospheric wave flux to the stratosphere (e.g. Domeisen et al., 2019). In accord, the atmosphere-only simulations (which by definition have no ENSO response) show NH polar vortex changes that have different seasonality in winters 1 and 3 to the coupled ocean simulations, with statistically significant strengthening of the stratospheric vortex found mainly in late winter periods (Fig. 5b). In addition, the stratospheric anomalies do not appear to propagate readily to the troposphere, and so any changes in sea-level pressure and surface temperature are much weaker and 230 mostly not statistically significant (Fig. 8; Fig. S4). This suggests that the modulation of ENSO and its teleconnections plays an important contribution to the NH polar vortex changes simulated after the eruption in the coupled ocean experiments. In addition, it is also likely that the use of atmosphere-only set up substantially constrains and dampens any surface responses, highlighting the need for caution when interpreting the inferred signatures, or their absence, of the Hunga eruption on 235 climate. Significant differences in the simulated NH stratospheric vortex responses between the coupled ocean and atmosphere-only simulations were also previously reported in the context of idealized high latitude volcanic eruptions (Guðlaugsdóttir et al., 2025).


The extratropical atmospheric circulation also exhibits substantial variability across timescales, and so we examine the role 240 of interannual variability and ensemble size for the detectability of the climate signals from the eruption (e.g. Maycock and Hitchcock, 2015; Bittner et al., 2016; Milinski et al., 2020 used in different contexts). Figures 9-10 show the range of possible early and late winter NH lower stratospheric wind and NAO changes that can be inferred for the first winter following the eruption if a smaller number of ensemble members than the full ensemble is used. In particular, for each reduced ensemble size, 2000 artificial ensemble means are created by randomly sampling the available 30 members with



replacement. The means, +/- 2 standard deviation confidence intervals as well as maximum and minimum values of the
245 resulting 2000 inferred responses are shown in each case.

Inspection of Figure 9 shows that if only 10 ensemble members are used, then in most cases the envelope of potential NH
high latitude “ensemble-mean” responses spans both positive and negative values. This highlights the challenge of isolating
250 the relatively weak Hunga impact from the much larger natural interannual variability of the Arctic winter stratosphere, and
underscores the need for caution when interpreting surface climate impacts of the eruption inferred both from observations
and/or reanalysis (which effectively constitute just one ensemble member) or from model simulations with insufficient
ensemble size. For instance, Kuchar et al (2025) used a 10-member ensemble of Hunga simulations with the SOCOLv4
model, and reported a late winter weakening of the NH polar vortex from the eruption, with the associated impacts on the
high latitude surface climate. While the results qualitatively agree with the CESM coupled experiment results here, our
255 results also show that 10 members is not enough to infer the footprint of the eruption on the NH winter vortex with any
confidence (assuming similar magnitude of both the response and natural variability). In addition, here the variability is
particularly large for the late winter vortex response (i.e. even larger than in early winter; compare left and right panels in
Fig. 9), consistent with added uncertainty in the vortex final warming date.

260 The confidence intervals around the mean response become narrower as more ensemble members are used. Nonetheless,
even if close to 30 members are used, the edge of the +/- 2 standard deviation envelope of the distribution tends to be close
to 0, and an individual, albeit an outlier, ensemble configuration usually exists (indicated by the max/min lines) that can
indicate an opposite response instead. This shows that while interannual variability is likely to have a first order influence on
the response inferred from an ensemble of smaller (e.g. 10 members) size, it can likely still have a non-negligible
265 contribution even if as much as 30 members are used. Given that the observational record of the real world response
constitutes effectively just one realization, this makes attribution of any regional climate response to the eruption virtually
impossible without using a large ensemble.



270 **Figure 5.** Evolution of the NH polar vortex. Shading: Time series of ensemble-mean changes in (a-b) zonal wind at 65°N [in units of m/s] and (c-d) geopotential height [in units of standard deviations] averaged over the polar cap (65°N-90°N) between the forced simulation and the control. The left panels are for the coupled ocean simulations and the right panels are for the atmosphere-only simulations. Stippling denotes statistical significance (defined as in Fig. 1). Contours indicate values in the control for reference.




Figure 6. Changes in the early NH winter (November-December-January) ensemble mean (top) sea-level pressure and (bottom) near-surface air temperature between the forced simulation and the control in the coupled ocean simulations for the first (i.e. 2022/2023), second (i.e. 2023/2024) and third (2024/2025) winter following the eruption (columns). Stippling denotes statistical significance (defined as in Fig. 1).




Figure 7. As in Fig. 6 but for the corresponding late NH winter (February-March-April) changes. See Fig. S4 in the Supplement
285 for the corresponding changes in the atmosphere-only simulations.




Figure 8. As in Fig. 6 but for the changes in the atmosphere-only simulations.




Figure 9. Detectability of the early winter (a,c) and late winter (b,d) NH stratospheric vortex (a,b) and NAO (c,d) response in the coupled ocean simulations in the first winter following the eruption (2022/2023). Black lines denote the mean response, and blue and red lines indicate the +/- 2 standard deviation and max/min ranges, respectively, of the possible responses obtained by randomly subsampling the ensemble with replacement to obtain 2000 artificial ensembles of each different ensemble size. See Fig. S5 for the corresponding changes in the atmosphere-only simulations.

6. Seasonal changes in the SH polar vortex and impacts on the extratropical surface climate


As in the NH, changes in the radiative heating and thus stratospheric temperatures as the result of enhanced aerosol and water vapor following the eruption could drive changes in the Antarctic stratospheric vortex and surface climate. In addition, changes in SH stratospheric winds are strongly correlated with changes in Antarctic lower stratospheric ozone via ozone-circulation feedbacks, and so any response in ozone can influence the polar vortex evolution (and vice versa). Figure 10 shows the evolution of changes in SH zonal winds at 60°S (a-b) and Antarctic geopotential heights (c-d) for 5 years following the eruption.



305

No robust change in the Antarctic polar vortex strength is found in the coupled ocean simulations in the first year following the eruption. Note, however, that our diagnostic of polar vortex strength does not capture any changes in jet position, for instance the equatorial shift of winter stratospheric jet discussed in Wang et al. (2023). While the atmosphere-only simulations show some statistically significant strengthening of the stratospheric winds at 60S in austral spring, the response 310 does not propagate down to the troposphere, which shows a negative Southern Annular Mode (SAM, Thompson and Wallace, 2000) like response in late austral spring and early summer, Fig. 11a, i.e. opposite of what is expected from a strengthened polar vortex. In the second year, however, both simulations suggest a strengthening of the Antarctic vortex in austral spring (i.e. late 2023). In the atmosphere-only simulations the response extends down to the troposphere, and at the 315 surface manifests in sea-level pressure as the positive phase of the SAM - i.e. reduced sea-level pressure over the Antarctic and increased pressure in the SH mid-latitudes (Fig. 11b), with associated cooling over the Antarctic continent and warming over the southernmost part of South America (Fig. 11e) over late austral spring and early summer (NDJ). Similar surface 320 signatures of a positive SAM like response have also been found in the context of impacts on Antarctic ozone depletion by man-made halogenated substances (e.g. Thomson et al., 2011; Keeble et al., 2014). In agreement, our simulations also show significant SH ozone reductions in the lower stratosphere, primarily in the mid-latitudes (as noted already in e.g. Wang et al., 325 2023) but also in the springtime Antarctic region (Fig. 12). These negative polar ozone anomalies are driven by the combination of enhanced heterogeneous halogen activation, both on sulfate aerosols in the lowermost stratosphere as well as on more ice clouds above it under enhanced water vapour (Bednarz et al. in prep), and are strongest in year 2 (i.e. 2023) due to the time it takes for Hunga aerosol and water vapour to reach the polar stratosphere (Section 2, Fig. 1). Since ozone 330 absorbs strongly the incoming solar radiation, the reduction in Antarctic ozone from the Hunga eruption cools the polar stratosphere (Fig. S1ab) and strengthens the polar vortex (Fig. 10a-b), and the stratospheric response then propagates down to the troposphere and affects surface climate.

Notably, in addition to having much stronger stratosphere-troposphere response, the atmosphere-only simulations show substantially larger variability of the Antarctic polar vortex - ozone response (as indicated by the significantly wider spread 335 of potential responses in Fig. 13). This could be because the ENSO changes in the coupled ocean simulations drive SH stratospheric teleconnection patterns that destructively interfere with the Hunga impacts. In particular, reanalysis and model data show that La Niña-like tropical Pacific anomalies are associated with a weaker SH springtime stratospheric polar vortex and its tropospheric extension (Stone et al. 2022), and this thus agrees with and can explain the much weaker Hunga-induced polar vortex strengthening in the coupled runs. In addition, while the strengthening of the polar vortex associated with the Antarctic ozone reduction in the coupled simulations is also evident in the third winter (i.e. late 2024), the response is not found in the atmosphere-only simulations (Fig. 10b). This may also be explained by the contribution of ENSO variability in the coupled runs, whereby the now El Niño-like response enhances the strengthening of the stratospheric vortex.



340

Figure 10. Evolution of the SH polar vortex. Shading: Timeseries of changes in (a-b) zonal wind at 60°S [in units of m/s] and (c-d) geopotential height [in units of standard deviations] averaged over the Antarctic polar cap (55°S-90°S) between the forced simulation and the control. The left panels are for the coupled ocean simulations and the right panels are for the atmosphere-only simulations. Stippling denotes regions where the response is statistically significant (defined as in Fig. 1).




Figure 11. Changes in late austral spring and early summer (November–December–January) (top) sea-level pressure and (bottom) near-surface air temperature between the forced simulation and the control in the atmosphere-only simulations for the first (i.e. 2022/2023), second (i.e. 2023/2024) and third (2024/2025) winter following the eruption (columns). Stippling denotes statistical significance (defined as in Fig. 1). See Fig. S6 in the Supplement for the corresponding changes in the coupled ocean simulations.

345

350





Figure 12. Timeseries of zonal mean ozone changes at 50 hPa between the forced simulation and the control for the coupled ocean (left) and atmosphere-only (right) simulations. Stippling denotes statistical significance (as in Fig. 1).



355

(a) ΔU (60S, 50hPa), NDJ winter 2, HUNGA_cpl



(b) ΔU (60S, 50hPa), NDJ winter 2, HUNGA_fix




Figure 13. Detectability of the changes in the 2023-2024 November-December-January SH lower stratospheric vortex in the coupled ocean (a) and atmosphere-only (b) simulations. Different lines as in Fig. 9.

7. Conclusions.

360 Observations and modelling studies of the Hunga eruption in January 2022 have linked the anomalous stratospheric aerosol and water vapor levels with statistically significant changes in stratospheric and mesospheric temperatures (Stocker, et al., 2024; Wang et al. 2023; Randel et al., 2024; Yu et al., 2023) and ozone (Zhang et al., 2024). In the troposphere, on the other hand, any direct radiatively-driven surface impacts of the eruption are likely to be small (Schoeberl et al., 2024; Quaglia, et al. in prep). However, whether the eruption could lead to surface climate changes via other, more indirect pathways remains

365 not well understood. Here we address this using large (30-member each) ensembles of CESM2(WACCM) earth system model simulations with interactive chemistry and aerosols forced both with and without SO₂ and H₂O injections; the simulations are also carried out using either the atmosphere-only or coupled-ocean configuration to investigate the role of atmosphere-ocean coupling.

370 We find some statistically significant extratropical circulation and regional climate responses to the Hunga eruption, particularly in the coupled ocean simulations. These arise because of the combination of the ENSO response to the eruption, and its teleconnections, in this model as well as to the associated polar vortex changes in both hemispheres. The modulation of ENSO manifests itself in the form of La Niña-like response in years 1-2 (2022-23), followed by an opposite, El Niño-like response in year 4 (2025). The La Niña-like response following the eruption is consistent with that previously inferred for

375 the past SH volcanic eruptions (Pausata et al. 2020, 2023; Ward et al. 2021) and SH wildfire aerosols (Fasullo et al. 2023), suggesting changes in the ITCZ following a hemispherically unsymmetrical forcing could be the main driver contributing to the ENSO response here. Our results also suggest that the eruption could have contributed to the anomalous persistence of



the La Niña-like conditions observed between 2021-2023 (e.g. Iwakiri et. al., 2023). The modulation of the ENSO variability following the eruption in the model also gives rise to a small global mean cooling around the same period, as well as a
380 number of regional surface temperature responses in the Pacific region that are generally consistent with those associated with anomalous Pacific SSTs and their teleconnections more generally (e.g. Domisen et al. 2019).

The simulations also show statistically significant stratospheric polar vortex responses in both hemispheres. These changes propagate down to the troposphere and affect surface climate. In the NH, these surface climate responses manifest
385 themselves as modulations of the North Atlantic Oscillation. The NH extratropical responses change sign between early and late winter, with generally stronger polar vortex and positive NAO-like response at the surface in early winter, and vice versa for late winter, and are simulated during three boreal winters following the eruption in the coupled run. The modulation of the polar vortex under the Hunga eruption could be driven by the changes in radiative heating and cooling under enhancement of stratospheric aerosols and water vapour (with increased stratospheric aerosols acting to increase tropical
390 lower stratospheric temperatures and meridional temperature gradients and hence acting to strengthen polar vortex, while increased H₂O acting to cool mid-to-upper stratosphere and reduce meridional T gradients at these altitudes, thereby acting to weaken polar vortex there). However, the simulated changes in the polar vortex, especially in the NH, are likely to arise not only because of radiative/temperature changes in the stratosphere due to the aerosols and water vapor, but also at least partly due to the modulation of ENSO and the associated changes in tropospheric wave flux to the stratosphere. In accord,
395 the atmosphere-only simulations show NH polar vortex changes that are significantly weaker and have different seasonality than those in the coupled ocean runs, and largely do not propagate down to the troposphere and affect surface climate. This suggests that the modulation of ENSO and its teleconnections plays an important contribution to the polar vortex changes simulated after the eruption.

400 For the SH response, the simulations show strengthening of the SH polar vortex in austral spring of 2023 that is likely at least partly related to the associated reduction in Antarctic ozone as increased stratospheric aerosols and water vapour reach the polar vortex. In the atmosphere-only case, the stratospheric response propagates down to the surface, leading to the pattern of changes in sea-level pressure projecting on the positive phase of the Southern Annular Mode, and some statistically significant near-surface temperature changes, including a small cooling over Antarctica. However, here the
405 influence of the ENSO modulation on the vortex in the coupled ocean runs likely acts deconstructively with the impacts driven directly in the stratosphere, and so the coupled ocean simulations show reduced variability of the polar vortex and its weaker response to the eruption than that found in the atmosphere-only simulations.

410 Finally, we examine the role of interannual variability and ensemble size for the detectability of the climate signals from the eruption. By randomly sub-sampling the ensemble we show the range of apparent surface climate responses can be inferred if using smaller ensemble sizes. We demonstrate that interannual variability is likely to have a first order influence on the



response inferred from an ensemble of smaller (e.g. 10 members) size, but that it can still have a non-negligible contribution even if as much as 30 members are used. The results thus highlight the need for caution when interpreting the surface climate impacts of the eruption inferred from insufficient ensemble size.

415

All in all, our study demonstrates that the Hunga eruption could have a non-negligible influence on regional surface climate, and discusses the mechanism via which such an influence could occur. However, the results also highlight that this forcing is relatively weak compared to interannual variability, is subject to model uncertainties in the representation of key processes (e.g. it is unclear whether the ENSO response is particular to the model sensitivities) as well as in some parts requires a large 420 number of ensemble members to confidently detect. This calls into question whether such an influence could be robustly detected in the single ‘realization’ of the real world. Still, studies show that current climate models tend to underestimate the signal-to-noise ratio and the predictable component of the forced response regionally (e.g. Scaife and Smith, 2018; Gillet et al., 2003; Williams et al. 2023), suggesting that the indirect climate response to the eruption seen in this study may not 425 adequately represent the real world response, and that more research is needed before definitive statements on the role of the eruption in contributing to the surface climate and weather events in the following years are made.

Acknowledgements

We would like to acknowledge high-performance computing support from Derecho and Casper, provided by the National Center for Atmospheric Research (NCAR) Computational and Information Systems Laboratory, sponsored by the National Science Foundation.

430 Support has been provided by the National Oceanic and Atmospheric Administration (NOAA) cooperative agreement NA22OAR4320151 and the NOAA Earth Radiative Budget (ERB) program. WY’s work is prepared by LLNL under Contract DE-AC52-07NA27344.

We thank Simone Tilmes for help setting up the simulations, and Mark Schoeberl for helpful comments on the study.

Authors Contributions

435 EMB conceptualized the study, analysed the data and wrote the manuscript, with the support from AHB. XW, EMB, ZZ and WY performed the simulations. XW and YZ designed the simulations. All authors contributed to discussion of the results and writing of the manuscript.

Conflict of Interest

EMB and MT are members of the editorial board of ACP.



440 **References:**

Azoulay, A., Schmidt, H., & Timmreck, C.: The Arctic polar vortex response to volcanic forcing of different strengths. Journal of Geophysical Research: Atmospheres, 126, e2020JD034450. <https://doi.org/10.1029/2020JD034450>, 2021.

Banzon, Viva &, Reynolds, Richard & National Center for Atmospheric Research Staff (Eds). Last modified 2022-09-09
445 "The Climate Data Guide: SST data: NOAA High-resolution (0.25x0.25) Blended Analysis of Daily SST and Ice, OISSTv2." Retrieved from <https://climatedataguide.ucar.edu/climate-data/sst-data-noaa-high-resolution-025x025-blended-analysis-daily-sst-and-ice-oisstv2> on 2025-04-04.

450 Bednarz et al.: "Diagnosing chemical and dynamical impacts of the Hunga Tonga eruption on stratospheric and tropospheric ozone", in prep.

Bittner, M., H. Schmidt, C. Timmreck, and F. Sienz: Using a large ensemble of simulations to assess the Northern Hemisphere stratospheric dynamical response to tropical volcanic eruptions and its uncertainty, Geophys. Res. Lett., 43, 9324–9332, doi:10.1002/2016GL070587, 2016.

455 Carn, S. A., et al. "Out of the blue: Volcanic SO₂ emissions during the 2021–2022 eruptions of Hunga Tonga—Hunga Ha'apai (Tonga)." Frontiers in Earth Science 10: 976962, 2022.

460 DallaSanta, K. and Polvani, L. M.: Volcanic stratospheric injections up to 160 Tg(S) yield a Eurasian winter warming indistinguishable from internal variability, Atmos. Chem. Phys., 22, 8843–8862, <https://doi.org/10.5194/acp-22-8843-2022>, 2022.

465 Danabasoglu, G., Lamarque, J.-F., Bacmeister, J., Bailey, D. A., DuVivier, A. K., Edwards, J., Emmons, L. K., Fasullo, J., Garcia, R., Gettelman, A., Hannay, C., Holland, M. M., Large, W. G., Lauritzen, P. H., Lawrence, D. M., Lenaerts, J. T. M., Lindsay, K., Lipscomb, W. H., Mills, M. J., Neale, R., Oleson, K. W., Otto-Bliesner, B., Phillips, A. S., Sacks, W., Tilmes, S., van Kampenhout, L., Vertenstein, M., Bertini, A., Dennis, J., Deser, C., Fischer, C., Fox-Kemper, B., Kay, J. E., Kinnison, D., Kushner, P. J., Larson, V. E., Long, M. C., Mickelson, S., Moore, J. K., Nienhouse, E., Polvani, L., Rasch, P. J., and Strand, W. G.: The Community Earth System Model Version 2 (CESM2), J. Adv. Model. Earth Sy., 12, e2019MS001916, <https://doi.org/10.1029/2019MS001916>, 2020.

470 Danabasoglu, G., Bates, S. C., Briegleb, B. P., Jayne, S. R., Jochum, M., Large, W. G., et al.: The CCSM4 ocean component. Journal of Climate, 25(5), 1361–1389. <https://doi.org/10.1175/JCLI-D-11-00091.1>, 2012.



475 Davis, N. A., Visioni, D., Garcia, R. R., Kinnison, D. E., Marsh, D. R., Mills, M. J., Richter, J. H., Tilmes, S., Bardeen, C.,
Gettelman, A., Glanville, A. A., MacMartin, D. G., Smith, A. K., and Vitt, F.: Climate, variability, and climate sensitivity of
“Middle Atmosphere” chemistry configurations of the Community Earth System Model Version 2, Whole Atmosphere
Community Climate Model Version 6 (CESM2(WACCM6)), *J. Adv. Model. Earth Sy.*, 15, e2022MS003579,
<https://doi.org/10.1029/2022MS003579>, 2023.

480 Deser, Clara, Isla R Simpson, Karen A McKinnon, and Adam. S Phillips. “The Northern Hemisphere Extra-Tropical
Atmospheric Circulation Response to ENSO: How Well Do We Know It and How Do We Evaluate Models Accordingly?”
Journal of Climate 30: 5059–82. <https://doi.org/10.1175/JCLI-D-16-0844.1>, 2017.

485 Deser, Clara, Isla R Simpson, Adam. S Phillips, and Karen A McKinnon. “How Well Do We Know ENSO’s Climate
Impacts over North America, and How Do We Evaluate Models Accordingly?” *Journal of Climate* 31: 4991–5014.
<https://doi.org/10.1175/JCLI-D-17-0783.1>, 2018.

490 Domeisen, D. I., Garfinkel, C. I., & Butler, A. H.: The teleconnection of El Niño Southern Oscillation to the stratosphere.
Reviews of Geophysics, 57, 5–47. <https://doi.org/10.1029/2018RG000596>, 2019.

495 Fang, Yingfei, James A. Screen, Xiaoming Hu, Shuheng Lin, Ned C. Williams, and Song Yang. “CMIP6 Models
Underestimate ENSO Teleconnections in the Southern Hemisphere.” *Geophysical Research Letters* 51, no. 18:
e2024GL110738. <https://doi.org/10.1029/2024GL110738>, 2024.

500 John T. Fasullo et al. ,A multiyear tropical Pacific cooling response to recent Australian wildfires in CESM2.Sci.
Adv.9,eadg1213.DOI:10.1126/sciadv.adg1213, 2023.

505 Gelaro, R., McCarty, W., Suárez, M.J., Todling, R., Molod, A., Takacs, L., Randles, C.A., Darmenov, A., Bosilovich, M.G.,
Reichle, R. and Wargan, K.: The modern-era retrospective analysis for research and applications, version 2 (MERRA-2).
Journal of climate, 30(14), pp.5419-5454, 2017.

Guðlaugsdóttir, H., Peings, Y., Zanchettin, D., and Magnusdottir, G.: Stratospheric circulation response to large Northern
Hemisphere high-latitude volcanic eruptions in a global climate model, *Atmos. Chem. Phys.*, 25, 3961–3980,
<https://doi.org/10.5194/acp-25-3961-2025>, 2025.



Gettelman, A., Mills, M. J., Kinnison, D. E., Garcia, R. R., Smith, A. K., Marsh, D. R., Tilmes, S., Vitt, F., Bardeen, C. G., McInerny, J., Liu, H.-L., Solomon, S. C., Polvani, L. M., Emmons, L. K., Lamarque, J.-F., Richter, J. H., Glanville, A. S., Bacmeister, J. T., Phillips, A. S., Neale, R. B., Simpson, I. R., DuVivier, A. K., Hodzic, A., and Randel, W. J.: The Whole Atmosphere Community Climate Model Version 6 (WACCM6), *J. Geophys. Res.-Atmos.*, 124, 12380–12403, 510 <https://doi.org/10.1029/2019JD030943>, 2019.

Gillett, N., Zwiers, F., Weaver, A. et al. Detection of human influence on sea-level pressure. *Nature* 422, 292–294, <https://doi.org/10.1038/nature01487>, 2003.

515 Gupta, A.K., Mittal, T., Fauria, K.E., Bennartz, R., Kok, J. F.: The January 2022 Hunga eruption cooled the southern hemisphere in 2022 and 2023. *Commun Earth Environ* 6, 240. <https://doi.org/10.1038/s43247-025-02181-9>, 2025.

Hoskins, B. J., and D. J. Karoly: The Steady Linear Response of a Spherical Atmosphere to Thermal and Orographic Forcing. *J. Atmos. Sci.*, 38, 1179–1196, [https://doi.org/10.1175/1520-0469\(1981\)038<1179:TSLROA>2.0.CO;2](https://doi.org/10.1175/1520-0469(1981)038<1179:TSLROA>2.0.CO;2), 1981.

520 Hurrell, J. W., Kushnir, Y., Ottersen, G. & Visbeck, M. An overview of the North Atlantic oscillation. *Geophys. Monogr. Ser.* 134, 1–36, 2003.

Iwakiri, T., Imada, Y., Takaya, Y., Kataoka, T., Tatebe, H., & Watanabe, M.: Triple-dip La Niña in 2020–23: North Pacific atmosphere drives 2nd year La Niña. *Geophysical Research Letters*, 50, e2023GL105763.

525 <https://doi.org/10.1029/2023GL105763>, 2023.

Jucker, M., C. Lucas, and D. Dutta: Long-Term Climate Impacts of Large Stratospheric Water Vapor Perturbations. *J. Climate*, 37, 4507–4521, <https://doi.org/10.1175/JCLI-D-23-0437.1>, 2024.

530 Keeble, J., Braesicke, P., Abraham, N. L., Roscoe, H. K., and Pyle, J. A.: The impact of polar stratospheric ozone loss on Southern Hemisphere stratospheric circulation and climate, *Atmos. Chem. Phys.*, 14, 13705–13717, <https://doi.org/10.5194/acp-14-13705-2014>, 2014.

Khaykin, S., Podglajen, A., Ploeger, F., Groß, J.-U., Tence, F., Bekki, S., Khlopenkov, K., Bedka, K., Rieger, L., Baron, A., 535 Godin-Beekmann, S., Legras, B., Sellitto, P., Sakai, T., Barnes, J., Uchino, O., Morino, I., Nagai, T., Wing, R., Baumgarten, G., Gerding, M., Duflot, V., Payen, G., Jumelet, J., Querel, R., Liley, B., Bourassa, A., Clouser, B., Feofilov, A., Hauchecorne, A., and Ravetta, F.: Global perturbation of stratospheric water and aerosol burden by Hunga eruption, *Commun. Earth Environ.*, 3, 316, <https://doi.org/10.1038/s43247-022-00652-x>, 2022.



540 Kuchar, A., Sukhodolov, T., Chiodo, G., Jörnemann, A., Kult-Herdin, J., Rozanov, E., and Rieder, H. H.: Modulation of the northern polar vortex by the Hunga Tonga–Hunga Ha'apai eruption and the associated surface response, *Atmos. Chem. Phys.*, 25, 3623–3634, <https://doi.org/10.5194/acp-25-3623-2025>, 2025.

545 Li, Q., Qian, Y., Luo, Y., Cao, L., Zhou, H., Yang, T., Si, F., & Liu, W.: Diffusion Height and Order of Sulfur Dioxide and Bromine Monoxide Plumes from the Hunga Tonga–Hunga Ha'apai Volcanic Eruption. *Remote Sensing*, 15(6), 1534. <https://doi.org/10.3390/rs15061534>, 2023.

550 Liu, X., Ma, P.-L., Wang, H., Tilmes, S., Singh, B., Easter, R. C., Ghan, S. J., and Rasch, P. J.: Description and evaluation of a new four-mode version of the Modal Aerosol Module (MAM4) within version 5.3 of the Community Atmosphere Model, *Geosci. Model Dev.*, 9, 505–522, <https://doi.org/10.5194/gmd-9-505-2016>, 2016.

555 Meinshausen, M., Nicholls, Z. R. J., Lewis, J., Gidden, M. J., Vogel, E., Freund, M., et al.: The shared socio-economic pathway (SSP) greenhouse gas concentrations and their extensions to 2,500. *Geoscientific Model Development*, 13, 3571–3605. <https://doi.org/10.5194/gmd-13-3571-2020>, 2020.

Milinski, S., Maher, N., and Olonscheck, D.: How large does a large ensemble need to be?, *Earth Syst. Dynam.*, 11, 885–901, <https://doi.org/10.5194/esd-11-885-2020>, 2020.

560 Millán, L., Santee, M. L., Lambert, A., Livesey, N. J., Werner, F., Schwartz, M. J., Pumphrey, H. C., Manney, G. L., Wang, Y., Su, H., Wu, L., Read, W. G., and Froidevaux, L.: The Hunga Tonga-Hunga Ha'apai Hydration of the Stratosphere, *Geophys. Res. Lett.*, 49, e2022GL099381, <https://doi.org/10.1029/2022GL099381>, 2022.

565 Mo, K. C., and R. E. Livezey: Tropical-Extratropical Geopotential Height Teleconnections during the Northern Hemisphere Winter. *Mon. Wea. Rev.*, 114, 2488–2515, [https://doi.org/10.1175/1520-0493\(1986\)114<2488:TEGHTD>2.0.CO;2](https://doi.org/10.1175/1520-0493(1986)114<2488:TEGHTD>2.0.CO;2), 1986.

570 Paik, S., Min, S.-K., Son, S.-W., Lim, E.-P., McGregor, S., An, S.-Il, Kug, J.-S., Yeh, S.-W.: Impact of volcanic eruptions on extratropical atmospheric circulations: review, revisit and future directions, *Environ. Res. Lett.*, 18, 063003, 2023.



Pausata, F. S. R., Zhao, Y., Zanchettin, D., Caballero, R., & Battisti, D. S.: Revisiting the mechanisms of ENSO response to tropical volcanic eruptions. *Geophysical Research Letters*, 50, e2022GL102183. <https://doi.org/10.1029/2022GL102183>, 2023.

575

Pausata, F. S. R., Zanchettin, D., Karamperidou, C., Caballero, R., and Battisti, D. S.: ITCZ shift and extratropical teleconnections drive ENSO response to volcanic eruptions, *Sci. Adv.*, 6, eaaz5006, <https://doi.org/10.1126/SCIAADV.AAZ5006>, 2020.

580 Polvani, L. M., Banerjee, A., and Schmidt, A.: Northern Hemisphere continental winter warming following the 1991 Mt. Pinatubo eruption: reconciling models and observations, *Atmos. Chem. Phys.*, 19, 6351–6366, <https://doi.org/10.5194/acp-19-6351-2019>, 2019.

585 Proud, Simon R., Andrew T. Prata, and Simeon Schmauß. "The January 2022 eruption of Hunga Tonga-Hunga Ha'apai volcano reached the mesosphere." *Science* 378.6619: 554-557, 2022.

Randel, W. J., Wang, X., Starr, J., Garcia, R. R., & Kinnison, D.: Long-term temperature impacts of the Hunga volcanic eruption in the stratosphere and above. *Geophysical Research Letters*, 51, e2024GL111500. <https://doi.org/10.1029/2024GL111500>, 2024.

590

Richter, J. H., Glanville, A. A., Edwards, J., Kauffman, B., Davis, N. A., Jaye, A., et al.: Subseasonal Earth system prediction with CESM2. *Weather and Forecasting*, 37(6), 797–815. <https://doi.org/10.1175/waf-d-21-0163.1>, 2022.

595 Scaife, A.A., Smith, D. A signal-to-noise paradox in climate science. *npj Clim Atmos Sci* 1, 28. <https://doi.org/10.1038/s41612-018-0038-4>, 2018.

Schoeberl, M. R., Wang, Y., Taha, G., Zawada, D. J., Ueyama, R., & Dessler, A.: Evolution of the climate forcing during the two years after the Hunga Tonga-Hunga Ha'apai eruption. *Journal of Geophysical Research: Atmospheres*, 129, e2024JD041296. <https://doi.org/10.1029/2024JD041296>, 2024.

600

Sellitto, P., Siddans, R., Belhadji, R., Carboni, E., Legras, B., Podglajen, A., et al.: Observing the SO₂ and sulfate aerosol plumes from the 2022 Hunga eruption with the Infrared Atmospheric Sounding Interferometer (IASI). *Geophysical Research Letters*, 51, e2023GL105565, 2024.



605 Smith, R., Jones, P., Briegleb, B., Bryan, F., Danabasoglu, G., Dennis, J., et al.: The parallel ocean program (POP) reference manual, ocean component of the community climate system model (CCSM). LANL Technical Report, LAUR-10-01853, 141., 2010.

610 Stocker, M., Steiner, A.K., Ladstädter, F. et al: Strong persistent cooling of the stratosphere after the Hunga eruption. Commun Earth Environ 5, 450 (2024). <https://doi.org/10.1038/s43247-024-01620-3>, 2024.

Stone, K. A., S. Solomon, D. W. J. Thompson, D. E. Kinnison, and J. C. Fyfe: On the Southern Hemisphere Stratospheric Response to ENSO and Its Impacts on Tropospheric Circulation. *J. Climate*, 35, 1963–1981, <https://doi.org/10.1175/JCLI-D-21-0250.1>, 2022.

615 Taha, G., Loughman, R., Colarco, P. R., Zhu, T., Thomason, L. W., & Jaross, G.: Tracking the 2022 Hunga Tonga-Hunga Ha'apai aerosol cloud in the upper and middle stratosphere using space-based observations. *Geophysical Research Letters*, 49, e2022GL100091. <https://doi.org/10.1029/2022GL100091>, 2022.

620 Thompson, D. W. J., Solomon, S., Kushner, P. J., England, M. H., Grise, K. M., and Karoly, D. J.: Signatures of the Antarctic ozone hole in Southern Hemisphere surface climate change, *Nat. Geosci.*, 4, 741–749, doi:10.1038/ngeo1296, 2011.

Thompson, D. W. J., & Wallace, J. M.: Annular modes in the extratropical circulation. Part I: Month-to-Month variability. 625 *Journal of Climate*, 13(5), 1000–1016. [https://doi.org/10.1175/1520-0442\(2000\)013<1000:amitec>2.0.co;2](https://doi.org/10.1175/1520-0442(2000)013<1000:amitec>2.0.co;2), 2000.

Trenberth, K. E.: The Definition of El Niño. *Bull. Amer. Meteor. Soc.*, 78, 2771–2778, [https://doi.org/10.1175/1520-0477\(1997\)078<2771:TDOENO>2.0.CO;2](https://doi.org/10.1175/1520-0477(1997)078<2771:TDOENO>2.0.CO;2), 1997.

630 Wang, X., Randel, W., Zhu, Y., Tilmes, S., Starr, J., Yu, W., et al.: Stratospheric climate anomalies and ozone loss caused by the Hunga Tonga-Hunga Ha'apai volcanic eruption. *Journal of Geophysical Research: Atmospheres*, 128(22), e2023JD039480. <https://doi.org/10.1029/2023JD039480>, 2023.

Ward, B., Pausata, F. S. R., and Maher, N.: The sensitivity of the El Niño–Southern Oscillation to volcanic aerosol spatial distribution in the MPI Grand Ensemble, *Earth Syst. Dynam.*, 12, 975–996, <https://doi.org/10.5194/esd-12-975-2021>, 2021.

635 Williams, N. C., Scaife, A. A., & Screen, J. A.: Underpredicted ENSO teleconnections in seasonal forecasts. *Geophysical Research Letters*, 50, e2022GL101689. <https://doi.org/10.1029/2022GL101689>, 2023.



Yook, S., Solomon, S., & Wang, X.: The impact of 2022 hunga tonga-hunga ha'apai (hunga) eruption on stratospheric
640 circulation and climate. *Journal of Geophysical Research: Atmospheres*, 130, e2024JD042943.
<https://doi.org/10.1029/2024JD042943>, 2025.

Yu, W., Garcia, R., Yue, J., Smith, A., Wang, X., Randel, W., et al.: Mesospheric temperature and circulation response to the Hunga Tonga-Hunga-Ha'apai volcanic eruption. *Journal of Geophysical Research: Atmospheres*, 128, e2023JD039636.
<https://doi.org/10.1029/2023JD039636>, 2023.

645

Zhang, J., Wang, P., Kinnison, D., Solomon, S., Guan, J., Stone, K., & Zhu, Y.: Stratospheric chlorine processing after the unprecedented Hunga Tonga eruption. *Geophysical Research Letters*, 51, e2024GL108649.
<https://doi.org/10.1029/2024GL108649>, 2024.

650 Zhu, Y., Portmann, R. W., Kinnison, D., Toon, O. B., Millán, L., Zhang, J., Vömel, H., Tilmes, S., Bardeen, C. G., Wang, X., Evan, S., Randel, W. J., and Rosenlof, K. H.: Stratospheric ozone depletion inside the volcanic plume shortly after the 2022 Hunga Tonga eruption, *Atmos. Chem. Phys.*, 23, 13355–13367, <https://doi.org/10.5194/acp-23-13355-2023>, 2023.

655 Zhu, Y., Akiyoshi, H., Aquila, V., Asher, E., Bednarz, E. M., Bekki, S., Brühl, C., Butler, A. H., Case, P., Chabriat, S., Chiodo, G., Clyne, M., Falletti, L., Colarco, P. R., Fleming, E., Jörnemann, A., Kovilakam, M., Koren, G., Kuchar, A., Lebas, N., Liang, Q., Liu, C.-C., Mann, G., Manyin, M., Marchand, M., Morgenstern, O., Newman, P., Oman, L. D., Østerstrøm, F., Peng, Y., Plummer, D., Quaglia, I., Randel, W., Rémy, S., Sekiya, T., Steenrod, S., Sukhodolov, T., Tilmes, S., Tsigaridis, K., Ueyama, R., Visioni, D., Wang, X., Watanabe, S., Yamashita, Y., Yu, P., Yu, W., Zhang, J., and Zhuo, Z.: Hunga Tonga-Hunga Ha'apai Volcano Impact Model Observation Comparison (HTHH-MOC) Project: Experiment Protocol and Model Descriptions, EGUsphere [preprint], <https://doi.org/10.5194/egusphere-2024-3412>, 2024.

665 Zhuo, Z., Wang, X., Zhu, Y., Bednarz, E. M., Fleming, E., Colarco, P. R., Watanabe, S., Plummer, D., Stenchikov, G., Randel, W., Bourassa, A., Aquila, V., Sekiya, T., Schoeberl, M. R., Tilmes, S., Yu, W., Zhang, J., Kushner, P. J., and Pausata, F. S. R.: Comparing Multi-Model Ensemble Simulations with Observations and Decadal Projections of Upper Atmospheric Variations Following the Hunga Eruption, EGUsphere [preprint], <https://doi.org/10.5194/egusphere-2025-1505>, 2025.