

Leveraging TROPOMI observations and WRF-GHG modeling towards improving methane emission assessments in India

Thara Anna Mathew^{1,2}, Dhanyalekshmi Pillai^{1,2}, Jithin Sukumaran^{1,2}, Monish Vijay Deshpande^{1,2,3}, Michael Buchwitz⁴, Oliver Schneising⁴, Vishnu Thilakan^{1,2,5}, Aparna Ravi^{1,2}, Sanjid Backer Kanakkassery^{1,2,6}, Advait J Vinod¹, Sivarajan Sijikumar⁷, Imran A Girach⁸, and S Suresh Babu⁷

¹Indian Institute of Science Education and Research Bhopal (IISERB), India

²Max Planck Partner Group (IISERB), Max Planck Society, Munich, Germany

³Now at the University of Michigan, Ann Harbor, Michigan, USA

⁴Institute of Environmental Physics (IUP), University of Bremen FB1, Bremen, Germany

⁵Now at Lund University, Lund, Sweden

⁶Now at Max Planck Institute for Biogeochemistry (MPI-BGC), Jena, Germany

⁷Space Physics Laboratory (SPL), Vikram Sarabhai Space Centre, Thiruvananthapuram, India

⁸Space Applications Centre (SAC), Indian Space Research Organization, Ahmedabad, India

Correspondence: Dhanyalekshmi Pillai (dhanya@iiserb.ac.in)

Abstract.

Atmospheric methane (CH₄) contributes to global warming and climate change. Multiple factors control its atmospheric growth rate, posing challenges for climate change mitigation in regions with limited observations, like India. In this study, we examine the potential of dry air column methane mixing ratio (XCH₄) observations from the TROPOspheric Monitoring Instrument (TROPOMI) in conjunction with the high-resolution Weather Research and Forecasting model with Greenhouse Gas module (WRF-GHG) to improve the annual CH₄ budget of India. In addition to an inversion framework, we present a spatiotemporal assessment of bottom-up Indian methane emissions and their influence on XCH₄, supplying the context needed for regional emission optimization. Our analysis demonstrates the potential of WRF-GHG to represent the atmospheric XCH₄ and CH₄ distributions, including seasonal patterns, albeit with non-negligible uncertainties when compared with satellite and ground-based observations for 2018 and 2019. We find that the WRF-GHG simulations tend to overestimate XCH₄ while underestimating near-surface CH₄ concentrations at the Thumba site. Our inversion analyses report annual CH₄ emissions ranging from 21.9 to 24.9 Tg with an uncertainty of 3.3 Tg (anthropogenic sources), implying an overestimation of 13 to 24 % by the EDGAR global inventory. Also, our estimates are approximately 19 % higher than those in the India Fourth Biennial Update Report (19.6 Tg) and close to the latest Global Methane Budget 2000-2020. Overall, this study demonstrates the usefulness of TROPOMI observations for assessing Indian CH₄ emissions and shows a way to improve our understanding of how regional processes can modulate atmospheric CH₄ mixing ratios. We highlight the need for expanded observational coverage and an improved carbon assimilation system over India to refine the methane budget in support of global climate goals.

1 Introduction

20 The concentration of atmospheric CO₂ has increased by more than 50 % of the pre-industrial levels, while that of CH₄ has increased by 150 % (Friedlingstein et al., 2024; Saunio et al., 2025). CH₄ is the most prevalent non-CO₂ greenhouse gas, with a warming potential 28 times that of CO₂ over 100 years and 84 times over 20 years. (Lee et al., 2023; Montzka et al., 2011; Saunio et al., 2016; Stocker et al., 2013) and an atmospheric lifetime of 9.1 ± 0.9 years (Zhou et al., 2023; Saunio et al., 2025). Starting from 2007, the concentration of CH₄ has proliferated from an annual global mean of 1775 ppb to 1921 ppb in 25 2024, with a total rise of 146 ppb, which denotes a huge overall growth since the start of industrialization. The warming of wetlands, an increase in the ruminant population, and a decline in biomass burning, previously masking the rise in isotopically negative fuel use, are some of the key factors that may have contributed to the recent surge in CH₄ concentrations (Nisbet et al., 2019). The observed decline in carbon isotope ratio ($\delta^{13}\text{CH}_4$) indicates a shift toward increasing biogenic CH₄ sources, such as microbial emissions from wetlands and agriculture (has a more negative $\delta^{13}\text{CH}_4$ signature) rather than fossil fuel or 30 biomass burning contributions (Skeie et al., 2023; Schaefer et al., 2016). The long-term trend in OH remains uncertain, with some studies suggesting increases (e.g. Stevenson et al. (2020)), others finding no significant trend (Thompson et al., 2024), and still others showing diverging results depending on methodology (Saunio et al., 2025).

The global stock-take under Article 14 of the Paris Agreement implies the responsibility of each party to prepare, communicate, and maintain the successive nationally determined contributions (NDCs) to climate action (EC 2014). A 30 % global 35 reduction in CH₄ emissions from 2020 to 2030 has been aimed at the Global Methane Pledge, launched at a 2021 meeting of the United Nations Framework Convention on Climate Change (UNFCCC). Being one of the significant contributors to greenhouse gas (GHG) emissions, India plays an essential role in the global GHG scenario. Still, it lacks sufficient long-term, continuous, and accurate observations of the GHG to quantify the sources and sinks (Zhang et al., 2014). The country has the largest cattle (including bovine) population in the world (Robinson et al., 2014). Along with this, its huge intense flood irrigation practices, 40 ever-increasing fuel demand, and large wetland extent (nearly 4.7 % of its total geographical area) contribute to its high CH₄ emission potential (Ganesan et al., 2017; Garg et al., 2011; Ministry of Environment and Change, 2015; Myhre et al., 2013b). CH₄ emissions from enteric fermentation account for about 44 % of the total CH₄ emissions of India's National GHG inventory 2020 (MoEFCC, 2024). The Emissions Database for Global Atmospheric Research (EDGAR) inventory provides a global sector-wise emission estimate for CH₄. However, the bottom-up approach of EDGAR inventory is limited by its accuracy and 45 temporal resolution owing to uncertainties in the data used and methodologies (e.g. uncertain emission factors, aggregation or interpolation errors and sector distribution). Besides, Indian wetland emissions also show inconsistency in estimations (~ 5 to 9 %) depending on the wetland model used (Bloom et al., 2017). Janardanan et al. (2024) reported large uncertainty in wetland

emission inventory data over the Indian domain based on satellite observations and models. Further, insufficient coverage of highly precise and accurate ground-based observations of CH₄ and inadequate access to emission reporting over the country
50 can lead to misrepresentations in global emission inventories.

Atmospheric concentration measurements contain integrated information on the underlying source-sink distribution. Therefore, integrating atmospheric mixing ratio measurements, flux information from bottom-up approaches, and transport model simulations can potentially enhance CH₄ estimates through inverse modeling (Bergamaschi et al., 2018) and independently evaluate reported flux estimates. Previous studies over India have been limited by coarse model resolution, incomplete representation of transport processes, or lack of high-resolution emission inventories. Due to these inadequate modeling systems and sparse ground measurements, limited studies have used atmospheric CH₄ observations to inform about CH₄ emission flux estimates across India. There is an urgent call for measurement that can sufficiently constrain regional emissions in modeling systems (Patra et al., 2016). Recent technological advancements in satellite remote-sensing enable high-resolution-high-density observations to be utilized for this inverse-based quantification when modeling techniques are adequately advanced (Myhre
55 et al., 2013a; Jacob et al., 2016; Alexe et al., 2015; Buchwitz et al., 2017; Liang et al., 2023; Lu et al., 2022). Ganesan et al. (2017) used a top-down approach to estimate India's CH₄ emission for 2010-2015. The above study used column-averaged observations of CH₄ from the Greenhouse Gases Observing Satellite (GOSAT) along with aircraft observations from Civil Aircraft for the Regular Investigation of the Atmosphere Based on an Instrument Container (CARIBIC) and a few surface measurements from Indian sites to calculate methane emissions by atmospheric inverse modeling. Since 2009, GOSAT has
60 measured the atmospheric column for CH₄ every three days at a 10 km diameter circle (Butz et al., 2011). Despite some limitations in temporal coverage, the spatial resolution of GOSAT observations is much better than its predecessor, the Scanning Imaging Absorption Spectrometer for Atmospheric Chartography (SCIAMACHY), which the research community has widely used (Butz et al., 2011; Yokota et al., 2009; Turner et al., 2015; Buchwitz et al., 2005; Schneising et al., 2011).

Since November 2017, the more recent TROPOspheric Monitoring Instrument (TROPOMI) on board the Copernicus Sentinel-
70 5 Precursor satellite provides much higher-density CH₄ observations at a high spatial resolution of $7 \times 7 \text{ km}^2$, upgraded to $5.5 \times 7 \text{ km}^2$ in August 2019 (Hu et al., 2018; Schneising et al., 2023). TROPOMI measures CH₄ at the $2.3 \mu\text{m}$ band, with a swath width of 2600 km (Jacob et al., 2016; Cusworth et al., 2018). These observations are expected to capture seasonal fluctuations, which, in turn, will give better insight into the source-sink characteristics and quantification. Hence, in the present study, we explore the potential of TROPOMI measurements in representing the distribution of CH₄ fluxes over the Indian region along-
75 side a spatio-seasonal analysis of the CH₄ bottom-up inventory information. We use TROPOMI XCH₄ for its dense, near-daily coverage over India, enabling higher comparability with the simulation of 10 km from our Weather Research and Forecasting

model coupled with the Chemistry and Greenhouse Gas module (WRF-GHG), which may effectively minimize the forward model-related uncertainties in the carbon assimilation system over India. The performance of this high-resolution model and the advantage of using highly resolved transport fields are previously reported in Thilakan et al. (2022) and Vellalassery et al. (2021). The assessment of the forward model, WRF-GHG, in the atmospheric boundary layer is performed by comparing the atmospheric CH₄ simulations with atmospheric measurements from a ground-based site. Finally, the annual CH₄ emission estimate is also derived for the period 2018-2019 by incorporating the TROPOMI measurements and WRF-GHG forward model in an atmospheric inversion algorithm. The overview of spatiotemporal analysis of methane-emission patterns across India from global bottom-up inventories, including agriculture, livestock, and fossil-fuel sectors and their impact on column-averaged methane mixing ratio enhancements, provides the context for improving or complementing current emission estimates over India.

The paper is organized as follows: Section 2 describes the data and methods used for the study. Section 3 presents data post-processing and inverse analysis, and Section 4 discusses the results of the study. The conclusions of the study are presented in Section 5.

2 Data and methods

In this section, we describe the measurements and techniques used for exploring the potential of TROPOMI measurements and the WRF-GHG atmospheric transport model in inferring CH₄ distribution over India. An inverse method has been devised to deduce the CH₄ fluxes over the Indian region by minimizing mismatches between TROPOMI CH₄ measurements and WRF-GHG mixing ratio simulations, thereby correcting the distribution of prior fluxes. Figure 1 shows the model domain with outlines of each geographical region (more details are given in the subsequent sections) considered in this study.

2.1 TROPOMI observations

The potential of TROPOMI Sentinel-5p to detect significant sources in the single overpass has been demonstrated in recent publications (Hu et al., 2018; de Gouw et al., 2020; Schneising et al., 2020; Chen et al., 2022; Jacob et al., 2022). We utilized TROPOMI CH₄ observations obtained through the Short Wave Infrared (SWIR) band, centered at approximately 2.3 μm. The atmospheric column-averaged CH₄ mixing ratio (XCH₄) is retrieved using the Weighting Function Modified Differential Optical Absorption Spectroscopy (WFM-DOAS) algorithm. The WFMD algorithm uses a least-squares approach based on scaling prior atmospheric vertical profiles to retrieve XCH₄ and XCO simultaneously (Buchwitz et al., 2007; Schneising et al., 2011). Here, we use the WFMD v1.8 algorithm, for which the efficiency has been validated using Total Carbon Column

Observing Network (TCCON) measurements, resulting in an improved random error (12.4 ppb) compared to the previous
105 versions (v1.5 and v1.2) (Schneising et al., 2023). For the analysis, we filtered flagged data and utilized only good-quality
retrievals represented by *xch4_quality_flag* = 0.

2.2 Modeling system for atmospheric XCH₄ mixing ratio simulations

We have used the Weather Research and Forecasting model coupled with the Chemistry and Greenhouse Gas module (hereafter
referred to as WRF-GHG) for atmospheric CH₄ transport simulations. The core component is the WRF model, based on
110 fully compressible, non-hydrostatic Eulerian equations on terrain-following vertical grids for simulating atmospheric transport
(Skamarock et al., 2008). The GHG-TRACER package allows the online passive tracer transport of CH₄ mixing ratio in the
atmosphere (Beck, 2011; Pillai et al., 2016). The input fluxes from each emission sector are separately provided as "tagged"
tracers when added to the first layer in the modeling system. This allows the decoupling of emission contributions to the total
atmospheric CH₄ mixing ratio. We have used the WRF-GHG 3.9.1.1 version with a horizontal resolution of 10×10 km²
115 (Lambert conformal conic projection grid) and an output temporal resolution of 1 hour. The model covers the Indian domain
with 307×407 grid points and 39 vertical levels. The fifth generation ECMWF reanalysis (ERA-5) data with a horizontal
resolution of $0.25^\circ \times 0.25^\circ$ and temporal resolution of 6 hours with 137 vertical levels are used as initial and boundary
conditions for meteorology (Hersbach et al. (2020); see Table 1). The model is re-initialized each day with

We used the Emission Database for Global Atmospheric Research (EDGAR v7.0; Crippa et al. (2024)), Global Fire Assim-
120 ilation System (GFAS v1.2; Kaiser et al. (2012)), and a global wetland CH₄ emissions and uncertainty dataset for atmospheric
chemical transport models (WetCHARTs 1.3.1; Bloom et al. (2021)) as prior emission fluxes to represent anthropogenic,
biomass burning and wetland emissions respectively. We applied temporal scaling factors to the annual EDGAR emissions
using step-function time profiles and converted them to 1-hour temporal resolution, following Kretschmer et al. (2014). ERA5
meteorology, in which a 6-hour spin-up time was configured. For CH₄ mixing ratio fields, initial and boundary conditions are
125 prescribed from the Copernicus Atmosphere Monitoring Service (CAMS re-analysis data). Meteorological fields are reinitial-
ized daily using ERA5 reanalysis, followed by a 6-hour spin-up period. In contrast, note that CH₄ tracer fields are initialized
only at the beginning of the simulation and the simulated background fields are continuously transported across successive
simulation days (i.e. not reinitialized daily). This approach ensures that methane concentrations reflect the cumulative influence
of emissions, preserving the temporal memory for inverse optimization of local surface fluxes. CAMS provides the simulated
130 atmospheric mixing ratios of CH₄, with a spatial resolution of $0.25^\circ \times 0.25^\circ$ and a temporal resolution of 6 hours on 60 vertical
levels (Inness et al. (2019); see Table 1). The model utilizes these initial fields to represent the background of the total mixing




Figure 1. (a) Outlines showing each geographical region of the Indian landmass according to Survey of India (2024) and (b) the topographical height contour of the model domain. CI stands for Central India, NEI for North East India, NI for North India, SI for South India, and IGP for the Indo Gangetic Plain regions.

ratios. Similar to emission flux contributions, we disentangled the background contribution in the model output to investigate its impacts separately. We have also considered the reported level of uncertainties in CAMS-simulated mixing ratios (e.g. Agustí-Panareda et al. (2023); Wang et al. (2023)) and applied a monthly scaling factor correction to our methane background
 135 (initialized by CAMS) to minimize the unrealistic representation of background contribution to methane mixing ratios. The scaling factor is applied uniformly across the domain to the CH₄ background, specific for each month (i.e., one scaling factor per month). These monthly background scaling factors range from 1% to 3%. The corrected WRF-GHG background mixing ratios are hereafter termed simply as background mixing ratios. GFAS data with 0.1° × 0.1° resolution represents biomass
 140 Resolution Imaging Spectroradiometer (MODIS) instrument aboard the Terra satellite.

WetCHARTs is an ensemble dataset that provides gridded emissions data from 2001 to 2019 at a resolution of 0.50° × 0.50° (Bloom et al., 2017), which were then re-gridded to 0.1° × 0.1° with a temporal resolution of 1 hour. The simulated total atmospheric CH₄ mixing ratio thus contains contributions from the initial fields as well as anthropogenic, biomass burning, and biogenic fields of CH₄. Table 1 summarizes the WRF-GHG model set-up used in this study. The general meteorological
 145 configuration for the WRF-GHG model set-up applied here is described in Thilakan et al. (2022).

Table 1. WRF-GHG model configuration.

Parameter	Details
Domain Configuration	Single domain with a horizontal resolution of 10 km; 39 vertical levels; 307 × 407 grid points
Vertical coordinates	Terrain-following hydrostatic pressure vertical coordinates
Basic equations	Non-hydrostatic; compressible
Grid type	Arakawa C grid
Time integration	Third-order Runge–Kutta split explicit
Spatial integration	Third- and fifth-order differencing for vertical and horizontal advection, respectively; both for momentum and scalars
Time step	60 s
Model top pressure	50 hPa
Physics schemes	Rapid Radiative Transfer Model (RRTM) for longwave and Dudhia for shortwave
<i>Radiation</i>	WRF single-moment three-class (WSM3) classic simple ice scheme
<i>Microphysics</i>	Yonsei University (YSU) scheme
<i>PBL</i>	Monin–Obukhov
<i>Surface layer</i>	NOAH land surface model (LSM)
<i>Land surface</i>	Grell–Freitas ensemble scheme
<i>Cumulus</i>	
Emission fields	
<i>Flux type</i>	Product Version Spatial res. Temporal res. Source/website Reference
<i>Anthropogenic</i>	EDGAR v7.0 10 km Annual https://edgar.jrc.ec.europa.eu/ (Crippa et al., 2024)
<i>Biomass burning</i>	GFAS v1.2 10 km Daily http://apps.ecmwf.int/datasets/data/cams-gfas/ (Kaiser et al., 2012)
<i>Biospheric</i>	WetChARTs V.1.3.1 0.5° Monthly https://daac.ornl.gov/cgi-bin/dsviewer.pl?ds_id=1915 (Bloom et al., 2017)
Initial and lateral boundary conditions	Product Version Spatial res. Temporal res. Source/website Reference
<i>Field</i>	ERA5 NA 25 km 1 h https://cds.climate.copernicus.eu/cdsapp#!/home (Hersbach et al., 2020)
<i>Meteorology</i>	ECMWF/CAMS EGG4 50 km 6 h http://atmosphere.copernicus.eu (Agusti-Panareda et al., 2023)
<i>Tracer</i>	

2.3 Ground-level observations

To assess the model's performance at the surface level, a comparative analysis was conducted using CH₄ *in situ* measurements from a ground-level pollution monitoring station in Thumba (8.5°N, 76.9°E) as denoted in Fig. 1 for 2018 & 2019. Located in southwestern India, Thumba is a tropical coastal station approximately 10 km northwest of Thiruvananthapuram and 500 m inland from the Arabian Sea. This site reflects local to regional influences, but cannot capture the full spatial variability across India. CH₄ concentrations were measured using a greenhouse gas analyzer (model: 911-0011-1001) by Los Gatos Research, USA, based on the off-axis integrated cavity output spectroscopy (OA-ICOS) method (Baer et al., 2002; Raju et al., 2022; Sijikumar et al., 2023; Uma et al., 2024). Air samples were collected from about 10 m above ground level (AGL) using the analyzer's internal pump. Calibration was performed periodically. However, it should be noted that the instrument can be sensitive to temperature, requiring frequent calibration, which was not regularly met. Measurements were recorded at 1-second intervals, with hourly averages used for subsequent analysis. CH₄ measurement uncertainty is 0.25 % (i.e. 5 ppb with respect to 2000 ppb of CH₄) and the reported precision (1 σ) is 1 ppb.

2.4 Data post-processing and inverse analysis

We regridded the daily total dry column mixing ratio of CH₄ from TROPOMI at 10 km \times 10 km resolution, covering the period from 2018 to 2019. From the hourly WRF-GHG CH₄ mixing ratio simulations generated, we sampled those corresponding to the TROPOMI overpass time for the model domain. The model top is set near \sim 50 hPa. When computing the column-averaged mole fractions from simulations, the model profile is extended above the model top utilizing satellite *a priori* profiles, ensuring the same sampled air column corresponding to the observation datasets, also following *a priori* profile weights when computing the column-averaged mole fraction (more details below). Here, we assume negligible model biases for the remaining atmospheric contribution above the model top compared to that of the tropospheric and lower stratospheric counterparts. Previous studies of stratospheric and tropospheric contributions to total column CH₄ demonstrated that the model biases in the stratospheric partial column tend to be smaller than those in the tropospheric partial column owing to the substantially higher tropospheric CH₄ contribution to the total column than stratospheric CH₄ (Wang et al., 2017). However, the assumption underlying the extension above the actual model top is a simplification of real-case conditions and may be considered when analyzing the results.

In the present study, the model simulations have not accounted for any impacts of stratospheric CH₄ chemistry reactions on mixing ratios. Ignoring chemical reactions is typically a valid assumption when considering the regional model domain and the considerably longer atmospheric lifetimes of target species, approximately 10 years for CH₄, than the simulation period.

Excluding the OH reactions has shown a negligible impact on annual CH₄ at the regional scale, resulting in smaller biases
175 than the measurement uncertainties (1 ppb for in situ measurements and 6 ppb for TCCON) and the typical magnitudes of the
observational bias in the inversion (Callewaert et al., 2025). However, atmospheric CH₄ is susceptible to chemical reactions in
the stratosphere, which warrants consideration in modelling and analyzing long time series such as decadal contributions.

To ensure a fair comparison with observations, we applied the satellite's averaging kernel (AK), as shown in Equation 1,
accounting for the satellite instrument's vertical sensitivity (Schneising et al., 2019). AK is proportional to the sensitivity profile
180 of the measurement that is weighted with the assumed tracer profiles and provides the relation between the retrieved and known
tracer profiles. i.e. Applying satellite AK to the model simulations at different vertical levels minimizes the mismatches owing
to the instrument vertical sensitivities to the column observations (Eskes and Boersma, 2003; Wang et al., 2023; Schneising,
2024).

We applied the AK to the modeled dry-air CH₄ profiles and derived the dry-air column-averaged mixing ratio of CH₄, Υ_{mod} ,
185 as follows:

$$\Upsilon_{\text{mod}} = \sum_l (\Upsilon_{\text{apr}}^l + A_l (\Upsilon_{\text{mod}}^l - \Upsilon_{\text{apr}}^l)) w_l \quad (1)$$

where l is the index of the vertical layer, A_l is the averaging kernel and Υ_{apr}^l is the *a priori* mole fraction of layer l , and Υ_{mod}^l
is the corresponding simulated mole fraction of layer l . w_l is the layer-dependent pressure weight.

Hence, Υ_{mod} (= XCH_{4,mod}) is used for the model-observation comparisons and inversion analyses. i.e., in this study, Υ_{mod}
190 represents WRF-GHG XCH₄ simulations.

2.4.1 Estimating optimized CH₄ flux over India

We performed a simple Bayesian inverse optimization to deduce the improved emission estimates over the Indian domain. The
inversion is designed in such a way that it describes the relationship between the mixing ratio observations and the surface
flux emission information (the unknown state) and *a priori* information available. This approach allows us to identify the class
195 of possible states consistent with the available information and to assign a probability density function (pdf) to them. The
quantities to be optimized, represented by the state vector \mathbf{x} with n elements $x_1, x_2, x_3, \dots, x_n$ correspond to the monthly,
state-wise emissions from (i) anthropogenic components (EDGAR) and (ii) the sum of anthropogenic (EDGAR) and biomass-
burning (GFAS) components, both (i) and (ii) optimized separately. We have omitted the wetland component here since it
contributed negligibly to the column-mixing-ratio enhancement. Here, n represents 36 state regions of India. The measured

200 quantities, represented by the measurement vector \mathbf{y} with m elements $y_1, y_2, y_3, \dots, y_m$, represent the column observations from TROPOMI over a month at $0.1^\circ \times 0.1^\circ$ spatial resolution. Here, m represents the total number of observations available in each political state.

The relationship between the measurement vector, \mathbf{y} , and the state vector, \mathbf{x} , can be written as:

$$\mathbf{y} = \mathbf{F}(\mathbf{x}) + \epsilon \quad (2)$$

205 where $\mathbf{F}(\mathbf{x})$ encapsulates the physics of the measurements as a function of the state vector, described here by our forward model, WRF-GHG, which includes forward transport and mapping of flux fields. The error term ϵ includes model error, representation error (sampling mismatch between the observations and the model), and measurement error.

Linearizing the forward model to a reference state yields:

$$\mathbf{F}(\mathbf{x}) = \mathbf{K}\mathbf{x} + \epsilon \quad (3)$$

210 Here, \mathbf{K} is the $m \times n$ Jacobian matrix, representing the sensitivity of the mixing ratio simulated by the forward model to the state vector. The elements of \mathbf{K} are thus: $k_{m,n} = \frac{\partial \mathbf{F}_m(\mathbf{x})}{\partial x_n}$

Since we have not implemented the adjoint model for our forward transport model, the Jacobian matrix is constructed using a finite-difference approach: the transport model (WRF-GHG) is applied to perturbed emissions, and sensitivities of the resulting mixing ratio simulations are derived. This perturbation-based Jacobian construction is mathematically equivalent
 215 to computing the response functions to state vectors. The Jacobian matrix in the present study ensures capturing near-field influence on observations and minimizes the over-interpretation of far-field influences. For robust flux estimations at regional and sub-regional scales, it is widely accepted that increasing reliance on near-field sensitivity minimizes the ill-posed solutions (Nesser et al., 2024; Munassar et al., 2023; Ware et al., 2019; Pillai et al., 2012; Gerbig et al., 2006).

The elements in the Jacobian matrix are derived as follows:

$$220 \quad k_{m,n} = \frac{(\mathcal{Y}_{pert})_m - \mathcal{Y}_m}{(\Phi_{pert})_n - \Phi_n} \quad (4)$$

The above derivation closely follows methods implemented in Pillai et al. (2016), Ye et al. (2020) and Kuhlmann et al. (2020). Here, \mathcal{Y} and \mathcal{Y}_{pert} are the 1-D representations of the column mixing ratio and the perturbed column mixing ratio, respectively,

both derived by our forward model, corresponding to each m elements in the measurement vector \mathbf{y} . Φ and Φ_{pert} represent emissions and perturbed emissions, respectively, corresponding to each n elements in the state vector \mathbf{x} . Note that perturbed
 225 mixing ratios are simulated by applying the transport operator (WRF-GHG) to perturbed emissions, as explained before. By our design, the elements in the \mathbf{y} correspond to the available column mixing ratio observations over a month at $0.1^\circ \times 0.1^\circ$ spatial resolution, and the elements in the \mathbf{x} refer to total monthly emissions of each political state.

In our implementation, we focus on anthropogenic fluxes and their contributions to atmospheric dry-air column mixing ratios. \mathbf{x}_A represents the prior fluxes, which consist of monthly anthropogenic (major contributions from enteric fermentation,
 230 agricultural soil, waste water handling, and fuel exploitation) and biomass burning emissions. The inversion optimizes the corresponding total emissions per state by minimizing the model-observation mismatches. The background contributions are removed from the observations to target optimizing the regional enhancement fluxes. i.e., the measurement vector \mathbf{y} consists of column observations from TROPOMI subtracted by simulated background mixing ratios over a month at $0.1^\circ \times 0.1^\circ$ spatial resolution.

235 The background mixing ratios are simulated by WRF-GHG, as explained in Sect. 2.2. We have considered measurement errors (including forward model errors) and prior errors: \mathbf{S}_e and \mathbf{S}_a represent measurement error and prior error covariance matrices, respectively. The measurement error covariance matrix \mathbf{S}_e consists of retrieval (\mathbf{y}) and the forward model ($\mathbf{F}(\mathbf{x})$) errors for methane. We assumed a prior emission uncertainty of 80% and a measurement uncertainty of 16 ppb, as adopted from Liang et al. (2023), calculated using the residual error method (Heald et al., 2004). We have not considered cross-correlations;
 240 hence, only the diagonal elements of the matrices \mathbf{S}_e and \mathbf{S}_a are non-zero.

Since multiple observational products are available and we may expect differences among those products, we assess product differences across our region using scientific, operational, and GOSAT-blended products. The additional data products are: the operational Sentinel-5P/TROPOMI Level-2 methane product provided by ESA/Copernicus (Copernicus Sentinel-5P, 2021), and the blended TROPOMI+GOSAT methane product available from April 2018 (Balasus et al., 2023). Further, we conduct
 245 an additional inversion by inflating the measurement uncertainty by more than 50 % (~ 25 ppb) to examine the influence of product differences on posterior flux uncertainties.

The solution to the inverse problem is obtained by minimizing the Bayesian scalar cost function $J(\mathbf{x})$ (Rodgers, 2000):

$$J(\mathbf{x}) = (\mathbf{x} - \mathbf{x}_A)^T \mathbf{S}_a^{-1} (\mathbf{x} - \mathbf{x}_A) + (\mathbf{y} - \mathbf{K}\mathbf{x})^T \mathbf{S}_e^{-1} (\mathbf{y} - \mathbf{K}\mathbf{x}_A) \quad (5)$$

where $\nabla_{\mathbf{x}}J(\mathbf{x}) = 0$, the optimal estimate $\hat{\mathbf{x}}$ is obtained (Rodgers, 2000). The state that maximizes the posterior pdf $P(\mathbf{x}|\mathbf{y})$ is the maximum a posteriori solution (MAP). The maximum a posteriori solution is obtained as follows:

$$\hat{\mathbf{x}} = \mathbf{x}_A + \left(\mathbf{K}^T \mathbf{S}_e^{-1} \mathbf{K} + \mathbf{S}_a^{-1} \right)^{-1} \mathbf{K}^T \mathbf{S}_e^{-1} (\mathbf{y} - \mathbf{K} \mathbf{x}_A) \quad (6)$$

Thus, $\hat{\mathbf{x}}$ represents the optimized spatially averaged monthly anthropogenic and biomass burning fluxes corresponding to each political state considered. The posterior error covariance matrix, denoted by $\hat{\mathbf{S}}$, is derived as follows.

$$\hat{\mathbf{S}} = \left(\mathbf{K}^T \mathbf{S}_e^{-1} \mathbf{K} + \mathbf{S}_a^{-1} \right)^{-1} \quad (7)$$

The error reduction for each month following the inversion procedure (\tilde{e}) is calculated as follows:

$$\tilde{e} = 1 - \frac{\sigma_{post}}{\sigma_{prior}} \quad (8)$$

where σ_{post} is derived as the square root of the diagonal elements of $\hat{\mathbf{S}}$. Similarly, σ_{prior} is obtained from the square root of the diagonal elements of the prior error covariance matrix.

The national budget for annual optimized fluxes $\hat{\mathbf{x}}_{annual}$ (in Tg y^{-1}) is derived as:

$$\hat{\mathbf{x}}_{annual} = \sum_{t=1}^{N_t} \sum_{s=1}^{N_s} \hat{\mathbf{x}}_{s,t}, \quad (9)$$

where $\hat{\mathbf{x}}_{s,t}$ represents the estimated value at each political state s for the corresponding month t . N_s is the total number of political states, and N_t is the total number of months considered.

3 Results and Discussion

3.1 Regional and sectoral distribution of CH₄ sources

The bottom-up inventories such as EDGAR v8.0 (latest release; Crippa et al. (2024)), WetCHARTs v1.3.1, and GFAS v1.2. relate the GHG emissions to the causative processes by considering emission activities and emission factors, thereby providing us with a "first guess" to identify the prominent sources (Miller and Michalak, 2017), although with inherent uncertainties. In this section, we present a detailed comparative assessment of the sectoral and regional distributions of different CH₄ sources (such as enteric fermentation, wastewater handling, rice agricultural land, wetlands, and biomass burning).



(a)**(b)**

Figure 2. (a) The percentage of contributions of different CH₄ sources towards total annual emission flux over the Indian domain, (b) monthly contribution (calculated from the EDGAR temporal profiles described in Crippa et al. (2020)) from each source.

270 Our sector-wise analysis of the bottom-up inventories shows that the enteric fermentation associated with the digestive process in cattle makes a significant contribution to CH₄ emissions in India (42.9 %), followed by wastewater treatment (19.2 %), agricultural soil (12.4 %), fuel exploitation (6.7 %), and wetland (5.2 %, excluding agriculture) (see Fig.2). The sources with significant seasonality include agriculture (see Fig.2 (b)) and biomass burning, also reported in Ganesan et al. (2017). Figure S1. (a)-(d) shows an annual average of the spatial distribution of CH₄ emissions for the four major emission sectors in

275 2018. Anthropogenic sources are expected to provide the bulk of India's CH₄ emissions, especially livestock, rice cultivation, and waste management (Maasackers et al., 2019). The annual CH₄ emissions corresponding to rice cultivation (emission from the sector 'Agricultural soil' as given by the EDGAR inventory) show a peak in values over the Indo-Gangetic Plain (IGP) region (Fig.S1 (c)). Several other studies have used GOSAT and other data sources to analyse the CH₄ emissions from rice paddies in India (Miller et al., 2019; Ganesan et al., 2017; Anand et al., 2005). Previous studies reported CH₄ emissions from

280 rice paddies in India of about 3.9 Tg yr⁻¹, with the bulk emitted between June and September (Janssens-Maenhout et al., 2011; Garg et al., 2011; Panigrahy et al., 2010). The sector-wise analysis of the EDGAR inventory for the year 2018 shows significant CH₄ highs on the eastern coast, including West Bengal and Odisha, which can be attributed to the large rice cultivation in these regions (Crippa et al., 2023). The analysis of monthly emissions from rice cultivation over the year 2018 indicates an increasing pattern in summer monsoon seasons, June to September (see "Agricultural soil" in Fig. 2 (b)), with the maximum in August




Figure 3. Seasonal mean emissions from (top panels) wetland, (middle panels) biomass burning, and (bottom panels) anthropogenic sources, separated for seasons, for each region as specified in Figure 1. Note: The ranges of the y-axes are not uniform in panels to improve visibility.

285 (16.9 %) and September (16.5 %) and the minimum in April (1.9 %); percentages are shares of the annual flux. There is a
smaller peak in February-March time, owing to winter rice cultivation, which comprises 14 % of total rice production in India
(Manjunath et al., 2006). Similarly, in the wetland emissions, we also see an increase in monsoon months (Fig. 2 b). The peak
wetland emissions are seen in July (24.6 %), followed by August (21.4 %), and the minimum in January (0.2 %). It has been
previously reported from satellite observations that the waterlogged areas increase nearly threefold from the beginning to the
290 end of the monsoon, resulting in higher wetland CH₄ emissions (Agarwal and Garg, 2009). The pre-monsoon CH₄ emission
(15.5 %) is higher than post-monsoon (6.2 %; see Fig. 2 (b), possibly due to higher temperatures during the pre-monsoon season
as described in Das et al. (2023). Our analysis of monthly emissions based on GFAS shows a peak due to biomass burning in
March (65.4 %), with the lowest burning reported in July (0.1 %) (Fig. 3 (e)-(h)). Other sources of CH₄, including fossil-fuel
emissions, enteric fermentation, and wastewater handling, have not shown considerable seasonal variability. A similar pattern
295 has also been observed from the 2019 anthropogenic and natural CH₄ emission analysis (figure not shown).

Further, the anthropogenic emissions remain the largest contributors to regional CH₄ emissions (see Fig. 3 & S2). The total
anthropogenic emissions, reported by the EDGAR bottom-up inventory (version 8), peak seasonally in the South India (SI)
region ($> 700 \times 10^{12} \text{ kg m}^{-2} \text{ s}^{-1}$), followed by IGP ($> 400 \times 10^{12} \text{ kg m}^{-2} \text{ s}^{-1}$), with the minimum emissions observed in North
India (NI; see Fig. 3). The peak is typically in October-November, followed by June-September, with a minimum in March-
300 May. Total anthropogenic emissions over India show a peak in October-November and a dip in March-May ($1410- 1090 \times 10^{12}$
 kg m^{-2}). While IGP shows high magnitude in spatial distribution (see Fig. S1 (a)-(d)), SI contributes more emissions due to
emission hotspots. Four hotspots identified (see Fig. S3) in SI include one city- Namakkal (11.25°N , 78.15°E ; HS1) and three
villages- Mandapaka Rural (16.75°N , 81.65°E ; HS2), Pasumamla (17.35°N , 78.65°E ; HS3), and Mulkalappalli (18.65°
 $^\circ \text{N}$, 79.55°E ; HS4). Excluding these hotspots (SI_A) shifts the highest emissions to IGP (see Fig. 3 (i)-(l)). Namakkal in Tamil
305 Nadu faces poultry waste deposition, potentially increasing methane emissions (Ramasamy and Manivel, 2019). Mandapaka in
Andhra Pradesh, known as the "rice bowl" of the region, contributes to higher agricultural rice emissions (Gururaj Katti et al.,
2002). Pasumamla in Telangana sees significant poultry waste dumping in landfills ([https://rangareddy.telangana.gov.in/animal-](https://rangareddy.telangana.gov.in/animal-husbandry)
[husbandry](https://rangareddy.telangana.gov.in/animal-husbandry), accessed 12 Feb 2025). Mulkappalli, a coal mining site in Telangana, contributes to the high CH₄ levels in the
inventory (<https://khammam.telangana.gov.in/economy>, accessed 12 Feb 2025). Figure S4 shows the methane hotspots over
310 the Indian domain from the coal mining sector as derived from the EDGAR inventory.

Based on WetCHARTs (version 1.3.1), natural wetland emissions are approximated at 1.7 Tg yr^{-1} , with peak emissions
occurring from June to September ($\sim 185 \times 10^{12} \text{ kg m}^{-2}$). The highest wetland emissions ($> 25 \times 10^{12} \text{ kg m}^{-2} \text{ s}^{-1}$) are seen over
the SI region in June-September, October-November, and March-May, with a peak of $\sim 50 \times 10^{12} \text{ kg m}^{-2} \text{ s}^{-1}$ (Fig. 3). These

peaks are likely due to increased waterlogged areas during the monsoon (Fig. 2b). Biomass burning emissions, smaller than
315 wetland and anthropogenic emissions, peak in March-May in most regions except the IGP, where crop residue burning occurs
from October to November (Deshpande et al. (2022, 2023)). The North East India (NEI) region shows the highest biomass
burning emissions in March-May ($\sim 47 \times 10^{12}$ kg m⁻²), likely due to slash-and-burn practices before planting (Deshpande
et al. (2023)). Total emissions, combining anthropogenic (EDGAR), wetland (WetCHARTs), and biomass burning (GFAS)
emissions, peak in October-November and June-September, with the SI region contributing the highest share (49.2 % and 46.4
320 %, respectively), and 52 % in March-May and 51.7 % in December-February (Fig.S5).

Though the above estimations give an overview of Indian CH₄ source contributions and their regional patterns, there have
been increased concerns about their accuracy due to methodological weaknesses and data gaps (Solazzo et al., 2021; Madrazo
et al., 2018). For example, the combined emissions from Oil, Gas, and Coal over the Indian region reported by Scarpelli et al.
(2025) is 1.8 Tg yr⁻¹, whereas EDGAR reported 2.2 Tg yr⁻¹. Also, bottom-up methods can overestimate or misinterpret emission
325 sources even at the global level (Saunois et al., 2016). Though inverse modeling can improve the CH₄ budget significantly, its
potential to minimize biases in the bottom-up models used as prior estimates may be limited by insufficient coverage of mixing
ratio observations and its inadequate representation in the forward models. Satellite instruments such as TROPOMI may aid
in high observation density with good spatial coverage (Palmer et al., 2021), the potential of which in the Indian context is
explored in further sections.

330 **3.2 Assessment of the forward-model performance against surface measurements**

As discussed in Section 2.3, we utilized hourly ground-based observations from a ground-based site, Thumba, to assess the
WRF-GHG performance in the planetary boundary layer. The lowest level (approximately 35.2 m) of WRF-GHG-simulated
CH₄ at Thumba is compared with surface-level observations of CH₄ and is presented here. Generally, the analysis indicated
a reasonable performance of WRF-GHG simulations. CH₄ mixing ratios are found to be lowest during the monsoon season
335 (June–August), increasing from early October and peaking during the post-monsoon and winter months (November–January;
see Fig. 4a). The maximum values are seen in December (~ 2100 ppb). The hourly observations show high variability (about
112.7 ppb), but ranging from 1817.4 to 2612.6 ppb for 2018-2019; see Fig. S6. Despite some discrepancies, the WRF-GHG
simulations broadly align with those highly fluctuating observation patterns, capturing about 56 % of the observed variabil-
ity. In October, monthly averaged WRF-GHG simulations and TROPOMI observations are strongly correlated, although their
340 absolute values differ substantially (Figure not shown). The mean difference between observations and simulations is 47 ppb,
but it shows large model-observation variability of up to 73.9 ppb (Fig. 4(a) and Fig. S6(a)). However, this large discrepancy




Figure 4. Monthly distribution of observed and WRF-GHG simulated CH₄ mixing ratios at Thumba for 2018 & 2019 for (a) all hours and (b) only 15:00 - 17:00 Local Time (IST hours) (25th and 75th quartiles; see the site location as denoted in Fig. 1). Note that the ranges of the y-axes are not uniform in panels to improve visibility.

between the model and observations can be attributed to the influence of fine-scale nocturnal coastal meteorological conditions at the measurement site, as reported in Kavitha et al. (2018). Considering only the afternoon hours, the model-observation differences are reduced to 6.4 ppb, with a maximum difference varying up to 28.1 ppb, capturing about 79 % of observed variability (Fig. 4(b) and Fig. S6(b)). The comparison has also been performed by removing the boundary contributions from CAMS (see Fig. S7), showing that enhancements correlate with observed variability ($R^2 = 0.48$). Thus, the above comparison suggests the potential of our model in representing the regional and seasonal variations. The above result is promising, confirming the usability of those afternoon measurements representing well-mixed atmospheric conditions, which can be utilized for future carbon assimilation systems in conjunction with a high-resolution forward modeling framework. As seen for all hours, WRF-GHG generally underestimates surface CH₄ mixing ratios (Fig. S6). Notably, the model-observation differences peaked in winter, owing to the unusually high variability seen in the observations during this period. The effect of enhanced vertical mixing can be seen in the summer months, causing low observed CH₄ magnitude and associated mixing ratio variability. Noteworthy is that the magnitude of observed CH₄ is found to be the smallest during the summer monsoon. While the shallow planetary boundary layer (PBL) in the winter accumulates the effect of surface emissions to the lower boundary, increased boundary layer mixing in the summer can cause lower CH₄ magnitude and variability. Further, Guha et al. (2018) and Metya




Figure 5. Spatial distribution for annual WRF-GHG simulated anthropogenic mixing ratio enhancement of XCH₄ (including biomass burning) for 2018.

et al. (2021) report that the influx of clean air from the Southern Hemisphere, carried by the monsoonal south-westerly winds, can influence the surface CH₄ to lower its concentration. The high rates of OH radical oxidation may also influence surface CH₄ mixing ratios (Lin et al., 2015).

Even though WRF-GHG has shown reasonable performance in evaluating ground-based observations from a complex site (located near the southernmost coastal boundary of the model domain), the model's robustness needs to be further examined across multiple locations in India when available. Those evaluations are particularly necessary for assessing our confidence in the derived posterior fluxes. Although we found that local fluxes had a more dominant contribution to the observed variability




Figure 6. Distribution of seasonal average of simulated mixing ratio enhancement ($XCH_{4,ant} + XCH_{4,bbu}$) over different regions of India in 2018. The box plot displays medians, interquartile ranges, and minimum and maximum values, with data points beyond 1.5 times the interquartile range represented as outliers.

than the background contributions (see Fig. S7), there can be non-negligible differences arising from the choice of global model products used for initialization. For instance, we conducted additional analysis comparing the CAMS EGG4 product, which was used for initialization in this study, with the inversion-optimized CAMS product (Bergamaschi et al., 2013). This comparison indicated a difference of approximately 13 ppb over India (see Fig. S8). While it is expected that only a small fraction (less than 10 to 15%) of these initial tracer differences will effectively converge to influence model simulations (e.g. Monteil et al. (2011)), the potential impact of various tracer initializations on model simulations over this region is not addressed in the current study. The above aspect is worth a future investigation through a dedicated model sensitivity study.

3.3 Anthropogenic XCH_4 mixing ratio enhancements

In this section, we discuss the mixing ratio enhancements in the atmospheric column in response to spatial and temporal distributions of regional sources for the period 2018-2019. i.e. by considering only contributions from the sum of anthropogenic and biomass-burning emission sources (mostly human-influenced in India, i.e. from agricultural residue burning and managed

fires) over the model domain and not using CAMS-derived background XCH₄ (see section 2.2). As mentioned in section 375 2.4.1, we have also omitted the wetland (biogenic) component here since it contributed negligibly to the column-mixing-ratio enhancement (see Figs. S9 & S10). The IGP region exhibits significant XCH₄ enhancements (from 27 ppb to 67 ppb) from regional sources attributed to anthropogenic and biomass-burning fluxes (see Fig. 5). Seasonally, the highest XCH₄ enhancements occur during winter (with a maximum of ~ 251 ppb in January; see Fig. S11) over India. The minimum enhancement for the whole Indian domain occurs during the monsoon season (June–September), likely due to a combination of higher bound- 380 ary layer heights and stronger winds, which enhance vertical and horizontal transport affecting column CH₄ concentrations. The concentrations may also be impacted by the seasonal changes in regional or larger fluxes (>1000 km); however, further investigation is needed to assess their contributions.

Figure 6 shows the regional variability of anthropogenic XCH₄ enhancements across different parts of India as shown in Fig. 1. The highest regional magnitudes and variability in mixing ratio enhancements occur in the winter season. Here, consistently 385 high magnitude in spatial distribution is found over the IGP region (with a median of ~ 50 ppb), showing maximum values over the SI region (~ 90 ppb) owing to emission hotspots. During winter, the NEI shows high XCH₄ enhancements, with values reaching up to 115 ppb (with a median value of ~ 60 ppb). Winter peaks in XCH₄ likely arise from stable atmospheric transport that carries and concentrates emissions from the preceding October–November period. From June to September, SI enhancements reach up to ~ 75 ppb (with a median value of ~ 25 ppb). A similar trend is seen for 2019 (Figure not shown). 390 SI exhibits the widest interquartile range with the lowest minimum value, a relatively low median, and the highest maximum in most seasons, indicating the influence of hotspot emissions, as discussed earlier in Section 4.1.

3.4 Comparison of modeled and observed total XCH₄

In this section, we present our comparisons of WRF-GHG simulations with TROPOMI observations of total XCH₄ in 2018, considering all months in which reasonable amounts of satellite measurements are available after filtering. The details of 395 filtering are provided in Schneising et al. (2023). TROPOMI observations show distinct seasonal variations in the large spatial domain (Fig. 7), possibly resulting from both atmospheric transport and surface emissions variations. Observations indicate highest values in the mean spatial distribution of XCH₄ from October to November within the range of ~ 1862 to 1870 ppb. These seasonal increments can be attributed to the combination of surface emissions, boundary layer height, and horizontal transport, which accumulates the effect of the distribution of tracers at lower atmospheric levels. These increased regional 400 emissions, especially from anthropogenic sources, are also seen in Fig. S2 & Fig. 3, which are typical for some parts of India, like the SI and IGP, during the October–November season. However, we cannot neglect the likelihood of bias in the observations




Figure 7. Spatial distribution of the TROPOMI Sentinel-5P measurements, averaged for (a)-(g) each available month in 2018. Some months are excluded due to insufficient data points, due to filtering using the quality flag as given in the data product.




Figure 8. Same as Fig. 7, but showing WRF-GHG simulations of total XCH₄

due to high aerosol loads, which impacts XCH₄ retrievals (Lorente et al., 2021; Hu et al., 2018; Pandey et al., 2019). All the months show significantly higher total XCH₄ mixing ratios over the IGP region in comparison with the other regions over India. The IGP region is more prone to biomass burning in October and November, causing more aerosols in the region. We




Figure 9. Monthly distribution of the difference between WRF-GHG simulations and TROPOMI Sentinel-5p retrievals of XCH₄ for each available month in 2018 and 2019 when sufficient observations are available (Outliers are not removed; instead, a 90 % winsorization (Wilcox, 2005) is applied for the outliers.)

405 have examined the particulate matter (PM_{2.5}) content using the MERRA database, which indicates a heavier aerosol content due to burning during the winter, not always necessarily peaking in October but during the December-February season (Figure not shown). There is a gradual increase in XCH₄ mixing ratios beginning from the winter month of January till March, with a slight dip in April and then a distinctly high increment in October (exceeding 1870 ppb) in October over the IGP (see Fig. 7. (a)-(f)).

410 We find that the WRF simulations generally overestimate the total XCH₄ mixing ratio over the Indian region compared to TROPOMI observations, peaking in winter months (maximum at 1883 ppb; see Fig. 8). The IGP emission hotspot is also pronounced in the EDGAR inventory (Fig. S1, see Sect. 4.1), suggesting the large impact of anthropogenic emissions on the observed total XCH₄. Further, the sectoral analysis of the EDGAR emission inventory and the consistency of the spatial pattern with TROPOMI observations indicate that the enhancement over the IGP hotspot can be attributed to anthropogenic emissions

415 from the large cattle population and agricultural activities, especially rice production. Similarly, high XCH₄ values observed along the eastern coast during October and November can be attributed to the agricultural soil emissions, as seen in Fig. S1c. Wetland emissions also peak on the eastern coast, but the emissions are not found to be high enough to affect the mixing ratio enhancement significantly (see Fig. S9 & S12). Table S1 shows the mean observed XCH₄ and the variability over the entire study domain for the non-monsoon months of 2018 and 2019.

420 In general, the WRF-GHG simulations tend to show high bias in the winter months (see Fig. 9). A definite and widespread underestimation by the model was found in October 2018. However, in 2019, WRF-GHG was almost able to capture the




Figure 10. Model-Observation mismatches before and after optimization: (a) Probability density for difference between observed (Obs.) and simulated (Sim.) concentrations, (b) monthly Mean Bias, (c) monthly RMSE and (d) R^2 between observations and simulations before (blue) and after (green) optimization. In the legend, the terms Prior Sim. and Post Sim. refer to simulations before and after optimization, respectively.

observed XCH_4 . In the summer months, the model shows patterns of overestimation in eastern India and underestimation in western India. These regional differences in patterns of XCH_4 can arise from heterogeneous sectorial distributions of surface emissions with seasonality that would have been misrepresented in the inventories in conjunction with the large-scale meteorological influences (e.g. southwest monsoon over India, Chandra et al. (2017)).

While the peak total column XCH_4 for TROPOMI falls in October (~ 1870 ppb), that of the WRF-GHG simulations is in November (~ 1883 ppb). However, it is noteworthy that both the model and observation indicate the XCH_4 peak in either of the winter months between October and February. The WRF-GHG simulations show a higher variability (standard deviation) than observations for each month. WRF-GHG overestimates the XCH_4 values with a bias of ~ 13 (29) ppb in 2018 (2019)(see Table S1).

3.5 National CH₄ budget estimation via inverse optimization

In this section, we present estimates of India's anthropogenic CH₄ budget for the period 2018–2019 derived through inverse optimization as described in Sect. 3.2. Two separate inversions were performed using identical model configurations and observational constraints: one including biomass-burning emissions from GFAS in addition to anthropogenic sources, and
435 another excluding biomass-burning emissions. Posterior emission estimates from both inversions are reported separately to quantify the impact of biomass burning on inferred anthropogenic CH₄ emissions.

The model-observation mismatches before and after optimization are shown in Fig. 10. The optimization significantly reduces mismatches in XCH₄ as expected for a robust inversion, resulting in a narrower distribution around zero. For example, in January 2018, the mean XCH₄ bias improves from -12.4 ppb (before optimization) to 5.2 ppb (after optimization). Simultane-
440 ously, the improved explanation of variance in statistics is found, indicating a better fit to the observed data after optimization (see Fig. 10).

The EDGAR emission inventory reports an annual mean CH₄ emission budget of 28.8 Tg yr⁻¹, and we assumed 80% uncertainty (23 Tg yr⁻¹) in our prior as discussed in section 3.1. The posterior annual emission estimate is 23.4 Tg, with the uncertainty reduced to 3.3 Tg. The percentage of error reduction (calculated using Eq. 8) for monthly posterior fluxes ranges
445 from 68 to 92 %. Our inverse model results indicate an overestimation of 13 to 24 % in the EDGAR inventory. Incorporating biomass burning emissions from GFAS has an impact of +0.3 Tg yr⁻¹ on both prior and posterior emission estimates over the Indian region.

As per the India Fourth Biennial Update Report (BUR4) submitted to the United Nations Framework Convention on Climate Change (UNFCCC) (MoEFCC, 2024), the CH₄ emission budget for India is approximately 19.6 Tg yr⁻¹, which is around 32 %
450 less than the EDGAR-reported emissions during the 2018-2019 period. Previous studies also reported an overestimation of the global emission inventories over India. For instance, Qu et al. (2021) reports 41–57 Tg yr⁻¹ anthropogenic CH₄ emission from India, which is significantly higher than our estimations. Also, Zhang et al. (2021) estimates Indian anthropogenic methane emissions of 33±0.6 Tg yr⁻¹, higher than this study estimates. However, the Global Methane Budget (2000-2017, Saunio et al. (2019)), based on top-down approaches using in-situ and GOSAT observations, suggests 25 Tg yr⁻¹ of anthropogenic
455 CH₄ emissions from India, but acknowledges large uncertainty ranges in their estimates. Also, bottom-up models' estimates that are compiled in Saunio et al. (2019) and Jackson et al. (2020) indicate a mean anthropogenic CH₄ emission of 21-24 Tg yr⁻¹ from India. The above two estimates align with our results, though we used independent observations and a different modeling approach. The recent updates on the Global Methane Budget (2000-2020, Saunio et al. (2025)) indicate anthropogenic methane emissions of 37-49 Tg yr⁻¹ for South Asia (including Afghanistan, Bangladesh, Bhutan, India, Nepal, Pakistan, and

460 Sri Lanka), in which around 21.7 Tg yr^{-1} are contributed from the Indian region (calculated using the data prescribed from Martinez et al. (2024)). Janardanan et al. (2024) reported the annual averaged (2009 - 2020) CH_4 emissions from anthropogenic sectors over the India as $24.2 \pm 2.1 \text{ Tg yr}^{-1}$ which is close to our results. The total CH_4 emissions derived from a combination of satellite data (GOSAT), surface and aircraft measurements, and the atmospheric transport model for 2010–2015 were found to be 22 Tg yr^{-1} , which is substantially lower than the emissions reported by the EDGAR v4.2 inventory (Ganesan et al., 2017).
465 On the other hand, Raju et al. (2022) reported that the CH_4 budget for peninsular India is 0.13 Tg yr^{-1} higher than EDGAR v6.0 inventory-based estimates for the period 2017-2018. These variations in emission reports emphasize the need to improve CH_4 emission estimation in India using more regional-specific information and robust methodologies. Our findings also highlight that top-down evaluations of emissions inventories are critical for implementing effective climate change mitigation strategies in countries like India, which are largely understudied and undersampled, leading to poor quantification of their contributions
470 in the context of global climate policies.

Although our primary emphasis is on national-scale inversion estimates, the inversion framework explicitly resolves emissions at the state level, with each political state constituting an element of the state vector. The spatial distribution of flux adjustments is shown in Fig. S13, which illustrates how the inversion adjusts emissions across the states. Quantitative assessments of prior and posterior estimates are also presented region-wise in Fig. S13.

475 We analyzed three TROPOMI XCH_4 products: scientific (Schneising et al., 2023), operational (Copernicus Sentinel-5P, 2021), and GOSAT-blended (Balasus et al., 2023). The results are presented in Figs. 11 and S14-S16. The above product comparison indicates that differences among these datasets over the region lie mostly in the range of 3 to 6 ppb at the monthly scale. Fig. 11 also indicates smaller spatial differences among these products across the Indian region. While these differences do not exceed the unresolved modeling errors, they can still influence inverse estimates, especially when fluxes are retrieved
480 at finer scales. In our inverse setup as well, differences in the products can introduce additional uncertainty into estimates. For instance, using the increased measurement uncertainty as explained in Sect. 2.4.1, we find that the national-scale posterior uncertainty for 2018 increases from 3.5 Tg yr^{-1} in the baseline inversion to 4.4 Tg yr^{-1} . While the mean posterior emission estimates remained unchanged in our case, an increase in posterior uncertainty underscores the importance of addressing differences in satellite retrievals and their error characterization in inverse modeling.

485 While these mean differences are small across the region, the comparisons indicate considerable variation in spatial coverage between WFMD and Operational data products. For instance, Fig. S15 shows the spatial patterns of XCH_4 using the Operational product, but with comparatively sparse spatial coverage compared to those using the WFMD product (Fig. S14). Our analysis does not account for differences in spatial coverage between products, as illustrated in Figures S15 and S14. These differences

may potentially affect the inverse estimates, and assessing their impact requires further study. A recent study also reveals
490 considerable impacts of differences in retrievals on the inverse-based European CH₄ emission estimates when utilizing those
TROPOMI XCH₄ products (Sicsik-Paré et al., 2025). While a detailed performance assessment of different observational
products -regarding their coverage, retrieval errors, and spatial inconsistencies- is beyond the scope of this study, we emphasize
the necessity to investigate the robustness of satellite-based inversions considering these differences, particularly at a finer scale.
This focused approach will thus pave the way for more consistent emission estimates.

495 Although we utilize high-density, high-quality, and high-resolution TROPOMI satellite retrievals together with a high-
resolution transport model, our inversion algorithm is limited by its dependence on the spatial distribution of emissions in
prior inventories. Our optimization adjusts the magnitude of prior emissions over the target region by utilizing additional in-
formation from independent measurements, but the present inverse modeling design has the limitation to minimize any flux
errors in the sub-scale spatial distribution. However, we expect that those spatial errors may have a minor impact on our an-
500 nual national estimates owing to our temporal and spatial averaging. We also acknowledge that the resolution of the retrieved
emissions (as defined by state vectors) is a limitation when considering the information content that can be deduced from
TROPOMI observations. For instance, Sicsik-Paré et al. (2025) demonstrated inversions over Europe using TROPOMI ob-
servations to retrieve fluxes at 0.5° × 0.5° resolution. The above-mentioned limitation of the present study stems from the
region being undersampled and understudied against independent observations, as well as the need to address the risk of in-
505 ferring overly confident flux retrievals. To retrieve the maximum information content from TROPOMI observations without
compromising the accuracy of the retrieved fluxes, further improvements in the demonstrated inversion strategy are required.
This includes adequate characterization of both forward model and observational errors against independent observations, as
well as conducting sensitivity tests to examine the representativeness of observations to retrieve high-resolution fluxes. The
above tasks thus warrant increased observational coverage and advanced inverse modeling techniques that properly account
510 for such errors and sensitivities, which are currently limited over the region, not addressed in the study; thus, demand a future
investigation in this direction.

The present analysis excluded wetland emissions from the optimization due to their negligible contribution to column en-
hancements. This is demonstrated in Figs. S9 and S10, which show that the simulated wetland emission enhancements in
the column are significantly smaller than anthropogenic enhancements. As a result, TROPOMI observations cannot be utilized
515 with our model setup to reliably differentiate wetland emission signals for accurate wetland emission estimates. Approximately
7.5 Tg wetland CH₄ emissions from the Indian region were reported in the Global Methane Budget 2000-2020 (Saunio et al.,
2024). Janardanan et al. (2024) used wetland prior emissions from the Global Methane Budget 2000-2017 (Saunio et al., 2019)




Figure 11. Spatial distribution of the difference between TROPOMI retrievals of XCH₄ (WFMD - Operational) for different months of 2019. Some months are omitted due to poor data coverage caused by filtering based on cloud pixels.

in their inversions and reported approximately 3.8 ± 0.16 Tg CH₄ emissions annually from Indian wetlands. At the same time, the BUR4 report (MoEFCC, 2024) has not included the wetland emission estimates, possibly due to inadequate data coverage.

520 Bernard et al. (2025) also discussed the limitations in modeling the wetland emissions from the tropical region due to the inadequacy of available measurements. The above level of estimation discrepancies calls for a country-specific wetland inventory that can also be used as reliable prior fluxes in future inverse modeling. Also, there could be a possible overlapping of natural and anthropogenic (agricultural fields) wetlands in the emission inventories used, which may overestimate the sectoral contribution of posterior fluxes (Zhang et al., 2014).

525 Another limitation could be that though the GFAS inventory includes agricultural residue burning, small fires that are common in smallholders for clearing the wastes and field preparation can be missed from prior inventories, as reported in Deshpande et al. (2022). Also, in this study, our focus is restricted to providing national-scale anthropogenic CH₄ emission estimates. While the inversion framework in principle allows for analysis at finer spatial or sectoral scales, here we intentionally report only the aggregated national totals, as our aim is to evaluate the feasibility of TROPOMI–WRF-GHG for constraining India’s methane

530 budget. A more detailed exploration of sectoral and regional signals is left to future work with more coverage of observations and the implementation of more advanced inverse modeling methods.

4 Conclusions

In this study, we investigate the potential of TROPOMI satellite observations along with a high-resolution atmospheric transport model, WRF-GHG, to represent the distribution of CH₄ emissions over the Indian region. Analysis of the bottom-up inventories shows enteric fermentation as the most significant contributor to CH₄ emissions in India (42.9 %), followed by wastewater treatment (19.2 %), agricultural soil (12.4 %), fuel exploitation (6.7 %), and wetlands (5.2 %, excluding agriculture). The above proportions highlight the considerable impact of anthropogenic sources on CH₄ accumulation in the atmosphere. As expected, CH₄ emissions from rice agriculture (August), wetlands (July), and biomass burning (March) exhibit distinct seasonal patterns. The bottom-up anthropogenic CH₄ emissions, and consequently the total atmospheric XCH₄ mixing ratios, have shown some peaks over South India due to a few prominent emission hotspots. This study characterizes regional and seasonal methane-emission patterns from global bottom-up inventories and assesses their possible influence on XCH₄ enhancements. The analysis identifies key uncertainty drivers such as the elevated anthropogenic emissions in the post-monsoon months, thereby guiding refinement of top-down CH₄ estimates across India.

The WRF-GHG simulations of XCH₄ mixing ratio enhancements indicate considerable contributions from anthropogenic and biomass burning emissions, particularly in the IGP region (from 27 to 67 ppb). The highest seasonal enhancements of anthropogenic XCH₄ occur during winter, influenced by agricultural emissions, biomass burning, and atmospheric winter transport. This inference aligns with previous studies (eg. Patra et al. (2011)) that show stronger vertical mixing during the summer, associated with higher boundary layers and faster wind speeds, may impact CH₄ columns. Both the observed and modeled total XCH₄ show significant peaks over the IGP region, with values ranging from ~1862 to ~1870 ppb during October-November. Though WRF-GHG remarkably captures atmospheric XCH₄ patterns, simulations generally overestimate XCH₄ levels compared to TROPOMI. The total XCH₄ along the eastern coast reflects the influence of agricultural soil emissions on column-averaged methane. Although wetland emissions peak in this region, their contribution to atmospheric mixing ratios is negligible. Our high-resolution model is capable of capturing surface CH₄ variability, especially for the well-mixed conditions, as confirmed by the ground-based CH₄ observations. However, this comparison is representative of only one station, though it is a complicated measurement location to be represented by the model owing to the influence of coastal meteorology. Such ground-based observations across India are essential for evaluating the full potential of high-resolution models in representing the atmospheric distribution of trace gases and to better constrain vertical transport processes and regional representativeness.

The inversion analysis using our high-resolution model and TROPOMI observations reports an annual mean anthropogenic CH₄ emission budget of $23.4 \pm 3.3 \text{ Tg yr}^{-1}$ (excluding biomass burning of 0.3 Tg yr^{-1}). Our estimations are 13 to 24 % lower

560 than the EDGAR emission estimates. At the same time, our estimate is 19 % higher than what the Government of India reported to the UNFCCC for the same period, but close to the latest Global Methane Budget 2000-2020.

We emphasize the critical need for robust reporting of CH₄ emissions from the Indian region in global emission inventories. Achieving this requires an enhanced network of ground-based atmospheric trace gas measurements and advancements in satellite capabilities, alongside advanced modeling techniques with adequate model error characterization. With the above
565 expansion, future research can decisively explore and evaluate various inverse techniques. By implementing methods such as the Ensemble Kalman Filter (EnKF) and 4D Variational Inversion (4D-Var), we can effectively manage highly resolved state vectors, leading to significantly improved emissions data at much finer scales over India. Additionally, we recommend inter-comparisons of TROPOMI-based inversions using various inversion frameworks and transport models over India, with the aim of identifying biases in the forward models and the inversion frameworks. Further, we encourage rigorous sensitivity testing
570 with TROPOMI inversions to assess the robustness of derived emissions, particularly with respect to differences in satellite products, coverage and sampling, as these factors can significantly influence inverse-based estimates. Overall, our analyses highlight that TROPOMI observations can provide valuable insights into CH₄ emissions, and the WRF-GHG model has the potential to be used in an assimilation system to refine emissions.

Code and data availability. The anthropogenic CH₄ emission inventories used in this study are downloaded from https://edgar.jrc.ec.europa.eu/archived_datasets (last access: March 2024) (Crippa et al., 2024). CAMS global biomass burning emission based on fire radiative power (GFAS) is accessed from Copernicus Atmosphere Monitoring Service (CAMS) Atmosphere Data Store, DOI: 10.24381/a05253c7 (last access : November 2023). The global wetland CH₄ emissions, WetCHARTs v1.3.1 is prescribed from <https://daac.ornl.gov/cgi-bin/> (last access: November 2023) (Bloom et al., 2017). The WRF source code is freely available and can be accessed from https://www2.mmm.ucar.edu/wrf/users/download/get_source.html (last access: October 2023). The TROPOMI/WFMD v1.8 product is made available via
580 https://www.iup.uni-bremen.de/carbon_ghg/products/tropomi_wfmd/ (last access: January 2025). The blended TROPOMI+GOSAT methane product is available at <https://registry.opendata.aws/blended-tropomi-gosat-methane> (last access: October 2025) (Balasus et al., 2023). The operational Sentinel-5P/TROPOMI Level-2 methane product is available at https://sentinels.copernicus.eu/data-products/-/asset_publisher/fp37fc19FN8F/content/tropomi-level-2-methane (last access: October 2025) (Hu et al., 2016).

Author contributions. DP designed the study, TAM and DP performed the model simulations, raw data analysis, and postprocessing, and
585 wrote the initial version of the manuscript. JS, MVD, VT, and AR contributed to data curation and figure preparation. MB and OS contributed

to data archival and processing. SBK and AJV contributed to editing. SS, IAG, and SB contributed to the ground-based data collection and pre-processing. All authors contributed to the data analysis, interpretation and writing.

Competing interests. The corresponding author has declared that none of the authors has any competing interests.

Disclaimer. Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims made in the text, published maps, institutional affiliations, or any other geographical representation in this paper. While Copernicus Publications makes every effort to include appropriate place names, the final responsibility lies with the authors.

Acknowledgements. This study has been supported by funding from the Indian Ministry of Education and the Max Planck Society in Germany, which has been allocated to IISERB. The University of Bremen team acknowledges funding from ESA via the project GHG-CCI+ (ESA contract no. 4000126450/19/I-NB) and the Bundesministerium für Bildung und Forschung within its project ITMS (grant no. 01 LK2103A). We acknowledge the support of IISERB's high-performance cluster system for computations, data analysis, and visualization. The TROPOMI/WFMD retrievals were performed on HPC facilities funded by the Deutsche Forschungsgemeinschaft (grant nos. INST 144/379-1 FUGG and INST 144/493-1 FUGG). This publication contains modified Copernicus Sentinel data (2018–2019). Sentinel-5 Precursor is an ESA mission implemented on behalf of the European Commission. The TROPOMI payload is a joint development by ESA and the Netherlands Space Office (NSO). The Sentinel-5 Precursor ground segment development has been funded by ESA and with national contributions from the Netherlands, Germany, and Belgium. Thara Anna Mathew acknowledges the financial support provided by the Prime Minister's Research Fellowship (PMRF) Scheme, which funded her PhD fellowship. Jithin Sukumaran acknowledges the Council of Scientific and Industrial Research (CSIR) funding for his PhD fellowship. Imran A Girach acknowledges Prabha R Nair, former scientist at SPL, for supporting the surface trace gas measurements at Thumba utilized in this study. Special thanks to Navaneetha Jayan for the help with the graphics. We greatly appreciate the anonymous reviewers for their feedback, which helped in improving the initial version of the manuscript.

- Agarwal, R. and Garg, J.: Methane emission modelling from wetlands and waterlogged areas using MODIS data, *Current Science*, pp. 36–40, 2009.
- Agustí-Panareda, A., Barré, J., Massart, S., Inness, A., Aben, I., Ades, M., Baier, B. C., Balsamo, G., Borsdorff, T., Bousserez, N., et al.: The CAMS greenhouse gas reanalysis from 2003 to 2020, *Atmospheric Chemistry and Physics*, 23, 3829–3859, <https://doi.org/10.5194/acp-23-3829-2023>, 2023.
- Alexe, M., Bergamaschi, P., Segers, A., Detmers, R., Butz, A., Hasekamp, O., Guerlet, S., Parker, R., Boesch, H., Frankenberg, C., et al.: Inverse modelling of CH₄ emissions for 2010–2011 using different satellite retrieval products from GOSAT and SCIAMACHY, *Atmospheric Chemistry and Physics*, 15, 113–133, <https://doi.org/10.5194/acp-15-113-2015>, 2015.
- Anand, S., Dahiya, R., Talyan, V., and Vrat, P.: Investigations of methane emissions from rice cultivation in Indian context, *Environment International*, 31, 469–482, <https://doi.org/10.1016/j.envint.2004.10.016>, 2005.
- Baer, D. S., Paul, J. B., Gupta, M., and O’keefe, A.: Sensitive absorption measurements in the near-infrared region using off-axis integrated-cavity-output spectroscopy, *Applied Physics B*, 75, 261–265, <https://doi.org/10.1007/s00340-002-0971-z>, 2002.
- Balagus, N., Jacob, D. J., Lorente, A., Maasakkers, J. D., Parker, R. J., Boesch, H., Chen, Z., Kelp, M. M., Nesser, H., and Varon, D. J.: A blended TROPOMI+GOSAT satellite data product for atmospheric methane using machine learning to correct retrieval biases, *Atmospheric Measurement Techniques*, 16, 3787–3807, <https://doi.org/10.5194/amt-16-3787-2023>, 2023.
- Beck, V.: *The WRF Greenhouse Gas Model (WRF-GHG)*, 2011.
- Bergamaschi, P., Houweling, S., Segers, A., Krol, M., Frankenberg, C., Scheepmaker, R., Dlugokencky, E., Wofsy, S. C., Kort, E., Sweeney, C., et al.: Atmospheric CH₄ in the first decade of the 21st century: Inverse modeling analysis using SCIAMACHY satellite retrievals and NOAA surface measurements, *Journal of Geophysical Research: Atmospheres*, 118, 7350–7369, 2013.
- Bergamaschi, P., Karstens, U., Manning, A. J., Saunois, M., Tsuruta, A., Berchet, A., Vermeulen, A. T., Arnold, T., Janssens-Maenhout, G., Hammer, S., et al.: Inverse modelling of European CH₄ emissions during 2006–2012 using different inverse models and reassessed atmospheric observations, *Atmospheric Chemistry and Physics*, 18, 901–920, <https://doi.org/10.5194/acp-18-901-2018>, 2018.
- Bernard, J., Salmon, E., Saunois, M., Peng, S., Serrano-Ortiz, P., Berchet, A., Gnanamoorthy, P., Jansen, J., and Ciais, P.: Satellite-based modeling of wetland methane emissions on a global scale (SatWetCH₄ 1.0), *Geoscientific Model Development*, 18, 863–883, <https://doi.org/10.5194/egusphere-2024-1331>, 2025.
- Bloom, A., Bowman, K., Lee, M., Turner, A., Schroeder, R., Worden, J., Weidner, R., McDonald, K., and Jacob, D.: CMS: global 0.5-deg wetland methane emissions and uncertainty (WetCHARTs v1. 3.1), ORNL DAAC, <https://doi.org/10.3334/ORNLDAAC/1915>, 2021.
- Bloom, A. A., Bowman, K. W., Lee, M., Turner, A. J., Schroeder, R., Worden, J. R., Weidner, R., McDonald, K. C., and Jacob, D. J.: A global wetland methane emissions and uncertainty dataset for atmospheric chemical transport models (WetCHARTs version 1.0), *Geoscientific Model Development*, 10, 2141–2156, <https://doi.org/10.5194/gmd-10-2141-2017>, 2017.

- Buchwitz, M., Khlystova, I., Bovensmann, H., and Burrows, J.: Three years of global carbon monoxide from SCIAMACHY: comparison with MOPITT and first results related to the detection of enhanced CO over cities, *Atmospheric Chemistry and Physics*, 7, 2399–2411, <https://doi.org/10.5194/acp-7-2399-2007>, 2007.
- 640 Buchwitz, M., Schneising, O., Reuter, M., Heymann, J., Krautwurst, S., Bovensmann, H., Burrows, J. P., Boesch, H., Parker, R. J., Somkuti, P., et al.: Satellite-derived methane hotspot emission estimates using a fast data-driven method, *Atmospheric Chemistry and Physics*, 17, 5751–5774, <https://doi.org/10.5194/acp-17-5751-2017>, 2017.
- Buchwitz, M. d., De Beek, R., Burrows, J., Bovensmann, H., Warneke, T., Notholt, J., Meirink, J., Goede, A., Bergamaschi, P., Körner, S., et al.: Atmospheric methane and carbon dioxide from SCIAMACHY satellite data: initial comparison with chemistry and transport models, *Atmospheric Chemistry and Physics*, 5, 941–962, <https://doi.org/10.5194/acp-5-941-2005>, 2005.
- 645 Butz, A., Guerlet, S., Hasekamp, O., Schepers, D., Galli, A., Aben, I., Frankenberg, C., Hartmann, J.-M., Tran, H., Kuze, A., et al.: Toward accurate CO₂ and CH₄ observations from GOSAT, *Geophysical Research Letters*, 38, <https://doi.org/10.1029/2011GL047888>, 2011.
- Callewaert, S., Zhou, M., Langerock, B., Wang, P., Wang, T., Mahieu, E., and De Mazière, M.: A WRF-Chem study of the greenhouse gas column and in situ surface concentrations observed in Xianghe, China – Part 1: Methane (CH₄), *Atmospheric Chemistry and Physics*, 25, 9519–9544, <https://doi.org/10.5194/acp-25-9519-2025>, 2025.
- 650 Chandra, N., Hayashida, S., Saeki, T., and Patra, P. K.: What controls the seasonal cycle of columnar methane observed by GOSAT over different regions in India?, *Atmospheric Chemistry and Physics*, 17, 12 633–12 643, <https://doi.org/10.5194/acp-17-12633-2017>, 2017.
- Chen, Z., Jacob, D. J., Nesser, H., Sulprizio, M. P., Lorente, A., Varon, D. J., Lu, X., Shen, L., Qu, Z., Penn, E., et al.: Methane emissions from China: a high-resolution inversion of TROPOMI satellite observations, *Atmospheric Chemistry and Physics*, 22, 10 809–10 826, <https://doi.org/10.5194/acp-22-10809-2022>, 2022.
- 655 Copernicus Sentinel-5P, .: Copernicus Sentinel-5P (processed by ESA), 2021, TROPOMI Level 2 Methane Total Column products. Version 02. European Space Agency, <https://doi.org/https://doi.org/10.5270/S5P-3p6lnwd>, 2021.
- Crippa, M., Solazzo, E., Huang, G., Guizzardi, D., Koffi, E., Muntean, M., Schieberle, C., Friedrich, R., and Janssens-Maenhout, G.: High resolution temporal profiles in the Emissions Database for Global Atmospheric Research, *Scientific data*, 7, 121, <https://doi.org/10.6084/m9.figshare.12052887>, 2020.
- 660 Crippa, M., Guizzardi, D., Pagani, F., Schiavina, M., Melchiorri, M., Pisoni, E., Graziosi, F., Muntean, M., Maes, J., Dijkstra, L., et al.: Insights on the spatial distribution of global, national and sub-national GHG emissions in edgarv8. 0, *Earth System Science Data Discussions*, 2023, 1–28, <https://doi.org/10.5194/essd-16-2811-2024>, 2023.
- Crippa, M., Guizzardi, D., Pagani, F., Schiavina, M., Melchiorri, M., Pisoni, E., Graziosi, F., Muntean, M., Maes, J., Dijkstra, L., et al.: Insights into the spatial distribution of global, national, and subnational greenhouse gas emissions in the Emissions Database for Global Atmospheric Research (EDGAR v8. 0), *Earth System Science Data*, 16, 2811–2830, <https://doi.org/10.5194/essd-16-2811-2024>, 2024.
- 665

- Cusworth, D. H., Jacob, D. J., Sheng, J.-X., Benmergui, J., Turner, A. J., Brandman, J., White, L., and Randles, C. A.: Detecting high-emitting methane sources in oil/gas fields using satellite observations, *Atmospheric Chemistry and Physics*, 18, 16 885–16 896, <https://doi.org/10.5194/acp-18-16885-2018>, 2018.
- Das, N., Chakraborty, R., Pal, S. C., Mondal, A., and Mandal, S.: A novel coupled framework for detecting hotspots of methane emission from the vulnerable Indian Sundarban mangrove ecosystem using data-driven models, *Science of The Total Environment*, 866, 161 319, <https://doi.org/10.1016/j.scitotenv.2022.161319>, 2023.
- de Gouw, J., Veefkind, J., Roosenbrand, E., Dix, B., Lin, J., Landgraf, J., and Levelt, P.: Daily Satellite Observations of Methane from Oil and Gas Production Regions in the United States, *Scientific Reports*, 10, <https://doi.org/10.1038/s41598-020-57678-4>, 2020.
- Deshpande, M. V., Pillai, D., and Jain, M.: Detecting and quantifying residue burning in smallholder systems: An integrated approach using Sentinel-2 data, *International Journal of Applied Earth Observation and Geoinformation*, 108, 102 761, <https://doi.org/10.1016/j.jag.2022.102761>, 2022.
- Deshpande, M. V., Kumar, N., Pillai, D., Krishna, V. V., and Jain, M.: Greenhouse gas emissions from agricultural residue burning have increased by 75% since 2011 across India, *Science of the Total Environment*, 904, 166 944, <https://doi.org/10.1016/j.scitotenv.2023.166944>, 2023.
- Eskes, H. and Boersma, K.: Averaging kernels for DOAS total-column satellite retrievals, *Atmospheric Chemistry and Physics*, 3, 1285–1291, <https://doi.org/10.5194/acp-3-1285-2003>, 2003.
- Friedlingstein, P., O’sullivan, M., Jones, M. W., Andrew, R. M., Hauck, J., Landschützer, P., Le Quéré, C., Li, H., Luijkx, I. T., Olsen, A., et al.: Global carbon budget 2024, *Earth System Science Data Discussions*, 2024, 1–133, 2024.
- Ganesan, A. L., Rigby, M., Lunt, M. F., Parker, R. J., Boesch, H., Goulding, N., Umezawa, T., Zahn, A., Chatterjee, A., Prinn, R. G., et al.: Atmospheric observations show accurate reporting and little growth in India’s methane emissions, *Nature Communications*, 8, 836, <https://doi.org/DOI: 10.1038/s41467-017-00994-7>, 2017.
- Garg, A., Kankal, B., and Shukla, P.: Methane emissions in India: Sub-regional and sectoral trends, *Atmospheric environment*, 45, 4922–4929, <https://doi.org/10.1016/j.atmosenv.2011.06.004>, 2011.
- Gerbig, C., Lin, J. C., Munger, J. W., and Wofsy, S. C.: What can tracer observations in the continental boundary layer tell us about surface-atmosphere fluxes?, *Atmospheric Chemistry and Physics*, 6, 539–554, <https://doi.org/10.5194/acp-6-539-2006>, 2006.
- Guha, T., Tiwari, Y. K., Valsala, V., Lin, X., Ramonet, M., Mahajan, A., Datye, A., and Kumar, K. R.: What controls the atmospheric methane seasonal variability over India?, *Atmospheric Environment*, 175, 83–91, <https://doi.org/10.1016/j.atmosenv.2017.11.042>, 2018.
- Gururaj Katti, G. K., Pasalu, I., Rao, P., Varma, N., and Krishnaiah, K.: Farmer’s participatory approach to improve pest management decision making in high production systems of rice in Andhra Pradesh-a case study., <https://doi.org/DOI: 10.1080/14735903.2022.2070340>, 2002.
- Heald, C. L., Jacob, D. J., Jones, D. B., Palmer, P. I., Logan, J. A., Streets, D., Sachse, G. W., Gille, J. C., Hoffman, R. N., and Nehr Korn, T.: Comparative inverse analysis of satellite (MOPITT) and aircraft (TRACE-P) observations to estimate Asian sources of carbon monoxide, *Journal of Geophysical Research: Atmospheres*, 109, <https://doi.org/10.1029/2004JD005185>, 2004.

- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., et al.: The ERA5 global reanalysis, *Quarterly Journal of the Royal Meteorological Society*, 146, 1999–2049, <https://doi.org/10.1002/qj.3803>, 2020.
- 700 Hu, H., Landgraf, J., Detmers, R., Borsdorff, T., Aan de Brugh, J., Aben, I., Butz, A., and Hasekamp, O.: Toward global mapping of methane with TROPOMI: First results and intersatellite comparison to GOSAT, *Geophysical Research Letters*, 45, 3682–3689, <https://doi.org/10.1002/2018GL077259>, 2018.
- Inness, A., Ades, M., Agustí-Panareda, A., Barré, J., Benedictow, A., Blechschmidt, A.-M., Dominguez, J. J., Engelen, R., Eskes, H., Fleming, J., et al.: The CAMS reanalysis of atmospheric composition, *Atmospheric Chemistry and Physics*, 19, 3515–3556, 2019.
- 705 Jackson, R. B., Saunio, M., Bousquet, P., Canadell, J. G., Poulter, B., Stavert, A. R., Bergamaschi, P., Niwa, Y., Segers, A., and Tsuruta, A.: Increasing anthropogenic methane emissions arise equally from agricultural and fossil fuel sources, *Environmental Research Letters*, 15, 071 002, <https://doi.org/10.1088/1748-9326/ab9ed2>, 2020.
- Jacob, D. J., Turner, A. J., Maasackers, J. D., Sheng, J., Sun, K., Liu, X., Chance, K., Aben, I., McKeever, J., and Frankenberg, C.: Satellite observations of atmospheric methane and their value for quantifying methane emissions, *Atmospheric Chemistry and Physics*, 16, 14 371–710 14 396, <https://doi.org/10.5194/acp-16-14371-2016>, 2016.
- Jacob, D. J., Varon, D. J., Cusworth, D. H., Dennison, P. E., Frankenberg, C., Gautam, R., Guanter, L., Kelley, J., McKeever, J., Ott, L. E., et al.: Quantifying methane emissions from the global scale down to point sources using satellite observations of atmospheric methane, *Atmospheric Chemistry and Physics*, 22, 9617–9646, <https://doi.org/10.5194/acp-22-9617-2022>, 2022.
- Janardanan, R., Maksyutov, S., Wang, F., Nayagam, L., Sahu, S. K., Mangaraj, P., Saunio, M., Lan, X., and Matsunaga, T.: Country-level 715 methane emissions and their sectoral trends during 2009–2020 estimated by high-resolution inversion of GOSAT and surface observations, *Environmental Research Letters*, 19, 034 007, <https://doi.org/DOI.10.1088/1748-9326/ad2436>, 2024.
- Janssens-Maenhout, G., Crippa, M., Guizzardi, D., Muntean, M., and Schaaf, E.: Emissions Database for Global Atmospheric Research (EDGAR), version v4.2 (time-series), <http://data.europa.eu/89h/jrc-edgar-emissiontimeseriesv42>, 2011.
- Kaiser, J., Heil, A., Andreae, M., Benedetti, A., Chubarova, N., Jones, L., Morcrette, J.-J., Razinger, M., Schultz, M., Suttie, M., et al.: 720 Biomass burning emissions estimated with a global fire assimilation system based on observed fire radiative power, *Biogeosciences*, 9, 527–554, <https://doi.org/10.5194/bg-9-527-2012>, 2012.
- Kavitha, M., Nair, P. R., Girach, I., Aneesh, S., Sijikumar, S., and Renju, R.: Diurnal and seasonal variations in surface methane at a tropical coastal station: Role of mesoscale meteorology, *Science of The Total Environment*, 631–632, 1472–1485, <https://doi.org/10.1016/j.scitotenv.2018.03.123>, 2018.
- 725 Kretschmer, R., Gerbig, C., Karstens, U., Biavati, G., Vermeulen, A., Vogel, F., Hammer, S., and Totsche, K.: Impact of optimized mixing heights on simulated regional atmospheric transport of CO₂, *Atmospheric Chemistry and Physics*, 14, 7149–7172, <https://doi.org/10.5194/acp-14-7149-2014>, 2014.
- Kuhlmann, G., Brunner, D., Broquet, G., and Meijer, Y.: Quantifying CO₂ emissions of a city with the Copernicus Anthropogenic CO₂ Monitoring satellite mission, *Atmospheric Measurement Techniques*, 13, 6733–6754, 2020.

- 730 Lee, H., Calvin, K., Dasgupta, D., Krinner, G., Mukherji, A., Thorne, P., Trisos, C., Romero, J., Aldunce, P., Barret, K., et al.: IPCC, 2023: Climate Change 2023: Synthesis Report, Summary for Policymakers. Contribution of Working Groups I, II and III to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change [Core Writing Team, H. Lee and J. Romero (eds.)]. IPCC, Geneva, Switzerland., 2023.
- Liang, R., Zhang, Y., Chen, W., Zhang, P., Liu, J., Chen, C., Mao, H., Shen, G., Qu, Z., Chen, Z., et al.: East Asian methane emissions inferred
735 from high-resolution inversions of GOSAT and TROPOMI observations: a comparative and evaluative analysis, *Atmospheric Chemistry and Physics*, 23, 8039–8057, <https://doi.org/10.5194/acp-23-8039-2023>, 2023.
- Lin, X., Indira, N., Ramonet, M., Delmotte, M., Ciais, P., Bhatt, B., Reddy, M., Angchuk, D., Balakrishnan, S., Jorphail, S., et al.: Long-lived atmospheric trace gases measurements in flask samples from three stations in India, *Atmospheric Chemistry and Physics*, 15, 9819–9849, <https://doi.org/10.5194/acp-15-9819-2015>, 2015.
- 740 Lorente, A., Borsdorff, T., Butz, A., Hasekamp, O., Schneider, A., Wu, L., Hase, F., Kivi, R., Wunch, D., Pollard, D. F., et al.: Methane retrieved from TROPOMI: improvement of the data product and validation of the first 2 years of measurements, *Atmospheric Measurement Techniques*, 14, 665–684, <https://doi.org/10.5194/amt-14-665-2021>, 2021.
- Lu, X., Jacob, D. J., Wang, H., Maasakkers, J. D., Zhang, Y., Scarpelli, T. R., Shen, L., Qu, Z., Sulprizio, M. P., Nesser, H., et al.: Methane
745 emissions in the United States, Canada, and Mexico: evaluation of national methane emission inventories and 2010–2017 sectoral trends by inverse analysis of in situ (GLOBALVIEWplus CH 4 ObsPack) and satellite (GOSAT) atmospheric observations, *Atmospheric Chemistry and Physics*, 22, 395–418, <https://doi.org/10.5194/acp-22-395-2022>, 2022.
- Maasakkers, J. D., Jacob, D. J., Sulprizio, M. P., Scarpelli, T. R., Nesser, H., Sheng, J.-X., Zhang, Y., Hersher, M., Bloom, A. A., Bowman, K. W., et al.: Global distribution of methane emissions, emission trends, and OH concentrations and trends inferred from an inversion of GOSAT satellite data for 2010–2015, *Atmospheric Chemistry and Physics*, 19, 7859–7881, <https://doi.org/10.5194/acp-19-7859-2019>,
750 2019.
- Madrazo, J., Clappier, A., Belalcazar, L. C., Cuesta, O., Contreras, H., and Golay, F.: Screening differences between a local inventory and the Emissions Database for Global Atmospheric Research (EDGAR), *Science of the Total Environment*, 631, 934–941, <https://doi.org/10.1016/j.scitotenv.2018.03.094>, 2018.
- Manjunath, K., Panigrahy, S., Kumari, K., Adhya, T., and Parihar, J.: Spatiotemporal modelling of methane flux from the rice fields of India
755 using remote sensing and GIS, *International Journal of Remote Sensing*, 27, 4701–4707, <https://doi.org/10.1080/01431160600702350>, 2006.
- Martinez, A., Saunio, M., Poulter, B., Bousquet, P., Canadell, J. G., Jackson, R. B., Dlugokencky, E. J., Ciais, P., Bastviken, D., Blake, D. R., Castaldi, S., Etiope, G., Gedney, N., Höglund-Isaksson, L., Hugelius, G., Ito, A., Kleinen, T., Krummel, P. B., Liu, L., McDonald, K. C., Melton, J. R., Müller, J., Murguia-Flores, F., Niwa, Y., Noce, S., Parker, R. J., Peng, C., Ramonet, M., Riley, W. J., Rosentreter, J. A., Segers, A., Smith, S. J., Tian, H., Tubiello, F. N., Tsuruta, A., Weber, T. S., Werf, G. R. v. d., Worthy, D., Yoshida, Y., Zhang, W.,
760 Zhang, Z., Zheng, B., Zhu, Q., Zhu, Q., and Zhuang, Q.: Supplemental data of the Global Carbon Project methane budget 2024 v1, 2024.

- Metya, A., Datye, A., Chakraborty, S., Tiwari, Y. K., Sarma, D., Bora, A., and Gogoi, N.: Diurnal and seasonal variability of CO₂ and CH₄ concentration in a semi-urban environment of western India, *Scientific reports*, 11, 2931, <https://doi.org/10.1038/s41598-021-82321-1>, 2021.
- 765 Miller, S. M. and Michalak, A. M.: Constraining sector-specific CO₂ and CH₄ emissions in the US, *Atmospheric Chemistry and Physics*, 17, 3963–3985, <https://doi.org/10.5194/acp-17-3963-2017>, 2017.
- Miller, S. M., Michalak, A. M., Detmers, R. G., Hasekamp, O. P., Bruhwiler, L. M., and Schwietzke, S.: China's coal mine methane regulations have not curbed growing emissions, *Nature Communications*, 10, 303, <https://doi.org/10.1038/s41467-018-07891-7>, 2019.
- Ministry of Environment, F. and Change, C.: India: First biennial update report to the United Nations framework convention on climate change, 2015.
- 770 MoEFCC, .: India: Fourth Biennial update report to the United Nations Framework Convention on Climate Change, Ministry of Environment, Forest and Climate Change, Government of India., 2024.
- Monteil, G., Houweling, S., Dlugokenky, E., Maenhout, G., Vaughn, B., White, J., and Rockmann, T.: Interpreting methane variations in the past two decades using measurements of CH₄ mixing ratio and isotopic composition, *Atmospheric Chemistry and Physics*, 11, 9141–9153, 2011.
- 775 Montzka, S. A., Dlugokenky, E. J., and Butler, J. H.: Non-CO₂ greenhouse gases and climate change, *Nature*, 476, 43–50, <https://doi.org/doi:10.1038/nature10322>, 2011.
- Munassar, S., Monteil, G., Scholze, M., Karstens, U., Rödenbeck, C., Koch, F.-T., Totsche, K. U., and Gerbig, C.: Why do inverse models disagree? A case study with two European CO₂ inversions, *Atmospheric Chemistry and Physics*, 23, 2813–2828, <https://doi.org/10.5194/acp-23-2813-2023>, 2023.
- 780 Myhre, G., Myhre, C. L., Samset, B., and Storelvmo, T.: Aerosols and their relation to global climate and climate sensitivity, *Nature Education Knowledge*, 4, 7, 2013a.
- Myhre, G., Shindell, D., Bréon, F.-M., Collins, W., Fuglestedt, J., Huang, J., Koch, D., Lamarque, J.-F., Lee, D., Mendoza, B., Nakajima, T., Robock, A., Stephens, G., Takemura, T., and Zhang, H.: Anthropogenic and natural radiative forcing, pp. 659–740, Cambridge University Press, Cambridge, UK, <https://doi.org/10.1017/CBO9781107415324.018>, 2013b.
- 785 Nesser, H., Jacob, D. J., Maasackers, J. D., Lorente, A., Chen, Z., Lu, X., Shen, L., Qu, Z., Sulprizio, M. P., Winter, M., Ma, S., Bloom, A. A., Worden, J. R., Stavins, R. N., and Randles, C. A.: High-resolution US methane emissions inferred from an inversion of 2019 TROPOMI satellite data: contributions from individual states, urban areas, and landfills, *Atmospheric Chemistry and Physics*, 24, 5069–5091, <https://doi.org/10.5194/acp-24-5069-2024>, 2024.
- 790 Nisbet, E. G., Manning, M., Dlugokenky, E., Fisher, R., Lowry, D., Michel, S., Myhre, C. L., Platt, S. M., Allen, G., Bousquet, P., et al.: Very strong atmospheric methane growth in the 4 years 2014–2017: Implications for the Paris Agreement, *Global Biogeochemical Cycles*, 33, 318–342, <https://doi.org/10.1029/2018GB006009>, 2019.

- Palmer, P. I., Feng, L., Lunt, M. F., Parker, R. J., Bösch, H., Lan, X., Lorente, A., and Borsdorff, T.: The added value of satellite observations of methane for understanding the contemporary methane budget, *Philosophical Transactions of the Royal Society A*, 379, 20210106, <https://doi.org/10.1098/rsta.2021.0106>, 2021.
- 795 Pandey, S., Gautam, R., Houweling, S., Van Der Gon, H. D., Sadavarte, P., Borsdorff, T., Hasekamp, O., Landgraf, J., Tol, P., Van Kempen, T., et al.: Satellite observations reveal extreme methane leakage from a natural gas well blowout, *Proceedings of the National Academy of Sciences*, 116, 26376–26381, <https://doi.org/10.1073/pnas.1908712116>, 2019.
- Panigrahy, S., Upadhyay, G., Ray, S. S., and Parihar, J. S.: Mapping of cropping system for the Indo-Gangetic plain using multi-date SPOT NDVI-VGT data, *Journal of the Indian Society of Remote Sensing*, 38, 627–632, <https://doi.org/10.1007/s12524-011-0059-5>, 2010.
- 800 Patra, P. K., Houweling, S., Krol, M., Bousquet, P., Belikov, D., Bergmann, D., Bian, H., Cameron-Smith, P., Chipperfield, M. P., Corbin, K., et al.: TransCom model simulations of CH₄ and related species: linking transport, surface flux and chemical loss with CH₄ variability in the troposphere and lower stratosphere, *Atmospheric Chemistry and Physics*, 11, 12813–12837, 2011.
- Patra, P. K., Saeki, T., Dlugokencky, E. J., Ishijima, K., Umezawa, T., Ito, A., Aoki, S., Morimoto, S., Kort, E. A., Crotwell, A., et al.: Regional methane emission estimation based on observed atmospheric concentrations (2002–2012), *Journal of the Meteorological Society of Japan. Ser. II*, 94, 91–113, <https://doi.org/10.2151/jmsj.2016-006>, 2016.
- 805 Pillai, D., Gerbig, C., Kretschmer, R., Beck, V., Karstens, U., Neininger, B., and Heimann, M.: Comparing Lagrangian and Eulerian models for CO₂ transport – a step towards Bayesian inverse modeling using WRF/STILT-VPRM, *Atmospheric Chemistry and Physics*, 12, 8979–8991, <https://doi.org/10.5194/acp-12-8979-2012>, 2012.
- 810 Pillai, D., Buchwitz, M., Gerbig, C., Koch, T., Reuter, M., Bovensmann, H., Marshall, J., and Burrows, J. P.: Tracking city CO₂ emissions from space using a high-resolution inverse modelling approach: a case study for Berlin, Germany, *Atmospheric Chemistry and Physics*, 16, 9591–9610, <https://doi.org/10.5194/acp-16-9591-2016>, 2016.
- Qu, Z., Jacob, D. J., Shen, L., Lu, X., Zhang, Y., Scarpelli, T. R., Nesser, H., Sulprizio, M. P., Maasakkers, J. D., Bloom, A. A., Worden, J. R., Parker, R. J., and Delgado, A. L.: Global distribution of methane emissions: a comparative inverse analysis of observations from the TROPOMI and GOSAT satellite instruments, *Atmospheric Chemistry and Physics*, 21, 14159–14175, <https://doi.org/10.5194/acp-21-14159-2021>, 2021.
- 815 Raju, A., Sijikumar, S., Valsala, V., Tiwari, Y. K., Halder, S., Girach, I., Jain, C. D., and Ratnam, M. V.: Regional estimation of methane emissions over the peninsular India using atmospheric inverse modelling, *Environmental Monitoring and Assessment*, 194, 647, <https://doi.org/10.1007/s10661-022-10323-1>, 2022.
- 820 Ramasamy, C. and Manivel, S.: An analysis of aspects of performance and difficulties of poultry farming in Namakkal, Tamilnadu, 2019.
- Robinson, T. P., Wint, G. R. W., Conchedda, G., Van Boeckel, T. P., Ercoli, V., Palamara, E., Cinardi, G., D’Aielli, L., Hay, S. I., and Gilbert, M.: Mapping the Global Distribution of Livestock, *PLoS ONE*, 9, e96084, <https://doi.org/10.1371/journal.pone.0096084>, 2014.
- Rodgers, C. D.: Inverse methods for atmospheric sounding: theory and practice, vol. 2, World scientific, <https://doi.org/10.1142/3171>, 2000.

- Saunois, M., Bousquet, P., Poulter, B., Peregón, A., Ciais, P., Canadell, J. G., Dlugokencky, E. J., Etiope, G., Bastviken, D., Houweling, S.,
825 et al.: The global methane budget: 2000–2012, *Earth System Science Data Discussions*, 2016, 1–79, <https://doi.org/10.5194/essd-8-697-2016>, 2016.
- Saunois, M., Stavert, A. R., Poulter, B., Bousquet, P., Canadell, J. G., Jackson, R. B., Raymond, P. A., Dlugokencky, E. J.,
Houweling, S., Patra, P. K., et al.: The global methane budget 2000–2017, *Earth System Science Data Discussions*, 2019, 1–136,
<https://doi.org/10.5194/essd-12-1561-2020>, 2019.
- 830 Saunois, M., Martinez, A., Poulter, B., Zhang, Z., Raymond, P., Regnier, P., Canadell, J. G., Jackson, R. B., Patra, P. K., Bousquet, P., et al.:
Global methane budget 2000–2020, *Earth System Science Data Discussions*, 2024, 1–147, <https://doi.org/10.5194/essd-2024-115>, 2024.
- Saunois, M., Martinez, A., Poulter, B., Zhang, Z., Raymond, P. A., Regnier, P., Canadell, J. G., Jackson, R. B., Patra, P. K., Bousquet, P.,
Ciais, P., Dlugokencky, E. J., Lan, X., Allen, G. H., Bastviken, D., Beerling, D. J., Belikov, D. A., Blake, D. R., Castaldi, S., Crippa,
M., Deemer, B. R., Dennison, F., Etiope, G., Gedney, N., Höglund-Isaksson, L., Holgerson, M. A., Hopcroft, P. O., Hugelius, G., Ito, A.,
835 Jain, A. K., Janardanan, R., Johnson, M. S., Kleinen, T., Krummel, P. B., Lauerwald, R., Li, T., Liu, X., McDonald, K. C., Melton, J. R.,
Mühle, J., Müller, J., Murguía-Flores, F., Niwa, Y., Noce, S., Pan, S., Parker, R. J., Peng, C., Ramonet, M., Riley, W. J., Rocher-Ros, G.,
Rosentreter, J. A., Sasakawa, M., Segers, A., Smith, S. J., Stanley, E. H., Thanwerdas, J., Tian, H., Tsuruta, A., Tubiello, F. N., Weber,
T. S., van der Werf, G. R., Worthy, D. E. J., Xi, Y., Yoshida, Y., Zhang, W., Zheng, B., Zhu, Q., Zhu, Q., and Zhuang, Q.: Global Methane
Budget 2000–2020, *Earth System Science Data*, 17, 1873–1958, <https://doi.org/10.5194/essd-17-1873-2025>, 2025.
- 840 Scarpelli, T. R., Roy, E., Jacob, D. J., Sulprizio, M. P., Tate, R. D., and Cusworth, D. H.: Using new geospatial data and 2020 fossil
fuel methane emissions for the Global Fuel Exploitation Inventory (GFEI) v3, *Earth System Science Data Discussions*, 2025, 1–23,
<https://doi.org/10.5194/essd-2024-552>, 2025.
- Schaefer, H., Fletcher, S. E. M., Veidt, C., Lassey, K. R., Brailsford, G. W., Bromley, T. M., Dlugokencky, E. J., Michel, S. E., Miller,
J. B., Levin, I., et al.: A 21st-century shift from fossil-fuel to biogenic methane emissions indicated by $^{13}\text{CH}_4$, *Science*, 352, 80–84,
845 [https://doi.org/DOI: 10.1126/science.aad2705](https://doi.org/DOI:10.1126/science.aad2705), 2016.
- Schneising, O.: Product User Guide (PUG) TROPOMI WFM-DOAS (TROPOMI/WFMD) XCH_4 , [https://admin.climate.esa.int/media/
documents/PUG_CRDP9_v2_GHG-CCI_CH4_S5P_WFMD_v1.8.pdf](https://admin.climate.esa.int/media/documents/PUG_CRDP9_v2_GHG-CCI_CH4_S5P_WFMD_v1.8.pdf), 2024.
- Schneising, O., Buchwitz, M., Reuter, M., Heymann, J., Bovensmann, H., and Burrows, J. P.: Long-term analysis of carbon dioxide
and methane column-averaged mole fractions retrieved from SCIAMACHY, *Atmospheric Chemistry and Physics*, 11, 2863–2880,
850 <https://doi.org/10.5194/acp-11-2863-2011>, 2011.
- Schneising, O., Buchwitz, M., Reuter, M., Bovensmann, H., Burrows, J. P., Borsdorff, T., Deutscher, N. M., Feist, D. G., Griffith, D. W., Hase,
F., et al.: A scientific algorithm to simultaneously retrieve carbon monoxide and methane from TROPOMI onboard Sentinel-5 Precursor,
Atmospheric Measurement Techniques, 12, 6771–6802, <https://doi.org/10.5194/amt-12-6771-2019>, 2019.

- Schneising, O., Buchwitz, M., Reuter, M., Vanselow, S., Bovensmann, H., and Burrows, J. P.: Remote sensing of methane leakage from natural gas and petroleum systems revisited, *Atmospheric Chemistry and Physics*, 20, 9169–9182, <https://doi.org/10.5194/acp-20-9169-2020>, 2020.
- Schneising, O., Buchwitz, M., Hachmeister, J., Vanselow, S., Reuter, M., Buschmann, M., Bovensmann, H., and Burrows, J. P.: Advances in retrieving XCH₄ and XCO from Sentinel-5 Precursor: improvements in the scientific TROPOMI/WFMD algorithm, *Atmospheric Measurement Techniques*, 16, 669–694, <https://doi.org/10.5194/amt-16-669-2023>, 2023.
- Sicsik-Paré, A., Fortems-Cheiney, A., Pison, I., Broquet, G., Opler, A., Potier, E., Martinez, A., Schneising, O., Buchwitz, M., Maasackers, J. D., et al.: Can we obtain consistent estimates of the emissions in Europe from three different CH₄ TROPOMI products?, *EGU sphere*, 2025, 1–48, 2025.
- Sijkumar, S., Raju, A., Valsala, V., Tiwari, Y., Girach, I., Jain, C. D., and Ratnam, M. V.: High-Resolution Bayesian Inversion of Carbon Dioxide Flux Over Peninsular India, *Atmospheric Environment*, 308, 119 868, <https://doi.org/10.1016/j.atmosenv.2023.119868>, 2023.
- Skamarock, W. C., Klemp, J. B., Dudhia, J., Gill, D. O., Barker, D. M., Duda, M. G., Huang, X., Wang, W., and Powers, J. G.: A description of the advanced research WRF, National Center for Atmospheric Research, Boulder, CO), Version, 3, <https://doi.org/10.5065/D68S4MVH>, 2008.
- Skeie, R. B., Hodnebrog, Ø., and Myhre, G.: Trends in atmospheric methane concentrations since 1990 were driven and modified by anthropogenic emissions, *Communications Earth & Environment*, 4, 317, <https://doi.org/10.1038/s43247-023-00969-1>, 2023.
- Solazzo, E., Crippa, M., Guizzardi, D., Muntean, M., Choulga, M., and Janssens-Maenhout, G.: Uncertainties in the Emissions Database for Global Atmospheric Research (EDGAR) emission inventory of greenhouse gases, *Atmospheric Chemistry and Physics*, 21, 5655–5683, <https://doi.org/10.5194/acp-21-5655-2021>, 2021.
- Stevenson, D. S., Zhao, A., Naik, V., O’connor, F. M., Tilmes, S., Zeng, G., Murray, L. T., Collins, W. J., Griffiths, P. T., Shim, S., et al.: Trends in global tropospheric hydroxyl radical and methane lifetime since 1850 from AerChemMIP, *Atmospheric Chemistry and Physics*, 20, 12 905–12 920, <https://doi.org/10.5194/acp-20-12905-2020>, 2020.
- Stocker, T. F., Qin, D., Plattner, G., Tignor, M., Allen, S., Boschung, J., Nauels, A., Xia, Y., Bex, V., Midgley, P., et al.: Contribution of working group I to the fifth assessment report of the intergovernmental panel on climate change, *Climate change*, 5, 1–1552, <https://doi.org/10.1017/CBO9781107415324.004>, 2013.
- Survey of India: Political map of India, <https://www.surveyofindia.gov.in/pages/political-map-of-india>, accessed on: 21 May 2024, 2024.
- Thilakan, V., Pillai, D., Gerbig, C., Galkowski, M., Ravi, A., and Anna Mathew, T.: Towards monitoring CO₂ source-sink distribution over India via inverse modelling: Quantifying the fine-scale spatiotemporal variability of atmospheric CO₂ mole fraction, *Atmospheric Chemistry and Physics Discussions*, 2022, 1–38, <https://doi.org/10.5194/acp-22-15287-2022>, 2022.
- Thompson, R. L., Montzka, S. A., Vollmer, M. K., Arduini, J., Crotwell, M., Krummel, P. B., Lunder, C., Mühle, J., O’doherly, S., Prinn, R. G., et al.: Estimation of the atmospheric hydroxyl radical oxidative capacity using multiple hydrofluorocarbons (HFCs), *Atmospheric Chemistry and Physics*, 24, 1415–1427, 2024.

- Turner, A., Jacob, D. J., Wecht, K. J., Maasackers, J. D., Lundgren, E., Andrews, A. E., Biraud, S. C., Boesch, H., Bowman, K. W., Deutscher, N. M., et al.: Estimating global and North American methane emissions with high spatial resolution using GOSAT satellite data, *Atmospheric Chemistry and Physics*, 15, 7049–7069, <https://doi.org/10.5194/acp-15-7049-2015>, 2015.
- Uma, K., Girach, I. A., Chandra, N., Patra, P. K., Kumar, N. K., and Nair, P. R.: CO₂ variability over a tropical coastal station in India: Synergy of observation and model, *Science of The Total Environment*, 957, 177 371, <https://doi.org/10.1016/j.scitotenv.2024.177371>, 2024.
- Vellalassery, A., Pillai, D., Marshall, J., Gerbig, C., Buchwitz, M., Schneising, O., and Ravi, A.: Using TROPOspheric Monitoring Instrument (TROPOMI) measurements and Weather Research and Forecasting (WRF) CO modelling to understand the contribution of meteorology and emissions to an extreme air pollution event in India, *Atmospheric Chemistry and Physics*, 21, 5393–5414, <https://doi.org/10.5194/acp-21-5393-2021>, 2021.
- Wang, Y., Yuan, Q., Li, T., Yang, Y., Zhou, S., and Zhang, L.: Seamless mapping of long-term (2010–2020) daily global XCO₂ and XCH₄ from the Greenhouse Gases Observing Satellite (GOSAT), Orbiting Carbon Observatory 2 (OCO-2), and CAMS global greenhouse gas reanalysis (CAMS-EGG4) with a spatiotemporally self-supervised fusion method, *Earth System Science Data*, 15, 3597–3622, <https://doi.org/10.5194/essd-15-3597-2023>, 2023.
- Wang, Z., Warneke, T., Deutscher, N. M., Notholt, J., Karstens, U., Saunio, M., Schneider, M., Sussmann, R., Sembhi, H., Griffith, D. W. T., Pollard, D. F., Kivi, R., Petri, C., Velazco, V. A., Ramonet, M., and Chen, H.: Contributions of the troposphere and stratosphere to CH₄ model biases, *Atmospheric Chemistry and Physics*, 17, 13 283–13 295, <https://doi.org/10.5194/acp-17-13283-2017>, 2017.
- Ware, J., Kort, E. A., Duren, R., Mueller, K. L., Verhulst, K., and Yadav, V.: Detecting Urban Emissions Changes and Events With a Near-Real-Time-Capable Inversion System, *Journal of Geophysical Research: Atmospheres*, 124, 5117–5130, <https://doi.org/10.1029/2018jd029224>, 2019.
- Wilcox, R.: *Trimming and Winsorization*, John Wiley Sons, Ltd, <https://doi.org/10.1002/0470011815.b2a15165>, 2005.
- Ye, X., Lauvaux, T., Kort, E. A., Oda, T., Feng, S., Lin, J. C., Yang, E. G., and Wu, D.: Constraining fossil fuel CO₂ emissions from urban area using OCO-2 observations of total column CO₂, *Journal of Geophysical Research: Atmospheres*, 125, e2019JD030 528, 2020.
- Yokota, T., Yoshida, Y., Eguchi, N., Ota, Y., Tanaka, T., Watanabe, H., and Maksyutov, S.: Global concentrations of CO₂ and CH₄ retrieved from GOSAT: First preliminary results, *Sola*, 5, 160–163, <https://doi.org/10.2151/sola.2009-041>, 2009.
- Zhang, X., Lee, X., Griffis, T. J., Baker, J. M., and Xiao, W.: Estimating regional greenhouse gas fluxes: an uncertainty analysis of planetary boundary layer techniques and bottom-up inventories, *Atmospheric chemistry and physics*, 14, 10 705–10 719, <https://doi.org/10.5194/acp-14-10705-2014>, 2014.
- Zhang, Y., Jacob, D. J., Lu, X., Maasackers, J. D., Scarpelli, T. R., Sheng, J.-X., Shen, L., Qu, Z., Sulprizio, M. P., Chang, J., Bloom, A. A., Ma, S., Worden, J., Parker, R. J., and Boesch, H.: Attribution of the accelerating increase in atmospheric methane during 2010–2018 by inverse analysis of GOSAT observations, *Atmospheric Chemistry and Physics*, 21, 3643–3666, <https://doi.org/10.5194/acp-21-3643-2021>, 2021.

Zhou, L., Warner, J., Nalli, N. R., Wei, Z., Oh, Y., Bruhwiler, L., Liu, X., Divakarla, M., Pryor, K., Kalluri, S., et al.: Spatiotemporal variability of global atmospheric methane observed from two decades of satellite hyperspectral infrared sounders, *Remote Sensing*, 15, 2992, 2023.