

GC Insights: “Sedimentary Rock!” – A web app that converts geological strata data into music

Toshiyuki Kitazawa¹, Kazuaki Aoki¹, Hiroaki Yasutomo², Takuma Kato²

¹Faculty of Geo-Environmental Science, Risho University, Kumagaya, Saitama, 360-0194, Japan

²SHIFTBRAIN Inc., Setagaya, Tokyo, 155-0033, Japan

Correspondence to: Toshiyuki Kitazawa (kitazawa@ris.ac.jp)

Abstract. A web app called “Sedimentary Rock!” has been developed to reveal geological strata to human hearing and rhythmic senses, much like music, in a straightforward manner. The web app was designed and opened for non-specialists in geology. A columnar section created on the app is converted to notes and played. The thickness of a layer is converted to the length of a note, and the grain size is converted to pitch level. We converted schematic columnar sections of several environments and real Pleistocene strata from Vietnam into music and surveyed the educational effect of the web app using questionnaires. Sedimentary Rock! makes learners understand that the strata consist of concepts of grain size and thickness of stratum, and provides a new intuitive experience to listen to geological history.

Formatted: Font: Bold

Deleted: similar to

Deleted: way

1 Introduction

Geological strata were formed by the upward stacking of sediments and record the geological history, the changes in time of the local or global environment. The most essential feature of strata is grain size, which is divided into gravel, sand, and mud with more detailed subdivisions, indicating the strength of the current that caused the deposition. Another is the thickness of the bed, which means water depth, sediment discharge, time length, etc. Primary information obtained from geological strata is generally interpreted visually. However, methods other than visual inspection can be used to interpret geological strata, such as verifying grain size by touch with the fingers and through physical surveys where strata cannot be directly observed underground. New techniques and perspectives can interpret strata, which may lead to discoveries and understanding. Our motivation for this study is into convert researchers' and educators' field data to music, and anyone can create music from natural records.

Attempts to make natural phenomena audible that we know as “sonification” converted from atom trap (Sturm, 2001), brainwaves (Sonifyer.org, 2006), the behavior and structure of molecules (Nakamura, 2012; Mahjour et al., 2023), colors (Harbisson, 2012), and molecular evolution (Martin et al., 2024) to sounds. In the geoscience field, some attempts at “musification” have been made, which involve creating music from geoscience data. The sonification of seismic signals for 2 days was presented in a video 10 days after the 2011 Tohoku Earthquake (Sonifyer.org, 2011). Volcanic geophysical information on seismic movements was collected for sound analysis in eruption dynamics that can be used to predict the future activity of a volcano (Avanzo et al., 2012). Climate data were transformed into acoustic signals, and videos playing the music were distributed to enhance the communication (Planetary Bands, 2015; St. George et al., 2017). By using simulated data instead of observational data, the climate model was musificated from time periods outside the recent past and possible future (de

Formatted: Font: Bold

Deleted:

Deleted: viewing

Deleted: seen

Deleted: to

Moved (insertion) [1]

Deleted:

45 Mora et al., 2020). Observed energy budget data were converted into sound and aligned more closely with traditional music (Nagai, 2024). Terrain relief was translated into acoustic representations for landform quantitative analysis in drainage basins (Lin et al., 2024). These attempts were made by geoscience specialists, and the performance or video of sound was shared with the audience, including non-specialists.

50 To reveal geological strata to human hearing and rhythmic senses, we have developed a web app, Sedimentary Rock! (Kitazawa et al., 2017), which plays music by converting strata thicknesses to note lengths and grain sizes to pitch levels. The web app was developed and made available not only to specialists in geology but also to non-specialists. The access has been approximately 1,000 since we opened the web app in 2017. It is helpful to understand the cyclicity and drastic changes in geological 55 history recorded in strata, as well as the variation in each sedimentary environment. We also surveyed the educational effect of the web app using questionnaires.

2 Sedimentary Rock!

The web app was created with JavaScript and works in web browsers for computers, smartphones, and tablets (Fig. 1a). The procedure when using the web app is to create and edit a single columnar section in the web app and then convert it to music. The web app may create columnar sections without playing music, and descriptions and remarks can be written in a memo box. The user must initially connect to the Risho University server (Kitazawa et al., 2017), but once the web app has been started up, columnar sections can be created, saved, and loaded, and music can be played, even offline, for as long 65 as the browser window is kept open. The web app will be helpful in education and outreach activities because it is easy for anyone to use freely.

3 Creating columnar sections

A columnar section is a geologic illustration that shows a vertical stack of rock types; vertical width 70 represents the thickness, and horizontal width often indicates the grain size. Using the web app, a columnar section is created by entering a single layer's thickness and grain size and adding more layers above or below. After the columnar section is created, layers can be freely selected and edited, and further layers can be inserted at selected positions. Thicknesses can be entered in units of m, cm, or mm, ranging from 1 mm to 100 m. In the rare case of a layer with a thickness of more than 100 m, multiple 75 100-m sections can be individually added. Grain sizes are entered as selections from eight types—“gravel”, “very coarse sand”, “coarse sand”, “medium sand”, “fine sand”, “very fine sand”, “silt”, and “clay”—or “none” (representing a gap or core damage). In a created columnar section, gravel is colored brown, sands are colored yellow, and mud is colored blue. The grain size can be varied between the bottom and top of a layer, which are then interpolated linearly. Thus, grading and reverse grading can 80 be represented. Most grain size variations can be represented by entering separate positions where variation trends change. Other functions include adjusting a columnar section's horizontal width and length to match the screen size or changing its visual impact. Black boundary lines can be displayed at bedding planes, although the default setting does not display these lines.

When information on the thicknesses and grain sizes of a created columnar section is saved, the 85 information is exported in the “Sedimentary Rock Data (SRD)” format that we developed, which is an image of a pattern of colored squares. SRD is an ordinary image file (PNG), which can be saved in an

Moved up [1]: George et al.,

Deleted: Especially in the geoscience field, some attempts at “musification” that make music from geoscience data such as volcano seismograms (Avanzo et al., 2010), earthquakes (Sonifyer.org, 2013), climate data (Planetary Bands, 2015; St.

Deleted: 2017), plasma waves in space (Meredith, 2019), climate model (de Mora et al., 2020), Anthropocene record (Perera, 2020), energy budget (Nagai, 2024), and terrain relief (Lin et al., 2024) are known.[†]

Deleted: very simply

Deleted: !,

Deleted: opened in 2018

Deleted: used.

Deleted: helps anyone

Deleted: or

Deleted: of

Deleted: and

Formatted: English (US)

Deleted: general

Deleted: <http://risho-es.jp/kitazawa/SedimentaryRock/>,

Deleted: Because it is easy for anyone to use freely, it

Deleted: A

Deleted: thickness

Deleted: muds are

110 image folder on a smartphone and sent to other people. A saved columnar section can be loaded into the web app by selecting the corresponding SRD file. A loaded columnar section can continue to be edited and, if saved again, saved as a new SRD file rather than overwriting the old one. As a result, Sedimentary Rock! is not only a musification app but also a helpful tool for making and presenting columnar sections and straightforwardly managing editable sedimentological data.

115 **4 Converting strata to music**

For the conversion of the thickness t (mm) of a layer to the length l (ms) of a note, it is specified that the thicker the layer, the longer the note. In order to handle strata with a wide range of thicknesses, the natural logarithm of the thickness is used, $l = 100\log t$. The thickness of each layer can be from 1 mm to 100 m, corresponding to note lengths from 0 to 1151 ms. The [A minor](#) pentatonic scale is used to convert from grain size to pitch level. It tends not to produce unpleasant sounds because it has no semitone intervals. Eight tones starting from A at 440 Hz are specified such that the smaller the grain size, the higher the pitch. That is, gravel is A, very coarse sand is C, coarse sand is D, medium sand is E, fine sand is G, very fine sand is high A, silt is high C, and clay is high D. “None” is represented by silence. [A graded bed or inverse-graded bed is expressed by glissando \(slides\) from one pitch to another.](#)

When the Play button is pressed, the columnar section, starting from the bottom, is converted to notes and played. This sequence of sounds can be listened to as music (whether it actually “rocks” is another matter), and the strata can be heard. The music would need improvement to sound pleasant, but the basic objective of revealing geological strata to human hearing and rhythmic senses is achieved.

130 We [shared](#) a video on how to use Sedimentary Rocks! and several examples of musification from schematic columnar sections of typical depositional environments of the delta, wave-dominated coast, tidal flat, and submarine fan ([Kitazawa, 2024](#)).

Deleted: t
...

Deleted: $l=100\log t$.

Formatted: Font: Italic



Deleted: C major

Deleted: (C-D-E-G-A-C)

Deleted: The C major pentatonic scale is often used in rock music and other genres; it

Deleted: share


Deleted: <https://www.youtube.com/@SedimentaryRock-e7r/videos>



145 **Figure 1.** (a) Interface of “Sedimentary Rock!”. A columnar section is created and edited by entering a
| single layer’s thickness and grain size. (b) Musification from the Middle Pleistocene tidal deposits. A
| columnar section on the left was created on Sedimentary Rock!, and another was modified from
| Kitazawa and Tateishi (2005).

150 **5 Musification of geological history**

A case study of musification from a columnar section of tidal deposits was done (Fig. 1b). The geological strata data were collected from the Middle Pleistocene Ba Mieu and Thu Duc Formations in southern Vietnam (Kitazawa et al., 2006). The Ba Mieu Formation on this outcrop is a tide-dominated delta succession of the paleo Dong Nai River during the highstand of MIS 5 (Kitazawa, 2007),



160 indicating stacks of two tidal sand bars and being buried by muddy tidal flat deposits (Kitazawa and Tateishi, 2005). Upward convex bar-shaped sediments of Units A and B consist of cross-bedded coarse-grained sediments transported by ebb tidal currents and mud layers (mud drapes) which formed in slack water during high or low tide. The tidal range estimated from around deposits is about 5 m. The music converted from the tidal sand bar deposits is a cyclic melody indicating upward-fining successions due to frequent depositional cycles of development and migration of tidal sand bars. The delta progradation resulted in the shallowing, weakening of tidal current, and deposition of the fine mud in the mud flat.

165 The longer and larger environmental change is converted to a gradually higher tone in the music.

The mud flat deposits are unconformably overlain by mixed flat deposits of the Thu Duc Formation, and the unconformity is amalgamated with a wave/tidal ravinement surface (W/TRS) that is typical in the seaward part of the paleo Dong Nai estuary during a transgression of MIS 5 (Kitazawa, 2007; Kitazawa and Murakoshi, 2016). Thus, there is a remarkable erosion and a loss of previously formed 170 records between the Ba Mieu and Thu Duc Formations. The drastic change in geological history is converted into an abrupt melody change.

Formatted: Font colour: Red

Deleted:

Deleted: a

Formatted: Font colour: Custom Colour (RGB(14,16,26)), English (US)

Formatted: Normal (Web)

185 Table 1. Questionnaire about the educational effects of Sedimentary Rock! (n=29)

Questionary	Strongly Agree (%)	Agree (%)	Neutral (%)	Disagree (%)	Strongly Disagree (%)
Working to make a columnar section deepens your understanding that the strata consist of concepts of grain size and thickness of the stratum.	76	21	3	0	0
The melody of the music deepens your understanding that the stacks of strata represent the geological history.	41	34	10	14	0
Changes in the melody deepen your understanding that grain size and thickness changes indicate environmental changes.	48	31	10	10	0
Adding hearing and rhythmic senses, as well as visual senses such as outcrop pictures or columnar sections, is useful to understand strata.	45	41	7	7	0

Questionary

Working to make a columnar section deepens your understanding that the strata consist of concepts of grain size and thickness of the stratum.

The melody of the music deepens your understanding that the stacks of strata represent the geological history.

Changes in the melody deepen your understanding that grain size and thickness changes indicate environmental changes.

Adding hearing and rhythmic senses, as well as visual senses such as outcrop pictures or columnar sections, is useful to understand strata.

Deleted:

190 Although the concept of Sedimentary Rock! is to reveal geological strata to human hearing, we have no doubt that visual inspection remains the most important way to interpret strata. For university students studying geology, the ideal way to use Sedimentary Rock! is to transcribe their own columnar sections or descriptions from fieldwork. Converting their own columnar sections into music would lead to new discoveries and a deeper understanding.

195 On the other hand, in the science class for younger students, the effort required to create their own columnar section is a bit high. Teachers may need to prepare columnar sections, save them as SRD files, and share them with the students. Even in that case, an experience to make or transcribe a columnar section by themselves is effective in understanding the most essential feature of strata: grain size and thickness.

200 In addition, Sedimentary Rock! has potential applications not only in education but also in outreach and understanding of geology and sedimentology for non-specialists. We also consider the potential benefits to users with disabilities. Although we have never had the opportunity to survey, people with visual impairments could interpret geological strata using the sound of Sedimentary Rock! Thus, we should continue to update the Sedimentary Rock! to enhance operations and sounding. There is room for improvement in the web app interface in the future, for example, by adding thickness and color scales, and making it easier to edit layers. On the musical side, we can also improve the timbre and tempo.

Deleted: new

7 Conclusions

Sedimentary Rock! is a tool that converts columnar sections into human hearing and rhythmic senses. It is easy for anyone to use, and educators can make learners understand that the strata consist of

concepts of grain size and thickness of the stratum by working to make a columnar section.

Furthermore, the learners understand that the stacks of strata represent the geological history, and the changes in grain size and thickness indicate the environmental changes not only with visual senses, such as outcrop pictures or columnar sections, but also in hearing and rhythmic senses. Sedimentary Rock! provides a new, intuitive experience to listen to the music of geological history.

Supplement link. The web app “Sedimentary Rock!” is available at: <http://rissho-es.jp/kitazawa/SedimentaryRock/>

220 The code is available at: <https://github.com/kazu404/SedimentaryRock.git>

Author contribution. All authors contributed to the design of the concepts of the web app “Sedimentary Rock!”. HiroakiY and TakumaK developed the code and created the Sedimentary Rock Data (SRD) format. KazuakiA administers the web app. ToshiyukiK collected and interpreted geological data. ToshiyukiK prepared the manuscript with contributions from all co-authors.

Competing interests. The authors declare that they have no conflict of interest.

Ethical statement. The questionnaires in this article have been reviewed by the Ethics Committee, 230 Institute of Environmental Science, Rissho University, and cleared of ethical problems.

Disclaimer. This article was written and published on volunteered time by the authors.

Acknowledgements. We thank Takuma Noda for his basic study on this paper. We thank cooperators 235 for the questionnaires.

Deleted: in

Review statement. This paper was edited by Sebastian G. Mutz and reviewed by Lee de Mora and an anonymous referee.

Deleted: XXX

240 References

Avanzo, S., Barbera, R., De Mattia, F., Rocca, G. L., Sorrentino, M., and Vicinanza, D.: Data Sonification of Volcano Seismograms and Sound/Timbre Reconstruction of Ancient Musical Instruments with Grid Infrastructures, *Procedia Computer Science*, 1, 397-406.

<https://doi.org/10.1016/j.procs.2010.04.043>, 2012.

Deleted: 2010

245 de Mora, L., Sellar, A. A., Yool, A., Palmieri, J., Smith, R. S., Kuhlbrodt, T., Parker, R. J., Walton, J., Blackford, J. C., and Jones, C. G.: Earth system music: music generated from the United Kingdom Earth System Model (UKESM1). *Geosci. Commun.* 3, 263–278. <https://doi.org/10.5194/gc-3-263-2020>, 2020.

Harbisson, N.: I listen to color: https://www.ted.com/talks/neil_harbisson_i_listen_to_color/transcript, 250 (last access: 9 September 2024), 2012.

Kitazawa, T.: Pleistocene macrotidal tide-dominated estuary-delta succession, along the Dong Nai River, southern Vietnam. *Sedimentary Geol.* 194, 115-140. <https://doi.org/10.1016/j.sedgeo.2006.05.016>, 2007.

Formatted: English (UK)

- Kitazawa, T.: [Sedimentary Rock! GEOLOGY into MUSIC: https://www.youtube.com/@SedimentaryRock-e7r/videos](https://www.youtube.com/@SedimentaryRock-e7r/videos), (last access: 13 September 2025), 2024.
- 260 Kitazawa, T., Aoki, K., Yasutomo, H., Kato, T.: [Sedimentary Rock!: http://risshe-es.jp/kitazawa/SedimentaryRock/](http://risshe-es.jp/kitazawa/SedimentaryRock/), (last access: 13 September 2025), 2017.
- Kitazawa, T. and Murakoshi, N.: Tidal ravinement surfaces in the Pleistocene macrotidal tide-dominated Dong Nai estuary, southern Vietnam. Contributions to Modern and Ancient Tidal Sedimentology, IAS Special Publication 47, 233–241. <https://doi.org/10.1002/9781119218395.ch13>, 2016.
- 265 Kitazawa, T. and Tateishi, M.: Geometry and preservation process of tidal sand bar deposits of Middle Pleistocene macrotidal tide-dominated delta succession, southern Vietnam. Journal of Sedimentological Society of Japan 61, 27–38. <https://doi.org/10.4096/jssj1995.61.27>, 2005.
- 270 Kitazawa, T., Nakagawa, T., Hashimoto, T., and Tateishi, M.: Stratigraphy and optically stimulated luminescence (OSL) dating of a Quaternary sequence along the Dong Nai River, southern Vietnam. J. Asian Earth Sci. 27, 788–804. <https://doi.org/10.1016/j.jseas.2005.08.001>, 2006.
- Lin, S., Yu, Y., Chen, N., Shen, R., and Wang, X.: Using music and music-based analysis to model and to classify terrain in geomorphology, Earth Surf. Process. Landforms, 49, 1544–1559. <https://doi.org/10.1002/esp.5785>, 2024.
- 275 Mahjour, B., Bench, J., Zhang, R., Frazier, J., and Cernak, T.: Molecular sonification for molecule to music information transfer, Digital Discovery, 2, 520–530. <https://doi.org/10.1039/d3dd00008g>, 2023.
- Martin, E.J., Knotts, S., Phillips, M., Weise, N., Meagher, T.R., and Barker, D.: Innovative communication of molecular evolution through sound: a biological sonification concert. Evo. Edu. Outreach, 17, 6, <https://doi.org/10.1186/s12052-024-00200-5>, 2024.
- 280 Nagai, H.: [String Quartet No. 1 “Polar Energy Budget” – Music composition using Earth observation data of polar regions](#), iScience, 27, 109622. <https://doi.org/10.1016/j.isci.2024.109622>, 2024.
- Nakamura, S.: The sound of chemistry: Converting molecular dynamics information to sounds!?, Gendai Kagaku (Chemistry Today), 5, 39–41, 2012.
- 285 Planetary Bands: Worming World: <https://planetbands.mystrikingly.com>, (last access: 9 September 2024), 2015.
- Sonifyer.org: [Sonification of Tohoku Earthquake / Sendai Coast, Japan, 2011/03/11](#): <https://www.youtube.com/watch?v=3PJxUPVz9Oo>, (last access: 17 September 2025), 2011.
- 290 St. George, S., Crawford, D., Reubold, T., and Giorgi, E.: Making Climate Data Sing: Using Music-like Sonifications to Convey a Key Climate Record. Bull. Am. Meteorol. Soc. 98, 23–27. <https://doi.org/10.1175/BAMS-D-15-00223.1>, 2017.
- Sturm, B. L.: Composing for an Ensemble of Atoms: The Metamorphosis of Scientific Experiment into Music. Organised Sound 6 (2), 131–145, <https://doi.org/10.1017/S1355771801002102>, 2001.

Deleted: Meredith, N.P.: Turning the sounds of space into art, Astron. Geophys., 60, 2.18–2.21. <https://doi.org/10.1093/astrogeo/atz097>, 2019. ↗

Deleted: Perera, J.: Anthropocene in C Major: <https://www.jamiperera.com/climate-data-sonification>, (last access: 9 September 2024), 2020. ↗

Deleted: sonifyer.org/sound/eeg/

Deleted: 9

Deleted: 2024, 2006. ↗
Sonifyer.org: <https://www.sonifyer.org/sound/erdbewegung/>, (last access: 9 September 2024), 2013